Galilei invariant theories. I. Constructions of indecomposable finite-dimensional representations of the homogeneous Galilei group: directly and via contractions¹

M. de Montigny^{a,b}, J. Niederle^c, and A.G. Nikitin^d

^aCampus Saint-Jean, University of Alberta, 8406 - 91 Street, Edmonton, Alberta, Canada T6C 4G9

^bTheoretical Physics Institute, University of Alberta, Edmonton, Alberta, Canada T6G 2J1

^cInstitute of Physics of the Academy of Sciences of the Czech Republic, Na Slovance 2, 18221 Prague 8, Czech Republic

^dInstitute of Mathematics, National Academy of Sciences of Ukraine, 3 Tereshchenkivs'ka Street, Kyiv-4, Ukraine, 01601

Abstract

All indecomposable finite-dimensional representations of the homogeneous Galilei group which when restricted to the rotation subgroup are decomposed to spin 0, 1/2 and 1 representations are constructed and classified. These representations are also obtained via contractions of the corresponding representations of the Lorentz group. Finally the obtained representations are used to derive a general Pauli anomalous interaction term and Darwin and spin-orbit couplings of a Galilean particle interacting with an external electric field.

¹E-mail: montigny@phys.ualberta.ca, niederle@fzu.cz, nikitin@imath.kiev.ua

1 Introduction

Unlike what most physicists might think of, the mathematical structure of the representations of the Galilei group is in many respects more complex and sophisticated than that of their relativistic counterparts. This is perhaps also the reason why unitary irreducible representations of the Poincaré group – the symmetry group of special theory of relativity – has been known nearly twenty years earlier than those of the Galilei group, even though the Galilei principle of relativity was discovered several centuries prior to the Einstein one.

The Galilei group and its representations are described in the Lévy-Leblond masterful survey [1] written almost thirty five years ago. They form the group-theoretical basis for description of various physical predictions in non-relativistic classical mechanics and electrodynamics and in non-relativistic quantum mechanics as well (see also a more recent review [2]). These predictions are generally more than just some approximations to the relativistic results since only comparison of the predictions based on the Galilei group and its representations with those on the Poincaré group can indicate which predictions have the origin in the non-relativistic and which in the relativistic one. For instance, it was shown in [3], [4] and by Lévy-Leblond in [1], [5], [6] that the concept of spin of particles and of magnetic moments of particles have the origin already in the Galilean non-relativistic quantum mechanics and not as stated in many textbooks as a consequence of relativistic effects.

In the Galilei invariant framework we shall consider first free particles and then particles interacting with external fields. The free particles can be described either by representations of the Galilei group (irreducible in the case of elementary particles and indecomposable for particles with internal structures) or equivalently, by Galilean invariant wave equations. For interacting particles the wave equations are more appropriate since they allow to introduce interactions.

There exist three approaches how to formulate the Galilei invariant wave equations. The first is based on the fact that the Galilei group and the Galilei invariant equations can be obtained from the Poincaré group and Poincaré invariant equations respectively by a limiting procedure – the so-called Inonü-Wigner contraction [7].

The second approach consists of writing first the Lorentz-invariant equations in the (4 + 1)-dimensional space-time, and then projecting them down to the (3 + 1)dimensional Newtonian space-time using the fact that the extended Galilei algebra in (3 + 1) dimensions is a subalgebra of the Poincaré algebra in (4 + 1) dimensions, for connection of representations of these algebras see [8]. This approach, referred here as a projecting one, has been developed in [9]-[11] (see also [12]).

The third way to construct Galilei invariant theories consists in searching for this theories directly using the requirement of Galilei invariance and knowledge of representations of the Galilei group. We shall show in this paper that, in many respects, this latest approach is the most powerful and comprehensive. Moreover, it allows us to construct such consistent Galilei-invariant equations of motion which are very difficult to derive using the contraction or projection methods. On the other hand, the direct search for wave equations invariant with respect to the inhomogeneous Galilei group presupposes a good knowledge of the indecomposable finite-dimensional representations of its homogeneous Galilei subgroup, which has not been available till now.

Our article serves the following four aims: (1) to describe all indecomposable finite-dimensional representations of the homogeneous Galilei group that are defined on spinor, scalar and vector representation spaces; (2) to specify all Galilei invariant bilinear forms, which facilitate derivation of various non-linear Galilei wave equations; (3) to specify the reducible representations of the Lorentz group which lead to the found representations of the Galilei group via contractions; (4) to determine the Galilean spinorial wave equations which include the Pauli anomalous terms.

In the next section we define indecomposable finite-dimensional representations of the homogeneous Galilei group in general.

In Sections 3 and 4 some of these representations are constructed explicitly, namely all those which when restricted to representations of the rotation subgroup of the homogeneous Galilei group are decomposed to spin 0, 1/2 and 1 representations. In addition we present here also the complete list of bilinear forms invariant with respect to all found representations of the Galilei group. Section 5 contains various examples of the Galilean vectors. In Section 6 we obtain the previously found representations of the homogeneous Galilei group via the Inönü-Wigner contraction of the corresponding representations of the Lorentz group. In Section 7 the found representations are used to deduce the most general form of the Pauli interaction term which can be added to the Galilei invariant equation for spinors. Subsection 7.3 presents a simple Galilean system which describes the Darwin and spin-orbital interactions of particles with an external electric field. Finally, Section 8 is devoted to discussions of the obtained results.

2 Definitions and properties of the Galilei group and its Lie algebra

The Galilei group G(1,3) consists of the following transformations of time variable tand of space variables $\mathbf{x} = (x_1, x_2, x_3)$:

$$\begin{aligned} t \to t' &= t + a, \\ \mathbf{x} \to \mathbf{x}' &= \mathbf{R}\mathbf{x} + \mathbf{v}t + \mathbf{b} \end{aligned} \tag{1}$$

where a, **b** and **v** are real parameters of time translation, space translations and pure Galilei transformations respectively, and matrix **R** specifies rotations determined by the three parameters $\theta_1, \theta_2, \theta_3$.

The Galilei group includes a subgroup leaving invariant a point $\mathbf{x} = (0, 0, 0)$ at time t = 0. It is formed by all space rotations and pure Galilei transformations, i.e.,

by transformations (1) with $a = \mathbf{b} \equiv 0$. This subgroup is said to be the homogeneous Galilei group HG(1,3). It is a semi-direct product of the three-parameter commutative group of pure Galilei transformations with the rotation group. Thus this group is not compact and does not have unitary finite-dimensional representations.

The Galilei group G(1,3) is a semidirect product of its Abelian subgroup generated by time and space translations with the homogeneous Galilei group HG(1,3).

Unitary representations of Galilei group which are ordinary ones were described by Inönü and Wigner [3] in 1952, whereas those which are ray by Bargmann [4] in 1954. A nice review of these representations can be found in [1], see also [13].

However, a decisive role in the description of various finite-component Galilei invariant equations is played by finite-dimensional representations of the homogeneous Galilei group HG(1,3). They where studied first apparently in paper [14] but have not been classified till now.

Let us remind that the representations of HG(1,3) induce ray representations of the Galilei group G(1,3) as well as ordinary representations of the extended Galilei group $G_m(1,3)$ which is a central extension of the Galilei group via a one-parameter subgroup. Both of them are realized in the space of (square integrable) *n*-component functions $\Psi(t, \mathbf{x})$ which for any transformation (1) cotransform in the following way [1]

$$\Psi(\mathbf{x},t) \to \Psi'(\mathbf{x}',t') = e^{\mathrm{i}f(\mathbf{x},t)}T\Psi(\mathbf{x},t)$$
(2)

where $\Psi(\mathbf{x}, t) = \text{column}(\Psi_1(\mathbf{x}, t), \Psi_2(\mathbf{x}, t), \cdots, \Psi_n(\mathbf{x}, t))$ are *n*-component vectors from the representation space, T are $n \times n$ numerical matrices depending on transformation parameters \mathbf{v} and $\boldsymbol{\theta} = (\theta_1, \theta_2, \theta_3)$,

$$f(\mathbf{x},t) = m\left(\mathbf{v}\cdot\mathbf{x} + \frac{v^2}{2}t + c\right)$$

is a phase which includes two parameters: m and c. For m = 0 the central extension is trivial and representations (2) are ordinary representations of G(1,3). For m different from zero the central extension is nontrivial, and transformations (2) realize ray representations of the Galilei group G(1,3) and ordinary representations of $G_m(1,3)$. Moreover, the transformation matrices T realize a finite-dimensional representations of the homogeneous Galilei group HG(1,3).

Let us mention that realizations (2) are precisely those which are used in quantum mechanics and quantum field theory. These realizations are also essential to construct wave equations invariant w.r.t. the Galilei group [12].

Taking expressions (2) corresponding to the infinitesimal transformations (1) and treating c as an additional transformation parameter we can calculate the 11-dimensional Lie algebra of the extended Galilei group. Basis elements of this algebra are of the following form

$$P_{0} = i\partial_{0}, \quad P_{a} = -i\partial_{a}, \quad M = Im$$

$$J_{a} = -i\varepsilon_{abc}x_{b}\partial_{c} + S_{a},$$

$$K_{a} = -ix_{0}\partial_{a} - mx_{a} + \eta_{a},$$
(3)

where indices a, b and c run over the values 1, 2, 3, I is the unit matrix, and S_a and η_a are matrices which satisfy the following commutation relations:

$$[S_a, S_b] = i\varepsilon_{abc}S_c,\tag{4}$$

$$[\eta_a, S_b] = i\varepsilon_{abc}\eta_c,\tag{5}$$

$$[\eta_a, \eta_b] = 0, \tag{6}$$

that is, they form a basis of the Lie algebra hg(1,3) of the homogeneous Galilei group HG(1,3). This algebra is isomorphic to the Lie algebra hg(1,3) of the Euclidean group.

Conditions (4) - (6) are necessary and sufficient in order generators (3) form a basis of a Lie algebra namely the extended Galilei algebra satisfying the following relations:

$$\begin{bmatrix}
[J_a, J_b] &= i\epsilon_{abc}J_c, & [J_a, K_b] &= i\epsilon_{abc}K_c, \\
[J_a, P_b] &= i\epsilon_{abc}P_c, & [K_a, P_0] &= iP_a, \\
[K_a, P_b] &= i\delta_{ab}M, & [K_a, K_b] &= 0, \\
[P_a, P_b] &= 0, & [P_0, P_a] &= 0, \\
[M, P_0] &= [M, P_a] &= & [M, J_a] &= [M, K_a] &= 0.
\end{bmatrix}$$
(7)

Unfortunately, the problem of a complete classification of non-equivalent finitedimensional realizations of algebra (4)-(6) appears to be an unsolvable 'wild' algebraic problem. However we shall show that this problem can be completely solved in the two important particular cases: for the purely spinor representations and vector representations.

3 Spinor representations

The Lie algebra hg(1,3), defined by relations (4)-(6), includes the subalgebra so(3) spanned on the basis elements S_1, S_2 and S_3 . Without loss of generality we suppose that representations of this subalgebra are Hermitian and completely reducible and shall search for representations of hg(1,3) in the so(3) basis in which the Casimir operator of so(3) is diagonal.

Irreducible representations of so(3) are labelled by integers or half-integers s. Let \tilde{s} be the highest value of s which appears in the decomposition of a reducible representation of so(3) subduced by the indecomposable representation of hg(1,3). We shall call the related carrier space of this representation of hg(1,3) the representation space of spin \tilde{s} .

Consider finite-dimensional indecomposable representations of algebra hg(1,3) of spin $\tilde{s} = \frac{1}{2}$. Then the corresponding matrices S_1, S_2 and S_3 can be decomposed to a direct sum of the irreducible representations D(1/2) of algebra so(3):

$$S_a = \frac{1}{2} I_{n \times n} \otimes \sigma_a,\tag{8}$$

where $I_{n \times n}$ is the $n \times n$ unit matrix with a finite n and σ_a are the usual Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

From (8) and (5), the generic form of the related matrices η_a is:

$$\eta_a = A_{n \times n} \otimes \sigma_a,\tag{9}$$

where $A_{n \times n}$ is an $n \times n$ matrix. The commutativity of matrices (9) leads to the nilpotency condition

$$A_{n\times n}^2 = 0. (10)$$

Thus, without any loss of generality, $A_{n \times n}$ may be expressed as a direct sum of 2×2 Jordan cells and zero matrix.

Because of (10) there exist only two different indecomposable representations of algebra hg(1,3) defined on spin 1/2 carrier space:

$$S_a = \frac{1}{2}\sigma_a, \qquad \eta_a = 0,$$

and

$$S_a = \frac{1}{2} \begin{pmatrix} \sigma_a & 0\\ 0 & \sigma_a \end{pmatrix}, \qquad \eta_a = \frac{1}{2} \begin{pmatrix} 0 & 0\\ \sigma_a & 0 \end{pmatrix}.$$
(11)

The corresponding vectors from the representation space are two-component spinors $\varphi(\mathbf{x}, t)$, as well as four-component bispinors $\Psi = \begin{pmatrix} \varphi_1(\mathbf{x}, t) \\ \varphi_2(\mathbf{x}, t) \end{pmatrix}$ with two-component φ_1 and φ_2 , respectively. When t and **x** undergo a Galilean transformation (1), then φ cotransforms as

$$\varphi(\mathbf{x},t) \to \varphi'(\mathbf{x}',t') = e^{\mathrm{i}m\left(\mathbf{v}\cdot\mathbf{x}+\mathbf{v}^2t/2+c\right)} \left(\cos\frac{\theta}{2} + i\frac{\boldsymbol{\sigma}\cdot\boldsymbol{\theta}}{\theta}\sin\frac{\theta}{2}\right)\varphi(\mathbf{x},t)$$

while the transformation law for components of Galilean bispinor is

$$\begin{split} \varphi_1(\mathbf{x},t) &\to \varphi_1'(\mathbf{x}',t') = e^{\mathrm{i}m\left(\mathbf{v}\cdot\mathbf{x}+\mathbf{v}^2t/2+c\right)} \left(\cos\frac{\theta}{2} + i\frac{\boldsymbol{\sigma}\cdot\boldsymbol{\theta}}{\theta}\sin\frac{\theta}{2}\right)\varphi_1(\mathbf{x},t), \\ \varphi_2(\mathbf{x},t) &\to \varphi_2'(\mathbf{x}',t') = e^{\mathrm{i}m\left(\mathbf{v}\cdot\mathbf{x}+\mathbf{v}^2t/2+c\right)} \left(\left(\cos\frac{\theta}{2} + i\frac{\boldsymbol{\sigma}\cdot\boldsymbol{\theta}}{\theta}\sin\frac{\theta}{2}\right)\varphi_2(\mathbf{x},t) \right. \\ &+ \left(\frac{\mathrm{i}}{2}\boldsymbol{\sigma} \cdot \mathbf{v}\cos\frac{\theta}{2} - \left(\boldsymbol{\theta}\cdot\mathbf{v} + i\boldsymbol{\sigma}\cdot\boldsymbol{\theta}\times\mathbf{v}\right)\frac{\sin\frac{\theta}{2}}{\theta}\right)\varphi_1(\mathbf{x},t)\right). \end{split}$$

We use the notation $\theta = \sqrt{\theta_1^2 + \theta_2^2 + \theta_3^2}$. Invariants of these transformations which are independent on t and **x** are arbitrary functions of $\varphi^{\dagger}\varphi$, $\varphi_1^{\dagger}\varphi_1$ and $\varphi_1^{\dagger}\varphi_2 + \varphi_2^{\dagger}\varphi_1$.

4 Vector representations

In this section, we examine the finite-dimensional indecomposable representations of the algebra hg(1,3) defined on vector, or spin-one, representation spaces. The corresponding matrices S_a can be expressed as direct sums of spin-one and spin-zero matrices:

$$S_a = \begin{pmatrix} I_{n \times n} \otimes s_a & \cdot \\ \cdot & \mathbf{0}_{m \times m} \end{pmatrix}.$$
(12)

The symbols $I_{n \times n}$ and $\mathbf{0}_{m \times m}$ denote the $n \times n$ unit matrix and $m \times m$ zero matrix respectively, s_a (a = 1, 2, 3) are 3×3 matrices of spin equal to one for which we choose the following realization:

$$s_{1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad s_{2} = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad s_{3} = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$
(13)

The general form of matrices η_a which satisfy relations (5) with matrices (12) is given by the following formulae (see, e.g., [15]):

$$\eta_a = \begin{pmatrix} A \otimes s_a & B \otimes k_a^{\dagger} \\ C \otimes k_a & \mathbf{0}_{m \times m} \end{pmatrix}$$
(14)

A, B and C are matrices of dimension $n \times n$, $n \times m$ and $m \times n$ respectively, k_a are 1×3 matrices of the form

$$k_1 = (i, 0, 0), \qquad k_2 = (0, i, 0), \qquad k_3 = (0, 0, i).$$
 (15)

The matrices (12) and (14) satisfy conditions (5) with any A, B and C. Substituting (14) into (6) and using the relations

$$s_a k_b^{\dagger} = i \varepsilon_{abc} k_c^{\dagger}, \ k_a s_b = i \varepsilon_{abc} k_c,$$

$$[s_a, s_b] = k_a^{\dagger} k_b - k_b^{\dagger} k_a = i \varepsilon_{abc} s_c,$$

$$k_a k_b^{\dagger} - k_b k_a^{\dagger} = 0$$

we obtain the following equations for matrices A, B and C:

$$A^2 + BC = 0, (16)$$

$$CA = 0, \quad AB = 0. \tag{17}$$

The solution of the matrix problem defined by equations (16) and (17) is relatively easily to handle. Namely, if we multiply equation (16) by A and use equation (17), then we obtain the condition $A^3 = 0$ so that A is a nilpotent matrix whose nilpotency index N satisfies the condition $N \leq 3$. This implies that BC is a nilpotent matrix with index of nilpotency equal to 2. Thus A can be represented as a direct sum of the Jordan cells of dimension 3, 2 and zero matrices. We can prove that in order to obtain indecomposable representations of algebra (4), (6) it is necessary to restrict ourselves to the case of the indecomposable matrices A, and so there are three possibilities

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \ A = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \ A = 0.$$
(18)

We can prove that for indecomposable matrices η dimension m and dimension n satisfy the condition $-1 \leq (n-m) \leq 2$, $n \leq 3$. Matrices η_a are defined up to a constant multiplier, which we fix imposing the condition

$$\operatorname{Trace}(\eta_{a}^{\dagger}\eta_{a}) \equiv \operatorname{Trace}(\eta_{1}^{\dagger}\eta_{1} + \eta_{2}^{\dagger}\eta_{2} + \eta_{3}^{\dagger}\eta_{3}) = 6(n-1) + 3(m+\delta_{3n}).$$
(19)

By going over all admissible values of n, m and using relations (18) we find that up to equivalence transformations there exist ten solutions of equations (17), which can be labelled by triplets of numbers n, m, λ where $\lambda = \frac{1}{3} \operatorname{Trace} \left((S_a S_a - 2) \eta_a^{\dagger} \eta_a \right)$ takes the values

$$\lambda = \begin{cases} 0 \text{ if } m = 0, \\ 1 \text{ if } m = 2 \text{ or } n - m = 2, \\ 0, 1 \text{ if } m = 1, n \neq 3. \end{cases}$$

We shall not present them here but write directly the corresponding matrices S_a (12) and η_a (14), which are collected in Table 1.

Any set of matrices S_a , η_a given in Table 1 yields a finite-dimensional indecomposable representation of algebra hg(1,3) which generates a representation of the extended Galilei algebra via (7). From our analysis, there exist ten indecomposable vector representations of hg(1,3).

The finite-transformations corresponding to these realizations are found by integrating the Lie equations for generators given in (7) and Table 1. For a specific representation $D(n, m, \lambda)$ they have the following forms.

• D(0,0,0): The related representation space is formed by a field ψ invariant under rotations and which transforms under a Galilean boost as

$$\psi \to e^{\mathrm{i}mf}\psi,$$
 (20)

where $f = \mathbf{v} \cdot \mathbf{x} + \mathbf{v}^2 t/2$. Such transformations keep invariant the bilinear form $I_1 = \psi^* \psi$.

• D(1,0,0): The Galilean 3-vectors $\mathbf{R} = \operatorname{column}(R_1, R_2, R_3)$ transforms as a vectors under rotations:

$$\mathbf{R} \to \mathbf{R}\cos\theta + \frac{\boldsymbol{\theta} \times \mathbf{R}}{R}\sin\theta + \frac{\boldsymbol{\theta}(\boldsymbol{\theta} \cdot \mathbf{R})}{\theta^2}(1 - \cos\theta),$$

$\begin{array}{c} \text{Representation} \\ D(n,m,\lambda) \end{array}$	S_a	η_a	
D(0,0,0)	0	0	
D(1,0,0)	s_a	$0_{3 imes 3}$	
D(1, 1, 0)	$\left(\begin{array}{cc} s_a & 0_{3\times 1} \\ 0_{1\times 3} & 0 \end{array}\right)$	$\left(\begin{array}{cc} 0_{3\times3} & 0_{3\times1} \\ k_a & 0 \end{array}\right)$	
D(1, 1, 1)	$\left(\begin{array}{cc} s_a & 0_{3\times 1} \\ 0_{1\times 3} & 0 \end{array}\right)$	$\left(egin{array}{ccc} 0_{3 imes 3} & k_a^\dagger \ 0_{1 imes 3} & 0 \end{array} ight)$	
D(1, 2, 1)	$\begin{pmatrix} s_a & 0_{1\times 3} & 0_{1\times 3} \\ 0_{3\times 1} & 0 & 0 \\ 0_{3\times 1} & 0 & 0 \end{pmatrix}$	$\left(\begin{array}{cccc} 0_{3\times 3} & k_a^{\dagger} & 0_{3\times 1} \\ 0_{1\times 3} & 0 & 0 \\ k_a & 0 & 0 \end{array}\right)$	
D(2, 0, 0)	$\left(egin{array}{cc} s_a & 0_{3 imes 3} \ 0_{3 imes 3} & s_a \end{array} ight)$	$\left(egin{array}{ccc} 0_{3 imes 3} & 0_{3 imes 3} \ s_a & 0_{3 imes 3} \end{array} ight)$	
D(2, 1, 0)	$ \begin{pmatrix} s_a & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & s_a & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & 0 \end{pmatrix} $	$ \begin{pmatrix} 0_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ s_a & 0_{3\times3} & 0_{3\times1} \\ k_a & 0_{1\times3} & 0 \end{pmatrix} $	
D(2, 1, 1)	$ \begin{pmatrix} s_a & 0_{3\times 3} & 0_{3\times 1} \\ 0_{3\times 3} & s_a & 0_{3\times 1} \\ 0_{1\times 3} & 0_{1\times 3} & 0 \end{pmatrix} $	$\left(\begin{array}{cccc} 0_{3\times3} & s_a & k_a^{\dagger} \\ 0_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & 0 \end{array}\right)$	
D(2, 2, 1)	$\begin{pmatrix} s_a & 0_{3\times 3} & 0_{3\times 1} & 0_{3\times 1} \\ 0_{3\times 3} & s_a & 0_{3\times 1} & 0_{3\times 1} \\ 0_{1\times 3} & 0_{1\times 3} & 0 & 0 \\ 0_{1\times 3} & 0_{1\times 3} & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0_{3\times3} & 0_{3\times3} & 0_{3\times1} & 0_{3\times1} \\ s_a & 0_{3\times3} & k_a^{\dagger} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & 0 & 0 \\ k_a & 0_{1\times3} & 0 & 0 \end{pmatrix}$	
D(3, 1, 1)	$\begin{pmatrix} s_a & 0_{3\times 3} & 0_{3\times 3} & 0_{3\times 1} \\ 0_{3\times 3} & s_a & 0_{3\times 3} & 0_{3\times 1} \\ 0_{3\times 3} & 0_{3\times 3} & s_a & 0_{3\times 1} \\ 0_{1\times 3} & 0_{1\times 3} & 0_{1\times 3} & 0 \end{pmatrix}$	$\begin{pmatrix} 0_{3\times3} & 0_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ s_a & 0_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & s_a & 0_{3\times3} & k_a^{\dagger} \\ -k_a & 0_{1\times3} & 0_{1\times3} & 0 \end{pmatrix}$	

Table 1: Vector representations: spin matrices S_a and boost matrices η_a where s_a and k_a are matrices (13) .

and via

$$\mathbf{R} \to e^{\mathrm{i}mf} \mathbf{R},$$
 (21)

under the Galilei boosts. Then, the bilinear form $I_2 = \mathbf{R}^* \cdot \mathbf{R}$ is invariant under Galilean transformations.

• D(1, 1, 0): Galilean 4-vector $\Psi_4 = (\mathbf{U}, \psi)$, where **U** is a vector and ψ is a scalar under the rotation transformations. The action of Galilean boosts on **U** and ψ can be written as

$$\begin{aligned} \mathbf{U} &\to e^{\mathrm{i}mf} (\mathbf{U} + \mathbf{v}\psi), \\ \psi &\to e^{\mathrm{i}mf}\psi. \end{aligned}$$
(22)

The corresponding invariant of Galilei group is of the form $I_1 = \psi^* \psi$.

• D(1, 1, 1): The second Galilean 4-vector $\tilde{\Psi}_4 = (\mathbf{R}, B)$, where \mathbf{R} and B transform under the Galilei boost as:

The invariant form for these (and rotation) transformations can be written as $I_2 = \mathbf{R}^* \cdot \mathbf{R}$.

D(1,2,1): The Galilean 5-vector Ψ₅ = (ψ, U, C), where ψ and C are scalars and U is a vector with respect to rotations. Under a Galilean boost, they transform as:

$$\begin{split} \psi &\to e^{\mathrm{i}mf}\psi, \\ \mathbf{U} &\to e^{\mathrm{i}mf}(\mathbf{U} + \mathbf{v}\psi), \\ C &\to e^{\mathrm{i}mf}(C + \mathbf{v} \cdot \mathbf{U} + \frac{1}{2}\mathbf{v}^{2}\psi), \end{split}$$
(24)

and the invariants of these transformations are $I_2 = \psi^* \psi$ and $I_3 = \psi^* C + \psi C^* - \mathbf{U}^* \mathbf{U}$.

• D(2,0,0): The Galilean 6-vectors (bi-vectors) $\Psi_6 = (\mathbf{R}, \mathbf{W})$, which under the Galilei boosts transform as:

$$\begin{aligned} \mathbf{R} &\to e^{\mathrm{i}mf} \mathbf{R}, \\ \mathbf{W} &\to e^{\mathrm{i}mf} (\mathbf{W} + \mathbf{v} \times \mathbf{R}), \end{aligned}$$
(25)

and the corresponding invariants are $I_2 = \mathbf{R}^* \cdot \mathbf{R}$ and $I_4 = \mathbf{R}^* \cdot \mathbf{W} + \mathbf{R} \cdot \mathbf{W}^*$.

• D(2, 1, 0): The Galilean 7-vectors $\Psi_7 = (\mathbf{R}, \mathbf{W}, B)$. Under the Galilei boost its components transform according to

$$\mathbf{R} \to e^{imf} \mathbf{R},
 B \to e^{imf} (B + \mathbf{v} \cdot \mathbf{R}),
 \mathbf{W} \to e^{imf} (\mathbf{W} + \mathbf{v} \times \mathbf{R}).$$
(26)

The corresponding invariants are the same as the previous case: $I_2 = \mathbf{R}^* \cdot \mathbf{R}$ and $I_4 = \mathbf{R}^* \cdot \mathbf{W} + \mathbf{R} \cdot \mathbf{W}^*$. • D(2,1,1): The second 7-vector $\tilde{\Psi}_7 = (\mathbf{K}, \mathbf{R}, \psi)$ with \mathbf{K} being a 3-vector which under the Galilei boost transforms as

$$\mathbf{K} \to e^{\mathrm{i}mf} \left(\mathbf{K} + \mathbf{v} \times \mathbf{R} + \mathbf{v} \psi \right), \tag{27}$$

and with \mathbf{R} , ψ which transform as:

$$\psi \to e^{\mathrm{i}mf}\psi, \ \mathbf{R} \to e^{\mathrm{i}mf}\mathbf{R}.$$
 (28)

The corresponding invariants are $I_1 = \psi^* \psi$ and $I_2 = \mathbf{R}^* \cdot \mathbf{R}$.

• D(2,2,1): Galilean 8-vector $\Psi_8 = (\mathbf{R}, \mathbf{K}, B, \psi)$, whose components with the Galilei boost transform as:

$$\psi \to e^{imf}\psi, \ \mathbf{R} \to e^{imf}\mathbf{R}, \mathbf{K} \to e^{imf}\left(\mathbf{K} + \mathbf{v} \times \mathbf{R} + \mathbf{v}\psi\right), B \to e^{imf}\left(B + \mathbf{v} \cdot \mathbf{R}\right).$$
(29)

The invariants of these transformations are $I_1 = \psi^* \psi$, $I_2 = \mathbf{R}^* \cdot \mathbf{R}$ and $I_5 = \psi^* B + \psi B^* - \mathbf{K} \cdot \mathbf{R}^* - \mathbf{K}^* \cdot \mathbf{R}$.

• D(3, 1, 2): The ten-vector fields $\Psi_{10} = (\mathbf{R}, \mathbf{W}, \mathbf{N}, B)$ combine three 3-vectors $\mathbf{R}, \mathbf{W}, \mathbf{N}$ and one scalar B. They cotransform under the boost transformations via

$$\mathbf{R} \to e^{\mathrm{i}m_{f}} \mathbf{R}, \\
\mathbf{W} \to e^{\mathrm{i}m_{f}} (\mathbf{W} + \mathbf{v} \times \mathbf{R}), \\
\mathbf{N} \to e^{\mathrm{i}m_{f}} \left(\mathbf{N} + \mathbf{v} \times \mathbf{W} + \mathbf{v}B + \mathbf{v}(\mathbf{v} \cdot \mathbf{R}) - \frac{1}{2}\mathbf{v}^{2}\mathbf{R} \right), \\
B \to e^{\mathrm{i}m_{f}} (B + \mathbf{v} \cdot \mathbf{R}).$$
(30)

The invariants of these transformations are $I_2 = \mathbf{R}^* \cdot \mathbf{R}$, $I_4 = \mathbf{R}^* \cdot \mathbf{W} + \mathbf{R} \cdot \mathbf{W}^*$ and $I_6 = B^*B + \mathbf{W} \cdot \mathbf{W}^* - \mathbf{R} \cdot \mathbf{N}^* - \mathbf{N} \cdot \mathbf{R}^*$. In Section 7, we shall utilize this vector in the Galilean electromagnetism.

Thus in addition to Galilean scalar A there exist nine Galilean vectors enumerated in the above items. We see that the number of such vectors is notably larger than in the case of the proper Lorentz group, when there are only three irreducible multiplets whose components transform as vectors or scalars under rotations, namely, a fourvector and self-dual and anti self-dual components of an antisymmetric tensor [16].

5 Examples of Galilean vectors

In the previous section, we have described finite-dimensional indecomposable Galilean vectors and present explicitly their group transformations and invariants. They form the main tool for constructing various Galilei invariant models. In particular, by using the realizations of matrices S_a and η_a given in Table 1, it is possible to describe all non-equivalent Galilei invariant first order partial differential equations for vector fields. Here we present some important examples of Galilean vectors.

<u>Example 1</u>. Generators P_0 and P_a of time and space translations defined by relations (7) and mass m form a Galilean 5-vector of type Ψ_5 provided we identify $P_0 \sim C, \mathbf{P} \sim \mathbf{U}, m \sim \psi$. For m = 0 this 5-vector is reduced to a 4-vector of type $\tilde{\Psi}_4$.

<u>Example 2</u>. Five-potential of Galilean electromagnetic field [17] $\hat{A} = (A_0, \mathbf{A}, A_4)$ with transformation law

$$\mathbf{A}' = \mathbf{A} + \mathbf{v}A_4,$$

$$A'_0 = A_0 + \mathbf{v} \cdot \mathbf{A} + \frac{1}{2}\mathbf{v}^2 A_4,$$

$$A'_4 = A_4$$
(31)

is an example of Galilean 5-vector field with zero mass, which is a carrier space of the representation D_5 described in Table 1.

<u>Example 3.</u> The field strength tensor of Galilean electromagnetic field

$$F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu} \tag{32}$$

where $\mu, \nu = 0, 1, 2, 3, 4$ and by definition $\partial_4 A_{\mu} = 0$, is the example of massless 10-vector which transforms in accordance with (30). The explicit relation between components of $F_{\mu\nu}$ and Ψ_{10} is given by the following formula:

$$F_{\mu\nu} = \begin{pmatrix} 0 & -N_1 & -N_2 & -N_3 & B \\ N_1 & 0 & W_3 & -W_2 & R_1 \\ N_2 & -W_3 & 0 & W_1 & R_2 \\ N_3 & W_2 & -W_1 & 0 & R_3 \\ -B & -R_1 & -R_2 & -R_3 & 0 \end{pmatrix}$$
(33)

where

$$B = \partial_0 A_4,$$

$$\mathbf{W} = \nabla \times \mathbf{A},$$

$$\mathbf{N} = \nabla A_0 - \partial_0 \mathbf{A},$$

$$\mathbf{R} = \nabla A_4.$$

(34)

Some subsets of components of $F_{\mu\nu}$ form carrier spaces for other representations of the Galilei group. For example, if $R_a = F_{4a} \equiv 0$ then the remaining components of $F_{\mu\nu}$ transform as a seven-vector of type Ψ_7 . The complete list of various vectors which can be formed using components of $F_{\mu\nu}$ is present in Section 5.

<u>Example</u> 4. As another example of a Galilean field can be considered the matrix 5-vector $D_5 = (\psi, U_1, U_2, U_3, C)$ with components

$$\psi = \beta_0 = \sqrt{2}(\gamma_0 + \gamma_4),$$

$$C = \beta_4 = \sqrt{2}(\gamma_0 - \gamma_4),$$

$$U_a = \beta_a = \gamma_a$$
(35)

where γ_{μ} are Dirac matrices, $\mu = 0, \ldots, 4$, a = 1, 2, 3. Such set of matrices commutes as a 5-vector with the Galilei generators (3) if we choose

$$S_a = \frac{1}{4} \varepsilon_{abc} \gamma_b \gamma_c, \eta_a = \frac{1}{2\sqrt{2}} \left(\gamma_0 + \gamma_4 \right) \gamma_a.$$
(36)

<u>Example 5.</u> As in the two previous examples, the matrix tensor

$$S_{\mu\nu} = \beta_{\mu}\beta_{\nu} - \beta_{\nu}\beta_{\mu} \tag{37}$$

transforms like $F_{\mu\nu}$, that is, as a 10-vector. Moreover, it is possible to form the following matrix vectors:

$$\tilde{\Psi}_{4}: (R_{a} = S_{4a}, B = S_{40}),
\tilde{\Psi}_{6}: (V_{a} = \frac{1}{2}\varepsilon_{abc}S_{bc}, W_{a} = S_{4a}),
\tilde{\Psi}_{7}: (K_{a} = S_{0a}, R_{a} = \frac{1}{2}\varepsilon_{abc}S_{bc}, A = S_{40}).$$
(38)

6 Contractions of representations of Lorentz algebra

It is well known that the Galilei group (algebra) and (some of) its representations can be obtained from the Poincaré group (algebra) and from its appropriate representations by a limiting procedure called "contraction". The process of contraction has, by now, an extensive literature. First it was proposed by Segal [18] and in more specific forms by Inönü and Wigner [7], by Saletan [19], by Doebner and Melsheimer [20] and many others, see the excellent review article of Lôhmus [21] and references cited therein, a more recent review can be found in [22].

The Lie algebra of a given Lie group is defined via commutation relations for basis elements. As shown by Cartan, whenever we change the basis by a non-singular transformation we come to algebra isomorphic to the original one. However, if the transformation is singular, a new algebra may be received, provided this transformation leads to well defined commutation relations for the transformed basis elements.

In the simplest case a *contraction* is a limit procedure which transforms an Ndimensional Lie algebra \mathcal{L} into an non-isomorphic Lie algebra \mathcal{L}' , also with N dimensions. The commutation relations of a *contracted Lie algebra* \mathcal{L}' are given by:

$$[x, y]' \equiv \lim_{\varepsilon \to \varepsilon_0} W_{\varepsilon}^{-1}([W_{\varepsilon}(x), W_{\varepsilon}(y)],$$
(39)

where $W_{\varepsilon} \in \mathrm{GL}(N,k)$ is a non-singular linear transformation of \mathcal{L} , with ε_0 being a singularity point of its inverse W_{ε}^{-1} .

The papers [18]-[22] indicate different ways of performing this (and more general) singular transformation and necessary and sufficient conditions that a given Lie algebra can be contracted into another one. However there is no a regular way to obtain

representations of the contracted algebra starting with the representations of the initial one. Namely, in contracting representations we meet the following difficulties of principle:

i) Contracting the faithful representation of a given Lie algebra we obtain in general a non-faithful representation of the resulting Lie algebra since a part of generators is represented trivially.

ii) The resulting (contracted) algebra is always non-compact. Hence any contraction of a Hermitian irreducible representation of some compact Lie algebra, which is always *finite*-dimensional, has to yield at the end to an *infinite*-dimensional Hermitian irreducible representation of the non-compact Lie algebra.

Inönü and Wigner in [7] mentioned possible ways of treating the difficulties (for details see paper [23] where contraction of representations of the de Sitter group was carried out), one of them will be used in what follows.

There is a simple contractions procedure (the Inönü-Wigner contraction) connecting the Lie algebra so(1,3) of Lorentz group with algebra hg(1,3). The related transformation W does not change basis elements of so(1,3) forming its subalgebra so(3) while the remaining basis elements are multiplied by a small parameter ε which tends to zero [7].

Here we find representations of the Lorentz group whose contraction makes it possible to obtain found realizations of homogeneous Galilei group. A specific feature of these representations is that they are in general case completely reducible while the corresponding contracted representations of Galilei algebra are indecomposable ones.

Let $S_{\mu\nu}$, $\mu, \nu = 0, 1, 2, 3$ are matrices realizing a representation of the Lorentz algebra, i.e., satisfying the relations

$$[S_{\mu\nu}, S_{\lambda\sigma}] = i(g_{\mu\lambda}S_{\nu\sigma} + g_{\nu\sigma}S_{\mu\lambda} - g_{\nu\lambda}S_{\mu\sigma} - g_{\mu\sigma}S_{\nu\lambda})$$
(40)

with $g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$. The contraction procedure consists in transition to a new basis

$$S_{ab} \to S_{ab}, \ S_{0a} \to \varepsilon S_{0a}$$

and simultaneous transformation of all basis elements $S_{\mu\nu} \to S'_{\mu\nu} = U S_{\mu\nu} U^{-1}$ with a matrix U depending on ε . Moreover, u should depend on ε in a tricky way, such that all the transformed generators $S'_{\mu\nu}$ are kept non-trivial when $\varepsilon \to 0$ [7].

We suppose that representations obtained by the contraction are indecomposable representations $D(n, m, \lambda)$ described in Section 4. To construct representations of Lorentz algebra which can be contracted to representations $D(n, m, \lambda)$ we use the following observation.

Lemma Let $\{S_a, \eta_a\}$ be an indecomposable set of matrices realizing one of representations D(1, 1, 0), D(1, 1, 1), D(1, 2, 1), D(2, 0, 0) or D(3, 1, 1) presented in Table 1. Then matrices

$$S_{ab} = \varepsilon_{abs} S_c, \ S_{0a} = \nu (\eta_a - \eta_a^{\dagger}) \tag{41}$$

where $\nu = 1$ for representations D(1, 1, 0), D(1, 1, 1) D(2, 0, 0) and $\nu = \frac{1}{\sqrt{2}}$ for representations D(1, 2, 1), D(3, 1, 1), form a basis of the Lie algebra of Lorentz group.

The proof is reduced to the direct verification, that for all basis elements S_a , η_a of homogeneous Galilei algebra the corresponding linear combinations (41) satisfy the relations (40), i.e., do form a basis of the Lorentz algebra.

For representations D(1,0,0), (D(2,1,1), D(2,1,0)) and D(2,2,1) the related matrices (41) do not form a basis of Lie algebra. Nevertheless it is possible to find the corresponding generators of the Lorentz algebra starting with its known representations of dimension 3×3 , 7×7 and 8×8 correspondingly. We choose these representations in the following forms:

$$S_{ab} = \varepsilon_{abc} s_c, \ S_{0a} = i s_a, \tag{42}$$

$$S_{ab} = \varepsilon_{abc} \begin{pmatrix} s_c & \mathbf{0}_{3\times3} & \mathbf{0}_{3\times1} \\ \mathbf{0}_{3\times3} & s_c & \mathbf{0}_{3\times1} \\ \mathbf{0}_{1\times3} & \mathbf{0}_{1\times3} & 0 \end{pmatrix}, \quad S_{0a} = \frac{1}{2} \begin{pmatrix} is_a & -s_a & i\sqrt{2}k_a^{\dagger} \\ s_a & is_a & -\sqrt{2}k_a^{\dagger} \\ i\sqrt{2}k_a & \sqrt{2}k_a & 0 \end{pmatrix}, \quad (43)$$

and

$$S_{ab} = \varepsilon_{abc} \begin{pmatrix} s_c & \mathbf{0}_{3\times3} & \mathbf{0}_{3\times1} & \mathbf{0}_{3\times1} \\ \mathbf{0}_{3\times3} & s_c & \mathbf{0}_{3\times1} & \mathbf{0}_{3\times1} \\ \mathbf{0}_{1\times3} & \mathbf{0}_{1\times3} & 0 & 0 \\ \mathbf{0}_{1\times3} & \mathbf{0}_{1\times3} & 0 & 0 \end{pmatrix}, \ S_{0a} = \frac{1}{2} \begin{pmatrix} is_a & -s_a & ik_a^{\dagger} & k_a^{\dagger} \\ s_a & is_a & -k_a^{\dagger} & ik_a^{\dagger} \\ ik_a & k_a & 0 & 0 \\ -k_a & ik_a & 0 & 0 \end{pmatrix}$$
(44)

where s_a and k_a are matrices given by equations (13) and (15).

Matrices (41)-(44) form basises of representations of Lorentz algebra which are in general reducible. These representations (together with the related realizations of homogeneous Galilei algebra given in Table 1) are enumerated in Table 2.

Thus we obtain the indecomposable representations of the Lie algebra of the homogeneous Galilei group starting with finite-dimension representations of the Lorentz algebra so(1,3) and applying the contraction procedure. Special features of this approach are summarized at the following items.

- To obtain *indecomposable* representations of algebra hg(1,3) found in Section 4 we were supposed to use *completely reducible* representations of the Lorentz algebra.
- It is possible to obtain different non-equivalent realizations of hg(1,3) starting with a given representation of so(1,3). For example, both realizations D_7 and D_8 can be obtained via contractions of the representation $D(\frac{1}{2}, \frac{1}{2}) \oplus D(1,0)$.
- It is possible to obtain a given representation of hg(1,3) via contraction of different representations of so(1,3). For example, above mentioned representation $D(\frac{1}{2},\frac{1}{2}) \oplus D(1,0)$ can be replaced by $D(\frac{1}{2},\frac{1}{2}) \oplus D(0,1)$, or, more generally, any representation D(m,n) of so(1,3) can be replaced by D(n,m).

Representation	Representations	Basis	Contracting matrix U
of algebra $hg(1,3)$	of algebra $so(1,3)$	elements	
		$S_{\mu\nu}$	
D(1, 0, 0)	D(1,0)	(42)	$I_{3 imes 3}$
D(1,1,0)	$D(rac{1}{2},rac{1}{2})$	(41)	$\left(egin{array}{ccc} I_{3 imes 3} & 0_{3 imes 1} \ 0_{1 imes 3} & arepsilon^{-1} \end{array} ight)$
D(1,1,1)	$D(\frac{1}{2},\frac{1}{2})$	(41)	$\left(egin{array}{ccc} I_{3 imes 3} & 0_{3 imes 1} \ 0_{1 imes 3} & arepsilon \end{array} ight)$
D(1,2,1)	$D(rac{1}{2},rac{1}{2})\oplus D(0,0)$	(41)	$\begin{pmatrix} I_{3\times3} & 0_{3\times1} & 0_{3\times1} \\ 0_{1\times3} & \varepsilon & 0 \\ 0_{1\times3} & 0 & \varepsilon^{-1} \end{pmatrix}$
D(2,0,0)	$D(0,1)\oplus D(1,0)$	(41)	$\left(\begin{array}{cc}I_{3\times3} & 0_{3\times3}\\ 0_{3\times3} & \varepsilon^{-1}I_{3\times3}\end{array}\right)$
D(2, 1, 0)	$D(rac{1}{2},rac{1}{2})\oplus D(1,0)$	(43)	$\begin{pmatrix} I_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & \varepsilon^{-1}I_{3\times3} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & \varepsilon^{-1} \end{pmatrix}$
D(2, 1, 1)	$D(rac{1}{2},rac{1}{2})\oplus D(1,0)$	(43)	$\begin{pmatrix} I_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & \varepsilon I_{3\times3} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & \varepsilon \end{pmatrix}$
D(2, 2, 1)	$egin{aligned} D(rac{1}{2},rac{1}{2})\oplus\ D(0,0)\oplus D(0,0) \end{aligned}$	(44)	$\begin{pmatrix} I_{3\times3} & 0_{3\times3} & 0_{3\times1} & 0_{3\times1} \\ 0_{3\times3} & \varepsilon^{-1}I_{3\times3} & 0_{1\times3} & 0_{1\times3} \\ 0_{1\times3} & 0_{1\times3} & 1 & 0 \\ 0_{1\times3} & 0_{1\times3} & 0 & \varepsilon^{-1} \end{pmatrix}$
D(3, 1, 1)	$D(0,1) \oplus D(1,0) \ \oplus D(rac{1}{2},rac{1}{2})$	(41)	$\begin{pmatrix} I_{3\times3} & 0_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & \varepsilon^{-1}I_{3\times3} & 0_{3\times3} & 0_{3\times1} \\ 0_{3\times3} & 0_{3\times3} & \varepsilon^{-2}I_{3\times3} & 0_{3\times1} \\ 0_{1\times3} & 0_{1\times3} & 0_{1\times3} & \varepsilon^{-1} \end{pmatrix}$

Table 2: Representations of Lorentz algebra and contracting matrices

7 Galilean linear spin one-half wave equation with Pauli anomalous interaction

The completed list of indecomposable spinor and vector representations of algebra hg(1,3) found in Section 3 and 4 can be used to construct Galilei invariant models for spinorial and vector fields both linear and nonlinear ones. In particular these representations can be applied to derive systems of equations invariant with respect to Galilei group.

In this section we apply them to describe possible Pauli type interactions for spinor field invariant with respect to Galilei group.

7.1 Reduction approach

Consider the Dirac equation describing a fermionic field $\Psi(x)$ coupled to a gauge field A_{μ} , and a Pauli anomalous term :

$$(\gamma_{\mu}\pi^{\mu} + k[\gamma^{\mu}, \gamma^{\nu}]F_{\mu\nu} - \lambda)\Psi(x) = 0, \qquad (45)$$

where

$$\pi_{\mu} \equiv p_{\mu} - qA_{\mu}$$

and the second term is the Pauli anomalous term.

A natural way to construct a Galilean analogue of equation (45) is to generalize it to the case of (4, 1)-dimensional space and then make the reduction discussed in Section 1. However in this case it is desirable to present a clear physical interpretation for all values obtained by the reduction. In addition, in this way we cannot obtain the most general Pauli interaction term. Thus we consider two possibilities: the reduction of equation defined in (4, 1)-dimensional space and a direct search for Pauli term invariant under the Galilei group.

In order to perform the reduction, it is sufficient to change in (45) the Dirac matrices to their Galilean analogues, i.e., to change $\gamma_{\mu} \rightarrow \beta_{\mu}$ where β_{μ} are defined by relations (35). In particular they may be chosen as [24]

$$\beta^{0} = \begin{pmatrix} 0 & 0 \\ -\sqrt{2} & 0 \end{pmatrix}, \qquad \beta^{a} = \begin{pmatrix} \sigma^{a} & 0 \\ 0 & -\sigma^{a} \end{pmatrix}, \qquad \beta^{4} = \begin{pmatrix} 0 & \sqrt{2} \\ 0 & 0 \end{pmatrix}.$$
(46)

If we substitute these matrices into equation (45) for $\psi = \begin{pmatrix} \phi \\ \chi \end{pmatrix}$ (where ϕ and χ are two component spinors) and use the notations (33), we find

$$(\boldsymbol{\sigma} \cdot \boldsymbol{\pi} - \lambda + 4ik\boldsymbol{\sigma} \cdot \mathbf{W} + 4kB]\varphi + \sqrt{2} (\pi_4 + 4k\boldsymbol{\theta} \cdot \mathbf{R})\chi = 0,$$

$$\sqrt{2} (\pi_0 - 4k\boldsymbol{\sigma} \cdot \mathbf{N})\varphi + (\boldsymbol{\sigma} \cdot \boldsymbol{\pi} + \lambda - 4ik\boldsymbol{\sigma} \cdot \mathbf{W} + 4kB)\chi = 0$$
(47)

where $\pi_4 = m - qA_4$. Solving the first of equations (47) for χ and substituting the solution into the second line we come to the Schrödinger equation with spin dependent potential which is not discussed here.

In the magnetic limit, defined as in reference [17]:

$$(\mathbf{A}_m, \phi_m) \hookrightarrow A_m = (A_0, \mathbf{A}_m, A_4) = (-\phi_m, \mathbf{A}_m, 0), \qquad (48)$$

we find

$$\mathbf{B}_m = \mathbf{W} = \nabla \times \mathbf{A}_m, \\ \mathbf{E}_m = \mathbf{N} = -\nabla \phi_m - \partial_t \mathbf{A}_m$$

Note that $\mathbf{R} = \mathbf{0}$ and B = 0, and Galilei transformations for $\mathbf{A}_{\mathbf{m}}$, ϕ_m , \mathbf{B}_m and \mathbf{E}_m have the form

$$\begin{aligned}
\mathbf{A}_m &\to \mathbf{A}_m, & \phi_m \to \phi_m - \mathbf{v} \cdot \mathbf{A}_m, \\
\mathbf{B}_m &\to \mathbf{B}_m, & \mathbf{E}_m \to \mathbf{E}_m + \mathbf{v} \times \mathbf{B}_m.
\end{aligned} \tag{49}$$

With these definitions, equation (47) becomes

$$(\boldsymbol{\sigma} \cdot (\boldsymbol{\pi} - q\mathbf{A}_m) - \lambda + 4ik\boldsymbol{\sigma} \cdot \mathbf{B}_m)\varphi + \sqrt{2} \ m\chi = 0,$$

$$\sqrt{2} \ (p_0 + q\phi_m - 4k\boldsymbol{\sigma} \cdot \mathbf{E}_m)\varphi + [\boldsymbol{\sigma} \cdot (\boldsymbol{\pi} - q\mathbf{A}_m) + \lambda - 4ik\boldsymbol{\sigma} \cdot \mathbf{B}_m]\chi = 0.$$

This may be rewritten as

$$\left[\gamma_{\mu}\pi^{\mu} - \lambda + 2k\left(\mathbf{S}\cdot\mathbf{B}_{m} + \boldsymbol{\eta}\cdot\mathbf{E}_{m}\right)\right]\psi = 0$$
(50)

where **S** and η are matrix vectors whose components are given in equation (36).

Thus we obtain in rather elegant way equation (50) which is manifestly Galilei invariant and describes both the minimal and anomalous couplings of particle of spin 1/2 with an external Galilean electromagnetic field of magnetic type. In the case k = 0, i.e., when only the minimal interaction is present, this equation is reduced to the Lévy-Leblond equation. However, it presents only one of many possibilities to introduce anomalous coupling into the Lévy-Leblond equation. A more general approach is presented in the following subsection.

7.2 Direct approach

Anomalous Pauli interaction is represented in equation (45) by the term $k[\gamma_{\mu}, \gamma_{\nu}]F^{\mu\nu}$, which is linear with respect to the electromagnetic field strength F. It is possible to show that the requirement of relativistic invariance defines this term in a unique way (up to a value of the coupling constant k).

In contrast with the above, in the Galilei-invariant approach there are more possibilities to introduce the anomalous interaction. In this subsection we use our knowledge of Galilean vector field (refer to Section 4) to find the most general form of anomalous interaction for Galilean spinors.

First we remind that there exist two types of massless Galilean 4-vector fields, i.e., Ψ_4 and $\tilde{\Psi}_4$, whose transformation properties are defined by equations (22) and

(23) with m = 0, and a 5-vector Ψ_5 which transforms as given in (31). In other words we have three types of potentials of external vector field which can be used to introduce the *minimal* interaction into the Lévy-Leblond equation. And they are potentials which generate field strengthes involved into anomalous interaction terms whose examples are present in the previous subsection.

We notice that in addition it is possible to introduce minimal and anomalous interactions with fields whose potentials are three-vectors or scalars. Moreover, the Galilei invariance condition admits some constrains for the potentials which generate additional possible anomalous interactions.

Let us start with the vector $\tilde{\Psi}_4$ which corresponds to the magnetic limit field considered in the previous section, and find the most general Galilean scalar matrix F linear in \mathbf{B}_m and \mathbf{E}_m . Expanding F via the complete set of matrices β_{μ} (35) and $S_{\mu\nu}$ (37) with well defined transformation properties and using (49) we easily find that $F = 2k (\mathbf{S} \cdot \mathbf{B}_m + \boldsymbol{\eta} \cdot \mathbf{E}_m) + g \boldsymbol{\eta} \cdot \mathbf{B}_m$ where k and g are arbitrary parameters. In other words, we come to the following Galilei-invariant equation for spinor field with the minimal and anomalous interaction:

$$\left[\gamma_{\mu}\pi^{\mu} - \lambda + 2k\left(\mathbf{S}\cdot\mathbf{B}_{m} + \boldsymbol{\eta}\cdot\mathbf{E}_{m}\right) + g\boldsymbol{\eta}\cdot\mathbf{B}_{m}\right]\psi = 0.$$
(51)

In contrast with (50), equation (51) includes two coupling constants, k and g.

The other Galilean limit, named 'electric limit' [6] for the electromagnetic field corresponds to the gauge fields of type Ψ_4 , i.e., the related vector-potential

$$(\mathbf{A}_e, \phi_e) \hookrightarrow A_e = (A_0, \mathbf{A}, A_4) = (0, \mathbf{A}_e, \phi_e), \qquad (52)$$

and field strengthes $\mathbf{B}_e = \nabla \times \mathbf{A}_e$, $\mathbf{E}_e = -\nabla \phi_e$, $\Phi = \partial_0 \phi$ transform as

$$\begin{aligned}
\mathbf{A}_{e} &\to \mathbf{A}_{e} + \mathbf{v}\phi_{e}, & \phi_{e} \to \phi_{e} \\
\mathbf{E}_{e} &\to \mathbf{E}_{e}, & \mathbf{B}_{e} \to \mathbf{B}_{e} - \mathbf{v} \times \mathbf{E}_{e}, & \Phi \to \Phi - \mathbf{v} \cdot \mathbf{E}_{e}.
\end{aligned}$$
(53)

Searching for the related Galilean Dirac equation with general Pauli interaction term and using the fact that the vectors $(\mathbf{B}_e, -\mathbf{E}_e)$ have the same transformation properties as $(\mathbf{E}_m, \mathbf{B}_m)$ we conclude that to achieve our goal it is sufficient to change $\mathbf{E}_m \to \mathbf{B}_e$ and $\mathbf{B}_m \to -\mathbf{E}_e$ in (51) and add the additional invariant term $\gamma_0 \Phi + \boldsymbol{\gamma} \cdot \mathbf{E}_e$. In addition, to keep the Galilei invariance we should introduce the minimal interaction in the following manner:

$$\pi_0 = i\partial_0,$$

$$\pi_a = -i\partial_a - q(A_e)_a$$

$$\pi_4 = m + q\phi_e$$

As a result we obtain

$$\left(\gamma_{\mu}\pi^{\mu} - \lambda + 2k\left(\boldsymbol{\eta}\cdot\mathbf{B}_{e} - \mathbf{S}\cdot\mathbf{E}_{e}\right) + g\boldsymbol{\eta}\cdot\mathbf{E}_{e} + r(\gamma_{0}\phi_{e} + \boldsymbol{\gamma}\cdot\mathbf{E}_{e})\right)\psi = 0.$$
(54)

Thus there exist at least two ways to describe anomalous interaction in the Galilei invariant approach, presented by equations (51) and (54). Equation (51) includes two coupling constants while in (54) the number of such constants is equal to 3.

Note that transformation laws (49) enable to impose in (51) the Galilei-invariant condition $\mathbf{A}_m = 0$ on the vector-potential. Thus there exist one more Galilei-invariant equation with anomalous interaction, namely

$$\left[\gamma_0(p_0 + q\phi) - \boldsymbol{\gamma} \cdot \mathbf{p} - \gamma_4 m + 2k\boldsymbol{\eta} \cdot \nabla\phi\right]\psi = 0.$$
(55)

Analogously, starting with (54) taking into account that transformations (53) are compatible with the condition $\phi_e = 0$ we come to one more Galilei invariant equation with Pauli interaction term, namely

$$(\gamma_0 \partial_0 - \boldsymbol{\gamma} \cdot \boldsymbol{\pi} - \gamma_4 m - \lambda + g \boldsymbol{\eta} \cdot \mathbf{B}) \psi = 0,$$
(56)

where $\boldsymbol{\pi} = -i\nabla - q\mathbf{A}, \ \mathbf{B} = \nabla \times \mathbf{A}.$

Gauge invariant fields $\mathbf{E} = -\nabla \phi$ and $\mathbf{B} = \nabla \times \mathbf{A}$ present in equations (55) and (56) can be interpreted as external electric and magnetic fields.

Consider also the case when the potential of external field form a 5-vector, and related field strengthes are given by formulae (32)-(34). We will not discuss possible physical interpretation of the corresponding ten-component vector field but mention that there exist a formal possibility to introduce the anomalous interaction with it by generalizing the Galilean Dirac equation to the following one:

$$(\gamma^{\mu}\pi_{\mu} + k[\gamma^{\mu}, \gamma^{\nu}]F_{\mu\nu} + g\eta_{a}R_{a} + r(S_{a}R_{a} + \eta_{a}U_{a}) - \lambda)\Psi(x) = 0,$$
(57)

with three coupling constants k, g r additional to q.

Like equation (45) with γ -matrices (46), equation (57) is Galilei-invariant but is quite more general.

Finally let the external field is defined by a five-vector potential satisfying the Galilei invariant constraint

$$\partial_a A_4 = 0, \tag{58}$$

then $F_{4a} \equiv 0$ and tensor $F_{\mu\nu}$ (33) is reduced to the seven-vector $\tilde{\Psi}_7 = (\mathbf{K}, \mathbf{R}, A)$ whose components are

$$\mathbf{K} = \nabla A_0 - \partial_0 \mathbf{A}, \ \mathbf{R} = \nabla \times \mathbf{A}, \ A = \partial_0 A_4.$$

To find the corresponding invariants linear in $F_{\mu\nu}$ it is necessary to construct invariant scalar products of vector functions (58) and matrices (38) belonging to $\tilde{\Psi}_7$. As a result we come to the following equation

$$\left(\gamma^{\mu}\pi_{\mu} + g\eta_{a}R_{a} + r(S_{a}R_{a} + \eta_{a}K_{a} + S^{04}A) + \nu\gamma_{0}A - \lambda\right)\psi(x) = 0,$$
 (59)

where g, r and ν are coupling constants. For A = 0 equation (59) is reduced to equation (51) which describe anomalous interaction of the Galilei particle of spin 1/2 with external field of magnetic type.

Formulae (51), (54), (55), (56), (57) and (59) present all non-equivalent Galilean invariant equations for spinor field, describing minimal and anomalous interactions with external gauge fields.

We see that the direct search for Galilei invariant Pauli interaction makes it possible to find more general coupling than the reduction method.

7.3 Galilean system with spin-orbit coupling

In this section we consider one of the described systems and discuss its physical content.

Let us start with equation (55) which describes interaction of the Galilean spinor particle with an external electric field. Choosing γ -matrices in the form (46) and denoting $k = -\frac{q\hat{k}}{4m}$ we write this equation componentwise

$$(\boldsymbol{\sigma} \cdot \mathbf{p} - \lambda)\varphi + \sqrt{2} \ m\chi = 0,$$

$$\sqrt{2} \ (p_0 + q\phi + \frac{q\hat{k}}{m}\boldsymbol{\sigma} \ \cdot \mathbf{E})\varphi + (\boldsymbol{\sigma} \cdot \mathbf{p} + \lambda)\chi = 0$$

where $\mathbf{E} = -\nabla \phi$.

Solving the first equation for χ and substituting the result into the second equation we obtain

$$L\varphi \equiv \left(p_0 - \frac{p^2 + \lambda^2}{2m} + q\phi - \frac{q\hat{k}}{m}\boldsymbol{\sigma} \cdot \mathbf{E}\right)\varphi = 0.$$
(60)

In other word we come to the Galilei invariant Schrödinger equation with a matrix potential.

To analyze the physical content of equation (60) we transform it to a more familiar form using the operator $U = \exp(-\frac{ki}{m}\boldsymbol{\sigma}\cdot\mathbf{p})$. Applying this operator to φ and L we obtain the equation $L'\varphi' = 0$ where $\varphi' = U\varphi$,

$$L' = ULU^{-1} = p_0 - \frac{p^2}{2m} + q\phi - \frac{\lambda^2}{2m} - \frac{q\hat{k}^2}{2m^2} \left(\boldsymbol{\sigma} \cdot \left(\mathbf{p} \times \mathbf{E} - \mathbf{E} \times \mathbf{p}\right) - \mathtt{div}\mathbf{E}\right) + \cdots$$

where the dots denote the terms of order $o(\frac{1}{m^3})$.

All terms in operator L' have exact physical meaning. In particular, the last two terms describe spin-orbit and Darwin coupling of a Galilean particle with an external field.

8 Discussion

The relativity principle is one of the corner stones of modern physics. Moreover, it should be applied while considering phenomena characterizing by circumlight velocities but also in the cases when the velocities are small in comparison with the velocity of light. In the latest case any well formulated physical theory should not be simple "non-relativistic", but to satisfy the Galilei relativity principle. In other words, the group of motion of the special relativity (i.e., the Lorentz group) should be replaced by the Galilei group.

It appears that a consistent use of Galilei group and its representations in many aspects is much more complicated than in the case of Poincaré group. In particular, description of finite- dimensional representations of the homogeneous Galilei group is a wild algebraic problem while such representations for the Lorentz group have been found long time ago.

The main goal of the present paper was to present the completed classification of indecomposable Galilean fields which transform as vectors or are scalars with respect to the rotation transformations. In other words, we present the classification of indecomposable finite dimension representations of the homogeneous Galilei group which being reduced to the rotation subgroup correspond to spins $s \leq 1$. In contrast with the fields with spin s > 1 the Galilean vectors field can be described completely. The results of this classification are presented in Section 3.

In contrast with the Lorentz vectors, the number of non-equivalent Galilean vectors appears to be rather extended. Namely, for the Lorentz group there exist the following indecomposable vector fields: four-vector, bi-vector (i.e., antisymmetric tensor of second rank) and two three-vectors which are nothing but a self-dual and anti self-dual parts of the antisymmetric tensor. In the case of Galilei group it is possible to indicate nine indecomposable vector fields.

We use our knowledge of vector representations of the Galilei group to describe all possible Pauli interactions for spinor fields, compatible with the Galilei invariance. The number of such interactions appears to be rather extended in contrast with the relativistic approach where this interaction is unique up to the coupling constant. We show that there exist such Galilei invariant systems which describe spin-orbit and Darwin couplings which are traditionally treated as pure relativistic effects.

It is generally accepted to think that Galilei group and its representations can be obtained easily starting with representations of the Lorentz group and making the Inönü-Wigner contraction [7]. We had shown that this procedure is not too straightforward in as much as starting with indecomposable representations of the Lorentz group we can obtain only a part of the corresponding representations of homogeneous Galilei group. On the other hand, it is possible to contract reducible representations of the Lorentz group to indecomposable representations of the homogeneous Galilei group.

The completed list of vector-scalar and spinor indecomposable representations representations presented in Sections 3 and 4 opens a way to describe all non-equivalent systems of Galilei invariant equations for scalar, spinor and vector fields. We plane to discuss these equations in detail in the following paper.

Ackowledgements

This publication is based on work sponsored by the Grant Agency of the Academy of Sciences of the Czech Republic under project Number A1010711. Partial financial support was provided by the Natural Sciences and Engineering Research Council (NSERC) of Canada.

References

- Lévy-Leblond J M 1971 Galilei group and galilean invariance, in *Group Theory* and Applications Ed. E.M. Loebl, Vol. II (New York: Academic) 221-99
- [2] Cassinelli G, De Vito E, Lahti P J and Levrero A 2004 The theory of Symmetry Action in Quantum Mechanics (Lect. Notes Phys. 654, Springer, Berlin)
- [3] Inönü E and Wigner E1952 Representations of the Galilei group Nuov. Cim. 9 B 705-718
- [4] Bargmann V 1954 1954 On unitary ray representations of continuous groups Ann. Math. 59 1-46
- [5] Lévy-Leblond J M 1967 Non-relativistic particles and wave equations Comm. Math. Phys. 6 286-311
- [6] Le Bellac M and Lévy-Leblond J M 1973 Galilean electromagnetism Nuov. Cim. 14 B 217-33
- [7] Inönü E and Wigner E P 1953 On the contraction of groups and their representations Proc. Nat. Acad. Sci. U.S. 39 510-24
- [8] Fushchich V I and Nikitin A G 1980 Reduction of the representations of the generalised Poincaré algebra by the Galilei algebra J. Phys. A: Math. Gen. 13 2319-30
- [9] Takahashi Y 1988 Towards the many-body theory with the Galilei invariance as a guide. I. Fortschr. Phys. 36 63-81

Takahashi Y 1988 Towards the many-body theory with the Galilei invariance as a guide. II. *Fortschr. Phys.* **36** 83-96

Takahashi Y 1987 An invitation to a Galilei invariant world, in Wandering in the Fields: Festschrift for Professor Kazahiko Nishijima on the Occasion of his Sixtieth Birthday Eds. K. Kwarabayashi and A. Ukawa (Singapore: World Scientific) 117-27

Omote M, Kamefuchi S, Takahashi Y and Ohnuki Y 1989 Galilean covariance and the Schrödinger equation *Fortschr. Phys.* **37** 933-50

 [10] Soper D E 1976 Classical Field Theory (New York: Wiley and Sons), Section 7.3 Pinski G 1968 Galilean tensor calculus J. Math. Phys. 9 1927-30

Künzle H P and Duval C 1994 Relativistic and nonrelativistic physical theories on five-dimensional space-time, in *Semantical Aspects of Spacetime Theories* Eds. U. Majer and H.J. Schmidt (Mannheim: BI-Wissenschaftsverlag) 113-129 (and the references therein)

- [11] Kapuścik E 1986 On the physical meaning of the Galilean space-time coordinates Acta Phys. Pol. B 17 569-75
 Kapuścik E 1980 On nonrelativistic gauge theories Nuov. Cim. 58 A 113-24
 Kapuścik E 1981 The non-relativistic space-time manifolds Acta Phys. Pol. B 12 81-6
- [12] Nikitin A G and Fuschich W I 1980 Equations of motion for particles of arbitrary spin invariant under the Galilei group *Theor. Math. Phys.* 44 584-92
- [13] Fushchich W I and Nikitin A G 1994 Symmetries of Equations in Quantum Mechanics (New York: Allerton Press)
- [14] George C and Lévi-Nahas M 1966 Finite-dimensional representations of some non-semisimple Lie algebras J. Math. Phys. 7 980-988
- [15] Gelfand I M, Minlos R A and Shapiro Z Ya 1963 Representations of the Rotation and Lorentz Groups and their Applications (New York: Pergamon Press)
- [16] Corson E M 1955 Introduction to Tensors, Spinors, and Relativistic Wave Equations (London: Blackie & Sons)
- [17] de Montigny M, Khanna F C and Santana A E 2003 Nonrelativistic wave equations with gauge fields Int. J. Theor. Phys. 42 649-71
- [18] Segal I E 1951 A class of operator algebras which are determined by groups Duke Math. J. 18 221-265.
- [19] Saletan E 1961 Contraction of Lie groups J. Math. Phys. 2 1-21
- [20] Doebner H D 1967 On a class of generalized group contractions Nuov. Cim. 49 A 306-311
- [21] Lôhmus J 1967 Contractions of Lie Groups Proceedings of the 1967 Summer School on Elementary Particle Physics, Otepää, Estonia
- [22] Weimar-Woods E 2000 Contractions, generalized Inonu-Wigner contractions and deformations of finite-dimensional Lie algebras *Rev. Math. Phys* 12 1505-1529.
- [23] Mickelsson J and Niederle J 1972 Contractions of representations of de Sitter groups Comm. Math. Phys. 27 167-180.
- [24] de Montigny M, Khanna F C, Santana A E and Santos E S 2001 Galilean covariance and the non-relativistic Bhabha equations J. Phys. A: Math. Gen. 34 8901-17