REPRESENTATIONS AND PROPERTIES OF GENERALIZED A_r **STATISTICS**

M. Daoud¹

Faculté des Sciences, Département de Physique, LPMC, Agadir, Morocco

Abstract

A generalization of A_r statistics is proposed and developed. The generalized A_r quantum statistics is completely specified by a set of Jacobson generators satisfying a set of triple algebraic relations. Fock-Hilbert representations and Bargmann-Fock realizations are derived.

 $^{^{1}}m_{-}daoud@hotmail.com$

1 Introduction

Various kinds of quantum statistics, different from Bose and Fermi statistics, have been introduced in the literature [1-11]. For example, in two space dimension, one can have a one parameter family of statistics (anyons) interpolating between bosons and fermions [4]. On the other hand, in three and higher space dimensions the parastatistics, developed by Green [1], constitute a natural extension of the usual Fermi and Bose statistics. In this generalization, the classical bilinear commutators or anti- commutators are replaced by certain triple relations. The trilinear commutation relations defining the parafermions are given by

$$[[f_i^+, f_j^-], f_k^-]] = -2\delta_{ik}f_j^-, \quad [[f_i^+, f_j^+], f_k^-]] = -2\delta_{ik}f_j^+ + 2\delta_{jk}f_i^+, \quad [[f_i^-, f_j^-], f_k^-]] = 0, \quad (1)$$

for i, j, k = 1, 2, ..., r and the triple relations defining the parabosons are defined as

$$[\{b_i^+, b_j^-\}, b_k^-]] = -2\delta_{ik}b_j^-, \quad [\{b_i^+, b_j^+\}, b_k^-]] = -2\delta_{ik}b_j^+ - 2\delta_{jk}b_i^+, \quad [\{b_i^-, b_j^-\}, b_k^-]] = 0.$$
(2)

It was established that the relations (1) are associated with the orthogonal Lie algebra $so(2r + 1) = B_r$ [12] and the relations (2) are connected to the orthosymplectic superalgebra osp(1/2r) = B(0,r) [13]. Recently, in the spirit of Palev works [6,10], a classification of generalized quantum statistics were derived for the classical Lie algebras A_r , B_r , C_r and D_r [11]. The increasing interest in generalized statistics stay essentially motivated by their possible relevance in the understanding of the theory of fractional quantum Hall effect [14-15], the superconductivity [16], and probably they can give some adequate description of black hole statistics [17].

In this paper, We shall rather focus our attention on a generalization of A_r statistics introduced in [6]. Let us discuss how to realize the generators and the quantum states, associated with the generalized A_r statistics, in a purely analytical way. This yields, in a natural way, coherent states which are interesting in many respects [18,19], particularly in the study of dynamical evolution of quantum systems.

The content of this paper is as follows. Generalized quantum statistics are approached from a set of Jacobson generators [20] (defined in section 2) satisfying certain triple relations. This generalization includes two fundamental sectors. A fermionic one reproducing the A_r statistics introduced in [6]. The second sector approached through this generalization is of bosonic kind. For each sector, we give the associated Fock space. An Hamiltonian is derived in terms of the Jacobson generators identified with creation and annihilation operators. In section 3, two non equivalents analytic realizations of the Fock space for the bosonic A_r statistics are performed. The first realization brings the so-called Gazeau-Klauder coherent states [21]. The second realization leads to the well known Klauder-Perelomov coherent states [18,19]. Differential actions of the Jacobson generators in both realizations are given. In section 4, we realize analytically the Fock space related to the A_r statistics of fermionic kind. In this case, we show that the Jacobson generators act on a over-complete set of coherent states similar to Klauder-Perelomov ones labeling the complex projective spaces \mathbb{CP}^r . Some concluding remarks close this paper.

2 The generalized A_r statistics

This section is devoted to the definition of the generalized A_r statistics viewed as Lie triple system. We give the corresponding Fock space and the Hamiltonian describing a quantum system obeying the generalized A_r statistics.

2.1 The algebra

To begin, let us recall the definition of Lie triple systems. A vector space with trilinear composition [x, y, z] is called Lie triple system if the following identities are satisfied:

$$\begin{split} [x,x,x] &= 0, \\ [x,y,z] + [y,z,x] + [z,x,y] = 0, \\ [x,y,[u,v,w]] &= [[x,y,u],v,w] + [u,[x,y,v],w] + [u,v,[x,y,w]] \end{split}$$

Following this definition, we will introduce the generalized A_r statistics as Lie triple system. For this end, we consider the set of 2r operators x_i^+ and $x_i^- = (x_i^+)^{\dagger}$, (i = 1, 2, ..., r). Inspired by the para-Fermi case [1] and the example of A_r statistics [6,10], these 2r operators should satisfy certain conditions and relations. First, the operators x_i^+ are mutually commuting. A similar statement holds for the operators x_i^- . They also satisfy the following triple relations

$$[[x_i^+, x_j^-], x_k^+]] = -\epsilon \delta_{jk} x_i^+ - \epsilon \delta_{ij} x_k^+$$
(3)

$$[[x_i^+, x_j^-], x_k^-]] = \epsilon \delta_{ik} x_j^- + \epsilon \delta_{ij} x_k^-$$

$$\tag{4}$$

where $\epsilon \in \mathbf{R}$. It is easy to see that the algebra \mathcal{A} (defined by means of the generators x_i^{\pm} and relations (3) and (4)) is closed under the ternary operation [x, y, z] = [[x, y], z] and define a Lie triple system. Note that for $\epsilon = -1$, the algebra \mathcal{A} reduces to one defining the A_r statistics discussed in [6]. It is also important to remember that the description (3) and (4) of algebras by means of minimal set of generators and relations was initiated by N. Jacobson [20]. Therefore, the elements x_i^{\pm} are referred to as Jacobson generators which will be identified later with creation and annihilation operators of a quantum system obeying to generalized A_r statistics. We redefine the generators of the algebra \mathcal{A} as $a_i^{\pm} = \frac{x_i^{\pm}}{\sqrt{|\epsilon|}}$. The triple relations (3) and (4) rewrite now as

$$[[a_i^+, a_j^-], a_k^+]] = -s\delta_{jk}a_i^+ - s\delta_{ij}a_k^+$$
(5)

$$[[a_i^+, a_j^-], a_k^-]] = s\delta_{ik}a_j^- + s\delta_{ij}a_k^-$$
(6)

where $s = \frac{\epsilon}{|\epsilon|}$ is the sign of the parameter ϵ and $[a_i^+, a_j^+] = [a_i^-, a_j^-] = 0$. This redefinition is more convenient for our investigation, in particular in determining the irreducible representation associated with the algebra \mathcal{A} . The sign of the parameter ϵ play an important in the representation of the algebra \mathcal{A} and consequently, one can obtain different microscopic and macroscopic statistical properties of the quantum system under consideration.

2.2 The Hamiltonian

To characterize a quantum gas obeying the generalized A_r statistics, we have to specify the Hamiltonian for the system. The operators a_i^{\pm} define creation and annihilation operators for a quantum mechanical system, described by an Hamiltonian H, when the Heisenberg equation of motion

$$[H, a_i^{\pm}] = \pm e_i a_i^{\pm} \tag{7}$$

is fulfilled. The quantities e_i are the energies of the modes i = 1, 2, ..., r. One can verify that if $|E\rangle$ is an eigenstate with energy E, $a_i^{\pm}|E\rangle$ are eigenvectors of H with energies $E \pm e_i$. In this respect, the operators a_i^{\pm} can be interpreted as ones creating or annihilating particles. To solve the consistency equation (7), we write the Hamiltonian H as

$$H = \sum_{i=1}^{r} e_i h_i \tag{8}$$

which seems to be a simple sum of "free" (non-interacting) Hamiltonians h_i . However, note that, in the quantum system under consideration, the statistical interactions occur and are encoded in the in the triple commutation relations (5) and (6). Using the structure relations of the algebra \mathcal{A} , the solution of the Heisenberg condition (7) is given by

$$h_i = \frac{s}{r+1} \left[(r+1)[a_i^-, a_i^+] - \sum_{j=1}^r [a_j^-, a_j^+] \right] + c \tag{9}$$

where the constant c will be defined later such that the fundamental state (vacuum) of the Hamiltonian H be zero energy.

2.3 The representation

We now consider an Hilbertean representation of the algebra \mathcal{A} . Let \mathcal{F} be the Hilbert-Fock space on which the generators of \mathcal{A} act. Since, the algebra \mathcal{A} is spanned by r pairs of Jacobson generators, it is natural to assume that the Fock space is given by

$$\mathcal{F} = \{ |n_1, n_2, \cdots, n_r \rangle , n_i \in \mathbf{N} \}.$$
(10)

The action of a_i^{\pm} , on \mathcal{F} , are defined as

$$a_i^{\pm}|n_1,\cdots,n_i,\cdots,n_r\rangle = \sqrt{F_i(n_1,\cdots,n_i\pm 1,\cdots,n_r)}|n_1,\cdots,n_i\pm 1,\cdots,n_r\rangle$$
(11)

in terms of the functions F_i called the structure functions which should be non-negatives so that all states are well defined. To determine the expressions of the functions F_i in terms of the quantum numbers n_1, n_2, \dots, n_r , let first assume the existence of a vacuum vector $a_i^- |0, 0, \dots, 0\rangle = 0$ for all $i = 1, 2, \dots, r$. This implies that the functions F_i can be written

$$F_i(n_1, \cdots, n_i, \cdots, n_r) = n_i \Delta F_i(n_1, \cdots, n_i, \cdots, n_r)$$
(12)

in a factorized form where the new functions ΔF_i are defined such that $\Delta F_i(n_1, \dots, n_i = 0, \dots, n_r) \neq 0$ for $i = 1, 2, \dots, r$. Furthermore, since the Jacobson operators satisfy the trilinear relations (5) and (6), these functions should be linear in the quantum numbers n_i :

$$\Delta F_i(n_1, \cdots, n_i, \cdots, n_r) = k_0 + (k_1 n_1 + k_2 n_2 + \cdots + k_r n_r).$$
(13)

Finally, using the relations $[a_i^+, a_j^+] = 0$ and $[[a_i^+, a_i^-], a_i^+]] = -2sa_i^+$, one verifies that $k_i = k_j$ and $k_i = s$, respectively. For convenience, we set $k_0 = k - \frac{1+s}{2}$ assumed to be a non vanishing integer. The actions of Jacobson generators on the states spanning the Hilbert-Fock space \mathcal{F} are now given by

$$a_{i}^{-}|n_{1},\cdots,n_{i},\cdots,n_{r}\rangle = \sqrt{n_{i}(k_{0}+s(n_{1}+n_{2}+\cdots+n_{r}))}|n_{1},\cdots,n_{i}-1,\cdots,n_{r}\rangle$$
(14)

$$a_i^+|n_1,\dots,n_i,\dots,n_r\rangle = \sqrt{(n_i+1)(k_0+s(n_1+n_2+\dots+n_r+1))}|n_1,\dots,n_i+1,\dots,n_r\rangle$$
(15)

The dimension of the irreducible representation space \mathcal{F} is determined by the condition:

$$k_0 + s(n_1 + n_2 + \dots + n_r) > 0.$$
(16)

It depends on the sign of the parameter s. It is clear that for s = 1, the Fock space \mathcal{F} is infinite dimension. However, for s = -1, there exists a finite number of states satisfying the condition $n_1 + n_2 + \cdots + n_r \leq k - 1$. The dimension is given, in this case, by $\frac{(k-1+r)!}{(k-1)!r!}$. This is exactly the dimension of the Fock representation of A_r statistics discussed in [6]. This conditionrestriction is closely related to so-called generalized exclusion Pauli principle according to which no more than k - 1 particles can be accommodated in the same quantum state. In this sense, for s = -1, the generalized A_r quantum statistics give statistics of fermionic behaviour. They will be termed here as fermionic A_r statistics and ones corresponding to s = 1 will be named bosonic A_r statistics.

Setting $c = \frac{r}{r+1}sk_0$ in (9) and using the equation (8) together with the actions of creation and annihilation operators (14-15), one has

$$H|n_1,\cdots,n_i,\cdots,n_r\rangle = \sum_{i=1}^r e_i n_i |n_1,\cdots,n_i,\cdots,n_r\rangle.$$
(17)

Finally, we point out one interesting property of the generalized A_r statistics. Introduce the operators $b_i^{\pm} = \frac{a_i^{\pm}}{\sqrt{k}}$ for $i = 1, 2, \dots, r$ and consider k very large. From equations (14) and (15), we obtain

$$b_i^-|n_1,\cdots,n_i,\cdots,n_r\rangle = \sqrt{n_i}|n_1,\cdots,n_i-1,\cdots,n_r\rangle$$
(18)

$$b_i^+ |n_1, \cdots, n_i, \cdots, n_r\rangle = \sqrt{n_i + 1} |n_1, \cdots, n_i + 1, \cdots, n_r\rangle.$$
 (19)

In this limit, the generalized A_r statistics (fermionic and bosonic ones) coincide with the Bose statistics and the Jacobson operators reduce to Bose ones (creation and annihilation operators of harmonic oscillators).

Beside the Fock representations discussed in this section, it is interesting to look for analytical realizations of the Fock spaces associated with the generalized A_r statistics. These realizations constitute an useful tool in connection with variational and path integral methods to describe the quantum dynamics of the system described by the Hamiltonian H.

3 Bargmann realizations of A_r bosonic statistics

In this section, we give two different realizations à la Bargmann using a suitably defined Hilbert space of entire analytic functions associated to A_r statistics of bosonic kind. In the first analytic realization, the Jacobson creation operators are realized as simple multiplication by some complex variables. In the second one, the Jacobson annihilation operators are represented as derivation with respect to variables of the analytic representation space. As by product, the first (resp. second) realization brings, in a natural way, the Gazeau-Klauder [21](resp. Klauder-Perelomov [18,19]) coherent states associated to a quantum mechanical system described by the Hamiltonian given by (8) and obeying the A_r statistics of bosonic kind.

3.1 Realization I

To begin, we ask for a realization in which the vector $|k; n_1, \dots, n_r\rangle$ is realized as powers of complex variables $\omega_1, \dots, \omega_r$

$$|k; n_1, \cdots, n_r\rangle \longrightarrow C_{k; n_1, \cdots, n_r} \omega_1^{n_1} \cdots \omega_r^{n_r}$$
 (20)

and Jacobson creation operators a_i^+ act as a simple multiplication by ω_i .

The coefficients $C_{k;n_1,\dots,n_r}$ occurring in (20) are determined in the following manner. Using the equation (15) and the realization (20), one has the recursion relation

$$C_{k;n_1,\dots,n_i,\dots,n_r} = ((n_i+1)(k+n_1+\dots+n_i+\dots+n_r))^{\frac{1}{2}}C_{k;n_1,\dots,n_i+1,\dots,n_r},$$
(21)

which can be easily solved

$$C_{k;n_1,\dots,n_i,\dots,n_r} = \left[\frac{(k-1+n-n_i)!}{n_i!(k-1+n)!}\right]^{\frac{1}{2}} C_{k;n_1,\dots,0,\dots,n_r}$$
(22)

where $n = n_1 + n_2 + \cdots + n_r$. Repeating this procedure for all $i = 1, 2, \cdots, r$ and setting $C_{k;0,\cdots,0} = 1$, we obtain

$$C_{k;n_1,\dots,n_i,\dots,n_r} = \left[\frac{(k-1)!}{n_1!\dots n_r!(k-1+n)!}\right]^{\frac{1}{2}}.$$
(23)

The differential realization of the operators $N_i \ (\neq a_i^+ a_i^-)$ acting on the Fock space \mathcal{F} as

$$N_i|k;n_1,\cdots,n_i\cdots,n_r\rangle = n_i|k;n_1,\cdots,n_i\cdots,n_r\rangle,$$
(24)

are given by

$$N_i \longrightarrow \omega_i \frac{d}{d\omega_i}.$$
 (25)

Using the realization (25), the actions of a_i^- on \mathcal{F} and the relations defining the algebra \mathcal{A} , one show that the annihilation operators act as

$$a_i^- \longrightarrow k \frac{d}{d\omega_i} + \omega_i \frac{d^2}{d^2\omega_i} + \frac{d}{d\omega_i} \sum_{i \neq j} \omega_j \frac{d}{d\omega_j}$$
 (26)

a second order differential on the Bargmann space generated by the set of complex variables $\omega_1, \dots, \omega_r$. A general vector

$$|\psi\rangle = \sum_{n_1,\dots,n_r} \psi_{n_1,\dots,n_r} |k; n_1,\dots,n_r\rangle$$
(27)

in the Fock space \mathcal{F} now is realized as follows

$$\psi(\omega_1,\cdots,\omega_r) = \sum_{n_1,\cdots,n_r} \psi_{n_1,\cdots,n_r} C_{k;n_1,\cdots,n_r} \omega_1^{n_1} \cdots \omega_r^{n_r}.$$
(28)

We define the inner product in this realization in the following form

$$\langle \psi' | \psi \rangle = \int d^2 \omega_1 \cdots d^2 \omega_r K(k; \omega_1, \cdots, \omega_r) \psi'^{\star}(\omega_1, \cdots, \omega_r) \psi(\omega_1, \cdots, \omega_r)$$
(29)

where $d^2\omega_i \equiv dRe\omega_i dIm\omega_i$ and the integration extends over the entire complex space $\mathbf{C}^{\mathbf{r}}$. To compute the measure function, appearing in the definition of the inner product (29), we identify $|\psi\rangle$ (resp. $|\psi'\rangle$) with the vector $|k; n_1, \dots, n_r\rangle$ (resp. $|k; n'_1, \dots, n'_r\rangle$). We also assume that it depends only on $\rho_i = |\omega_i|$ for $i = 1, \dots, r$. This hypothesis, usually used in the moment problems, is called the isotropy condition. It is a simple matter of computation to show that the function $K(k; \rho_1, \dots, \rho_r)$ satisfy the integral equation

$$(2\pi)^r \int_0^\infty \cdots \int_0^\infty d\rho_1 \cdots d\rho_r K(k;\rho_1,\cdots,\rho_r) |\rho_1|^{2n_1+1} \cdots |\rho_r|^{2n_r+1} = \frac{n_1! \cdots n_r! (k-1+n)!}{(k-1)!}.$$
(30)

A solution of this equation exists [22](see a nice proof in [23]) in term of the Bessel function

$$K(k;R) = \frac{2}{\pi^r (k-1)!} R^{k-r} K_{k-r}(2R)$$
(31)

where $R^2 = \rho_1^2 + \cdots + \rho_r^2$. Note that the analytic function $\psi(\omega_1, \cdots, \omega_r)$ can be viewed as the inner product of the ket $|\psi\rangle$ with a bra $\langle k; \omega_1^{\star}, \cdots, \omega_r^{\star}|$ labeled by the complex conjugate of the variables $\omega_1, \cdots, \omega_r$

$$\psi(\omega_1, \cdots, \omega_r) = \mathcal{N}\langle k; \omega_1^\star, \cdots, \omega_r^\star | \psi \rangle$$
(32)

where $\mathcal{N} \equiv \mathcal{N}(|\omega_1|, \cdots, |\omega_r|)$ stands for a normalization constant. As particular case, we take $|\psi\rangle = |k; n_1, \cdots, n_r\rangle$ to obtain

$$\langle k; \omega_1^{\star}, \cdots, \omega_r^{\star} | k; n_1, \cdots, n_r \rangle = \mathcal{N}^{-1} C_{k; n_1, \cdots, n_r} \omega_1^{n_1} \cdots \omega_r^{n_r}.$$
(33)

It follows that the normalized states $|k; \omega_1, \cdots, \omega_r\rangle$ are expressed as

$$|k;\omega_{1},\cdots,\omega_{r}\rangle = \mathcal{N}^{-1}\sum_{n_{1}=0}^{\infty}\cdots\sum_{n_{r}=0}^{\infty}\left[\frac{(k-1)!}{n_{1}!\cdots n_{r}!(k-1+n)!}\right]^{\frac{1}{2}}\omega_{1}^{n_{1}}\cdots\omega_{r}^{n_{r}}|k;n_{1},\cdots,n_{r}\rangle$$
(34)

where

$$\mathcal{N}^{2}(|\omega_{1}|,\cdots,|\omega_{r}|) = \sum_{n_{1}=0}^{\infty} \cdots \sum_{n_{r}=0}^{\infty} \frac{(k-1)!}{n_{1}!\cdots n_{r}!(k-1+n)!} |\omega_{1}|^{2n_{1}}\cdots |\omega_{r}|^{2n_{r}}.$$
 (35)

It may be noted that the states $|k; \omega_1, \dots, \omega_r\rangle$ are not orthogonal. It is also interesting to remark that they are eigenvectors of the Jacobson operators a_i^- with the eigenvalue ω_i . In this sense, the states $|k; \omega_1, \dots, \omega_r\rangle$ can be considered as Gazeau-Klauder coherent states associated with a quantum mechanical system whose Hamiltonian is given by (8) and obeying the A_r bosonic statistics. Finally, note that this first realization is possible because the Fock space is infinite dimension.

3.2 Realization II

There exists a second (non equivalent) Bargmann realization associated with A_r bosonic systems. In this realization the annihilation operators act as derivation with respect to complex variables z_i

$$a_i^- \longrightarrow \frac{d}{dz_i}$$
 (36)

belonging to the complex domain $\mathcal{D} = \{(z_1, z_2, \cdots, z_r) : |z_1|^2 + |z_2|^2 + \cdots + |z_r|^2 < 1\}$. The reason of this condition will explained in the sequel of this section. The elements of the Fock space are realized as follows

$$|k; n_1, \cdots, n_r\rangle \longrightarrow C_{k; n_1, \cdots, n_r} z_1^{n_1} \cdots z_r^{n_r}$$
(37)

Using the action of the annihilation operators on the Fock space \mathcal{F} and the correspondence (37), one obtain the following recursion formula

$$\sqrt{k-1+n_1+\dots+n_i+\dots+n_r}C_{k;n_1,\dots,n_i-1,\dots,n_r} = \sqrt{n_i}C_{k;n_1,\dots,n_i,\dots,n_r}$$
(38)

which can be solved in a similar manner that one used above (Eq. 21). As result, we obtain

$$C_{k;n_1,\cdots,n_i,\cdots,n_r} = \left[\frac{(k-1+n)!}{n_1!\cdots n_r!(k-1)!}\right]^{\frac{1}{2}}$$
(39)

where $n = n_1 + n_2 + \cdots + n_r$. Having the expression of the coefficients C, one can determine the differential action of the Jacobson creation operators. Indeed, using the actions of the generators a_i^+ on the Fock space, we show that

$$a_i^+ \longrightarrow kz_i + z_i \sum_{j=1}^r z_j \frac{d}{dz_j}.$$
 (40)

In this realization, the Jacobson generators act as first order linear differential operators. Here also, we realize a general vector of the Fock space \mathcal{F}

$$|\phi\rangle = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots \sum_{n_r=0}^{\infty} \phi_{n_1,n_2\cdots,n_r} |k;n_1,n_2,\cdots,n_r\rangle$$

as

$$\phi(z_1, z_2 \cdots, z_r) = \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots \sum_{n_r=0}^{\infty} \phi_{n_1, n_2 \cdots, n_r} C_{k; n_1, n_2 \cdots, n_r} z_1^{n_1} z_2^{n_2} \cdots z_r^{n_r}.$$
 (41)

The inner product of two functions ϕ and ϕ' is defined now as follows

$$\langle \phi' | \phi \rangle = \int \int \cdots \int d^2 z_1 d^2 z_2 \cdots d^2 z_r \Sigma(k; z_1, z_2 \cdots, z_r) \phi'^{\star}(z_1, z_2 \cdots, z_r) \phi(z_1, z_2 \cdots, z_r)$$
(42)

where the integration is carried out the complex domain \mathcal{D} . The computation of the integration measure Σ , assumed to be isotropic, can be performed by choosing $|\phi\rangle = |k; n_1, n_2, \dots, n_r\rangle$ and $|\phi'\rangle = |k; n'_1, n'_2, \dots, n'_r\rangle$. So, one has to look for a solution of the following moment equation

$$\int \int \cdots \int d\varrho_1 d\varrho_2 \cdots d\varrho_r \Sigma(k; \varrho_1, \varrho_2, \cdots, \varrho_r) \varrho_1^{2n_1+1} \varrho_2^{2n_2+1} \cdots \varrho_r^{2n_r+1} = \frac{n_1! n_2! \cdots n_r! (k-1)!}{(2\pi)^r (k-1+n)!}$$
(43)

where $n = n_1 + n_2 + \cdots + n_r$ and $\varrho_i = |z_i|$. To find the function satisfying the equation (43), we use following result

$$\int_{0}^{1} \varrho_{1}^{n_{1}} d\varrho_{1} \int_{0}^{1-\varrho_{1}} \varrho_{2}^{n_{2}} d\varrho_{2} \cdots \int_{0}^{1-\varrho_{1}-\varrho_{2}-\dots-\varrho_{r-1}} \varrho_{r}^{n_{r}} (1-\varrho_{1}-\varrho_{2}-\dots-\varrho_{r})^{k-r-1} d\varrho_{r}$$

$$= \frac{n_{1}!n_{2}!\cdots n_{r}!(k-1)!}{(k-1)!(k-r)(k-r+1)\cdots(k-1)}$$
(44)

which can be easily verified. The measure is then given by

$$\Sigma(k;\varrho_1,\varrho_2,\cdots,\varrho_r) = \pi^{-r} [1 - (\varrho_1^2 + \varrho_2^2 + \cdots + \varrho_r^2)]^{k-r-1} (k-r)(k-r+1)\cdots(k-1).$$
(45)

At this stage, one can write the function $\phi(z_1, z_2 \cdots, z_r)$ as the product of the state $|\phi\rangle$ with some ket $|k; z_1^*, z_2^* \cdots, z_r^*\rangle$ labeled by the complex conjugate of the variables z_1, z_2, \cdots, z_r .

$$\phi(z_1, z_2, \cdots, z_r) = \mathcal{N}\langle k; z_1^*, z_2^*, \cdots, z_r^* | \phi \rangle.$$
(46)

Taking $|\phi\rangle = |k; n_1, n_2, \cdots, n_r\rangle$, we obtain

$$|k; z_1, z_2, \cdots, z_r\rangle = \mathcal{N}^{-1} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \cdots \sum_{n_r=0}^{\infty} \left[\frac{(k-1+n)!}{n_1! \cdots n_r! (k-1)!} \right]^{\frac{1}{2}} z_1^{n_1} z_2^{n_2} \cdots z_r^{n_r} |k; n_1, n_2, \cdots, n_r\rangle.$$
(47)

The expansion in (47) converges when the the complex variables z_1, z_2, \dots, z_r are in the *r*dimensional unit ball $|z_1|^2 + |z_2|^2 + \dots + |z_r|^2 < 1$. This explain the above hypothesis according to which z_1, z_2, \dots, z_r should be in the complex domain \mathcal{D} . The normalization condition of the states (47) gives the following expression for the normalization constant

$$\mathcal{N} = (1 - |z_1|^2 - |z_2|^2 - \dots - |z_r|^2)^{\frac{k}{2}}.$$
(48)

It is important to remark that the states (47) are continuous in the labeling, constitute an over complete set and then are coherent in the Klauder-Perelomov sense [18,19]. These states can be used in the investigation of quantum Hall systems in higher dimensional ball [24,25].

4 Bargmann realization of A_r fermionic statistics

We are going to construct the analytic realization of the irreducible representation, related to A_r fermionic statistics (s = -1) characterized by the so-called the generalized Pauli principle. As we mentioned before, because the Fock space is finite dimension, the Jacobson creation operators can not be represented by a multiplication by some complex variable and only one analytic realization can be established. It corresponds to one in which the generators a_i^- act as

$$a_i^- \longrightarrow \frac{d}{d\zeta_i}$$
 (49)

in the space of the polynomials of the form $C_{k;n_1,n_2,\dots,n_r}\zeta_1^{n_1}\zeta_2^{n_2}\cdots\zeta_r^{n_r}$ defined in the *r*-dimensional space **C**^{**r**} of complex lines $(\zeta_1, \zeta_2, \dots, \zeta_r)$ with the correspondence

$$|k; n_1, n_2, \cdots, n_r\rangle \longrightarrow C_{k; n_1, n_2, \cdots, n_r} \zeta_1^{n_1} \zeta_2^{n_2} \cdots \zeta_r^{n_r}.$$
(50)

Following similar analysis as above, one can see that the coefficients in (50) satisfy the recurrence formula

$$\sqrt{n_i}C_{k;n_1,n_2,\cdots,n_i,\cdots,n_r} = \sqrt{k-n}C_{k;n_1,n_2,\cdots,n_i-1,\cdots,n_r}$$
(51)

where $n = n_1 + n_2 + \cdots + n_r$. Solution of this equation, for all $i = 1, 2, \cdots, r$, is given by

$$C_{k;n_1,n_2,\cdots,n_r} = \left[\frac{(k-1)!}{n_1!n_2!\cdots n_r!(k-1-r)}\right]^{\frac{1}{2}}$$
(52)

Using the action of the creation operators on the Fock space, it is not difficult to verify that they act

$$a_i^+ \longrightarrow (k-1)\zeta_i - \zeta_i \sum_{j=1}^r \zeta_j \frac{d}{d\zeta_i}$$
(53)

as first order differential operators. As in the previous realizations, the existence of a measure $\sigma(k; \zeta_1, \zeta_2, \dots, \zeta_r)$ ensures the definition of inner product of two functions. To determine this measure, one use the orthogonality of the Fock states $|k; n_1, n_2, \dots, n_r\rangle$. It follows that it satisfies

$$\int \int \cdots \int d^2 \zeta_1 d^2 \zeta_2 \cdots d^2 \zeta_r \sigma(k; \zeta_1, \zeta_2, \cdots, \zeta_r) C_{k;n_1, n_2, \cdots, n_r} C_{k;n_1', n_2', \cdots, n_r'} \zeta_1^{n_1} \zeta_1^{n_2} \zeta_2^{n_2'} \cdots \zeta_r^{n_r} \zeta_r^{n_r'}$$

$$= \delta_{n_1, n_1'} \delta_{n_2, n_2'} \cdots \delta_{n_r, n_r'}.$$
(54)

Setting $\zeta_i = |\zeta_i|e^{i\theta}$ and assuming the isotropy condition, the last equation becomes

$$\int_{0}^{\infty} \int_{0}^{\infty} \cdots \int_{0}^{\infty} dx_{1} dx_{2} \cdots dx_{r} \mu(k, x_{1}, x_{2}, \cdots, x_{r}) x_{1}^{n_{1}} x_{2}^{n_{2}} \cdots x_{r}^{n_{r}}$$
$$= \frac{n_{1}! n_{2}! \cdots n_{r}! (k-1-n)!}{(k-1)!}$$
(55)

where $\mu \equiv \pi^r \sigma$ and $x_i = |\zeta_i|^2$. Using the Mellin inverse transform [22], we obtain

$$\mu(k, x_1, x_2, \cdots, x_r) = \frac{(k-1+r)!}{(k-1)!} (1+x_1+x_2+\cdots+x_r)^{-(k+r)}.$$
(56)

Any function $f(\zeta_1, \zeta_2, \dots, \zeta_r)$ can be written in the following form

$$f(\zeta_1, \zeta_2, \cdots, \zeta_r) = \mathcal{N}\langle k; \zeta_1^*, \zeta_2^*, \cdots, \zeta_r^* | f \rangle$$
(57)

where $|f\rangle$ is a generic element of the Fock space with s = -1 and the normalization constant is given by

$$\mathcal{N}(|\zeta_1|^2, |\zeta_2|^2, \cdots, |\zeta_r|^2) = (1 + |\zeta_1|^2 + |\zeta_2|^2 + \cdots + |\zeta_r|^2)^{-\frac{k-1}{2}}.$$
(58)

It is remarkable to notice that the states $|k; \zeta_1, \zeta_2, \dots, \zeta_r\rangle$ are nothing but the coherent states for the complex projective space \mathbb{CP}^r which emerges, in a natural way, in realizing analytically the quantum states of a system obeying fermionic A_r statistics. It is also interesting to mention that the states $|k; \zeta_1, \zeta_2, \dots, \zeta_r\rangle$ were used in the description of quantum Hall systems in higher dimension complex projective spaces [24]. I this respect, the generalized quantum A_r statistics can be applied in the description of quantum Hall effect in \mathbb{CP}^r spaces.

5 Concluding remarks

The main ingredient for the present paper is the definition of generalized A_r statistics. We have obtained the Fock representation associated to bosonic (s = 1) and fermionic (s = -1) A_r statistics. We have developed the Bargmann realizations of Fock spaces and the differential action of Jacobson generators. We shown that the analytical representations lead to overcomplete sets identified as Klauder-Perelomov and Gazeau-Klauder coherent states associated with a quantum system described by the Hamiltonian (8). The present work can be extended to quantum statistics associated with classical Lie algebras B_r , C_r and D_r . It will also be interesting to study the macroscopic properties of such statistics. Another possible extension concerns the connection of the generalized A_r statistics and the quantum Hall effect in higher dimension spaces.

References

- [1] H.S. Green, Phys. Rev. 90 (1953) 270.
- [2] O.W. Greenberg and A.M.L. Messiah, Phys. Rev. B 138 (1965) 1155;
- J. Math. Phys. 6 (1965) 500.
- [3] Y. Ohnuki and S. Kamefuchi, *Quantum field theory and parastatistics* (Springer, Berlin, 1982).
- [4] J.M. Leinaas and J. Myrheim, Nuovo Cimento B 37 (1977) 1.
- [5] F. Wilczek, Phys. Rev. Lett. 49 (1982) 957.
- [6] T.D. Palev, Lie algebraic aspects of quantum statistics. Unitary quantization (A-quantization), Preprint JINR E17-10550 (1977) and hep-th/9705032.
- [7] O.W. Greenberg, Phys. Rev. Lett. 64 (1990) 705; Phys. Rev. D 43(1991) 4111.

[8] M. Daoud and M. Kibler, Phys. Lett. A 321 (2004) 147; Phys. Part. Nuclei (Suppl. 1) 33 (2002) S43.

[9] F.D.M. Haldane, Phys. Rev. Lett. 67 (1991) 937.

[10] T.D. Palev, Czech. J. Phys. B 29 (1979) 91; Rep. Math. Phys. 18 (1980) 177; Rep.

Math. Phys. 18 (1980) 129; J. Math. Phys 21 (1980) 1293.

[11] N.I. Stoilova and J. Van der Jeugt, preprint math-phys/0409002.

[12] S. Kamefuchi and Y. Takahashi, Nucl. Phys. 36 (1962) 177; C. Ryan and E.C.G. Sudarshan, Nucl. Phys. 47 (1963) 207.

- [13] V.G Kac, Adv. Math. 26 (1977) 8.
- [14] R.B. Laughlin, Phys. Rev. Lett. 50 (1983) 1395.

[15] B.I. Halperin, Phys. Rev. Lett. 52 (1984) 1583.

[16] R.B. Laughlin, Phys. Rev. Lett. 60 (1988) 2677.

[17] A. Strominger, Phys. Rev. Lett. 71 (1993) 3397.

[18] J.R. Klauder and B.S Skagerstam, *Coherent states-Applications in Physics and Mathematical Physics* (World Scientific, Singapore, 1985).

[19] A. Perelomov, *Generalized Coherent States and their Applications*, Texts and Monographs in Physics, (Spinger-Verlag, 1986).

[20] N. Jacobson, Amer. J. Math. 71 (1949) 149.

[21] J.R. Klauder, J. Phys. A: Math. Gen.29 (1996) L 293; J-P. Gazeau and J-R. Klauder,
 J. Phys. A: Math. Gen.32 (1999) 123.

 $[\mathbf{22}]$ H. Bateman, Table of integral transforms, Vol $\underline{1}$ (1954), Ed A. Erdélyi (New York, Mc Graw Hill) .

[23] K. Fujii and K. Funahashi, preprint quant-phys/9704011.

[24] D. Karabali and V.P. Nair, Nucl. Phys. B 641(2002) 533; Nucl. Phys. B 679 (2004) 427.

[25] A. Jellal, Nucl. Phys. B 725 (2005) 554.