

Strict quantization of coadjoint orbits

N.P. Landsman*

Korteweg-de Vries Institute for Mathematics
University of Amsterdam
Plantage Muidergracht 24
1018 TV AMSTERDAM, THE NETHERLANDS

email: npl@wins.uva.nl

August 16, 2018

Abstract

A strict quantization of a compact symplectic manifold S on a subset $I \subseteq \mathbb{R}$, containing 0 as an accumulation point, is defined as a continuous field of C^* -algebras $\{\mathfrak{A}_{\hbar}\}_{\hbar \in I}$, with $\mathfrak{A}_0 = C_0(S)$, and a set of continuous cross-sections $\{\mathcal{Q}(f)\}_{f \in C^\infty(S)}$ for which $\mathcal{Q}_0(f) = f$. Here $\mathcal{Q}_{\hbar}(f^*) = \mathcal{Q}_{\hbar}(f)^*$ for all $\hbar \in I$, whereas for $\hbar \rightarrow 0$ one requires that $i[\mathcal{Q}_{\hbar}(f), \mathcal{Q}_{\hbar}(g)]/\hbar \rightarrow \mathcal{Q}_{\hbar}(\{f, g\})$ in norm. We discuss general conditions which guarantee that a (deformation) quantization in a more physical sense leads to one in the above sense.

Using ideas of Berezin, Lieb, Simon, and others, we construct a strict quantization of an arbitrary integral coadjoint orbit \mathcal{O}_λ of a compact connected Lie group G , associated to a highest weight λ . Here $I = 0 \cup 1/\mathbb{N}$, so that $\hbar = 1/k$, $k \in \mathbb{N}$, and $\mathfrak{A}_{1/k}$ is defined as the C^* -algebra of all matrices on the finite-dimensional Hilbert space $\mathcal{V}_{k\lambda}$ carrying the irreducible representation $U_{k\lambda}(G)$ with highest weight $k\lambda$. The quantization maps $\mathcal{Q}_{1/k}(f)$ are constructed from coherent states in $\mathcal{V}_{k\lambda}$, and have the special feature of being positive maps.

1998 PACS: 02.20Qs, 03.65Db, 02.40Vh, 02.30Tb

*Supported by a fellowship from the Royal Netherlands Academy of Arts and Sciences (KNAW)

I. Introduction

The aim of this paper is to combine a number of what appear to the author to be good ideas in mathematical physics, whose interplay has so far not sufficiently been studied. Firstly, there now exists a satisfying C^* -algebraic definition of quantization, which enables one to link Poisson and symplectic geometry [1, 2] with non-commutative geometry [3]. In particular, the geometric theory of classical mechanics and reduction is thereby related to the C^* -algebraic formulation of quantum mechanics and induction [4]. The main mathematical idea of this definition goes back to Rieffel [5], who showed how deformation quantization makes sense in an operator-algebraic context. The physical postulates, of course, may be traced back to Dirac. In this definition, like in conventional ('formal') deformation quantization, it is crucial that one studies the quantum theory for a family of values of Planck's constant \hbar [6].

Secondly, one has the well-known connection between coadjoint orbits of certain Lie groups, unitary representations, and geometric quantization [7]. This connection works particularly well for either compact or exponential nilpotent Lie groups. Here one keeps \hbar fixed.

Our third source of inspiration is the work of Lieb [8], Simon [9], and others on the classical limit of quantum spin systems and their generalizations to arbitrary compact Lie groups. This work is closely related to Perelomov's coherent states [10, 11], as well as to Berezin's approach to quantization and the classical limit [6].

Historical comments and extensive references concerning these ideas may be found in [4].

Using the second and the third group of ideas at both a conceptual and a technical level, we will construct a C^* -algebraic quantization of an arbitrary integral coadjoint orbit of a compact connected Lie group. A different approach to this problem has recently been considered in [12]; also see [13].

Section 2 contains a general C^* -algebraic definition of quantization. A quantization satisfying this definition is called 'strict'. We present criteria on a given quantization which guarantee that the postulates in the definition are met. It is on this basis that we will actually construct the quantizations in this paper. Section 3 describes our approach to coherent states and Berezin quantization, and develops conditions under which the Berezin quantization constructed from a family of coherent states is strict. This is done in terms of a so-called pure state quantization, which, in a heuristic sense, is dual to a C^* -algebraic quantization. The material in sections 2 and 3 is model-independent, and should be relevant to quantization theory in general.

Section 4 is a brief review of Perelomov coherent states, coadjoint orbits of compact Lie groups, the momentum map, and the connection between these concepts. The material in this section is not new, but is worth summarizing in preparation for our main results. Section 5 contains the two principal theorems of this paper. The first states that Perelomov's coherent states for compact connected Lie groups define pure state quantizations of particular coadjoint orbits. The second states that such pure state quantizations lead to strict quantizations of the coadjoint orbits in question.

II. On quantization

The central notion in C^* -algebraic quantization theory is that of a continuous field of C^* -algebras [14]. For our purposes the following reformulation is useful [15].

Definition 1 *A continuous field of C^* -algebras $(\mathfrak{C}, \{\mathfrak{A}_x, \varphi_x\}_{x \in X})$ over a locally compact Hausdorff space X consists of a C^* -algebra \mathfrak{C} , a collection of C^* -algebras $\{\mathfrak{A}_x\}_{x \in X}$, and a set $\{\varphi_x : \mathfrak{C} \rightarrow \mathfrak{A}_x\}_{x \in X}$ of surjective $*$ -homomorphisms, such that for all $A \in \mathfrak{C}$:*

1. *the function $x \rightarrow \|\varphi_x(A)\|$ is in $C_0(X)$;*
2. *one has $\|A\| = \sup_{x \in X} \|\varphi_x(A)\|$;*
3. *for any $f \in C_0(X)$ there is an element $fA \in \mathfrak{C}$ for which $\varphi_x(fA) = f(x)\varphi_x(A)$ for all $x \in X$.*

Here $C_0(X)$ is the space of continuous functions on X which vanish at infinity. Note that it is sufficient that each φ_x has dense image, since the image of a $*$ -homomorphism between C^* -algebras is automatically closed. The continuous cross-sections of the field in the sense of [14] consist of those elements $\{A_x\}_{x \in X}$ of $\prod_{x \in X} \mathfrak{A}_x$ for which there is a $A \in \mathfrak{C}$ such that $A_x = \varphi_x(A)$ for all $x \in X$.

The following definition of quantization, which is a slight reformulation of a definition in [16], seems to combine the best of previous definitions in this direction in [5] and [17].

Definition 2 *Let $I \subseteq \mathbb{R}$ contain 0 as an accumulation point. A strict quantization of a Poisson manifold P on I consists of*

1. a continuous field of C^* -algebras $(\mathfrak{C}, \{\mathfrak{A}_\hbar, \varphi_\hbar\}_{\hbar \in I})$ over I , with $\mathfrak{A}_0 = C_0(P)$;
2. a dense $*$ -subalgebra $\tilde{\mathfrak{A}}_0$ of $C_0(P)$ on which the Poisson bracket is defined, and which is closed under taking Poisson brackets (so that $\tilde{\mathfrak{A}}_0$ is a complex Poisson algebra);
3. a linear map $\mathcal{Q} : \tilde{\mathfrak{A}}_0 \rightarrow \mathfrak{C}$ which (with $\mathcal{Q}_\hbar(f) \equiv \varphi_\hbar(\mathcal{Q}(f))$) for all $f \in \tilde{\mathfrak{A}}_0$ and $\hbar \in I$ satisfies

$$\mathcal{Q}_0(f) = f, \quad (1)$$

$$\mathcal{Q}_\hbar(f^*) = \mathcal{Q}_\hbar(f)^*, \quad (2)$$

and for all $f, g \in \tilde{\mathfrak{A}}_0$ satisfies Dirac's condition

$$\lim_{\hbar \rightarrow 0} \left\| \frac{i}{\hbar} [\mathcal{Q}_\hbar(f), \mathcal{Q}_\hbar(g)] - \mathcal{Q}_\hbar(\{f, g\}) \right\| = 0. \quad (3)$$

Elements of I are interpreted as possible values of Planck's constant \hbar , and \mathfrak{A}_\hbar is the quantum algebra of observables of the theory at the given value of $\hbar \neq 0$. For real-valued f , the operator $\mathcal{Q}_\hbar(f)$ is the quantum observable associated to the classical observable f . This interpretation is possible because of condition (2) in Definition 2.

In the examples of the present paper, the Poisson manifold P will be a compact symplectic manifold S , and we will choose $\tilde{\mathfrak{A}}_0 = C^\infty(S)$. It will be a special feature that \mathcal{Q} is defined on all of \mathfrak{A}_0 ; the condition (3) of course makes sense on $\tilde{\mathfrak{A}}_0$ only.

The connection between Definition 2 and the more physically and historically oriented definition of quantization proposed in [17, 4] is established in Proposition 1 below. This requires two lemmas, the first of which, as a bonus, entails the equivalence between the definition of a continuous field given above, and the one given by Dixmier [14] (restricted to the case that the base space is locally compact).

Lemma 1 *The C^* -algebra \mathfrak{C} of (sections of) a continuous field is locally uniformly closed. That is, if $A \in \prod_x \mathfrak{A}_x$ is such that for every $y \in X$ and every $\epsilon > 0$ there exists a $B^y \in \mathfrak{C}$ and a neighbourhood \mathcal{N}^y of y in which $\|A_x - B_x^y\| < \epsilon$ for all $x \in \mathcal{N}^y$, and also $\lim_{x \rightarrow \infty} \|A_x\| = 0$, then $A \in \mathfrak{C}$.*

Alternatively, if the function $x \rightarrow \|A_x - C_x\|$ lies in $C_0(X)$ for each $C \in \mathfrak{C}$, then $A \in \mathfrak{C}$.

In the situation of the first part, there is a compact set $K \subseteq X$ for which $\|A_x\| < \epsilon$ outside K , as well as a finite subcover $\{\mathcal{N}^{x_1}, \dots, \mathcal{N}^{x_n}\}$ of K . Taking a partition of unity $\{u_i\}$ on K subordinate to this subcover, the operator $B = \sum_i u_i B^{x_i}$ lies in \mathfrak{C} because of Definition 1.3, and satisfies $\sup_{x \in X} \|A_x - B_x\| < \epsilon$. Hence $A \in \mathfrak{C}$ by Definition 1.2 and the completeness of \mathfrak{C} .

Given any $A \in \prod_x \mathfrak{A}_x$ and $y \in X$, because φ_y is surjective there is a $B^y \in \mathfrak{C}$ so that $A_y = B_y^y$. The assumption in the second part of the lemma then implies that the conditions in the first part are satisfied, so that $A \in \mathfrak{C}$. ■

The next lemma adapts Props. 10.2.3 and 10.3.2 in [14], which relate to Dixmier's own definition of a continuous field, to Definition 1.

Lemma 2 *Suppose one has a family $\{\mathfrak{A}_x\}_{x \in X}$ of C^* -algebras indexed by a locally compact Hausdorff space X , as well as a subset $\mathfrak{C} \subseteq \prod_x \mathfrak{A}_x$ which satisfies the following conditions:*

1. the set $\{A_x \mid A \in \tilde{\mathfrak{C}}\}$ is dense in \mathfrak{A}_x for each $x \in X$;
2. the function $x \rightarrow \|A_x\|$ is in $C_0(X)$ for each $A \in \tilde{\mathfrak{C}}$;
3. the set $\tilde{\mathfrak{C}}$ is closed under pointwise scalar multiplication, addition, adjointing, and operator multiplication.

There exists a unique continuous field of C^* -algebras $(\mathfrak{C}, \{\mathfrak{A}_x, \varphi_x\}_{x \in X})$ whose collection of continuous cross-sections contains $\tilde{\mathfrak{C}}$. Firstly, as a set \mathfrak{C} consists of all $A \in \prod_x \mathfrak{A}_x$ for which the function $x \rightarrow \|A_x - C_x\|$ lies in $C_0(X)$ for each $C \in \tilde{\mathfrak{C}}$. This set is regarded as a C^* -algebra under the pointwise operations listed in item 3 above, and the norm defined in Definition 1.2. Secondly, $\varphi_x(A) = A_x$ is the evaluation map.

We first show that if $A \in \prod_x \mathfrak{A}_x$ is such that for every $x_0 \in X$ and every $\epsilon > 0$ there exists a $B \in \mathfrak{C}$ and a neighbourhood \mathcal{N} of x_0 such that $\|A_x - B_x\| < \epsilon$ for all $x \in \mathcal{N}$, and also $\lim_{x \rightarrow \infty} \|A_x\| = 0$, then $A \in \mathfrak{C}$. Indeed, take $C \in \tilde{\mathfrak{C}}$ arbitrary, and define the functions $f_{AC} : x \rightarrow \|A_x - C_x\|$ and $f_{BC} : x \rightarrow \|B_x - C_x\|$. Using the inequality

$$|(\|X\| - \|Y\|)| \leq \|X - Y\|, \quad (4)$$

one finds $|f_{AC}(x) - f_{BC}(x)| < \epsilon$ for all $x \in \mathcal{N}$. By assumption, f_{BC} is continuous, so that $|f_{BC}(x) - f_{BC}(x_0)| < \epsilon$ for all x in some neighbourhood \mathcal{N}' of x_0 . Combining the two inequalities yields $|f_{AC}(x) - f_{AC}(x_0)| < 3\epsilon$ for all $x \in \mathcal{N} \cap \mathcal{N}'$. Hence f_{AC} is continuous at x_0 , which was arbitrary, so that $A \in \mathfrak{C}$ by definition of \mathfrak{C} .

Using this property, it is easily shown that \mathfrak{C} is a C^* -algebra, and that condition 3 in Definition 1 is satisfied. It is clear from Definition 1.1 and the definition of \mathfrak{C} that \mathfrak{C} is maximal. On the other hand, according to the second part of Lemma 1, \mathfrak{C} is minimal, so that it is unique. \blacksquare

Adding an assumption satisfied by the examples in this paper, we are now in a position to relate Definitions 1 and 2.

Proposition 1 *Suppose one has a Poisson manifold P and a family $\{\mathfrak{A}_{\hbar}\}_{\hbar \in I}$ of C^* -algebras, with $\tilde{\mathfrak{A}}_0 \subset \mathfrak{A}_0 = C_0(P)$ as in Definition 2, and a collection of linear maps $\{\mathcal{Q}_{\hbar} : \tilde{\mathfrak{A}}_0 \rightarrow \mathfrak{A}_{\hbar}\}_{\hbar \in I}$ satisfying (1), (2), (3), as well as*

$$\lim_{\hbar \rightarrow 0} \|\frac{1}{2}(\mathcal{Q}_{\hbar}(f)\mathcal{Q}_{\hbar}(g) + \mathcal{Q}_{\hbar}(g)\mathcal{Q}_{\hbar}(f)) - \mathcal{Q}_{\hbar}(fg)\| = 0; \quad (5)$$

$$\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f)\| = \|f\|_{\infty} \quad (6)$$

for all $f, g \in \tilde{\mathfrak{A}}_0$, and finally the completeness condition that the collection $\{\mathcal{Q}_{\hbar}(f) \mid f \in \tilde{\mathfrak{A}}_0\}$ be dense in \mathfrak{A}_{\hbar} for each $\hbar \in I$. Furthermore, assume that \hbar assumes discrete values.

There exists a unique continuous field of C^* -algebras $(\mathfrak{C}, \{\mathfrak{A}_{\hbar}, \varphi_{\hbar}\}_{\hbar \in I})$ whose collection of continuous cross-sections $\{\varphi_{\hbar}(A)\}_{\hbar \in I}$, $A \in \mathfrak{C}$, contains all maps $\{\mathcal{Q}_{\hbar}(f)\}_{\hbar \in I}$, $f \in \tilde{\mathfrak{A}}_0$.

One defines $\tilde{\mathfrak{C}} \subset \prod_{\hbar} \mathfrak{A}_{\hbar}$ as the complex linear span of all expressions of the form $\hbar \rightarrow \mathcal{Q}_{\hbar}(f_1) \dots \mathcal{Q}_{\hbar}(f_n)$, where $f_i \in \tilde{\mathfrak{A}}_0$.

Given that I is discrete, the continuity of each function $F_{f_1, \dots, f_n} : \hbar \rightarrow \|\mathcal{Q}_{\hbar}(f_1) \dots \mathcal{Q}_{\hbar}(f_n)\|$ away from 0 is trivial. Now note that (5) and (3) imply that

$$\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f)\mathcal{Q}_{\hbar}(g) - \mathcal{Q}_{\hbar}(fg)\| = 0 \quad (7)$$

for all $f, g \in \tilde{\mathfrak{A}}_0$. Hence $\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f_1)\mathcal{Q}_{\hbar}(f_2) \dots \mathcal{Q}_{\hbar}(f_n) - \mathcal{Q}_{\hbar}(f_1 \dots f_n)\| = 0$, and by induction $\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f_1) \dots \mathcal{Q}_{\hbar}(f_n) - \mathcal{Q}_{\hbar}(f_1 \dots f_n)\| = 0$. Eq. (4) then yields $\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f_1) \dots \mathcal{Q}_{\hbar}(f_n)\| - \|\mathcal{Q}_{\hbar}(f_1 \dots f_n)\| = 0$, so that finally $\lim_{\hbar \rightarrow 0} \|\mathcal{Q}_{\hbar}(f_1) \dots \mathcal{Q}_{\hbar}(f_n)\| = \|f_1 \dots f_n\|_{\infty}$ by (6). Because of (1), this shows that F_{f_1, \dots, f_n} is continuous at $\hbar = 0$.

Hence one is in the situation of Lemma 2, and the claim follows. \blacksquare

Although it is irrelevant for the present paper, we note that Proposition 1 equally well holds (with a different proof) if the set I is not discrete, provided that all \mathfrak{A}_{\hbar} are identical for $\hbar \neq 0$, and in addition the function $\hbar \rightarrow \mathcal{Q}_{\hbar}(f)$ is continuous for each $f \in \tilde{\mathfrak{A}}_0$ [4].

III. Quantization of pure states and Berezin quantization

A Hilbert space \mathcal{H} is a symplectic manifold, with symplectic form $\omega(\Phi, \Omega) = 2\text{Im}(\Phi, \Omega)$ (here the inner product (\cdot, \cdot) on \mathcal{H} is linear in the second entry) [1, 2]. This form is invariant under the standard action $\exp(i\alpha) : \Psi \rightarrow \exp(i\alpha)\Psi$ of $U(1)$ on \mathcal{H} , so that the quotient $\mathcal{H}^*/U(1)$ is a Poisson manifold (here $\mathcal{H}^* = \mathcal{H} \setminus \{0\}$). The symplectic leaves [2] of $\mathcal{H}^*/U(1)$ are the spaces $S_r = \mathcal{H}_r/U(1)$, where $\mathcal{H}_r = \{\Psi \in \mathcal{H} \mid (\Psi, \Psi) = r^2\}$. In particular, the projective space $\mathbb{P}\mathcal{H}$ may be identified with S_1 , and is therefore a symplectic manifold, with symplectic form $\omega_{\mathbb{P}\mathcal{H}}$.

In addition, $\mathbb{P}\mathcal{H}$ is equipped with a transition probability $p : \mathbb{P}\mathcal{H} \times \mathbb{P}\mathcal{H} \rightarrow [0, 1]$, given by $p(\psi, \varphi) = |(\Psi, \Phi)|^2$; here Ψ and Φ are arbitrary lifts of ψ and φ to unit vectors in \mathcal{H} . Equipped with these transition probabilities and with the Poisson bracket defined by $\omega_{\mathbb{P}\mathcal{H}}$, the manifold $\mathbb{P}\mathcal{H}$ is the pure state space of a quantum system without superselection rules. See [4] and refs. therein.

The pure state space of a classical system is a symplectic manifold (S, ω_S) , supporting the Liouville measure μ_L . Such a classical pure state space may be seen as carrying the ‘classical’ transition probability p_0 , defined by $p_0(\rho, \sigma) = \delta_{\rho\sigma}$.

Definition 3 *Let $I \subseteq \mathbb{R}$ be as in Definition 2, and put $I_0 = I \setminus \{0\}$. A pure state quantization of a symplectic manifold (S, ω) consists of a collection of Hilbert spaces $\{\mathcal{H}_{\hbar}\}_{\hbar \in I_0}$ and a collection of smooth injections $\{q_{\hbar} : S \rightarrow \mathbb{P}\mathcal{H}_{\hbar}\}_{\hbar \in I_0}$, for which the following requirements are satisfied.*

1. *There exists a positive function $c : I_0 \rightarrow \mathbb{R} \setminus \{0\}$ such that for all $\hbar \in I_0$ and all $\psi \in \mathbb{P}\mathcal{H}_{\hbar}$ one has*

$$c(\hbar) \int_S d\mu_L(\sigma) p(q_{\hbar}(\sigma), \psi) = 1. \quad (8)$$

2. *For all fixed $f \in C_c(S)$ and $\rho \in S$ the function*

$$\hbar \rightarrow \int_S d\mu_L(\sigma) p(q_{\hbar}(\rho), q_{\hbar}(\sigma)) f(\sigma)$$

is continuous on I_0 and satisfies

$$\lim_{\hbar \rightarrow 0} c(\hbar) \int_S d\mu_L(\sigma) p(q_{\hbar}(\rho), q_{\hbar}(\sigma)) f(\sigma) = f(\rho). \quad (9)$$

3. *for each $\hbar \in I_0$, the map q_{\hbar} is a symplectomorphism, that is,*

$$q_{\hbar}^* \omega_{\mathbb{P}\mathcal{H}} = \omega_S. \quad (10)$$

Since $f \in C_c(S)$, the continuity requirement on I_0 in this condition is equivalent to the continuity of the function $\hbar \rightarrow p(q_{\hbar}(\rho), q_{\hbar}(\sigma))$ for fixed ρ and σ . Using Urysohn’s lemma, it is not difficult to show [4] that (9) and (8) imply

$$\lim_{\hbar \rightarrow 0} p(q_{\hbar}(\rho), q_{\hbar}(\sigma)) = \delta_{\rho\sigma}. \quad (11)$$

That is, in quantizing pure states the quantum-mechanical transition probabilities should converge to the classical ones for $\hbar \rightarrow 0$.

More generally, one may replace the Liouville measure μ_L in Definition 3 by a family of measures μ_{\hbar} , and substitute an appropriate limiting condition for (10); see [4]. This generality is not needed for the present paper. The author is indebted to E. Hawkins for this remark.

A pure state quantization naturally leads to the quantization of observables.

Definition 4 *Let $\{\mathcal{H}_{\hbar}, q_{\hbar}\}_{\hbar \in I_0}$ be a pure state quantization of a symplectic manifold S . The Berezin quantization of a function $f \in L^\infty(S)$ is the family of operators $\{Q_{\hbar}^B(f)\}_{\hbar \in I_0}$, where $Q_{\hbar}^B(f) \in \mathfrak{B}(\mathcal{H}_{\hbar})$ is defined by the weak integral*

$$Q_{\hbar}^B(f) = c(\hbar) \int_S d\mu_L(\sigma) f(\sigma) [q_{\hbar}(\sigma)]. \quad (12)$$

Here $[q_{\hbar}(\sigma)]$ is the projection onto the one-dimensional subspace in \mathcal{H}_{\hbar} whose image in $\mathbb{P}\mathcal{H}_{\hbar}$ is $q_{\hbar}(\sigma)$. In case that $f \in L^1(S) \cap L^\infty(S)$, the integral is an ordinary Lebesgue integral (with values in a Banach space).

A number of properties of \mathcal{Q}_\hbar^B are immediately evident (cf. [9]): \mathcal{Q}_\hbar^B is positive (that is, $f \geq 0$ almost everywhere on S implies $\mathcal{Q}_\hbar^B(f) \geq 0$ in $\mathfrak{B}(\mathcal{H}_\hbar)$), and if f is real-valued then $\mathcal{Q}_\hbar^B(f)$ is self-adjoint. Moreover, $\mathcal{Q}_\hbar^B(f)$ is bounded, with

$$\| \mathcal{Q}_\hbar^B(f) \| \leq \| f \|_\infty . \quad (13)$$

To prove the last property, let A be a bounded symmetric operator such that $|(\Psi, A\Psi)| \leq c \| \Psi \|^2$ for some $c > 0$, and all Ψ . One then replaces Ψ by $\Psi \pm A\Psi/c$, and subtracts the two inequalities thus obtained. This implies the inequality $\| A\Psi \| \leq c \| \Psi \|$, showing that A is bounded with norm $\leq c$. This argument with (8) implies (13).

A Berezin quantization is not necessarily strict when restricted to, say, $C_c^\infty(S)$; even (10) does not imply (3). At the present abstract level, all that can be inferred is the following.

Proposition 2 *Let $f \in C_0(S)$, and assume that I is discrete. Then*

$$\lim_{\hbar \rightarrow 0} \| \mathcal{Q}_\hbar^B(f) \| = \| f \|_\infty . \quad (14)$$

We initially assume that $f \in L^1(S) \cap C_0(S)$, and at the end extend the result to $f \in C_0(S)$ using the continuity of \mathcal{Q}_\hbar^B . Eq. (13) implies

$$\limsup_{\hbar \rightarrow 0} \| \mathcal{Q}_\hbar^B(f) \| \leq \| f \|_\infty . \quad (15)$$

On the other hand, for $f \in C_0(S)$ we can find $\rho \in S$ for which $\| f \|_\infty = |f(\rho)|$. Now for any unit vector $\Psi \in \mathcal{H}_\hbar$, eq. (12) implies that

$$(\Psi, \mathcal{Q}_\hbar^B(f)\Psi) = c(\hbar) \int_S d\mu_L(\sigma) p(q_\hbar(\sigma), \psi) f(\sigma), \quad (16)$$

where ψ is the projection of Ψ to $\mathbb{P}\mathcal{H}_\hbar$. We now use the obvious inequality $\| \mathcal{Q}_\hbar^B(f) \| \geq |(\Psi, \mathcal{Q}_\hbar^B(f)\Psi)|$, take Ψ to be a lift of $q_\hbar(\rho)$, and use (9) to find

$$\liminf_{\hbar \rightarrow 0} \| \mathcal{Q}_\hbar^B(f) \| \geq \| f \|_\infty . \quad (17)$$

Combining this with (15), eq. (14) follows. ■

IV. Coadjoint orbits and Perelomov's coherent states

Let G be a Lie group, with Lie algebra \mathfrak{g} . The dual \mathfrak{g}^* of \mathfrak{g} is a Poisson manifold under the so-called Lie-Poisson bracket [2, 4]

$$\{f, g\}(\theta) = -\theta([df_\theta, dg_\theta]); \quad (18)$$

here the differential df_θ of $f \in C^\infty(\mathfrak{g}^*)$ at $\theta \in \mathfrak{g}^*$, which is a linear map from $T_\theta\mathfrak{g}^* \simeq \mathfrak{g}^*$ to \mathbb{R} , is identified with an element of $\mathfrak{g} \simeq \mathfrak{g}^{**}$, so that the right-hand side of (18) is the Lie bracket in \mathfrak{g} .

The symplectic leaves of \mathfrak{g}^* with respect to the Lie-Poisson structure are the coadjoint orbits of G [2, 4]. This endows each coadjoint orbit \mathcal{O} with the so-called Lie-Kirillov symplectic structure. It is clear that $\theta \in \mathfrak{g}^*$ satisfies $\theta([X, Y]) = 0$ for all $X, Y \in \mathfrak{g}_\theta$, where \mathfrak{g}_θ is the Lie algebra of the stabilizer G_θ of θ under the coadjoint action. In other words, $\theta : \mathfrak{g}_\theta \rightarrow \mathbb{R}$ is a Lie algebra homomorphism.

Definition 5 *A coadjoint orbit $\mathcal{O} \in \mathfrak{g}^*$ is called integral if for some (hence all) $\theta \in \mathcal{O}$ the functional $\theta \upharpoonright \mathfrak{g}_\theta$ exponentiates to a character of G_θ .*

In other words, θ is integral iff there is a character U_θ of G_θ such that $\theta = idU_\theta$ on \mathfrak{g}_θ . If this holds for one $\theta \in \mathcal{O}$ it holds for all, since one has $U_{\text{Co}(x)\theta} = U_\theta \circ \text{Ad}(x^{-1})$. Here and in what follows, Co and Ad stand for the coadjoint and the adjoint action of G on \mathfrak{g}^* and on \mathfrak{g} , respectively.

In the remainder of this paper, G is a compact connected Lie group. We assume familiarity with the standard Cartan-Weyl description of the unitary irreducible representations of G [18].

We fix a maximal torus $T \subset G$. The coadjoint orbits of T in the dual \mathfrak{t}^* of the Lie algebra \mathfrak{t} of T are points. The integral coadjoint orbits of T form the weight lattice $\Lambda \subset \mathfrak{t}^*$. The Weyl group $W = N(T)/T$ (where $N(T)$ is the normalizer of T) acts on T by conjugation. The derivative of this action is a W -action on \mathfrak{t} , whose dual action on \mathfrak{t}^* is the projection of the coadjoint action of $N(T)$ to $N(T)/T$. This action maps the weight lattice Λ into itself.

A functional $\lambda \in \mathfrak{t}^*$ is regular when $w\lambda = \lambda$ for $w \in W$ implies $w = e$ (and singular otherwise); this defines the sets \mathfrak{t}_r^* and $\Lambda_r = \mathfrak{t}_r^* \cap \Lambda$ of regular elements and regular weights in \mathfrak{t}^* , respectively. A Weyl chamber is a connected component C of \mathfrak{t}_r^* , and thereby forms an open convex cone in \mathfrak{t}^* . Singular weights clearly lie on the boundary of some Weyl chamber. One singles out an arbitrary Weyl chamber C_d , and declares a weight dominant if it lies in the closure \overline{C}_d . The point is now that each W -orbit intersects a given closed Weyl chamber \overline{C} in exactly one point.

Combining the Cartan-Weyl theory with that of Kostant [7], one obtains a number of parametrizations of the unitary dual of G (i.e., the set of unitary irreducible representations of G modulo unitary equivalence).

Proposition 3 *There exist bijective correspondences between the unitary dual \hat{G} of G , the set of W -orbits in Λ , the set $\Lambda_d = \Lambda \cap \overline{C}_d$ of dominant weights, and the set of integral coadjoint orbits in \mathfrak{g}^* . The latter set is isomorphic to \mathfrak{t}^*/W . In other words, one has*

$$\hat{G} \simeq \Lambda/W \simeq \Lambda_d \simeq (\mathfrak{g}^*/W)_{\text{integral}}. \quad (19)$$

Though exceedingly well-known, we recall the explicit form of the bijection $\hat{G} \simeq \Lambda_d$ in Proposition 3, as it will play an important role in the proof of our main result. We use the standard Cartan-Weyl basis $\{H_j\}_{j=1,\dots,r} \cup \{E_\alpha, E_{-\alpha}\}_{\alpha \in \Delta^+}$ of $\mathfrak{g}_{\mathbb{C}}$, where r is the rank of G , and Δ^+ is the set of positive roots (relative to a choice of C_d). The vectors H_i lie in \mathfrak{t} , and the $E_{\pm\alpha}$ are eigenvectors of each H_i under restriction from \mathfrak{g} to \mathfrak{t} of the adjoint representation of \mathfrak{g} on $\mathfrak{g}_{\mathbb{C}}$.

A Hilbert space \mathcal{V}_λ carrying an irreducible representation $U_\lambda(G)$ corresponding to a dominant weight $\lambda \in \Lambda_d$ has a highest weight vector Ψ_λ of norm 1, unique up to a phase, such that

$$dU_\lambda(H_j)\Psi_\lambda = -i\lambda(H_j)\Psi_\lambda \quad (20)$$

for $j = 1, \dots, r$, whereas for all $\alpha \in \Delta^+$ one has

$$dU_\lambda(E_\alpha)\Psi_\lambda = 0. \quad (21)$$

Since $dU_\lambda(E_\alpha)^* = -dU_\lambda(E_{-\alpha})$, it follows that

$$(\Psi_\lambda, dU_\lambda(E_\alpha)\Psi_\lambda) = 0 \quad (22)$$

for all $\alpha \in \Delta$.

The association of a coadjoint orbit in \mathfrak{g}^* with a W -orbit in \mathfrak{t}^* is as follows. We write $\mathfrak{t}_{\mathbb{C}}^\perp$ for the span of all $E_{\pm\alpha}$, and $\mathfrak{t}^\perp = \mathfrak{t}_{\mathbb{C}}^\perp \cap \mathfrak{g}$. Given a weight $\lambda \in \mathfrak{t}^*$, putting $\theta(\lambda) = 0$ on \mathfrak{t}^\perp and $\theta(\lambda) = \lambda$ on \mathfrak{t} , one obtains an extension $\theta(\lambda) \in \mathfrak{g}^*$ of λ . Thus the coadjoint orbit $\mathcal{O}_\lambda = \mathcal{O}_{\theta(\lambda)}$ associated to λ is the coadjoint orbit through $\theta(\lambda)$; it is obvious from the definition of the W -action on \mathfrak{t}^* that all points of W -orbit of λ are mapped into \mathcal{O}_λ . We will use the label λ to denote a dominant weight in $\Lambda_d \subset \mathfrak{t}^*$, as well as the corresponding element $\theta(\lambda)$ in the coadjoint orbit $\mathcal{O}_\lambda \subset \mathfrak{g}^*$.

The correspondence $\Lambda_d \simeq \mathfrak{g}^*/G$ is most easily described in terms of the momentum map [1, 2]. The representation $U_\lambda(G)$ on \mathcal{V}_λ quotients to a G -action \tilde{U}_λ on the projective space $\mathbb{P}\mathcal{V}_\lambda$. Since U_λ is unitary, this quotient action may be computed from the action of $U_\lambda(G)$ on unit vectors in \mathcal{V}_λ . As explained at the beginning of section 3, $\mathbb{P}\mathcal{V}_\lambda$ is a symplectic manifold. It is clear from the definition of the symplectic structure on $\mathbb{P}\mathcal{V}_\lambda$ and the unitarity of U_λ that the reduced G -action \tilde{U}_λ preserves the symplectic form $\omega_{\mathbb{P}\mathcal{V}_\lambda}$. More strongly, this action admits an equivariant momentum map $J : \mathbb{P}\mathcal{V}_\lambda \rightarrow \mathfrak{g}^*$, given by

$$J_X(\psi) = i(\Psi, dU_\lambda(X)\Psi), \quad (23)$$

As always, the unit vector $\Psi \in \mathcal{V}_\lambda$ is an arbitrary lift of $\psi \in \mathbb{P}\mathcal{V}_\lambda$.

Proposition 4 *The coadjoint orbit \mathcal{O}_λ corresponding to an irreducible representation U_λ with highest weight vector Ψ_λ contains $J\psi_\lambda$. In fact, the restriction of J to $\mathbb{P}U_\lambda(G)\Psi_\lambda$ is a symplectomorphism onto \mathcal{O}_λ .*

Here it is understood that $\mathbb{P}U_\lambda(G)\Psi_\lambda$ inherits the usual symplectic structure of $\mathbb{P}\mathcal{V}_\lambda$, and that \mathcal{O}_λ is endowed with the Lie-Kirillov symplectic form. For convenience, as well as for later reference, we include a proof of this fundamental result (also cf. [1] or [4]).

Eq. (23), (20), (21), and (22) imply that $\langle J(\psi_\lambda), X \rangle$ equals $\lambda(X)$ for $X \in \mathfrak{t}$ and equals 0 for $X \in \mathfrak{t}^\perp$. Hence $J(\psi_\lambda)$ is precisely the element $\theta(\lambda) \in \mathfrak{g}^*$ discussed after (22), proving the first claim.

By (23), the stability group $G_{J(\psi_\lambda)}$ of $J(\psi_\lambda)$ consists of those $x \in G$ for which

$$(U_\lambda(x)\Psi_\lambda, dU_\lambda(Y)U_\lambda(x)\Psi_\lambda) = (\Psi_\lambda, dU_\lambda(Y)\Psi_\lambda) \quad (24)$$

for all $Y \in \mathfrak{g}$. Since U_λ is irreducible this implies that Ψ_λ and $U_\lambda(x)\Psi_\lambda$ define the same element of $\mathbb{P}\mathcal{H}_\lambda$, proving that $G_{J(\psi_\lambda)} \subseteq G_{\psi_\lambda}$. The opposite inclusion is trivial from the equivariance of J . \blacksquare

The unit vectors in \mathcal{V}_λ of the form $U_\lambda(x)\Psi_\lambda$, where $x \in G$, are Perelomov's coherent states [10, 11], which are parametrized by G . It should be clear by now that we are not interested in these states themselves, but in their projections to $\mathbb{P}\mathcal{V}_\lambda$. It follows from Proposition 4 that the ensuing family is parametrized by the coadjoint orbit \mathcal{O}_λ . Remarkably, the coadjoint orbits of maximal dimension among all coadjoint orbits \mathfrak{g} , which are the ones for which $G_\lambda = T$, are precisely the ones labeled by a regular weight λ . The ones of smaller dimension, for which G_λ properly contains T , are labeled by a singular weight.

V. Berezin quantization of coadjoint orbits

Proposition 4 shows how to construct the coadjoint orbit \mathcal{O}_λ corresponding to a given unitary irreducible representation U_λ by Proposition 3. The converse passage is accomplished by the Borel-Weil-Bott theory, which may be seen as a special case of geometric quantization [7]. In what follows we assume the correspondence in question to be given, and instead are interested in constructing an entire family of representations of G that are in some sense associated to a given coadjoint orbit. The fundamental idea of rescaling the label of an irreducible representation by multiplying with $1/\hbar$ (which accordingly has to be quantized in the compact case) is due to Berezin [1975a]; also cf. Perelomov [1986]. In a more intuitive setting, this rescaling was explicit in the early years of quantum mechanics, and seems to comprise one of the faces of Bohr's correspondence principle. The idea was further developed in the context of the classical limit of quantum mechanics by Lieb [8] and Simon [9].

Theorem 1 *Let G be a compact connected Lie group, and \mathcal{O}_λ an integral coadjoint orbit (cf. 5), corresponding to a highest weight $\lambda \in \Lambda_d$. For $\hbar = 1/k$, $k \in \mathbb{N}$, define $\mathcal{H}_\hbar = \mathcal{V}_{\lambda/\hbar}$, i.e., the carrier space of the irreducible representation $U_{\lambda/\hbar}(G)$ with highest weight $\lambda/\hbar = k\lambda$.*

Let $\tau : \mathcal{H}_\hbar \rightarrow \mathbb{P}\mathcal{H}_\hbar$ be the canonical projection. The map $q_\hbar : \mathcal{O}_\lambda \rightarrow \mathbb{P}\mathcal{H}_\hbar$, given by

$$q_\hbar(\text{Co}(x)\lambda) = \tau(U_{\lambda/\hbar}(x)\Psi_{\lambda/\hbar}), \quad (25)$$

is well defined and injective, and provides a pure state quantization of \mathcal{O}_λ (equipped with minus the Lie-Kirillov symplectic structure) on $I = 1/\mathbb{N} \cup 0$.

One should note here that $k\lambda \in \overline{\mathcal{C}}_d$ when $\lambda \in \overline{\mathcal{C}}_d$, since Weyl chambers are convex cones. The map q_\hbar is well-defined and injective by the equation $G_{k\lambda} = G_\lambda$ plus the argument on stability groups used in the proof of Proposition 4. In fact, if we define $J_\hbar : \mathbb{P}\mathcal{H}_\hbar \rightarrow \mathfrak{g}^*$ by (23) with the right-hand side divided by k , it follows from 4 that J_\hbar takes values in \mathcal{O}_λ and is a left-inverse of q_\hbar .

We start from the fact that the Haar measure on G (with total mass 1) pushes forward to the Liouville measure derived from the Lie symplectic structure under the canonical projection

$G \rightarrow \mathcal{O}_\lambda \simeq G/G_\lambda$. Using the invariance of the Haar measure and the unitarity of $U_{k\lambda}$, we then have

$$\int_{\mathcal{O}_\lambda} d\mu_L(\sigma) p(q_{\hbar}(\rho), q_{\hbar}(\sigma)) f(\sigma) = \int_G dx |(\Psi_{k\lambda}, U_{k\lambda}(x)\Psi_{k\lambda})|^2 f_\lambda(yx) \quad (26)$$

for all $f \in C(\mathcal{O}_\lambda)$, where $f_\lambda = \tau_{G \rightarrow G/G_\lambda}^* f \in C(G)$ is a right- G_λ -invariant function, and y is such that $\tau_{G \rightarrow G/G_\lambda}(y) = \rho \in \mathcal{O}_\lambda$. Choosing $f = 1$, the orthogonality relations for compact groups then imply (8) with

$$c(\hbar) = d_{\lambda/\hbar}, \quad (27)$$

i.e., the dimension of $\mathcal{H}_\hbar = \mathcal{V}_{\lambda/\hbar}$.

Eq. (10) follows from Proposition 4. To prove (9) we use a result of Gilmore [19].

Lemma 3 *Let λ_i be dominant weights with highest weight representations and vectors U_{λ_i} and Ψ_{λ_i} , respectively ($i = 1, 2$). Then for each $x \in G$*

$$(\Psi_{\lambda_1}, U_{\lambda_1}(x)\Psi_{\lambda_1}) \cdot (\Psi_{\lambda_2}, U_{\lambda_2}(x)\Psi_{\lambda_2}) = (\Psi_{\lambda_1+\lambda_2}, U_{\lambda_1+\lambda_2}(x)\Psi_{\lambda_1+\lambda_2}). \quad (28)$$

This is immediate from (20), (21), and the connectedness of G . ■

This lemma implies that

$$(\Psi_{k\lambda}, U_{k\lambda}(x)\Psi_{k\lambda}) = (\Psi_\lambda, U_\lambda(x)\Psi_\lambda)^k. \quad (29)$$

Using (26), we can write the left-hand side of (9) as

$$\lim_{\hbar \rightarrow 0} \int_{\mathcal{O}_\lambda} d\mu_L(\sigma) p(\rho_\hbar, \sigma_\hbar) f(\sigma) = \lim_{k \rightarrow \infty} \int_G d\mu_k(x) f_\lambda(yx),$$

where μ_k is a probability measure on G defined by

$$d\mu_k(x) = d_{k\lambda} dx |(\Psi_\lambda, U_\lambda(x)\Psi_\lambda)|^{2k}. \quad (30)$$

It is obvious that each μ_k is right- G_λ -invariant. It follows from (21), (22), and the fact that the exponential map is surjective for compact Lie groups, that $|(\Psi_\lambda, U_\lambda(x)\Psi_\lambda)|$, which is evidently ≤ 1 , equals 1 iff $x \in G_\lambda$. Hence for large k the support of μ_k is increasingly concentrated on G_λ . This suggests that

$$\lim_{k \rightarrow \infty} \mu_k(f) = \int_{G_\lambda} dh f(h) \quad (31)$$

for all $f \in C(G)$, where dh is the normalized Haar measure on G_λ . This is indeed the case, as proved by Duffield [20] (related results may be found in [8] and [9]). For the right- G_λ -invariant function $f_\lambda \in C(G)^{G_\lambda}$ one therefore obtains

$$\lim_{k \rightarrow \infty} \mu_k(f_\lambda) = f_\lambda(e).$$

This proves (9), which finishes the proof of Theorem 1. ■

The Berezin quantization \mathcal{Q}_\hbar^B associated with the pure state quantization in Theorem 1 is defined on $\tilde{\mathfrak{A}}_0 = C^\infty(\mathcal{O}_\lambda)$. By (12), one has

$$\mathcal{Q}_{1/k}^B(f) = d_{k\lambda} \int_G dx f_\lambda(x) [U_{k\lambda}(x)\Psi_{k\lambda}]; \quad (32)$$

this is an element of $\mathfrak{A}_{1/k} = \mathfrak{M}_{d_{k\lambda}}(\mathbb{C})$ (the C^* -algebra of complex $d_{k\lambda} \times d_{k\lambda}$ matrices). Recall that $[\Phi]$ is the projection onto $\mathbb{C}\Phi$.

The most important property of \mathcal{Q}_\hbar^B is its G -equivariance. For $x \in G$ we write

$$\alpha_x^{1/k}(A) = U_{k\lambda}(x) A U_{k\lambda}(x)^*, \quad (33)$$

where $A \in \mathfrak{A}_{1/k}$, and for $f \in \mathfrak{A}_0 = C(\mathcal{O}_\lambda)$ we put

$$\alpha_x^0(f) = \text{Co}(x^{-1})^* f. \quad (34)$$

Proposition 5 For all $k \in \mathbb{N}$, $x \in G$, and $f \in L^\infty(\mathcal{O}_\lambda)$ one has

$$\mathcal{Q}_{1/k}^B(\alpha_x^0(f)) = \alpha_x^{1/k}(\mathcal{Q}_{1/k}^B(f)). \quad (35)$$

This is immediate from (32), (34), the fact that $(\text{Co}(y)^*f)_\lambda = L_y^*f_\lambda$, the right-invariance of the Haar measure, and (33). \blacksquare

We are now ready for the main result of this paper.

Theorem 2 The Berezin quantization (32), defined on the space $C^\infty(\mathcal{O}_\lambda)$, is strict.

We show that the assumptions in Proposition 1 are met. Eq. (6) follows from Theorem 1 and Proposition 2. The completeness condition is an easy consequence of Schur's lemma and the irreducibility of $U_{k\lambda}$.

We will now prove the remaining conditions (5) and (3), using the notation of the proof of Theorem 1. We pick a unit vector Φ_k in each $\mathcal{V}_{k\lambda}$, and use the invariance of the Haar measure and the orthogonality relations for compact groups to write

$$\left(\Phi_k, (\mathcal{Q}_{1/k}^B(f)\mathcal{Q}_{1/k}^B(g) - \mathcal{Q}_{1/k}^B(fg))\Phi_k \right) = d_{k\lambda} \int_G dx f_\lambda(x) (\Phi_k, U_{k\lambda}(x)\Psi_{k\lambda}) I_k(x), \quad (36)$$

where

$$I_k(x) = d_{k\lambda} \int_G dy (\Psi_{k\lambda}, U_{k\lambda}(y)\Psi_{k\lambda}) F_\lambda^x(y), \quad (37)$$

in which we have abbreviated

$$F_\lambda^x(y) = (U_{k\lambda}(xy)\Psi_{k\lambda}, \Phi_k)[g_\lambda(xy) - g_\lambda(x)]. \quad (38)$$

In the notation used after (26), the function F_λ^x on G corresponds to a function F^x on \mathcal{O}_λ .

Using (29), we can write $(\Psi_{k\lambda}, U_{k\lambda}(y)\Psi_{k\lambda}) = \exp(-kS_\lambda(y))$, where $S_\lambda(y) = -\log(\Psi_\lambda, U_\lambda(y)\Psi_\lambda)$ (in view of the exponentiation, the choice of the branch cut of the logarithm is irrelevant). The function S_λ is right- G_λ -invariant; we denote the corresponding function on G/G_λ by S . We identify G/G_λ with \mathcal{O}_λ , so that the coset $[G_\lambda] \in G/G_\lambda$ is identified with $\lambda \in \mathcal{O}_\lambda$.

Putting $S_\lambda^+(y) = -\log|(\Psi_\lambda, U_\lambda(y)\Psi_\lambda)|$, the absolute value of $\exp(-kS)$ is $\exp(-kS^+)$. As in the argument preceding (31) we see that S^+ takes values in $[0, \infty]$ and assumes its unique absolute minimum 0 at λ . Since F_λ^x in (37) is bounded, a standard argument implies that to $O(\exp(-k))$ we may replace the integration over G/G_λ by one over any neighbourhood of λ .

We identify $T_\lambda\mathcal{O}_\lambda$ with $\mathfrak{g}/\mathfrak{g}_\lambda$, and use complex co-ordinates $\{z_\alpha, \bar{z}_\alpha\}_{\alpha \in \Delta_\lambda^+}$, where Δ_λ^+ consists of those positive roots for which $(\lambda, \alpha) \neq 0$. By the definition of a highest weight, this implies that $(\lambda, \alpha) > 0$ for all $\alpha \in \Delta_\lambda^+$. The co-ordinates $(z_\alpha, \bar{z}_\alpha)$ correspond to the point in \mathcal{O}_λ given by

$$\text{Co} \left[\text{Exp} \left(i \sum_{\alpha \in \Delta_\lambda^+} (z_\alpha E_\alpha - \bar{z}_\alpha E_{-\alpha}) \right) \right] \lambda.$$

A simple computation, using the Cartan-Weyl form of the commutation relations in \mathfrak{g} [18] and (20) - (22), leads to

$$S(z_\alpha, \bar{z}_\alpha) = \sum_{\alpha \in \Delta_\lambda^+} (\lambda, \alpha) z_\alpha \bar{z}_\alpha + O(|z|^3). \quad (39)$$

Hence to $O(\exp(-k))$ we may approximate $I_k(x)$ by

$$d_{k\lambda} \int_{\mathfrak{g}/\mathfrak{g}_\lambda} \left(\prod_{\alpha \in \Delta_\lambda^+} \frac{dz_\alpha d\bar{z}_\alpha}{2\pi} \right) J(z_\alpha, \bar{z}_\alpha) e^{-k \sum_{\alpha \in \Delta_\lambda^+} (\lambda, \alpha) z_\alpha \bar{z}_\alpha} F^x(z_\alpha, \bar{z}_\alpha),$$

where J is a Jacobian, and F_λ^x has been extended to $\mathfrak{g}/\mathfrak{g}_\lambda$ by, say, the exponential map. If we omit the factor [...] in (38), the integral (37) can be done, using the orthogonality relations for

compact groups. On the other hand, we can compute the above integral to lowest order in the steepest descent approximation; this avoids the need to compute $J(0)$. Comparing the results computes the prefactor in the steepest descent approximation as unity. As a by-product we obtain the asymptotic expression for $k \rightarrow \infty$

$$d_{k\lambda} \sim \frac{\prod_{\alpha \in \Delta^+} (\lambda, \alpha)}{J(0)} k^{\frac{1}{2} \dim(\mathcal{O}_\lambda)}, \quad (40)$$

where $\dim(\mathcal{O}_\lambda)$ is given by

$$\dim(\mathcal{O}_\lambda) = \dim(\mathfrak{g}) - \dim(\mathfrak{t}) - 2 \text{Card} \{ \alpha \in \Delta^+ | (\lambda, \alpha) = 0 \}. \quad (41)$$

Comparison with the Weyl dimension formula [18] then yields $J(0) = \prod_{\alpha \in \Delta^+} (\alpha, \delta)$, where $\delta = \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha$. Eq. (41) follows from the decomposition

$$\mathfrak{g}_\lambda = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Delta^+ | (\lambda, \alpha) = 0} \mathfrak{g}_\alpha^{\mathbb{R}}, \quad (42)$$

where $\mathfrak{g}_\alpha^{\mathbb{R}} = \mathfrak{g} \cap (\mathbb{C}E_\alpha \oplus \mathbb{C}E_{-\alpha})$. This decomposition is easily derived from the proof of Proposition 4, the Cartan-Weyl commutation relations, and the fact that $\lambda(E_\alpha) = 0$ for all $\alpha \in \Delta$.

Thus the steepest descent approximation to the above integral, and therefore to (37), reads

$$I_k(x) = \sum_{l=0}^N \frac{1}{l!} D^l (JF^x)(0) + O(k^{-N-1}), \quad (43)$$

where, abbreviating $\partial_\alpha = \partial/\partial z_\alpha$ and $\bar{\partial}_\alpha = \bar{\partial}/\bar{\partial} z_\alpha$, we have put

$$D = \sum_{\alpha \in \Delta^+} \frac{1}{(\lambda, \alpha)} \partial_\alpha \bar{\partial}_\alpha. \quad (44)$$

Substituting this expansion in (36) we see that

$$\left(\Phi_k, (\mathcal{Q}_{1/k}^B(f) \mathcal{Q}_{1/k}^B(g) - \mathcal{Q}_{1/k}^B(fg)) \Phi_k \right) = O(1/k). \quad (45)$$

To analyze the remainder of $O(1/k)$ we note that the l 'th term in the expansion leads to an x -integrand in (36) of the form

$$(U_{k\lambda}(x^{-1}) \Phi_k, \Psi_{k\lambda}) \overline{(U_{k\lambda}(x^{-1}) \Phi_k, \Psi^{(l_1)})} f \partial^{l_2} \bar{\partial}^{l_3} g_\lambda(x),$$

where $l_i \leq l$ and the $\Psi^{(l_1)}$ are given by the action of products of $dU_{k\lambda}(E_\alpha)$ and $dU_{k\lambda}(E_{-\alpha})$ on $\Psi_{k\lambda}$. The important point is now that the orthogonality relations (applied to the x -integration) then imply that the $O(k^{-N-1})$ term is bounded by $C \|\Phi_k\|^2 / k^{N+1}$ for some constant C . Hence (5) follows.

To prove (3) we need the $l = 1$ term in (43). We substitute (44), and perform some partial integrations in the remaining x -integral (using the invariance of the Haar measure). We abbreviate $A = (\Phi_k, U_{k\lambda}(x) \Psi_{k\lambda})$; then (21) implies that $\partial_\alpha A$ and $\bar{\partial}_\alpha \bar{A}$ vanish at $z_\alpha = \bar{z}_\alpha = 0$. Terms of the form $\partial_\alpha \bar{\partial}_\alpha A$ (or \bar{A}) drop out in the commutator, as do contributions from J (whose first derivatives at 0 already vanish identically). What remains is

$$\left(\Phi_k, ik [\mathcal{Q}_{1/k}^B(f), \mathcal{Q}_{1/k}^B(g)] - \mathcal{Q}_{1/k}^B(\{f, g\}) \Phi_k \right) = O(1/k), \quad (46)$$

where, in the realization of f, g as G_λ -invariant functions f_λ, g_λ on G ,

$$\{f_\lambda, g_\lambda\} = -i \sum_{\alpha \in \Delta_\lambda} \frac{1}{(\lambda, \alpha)} \xi_\alpha^L f_\lambda \xi_{-\alpha}^L g_\lambda. \quad (47)$$

Here the left-invariant vector fields $\xi_{\pm\alpha}^L$ on G are defined by first expressing $E_{\pm\alpha} \in \mathfrak{g}_{\mathbb{C}}$ in terms of elements of \mathfrak{g} , and then using the usual definition $\xi_X^L f(y) = df(y\text{Exp}(tX))/dt(t=0)$. Also, $\Delta_\lambda = \Delta_\lambda^+ \cup \Delta_\lambda^-$, i.e., the set of all roots α for which $(\lambda, \alpha) \neq 0$.

To finish the proof, we remark that (47) is precisely the Lie-Kirillov Poisson bracket on \mathcal{O}_λ ; this may be verified at the point $\lambda \in \mathcal{O}_\lambda$ (or $e \in G$) by direct computation from (18), from which the general statement follows by the G -invariance of the Poisson structure.

It is manifest that the right-hand side of (47) is left- G -invariant if f_λ and g_λ are. To prove the right- G_λ -invariance of (47), it is sufficient to establish that it is right-invariant under the derived action of the Lie algebra \mathfrak{g}_λ , for G_λ is connected. Recall that $[E_\alpha, E_\beta] = N_{\alpha,\beta}E_{\alpha+\beta}$, where $\beta \neq -\alpha$. We now need the identity $N_{-\alpha-\beta,\beta} = -N_{\alpha,\beta}$ (where $\beta \neq \pm\alpha$), which follows from the $\text{Ad}(\mathfrak{g})$ -invariance of the inner product on $\mathfrak{g}_{\mathbb{C}}$, combined with the normalization of the E_α . The right-invariance of (47) under \mathfrak{g}_λ follows by combining this identity with (42) and the Cartan-Weyl commutation relations.

The higher-order terms in (46) are dealt with as in the above proof of (5). This proves (3), finishing the proof of Theorem 2. ■

Finally, we remark that the results in this section have an obvious yet somewhat cumbersome generalization: if the orbit \mathcal{O}_λ is not integral, but such that $\mathcal{O}_{\lambda/c}$ is integral for some $c \in \mathbb{R} \setminus \{0\}$, we can construct a strict quantization for the values $\hbar = c/k$, $k \in \mathbb{N}$.

References

- [1] V. Guillemin and S. Sternberg *Symplectic Techniques in Physics* (Cambridge University Press, Cambridge 1984).
- [2] J.E. Marsden and T.S. Ratiu, *Introduction to Mechanics and Symmetry* (Springer, New York, 1994).
- [3] A. Connes, *Noncommutative Geometry*. (Academic Press, San Diego, 1994).
- [4] N.P. Landsman, *Mathematical Topics between Classical and Quantum Mechanics* (Springer, New York, 1998)
- [5] M.A. Rieffel, “Deformation quantization of Heisenberg manifolds”, *Commun. Math. Phys.* **122**, 531-562 (1989).
- [6] F.A. Berezin, “General concept of quantization”, *Commun. Math. Phys.* **40**, 153-174 (1975).
- [7] B. Kostant, “Quantization and unitary representations” *Lect. Notes Math.* **170**, 87-208 (1970).
- [8] E.H. Lieb, “The classical limit of quantum spin systems”, *Commun. Math. Phys.* **62**, 327-340 (1973) .
- [9] B. Simon, “The classical limit of quantum partition functions” *Commun. Math. Phys.* **71**, 247-276 (1980).
- [10] A.M. Perelomov, “Coherent states for arbitrary Lie groups”, *Commun. Math. Phys.* **26**, 222-236 (1972).
- [11] A. Perelomov, *Generalized Coherent States and their Applications* (Springer, Berlin, 1986).
- [12] E. Hawkins, “Quantization of equivariant vector bundles”, e-print q-alg/9708030.
- [13] D. Barmoshe and M.S. Marinov, “Realization of compact Lie algebras in Kähler manifolds”, *J. Phys.* **A27**, 6287-6298 (1994).
- [14] J. Dixmier, *C*-Algebras* (North-Holland, Amsterdam, 1977).
- [15] E. Kirchberg and S. Wassermann, “Operations on continuous bundles of C*-algebras”, *Math. Ann.* **303**, 677-697 (1995).
- [16] M.A. Rieffel, “Quantization and operator algebras”, in *Meeting with the Platypus. Proc. XIIIth Int. Congress of Mathematical Physics (Brisbane 1997)*, eds. A.J. Bracken, D. De Wit, M. Gould, P. Pearce (International Press, Singapore, 1998).
- [17] N.P. Landsman, “Strict deformation quantization of a particle in external gravitational and Yang-Mills fields”, *J. Geom. Phys.* **12**, 93-132 (1993).
- [18] B. Simon, *Representations of Finite and Compact Groups* (American Mathematical Society, Providence, 1996).
- [19] R. Gilmore, “The classical limit of quantum nonspin systems”, *J. Math. Phys.* **20**, 891-893 (1979).
- [20] N.G. Duffield, “Classical and thermodynamic limits for generalized quantum spin systems”, *Commun. Math. Phys.* **127**, 27-39 (1990).