

# FUNDAMENTAL GROUPS OF RATIONALLY CONNECTED VARIETIES

JÁNOS KOLLÁR

## 1. INTRODUCTION

Let  $X$  be a smooth, projective, unirational variety and  $U \subset X$  an open set. The aim of this paper is to find a smooth rational curve  $C \subset X$  such that the fundamental group of  $C \cap U$  surjects onto the fundamental group of  $U$ . Following the methods of [Kollár99] and [Colliot-Thélène00], a positive answer over  $\mathbb{C}$  translates to a positive answer over any  $p$ -adic field. This gives a rather geometric proof of the theorem of [Harbater87] about the existence of Galois covers of the line over  $p$ -adic fields (1.4). We also obtain a slight generalization of the results of [Colliot-Thélène00] about the existence of certain torsors over open subsets of the line over  $p$ -adic fields (1.6).

If  $U = X$  then  $\pi_1(X)$  is trivial (cf. (2.3)), thus any rational curve  $C$  will do. If  $X \setminus U$  is a divisor with normal crossings and  $C$  intersects every irreducible component of  $X \setminus U$  transversally, then the *normal* subgroup of  $\pi_1(U)$  generated by the image of  $\pi_1(C \cap U)$  equals  $\pi_1(U)$  by a simple argument. (See, for instance, the beginning of (4.2).) It is also not hard to produce rational curves  $C$  such that the image of  $\pi_1(C \cap U)$  has finite index in  $\pi_1(U)$  (cf. (3.3)). These results suggest that we are very close to a complete answer, but surjectivity is not obvious. Differences between surjectivity and finiteness of the index appear in many similar situation; see, for instance, [Kollár95, Part I] or [Napier-Ramachandran98].

The present proof relies on the machinery of rationally connected varieties developed in the papers [Ko-Mi-Mo92a, Ko-Mi-Mo92b, Ko-Mi-Mo92c]. The relevant facts are recalled in Section 2.

The main geometric result is the following:

**Theorem 1.1.** *Let  $K$  be an algebraically closed field of characteristic zero and  $X$  a smooth, projective variety over  $K$  which is rationally connected (2.1). Let  $U \subset X$  be an open subset and  $x_0 \in U$  a point. Then there is an open subset  $0 \in V \subset \mathbb{A}^1$  and a morphism  $f : V \rightarrow U$*

such that  $f(0) = x_0$  and

$$\pi_1(V, 0) \twoheadrightarrow \pi_1(U, x_0) \quad \text{is surjective.}$$

Moreover, we can assume that the following also hold:

1.  $H^1(\mathbb{P}^1, \bar{f}^*T_X(-2)) = 0$  where  $\bar{f} : \mathbb{P}^1 \rightarrow X$  is the unique extension of  $f$ .
2.  $\bar{f}$  is an embedding if  $\dim X \geq 3$  and an immersion if  $\dim X = 2$ .

**Corollary 1.2.** *Let  $K$  be a  $p$ -adic field and  $X$  a smooth, projective variety over  $K$  which is rationally connected over  $\bar{K}$ . Let  $U \subset X$  be an open subset and  $x_0 \in U(K)$  a point. Then there is an open subset  $0 \in V \subset \mathbb{A}^1$  and a morphism  $f : V \rightarrow U$  (all defined over  $K$ ) such that  $f(0) = x_0$  and*

$$\pi_1(V, 0) \twoheadrightarrow \pi_1(U, x_0) \quad \text{is surjective,}$$

where  $\pi_1$  here denotes the algebraic fundamental group.

**Remark 1.3.** More generally, (1.2) holds for any field  $K$  of characteristic zero such that every curve with a smooth  $K$ -point contains a Zariski dense set of  $K$ -points. Characterizations of this property are given in [Pop96, 1.1]. The following are some interesting classes of such fields:

1. Fields complete with respect to a discrete valuation.
2. Quotient fields of local Henselian domains.
3.  $\mathbb{R}$  and all real closed fields.
4. Pseudo algebraically closed fields, cf. [Fried-Jarden86, Chap. 10]

**Corollary 1.4.** [Harbater87] *Let  $G$  be a finite group and  $K$  a field of characteristic zero as in (1.3). Then there is a Galois cover  $g : C \rightarrow \mathbb{P}_K^1$  with Galois group  $G$  such that  $C$  is geometrically irreducible and  $g^{-1}(0:1) \cong G$ .*

Proof. Let  $G \subset GL(n, K)$  be a faithful representation. Set  $U = GL(n)/G$  with quotient map  $h : GL(n) \rightarrow U$  and let  $x_0$  be the image of the identity matrix. Then  $U$  is unirational, thus by (1.2) there is a  $0 \in V \subset \mathbb{A}^1$  and a morphism  $f : V \rightarrow U$  such that  $\pi_1(V) \twoheadrightarrow \pi_1(U)$  is onto.  $h : GL(n) \rightarrow U$  is étale and proper, thus it corresponds to a quotient  $\pi_1(U) \twoheadrightarrow G$ . The fiber product  $W := GL(n) \times_U V \rightarrow V$  corresponds to the surjective homomorphism

$$\pi_1(V) \twoheadrightarrow \pi_1(U) \twoheadrightarrow G.$$

Thus  $W$  is connected and  $W \rightarrow V$  is a Galois cover with Galois group  $G$ . Since  $W$  has a  $K$ -point, it is also geometrically connected. The

preimage of  $0 \in V$  is isomorphic to  $G$  (the disjoint union of  $|G|$  copies of  $\text{Spec}_K$ ).  $W \rightarrow V$  can be extended to a (ramified) Galois cover of the whole  $\mathbb{P}_K^1$ .  $\square$

**Remark 1.5.** The above proof works in positive characteristic if we know that for every subgroup  $H < G$  the quotient  $GL(n)/H$  has a smooth compactification.

The following result was proved by [Colliot-Thélène00] for finite groups, which is probably the most important for applications.

**Corollary 1.6.** *Let  $K$  be a field of characteristic zero as in (1.3),  $G$  a linear algebraic group scheme over  $K$  and  $A$  a principal homogeneous  $G$ -space. Then there is an open set  $0 \in V \subset \mathbb{A}_K^1$  and a geometrically irreducible  $G$ -torsor  $g : W \rightarrow V$  such that  $g^{-1}(0) \cong A$  (as a  $G$ -space).*

Proof. Assume that  $G$  acts on  $A$  from the left and choose an embedding  $G \subset GL(n)$  over  $K$ .  $A \times GL(n)$  has a diagonal left action by  $G$  and a right action by  $GL(n)$  acting only on  $GL(n)$ .

The right  $G$ -action makes the morphism  $h : G \backslash (A \times GL(n)) \rightarrow G \backslash (A \times GL(n))/G =: U$  into a  $G$ -torsor. Let  $x_0 \in U$  be the image of  $G \backslash (A \times G)$ . The fiber of  $h$  over  $x_0$  is isomorphic to  $A$ . Let  $G^0$  be the connected component of  $G$ . Then  $G \backslash (A \times GL(n)) \rightarrow G \backslash (A \times GL(n))/G^0$  is smooth with connected fibers and  $G \backslash (A \times GL(n))/G^0 \rightarrow U$  is étale and proper. Let  $0 \in V \subset \mathbb{A}^1$  and  $f : V \rightarrow U$  be as in the proof of (1.4). Then  $W := (G \backslash (A \times GL(n))) \times_U V$  works.  $\square$

A geometric application is the following:

**Corollary 1.7.** *For every  $2 \leq g \leq 13$  there is an open set  $0 \in V_g \subset \mathbb{C}$  and a smooth proper morphism with genus  $g$  fibers  $S_g \rightarrow V_g$  such that the image of the monodromy representation is the full Teichmüller group.*

Proof. The moduli of curves is unirational for  $g \leq 13$ . Apply (1.1) to the open subset of curves without automorphisms  $U_g \subset M_g$ .  $\square$

## 2. RATIONALLY CONNECTED VARIETIES AND MORPHISMS OF RATIONAL CURVES

Rationally connected varieties were introduced in [Ko-Mi-Mo92b] as a higher dimensional generalization of rational and unirational varieties. A surface is rationally connected iff it is rational. In higher dimensions rationality and unirationality are very hard to check. The notion of rational connectedness concentrates on rational curves on a variety. The following characterizations were developed in the papers [Ko-Mi-Mo92b], [Kollár96, IV.3], [Kollár98, 4.1.2].

**Definition–Theorem 2.1.** Let  $K$  be an algebraically closed field of characteristic zero. A smooth proper variety  $X$  over  $K$  is called *rationally connected* if it satisfies any of the following equivalent properties:

1. There is an open subset  $\emptyset \neq X^0 \subset X$ , such that for every  $x_1, x_2 \in X^0$ , there is a morphism  $f : \mathbb{P}^1 \rightarrow X$  satisfying  $x_1, x_2 \in f(\mathbb{P}^1)$ .
2. For every  $x_1, \dots, x_n \in X$ , there is a morphism  $f : \mathbb{P}^1 \rightarrow X$  satisfying  $x_1, \dots, x_n \in f(\mathbb{P}^1)$ .
3. There is a morphism  $f : \mathbb{P}^1 \rightarrow X$  such that  $H^1(\mathbb{P}^1, f^*T_X(-2)) = 0$ . (This is equivalent to  $f^*T_X$  being ample.)
4. There is a variety  $P$  and a dominant morphism  $F : \mathbb{P}^1 \times P \rightarrow X$  such that  $F((0:1) \times P)$  is a point. We can also assume that  $H^1(\mathbb{P}^1, F_p^*T_X(-2)) = 0$  for every  $p$  where  $F_p := F|_{\mathbb{P}^1 \times \{p\}}$ .
5. Let  $z_1, \dots, z_n \in \mathbb{P}^1$  be distinct points and  $m_1, \dots, m_n$  natural numbers. For each  $i = 1, \dots, n$  let  $f_i : \text{Spec } K[t]/(t^{m_i}) \rightarrow X$  be a morphism. Then there is a variety  $P$  and a dominant morphism  $F : \mathbb{P}^1 \times P \rightarrow X$  such that
  - (a) the Taylor series of  $F_p$  at  $z_i$  coincides with  $f_i$  up to order  $m_i$  for every  $i$  and  $p \in P$ ,
  - (b)  $H^1(\mathbb{P}^1, F_p^*T_X(-\sum m_i)) = 0$  for every  $i$  and  $p$ .  $\square$

Another easy result that we need is the following.

**Lemma 2.2.** (cf. [Kollár96, II.3.5.4, II.3.10.1 and II.3.11]) *Let  $X, P$  be smooth varieties and  $F : \mathbb{P}^1 \times P \rightarrow X$  a dominant morphism such that  $F((0:1) \times P)$  is a point. Then there is a dense open set  $P^0 \subset P$  such that  $H^1(\mathbb{P}^1, F_p^*T_X(-2)) = 0$  for every  $p \in P^0$  where  $F_p := F|_{\mathbb{P}^1 \times \{p\}}$ .*

*Conversely, let  $f : \mathbb{P}^1 \rightarrow X$  be a morphism such that  $H^1(\mathbb{P}^1, f^*T_X(-2)) = 0$ . Then there is a pointed variety  $p_0 \in P$  and a dominant morphism  $F : \mathbb{P}^1 \times P \rightarrow X$  such that  $F((0:1) \times P)$  is a point,  $F$  is smooth away from  $(0:1) \times P$  and  $F_{p_0} = f$ .  $\square$*

The following result was proved by [Serre59] for unirational varieties and by [Campana91, Ko-Mi-Mo92c] in general.

**Proposition 2.3.** *A smooth, proper, rationally connected variety is simply connected.  $\square$*

In the course of the proof we repeatedly encounter the following situation. We have morphisms  $f^i : \mathbb{P}^1 \rightarrow X$  each passing through the same point  $x_0 \in X$ . We would like to have a family of morphisms  $f_t : \mathbb{P}^1 \rightarrow X$  such that the union of the maps  $f^i$  can be considered as the limit of the maps  $f_t$  as  $t \mapsto 0$ . The following lemma is a technical formulation of this idea. Its statement is a bit complicated since we also want to keep track of the field over which the  $f_t$  are defined.

**Lemma 2.4.** (cf. [Kollár99, 3.2]) *Let  $K$  be a field and  $x_0 \in X$  a smooth, proper, pointed  $K$ -scheme. Let  $S$  be a zero dimensional reduced  $K$ -scheme and  $f_0 : \mathbb{P}_S^1 \rightarrow X$  a morphism such that*

1.  $H^1(\mathbb{P}_S^1, f_0^* T_X(-2)) = 0$ , and
2.  $f_0(S \times \{(0:1)\}) = \{x_0\}$ .

*Then there are*

3. *a smooth pointed curve  $0 \in D$  over  $K$ ,*
4. *a smooth surface  $Y$  with a proper morphism  $h : Y \rightarrow D$  and a section  $B \subset Y$  of  $h$ , and*
5. *a morphism  $F : Y \rightarrow X$ ,*

*such that*

6.  *$h^{-1}(0)$  is the union of  $\mathbb{P}_S^1$  with a copy  $B_0$  of  $\mathbb{P}_K^1$  such that  $B_0 \cap \mathbb{P}_S^1 = S \times \{(0:1)\}$  and  $B_0 \cap B$  is a single point.*
7.  *$F$  restricted to  $\mathbb{P}_S^1$  coincides with  $f_0$  and  $F(B_0 \cup B) = \{x_0\}$ ,*
8.  *$h^{-1}(D^0) \cong \mathbb{P}_K^1 \times D^0$ , where  $D^0 := D \setminus \{0\}$ ,*
9.  *$H^1(\mathbb{P}_d^1, F_d^* T_X(-2)) = 0$  for every  $d \in D^0$  where  $F_d := F|_{h^{-1}(d)}$ .*

*Proof.* Let us start with any curve  $0 \in D'$  and  $Y' := \mathbb{P}^1 \times D'$ . Let  $S' \subset \mathbb{P}^1 \times \{0\}$  be a subscheme isomorphic to  $S$  and  $Y''$  the blow up of  $S' \subset D'$  with projection  $h' : Y'' \rightarrow D'$ . We can define a morphism  $f' : (h')^{-1}(0) \rightarrow X$  by setting  $f'$  to be  $f_0$  on the exceptional divisor of  $Y'' \rightarrow Y'$  and the constant morphism to  $\{x_0\}$  on the birational transform of  $\mathbb{P}^1 \times \{0\}$ . Fix any section  $B' \subset Y'$  which does not pass through  $S'$ . We are done if  $f'$  can be extended to  $F' : Y'' \rightarrow X$  as required. In general this is not possible, but such an extension exists after a suitable étale base change  $(0 \in D) \rightarrow (0 \in D')$ . This is proved in [Ko-Mi-Mo92b, 1.2] and [Kollár99, 2.2].  $\square$

### 3. FUNDAMENTAL GROUPS OF FIBERS OF MORPHISMS

We need some easy results about the variation of fundamental groups for fibers of nonproper morphisms.

**Lemma 3.1.** *Let  $K$  be an algebraically closed field of characteristic zero,  $\bar{Z}, D$  irreducible  $K$ -varieties and  $f : \bar{Z} \rightarrow D$  a smooth and proper morphism with connected fibers. Let  $Z \subset \bar{Z}$  be an open subset such that  $(\bar{Z} \setminus Z) \rightarrow D$  is smooth. Let  $z_0 \in Z(K)$  be a  $K$ -point,  $d_0 = f(z_0)$  and  $Z_0$  the fiber of  $Z \rightarrow D$  through  $z_0$ . Then there is an exact sequence*

$$\pi_1(Z_0, z_0) \rightarrow \pi_1(Z, z_0) \rightarrow \pi_1(D, d_0) \rightarrow 1.$$

*Proof.* Over  $\mathbb{C}$  the fibration  $Z(\mathbb{C}) \rightarrow D(\mathbb{C})$  is a topological fiber bundle, thus we have the above exact sequence. To settle the algebraic case, let  $Y \rightarrow Z$  be any connected finite degree étale cover and extend

it to a finite morphism  $\bar{Y} \rightarrow \bar{Z}$  where  $\bar{Y}$  is normal. Since  $\bar{Z} \setminus Z$  is smooth over  $D$ , the same holds for  $\bar{Y} \rightarrow D$ . (This is a special case of Abhyankar's lemma, cf. [SGA1, XIII.5.2].) The generic fiber of  $\bar{Y} \rightarrow D$  is irreducible. Since  $\bar{Y} \rightarrow D$  is smooth and proper, every fiber is irreducible. This is equivalent to the exactness of the above sequence.  $\square$

The following technical lemma is an upper semi-continuity statement for the fundamental groups of fibers of nonproper morphisms.

**Lemma 3.2.** *Let  $K$  be an algebraically closed field of characteristic zero,  $W$  a normal surface over  $K$ ,  $f : W \rightarrow D$  a (not necessarily proper) morphism to a curve with connected fibers. Let  $B \subset W$  be a connected subset, one of whose irreducible components is a section of  $f$ . Let  $d_0 \in D$  be a  $K$ -point and  $C_0$  an irreducible component of  $f^{-1}(d_0)$  with a  $K$ -point  $b_0 \in C_0 \cap B$ . Let  $x_0 \in U$  be a pointed  $K$ -scheme and  $h : W \rightarrow U$  a morphism such that  $h(B) = \{x_0\}$ . Then there is an open subset  $D^0 \subset D$  such that for every  $d \in D^0$  and for every  $K$ -point  $b_d \in C_d \cap B$ ,*

$$\mathrm{im}[\pi_1(C_d, b_d) \rightarrow \pi_1(U, x_0)] \supset \mathrm{im}[\pi_1(C_0, b_0) \rightarrow \pi_1(U, x_0)].$$

*Proof.* Choose a normal compactification  $\bar{f} : \bar{W} \rightarrow D$ . Let  $D^0 \subset D$  be any open subset such that  $\bar{f}$  is smooth with irreducible fibers over  $D^0$  and  $\bar{W} \setminus W \rightarrow D$  is unramified over  $D^0$ . Set  $W^0 := f^{-1}(D^0)$ .  $\pi_1(C_0, b_0) \rightarrow \pi_1(U, x_0)$  factors through  $\pi_1(W, b_0) \rightarrow \pi_1(U, x_0)$  and it has the same image as  $\pi_1(W, b) \rightarrow \pi_1(U, x_0)$  for any  $b \in B(K)$ .  $\pi_1(W^0, b) \rightarrow \pi_1(W, b)$  is surjective by (3.3). By (3.1) there is an exact sequence

$$\pi_1(C_d, b_d) \rightarrow \pi_1(W_0, b_d) \rightarrow \pi_1(D^0, d) \rightarrow 1.$$

Let  $B^0 \subset B \cap W^0$  be a section of  $f$ . Then  $\pi_1(B^0, b_d)$  maps onto  $\pi_1(D^0, d)$  and the image of  $\pi_1(B^0, b_d)$  in  $\pi_1(U, x_0)$  is trivial. Thus

$$\mathrm{im}[\pi_1(C_d, b_d) \rightarrow \pi_1(U, x_0)] = \mathrm{im}[\pi_1(W^0, b_d) \rightarrow \pi_1(U, x_0)]$$

and we are done.  $\square$

**Lemma 3.3.** *(cf. [SGA1, IX.5.6], [Campana91, 1.3]). Let  $X, Y$  be normal varieties,  $x_0 \in X$  a closed point and  $f : X \rightarrow Y$  a dominant morphism. Then  $\mathrm{im}[\pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))]$  has finite index in  $\pi_1(Y, f(x_0))$ . If  $f$  is an open immersion then  $\pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$  is surjective.  $\square$*

## 4. PROOF OF THE MAIN RESULTS

The theory of free morphisms of curves (cf. [Kollár96, II.3]) suggests that morphisms  $f : \mathbb{P}^1 \rightarrow X$  such that  $H^1(\mathbb{P}^1, f^*T_X(-2)) = 0$  behave rather predictably. Therefore we concentrate on such morphisms. First we establish that there is a unique maximal subgroup of  $\pi_1(U)$  obtainable from such a morphism.

**Lemma 4.1.** *Let  $X$  be a smooth, proper, rationally connected variety over an algebraically closed field of characteristic zero. Let  $U \subset X$  be an open set and  $x_0 \in U$  a point. Then there is a unique finite index subgroup  $H < \pi_1(U, x_0)$  with the following properties:*

1. *There is a morphism  $f : \mathbb{P}^1 \rightarrow X$  such that  $H^1(\mathbb{P}^1, f^*T_X(-2)) = 0$ ,  $f(0:1) = x_0$  and  $H = \text{im}[\pi_1(f^{-1}(U), (0:1)) \rightarrow \pi_1(U, x_0)]$ .*
2. *Let  $g$  be any morphism  $g : \mathbb{P}^1 \rightarrow X$  such that  $g(0:1) = x_0$  and  $H^1(\mathbb{P}^1, g^*T_X(-2)) = 0$ . Then  $H \supset \text{im}[\pi_1(g^{-1}(U), (0:1)) \rightarrow \pi_1(U, x_0)]$ .*

Proof. First we find one morphism as in (4.1.2) such that

$$\text{im}[\pi_1(g^{-1}(U), (0:1)) \rightarrow \pi_1(U, x_0)]$$

has finite index in  $\pi_1(U, x_0)$ .

Let  $F : \mathbb{P}^1 \times P \rightarrow X$  be as in (2.1.4). Let  $P^0 \subset P$  be an open subset such that  $F^{-1}(X \setminus U) \rightarrow P$  is étale over  $P^0$ . Pick any point  $p \in P^0$ . By (3.1) there is an exact sequence

$$\begin{aligned} \pi_1(F_p^{-1}(U), (0:1) \times \{p\}) &\rightarrow \pi_1((\mathbb{P}^1 \times P^0) \cap F^{-1}(U), (0:1) \times \{p\}) \\ &\rightarrow \pi_1(P^0, p) \rightarrow 1. \end{aligned}$$

The section  $(0:1) \times P^0$  is mapped to a point by  $F$ , thus

$$\pi_1(F_p^{-1}(U), (0:1) \times \{p\}) \quad \text{and} \quad \pi_1((\mathbb{P}^1 \times P^0) \cap F^{-1}(U), (0:1) \times \{p\})$$

have the same image in  $\pi_1(U, x_0)$ . The latter image has finite index by (3.3), thus  $g := F_p : \mathbb{P}^1 \rightarrow X$  is as desired.

In order to finish, it is sufficient to prove that if  $g_1, g_2 : \mathbb{P}^1 \rightarrow X$  are as in (4.1.2) then there is a third morphism  $g : \mathbb{P}^1 \rightarrow X$  such that

$$\text{im}[\pi_1(g^{-1}(U), (0:1)) \rightarrow \pi_1(U, x_0)] \supset \text{im}[\pi_1(g_i^{-1}(U), (0:1)) \rightarrow \pi_1(U, x_0)]$$

for  $i = 1, 2$ . To do this, let  $S = \{1, 2\}$  be a 2 point scheme and take  $f_0 : \mathbb{P}_S^1 \rightarrow X$  to be  $g_i$  on  $\{i\} \times \mathbb{P}^1$ . Construct  $h : Y \rightarrow D$  and  $F : Y \rightarrow X$  as in (2.4). Set  $W := F^{-1}(U)$  and apply (3.2) twice for  $i = 1, 2$  with  $C_0 := f_i^{-1}(U)$ . For any  $d \in D^0$ ,  $g := F_d$  has the required property.  $\square$

## 4.2 (Proof of (1.1)).

Let  $H < \pi_1(U, x_0)$  be the subgroup obtained in (4.1). We are done if  $H = \pi_1(U, x_0)$ . Otherwise there is a corresponding irreducible étale

cover  $(x'_0 \in U') \rightarrow (x_0 \in U)$ . Let  $w : X' \rightarrow X$  be the normalization of  $X$  in the function field of  $U'$ .  $X$  is simply connected by (2.3), hence by the purity of branch loci (cf. [SGA1, X.3.1]) there is a divisor  $D' \subset X'$  such that  $w$  ramifies along  $D'$ . Let  $D \subset X$  be the image of  $D'$ . By construction  $D \subset X \setminus U$ . We derive a contradiction as follows.

Let  $f : \mathbb{P}^1 \rightarrow X$  be a morphism such that  $f(0:1) = x_0$  and  $f(\mathbb{P}^1)$  intersects  $D$  transversally at a point  $x_1 := f(1:1)$ . This implies that the local fundamental group of  $\mathbb{P}^1 \setminus \{(1:1)\}$  at  $(1:1)$  surjects onto the local fundamental group of  $X \setminus D$  at  $x_1$ . (See [Grothendieck-Murre71] for the local fundamental group of a divisor in a variety. There it is called the fundamental group of the formal neighborhood of a divisor.) Therefore, if  $f$  lifts to  $f' : \mathbb{P}^1 \rightarrow X'$  then  $X' \rightarrow X$  is étale at  $f'(1:1)$ . If  $X' \rightarrow X$  is a Galois extension then we already have a contradiction since  $X' \rightarrow X$  ramifies everywhere above  $D$ . In the non-Galois case  $p$  may be unramified along some of the irreducible components of  $p^{-1}(D)$ , and we only get a contradiction if the image of  $f'$  intersects  $D'$ .

If  $X' \rightarrow X$  is not Galois, we need to proceed in a somewhat round-about way. First I give the outlines; a precise version is given afterwards.

It is clear from the definition that there are many maps  $\mathbb{P}^1 \rightarrow X'$ , thus  $X'$  (or rather any desingularization of  $X'$ ) should be rationally connected. This indeed follows from (4.3) applied to any  $F : \mathbb{P}^1 \times P \rightarrow X$  as in (2.1.4). Thus by (2.1.5) there is a morphism  $f' : \mathbb{P}^1 \rightarrow X'$  which passes through  $x'_0$  and intersects  $D'$  at a smooth point  $x'_1$ . We obtain a contradiction if  $w \circ f'$  is the limit of a sequence of maps  $f_t : \mathbb{P}^1 \rightarrow X$  such that

1.  $f_t$  lifts to  $f'_t : \mathbb{P}^1 \rightarrow X'$  and  $f'$  is the limit of the maps  $f'_t$ ,
2. the image of  $f_t$  intersects  $D$  transversally.

First we prove a general lifting property for families of morphisms and then we proceed to construct the morphism  $f'$ .

**Lemma 4.3.** *Let  $V$  be a normal variety and  $G : \mathbb{P}^1 \times V \rightarrow X$  a morphism such that  $G((0:1) \times V) = \{x_0\}$ . Assume that  $H^1(\mathbb{P}^1, G_v^* T_X(-2)) = 0$  for some  $v \in V$ . Then  $G$  can be lifted to  $G' : \mathbb{P}^1 \times V \rightarrow X'$ .*

*Proof.* First we show that such a lifting exists over an open subset of  $\mathbb{P}^1 \times V$ . Choose  $V^0 \subset V$  such that  $H^1(\mathbb{P}^1, G_v^* T_X(-2)) = 0$  for every  $v \in V^0$  (this is possible by the upper semi continuity of cohomology groups) and such that we have an exact sequence

$$\pi_1(G_v^{-1}(U), (0:1) \times v) \rightarrow \pi_1((G^0)^{-1}(U), (0:1) \times v) \rightarrow \pi_1(V^0, v) \rightarrow 1$$

for every  $v \in V^0$  where  $G^0 := G|_{\mathbb{P}^1 \times V^0}$  (this is possible by (3.1)). Then

$$\text{im}[\pi_1((G^0)^{-1}(U), (0:1) \times v) \rightarrow \pi_1(U, x_0)] \subset H,$$



thus  $G^0|_{(G^0)^{-1}(U)}$  can be lifted to  $G' : (G^0)^{-1}(U) \rightarrow U'$ . This extends to a rational map  $G' : \mathbb{P}^1 \times V^0 \dashrightarrow X'$ .

Next let  $\Gamma \subset (\mathbb{P}^1 \times V) \times X$  be the graph of  $G$  and  $\Gamma^* \subset (\mathbb{P}^1 \times V) \times X'$  its preimage. The rational lifting  $G'$  corresponds to an irreducible component  $\Gamma' \subset \Gamma^*$  such that the projection  $\Gamma' \rightarrow (\mathbb{P}^1 \times V)$  is birational. Since  $X' \rightarrow X$  is finite, so is  $\Gamma^* \rightarrow \Gamma$ . Thus  $\Gamma' \rightarrow \Gamma \rightarrow \mathbb{P}^1 \times V$  is finite and birational, hence an isomorphism.  $\square$

Let  $\phi : X'' \rightarrow X'$  be any desingularization and  $x'_1 \in D'$  a point such that  $D'$  and  $X'$  are smooth at  $x'_1$  and  $\phi^{-1}$  is a local isomorphism near  $x'_1$ . By (2.1.5) there is a dominant morphism  $F : \mathbb{P}^1 \times P \rightarrow X'$  such that

1.  $F((0:1) \times P) = \{x'_0\}$ ,
2.  $F((1:1) \times P) = \{x'_1\}$ , and
3. the image of  $F_p : \mathbb{P}^1 \rightarrow X'$  is transversal to  $D'$  at  $x'_1$  for every  $p \in P$ .

Let us now consider the dominant morphism  $w \circ F : \mathbb{P}^1 \times P \rightarrow X$ . By the first part of (2.2), there is a  $p_0 \in P$  such that  $H^1(\mathbb{P}^1, (w \circ F_{p_0})^* T_X(-2)) = 0$ . Thus by the second part of (2.2) there is a pointed variety  $q_0 \in Q$  and a morphism  $G : \mathbb{P}^1 \times Q \rightarrow X$  such that  $G_{q_0} = w \circ F_{p_0}$  and  $G$  is smooth away from  $(0:1) \times Q$ . In particular,  $G^{-1}(D) \subset \mathbb{P}^1 \times Q$  is a generically smooth divisor, hence there is a dense open set  $Q^0 \subset Q$  such that the projection  $G^{-1}(D) \rightarrow Q$  is smooth over  $Q^0$ . This means that the image of  $G_q : \mathbb{P}^1 \rightarrow X$  intersects  $D$  transversally for every  $q \in Q^0$ . (Note that  $G^{-1}(D)$  denotes the inverse image scheme.)

By (4.3),  $G$  can be lifted to  $G' : \mathbb{P}^1 \times Q \rightarrow X'$ . On  $\mathbb{P}^1 \times \{q_0\}$  the lifting agrees with  $F_{p_0}$ , hence  $G'(\mathbb{P}^1 \times \{q_0\})$  intersects  $D'$  at the point  $x'_1$ . This implies that  $(G')^{-1}(D')$  is a divisor and so by shrinking  $Q$  we may assume that the image of  $G'_q$  intersects  $D'$  for every  $q \in Q$ . Thus  $G_q = w \circ G'_q$  is never transversal to  $D$ , a contradiction.

Condition (1.1.1) holds by construction and a general choice of  $f$  satisfies (1.1.2) by [Kollár96, II.3.14].  $\square$

#### 4.4 (Proof of (1.2)).

Pick  $f_1 : V_1 \rightarrow U_{\bar{K}}$  defined over  $\bar{K}$  such that  $\pi_1(V_1, 0) \rightarrow \pi_1(U_{\bar{K}}, x_0)$  is surjective and  $H^1(\mathbb{P}^1, \bar{f}_1^* T_X(-2)) = 0$ .  $f_1$  is defined over a finite Galois extension  $L \supset K$ ; let  $f_i : V_i \rightarrow U_{\bar{K}}$  be its conjugates. Each of these extends to a morphism  $\bar{f}_i : \mathbb{P}^1 \rightarrow X_{\bar{K}}$ . Let  $S = \text{Spec}_K L$ . We can view the morphisms  $f_i$  as one morphism  $f_0 : \mathbb{P}_S^1 \rightarrow X$  defined over  $K$ . By (2.4) we obtain  $h : Y \rightarrow D$  and  $F : Y \rightarrow X$ , all defined over  $K$ . Let  $d \in D^0(\bar{K})$  be any point. Then by (3.2) we see that

$$\text{im}[\pi_1(Y_d, 0) \xrightarrow{F_d} \pi_1(U_{\bar{K}}, x_0)] \supset \text{im}[\pi_1(V_1, 0) \xrightarrow{f_1} \pi_1(U_{\bar{K}}, x_0)],$$

and the latter image is  $\pi_1(U_{\bar{K}}, x_0)$  by assumption.  $D^0$  is a Zariski open set in a curve  $D$  with a smooth  $K$ -point  $0$ . By (1.3) this implies that  $D(K)$  is dense in  $D$ , hence  $D^0(K) \neq \emptyset$ . By choosing  $d \in D^0(K)$  we obtain an open subset  $0 \in V \subset \mathbb{A}^1$  and a morphism  $f : V \rightarrow U$  (all defined over  $K$ ) such that  $f(0) = x_0$  and

$$\pi_1(V_{\bar{K}}, 0) \twoheadrightarrow \pi_1(U_{\bar{K}}, x_0) \quad \text{is surjective.}$$

The fundamental group of a  $K$ -scheme  $W$  is related to the fundamental group of  $W_{\bar{K}}$  by the exact sequence (cf. [SGA1, IX.6.1])

$$1 \rightarrow \pi_1(W_{\bar{K}}, 0) \rightarrow \pi_1(W, 0) \rightarrow \text{Gal}(\bar{K}/K) \rightarrow 1.$$

This implies that

$$\pi_1(V, 0) \twoheadrightarrow \pi_1(U, x_0) \quad \text{is also surjective.}$$

□

**Example 4.5.** For every  $n \geq 4$  here I give an example of a rational threefold  $X$ , a normal crossing divisor  $F \subset X$  and a smooth rational curve  $B \subset X$  such that  $B$  intersects  $F$  everywhere transversally,  $B$  intersects every irreducible component of  $F$  and the image of  $\pi_1(B \setminus F) \rightarrow \pi_1(X \setminus F)$  has index  $n$  in  $\pi_1(X \setminus F)$ .

Let us start with a similar surface example. Let  $g_1 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a degree  $n$  morphism with critical points  $x_1, \dots, x_{2n-2} \in \mathbb{P}^1$  and different critical values  $y_1, \dots, y_{2n-2}$ . Choose 3 other points  $x_{2n-1}, x_{2n}, x_{2n+1} \in \mathbb{P}^1$  such that  $g_1(x_{2n-1})$  and  $g_1(x_{2n})$  are different critical values of  $g_1$  and  $g_1(x_{2n+1})$  is not a critical value of  $g_1$ . Let  $g_2 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a morphism with  $g_2(x_i) = x_i$  for  $i \leq 2n+1$ . Consider the morphism  $h : (g_2, g_1) : \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$

Set  $D = \mathbb{P}^1 \times \{y_1, \dots, y_{2n-2}\}$  and  $U = \mathbb{P}^1 \times \mathbb{P}^1 \setminus D$ .

$\text{im } h$  intersects every irreducible component of  $D$  transversally at  $n-2$  points and the image of  $\pi_1(h^{-1}(U)) \rightarrow \pi_1(U)$  has index  $n$  in  $\pi_1(U)$ .

Here  $\text{im } h$  is also tangent to every irreducible component of  $D$ . The tangencies can be resolved by  $2n$  blow ups, but then the birational transform of  $\text{im } h$  does not intersect every boundary component. To remedy this situation, take another morphism  $g_3 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  with  $g_3(x_i) = x_i$  for  $i \leq 2n+1$ . Set  $Y := \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ ,  $F = D \times \mathbb{P}^1$  and  $H := (g_3, g_2, g_1) : \mathbb{P}^1 \rightarrow Y$ . Again the only problem is that  $\text{im } H$  is tangent to every irreducible component of  $F$ . These can be resolved by blowing up two suitable smooth curves. First we take a smooth curve which passes through every point of tangency and also through the point  $H(x_{2n-1})$ . After blow up, the birational transform of  $\text{im } H$  intersects every boundary component transversally but above each point

there is a point common to two boundary components and to the birational transform of  $\text{im } H$ . Next take a smooth curve which passes through all these points with a general tangent direction there and also through  $H(x_{2n})$ . We can also assume that neither of the two curves passes through  $H(x_{2n+1})$ . Doing two such blow ups creates two new boundary components and the birational transform of  $\text{im } H$  intersects both of them. The fundamental group computation is unchanged.

Varying  $g_2, g_3$  we obtain many morphisms all of which pass through the point  $H(x_{2n+1})$ .

**Acknowledgments .** I thank J.-L. Colliot-Thélène and D. Madore for many helpful comments and references. Partial financial support was provided by the NSF under grant number DMS-9970855.

## REFERENCES

- Campana91. F. Campana, On twistor spaces of the class  $C$ , J. Diff. Geom. 33 (1991) 541-549
- Colliot-Thélène00. J.-L. Colliot-Thélène, Rational connectedness and Galois covers of the projective line, Annals of Math. 151 (2000) 359-373
- Fried-Jarden86. M. Fried and M. Jarden, Field arithmetic, Springer-Verlag, 1986
- Grothendieck-Murre71. A. Grothendieck and J. Murre, The tame fundamental group of a formal neighbourhood of a divisor with normal crossings on a scheme, Springer Lecture Notes vol. 208 (1971)
- Harbater87. D. Harbater, Galois coverings of the arithmetic line, Number Theory Seminar, New York 1984/85, Springer Lecture Notes vol. 1240, 1987, 165-195,
- Hartshorne77. R. Hartshorne, Algebraic Geometry, Springer, 1977
- Kollár95. J. Kollár, Shafarevich maps and automorphic forms, Princeton Univ. Press, 1995
- Kollár96. J. Kollár, Rational Curves on Algebraic Varieties, Springer Verlag (1996) Ergebnisse der Math. vol. 32
- Kollár98. J. Kollár, Low degree polynomial equations, European Congress of Math. Birkhäuser, 1998, 255-288
- Kollár99. J. Kollár, Rationally connected varieties over local fields, Annals of Math. 150 (1999) 357-367
- Ko-Mi-Mo92a. J. Kollár, Y. Miyaoka and S. Mori, Rational Curves on Fano Varieties, Proc. Alg. Geom. Conf. Trento, Springer Lecture Notes 1515 (1992) 100-105
- Ko-Mi-Mo92b. J. Kollár, Y. Miyaoka and S. Mori, Rationally Connected Varieties, J. Alg. Geom. 1 (1992 ) 429-448
- Ko-Mi-Mo92c. J. Kollár, Y. Miyaoka and S. Mori, Rational Connectedness and Boundedness of Fano Manifolds, J. Diff. Geom. 36 (1992) 765-769
- Napier-Ramachandran98. T. Napier and M. Ramachandran, The  $L^2$   $\bar{\partial}$ -method, weak Lefschetz theorems, and the topology of Kähler manifolds, J. Amer. Math. Soc. 11 (1998) 375-396

- Pop96. F. Pop, Embedding problems over large fields, *Annals of Math*, 144 (1996) 1-34
- Serre59. J.-P. Serre, On the fundamental group of a unirational variety, *J. London Math. Soc.* 34 (1959) 481-484
- SGA1. A. Grothendieck, *Revêtements étales et groupes fondamentaux*, Springer Lecture Notes vol. 224, 1971

Princeton University, Princeton NJ 08544-1000

`kollar@math.princeton.edu`