Study on the family of K3 surfaces induced from the lattice $(D_4)^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle$

K.Koike, H.Shiga, N.Takayama and T.Tsutsui

Mar. 18, 2000

Mathematics Subject Classifications (1991): 14J28, 33C70, 14K99

0 Introduction

In 1992 K.Matsumoto, T.Sasaki and M. Yoshida [8] studied the period mapping for a family of K3 surfaces of type (3,6), that is the family of double sextic surface over \mathbf{P}^2 ramified along 6 lines in general position, and Matsumoto [6] gave the description of the inverse mapping in terms of theta constants. It gives the modular mapping for the 4 dimensional Shimura variety in the Siegel upper half space \mathcal{H}_4 derived from the family of 4-dimensional abelian varieties with generalized complex multiplication by $\sqrt{-1}$. So we call it MSY modular mapping. Shiga showed an arithmetic application of MSY modular mapping in [15]. This story is the consequence of the eventual coincidence of 2 different bounded symmetric domains between D_{IV}^4 and $I_{2,2}$. There are a few (finite) such exceptional coincidences. The highest one is the (analytic) equivalence between D_{IV}^6 and $D_{II}(4)$ (in terms of Lie algebra $so(2,6; \mathbf{R}) \cong so(4, \mathbf{H})$) and it contains the above coincidence of MSY case.

Our present study is the first step to get the extended model of the MSY modular mapping using this equivalence.

Let us consider the rank 14 lattice $P = D_4^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle$. We define a K3 surface S of type P with the property that $P \subset \operatorname{Pic}(S)$ (see Definition 2.1), where $\operatorname{Pic}(S)$ indicates the Picard lattice of S. In this article we study the family of K3 surfaces of type P with a certain fixed multi-polarization. And we do not discuss the representation of the inverse of the period mapping.

We mention that our family is already appeared in the work [7] (in section 7) and they obtained the differential equation coming from this family standing on a different view point.

Throughout this article we work on the field **C**.

In Section 1 we study the family \mathcal{F} of double covering surfaces over $\mathbf{P}^1 \times \mathbf{P}^1$ branching along 4 bidegree (1,1) curves imposed with a certain generality condition. The element of \mathcal{F} is called a double 4*H* surface (see Definition 2.2). Such a surface is given in the affine form

$$S = S(x) : w^{2} = \prod_{k=1}^{4} (x_{1}^{(k)}st + x_{2}^{(k)}s + x_{3}^{(k)}t + x_{4}^{(k)}),$$

where we use the notation

$$x_k = \begin{pmatrix} x_1^{(k)} & x_2^{(k)} \\ x_3^{(k)} & x_4^{(k)} \end{pmatrix} \in M(2, \mathbf{C}).$$

We set up the view point that a general member S of \mathcal{F} to be an elliptic fibred surface (Proposition 1.1). It becomes to be a K3 surface. Then we construct the basis of the transcendental lattice $\operatorname{Tra}(S)$ of a general member S. Fortiori we know the structures of the Picard lattice $\operatorname{Pic}(S)$:

Theorem 1

For a general member S of \mathcal{F} it holds

$$\operatorname{Tra}(S) \cong U(2)^2 \oplus \langle -2 \rangle^4.$$

By making its orthogonal complement in the K3 lattice L we get the Picard lattice

$$\operatorname{Pic}(S) \cong D_4^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle.$$

In Section 2 we show that a K3 surface X of type P can be realized as a double covering surface studied in Section 1 provided a certain fixed polarization (Theorem 2).

In Section 3 we define a fixed marking of S of a K3 surface of type P. First we study the period domain for the family of such marked surfaces. That is a 6 dimensional domain given in the form

$$D^{+} = \{\eta = [\eta_{1}, \cdots, \eta_{8}] \in \mathbf{P}^{7} : {}^{t}\eta A\eta = 0, {}^{t}\overline{\eta}A\eta > 0, \Im(\eta_{3}/\eta_{1}) > 0\},\$$

where $A = U(2) \oplus U(2) \oplus (-2I_4)$. It is a bounded symmetric domain of type IV. Next we determine the modular group for the isomorphism classes of the marked surfaces. It is given as the principal congruence subgroup $G^+(2)$ of level 2 in the full group

$$G^{+} = \{g \in PGL(8, \mathbf{Z}) : {}^{t}gAg = A, g(D^{+}) = D^{+}\}\$$

of the positive isometries for the lattice $U(2)^2 \oplus \langle -2 \rangle^4$. The exact statement is given in Theorem 3.

In Section 4 we consider a general form of the period

$$u(x) = \int_C \Omega = \int_C \prod_{k=1}^4 (x_1^{(k)}st + x_2^{(k)}s + x_3^{(k)}t + x_4^{(k)})^{-1/2}ds \wedge dt,$$

where C indicates an element of $H_2(S(x), \mathbb{Z})$. We describe the differential equation for $\int_C \Omega$ as a multi-valued analytic function of 16 variables $x = (x_1, \dots, x_4)$, where we use the notation

$$x_k = \begin{pmatrix} x_1^{(k)} & x_2^{(k)} \\ x_3^{(k)} & x_4^{(k)} \end{pmatrix} \in M(2, \mathbf{C}).$$

That is a certain type of GKZ hypergeometric differential equation (Proposition 4.1 and 4.2) that is not so called Aomoto-Gelfond type (compare with [7]). We show the regularity of the system outside some divisor in the parameter space as stated in Theorem 4. We show the regular holonomicity of the system and make the calculation of the rank based on the theory of [14]. The exact statement is given in Theorem 5. We determine the monodromy group for our system in Theorem 6.

In Section 5 we construct the family $\{KS(\eta) : \eta \in D^+\}$ of Kuga-Satake varieties corresponding to \mathcal{F} . That becomes to be equivalent with the family of 8 dimensional Abelian varieties with the Hamilton quaternion endomorphism structure (Theorem 7).

1 Lattice structure of a double 4H surface

1.1 Setting up the situation

We consider an algebraic surface S' obtained as a double cover over $\mathbf{P}^1 \times \mathbf{P}^1$ ramifying along 4 different rational curves H_1, H_2, H_3, H_4 of bidegree (1,1). Here we suppose the following generality condition:

- (g1) H_i (*i* = 1, 2, 3, 4) is irreducible,
- (g2) $H_i \cap H_j$ $(i \neq j)$ consists of 2 different points,
- (g3) For any different 3 indices i, j, k we have $H_i \cap H_j \cap H_k = \emptyset$.

We denote π the projection $S' \to \mathbf{P}^1 \times \mathbf{P}^1$. Set $L_i = \pi^* H_i$. The surface S' has 12 singular points of type A_1 corresponding to the intersections $L_i \cap L_j$ $(i \neq j)$. By the desingularization procedure we get a K3 surface S.

If we have the condition

(e1) the algebraic variety S' has at most simple singularities

instead of (g1), (g2), (g3), we obtain a K3 surface by the same procedure. Henceforth we describe the curve $H_k = H(x^k)$ (k = 1, 2, 3, 4) in the form

$$(s,1)\begin{pmatrix} x_1^{(k)} & x_2^{(k)} \\ x_3^{(k)} & x_4^{(k)} \end{pmatrix} \begin{pmatrix} t \\ 1 \end{pmatrix} = 0$$

with a matrix

$$x^{k} = \begin{pmatrix} x_{1}^{(k)} & x_{2}^{(k)} \\ x_{3}^{(k)} & x_{4}^{(k)} \end{pmatrix} \in \mathbf{M}(2, \mathbf{C})$$

and an affine coordinate (s,t) of $\mathbf{P}^1 \times \mathbf{P}^1$. Let S' = S'(x) be the algebraic variety obtained from $H(x^1) \cup \cdots \cup H(x^4)$ by the above way, and let S(x) be its desingularization. Set

$$X^{\circ} = \{ x = (x^1, x^2, x^3, x^4) \in \mathcal{M}(2, \mathbb{C})^4 : H(x^1), \cdots, H(x^4) \text{ satisfy } (g1), (g2), (g3) \},\$$

and set

$$X' = \{x = (x^1, x^2, x^3, x^4) \in \mathcal{M}(2, \mathbb{C})^4 : S' \text{ has at most simple singularities} \}.$$

Definition 1.1. We call a double 4H surface the K3 surface S obtained as $S = S(x), x \in X^{\circ}$. An extended double 4H surface is a K3 surface obtained as $S = S(x), x \in X'$. Let \mathcal{F} denote the totality of double 4H surfaces:

$$\{S(x): x \in X^{\circ}\}$$

We use the following notations.

L: the K3 lattice $E_8^2 \oplus U^3$, where E_8 denotes the negative definite even unimodular lattice of rank 8 and U denotes the hyperbolic lattice $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The basis of L is fixed once and always. Pic(S): the Picard lattice of S,

Tra(S): the transcendental lattice of S. That is defined as the orthogonal complement of Pic(S) in $H^2(S, \mathbb{Z})$

Remark 1.1. We consider X° to be the parameter space of \mathcal{F} , but it contains many abundant parameters. Our family \mathcal{F} contains 6 essential parameters. We shall discuss the problem of abundance in Section 2 and Section 3.

1.2 Elliptic fibration

In this section we determine $\operatorname{Tra}(S)$ and $\operatorname{Pic}(S)$ for a general member of \mathcal{F} . For this work we always consider the problem in the dual lattice $H_2(S, \mathbb{Z})$ and we observe a special member of \mathcal{F} .

Let s, t be the affine coordinates of $\mathbf{P}^1 \times \mathbf{P}^1$. Henceforth we denote the s-space (t-space) by $\mathbf{P}^1(s)(\mathbf{P}^1(t))$, respectively. Set

$$f_1(s) = s$$
, $f_2(s) = \frac{16}{s}$, $f_3(s) = \frac{5s+2}{2s+5}$, $f_4(s) = \frac{3s+64}{4s+27}$

Consider the following four (1,1) curves :

$$H_i: t = f_i(s) \ (i = 1, 2, 3, 4),$$

and make the double 4H surface :

$$S_0: w^2 = \prod_{i=1}^4 (t - f_i(s))$$

that is derived from the above system $\{H_1, \dots, H_4\}$. Let π_1 be the projection from S_0 to $\mathbf{P}^1(s)$. For a generic point on $\mathbf{P}^1(s)$ we obtain an elliptic curve $\pi^{-1}(s)$. So we get an elliptic surface $(S_0, \pi_1, \mathbf{P}^1)$.

It is easy to observe the following.

Proposition 1.1. The elliptic surface $(S_0, \pi_1, \mathbf{P}^1)$ has 12 singular fibres corresponding to the intersections $H_i \cap H_j$ $(i \neq j)$. Every singular fibre is of type I_2 (according to the Kodaira classification of the singular fibres). These are situated over real points :

$$s = \pm 4 \quad (= H_1 \cap H_2), \pm 1 \quad (= H_1 \cap H_3), 2 \text{ and } -8 \quad (= H_1 \cap H_4), \\ -2 \text{ and } 8 \quad (= H_2 \cap H_3), \pm 12 \quad (= H_3 \cap H_4), \pm \sqrt{19} \quad (= H_3 \cap H_4).$$

We denote these 12 points by s_j $(j = 1, \dots, 12)$ according to the ascending order and set $Sing = \{s_1, \dots, s_{12}\}.$

We fix a base point $s_0 = 4\sqrt{-1}$ and set $E_0 = \pi_1^{-1}(s_0)$. It is given by

$$w^{2} = (t - f_{1}(4i))(t - f_{2}(4i))(t - f_{3}(4i))(t - f_{4}(4i)).$$

So this is a double cover over $\mathbf{P}^1(t)$ branched at 4 points $t = f_k(4i)$ (k = 1, 2, 3, 4). On E_0 we take a basis α_1, α_2 of $H_1(E_0, \mathbf{Z})$ with the intersection multiplicity $\alpha_1 \cdot \alpha_2 = 1$ as in Fig.1.

The fundamental group $\pi_1(\mathbf{P}^1 - \operatorname{Sing}, *)$ acts on $H_1(E_0, \mathbf{Z})$ as the monodromy group. We describe the generator system for it. Let γ_i $(i = 1, \dots, 12)$ be a loop starting from s_0 and goes around s_i in the positive sense on the upper half plain except the circuit around s_i (see Fig.2). By a direct observation we can calculate the monodromy transformation $T_j = T(\gamma_j)$ of $H_1(E_0, \mathbf{Z})$ along γ_j with respect to the basis $\{\alpha_1, \alpha_2\}$.

Proposition 1.2. The circuit matrices for $T_j = T(\gamma_j)$ are given by the following table. Here T_j acts from right to the system $\{\alpha_1, \alpha_2\}$.

s_j	$\pm 4, \pm \sqrt{19}$	$\pm 1, \pm 12$	8, -2	2, -8		
T_j	$ \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} $	$ \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix} $	$\begin{pmatrix} -1 & 2 \\ -2 & 3 \end{pmatrix}$	$ \begin{pmatrix} 3 & 2 \\ -2 & -1 \end{pmatrix} $		
vanishing cycle	α_1	α_2	$\alpha_1 + \alpha_2$	$\alpha_1 - \alpha_2$		



Figure 1: double covering of $\mathbf{P}^1(t)$



Figure 2: generator of $\pi_1(\mathbf{P}^1(s) - Sing)$

We choose a fixed point <u>s</u> in the lower half plane of $\mathbf{P}^1(s)$. Make cut lines c_i by the line segment between <u>s</u> and s_i $(i = 1, \dots, 12)$. By restricting $(S_0, \pi_1, \mathbf{P}^1)$ on $\mathbf{P}^1 - \bigcup_{i=1}^{12} c_i$ we have a topologically trivial fibration. So we can determine the cycles α_1, α_2 of $H_1(\pi^{-1}(s), \mathbf{Z})$ for any $s \in \mathbf{P}^1 - \bigcup_{i=1}^{12} c_i$ using this trivialization. If we make the continuation of the system $\{\alpha_1, \alpha_2\}$ passing through the line c_i from left to right, It is transformed according to Table 1.

1.3 Two systems $\{\Gamma_i\}$ and $\{C_i\}$ of $H_2(S_0, \mathbb{Z})$

Let r be an oriented arc on $\mathbf{P}^1(s)$ starting from s_0 . We make a 2-chain on S_0 by the continuation of a starting 1-cycle $\alpha \in H_1(E_0, \mathbf{Z})$ along r, and denote it by

 $r \times \alpha$.

Here we define its orientation as the ordered pair of the ones of r and α . Using this notation we make a system $\{\Gamma_1, \dots, \Gamma_8\}$ of 8 elements in $H_2(S_0, \mathbb{Z})$ (see Fig.3, Fig.4, Fig.5, Fig.6):

$$\begin{split} &\Gamma_{1} = \gamma(7)^{-1}\gamma(8)^{-1}\gamma(9)^{-1}\gamma(11)^{-1}\gamma(12)^{-1} \times \alpha_{1}, \\ &\Gamma_{2} = \gamma(4)^{-1}\gamma(5)^{-1}\gamma(6)^{-1}\gamma(7)^{-1}\gamma(8)^{-1}\gamma(9)^{-1} \times \alpha_{2}, \\ &\Gamma_{3} = \gamma(5)^{-1}\gamma(6)^{-1}\gamma(7)^{-1}\gamma(8)^{-1} \times \alpha_{1}, \\ &\Gamma_{4} = \gamma(2)^{-1}\gamma(3)^{-1}\gamma(4)^{-1}\gamma(5)^{-1} \times \alpha_{2}, \\ &\Gamma_{5} = \gamma(12) \times (\alpha_{1} + \alpha_{2}) + \gamma(10)^{-1}\gamma(11)^{-1}\gamma(12)^{-1} \times (-\alpha_{2}), \\ &\Gamma_{6} = \gamma(6) \times (\alpha_{1} + \alpha_{2}) + \gamma(4)^{-1}\gamma(5)^{-1}\gamma(6)^{-1} \times (-\alpha_{2}), \\ &\Gamma_{7} = \gamma(9) \times (\alpha_{1} - \alpha_{2}) + \gamma(7)^{-1}\gamma(8)^{-1}\gamma(9)^{-1} \times (-\alpha_{1}), \\ &\Gamma_{8} = \gamma(3) \times (\alpha_{1} - \alpha_{2}) + \gamma(1)^{-1}\gamma(2)^{-1}\gamma(3)^{-1} \times (-\alpha_{1}). \end{split}$$

where the composite arcs are performed from right to left.



Figure 3: Γ_1 (these are homologically equivalent)



Figure 4: Γ_2



Figure 5: $\Gamma_3(\text{left}), \quad \Gamma_4(\text{right})$

We note that any of $\{\Gamma_1, \dots, \Gamma_8\}$ is a 2-cycle according to the monodromy action in Table 1. So we regard them a system in $H_2(S_0, \mathbb{Z})$. Let $L(\Gamma)$ denote the sublattice $\oplus \mathbb{Z}\Gamma_i$.

Next we construct another system that is dual to $L(\Gamma)$. Let r(i) be the oriented line segment from s_0 to s_i . Using this notation we construct 8 elements in $H_2(S_0, \mathbb{Z})$ (see Fig.7):

$$\begin{split} C_1 &= (r(6) - r(7)) \times \alpha_2, \\ C_2 &= (r(9) - r(10)) \times \alpha_1, \\ C_3 &= (r(7) - r(12)) \times \alpha_2, \\ C_4 &= (r(4) - r(9)) \times \alpha_1, \\ C_5 &= r(10) \times \alpha_1 - r(11) \times (\alpha_1 + \alpha_2) - r(12) \times \alpha_2, \\ C_6 &= r(4) \times \alpha_1 - r(5) \times (\alpha_1 + \alpha_2) - r(6) \times \alpha_2, \\ C_7 &= -r(7) \times \alpha_2 - r(8) \times (\alpha_1 - \alpha_2) - r(9) \times \alpha_1, \\ C_8 &= -r(1) \times \alpha_2 - r(2) \times (\alpha_1 - \alpha_2) - r(3) \times \alpha_1. \end{split}$$

We note that any element of $\{C_1, \dots, C_8\}$ ends with a vanishing cycle at the terminal point



Figure 7: $C_1(\text{left}), \quad C_5(\text{right})$

of the base arc. So it is a 2-cycle on S_0 . Let L(C) denote the sublattice $\oplus \mathbb{Z}C_i$.

Proposition 1.3. We have the matrices of the intersection numbers $\Gamma_i \cdot \Gamma_j$ and $\Gamma_i \cdot C_j$ as follows :

[Proof]. We can calculate all the intersection numbers by a direct observation of the cycles. For example we consider $\Gamma_1 \cdot \Gamma_2$. They have 2 intersections on a vertical line between 4 and $\sqrt{19}$ of Γ_2 saying p_1, p_2 (see Fig.8). At the upper intersection p_1 the base arcs intersect with a negative sign, and the fibre cycles α_1, α_2 intersect with a positive sign. By considering the definition of the orientation on Γ_i we get there intersection number +1. The situation is quite



Figure 8: *intersection*

the same for the intersection in the lower half plane. So we have $\Gamma_1 \cdot \Gamma_2 = 2$. We can proceed this type of calculation until we get the full intersection matrices.

q.e.d.

Remark 1.2. 1) By using the conventional notations $U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $U(k) = \begin{pmatrix} 0 & k \\ k & 0 \end{pmatrix}$, we have

$$(\Gamma_i \cdot \Gamma_j) = U(2) \oplus U(2) \oplus \langle -2 \rangle^4.$$

2) We note that $(\Gamma_i \cdot C_j)$ is a unimodular matrix. So we can find a generator system $\{\check{\Gamma}_1, \cdots, \check{\Gamma}_8\}$ of L(C) with $\Gamma_i \cdot \check{\Gamma}_j = \delta_{ij}$.

Lemma 1.1. $L(\Gamma)$ and L(C) are nondegenerate rank 8 lattices. $L(\Gamma)$ is a primitive sublattice in $H_2(S_0, \mathbf{Z}) \cong L$.

Proof]. The first statement is a direct consequence of the fact that $(\Gamma_i \cdot C_j)$ is a unimodular matrix.

Suppose $L(\Gamma)$ is not primitive. It means that there is an element $\lambda \in L - L(\Gamma)$ with $k\lambda \in L(\Gamma)$ for some integer k.Put

$$k\lambda = \sum_{i=1}^{8} k_i \Gamma_i.$$

Then we have

$$k(\lambda \cdot \check{\Gamma}_i) = (k\lambda) \cdot \check{\Gamma}_i = k_i.$$

It indicates $k \mid k_i \ (i = 1, \dots, 8)$. So it is deduced $\lambda \in L(\Gamma)$. This is a contradiction.

q.e.d

1.4 The algebraic sublattice

For the moment we study the divisors on the reference surface S_0 . Let P_{ij}^{\pm} denote two intersections $H_i \cap H_j$ $(1 \le i < j \le 4)$, here we distinguish them by the signature of the *s*-coordinate (see Proposition 1.1). The surface S_0 has an exceptional curve corresponding to every intersection P_{ij}^{\pm} $(1 \le i < j \le 4)$. We denote them by E_{ij}^{\pm} . Let us consider the projection $\pi: S \to \mathbf{P}^1 \times \mathbf{P}^1$. Let F_s denote the pull back $\pi^*(\mathbf{P}^1(s)) \times \{t\}$ of the generic s-coordinate line, and let F_t denote the pull back of the generic t-coordinate line. Set

$$G_i = \frac{1}{2} (\pi^* H_i - \sum_{j \neq i} (E_{ij}^+ + E_{ij}^-)), \ (i = 1, 2, 3, 4).$$

This is the reduced divisor coming from the 4 sections H_i of the elliptic fibration $(S_0, \pi_1, \mathbf{P}^1(s))$.By an easy observation we have the following.

Lemma 1.2. We have intersection numbers among E_{ij}^{\pm} , G_i , F_s , F_t :

$$\begin{array}{ll} G_i \cdot G_i = -2, & F_s \cdot F_t = 2, & G_i \cdot F_s = G_i \cdot F_t = 1, \\ G_i \cdot E_{ij}^+ = G_i \cdot E_{ij}^- = 1, & E_{ij}^+ \cdot E_{ij}^+ = E_{ij}^- \cdot E_{ij}^- = -2, \end{array}$$

and all the other intersections are 0. We let L(div) denote the sublattice of $H_2(S_0, \mathbf{Z}) = L$ generated by $E_{ij}^{\pm}, G_i, F_s, F_t$.

Remark 1.3. If we observe the construction of Γ_i , we can show that $L(div) \subset L(\Gamma)^{\perp}$. Where \perp indicates the orthogonal complement in the full lattice $H_2(S_0, \mathbb{Z}) \cong L$.

Proposition 1.4. We have

$$L(\Gamma)^{\perp} = L(div) = \langle E_{ij}^{\pm}, G_i, F_s, F_t \rangle$$

$$\cong (D_4)^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle$$

on the reference surface S_0 .

Proof]. Set the sublattices L_i (i = 1, 2, 3) as follows:

$$L_{1} = \langle G_{1}, E_{14}^{+}, E_{14}^{-}, F_{t} - E_{23}^{+} \rangle, L_{2} = \langle G_{2}, E_{24}^{+}, E_{24}^{-}, F_{t} - E_{13}^{+} \rangle, L_{3} = \langle G_{3}, E_{34}^{+}, E_{34}^{-}, F_{t} - E_{12}^{+} \rangle.$$

They are isometric with the lattice

$$D_4 = \begin{pmatrix} -2 & 1 & 1 & 1\\ 1 & -2 & 0 & 0\\ 1 & 0 & -2 & 0\\ 1 & 0 & 0 & -2 \end{pmatrix},$$

and they are perpendicular each others. Set

$$\Delta_1 = F_s + F_t - E_{12}^+ - E_{13}^+ - E_{23}^+,$$

$$\Delta_2 = G_1 + G_2 + G_3 - G_4 + F_s + 3F_t - E_{12}^+ - E_{13}^+ - E_{23}^+,$$

then we have

$$\Delta_1 \cdot \Delta_1 = -2, \Delta_1 \cdot \Delta_2 = 0, \Delta_2 \cdot \Delta_2 = 2.$$

Now we put

$$P = L_1 \oplus L_2 \oplus L_3 \oplus \langle \Delta_1 \rangle \oplus \langle \Delta_2 \rangle.$$

Then it holds

$$P \subset L(div) \subset L(\Gamma)^{\perp},$$

rank $P = 14$, discr $P = -2^8$.

On the other hand we have discr $L(\Gamma) = 2^8$. By considering the fact that L is unimodular and $L(\Gamma)$ is a primitive sublattice, we get the conclusion that $P = L(div) = L(\Gamma)^{\perp}$.

q.e.d.

Now we extend Proposition 1.4 to the general situation.

Theorem 1. Let S be a double 4H surface. If we have rank Pic(S) = 14, then it holds

$$Tra(S) \cong U(2) \oplus U(2) \oplus (-2I_4),$$

$$Pic(S) \cong D_4 \oplus D_4 \oplus D_4 \oplus \langle -2 \rangle \oplus \langle 2 \rangle.$$

[Proof]. We note that the family $\mathcal{F} = \{S(x)\}$ is fibred over the parameter space of $M(2, \mathbb{C})^4$. This fibration is locally topological trivial on a Zariski open subset in $X^\circ \subset M(2, \mathbb{C})^4$. Using this trivialization we can proceed the same argument for a general member as for the specialized element S_0 .In fact always L(div) and P are defined as a sublattice in $\operatorname{Pic}(S(x))$. We have the systems $L(\Gamma)$ and L(C) also. So Proposition 1.4 is valid for every S of \mathcal{F} . If the condition is satisfied, then $L(\Gamma)$ cannot contain any divisor class. Hence we obtain the required conclusion. q.e.d

Remark 1.4. As we will see in Section 2, we have rank Pic(S) = 14 for a general member of \mathcal{F} .

2 Family \mathcal{F} as that of lattice K3 surfaces

2.1 General properties

In this section we show the converse of Theorem 1. Before starting this argument we state some general properties of a K3 surface.

Lemma 2.1. Let S be a K3 surface that is given by a minimal nonsingular model.

1) For an irreducible curve C on S, $C^2 \ge 0$ or $C^2 = -2$.

2) If $D \in Pic(S)$ satisfies $D^2 \ge -2$, then either D or -D is an effective class (that is given by an effective divisor).

This is deduced from the Riemann-Roch theorem by a routine argument.

Definition 2.1. We call an element $D \in Pic(S)$ is nef when it holds $D \cdot C \ge 0$ for any effective class C.

Proposition 2.1 (Pjateckii-Šapiro and Šafarevič [12]). Let S be a K3 surface, then we have the following:

1) Suppose $x \in \text{Pic}(S)$ satisfies $x \neq 0$, $x^2 = 0$. Then there exists an isometry γ of Pic(S) such that $\gamma(x)$ becomes to be effective and nef.

2) Suppose $x \in \text{Pic}(S)$ is effective, nef and $x^2 = 0$ then x is a multiple class of an elliptic curve (i.e. x = m[E] for a certain $m \in \mathbf{N}$ and an elliptic curve E on S).

3) The linear system of an elliptic curve C on X determines an elliptic fibration $S \to \mathbf{P}^1$.

2.2 A lattice K3 surface and its realization as a double 4H surface

Let us consider the lattice

$$P = D_4^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle, \tag{2.1}$$

and we let (*, *) denote the bilinear form on an abstract lattice P.

Definition 2.2. We say a K3 surfaces S is of type P if

(a) S admits an embedding $P \hookrightarrow \operatorname{Pic}(S)$,

and S is of exact type P if

(a') S admits an isomorphism $P \cong Pic(S)$.

By the argument in Section 1 we can find the generator system of P:

$$f_s, \quad f_t, \quad g_i \quad (i = 1, \cdots, 4), \quad e_{ij}^+, \quad e_{ij}^- \quad (i \neq j, 1 \le i, j \le 4),$$

$$(2.2)$$

with the following properties among them :

(p1)

$$\begin{array}{l} (g_i,g_i) = -2, \quad (f_s,f_t) = 2, \quad (g_i,f_s) = (g_i,f_t) = 1 \\ (g_i,e_{ij}^+) = (g_i,e_{ij}^-) = 1, \quad (e_{ij}^+,e_{ij}^+) = (e_{ij}^-,e_{ij}^-) = -2, \end{array}$$

and all other intersections equal 0,

(p2)

$$f_s + f_t = 2g_i + \sum_{j \neq i} (e_{ij}^+ + e_{ij}^-)$$

We note that this system is determined uniquely up to isometries of P.

Theorem 2. Let S be a K3 surface of type P with the property:

(n) f_s, f_t are nef classes under the identification in (a).

Then S has a representation as an extended double 4H surface.

When S is of exact type P with the above condition (n), S has a representation as a double 4H surface.

[Proof]. Always we regard the element of P under the identification in (a) or (a'). We show the first assertion.

(Claim 1) We can assume that every e_{ij}^{\pm} is effective.

Let e be one of the twelve elements e_{ij}^{\pm} . Since $e \cdot e = -2$, either e or -e is effective. If -e is effective then we perform the reflection γ on $\operatorname{Pic}(S)$ determined by e. And we can use

$$\gamma(f_s), \quad \gamma(f_t), \quad \gamma(g_i), \quad \gamma(e_{ij}^{\pm})$$

instead of the system (2.2). According to the orthogonality of the system $\{e_{ij}^{\pm}\}\$ we can iterate this procedure until we get the required effective system. Since we have $\gamma(f_s) = f_s, \gamma(f_t) = f_t$, the nef property for f_s, f_t is always satisfied throughout this process.

(Claim 2) We can find a double covering S', that is birationally equivalent with S,

$$\pi = (\pi_1, \pi_2) : S' \to \mathbf{P}^1 \times \mathbf{P}^1,$$

ramified along a bidegree (4, 4) curve B. S' has at most simple singularities coming from the singular points of B.

By Proposition 2.1 f_s , f_t are multiple classes of elliptic curve. Observing the assumption $g_i \cdot f_s = g_i \cdot f_t = 1$ we know that they are reduced classes. Again by Proposition 2.1 f_s and f_t determine two different elliptic fibrations $\pi_1 : X \to \mathbf{P}^1$ and $\pi_2 : X \to \mathbf{P}^1$, respectively. Set

$$\pi = (\pi_1, \pi_2) : S \to \mathbf{P}^1 \times \mathbf{P}^1.$$

This map is surjective and of degree 2, because we have $\pi_1^{-1}(x) \cdot \pi_1^{-1}(y) = f_s \cdot f_t = 2$ for any $(x, y) \in \mathbf{P}^1 \times \mathbf{P}^1$. Let L_s, L_t be two lines $\mathbf{P}^1 \times \{\infty\}$ and $\{\infty\} \times \mathbf{P}^1$, respectively. So we obtain a double covering $\pi = (\pi_1, \pi_2) : S \to \mathbf{P}^1 \times \mathbf{P}^1$. Generic fibers $\pi_1^*(x)$ and $\pi_2^*(y)$ are elliptic curves realized as double coverings over \mathbf{P}^1 . So each of them has four branch points. Consequently, the branch locus B of π is a curve of degree (4, 4). Since S is a K3 surface, the canonical class $K_S = 0$. Hence B has at most simple singularities.

(Claim 3) Let G_i be the effective divisor representing g_i . Then $H_i = \pi_*G_i$ is a (1, 1)-curve (i = 1, 2, 3, 4).

Since $g_i \cdot g_i = -2$, either g_i or $-g_i$ is effective. But f_s is nef and has intersection $f_s \cdot g_i = 1$, then we know that g_i is effective. By the projection formula, we obtain

$$\pi_*G_i \cdot L_s = G_i \cdot \pi^*L_s = g_i \cdot f_s = 1.$$

By the same argument, we get $\pi_*G_i \cdot L_t = 1$ also. Hence we obtain the required property.

(Claim 4) The effective class e_{ij}^{\pm} is obtained by an exceptional divisor coming from the singularity of B.

Let E_{ij}^{\pm} be the effective divisor representing e_{ij}^{\pm} . Since $(e_{ij}^{\pm}, f_s) = (e_{ij}^{\pm}, f_t) = 0$ we get $\pi_* E_{ij}^{\pm} = 0$. It indicates that every E_{ij}^{\pm} is an exceptional divisor derived from the singularity of B.

(Claim 5). We have $B = H_1 + H_2 + H_3 + H_4$

We consider the (1,1) curve $H_i = \pi_* G_i$. As a divisor class $\pi^* H_i$ equals to $f_s + f_t$. By the starting assumption (p2), $\pi^* H_i$ and $2G_i + \sum_{j \neq i} (E_{ij}^+ + E_{ij}^-)$ are linearly equivalent. So there exist a principal divisor D such that

$$\pi^* H_i = 2G_i + \sum_{j \neq i} (E_{ij}^+ + E_{ij}^-) + D.$$

Then

$$2H_i = \pi_* \pi^* H_i = 2\pi_* G_i + \pi_* D = 2H_i + \pi_* D.$$

This is an equality as divisors its elves (not as divisor classes), so we get $\pi_*D = 0$. This implies that D is an sum of exceptional divisors. But D is principal, so we must have D = 0. By observing the equality

$$\pi^* H_i = 2G_i + \sum_{j \neq i} (E_{ij}^+ + E_{ij}^-)$$

we know that $\pi^* H_i$ has six components of the form E_{ij}^{\pm} , note that we don't have any cancellation with an effective divisor G_i . Hence H_i meets six different singular points on B. The sum of intersection numbers at these double points exceeds the number $(B, H_i) = 8$. In case H_i is irreducible it must be an irreducible component of B. In case we have $H_i = \ell_1 + \ell_2$ with a (0,1) curve and a (1,0) curve, at least one component, saying ℓ_1 , is contained in B. If ℓ_2 is not contained in B, we have $\ell_2 \cdot (B - \ell_1) = 3$. So ℓ_2 contains only one $\pi(E_{ij}^{\pm})$. Namely ℓ_1 contains 5 others. Observing $\ell_1 \cdot (B - \ell_1) = 4$ we know that is too many. So we have $H_i \subset B$. Hence we obtained the claim.

According to the above arguments we obtained the double covering $\pi : S \to \mathbf{P}^1 \times \mathbf{P}^1$. realizing the extended double 4H surface.

Next let us prove the second assertion using the condition (a'). It is enough to show (Claim 6) S' has exactly 12 singular points of type A_1 on π^*B .

Let L(exc) be the sublattice generated by the system $\{e_{ij}^{\pm}\}$ in P = Pic(S), this is isometric with $\langle -2 \rangle^{12}$. Suppose $x \in P$ is a class of irreducible (-2)-curve with $\pi_* x = 0$. We can describe

x in the form

$$x = m_s f_s + m_t f_t + e + \sum_{i=1}^4 m_i g_i, \quad e \in L(exc)$$

with the coefficients $m_s, m_t, m_i \in \mathbf{Z}$. By assumption it holds $\pi_* x \cdot L_s = \pi_* x \cdot L_t = 0$, so we obtain

$$m_s = m_t$$
, $2m_s + m_1 + m_2 + m_3 + m_4 = 0$.

According to the starting condition (p1), (p2), we have

$$2(x-e) = 2(m_s(f_s + f_t) + m_1g_i + m_2g_2 + m_3g_3 + m_4g_4)$$

=
$$\sum_{i=1}^4 m_i(2g_i - f_s - f_t)$$

=
$$-\sum_{i=1}^4 m_i \sum_{j \neq i} (e_{ij}^+ + e_{ij}^-)$$

So we have $2x \in L(excep)$. By assumption we have (2x, 2x) = -8. But such an element in L(exc) should be the form $\pm 2e_{ij}^{\pm}$ or $\pm e_1 \pm e_2 \pm e_3 \pm e_4$. Where $\{e_i \ (i = 1, 2, 3, 4)\}$ are distinct four elements of $\{e_{ij}^{\pm}\}$. In the case $2x = \pm 2e_{ij}^{\pm}$, we conclude $x \in L(excep)$.

In the later case, we obtain

$$\pm e_1 \pm e_2 \pm e_3 \pm e_4 - 2e = -\sum_{i=1}^4 m_i \sum_{j \neq i} (e_{ij}^+ + e_{ij}^-).$$

Observe the right hand side, the number of the odd coefficients of e_{ij}^{\pm} should be one of 0, 6, 8. It does not attained in the left hand. So this case does not happen. Hence we obtained the claim.

q.e.d

3 Modular group and marking with a multi-polarization

3.1 Congruence subgroup G(2)

Let L be the K3 lattice with the fixed basis as defined in Section 1 and (,) be the corresponding bilinear form. We consider the sublattice $P = (D_4)^3 \oplus \langle -2 \rangle \oplus \langle 2 \rangle$ with the fixed generator system $\{f_s, f_t, e_{ij}^{\pm}, g_i\}$ as in Section 2. Set $T = P^{\perp} = U(2)^2 \oplus \langle -2 \rangle^4$ with the fixed basis so that $A = U(2) \oplus U(2) \oplus (-2I_4)$ be the intersection form. We denote the group of isometries of a lattice M by O(M). Set G = O(T) and

$$O(L, P) = \{ g \in O(L) : g(x) = x \text{ for } \forall x \in P \}.$$

Note that we have g(T) = T for $g \in O(L, P)$. So we can regard O(L, P) as a subgroup of G. Now G is given as

$$G = \{g \in M(8, \mathbf{Z}) : \quad {}^{t}gAg = A\}.$$

Set

$$G(2) = \{ g \in G : g \equiv I \mod 2 \}.$$

Let $\alpha \in T$ be a (-2)-element, that is $(\alpha, \alpha) = -2$. It determines a reflection $\gamma_{\alpha} \in O(L, P)$ by putting

$$\gamma_{\alpha}(x) = x + (x, \alpha)\alpha.$$

Remark 3.1. According to [5] and [9], we know that G(2) is generated by the reflections of the above form γ_{α} . So it holds $G(2) \subset O(L, P)$.

Proposition 3.1. We have

$$G(2) = \mathcal{O}(L, P)$$

[Proof]. It is enough to show $O(L, P) \subset G(2)$. Here we use the notations and results in [10]. Let $q_T : A_T \to \mathbf{Q}$ (resp. q_P) be the discriminant form of T (resp. P), where $A_T = T^*/T$.

(Fact): $H = L/T \oplus P$ is a maximal isotropy subgroup for $q_T \oplus q_P$ in $A_T \oplus A_P$. Namely it holds $(q_T \oplus q_P) \mid_H \equiv 0$ and H is maximal with this property. Moreover the canonical projections $H \to A_T$ and $H \to A_P$ induce the isomorphism

$$A_T \cong H \cong A_P.$$

Hence we have

$$O(L, P) \hookrightarrow Ker\{G \to O(q_T)\}$$

, where $G \to O(q_T)$ indicates the canonical map. We can easily show that

$$G(2) = Ker\{G \to \mathcal{O}(q_T)\}.$$

q.e.d

3.2 Marking with a multi-polarization

Definition 3.1. Let S be a K3 surface with an embedding $P \hookrightarrow \text{Pic}(S)$. The triple (S, φ, P) is a P-marking of S provided

- 1) $\varphi: H_2(S, \mathbf{Z}) \to L$ is an isometry,
- 2) $\varphi^{-1}(f_s), \varphi^{-1}(f_t), \varphi^{-1}(e_{ij}^{\pm}), \varphi^{-1}(g_i)$ are effective and $\varphi^{-1}(f_s), \varphi^{-1}(f_t)$ are nef.

According to Theorem 2 we have the double covering representation $\pi: S \to \mathbf{P}^1 \times \mathbf{P}^1$ and the class

$$\pi_*(\varphi^{-1}(g_1) + \dots + \varphi^{-1}(g_4))$$

gives the class of the ramification divisor.

Definition 3.2. Let (S, φ, P) and (S', φ', P) be *P*-markings of *S* and *S'*, respectively. An isomorphism $\rho: S \to S'$ is an isomorphism between these markings provided $\varphi = \varphi' \circ \rho_*$.

Remark 3.2. Because we have $\rho_*\varphi^{-1}(f_s) = (\varphi')^{-1}(f_s)$ and $\rho_*\varphi^{-1}(f_t) = (\varphi')^{-1}(f_t)$, such an isomorphism ρ preserves the covering structure over $\mathbf{P}^1 \times \mathbf{P}^1$.

Namely we have the following commutative diagram with an element $\sigma \in PGL_2(\mathbb{C}) \times PGL_2(\mathbb{C})$:



Definition 3.3. Let $H = (H_1, H_2, H_3, H_4)$, $H' = (H'_1, H'_2, H'_3, H'_4)$ be ordered sets of four curves of bidegree (1,1). We say that H and H' are equivalent if there exists $\sigma \in PGL_2(\mathbf{C}) \times PGL_2(\mathbf{C})$ such that $\sigma(H_i) = H'_i$ (i = 1, 2, 3, 4).

Remark 3.3. Let (S, φ, P) and (S', φ', P) be *P*-markings of *S* and *S'*, respectively.

(1) If these two P-markings are isomorphic then we have the same equivalent class of the ordered sets $H = (H_1, H_2, H_3, H_4)$.

(2) But we don't have the converse of (1).

3.3 Modular group

Let $\{\Gamma_1, \dots, \Gamma_8\}$ be the basis of T such that $(\Gamma_i, \Gamma_j) = A$. We have elements $\{\check{\Gamma}_1, \dots, \check{\Gamma}_8\}$ in L such that $(\check{\Gamma}_i, \Gamma_j) = \delta_{ij}$. The system $\{\check{\Gamma}_i\}$ is uniquely determined modulo P.

Let (S, φ, P) be a *P*-marking of a surface *S*. We consider the period

$$\left[\int_{\varphi^{-1}(\check{\Gamma}_1)}\Omega:\cdots:\int_{\varphi^{-1}(\check{\Gamma}_8)}\Omega\right]\in\mathbf{P}^7$$

where Ω is a holomorphic 2-form on S. The bilinear relation

$$\int_{S} \Omega \wedge \Omega = 0, \quad \int_{S} \bar{\Omega} \wedge \Omega > 0$$

implies that the period belongs to the domain

$$D = \{ \eta \in \mathbf{P}^7 : \quad {}^t \eta A \eta = 0, \quad {}^t \bar{\eta} A \eta > 0 \}.$$

The domain D has two connected components

$$D = D^+ \cup D^-, \quad D^\pm = \{(\eta_1, \cdots, \eta_8) \in D : \pm \operatorname{Im}(\eta_3/\eta_1) > 0\}$$

and we can take D^+ as the period domain for the family of isomorphism classes of marked surfaces $\{(S(x),\varphi,P):\ x\in X'\}$.

Remark 3.4. Two domains D^{\pm} are complex conjugate. That is

$$\left[\int_{\varphi^{-1}(\check{\Gamma}_1)}\Omega:\cdots:\int_{\varphi^{-1}(\check{\Gamma}_8)}\Omega\right]\in D^+\quad\Leftrightarrow\quad \left[\int_{\varphi^{-1}(\check{\Gamma}_1)}\bar{\Omega}:\cdots:\int_{\varphi^{-1}(\check{\Gamma}_8)}\bar{\Omega}\right]\in D^-.$$

The group G acts on the domain D. Set

$$G(2)^+ = \{g \in G(2) : g(D^+) = D^+\}.$$

It is a subgroup of G(2) with index 2.

Theorem 3. Let (S, φ, P) and (S', φ', l'_s, P) be *P*-markings of extended double 4H surfaces S and S', respectively. Let $\eta, \eta' \in D^+$ be corresponding periods. Then these markings are isomorphic if and only if

$$g(\eta) = \eta'$$

for some $g \in G(2)^+$ (= O⁺(L, P) := { $g \in O(L, P) : g(D^+) = D^+$ }).

Proof. Assume $g(\eta) = \eta'$ for some $g \in G(2)^+$. According to Proposition 3.1 there exist an element $\hat{g} \in O(L, P)$ such that $\hat{g}|_T = g$. Then we have

$$\left[\int_{\varphi^{-1}\circ\hat{g}(\check{\Gamma}_1)}\Omega:\cdots:\int_{\varphi^{-1}\circ\hat{g}(\check{\Gamma}_8)}\Omega\right]=\left[\int_{(\varphi')^{-1}(\check{\Gamma}_1)}\Omega':\cdots:\int_{(\varphi')^{-1}(\check{\Gamma}_8)}\Omega'\right],$$

where Ω' indicates the holomorphic form on S'. Consider the composition

$$f = (\varphi')^{-1} \circ \hat{g}^{-1} \circ \varphi : \mathrm{H}_2(S, \mathbf{Z}) \longrightarrow L \longrightarrow \mathrm{H}_2(S', \mathbf{Z}).$$

The above composite isomorphism induces the dual map

$$f^*: \mathrm{H}^2(S', \mathbf{Z}) \longrightarrow \mathrm{H}^2(S, \mathbf{Z})$$

with $f^*(\mathrm{H}^{2,0}(S')) = \mathrm{H}^{2,0}(S)$. Moreover, f^* preserves ample classes. Hence the Torelli theorem for K3 surfaces assures that there exists the unique isomorphism $\rho : S \to S'$ such that $\rho_* = f$. It is obvious that ρ is an isomorphism of marked surfaces. The converse is derived by the same argument.

4 The hypergeometric differential equation for the periods

Let us take an element $C \in H_2(S_0, \mathbb{Z})$, and let $C(x) \in H_2(S(x), \mathbb{Z})$ denote its continuation to $x = (x^1, x^2, x^3, x^4) \in X^0$ (that is multivalued and depends on the paths to x in X°). Now we investigate the differential equation for the period

$$u(x) = \int_{C(x)} \Omega = \iint_{C(x)} \{ \prod_{p=1}^{4} (x_{11}^p \xi_1 \eta_1 + x_{12}^p \xi_1 \eta_2 + x_{21}^p \xi_2 \eta_1 + x_{22}^p \xi_2 \eta_2)^{-1/2} \} \omega$$

$$\omega = (\xi_1 d\xi_2 - \xi_2 d\xi_1) \wedge (\eta_1 d\eta_2 - \eta_2 d\eta_1), \qquad (4.1)$$

where $[\xi_1, \xi_2]$ and $[\eta_1, \eta_2]$ denote the homogeneous coordinates of $\mathbf{P}^1(s)$ and $\mathbf{P}^1(t)$, respectively. Note that this integral does not depend on the affine representatives of the homogeneous coordinate of $\mathbf{P}^1(s)$ and $\mathbf{P}^1(t)$.

There are left and right actions of $GL(2, \mathbb{C})$ and a multiplicative action of $(\mathbb{C}^*)^4$ on X^0 . They induce the following actions on the period u(x):

$$u(g \cdot x) = u(gx^{1}, \cdots, gx^{4}) \quad g \in GL(2, \mathbb{C}),$$
$$u(x \cdot h) = u(x^{1t}h, \cdots, x^{4t}h) \quad h \in GL(2, \mathbb{C}),$$
$$u(\lambda \circ x) = u(\lambda_{1}x^{1}, \cdots, \lambda_{4}x^{4}) \quad \lambda = (\lambda_{1}, \cdots, \lambda_{4}) \in (\mathbb{C}^{*})^{4}$$

Lemma 4.1. We have the following equalities.

(1)
$$u(\lambda \circ x) = \prod_{p=1}^{4} \lambda_p^{a_p - 1} u(x).$$
 (4.2)

(2)
$$u(g \cdot x) = \frac{1}{\det(g)}u(x), \quad u(x \cdot h) = \frac{1}{\det(h)}u(x).$$
 (4.3)

Proposition 4.1. The integral u(x) satisfies the following systems :

$$\sum_{1 \le j,k \le 2} x_{jk}^p \frac{\partial u}{\partial x_{jk}^p} = -\frac{1}{2} u(x) \quad (p = 1, 2, 3, 4), \tag{4.4}$$

$$E_1: \quad \sum_{p=1}^{4} \sum_{j=1}^{2} x_{lj}^p \frac{\partial u}{\partial x_{mj}^p} = -\delta_{lm} u(x) \quad (\ell, m \in \{1, 2\}), \tag{4.5}$$

$$\sum_{p=1}^{4} \sum_{j=1}^{2} x_{jl}^{p} \frac{\partial u}{\partial x_{jm}^{p}} = -\delta_{lm} u(x) \quad (\ell, m \in \{1, 2\}),$$
(4.6)

$$E_2: \quad \frac{\partial^2 u}{\partial x_{ij}^q \partial x_{k\ell}^p} = \frac{\partial^2 u}{\partial x_{k\ell}^q \partial x_{ij}^p} \quad (i, j, k, \ell \in \{1, 2\}, p, q \in \{1, 2, 3, 4\}), \tag{4.7}$$

$$\frac{\partial^2 u}{\partial x_{11}^q \partial x_{22}^p} = \frac{\partial^2 u}{\partial x_{21}^q \partial x_{12}^p} \quad (p, q \in \{1, 2, 3, 4\}).$$

$$(4.8)$$

[Proof] Differentiate the first equality (4.2) in Lemma 4.1 with respect to λ_i and put $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4) = (1, 1, 1, 1)$. Then we get the first equality (4.4).

Differentiate the second equality (4.3) in Lemma 3.1 for the left action with respect to the *ij*-component g_{ij} of g and put $g = I_2$. Then we get the second equality (4.5). We get the third equality by using the equality for the right action of PGL(2, **C**) with the same method. The system E_2 is deduced from the direct computation and the commutativity of the partial differentiations and the integral.

Now we show the system $E_1 + E_2$ is a holonomic system on X^0 . We consider the variety B in the cotangent bundle $T^*(X^\circ)$ defined by

$$\sum_{1 \le j,k \le 2} x_{jk}^p \xi_{jk}^p = 0 \quad (p = 1, 2, 3, 4), \tag{4.9}$$

$$\sum_{p=1}^{4} \sum_{j=1}^{2} x_{\ell j}^{p} \xi_{m j}^{p} = 0 \quad (\ell, m \in \{1, 2\}),$$
(4.10)

$$\sum_{p=1}^{4} \sum_{j=1}^{2} x_{j\ell}^{p} \xi_{jm}^{p} = 0 \quad (\ell, m \in \{1, 2\}),$$
(4.11)

$$\xi_{ij}^{q}\xi_{k\ell}^{p} - \xi_{k\ell}^{q}\xi_{ij}^{p} = 0 \quad (i, j, k, \ell \in \{1, 2\}, p, q \in \{1, 2, 3, 4\}),$$

$$(4.12)$$

$$\xi_{11}^q \xi_{22}^p - \xi_{21}^q \xi_{12}^p = 0 \quad (p, q \in \{1, 2, 3, 4\}), \tag{4.13}$$

where ξ_{jk}^p stands for $\frac{\partial u}{\partial x_{ij}^k}$. We regard *B* as a fiber space over X° in the product of X° and a space of symbols $\xi = (\xi^1, \xi^2, \xi^3, \xi^4)$ for

$$\xi^p = \begin{pmatrix} \xi^p_{11} & \xi^p_{12} \\ \xi^p_{21} & \xi^p_{22} \end{pmatrix}.$$

The characteristic variety itself does not necessarily coincide with B, but it contains B, which we will call the fake characteristic variety. We will prove that the fake characteristic variety consists of only $\{0\}$ (zero-section) on X^0 . It implies that any solution of $E_1 + E_2$ on a simply connected domain in X^0 is holomorphic.

Let us fix an arbitrary point

$$x = (x^1, x^2, x^3, x^4) \in X^0, \quad x^p = \begin{pmatrix} x^p_{11} & x^p_{12} \\ x^p_{21} & x^p_{22} \end{pmatrix} \quad (p = 1, 2, 3, 4).$$

Put

$$H_p: \quad f_p = x_{11}^p \xi_1 \eta_1 + x_{12}^p \xi_1 \eta_2 + x_{21}^p \xi_2 \eta_1 + x_{22}^p \xi_2 \eta_2 = 0,$$

and set

$$D(pq) = \text{Det}(x^p) \text{ Trace}((x^p)^{-1}x^q) = x_{11}^p x_{22}^q + x_{22}^p x_{11}^q - x_{12}^p x_{21}^q - x_{21}^p x_{12}^q.$$

We set

$$M_{pqr} = \begin{pmatrix} x_{11}^p & x_{12}^p & x_{21}^p & x_{22}^p \\ x_{11}^q & x_{12}^q & x_{21}^q & x_{22}^q \\ x_{11}^r & x_{12}^r & x_{21}^r & x_{22}^r \end{pmatrix}, \quad p,q,r \in \{1,2,3,4\},$$

and let (ijk) denote the 3×3 minor determinant of M_{pqr} induced from the i, j, k-th column vectors. Put

$$D(pqr) = (234)(123) - (134)(124).$$

Lemma 4.2. We have the following :

(1) $D(pq) = D(qp), \quad D(pp) = 2\det(x^p)$

(2) H_p is irreducible if and only if

 $D(pp) \neq 0$

(3) Let H_p and H_q be different and both irreducible. Then they have different intersection points if and only if

$$D(pq)^2 - D(pp)D(qq) \neq 0.$$

(4) We have $H_p \cap H_q \cap H_r = \emptyset$ if and only if $D(pqr) \neq 0$.

[Proof].

The first two claims are obvious. So we consider the third statement. H_p is expressed in the form

$$\frac{\xi_2}{\xi_1} = -\frac{x_{11}^p \eta_1 + x_{12}^p \eta_2}{x_{21}^p \eta_1 + x_{22}^p \eta_2}.$$

So we obtain the intersections $H_p \cap H_q$ from

$$\frac{x_{11}^p \eta_1 + x_{12}^p \eta_2}{x_{21}^p \eta_1 + x_{22}^p \eta_2} = \frac{x_{11}^q \eta_1 + x_{12}^q \eta_2}{x_{21}^q \eta_1 + x_{22}^q \eta_2}.$$

Then the intersection comes from the eigen vector of $(x_p)^{-1}x^q$. Hence we get the required condition.

Next we examine the last statement. Generally we have

$$M_{pqr} \begin{pmatrix} (234) \\ -(134) \\ (124) \\ -(123) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

Note that we have $H_p \cap H_q \cap H_r \neq \emptyset$ if and only if there exists a solution $\zeta = {}^t(\zeta_1, \zeta_2, \zeta_3, \zeta_4)$ with $\zeta_1 \zeta_4 - \zeta_2 \zeta_3 = 0$ for

$$M_{pqr} \begin{pmatrix} \zeta_1 \\ \zeta_2 \\ \zeta_3 \\ \zeta_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

If rank $M_{pqr} < 3$ we can find easily such a solution ζ . In case rank $M_{pqr} = 3$ we have 1 dimensional solution space for $M_{pqr}\zeta = 0$. So t((234), -(134), (124), -(123)) becomes a required solution only when D(pqr) = 0.

q.e.d

Theorem 4. Let $x = (x^1, x^2, x^3, x^4) \in M(2, \mathbb{C})^4$ be a point on X^0 , namely x satisfies the conditions:

(g1). Any H_p is irreducible i.e. $D(pp) \neq 0$,

(g2). H_p and H_q have 2 different intersections i.e. $D(pq)^2 - D(pp)D(qq) \neq 0$ for any $i \neq j$

(g3). $H_p \cap H_q \cap H_r = \emptyset$ i.e $D(pqr) \neq 0$ for any $(p-q)(q-r)(r-p) \neq 0$

Then any local solution of the system $E_1 + E_2$ around the point $x \in X^0$ is locally holomorphic.

[Proof]. We show $B_x = \{0\}$ in several steps.

(Step1). Let us consider the Segre embedding

$$\psi: \mathbf{P}^1 \times \mathbf{P}^1 \longrightarrow \mathbf{P}^3,$$

that is defined by

$$\begin{bmatrix} \xi_1, \xi_2 \end{bmatrix} \times \begin{bmatrix} \eta_1, \eta_2 \end{bmatrix} \mapsto \begin{pmatrix} \xi_1 \eta_1 & \xi_1 \eta_2 \\ \xi_2 \eta_1 & \xi_2 \eta_2 \end{pmatrix}$$

Note that the point $\begin{pmatrix} \xi_{11} & \xi_{12} \\ \xi_{21} & \xi_{22} \end{pmatrix} \in \mathbf{P}^3$ belongs to $\operatorname{Im}(\psi)$ if and only if $\xi_{11}\xi_{22} - \xi_{12}\xi_{21} = 0$. Let Δ denote the diagonal map

$$\Delta: \mathbf{P}^3 \longrightarrow \mathbf{P}^3 \times \mathbf{P}^3 \times \mathbf{P}^3 \times \mathbf{P}^3, \qquad P \mapsto (P, P, P, P).$$

 Set

$$\xi^p = \begin{pmatrix} \xi^p_{11} & \xi^p_{12} \\ \xi^p_{21} & \xi^p_{22} \end{pmatrix},$$

and regard $\xi = (\xi^1, \xi^2, \xi^3, \xi^4)$ as a homogeneous coordinate on $\mathbf{P}^3 \times \mathbf{P}^3 \times \mathbf{P}^3 \times \mathbf{P}^3$. Then the system E_2 in Proposition 4.1 determines exactly the image of $\Delta \circ \psi$.

(Step 2). Let $\xi = (\xi^1, \xi^2, \xi^3, \xi^4)$ be a point on B_x . Then at least two of $\{\xi^p\}$ should be O. Suppose in contrary three of them , saying ξ^1, ξ^2, ξ^3 , are not equal to O. According to the argument in Step 1 they have the pull backs in $\mathbf{P}^1 \times \mathbf{P}^1$, and these are the same point $P = ([\xi_1, \xi_2], [\eta_1, \eta_2])$. Replace ξ^p by $\psi(P)$ in (4.9), which yields $P \in H_p$. Hence we obtain $P \in H_1 \cap H_2 \cap H_3$. This contradicts (g3).

(Step 3). Let $\xi = (\xi^1, \xi^2, \xi^3, \xi^4)$ be a point on B_x . If three of $\{\xi^p\}$ are O, then all of them equal O.

Suppose $\xi^2 = \xi^3 = \xi^4 = 0$. Then the equation (4.10) reduces to

$$x^{1t}\xi^1 = O.$$

Because x^1 is assumed to be invertible, we obtain $\xi^1 = O$.

(Step 4). We don't have the case $\xi^1 \neq 0, \xi^2 \neq 0, \xi^3 = \xi^4 = 0$.

Suppose it happens, then we have $\xi^2 = c\xi^1$ for some constant $c \neq 0$. The two equations (4.10) and (4.11) are expressed in the form

$$x^{1} {}^{t}\xi^{1} + x^{2} {}^{t}\xi^{2} = 0, \quad {}^{t}\xi^{1} x^{1} + {}^{t}\xi^{2} x^{2} = 0.$$

Because we supposed x^2 to be invertible, we have

$$(x^2)^{-1} x^1 {}^t \xi^1 = {}^t \xi^1 x^1 (x^2)^{-1}.$$

So we get 4 linear equations for 4 unknowns $\xi_{11}^1, \xi_{12}^1, \xi_{21}^1, \xi_{22}^1$. Together with the 2 equations coming from the first equation (4.9) for p = 1, 2 we obtain the system of linear equations

$$M\begin{pmatrix} \xi_{11}^1\\ \xi_{12}^1\\ \xi_{21}^1\\ \xi_{22}^1 \end{pmatrix} = \begin{pmatrix} 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0 \end{pmatrix} \qquad (*).$$

By a direct calculation we have

$$M = \begin{pmatrix} x_{12}^1 x_{21}^2 - x_{21}^1 x_{12}^2 & x_{12}^1 x_{22}^2 - x_{22}^1 x_{12}^2 & x_{22}^1 x_{21}^2 - x_{21}^1 x_{22}^2 & 0 \\ x_{11}^1 x_{12}^2 - x_{12}^1 x_{11}^2 & 0 & x_{11}^1 x_{22}^2 - x_{22}^1 x_{11}^2 & x_{12}^1 x_{22}^2 - x_{22}^1 x_{12}^2 \\ x_{21}^1 x_{11}^2 - x_{11}^1 x_{21}^2 & x_{22}^1 x_{11}^2 - x_{11}^1 x_{22}^2 & 0 & x_{22}^1 x_{21}^2 - x_{12}^1 x_{22}^2 \\ 0 & x_{11}^1 x_{12}^2 - x_{12}^1 x_{11}^2 & x_{21}^1 x_{11}^2 - x_{11}^1 x_{22}^2 & x_{21}^1 x_{12}^2 - x_{12}^1 x_{21}^2 \\ x_{11}^1 & x_{12}^1 & x_{21}^1 x_{21}^1 - x_{11}^1 x_{21}^2 & x_{21}^1 x_{22}^2 - x_{12}^1 x_{21}^2 \\ x_{11}^2 & x_{12}^2 & x_{21}^2 & x_{22}^2 \end{pmatrix}$$

Let (i, j, k, ℓ) denote the 4 × 4 minor of M obtained by taking i, j, k, ℓ -th row vectors. By assumption the system (*) has a nontrivial solution ξ^1 , so every (i, j, k, ℓ) should be 0. By a direct calculation we have the following:

$$\begin{split} (1256) &= (x_{12}^1 x_{22}^2 - x_{22}^1 x_{12}^2) (D(12)^2 - D(11)D(22)), \\ (1356) &= (x_{22}^1 x_{21}^2 - x_{21}^1 x_{22}^2) (D(12)^2 - D(11)D(22)), \\ (1456) &= (x_{21}^1 x_{12}^2 - x_{12}^1 x_{21}^2) (D(12)^2 - D(11)D(22)), \\ (2356) &= (x_{11}^1 x_{22}^2 - x_{22}^1 x_{11}^2) (D(12)^2 - D(11)D(22)), \\ (2456) &= (x_{12}^1 x_{11}^2 - x_{11}^1 x_{12}^2) (D(12)^2 - D(11)D(22)), \\ (3456) &= (x_{11}^1 x_{21}^2 - x_{21}^1 x_{11}^2) (D(12)^2 - D(11)D(22)), \end{split}$$

and other 4×4 minors are 0. Here recall the 2nd assumption for x. So we have $(D(12)^2 - D(11)D(22))$ is not 0. Hence we obtain

$$x_{ij}^1 x_{k\ell}^2 = x_{ij}^2 x_{k\ell}^1$$

for any indices $i, j, k, \ell \in \{1, 2\}$. It means $x^2 = cx^1$ and $H_1 = H_2$. This is a contradiction.

By the combination of the above arguments we obtain $B_x = \{0\}$.

q.e.d.

The system of partial differential equations $E_1 + E_2$ is closely related to the GKZ hypergeometric system introduced by Gel'fand, Kapranov and Zelevinski. Let us explain the relation and evaluate the dimension of the solution space of $E_1 + E_2$.

Define a set of operators $E_1(\ell, m)$ and $E'_1(\ell, m)$ by

$$E(\ell,m) : \sum_{p,j} x_{\ell j}^p \partial_{m j}^p + \delta_{\ell m},$$

$$E'(\ell,m) : \sum_{p,j} x_{j\ell}^p \partial_{jm}^p + \delta_{\ell m}.$$

Let D be the Weyl algebra

 $\mathbf{C}\langle x_{11}^1, x_{12}^1, x_{21}^1, x_{22}^1, x_{11}^2, \dots, x_{22}^4, \partial_{11}^1, \partial_{12}^1, \partial_{21}^1, \partial_{22}^1, \partial_{11}^2, \dots, \partial_{22}^4\rangle.$

Consider the GKZ-hypergeometric ideal $H_A(\beta)$ in D associated to the matrix

A =	(1)	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0 \
	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0	0
	0	0	0	0	0	0	0	0	1	1	1	1	0	0	0	0
	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1
	1	1	0	0	1	1	0	0	1	1	0	0	1	1	0	0
	$\setminus 1$	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0 /

and $\beta = (-1/2, -1/2, -1/2, -1/2, -1, -1).$

Proposition 4.2. Our system of partial differential equations E_1 and E_2 consists of three groups of operators

$$A \cdot \theta - {}^{t}\beta,$$

$$I_{A} = \{\partial^{u} - \partial^{v} \mid Au = Av, \ u, v \in \mathbf{N}_{0}^{16}\},$$

$$E_{1}(\ell, m) \text{ and } E'_{1}(\ell, m)(\ell \neq m).$$

Here, we denote by θ the column vector $(x_{ik}^i \partial_{ik}^i)$ of Euler operators of length 16.

The GKZ-hypergeometric ideal $H_A(\beta)$ is the left ideal in D generated by $A \cdot \theta - {}^t\beta$ and I_A . Since the toric ideal I_A is homogeneous, the D-module $D/H_A(\beta)$ is regular holonomic on $X = \mathbf{C}^{16}$ (Hotta's theorem, see, e.g., [14, p.82]). Our toric ideal I_A satisfies the following properties:

- 1. The initial ideal of I_A with respect to the reverse lexicographic order is generated by square free monomials.
- 2. The toric ideal is Cohen-Macaulay.
- 3. The multiplicity of I_A is 20.
- 4. The variety $V(I_A)$ admits a natural action of $(\mathbf{C}^*)^4$ and $V(I_A)/(\mathbf{C}^*)^4$ is isomorphic to $\operatorname{Im}(\Delta \circ \psi)$ in Theorem 4.

The first statement can be easily checked by Buchberger's criterion. The second fact follows from the first (see, e.g., [14, p.153]). The third fact can be shown by computing the Hilbert polynomial of I_A on computers. Therefore, by theorems due to Gel'fand, Kapranov and Zelevinsky, the rank of the solution space of $H_A(\beta)$ is 20 and the singular locus agrees with the zero set of the principal A-determinant (see, e.g., [14, p.173]).

We denote by E the left ideal in D generated by first order operators $E_1(\ell, m)$, $(\ell \neq m)$ and the GKZ-hypergeometric ideal $H_A(\beta)$.

Theorem 5. The D-module D/E is regular holonomic on X, and the rank of the solution space of $E = E_1 + E_2$ is equal to 8.

[Proof]. Since $H_A(\beta) \subseteq E$ holds and the GKZ system is regular holonomic, the *D*-module D/E is also regular holonomic on *X*. The differential operators $E_1(\ell, m)$, $(\ell \neq m)$ are used to extract the space of the period maps from the 20 dimensional solution space of the GKZ system.

Next let us show the second statement. Already in Section 3 we constructed 8 independent periods. So it is enough to say that "at most 8 dimensional". Although it is possible in

principle to evaluate the rank by computer and Oaku's algorithm (see, e.g., [14, p.31]), we could not evaluate it because of an exhaustion of memory. So, we try to find sufficiently many initial terms for the left ideal E and a suitable weight. For this system, we chose a weight $w^1 = 1 \cdot w$, $w^2 = 4 \cdot w$, $w^3 = 9 \cdot w$, $w^4 = 16 \cdot w$, $w = \begin{pmatrix} 1 & 2 \\ 0 & 4 \end{pmatrix}$. Put $W = (w^1, w^2, w^3, w^4)$. The initial ideal generated by $in_{(-W,W)}(E_1)$ and $in_{(-W,W)}(E_2)$ has the rank 20, which means that we do not have sufficiently many initial terms. We computed a partial Gröbner basis of E_1 and E_2 with the weight (-W, W) up to the degree 7 in the homogenized Weyl algebra. The ideal generated by the initial terms of the partial Gröbner basis has rank 8. It follows from Theorem 2.5.1 of [14] and the argument of the regular holonomicity that the rank is bounded by 8.

q.e.d.

According to Theorem 4 and Theorem 5 the solution space of $E_1 + E_2$ looks like a vector bundle over X° of rank 8. Let us take a *P*-marking $(S(x_0), \varphi, P)$, then we can choose the basis for the solution space of $E_1 + E_2$

$$\{\int_{\varphi^{-1}(\check{\Gamma}_1)}\Omega,\cdots,\int_{\varphi^{-1}(\check{\Gamma}_8)}\Omega\}$$

at $x_0 \in X^\circ$. So the system $E_1 + E_2$ induces a representation of $\pi_1(X^\circ, x_0)$ over $GL(8, \mathbb{Z})$.

Definition 4.1. The monodromy group Mono for (\mathcal{F}, X°) is the image of this representation.

Remark 4.1. Set $\Sigma = (GL(2, \mathbb{C}) \setminus X^{\circ}/GL(2, \mathbb{C}))/(\mathbb{C}^{*})^{4}$. According to Lemma 4.1 we have the same period along the orbit of the actions of $GL(2, \mathbb{C})$ and $(\mathbb{C}^{*})^{4}$. So the above monodromy representation reduces to that of $\pi_{1}(\Sigma, *)$.

Let B_r be the sublattice of L generated by the elements $f_s, f_t, g_i \ (i = 1, \dots, 4)$. Put

$$O(L, Br) = \{g \in O(L) : g(P) = P, g \mid_{Br} = \mathrm{id}\}.$$

We define

$$O^+(L, Br) = \{ g \in O(L, Br) : g \mid_T (D^+) = D^+ \}.$$

The group $\pi_1(X^\circ, x_0)$ acts on $L \cong H_2(S(x_0), \mathbb{Z})$ by the natural way, and the branch locus is fixed under this action. So Mono is identified with a certain subgroup of $O^+(L, Br)$.

Lemma 4.3. Let (S_1, φ_1, P) and (S_2, φ_2, P) be *P*-markings, and let x_1, x_2 be corresponding points on X° . Suppose there is an isomorphism $\rho : S_1 \mapsto S_2$ such that we have $\varphi_2 \circ \rho_* \circ \varphi_1^{-1} \in O^+(L, Br)$. Then the points x_1 and x_2 determine the same point on $\Sigma = (GL(2, \mathbb{C}) \setminus X^\circ/GL(2, \mathbb{C}))/(\mathbb{C}^*)^4$.

[Proof].

Because we have $\rho_*\varphi_1^{-1}(f_s) = \varphi_2^{-1}(f_s)$ and $\rho_*\varphi_1^{-1}(f_s) = \varphi_2^{-1}(f_s)$, the isomorphism ρ preserves the covering structure over $\mathbf{P}^1 \times \mathbf{P}^1$. Moreover we have $\rho_* \circ \varphi_1^{-1}(g_i) = \varphi_2^{-1}(g_i)$ (i = 1, 2, 3, 4). Hence ρ preserves the branch locus with the same numbering order. So x_1 and x_2 have the required property.

q.e.d.

Theorem 6. We have

$$O^+(L, Br) \mid_T =$$
Mono.

[Proof]. It is enough to show that $O^+(L, Br) |_T \subset$ Mono. Let us fix a *P*-marking (S_0, φ_0, P) corresponding to the initial point $x_0 \in X^\circ$. Let g be an arbitrary element of $O^+(L, Br)$, and put $g' = g |_T$. Let $\eta_0 \in D^+$ be the period point determined by (S_0, φ_0, P) , and set $\eta_1 = g'(\eta_0)$. Let us take an oriented arc r in D^+ that starts from η_0 and ends at η_1 . Because of the surjectivity of the period map, we can find a *P*-marking $(S(\eta), \varphi(\eta), P)$ for every point η on r. Set (S_1, φ_1, P) be the terminal marking and let x_1 be the corresponding point on X° . Because of the injectivity of the period map we have $(S_1, \varphi_1, P) = (S_1, \varphi_1 \circ g^{-1}, P)$. Then by virtue of the above Lemma x_0 and x_1 determine the same point on Σ . Hence we get unique closed arc γ in Σ corresponding to r. It means that g' is the monodromy transformation coming from the arc γ .

q.e.d.

5 Periods and Kuga-Satake varieties

In this section we construct the abelian variety attached to the extended double 4H surface starting from its period. The reader will find the method in [16] and [13]. The detailed calculation and argument is exposed in [4] also.

Let us consider the lattice T defined by the intersection matrix $A = U(2) \oplus U(2) \oplus (-2I_4)$ and put $V_k = T \otimes k$ ($k = \mathbf{R}$ or \mathbf{Q}). Let Q(x) denote the quadratic form on T and at the same time on V_k . Let Tens(T) and $Tens(V_k)$ be the corresponding tensor algebras. And we let $Tens^+(T)$ and $Tens^+(V_k)$ denote the subalgebras composed of the parts with even degree in Tens(T) and $Tens(V_k)$ respectively. We consider the two sided ideal I in $Tens^+(V_k)$ generated by the elements of the form $x \otimes x - Q(x)$ for $x \in V_k$, and the ideal $I_{\mathbf{Z}}$ in Tens(T) by the same manner. The corresponding even Clifford algebra is defined by

$$C^+(V_k, Q) = Tens^+(V_k)/I.$$

By the same manner we define the even Clifford algebra over \mathbf{Z} by

$$C^+(T,Q) = Tens^+(T)/I_{\mathbf{Z}}.$$

We note that $C^+(V_{\mathbf{R}}, Q)$ is a 128 dimensional real vector space and $C^+(T, Q)$ is a lattice in it. So we obtain a real torus

$$\mathcal{T}_{\mathbf{R}} = C^+(V_{\mathbf{R}}, Q) / C^+(T, Q).$$

Let \mathbf{F} denote the quaternion algebra

$$\mathbf{Q} \oplus \mathbf{Q}i \oplus \mathbf{Q}j \oplus \mathbf{Q}ij$$

with $i^2 = j^2 = -1$. By some routine calculations of Clifford algebra we obtain the following.

Proposition 5.1. We have the isomorphism of algebras $C^+(V_{\mathbf{Q}}, Q) \cong M(4, \mathbf{F}) \oplus M(4, \mathbf{F})$.

Let a complex vector $\underline{\eta} = (\eta_1, \dots, \eta_8)$ be a representative of a point $\eta = [\eta_1, \dots, \eta_8] \in D^+$. So it has an ambiguity of the multiplication by a non zero complex number. Put $\underline{\eta} = s + it$ $(s, t \in \mathbb{R}^8)$. If we impose the condition $(st)^2 = -1$ in $C^+(V_{\mathbf{R}}, Q)$, the representative is uniquely determined up to a multiplication by a complex unit. We denote it by

$$\eta = m_1(\eta) + im_2(\eta),$$

and put

$$m(\eta) = m_1(\eta)m_2(\eta).$$

It is uniquely determined by η without any ambiguity. According to the imposed condition the element $m(\eta) \in C^+(V_{\mathbf{R}}, Q)$ defines a complex structure on $C^+(V_{\mathbf{R}}, Q)$ by the left action. It induces a complex structure on the real torus $\mathcal{T}_{\mathbf{R}}$ also. We denote this complex torus by $(T, m(\eta))$.

Let $\{\varepsilon_1, \dots, \varepsilon_8\}$ be the basis of T with the intersection matrix $U(2) \oplus U(2) \oplus (-2I_4)$. And let $\{e_1, \dots, e_8\}$ be a orthonormal basis of V given by

$$(e_1, \cdots, e_8) = (\varepsilon_1, \cdots, \varepsilon) \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & 0 & 0\\ 0 & 0 & \frac{1}{2} & -\frac{1}{2}\\ \frac{1}{2} & -\frac{1}{2} & 0 & 0\\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \oplus (I_4)).$$

Then the corresponding intersection matrix takes the form $I_2 \oplus (-I_2) \oplus (-2I_4)$.

Let ι be an involution on $C^+(V,Q)$ induced from the transformation

$$\iota: e_{i_1} \otimes e_{i_2} \otimes \cdots \otimes e_{i_k} \mapsto e_{i_k} \otimes \cdots \otimes e_{i_2} \otimes e_{i_1}$$

for the basis. Set $\alpha = 4e_2e_1$. According to the method in [St] we know that

$$E(x,y) = tr(\alpha x^{\iota} y)$$

determines a Riemann form on $(L, m(\eta))$. We denote this abelian variety by $A^+(\eta)$, that is so called the Kuga-Satake variety attached to the K3 surface corresponding to the period η . In this way we can construct a family of abelian varieties

$$\mathcal{A}^{+} = \{ A^{+}(\eta) : \eta \in D^{+} \}$$

induced from the lattice T parameterized by the domain D^+ . We can construct the "conjugate family"

$$\mathcal{A}^- = \{ A^-(\eta) : \eta \in D^- \}$$

parameterized by

$$D^{-} = \{ \eta = [\eta_1, \cdots, \eta_8] : {}^t \eta A \eta = 0, \; {}^t \overline{\eta} A \eta > 0, \; \Im(\eta_3/\eta_1) < 0 \}$$

by the same procedure with the Riemann form $E^{-}(x,y) = -\operatorname{tr}(\alpha x^{\iota} y)$. The right action of $C^{+}(V_{\mathbf{Q}}, Q)$ on $(V, T, m(\eta))$ commutes with the left action of $\alpha(\eta)$. So we have

$$C^+(T_{\mathbf{Q}}) \subset \operatorname{End}(\mathrm{A}^{\pm}(\eta)) \otimes \mathbf{Q}$$

for any $A^{\pm}(\eta)$. For a general member $\eta \in \mathcal{D}^+$, the endmorphism ring is given by

$$\operatorname{End}_{\mathbf{Q}}(\mathbf{A}(\eta)) = \operatorname{End}(\mathbf{A}(\eta)) \otimes \mathbf{Q} \cong \mathrm{C}^+(\mathrm{V}_{\mathbf{Q}}).$$

According to Proposition 5.1 we obtain

Theorem 7. For a general member $\eta \in \mathcal{D}^+$, $A^+(\eta)$ is isogeneous to a product of abelian varieties $(A_1(\eta) \times A_2(\eta))^4$ where $A_1(\eta)$ and $A_2(\eta)$ are 8-dimensional simple abelian varieties with $\operatorname{End}_{\mathbf{Q}}(A_i(\eta)) = \mathbf{F}$ (i = 1, 2).

Remark 5.1. Here we describe the relation between $A_1(\eta)$ and $A_2(\eta)$. Now we define the linear involution * on $V_{\mathbf{R}}$ by

$$e_1^* = -e_1$$
 and $e_i^* = e_i$ $(i = 2, \dots, 8)$.

It can be extended on $C^+(V_{\mathbf{R}}, Q)$ as an autmorphism of algebra. We define an involution σ on \mathcal{D} :

$$\sigma: \mathcal{D} \longrightarrow \mathcal{D}, \quad (\eta_1, \cdots, \eta_8) \mapsto (-\eta_2, -\eta_1, \eta_3, \cdots \eta_8).$$

So we have $\mathcal{D}^{\sigma}_{+} = \mathcal{D}_{-}$. It is easy to check that we have

$$A_2(\eta) \sim A_1(\eta^{\sigma}), \quad A_1(\eta) \sim A_2(\eta^{\sigma}),$$

where \sim indicates the isogenous relation.

References

- [1] W. Barth, C. Peter and A. van de Ven, Compact complex surfaces, Springer(1984).
- [2] E. Freitag and C. F. Hermann, Some modular varieties of low dimension, preprint.
- [3] K. Kodaira, On analylic surfaces II, Ann. Math., 77(1963), 563 626.
- [4] K. Koike, The Kuga Satake variety attached to the double covering of $\mathbf{P}^1 \times \mathbf{P}^1$ branched along four curves of bidegree (1, 1), preprint.
- [5] M. Kitazume, K. Koike, and H. Shiga, Notes on a certain integral orthogonal group and its quotient by the principal congruence subgroup of level 2, to be appeared in Kyushu J. Math. .
- [6] K. Matsumoto, Theta functions on the bounded symmetric domain of type $I_{2,2}$ and the period map of a 4 parameter family of K3 surfaces, Math. Ann., 295(1993), 383 409.
- [7] K. Matsumoto and T. Sasaki, On the system of differential equations associated with a quadric and hyperplanes, Kyushu J. Math., 50(1996), 93 131.
- [8] K. Matsumoto, T. Sasaki and M. Yoshida, The monodromy of the period map of a 4 parameter family of K3 surfaces and the hypergeometric function of type (3,6), Int. J. Math., 3(1992), 1 - 164.
- [9] K. Matsumoto and M. Yoshida, Certain reflection groups acting on symmetric domains of type IV, Mem. Fac. Sci. Kyushu Univ., Ser. A, vol. 46(1992), 115 - 128.
- [10] V. V. Nikulin, Integral symmetric bilinear forms and some of their applications, Math. USSR Izv., 14(1980), 103 - 166.
- [11] K. Paranjape, Abelian varieties associated to certain K3 surfaces, Compositio Math., 68(1988), 11 - 22.
- [12] I. I. Pjateckii Šapiro and Šafarevič, A Torelli theorem for algebraic surfaces of type K3, Math. USSR Izv., 5(1971), No.3, 547 - 588.

- [13] I. Satake, Clifford algebras and families of Abelian varieties, Nagoya Math.J. 27(1966), 435-466.
- [14] M. Saito, B. Sturmfels and N. Takayama, Gröbner Deformations of Hypergeometric Differential Equations. Algorithms and Computation in Mathematics, Vol. 6, Springer(1999).
- [15] H. Shiga, On the arithmetic properties of the modular function induced from the hypergeometric differential equation of type (3,6), Proc. 7th Int. Coll. Differential Equations Provdiv 1996, 381 - 388.
- [16] B. van Geemen, Kuga Satake varieties and the Hodge conjecture, To appear in Proc. NATO ASI and CRM Summer School Banff 1998.

T. Tsutsui

Department of Mathematics and Informatics Faculty of Science Chiba University Yayoi-cho Inage-ku Chiba 263-8522, JAPAN

K. Koike, H. Shiga

Graduate School of Science Chiba University Yayoi-cho Inage-ku Chiba 263-8522, JAPAN

N. Takayama

Department of Mathematics Faculty of Science Kobe University Rokko Kobe 657-8501, JAPAN