Universal Similarity Factorization Equalities over Complex Clifford Algebras

Yongge Tian Department of Mathematics and Statistics Queen's University Kingston, Ontario, Canada K7L 3N6 e-mail:ytian@mast.queensu.ca

Abstract

A set of valuable universal similarity factorization equalities are established over complex Clifford algebras C_n . Through them matrix representations of complex Clifford algebras C_n can directly be derived, and their properties can easily be determined.

AMS Subject Classification: 15A23, 15A66. Keywords: Clifford algebras, matrix representations, universal similarity factorizations.

1 Introduction

Let C_n be the complex Clifford algebra, with the identity 1, defined on n generators e_1, e_2, \dots, e_n subject to the multiplication laws:

$$e_i^2 = -1, \quad i = 1, 2, \cdots, n,$$
 (1)

$$e_i e_j + e_j e_i = 0, \quad i \neq j, \ i, j = 1, 2, \cdots, n,$$
(2)

and $e_1e_2\cdots e_n \neq \pm 1$. In that case \mathcal{C}_n is spanned as a 2^n -dimensional vector space with 2^n basis $\{e_A\}$, where the multi index A ranges all naturally ordered subsets of the first positive integer set $\{1, 2, \dots, n\}$; the basis element e_A , where $A = (i_1, i_2, \dots, i_k)$ with $1 \leq i_1 < i_2 < \dots < i_k \leq n$, is defined as the product

$$e_A \equiv e_{(i_1, i_2, \cdots, i_k)} = e_{i_1} e_{i_2} \cdots e_{i_k}, \quad e_{A=\emptyset} = 1.$$

For simplicity, the volume element $e_{12\dots n} = e_1 e_2 \cdots e_n$ of \mathcal{C}_n will be denoted by $e_{[n]}$ in the sequel. The square of the volume element is

$$e_{[n]}^2 = (-1)^{\frac{1}{2}n(n-1)} e_1^2 e_2^2 \cdots e_n^2.$$

In that case, all $a \in \mathcal{C}_n$ can be expressed as

$$a = \sum_{A} a_A e_A, \quad a_A \in \mathcal{C},$$

where A ranges all naturally ordered subsets of $\{1, 2, \dots, n\}$. We shall adopt the following notation from now on: $C_n := C\{e_1, \dots, e_n\}$.

Clifford algebras (real or complex) have been studied for many years and their algebraic properties are well known. In particular, all Clifford algebras are classified as matrix algebras, or as direct sums of matrix algebras over the fields of real or complex numbers, or the quaternion ring (see, e.g., [2, 3, 5, 7, 8, 9, 10, 11]). For complex Clifford algebras C_n , it is well known that they can faithfully be realized as certain matrix algebras over C, and the general algebraic isomorphism is

$$C_n \simeq \begin{cases} \mathcal{C}(2^{\frac{n}{2}}) & \text{if } n \text{ is even,} \\ {}^2\mathcal{C}(2^{\frac{n-1}{2}}) & \text{if } n \text{ is odd,} \end{cases}$$
(3)

where $\mathcal{C}(s)$ stands for the $s \times s$ total complex matrix algebra, and ${}^{2}\mathcal{C}(s)$ stands for the complex matrix algebra

$${}^{2}\mathcal{C}(s) = \left\{ \left[\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right] \middle| A, B \in \mathcal{C}(s) \right\}.$$

In this article we improve this relationship to a new level by establishing a set of valuable universal similarity factorization equalities between elements of C_n and complex matrices over $C(2^{\frac{n}{2}})$ or $C(2^{\frac{n-1}{2}})$ for all n. Through these universal factorization equalities the complex matrix representations of elements in C_n can explicitly be established. Based on them various results in the complex matrix theory can directly be extended to complex Clifford numbers. Moreover, one can also easily develop matrix analysis over complex Clifford algebras.

We first establish some basic universal similarity factorization equalities for elements in Clifford algebras with dimensions 2 and 4, which will be directly applied to establish some more general results for C_n .

Lemma 1. Let \mathcal{F} be an algebraically closed field and $\mathcal{F}_1 = \mathcal{F}\{e \mid e^2 = u\}$ be a Clifford algebra defined on a generator e with $e^2 = u \in \mathcal{F}$ and $u \neq 0$. Then all $a \in \mathcal{F}_1$ can be written as $a = a_0 + a_1 e$, where $a_0, a_1 \in \mathcal{F}$. Moreover, define $\overline{a} = a_0 - a_1 e$. In that case, a and \overline{a} satisfy the following universal similarity factorization equality

$$P\begin{bmatrix}a & 0\\ 0 & \overline{a}\end{bmatrix}P^{-1} = \begin{bmatrix}a_0 + \sqrt{u}a_1 & 0\\ 0 & a_0 - \sqrt{u}a_1\end{bmatrix},\tag{4}$$

where P and P^{-1} have the following universal forms (no relation with a):

$$P = \frac{1}{2} \begin{bmatrix} 1 + \frac{1}{\sqrt{u}}e & -(\sqrt{u} - e) \\ \frac{1}{u}(\sqrt{u} - e) & 1 + \frac{1}{\sqrt{u}}e \end{bmatrix},$$

$$P^{-1} = \frac{1}{2} \begin{bmatrix} 1 + \frac{1}{\sqrt{u}}e & \sqrt{u} - e \\ -\frac{1}{u}(\sqrt{u} - e) & 1 + \frac{1}{\sqrt{u}}e \end{bmatrix}.$$
 (5)

Proof. Note that $e^2 = u$. It is easy to verify that

$$\begin{bmatrix} 1 & e \\ e^{-1} & -1 \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & \overline{a} \end{bmatrix} \begin{bmatrix} 1 & e \\ e^{-1} & -1 \end{bmatrix} = \begin{bmatrix} a_0 & ua_1 \\ a_1 & a_0\overline{a} \end{bmatrix}.$$

On the other hand, it is also easy to verify that

$$\begin{bmatrix} 1 & \sqrt{u} \\ \frac{1}{\sqrt{u}} & -1 \end{bmatrix} \begin{bmatrix} a_0 + ua_1 & 0 \\ 0 & a_0 - ua_1 \end{bmatrix} \begin{bmatrix} 1 & \sqrt{u} \\ \frac{1}{\sqrt{u}} & -1 \end{bmatrix} = \begin{bmatrix} a_0 & ua_1 \\ a_1 & a_0\overline{a} \end{bmatrix}$$

Combining the above two equalities yields (4) and (5). \Box

Lemma 2. Let $M_2(\mathcal{F})$ be the 2×2 total matrix algebra over an arbitrary field \mathcal{F} with its basis satisfying the multiplication rules

$$\tau_{pq}\tau_{st} = \begin{cases} \tau_{pt}, & q = s, \\ 0, & q \neq s, \end{cases}$$
(6)

for p, q, s, t = 1, 2. Then all $a = a_{11}\tau_{11} + a_{12}\tau_{12} + a_{21}\tau_{21} + a_{22}\tau_{22} \in M_2(\mathcal{F})$, where $a_{pq} \in \mathcal{F}$, satisfy the following universal similarity factorization equality

$$Q\begin{bmatrix} a & 0\\ 0 & a \end{bmatrix} Q^{-1} = \begin{bmatrix} a_{11} & a_{12}\\ a_{21} & a_{22} \end{bmatrix},$$
 (7)

where Q has the following universal form

$$Q = Q^{-1} = \begin{bmatrix} \tau_{11} & \tau_{21} \\ \tau_{12} & \tau_{22} \end{bmatrix}.$$
 (8)

Proof. Follows directly from a verification. \Box

Lemma 3. Let \mathcal{F} be an algebraically closed field of characteristic not two, and $\mathcal{F}_2 = \mathcal{F}\{e_1, e_2\}$ be a Clifford algebra defined on two generators e_1, e_2 with $e_1^2 = u \in \mathcal{F}, e_2^2 = v \in \mathcal{F}$ and $u \neq 0, v \neq 0$. Then all $a \in \mathcal{F}_2$ can be written as

$$a = a_0 + a_1 e_1 + a_2 e_2 + a_3 e_{12}, (9)$$

where $a_0, \ldots, a_3 \in \mathcal{F}$. In that case, aI_2 satisfies the following universal similarity factorization equalities

$$R\begin{bmatrix} a & 0\\ 0 & a \end{bmatrix} R^{-1} = \begin{bmatrix} a_0 + \sqrt{u}a_1 & v(a_2 + \sqrt{u}a_3)\\ a_2 - \sqrt{u}a_3 & a_0 - \sqrt{u}a_1 \end{bmatrix},$$
(10)

and

$$T\begin{bmatrix} a & 0\\ 0 & a \end{bmatrix} T^{-1} = \begin{bmatrix} a_0 + \sqrt{v}a_1 & u(a_1 - \sqrt{v}a_3)\\ a_1 + \sqrt{v}a_3 & a_0 - \sqrt{v}a_1 \end{bmatrix},$$
(11)

where

$$R = R^{-1} = \frac{1}{2} \begin{bmatrix} 1 + \frac{1}{\sqrt{u}}e_1 & e_2 - \frac{1}{\sqrt{u}}e_{12} \\ \frac{1}{v}(e_2 + \frac{1}{\sqrt{u}}e_{12}) & 1 - \frac{1}{\sqrt{u}}e_1 \end{bmatrix},$$
(12)

and

$$T = T^{-1} = \frac{1}{2} \begin{bmatrix} 1 + \frac{1}{\sqrt{v}}e_2 & e_1 + \frac{1}{\sqrt{v}}e_{12} \\ \frac{1}{u}(e_1 - \frac{1}{\sqrt{v}}e_{12}) & 1 - \frac{1}{\sqrt{v}}e_2 \end{bmatrix}.$$
 (13)

Proof. By Lemma 2, we take the change of basis of \mathcal{F}_2 as follows

$$\tau_{11} = \frac{1}{2} \left(1 + \frac{1}{\sqrt{u}} e_1 \right), \qquad \tau_{12} = \frac{1}{2v} \left(e_2 + \frac{1}{\sqrt{u}} e_{12} \right), \qquad (14a)$$

$$\tau_{21} = \frac{1}{2} \left(e_2 - \frac{1}{\sqrt{u}} e_{12} \right), \qquad \qquad \tau_{22} = \frac{1}{2} \left(1 - \frac{1}{\sqrt{u}} e_1 \right). \tag{14b}$$

Then it is not hard to verify that this new basis satisfies the multiplication rules in (6). In this new basis, any $a = a_0 + a_1e_1 + a_2e_2 + a_3e_{12} \in \mathcal{F}_2$ can be rewritten as

$$a = (a_0 + \sqrt{u}a_1)\tau_{11} + (va_2 + \sqrt{u}a_3)\tau_{12} + (a_2 - \sqrt{u}a_3)\tau_{21} + (a_0 - \sqrt{u}a_1)\tau_{22}.$$

Substituting this and (14) into (7) and (8), we can obtain (10) and (12). By the similar approach, we can get (11) and (13). \Box

2 Main results

Notice that C is an algebraically closed field. We then can establish a set of universal similarity factorization equalities for elements in C_n on the basis of Lemmas 1 and 3.

Theorem 4. Let $a \in C_1 = C\{e\}$ be given. Then *a* can be written as $a = a_0 + a_1 e$, where $a_0, a_1 \in C = \{x + iy \mid x, y \in \mathcal{R}\}$. Moreover denote by $\overline{a} = a_0 - a_1 e$, called the conjugate of *a*. Then *a* and \overline{a} satisfy the following universal similarity factorization equality

$$P_1 \begin{bmatrix} a & 0 \\ 0 & \overline{a} \end{bmatrix} P_1^{-1} = \begin{bmatrix} a_0 + a_1 i & 0 \\ 0 & a_0 - a_1 i \end{bmatrix},$$
(15)

where P_1 and P_1^{-1} have the following universal forms (no relation with a):

$$P_1 = \frac{1}{2} \begin{bmatrix} 1 - ie & -(i - e) \\ -(i - e) & 1 - ie \end{bmatrix}, \quad P_1^{-1} = \frac{1}{2} \begin{bmatrix} 1 - ie & i - e \\ i - e & 1 - ie \end{bmatrix}.$$
 (16)

Proof. Let $\mathcal{F} = \mathcal{C}$ and u = -1 as in Lemma 1. Then (15) and (16) follow directly from (4) and (5). \Box

It is easy to verify that

$$\overline{\overline{a}} = a, \quad \overline{a+b} = \overline{a} + \overline{b}, \quad \overline{ab} = \overline{a}\overline{b}, \quad \overline{\lambda a} = \overline{a\lambda} = \lambda \overline{a}.$$
(17)

hold for all $a, b \in C_1$ and $\lambda \in C$. According to (15), we define a map from C_1 to the double field $C \oplus C$ by

$$\phi_1: a = a_0 + a_1 e \in \mathcal{C}_1 \longrightarrow \begin{bmatrix} a_0 + a_1 i & 0\\ 0 & a_0 - a_1 i \end{bmatrix} \in \mathcal{C} \oplus \mathcal{C}.$$
(18)

Then it is easy to derive from (15) and (17) the following properties.

Corollary 5. Let $a = a_0 + a_1 e, b = b_0 + b_1 e \in C_1, \lambda \in C$ be given, and ϕ_1 be defined by (18). Then

(a)
$$a = b \iff \phi_1(a) = \phi_1(b).$$

(b)
$$\phi_1(a+b) = \phi_1(a) + \phi_1(b), \ \phi_1(ab) = \phi_1(a)\phi_1(b), \ \phi_1(\lambda a) = \lambda\phi_1(a), \ \phi_1(1) = I_2$$

- (c) $\phi_1(\overline{a}) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \phi_1(a) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$
- (d) Denote $a^{\#} = \overline{a_0} \overline{a_1}e$, then $\phi_1(a^{\#}) = \phi_1^*(a)$, the conjugate transpose of the complex matrix $\phi_1(a)$.
- (e) $a = \frac{1}{4}[1 ie, i e]\phi_1(a)[1 ie, e i]^T$.
- (f) det $[\phi_1(a)] = a_0^2 + a_1^2$.
- (g) a is invertible $\iff \phi_1(a)$ is invertible, in which case $\phi_1(a^{-1}) = \phi_1^{-1}(a)$.

The properties in Corollary 5(a) and (b) show that through the bijective map (18), the Clifford algebra C_1 is algebraically isomorphic to the double field $\mathcal{C} \oplus \mathcal{C}$, and $\phi_1(a)$ is a faithful matrix representation of a in $\mathcal{C} \oplus \mathcal{C}$.

Notice that P_1 and P_1^{-1} in (15) have no relation with a. Thus the equality in Theorem 4 can also be extended to all matrices over the complex Clifford algebra C_1 .

Theorem 6. Let $A = A_0 + A_1 e \in C_1^{m \times n}$ be given, where $A_0, A_1 \in C^{m \times n}$. Then A and its conjugate $\overline{A} = A_0 - A_1 e$ satisfy the following universal factorization equality

$$J_{2m} \begin{bmatrix} A & 0\\ 0 & \overline{A} \end{bmatrix} J_{2n}^{-1} = \begin{bmatrix} A_0 + A_1 i & 0\\ 0 & A_0 - A_1 i \end{bmatrix},$$
(19)

where

$$J_{2m} = \frac{1}{2} \begin{bmatrix} (1-ie)I_m & -(i-e)I_m \\ -(i-e)I_m & (1-ie_1)I_m \end{bmatrix},$$

$$J_{2n}^{-1} = \frac{1}{2} \begin{bmatrix} (1-ie_1)I_n & (i-e)I_n \\ (i-e)I_n & (1-ie)I_n \end{bmatrix}.$$
 (20)

In particular, if m = n, then (19) becomes a universal similarity factorization equality over C_1 .

It is easy to verify that for any matrices $A, B \in \mathcal{C}_1^{m \times n}, C \in \mathcal{C}_1^{n \times p}$, and $\lambda \in \mathcal{C}$

$$\overline{\overline{A}} = A, \quad \overline{\overline{A} + B} = \overline{\overline{A}} + \overline{B}, \quad \overline{\overline{AC}} = \overline{\overline{A}} \overline{\overline{C}}, \quad \overline{\overline{\lambda}A} = \overline{\overline{A\lambda}} = \overline{\overline{\lambda}A}.$$
(21)

Now according to (19), we define the complex matrix representation of a matrix $A = A_0 + A_1 e \in C_1^{m \times n}$ by $\Phi_1(A) = \begin{bmatrix} A_0 + A_1 i & 0 \\ 0 & A_0 - A_1 i \end{bmatrix}$. Then the following properties can be easily derived from (19) and (21).

Corollary 7. Let $A, B \in \mathcal{C}_1^{m \times n}, C \in \mathcal{C}_1^{n \times p}$ and $\lambda \in \mathcal{C}$ be given. Then

- (a) $A = B \iff \Phi_1(A) = \Phi_1(B).$
- (b) $\Phi_1(A+B) = \Phi_1(A) + \Phi_1(B).$
- (c) $\Phi_1(AC) = \Phi_1(A)\Phi_1(C), \ \Phi_1(\lambda A) = \lambda \Phi_1(A), \ \Phi_1(I_m) = I_{2m}.$
- (d) $\Phi_1(\overline{A}) = \begin{bmatrix} 0 & I_m \\ I_m & 0 \end{bmatrix} \Phi_1(A) \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}.$
- (e) Let $A = A_0 + A_1 e$ and denote $A^{\#} = A_0^* A_1^* e$, where A_0^* and A_1^* are the conjugate transposes of the complex matrices A_0 and A_1 . Then $\Phi_1(A^{\#}) = \Phi_1^*(A)$, the conjugate transpose of the complex matrix $\Phi_1(A)$.
- (f) $A = \frac{1}{4} [(1 ie)I_m, (i e)I_m] \Phi_1(A) [(1 ie)I_n, (e i)I_n]^T.$
- (g) A is invertible $\iff \Phi_1(A)$ is invertible, in which case $\Phi_1(A^{-1}) = \Phi_1^{-1}(A)$.
- (h) $p_A(A) = 0$, where $p_A(x) = \det[xI_{2m} \Phi_1(A)]$.

Theorem 8. Let $a \in C_2 = C\{e_1, e_2\}$, the complex quaternion algebra. Then a can be written as

$$a = a_0 + a_1 e_1 + a_2 e_2 + a_3 e_{12},$$

where $a_0, \ldots, a_3 \in C$. In that case, aI_2 satisfies the following universal similarity factorization equality

$$P_2 \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix} P_2^{-1} = \begin{bmatrix} a_0 + a_1 i & -(a_2 + a_3 i) \\ a_2 - a_3 i & a_0 - a_1 i \end{bmatrix},$$
(22)

where P_2 has the universal form

$$P_2 = P_2^{-1} = \frac{1}{2} \begin{bmatrix} 1 - ie_1 & e_2 + ie_{12} \\ -e_2 + ie_{12} & 1 + ie_1 \end{bmatrix}.$$
 (23)

Proof. Let $\mathcal{F} = \mathcal{C}$ and u = v = -1 in Lemma 3. Then (22) and (23) follow from (10) and (12). \Box

According to (22), define a map from C_2 to the 2 × 2 complex matrix algebra $C^{2\times 2}$ by

$$\phi_2 : a = a_0 + a_1 e_1 + a_2 e_2 + a_3 e_{12} \in \mathcal{C}_2 \longrightarrow \begin{bmatrix} a_0 + a_1 i & -(a_2 + a_3 i) \\ a_2 - a_3 i & a_0 - a_1 i \end{bmatrix} \in \mathcal{C}^{2 \times 2}.$$
(24)

Then we can easily derive from (22) the following properties.

Corollary 9. Let $a, b \in C_2 = C\{e_1, e_2\}$ and $\lambda \in C$ be given. Then

- (a) $a = b \iff \phi_2(a) = \phi_2(b).$
- (b) $\phi_2(a+b) = \phi_2(a) + \phi_2(b), \ \phi_2(ab) = \phi_2(a)\phi_2(b), \ \phi_2(\lambda a) = \lambda\phi_2(a), \ \phi_2(1) = I_2.$
- (c) Let $a = a_0 + a_1e_1 + a_2e_2 + a_3e_{12} \in C_2$, and denote $a^{\#} = \overline{a_0} \overline{a_1}e_1 \overline{a_2}e_2 \overline{a_3}e_{12}$. Then $\phi_2(a^{\#}) = \phi_2^*(a)$, the conjugate transpose of the complex matrix $\phi_2(a)$.

- (d) $a = \frac{1}{4} [1 ie, e_2 + ie_{12}] \phi_2(a) [1 ie, -e_2 + ie_{12}]^T.$
- (e) $det[\phi_2(a)] = a_0^2 + a_1^2 + a_2^2 + a_3^2$.
- (f) a is invertible $\iff \phi_2(a)$ is invertible, in which case $\phi_2(a^{-1}) = \phi_2^{-1}(a)$.
- (g) The two elements a and b in C_2 are similar, i.e., there is an invertible $x \in C_2$ such that ax = xb if and only if the two complex matrices $\phi_2(a)$ and $\phi_2(b)$ are similar over C.

The properties in Corollary 9(a) and (b) clearly show that through the bijective map (24) the Clifford algebra C_2 , i.e., the complex quaternion algebra, is algebraically isomorphic to the complex matrix algebra $C^{2\times 2}$, and $\phi_2(a)$ is a faithful matrix representation of a in $C^{2\times 2}$.

Notice that P_2 and P_2^{-1} in (23) have no relation to a. Thus the equality in Theorem 8 can also be extended to all matrices over C_2 .

Theorem 10. Let $A = A_0 + A_1e_1 + A_2e_2 + A_3e_{12} \in \mathcal{C}_2^{m \times n} = \mathcal{C}^{m \times n}\{e_1, e_2\}$ be given where $A_0, \ldots, A_3 \in \mathcal{C}^{m \times n}$. Then A satisfies the following universal factorization equality

$$K_{2m} \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} K_{2n}^{-1} = \begin{bmatrix} A_0 + A_1 i & -(A_2 + A_3 i) \\ A_2 - A_3 i & A_0 - A_1 i \end{bmatrix},$$
(25)

where

$$K_{2t} = K_{2t}^{-1} = \frac{1}{2} \begin{bmatrix} (1 - ie_1)I_t & (e_2 + ie_{12})I_t \\ (-e_2 + ie_{12})I_t & (1 + ie_1)I_t \end{bmatrix}.$$
(26)

In particular, if m = n, then (25) becomes a universal similarity factorization equality over C_2 .

According to (25), we define the complex representation of a matrix $A = A_0 + A_1 e \in C_2^{m \times n}$ by $\Phi_2(A) = \begin{bmatrix} A_0 + A_1 i & -(A_2 + A_3 i) \\ A_2 - A_3 i & A_0 - A_1 i \end{bmatrix}$. Then the following properties are easy to verify by (25).

Corollary 11. Let $A, B \in \mathcal{C}_2^{m \times n}, C \in \mathcal{C}_2^{n \times p}$, and $\lambda \in \mathcal{C}$ be given. Then

(a)
$$A = B \iff \Phi_2(A) = \Phi_2(B)$$

(b)
$$\Phi_2(A+B) = \Phi_2(A) + \Phi_2(B).$$

(c)
$$\Phi_2(AC) = \Phi_2(A)\Phi_2(C), \ \Phi_2(\lambda A) = \lambda \Phi_2(A), \ \Phi_2(I_m) = I_{2m}.$$

(d) Let $A = A_0 + A_1e_1 + A_2e_2 + A_3e_{12}$ and denote $A^{\#} = A_0^* - A_1^*e_1 - A_2^*e_2 - A_3^*e_{12}$, then $\Phi_2(A^{\#}) = \Phi_2^*(A)$, the conjugate transpose of the complex matrix $\Phi_2(A)$.

(e)
$$A = \frac{1}{4} [(1 - ie_1)I_m, (e_2 + ie_{12})I_m] \Phi_2(A) [(1 - ie_1)I_n, (-e_2 + ie_{12})I_n]^T$$

- (f) A is invertible $\iff \Phi_2(A)$ is invertible, in which case $\Phi_2(A^{-1}) = \Phi_2^{-1}(A)$.
- (g) $p_A(A) = 0$, where $p_A(\lambda) = \det[\lambda I_{2m} \Phi_2(A)]$, the characteristic polynomial of $\Phi_2(A)$.

(h) Two square matrices A and B are similar over C_2 , i.e., there is an invertible matrix X over C_2 such that AX = XB if and only if $\Phi_2(A)$ and $\Phi_2(B)$ are similar over C.

In the next several results we only present the basic universal similarity factorization equalities without listing their operation properties and their extensions to matrices over C_n .

Theorem 12. Let $a \in C_3 = C\{e_1, e_2, e_3\}$ be given. Then a can factor as

$$a = a_0 + a_1 e_{[3]},\tag{27}$$

where

$$a_0, a_1 \in \mathcal{C}_2 = \mathcal{C}\{e_1, e_2\}, e_{[3]}^2 = 1.$$

Moreover, define $\overline{a} = a_0 - a_1 e_{[3]}$. In that case, the diagonal matrix $D_a = \text{diag}(aI_2, \overline{a}I_2)$ satisfies the following universal similarity factorization equality

$$P_{3}D_{a}P_{3}^{-1} = \begin{bmatrix} \phi_{2}(a_{0}) + \phi_{2}(a_{1}) & 0\\ 0 & \phi_{2}(a_{0}) - \phi_{2}(a_{1}) \end{bmatrix}$$
$$:= \phi_{3}(a) \in {}^{2}\mathcal{C}^{2 \times 2}, \tag{28}$$

where $\phi_2(a_t)$, t = 0, 1, is the matrix representation of a_t in $\mathcal{C}^{2 \times 2}$ defined in (24) and

$$P_3 = \frac{1}{2} \begin{bmatrix} (1+e_{[3]})P_2 & -(1-e_{[3]})P_2 \\ (1-e_{[3]})P_2 & (1+e_{[3]})P_2 \end{bmatrix},$$
(29)

$$P_3^{-1} = \frac{1}{2} \begin{bmatrix} P_2^{-1}(1+e_{[3]}) & P_2^{-1}(1-e_{[3]}) \\ -P_2^{-1}(1-e_{[3]}) & P_2^{-1}(1+e_{[3]}) \end{bmatrix},$$
(30)

where P_2 and P_2^{-1} are given by (23).

Proof. Notice that $be_{[3]} = e_{[3]}b$ holds for all $b \in C_2 = C\{e_1, e_2\}$. We have by applying (22) to a in (27) that

$$P_2(aI_2)P_2^{-1} = P_2(a_0I_2)P_2^{-1} + P_2(a_1I_2)P_2^{-1}e_{[3]} = \phi_2(a_0) + \phi_2(a_1)e_{[3]} := \psi(a),$$

and

$$P_2(\overline{a}I_2)P_2^{-1} = \phi_2(a_0) - \phi_2(a_1)e_{[3]} := \psi(\overline{a}).$$

Next we build, according to Lemma 1, a matrix and its inverse as follows

$$V = \frac{1}{2} \begin{bmatrix} (1+e_{[3]})I_2 & -(1-e_{[3]})I_2 \\ (1-e_{[3]})I_2 & (1+e_{[3]})I_2 \end{bmatrix},$$
$$V^{-1} = \frac{1}{2} \begin{bmatrix} (1+e_{[3]})I_2 & (1-e_{[3]})I_2 \\ -(1-e_{[3]})I_2 & (1+e_{[3]})I_2 \end{bmatrix}.$$

and then calculate to get

$$V\begin{bmatrix} \psi(a) & 0\\ 0 & \psi(\overline{a}) \end{bmatrix} V^{-1} = \begin{bmatrix} \phi_2(a_0) + \phi_2(a_1) & 0\\ 0 & \phi_2(a_0) - \phi_2(a_1) \end{bmatrix}.$$

Finally substituting $\psi(a) = P_2(aI_2)P_2^{-1}$ and $\psi(\overline{a}) = P_2(\overline{a}I_2)P_2^{-1}$ into the left-hand side of the above equality yields (28), (29), and (30).

Theorem 13. Let $a \in C_4 = C\{e_1, e_2, e_3, e_4\}$ be given. Then a can factor as

$$a = a_0 + a_1 e_{123} + a_2 e_{124} + a_3 e_{43} = a_0 + e_{123} a_1 + e_{124} a_2 + e_{43} a_3,$$
(31)

where

$$a_0, a_1, a_2, a_3 \in \mathcal{C}_2 = \mathcal{C}\{e_1, e_2\},$$

 $e_{123}^2 = 1, \quad e_{124}^2 = 1, \quad e_{43} = e_{123}e_{124} = -e_{124}e_{123}.$

In that case, aI_4 satisfies the following universal similarity factorization equality

$$P_4(aI_4)P_4^{-1} = \begin{bmatrix} \phi_2(a_0) + \phi_2(a_1) & \phi_2(a_2) + \phi_2(a_3) \\ \phi_2(a_2) - \phi_2(a_3) & \phi_2(a_0) - \phi_2(a_1) \end{bmatrix}$$

$$:= \phi_4(a) \in \mathcal{C}^{4 \times 4}, \tag{32}$$

where $\phi_2(a_t), t = 0, \dots, 3$, is the matrix representation of a_t in $\mathcal{C}^{2 \times 2}$ defined in (24) and

$$P_4 = P_4^{-1} = \frac{1}{2} \begin{bmatrix} (1+e_{[3]})P_2 & (e_{124}-e_{43})P_2 \\ (e_{124}+e_{43})P_2 & (1-e_{[3]})P_2 \end{bmatrix},$$
(33)

where P_2 is given in (23).

Proof. Note that the commutative rules $be_{123} = e_{123}b$, $be_{124} = e_{124}b$, $be_{43} = e_{43}b$ hold for all $b \in C_2 = C\{e_1, e_2\}$. Thus, it follows from (22) that

$$P_{2}(aI_{2})P_{2}^{-1}$$

$$= P_{2}(a_{0}I_{2})P_{2}^{-1} + P_{2}(a_{1}I_{2})P_{2}^{-1}e_{123} + P_{2}(a_{2}I_{2})P_{2}^{-1}e_{124} + P_{2}(a_{3}I_{2})P_{2}^{-1}e_{43}$$

$$= \phi_{2}(a_{0}) + \phi_{2}(a_{1})e_{123} + \phi_{2}(a_{2})e_{124} + \phi_{2}(a_{3})e_{43}$$

$$:= \psi(a).$$

Next building, according to Lemma 3, a matrix and its inverse as follows

$$V = V^{-1} = \frac{1}{2} \begin{bmatrix} (1+e_{123})I_2 & (e_{124}-e_{43})I_2 \\ (e_{124}+e_{43})I_2 & (1-e_{123})I_2 \end{bmatrix},$$

and applying them to $\psi(a)$ given above, we obtain

$$V\begin{bmatrix} \psi(a) & 0\\ 0 & \psi(a) \end{bmatrix} V^{-1} = \begin{bmatrix} \phi_2(a_0) + \phi_2(a_1) & \phi_2(a_2) + \phi_2(a_3)\\ \phi_2(a_2) - \phi_2(a_3) & \phi_2(a_0) - \phi_2(a_1) \end{bmatrix}.$$

Finally substituting $\psi(a) = P_2(aI_2)P_2^{-1}$ into its left-hand side yields (32) and (33).

By induction, we have the following two general results.

Theorem 14. Suppose that there is an independent invertible matrix P_n over $C_n = C\{e_1, \ldots, e_n\}$ with n even such that

$$P_n(aI_{2^{\frac{n}{2}}})P_n^{-1} := \phi_n(a) \in \mathcal{C}^{2^{\frac{n}{2}} \times 2^{\frac{n}{2}}} = \mathcal{C}(2^{\frac{n}{2}})$$
(34)

holds for all $a \in C_n$. Now let $a \in C_{n+1} = C\{e_1, \ldots, e_{n+1}\}$. Then a can factor as

$$a = a_0 + a_1 e_{[n+1]} = a_0 + e_{[n+1]} a_1,$$
(35)

where

$$a_0, a_1 \in \mathcal{C}_n = \mathcal{C}\{e_1, \dots, e_n\}, \quad e_{[n+1]}^2 = (-1)^{\frac{1}{2}(n+1)(n+2)} := r.$$

Moreover define $\overline{a} = a_0 - a_1 e_{[n+1]}$. In that case, $D_a = \text{diag}(aI_{2\frac{n}{2}}, \overline{a}I_{2\frac{n}{2}})$ satisfies the following universal similarity factorization equality

$$P_{n+1}D_a P_{n+1}^{-1} = \begin{bmatrix} \phi_n(a_0) + \sqrt{r}\phi_n(a_1) & 0\\ 0 & \phi_n(a_0) - \sqrt{r}\phi_n(a_1) \end{bmatrix}$$
$$:= \phi_{n+1}(a) \in {}^2\mathcal{C}(2^{\frac{n}{2}}), \tag{36}$$

where

$$P_{n+1} = \frac{1}{2} \begin{bmatrix} (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) P_n & -(\sqrt{r} - e_{[n+1]}) P_n \\ \frac{1}{r} (\sqrt{r} - e_{[n+1]}) P_n & (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) P_n \end{bmatrix},$$
(37)

and

$$P_{n+1}^{-1} = \frac{1}{2} \begin{bmatrix} P_n^{-1}(1 + \frac{1}{\sqrt{r}}e_{[n+1]}) & P_n^{-1}(\sqrt{r} - e_{[n+1]}) \\ -P_n^{-1}\frac{1}{r}(\sqrt{r} - e_{[n+1]}) & P_n^{-1}(1 + \frac{1}{\sqrt{r}}e_{[n+1]}) \end{bmatrix}.$$
(38)

Proof. Note that the commutative rule $be_{[n+1]} = e_{[n+1]}b$ holds for all $b \in C_n = C\{e_1, \ldots, e_n\}$. By applying (34) to (35), we obtain

$$P_n(aI_{2^{\frac{n}{2}}})P_n^{-1} = P_n(aI_{2^{\frac{n}{2}}})P_n^{-1} + P_n(a_1I_{2^{\frac{n}{2}}})P_n^{-1}e_{[n+1]}$$
$$= \phi_n(a_0) + \phi_n(a_1)e_{[n+1]} := \psi(a).$$

and

$$P_n(\overline{a}I_{2^{\frac{n}{2}}})P_n^{-1} = \phi_n(a_0) - \phi_n(a_1)e_{[n+1]} := \psi(\overline{a}).$$

Next, setting

$$\begin{split} V &= \frac{1}{2} \left[\begin{array}{cc} (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) I_{2\frac{n}{2}} & -(\sqrt{r} - e_{[n+1]}) I_{2\frac{n}{2}} \\ \frac{1}{\sqrt{r}} (\sqrt{r} - e_{[n+1]}) I_{2\frac{n}{2}} & (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) I_{2\frac{n}{2}} \end{array} \right], \\ V^{-1} &= \frac{1}{2} \left[\begin{array}{cc} (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) I_{2\frac{n}{2}} & (\sqrt{r} - e_{[n+1]}) I_{2\frac{n}{2}} \\ -\frac{1}{r} (\sqrt{r} - e_{[n+1]}) I_{2\frac{n}{2}} & (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) I_{2\frac{n}{2}} \end{array} \right], \end{split}$$

and applying them to $D_a = \operatorname{diag}(\psi(a), \psi(\overline{a}))$ we get

$$V\begin{bmatrix} \psi(a) & 0\\ 0 & \psi(\overline{a}) \end{bmatrix} = \begin{bmatrix} \phi_n(a_0) + \sqrt{r}\phi_n(a_1) & 0\\ 0 & \phi_n(a_0) - \sqrt{r}\phi_n(a_1) \end{bmatrix}.$$

Finally, substituting $\psi(a) = P_n(aI_{2\frac{n}{2}})P_n^{-1}$ and $\psi(\overline{a}) = P_n(\overline{a}I_{2\frac{n}{2}})P_n^{-1}$ into its left-hand side yields the desired results. \Box

Theorem 15. Suppose that there is an independent invertible matrix P_n over $C_n = C\{e_1, \ldots, e_n\}$ with n even such that

$$P_n(aI_{2^{\frac{n}{2}}})P_n^{-1} := \phi_n(a) \in \mathcal{C}(2^{\frac{n}{2}})$$
(39)

holds for all $a \in C_n$. Now, let $a \in C_{n+2} = C\{e_1, \ldots, e_{n+2}\}$. Then it can factor as

$$a = a_0 + a_1 e_{[n]} e_{n+1} + a_2 e_{[n]} e_{n+2} + a_3 \mu_{n+2}, \tag{40}$$

where

$$a_0, a_1, a_2, a_3 \in \mathcal{C}_n = \mathcal{C}\{e_1, \dots, e_n\},$$

$$(e_{[n]}e_{n+1})^2 = (e_{[n]}e_{n+2})^2 = (-1)^{\frac{1}{2}(n+1)(n+2)} = r,$$

$$\mu_{n+2} = (e_{[n]}e_{n+1})(e_{[n]}e_{n+2}) = -(e_{[n]}e_{n+2})(e_{[n]}e_{n+1}).$$

In that case, $aI_{2^{\frac{n+1}{2}}}$ satisfies the following universal similarity factorization equality

$$P_{n+2}(aI_{2^{\frac{n+1}{2}}})P_{n+2}^{-1} = \begin{bmatrix} \phi_n(a_0) + \sqrt{r}\phi_n(a_1) & r[\phi_n(a_2) + \sqrt{r}\phi_n(a_3)] \\ \phi_n(a_2) - \sqrt{r}\phi_n(a_3) & \phi_n(a_0) - \sqrt{r}\phi_n(a_1) \end{bmatrix}$$
$$:= \phi_{n+2}(a) \in \mathcal{C}(2^{\frac{n+1}{2}}),$$

where

$$\begin{split} P_{n+2} &= \frac{1}{2} \begin{bmatrix} (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) P_n & (e_{[n]} e_{n+2} - \frac{1}{\sqrt{r}} \mu_{n+2}) P_n \\ \frac{1}{r} (e_{[n]} e_{n+2} + \frac{1}{\sqrt{r}} \mu_{n+2}) P_n & (1 - \frac{1}{\sqrt{r}} e_{[n+1]}) P_n \end{bmatrix}, \\ P_{n+2}^{-1} &= \frac{1}{2} \begin{bmatrix} P_n^{-1} (1 + \frac{1}{\sqrt{r}} e_{[n+1]}) & P_n^{-1} (e_{[n]} e_{n+2} - \frac{1}{\sqrt{r}} \mu_{n+2}) \\ P_n^{-1} \frac{1}{r} (e_{[n]} e_{n+2} + \frac{1}{\sqrt{r}} \mu_{n+2}) & P_n^{-1} (1 - \frac{1}{\sqrt{r}} e_{[n+1]}) \end{bmatrix}, \end{split}$$

The proof of this result is analogous to that of Theorem 13, and is therefore omitted here.

3 Conclusions

In this article we have established a set of universal similarity factorization equalities for elements over the complex Clifford algebra C_n . These equalities reveal two basic facts about C_n :

- (i) Each element a in C_n has a complex matrix representation $\phi_n(a)$. Moreover, a diagonal matrix constructed by the element a is uniformly similar to its complex matrix representation $\phi_n(a)$.
- (ii) Conversely, each element a in C_n could be regarded as an eigenvalue of its complex representation matrix $\phi_n(a)$. In other words, all complex matrices with the form $\phi_n(a)$ can uniformly be diagonalized over C_n .

Based on the above two facts, one easily see that almost all known results in complex matrix theory can be extended to complex Clifford algebras. On the other hand, some problems related to complex matrices can also transform to the problems related to Clifford numbers. One such a problem (see [1]) is concerning the exponential e^A of a complex matrix A. In fact, we can see from Theorems 14 and 15 that for any $a \in C_n$, there is

$$P_n(e^a I)P_n^{-1} = e^{\phi_n(a)}$$

or

$$P_n \operatorname{diag}(e^a I, e^{\overline{a}} I) P_n^{-1} = e^{\phi_n(a)},$$

which implies that the exponential of a complex matrix $\phi_n(a)$ can be determined by the exponential of its corresponding Clifford numbers.

References

- R. Abłamowicz, Matrix exponential via Clifford algebras, J. Nonlinear Math. Physics 5(1998), 294–313.
- [2] M. F. Atiyah, R. Bott and A. Shapiro, Clifford Modules, Topology 3(1961), 3–38.
- [3] G. N. Hile and P. Lounesto, Matrix representations of Clifford algebras, Linear Algebra Appl. 128(1990), 51–63.
- [4] T. Y. Lam, The Algebraic Theory of Quadratic Forms, W. A. Benjamin, Reading, Mass., 1973.
- [5] M. Karoubi, K-Theory, Springer-Verlag, Berlin, 1978.
- [6] M. Marcus, Finite Dimensional Multilinear Algebra (Part II), Marcel Dekker Inc., New York, 1975.
- [7] S. Okubo, Real representations of finite Clifford algebras. I. classification, J. Math. Phys. 32(1991), 1657-1668.
- [8] S. Okubo, Real representations of finite Clifford algebras. II. explicit construction and pseudooctonion, J. Math. Phys. 32(1991), 1669–1673.
- [9] S. Okubo, Representations of Clifford algebras and its applications, Math. Japn. 41(1995), 59–79.
- [10] I. R. Porteous, Topological Geometry, Cambridge U. P., Cambridge, 1981.
- [11] I. R. Porteous, Clifford Algebras and the Classical Groups, Cambridge U. P., Cambridge, 1995.
- [12] Y. Tian, Universal similarity factorization equalities over real Clifford algebras, Adv. Appl. Clifford Algebras 8(1998), 365–402.
- [13] V. V. Varlamov, Modulo 2 periodicity of complex Clifford algebras and electromagnetic field, preprint, 1997.