

SEQUENCES IN NON-COMMUTATIVE L^p -SPACES

NARCISSE RANDRIANANTOANINA

ABSTRACT. Let \mathcal{M} be a semi-finite von Neumann algebra equipped with a distinguished faithful, normal, semi-finite trace τ . We introduce the notion of equi-integrability in non-commutative spaces and show that if a rearrangement invariant quasi-Banach function space E on the positive semi-axis is α -convex with constant 1 and satisfies a non-trivial lower q -estimate with constant 1, then the corresponding non-commutative space of measurable operators $E(\mathcal{M}, \tau)$ has the following property: every bounded sequence in $E(\mathcal{M}, \tau)$ has a subsequence that splits into a E -equi-integrable sequence and a sequence with pairwise disjoint projection supports. This result extends the well known Kadec-Pełczyński subsequence decomposition for Banach lattices to non-commutative spaces. As applications, we prove that for $1 \leq p < \infty$, every subspace of $L^p(\mathcal{M}, \tau)$ either contains almost isometric copies of ℓ^p or is strongly embedded in $L^p(\mathcal{M}, \tau)$.

1. INTRODUCTION

In [11], Kadec and Pełczyński proved that if $1 \leq p < \infty$ then every bounded sequence $\{f_n\}_{n=1}^\infty$ in $L^p[0, 1]$ has a subsequence that can be decomposed into two extreme sequences $\{g_k\}_{k=1}^\infty$ and $\{h_k\}_{k=1}^\infty$, where the h_k 's are pairwise disjoint and the g_k 's are L_p -equi-integrable that is $\lim_{m(A) \rightarrow 0} \sup_k \|\chi_A g_k\|_p \rightarrow 0$ and $h_k \perp g_k$ for every $k \geq 1$. This result was used to study different structures of subspaces of $L^p[0, 1]$. Later, the same decomposition property was proved for larger classes of Banach function spaces (see [9] for Orlicz spaces with Δ_2 -condition and q -concave lattices, [10] for some symmetric spaces). There are however Banach lattices with sequences for which the above decomposition is not possible. Examples of reflexive, p -convex Banach lattices without the subsequence decomposition can be found in a paper of Figiel *et al* [9]. Subsequently, Weis [20] characterized, in terms of uniform order continuity conditions and ultrapowers, all Banach lattices where such property is possible. For the case of rearrangement invariant function spaces, the order-continuous spaces, in which the above decomposition works, were fully characterized as those that have the Fatou property (equivalently, those that contains no subspace isomorphic to c_0). In [7], a version of Kadec-Pełczyński decomposition was considered for preduals of semi-finite von Neumann algebras.

1991 *Mathematics Subject Classification.* 46L50, 47D15.

Key words and phrases. von Neumann algebras, function spaces, symmetric spaces of operators.

Supported in part by NSF grant DMS-9703789.

It is the intention of the present paper to give an extension of the Kadec-Pełczyński decomposition stated above to the case of general non-commutative symmetric spaces of measurable operators. Let \mathcal{M} be a von Neumann algebra, equipped with a distinguished faithful, normal, semi-finite trace τ and E be a rearrangement invariant Banach function space on $[0, 1]$ or the half line $(0, \infty)$. We define equi-integrability in the non-commutative setting as generalizations of Akemann's characterization of weak compactness on preduals of von Neumann algebras. Using such notion, we provide an analogue of the Kadec-Pełczyński decomposition for non-commutative spaces. Namely, we proved that if E is order continuous and satisfies the Fatou property then the corresponding symmetric space of measurable operators $E(\mathcal{M}, \tau)$ has the subsequence splitting property. Our approach allows ones to consider more general spaces such as quasi-Banach rearrangement invariant spaces that are α -convex with constant 1 and satisfy non trivial q -lower estimate with constant 1. In particular splitting of bounded sequences is valid in non-commutative L^p -spaces for $0 < p < \infty$. It should be noted that Sukochev [18] obtain a similar result for the case of finite von Neumann algebras. We also remark that since c_0 fails the subsequence splitting property, our result for the case where E is a Banach space case is the best possible.

As application of the main result, we study the structure of subspaces of $L^p(\mathcal{M}, \tau)$ for $1 < p < \infty$ which generalizes the case $p = 1$ treated in [7].

We refer to [12] and [19] for general information concerning von Neumann algebras as well as non-commutative integration, to [17] and [14] for Banach lattice theory.

2. DEFINITIONS AND PRELIMINARY RESULTS

Throughout, H is a given Hilbert space and $\mathcal{M} \subset \mathcal{B}(H)$ denotes a semi-finite von Neumann algebra with a distinguished normal, faithful semi-finite trace τ . The identity in \mathcal{M} will be denoted by $\mathbf{1}$ and \mathcal{M}_p will stand for the set of all (self adjoint) projections in \mathcal{M} . A closed and densely defined operator a on H is said to be affiliated with \mathcal{M} if $ua = au$ for all unitary operator u in the commutant \mathcal{M}' of \mathcal{M} .

A closed and densely defined operator x , affiliated with \mathcal{M} , is called τ -measurable if for every $\varepsilon > 0$, there exists an orthogonal projection $p \in \mathcal{M}$ such that $p(H) \subseteq \text{dom}(x)$, $\tau(\mathbf{1} - p) \leq \varepsilon$ and $xp \in \mathcal{M}$. The set of all τ -measurable operators will be denoted by $\widetilde{\mathcal{M}}$. The set $\widetilde{\mathcal{M}}$ is a $*$ -algebra with respect to the strong sum, the strong product and the adjoint operation. Given a self-adjoint operator x in $\widetilde{\mathcal{M}}$, we denote by $e^x(\cdot)$ its spectral measure. Recall that $e^{|x|}(B) \in \mathcal{M}$ for all Borel sets $B \subseteq \mathbb{R}$ and $x \in \widetilde{\mathcal{M}}$. For fixed $x \in \widetilde{\mathcal{M}}$ and $t \geq 0$, we define

$$\mu_t(x) = \inf \{s \geq 0 : \tau(e^{|x|}(s, \infty)) \leq t\}.$$

The function $\mu_{(\cdot)}(x) : [0, \infty) \rightarrow [0, \infty]$ is called the generalized singular value function (or decreasing rearrangement) of x . We note that $\mu_t(x) < \infty$ for every $t > 0$. For a complete

study of $\mu(\cdot)$, we refer the reader to [8]. The topology defined by the metric on $\widetilde{\mathcal{M}}$ obtained by setting

$$d(x, y) = \inf \{t \geq 0 : \mu_t(x - y) \leq t\}, \quad \text{for } x, y \in \widetilde{\mathcal{M}},$$

is called the *measure topology*. It is well-known that a net $(x_\alpha)_{\alpha \in I}$ in $\widetilde{\mathcal{M}}$ converge to $x \in \widetilde{\mathcal{M}}$ in measure topology if and only if for every $\varepsilon > 0$, $\delta > 0$, there exists $\alpha_0 \in I$ such that whenever $\alpha \geq \alpha_0$, there exists a projection $p \in \mathcal{M}_p$ such that

$$\|(x_\alpha - x)p\|_{\mathcal{M}} < \varepsilon \quad \text{and} \quad \tau(\mathbf{1} - p) < \delta.$$

It was shown in [15] that $(\widetilde{\mathcal{M}}, d)$ is a complete metric space.

Recall that if we consider $\mathcal{M} = L^\infty(\mathbb{R}^+, m)$, where m is the Lebesgue measure on \mathbb{R}^+ then \mathcal{M} is an abelian von Neumann algebra acting on $L^2(\mathbb{R}^+, m)$ via multiplication, with the trace being the integration with respect to m , then $\widetilde{\mathcal{M}} = L^0(\mathbb{R}^+, m)$ (the usual space of all measurable functions on \mathbb{R}^+) and the generalized singular value $\mu(f)$ is precisely the decreasing rearrangement of the function $|f|$ (usually denoted by f^* in Banach lattice theory).

Definition 2.1. *A symmetric quasi-Banach function space on \mathbb{R}^+ is a quasi-Banach lattice E of measurable functions with the following properties:*

- (i) E is an order ideal in $L^0(\mathbb{R}^+, m)$;
- (ii) E is rearrangement invariant in the sense of [14] (p .114);
- (iii) E contains all finitely supported simple functions.

Unless stated otherwise, E will always denote a symmetric quasi-Banach function space on \mathbb{R}^+ . We define the symmetric space of measurable operators $E(\mathcal{M}, \tau)$ by setting

$$E(\mathcal{M}, \tau) := \left\{ x \in \widetilde{\mathcal{M}} : \mu(x) \in E \right\}$$

and

$$\|x\|_{E(\mathcal{M}, \tau)} = \|\mu(x)\|_E \quad \text{for all } x \in E(\mathcal{M}, \tau).$$

It is shown in [21] (Lemma 4.1) that if E is α -convex (for some $0 < \alpha \leq 1$) with constant 1, then $\|\cdot\|_{E(\mathcal{M}, \tau)}$ is an α -norm that is for every $x, y \in E(\mathcal{M}, \tau)$,

$$\|x + y\|_{E(\mathcal{M}, \tau)}^\alpha \leq \|x\|_{E(\mathcal{M}, \tau)}^\alpha + \|y\|_{E(\mathcal{M}, \tau)}^\alpha.$$

Equipped with $\|\cdot\|_{E(\mathcal{M}, \tau)}$, the space $E(\mathcal{M}, \tau)$ is a α -Banach space. The space $E(\mathcal{M}, \tau)$ is often referred to as the non-commutative analogue of the function space E . We remark that if $0 < p < \infty$ and $E = L^p(\mathbb{R}^+, m)$ then $E(\mathcal{M}, \tau)$ coincides with the usual non-commutative L^p -space associated to the semi-finite von Neumann algebra \mathcal{M} . Also if $E = L^\infty(\mathbb{R}^+, m)$, then $L^\infty(\mathcal{M}, \tau)$ is the von Neumann algebra \mathcal{M} . We refer to [4], [5] and [21] for some background on the space $E(\mathcal{M}, \tau)$.

Definition 2.2. A quasi-Banach function space E is said to satisfy a lower q -estimate if there exists a positive constant $C > 0$ such that for all finite sequences $\{x_n\}$ of mutually disjoint elements in E ,

$$\left(\sum \|x_n\|^q\right)^{\frac{1}{q}} \leq C \left\|\sum x_n\right\|.$$

The least such constant C is called the constant of the lower q -estimate. Recall that if E is a quasi-Banach function space and $1 < p < \infty$,

$$E^{(p)} = \{x \in L^0(\mathbb{R}^+, m); |x|^p \in E\} \text{ with } \|x\|_{E^{(p)}} = \left\| |x|^p \right\|_E^{\frac{1}{p}}.$$

We will need the following known result. A proof can be found in [6].

Proposition 2.3. Assume that E is order-continuous and α -convex with constant 1 for some $0 < \alpha \leq 1$.

- (i) If $x \in E(\mathcal{M}, \tau)$ and $e \leq f$ are projections in \mathcal{M} then $\|exe\|_{E(\mathcal{M}, \tau)} \leq \|xf\|_{E(\mathcal{M}, \tau)}$;
- (ii) If $x \in E(\mathcal{M}, \tau)$ and $e_\beta \downarrow_\beta 0$ is a net of projections in \mathcal{M} then $\|xe_\beta\|_{E(\mathcal{M}, \tau)} \downarrow_\beta 0$.

The following definition isolates the main topic of this paper.

Definition 2.4. Let E be a quasi-Banach function space on \mathbb{R}^+ and K be a bounded subset of $E(\mathcal{M}, \tau)$. We will say that K is **E -equi-integrable** if $\lim_{n \rightarrow \infty} \sup_{x \in K} \|e_n x e_n\|_{E(\mathcal{M}, \tau)} = 0$ for every decreasing sequence $\{e_n\}_{n=1}^\infty$ of projections with $e_n \downarrow_n 0$.

We remark, from Proposition 2.3, that since $\{e_n\}_{n=1}^\infty$ is decreasing, so is the sequence $\left\{ \sup_{x \in K} \|e_n x e_n\|_{E(\mathcal{M}, \tau)} \right\}_{n=1}^\infty$ and therefore the limit in the definition above always exists. This definition was motivated by the commutative case on one hand and the characterization of weakly compact subsets of $L^1(\mathcal{M}, \tau)$ by Akemann [1] (see also [19] p.150) on the other. Using this terminology, Akemann's characterization can be stated as in the commutative case: relatively weakly compact subsets of $L^1(\mathcal{M}, \tau)$ are exactly the equi-integrable sets. Such characterization is not valid in general. For $1 < p < \infty$, any set of normalized disjoint sequence cannot be L^p -integrable but since L^p is reflexive, such set is relatively weakly compact.

On the next proposition, we will show that for the general case, one implication always holds.

Proposition 2.5. Assume that E is an order-continuous symmetric Banach function space and K is a E -equi-integrable set in $E(\mathcal{M}, \tau)$ then K is relatively weakly compact.

The proposition will be proved in several steps. Recall that if E is a symmetric Banach function space, then $E(\mathcal{M}, \tau)$ is a subset of $L^1(\mathcal{M}, \tau) + \mathcal{M}$ and therefore if p is a projection in $L^1(\mathcal{M}, \tau) \cap \mathcal{M}$ and K is a subset of $E(\mathcal{M}, \tau)$, then pK and Kp are subsets of $L^1(\mathcal{M}, \tau)$.

Lemma 2.6. *Let p be a projection in $L^1(\mathcal{M}, \tau) \cap \mathcal{M}$ and K be a E -equi-integrable subset of $E(\mathcal{M}, \tau)$. The sets pKp and $pK(1-p)$ are relatively weakly compact in $L^1(\mathcal{M}, \tau)$.*

To see this lemma, it is enough to check that these sets are L^1 -equi-integrable. Let $T : E(\mathcal{M}, \tau) \rightarrow L^1(\mathcal{M}, \tau)$ be the linear map defined by $x \rightarrow Tx = pxp$. This map is well-defined and one can deduce from the closed graph theorem that it is bounded. Let $\{e_n\}_{n=1}^\infty$ be a sequence of projections with $e_n \downarrow 0$. For each $n \geq 1$, set f_n to be the right support projection of $e_n p$. By the definition of support projections, $f_n \leq p$. So $\{f_n\}_{n=1}^\infty$ is a sequence of finite projections. We also note that (see for instance the proof of [19, Proposition 1.6 p.292]),

$$f_n = e_n \vee (\mathbf{1} - p) - (\mathbf{1} - p)$$

and by Kaplansky formula (see for instance [12, Theorem 6.1.6 p.403]),

$$f_n \sim e_n - e_n \wedge (\mathbf{1} - p).$$

Since $\tau(f_n) = \tau(e_n - e_n \wedge (\mathbf{1} - p)) \leq \tau(p)$ and $\{e_n - e_n \wedge (\mathbf{1} - p)\}_{n=1}^\infty$ converges to zero, $\{f_n\}_{n=1}^\infty$ converges to zero. Now since the f_n 's are finite projections, we conclude that if $g_n = \bigwedge_{k \geq n} f_k$, then $\{g_n\}_{n=1}^\infty$ converges to zero. Therefore, for every $x \in K$,

$$\begin{aligned} \|e_n p x p e_n\|_1 &= \|e_n p (g_n x g_n) p e_n\|_1 \\ &\leq \|p (g_n x g_n) p\|_1 \\ &\leq \|T\| \cdot \|g_n x g_n\|_{E(\mathcal{M}, \tau)}. \end{aligned}$$

Since K is E -equi-integrable, one obtain that

$$\lim_{n \rightarrow \infty} \sup_{y \in pKp} \|e_n y e_n\|_1 \leq \|T\| \cdot \lim_{n \rightarrow \infty} \sup_{x \in K} \|g_n x g_n\|_{E(\mathcal{M}, \tau)} = 0$$

which concludes that pKp is relatively weakly compact in $L^1(\mathcal{M}, \tau)$.

For $pK(\mathbf{1} - p)$, set $S : E(\mathcal{M}, \tau) \rightarrow L^1(\mathcal{M}, \tau)$ be the map defined by $x \rightarrow Sx = px(\mathbf{1} - p)$. As above, S is bounded. Let $\{e_n\}_{n=1}^\infty$ and $\{g_n\}_{n=1}^\infty$ be sequences of projections as described above. For each $n \geq 1$, let s_n be the left support projection of $(\mathbf{1} - p)e_n$. Then $s_n = e_n \vee p - p$ for every $n \geq 1$ and the sequence $\{s_n\}_{n=1}^\infty$ is decreasing. It is claimed that $s_n \downarrow 0$.

For this, it is enough to check that $e_n \vee p \downarrow p$. In fact, $e_n \vee p - e_n \sim p - e_n \wedge p$ and the sequence defined by the right hand side of the equivalence converges to p which implies that

$$\lim_{n \rightarrow \infty} \tau(e_n \vee p - e_n) = \lim_{n \rightarrow \infty} \tau(p - e_n \wedge p) = \tau(p)$$

which gives

$$\lim_{n \rightarrow \infty} \tau(e_n \vee p - p - e_n) = 0.$$

But since $(e_n \vee p - p - e_n)^2 = (e_n \vee p - p - e_n) + e_n p + p e_n$, we can conclude that

$$\lim_{n \rightarrow \infty} \|e_n \vee p - p - e_n\|_2 = 0.$$

From this, we get (by passing to a subsequence if necessary) that $\{e_n \vee p - p - e_n\}_{n=1}^\infty$ converges to zero in measure. Similarly, $\{e_n \vee p - p - pe_n\}_{n=1}^\infty$ converges to zero in measure so $e_n \vee p \downarrow_n p$ hence $s_n \downarrow_n 0$.

To conclude the proof of Lemma 2.6, note that $g_n \perp s_n$ so $g_n \vee s_n = g_n + s_n$. In particular, $g_n \vee s_n \downarrow_n 0$ and we get that

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{y \in pK(1-p)} \|e_n y e_n\|_1 &= \lim_{n \rightarrow \infty} \sup_{x \in K} \|e_n p x (\mathbf{1} - p) e_n\|_1 \\ &= \lim_{n \rightarrow \infty} \sup_{x \in K} \|e_n p (g_n \vee s_n) x (g_n \vee s_n) (\mathbf{1} - p) e_n\|_1 \\ &\leq \lim_{n \rightarrow \infty} \sup_{x \in K} \|p (g_n \vee s_n) x (g_n \vee s_n) (\mathbf{1} - p)\|_1 \\ &\leq \|S\| \cdot \lim_{n \rightarrow \infty} \sup_{x \in K} \|(g_n \vee s_n) x (g_n \vee s_n)\|_{E(\mathcal{M}, \tau)} = 0. \end{aligned}$$

The proof is complete.

Lemma 2.7. *Let p and K be as in Lemma 2.6. Then pK is relatively weakly compact in $E(\mathcal{M}, \tau)$.*

Note first that pK is E -equi-integrable. This can be seen by applying the series of argument used in Lemma 2.6, considering the operators T and S as maps from $E(\mathcal{M}, \tau)$ into $E(\mathcal{M}, \tau)$.

Let $\{px_n\}_{n=1}^\infty$ be a bounded sequence in pK . From Lemma reffinitecompact1, we can assume that $\{px_n\}_{n=1}^\infty$ is weakly convergent in $L^1(\mathcal{M}, \tau)$. Fix $\varphi \in E^*(\mathcal{M}, \tau)_+$ and let $\varphi = \int_0^\infty t \, de_t$ be its spectral decomposition. For each $k \geq 1$, set $q_k := e^\varphi((0, k))$. We remark that $\varphi q_k = q_k \varphi \in \mathcal{M}$. For $m, n \in \mathbb{N}$,

$$\begin{aligned} \langle \varphi, px_n - px_m \rangle &= \langle \varphi - \varphi q_k, px_n - px_m \rangle + \langle \varphi q_k, px_n - px_m \rangle \\ &= \langle \varphi(\mathbf{1} - q_k), px_n - px_m \rangle + \langle \varphi q_k, px_n - px_m \rangle \\ &= \langle (\mathbf{1} - q_k) \varphi (\mathbf{1} - q_k), px_n - px_m \rangle + \langle \varphi q_k, px_n - px_m \rangle. \end{aligned}$$

This gives

$$|\langle \varphi, px_n - px_m \rangle| \leq |\tau(\varphi(\mathbf{1} - q_k)(px_n - px_m)(\mathbf{1} - q_k))| + |\langle \varphi q_k, px_n - px_m \rangle|.$$

Since φq_k belongs to \mathcal{M} ,

$$\limsup_{n, m \rightarrow \infty} |\langle \varphi q_k, px_n - px_m \rangle| \leq 2 \|\varphi\|_{E^*(\mathcal{M}, \tau)} \cdot \sup_{a \in K} \|(\mathbf{1} - q_k) p a (\mathbf{1} - q_k)\|_{E(\mathcal{M}, \tau)}.$$

Now since $\mathbf{1} - q_k \downarrow_k 0$ and pK is E -equi-integrable, we obtain that

$$\lim_{n, m \rightarrow \infty} \langle \varphi q_k, px_n - px_m \rangle = 0.$$

The lemma is proved.

To complete the proof of Proposition 2.5, let $\{p_k\}_{k=1}^\infty$ be a sequence of projections that increases to $\mathbf{1}$ and $\tau(p_k) < \infty$ and fix $\varepsilon > 0$. Choose $k_0 \geq 1$ such that

$$\sup_{a \in K} \|(\mathbf{1} - p_{k_0}) a (\mathbf{1} - p_{k_0})\|_{E(\mathcal{M}, \tau)} \leq \varepsilon.$$

We have $K = p_{k_0}K + (\mathbf{1} - p_{k_0})Kp_{k_0} + (\mathbf{1} - p_{k_0})K(\mathbf{1} - p_{k_0})$ which implies that

$$K \subset p_{k_0}K + (\mathbf{1} - p_{k_0})Kp_{k_0} + \varepsilon B_{E(\mathcal{M}, \tau)}$$

where $B_{E(\mathcal{M}, \tau)}$ denotes the closed unit ball of $E(\mathcal{M}, \tau)$. From Lemma 2.7, the sets $p_{k_0}K$ and $(\mathbf{1} - p_{k_0})Kp_{k_0}$ are relatively weakly compact which concludes that K is relatively weakly compact. The proof is complete. \square

Remark 2.8. *If $\tau(\mathbf{1}) < \infty$, the proof above can be considerably shortened. In this case, $E(\mathcal{M}, \tau) \subset L^1(\mathcal{M}, \tau)$ so if K is E -equi-integrable, then it is relatively weakly compact in $L^1(\mathcal{M}, \tau)$ and one can argue as in the last part of Lemma 2.7 to conclude that K is relatively weakly compact in $E(\mathcal{M}, \tau)$.*

The following proposition should be compared with [3](Theorem 5.1 and Theorem 5.2). It generalizes well known property of equi-integrable sets in function spaces to the non-commutative settings.

Proposition 2.9. *Let E be a symmetric quasi-Banach space function space and K be a E -equi-integrable subset of $E(\mathcal{M}, \tau)$. For each sequence $\{x_n\}_{n=1}^\infty$ in K and $x \in \overline{K}$, the following are equivalent:*

- (a) $\lim_{n \rightarrow \infty} \|x_n - x\|_{E(\mathcal{M}, \tau)} = 0$;
- (b) $\{x_n\}_{n=1}^\infty$ converges to x in measure (as $n \rightarrow \infty$).

Proof. The implication (a) \Rightarrow (b) is trivial. For (b) \Rightarrow (a), we will assume that $x = 0$. Recall that there exists $0 < \alpha \leq 1$, such that $\mathcal{M} \cap L^\alpha(\mathcal{M}, \tau) \subset E(\mathcal{M}, \tau) \subset \mathcal{M} + L^\alpha(\mathcal{M}, \tau)$ with $\|x\|_{\mathcal{M} + L^\alpha(\mathcal{M}, \tau)} \leq \|x\|_{E(\mathcal{M}, \tau)} \leq 2\|x\|_{\mathcal{M} \cap L^\alpha(\mathcal{M}, \tau)}$ for every $x \in \mathcal{M} \cap L^\alpha(\mathcal{M}, \tau)$. The proposition will be proved by showing the following lemma:

Lemma 2.10. *For every $p \in \mathcal{M}_p$ with $\tau(p) < \infty$, $\lim_{n \rightarrow \infty} \|x_n p\|_{E(\mathcal{M}, \tau)} = \lim_{n \rightarrow \infty} \|p x_n\|_{E(\mathcal{M}, \tau)} = 0$.*

To see this lemma, fix $\varepsilon > 0$ and let $C = \max\{1, \tau(p)\}$. Since K is equi-integrable, there exists $\delta > 0$ such that whenever $q \in \mathcal{M}_p$ satisfies $\tau(q) < \delta$, then for every $n \in \mathbb{N}$, $\|q x_n\|_{E(\mathcal{M}, \tau)} \leq \varepsilon / (2)^{1/\alpha}$. Since both $\{x_n\}_{n=1}^\infty$ and $\{x_n^*\}_{n=1}^\infty$ converge to zero in measure, one can choose $n_0 \geq 1$ such that for each $n \geq n_0$, there exists a projection $p_n \in \mathcal{M}_p$ with $\tau(\mathbf{1} - p_n) < \delta$,

$$\|x_n p_n\|_{\mathcal{M}} < \frac{\varepsilon}{2[4C]^{1/\alpha}}$$

and

$$\|x_n^* p_n\|_{\mathcal{M}} < \frac{\varepsilon}{2[4C]^{1/\alpha}}.$$

For $n \geq n_0$,

$$\begin{aligned}
\|x_n p\|_{E(\mathcal{M}, \tau)}^\alpha &\leq \|x_n p_n p\|_{E(\mathcal{M}, \tau)}^\alpha + \|p_n x_n (\mathbf{1} - p_n) p\|_{E(\mathcal{M}, \tau)}^\alpha + \|(\mathbf{1} - p_n) x_n (\mathbf{1} - p_n) p\|_{E(\mathcal{M}, \tau)}^\alpha \\
&\leq 2^\alpha \max \left\{ \|x_n p_n\|_{\mathcal{M}}^\alpha, \|x_n p_n p\|_{L^\alpha(\mathcal{M}, \tau)}^\alpha \right\} \\
&\quad + 2^\alpha \max \left\{ \|x_n^* p_n\|_{\mathcal{M}}^\alpha, \|p(\mathbf{1} - p_n) x_n^* p_n\|_{L^\alpha(\mathcal{M}, \tau)}^\alpha \right\} + \|(\mathbf{1} - p_n) x_n (\mathbf{1} - p_n)\|_{E(\mathcal{M}, \tau)}^\alpha \\
&\leq 2 \cdot 2^\alpha \max \left\{ \varepsilon^\alpha / 2^\alpha 4C, (\varepsilon^\alpha / 2^\alpha 4C) \tau(p) \right\} + \varepsilon^\alpha / 2 \\
&\leq \varepsilon^\alpha.
\end{aligned}$$

A similar estimate works for $\{x_n^* p\}_{n=1}^\infty$. The lemma is verified.

To complete the proof of Proposition 2.9, choose a mutually disjoint family $\{e_i\}_{i \in I}$ of projections in \mathcal{M} with $\sum_{i \in I} e_i = \mathbf{1}$ for the strong operator topology and $\tau(e_i) < \infty$ for all $i \in I$. Using a similar argument as in [21], one can get an at most countable subset $\{e_k\}_{k=1}^\infty$ of $\{e_i\}_{i \in I}$ such that for each e_i outside of $\{e_k\}_{k=1}^\infty$, $e_i x_n = x_n e_i = 0$ for every $n \in \mathbb{N}$. Let $e = \sum_{k \in \mathbb{N}} e_k$. Replacing \mathcal{M} by $e \mathcal{M} e$ and τ by its restriction on $e \mathcal{M} e$, we may assume that $e = \mathbf{1}$. Let $p_n = \sum_{k \geq n} e_k$. It is clear that $p_n \downarrow_n 0$ and $\tau(\mathbf{1} - p_n) < \infty$ for every $n \in \mathbb{N}$. Fix $\varepsilon > 0$ and choose $n_0 \geq 1$ such that

$$\sup_{n \in \mathbb{N}} \|p_{n_0} x_n p_{n_0}\|_{E(\mathcal{M}, \tau)} \leq \varepsilon.$$

We get that

$$\limsup_{n \rightarrow \infty} \|x_n\|_{E(\mathcal{M}, \tau)}^\alpha \leq \lim_{n \rightarrow \infty} \|x_n (\mathbf{1} - p_{n_0})\|_{E(\mathcal{M}, \tau)}^\alpha + \lim_{n \rightarrow \infty} \|x_n^* (\mathbf{1} - p_{n_0})\|_{E(\mathcal{M}, \tau)}^\alpha + \varepsilon = \varepsilon$$

and since ε is arbitrary, the proof is complete. \square

For the rest of this section, we collect some results that will be useful in the later sections of the paper.

The inequality given below can be viewed as the analogue of the well-known fact on normal functional on von Neumann algebra, $|\varphi(a)|^2 \leq \|\varphi\| \cdot |\varphi|(aa^*)$ whenever $a \in \mathcal{M}$ and $\varphi \in \mathcal{M}_*$ [19, Proposition 4.6 p. 146], to the general case of symmetric spaces of measurable operators.

Proposition 2.11. *Let $x \in E(\mathcal{M}, \tau)$ and $y \in \mathcal{M}$ then*

$$\|xy\|_{E(\mathcal{M}, \tau)} \leq \| |x|y \|_{E(\mathcal{M}, \tau)} \leq \|x\|_{E(\mathcal{M}, \tau)}^{\frac{1}{2}} \cdot \|y^* |x|y\|_{E(\mathcal{M}, \tau)}^{\frac{1}{2}}.$$

Proof. Let $x = u|x|$ be the polar decomposition of x . Then $\|xy\|_{E(\mathcal{M}, \tau)} = \|u|x|y\|_{E(\mathcal{M}, \tau)} \leq \|u\|_\infty \cdot \| |x|y \|_{E(\mathcal{M}, \tau)}$. Also $\| |x|y \|_{E(\mathcal{M}, \tau)} = \left\| |x|^{\frac{1}{2}} |x|^{\frac{1}{2}} y \right\|_{E(\mathcal{M}, \tau)}$ and using Hölder's inequality,

$$\begin{aligned}
\| |x|y \|_{E(\mathcal{M}, \tau)} &\leq \left\| |x|^{\frac{1}{2}} \right\|_{E^{(2)}(\mathcal{M}, \tau)} \cdot \left\| |x|^{\frac{1}{2}} y \right\|_{E^{(2)}(\mathcal{M}, \tau)} \\
&= \|x\|_{E(\mathcal{M}, \tau)}^{\frac{1}{2}} \cdot \|y^* |x|y\|_{E(\mathcal{M}, \tau)}^{\frac{1}{2}}.
\end{aligned}$$

□

Remark 2.12. Let K be a bounded subset of $E(\mathcal{M}, \tau)$. If we set $|K| := \{|a| : a \in K\}$, then it is clear from Proposition 2.11 that if $|K|$ is E -equi-integrable then for every decreasing projections $e_n \downarrow_n 0$, $\limsup_{n \rightarrow \infty} \sup_{x \in K} \|xe_n\|_{E(\mathcal{M}, \tau)} = \limsup_{n \rightarrow \infty} \sup_{x \in K} \|e_n x\|_{E(\mathcal{M}, \tau)} = 0$. In particular if $|K|$ is E -equi-integrable then so is K .

Proposition 2.13. Assume that E is α -convex with constant 1 for some $0 < \alpha \leq 1$. Let $\{p_n\}_{n=1}^\infty$ be a sequence of decreasing projections in \mathcal{M} and K be a bounded subset of $E(\mathcal{M}, \tau)$ such that:

(i) $p_n \downarrow_n 0$;

(ii) For each $n \geq 1$, the sets $(\mathbf{1} - p_n)K$ and $|K(\mathbf{1} - p_n)|$ are E -equi-integrable.

Then K is E -equi-integrable if and only if $\limsup_{n \rightarrow \infty} \sup_{a \in K} \|p_n a p_n\|_{E(\mathcal{M}, \tau)} = 0$.

Proof. We will show the non trivial implication. Fix $f_k \downarrow_k 0$, a sequence in \mathcal{M}_p . We need to show that $\limsup_{k \rightarrow \infty} \sup_{a \in K} \|f_k a f_k\|_{E(\mathcal{M}, \tau)} = 0$.

We will assume without loss of generality that K is a subset of the unit ball of $E(\mathcal{M}, \tau)$. For every $a \in K$,

$$\begin{aligned} f_k a f_k &= f_k(\mathbf{1} - p_n) a f_k + f_k p_n a f_k \\ &= f_k(\mathbf{1} - p_n) a f_k + f_k p_n a (\mathbf{1} - p_n) f_k + f_k p_n a p_n f_k. \end{aligned}$$

Since $E(\mathcal{M}, \tau)$ is α -convex, we get:

$$\begin{aligned} \|f_k a f_k\|_{E(\mathcal{M}, \tau)}^\alpha &\leq \|f_k(\mathbf{1} - p_n) a f_k\|_{E(\mathcal{M}, \tau)}^\alpha + \|f_k p_n a (\mathbf{1} - p_n) f_k\|_{E(\mathcal{M}, \tau)}^\alpha + \|f_k p_n a p_n f_k\|_{E(\mathcal{M}, \tau)}^\alpha \\ &\leq \|f_k(\mathbf{1} - p_n) a f_k\|_{E(\mathcal{M}, \tau)}^\alpha + \|a(\mathbf{1} - p_n) f_k\|_{E(\mathcal{M}, \tau)}^\alpha + \|p_n a p_n\|_{E(\mathcal{M}, \tau)}^\alpha. \end{aligned}$$

Using Proposition 2.11 on the second term, we have

$$\begin{aligned} \|f_k a f_k\|_{E(\mathcal{M}, \tau)}^\alpha &\leq \|f_k(\mathbf{1} - p_n) a f_k\|_{E(\mathcal{M}, \tau)}^\alpha + \|a(\mathbf{1} - p_n)\|_{E(\mathcal{M}, \tau)}^{\frac{\alpha}{2}} \cdot \|f_k |a(\mathbf{1} - p_n)| f_k\|_{E(\mathcal{M}, \tau)}^{\frac{\alpha}{2}} \\ &\quad + \|p_n a p_n\|_{E(\mathcal{M}, \tau)}^\alpha. \end{aligned}$$

Let $\varepsilon > 0$, choose n_0 large enough so that $\sup_{a \in K} \|p_{n_0} a p_{n_0}\|_{E(\mathcal{M}, \tau)} < \varepsilon$. We conclude that

$$\begin{aligned} \limsup_{k \rightarrow \infty} \sup_{a \in K} \|f_k a f_k\|_{E(\mathcal{M}, \tau)}^\alpha &\leq \limsup_{k \rightarrow \infty} \sup_{a \in K} \|f_k(\mathbf{1} - p_{n_0}) a f_k\|_{E(\mathcal{M}, \tau)}^\alpha \\ &\quad + \limsup_{k \rightarrow \infty} \sup_{a \in K} \|f_k |a(\mathbf{1} - p_{n_0})| f_k\|_{E(\mathcal{M}, \tau)}^{\frac{\alpha}{2}} + \varepsilon^\alpha. \end{aligned}$$

By (ii), the first two terms converge to zero so $\limsup_{k \rightarrow \infty} \sup_{a \in K} \|f_k a f_k\|_{E(\mathcal{M}, \tau)} \leq \varepsilon$ and since ε is arbitrary, the proof is complete. □

The next proposition can be found in [6] (Proposition 2.5).

Proposition 2.14. *Assume that E is α -convex with constant 1 for some $0 < \alpha \leq 1$ and satisfies a lower q -estimate with constant 1 for some finite $q \geq \alpha$. If $k = 2q/\alpha$, then for all $y \in E(\mathcal{M}, \tau)$, for all projections $e, f \in \mathcal{M}$ with $e + f = 1$ and $\tau(e) < \infty$, it follows that*

$$\|eye\|_{E(\mathcal{M}, \tau)}^k + \|eyf\|_{E(\mathcal{M}, \tau)}^k + \|fye\|_{E(\mathcal{M}, \tau)}^k + \|fyf\|_{E(\mathcal{M}, \tau)}^k \leq \|y\|_{E(\mathcal{M}, \tau)}^k.$$

The proof of the next lemma is just a notational adjustment of the proof of a lemma from the L^1 -case [7] so we will leave the details to the interested readers.

Lemma 2.15. *Let $\{x_k\}_{k=1}^\infty$ be a bounded sequence in $E(\mathcal{M}, \tau)$ and $\{e_n\}_{n=1}^\infty$ be a decreasing sequence of projections in \mathcal{M} such that $e_n \downarrow_n 0$. Assume that $\lim_{n \rightarrow \infty} \sup_{k \in \mathbb{N}} \|e_n x_k e_n\|_{E(\mathcal{M}, \tau)} = \gamma > 0$ then there exists a subsequence $\{x_{k_n}\}_{n=1}^\infty$ so that $\lim_{n \rightarrow \infty} \|e_n x_{k_n} e_n\|_{E(\mathcal{M}, \tau)} = \gamma$.*

3. KADEC-PEŁCZYŃSKI THEOREM FOR SYMMETRIC SPACES

The main result of the present article is the following theorem.

Theorem 3.1. *Let E be an order continuous symmetric quasi-Banach function space in \mathbb{R}^+ that is α -convex with constant 1 for some $0 < \alpha \leq 1$ and suppose that E satisfies a lower q -estimate with constant 1 for some $q \geq \alpha$.*

Let $\{x_n\}_{n=1}^\infty$ be a bounded sequence in $E(\mathcal{M}, \tau)$ then there exists a subsequence $\{x_{n_k}\}_{k=1}^\infty$ of $\{x_n\}_{n=1}^\infty$, bounded sequences $\{y_k\}_{k=1}^\infty$ and $\{z_k\}_{k=1}^\infty$ in $E(\mathcal{M}, \tau)$ and a decreasing sequence of projections $p_k \downarrow_n 0$ in \mathcal{M} such that:

- (i) $x_{n_k} = y_k + z_k$ for all $k \geq 1$;
- (ii) $\{y_k : k \geq 1\}$ is E -equi-integrable and $p_k y_k p_k = 0$ for all $k \geq 1$;
- (iii) $\{z_k\}_{k=1}^\infty$ is such that $p_k z_k p_k = z_k$ for all $k \geq 1$.

The proof will be divided into several steps. Without loss of generality, we will assume that the sequence $\{x_k\}_{k=1}^\infty$ is a subset of the unit ball of $E(\mathcal{M}, \tau)$. Since we are dealing with sequences, we can and do assume without loss of generality that \mathcal{M} is countably decomposable (see [21] and the proof of Proposition 2.9 above for the details of such reduction).

Set $\mathcal{D}_1 := \{\{e_n\}_{n=1}^\infty \subset \mathcal{M}_p : e_n \downarrow_n 0 \text{ and } \tau(e_1) < \infty\}$ and consider

$$\delta := \sup \left\{ \lim_{n \rightarrow \infty} \sup_{k \in \mathbb{N}} \|e_n |x_k| e_n\|_{E(\mathcal{M}, \tau)} : \{e_n\}_{n=1}^\infty \in \mathcal{D}_1 \right\}.$$

As in [7], one can show that δ is attained and using Lemma 2.15, one can choose a subsequence of $\{x_n\}_{n=1}^\infty$ (which we will denote again by $\{x_n\}_{n=1}^\infty$) and $\{e_n\}_{n=1}^\infty \in \mathcal{D}_1$ such that

$$(3.1) \quad \delta = \lim_{n \rightarrow \infty} \|e_n |x_n| e_n\|_{E(\mathcal{M}, \tau)}$$

and

$$(3.2) \quad \delta^* = \lim_{n \rightarrow \infty} \|e_n |x_n^*| e_n\|_{E(\mathcal{M}, \tau)} = \sup \left\{ \lim_{n \rightarrow \infty} \sup_k \|q_n |x_k^*| q_n\|_{E(\mathcal{M}, \tau)} ; \{q_n\}_{n=1}^\infty \in \mathcal{D}_1 \right\}.$$

For each $n \geq 1$, set $v_n := x_n - e_n x_n e_n$ and let $V := \{v_n : n \geq 1\}$.

Lemma 3.2. *There exists a sequence of projections $\{g_n\}_{n=1}^\infty$ in \mathcal{M} with:*

- (1) For every $n \geq 1$, $g_n \leq \mathbf{1} - e_n$;
- (2) $\tau(g_n) < \infty$, in particular g_n is a finite projection;
- (3) $g_n \uparrow^n \mathbf{1}$.

Proof. The lemma can be obtained inductively. Since \mathcal{M} is countably decomposable, there exists φ_0 a faithful normal state in \mathcal{M}_* . Since $\mathbf{1} - e_n$ is a semifinite projection, there exists a sequence of projections $\{g_j^{(n)}\}_{j=1}^\infty$ with $\tau(g_j^{(n)}) < \infty$ for every $j \geq 1$ and $g_j^{(n)} \uparrow^j \mathbf{1} - e_n$. One can choose $j_n \geq 1$ such that $\varphi_0(\mathbf{1} - e_n) - \varphi_0(g_{j_n}^{(n)}) < 1/n$. Set

$$\begin{cases} g_n := g_{j_1}^{(1)} & \text{for } n = 1 \\ g_n = g_{j_n}^{(n)} \vee g_{n-1} & \text{for } n > 1. \end{cases}$$

It is easy to verify that $\{g_n\}_{n=1}^\infty$ satisfies the requirements of the lemma. \square

For each $n \geq 1$, let $p_n = \mathbf{1} - g_n$. Clearly $p_n \downarrow_n 0$, $\mathbf{1} - p_n$ is a finite projection and $p_n \geq e_n$ for each $n \geq 1$.

Lemma 3.3. *For each $n \geq 1$, the sets $|V(\mathbf{1} - p_n)|$ and $|(\mathbf{1} - p_n)V|$ are E -equi-integrable.*

Proof. Let us show that for every $n \geq 1$, $|V(\mathbf{1} - p_n)|$ is an E -equi-integrable set. Assume that there exists $k_0 \geq 1$ such that $|V(\mathbf{1} - p_{k_0})|$ is not E -equi-integrable. By definition, there exists a decreasing sequence of projections $q_n \downarrow_n 0$ such that $\limsup_{n \rightarrow \infty} \sup_{a \in |V(\mathbf{1} - p_{k_0})|} \|q_n a q_n\|_{E(\mathcal{M}, \tau)} > 0$, that is

$$\limsup_{n \rightarrow \infty} \sup_{m \in \mathbb{N}} \|q_n |v_m(\mathbf{1} - p_{k_0})| q_n\|_{E(\mathcal{M}, \tau)} > 0.$$

Choose a strictly increasing sequence $\{m_n\}_{n=1}^\infty$ of \mathbb{N} such that

$$\lim_{n \rightarrow \infty} \|q_n |v_{m_n}(\mathbf{1} - p_{k_0})| q_n\|_{E(\mathcal{M}, \tau)} > 0.$$

Let u_{n, k_0} be a bounded operator such that $|v_{m_n}(\mathbf{1} - p_{k_0})| = u_{n, k_0} v_{m_n}(\mathbf{1} - p_{k_0})$. We get that

$$\begin{aligned} \|q_n |v_{m_n}(\mathbf{1} - p_{k_0})| q_n\|_{E(\mathcal{M}, \tau)} &= \|q_n u_{n, k_0} v_{m_n}(\mathbf{1} - p_{k_0}) q_n\|_{E(\mathcal{M}, \tau)} \\ &= \|q_n u_{n, k_0} [x_{m_n} - e_{m_n} x_{m_n} e_{m_n}] (\mathbf{1} - p_{k_0}) q_n\|_{E(\mathcal{M}, \tau)}. \end{aligned}$$

We recall that $e_{k_0} \leq p_{k_0}$ and since $\{e_n\}_{n=1}^\infty$ is decreasing, for $m_n \geq k_0$, $e_{m_n} \leq p_{k_0}$ and therefore $e_{m_n}(\mathbf{1} - p_{k_0}) = 0$ and since $\|q_n u_{n, k_0}\|_\infty \leq 1$, we obtain that for n large enough,

$$\begin{aligned} \|q_n |v_{m_n}(\mathbf{1} - p_{k_0})| q_n\|_{E(\mathcal{M}, \tau)} &= \|q_n u_{n, k_0} (x_{m_n}) (\mathbf{1} - p_{k_0}) q_n\|_{E(\mathcal{M}, \tau)} \\ &\leq \|x_{m_n} (\mathbf{1} - p_{k_0}) q_n\|_{E(\mathcal{M}, \tau)}. \end{aligned}$$

Using Proposition 2.11, with $x = x_{m_n}$ and $y = (\mathbf{1} - p_{k_0})q_n$, we get

$$\begin{aligned} \|q_n|v_{m_n}(\mathbf{1} - p_{k_0})|p_n\|_{E(\mathcal{M},\tau)} &\leq \|x_{m_n}\|_{E(\mathcal{M},\tau)}^{\frac{1}{2}} \cdot \|q_n(\mathbf{1} - p_{k_0})|x_{m_n}|(\mathbf{1} - p_{k_0})q_n\|_{E(\mathcal{M},\tau)}^{\frac{1}{2}} \\ &\leq \|q_n(\mathbf{1} - p_{k_0})|x_{m_n}|(\mathbf{1} - p_{k_0})q_n\|_{E(\mathcal{M},\tau)}^{\frac{1}{2}}. \end{aligned}$$

This implies that

$$\limsup_{n \rightarrow \infty} \|q_n(\mathbf{1} - p_{k_0})|x_{m_n}|(\mathbf{1} - p_{k_0})q_n\|_{E(\mathcal{M},\tau)} > 0.$$

Let s_n be the left support projection of $(\mathbf{1} - p_{k_0})q_n$ (this is equal to the right support projection of $q_n(\mathbf{1} - p_{k_0})$). We have

$$\begin{aligned} \|q_n(\mathbf{1} - p_{k_0})|x_{m_n}|(\mathbf{1} - p_{k_0})q_n\|_{E(\mathcal{M},\tau)} &= \|q_n(\mathbf{1} - p_{k_0})s_n|x_{m_n}|s_n(\mathbf{1} - p_{k_0})q_n\|_{E(\mathcal{M},\tau)} \\ &\leq \|s_n|x_{m_n}|s_n\|_{E(\mathcal{M},\tau)}. \end{aligned}$$

By the definition of support projection, $s_n \leq (\mathbf{1} - p_{k_0})$ for every $n \geq 1$, so $\{s_n\}_{n=1}^{\infty}$ is a sequence of finite projections. As in proof of Lemma 2.6, we note that $s_n = q_n \vee p_{k_0} - p_{k_0}$ and as before, $s_n \sim q_n - q_n \wedge p_{k_0}$. Now since $q_n \downarrow_n 0$, $q_n - q_n \wedge p_{k_0} \downarrow_n 0$ hence $\tau(s_n) = \tau(q_n - q_n \wedge p_{k_0})$ converges to zero which implies that $s_n \downarrow_n 0$. Therefore, $\{s_n\}_{n=1}^{\infty} \in \mathcal{D}_1$.

In summary, we get $\{s_n\}_{n=1}^{\infty} \in \mathcal{D}_1$ with $s_n \leq \mathbf{1} - p_{k_0}$ for each $n \geq 1$ and for some $\gamma > 0$,

$$(3.3) \quad \limsup_{n \rightarrow \infty} \|s_n|x_{m_n}|s_n\| = \gamma.$$

Let $f_n := s_n \vee e_{m_n}$.

For each $m_n \geq k_0$, $s_n \leq \mathbf{1} - p_{k_0} \leq \mathbf{1} - e_{k_0}$ so $s_n \perp e_{m_n}$ hence $f_n = s_n + e_{m_n}$. In particular $\{f_n\}_{n=1}^{\infty} \in \mathcal{D}_1$.

Using Proposition 2.14 (it applies since $\tau(f_n) < \infty$),

$$\begin{aligned} \|f_n|x_{m_n}|f_n\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} &\geq \|s_n|x_{m_n}|s_n\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} + \|e_{m_n}|x_{m_n}|e_{m_n}\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} \\ &\quad + \|s_n|x_{m_n}|e_{m_n}\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} + \|e_{m_n}|x_{m_n}|e_{m_n}\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} \\ &\geq \|s_n|x_{m_n}|s_n\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}} + \|e_{m_n}|x_{m_n}|e_{m_n}\|_{E(\mathcal{M},\tau)}^{\frac{2q}{\alpha}}. \end{aligned}$$

Taking the limit as n tends to ∞ , one gets from (3.1) and (3.3) that $\delta^{\frac{2q}{\alpha}} \geq \gamma^{\frac{2q}{\alpha}} + \delta^{\frac{2q}{\alpha}}$. This is a contradiction since $\gamma > 0$.

We conclude that for every $n \geq 1$, the set $|V(\mathbf{1} - p_n)|$ is an E -equi-integrable set.

For the case of $|(\mathbf{1} - p_n)V|$, it is enough to repeat the argument above for $V^*(\mathbf{1} - p_n)$ using the definition of δ^* (instead of δ). Details are left to the reader. This ends the proof of the lemma. \square

We will proceed to the proof of Theorem 3.1. Consider two cases.

Case 1: Assume that V is E -equi-integrable.

It is enough to set $y_n = x_{m_n} - e_{m_n}x_{m_n}e_{m_n}$ and $z_n = e_{m_n}x_{m_n}e_{m_n}$.

Case 2: Assume that V is not E -equi-integrable.

Proposition 2.13 and Lemma 3.3 imply that there exists $\nu > 0$ such that

$$\limsup_{n \rightarrow \infty} \sup_{v \in V} \|p_n v p_n\|_{E(\mathcal{M}, \tau)} = \nu > 0.$$

Choose a subsequence $\{v_{n_k}\}_{k=1}^\infty$ such that

$$(3.4) \quad \lim_{k \rightarrow \infty} \|p_k v_{n_k} p_k\|_{E(\mathcal{M}, \tau)} = \nu > 0.$$

For each $k \geq 1$, let $w_k := v_{n_k} - p_k v_{n_k} p_k$ and set

$$W := \{w_k : k \geq 1\}.$$

Lemma 3.4. *The set W is E -equi-integrable.*

Proof. We note first that if $k \geq n$, then $(\mathbf{1} - p_n)w_k = (\mathbf{1} - p_n)v_k$ and $w_k(\mathbf{1} - p_n) = v_k(\mathbf{1} - p_n)$ so for fixed $n \geq 1$, $(\mathbf{1} - p_n)W = \{(\mathbf{1} - p_n)w_k : k < n\} \cup \{(\mathbf{1} - p_n)v_k : k \geq n\}$. Similarly, $W(\mathbf{1} - p_n) = \{w_k(\mathbf{1} - p_n) : k < n\} \cup \{v_k(\mathbf{1} - p_n) : k \geq n\}$.

Lemma 3.3 implies that for every $n \geq 1$, both $|W(\mathbf{1} - p_n)|$ and $(\mathbf{1} - p_n)W$ are E -equi-integrable sets. Therefore, if W is not E -equi-integrable, there would be a subsequence $\{w_{k(j)}\}_{j=1}^\infty$ of $\{w_k\}_{k=1}^\infty$ and $\varepsilon > 0$ such that

$$(3.5) \quad \lim_{j \rightarrow \infty} \|p_j w_{k(j)} p_j\|_{E(\mathcal{M}, \tau)} = \varepsilon.$$

Using Proposition 2.11 on $v_{n_{k(j)}}$ and $p_j = (p_j - p_{k(j)}) + p_{k(j)}$, we obtain:

$$\begin{aligned} \left\| p_j v_{n_{k(j)}} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} &\geq \left\| (p_j - p_{k(j)}) v_{n_{k(j)}} (p_j - p_{k(j)}) \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \left\| p_{k(j)} v_{n_{k(j)}} (p_j - p_{k(j)}) \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \\ &\quad \left\| (p_j - p_{k(j)}) v_{n_{k(j)}} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \left\| (p_j - p_{k(j)}) v_{n_{k(j)}} (p_j - p_{k(j)}) \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}}. \end{aligned}$$

Taking into account the identities, $(p_j - p_{k(j)}) v_{n_{k(j)}} (p_j - p_{k(j)}) = (p_j - p_{k(j)}) w_{k(j)} (p_j - p_{k(j)})$, $(p_j - p_{k(j)}) v_{n_{k(j)}} p_{k(j)} = p_j w_{k(j)} p_{k(j)}$ and $p_{k(j)} v_{n_{k(j)}} (p_j - p_{k(j)}) = p_{k(j)} w_{k(j)} p_j$, one can deduce that,

$$\begin{aligned} \left\| p_j v_{n_{k(j)}} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} &\geq \left\| (p_j - p_{k(j)}) w_{k(j)} (p_j - p_{k(j)}) \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \left\| p_j w_{k(j)} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \\ &\quad \left\| p_{k(j)} w_{k(j)} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \left\| p_{k(j)} v_{n_{k(j)}} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}}. \end{aligned}$$

Let $C(q, \alpha)$ be the norm of the identity map from $\ell_3^{\frac{2q}{\alpha}}$ onto ℓ_3^α , where $\ell_3^{\frac{2q}{\alpha}}$ (resp. ℓ_3^α) denotes the 3-dimensional $\ell^{\frac{2q}{\alpha}}$ -space (resp. ℓ^α -space). We have

$$\begin{aligned} \left\| p_j v_{n_{k(j)}} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} &\geq C(q, \alpha)^{\frac{2q}{\alpha}} \left[\left(\left\| (p_j - p_{k(j)}) w_{k(j)} (p_j - p_{k(j)}) \right\|_{E(\mathcal{M}, \tau)}^\alpha + \right. \right. \\ &\quad \left. \left. \left\| p_j w_{k(j)} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^\alpha + \left\| p_{k(j)} w_{k(j)} p_j \right\|_{E(\mathcal{M}, \tau)}^\alpha \right)^{\frac{1}{\alpha}} \right]^{\frac{2q}{\alpha}} + \\ &\quad \left\| p_{k(j)} v_{n_{k(j)}} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}}. \end{aligned}$$

We remark that $p_j w_{k(j)} p_j = (p_j - p_{k(j)}) w_{k(j)} (p_j - p_{k(j)}) + p_j w_{k(j)} p_{k(j)} + p_{k(j)} w_{k(j)} p_j$ and since $E(\mathcal{M}, \tau)$ is α -convex (with constant 1), the above inequality implies

$$\left\| p_j v_{n_{k(j)}} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} \geq C(q, \alpha)^{\frac{2q}{\alpha}} \left\| p_j w_{k(j)} p_j \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}} + \left\| p_{k(j)} v_{n_{k(j)}} p_{k(j)} \right\|_{E(\mathcal{M}, \tau)}^{\frac{2q}{\alpha}},$$

and taking the limit as $j \rightarrow \infty$, we get from (3.4) and (3.5) that

$$\nu^{\frac{2q}{\alpha}} \geq C(q, \alpha)^{\frac{2q}{\alpha}} \varepsilon^{\frac{2q}{\alpha}} + \nu^{\frac{2q}{\alpha}}.$$

This is a contradiction since $\varepsilon > 0$, so W is a E -equi-integrable set. The lemma is proved.

To complete the proof of Theorem 3.1, we note that $W = \{x_{n_k} - p_k x_{n_k} p_k : k \geq 1\}$ so if we set $y_k = x_{n_k} - p_k x_{n_k} p_k$ and $z_k = p_k x_{n_k} p_k$. The proof of is complete. \square

Remarks 3.5. (1) If $\mathcal{M} = \mathcal{B}(\ell^2)$ with the usual trace, then every projection of finite trace is a finite rank projection so in the proof above, $\delta = \delta^* = 0$. In the particular case of unitary matrix space C_E where E is a symmetric sequence space, one proceed directly to Case 2 by setting $W := \{x_n - p_n x_n p_n; n \geq 1\}$ where $\{p_n\}_{n=1}^\infty$ is an arbitrary sequence of projections satisfying: $p_n \downarrow_n 0$ and for every $n \geq 1$, $1 - p_n$ is a finite projection.

(2) If \mathcal{M} is a finite von Neumann algebra with a normalized finite trace τ and E is a symmetric space on $[0, 1]$ satisfying the assumptions of Theorem 3.1, it is enough to take $p_n = e_n$ (i.e $g_n = \mathbf{1} - e_n$ on Lemma 3.2) and conclude immediately as in Lemma 3.3 that V is E -equi-integrable.

(3) In the proof above, it is clear that the projections $\{p_k\}_{k=1}^\infty$ are such that either $\tau(p_1) < \infty$ or $\tau(\mathbf{1} - p_k) < \infty$ for all $k \geq 1$. In fact, the argument above shows that if $\{e_n\}_{n=1}^\infty$ is a sequence in \mathcal{D}_1 that attained the quantities δ and δ^* , then any sequence of projections satisfying $p_n \downarrow_n 0$, $e_n \leq p_n$ for each $n \geq 1$ and $\tau(\mathbf{1} - p_n) < \infty$ for each $n \geq 1$, would satisfy the conclusion of Theorem 3.1.

The following extension shows that if one considers finitely many bounded sequences in $E(\mathcal{M}, \tau)$, one can choose a single sequence of projections that works for each sequence.

Corollary 3.6. *If \mathcal{M} and E are as in Theorem 3.1 and $\{x^{(1)}\}_{n=1}^\infty, \{x^{(2)}\}_{n=1}^\infty, \dots, \{x^{(j_0)}\}_{n=1}^\infty$ be finitely many bounded sequences in $E(\mathcal{M}, \tau)$. Then there exist a strictly increasing sequence $\{n_k\}_{k=1}^\infty$ of \mathbb{N} and a sequence of decreasing projections $p_k \downarrow_k 0$ in \mathcal{M} such that for each $1 \leq j \leq j_0$, the set $\{x_{n_k}^{(j)} - p_k x_{n_k}^{(j)} p_k : k \geq 1\}$ is E -equi-integrable.*

Proof. For $1 \leq j \leq j_0$, we set, as in the proof of Theorem 3.1,

$$\delta_j := \sup \left\{ \limsup_{n \rightarrow \infty} \sup_{k \in \mathbb{N}} \left\| e_n |x_k^{(j)}| e_n \right\|_{E(\mathcal{M}, \tau)} : \{e_n\}_{n=1}^\infty \in \mathcal{D}_1 \right\}.$$

One can choose a strictly increasing sequence $\{n_k\}_{k=1}^\infty$ in \mathbb{N} such that for each $1 \leq j \leq j_0$, there exists a sequence $\{e_k^{(j)}\}_{k=1}^\infty \in \mathcal{D}_1$ with

$$\delta_j = \lim_{k \rightarrow \infty} \left\| e_k^{(j)} |x_{n_k}^{(j)}| e_k^{(j)} \right\|_{E(\mathcal{M}, \tau)}$$

and

$$\delta_j^* = \lim_{k \rightarrow \infty} \left\| e_k^{(j)} |x_{n_k}^{(j)}| * |e_k^{(j)}| \right\|_{E(\mathcal{M}, \tau)} = \sup \left\{ \limsup_{n \rightarrow \infty} \sup_k \left\| q_n |x_{n_k}^{(j)*}| q_n \right\|_{E(\mathcal{M}, \tau)} ; \{q_n\}_{n=1}^\infty \in \mathcal{D}_1 \right\}.$$

For every $k \geq 1$, set $e_k := \vee_{1 \leq j \leq j_0} e_k^{(j)}$. Since $\tau(e_k) \leq \sum_{j=1}^{j_0} \tau(e_k^{(j)})$, it is clear that the sequence $\{e_k\}_{k=1}^\infty$ belongs to \mathcal{D}_1 and each of the δ_j 's and δ_j^* 's are attained at $\{e_k\}_{k=1}^\infty$. One can complete the proof by proceeding as in the proof of Theorem 3.1, simultaneously on the finite set of sequences and the fixed $\{e_k\}_{k=1}^\infty$. \square

Our next result shows that the decreasing projections in the decomposition can be replaced by mutually disjoint projections.

Theorem 3.7. *Let E be an order continuous quasi-Banach function space as in Theorem 3.1. Let $\{x_n\}_{n=1}^\infty$ be a bounded sequence in $E(\mathcal{M}, \tau)$ then there exists a subsequence $\{x_{n_k}\}_{k=1}^\infty$ of $\{x_n\}_{n=1}^\infty$, bounded sequences $\{\varphi_k\}_{k=1}^\infty$ and $\{\zeta_k\}_{k=1}^\infty$ in $E(\mathcal{M}, \tau)$ and mutually disjoint sequence of projections $\{e_k\}_{k=1}^\infty$ such that:*

- (i) $x_{n_k} = \varphi_k + \zeta_k$ for all $k \geq 1$;
- (ii) $\{\varphi_k : k \geq 1\}$ is E -equi-integrable and $e_k \varphi_k e_k = 0$ for all $k \geq 1$;
- (iii) $\{\zeta_k\}_{k=1}^\infty$ is such that $e_k \zeta_k e_k = \zeta_k$ for all $k \geq 1$.

Proof. Let $\{x_n\}_{n=1}^\infty$ be a bounded sequence in $E(\mathcal{M}, \tau)$ and suppose (by taking a subsequence if necessary), $x_n = y_n + z_n$ with $p_n y_n p_n = 0$, the set $\{y_n : n \geq 1\}$ is E -equi-integrable and $p_n z_n p_n = z_n$ for all $n \geq 1$, be the decomposition of $\{x_n\}_{n=1}^\infty$ as in Theorem 3.1.

Let $n_1 = 1$. Since $p_n \downarrow_n 0$ and

$$p_1 z_1 p_1 - (p_1 - p_n) z_1 (p_1 - p_n) = p_n z_1 p_1 + p_1 z_1 p_n - p_n z_1 p_n,$$

Proposition reorder-continuity(part(ii)) shows that

$$\lim_{n \rightarrow \infty} \|p_1 z_1 p_1 - (p_1 - p_n) z_1 (p_1 - p_n)\|_{E(\mathcal{M}, \tau)} = 0.$$

Choose $n_2 > n_1 = 1$ such that

$$\|p_1 z_1 p_1 - (p_1 - p_{n_2}) z_1 (p_1 - p_{n_2})\|_{E(\mathcal{M}, \tau)} < \frac{1}{2}.$$

Inductively, one can construct $n_1 < n_2 < \dots < n_k < \dots$ such that

$$\|p_{n_k} z_{n_k} p_{n_k} - (p_{n_k} - p_{n_{k+1}}) z_{n_k} (p_{n_k} - p_{n_{k+1}})\|_{E(\mathcal{M}, \tau)} < \frac{1}{2^k}.$$

Since $z_n = p_n z_n p_n$ for every $n \geq 1$, one gets

$$\|z_{n_k} - (p_{n_k} - p_{n_{k+1}}) z_{n_k} (p_{n_k} - p_{n_{k+1}})\|_{E(\mathcal{M}, \tau)} < \frac{1}{2^k}.$$

For every $k \geq 1$, set

$$\begin{aligned} e_k &:= p_{n_k} - p_{n_{k+1}} \\ \zeta_k &:= (p_{n_k} - p_{n_{k+1}}) z_{n_k} (p_{n_k} - p_{n_{k+1}}) \\ \varphi_k &:= y_{n_k} + [z_{n_k} - e_k z_{n_k} e_k]. \end{aligned}$$

Since $\{y_{n_k} : k \geq 1\}$ is a E -equi-integrable set and $\lim_{k \rightarrow \infty} \|z_{n_k} - e_k z_{n_k} e_k\|_{E(\mathcal{M}, \tau)} = 0$, it is clear that $\{\varphi_k : k \geq 1\}$ is E -equi-integrable. Also $\{e_k\}_{k=1}^{\infty}$ is mutually disjoint. The proof is complete. \square

Corollary 3.8. *Let E be an order-continuous symmetric Banach function space on \mathbb{R}^+ with the Fatou property. Let $\{x_n\}_{n=1}^{\infty}$ be a bounded sequence in $E(\mathcal{M}, \tau)$ then there exists a subsequence $\{x_{n_k}\}_{k=1}^{\infty}$ of $\{x_n\}_{n=1}^{\infty}$, bounded sequences $\{\varphi_k\}_{k=1}^{\infty}$ and $\{\zeta_k\}_{k=1}^{\infty}$ in $E(\mathcal{M}, \tau)$ and mutually disjoint sequence of projections $\{e_k\}_{k=1}^{\infty}$ such that:*

- (i) $x_{n_k} = \varphi_k + \zeta_k$ for all $k \geq 1$;
- (ii) $\{\varphi_k : k \geq 1\}$ is E -equi-integrable and $e_k \varphi_k e_k = 0$ for all $k \geq 1$;
- (iii) $\{\zeta_k\}_{k=1}^{\infty}$ is such that $e_k \zeta_k e_k = \zeta_k$ for all $k \geq 1$.

Proof. Assume that E has the Fatou property (equivalently E does not contain c_0). Since E is symmetric, $E \not\supset c_0$ is equivalent to E not containing ℓ_{∞}^n uniformly, and therefore E satisfies the q -lower estimate for some q and one can renorm E so that it satisfies the lower q -estimate of constant 1. All of these facts can be found in [14]. \square

The proof of Theorem 3.1 can be adjusted to obtain decompositions where the projections are taken only on one side, that is, the following result follows:

Corollary 3.9. *Let E be an order continuous quasi-Banach function space in \mathbb{R}^+ that is α -convex with constant 1 for some $0 < \alpha \leq 1$ and suppose that E satisfies a lower q -estimate with constant 1 for some $q \geq \alpha$. Let $\{x_n\}_{n=1}^\infty$ be a bounded sequence in $E(\mathcal{M}, \tau)$ then there exist a subsequence $\{x_{n_k}\}_{k=1}^\infty$ of $\{x_n\}_{n=1}^\infty$, bounded sequences $\{y_k\}_{k=1}^\infty$ and $\{z_k\}_{k=1}^\infty$ in $E(\mathcal{M}, \tau)$ and decreasing projections $e_k \downarrow_n 0$ in \mathcal{M} such that:*

- (i) $x_{n_k} = y_k + z_k$ for all $k \geq 1$;
- (ii) $e_k y_k = 0$ for all $k \geq 1$ and $\lim_{n \rightarrow \infty} \sup_{k \geq 1} \|f_n y_k\|_{E(\mathcal{M}, \tau)} = 0$ for every $f_n \downarrow_n 0$.
- (iii) $\{z_k\}_{k=1}^\infty$ is such that $e_k z_k = z_k$ for all $k \geq 1$.

Definition 3.10. *A subspace X of $L^p(\mathcal{M}, \tau)$ is called strongly embedded into $L^p(\mathcal{M}, \tau)$ if the L^p and the measure topologies on X coincide.*

The following result is a direct application of Proposition 2.9 and Theorem 3.7.

Theorem 3.11. *Let $1 \leq p < \infty$. Every subspace of $L^p(\mathcal{M}, \tau)$ either contains almost isometric copies of ℓ^p or is strongly embedded in $L^p(\mathcal{M}, \tau)$.*

The next corollary should be compared with [18, Theorem 2.4].

Corollary 3.12. *Assume that \mathcal{M} is finite and $p > 2$. Every subspace of $L^p(\mathcal{M}, \tau)$ either contains almost isometric copies of ℓ^p or is isomorphic to a Hilbert space.*

For the commutative case, the space ℓ^p can not be strongly embedded in $L^p[0, 1]$ for $0 < p < 2$. This is due to Kalton [13] for $0 < p < 1$ and Rosenthal [16] for the case $1 \leq p < 2$ (see also [2] for another approach). A non-commutative analogue should be of interest.

Problem: *Let \mathcal{M} be a semifinite von Neumann algebra and $0 < p < 2$. Does ℓ^p strongly embed into $L^p(\mathcal{M}, \tau)$?*

REFERENCES

- [1] C.A. Akemann, *The dual space of an operator algebra*, Trans. Amer. Math. Soc. **126** (1967), 286–302.
- [2] N.L. Carothers and S.I. Dilworth, *Subspaces of $L_{p,q}$* , Proc. Amer. Math. Soc. **104** (1988), 537–545.
- [3] V.I. Chilin and F.A. Sukochev, *Weak convergence in non-commutative symmetric spaces*, J. Op. Th. **31** (1994), 35–65.
- [4] P.G. Dodds, T.K. Dodds, and B. de Pagter, *Non-commutative Banach function spaces*, Math. Zeit. **201** (1989), 583–597.
- [5] ———, *Non-commutative Köthe duality*, Trans. Amer. Math. Soc. **339** (1993), 717–750.
- [6] P.G. Dodds, T.K. Dodds, P. N. Dowling, C. Lennard, and F. A. Sukochev, *A uniform Kadec-Klee property for symmetric operator spaces*, Math. Proc. Camb. Phil. Soc. **118** (1995), 487–502.
- [7] P.G. Dodds, T.K. Dodds, P.N. Dowling, C. Lennard, N. Randrianantoanina, and F.A. Sukochev, *Subspaces of preduals of von Neumann algebras*, In preparation.
- [8] T. Fack and H. Kosaki, *Generalized s -numbers of τ -measurable operators*, Pac. J. Math. **123** (1986), 269–300.

- [9] T. Figiel, N. Ghoussoub, and W. B. Johnson, *On structures of non weakly compact operators on Banach lattices*, Math. Ann. **257** (1981), 317–334.
- [10] W. B. Johnson, B. Maurey, G. Schechtman, and L. Tzafriri, *Symmetric structures in Banach spaces*, Mem. Amer. Math. Soc. **217** (1979).
- [11] M. Kadec and A. Pelczyński, *Bases, lacunary sequences and complemented subspaces of L_p* , Studia Math. **21** (1961/1962), 161–176.
- [12] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras II*, first ed., vol. **2**, Academic Press, 1986.
- [13] N. Kalton, *Linear operators on L_p for $0 < p < 1$* , Trans. Amer. Math. Soc. **259** (1980), 319–355.
- [14] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II*, first ed., Modern Surveys In Math. , vol. **97**, Springer-Verlag, Berlin-Heidelberg-New York, 1979.
- [15] E. Nelson, *Notes on non-commutative integration*, J. Funct. Anal. **15** (1974), 103–116.
- [16] H.P. Rosenthal, *On subspaces of L_p* , Ann. Math. **97** (1973), 344–373.
- [17] H. H. Schaefer, *Banach lattices and positive operators*, first ed., Springer-Verlag, 1974.
- [18] K. A. Sukochev, *Non-isomorphism of L_p -spaces associated with finite and infinite von Neumann algebras*, Proc. Amer. Math. Soc. **124** (1996), 1517–1527.
- [19] M. Takesaki, *Theory of operator algebras I*, Springer-Verlag, New-York, Heidelberg, Berlin, 1979.
- [20] L. W. Weis, *Banach lattices with subsequence splitting property*, Proc. Amer. Math. Soc. **105** (1989), 87–96.
- [21] Q. Xu, *Analytic functions with values in lattices and symmetric spaces of measurable operators*, Math. Proc. Camb. Phil. Soc. **109** (1991), 541–563.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MIAMI UNIVERSITY, OXFORD, OHIO 45056
E-mail address: randrin@muohio.edu