COMPACT RANGE PROPERTY AND OPERATORS ON C*-ALGEBRAS

NARCISSE RANDRIANANTOANINA

ABSTRACT. We prove that a Banach space E has the compact range property (CRP) if and only if for any given C^* -algebra \mathcal{A} , every absolutely summing operator from \mathcal{A} into E is compact. Related results for p-summing operators (0) are also discussed as well as $operators on non-commutative <math>L^1$ -spaces and C^* -summing operators.

1. INTRODUCTION

A Banach space E is said to have the compact range property (CRP) if every E-valued countably additive measure of bounded variation has compact range. It is well known that every Banach space with the Radon-Nikodym property (RNP) has the (CRP) and for dual Banach spaces, the (CRP) were completely characterized as those whose predual do not contain any copies ℓ^1 . For more in depth discussions on Banach spaces with the (CRP), we refer to [9].

The following characterization can be found in [9]: A Banach space E has the (CRP) if and only if every 1-summing operator from C[0, 1] into E is compact. Since C[0, 1] is a (commutative) C^* -algebra, it is a natural question whether C[0, 1] can be replaced by any C^* -algebras. Let us recall that in [7], it was shown that if X is a Banach space that does not contain any copies of ℓ^1 then any 1-summing operators from any given C^* -algebra into X^* is compact; hinting that, as in commutative case, the (CRP) is the right condition to provide compactness. The present note is an improvement of [7]. Our main result confirms that, if \mathcal{A} is a C^* -algebra and E is a Banach space that has the (CRP) then every 1-summing from \mathcal{A} into E is compact. Our proof relies on factorizations of summing operators used in [7] and properties of integral operators.

There is another well kown characterization of spaces with the (CRP) in terms of operators defined on $L^1[0, 1]$: a Banach space E has the (CRP) if and only every operator T from $L^1[0, 1]$ into E is Dunford Pettis (completely continuous) thus the (CRP) is also referred to as the *complete continuity property* (CCP). Unlike the 1-summing operators on C^* -algebras,

¹⁹⁹¹ Mathematics Subject Classification. 46L50,47D15.

Key words and phrases. C^* -algebras, Vector measures.

Supported in part by NSF Grant DMS-9703789.

operators defined on non-commutative L^1 -spaces do not behave the same way as those defined on $L^1[0, 1]$ do. In the last section of this note, we will discuss these operators along with C^* -summing operators studied by Pisier in [6].

Our terminology and notation are standard as may be found in [2] and [4] for Banach spaces, [5] and [8] for C^* -algebras and operator algebras.

2. PRELIMINARIES

In this section, we recall some definitions.

Definition 1. Let X and Y be Banach spaces and $0 . An operator <math>T : X \to Y$ is said to be p-summing if there is a constant C such that for any finite sequence (x_1, x_2, \ldots, x_n) of X, one has

$$\left(\sum_{i=1}^{n} \|Tx_i\|^p\right)^{\frac{1}{p}} \le C \sup\left\{\left(\sum_{i=1}^{n} |\langle x_i, x^* \rangle|^p\right)^{\frac{1}{p}}; \ x^* \in X^*, \|x^*\| \le 1\right\}$$

The smallest constant C for which the above inequality holds is denoted by $\pi_p(T)$ and is called the *p*-summing norm of T.

Definition 2. We say that an operator $T : X \to Y$ is an integral operator if it admits a factorization:



where *i* is the natural inclusion from *Y* into Y^{**} , μ is a probability measure on a compact space *K*, *J* is the natural inclusion and α and β are bounded linear operators.

We define the integral norm $i(T) := \inf \{ \|\alpha\| \cdot \|\beta\| \}$ where the infinum is taken over all such factorizations.

Similarly, we shall say that T is *strictly integral* if T is integral and on the factorization above β takes its values in Y.

It is well known that integral operators are 1-summing but the converse is not true.

If X = C(K) where K is a compact Hausdorff space then it is well known that every 1-summing operator from X into Y is integral.

For more details on the different properties of the classes of operators involved, we refer to [3].

The following simple fact will be needed in the sequel.

Proposition 3. Let $T: X \to Y$ be a strictly integral operator. If Y has the (CRP) then T is compact.

Proof. The operator T has a factorization $T = \beta \circ J \circ \alpha$ where $\alpha : X \to L^{\infty}(\mu), J : L^{\infty}(\mu) \to L^{1}(\mu)$ and $\beta : L^{1}(\mu) \to Y$ are as in the above definition. Note that J is 1-summing so $\beta \circ J : L^{\infty}(\mu) \to Y$ is 1-summing and since $L^{\infty}(\mu)$ is a C(K)-space and Y has the (CRP), $\beta \circ J$ (and hence T) is compact.

We recall that a von Neumann algebra \mathcal{M} is said to be σ -finite if the identity is countably decomposable equivalently if there exist a faithful state $\varphi \in \mathcal{M}_*$. As is customary, for every functional $\varphi \in \mathcal{M}_*$ and $x \in \mathcal{M}, x\varphi$ (resp. φx) denotes the normal functional $y \to \varphi(yx)$ (resp. $y \to \varphi(xy)$).

3. MAIN RESULT

Theorem 4. For a Banach space E, the following are equivalent:

- (1) E has the CRP;
- (2) Every 1-summing operator $T: C[0,1] \to E$ is compact;
- (3) For any given C^* -algebra \mathcal{A} , every 1-summing operator $T: \mathcal{A} \to E$ is compact.

The equivalence $(1) \Leftrightarrow (2)$ is well known, we refer to [4], [9] for more details. Clearly $(3) \Rightarrow$ (2) so what we need to show is $(1) \Rightarrow (3)$. For this, it is enough to consider the following particular case (see [7] for this reduction).

Proposition 5. Let E be a Banach space with the (CRP) and \mathcal{M} be a σ -finite von Neumann algebra. If $T : \mathcal{M} \to E$ is 1-summing and is weak^{*} to weakly continuous then T is compact.

Proof. The proof is a refinement of the argument used in Proposition 3.2 of [7]. We will include most of the details for completeness. Without loss of generality, we can and do assume that E is separable.

Let $\delta > 0$. From Lemma 2.3 of [7],

$$||Tx|| \le 2(1+\delta)\pi_1(T)||xf + fx||_{\mathcal{M}_*} \quad \text{for every } x \in \mathcal{M},$$

where f is a faithful normal state in \mathcal{M}_* . If $L^2(f)$ is completion of the prehilbertian space $(\mathcal{M}, \langle \cdot, \cdot \rangle)$ where $\langle x, y \rangle = f(\frac{xy^* + y^*x}{2})$ then we have the following factorization:



where J is the inclusion map, $\theta(Jx) = \langle \cdot, J(x^*) \rangle$ for every $x \in \mathcal{M}$ and $L(\frac{xf+fx}{2}) = Tx$. We recall that L is a well defined bounded linear map since $\{xf + fx; x \in \mathcal{M}\}$ is dense in \mathcal{M}_* and $\|L(xf + fx)\| \leq 4(1 + \delta)\pi_1(T)\|xf + fx\|_{\mathcal{M}_*}$. Let $S := J^* \circ \theta \circ J$.

Claim: $J \circ L^* : E^* \to L^2(f)$ is compact.

For this, let us consider $L^* : E^* \longrightarrow \mathcal{M}$. Since E is separable, it is isometric to a subspace of C[0, 1]. Let I_E be the isometric embedding of E in C[0, 1] and i be the natural inclusion of C[0, 1] into $C[0, 1]^{**}$.

Define the following map \widetilde{T} from \mathcal{M} into C[0,1] by setting $\widetilde{T} = \overline{I_E \circ T(x^*)}$ for every $x \in \mathcal{M}$. (Here, \overline{f} is the map $t \to \overline{f(t)}$ for $f \in C[0,1]$ with $\overline{f(t)}$ being the conjugate of the complex number f(t)).

Clearly, \widetilde{T} is linear and bounded and it can be shown that \widetilde{T} is 1-summing and is weak^{*} to weakly continuous. In fact, if (x_1, x_2, \ldots, x_n) is a finite sequence in \mathcal{M} then

$$\sum_{i=1}^{n} \|\widetilde{T}x_{i}\| = \sum_{i=1}^{n} \|\overline{I_{E} \circ T(x_{i}^{*})}\|$$
$$= \sum_{i=1}^{n} \|I_{E} \circ T(x_{i}^{*})\|$$
$$= \sum_{i=1}^{n} \|T(x_{i}^{*})\|$$
$$\leq \pi_{1}(T) \sup\left\{\sum_{i=1}^{n} |\langle x_{1}^{*}, \varphi \rangle|, \ \varphi \in \mathcal{M}^{*}, \ \|\varphi\| \leq 1\right\}$$
$$\leq \pi_{1}(T) \sup\left\{\sum_{i=1}^{n} |\langle x_{i}, \varphi^{*} \rangle|, \ \varphi \in \mathcal{M}^{*}, \ \|\varphi\| \leq 1\right\}$$

so \widetilde{T} is 1-summing with $\pi_1(\widetilde{T}) \leq \pi_1(T)$. Moreover if $(x_{\alpha})_{\alpha}$ is a net that converges to zero weak^{*} in \mathcal{M} so does the net $(x_{\alpha}^*)_{\alpha}$ and since T is weak^{*} to weakly continuous, $(T(x_{\alpha}^*))_{\alpha}$ converges to zero weakly in E and hence $(\widetilde{T}(x_{\alpha}))_{\alpha}$ is weakly null which shows that \widetilde{T} is weak^{*} to weakly continuous.

To complete the proof, consider $E^* \xrightarrow{L^*} \mathcal{M} \xrightarrow{i \circ \widetilde{T}} C[0,1]^{**}$.

Since $C[0,1]^{**}$ has the Hahn-Banach extension property and $i \circ \widetilde{T}$ is 1-summing, $i \circ \widetilde{T}$ is an integral operator. Let $K : C[0,1]^* \longrightarrow \mathcal{M}_*$ such that $K^* = i \circ \widetilde{T}$ (such operator exists since $i \circ \widetilde{T}$ is weak^{*} to weakly continuous); K is integral ([3]) and since \mathcal{M}_* is a complemented subspace of its bidual \mathcal{M}^* (see for instance [8]), K is strictly integral and therefore $L \circ K : C[0,1]^* \to E$ is strictly integral and by Proposition 3, $L \circ K$ (and hence $(L \circ K)^* = i \circ \widetilde{T} \circ L^*$) is compact.

Let (U_n) be a bounded sequence in E^* . There exists a subsequence (U_{n_k}) so that $(i \circ \widetilde{T} \circ L^*(U_{n_k}))_k$ is norm convergent in $C[0, 1]^{**}$. Since *i* and I_E are isometries, we get that $(T \circ L^*(U_{n_k}))_k$ is norm convergent so

 $\lim_{k,m} ||T(L^*(U_{n_k})^*) - T(L^*(U_{n_m})^*)|| = 0.$ As in [7], we get

$$\lim_{k,m} \langle T(L^*(U_{n_k})^*) - T(L^*(U_{n_m})^*), U_{n_k} - U_{n_m} \rangle = \lim_{k,m} \|J \circ L^*(U_{n_k} - U_{n_m})\|_{L^2(f)}^2 = 0$$

which proves that $(J \circ L^*(U_{n_k}))_k$ is norm-convergent in $L^2(f)$. The proof is complete. \Box

Theorem 6. Let \mathcal{A} be a C^* -algebra, E be a Banach space and $0 . Every p-summing operator from <math>\mathcal{A}$ into E is compact.

Proof. Let $T : \mathcal{A} \to E$ be an operator with $\pi_p(T) < \infty$. One can choose, by the Pietsch Factorization Theorem, a probability space (Ω, Σ, μ) such that



where S is a subspace of $L^{\infty}(\mu)$, S_p is the closure of S in $L^p(\mu)$ and i_p is the restriction of the natural inclusion j_p .

Denote by S_1 the closure of S in $L^1(\mu)$, by i_1 the restriction of the natural inclusion and $i_{1,p}$ the natural inclusion of S_1 into S_p .

Claim: $\widetilde{T} \circ i_{1,p} : S_1 \to E$ is weakly compact.

To see this, let $(f_n)_n$ be a bounded sequence in $S_1 \subset L^1(\mu)$. By Komlòs's Theorem, there exists a subsequence $(f_{n_k})_k$ and a function $f \in L^1(\mu)$ such that $\lim_{m\to\infty} \frac{1}{m} \sum_{k=1}^m f_{n_k}(\omega) =$

 $f(\omega)$ for a.e. $\omega \in \Omega$. Since 0 ,

$$\lim_{m \to \infty} \|\frac{1}{m} \sum_{k=1}^{m} f_{n_k} - f\|_p = 0.$$

This shows that $f \in S_p$ and $\left(\widetilde{T} \circ i_{1,p} \left(\frac{1}{m} \sum_{k=1}^m f_{n_k}\right)\right)_m$ converges to $\widetilde{T}(f)$ in E and the claim follows.

Using the factorization of weakly compact operator [1], $i_{1,p} \circ \widetilde{T}$ factors through a reflexive space and since $i_1 \circ J$ is 1-summing, the theorem follows from Theorem 4.

4. Concluding remarks

Let us recall some definitions

Definition 7. Let X and Y be Banach spaces. An operator $T : X \to Y$ is called Dunford-Pettis if T sends weakly compact sets into norm compact sets.

The following class of operators was introduced by Pisier in [6] as extension of the *p*-summing operators in the setting of C^* -algebras.

Definition 8. Let \mathcal{A} be a C^* -algebra and E be a Banach space, $0 . An operator <math>T : \mathcal{A} \to E$ is said to be p-C*-summing if there exists a constant C such that for any finite sequence (A_1, \ldots, A_n) of Hermitian elements of \mathcal{A} , one has

$$\left(\sum_{i=1}^{n} \|T(A_i)\|^p\right)^{\frac{1}{p}} \le C \left\| \left(\sum_{i=1}^{n} |A_i|^p\right)^{1/p} \right\|_{\mathcal{A}}.$$

Let \mathcal{M} be a finite von Neumann algebra with a faithful tracial state τ and let J be the canonical inclusion map from \mathcal{M} into $L^1(\mathcal{M}, \tau)$. As in the commutative case, we have the following :

Proposition 9. Let E be a Banach space and $T : L^1(\mathcal{M}, \tau) \to E$ a bounded linear map. Then the following are equivalent:

- (i) T is Dunford-Pettis;
- (ii) $T \circ J$ is compact.

Proof. (i) \implies (ii) is trivial. For the converse, let $(a_n)_n$ be a weakly null sequence in the unit ball of $L^1(\mathcal{M}, \tau)$. It is clear that $(a_n^*)_n$ is also weakly null so without loss of generality, we can

assume that $(a_n)_n$ is a sequence of self-adjoint operators. For each $n \ge 1$, set $a_n = \int_{-\infty}^{\infty} t \, de_t^{(n)}$ the spectral decomposition of a_n and for every $N \ge 1$, let

$$p_{n,N} = \int_{-N}^{N} 1 \ de_t^{(n)}.$$

It is clear that for every $n \ge 1$ and $N \ge 1$,

$$\tau(\mathbf{1} - p_{n,N}) = \tau(\int_{\{|t| > N\}} 1 \ de_t^{(n)}) \le \frac{1}{N} \tau(|a_n|).$$

By the Akeman's characterization of relatively weakly compact subset in $L^1(\mathcal{M}, \tau)$ (see for instance [8] Theorem 5.4 p.149), we conclude that for any given $\epsilon > 0$, there is $N_0 \ge 1$ such that for every $n \ge 1$, $||a_n(\mathbf{1} - p_{n,N_0})|| \le \epsilon$. Moreover $(a_n p_{n,N_0})_n$ is a bounded sequence in \mathcal{M} and since $T \circ J$ is compact, there is a compact subset K_{ϵ} of E such that $\{T(a_n); n \in \mathbb{N}\} \subset$ $K_{\epsilon} + \epsilon B_E$. The proof is complete.

Fix a type II_1 von Neumann algebra \mathcal{M} such that \mathcal{M} contains a complemented copy of a Hilbert space H. The space H is reflexive (and therefore has (CRP)) but the projection map P from $L^1(\mathcal{M}, \tau)$ onto H can not be Dunford-Pettis.

A very well known property of *p*-summing operators is that they are Dunford-Pettis. This in not the case for C^* -summing operators in general. By Proposition 9, $P \circ J$ is not compact. We remark that the argument used in [7] requires only that the operator is C^* -summing and Dunford-Pettis hence since J is clearly C^* -summing and $P \circ J$ is not compact, $P \circ J$ should not be Dunford-Pettis.

Acknowledgments. The author wishes to express his thanks to Patrick Dowling for helpful discussions regarding this note.

References

- W. J. Davis, T. Figiel, W. B. Johnson, and A. Pełczyński, *Factoring weakly compact operators*, J. Funct. Anal. 17 (1974), 311–327.
- J. Diestel, Sequences and series in Banach spaces, first ed., Graduate Text in Mathematics, vol. 92, Springer Verlag, New York, (1984).
- [3] J. Diestel, H. Jarchow, and A. Tonge, Absolutely summing operators, vol. 43, Cambridge University Press, (1995).
- [4] J. Diestel and J.J. Uhl, Jr., Vector measures, Math Surveys, vol. 15, AMS, Providence, RI, (1977).
- [5] R. V. Kadison and J. R. Ringrose, Fundamentals of the theory of operator algebras II, first ed., vol. 2, Academic Press, (1986).

 $\overline{7}$

- [6] G. Pisier, Grothendieck's theorem for non-commutative C*-algebras with appendix on Grothendieck's constants, J. Funct. Anal. 29 (1978), 397–415.
- [7] N. Randrianantoanina, Absolutely summing operators on non-commutative C*-algebras and applications, Houston J. Math. 25 (1999), 745–756.
- [8] M. Takesaki, Theory of operator algebras I, Springer-Verlag, New-York, Heidelberg, Berlin, (1979).
- [9] M. Talagrand, Pettis integral and measure theory, Mem. Amer. Math. Soc. 51 ((1984)), 307.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MIAMI UNIVERSITY, OXFORD, OHIO 45056 *E-mail address*: randrin@muohio.edu