

THE GERBE OF HIGGS BUNDLES

R. DONAGI AND D. GAITSGORY

0. INTRODUCTION

0.1. Let G be a connected reductive group and let X be a scheme. A Higgs bundle over X is a principal G -bundle plus some additional data.

It is especially easy to spell out the definition when $G = GL(n)$. In this case a G -bundle is the same as a vector bundle E over X , and a Higgs structure on it is a subbundle of commutative associative algebras $\mathfrak{c}_X \subset \text{End}_{\mathcal{O}_X}(E)$, which has rank n over X and such that \mathfrak{c}_X is locally generated by one section. In this case the relative spectrum of \mathfrak{c}_X over X is a flat n -sheeted cover of X , called the *spectral cover* corresponding to our Higgs bundle and we will denote it by \overline{X} .

One can ask the following question: how can we classify Higgs bundles with the given spectral cover? The answer is simple: these are in a bijection with line bundles on \overline{X} . Thus, by asking not just for principal G -bundles, but rather for G -bundles endowed with a Higgs structure with a fixed “spectral data”, we go from a *non-abelian* problem to an *abelian* one.

This idea of abelianization has its source in quantum field theory and has been extensively exploited by both physicists and mathematicians. However, perhaps a more common notion is not that of a Higgs bundle, but rather of a K -valued Higgs bundle, where K is a line bundle on X . By definition, a K -valued Higgs bundle for $GL(n)$ is a vector bundle plus a *regular* endomorphism $\varphi : E \rightarrow E \otimes K$, called the Higgs field. In this case, the sheaf \mathfrak{c}_X can be recovered as the centralizer of φ . (When X is a curve and K is the canonical bundle, the corresponding objects (for $GL(n)$ as well as for other reductive groups) have been studied in [14], [3] and [11]. The case of a more general line bundle K over a curve was considered among other works in [1], [2] when $X = \mathbb{P}^1$ and in [4], [10], [15] and [17] for an arbitrary genus. The case of a higher-dimensional X was considered in [18].)

Our philosophy is to think of a K -valued Higgs bundle as involving two separate types of data. One is the spectral data, which amounts to specifying the basis of “eigenvectors” of the Higgs field, i.e. \overline{X} . The other data corresponds to “eigenvalues”, which in our case is a section of $\mathfrak{c}_x \otimes K$. Our point is that the second part of the data is irrelevant for the abelianization process, so we focus on the “eigenvectors” encoded in the abstract Higgs bundle. One obvious advantage of this abstract approach is that it allows the bundle K of “values” to be replaced by various other objects, such as elliptic fibrations. In fact, this elliptic version of the theory has recently found applications to string dualities, cf. [9, 12].

0.2. The obvious question now is how to generalize the above discussion for other reductive groups. It turns out that the notion of an abstract Higgs bundle is quite

easy to generalize. Namely, a Higgs bundle is a pair (E_G, \mathfrak{c}_X) , where E_G is a principal G -bundle over X and \mathfrak{c}_X is a subbundle of the associated bundle of Lie algebras \mathfrak{g}_{E_G} , whose fibers are *regular centralizers*. The precise definition is given in Sect. 2.

The situation is less transparent with spectral covers. In fact, we do not know a good definition of a spectral cover that would work for any G and reproduce for $GL(n)$ the old object.

Instead, we introduce the notion of a *cameral cover*. By definition, the latter is a finite flat map $p : \tilde{X} \rightarrow X$ such that the Weyl group W of G acts on \tilde{X} and certain restrictions on the ramification behaviour are satisfied (cf. Sect. 2). When $G = GL(n)$ this notion is *different* from that of a spectral cover and the relation between the two is studied in detail in Sect. 9 (cf. [7]).

It turns out that every Higgs bundle determines in a canonical way a cameral cover and one can pose the problem of classification of Higgs bundles with a given cameral cover. This is exactly what the present paper is about. Here is a rough outline of the results:

First, we postulate that the above classification problem must be posed in the categorical framework. Namely, one should aim to describe the category $\text{Higgs}_{\tilde{X}}(X)$ of Higgs bundles together with an isomorphism of the induced cameral cover with \tilde{X} . Our first result shows that this classification problem is indeed abelian.

Namely, starting from \tilde{X} we define a sheaf of groups $T_{\tilde{X}}$. We assert in Theorem 4.4 that $\text{Higgs}_{\tilde{X}}(X)$ is a *gerbe* bound by the Picard category of $T_{\tilde{X}}$ -torsors. (The corresponding notions are reviewed for the reader's convenience in Sect. 3.) In particular, it follows that the set of isomorphism classes of Higgs bundles with a given cameral cover carries a simply-transitive action of the group $H^1(X, T_{\tilde{X}})$, which is a generalized Prym variety (cf. [8]).

However, Theorem 4.4 as it is, cannot be completely satisfactory, as it does not describe which $T_{\tilde{X}}$ -gerbe we are getting in this way. Our main result, Theorem 6.4, is the answer to the latter question. It describes the category $\text{Higgs}_{\tilde{X}}(X)$ completely in terms of T -bundles on \tilde{X} (here T is the Cartan group of G), the W -action, and the ramification pattern of \tilde{X} over X .

We should note that the final form of the answer is surprisingly complicated. The main source of troubles comes from the presense in G of non-primitive coroots (cf. [19]). (From the classification of reductive groups we know that this can occur only when G has $SO(2n+1)$ as a direct factor.) In Sect. 8 we will work out explicitly the examples of $G = SL(2)$ and $G = PGL(2)$ which, we hope, will help the reader understand the general structure of the result.

0.3. Although we formulate our results in terms of Higgs bundles on arbitrary schemes, most of our work is the analysis of various varieties attached to the group G . The most important of them are the varieties $\overline{G/N}$ and $\overline{G/T}$, where $\overline{G/T}$ parametrizes *regular centralizers* in the Lie algebra \mathfrak{g} and $\overline{G/T}$ is a certain canonical W -cover of $\overline{G/N}$ (cf. Sect. 1 and Sect. 10).

In particular, over $\overline{G/N}$ we have the tautological group-scheme \mathcal{C} of centralizers. One of the key points of this paper is that this group-scheme can be completely recovered

by looking at the ramification pattern of $\overline{G/T}$ over $\overline{G/N}$. This is our Theorem 11.6, cf. Sect. 11. We emphasize that it is the phenomenon described in Theorem 11.6 which is “responsible” for the abelianization.

0.4. What we said above summarizes the key ideas of the paper. The reader will notice that Sections 1-6 are devoted to the formulation of the main results concerning Higgs bundles and cameral covers, in Sections 7-9 we work out some examples, in Sections 10-13 we prove the basic structure results about $\overline{G/N}$ and, finally, in Sections 14-16 we finish the proofs of Theorem 4.4 and Theorem 6.4.

0.5. **Notation.** We work throughout with a fixed connected reductive group G over \mathbb{C} and we let \mathfrak{g} denote its Lie algebra. We fix a Borel subgroup $B \subset G$ and denote by $\mathcal{F}l$ the flag variety G/B . By definition, $\mathcal{F}l$ classifies Borel subalgebras in \mathfrak{g} .

Let U be the unipotent radical of B and T the Cartan quotient B/U ; we will fix a splitting $T \rightarrow B$. We will denote by \mathfrak{b} and \mathfrak{t} the Lie algebras of B and T , respectively. The rank r of G is by definition the dimension of T .

By N we will denote the normalizer of T and by W the Weyl group N/T .

The set of positive roots will be denoted by Δ^+ . For a root α , let $s_\alpha \in W$ be the corresponding reflection. The set of simple roots we will denote by I . For $i \in I$, we will use the notation s_i instead of s_{α_i} .

1. REGULAR CENTRALIZERS

1.1. Recall that an element $x \in \mathfrak{g}$ is called *regular* if its centralizer $Z_{\mathfrak{g}}(x)$ has the smallest possible dimension, namely r (the rank of \mathfrak{g}). Note that with this definition, a regular element need not be semisimple. The set of all regular elements forms an open subvariety of \mathfrak{g} , which we will denote by \mathfrak{g}_{reg} .

A Lie subalgebra $\mathfrak{a} \subset \mathfrak{g}$ is called a *regular centralizer* if $\mathfrak{a} = Z_{\mathfrak{g}}(x)$ for some $x \in \mathfrak{g}_{reg}$. Note that such \mathfrak{a} is automatically abelian. Our first goal is to introduce a variety which parametrizes all regular centralizers in \mathfrak{g} .

1.2. Let Ab^r be the closed sub-variety in the Grassmannian $Gr_{\mathfrak{g}}^r$ that classifies abelian subalgebras in \mathfrak{g} of dimension r . Let $\Gamma \subset \text{Ab}^r \times \mathfrak{g}$ be the incidence correspondence, i.e. the closed subvariety defined by the condition:

$$(\mathfrak{a}, x) \in \Gamma \text{ if } x \in \mathfrak{a}.$$

Let Γ_{reg} be the intersection $\Gamma \cap (\text{Ab}^r \times \mathfrak{g}_{reg})$.

Proposition 1.3. *There is a smooth morphism $\phi : \mathfrak{g}_{reg} \rightarrow \text{Ab}^r$ whose graph is Γ_{reg} .*

The proof is postponed until Sect. 10

Let $\overline{G/N}$ denote the image of the map ϕ . The above proposition implies that $\overline{G/N}$ is smooth and irreducible. It is clear that \mathbb{C} -points of $\overline{G/N}$ are exactly the regular centralizers in \mathfrak{g} .

By definition, the group G acts on both Ab^r and \mathfrak{g}_{reg} . Therefore, the variety $\overline{G/N}$ acquires a natural G -action and the map ϕ is G -equivariant.

Consider the quotient G/N ; it classifies Cartan subalgebras in \mathfrak{g} . These are the centralizers in \mathfrak{g} of regular semisimple elements. Hence, G/N embeds into $\overline{G/N}$ as

an open subvariety. Obviously, the action of G on G/N by left multiplication is the restriction of its action on $\overline{G/N}$.

1.4. Consider the closed subvariety of $\overline{G/N} \times \mathcal{Fl}$ defined by the condition: for $\mathfrak{a} \in \overline{G/N}$ and $\mathfrak{b}' \in \mathcal{Fl}$,

$$(\mathfrak{a}, \mathfrak{b}') \in \overline{G/T} \text{ if } \mathfrak{a} \subset \mathfrak{b}'.$$

We will denote this variety by $\overline{G/T}$ and the natural projection $\overline{G/T} \rightarrow \overline{G/N}$ by π . It follows from the definitions that we have a natural G -action on $\overline{G/T}$.

The quotient G/T can clearly be identified with the open sub-scheme $\pi^{-1}(G/N)$ of $\overline{G/T}$. We have a natural action of the Weyl group $W = T \backslash N$ on G/T ; this action is free and the quotient can be identified with G/N .

In what follows, by a W -cover of a scheme X we will mean a finite flat scheme $p : \tilde{X} \rightarrow X$, acted on by W such that $p_* \mathcal{O}_{\tilde{X}}$ is locally isomorphic, as a coherent sheaf with a W -action, to $\mathcal{O}_X \otimes \mathbb{C}[W]$. A basic example is $\mathfrak{t} \rightarrow \mathfrak{t}/W$.

The following assertion will be proven in Sect. 10

Proposition 1.5. *The variety $\overline{G/T}$ is smooth and connected. The W -action on G/T extends to the whole of $\overline{G/T}$ and it makes the latter a W -cover of $\overline{G/N}$. Moreover, the two W -covers $\overline{G/T} \rightarrow \overline{G/N}$ and $\mathfrak{t} \rightarrow \mathfrak{t}/W$ are étale-locally isomorphic.*

1.6. Here is an explicit description of $\overline{G/N}$ and $\overline{G/T}$ for $G = SL(2)$. In this case $\overline{G/N}$ is the space of all lines in \mathfrak{g} , i.e. $\overline{G/N} \simeq \mathbb{P}^2$. We have a natural map $\overline{G/T} \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$, where the first projection is the natural map $\overline{G/T} \rightarrow \mathcal{Fl} \simeq \mathbb{P}^1$ and the second projection is a composition of the first one with the action of $-1 \in S_2 \simeq W$ on $\overline{G/T}$.

It is easy to see that this map is an isomorphism. Under the identification, $\pi : \overline{G/T} \rightarrow \overline{G/N}$ is the symmetrization map $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2$.

1.7. **G -orbits.** For each root α , let $D^\alpha \subset \overline{G/T}$ denote the fixed point set of s_α on $\overline{G/T}$. This is a smooth codimension 1 subscheme of $\overline{G/T}$. Indeed, using the étale-local isomorphism between $\overline{G/T} \rightarrow \overline{G/N}$ and $\mathfrak{t} \rightarrow \mathfrak{t}/W$ given in Proposition 1.5, it is enough to prove this statement on \mathfrak{t} . However, \mathfrak{t}^{s_α} is just the corresponding root hyperplane $\mathfrak{t}^\alpha \in \mathfrak{t}$.

Proposition 1.8. *The G -orbits in $\overline{G/T}$ are precisely the locally closed subsets*

$$D^{\Delta'} := \bigcap_{\alpha \in \Delta'} (D^\alpha) \setminus \bigcup_{\beta \notin \Delta'} (D^\beta)$$

where $\Delta' \subset \Delta$ is a subset of the set of roots, closed under linear combinations. The G -orbits in $\overline{G/N}$ are the images of the $D^{\Delta'}$; they are indexed by the Δ' modulo the action of W .

The proof will be given in Sect. 10.

2. HIGGS BUNDLES AND CAMERAL COVERS

2.1. A family of Cartan subalgebras parametrized by a scheme X is given by a map from X to G/N . Equivalently, it is given by a G -equivariant map from the trivial G -bundle over X to G/N . An advantage of this latter description is that there is a

natural way to twist it: given any principal G -bundle E_G over X , we specify a family of Cartan subalgebras *in the adjoint bundle* $\mathfrak{g}_{E_G} := E_G \times_G \mathfrak{g}$ by a G -equivariant map from E_G to the variety G/N . By generalizing this, we define:

Definition 2.2. *A Higgs bundle over a scheme X is a pair (E_G, σ) , where E_G is a principal G -bundle over X and σ is a G -equivariant map $\sigma : E_G \rightarrow \overline{G/N}$.*

Therefore, according to Proposition 1.3, a Higgs structure in a given G -bundle E_G is the same as a vector subbundle \mathfrak{c}_X of \mathfrak{g}_{E_G} of rank r such that $[\mathfrak{c}_X, \mathfrak{c}_X] = 0$ and such that locally in the étale topology \mathfrak{c}_X is the sheaf of centralizers of a section of $E_G \times_G \mathfrak{g}_{reg}$.

The restriction of a Higgs bundle to an open subset $U \subset X$ over which E_G is trivialized can be specified more simply by a map $U \rightarrow \overline{G/N}$. In particular, the *universal Higgs bundle* over $\overline{G/N}$ corresponds to the identity map $\overline{G/N} \rightarrow \overline{G/N}$.

Higgs bundles over X form a category, denoted $\text{Higgs}(X)$:
An element of $\text{Hom}((E_G^1, \sigma^1), (E_G^2, \sigma^2))$ is by definition a G -bundle map $s : E_G^1 \rightarrow E_G^2$ such that $\sigma^2 \circ s = \sigma^1$.

2.3. Now we will introduce our second basic object.

Definition 2.4. *A cameral cover of X is a W -cover $\tilde{X} \rightarrow X$, such that locally in the étale topology, \tilde{X} is a pull-back of the W -cover $\mathfrak{t} \rightarrow \mathfrak{t}/W$.*

As an example, we note that any W -cover is cameral when $G = \text{SL}(2)$, i.e. $W = S_2$. On the other hand, not every $W = S_3$ -cover is cameral: the stabilizer of each point must be a Weyl subgroup of W , so, for example, an A_3 stabilizer is not allowed.

2.5. Cameral covers form a category in a natural way, denoted $\text{Cam}(X)$. By definition, $\text{Hom}(\tilde{X}^1, \tilde{X}^2)$ consists of all W -equivariant isomorphisms $\tilde{X}^1 \rightarrow \tilde{X}^2$.

Proposition 2.6. *There is a natural functor $F : \text{Higgs}(X) \rightarrow \text{Cam}(X)$.*

Proof. Any map $\sigma : E_G \rightarrow \overline{G/N}$ determines a cameral cover \tilde{E}_G of E_G , namely $\overline{G/T} \times_{\overline{G/N}} E_G$. For a Higgs bundle, which involves a G -equivariant map σ , the cover $\tilde{E}_G \rightarrow E_G$ is itself G -equivariant, so by descent theory, it is pulled back from a unique cameral cover $\tilde{X} \rightarrow X$.

Clearly, the assignment $(E_G, \sigma) \mapsto \tilde{X}$ constructed above is functorial. □

Over an open set $U \subset X$ where E_G is trivialized, the restriction $\tilde{U} \rightarrow U$ of the cameral cover is given in terms of σ as: $\overline{G/T} \times_{\overline{G/N}} U$. For example, applying this to the universal Higgs bundle over $\overline{G/N}$ gives the cameral cover $\overline{G/T} \rightarrow \overline{G/N}$. For this reason we refer in this paper to $\overline{G/T} \rightarrow \overline{G/N}$ (rather than $\mathfrak{t} \rightarrow \mathfrak{t}/W$) as the *universal cameral cover*.

2.7. Let us now fix a cameral cover \tilde{X} . Let $\mathrm{Higgs}_{\tilde{X}}(X)$ denote the category-fiber of the above functor $F : \mathrm{Higgs}(X) \rightarrow \mathrm{Cam}(X)$ over \tilde{X} .

In other words, the objects of $\mathrm{Higgs}_{\tilde{X}}(X)$ are pairs

$$((E_G, \sigma) \in \mathrm{Higgs}(X), t : F(E_G, \sigma) \simeq \tilde{X})$$

and $\mathrm{Hom}((E_G^1, \sigma^1, t^1), (E_G^2, \sigma^2, t^2))$ is the set of all bundle maps $s : E_G^1 \rightarrow E_G^2$ with $\sigma^2 \circ s = \sigma^1$ and such that the composition

$$\tilde{X} \xrightarrow{(t^1)^{-1}} F(E_G^1, \sigma^1) \rightarrow F(E_G^2, \sigma^2) \xrightarrow{t^2} \tilde{X}$$

is the identity automorphism of \tilde{X} .

The goal of this paper is to describe explicitly the category $\mathrm{Higgs}_{\tilde{X}}(X)$ in terms of the W -action on \tilde{X} .

3. GERBES

3.1. Since the objects we study have automorphisms, it is difficult to describe them adequately without the use of some categorical language. Specifically, our description requires the notion of an A -gerbe, where A is a sheaf of abelian groups on X . This is a particularly useful case of the more general notion of a gerbe over a sheaf of Picard categories. In this section we review the corresponding definitions. For more details, the reader is referred to [13] or [5]

Let $\mathrm{Sch}_{et}(X)$ denote the big étale site over X . (By definition, $\mathrm{Sch}_{et}(X)$ is the category of all schemes over X and the covering maps are surjective étale morphisms.)

3.2. Recall that a presheaf \mathcal{Q} of categories on $\mathrm{Sch}_{et}(X)$ assigns to every object $U \rightarrow X$ in $\mathrm{Sch}_{et}(X)$ a category $\mathcal{Q}(U)$ and to every morphism $f : U_1 \rightarrow U_2$ in $\mathrm{Sch}_{et}(X)$ a functor $f_{\mathcal{Q}}^* : \mathcal{Q}(U_2) \rightarrow \mathcal{Q}(U_1)$. Moreover, for every composition $U_1 \xrightarrow{f} U_2 \xrightarrow{g} U_3$ there should be a natural transformation $f_{\mathcal{Q}}^* \circ g_{\mathcal{Q}}^* \rightarrow (g \circ f)_{\mathcal{Q}}^*$, such that an obvious compatibility relation for three-fold compositions holds.

A presheaf \mathcal{Q} of categories on $\mathrm{Sch}_{et}(X)$ is said to be a sheaf of categories (or a stack) if the following two axioms hold:

Axiom SC-1. For $U \rightarrow X$ in $\mathrm{Sch}_{et}(X)$ and a pair of objects $C_1, C_2 \in \mathcal{Q}(U)$, the presheaf of sets on $\mathrm{Sch}_{et}(U)$ that assigns to $f : U' \rightarrow U$ the set

$$\mathrm{Hom}_{\mathcal{Q}(U')}(f_{\mathcal{Q}}^*(C_1), f_{\mathcal{Q}}^*(C_2))$$

is a sheaf.

Axiom SC-2. If $f : U' \rightarrow U$ is a covering, then the category $\mathcal{Q}(U)$ is equivalent to the category of descent data on $\mathcal{Q}(U')$ with respect to f (i.e. every descent data on $\mathcal{Q}(U')$ with respect to f is canonically effective, cf. [5], p. 221).

3.3. Here is our main example of a sheaf of categories. Fix a cameral cover $\tilde{X} \rightarrow X$. For every object $U \in \text{Sch}_{et}(X)$ write $\tilde{U} := U \times_X \tilde{X}$, which is a cameral cover of U .

We define the presheaf of categories $\text{Higgs}_{\tilde{X}}$ by $\text{Higgs}_{\tilde{X}}(U) := \text{Higgs}_{\tilde{U}}(U)$ (the functors $\text{Higgs}_{\tilde{X}}(U) \rightarrow \text{Higgs}_{\tilde{X}}(U')$ for $U' \rightarrow U$ and the corresponding natural transformations are defined in a natural way).

The following is an easy exercise in descent theory:

Lemma 3.4. $\text{Higgs}_{\tilde{X}}$ satisfies SC-1 and SC-2.

3.5. Recall that a Picard category is a groupoid endowed with a structure of a tensor category, in which every object is invertible. A basic example (and the source of the name) is the category of line bundles over a scheme.

A sheaf of categories \mathcal{P} is said to be a sheaf of Picard categories if for every $(U \rightarrow X) \in \text{Sch}_{et}(X)$, $\mathcal{P}(U)$ is endowed with a structure of a Picard category such that the pull-back functors f_Q^* are compatible with the tensor structure in an appropriate sense. If \mathcal{P}_1 and \mathcal{P}_2 are two sheaves of Picard categories one defines (in a straightforward fashion) a notion of a tensor functor between them.

A typical and the most important example of a sheaf of Picard categories can be constructed as follows:

Let \mathcal{A} be a sheaf of abelian groups over $\text{Sch}_{et}(X)$. For an object $f : U \rightarrow X$ of $\text{Sch}_{et}(X)$ let $\text{Tors}_{\mathcal{A}}(U)$ denote the category of $\mathcal{A}|_U$ -torsors on U . This is a Picard category and it is easy to see that the assignment $U \rightarrow \text{Tors}_{\mathcal{A}}(U)$ defines a sheaf of Picard categories on $\text{Sch}_{et}(X)$ which we will denote by $\text{Tors}_{\mathcal{A}}$.

3.6. A category \mathcal{Q} is said to be a *gerbe bound by the Picard category* \mathcal{P} , if \mathcal{P} acts on \mathcal{Q} as a tensor category and for any object $C \in \mathcal{Q}$ the functor $\mathcal{P} \rightarrow \mathcal{Q}$ given by

$$P \in \mathcal{P} \implies \text{Action}(P, C) \in \mathcal{Q}$$

is an equivalence.

Now, if \mathcal{P} is a sheaf of Picard categories and \mathcal{Q} is another sheaf of categories we say that \mathcal{Q} is a *gerbe bound by the sheaf of Picard categories* \mathcal{P} , if the following holds:

- For every $(U \rightarrow X) \in \text{Sch}_{et}(X)$, $\mathcal{Q}(U)$ has a structure of a gerbe bound by $\mathcal{P}(U)$. This structure is compatible with the pull-back functors $f_{\mathcal{P}}^*$ and $f_{\mathcal{Q}}^*$.
- There exists a covering $U \rightarrow X$, such that $\mathcal{Q}(U)$ is non-empty.

3.7. The basic example of a gerbe bound by an arbitrary sheaf of Picard categories \mathcal{P} , is \mathcal{P} itself. Here is a less trivial example:

Fix a short exact sequence $0 \rightarrow \mathcal{A} \rightarrow \mathcal{A}'' \rightarrow \mathcal{A}' \rightarrow 0$ of sheaves of abelian groups on X and let $\tau_{\mathcal{A}'}$ be an \mathcal{A}' -torsor over X . We introduce a sheaf of categories \mathcal{Q} as follows. For $U \in \text{Sch}_{et}(X)$, $\mathcal{Q}(U)$ is the category of all “liftings” of $\tau_{\mathcal{A}'}|_U$ to an $\mathcal{A}''|_U$ -torsor. It is easy to check that \mathcal{Q} is a gerbe bound by $\mathcal{P} = \text{Tors}_{\mathcal{A}}$.

In fact, gerbes bound by $\text{Tors}_{\mathcal{A}}$ can be classified cohomologically:

Lemma 3.8. *There is a bijection between the set of equivalence classes of gerbes bound by $\text{Tors}_{\mathcal{A}}$ and $H^2(X, \mathcal{A})$. For a given gerbe \mathcal{Q} the corresponding class in $H^2(X, \mathcal{A})$ vanishes if and only if the category $\mathcal{Q}(X)$ of “global sections” is non-empty.*

In the above example, the class on $H^2(X, \mathcal{A})$ corresponds to the image of the class of $\tau_{\mathcal{A}'}$ under the boundary map $H^1(X, \mathcal{A}') \rightarrow H^2(X, \mathcal{A})$.

4. Higgs $_{\tilde{X}}$ IS A GERBE

4.1. Given a cameral cover $\tilde{X} \rightarrow X$, let $\overline{T}_{\tilde{X}}$ be the sheaf “of W -equivariant maps $\tilde{X} \rightarrow T$ ” on the étale site over X . More precisely, for $U \in \text{Sch}_{\text{ét}}(X)$, $\overline{T}_{\tilde{X}}(U) = \text{Hom}_W(\tilde{U}, T)$, where \tilde{U} is the induced cameral cover of U and the subscript “ W ” means maps respecting the W -action.

However, we need a slightly smaller sheaf.

4.2. Let D_X^α (for each positive root α) be the fixed point scheme of the reflection s_α acting on \tilde{X} . Locally, this is the pullback of the *universal ramification divisor*, i.e. $D^\alpha \subset \overline{G}/\overline{T}$.

Let α be a root of G , considered as a homomorphism $\alpha : T \rightarrow \mathbb{G}_m$. Then any section t of $\overline{T}_{\tilde{X}}$ determines a function $\alpha \circ t$ which goes to its own inverse under the reflection s_α . In particular, its restriction to the ramification locus D_X^α *equals* its inverse, so it equals ± 1 . The subsheaf $T_{\tilde{X}} \subset \overline{T}_{\tilde{X}}$ is defined by the following condition:

$$T_{\tilde{X}}(U) := \{t \in \overline{T}_{\tilde{X}}(U) \mid (\alpha \circ t)|_{D_X^\alpha} = +1 \text{ for each root } \alpha\}. \quad (*)$$

By construction, $\overline{T}_{\tilde{X}}/T_{\tilde{X}}$ is a \mathbb{Z}_2 -torsion sheaf. Note, in addition, that it suffices to impose condition $(*)$ for one representative of each orbit of W on the set of roots.

Remark. Recall that a coroot $\check{\alpha} : \mathbb{G}_m \rightarrow T$ is called primitive if $\ker(\check{\alpha}) = 1$ (this is equivalent to saying that $\check{\alpha}$ is a primitive element of the lattice of cocharacters of T .) It is clear that condition $(*)$ holds automatically for roots whose corresponding coroots are primitive. For example, when the derived group of G is simply-connected, all coroots are primitive, i.e. $(*)$ is automatic and $T_{\tilde{X}} = \overline{T}_{\tilde{X}}$. In fact, G has non-primitive coroots if and only if it contains $SO(2n+1)$ (e.g. $PGL(2) = SO(3)$) as a direct factor, as is easily seen from the classification of Dynkin diagrams.

4.3. Our first result can be stated as:

Theorem 4.4. *Higgs $_{\tilde{X}}$ is a gerbe bound by $\text{Tors}_{T_{\tilde{X}}}$.*

Let us list several corollaries of this theorem:

Corollary 4.5. *To a cameral cover \tilde{X} there corresponds a class in $H^2(X, T_{\tilde{X}})$, which vanishes if and only if \tilde{X} is the cameral cover corresponding to some Higgs bundle.*

This immediate from Lemma 3.8.

Corollary 4.6. *Suppose $\text{Higgs}_{\tilde{X}}(X)$ is non-empty. The set of isomorphism classes of objects in this category carries a simply-transitive action of $H^1(X, T_{\tilde{X}})$. The group of automorphisms of every object is canonically isomorphic to $T_{\tilde{X}}(X)$.*

5. RAMIFICATION

5.1. We now proceed to the formulation of our main result, Theorem 6.4, which describes the category $\text{Higgs}_{\tilde{X}}(X)$ completely in terms of \tilde{X} . For that purpose, we need to introduce some further notation that has to do with the ramification pattern of \tilde{X} over X .

5.2. For each root α we will define a line bundle R_X^α on \tilde{X} . Assume first that \tilde{X} is integral. In this case the subscheme $D_X^\alpha \subset \tilde{X}$ is a Cartier divisor, because locally it is the pull-back of $D^\alpha \subset \overline{G/N}$. We set $R_X^\alpha = \mathcal{O}(D_X^\alpha)$.

When \tilde{X} is arbitrary we proceed as follows. The construction is local, so we may assume that X , and hence also \tilde{X} , is affine. Let I_X^α be a coherent sheaf on \tilde{X} generated by symbols $\{g\}$, for $g \in \mathcal{O}_{\tilde{X}} \mid s_\alpha(g) = -g$ that satisfy the relations:

$$f \cdot \{g\} = \{f \cdot g\} \text{ for all } f \text{ such that } s_\alpha(f) = f.$$

Locally, I_X^α is the pull-back of the sheaf of ideals of the subscheme $D^\alpha \in \overline{G/N}$. Hence, I_X^α is a line bundle. We have a natural map $I_X^\alpha \rightarrow \mathcal{O}_{\tilde{X}}$ that sends $\{g\} \mapsto g$ and, by construction, its cokernel is $\mathcal{O}_{D_X^\alpha}$.

We define the line bundle R_X^α as the inverse of I_X^α . We have a canonical section $\mathcal{O}_{\tilde{X}} \rightarrow R_X^\alpha$ whose set of zeroes is the subscheme D_X^α .

5.3. Consider the T -bundle $\mathcal{R}_X^\alpha := \check{\alpha}(R_X^\alpha)$ (i.e. \mathcal{R}_X^α is induced from R_X^α by means of the homomorphism $\check{\alpha} : \mathbb{G}_m \rightarrow \check{T}$).

For an element $w \in W$ we introduce the T -bundle \mathcal{R}_X^w on \tilde{X} as

$$\mathcal{R}_X^w := \bigotimes_{\alpha} \mathcal{R}_X^\alpha,$$

where α runs over those positive roots for which $w(\alpha)$ is negative. For example, for $w = s_i$ (a simple reflection), $\mathcal{R}_X^{s_i} \simeq \mathcal{R}_X^{\alpha_i}$.

Observe that given a T -bundle \mathcal{L} on \tilde{X} and an element $w \in W$, there are two ways to produce a new T -bundle: we can pull back by w acting as an automorphism of \tilde{X} , or we can conjugate the T -action by w . We will *always* write $w^*(\mathcal{L})$ for the combination of *both* actions. For example, for $G = SL(2)$, the T -bundle \mathcal{L} is equivalent to a line bundle L . The two individual actions on \mathcal{L} of the non-trivial element $-1 \in S_2 = W$ send L to $(-1)^*(L)$ and L^{-1} , respectively, while $(-1)^*(\mathcal{L})$ corresponds to the line bundle $(-1)^*(L^{-1})$. In particular, we have: $w^*(\mathcal{R}_X^\alpha) \simeq \mathcal{R}_X^{w^{-1}(\alpha)}$.

Lemma 5.4. *There is a canonical isomorphism $\mathcal{R}_X^{w_1 \cdot w_2} \xrightarrow{\varpi(w_1, w_2)} w_1^*(\mathcal{R}_X^{w_2}) \otimes \mathcal{R}_X^{w_1}$.*

The proof follows immediately from the definition of \mathcal{R}_X^w . The following proposition is necessary for the formulation of Theorem 6.4.

Proposition 5.5. *Let α_i be a simple root and let $w \in W$ be such that $w(\alpha_i) = \alpha_j$ (another positive simple root). Then, the line bundle $\alpha_i(\mathcal{R}_X^w)|_{D_X^{\alpha_i}}$ admits a canonical trivialization.*

Proof. Let us observe first that, since we are using only roots rather than arbitrary weights, it is sufficient to consider the case when $[G, G]$ is simply-connected.

We have $w \cdot s_i = s_j \cdot w$, hence, by Lemma 5.4

$$s_i^*(\mathcal{R}_X^w) \otimes \mathcal{R}_X^{s_i} \simeq \mathcal{R}_X^{w \cdot s_i} \simeq \mathcal{R}_X^{s_j \cdot w} \simeq w^*(\mathcal{R}_X^{s_j}) \otimes \mathcal{R}_X^w.$$

However, by definition $w^*(\mathcal{R}_X^{s_j}) \simeq \mathcal{R}_X^{s_i}$, so we obtain that $s_i^*(\mathcal{R}_X^w) \simeq \mathcal{R}_X^w$. By restricting to $D_X^{\alpha_i}$, we obtain $\check{\alpha}_i(\alpha_i(\mathcal{R}_X^w)) \simeq \check{\alpha}_i(\mathcal{O}_{D_X^{\alpha_i}})$.

Since $[G, G]$ is simply-connected, every coroot is primitive. Therefore, there exists a weight λ , such that $\lambda \circ \check{\alpha}_i = \text{id} : \mathbb{G}_m \rightarrow \mathbb{G}_m$. By applying λ to the above isomorphism $\check{\alpha}_i(\alpha_i(\mathcal{R}_X^w)) \simeq \check{\alpha}_i(\mathcal{O}_{D_X^{\alpha_i}})$, we obtain an isomorphism $\alpha_i(\mathcal{R}_X^w) \xrightarrow{\text{isom}^\lambda} \mathcal{O}_{D_X^{\alpha_i}}$.

Now it only remains to check that this isomorphism is independent of the choice of λ . However, since the \mathcal{R}_X^w 's are locally pull-backs of the corresponding T -bundles on $\overline{G}/\overline{T}$, it suffices to consider the universal situation, namely the case $X = \overline{G}/\overline{N}$.

In the latter case, the T -bundle $\mathcal{R}^w|_{D^{\alpha_i}}$ itself is trivialized over an open dense part of D^{α_i} , namely over $D^{\alpha_i} - \bigcup_{\alpha \neq \alpha_i} (D^\alpha \cap D^{\alpha_i})$. This is because α_i is not among the set of roots which become negative under the action of w . In particular, we obtain an isomorphism $\alpha_i(\mathcal{R}^w) \xrightarrow{\text{isom}' } \mathcal{O}_{D^{\alpha_i}}$ over $D^{\alpha_i} - \bigcup_{\alpha \neq \alpha_i} (D^\alpha \cap D^{\alpha_i})$.

Moreover, it is easy to see that for any λ as above, the isomorphisms isom^λ and isom' coincide. In particular, isom^λ is independent of λ over $D^{\alpha_i} - \bigcup_{\alpha \neq \alpha_i} (D^\alpha \cap D^{\alpha_i})$ and hence over the whole of D^{α_i} , which is what we need. \square

The following notions will be used in the formulation of Theorem 6.4.

Definition 5.6. Let \mathcal{L}_0 be a T -bundle on \tilde{X} . We say that it is weakly W -equivariant if for every w there exists an isomorphism $w^*(\mathcal{L}_0) \rightarrow \mathcal{L}_0$.

For a weakly W -equivariant T -bundle, let $\text{Aut}(\mathcal{L}_0)$ be the group whose elements are pairs: an element $w \in W$ plus an isomorphism $w^*(\mathcal{L}_0) \rightarrow \mathcal{L}_0$. By definition, $\text{Aut}(\mathcal{L}_0)$ fits into a short exact sequence:

$$1 \rightarrow \text{Hom}(\tilde{X}, T) \rightarrow \text{Aut}(\mathcal{L}_0) \rightarrow W \rightarrow 1.$$

Definition 5.7. A strongly W -equivariant T -bundle is a weakly W -equivariant T -bundle \mathcal{L}_0 plus a choice of a splitting $\gamma_0 : W \rightarrow \text{Aut}(\mathcal{L}_0)$.

Definition 5.8. A T -bundle on \tilde{X} is called weakly R -twisted W -equivariant if for every $w \in W$ there exists an isomorphism $w^*(\mathcal{L}) \otimes \mathcal{R}_X^w \simeq \mathcal{L}$.

For a weakly R -twisted W -equivariant T -bundle \mathcal{L} we introduce the group $\text{Aut}_{\mathcal{R}}(\mathcal{L})$. Its elements are pairs $w \in W$ and an isomorphism $w^*(\mathcal{L}) \otimes \mathcal{R}_X^w \simeq \mathcal{L}$. The group law is defined via the isomorphism $\varpi(w_1, w_2)$ of Lemma 5.4. By definition, $\text{Aut}_{\mathcal{R}}(\mathcal{L})$ is also an extension of W by means of $\text{Hom}(\tilde{X}, T)$.

6. THE MAIN RESULT

6.1. We need one more piece of notation. For a simple root α_i , let M_i be the corresponding minimal Levi subgroup. Under the projection $N \rightarrow W$, the intersection $N \cap [M_i, M_i]$ surjects onto $\langle s_i \rangle \simeq S_2$. Let \mathcal{N}_i denote the preimage of s_i in $N \cap [M_i, M_i]$.

By definition, if n_i and n'_i are two elements in \mathcal{N}_i , there exists $c \in \mathbb{G}_m$ such that $n'_i = \check{\alpha}_i(c) \cdot n_i$.

6.2. For a cameral cover $\tilde{X} \rightarrow X$ we introduce the category $\text{Higgs}'_{\tilde{X}}(X)$ as follows. Its objects consist of

- A weakly R -twisted W -equivariant T -bundle \mathcal{L} on \tilde{X} .
- A map of short exact sequences:

$$\begin{array}{ccccccccc} 1 & \longrightarrow & T & \longrightarrow & N & \longrightarrow & W & \longrightarrow & 1 \\ \downarrow & & \text{natural map} \downarrow & & \gamma \downarrow & & \text{id} \downarrow & & \downarrow \\ 1 & \longrightarrow & \text{Hom}(\tilde{X}, T) & \longrightarrow & \text{Aut}_{\mathcal{R}}(\mathcal{L}) & \longrightarrow & W & \longrightarrow & 1 \end{array}$$

- For each simple root α_i and an element $n_i \in \mathcal{N}_i$, an isomorphism of line bundles on $D_X^{\alpha_i}$

$$\beta_i(n_i) : \alpha_i(\mathcal{L})|_{D_X^{\alpha_i}} \simeq R_X^{\alpha_i}|_{D_X^{\alpha_i}}.$$

These data must satisfy three compatibility conditions:

- (1) If $n'_i = \check{\alpha}_i(c) \cdot n_i$ for $c \in \mathbb{G}_m$, then $\beta_i(n'_i) = c \cdot \beta_i(n_i)$.
- (2) Let α_i be again a simple root and $n_i \in \mathcal{N}_i$. Consider the isomorphism

$$\gamma(n_i) : s_i^*(\mathcal{L}) \otimes \mathcal{R}_X^{s_i} \simeq \mathcal{L}.$$

When we restrict it to $D_X^{\alpha_i}$ it induces an isomorphism

$$\check{\alpha}_i(\alpha_i(\mathcal{L})|_{D_X^{\alpha_i}}) \simeq \check{\alpha}_i(R_X^{\alpha_i}|_{D_X^{\alpha_i}}),$$

by the definition of $\mathcal{R}_X^{s_i}$. We need that this isomorphism coincides with $\check{\alpha}_i(\beta_i(n_i))$.

- (3) Let α_i and α_j be two simple roots and let $w \in W$ be such that $w(\alpha_i) = \alpha_j$. Let $\tilde{w} \in N$ be an element that projects to w , and n_j be an element of \mathcal{N}_j . By pulling back the isomorphism $\beta_j(n_j)$ with respect to w , we obtain an isomorphism $\alpha_i(w^*(\mathcal{L}))|_{D_X^{\alpha_i}} \simeq R_X^{\alpha_i}|_{D_X^{\alpha_i}}$. In addition, the isomorphisms induced by $\gamma(\tilde{w})$ and Proposition 5.5 lead to a sequence of isomorphisms:

$$\alpha_i(\mathcal{L})|_{D_X^{\alpha_i}} \xrightarrow{\gamma(\tilde{w})} \alpha_i(w^*(\mathcal{L}))|_{D_X^{\alpha_i}} \otimes \alpha_i(\mathcal{R}_X^w)|_{D_X^{\alpha_i}} \xrightarrow{\text{Proposition 5.5}} \alpha_i(w^*(\mathcal{L}))|_{D_X^{\alpha_i}}.$$

By composing the two, we obtain an isomorphism $\alpha_i(\mathcal{L})|_{D_X^{\alpha_i}} \simeq R_X^{\alpha_i}|_{D_X^{\alpha_i}}$ and our condition is that it coincides with $\beta_i(n_i)$, where $n_i = \tilde{w}^{-1} \cdot n_j \cdot \tilde{w} \in \mathcal{N}_i$.

This concludes the definition of objects of $\text{Higgs}'_{\tilde{X}}(X)$. Morphisms between $(\mathcal{L}, \gamma, \beta_i)$ and $(\mathcal{L}^1, \gamma^1, \beta_i^1)$ are T -bundle isomorphism maps $\mathcal{L}^1 \rightarrow \mathcal{L}$, which intertwine in the obvious sense γ with γ^1 and β_i with β_i^1 .

6.3. It is easy to see that $\text{Higgs}'_{\tilde{X}}(X)$ can be naturally sheafified. Namely, we define the presheaf of categories $\text{Higgs}'_{\tilde{X}}$ by setting for $U \in \text{Sch}_{et}(X)$, $\text{Higgs}'_{\tilde{X}}(U) := \text{Higgs}'_{\tilde{U}}(U)$. The pull-back functors are defined in an evident manner and it is easy to see that $\text{Higgs}'_{\tilde{X}}$ satisfies SC-1 and SC-2.

Our main result is:

Theorem 6.4. *The sheaves of categories $\text{Higgs}'_{\tilde{X}}$ and $\text{Higgs}_{\tilde{X}}$ are naturally equivalent.*

In particular, we obtain that $\text{Higgs}_{\tilde{X}}(X)$ is equivalent to $\text{Higgs}'_{\tilde{X}}(X)$.

6.5. **Variation.** Assume that all coroots in G are primitive, i.e. for every α , the corresponding 1-parameter subgroup maps injectively into T .

We claim that the definition of $\text{Higgs}'_{\tilde{X}}(X)$ is equivalent to the following (simplified) one. We introduce the category $\text{Higgs}''_{\tilde{X}}(X)$ as follows:

Objects of $\text{Higgs}''_{\tilde{X}}(X)$ are pairs

- A weakly R -twisted W -equivariant T -bundle \mathcal{L} on \tilde{X} .
- A map of short exact sequences:

$$\begin{array}{ccccccccc} 1 & \longrightarrow & T & \longrightarrow & N & \longrightarrow & W & \longrightarrow & 1 \\ \downarrow & & \downarrow \text{natural map} & & \downarrow \gamma & & \downarrow \text{id} & & \downarrow \\ 1 & \longrightarrow & \text{Hom}(\tilde{X}, T) & \longrightarrow & \text{Aut}_{\mathcal{R}}(\mathcal{L}) & \longrightarrow & W & \longrightarrow & 1, \end{array}$$

such that the following condition holds:

(1') Let λ be a weight of T such that $\langle \lambda, \check{\alpha}_i \rangle = 0$, which implies that $\lambda(\mathcal{L})|_{D_X^{\alpha_i}} \simeq \lambda(s_i^*(\mathcal{L}) \otimes \mathcal{R}_X^{s_i})|_{D_X^{\alpha_i}}$. Our condition is that for every $n_i \in \mathcal{N}_i$ the composition

$$\lambda(\mathcal{L})|_{D_X^{\alpha_i}} \simeq \lambda(s_i^*(\mathcal{L}) \otimes \mathcal{R}_X^{s_i})|_{D_X^{\alpha_i}} \xrightarrow{\gamma(n_i)} \lambda(\mathcal{L})|_{D_X^{\alpha_i}}$$

is the identity map.

Morphisms between (\mathcal{L}, γ) and $(\mathcal{L}^1, \gamma^1)$ are T -bundle maps, which intertwine between γ and γ^1 .

Let us show that $\text{Higgs}'_{\tilde{X}}(X)$ and $\text{Higgs}''_{\tilde{X}}(X)$ are naturally equivalent. Indeed, if we have an object $(\mathcal{L}, \gamma, \beta_i) \in \text{Higgs}'_{\tilde{X}}(X)$, the corresponding object of $\text{Higgs}''_{\tilde{X}}(X)$ is obtained by just forgetting the β_i 's.

Conversely, if $(\mathcal{L}, \gamma) \in \text{Higgs}''_{\tilde{X}}(X)$, we reconstruct the β_i 's as follows:

For a simple root α_i and $n_i \in \mathcal{N}_i$ consider the isomorphism $\gamma(n_i)$ restricted to $D_X^{\alpha_i}$. It yields an isomorphism

$$\check{\alpha}_i(\alpha_i(\mathcal{L}))|_{D_X^{\alpha_i}} \simeq \check{\alpha}_i(R_X^{\alpha_i})|_{D_X^{\alpha_i}}.$$

Since $\check{\alpha}_i$ is primitive, there exists a weight λ' with $\langle \lambda', \check{\alpha}_i \rangle = 1$. By evaluating λ on the above isomorphism, we obtain the required identification $\beta_i(n_i) : \alpha_i(\mathcal{L})|_{D_X^{\alpha_i}} \rightarrow R_X^{\alpha_i}|_{D_X^{\alpha_i}}$. This isomorphism does not depend on the choice of λ' because of our condition (1') on γ .

The fact that conditions (1) and (2) hold follows from the construction. Condition (3) follows from the way in which we build the isomorphism of Proposition 5.5.

7. THE UNIVERSAL EXAMPLE

7.1. In the category $\text{Higgs}_{\overline{G/T}}(\overline{G/N})$ there is a canonical tautological object. One of the main steps in the proof of Theorem 6.4 is to exhibit the corresponding canonical object in $\text{Higgs}'_{\overline{G/T}}(\overline{G/N})$. This is our goal in this section.

7.2. Consider the canonical T -bundle $\mathcal{L}_{\mathcal{F}} = G/U$ over $\mathcal{F}l = G/B$ and let us denote by \mathcal{L}_{can} its pull-back to $\overline{G/T}$ under the natural projection $\overline{G/T} \rightarrow \mathcal{F}l$. This will be the first piece in the data $(\mathcal{L}_{can}, \gamma_{can}, \beta_{i,can})$.

When we restrict \mathcal{L}_{can} to $G/T \subset \overline{G/T}$, it becomes identified with $G \rightarrow G/T$. Hence for every element $\tilde{w} \in N$ that projects to $w \in W$, we obtain an isomorphism $\gamma_{can}(n) : w^*(\mathcal{L}_{can}) \simeq \mathcal{L}_{can}$ over G/T , given by right multiplication by \tilde{w}^{-1} on G .

However, when extended to the whole of $\overline{G/T}$, the above identification is meromorphic and the configuration of its zeroes and poles is given by a divisor on $\overline{G/T}$ with values in the cocharacter lattice of T .

Theorem 7.3. *For a simple reflection s_i , the divisor of the above meromorphic map $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ is given by $-\check{\alpha}_i(D^{\alpha_i})$.*

The proof will be given in Sect. 15.

Since $\mathcal{R}_X^{w_1 \cdot w_2} \simeq w_2^*(\mathcal{R}_X^{w_1}) \otimes \mathcal{R}_X^{w_2}$, Theorem 7.3 implies that for any element $w \in W$, the divisor of zeroes/poles of the above meromorphic map $w^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ coincides with $\mathcal{R}_{\overline{G/T}}^w$. Hence, we obtain the data of $\gamma_{can} : N \rightarrow \text{Aut}_R(\mathcal{L}_{can})$.

Finally, we have to specify the data of $\beta_{i,can}$ and check the compatibility conditions. Let us first consider the case when $[G, G]$ is simply connected. As was explained in Sect. 6.5, in this case the data of $\beta_{i,can}$ can be recovered from γ_{can} , once we check that condition (1') holds.

Thus, let α_i be a simple root and let λ be a weight orthogonal to $\check{\alpha}_i$. It suffices to check condition (1') at the generic point of D^{α_i} . Let M_i be the corresponding minimal Levi subgroup. We have a closed embedding $\overline{M_i/T} \subset \overline{G/T}$ (cf. Sect. 10.5) and its orbit under the G -action is the open subset of $\overline{G/T}$ equal to $G/T \cup (D^{\alpha_i} - \bigcup_{\alpha \neq \alpha_i} (D^{\alpha} \cap D^{\alpha_i}))$.

In particular, it contains a dense subset of D^{α_i} .

Since all our constructions were G -equivariant, this implies that condition (1') for α_i is equivalent to the corresponding statement for M_i . Moreover, we can replace M_i by an isogenous group, namely $[M_i, M_i] \times Z(M_i)$. However, in the latter case our compatibility condition becomes obvious, as λ factors through $Z(M_i)$.

Now, let G be arbitrary. Choose an isogeny $G' \rightarrow G$ such that $[G', G']$ is simply-connected. The varieties $\overline{G/T}$ and $\overline{G'/T'}$ are canonically identified and the T -bundle \mathcal{L}_{can} is induced from the T' -bundle \mathcal{L}'_{can} under $T' \rightarrow T$. Therefore, once we know the data of $\beta'_{i,can}$ for \mathcal{L}'_{can} that satisfies the compatibility conditions, it produces the corresponding data for \mathcal{L}_{can} .

Thus, we have constructed a canonical G -equivariant object of $\text{Higgs}'_{\overline{G/T}}$ over $\overline{G/N}$.

8. SOME EXAMPLES

8.1. The unramified situation. We call a Higgs bundle (E_G, σ) unramified if σ maps E_G to G/N . The category of unramified Higgs bundles is therefore equivalent to the category of principal N -bundles.

The functor $F : \text{Higgs}(X) \rightarrow \text{Cam}(X)$ sends an N -bundle E_N to $\tilde{X} := T \setminus E_N$, which is a principal W -bundle over X (i.e. an étale W -cover).

In this case the assertion of Theorem 6.4 is quite evident.

8.2. Now we will analyze the case $G = SL(2)$. Fix an S_2 -cover $p : \tilde{X} \rightarrow X$ and consider the subsheaf of $p_*(\mathcal{O}_{\tilde{X}})$ consisting of S_2 -anti-invariants. We will denote it by \mathfrak{c}_X .

It is easy to see that the category $\text{Higgs}_{\tilde{X}}(X)$ is canonically equivalent to the category of pairs (L, γ') , where L is a line bundle on \tilde{X} and γ' is an isomorphism $\det(p_*(L)) \simeq \mathcal{O}_X$.

Let $D_X \subset \tilde{X}$ be the ramification divisor and let R_X be the corresponding line bundle (cf. Sect. 5). It is easy to see that the category $\text{Higgs}'_{\tilde{X}}(X)$ (which in our case is equivalent to its simplified version $\text{Higgs}''_{\tilde{X}}(X)$) consists of pairs (L, γ) , where L is a line bundle on \tilde{X} and γ is an isomorphism $(-1)^*(L^{-1}) \otimes R_X \simeq L$ such that the composition

$$L \otimes (-1)^*(R_X) \simeq (-1)^*((-1)^*(L^{-1}) \otimes R_X) \xrightarrow{(-1)^*(\gamma)} (-1)^*(L) \xrightarrow{\gamma} L \otimes (-1)^*(R_X)$$

is minus the identity map.

Let us visualize the equivalence $\text{Higgs}_{\tilde{X}}(X) \simeq \text{Higgs}'_{\tilde{X}}(X)$ of Theorem 6.4 in this case. Indeed, for any line bundle L on \tilde{X} we have a canonical S_2 -equivariant isomorphism

$$p^*(\det(p_*(L))) \otimes R_X \simeq L \otimes (-1)^*(L).$$

Therefore, a data of γ' defines the data of γ , and it is easy to see that this sets up an equivalence.

8.3. Let us now consider the case $G = PGL(2)$. In this case the only coroot is non-primitive, so one has to work a little harder.

By definition, objects of $\text{Higgs}''_{\tilde{X}}(X)$ are the following data:

- A line bundle L on \tilde{X} .
- An S_2 -equivariant isomorphism of line bundles $\gamma : L \otimes (-1)^*(L) \simeq R_X^{\otimes 2}$.
- An identification $\beta : L|_{D_X} \rightarrow R_X|_{D_X}$, which is compatible in the obvious sense with the restriction of γ to D_X .

Let us make the statement of Theorem 6.4 explicit in this case too. Starting from an object (E_G, σ, t) in $\text{Higgs}_{\tilde{X}}(X)$ we can locally choose a principal $SL(2)$ -bundle E_G^1 , which induces E_G . Then (E_G^1, σ, t) is an $SL(2)$ -Higgs bundle. Using the above analysis for $SL(2)$, we can attach to it a pair (L^1, γ^1) , where L^1 is a line bundle on \tilde{X} and $\gamma^1 : (-1)^*((L^1)^{-1}) \otimes R_X \simeq L^1$.

The corresponding object of $\text{Higgs}'_{\tilde{X}}(X)$ is constructed as follows. We define the line bundle L as $(L^1)^{\otimes 2}$ and $\gamma := (\gamma^1)^{\otimes 2}$. The data of β comes from the sequence of

isomorphisms

$$(L^1)^{-1} \otimes R_X|_{D^X} \simeq (-1)^*((L^1)^{-1}) \otimes R_X|_{D^X} \stackrel{\gamma^1}{\simeq} L^1|_{D^X}.$$

If we choose a different lifting of E_G to an $SL(2)$ -bundle, the corresponding L^1 will be modified by tensoring with $p^*(L^0)$, where L^0 is a line bundle on X with $(L^0)^{\otimes 2} \simeq \mathcal{O}$, which will not affect the resulting (L, γ, β) .

It is an easy exercise to check that the above construction defines an equivalence of categories.

9. SPECTRAL COVERS VERSUS CAMERAL COVERS FOR $G = GL(n)$

9.1. Observe first that a regular centralizer in $\mathfrak{gl}(n)$ is the same as an n -dimensional associative and commutative subalgebra in $\text{Mat}(n, n)$ generated by one element.

Definition 9.2. *An n -sheeted spectral cover of a scheme X is a finite flat scheme $p : \overline{X} \rightarrow X$ such that $p_*(\mathcal{O}_{\overline{X}})$ has rank n and is locally uni-generated as a sheaf of algebras.*

Thus, a Higgs bundle for $\mathfrak{gl}(n)$ is the same as a rank- n vector bundle E and an n -sheeted spectral cover $\overline{X} \rightarrow X$ with an embedding of bundles of algebras $\mathcal{O}_{\overline{X}} \hookrightarrow \text{End}_{\mathcal{O}_X}(E)$. This is equivalent to saying that E is a line bundle over \overline{X} .

In this section we will analyze the connection of this description of Higgs bundles for $GL(n)$ with the one given by Theorem 6.4. The starting point is the observation that the category of S_n -cameral covers of X is naturally equivalent to the category of n -sheeted spectral covers. Let us describe the functors in both directions:

Given an S_n -cameral cover $\tilde{X} \rightarrow X$, we define the scheme $\widetilde{\overline{X}}$ as $S_{n-1} \backslash \tilde{X}$. Conversely, given an n -sheeted spectral cover $\overline{X} \rightarrow X$ we define $\widetilde{\overline{X}}$ as a scheme that represents the functor that attaches to a scheme S the set of data consisting of

$$(\text{A map } S \rightarrow X \text{ and } n \text{ sections } t_i : S \rightarrow \overline{S} := \overline{X} \times_X S),$$

such that for any function f on \overline{S} the characteristic polynomial of its action on $p_*(\mathcal{O}_{\overline{X}})$ by multiplication equals $\prod_i (Y - f \circ t_i)$, where Y is an indeterminate.

It is easy to see that this functor is indeed representable by a scheme finite over X . The group S_n acts on $\widetilde{\overline{X}}$ by permuting the t_i 's.

Proposition 9.3. *The functors $\tilde{X} \rightarrow \widetilde{\overline{X}}$ and $\overline{X} \rightarrow \widetilde{\overline{X}}$ send cameral covers to spectral covers and spectral covers to cameral covers, respectively. Moreover, they are inverses of one another.*

Proof. Let us consider first the universal situation: $X_0 = \text{Spec}(\mathbb{C}[a_1, \dots, a_n])$, $\tilde{X}_0 = \text{Spec}(\mathbb{C}[x_1, \dots, x_n])$, where the x_i 's satisfy

$$\prod_i (Y - x_i) = Y^n + a_{n-1} \cdot Y^{n-1} + \dots + a_1 \cdot Y + a_0,$$

and $\overline{X}_0 = \text{Spec}(\mathbb{C}[x_1, a_1, \dots, a_n])$, where x_1 satisfies

$$x_1^n + a_{n-1} \cdot x_1^{n-1} + \dots + a_1 \cdot x_1 + a_0.$$

The natural maps $\tilde{X}_0 \rightarrow X_0$ and $\overline{X}_0 \rightarrow X_0$ are a cameral and a spectral cover, respectively and it is easy to see that in this case $\overline{\tilde{X}}_0 \simeq \overline{X}_0$ and $\tilde{\tilde{X}}_0 \simeq \tilde{X}_0$.

This proves the first assertion of the proposition. Indeed, any cameral (resp., spectral) cover is locally induced from \tilde{X}_0 (resp., \overline{X}_0).

For a spectral cover \overline{X} there is a natural map $\overline{\tilde{X}} \rightarrow \overline{X}$, that attaches to a map $S \rightarrow \overline{\tilde{X}}$ given by an n -tuple $\{t_1, \dots, t_n\}$ of maps $t_i : S \rightarrow \overline{S}$ the composition $S \xrightarrow{t_n} \overline{S} \rightarrow \overline{X}$. The resulting map $\overline{\tilde{X}} \rightarrow \overline{X}$ is an isomorphism, because this is so in the universal situation, i.e. for $\tilde{X}_0 \rightarrow X_0$.

Similarly, we have n maps $\tilde{X} \rightarrow \overline{\tilde{X}}$ which correspond to the natural map $S_n/S_{n-1} \times \tilde{X} \rightarrow \overline{\tilde{X}}$. We claim that they define an isomorphism $\tilde{X} \rightarrow \overline{\tilde{X}}$.

Indeed, both the fact that these maps satisfy the condition on the characteristic polynomial and that the resulting map is an isomorphism follow from the corresponding facts for \tilde{X}_0 . □

9.4. Thus, fixing a spectral cover and fixing an S_n -cameral cover amounts to the same thing. Now, Theorem 6.4 implies that the category $\text{Higgs}'_{\tilde{X}}(X)$ is equivalent to the category of line bundles on the corresponding spectral cover \overline{X} . We would like to explain how to see this equivalence explicitly.

We start with the following observation:

Let $\tilde{X} \rightarrow X$ be an S_n -cameral cover and let $\text{Pic}_{\tilde{X},n}(X)$ be the groupoid of S_n -equivariant line bundles L on \tilde{X} for which the following condition holds:

For every reflection $s_{i,j} \in S_n$ the isomorphism

$$s_{i,j}^*(L) \rightarrow L$$

is the identity map on the fixed-point set of $s_{i,j}$ in \tilde{X} .

Proposition 9.5. *The pull-back functor establishes an equivalence between the category of line bundles on X and $\text{Pic}_{\tilde{X},n}(X)$.*

Let us see first how this proposition implies what we need:

The natural map $\tilde{X} \rightarrow \overline{X}$ is itself an S_{n-1} -cameral cover. On the one hand, by applying the above proposition to this map we obtain that the category of line bundles on \overline{X} is equivalent to $\text{Pic}_{\tilde{X},n-1}(\overline{X})$.

On the other hand, we claim that $\text{Pic}_{\tilde{X},n-1}(\overline{X})$ is equivalent to $\text{Higgs}''_{\tilde{X}}(X)$.

Indeed, let us identify the Cartan group of $GL(n)$ with the product of n copies of \mathbb{G}_m and let $\lambda_n : T \rightarrow \mathbb{G}_m$ be the weight corresponding to the last coordinate. Then a functor $\text{Higgs}''_{\tilde{X}}(X) \rightarrow \text{Pic}_{\tilde{X},n-1}(\overline{X})$ is given by $(\mathcal{L}, \gamma) \rightarrow L := \lambda_n(\mathcal{L})$. It is easy to see that this is indeed an equivalence.

9.6. Now let us prove Proposition 9.5. The argument will be a prototype of the one we are going to use to prove Theorem 6.4.

Given an object $L \in \text{Pic}_{\tilde{X},n}(X)$ and a point $x \in X$ we must find an étale neighbourhood of x such that, when restricted to the preimage of this neighbourhood, L becomes isomorphic to the unit object in $\text{Pic}_{\tilde{X},n}(X)$ (i.e. the one for which $L = \mathcal{O}_{\tilde{X}}$ with the tautological S_n -structure).

First, it is easy to reduce the statement to the case when the ramification over x is the maximal possible, i.e. when x has only one geometric preimage \tilde{x} in \tilde{X} . Further, we can assume that X (and therefore also \tilde{X}) is a spectrum of a local ring.

Choose some trivialization of L . Its discrepancy with the S_n -equivariant structure is a 1-cocycle $S_n \rightarrow \mathbb{G}_m$. We must show that this cocycle is homologous to 0.

Let K denote the kernel of the map $\text{Hom}(\tilde{X}, \mathbb{G}_m) \rightarrow \mathbb{G}_m$ given by the evaluation at \tilde{x} . Our condition on L implies that the above cocycle $S_n \rightarrow \mathbb{G}_m$ takes values in K . However, since \tilde{X} is local, K is divisible and torsion-free. Hence $H^1(S_n, K) = 0$, so our cocycle is cohomologically trivial.

10. THE STRUCTURE OF $\overline{G/N}$

10.1. The rest of the paper is devoted to the proofs of various results announced in the previous sections. We start with the proof of Proposition 1.3.

Proof. First we need to show that the map $\phi : \mathfrak{g}_{reg} \rightarrow \text{Ab}^r$ is well-defined, which is equivalent to saying that the projection $\Gamma_{reg} \rightarrow \mathfrak{g}_{reg}$ is an isomorphism.

Since the latter projection is proper and \mathfrak{g}_{reg} is reduced, it is enough to show that the scheme-theoretic preimage in Γ_{reg} of every $x \in \mathfrak{g}_{reg}$ is isomorphic to $\text{Spec}(\mathbb{C})$.

This is clear on the level of \mathbb{C} points, since by definition of regular elements, the only abelian r -dimensional subalgebra in \mathfrak{g} that contains x is its own centralizer.

For $\mathfrak{a} \in \text{Ab}^r$, the tangent space $T_{\mathfrak{a}}(\text{Ab}^r)$ can be identified with the space of maps $T : \mathfrak{a} \rightarrow \mathfrak{g}/\mathfrak{a}$ that satisfy:

$$\forall y_1, y_2 \in \mathfrak{a}, [T(y_1), y_2] + [y_1, T(y_2)] = 0 \in \mathfrak{g}.$$

We claim that the tangent space to $\Gamma_{reg} \cap (\text{Ab}^r \times x)$ at $\mathfrak{a} \times x$ is zero. Indeed, this is the space of maps $T : \mathfrak{a} \rightarrow \mathfrak{g}/\mathfrak{a}$ as above, for which, moreover $[T(y), x] = 0$, $\forall y \in \mathfrak{a}$. However, since $\mathfrak{a} = Z_{\mathfrak{g}}(x)$, any such T is identically zero.

This implies that $T_{\mathfrak{a} \times x}(\Gamma_{reg} \cap \text{Ab}^r \times x) = 0$ which means that $\Gamma_{reg} \cap (\text{Ab}^r \times x)$ is reduced, i.e. $\simeq \text{Spec}(\mathbb{C})$.

Now let us show that ϕ is smooth. Let $\mathfrak{a} \in \text{Ab}^r$ be equal to $\phi(x)$. Using the above description of the tangent space to Ab^r it is easy to see that $d\phi$ sends an element $u \in \mathfrak{g} \simeq T_x(\mathfrak{g}_{reg})$ to the unique map $T : \mathfrak{a} \rightarrow \mathfrak{g}/\mathfrak{a}$ that satisfies:

$$[T(y), x] + [y, u] = 0, \quad \forall y \in \mathfrak{a}.$$

Consider now the map $\mathbf{ev} : T_{\mathfrak{a}}(\text{Ab}^r) \rightarrow \mathfrak{g}/\mathfrak{a}$ given by $T \mapsto T(x)$. The above description of $d\phi$ implies that the composition

$$\mathfrak{g} \simeq T_x(\mathfrak{g}_{reg}) \xrightarrow{d\phi} T_{\mathfrak{a}}(\text{Ab}^r) \xrightarrow{\mathbf{ev}} \mathfrak{g}/\mathfrak{a}$$

coincides with the tautological projection $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{a}$.

However, since x is regular, the fact that $[T(x), y] = -[x, T(y)]$, $\forall y \in \mathfrak{a}$ implies that \mathbf{ev} is an injection. We conclude that \mathbf{ev} is an isomorphism, hence $\text{Im}(\phi)$ is contained in the smooth locus of Ab^r . Furthermore, $d\phi$ is surjective, so ϕ is smooth as claimed. \square

10.2. Let $\tilde{\mathfrak{g}}$ be the closed sub-variety in $\mathfrak{g} \times \mathcal{Fl}$ defined by the condition: $(x, \mathfrak{b}') \in \tilde{\mathfrak{g}}$ if $x \in \mathfrak{b}'$. Let $\tilde{\mathfrak{g}}_{reg}$ denote the intersection $\tilde{\mathfrak{g}} \cap (\mathfrak{g}_{reg} \times \mathcal{Fl})$ and let $\tilde{\pi}$ denote the projection $\tilde{\mathfrak{g}}_{reg} \rightarrow \mathfrak{g}_{reg}$. It is clear that as a variety, $\tilde{\mathfrak{g}}_{reg}$ is smooth and connected, since it is an open subset in a vector bundle over \mathcal{Fl} .

Proposition 10.3. *There exists a natural G -invariant map $\tilde{\phi} : \tilde{\mathfrak{g}}_{reg} \rightarrow \overline{G/T}$, such that the following square is Cartesian:*

$$\begin{array}{ccc} \tilde{\mathfrak{g}}_{reg} & \xrightarrow{\tilde{\phi}} & \overline{G/T} \\ \tilde{\pi} \downarrow & & \pi \downarrow \\ \mathfrak{g}_{reg} & \xrightarrow{\phi} & \overline{G/N} \end{array}$$

Proof. Consider the fibered product $\overline{G/T} \times_{\overline{G/N}} \mathfrak{g}_{reg}$. By definition of $\overline{G/T}$, there is a closed embedding

$$\overline{G/T} \times_{\overline{G/N}} \mathfrak{g}_{reg} \rightarrow \tilde{\mathfrak{g}}_{reg}$$

that sends a triple $(\mathfrak{a} \in \overline{G/N}, \mathfrak{b}' \in \mathcal{Fl}, x \in \mathfrak{g}_{reg}) \in \overline{G/T} \times_{\overline{G/N}} \mathfrak{g}_{reg}$ to $(x, \mathfrak{b}') \in \tilde{\mathfrak{g}}_{reg}$.

We claim that this embedding is in fact an isomorphism. Indeed, the statement is obvious over the preimage in $\tilde{\mathfrak{g}}_{reg}$ of the regular semisimple locus of \mathfrak{g} . Therefore, the two schemes coincide at the generic point of $\tilde{\mathfrak{g}}_{reg}$. This implies what we need, since $\tilde{\mathfrak{g}}_{reg}$ is reduced. \square

Now we are ready to prove Proposition 1.5:

Proof. The map $\tilde{\pi} : \tilde{\mathfrak{g}}_{reg} \rightarrow \overline{G/T}$ is smooth, since it is a base change of a smooth map. Hence, the fact that $\tilde{\mathfrak{g}}_{reg}$ is smooth and connected implies that $\overline{G/T}$ has the same properties.

A well-known theorem of Kostant (cf. [16] or [6], p. 277) says that the restriction of the Chevalley map $\mathfrak{g} \rightarrow \mathfrak{t}/W$ to \mathfrak{g}_{reg} is smooth and that it gives rise to a Cartesian square:

$$\begin{array}{ccc} \tilde{\mathfrak{g}}_{reg} & \longrightarrow & \mathfrak{t} \\ \downarrow & & \downarrow \\ \mathfrak{g}_{reg} & \longrightarrow & \mathfrak{t}/W \end{array}$$

Therefore, the natural action of W on the preimage in $\tilde{\mathfrak{g}}$ of the regular semisimple locus in \mathfrak{g} extends to the whole of $\tilde{\mathfrak{g}}_{reg}$. The same is true for $\overline{G/T}$, because the map

$\tilde{\phi}$ is flat and surjective. The étale local isomorphism follows from comparison of our Cartesian square with that of Proposition 10.3. \square

10.4. Now let us prove Proposition 1.8.

Proof. Let Δ' be as in the formulation of the proposition. Consider an element $t \in \mathfrak{t}$ such that $\alpha(t) = 0$ for $\alpha \in \Delta'$ and $\beta(t) \neq 0$ for $\beta \notin \Delta'$.

In this case $\mathfrak{m} := Z_{\mathfrak{g}}(t)$ is a Levi subalgebra of \mathfrak{g} and let M be the corresponding Levi subgroup. It is well-known that $\mathfrak{m} \cap \mathfrak{b}$ is a Borel subalgebra in \mathfrak{m} . Let u be an element in the unipotent radical of $\mathfrak{m} \cap \mathfrak{b}$, which is regular with respect to M .

We then see that $x = t + u$ is a regular element in \mathfrak{g} , since $Z_{\mathfrak{g}}(x) = Z_{\mathfrak{m}}(u)$. It is known that if a Borel subalgebra contains a regular element, then it also contains its centralizer (cf. Lemma 11.5). Therefore, $(Z_{\mathfrak{m}}(u), \mathfrak{b}) \in \overline{G/T}$. Moreover, it is easy to see that every pair $(\mathfrak{a}, \mathfrak{b}') \in \overline{G/T}$ is G -conjugate to one of the above form.

To conclude the proof, it remains to show that $(Z_{\mathfrak{m}}(u), \mathfrak{b}) \in \bigcap_{\alpha \in \Delta'} (D^\alpha) \setminus \bigcup_{\beta \notin \Delta'} (D^\beta)$.

For that, it suffices to show that the image of $(t + u, \mathfrak{b})$ as above under $\tilde{\mathfrak{g}}_{reg} \rightarrow \mathfrak{t}$ belongs to the corresponding locus of \mathfrak{t} . However, the above image is just t , which makes the assertion obvious. \square

10.5. **Levi subgroups.** Let $J \subset I$ be a subset. It defines a root subsystem Δ_J and let M_J (resp., $P_J \subset G$, $W_J \subset W$) denote the corresponding standard Levi subgroup (resp., standard parabolic, Weyl subgroup). Let N_{M_J} be the intersection $M_J \cap N$, which is the normalizer of T in M_J .

It is easy to see that the natural map $M_J/N_{M_J} \rightarrow G/N$ extends to a map $i_J : \overline{M_J/N_{M_J}} \rightarrow \overline{G/N}$. In fact, $\overline{M_J/N_{M_J}}$ is a closed sub-variety of $\overline{G/N}$ which corresponds to $\{\mathfrak{a} \in \overline{G/N} \mid \mathfrak{a} \subset \mathfrak{m}_J\}$.

Proposition 10.6. *There is a canonical W -equivariant isomorphism:*

$$\tilde{i}_J : W \times^{W_J} \overline{M_J/T} \simeq \overline{M_J/N_{M_J}} \times_{\overline{G/N}} \overline{G/T}.$$

Proof. First, we have a natural closed embedding

$$\overline{M_J/T} \rightarrow \overline{M_J/N_{M_J}} \times_{\overline{G/N}} \overline{G/T} \subset \overline{G/T}.$$

Its image consists of pairs $(\mathfrak{a}, \mathfrak{b}') \in \overline{G/T}$ such that $\mathfrak{a} \subset \mathfrak{m}_J$ and $\mathfrak{b}' \subset \mathfrak{p}_J := \text{Lie}(P_J)$.

This map is compatible with the W_J -action. Hence, it extends to a finite map

$$\tilde{i}_J : W \times^{W_J} \overline{M_J/T} \rightarrow \overline{M_J/N_{M_J}} \times_{\overline{G/N}} \overline{G/T}.$$

Since both varieties are smooth, in order to prove that \tilde{i}_J is an isomorphism, it suffices to do so over the open part, i.e. over M_J/N_{M_J} . However, in the latter case, the assertion becomes obvious. \square

It is easy to see that the G -orbit of $\overline{M_J/N_{M_J}} \times_{\overline{G/N}} \overline{G/T} \subset \overline{G/T}$ (resp., $\overline{M_J/T} \subset \overline{G/T}$) is the union of those $D^{\Delta'}$ for which Δ' is W -conjugate to a subset of Δ_J (resp., $\Delta' \subset \Delta_J$.)

11. THE GROUP-SCHEME OF CENTRALIZERS

11.1. In this section we will formulate two basic theorems, Theorem 11.6 and Theorem 11.8, which will be used for the proof of our main result.

Consider the constant group-scheme $G \times \overline{G/N}$ over $\overline{G/N}$ and let $\mathcal{C} \subset G \times \overline{G/N}$ be its closed group-subscheme of “centralizers”. In other words, \mathcal{C} is defined by the condition that $(g \in G, \mathfrak{a} \in \overline{G/N}) \in \mathcal{C}$ if g commutes with \mathfrak{a} . Clearly, \mathcal{C} is equivariant with respect to the G -action on $\overline{G/N}$.

Note that the corresponding sheaf \mathfrak{c} of Lie algebras can be identified with the tautological rank r vector bundle over $\overline{G/N}$ which comes from the embedding $\overline{G/N} \subset Gr_{\mathfrak{g}}^r$.

Proposition 11.2. *The group-scheme \mathcal{C} is commutative and smooth over $\overline{G/N}$ and is irreducible as a variety.*

Proof. Let \mathcal{C}' be the group-subscheme of $G \times \mathfrak{g}_{reg}$ over \mathfrak{g}_{reg} defined by the condition:

$$\mathcal{C}' := \{(g, x) \in G \times \mathfrak{g}_{reg} \mid \text{Ad}_g(x) = x\}.$$

First, let us show that \mathcal{C}' is commutative and smooth over \mathfrak{g}_{reg} .

Let (g, x) be a \mathbb{C} -point of \mathcal{C}' . The tangent space to \mathcal{C}' at (g, x) consists of pairs $(\xi, y) \in \mathfrak{g} \times \mathfrak{g}$ such that $[x, \xi] = \text{Ad}_g(y) - y$. The differential of the map $\mathcal{C}' \rightarrow \mathfrak{g}_{reg}$ sends (ξ, y) to y . We claim that it is surjective.

It is known that if G is of adjoint type, then the centralizer of every regular element is connected (in particular, each $Z_G(x)$ is commutative). Therefore, $\text{Span}_{g \in Z_G(x)} (\text{Ad}_g(y) - y) =$

$\text{Im}(\text{ad}_{Z_G(x)})$. However, the latter, as we saw in the proof of Proposition 1.3, coincides with $\text{Im}(\text{ad}_x)$, since x is regular.

To prove that \mathcal{C}' is smooth over \mathfrak{g}_{reg} , it remains to observe that the fibers of \mathcal{C}' are smooth (since they are algebraic groups in char.0) and all have dimension r , by definition of \mathfrak{g}_{reg} . The fact that \mathcal{C}' is commutative was established in the course of the above argument.

Now let us prove the assertion for \mathcal{C} . We have a natural closed embedding $\mathcal{C} \times_{\overline{G/N}} \mathfrak{g}_{reg} \rightarrow \mathcal{C}'$, which is an isomorphism over the regular semisimple locus of \mathfrak{g}_{reg} . Hence, it is an isomorphism, because \mathcal{C}' is reduced. Therefore, since the map $\phi : \mathfrak{g}_{reg} \rightarrow \overline{G/N}$ is flat and surjective, this shows that \mathcal{C} is commutative and smooth over $\overline{G/N}$. It is irreducible, because this is obviously true over G/N . \square

11.3. Now we will introduce another group-scheme over $\overline{G/N}$, seemingly of a different nature.

Consider the contravariant functor Schemes \Rightarrow Sets which assigns to a scheme S the set of pairs

(A map $S \rightarrow \overline{G/N}$, a W -equivariant map $\tilde{S} := S \times_{\overline{G/N}} \overline{G/T} \rightarrow T$).

It is easy to see that this functor is representable by an abelian group-scheme over $\overline{G/N}$, which we will denote by $\overline{\mathbb{T}}$. Therefore, once $S \rightarrow \overline{G/N}$ is fixed, $\mathrm{Hom}_{\overline{G/N}}(S, \overline{\mathbb{T}}) \simeq \Gamma(S, \overline{T}_{\tilde{S}})$. Clearly, the G -action on $\overline{G/T}$ gives rise to a G -action on $\overline{\mathbb{T}}$.

We define the open group-subscheme \mathbb{T} of $\overline{\mathbb{T}}$ by the following condition (**):

$\mathrm{Hom}(S, \mathbb{T})$ consists of those pairs $(S \rightarrow \overline{G/N}, \tilde{S} \rightarrow T)$ as above, for which for every root α the composition

$$S \times_{\overline{G/N}} D^\alpha \hookrightarrow \tilde{S} \rightarrow T \xrightarrow{\alpha} \mathbb{G}_m$$

avoids $-1 \in \mathbb{G}_m$.

Since for any map $S \rightarrow \mathbb{T}$, the above composition takes values in $\pm 1 \subset \mathbb{G}_m$, condition (**) is equivalent to condition (*) in the definition of the sheaf $T_{\tilde{S}}$ (cf. Sect. 4.2). In other words, for a fixed map $S \rightarrow \overline{G/N}$, $\mathrm{Hom}_{\overline{G/N}}(S, \mathbb{T}) \simeq \Gamma(S, T_{\tilde{S}})$.

Remark. As we will see later, \mathbb{T} is not only open in $\overline{\mathbb{T}}$, but is in fact a connected component there.

11.4. A remarkable fact is that the group-schemes \mathcal{C} and \mathbb{T} are canonically isomorphic. Here we will construct a map between them in one direction.

Let \mathcal{B} denote the universal group-scheme of Borel subgroups over $\mathcal{F}\ell$. Let us denote by $\tilde{\mathcal{B}}$ its pull-back to $\overline{G/T}$. In addition, let us denote by $\tilde{\mathcal{C}}$ the pull-back of \mathcal{C} to $\overline{G/T}$.

Both $\tilde{\mathcal{B}}$ and $\tilde{\mathcal{C}}$ are group-subschemas of the constant group-scheme $G \times \overline{G/T}$.

Lemma 11.5. *$\tilde{\mathcal{C}}$ is a closed group-subscheme of $\tilde{\mathcal{B}}$.*

Indeed, since $\tilde{\mathcal{C}}$ is reduced and irreducible, it suffices to check that over G/N , $\tilde{\mathcal{C}}$ is contained in $\tilde{\mathcal{B}}$. However, this is obvious.

We have a natural projection $\mathcal{B} \rightarrow T \times \mathcal{F}\ell$. By composing it with the inclusion of Lemma 11.5, we obtain a map

$$\mathcal{C} \times_{\overline{G/N}} \overline{G/T} \rightarrow T.$$

This map respects the group law on \mathcal{C} and T and commutes with the W -action. (This is because it suffices to check both facts after the restriction to G/N , where they become obvious.)

Hence, we obtain a homomorphism of group-schemes $\mathcal{C} \rightarrow \overline{\mathbb{T}}$.

Theorem 11.6. *The above map $\mathcal{C} \rightarrow \overline{\mathbb{T}}$ defines an isomorphism $\chi : \mathcal{C} \rightarrow \mathbb{T}$.*

The proof will be given in the next section.

11.7. Now we will formulate the second key result which will be used in the proof of Theorem 4.4.

Consider the functor that assigns to a scheme S the set of triples $(\overline{G/N}_S^1, \overline{G/N}_S^2, \nu)$, where $\overline{G/N}_S^1$ and $\overline{G/N}_S^2$ are two S -points of $\overline{G/N}$ and ν is a W -equivariant isomorphism

$$\nu : \tilde{S}^1 \rightarrow \tilde{S}^2,$$

where \tilde{S}^i is the W -cover of S induced by $\overline{G/N}_S^i$ from $\pi : \overline{G/T} \rightarrow \overline{G/N}$.

It is easy to see that this functor is representable. Let \mathcal{H} denote the representing scheme. Since the W -cover $\overline{G/T} \rightarrow \overline{G/N}$ is G -equivariant, we obtain a natural map $\xi : G \times \overline{G/N} \rightarrow \mathcal{H}$ which covers the map $G \times \overline{G/N} \xrightarrow{\text{Action} \times \text{id}} \overline{G/N} \times \overline{G/N}$.

Theorem 11.8. *The above map $\xi : G \times \overline{G/N} \rightarrow \mathcal{H}$ is smooth and surjective.*

This theorem will be proven in Sect. 13.

11.9. The scheme \mathcal{H} lives over $\overline{G/N} \times \overline{G/N}$. Let \mathcal{H}_Δ denote its restriction to the diagonal. By definition, \mathcal{H}_Δ is a group-scheme over $\overline{G/N}$ which represents the functor of W -equivariant automorphisms of $\overline{G/T}$ over $\overline{G/N}$.

Let $\text{St} \subset G \times \overline{G/N}$ be the closed group-subscheme of stabilizers, i.e.

$$(g, \mathfrak{a}) \in \text{St} \text{ if } \text{Ad}_g(\mathfrak{a}) = \mathfrak{a}.$$

Obviously, \mathcal{C} is a closed normal group-subscheme of St .

The map $\xi : G \times \overline{G/N} \rightarrow \mathcal{H}$ gives rise to a map $\xi_\Delta : \text{St} \rightarrow \mathcal{H}_\Delta$.

Proposition 11.10. *\mathcal{H}_Δ represents the quotient group-scheme St/\mathcal{C} .*

Proof. Theorem 11.8 implies that the map $\xi_\Delta : \text{St} \rightarrow \mathcal{H}_\Delta$ is smooth and surjective. Therefore, all we have to show is that if $S \rightarrow \text{St}$ is a map such that the induced automorphism of \tilde{S} is trivial, then S maps to \mathcal{C} .

Observe that \mathcal{H}_Δ acts on \mathbb{T} via its action on $\overline{G/T}$. Since the isomorphism $\chi : \mathcal{C} \rightarrow \mathbb{T}$ is G -equivariant, we obtain a commutative diagram of actions:

$$\begin{array}{ccc} \text{St} \times_{\overline{G/N}} \mathcal{C} & \longrightarrow & \mathcal{C} \\ \xi_\Delta \times \chi \downarrow & & \chi \downarrow \\ \mathcal{H}_\Delta \times_{\overline{G/N}} \mathbb{T} & \longrightarrow & \mathbb{T}, \end{array}$$

where the top horizontal arrow is the adjoint action.

Therefore, if a map $S \rightarrow \text{St}$ induces the trivial automorphism of \tilde{S} , its adjoint action on \mathcal{C} is trivial too. But this means that it factors through \mathcal{C} . □

Similarly, one shows:

Corollary 11.11. *The scheme \mathcal{H} represents the quotient group-scheme $(G \times \overline{G/N})/\mathcal{C}$.*

12. PROOF OF THEOREM 11.6

12.1. We start by establishing a result on compatibility of our objects with restrictions to Levi subgroups. We then verify the Lie-algebraic version of the theorem by restricting to an $\mathfrak{sl}(2)$ subalgebra, and finally we refine this to prove the desired group-theoretic version.

12.2. Let $M = M_J$ be a standard Levi subgroup of G (cf. Sect. 10.5) and let \mathcal{C}_M be the corresponding sheaf of centralizers over $\overline{M/N_M}$.

On the one hand, there is a natural closed embedding

$$\mathcal{C}_M \hookrightarrow i_J^*(\mathcal{C}) := \overline{M/N_M} \times_{\overline{G/N}} \mathcal{C}.$$

On the other hand, we have the group scheme $\overline{\mathbb{T}}$ over $\overline{G/N}$, as well as the group-scheme $\overline{\mathbb{T}}_M$ over $\overline{M/N_M}$. This time, by Proposition 10.6, we have a canonical isomorphism

$$\overline{\mathbb{T}}_M \simeq i_J^*(\overline{\mathbb{T}}) := \overline{M/N_M} \times_{\overline{G/N}} \overline{\mathbb{T}}.$$

Moreover, it induces an isomorphism $\mathbb{T}_M \simeq i_J^*(\mathbb{T})$, since if a root α is not W -conjugate to a root in M , then s_α has no fixed points on $W \times \overline{M/T}^{W_J}$.

Proposition 12.3. *The map $\mathcal{C}_M \rightarrow i_M^*(\mathcal{C})$ is an isomorphism. Moreover, the diagram*

$$\begin{array}{ccc} \mathcal{C}_M & \xrightarrow{\chi_M} & \overline{\mathbb{T}}_M \\ \downarrow & & \downarrow \\ i_J^*(\mathcal{C}) & \xrightarrow{i_J^*(\chi)} & i_J^*(\overline{\mathbb{T}}) \end{array}$$

is commutative.

Proof. The map $\mathcal{C}_M \hookrightarrow i_M^*(\mathcal{C})$ is an isomorphism because it is a closed embedding and at the same time an isomorphism over the generic point of $\overline{M/N_M}$. Commutativity of the diagram can be checked over the preimage of $\overline{M/N_M}$, in which case it becomes obvious. □

12.4. We will now prove the assertion of Theorem 11.6 on the Lie-algebra level.

Let \mathfrak{t} denote the sheaf of Lie algebras corresponding to $\overline{\mathbb{T}}$. Obviously, it is isomorphic to $\text{Lie}(\mathbb{T})$ as well. By definition, we have: $\mathfrak{t} \simeq (\mathfrak{t} \otimes \pi_*(\mathcal{O}_{\overline{G/T}}))^W$. Since $\pi_*(\mathcal{O}_{\overline{G/T}})$ is locally isomorphic to $\mathcal{O}_{\overline{G/N}} \otimes \mathbb{C}[W]$, \mathfrak{t} is a vector bundle of rank r over $\overline{G/N}$.

Proposition 12.5. *The map*

$$d\chi : \mathfrak{c} \rightarrow \mathfrak{t}$$

is an isomorphism.

Proof. The proof will consist of two steps. The first step will be a reduction to the case of $SL(2)$ and the second one will be a proof of the assertion for $SL(2)$.

Step 1.

Both \mathfrak{c} and \mathfrak{t} are vector bundles of rank r over $\overline{G/N}$ and the map $d\chi$ is clearly an isomorphism over G/N . Since the variety $\overline{G/N}$ is smooth, it remains to show that $d\chi$ is an isomorphism on an open subset of $\overline{G/N}$ whose complement has codimension at least 2.

It follows from Sect. 10.5 that such an open subset is formed by the union of the G -orbits of the images of $i_J(\overline{M_J/N_{M_J}})$, where $J = \{\alpha_j\}$ for all simple roots α_j .

Therefore, by G -equivariance and by Proposition 12.3, it suffices to show that the map

$$d\chi_{M_J} : \mathfrak{c}_{M_J} \rightarrow \mathfrak{t}_{M_J}$$

is an isomorphism. This reduces us to the case when G is a reductive group of semi-simple rank 1.

Moreover, the statement is clearly invariant under isogenies, so we may replace G by $Z^0(G) \times [G, G]$. Clearly, the assertion in such a case is equivalent to the one for $[G, G]$, which in turn can be replaced by $SL(2)$.

Step 2.

For $G = SL(2)$, the variety $\overline{G/N}$ can be identified with \mathbb{P}^2 in such a way that the sheaf \mathfrak{c} goes over to $\mathcal{O}(-1)$. Moreover, $\overline{G/T} \rightarrow \overline{G/N}$ can be identified with the S_2 -cover $\pi : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^2$.

To prove the assertion, it is enough to show that \mathfrak{t} has degree -1 , since any non-zero map between two line bundles of the same degree is automatically an isomorphism.

By definition, \mathfrak{t} is the $\mathcal{O}(\mathbb{P}^2)$ -module of anti-invariants of S_2 in $\pi_*(\mathcal{O}(\mathbb{P}^1 \times \mathbb{P}^1))$. Therefore,

$$\mathfrak{t} \simeq \det(\pi_*(\mathcal{O}(\mathbb{P}^1 \times \mathbb{P}^1))) \simeq \mathcal{O}(-1).$$

□

12.6. Now we will check that the map χ induces an isomorphism between \mathbb{C} -points of \mathcal{C} and \mathbb{T} . Evidently, this assertion, combined with Proposition 12.5 and Proposition 11.2, implies Theorem 11.6.

Let $\mathfrak{a} \in \overline{G/N}$ be the centralizer of a regular element $x \in \mathfrak{g}$. As we saw in the proof of Proposition 11.2, on the one hand, the fiber of \mathcal{C} at $\mathfrak{a} = Z_{\mathfrak{g}}(x)$ can be identified with $Z_G(x)$. On the other hand, the fiber of \mathbb{T} at \mathfrak{a} can be identified with $\text{Hom}_W(\mathcal{F}l^x, T)^{**}$, where $\mathcal{F}l^x$ is the fixed point scheme of the vector field induced by x on $\mathcal{F}l$ and the super-script $**$ corresponds to the $(**)$ condition in the definition of \mathbb{T} .

Let $x = x^{ss} + x^{nil}$ be the Jordan decomposition of x . We can assume that $Z_{\mathfrak{g}}(x^{ss})$ is a standard Levi subalgebra \mathfrak{m} and x^{nil} is a regular nilpotent element in \mathfrak{m} . Using Proposition 12.3, we can replace G by M and hence we can assume that x^{ss} is a central element in \mathfrak{g} .

There are natural embeddings $Z(G) \times \overline{G/N} \rightarrow \mathcal{C}$ and $Z(G) \times \overline{G/N} \rightarrow \mathbb{T}$, which make the diagram

$$\begin{array}{ccc} Z(G) \times \overline{G/N} & \longrightarrow & \mathcal{C} \\ \text{id} \downarrow & & \chi \downarrow \\ Z(G) \times \overline{G/N} & \longrightarrow & \mathbb{T} \end{array}$$

commute.

Proposition 12.7. *Let x be a regular nilpotent element and let*

$$Z_G(x) = Z_G(x)^{ss} \times Z_G(x)^{nil}$$

$$\mathrm{Hom}_W(\mathcal{F}l^x, T)^{**} = \mathrm{Hom}_W(\mathcal{F}l^x, T)^{ss,**} \times \mathrm{Hom}_W(\mathcal{F}l^x, T)^{nil,**}$$

be the Jordan decompositions of the fibers of \mathcal{C} and \mathbb{T} at $Z_{\mathfrak{g}}(x)$. Then the embedding of $Z(G)$ induces isomorphisms:

$$Z(G) \simeq Z_G(x)^{ss} \text{ and } Z(G) \simeq \mathrm{Hom}_W(\mathcal{F}l^x, T)^{ss,**}.$$

It is clear, first of all, that this proposition implies the theorem. Indeed, it is enough to show that χ induces an isomorphism $Z_G(x)^{nil} \rightarrow \mathrm{Mor}_W(\mathcal{F}l^x, T)^{nil,**}$. But since these groups are unipotent, our assertion follows from the corresponding assertion on the Lie-algebra level, which has been proven before.

Proof. The fact that $Z(G) \simeq Z_G(x)^{ss}$ is an immediate consequence of the fact that in a group of adjoint type centralizers of regular elements are connected.

To prove that $Z(G) \simeq \mathrm{Hom}_W(\mathcal{F}l^x, T)^{ss,**}$, let us observe that if x is a regular nilpotent element, $\mathcal{F}l^x$ is a local non-reduced scheme. Its closed point, viewed as a point of $\overline{G/T}$, belongs to the intersection of all the D^α 's.

Let $\mathrm{Hom}_W(\mathcal{F}l^x, T)_1$ be the sub-group of $\mathrm{Hom}_W(\mathcal{F}l^x, T)$ which corresponds to maps $\mathcal{F}l^x \rightarrow T$ that send the closed point of $\mathcal{F}l^x$ to the identity in T . Clearly, $\mathrm{Hom}_W(\mathcal{F}l^x, T)_1$ is unipotent and $\mathrm{Hom}_W(\mathcal{F}l^x, T) \simeq \mathrm{Hom}_W(\mathcal{F}l^x, T)_1 \times T^W$ is the Jordan decomposition of $\mathrm{Hom}_W(\mathcal{F}l^x, T)$.

The proof is concluded by the observation that

$$Z(G) = \{t \in T^W \mid \alpha(t) = 1, \forall \alpha \in \Delta\},$$

which is exactly the $(**)$ condition. □

12.8. The following will not be used in the sequel, but is important as an observation. It follows From Theorem 11.6 that \mathbb{T} is irreducible as a variety. We claim that it coincides in fact with the neutral connected component of $\overline{\mathbb{T}}$.

Proof. Indeed, let us choose an isogeny $G' \rightarrow G$ such that the derived group of G' is simply connected. For G' condition $(**)$ is superfluous, which means that $\overline{\mathbb{T}'} \simeq \mathbb{T}'$.

On the one hand, it is easy to see that the natural map $\overline{\mathbb{T}'} \rightarrow \overline{\mathbb{T}}$ is finite. On the other hand, Theorem 11.6 implies that it factors as

$$\overline{\mathbb{T}'} \simeq \mathbb{T}' \rightarrow \mathbb{T} \hookrightarrow \overline{\mathbb{T}},$$

which means that \mathbb{T} is closed in $\overline{\mathbb{T}}$. Hence, \mathbb{T} is simultaneously open and closed in $\overline{\mathbb{T}}$, which means that it is a connected component there. □

13. PROOF OF THEOREM 11.8

13.1. We will need an additional property of the isomorphism χ .

By definition, we have a canonical W -equivariant map

$$\mathfrak{t} \times_{\overline{G/N}} \overline{G/T} \rightarrow \mathfrak{t},$$

hence we obtain a map $\mathfrak{t} \rightarrow \mathfrak{t}/W$.

Lemma 13.2. *The above map coincides with the composition*

$$\mathfrak{t} \xrightarrow{\chi^{-1}} \mathfrak{c} \hookrightarrow \mathfrak{g} \times \overline{G/N} \rightarrow \mathfrak{g} \rightarrow \mathfrak{t}/W,$$

where the last arrow is the Chevalley map.

The proof follows from the fact that the two maps coincide over G/N .

13.3. To prove the theorem, we need to show that any map $S \rightarrow \mathcal{H}$ can be lifted, locally in the étale topology, to a map $S \rightarrow G \times \overline{G/N}$.

Thus, let \mathfrak{a}^1 and \mathfrak{a}^2 be two S -points of $\overline{G/N}$ and let $\nu : \tilde{S}^1 \rightarrow \tilde{S}^2$ be an isomorphism between the corresponding cameral covers. The maps \mathfrak{a}^i give rise to vector subbundles $\mathfrak{c}_S^i \subset \mathfrak{g} \otimes \mathcal{O}_S$, and Theorem 11.6 implies that

$$\mathfrak{c}_S^i \simeq \mathrm{Hom}_{W, \mathcal{O}_S}(\mathfrak{t}^*, \mathcal{O}_{\tilde{S}^i}), \quad i = 1, 2.$$

Therefore, the data of ν defines an isomorphism of vector bundles $\nu' : \mathfrak{c}_S^1 \rightarrow \mathfrak{c}_S^2$.

By Proposition 1.3 we can find a section $x_S^1 \in \mathfrak{c}_S^1$, such that $\mathfrak{c}_S^1 = Z_{\mathfrak{g}}(x_S^1)$. Let $x_S^2 \in \mathfrak{c}_S^2$ be the image of x_S^1 under ν' . By making the choice of x_S^1 sufficiently generic, we can assume that x_S^2 is regular, i.e. that $\mathfrak{c}_S^2 = Z_{\mathfrak{g}}(x_S^2)$.

Consider x_S^i , $i = 1, 2$ as maps $S \rightarrow \mathfrak{g}_{reg}$. Lemma 13.2 implies that their compositions with the Chevalley map

$$S \xrightarrow{x_S^i} \mathfrak{g}_{reg} \rightarrow \mathfrak{t}/W$$

coincide. Now, we have the following general assertion that follows from smoothness of the Chevalley map restricted to \mathfrak{g}_{reg} :

Lemma 13.4. *The adjoint action map $G \times \mathfrak{g}_{reg} \rightarrow \mathfrak{g}_{reg} \times_{\mathfrak{t}/W} \mathfrak{g}_{reg}$ is smooth and surjective.*

Therefore, locally there exists a map $g_S : S \rightarrow G$ that conjugates x_S^1 to x_S^2 . Then this map conjugates \mathfrak{c}_S^1 to \mathfrak{c}_S^2 , which is what we had to prove.

13.5. **Complements.** We conclude this section by two remarks regarding the assertions of Theorem 11.6 and Theorem 11.8.

First, let us fix a \mathbb{C} -point $\mathfrak{a} \in \overline{G/N}$ and let $\varphi : \mathcal{H}^{\mathfrak{a}} \rightarrow \mathfrak{t}$ be a W -equivariant map, which according to Theorem 11.6, is the same as an element $x_{\varphi} \in \mathfrak{a} = \mathfrak{c}_{\mathfrak{a}}$. One may wonder: how can one express the condition that x_{φ} is a regular element of \mathfrak{g}_{reg} in terms of φ ?

We claim that the necessary and sufficient condition is that $\varphi : \mathcal{H}^{\mathfrak{a}} \rightarrow \mathfrak{t}$ is a scheme-theoretic embedding. Here is a proof:

Proof. First, one easily reduces the assertion to the case when \mathfrak{a} is the centralizer of a regular nilpotent element, and in particular, consists of nilpotent elements.

Let $\mathrm{St}_G(\mathfrak{a})$ be the normalizer of \mathfrak{a} . It is easy to see that x_φ is regular if and only if its centralizer in $\mathrm{St}_G(\mathfrak{a})$ coincides with $\mathcal{C}_{\mathfrak{a}}$.

By Proposition 11.10, the quotient $\mathrm{St}_G(\mathfrak{a})/\mathcal{C}_{\mathfrak{a}}$ maps isomorphically to the group of W -equivariant automorphisms of $\mathcal{F}^{\mathfrak{a}}$. Therefore, if φ is an embedding, the fact that $\mathrm{Ad}_n(x_\varphi) = x_\varphi$ for some $n \in \mathrm{St}_G(\mathfrak{a})$ implies that Ad_n acts trivially on $\mathcal{F}^{\mathfrak{a}}$, and hence n belongs to $\mathcal{C}_{\mathfrak{a}}$.

Conversely, since \mathfrak{a} consists of nilpotent elements, $\mathfrak{a} \cap \mathfrak{g}_{\mathrm{reg}}$ is the only $\mathrm{St}_G(\mathfrak{a})$ -invariant open subset of \mathfrak{a} consisting of regular elements only. However, the locus of φ that are embeddings is clearly such a subset. \square

Secondly, let us see how Corollary 11.11 is related to Proposition 1.8:

Let \mathfrak{a}_1 and \mathfrak{a}_2 be two \mathbb{C} -points of $\overline{G/N}$. Corollary 11.11 says that they are G -conjugate if and only if $\pi^{-1}(\mathfrak{a}_1) \simeq \pi^{-1}(\mathfrak{a}_2)$ as W -schemes. The condition of Proposition 1.8 is seemingly weaker (but in fact, equivalent): it implies that \mathfrak{a}_1 and \mathfrak{a}_2 are G -conjugate if and only if $(\pi^{-1}(\mathfrak{a}_1))_{\mathrm{red}} \simeq (\pi^{-1}(\mathfrak{a}_2))_{\mathrm{red}}$ as W -schemes.

14. PROOF OF THEOREM 4.4

14.1. We are going to deduce our theorem from Theorem 11.6 and Theorem 11.8 combined with the following “abstract nonsense” observation:

Lemma 14.2. *Let \mathcal{Q} be a sheaf of groupoids on $\mathrm{Sch}_{\mathrm{et}}(X)$. Suppose that for every $(U \rightarrow X) \in \mathrm{Sch}_{\mathrm{et}}(X)$ and every $C \in \mathcal{Q}(U)$, we are given an isomorphism $\mathrm{Aut}_{\mathcal{Q}(U)}(C) \simeq \mathcal{A}(U)$ such that the following conditions hold:*

- (0) *There exists a covering $U \rightarrow X$ such that $\mathcal{Q}(U)$ is non-empty.*
- (1) *If $C_1 \rightarrow C_2$ is an isomorphism between two objects in $\mathcal{Q}(U)$, then the induced isomorphism $\mathrm{Aut}_{\mathcal{Q}(U)}(C_1) \simeq \mathrm{Aut}_{\mathcal{Q}(U)}(C_2)$ is compatible with the identification of both sides with $\mathcal{A}(U)$.*
- (2) *If $f : U' \rightarrow U$ is a morphism in $\mathrm{Sch}_{\mathrm{et}}(X)$ and $C \in \mathcal{Q}(U)$, then the map*

$$f_{\mathcal{Q}}^* : \mathrm{Aut}_{\mathcal{Q}(U)}(C) \rightarrow \mathrm{Aut}_{\mathcal{Q}(U')}(f_{\mathcal{Q}}^*(C))$$

is compatible with the restriction map $\mathcal{A}(U) \rightarrow \mathcal{A}(U')$.

- (3) *For any $U \in \mathrm{Sch}_{\mathrm{et}}(X)$ and any two $C_1, C_2 \in \mathcal{Q}(U)$, there exist a covering $f : U' \rightarrow U$ such that the objects $f_{\mathcal{Q}}^*(C_1)$ and $f_{\mathcal{Q}}^*(C_2)$ of $\mathcal{Q}(U')$ are isomorphic.*

Then \mathcal{Q} has a canonical structure of a gerbe bound by $\mathrm{Tors}_{\mathcal{A}}$.

14.3. We claim that $\mathrm{Higgs}_{\tilde{X}}$ satisfies the conditions of this lemma. Condition (0) is a tautology: locally the cameral cover $\tilde{X} \rightarrow X$ is induced from the universal one by means of a map $X \rightarrow \overline{G/N}$.

Let (E_G, σ, t) be an object of $\mathrm{Higgs}_{\tilde{X}}(U)$. We must construct an isomorphism

$$\mathrm{Aut}_{\mathrm{Higgs}_{\tilde{X}}(U)}(E_G, \sigma, t) \simeq T_{\tilde{X}}(U).$$

Let us first assume that E_G is trivialized and our Higgs bundle corresponds to a map $U \rightarrow \overline{G/N}$ such that $\tilde{U} \xrightarrow{t} \overline{G/T} \times_{\overline{G/N}} U$.

In this case, an automorphism of (E_G, σ) as an object of $\text{Higgs}(U)$ is the same as a map $U \rightarrow \text{St}$ (cf. Sect. 11.9) that covers the given map $X \rightarrow \overline{G/N}$. Now, Proposition 11.10 implies that this automorphism belongs to $\text{Aut}_{\text{Higgs}_{\tilde{X}}(U)}(E_G, \sigma, t)$ if and only if the above map factors as $U \rightarrow \mathcal{C} \rightarrow \text{St}$.

Now we apply Theorem 11.6 which says that $\text{Hom}_{\overline{G/N}}(U, \mathcal{C}) = \text{Hom}_{\overline{G/N}}(U, \mathbb{T}) = T_{\tilde{X}}(U)$.

The fact that the map $\chi : \mathcal{C} \rightarrow \mathbb{T}$ is G -equivariant implies that our isomorphism between $\text{Aut}_{\text{Higgs}_{\tilde{X}}(U)}(E_G, \sigma, t)$ and $T_{\tilde{X}}(U)$ is independent of the choice of a trivialization of E_G . In particular, by SC-1, it defines the required isomorphism for all E_G . The fact that conditions (1) and (2) are satisfied is automatic from the construction.

Finally, let us check condition (3). Let (E_G^1, σ^1, t^1) and (E_G^2, σ^2, t^2) be two objects of $\text{Higgs}_{\tilde{U}}(U)$. Without restricting the generality we can assume that both E_G^1 and E_G^2 are trivialized.

In this case, the data of $(\sigma^1, \sigma^2, t^1 \circ (t^2)^{-1})$ defines a U -point of the scheme \mathcal{H} . By Theorem 11.8 we can locally find a map $g_U : U \rightarrow G$ which conjugates (σ^1, t^1) to (σ^2, t^2) . We can regard g_U as a gauge transformation, i.e. a map $E_G^1 \rightarrow E_G^2$, which defines an isomorphism between (E_G^1, σ^1, t^1) and (E_G^2, σ^2, t^2) .

Thus, Theorem 4.4 is proved.

15. PROOF OF THEOREM 7.3

15.1. It remains to prove Theorem 6.4. In this section we will prove Theorem 7.3 which takes care of the universal situation.

15.2. **Step 1.** First we show that our map $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ is an isomorphism off D^{α_i} . To do that let us analyze more closely the situation described in Sect. 10.5.

Let $\Delta_J \subset \Delta$ be a root subsystem and let $M = M_J$ be the corresponding standard Levi subgroup. Let $\mathcal{F}l_M$ denote the flag variety of M and $B_M = B \cap M$, $U_M = U \cap M$.

It is well-known that there exists a canonical closed embedding $W/W_M \times \mathcal{F}l_M \rightarrow \mathcal{F}l$:

A point $\mathfrak{b}' \in \mathcal{F}l$ belongs to $w \times \mathcal{F}l_M$, if and only if \mathfrak{b}' is in relative position w with respect to $P = P_J$ (this makes sense, as P -orbits on $\mathcal{F}l$ are parametrized exactly by W/W_M) and $\mathfrak{b}' \cap \mathfrak{p}$ is a Borel subalgebra in M .

Consider the restriction of the canonical T -bundle $\mathcal{L}_{\mathcal{F}l}$ to $W/W_M \times \mathcal{F}l_M$. It is easy to see that its further restriction to the connected component $1 \times \mathcal{F}l_M$ identifies with $\mathcal{L}_{\mathcal{F}l_M}$.

Let $w \in W$ be a *minimal* representative of its coset in W/W_M . The action of w defines a map $1 \times \mathcal{F}l_M \rightarrow w \times \mathcal{F}l_M$. Let us consider the pull-back $w^*(\mathcal{L}_{\mathcal{F}l}|_{w \times \mathcal{F}l_M})$ as a T -bundle on $1 \times \mathcal{F}l_M = \mathcal{F}l_M$. Let $\tilde{w} \in N$ be an element that projects to $w \in W$.

Proposition 15.3. *We have a canonical M -equivariant isomorphism*

$$w^*(\mathcal{L}_{\mathcal{F}l}|_{w \times \mathcal{F}l_M}) \simeq \mathcal{L}_{\mathcal{F}l_M}.$$

Proof. Recall that by definition, the fiber of $\mathcal{L}_{\mathcal{F}}$ at a point $\mathfrak{b}' \in \mathcal{F}$ is

$$\{g \in G \mid \text{Ad}_{g^{-1}}(\mathfrak{b}') = \mathfrak{b}\}/U,$$

where the T -action is right multiplication. Similarly, the fiber of $\mathcal{L}_{\mathcal{F}_M}$ at $\mathfrak{b}' \in 1 \times \mathcal{F}_M$ can be canonically identified with $\{m \in M \mid \text{Ad}_{m^{-1}}(\mathfrak{b}') = \mathfrak{b}\}/U_M$.

Since w is minimal, $\text{Ad}_{\tilde{w}}(B_M) \subset B$ and if $\text{Ad}_m(\mathfrak{b})$ is a point of $1 \times \mathcal{F}_M$, then the corresponding point of $w \times \mathcal{F}_M$ equals $\text{Ad}_{m \cdot \tilde{w}^{-1}}(\mathfrak{b})$.

Therefore, the map $m \mapsto m \cdot \tilde{w}^{-1}$ sets up a bijection between the fiber of $\mathcal{L}_{\mathcal{F}_M}$ at $\mathfrak{b}' \in 1 \times \mathcal{F}_M$ and the fiber of $w^*(\mathcal{L}_{\mathcal{F}}|_{w \times \mathcal{F}_M})$ at $w \times \mathfrak{b}'$, such that the T -actions are intertwined up to the action of w , which is what we had to prove. \square

15.4. Let w be as above. Consider the map

$$\overline{M/T} \xrightarrow{\text{Proposition 10.6}} \overline{G/T} \xrightarrow{w} \overline{G/T}.$$

The fact that $\text{Ad}_{\tilde{w}^{-1}}(B) \cap M = B_M$ implies that the above map coincides with

$$\overline{M/T} \rightarrow 1 \times \mathcal{F}_M \xrightarrow{w} w \times \mathcal{F}_M \rightarrow \mathcal{F}.$$

Therefore, from Proposition 15.3 we obtain an isomorphism

$$\gamma'_{can}(\tilde{w}) : w^*(\mathcal{L}_{can})|_{\overline{M/T}} \simeq \mathcal{L}_{can}|_{\overline{M/T}}.$$

Moreover, it is easy to see that the above isomorphism coincides with the restriction to $\overline{M/T}$ of the (meromorphic) isomorphism $\gamma_{can}(\tilde{w})$. In particular, the *a priori* meromorphic isomorphism $\gamma_{can}(\tilde{w})$ is regular on $\overline{M/T}$.

Let us now go back to the situation of the theorem. We must check that the meromorphic map $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ has no poles along D^α if $\alpha \neq \alpha_i$. Choose a minimal Levi subgroup M_j such that $w(\alpha_j) = \alpha$ for some $w \in W$.

Then the fact that $\alpha \neq \alpha_i$ implies that both w and $s_i \cdot w$ are minimal representatives of the corresponding cosets in $W/\langle s_j \rangle$. Then the above discussion shows that $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ has no poles on $w \times \overline{M_j/T}$.

This proves what we need, since the G -orbit of $w \times \overline{M_j/T}$ contains an open part of D^α (cf. Proposition 1.8).

15.5. Step 2. Thus, we have shown that the poles of the map $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ can occur only on D^{α_i} . Let M_i be the corresponding minimal Levi subgroup. As we have seen before, there is a natural embedding $\overline{M_i/T} \rightarrow \overline{G/T}$, and \mathcal{L}_{can} restricts to the corresponding T -bundle on $\overline{M_i/T}$.

Since $s_i^*(\mathcal{L}_{can}) \rightarrow \mathcal{L}_{can}$ is G -equivariant, to determine the contribution of the divisor D^{α_i} , it is enough to perform the corresponding calculation for M_i . The latter case easily reduces to $SL(2)$.

For $SL(2)$, $\overline{G/T} \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{L}_{can} \simeq \mathcal{O}(1) \boxtimes \mathcal{O}$. Moreover, $-1 \in S_2 = W$ acts on $\overline{G/T}$ by swapping the two \mathbb{P}^1 factors, with the fixed-point locus $\overline{G/T}^{-1}$ being the diagonal \mathbb{P}^1 . Hence, $(-1)^*(\mathcal{L}_{can}) \simeq \mathcal{O} \boxtimes \mathcal{O}(-1)$.

Therefore, we have a meromorphic map between $\mathcal{O} \boxtimes \mathcal{O}(-1)$ and $\mathcal{O}(1) \boxtimes \mathcal{O}$, which is allowed to have zeroes and poles only on the diagonal. Then it must have a zero of order 1, by degree considerations.

16. PROOF OF THEOREM 6.4

16.1. Finally, we are ready to complete the proof of the main result. First, we claim that there is a natural functor $\Upsilon : \text{Higgs}'_{\tilde{X}} \rightarrow \text{Higgs}'_{\tilde{X}}$:

Let (E_G, σ, t) be an object of $\text{Higgs}'_{\tilde{X}}(U)$, where $\sigma : E_G \rightarrow \overline{G/N}$ is a G -equivariant map. We can pull-back the universal object of $\text{Higgs}'_{\overline{G/T}}(\overline{G/N})$ (cf. Sect. 7) and obtain a G -equivariant object of $\text{Higgs}'_{\tilde{E}_G}(E_G)$, where \tilde{E}_G is the induced cameral cover of E_G .

By descent, it gives rise to an object of $\text{Higgs}'_{\tilde{X}}(U)$ and this assignment is clearly a functor between sheaves of categories.

The key fact now is that $\text{Higgs}'_{\tilde{X}}$ is also a gerbe bound by $\text{Tors}_{T_{\tilde{X}}}$. Condition (0) of Lemma 14.2 follows from the mere existence of the functor Υ and the fact that $\text{Higgs}'_{\tilde{X}}$ satisfies condition (0).

Let $(\mathcal{L}, \gamma, \beta_i)$ be an object of $\text{Higgs}'_{\tilde{X}}(U)$. We must identify the group of its automorphisms with $T_{\tilde{X}}(U)$.

By definition, this group consists of T -bundle automorphisms, which respect the data of γ and β_i . However, a T -bundle map $\mathcal{L} \rightarrow \mathcal{L}$ is the same as a map $\tilde{U} \rightarrow T$ and compatibility with γ implies that this map is W -equivariant. Therefore, we obtain a section of $\overline{T}_{\tilde{X}}(U)$. Now, compatibility with β_i is exactly condition (*). (Recall that it suffices to impose condition (*) for one representative in every W -orbit on the set of roots. In particular, it is sufficient to impose it for simple roots only.)

It is easy to see that conditions (1) and (2) hold for the above identification of $\text{Aut}_{\text{Higgs}'_{\tilde{X}}(U)}(\mathcal{L}, \gamma, \beta_i) \simeq T_{\tilde{X}}(U)$. In addition, it follows from the construction of χ , that $\Upsilon : \text{Higgs}'_{\tilde{X}} \rightarrow \text{Higgs}'_{\tilde{X}}$ respects the identifications of groups of automorphisms of objects with $T_{\tilde{X}}(U)$.

Assume for a moment that condition (3) of Lemma 14.2 has been checked. We claim, that this already implies Theorem 6.4, because of the following general fact:

Lemma 16.2. *Let \mathcal{Q}_1 and \mathcal{Q}_2 be two gerbes bound by $\text{Tors}_{\mathcal{A}}$ and let $\Upsilon : \mathcal{Q}_1 \rightarrow \mathcal{Q}_2$ be a functor between the corresponding sheaves of categories. Assume that for every $U \in \text{Sch}_{\text{et}}(X)$ and $C \in \mathcal{Q}_1(U)$ we have a commutative square:*

$$\begin{array}{ccc} \mathcal{A}(U) & \longrightarrow & \text{Aut}_{\mathcal{Q}_1(U)}(C) \\ \text{id} \downarrow & & \Upsilon \downarrow \\ \mathcal{A}(U) & \longrightarrow & \text{Aut}_{\mathcal{Q}_2(U)}(\Upsilon(C)). \end{array}$$

Then Υ is an equivalence of $\text{Tors}_{\mathcal{A}}$ -gerbes.

16.3. It remains, therefore, to prove that every two objects of $\text{Higgs}'_{\tilde{X}}(U)$ are locally isomorphic. For that purpose we will introduce a sheaf of Picard categories $\text{Tors}'_{T_{\tilde{X}}}$, which will be the “homogeneous” version of $\text{Higgs}'_{\tilde{X}}(X)$.

Objects of $\text{Tors}'_{T_{\tilde{X}}}(U)$ are triples

$$(\mathcal{L}_0, \gamma_0, \beta_{i,0}),$$

where $(\mathcal{L}_0, \gamma_0)$ is a strongly W -equivariant T -bundle on \tilde{U} and each $\beta_{i,0}$ is a trivialization of $\alpha_i(\mathcal{L}_0)|_{D_U^{\alpha_i}}$.

The following compatibility conditions must hold:

(1) For a simple root α_i , the data of $\gamma_0(s_i) : s_i^*(\mathcal{L}_0) \simeq \mathcal{L}_0$ defines, after restriction to $D_U^{\alpha_i}$, a trivialization

$$\check{\alpha}_i(\alpha_i(\mathcal{L}_0)|_{D_U^{\alpha_i}}) \simeq \check{\alpha}_i(\mathcal{O}_{D_U^{\alpha_i}}).$$

We need that this trivialization coincides with $\check{\alpha}_i(\beta_{i,0})$.

(2) Assume that $w \in W$ conjugates a simple root α_i to another simple root α_j . The pull-back of $\beta_{j,0}$ under w is a trivialization of $\alpha_i(w^*(\mathcal{L}_0))|_{D_U^{\alpha_i}}$, which via $\gamma_0(w)$ defines a trivialization of $\alpha_i(\mathcal{L}_0)$. Our condition is that this trivialization coincides with $\beta_{i,0}$.

Morphisms in $\text{Tors}'_{T_{\tilde{X}}}(U)$ are by definition maps between strongly W -equivariant T -bundles, compatible with the data of $\beta_{i,0}$.

If $(\mathcal{L}_0^1, \gamma_0^1, \beta_{i,0}^1)$ and $(\mathcal{L}_0^2, \gamma_0^2, \beta_{i,0}^2)$ are two objects of $\text{Tors}'_{T_{\tilde{X}}}(U)$ we can form their tensor product $(\mathcal{L}_0^1 \otimes \mathcal{L}_0^2, \gamma_0^1 \otimes \gamma_0^2, \beta_{i,0}^1 \otimes \beta_{i,0}^2)$ which will be a new object of $\text{Tors}'_{T_{\tilde{X}}}(U)$. Moreover, if $(\mathcal{L}_0, \gamma_0, \beta_{i,0})$ is an object of $\text{Tors}'_{T_{\tilde{X}}}(U)$ and $(\mathcal{L}, \gamma, \beta_i)$ is an object of $\text{Higgs}'_{T_{\tilde{X}}}(U)$, we can take their tensor product and obtain another object of $\text{Higgs}'_{T_{\tilde{X}}}(U)$.

It is easy to see that the above constructions define on $\text{Tors}'_{T_{\tilde{X}}}$ a structure of a sheaf of Picard categories and on $\text{Higgs}'_{T_{\tilde{X}}}$ a structure of a gerbe bound by it.

Therefore, to prove that every two objects of $\text{Higgs}'_{T_{\tilde{X}}}(U)$ are locally isomorphic, it is enough to show that any object of $\text{Tors}'_{T_{\tilde{X}}}(U)$ is locally isomorphic to the unit object, i.e. to the one with \mathcal{L}_0 being the *trivial* T -bundle and $(\gamma_0, \beta_{i,0})$ being the tautological maps. The last assertion is equivalent to saying that $\text{Tors}'_{T_{\tilde{X}}}$ is equivalent as a sheaf of Picard categories to $\text{Tors}_{T_{\tilde{X}}}$. The proof is given below.

16.4. Step 1. Without restricting the generality, we can assume that $U = X$ and we must find an étale covering $X' \rightarrow X$, over which a given object $(\mathcal{L}_0, \gamma_0, \beta_{i,0})$ becomes isomorphic to the trivial one.

Fix a \mathbb{C} -point $x \in X$. First, we will reduce our situation to the case when the ramification over x is maximal possible, i.e. when x belongs to the image of $\bigcap_{\alpha} D_X^{\alpha}$, where the intersection is taken over all roots of G .

After an étale localization we can assume that we have a map $X \rightarrow \mathfrak{t}/W$ so that $\tilde{X} = X \times_{\mathfrak{t}/W} \mathfrak{t}$. Let t be a point in \mathfrak{t} which has the same image in \mathfrak{t}/W as x .

By conjugating t , we can assume that there exists $J \subset I$ such that $\alpha_j(t) = 0$ for $j \in \Delta_J$ and $\beta(t) \neq 0$ for $\beta \notin \Delta_J$.

We have a Cartesian square

$$\begin{array}{ccc} W \times^{W_J} (\mathfrak{t} \setminus \bigcup_{\beta \notin \Delta_J} \mathfrak{t}^\beta) & \longrightarrow & \mathfrak{t} \setminus \bigcup_{\beta \notin \Delta_J} \mathfrak{t}^\beta \\ \downarrow & & \downarrow \\ \mathfrak{t}/W_J & \longrightarrow & \mathfrak{t}/W, \end{array}$$

in particular, the map $\mathfrak{t} \rightarrow \mathfrak{t}/W$ is étale in a neighbourhood of the image of t in \mathfrak{t}/W .

Therefore, the base change $X \Rightarrow X' := X \times_{\mathfrak{t}/W} \mathfrak{t}/W_J$ is étale in a neighbourhood of

x . This reduces us to the situation, when the W -cover \tilde{X} is induced from a W_J -cover, i.e. $\tilde{X} \simeq W \times^{W_J} \tilde{X}_J$.

By restricting $(\mathcal{L}_0, \gamma_0, \beta_{i,0})$ to \tilde{X}_J we obtain an object of $\text{Tors}'_{T_{\tilde{X}_J}}(X)$ with respect to W_J . Moreover, it is easy to see that this establishes an equivalence between $\text{Tors}'_{T_{\tilde{X}}}$ and $\text{Tors}'_{T_{\tilde{X}_J}}$, thereby reducing us to the situation when $\Delta_J = \Delta$.

16.5. Step 2. According to Step 1, we may assume that there exists a unique geometric point $\tilde{x} \in \tilde{X}$ over x . To prove the assertion of the proposition, we can replace X by the spectrum of the local ring of X at x . In this case all the D_X^α 's and \tilde{X} are local too.

Let us choose a trivialization of our line bundle \mathcal{L}_0 , subject only to the condition that it is compatible with the data of $\beta_{i,0}$ at \tilde{x} for every simple root α_i . We must show that this trivialization can be modified so that it will be compatible with the structure on \mathcal{L}_0 of a W -equivariant T -bundle, i.e. with the data of γ_0 . (The argument given below mimics the proof of Proposition 9.5).

The discrepancy between our initial trivialization and γ_0 is given by a 1-cocycle $\mu : W \rightarrow \text{Hom}(\tilde{X}, T)$.

The evaluation at \tilde{x} gives rise to a surjection of W -modules: $\text{Hom}(\tilde{X}, T) \rightarrow T$. Thus we obtain a short exact sequence:

$$0 \rightarrow K \rightarrow \text{Hom}(\tilde{X}, T) \rightarrow T \rightarrow 0,$$

where K consists of maps $\tilde{X} \rightarrow T$ which have value 1 at \tilde{x} .

Now, our condition on the trivialization (i.e. its compatibility with $\beta_{i,0}$) and condition (1) in the definition of $\text{Tors}'_{T_{\tilde{X}}}$ imply that $\mu(s_i) \in K$ for every simple reflection s_i . Hence, μ takes values in K .

However, since \tilde{X} is local, K is torsion-free and divisible! Hence, $H^1(W, K) = 0$.

Therefore, we can choose a trivialization of \mathcal{L}_0 which respects the W -equivariant structure and the data of $\beta_{i,0}$ at \tilde{x} . But this implies that it is compatible with the data of $\beta_{i,0}$ on the entire $D_X^{\alpha_i}$, $\forall i \in I$.

Indeed, a possible discrepancy takes values in ± 1 , and its value is constant along every connected component of $D_X^{\alpha_i}$. However, by construction, each $D_X^{\alpha_i}$ is local with \tilde{x} being its unique closed point.

Thus, the proof of Theorem 6.4 is now complete.

REFERENCES

- [1] M. R. Adams, J. Harnad and J. Hurtubise, *Isospectral Hamiltonian flows in finite and infinite dimensions II: Integration of flows*, Com. Math. Phys. **134** (1990), pp. 555-585.
- [2] A. Beauville, *Jacobiennes des courbes spectrales et systèmes hamiltoniens complètement intégrables*, Acta Math. **164** (1990), pp. 211-235.
- [3] A. Beilinson, D. Kazhdan, *Flat Projective Connections*, unpublished (1990).
- [4] F. Bottacin, *Symplectic geometry on moduli spaces of stable pairs*, Thesis, Orsay, 1992.
- [5] P. Deligne and J. S. Milne, *Tannakian categories*, in: Hodge Cycles, Motives, and Shimura varieties, LNM **900** (1982), pp. 101-229.
- [6] J. Dixmier, *Enveloping algebras*, North-Holland Publ. Comp., Amsterdam-New York-Oxford (1977).
- [7] R. Donagi, *Decomposition of spectral covers*, in: Journées de Géométrie Algébrique D'Orsay, Astérisque **218** (1993), pp. 145-175.
- [8] R. Donagi, *Spectral covers*, in: Current topics in complex algebraic geometry, Math. Sci. Res. Inst. Publ. **28** (Berkeley, CA 1992), pp. 65-86.
- [9] R. Donagi, *Principal bundles on elliptic fibrations*, Asian J. Math. **1** (1997), pp. 214-223.
- [10] R. Donagi and E. Markman, *Spectral covers, algebraically completely integrable Hamiltonian systems, and moduli of bundles*, in: Integrable Systems and Quantum Groups, LNM **1620** (1996), pp. 1-119.
- [11] G. Faltings, *Stable G-bundles and Projective Connections*, Jour. Alg. Geo **2** (1993), pp. 507-568.
- [12] R. Friedman, J. Morgan, E. Witten, *Vector Bundles and F-Theory*, Comm. Math. Phys. **187** (1997) pp. 679-743.
- [13] J. Giraud, *Cohomologie non abélienne*, Springer-Verlag, Berlin-New York (1971).
- [14] N. Hitchin, *Stable bundles and integrable systems*, Duke Math. J. **54** (1987), pp. 91-114.
- [15] V. Kanev, *Spectral curves, simple Lie algebras and Prym-Tjurin varieties*, Proc. Symp. Pure Math. **49** (1989), Part I, 627-645.
- [16] B. Kostant, *Lie group representations on polynomial ring*, Amer. J. Math. **81** (1963), pp. 327-404.
- [17] E. Markman, *Spectral curves and integrable systems*, Comp. Math. **93** (1994), pp. 255-290.
- [18] C. Simpson, *Some families of local systems over smooth projective varieties*, Ann. of Math. **138** (1993), pp. 337-425.
- [19] R. Scognamiglio, *Prym-Tjurin Varieties and the Hitchin Map*, Math. Ann. **303** (1995), pp. 47-62.