

SOME ISOMORPHISMS FOR THE BRAUER GROUPS OF A HOPF ALGEBRA

Giovanna Carnovale

Department of Mathematics and Computer Science
Universitaire Instelling Antwerpen
Universiteitsplein 1 - 2610 Wilrijk - Belgium
email: gcarno@uia.ua.ac.be

Abstract

Using equivalences of categories we provide isomorphisms between the Brauer groups of different Hopf algebras. As an example, we show that when k is a field of characteristic different from 2 the Brauer groups $BC(k, H_4, r_t)$ for every dual quasitriangular structure r_t on Sweedler's Hopf algebra H_4 are all isomorphic and abelian. We provide an isomorphism between the Brauer group of a Hopf algebra H and the Brauer group of the dual Hopf algebra H^* generalizing a result of Tilborghs. Finally we compare the Brauer groups of H and of its opposite and co-opposite Hopf algebras.

1 Introduction

There have been given several generalizations of the Brauer group of a field k , due among others to Wall, Long, Van Oystaeyen, Caenepeel and Zhang. In particular, Caenepeel, Van Oystaeyen and Zhang defined in [1] the Brauer group $BQ(H, k)$ of a Hopf algebra H with bijective antipode. This is a special case of Brauer group of a braided monoidal category (see [18]: the Brauer group of a symmetric monoidal category had been defined by B. Pareigis in [15]). Here the category is that of left modules of Drinfel'd quantum double (see [5] and [11]) of a finitely generated projective Hopf algebra H with bijective antipode.

$BQ(H, k)$ generalizes the Brauer-Long group of a commutative and cocommutative Hopf algebra defined by Long in [10]. In fact it is shown in [1] that when H is a commutative cocommutative Hopf algebra, the Brauer group of H and the Brauer-Long group of H are (anti-)isomorphic.

Another example of Brauer group of a braided monoidal category is the Brauer group $BC(k, H, r)$ where H is a dual quasitriangular Hopf algebra with universal R -form r . In this case the category is that of right H -comodules.

By results in [3] and results in [13] (the dual version can be found in [6] and the survey book [8] in the context of quasi Hopf algebras) twisting the algebra structure of H by a 2-cocycle σ provides a new dual quasitriangular Hopf algebra ${}_{\sigma}H_{\sigma^{-1}}$ whose comodule category is equivalent to that of H . Hence the two Brauer groups $BC(k, H, r)$ and $BC(k, {}_{\sigma}H_{\sigma^{-1}}, r_{\sigma})$ will be isomorphic. In this paper we use this result in order to show that although the universal R -forms r_t for $t \in k$ of Sweedler's four dimensional Hopf algebra H_4 are not all isomorphic (this fact was proved by Radford in [16]), the Brauer groups $BC(k, H, r_t)$ are all isomorphic for every $t \in k$. This result is achieved by finding a suitable cocycle which does not change the algebra structure of H_4 but changes the universal R -form r_t into r_0 . Since the Brauer group $BC(k, H_4, r_0)$ (or equivalently, $BM(k, H_4, R_0)$ the Brauer group of the braided category of H_4 -modules with braiding given by the triangular R -matrix R_0) has been computed in [19], the computation of all BC 's for H_4 is accomplished.

Cocycle twisting also provides an isomorphism between the groups $BC(k, H, r)$ and $BC(k, H^{op}, r\tau)$ where τ is the usual flip and H^{op} denotes the Hopf algebra with opposite product.

In the case of a commutative and cocommutative finitely generated and projective Hopf algebra H , it was shown by F. Tilborghs in [17] that the Brauer-Long group of H is (anti-)isomorphic to the Brauer-Long group of H^* .

Hence for H commutative and cocommutative, finitely generated and projective one has an isomorphism between the Brauer group BQ of H and the Brauer group BQ of H^* . We shall use an isomorphism of Radford involving $D(H)$ and $D(H^*)$ together with the results about the Brauer group BC of the opposite Hopf algebra in order to generalize Tilborghs' result to the case of finitely generated projective Hopf algebras with a bijective antipode. Everything boils down to the fact that there is an anti-equivalence of braided categories between the category of Yetter-Drinfel'd H -modules (or crossed bimodules, or Quantum Yang-Baxter modules) and that of Yetter-Drinfel'd H^* -modules given by the usual duality functor mapping left H -modules to right H^* -comodules and right H -comodules to left H^* -modules. Equivalently, there is an anti-equivalence of braided categories between left $D(H)$ -modules and left $D(H^*)$ -modules, where $D(H)$ denotes Drinfel'd quantum double of H .

At the end of the paper we compare $BQ(k, H)$ and $BQ(k, H^{op})$. We show that there is always a map from $BQ(k, H)$ to $BC(k, D(H^{op}), \tau R_{D(H^{op})}^{-1})$. If $D(H)$ were triangular with respect to its standard R -matrix, then there would be an isomor-

phism $BQ(k, H) \simeq BQ(k, H^{op})$ but for nontrivial H the quantum double $D(H)$ cannot be triangular with respect to its standard R -matrix.

The paper is organized as follows. In Section 2 the construction of the Brauer group of a braided monoidal category is recalled. The particular case of the Brauer group(s) of a Hopf algebra are included in a subsection. In Section 3 the theory of cocycle twists of a Hopf algebra is used in order to get isomorphisms between the groups BC of Hopf algebras related by a twist. The main results of the paper are to be found in Sections 4 and 5. In Section 4 I apply the isomorphism above described to the particular example of Sweedler's four dimensional Hopf algebra H_4 in order to obtain $BC(k, H_4, r_t)$ for every $t \in k$. In Section 5, I provide an isomorphism between the Brauer group BQ of a Hopf algebra H with the Brauer group BQ of its dual and I compare it to the Brauer groups related to its opposite Hopf algebra.

2 The Brauer group of a braided category

The Brauer group of a braided monoidal category was defined in [18] and this definition contains all known Brauer groups. The case of a symmetric category had been already treated in [15] and in the symmetric case the Brauer group is abelian. Here we give a short account of the general construction.

Let \mathcal{C} denote a braided monoidal category with \otimes , ψ and I respectively the tensor, braiding and identity object. For objects P and Q in \mathcal{C} if the functor $\mathcal{C}(- \otimes P, Q)$ is representable one denotes the representing object by $[P, Q]$. Similarly if the functor $\mathcal{C}(P \otimes -, Q)$ is representable, the representing object is denoted by $\{P, Q\}$. An object A in \mathcal{C} is an algebra if there are morphisms $m: A \otimes A \rightarrow A$ (product) and $\eta: I \rightarrow A$ (unit) satisfying associativity and unitary conditions. The \mathcal{C} -opposite algebra \bar{A} is then defined as A as object but with product $\bar{m} := m \circ \psi$ and same unit. The tensor product of two algebras A and B in \mathcal{C} becomes an algebra in \mathcal{C} denoted by $A \# B$ with product $(m_A \otimes m_B) \circ (\text{id} \otimes \psi \otimes \text{id})$. For more details, see [13] and references therein. In particular, $[P, P]$ and $\{P, P\}$ are algebras in \mathcal{C} .

For an algebra A in \mathcal{C} one has the two maps:

$$F: \mathcal{C}(X, A \# \bar{A}) \rightarrow \mathcal{C}(X, [A, A]) \quad (2.1)$$

$$F(a \# \bar{b}) < d > = a \bar{m}(b \otimes d) \in \mathcal{C}(X \otimes Y, A) \quad (2.2)$$

for every $d \in \mathcal{C}(Y, A)$ where X and Y are objects in \mathcal{C} and $< d >$ stands for "evaluation" at d and

$$G: \mathcal{C}(X, \bar{A} \# A) \rightarrow \mathcal{C}(X, \{A, A\}) \quad (2.3)$$

$$< d > G(\bar{a} \# b) = \bar{m}(d \otimes a)b \in \mathcal{C}(X \otimes Y, A) \quad (2.4)$$

for every $d \in \mathcal{C}(Y, A)$ where X and Y are objects in \mathcal{C} and $\langle d \rangle$ stands for “evaluation” at d . An algebra in \mathcal{C} is called \mathcal{C} -Azumaya if F and G are isomorphisms and A is faithfully projective in \mathcal{C} (see [15] or [18] for the definition of faithfully projective). It turns out that: the product of two \mathcal{C} -Azumaya algebras is \mathcal{C} -Azumaya; the opposite algebra of a \mathcal{C} -Azumaya algebra is \mathcal{C} -Azumaya. If P is faithfully projective, then $[P, P]$ is also \mathcal{C} -Azumaya. One can define an equivalence relation on the set of \mathcal{C} -Azumaya algebras: $A \sim B$ if there exist faithfully projective objects M and N such that

$$A\#[M, M] \simeq B\#[N, N]. \quad (2.5)$$

It is proved in [18] that this is indeed an equivalence relation and that the set of equivalence classes becomes a group $Br(\mathcal{C})$ with product induced by $\#$. The inverse of a class represented by an algebra A will be the class represented by the algebra \bar{A} .

In [18] the second Brauer group $Br'(\mathcal{C})$ was also defined if \mathcal{C} satisfies some extra conditions. Under general conditions there is a group homomorphism $Br'(\mathcal{C}) \rightarrow Br(\mathcal{C})$ which is very often a monomorphism and it is the identity if the unit object I is projective. The elements of $Br'(\mathcal{C})$ are classes of separable \mathcal{C} -Azumaya algebras in the category (see [15] or [18] for the definition).

Remark 2.1 It is clear that if we replace a category \mathcal{C} by its opposite category \mathcal{C}^{op} with $A \otimes^{op} B := B \otimes A$ for two objects in \mathcal{C}^{op} , $f \otimes^{op} g := g \otimes f$ for two morphisms f and g , and $\psi_{AB}^{op} := \psi_{BA}$, then the map

$$\alpha: Br(\mathcal{C}) \rightarrow Br(\mathcal{C}^{op})$$

$$[A] \mapsto [\bar{A}]_{op}$$

where $[A]$ denotes the class of A and $[A]_{op}$ denotes the equivalence class of A in \mathcal{C}^{op} is a well-defined isomorphism of groups which induces also an isomorphism between $Br'(\mathcal{C})$ and $Br'(\mathcal{C}^{op})$. In fact it is clear that A is \mathcal{C} -Azumaya iff \bar{A} is \mathcal{C}^{op} -Azumaya and $A \sim_{\mathcal{C}} B$ iff $\bar{A} \sim_{\mathcal{C}^{op}} \bar{B}$ iff $A \sim_{\mathcal{C}^{op}} B$.

2.1 The Brauer groups of a Hopf algebra

From now on k will denote a commutative ring. Every k -module M will be assumed to be finitely generated, projective and faithful. This implies the existence of “dual bases” $\{m_i\} \subset M$ and $\{m_j^*\} \subset M^*$ for which $\sum m_j^*(m)m_j = m$ for every $m \in M$.

All tensor product will be intended to be over k .

All Hopf algebras will be assumed to be finitely generated, projective and with bijective antipode S . $\langle -, - \rangle$ shall always denote evaluation between H^* and

H . Since H is finitely generated and projective, H^* is also a finitely generated projective Hopf algebra.

A k -module M is called faithfully projective (progenerator) if there exist elements $m_j^* \in M^*$ and $m_j \in M$ for which $\sum m_j^*(m_j) = 1$.

H^{op} shall denote the Hopf algebra with opposite product and H^{cop} shall denote the Hopf algebra with opposite coproduct.

The Brauer group $BQ(k, H)$ of a Hopf algebra H over k is a particular case of Brauer group of a braided monoidal category where the category is that of Yetter-Drinfel'd H -modules.

Definition 2.2 *A Yetter-Drinfel'd H -module M is a k -module which is a left H -module and a right H -comodule satisfying the compatibility condition*

$$\sum h_{(1)} \cdot m_{(0)} \otimes h_{(2)} m_{(1)} = \sum (h_{(2)} \cdot m)_{(0)} \otimes (h_{(2)} \cdot m)_{(1)} h_{(1)} \quad (2.6)$$

for every $h \in H$ and $m \in M$. In the formula \cdot denotes the H -action on M , $\chi(m) = \sum m_{(0)} \otimes m_{(1)} \in M \otimes H$ is the right comodule structure map and $\Delta(h) = \sum h_{(1)} \otimes h_{(2)}$ denotes the coproduct in H .

Remark 2.3 A Yetter-Drinfel'd module is also sometimes called a crossed bi-module (see [11]) or a Quantum Yang-Baxter H -module (see [9]).

It is a result of S. Majid in [11] that the category of Yetter-Drinfel'd H -modules is equivalent to the category of left $D(H)$ -modules where $D(H)$ is the Drinfel'd double of H defined in [5].

In order to fix notation we recall that $D(H)$ is a quasitriangular Hopf algebra whose underlying coalgebra is $H^{*,cop} \otimes H$, with product

$$(\xi \otimes a)(\eta \otimes b) = \sum \xi \eta_{(2)} \otimes a_{(2)} b < \eta_{(1)}, S^{-1}(a_{(3)}) > < \eta_{(3)}, a_{(3)} >$$

and with R -matrix $\sum (\varepsilon \otimes h_i) \otimes (h_i^* \otimes 1)$ where $\{h_i\}$ and $\{h_i^*\}$ are dual bases in H and H^* .

In this notation the H -action \cdot and the H^* -action $h^* \rightharpoonup m = (\text{id} \otimes < h^*, - >) \chi$ together with the compatibility condition define a $D(H)$ -module structure on a Yetter-Drinfel'd module M and viceversa because H and H^* are subalgebras of $D(H)$. We shall denote the $D(H)$ -action by \triangleright .

It is well-known that the tensor product of two Yetter-Drinfel'd H -modules M and N can be naturally equipped of a Yetter-Drinfel'd H -module structure denoted by $M \tilde{\otimes} N$ as follows:

$$\chi(a \otimes b) = \sum a_{(0)} \otimes b_{(0)} \otimes b_{(1)} a_{(1)} \quad (2.7)$$

and

$$h \cdot (a \otimes b) = \sum h_{(1)} \cdot a \otimes h_{(2)} \cdot b. \quad (2.8)$$

The category $\mathcal{YD}(H)$ of Yetter-Drinfel'd modules (with module and comodule morphisms) together with $\tilde{\otimes}$ becomes a monoidal category. Moreover, there is an isomorphism of Yetter-Drinfel'd H -modules between $M \tilde{\otimes} N$ and $N \tilde{\otimes} M$ given by

$$\phi_{MN}(m \tilde{\otimes} n) = \sum n_{(0)} \tilde{\otimes} n_{(1)} \cdot m \quad (2.9)$$

which makes of the category of Yetter-Drinfel'd H -modules a braided monoidal category. The mentioned equivalence of $\mathcal{YD}(H)$ and ${}_{D(H)}\mathcal{M}$ is an equivalence of braided monoidal categories. In terms of $D(H)$ -modules, if $\mathcal{R} = \sum \mathcal{R}_1 \otimes \mathcal{R}_2$ is the standard R -matrix for $D(H)$ the braiding is nothing but

$$\phi_{MN}(m \tilde{\otimes} n) = \sum \mathcal{R}_2 \triangleright n \tilde{\otimes} \mathcal{R}_1 \triangleright m \quad (2.10)$$

In this setting for a Yetter-Drinfel'd module P , $[P, P]$ and $\{P, P\}$ are the usual $End(P)$ and $End(P)^{op}$ respectively, equipped with the Yetter-Drinfel'd module structures:

$$(h \cdot f)(m) = \sum h_{(1)} \cdot f(S(h_{(2)} \cdot m)); \quad (2.11)$$

$$\chi(f)(m) = \sum f(m_{(0)})_{(0)} \otimes S^{-1}(m_{(1)}) f(m_{(0)})_{(1)} \quad (2.12)$$

for every $m \in M$ and $f \in End(M)$ for $End(M)$ and

$$(h \cdot' f)(m) = \sum h_{(2)} \cdot f(S^{-1}(h_{(1)}) \cdot m); \quad (2.13)$$

$$\chi'(f)(m) = \sum f(m_{(0)})_{(0)} \otimes f(m_{(0)})_{(1)} S(m_{(1)}) \quad (2.14)$$

for every $m \in M$ and $f \in End(M)$ for $End(M)^{op}$. Those constructions come from two possible natural Yetter-Drinfel'd module structures on $[P, I] = P^*$.

An algebra in $\mathcal{YD}(H)$ is called a Yetter-Drinfel'd H -module algebra and it corresponds to a $D(H)$ -module algebra:

Definition 2.4 *A Yetter-Drinfel'd H -module algebra M is an algebra having the structure of a Yetter-Drinfel'd H -module and such that the module and comodule structure make of M a left H -module algebra and a right H^{op} -comodule algebra.*

The tensor product $A \tilde{\otimes} B$ of two Yetter-Drinfel'd H -module algebras A and B becomes a Yetter-Drinfel'd module algebra denoted by $A \# B$ with product given by $(a \# c)(b \# d) := a\phi(c \otimes b)d = \sum ab_{(0)} \# (b_{(1)} \cdot c)d$.

The maps F and G become then:

$$F: A \# \bar{A} \rightarrow End(A) \quad (2.15)$$

$$F(a \# \bar{b})(c) = \sum ac_{(0)}(c_{(1)} \cdot b) = m(a \otimes \phi_{AA}(b \otimes c)) \quad (2.16)$$

and

$$G: \bar{A} \# A \rightarrow \text{End}(A)^{op} \quad (2.17)$$

$$G(\bar{a} \# b)(c) = \sum a_{(0)}(a_{(1)} \cdot c)b = m(\phi_{AA}\tau \otimes \text{id})(\text{id} \otimes \tau)(a \otimes b \otimes c) \quad (2.18)$$

for every a, b and c in A . It had already been proved in [1] that F and G are Yetter-Drinfel'd H -module algebra maps. A Yetter-Drinfel'd H -module algebra A is H -Azumaya if F and G are isomorphisms. In this case $A \sim B$ if there exist faithfully projective Yetter-Drinfel'd H -modules M and N such that $A \# \text{End}(M) \simeq B \# \text{End}(N)$ as Yetter-Drinfel'd H -module algebras. $Br(\mathcal{YD}(H))$ is usually denoted by $BQ(k, H)$.

An H -Azumaya algebra is said to be strongly H -Azumaya if k is a direct summand of A as Yetter-Drinfel'd H -module. The second Brauer group $Br'(\mathcal{YD}(H))$ is usually denoted $BQS(k, H)$ and it coincides with $BQ(k, H)$ if H is semisimple-like and cosemisimple-like (see Prop. 2.27 in [2]). In general $BQS(k, H)$ is the subgroup whose elements are classes parametrized by strongly H -Azumaya algebras.

If H is a quasitriangular Hopf algebra with universal R matrix $R = \sum R_1 \otimes R_2$ it is well known that to every module (algebra) A one can associate a right H^{op} -comodule (algebra) structure on A given by

$$\chi_R(a) = \sum R_2 \cdot a \otimes R_1 \quad (2.19)$$

obtaining a Yetter-Drinfel'd H -module algebra. In this case the category ${}_H\mathcal{M}$ of left H -modules with H -module maps is a full subcategory of $\mathcal{YD}(H)$. $Br({}_H\mathcal{M})$ is then a subgroup of $BQ(k, H)$ and it is usually denoted by $BM(k, H, R)$. It is the subgroup of $BQ(k, H)$ whose elements are represented by a Yetter-Drinfel'd module algebras whose comodule structure is defined by (2.19).

Dually, if H is a dual quasitriangular Hopf algebra with universal R -form r , to a right H^{op} -module (algebra) A one can associate a left H -module (algebra) structure on A given by

$$h \cdot a = \sum a_{(0)}r(h \otimes a_{(1)}) \quad (2.20)$$

obtaining a Yetter-Drinfel'd module algebra. The category \mathcal{M}^H of right H -comodules with H -comodule maps is a full subcategory of $\mathcal{YD}(H)$. $Br(\mathcal{M}^H)$ is then a subgroup of $BQ(k, H)$ denoted by $BC(k, H, r)$. It is the subgroup of $BQ(k, H)$ whose elements are represented by a Yetter-Drinfel'd module algebras whose module structure is defined by (2.20).

It is well-known that $BQ(k, H) \simeq BC(k, D(H)^*, r) \simeq BM(k, D(H), \mathcal{R})$ where $D(H)$ is the Drinfel'd double of H and \mathcal{R} is its standard R -matrix. Hence it is enough to study $BC(k, H, r)$ for a dual quasitriangular Hopf algebra.

3 An equivalence of categories

In this section we show a few isomorphisms for the group BC of a dual quasitriangular Hopf algebra. Everything can be dualized, considering quasitriangular Hopf algebras and BM . We leave this task to the reader.

Let H be a bialgebra over k and let B be a left (resp. right) H -comodule algebra with comodule map χ . A left (resp. right) 2-cocycle σ is a linear map $\sigma: H \otimes H \rightarrow k$ satisfying

- $\sum \sigma(k_{(1)} \otimes m_{(1)}) \sigma(h \otimes k_{(2)} m_{(2)}) = \sum \sigma(h_{(1)} \otimes k_{(1)}) \sigma(h_{(2)} k_{(2)} \otimes m)$
 (resp. $\sum \sigma(k_{(2)} \otimes m_{(2)}) \sigma(h \otimes k_{(1)} m_{(1)}) = \sum \sigma(h_{(2)} \otimes k_{(2)}) \sigma(h_{(1)} k_{(1)} \otimes m)$)
- $\sigma(h \otimes 1) = \sigma(1 \otimes h) = \varepsilon(h) \quad \forall h, k, m \in H$

Then, the σ -left (resp. σ -right) twisted comodule ${}_{\sigma}B$ (resp. B_{σ}) is an algebra with the same underlying vector space as B , and product given by:

$$\bar{a} \cdot \bar{b} = \sum \sigma(a_{(1)} \otimes b_{(1)}) \overline{a_{(0)} b_{(0)}} \quad (3.1)$$

if $\chi(a) = \sum a_{(1)} \otimes a_{(0)}$, and $\chi(b) = \sum b_{(1)} \otimes b_{(0)} \in H \otimes B$ for $a, b \in B$;

$$\text{(resp. } \bar{a} \cdot \bar{b} = \sum \overline{a_{(0)} b_{(0)}} \sigma(a_{(1)} \otimes b_{(1)}) \text{)} \quad (3.2)$$

if $\chi(a) = \sum a_{(0)} \otimes a_{(1)}$, and $\chi(b) = \sum b_{(0)} \otimes b_{(1)} \in B \otimes H$), where $a \mapsto \bar{a}$ denotes the identification of vector spaces (see for instance [14], Paragraph 7.5).

If B is a bialgebra and σ is a left 2-cocycle one can perform such a twist to B , viewed as a left (resp. right) B -comodule algebra. If σ is a convolution invertible left 2-cocycle, then σ^{-1} is a right 2-cocycle. It is well-known (see for instance [13], [3] or [4]) that the double twist ${}_{\sigma}B_{\sigma^{-1}}$ with the same coproduct of B is again a bialgebra and if B is a Hopf algebra, then ${}_{\sigma}B_{\sigma^{-1}}$ is also a Hopf algebra with antipode S_{σ} given by $(u \otimes S \otimes u^{-1})(\Delta \otimes \text{id})\Delta$. Here $u \in {}_{\sigma}B_{\sigma^{-1}}^* \simeq B^*$ is the linear functional given by $u = \sigma(\text{id} \otimes S)\Delta$.

The following facts are also well-known (see for instance [13] and references therein or, for a dual version involving the case of Drinfel'd quasi Hopf algebras, [8] and references therein):

- The category of right H -comodules is equivalent to the category of right ${}_{\sigma}H_{\sigma^{-1}}$ -comodules as monoidal category. In this case though we need the comodule structure of a tensor product to be

$$m \otimes n \rightarrow \sum m_{(0)} \otimes n_{(0)} \otimes m_{(1)} n_{(1)}. \quad (3.3)$$

The monoidal functor \mathcal{F} is the identity on objects (the coproduct is unchanged) and the natural transformation $\eta: \mathcal{F}(M) \otimes \mathcal{F}(N) \rightarrow \mathcal{F}(M \otimes N)$

$$\eta(m \otimes n) = \sum m_{(0)} \otimes n_{(0)} \sigma^{-1}(m_{(1)} \otimes n_{(1)}). \quad (3.4)$$

The compatibility conditions for η follow by the cocycle condition on σ .

- If H is dual quasitriangular with universal R -form r , then ${}_{\sigma}H_{\sigma^{-1}}$ is also dual quasitriangular with universal R -form given by $r_{\sigma} = (\sigma\tau) * r * \sigma^{-1}$.
- The categories \mathcal{M}^H and $\mathcal{M}^{{}_{\sigma}H_{\sigma^{-1}}}$ are equivalent braided monoidal categories. In this case one uses the braiding

$$\phi_{MN}(m \otimes n) \mapsto \sum n_{(0)} \otimes m_{(0)} r(m_{(1)} \otimes n_{(1)}) \quad (3.5)$$

compatible with the different definition of comodule structure on $M \otimes N$. One can check that the braiding ϕ_{MN} is respected by the functor \mathcal{F} and the natural transformation η mentioned above.

- The universal R -forms r and $r^{-1}\tau$ of a dual quasitriangular Hopf algebra H are 2-cocycles and ${}_rH_{r^{-1}} = H^{op} = {}_{r^{-1}\tau}H_{r\tau}$. The new universal R -form is in both cases $r\tau$.

It is a straightforward check that if H is a bialgebra with coproduct Δ , and if σ is a 2-cocycle for H , then $\sigma\tau$ is a 2-cocycle for H^{op} and $({}_{\sigma}H_{\sigma^{-1}})^{op} \simeq_{\sigma\tau} (H^{op})_{\sigma^{-1}\tau}$. Hence, repeating the discussion above for H^{op} the category of right H -comodules with tensor product structure

$$m \otimes n \mapsto \sum m_{(0)} \otimes n_{(0)} \otimes n_{(1)} m_{(1)}$$

and braiding

$$m \otimes n \mapsto \sum n_{(0)} \otimes m_{(0)} r(n_{(1)} \otimes m_{(1)})$$

is equivalent, as braided monoidal category, to that of ${}_{\sigma}H_{\sigma^{-1}}$ -comodules with tensor product structure

$$m \otimes n \mapsto \sum m_{(0)} \otimes n_{(0)} \otimes \sigma(n_{(1)} \otimes m_{(1)}) n_{(2)} m_{(2)} \sigma^{-1}(n_{(3)} \otimes m_{(3)})$$

and braiding

$$m \otimes n \mapsto \sum n_{(0)} \otimes m_{(0)} \sigma(m_{(1)} \otimes n_{(1)}) r(n_{(2)} \otimes m_{(2)}) \sigma^{-1}(n_{(3)} \otimes m_{(3)}).$$

The natural transformation η is in this case:

$$m \otimes n \mapsto \sum m_{(0)} \otimes n_{(0)} \sigma^{-1}(n_{(1)} \otimes m_{(1)}). \quad (3.6)$$

An algebra A in \mathcal{M}^H is mapped by the functor to the algebra $A_{\sigma^{-1}\tau}$ with product $m_A \circ \eta_{A,A}$ i.e. $a \cdot b := \sum a_{(0)} b_{(0)} \sigma^{-1}(b_{(1)} \otimes a_{(1)})$. Hence we have:

Proposition 3.1 *Let H be a (faithfully projective) dual quasitriangular Hopf algebra with universal R -form r . Let σ be an invertible 2-cocycle.*

Then $BC(k, H, r) \simeq BC(k, {}_{\sigma}H_{\sigma^{-1}}, r_{\sigma})$. The class of an H -Azumaya algebra A is mapped to the class of $A_{\sigma^{-1}\tau}$. In particular, $BC(k, H, r) \simeq BC(k, H^{op}, r\tau)$.

Proof: It follows by the above observations. Observe that twisting H by $r^{-1}\tau$ implies that $[A]$ is mapped to $[\overline{A^{op}}]$. \square

Remark 3.2 The result about H^{op} could be obtained also by checking that the categories of right H^{op} comodules and of right H comodules are anti-equivalent braided categories. Then the anti-isomorphism between the two Brauer groups is given on representatives by $A \mapsto A^{op}$.

4 An example: $BC(k, H_4, r_t)$

Let k be a field of characteristic different from 2 and let H_4 denote Sweedler's four dimensional Hopf algebra over k generated by g and h such that $g^2 = 1$, $h^2 = 0$ and $gh + hg = 0$. As far as the coproduct Δ is concerned, g is grouplike and h is twisted-primitive with $\Delta(h) = h \otimes g + 1 \otimes h$. The antipode S is such that $S(g) = g$ and $S(h) = gh$. It is well-known that H_4^* is isomorphic to H_4 . An isomorphism is obtained sending g to $f_1 - f_g$ and h to $f_h + f_{gh}$ where f_x denotes the dual element of $x \in H_4$. It is also well-known that H_4 has a family of universal R -forms (and, dually of universal R -matrices) parametrized by the elements in k . The universal R -forms r_t ($t \in k$) were firstly found by Radford in [16] and are determined by the axioms of an R -form together with

$$r_t(1 \otimes x) = r_t(x \otimes 1) = \varepsilon(x)$$

$$r_t(g \otimes g) = -1 \quad r_t(g \otimes h) = r_t(h \otimes g) = r_t(g \otimes gh) = r_t(gh \otimes g) = 0$$

$$r_t(gh \otimes h) = r_t(h \otimes h) = r_t(gh \otimes gh) = -r_t(h \otimes gh) = t.$$

A first observation is that all those structures are cotriangular, i.e. $r_t * (r_t\tau) = \varepsilon \otimes \varepsilon = (r_t\tau) * r_t$. This means that for every a and b in H_4 one has:

$$\sum r_t(a_{(1)} \otimes b_{(1)})r_t(b_{(2)} \otimes a_{(1)}) = \varepsilon(a)\varepsilon(b) = \sum r_t(b_{(1)} \otimes a_{(1)})r_t(a_{(2)} \otimes b_{(2)}) \quad (4.1)$$

By the symmetry in the formula if we interchange a and b it is enough to check the left hand side condition on pairs of basis elements. We have various cases depending on the coproduct of a and b :

- a and b are both grouplike elements of the basis (i.e. 1 or g). The left hand side of (4.1) reads $r_t(a \otimes b)r_t(b \otimes a)$. This is 1^2 for the pairs $(1, g)$, $(g, 1)$ and $(1, 1)$ and it is $(-1)^2$ when $a = b = g$. In all cases this is equal to $\varepsilon(a)\varepsilon(b)$.

- One element in $\{a, b\}$ is grouplike and the other is twisted primitive (i.e. h or gh). Then each summand of the left hand side of (4.1) will contain an expression of type $r_t(x \otimes y)$ with x grouplike and y twisted primitive. Therefore each summand is zero. For instance

$$r_t * r_t \tau(h \otimes g) = r_t(h \otimes g)r_t(g \otimes g) + r_t(1 \otimes g)r_t(g \otimes h) = 0 = \varepsilon(g)\varepsilon(h).$$

Hence $\sum r_t(a_{(1)} \otimes b_{(1)})r_t(b_{(2)} \otimes a_{(1)}) = 0 = \varepsilon(a)\varepsilon(b)$.

- a and b are both twisted primitives. Then the only nonzero terms in the sum on the left hand side of (4.1) appear when $a_{(1)}$ and $b_{(1)}$ are both twisted primitives, or when $a_{(1)}$ and $b_{(1)}$ are both grouplikes. If $a = b = h$ the expression becomes

$$r_t * r_t \tau(h \otimes h) = r_t(h \otimes h)r_t(g \otimes g) + r_t(1 \otimes 1)r_t(h \otimes h) = -t + t = \varepsilon(h)^2.$$

If $a = b = gh$ the expression becomes

$$r_t(gh \otimes gh)r_t(1 \otimes 1) + r_t(g \otimes g)r_t(gh \otimes gh) = -t + t = \varepsilon(gh)\varepsilon(gh).$$

If $a = h$ and $b = gh$ the expression becomes

$$r_t(h \otimes gh)r_t(1 \otimes g) + r_t(1 \otimes g)r_t(gh \otimes h) = t - t = \varepsilon(gh)\varepsilon(h).$$

Finally if $a = gh$ and $b = h$ the expression is

$$r_t * r_t \tau(gh \otimes h) = r_t(gh \otimes h)r_t(g \otimes 1) + r_t(g \otimes 1)r_t(h \otimes gh) = t - t = \varepsilon(h)\varepsilon(gh).$$

Hence H_4 is cotriangular for every universal R -form r_t . Therefore $BC(k, H_4, r_t) \simeq BC(k, H^{op}, r_t^{-1}) \simeq BC(k, H, r_t^{-1}\tau)$ and it is an abelian group.

Dually, one can check that (H_4, R_t) is triangular for every universal R -matrix R_t . Although triangularity of R_t follows by the previous result we sketch the proof for sake of completeness because we have not met this result in the literature before. The family of R -matrices (see [16] or [9]) is given by

$$R_t = \frac{1}{2}(1 \otimes 1 + 1 \otimes g + g \otimes 1 - g \otimes g) + \frac{t}{2}(h \otimes h + h \otimes gh + gh \otimes gh - gh \otimes h)$$

for $t \in k$. $R_0 = \tau R_0$ and $R_0^2 = 1 \otimes 1$ because it corresponds to a Hopf involution $f_R: H_4^* \rightarrow H_4^{cop}$ where $f_R(\xi) = (\langle \xi, - \rangle \otimes \text{id})(R_0)$ (see [9]). Hence (H_4, R_0) is triangular. Put $R_t = R_0 + R'_t$. (H_4, R_t) is triangular if $(\tau R_t)R_t = 1 \otimes 1 = R_t(\tau R_t)$. Since τ is an algebra isomorphism it is enough to check the relation for $(\tau R_t)R_t$. This expression is equal to

$$(\tau R_0)R_0 + (\tau R_0)R'_t + (\tau R'_t)R'_0 + (\tau R'_t)R'_t = 1 \otimes 1 + (\tau R_0)R'_t + (\tau R'_t)R'_0.$$

$(\tau R'_t)R'_t = 0$ because $h^2 = 0$ appears in every component. Checking that $(\tau R_0)R'_t + (\tau R'_t)R'_0 = 0$ is a straightforward computation that we leave to the reader.

The group $BM(k, H_4, R_0)$ has been computed by Van Oystaeyen and Zhang in [19]. This group is isomorphic to $BC(k, H_4, r_0)$ because the universal R -matrix R_0 goes over to the universal R -form r_0 under the isomorphism $H_4 \rightarrow H_4^*$ previously given.

We want to show here that $BC(k, H_4, r_t) \simeq BC(k, H, r_0)$ for every $t \in k$, hence that $BM(k, H_4, R_0) \simeq BM(k, H_4, R_s)$ for every $s \in k$. We shall do this by providing for every $t \in k$ there exists a suitable element $\sigma_t \in (H_4 \otimes H_4)^*$ such that

- σ_t is a left 2-cocycle for H_4 ;
- σ_t is invertible;
- the twisted product in ${}_{\sigma_t}(H_4)_{\sigma_t^{-1}}$ coincides with the product in H_4 ;
- $\sigma_t \tau * r_t * \sigma_t^{-1} = r_0$.

The functional σ_t is defined on the basis elements of $H_4 \otimes H_4$ as follows:

$$\sigma_t(x \otimes 1) = \sigma_t(1 \otimes x) = \varepsilon(x) \quad \text{for every } x \in H_4.$$

$$\sigma_t(g \otimes g) = 1$$

$$\sigma_t(h \otimes h) = \sigma_t(gh \otimes h) = -\sigma_t(h \otimes gh) = -\sigma_t(gh \otimes gh) = \frac{t}{2}$$

and $\sigma_t(x \otimes y) = 0$ whenever x is grouplike and y twisted primitive or the other way around.

It is a 2-cocycle if for every triple of basis elements k, a, m there holds:

$$\sum \sigma_t(k_{(1)} \otimes m_{(1)}) \sigma_t(a \otimes k_{(2)} m_{(2)}) = \sum \sigma_t(a_{(1)} \otimes k_{(1)}) \sigma_t(a_{(2)} k_{(2)} \otimes m).$$

If one of the elements is 1 then the condition is verified because σ_t is unitary (i.e. it coincides with ε on $1 \otimes x$ and $x \otimes 1$). Hence we have to check the condition on all triples of elements in $\{g, h, gh\}$.

Then we have different cases depending on how often g appears in the triple.

1. g appears 3 times.

Then we have

$$\sigma_t(g \otimes g) \sigma_t(g \otimes 1) = 1 = \sigma_t(g \otimes g) \sigma_t(1 \otimes g)$$

2. g appears twice in the triple:

If $g = a = m$ and k is twisted primitive the condition becomes

$$\sum \sigma_t(k_{(1)} \otimes g) \sigma_t(g \otimes k_{(2)} g) = \sum \sigma_t(g \otimes k_{(1)}) \sigma_t(g k_{(2)} \otimes g).$$

Since k is twisted primitive, in every summand $k_{(1)}$ and $k_{(2)}$ can never be both grouplikes, hence both sums are 0. The cases $g = a = k$ and $g = k = m$ are checked similarly.

3. g appears once in the triple:

If $g = m$ then the condition reads

$$\sum \sigma_t(k_{(1)} \otimes g) \sigma_t(a \otimes k_{(2)} g) = \sum \sigma_t(a_{(1)} \otimes k_{(1)}) \sigma_t(a_{(2)} k_{(2)} \otimes g).$$

The only nonzero component of the left hand side appears when $k_{(1)}$ is grouplike and $k_{(2)} = k$, hence the left hand side is equal to $\sigma_t(a \otimes kg)$. The only nonzero component of the right hand side is when $a_{(2)} k_{(2)}$ is grouplike, i.e. when both $a_{(2)}$ and $k_{(2)}$ are grouplikes, hence $a_{(1)} = a$, $k_{(1)} = k$ and the right hand side becomes $\sigma_t(a \otimes k)$. It is straightforward to check that for the twisted primitives a and k in H_4 there holds: $\sigma_t(a \otimes kg) = \sigma_t(a \otimes k)$.

If $g = a$ by similar computations the left hand side becomes $\sigma_t(k \otimes m)$, with k and m twisted primitives. The right hand side becomes $\sigma_t(gk \otimes m) = \sigma_t(k \otimes m)$ for k and m twisted primitives.

If $g = k$ the right hand side is $\sigma_t(a \otimes gm)$ and the left hand side is $\sigma_t(ag \otimes m)$. Again they coincide for a and m twisted primitives.

4. g does not appear in the triple.

Then $\sum \sigma_t(k_{(1)} \otimes m_{(1)}) \sigma_t(a \otimes k_{(2)} m_{(2)}) = 0$ because if $k_{(2)}$ and $m_{(2)}$ are both twisted primitives, their product involves an h^2 hence it is zero, if one is twisted primitive and the other grouplike, then $\sigma_t(k_{(1)} \otimes m_{(1)}) = 0$ and if they are both grouplikes $\sigma_t(a \otimes k_{(2)} m_{(2)}) = 0$. Similarly one shows that the right hand side is also equal to zero.

Hence σ_t is a left 2-cocycle. Similarly one can prove that σ_t is also a right 2-cocycle.

We claim that $\nu_t \in (H_4 \otimes H_4)^*$ is the convolution inverse of σ_t , where ν_t is defined on the basis elements as follows:

$$\nu_t(x \otimes 1) = \nu_t(1 \otimes x) = \varepsilon(x) \quad \text{for every } x \in H_4.$$

$$\nu_t(g \otimes g) = 1$$

$$\nu_t(h \otimes h) = \nu_t(gh \otimes h) = -\nu_t(h \otimes gh) = -\nu_t(gh \otimes gh) = -\frac{t}{2}$$

and $\nu(x \otimes y) = 0$ whenever x is grouplike and y is twisted primitive or the other way around.

We show that $\sigma_t * \nu_t = \varepsilon \otimes \varepsilon = \nu_t * \sigma_t$. This means that we have to show that for every pair of basis elements a and b in H_4 one has

$$\sum \sigma_t(a_{(1)} \otimes b_{(1)}) \nu_t(a_{(2)} \otimes b_{(2)}) = \varepsilon(a) \varepsilon(b) = \sum \nu_t(a_{(1)} \otimes b_{(1)}) \sigma_t(a_{(2)} \otimes b_{(2)}).$$

Again we divide the different cases:

- a and b are both grouplikes i.e. $a, b \in \{1, g\}$. Then the expression becomes

$$\sigma_t(a \otimes b) \nu_t(a \otimes b) = 1 \cdot 1 = \varepsilon(a) \varepsilon(b) = \nu_t(a \otimes b) \sigma_t(a \otimes b)$$

- a and b are one grouplike and the other twisted primitive. Then the expressions are always zero because in each summand there will be either a $\sigma_t(x \otimes y)$ or a $\nu_t(x \otimes y)$ with one element grouplike and the other twisted primitive. Hence

$$\sum \sigma_t(a_{(1)} \otimes b_{(1)}) \nu_t(a_{(2)} \otimes b_{(2)}) = 0 = \varepsilon(a) \varepsilon(b) = \sum \nu_t(a_{(1)} \otimes b_{(1)}) \sigma_t(a_{(2)} \otimes b_{(2)})$$

- a and b are both twisted primitive. The only nonzero terms in the sums will be those where both $a_{(1)}$ and $b_{(1)}$ are both grouplikes or both twisted primitives. We have for $a = h = b$:

$$\sigma_t * \nu_t(h \otimes h) = \sigma_t(1 \otimes 1) \nu_t(h \otimes h) + \sigma_t(h \otimes h) \nu_t(g \otimes g) + 0 = 0 = \varepsilon(h)^2$$

$$\nu_t * \sigma_t(h \otimes h) = \nu_t(1 \otimes 1) \sigma_t(h \otimes h) + \nu_t(h \otimes h) \sigma_t(g \otimes g) = 0 = \varepsilon(h)^2.$$

For $a = h$ and $b = gh$:

$$\sigma_t * \nu_t(h \otimes gh) = \sigma_t(1 \otimes g) \nu_t(h \otimes gh) + \sigma_t(h \otimes gh) \nu_t(g \otimes 1) = 0 = \varepsilon(h) \varepsilon(gh)$$

$$\nu_t * \sigma_t(h \otimes gh) = \nu_t(1 \otimes g) \sigma_t(h \otimes gh) + \nu_t(h \otimes gh) \sigma_t(g \otimes 1) = 0 = \varepsilon(h) \varepsilon(gh).$$

For $a = gh$ and $b = h$:

$$\sigma_t * \nu_t(gh \otimes h) = \sigma_t(gh \otimes h) \nu_t(1 \otimes g) + \sigma_t(1 \otimes g) \nu_t(gh \otimes h) = 0 = \varepsilon(gh) \varepsilon(h)$$

$$\nu_t * \sigma_t(gh \otimes h) = \nu_t(gh \otimes h) \sigma_t(1 \otimes g) + \nu_t(1 \otimes g) \sigma_t(gh \otimes h) = 0 = \varepsilon(gh) \varepsilon(h).$$

Finally for $a = b = gh$

$$\sigma_t * \nu_t(gh \otimes gh) = \sigma_t(g \otimes g) \nu_t(gh \otimes gh) + \sigma_t(gh \otimes gh) \nu_t(1 \otimes 1) = 0 = \varepsilon(gh)^2$$

$$\nu_t * \sigma_t(gh \otimes gh) = \nu_t(g \otimes g) \sigma_t(gh \otimes gh) + \nu_t(gh \otimes gh) \sigma_t(1 \otimes 1) = 0 = \varepsilon(gh)^2$$

So $\nu_t = \sigma_t^{-1}$. Hence it makes sense to compute the product in ${}_{\sigma_t}H_{\sigma_t^{-1}}$. We shall see that the product in ${}_{\sigma_t}H_{\sigma_t^{-1}}$ coincides with the product in H_4 . We check this fact on products of the the generators h and g : the other products follow by associativity.

$$\begin{aligned}
\bar{g} \cdot \bar{g} &= \sigma_t(g \otimes g) \bar{g}^2 \sigma_t^{-1}(g \otimes g) = 1; \\
\bar{g} \cdot \bar{h} &= \sigma_t(g \otimes 1) \bar{g} \sigma_t^{-1}(g \otimes h) + \sigma_t(g \otimes 1) \bar{g} h \sigma_t^{-1}(g \otimes g) + \\
&\quad \sigma_t(g \otimes h) \bar{g}^2 \sigma_t^{-1}(g \otimes g) = \bar{g} h; \\
\bar{h} \cdot \bar{g} &= 0 + \sigma_t(1 \otimes g) \bar{h} g \sigma_t^{-1}(g \otimes g) = \bar{h} g; \\
\bar{h} \cdot \bar{h} &= \sigma_t(1 \otimes 1) \bar{1} \sigma_t^{-1}(h \otimes h) + \sigma_t(1 \otimes 1) \bar{h} \sigma_t^{-1}(h \otimes g) + \\
&\quad \sigma_t(1 \otimes h) \bar{g} \sigma_t^{-1}(h \otimes g) + \sigma_t(1 \otimes 1) \bar{h} \sigma_t^{-1}(g \otimes h) + \sigma_t(1 \otimes 1) \bar{h}^2 \sigma_t^{-1}(g \otimes g) = \\
&\quad \sigma_t(1 \otimes h) \bar{h} g \sigma_t^{-1}(g \otimes g) + \sigma_t(h \otimes 1) \bar{g} \sigma_t^{-1}(g \otimes h) + \sigma_t(h \otimes 1) \bar{g} h \sigma_t^{-1}(g \otimes g) + \\
&\quad \sigma_t(h \otimes h) \bar{g}^2 \sigma_t^{-1}(g \otimes g) = -\frac{t}{2} + 0 + \frac{t}{2} = 0.
\end{aligned}$$

Hence the product in ${}_{\sigma_t}(H_4)_{\sigma_t^{-1}}$ coincides with the product in H_4 .

Now we compute how r_s changes under twisting, i.e. $\sigma_t \tau * r_s * \sigma_t^{-1}$ for every s and t in k . We shall prove that this is equal to r_{t-s} .

It is known that $\sigma_t \tau * r_s * \sigma_t^{-1}$ is a universal R -form for the twist of H_4 , hence for H_4 . By the properties of an R -form:

$$r(a \otimes 1) = r(1 \otimes a) = \varepsilon(a) \quad \text{and}$$

$$r(ab \otimes c) = \sum r(a \otimes c_{(1)}) r(b \otimes c_{(2)}) \quad \text{for every } a, b \text{ and } c \in H_4$$

it is enough to check the equality when the first argument is h or g . Moreover, since $\Delta(h)$ and $\Delta(g)$ can be expressed in terms of tensor products of h , g and 1 , the property of any R -form r

$$r(a \otimes bc) = \sum r(a_{(2)} \otimes b) r(a_{(1)} \otimes c) \quad \text{for every } a, b \text{ and } c \in H_4$$

implies that it is enough to check that the two forms coincide on $h \otimes h$, $g \otimes g$, $g \otimes h$ and $h \otimes g$. Then

$$\sigma_t \tau * r_s * \sigma_t^{-1}(g \otimes g) = r_{t-s}(g \otimes g) = -1;$$

$$\sigma_t \tau * r_s * \sigma_t^{-1}(g \otimes h) = \sigma_t \tau * r_s * \sigma_t^{-1}(h \otimes g) = 0 = r_{t-s}(h \otimes g) = r_{t-s}(g \otimes h)$$

because every summand will involve one of the forms evaluated at a pair composed by the grouplike g and the twisted primitive h .

$$\sigma_t \tau * r_s * \sigma_t^{-1}(h \otimes h) = \sigma_t(h \otimes h) r_s(g \otimes g) \sigma_t^{-1}(g \otimes g) + \sigma_t(1 \otimes h) r_s(g \otimes h) \sigma_t^{-1}(g \otimes g) +$$

$$\begin{aligned}
& \sigma_t(1 \otimes h)r_s(g \otimes 1)\sigma_t^{-1}(g \otimes h) + \sigma_t(h \otimes 1)r_s(h \otimes g)\sigma_t^{-1}(g \otimes g) + \\
& \sigma_t(1 \otimes 1)r_s(h \otimes h)\sigma_t^{-1}(g \otimes g) + \sigma_t(1 \otimes 1)r_s(h \otimes 1)\sigma_t^{-1}(g \otimes h) + \\
& \sigma_t(h \otimes 1)r_s(1 \otimes g)\sigma_t^{-1}(h \otimes g) + \sigma_t(1 \otimes 1)r_s(1 \otimes h)\sigma_t^{-1}(h \otimes g) + \\
& \sigma_t(1 \otimes 1)r_s(1 \otimes 1)\sigma_t^{-1}(h \otimes h) = -\sigma_t(h \otimes h) + r_s(h \otimes h) + \sigma_t^{-1}(h \otimes h) = \\
& -t + s = r_{t-s}(h \otimes h).
\end{aligned}$$

In particular, for $s = t$ one has $\sigma_s \tau * r_s * \sigma_s^{-1} = r_0$.

Remark 4.1 It is a result by Majid (see [13] or [12]) that if two 2-cocycles are cohomologous then their corresponding twisted Hopf algebras are isomorphic. If the Hopf algebra involved is dual quasitriangular then the corresponding twisted Hopf algebras will be isomorphic as dual quasitriangular Hopf algebras. In particular if a 2-cocycle is a coboundary (i.e. it is cohomologous to $\varepsilon \otimes \varepsilon$) one obtains the same dual quasitriangular Hopf algebra he started with. The computations above show that the converse is not true, i.e. there are 2-cocycles which are not coboundaries for which the twist does not change the Hopf algebra structure. For $t \neq 0$ σ_t is not a coboundary because otherwise (H_4, r_t) would be isomorphic to (H_4, r_0) as dual triangular Hopf algebras which is never true by the results in [16].

We have proved the following

Proposition 4.2 *Let k be a field of characteristic different from 2. Let H_4 be Sweedler's Hopf algebra, and r_t , $t \in k$ be its universal R -forms. Then for every $t \in k$, $BC(k, H_4, r_t) \simeq BC(k, H_4, r_0)$. The isomorphism maps the class of A in $BC(k, H_4, r_0)$ to the class of $A_{\sigma_t \tau}$ in $BC(k, H_4, r_t)$. \square*

Remark 4.3 If k is algebraically closed, the isomorphism between $BC(k, H_4, r_s)$ and $BC(k, H_4, r_t)$ for $st \neq 0$ could be deduced *a priori* by the fact that all the dual quasitriangular structures are isomorphic for $st \neq 0$, see [16]. But $(H_4, r_s) \not\simeq (H_4, r_0)$ for $s \neq 0$. Still, we could show that the corresponding Brauer groups are isomorphic.

Remark 4.4 Proposition 4.2 combined with the results in [19] and self duality of H_4 provide a full description of $BC(k, H_4, r_t)$ and $BM(k, H_4, R_t)$. The computation of the full Brauer group $BQ(k, H_4)$ and the determination of how the different copies of BC fit into BQ in this case is still an open problem.

5 The Brauer groups of H^* and H^{op}

In this section we investigate the relation between $BQ(k, H)$ and $BQ(k, H^*)$ and between $B(k, H)$ and $B(k, H^{op})$. We start with a variation of a Lemma to be found in [16].

Lemma 5.1 *For a finitely generated projective Hopf algebra H over k the usual flip τ defines an isomorphism between $D(H)$ and $D(H^{op, cop, *})^{op}$. The standard R -matrix \mathcal{R} is mapped to $(\tau_{13} \otimes \tau_{24})\mathcal{R}'$ where \mathcal{R}' is the standard R -matrix of $D(H^{op, cop, *})$. In particular, if the antipode S of H is bijective, then $\tau(S^{*-1} \otimes S)$ defines an isomorphism between $D(H)$ and $D(H^*)^{op}$ mapping the universal R -matrix \mathcal{R} to $\tau_{D(H^*), D(H^*)}\mathcal{R}'$. \square*

This Lemma implies that the categories of left modules of the pairs $(D(H), \mathcal{R})$ and $(D(H^*)^{op}, \mathcal{R}')$ are equivalent. Hence the categories of left modules of $D(H)$ and $D(H^*)$ are anti-equivalent braided monoidal categories. Dualizing the above result and using the discussion in the previous sections and Remark 3.2 we have:

Proposition 5.2 *Let H be a faithfully projective Hopf algebra with bijective antipode S . Then $BQ(k, H) \simeq BQ(k, H^*)$.*

Proof: $BQ(k, H)$ is isomorphic to:

$$BC(k, D(H)^*, r) \simeq BC(k, D(H^*)^{op*}, r'\tau) \simeq BC(k, D(H^*)^{*cop}, r'\tau).$$

Applying the antipode of $D(H^*)^*$ and observing that the R -form $r'\tau$ remains unchanged one has:

$$BQ(k, H) \simeq BC(k, D(H^*)^{*op}, r'\tau) \simeq BC(k, D(H^*)^*, r') \simeq BQ(k, H^*).$$

\square

Observe that the above is in fact an anti-isomorphism. We describe it more explicitly generalizing the main result in [17]. We give it in terms of Yetter-Drinfel'd modules and in terms of modules for the Drinfel'd double.

Suppose (M, \cdot, χ) is a Yetter-Drinfel'd H -module. Then (M, \triangleright) is a $D(H)$ -module. The pull-back of this action along the map $(\tau(S^{*-1} \otimes S))^{-1} = (S^* \otimes S^{-1})\tau$ defines on M a $D(H^*)^{op}$ -module structure that is given by $(a \otimes \varepsilon).m = S^{-1}(a) \cdot m$ for elements of the subalgebra $H^{op} \otimes \varepsilon$ and by $(1 \otimes \xi).m = S^*(\xi) \rightharpoonup m$ for elements of the subalgebra $1 \otimes H^{*, op}$. The antipode of $D(H^*)^{op}$ is $S^{-1} \otimes \text{id}$ on $\varepsilon \otimes H^{op}$ and $\text{id} \otimes S^*$ on $1 \otimes H^{*, op}$. Using the action defined in Remark 3.2 the H^* -action on M is given by \cdot and \rightharpoonup . Therefore the map in Proposition 5.2 sends a Yetter-Drinfel'd H -module (M, \cdot, χ) to the Yetter-Drinfel'd H^* -module $(M, \rho, \rightharpoonup)$ where $\rho(m) = \sum m_{(0*)} \otimes m_{(1*)} \in M \otimes H^*$ is such that for every

$l \in H \simeq H^{**}$ one has $(\text{id} \otimes \langle -, l \rangle) \rho(m) = l \cdot m$. This is possible because M is rational.

In other words, the functor from objects in $\mathcal{YD}(H)$ to objects in $\mathcal{YD}(H^*)$ is given by the standard functors from the category of right H -comodules \mathcal{M}^H to the category of left H^* -modules ${}_{H^*}\mathcal{M}$ and from the category of left H -modules ${}_H\mathcal{M}$ to the category of right H^* -modules \mathcal{M}^{H^*} associating to a right comodule (M, χ) the left H^* -module (M, \rightharpoonup) where $h^* \rightharpoonup m = (\text{id} \otimes \langle h^*, - \rangle) \chi(m)$ and to the H -module (M, \cdot) the right H^* -comodule (M, ρ) where $\rho(m) = \sum m_{(0*)} \otimes m_{(1*)}$ as above. One can also check directly that this functor \mathcal{D} maps Yetter-Drinfel'd modules of H to Yetter-Drinfel'd modules of H^* and that it defines an anti-equivalence of braided categories. For instance, the compatibility condition between ρ and \rightharpoonup for the H^* module (M, \rightharpoonup) and comodule (M, ρ) follows from

$$\begin{aligned} & (\text{id} \otimes \langle -, l \rangle) \sum h_{(1)}^* \rightharpoonup m_{(0*)} \otimes h_{(2)}^* m_{(1*)} \\ &= \sum h_{(1)}^* \rightharpoonup (m_{(0*)} \langle m_{(1*)}, l_{(2)} \rangle) \langle h_{(2)}^*, l_{(1)} \rangle \\ &= \sum h_{(1)}^* \rightharpoonup (l_{(2)} \cdot m) \langle h_{(2)}^*, l_{(1)} \rangle \\ &= \sum (l_{(2)} \cdot m)_{(0)} \langle h_{(1)}^*, (l_{(2)} \cdot m)_{(1)} \rangle \langle h_{(2)}^*, l_{(1)} \rangle \\ &= (\text{id} \otimes \langle h^*, - \rangle) \sum (l_{(2)} \cdot m)_{(0)} \otimes (l_{(2)} \cdot m)_{(1)} l_{(1)} \end{aligned}$$

for any $h^* \in H^*$, $m \in M$ and $l \in H \simeq H^{**}$ together with the compatibility condition for the Yetter-Drinfel'd H -module M .

We have seen that \mathcal{D} defines a monoidal functor from $\mathcal{YD}(H)$ to $\mathcal{YD}(H^*)^{op}$, the monoidal category whose tensor structure is given by $\tilde{\otimes}^{op}$, where $M \tilde{\otimes}_*^{op} N = N \tilde{\otimes}_* M$ is the opposite product of N and M viewed as Yetter-Drinfel'd H^* -modules. The family of natural isomorphisms

$$\tau(M, N): \mathcal{D}(M) \tilde{\otimes}_*^{op} \mathcal{D}(N) = \mathcal{D}(N) \tilde{\otimes}_* \mathcal{D}(M) \rightarrow \mathcal{D}(M \tilde{\otimes} N)$$

compatible with \mathcal{D} and the tensor products is given by the usual flips τ . In fact one can also check directly that τ is a module and comodule isomorphism. For instance for M and N objects in $\mathcal{YD}(H)$ the left H^* -module structure on $M \tilde{\otimes} N$ is

$$h^* \rightharpoonup \tau(n \tilde{\otimes} m) = \sum m_{(0)} \otimes n_{(0)} \langle h^*, n_{(1)} m_{(1)} \rangle \quad (5.1)$$

$$= \sum m_{(0)} \langle h_{(2)}^*, m_{(1)} \rangle \otimes n_{(0)} \langle h_{(1)}^*, n_{(1)} \rangle \quad (5.2)$$

$$= \sum (h_{(2)}^* \rightharpoonup_M m) \otimes (h_{(1)}^* \rightharpoonup_N n) \quad (5.3)$$

where \rightarrow_M and \rightarrow_N denote the H^* left module structure associated to the right H -comodule structure of M and N . The right H^* -comodule structure ρ of $M \tilde{\otimes} N$ is such that for every $l \in H$ one has

$$\begin{aligned} (\text{id}^{\otimes 2} \otimes \langle -, l \rangle) \rho(m \tilde{\otimes} n) &= l \cdot (m \tilde{\otimes} n) = \\ \sum (l_{(1)} \cdot m) \tilde{\otimes} (l_{(2)} \cdot n) &= \sum (\text{id} \otimes l_{(1)}) (\rho_M(m)) \otimes (\text{id} \otimes l_{(2)}) (\rho_N(n)) \\ &= (\text{id}^{\otimes 2} \otimes \langle -, l \rangle) \sum m_{(0*)} \otimes n_{(0*)} \otimes m_{(1*)} n_{(1*)} \end{aligned}$$

where $\rho_M(m) = \sum m_{(0*)} \otimes m_{(1*)}$ and $\rho_N(n) = \sum n_{(0*)} \otimes n_{(1*)}$, hence τ is a comodule isomorphism.

One can also see directly that \mathcal{D} and τ are compatible with the braidings ϕ in $\mathcal{YD}(H)$ and ψ_{MN}^{op} in $\mathcal{YD}(H^*)^{op}$ where ψ_{MN}^{op} is $\psi_{NM}: N \tilde{\otimes}_* M \rightarrow M \tilde{\otimes}_* N$ as Yetter-Drinfel'd H^* -modules. In fact for any pair of objects M and N in $\mathcal{YD}(H)$ and for $n \otimes m \in \mathcal{D}(M) \tilde{\otimes}^{op} \mathcal{D}(N)$,

$$\mathcal{D}(\phi_{MN}) \tau_{NM}(n \otimes m) = \phi_{MN}(m \otimes n) = \sum n_{(0)} \otimes n_{(1)} \cdot m$$

coincides with

$$\tau_{MN} \circ \psi_{MN}^{op}(n \otimes m) = \tau_{MN} \sum m_{(0*)} \otimes \langle m_{(1*)}, n_{(1)} \rangle n_{(0)} = \sum n_{(0)} \otimes n_{(1)} \cdot m.$$

In terms of \mathcal{D} and τ one sees that if A is a Yetter-Drinfel'd H -module algebra, the product in $\mathcal{D}(A)$ is given by $\mathcal{D}(m_A) \tau_{A,A}$, where m_A is the product in A , i.e. A equipped with the opposite product is a Yetter-Drinfel'd H^* -module algebra as we had seen before. We have that τ defines an algebra isomorphism between $\mathcal{D}(B \# A) = (B \# A)^{op}$ and $\mathcal{D}(B)^{op} \#_*^{op} \mathcal{D}(A) \simeq A^{op} \#_* B^{op}$ where $\#_*$ the tensor product of Yetter-Drinfel'd H^* -module algebras. This can be checked directly for every $a, c \in A$ and $b, d \in B$:

$$\begin{aligned} m_{(B \# A)^{op}}(\tau \otimes \tau)((a \#_* b) \otimes (c \#_* d)) &= d \phi_{AB}(c \otimes b) a = \\ \tau \left(\sum a \circ \tau \phi_{AB}(c \otimes b) \circ d \right) &= \tau(a \circ (\psi_{BA} \tau(c \otimes b)) \circ d) \\ = \tau(a \circ (\psi_{BA}(b \otimes c)) \circ d) &= \tau m_{A^{op} \#_* B^{op}}((a \#_* b) \otimes (c \#_* d)) \end{aligned}$$

where \circ denotes the opposite product in A and B and the interchange between $\tau \phi_{AB}$ and $\phi_{BA} \tau$ follows by the compatibility of \mathcal{D} and τ with the braidings.

Proposition 5.3 *Let H be a finitely generated projective Hopf algebra over k with bijective antipode. Then the isomorphism between $BQ(k, H)$ and $BQ(k, H^*)$ of Proposition 5.2 maps $[A] \mapsto [\mathcal{D}(A)]_*$. Here $[B]_*$ denotes the equivalence class of B as H^* -Azumaya algebra. \square*

For sake of completeness we show also the (functorial) reciprocity for the opposite algebra, F and G with respect to \mathcal{D} . $\overline{\mathcal{D}(\bar{A})}$ has product $m_{\mathcal{D}(A)}\psi_{AA}^{op} = m_A\tau\psi_{AA} = m_A\phi_{AA}\tau = m_{\mathcal{D}(\bar{A})}$. Hence $\mathcal{D}(\bar{A}) \simeq \overline{\mathcal{D}(\bar{A})}$. As far as the canonical maps in $\mathcal{YD}(H^*)$ F_* and G_* are concerned, F_* is the composite:

$$\mathcal{D}(A) \#_*^{op} \overline{\mathcal{D}(\bar{A})} = \mathcal{D}(A) \#_* \mathcal{D}(\bar{A}) \xrightarrow{\tau} \mathcal{D}(A \# \bar{A}) \xrightarrow{\mathcal{D}(F)} \mathcal{D}(\text{End}(A)) = \text{End}(\mathcal{D}(A))^{op}$$

and similarly for G_*

$$\overline{\mathcal{D}(\bar{A})} \#_*^{op} \mathcal{D}(A) = \mathcal{D}(\bar{A}) \#_* \mathcal{D}(A) \xrightarrow{\tau} \mathcal{D}(\bar{A} \# A) \xrightarrow{\mathcal{D}(G)} \mathcal{D}(\text{End}(A)^{op}) = \text{End}(\mathcal{D}(A)).$$

It is a small exercise in sigma notation to check that $F_*(a \#_* b)(c) = G_*(\bar{b} \#_* a)(c)$, $G_*(\bar{b} \#_* a)(c) = F_*(a \#_* \bar{b})(c)$ and that $\mathcal{D}(\text{End}(A))$ is indeed $\text{End}(A)^{op}$ with the natural Yetter-Drinfel'd H^* -module structure.

Corollary 5.4 *For a finitely generated projective Hopf algebra H over a commutative ring k , with bijective antipode the isomorphism in Proposition 5.3 induces an isomorphism between $BQS(k, H)$ and $BQS(k, H^*)$.*

Proof: It is immediate to check that k is a direct summand of A as a Yetter-Drinfel'd H -module if and only if k is a direct summand of A as a Yetter-Drinfel'd H^* -module. \square

Corollary 5.5 *If H is commutative or cocommutative then all the “associated” Hopf algebras (opposite, co-opposite, opposite of the dual, co-opposite of the dual, etcetera) have the same Brauer group.* \square

It is an interesting question whether an isomorphism between $BQ(k, H)$ and $BQ(k, H^{op})$ holds in general. Here follow some observations concerning this question.

Lemma 5.6 *The linear map*

$$S_H^* \otimes \text{id}: D(H^{cop}) \rightarrow D(H)^{cop}$$

is a Hopf algebra isomorphism. If for $D(H)^{cop}$ we have the R -matrix $\tau\mathcal{R}_{D(H)}$ then the corresponding R -matrix in $D(H^{cop})$ is $\tau\mathcal{R}_{D(H^{cop})}^{-1}$.

Proof: Direct computation. \square

Proposition 5.7 *For any finitely generated projective Hopf algebra H , there is an isomorphism between $BQ(k, H)$ and $BM(k, D(H^{cop}), \tau\mathcal{R}_{D(H^{cop})}^{-1})$.*

Proof: One has

$$BQ(k, H) \simeq BM(k, D(H), R) \simeq BM(k, D(H)^{cop}, \tau\mathcal{R})$$

where the second isomorphism is clear because the algebra structure is unchanged. By Lemma 5.6

$$BQ(k, H) \simeq BM(k, D(H^{cop}), \tau\mathcal{R}_{D(H^{cop})}^{-1}) \simeq BM(k, D(H^{op}), \tau\mathcal{R}_{D(H^{op})}^{-1})$$

where the second isomorphism is induced by the isomorphism $S_H^{-1} \otimes S_H^*$ from $D(H^{op})$ to $D(H^{cop})$ that leaves the R -matrix invariant. \square Under the above series of maps (coming from equivalences of categories) the Yetter-Drinfel'd H -module (M, \cdot, χ) is mapped to the Yetter-Drinfel'd H^{op} -module (M, \rightarrow, χ) with $h \rightarrow m := S(h) \cdot m$ where S is the antipode of H . In fact

$$\chi(h \rightarrow m) = \chi(Sh \cdot m) = \sum (Sh)_{(2)} \cdot m_{(0)} \otimes (Sh)_{(3)} m_{(1)} S^{-1}(Sh)_{(1)} =$$

$$\sum S(h)_{(2)} \cdot m_{(0)} \otimes S(h_{(1)}) m_{(1)} h_{(3)} = \sum S(h)_{(2)} \cdot m_{(0)} \otimes h_{(3)} \circ m_{(1)} \circ S_{H^{op}}^{-1}(h_{(1)})$$

where \circ denotes the opposite product. This construction defines a functor which, together with the natural transformation τ gives an anti-equivalence of monoidal categories. The natural isomorphism τ would respect the braiding if for every Yetter-Drinfel'd module M and N , for every $m \in M$ and $n \in N$ one had

$$\sum m_{(0)} \otimes m_{(1)} \cdot n = \sum (Sn_{(1)}) \cdot m \otimes n_{(0)}.$$

In terms of modules over Drinfel'd quantum double of H ,

$$\sum m_{(0)} \otimes m_{(1)} \cdot n = \tau(\mathcal{R} \triangleright (n \otimes m))$$

if $\mathcal{R} = \sum \mathcal{R}_1 \otimes \mathcal{R}_2$ is the canonical \mathcal{R} -matrix for $D(H)$. If $P = \sum P_1 \otimes P_2$ is $\tau_{D(H), D(H)} \mathcal{R}^{-1} = \tau_{D(H), D(H)} (\text{id} \otimes S_{D(H)}^{-1}) \mathcal{R}$, then

$$\sum (Sn_{(1)}) \cdot m \otimes n_{(0)} = \tau(P \triangleright (n \otimes m)).$$

Hence τ would respect the braidings if $D(H)$ were triangular. Then $BQ(k, H)$ would be isomorphic to $BQ(k, H^{op})$. However, for k a field and H nontrivial, $(D(H), \mathcal{R})$ is never triangular. An easy way to see this is given by the theory of the exponent of a finite-dimensional Hopf algebra introduced by P. Etingof and S. Gelaki in [7]. The authors show the so-called exponent of a Hopf algebra H is equal to the order of $(\tau\mathcal{R})\mathcal{R}$ in $D(H)^{\otimes 2}$ and also equal to the order of the element $u = m(S_{D(H)} \otimes \text{id}^{\otimes 2})\tau(\mathcal{R}) = \sum S^{*-1}(h_j^*) \otimes h_j$ in $D(H)$. Here $\{h_j^*\}_{j \in J}$

and $\{h_j\}_{j \in J}$ are dual bases and S^* is the antipode of H^* . It is clear then that $(\tau\mathcal{R})\mathcal{R} \neq \varepsilon \otimes 1 \otimes \varepsilon \otimes 1$ because $u \neq \varepsilon \otimes 1$.

Observe that triangularity of \mathcal{R} is not a necessary condition for $BQ(k, H) \simeq BQ(k, H^{op})$ to hold. This is clear by looking at the case of H commutative. In particular Proposition 5.7 implies that if H is commutative then there is an isomorphism between $BC(k, D(H), \mathcal{R})$ and $BC(k, D(H), \tau\mathcal{R}^{-1})$.

ACKNOWLEDGEMENTS

This work has been carried over at the University of Antwerp where the author had a post-doc position financed by the EC network “Algebraic Lie Representations” contract ERB-FMRX-CT97-0100. The author wishes to thank University of Antwerpen for the warm hospitality and Professor Fred Van Oystaeyen, Dr. Juan Cuadra and Dr. Yinhua Zhang for valuable comments and useful discussions.

References

- [1] S. Caenepeel, F. Van Oystaeyen and Y. H. Zhang, Quantum Yang-Baxter Module Algebras, K-theory **8(3)** (1994) 231-255.
- [2] S. Caenepeel, F. Van Oystaeyen and Y. H. Zhang, The Brauer group of Yetter-Drinfel’d module algebras, Trans. Amer. Math. Soc. **349(9)** (1997) 3737-3771.
- [3] Y. Doi, Braided bialgebras and quadratic bialgebras, Comm. Algebra, **21(5)** (1993) 1731-1749.
- [4] Y. Doi and M. Takeuchi, Multiplication alteration by 2-cocycles - The quantum version, Comm. Algebra **22(14)**(1994) 5715-5732.
- [5] V.G. Drinfeld, Quantum Groups, Proc. ICM at Berkeley, ed. A. Gleason, AMS (1987).
- [6] V. G. Drinfel’d, On quasitriangular quasi Hopf algebras and a group closely connected to $Gal(\overline{\mathbf{Q}}/\mathbf{Q})$, Leningrad Math. J. **2** (1991) 829-860.
- [7] P. Etingof and S. Gelaki, On the exponent of finite-dimensional Hopf algebras Math. Res. Lett. **6(2)**(1999) 131-140.

- [8] C. Kassel, Quantum Groups, GTM 155, Springer-Verlag (1995).
- [9] L. A. Lambe and D. E. Radford, Introduction to the Quantum Yang-Baxter equation and Quantum Groups: An Algebraic Approach, Kluwer Academic Publishers (1997).
- [10] R. Long, The Brauer group of dimodule algebras, J. of Algebra **30** (1974) 559-601.
- [11] S. Majid, Doubles of quasitriangular Hopf algebras, Comm. Algebra **19** (1991) 3061-3073.
- [12] S. Majid, Cross product quantisation, nonabelian cohomology and twisting of Hopf algebras, In: Generalized symmetries in physics (Clausthal, 1993), World Sci. Publishing, River Edge, NJ, (1994) 13-41.
- [13] S. Majid, Foundations of Quantum Group Theory, Cambridge University Press (1995).
- [14] S. Montgomery, Hopf algebras and their actions on rings, CBMS 28, AMS (1993).
- [15] B. Pareigis, Non-additive ring and module theory IV- The Brauer group of a symmetric monoidal category, in: Brauer Groups, ed. D. Zelinski, LNM, **549** Springer-Verlag (1976) 112-133.
- [16] D. E. Radford, Minimal quasitriangular Hopf algebras, J. of Algebra, **157(2)** (1993) 285-315.
- [17] F. Tilborghs, An anti-homomorphism for the Brauer-Long group, Math. J. Okayama Univ. **32** (1990) 43-52.
- [18] F. Van Oystaeyen and Y. H. Zhang, The Brauer group of a braided monoidal category, J. of Algebra **202** (1998) 96-128.
- [19] F. Van Oystaeyen and Y. H. Zhang, The Brauer group of Sweedler's Hopf algebra H_4 , Proc. AMS to appear, (1999).