ON REPRESENTATIONS OF INTEGERS BY INDEFINITE TERNARY QUADRATIC FORMS

Mikhail Borovoi

ABSTRACT. Let f be an indefinite ternary integral quadratic form and let q be a nonzero integer such that $-q\det(f)$ is not a square. Let N(T, f, q) denote the number of integral solutions of the equation f(x) = q where x lies in the ball of radius T centered at the origin. We are interested in the asymptotic behavior of N(T, f, q) as $T \to \infty$. We deduce from the results of our joint paper with Z.Rudnick that $N(T, f, q) \sim cE_{HL}(T, f, q)$ as $T \to \infty$, where $E_{HL}(T, f, q)$ is the Hardy-Littlewood expectation (the product of local densities) and $0 \le c \le 2$. We give examples of f and q such that c takes values 0,1,2.

0. INTRODUCTION

Let f be a nondegenerate indefinite integral-matrix quadratic form of n variables:

$$f(x_1, \dots, x_n) = \sum_{i=1, j=1}^n a_{ij} x_i x_j, \quad a_{ij} \in \mathbf{Z}, \quad a_{ij} = a_{ji}.$$

Let $q \in \mathbf{Z}$, $q \neq 0$. Let $W = \mathbf{Q}^n$. Consider the affine quadric X in W defined by the equation

$$f(x_1,\ldots,x_n)=q$$

We wish to count the representations of q by the quadratic form f, that is the integer points of X.

Since f is indefinite, the set $X(\mathbf{Z})$ can be infinite. We fix a Euclidean norm $|\cdot|$ on \mathbf{R}^n . Consider the counting function

$$N(T, X) = \#\{x \in X(\mathbf{Z}) : |x| \le T\}$$

where $T \in \mathbf{R}$, T > 0. We are interested in the asymptotic behavior of N(T, X) as $T \to \infty$.

When $n \ge 4$, the counting function N(T, X) can be approximated by the product of local densities. For a prime p set

$$\mu_p(X) = \lim_{k \to \infty} \frac{\# X(\mathbf{Z}/p^k \mathbf{Z})}{(p^k)^{n-1}} \,.$$

¹⁹⁹¹ Mathematics Subject Classification. Primary 11E12.

Key words and phrases. Ternary quadratic forms.

Partially supported by the Hermann Minkowski Center for Geometry

For almost all p it suffices to take k = 1:

$$\mu_p = \frac{\#X(\mathbf{F}_p)}{p^{n-1}} \,.$$

Set $\mathfrak{S}(X) = \prod_p \mu_p(X)$, this product converges absolutely (for $n \ge 4$), it is called the singular series. Set

$$\mu_{\infty}(T,X) = \lim_{\varepsilon \to 0} \frac{\operatorname{Vol}\{x \in \mathbf{R}^n : |x| \le T, |f(x) - q| < \varepsilon/2, \}}{\varepsilon},$$

it is called the singular integral.

Theorem. For $n \ge 4$

$$N(T,X) \sim \mathfrak{S}(X)\mu_{\infty}(T,X) \text{ as } T \to \infty.$$

This theorem follows from results of [BR], 6.4 (based on analytical results of [DRS], [EM], [EMS]). For certain non-Euclidean norms it was earlier proved by the Hardy-Littlewood circle method, cf. [Da], [Est].

We are interested here in the case n = 3, a ternary quadratic form. This case is beyond the range of the Hardy-Littlewood circle method. Set $D = \det(a_{ij})$. We assume that -qD is not a square. Then the product $\mathfrak{S}(X) = \prod \mu_p(X)$ conditionally converges (see Sect. 1 below), but in general N(T, X) is not asymptotically $\mathfrak{S}(X)\mu_{\infty}(T, X)$. From results of [BR] it follows that

$$N(T,X) \sim c_X \mathfrak{S}(X) \mu_{\infty}(T,X)$$
 as $T \to \infty$

with $0 \le c_X \le 2$, see details in Subsection 1.5 below. We wish to know what values can take c_X .

A case when $c_X = 0$ was already known to Siegel, see also [BR], 6.4.1. Consider the quadratic form

$$f_1(x_1, x_2, x_3) = -9x_1^2 + 2x_1x_2 + 7x_2^2 + 2x_3^2,$$

and take q = 1. Let X be defined by $f_1(x) = q$. Then f_1 does not represent 1 over \mathbf{Z} , so N(T, X) = 0 for all T. On the other hand, f_1 represents 1 over \mathbf{R} and over \mathbf{Z}_p for all p, and $\mathfrak{S}(X)\mu_{\infty}(T, X) \to \infty$ as $T \to \infty$. Thus $c_X = 0$ (see details in Sect. 2).

We show that c_X can take value 2. Recall that two integral quadratic forms f, f' are in the same genus, if they are equivalent over \mathbf{R} and over \mathbf{Z}_p for every prime p, cf. e.g. [Ca].

Theorem 0.1. Let f be an indefinite integral-matrix ternary quadratic form, $q \in \mathbb{Z}$, $q \neq 0$, and let X be the affine quadric defined by the equation f(x) = q. Assume that f represents q over \mathbb{Z} and there exists a quadratic form f' in the genus of f, such that f' does not represent q over \mathbb{Z} . Then $c_X = 2$:

$$N(T,X) \sim 2\mathfrak{S}(X)\mu_{\infty}(T,X) \text{ as } T \to \infty.$$

Theorem 0.1 will be proved in Sect. 2

Example 0.1.1. Let $f_2(x_1, x_2, x_3) = -x_1^2 + 64x_2^2 + 2x_3^2$, q = 1. Then f_2 represents 1 ($f_2(1, 0, 1) = 1$) and the quadratic form f_1 considered above is in the genus of f_2 (cf. [CS], 15.6). The form f_1 does not represent 1. Take $|x| = (x_1^2 + 64x_2^2 + 2x_3^2)^{1/2}$. By Theorem 0.1 $c_X = 2$ for the variety $X : f_2(x) = 1$. Analytic and numeric calculations give $2\mathfrak{S}(X)\mu_{\infty}(T, X) \sim 0.794T$. On the other hand, numeric calculations give for T = 10,000 N(T, X)/T = 0.8024.

We also show that c_X can take the value 1.

Theorem 0.2. Let f be an indefinite integral-matrix ternary quadratic form, $q \in \mathbb{Z}$, $q \neq 0$, and let X be the affine quadric defined by the equation f(x) = q. Assume that $X(\mathbf{R})$ is two-sheeted (has two connected components). Then $c_X = 1$:

$$N(T,X) \sim \mathfrak{S}(X)\mu_{\infty}(T,X) \text{ as } T \to \infty.$$

Theorem 0.2 will be proved in Sect. 4.

Example 0.2.1. Let f_2 and |x| be as in Example 0.1.1, q = -1, $X : f_2(x) = q$. Then $X(\mathbf{R})$ has two connected components, and by Theorem 0.2 $c_X = 1$. Analytic and numeric calculations give $\mathfrak{S}(X)\mu_{\infty}(T,X) \sim 0.7065T$. On the other hand, numeric calculations give for T = 10,000 N(T,X)/T = 0.7048.

Question 0.3. Can c_X take values other than 0, 1, 2?

Remark 0.4. It seems that Theorems 0.1 and 0.2 also can be proved using a result of Kneser ([Kn], Satz 2) together with Siegel's weight formula [Si] and the results of [DRS], [EM], [EMS].

The plan of the paper is the following. In Section 1 we expose results of [BR] in the case of 2-dimensional affine quadrics. In Section 2 we treat in detail the example of $c_X = 0$. In Section 3 we prove Theorem 0.1. In Section 4 we prove Theorem 0.2.

Acknowledgements. This paper was partly written when the author was visiting Sonderforschungsbereich 343 "Diskrete Strukturen in der Mathematik" at Bielefeld University, and I am grateful to SFB 343 for hospitality and support. I thank Rainer Schulze-Pillot and John S. Hsia for useful e-mail correspondence. I am grateful to Zeév Rudnick for useful discussions and help in analytic calculations.

1. Results of [BR] in the case of ternary quadratic forms

1.0. Let f be an indefinite ternary integral-matrix quadratic form

$$f(x_1, x_2, x_3) = \sum_{i,j=1}^3 a_{ij} x_i x_j, \quad a_{ij} \in \mathbf{Z}, \quad a_{ij} = a_{ji}$$

Let $q \in \mathbf{Z}, q \neq 0$. Let $D = \det(a_{ij})$. We assume that -qD is not a square.

Let $W = \mathbf{Q}^3$ and let X denote the affine variety in W defined by the equation f(x) = q, where $x = (x_1, x_2, x_3)$. We assume that X has a **Q**-point x^0 . Set G = Spin(W, f), the spinor group of f. Then G acts on W on the left, and X is an orbit (a homogeneous group) of G.

MIKHAIL BOROVOI

1.1. Rational points in adelic orbits. Let \mathbf{A} denote the adèle ring of \mathbf{Q} . The group $G(\mathbf{A})$ acts on $X(\mathbf{A})$; let $\mathcal{O}_{\mathbf{A}}$ be an orbit. We are interested whether $\mathcal{O}_{\mathbf{A}}$ has a \mathbf{Q} -rational point.

Let W' denote the orthogonal complement of x^0 in W, and let f' denote the restriction of f to W'. Let H be the stabilizer of x^0 in G, then H = Spin(W', f'). Since dim W' = 2, the group H is a one-dimensional torus.

We have det f' = D/q, so up to multiplication by a square det f' = qD. It follows that up to multiplication by a scalar, f' is equivalent to the quadratic form $u^2 + qDv^2$. Set $K = \mathbf{Q}(\sqrt{-qD})$, then K is a quadratic extension of \mathbf{Q} , because -qD is not a square. The torus H is anisotropic over \mathbf{Q} (because -qD is not a square), and H splits over K. Let $X_*(H_K)$ denote the cocharacter group of $H_K, X_*(H_K) = \operatorname{Hom}(\mathbb{G}_{m,K}, H_K)$; then $X_*(H_K) \simeq \mathbf{Z}$. The non-neutral element of $\operatorname{Gal}(K/\mathbf{Q})$ acts on $X_*(H_K)$ by multiplication by -1.

Let $\mathcal{O}_{\mathbf{A}}$ be an orbit of $G(\mathbf{A})$ in $X(\mathbf{A})$, $\mathcal{O}_{\mathbf{A}} = \prod \mathcal{O}_v$ where \mathcal{O}_v is an orbit of $G(\mathbf{Q}_v)$ in $X(\mathbf{Q}_v)$, v runs over the places of \mathbf{Q} , and \mathbf{Q}_v denotes the completion of \mathbf{Q} at v. We define local invariants $\nu_v(\mathcal{O}_v) = \pm 1$. If $\mathcal{O}_v = G(\mathbf{Q}_v) \cdot x^0$, then we set $\nu_v(\mathcal{O}_v) = +1$, if not, we set $\nu_v(\mathcal{O}_v) = -1$. Then $\nu_v(\mathcal{O}_v) = +1$ for almost all v. We define $\nu(\mathcal{O}_{\mathbf{A}}) = \prod \nu_v(\mathcal{O}_v)$ where $\mathcal{O}_{\mathbf{A}} = \prod \mathcal{O}_v$. Note that the local invariants $\nu_v(\mathcal{O}_v)$ depend on the choice of the rational point $x^0 \in X(\mathbf{Q})$; one can prove, however, that their product $\nu(\mathcal{O}_{\mathbf{A}})$ does not depend on x^0 .

Let $x \in X(\mathbf{A})$. We set $\nu(x) = \nu(G(\mathbf{A}) \cdot x)$. Then $\nu(x)$ takes values ± 1 ; it is a locally constant function on $X(\mathbf{A})$, because the orbits of $G(\mathbf{A})$ are open in $X(\mathbf{A})$.

For $x \in X(\mathbf{A})$ define $\delta(x) = \nu(x) + 1$. In other words, if $\nu(x) = -1$ then $\delta(x) = 0$, and if $\nu(x) = +1$ then $\delta(x) = 2$. Then δ is a locally constant function on $X(\mathbf{A})$.

Theorem 1.1.1. An orbit $\mathcal{O}_{\mathbf{A}}$ of $G(\mathbf{A})$ in $X(\mathbf{A})$ has a **Q**-rational point if and only if $\nu(\mathcal{O}_{\mathbf{A}}) = +1$.

Below we will deduce Theorem 1.1.1 from [BR], Thm. 3.6.

1.2. Proof of Theorem 1.1.1. For a torus T over a field k of characteristic 0 we define a finite abelian group C(T) as follows:

$$C(T) = (X_*(T_{\bar{k}})_{\operatorname{Gal}(\bar{k}/k)})_{\operatorname{tors}}$$

where \bar{k} is a fixed algebraic closure of k, $X_*(T_{\bar{k}})_{\operatorname{Gal}(\bar{k}/k)}$ denotes the group of coinvariants, and $(\cdot)_{\operatorname{tors}}$ denotes the torsion subgroup. If k is a number field and k_v is the completion of k at a place v, then we define $C_v(T) = C(T_{k_v})$. There is a canonical map $i_v: C_v(T) \to C(T)$ induced by an inclusion $\operatorname{Gal}(\bar{k}_v/k_v) \to \operatorname{Gal}(\bar{k}/k)$. These definitions were given for connected reductive groups (not only for tori) by Kottwitz [Ko], see also [BR], 3.4. Kottwitz writes A(T) instead of C(T).

We compute C(H) for our one-dimensional torus H over **Q**. Clearly

$$C(H) = (X_*(H_K)_{\operatorname{Gal}(K/\mathbf{Q})})_{\operatorname{tors}} = \mathbf{Z}/2\mathbf{Z}$$

We have $C_v(H) = 1$ if $K \otimes \mathbf{Q}_v$ splits, and $C_v(H) \simeq \mathbf{Z}/2\mathbf{Z}$ if $K \otimes \mathbf{Q}_v$ is a field. The map i_v is injective for any v.

We now define the local invariants $\kappa_v(\mathcal{O}_v)$ as in [BR], where \mathcal{O}_v is an orbit of $G(\mathbf{Q}_v)$ in $X(\mathbf{Q}_v)$. The set of orbits of $G(\mathbf{Q}_v)$ in $X(\mathbf{Q}_v)$ is in canonical bijection with $\log |U|^2(\mathbf{Q}_v) = U^2(\mathbf{Q}_v)$ of [Sel] 1.5.4 Core 1 of Prop. 26. Hence \mathcal{O}_v defines

a cohomology class $\xi_v \in H^1(\mathbf{Q}_v, H)$. The local Tate–Nakayama duality for tori defines a canonical homomorphism $\beta_v: H^1(\mathbf{Q}_v, H) \to C_v(H)$, see [Ko], Thm. 1.2. (Kottwitz defines the map β_v in a more general setting, when H is any connected reductive group over a number field.) The homomorphism β_v is an isomorphism for any v. We set $\kappa_v(\mathcal{O}_v) = \beta_v(\xi_v)$. Note that if $\mathcal{O}_v = G(\mathbf{Q}_v) \cdot x^0$, then $\xi_v = 0$ and $\kappa_v(\mathcal{O}_v) = 0$; if $\mathcal{O}_v \neq G(\mathbf{Q}_v) \cdot x^0$, then $\xi_v \neq 0$ and $\kappa_v(\mathcal{O}_v) = 1$.

We define the Kottwitz invariant $\kappa(\mathcal{O}_{\mathbf{A}})$ of an orbit $\mathcal{O}_{\mathbf{A}} = \prod \mathcal{O}_v$ of $G(\mathbf{A})$ in $X(\mathbf{A})$ by $\kappa(\mathcal{O}_{\mathbf{A}}) = \sum_v i_v(\kappa_v(\mathcal{O}_v))$. We identify C(H) with $\mathbf{Z}/2\mathbf{Z}$, and $C_v(H)$ with a subgroup of $\mathbf{Z}/2\mathbf{Z}$. With this identifications $\kappa(\mathcal{O}_{\mathbf{A}}) = \sum \kappa_v(\mathcal{O}_v)$.

We prefer the multiplicative rather than additive notation. Instead of $\mathbb{Z}/2\mathbb{Z}$ we consider the group $\{+1, -1\}$, and set

$$\nu_{v}(\mathcal{O}_{v}) = (-1)^{\kappa_{v}(\mathcal{O}_{v})}, \ \nu(\mathcal{O}_{\mathbf{A}}) = (-1)^{\kappa(\mathcal{O}_{\mathbf{A}})}.$$

Here $\nu_v(\mathcal{O}_v)$ and $\nu(\mathcal{O}_{\mathbf{A}})$ take values ± 1 . We have $\nu(\mathcal{O}_{\mathbf{A}}) = \prod \nu_v(\mathcal{O}_v)$. Since $\kappa_v(\mathcal{O}_v) = 0$ if and only if $\mathcal{O}_v = G(\mathbf{Q}_v) \cdot x^0$, we see that $\nu_v(\mathcal{O}_v) = +1$ if and only if $\mathcal{O}_v = G(\mathbf{Q}_v) \cdot x^0$. Hence our $\nu_v(\mathcal{O}_v)$ and $\nu(\mathcal{O}_{\mathbf{A}})$ coincide with $\nu_v(\mathcal{O}_v)$ and $\nu(\mathcal{O}_{\mathbf{A}})$ introduced in 1.1.

By Thm. 3.6 of [BR] an adelic orbit $\mathcal{O}_{\mathbf{A}}$ contains **Q**-rational points if and only if $\kappa(\mathcal{O}_{\mathbf{A}}) = 0$. With our multiplicative notation $\kappa(\mathcal{O}_{\mathbf{A}}) = 0$ if and only if $\nu(\mathcal{O}_{\mathbf{A}}) = +1$. Thus $\mathcal{O}_{\mathbf{A}}$ contains **Q**-points if and only if $\nu(\mathcal{O}_{\mathbf{A}}) = +1$. We have deduced Thm. 1.1.1 from [BR], Thm. 3.6. \Box

1.3. Tamagawa measure. We define a gauge form on X, i.e. a regular differential form $\omega \in \Lambda^2(X)$ without zeroes. Recall that X is defined by the equation f(x) = q. Choose a differential form $\mu \in \Lambda^2(W)$ such that $\mu \wedge df = dx_1 \wedge dx_2 \wedge dx_3$, where x_1, x_2, x_3 are the coordinates in $W = \mathbf{Q}^3$. Let $\omega = \mu|_X$, the restriction of μ to X. Then ω is a gauge form on X, cf. [BR], 1.3, and it does not depend on the choice of μ . The gauge form ω is G-invariant, because there exists a G-invariant gauge form on X, cf. [BR], 1.4, and a gauge form on X is unique up to a scalar multiple, cf. [BR], Cor. 1.5.4.

For any place v of \mathbf{Q} one associates with ω a local measure m_v on $X(\mathbf{Q}_v)$, cf. [We], 2.2. We show how to define a Tamagawa measure on $X(\mathbf{A})$, following [BR], 1.6.2.

We have by [BR], 1.8.1, $\mu_p(X) = m_p(X(\mathbf{Z}_p))$, where $\mu_p(X)$ is defined in the Introduction. By [We], Thm. 2.2.5, for almost all p we have $m_p(X(\mathbf{Z}_p)) = \#X(\mathbf{F}_p)$.

We compute $\#X(\mathbf{F}_p)$. The group $\mathrm{SO}(f)(\mathbf{F}_p)$ acts on $X(\mathbf{F}_p)$ with stabilizer $\mathrm{SO}(f')(\mathbf{F}_p)$, where $\mathrm{SO}(f')(\mathbf{F}_p)$ is defined for almost all p. This action is transitive by Witt's theorem. Thus $\#X(\mathbf{F}_p) = \#\mathrm{SO}(f)(\mathbf{F}_p)/\#\mathrm{SO}(f')(\mathbf{F}_p)$. By [A], III-6,

$$#SO(f)(\mathbf{F}_p) = p(p^2 - 1), \quad #SO(f')(\mathbf{F}_p) = p - \chi(p),$$

where $\chi(p) = -1$ if $f' \mod p$ does not represent 0, and $\chi(p) = +1$ if $f' \mod p$ represents 0. We have $\chi(p) = \left(\frac{-qD}{p}\right)$. We obtain for $p \nmid qD$

$$\#X(\mathbf{F}_p) = \frac{p(p^2 - 1)}{p - \chi(p)}, \quad \mu_p(X) = \frac{\#X(\mathbf{F}_p)}{p^2} = \frac{1 - 1/p^2}{1 - \chi(p)/p}.$$

For p|qD set $\chi(p) = 0$. We define

$$L_p(s,\chi) = (1 - \chi(p)p^{-s})^{-1}, \quad L(s,\chi) = \prod L_p(s,\chi)$$

where s is a complex variable. We set

$$\lambda_p = L_p(1,\chi)^{-1} = 1 - \frac{\chi(p)}{p}, \quad r = L(1,\chi)^{-1}.$$

Then the product $\prod_p (\lambda_p^{-1} \mu_p)$ converges absolutely, hence the family (λ_p) is a family of convergence factors in the sense of [We], 2.3. We define, as in [BR], 1.6.2, the measures

$$m_f = r^{-1} \prod_p (\lambda_p^{-1} m_p), \quad m = m_\infty m_f ,$$

then m_f is a measure on $X(\mathbf{A}_f)$ (where \mathbf{A}_f is the ring of finite adèles) and m is a measure on $X(\mathbf{A})$. We call m the Tamagawa measure on $X(\mathbf{A})$.

1.4. Counting integer points. For T > 0 set $X(\mathbf{R})^T = \{x \in X(\mathbf{R}) : |x| \le T\}.$

Theorem 1.4.1.

$$N(T,X) \sim \int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm.$$

In other words,

(1.4.1)
$$N(T,X) \sim 2m(\{x \in X(\mathbf{R})^T \times X(\hat{\mathbf{Z}}) : \nu(x) = +1\}).$$

Theorem 1.4.1 follows from [BR], Thm. 5.3 (cf. [BR], 6.4 and [BR], Def. 2.3). For comparison note that

(1.4.2)
$$m(X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})) = m_{\infty}(X(\mathbf{R})^T)m_f(X(\hat{\mathbf{Z}})) = \mu_{\infty}(T,X)\mathfrak{S}(X),$$

cf. [BR], 1.8.

The following lemma will be used in the proof of Theorem 0.1.

Lemma 1.4.2. Assume that there exists $y \in X(\mathbf{R} \times \hat{\mathbf{Z}})$ such that $\nu(y) = +1$. Then the set $X(\mathbf{Z})$ is infinite.

Proof. Since ν is a locally constant function on $X(\mathbf{A})$, there exists an open subset $\mathcal{U}_f \in X(\hat{\mathbf{Z}})$ and an orbit \mathcal{U}_∞ of $G(\mathbf{R})$ in $X(\mathbf{R})$ such that $\nu(x) = +1$ for all $x \in \mathcal{U}_\infty \times \mathcal{U}_f$. Set $\mathcal{U}_\infty^T = \{x \in \mathcal{U}_\infty : |x| \leq T\}$, then $m_\infty(\mathcal{U}_\infty^T) \to \infty$ as $T \to \infty$. We have

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm \ge \int_{\mathcal{U}_{\infty}^T \times \mathcal{U}_f} \delta(x) dm = 2m_{\infty}(\mathcal{U}_{\infty}^T) m_f(\mathcal{U}_f) \ .$$

Since $2m_{\infty}(\mathcal{U}_{\infty}^T)m_f(\mathcal{U}_f) \to \infty$ as $T \to \infty$, we see that

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm \to \infty \text{ as } T \to \infty,$$

and by Theorem 1.4.1 $N(T, X) \to \infty$. Hence $X(\mathbf{Z})$ is infinite. \Box

1 E The constant of Here we prove the following regult

Proposition 1.5.1.

$$N(T,X) \sim c_X \mathfrak{S}(X) \mu_{\infty}(T,X) \text{ as } T \to \infty$$

with some constant c_X , $0 \le c_X \le 2$.

Proof. If $X(\mathbf{R})$ has two connected components, then by Theorem 0.2 (which we will prove in Sect. 3 below), $N(T, X) \sim \mathfrak{S}(X)\mu_{\infty}(T, X)$, so the proposition holds with $c_X = 1$.

If $X(\mathbf{R})$ has one connected component, then $X(\mathbf{R})$ consists of one $G(\mathbf{R})$ -orbit and $\nu_{\infty}(X(\mathbf{R})) = +1$. For an orbit $\mathcal{O}_f = \prod \mathcal{O}_p$ of $G(\mathbf{A}_f)$ in $X(\mathbf{A}_f)$ we set $\nu_f(\mathcal{O}_f) = \prod_p \nu_p(\mathcal{O}_p)$. We regard ν_f as a locally constant function on $X(\mathbf{A}_f)$ taking values ± 1 . We have

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm = 2m_{\infty}(X(\mathbf{R})^T) m_f(X(\hat{\mathbf{Z}})_+)$$

where $X(\hat{\mathbf{Z}})_+ = \{x_f \in X(\hat{\mathbf{Z}}) : \nu_f(x_f) = +1\}$. Set $c_X = 2m_f(X(\hat{\mathbf{Z}})_+)/m_f(X(\hat{\mathbf{Z}}))$, then $0 \le c_X \le 2$ and

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm = c_X m_\infty(X(\mathbf{R})^T) m_f(X(\hat{\mathbf{Z}})) = c_X \mu_\infty(T, X) \mathfrak{S}(X).$$

Using Theorem 1.4.1, we see that

$$N(T,X) \sim c_X \mu_{\infty}(T,X) \mathfrak{S}(X)$$
 as $T \to \infty$.

2. An example of
$$c_X = 0$$

Let

$$f_1(x_1, x_2, x_3) = -9x_1^2 + 2x_1x_2 + 7x_2^2 + 2x_3^2, \ q = 1.$$

This example was mentioned by Siegel and later mentioned in [BR], 6.4.1. Here we provide a detailed exposition.

Consider the variety X defined by the equation $f_1(x) = q$. We have $f_1(-\frac{1}{2}, \frac{1}{2}, 1) = 1$. It follows that f_1 represents 1 over **R** and over \mathbf{Z}_p for p > 2.

We have $f_1(4, 1, 1) = -127 \equiv 1 \pmod{2^7}$. We prove that f_1 represents 1 over \mathbb{Z}_2 . Define a polynomial of one variable $F(Y) = f_1(4, 1, Y) - 1$, $F \in \mathbb{Z}_2[Y]$. Then $F(1) = -2^7$, $|F(1)|_2 = 2^{-7}$, F'(Y) = 4Y, $|F'(1)^2|_2 = 2^{-4}$, $|F(1)|_2 < |F'(1)^2|_2$. By Hensel's lemma (cf. [La], II-§2, Prop. 2) F has a root in \mathbb{Z}_2 . Thus f_1 represents 1 over \mathbb{Z}_2 .

Now we prove that f_1 does not represent 1 over **Z**. I know the following elementary proof from D. Zagier.

We prove the assertion by contradiction. Assume on the contrary that

$$-9x_1^2 + 2x_1x_2 + 7x_2^2 + 2x_3^2 = 1$$
for some $x_1, x_2, x_3 \in \mathbf{Z}.$

We may write this equation as follows:

$$2\pi^2$$
 1 (m m)² + 9(m m)(m + m)

MIKHAIL BOROVOI

The left hand side is odd, hence $x_1 - x_2$ is odd and therefore $x_1 + x_2$ is odd. We have $(x_1 - x_2)^2 \equiv 1 \pmod{8}$. Hence the right hand side is congruent to 1 (mod 8). We see that x_3 is odd, hence $2x_3^2 - 1 \equiv 1 \pmod{16}$. But

$$8(x_1 - x_2)(x_1 + x_2) \equiv 8 \pmod{16}.$$

It follows that

$$(x_1 - x_2)^2 \equiv 9 \pmod{16}$$

 $x_1 - x_2 \equiv \pm 3 \pmod{8}.$

Therefore $x_1 - x_2$ must have a prime factor $p \equiv \pm 3 \pmod{8}$. Hence $2x_3^2 - 1$ has a prime factor $p \equiv \pm 3 \pmod{8}$. On the other hand, if $p|(2x_3^2 - 1)$, then

$$2x_3^2 \equiv 1 \pmod{p}$$

and 2 is a square modulo p, $\left(\frac{2}{p}\right) = 1$. By the quadratic reciprocity law $p \equiv \pm 1$ (mod 8). Contradiction. We have proved that f_1 does not represent 1 over \mathbf{Z} , hence N(T, X) = 0 for all T.

On the other hand,

$$\mathfrak{S}(X)\mu_{\infty}(T,X) = m_f(X(\mathbf{\hat{Z}}))m_{\infty}(X(\mathbf{R})^T).$$

Since $X(\hat{\mathbf{Z}})$ is a non-empty open subset in $X(\mathbf{A}_f)$, $m_f(X(\hat{\mathbf{Z}})) > 0$. Now the measure $m_{\infty}(X(R)^T) \to \infty$ as $T \to \infty$. Hence $\mathfrak{S}(X)\mu_{\infty}(T,X) \to \infty$ as $T \to \infty$, and thus $c_X = 0$.

3. Proof of Theorem 0.1

Lemma 3.1. Let k be a field, $\operatorname{char}(k) \neq 2$, and let V be a finite-dimensional vector space over k. Let f be a non-degenerate quadratic form on V. Let $u \in \operatorname{GL}(V)(k)$, $f' = u^*f$. Then the map $y \mapsto uy: V \to V$ takes the orbits of $\operatorname{Spin}(f)(k)$ in V to the orbits of $\operatorname{Spin}(f')(k)$.

Proof. Let $x \in V$, $f(x) \neq 0$. The reflection (symmetry) $r_x = r_{f,x} \colon V \to V$ is defined by

$$r_x(y) = y - \frac{2B(x,y)}{f(x)}x, \quad y \in V,$$

where B is the symmetric bilinear form on V associated with f. Every $s \in SO(f)(k)$ can be written as

 $(3.1) s = r_{x_1} \cdots r_{x_l}$

cf. [OM], Thm. 43:3. The spinor norm $\theta(s)$ of s is defined by

$$\theta(s) = f(x_1) \cdots f(x_l) \pmod{k^{*2}} \in k^*/k^{*2}$$

and it does not depend on the choice of the representation (3.1), cf. [OM], §55. Let $\Theta(f)$ denote the image of Spin(f)(k) in SO(f)(k). Then $s \in \text{SO}(f)(k)$ is contained in $\Theta(f)$ if and only if $\theta(s) = 1$, cf. [Se], III-3.2 or [Ca], Ch. 10, Thm. 3.3.

Now let u, f' be as above. Then $r_{f',ux} = ur_{f,x}u^{-1}$, f'(ux) = f(x), and so $\theta_{f'}(usu^{-1}) = \theta_f(s)$. We conclude that $u\Theta(f)u^{-1} = \Theta(f')$ and that the map $y \mapsto uy$ takes the orbits of $\Theta(f)$ in V to the orbits of $\Theta(f')$. \Box

Let f, f' be integral-matrix quadratic forms on \mathbb{Z}^n and assume that f' is in the genus of f. Then there exists $u \in \operatorname{GL}_n(\mathbb{R} \times \hat{\mathbb{Z}})$ such that $f'(x) = f(u^{-1}x)$ for $x \in \mathbb{A}^n$. Let $q \in \mathbb{Z}, q \neq 0$. Let X denote the affine quadric f(x) = q, and X' denote the sum drip f'(x) = g.

Lemma 3.2. The map $x \mapsto ux: \mathbf{A}^n \to \mathbf{A}^n$ takes $X(\mathbf{R} \times \mathbf{Z})$ to $X'(\mathbf{R} \times \mathbf{Z})$ and takes orbits of $\text{Spin}(f)(\mathbf{A})$ in $X(\mathbf{A})$ to orbits of $\text{Spin}(f')(\mathbf{A})$ in $X'(\mathbf{A})$.

Proof. Let A denote the matrix of f, and A' denote the matrix of f'. We have

$$(u^{-1})^t A u^{-1} = A$$
$$A = u^t A' u .$$

The variety X is defined by the equation $x^t A x = q$, and X' is defined by $x^t A' x = q$. One can easily check that the map $x \mapsto ux$ takes $X(\mathbf{R} \times \hat{\mathbf{Z}})$ to $X'(\mathbf{R} \times \hat{\mathbf{Z}})$ and $X(\mathbf{A})$ to $X'(\mathbf{A})$.

In order to prove that the map $x \mapsto ux: X(\mathbf{A}) \to X'(\mathbf{A})$ takes the orbits of $\operatorname{Spin}(f)(\mathbf{A})$ to the orbits of $\operatorname{Spin}(f')(\mathbf{A})$, it suffices to prove that the map $x \mapsto u_v x: X(\mathbf{Q}_v) \to X'(\mathbf{Q}_v)$ takes the orbits of $\operatorname{Spin}(f)(\mathbf{Q}_v)$ to the orbits of $\operatorname{Spin}(f')(\mathbf{Q}_v)$ for every v, where u_v is the v-component of u. This last assertion follows from Lemma 3.1. \Box

Proposition 3.3. Let f' and q be as in Theorem 0.1, in particular f' represents q over \mathbf{Z}_v for any v (we set $\mathbf{Z}_{\infty} = \mathbf{R}$), but not over \mathbf{Z} . Let X' be the quadric defined by f'(x) = q. Then $X'(\mathbf{R} \times \hat{\mathbf{Z}})$ is contained in one orbit of $\operatorname{Spin}(f')(\mathbf{A})$.

Proof. Set G' = Spin(f'). We prove that $X'(\mathbf{Z}_v)$ is contained in one orbit of $G'(\mathbf{Q}_v)$ for every v by contradiction. Assume on the contrary that for some $v X'(\mathbf{Z}_v)$ has nontrivial intersection with two orbits of $G'(\mathbf{Q}_v)$. Then ν_v takes both values +1 and -1 on $X'(\mathbf{Z}_v)$. It follows that ν takes both values +1 and -1 on $X'(\mathbf{R} \times \hat{\mathbf{Z}})$. Hence by Lemma 1.4.2 X' has infinitely many \mathbf{Z} -points. This contradicts to the assumption that f' does not represent q over \mathbf{Z} . \Box

Proof of Theorem 0.1. Let $u \in \operatorname{GL}_3(\mathbf{R} \times \hat{\mathbf{Z}})$ be such that $f'(x) = f(u^{-1}x)$. Let X, X' be as in the beginning of this section, in particular X' has no \mathbf{Z} -points. By Prop. 3.3 $X'(\mathbf{R} \times \hat{\mathbf{Z}})$ is contained in one orbit of $\operatorname{Spin}(f')(\mathbf{A})$. It follows from Lemma 3.2 that $X(\mathbf{R} \times \hat{\mathbf{Z}})$ is contained in one orbit of $\operatorname{Spin}(f)(\mathbf{A})$. Since f represents q over \mathbf{Z} , this orbit has \mathbf{Q} -rational points, and ν equals +1 on $X(\mathbf{R} \times \hat{\mathbf{Z}})$. Thus δ equals 2 on $X(\mathbf{R} \times \hat{\mathbf{Z}})$, and by Formulas (1.4.1) and (1.4.2) $N(T, X) \sim 2\mathfrak{S}(X)\mu_{\infty}(T, X)$. \Box

4. Proof of Theorem 0.2

We prove Theorem 0.2. We define an involution τ_{∞} of $X(\mathbf{R})$ by $\tau_{\infty}(x) = -x$, $x \in X(\mathbf{R}) \subset \mathbf{R}^3$. Since f(x) = f(-x), τ_{∞} is well defined, i.e takes $X(\mathbf{R})$ to itself. Since |-x| = |x|, τ_{∞} takes $X(\mathbf{R})^T$ to itself. We define an involution τ of $X(\mathbf{A})$ by defining τ as τ_{∞} on $X(\mathbf{R})$ and as 1 on $X(\mathbf{Q}_p)$ for all prime p. Then τ respects the Tamagawa measure m on $X(\mathbf{A})$.

By assumption $X(\mathbf{R})$ has two connected components. These are two orbits of $\operatorname{Spin}(f)(\mathbf{R})$. The involution τ_{∞} of $X(\mathbf{R})$ interchanges these two orbits. Thus we have

(4.1)
$$\nu_{\infty}(\tau_{\infty}(x_{\infty})) = -\nu_{\infty}(x_{\infty}) \text{ for all } x_{\infty} \in X(\mathbf{R})$$
$$\nu(\tau(x)) = -\nu(x) \text{ for all } x \in X(\mathbf{A})$$

Let $X(\mathbf{R})_1$ and $X(\mathbf{R})_2$ be the two connected components of $X(\mathbf{R})$. Set

$$\mathbf{V}(\mathbf{p})^T = \mathbf{V}(\mathbf{p}) \cap \mathbf{V}(\mathbf{p})^T = \mathbf{V}(\mathbf{p})^T = \mathbf{V}(\mathbf{p}) \cap \mathbf{V}(\mathbf{p})^T$$

Then τ interchanges $X(\mathbf{R})_1^T \times X(\hat{\mathbf{Z}})$ and $X(\mathbf{R})_2^T \times X(\hat{\mathbf{Z}})$. From Formula (4.1) we have

$$\int_{X(\mathbf{R})_1^T \times X(\hat{\mathbf{Z}})} \nu(x) dm = -\int_{X(\mathbf{R})_2^T \times X(\hat{\mathbf{Z}})} \nu(x) dm,$$

hence

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \nu(x) dm = 0.$$

Since $\delta(x) = \nu(x) + 1$, we obtain

$$\int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm = \int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} dm = m(X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})) = \mathfrak{S}(X)\mu_{\infty}(T,X).$$

By Theorem 1.4.1

$$N(T,X) \sim \int_{X(\mathbf{R})^T \times X(\hat{\mathbf{Z}})} \delta(x) dm.$$

Thus $N(T,X) \sim \mathfrak{S}(X)\mu_{\infty}(T,X)$ as $T \to \infty$, i.e. $c_X = 1$. \Box

References

- [A] E. Artin, *Geometric Algebra*, Interscience Publishers, New York, 1957.
- [BR] M. Borovoi, Z. Rudnick, Hardy-Littlewood varieties and semisimple groups, Invent. Math. 111 (1995), 37–66.
- [Ca] J. W. S. Cassels, *Rational Quadratic Forms*, Academic Press, London, 1978.
- [CS] J. H. Conway and N. J. A. Sloane, Sphere Packings, Lattices and Groups, 2nd edition, Springer-Verlag, New York, 1993.
- [**Da**] H. Davenport, Analytic Methods for Diophantine Equations and Diophantine Inequalities, Ann Arbor Publishers, Ann Arbor, Michigan, 1962.
- [DRS] W. Duke, Z. Rudnick and P. Sarnak, Density of integer points on affine homogeneous varieties, Duke Math. Jour. 71 (1993), 143–179.
- [EM] A. Eskin and C. McMullen, Mixing, counting, and equidistribution in Lie groups, Duke Math. J. 71 (1993), 181–209.
- [EMS] A. Eskin, S. Mozes and N. Shah, Unipotent flows and counting lattice points on homogeneous spaces, Ann. of Math. (2) 143 (1996), 253–299.
- [Est] T. Estermann, A new application of the Hardy-Littlewood-Kloosterman method, Proc. London Math. Soc. 12 (1962), 425–444.
- [Kn] M. Kneser, Darstellungsmaße indefiniter quadratischer Formen, Math. Z. 77 (1961), 188–194.
- [Ko] R. E. Kottwitz, Stable trace formula: elliptic singular terms, Math. Ann. 275 (1986), 365–399.
- [La] S. Lang, Algebraic Number Theory, Addison-Wesley, Reading, Mass., 1970.
- [OM] O. T. O'Meara, Introduction to Quadratic Forms, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
- [Se] J.-P. Serre, Cohomologie galoisienne, Lecture Notes in Math., vol. 5, Springer-Verlag, Berlin-Heidelberg-New York, 1965.
- [Si] C. L. Siegel, Uber die analytische Theorie der quadratischen Formen II, Ann. of Math.
 37 (1936), 230–263.
- [We] A. Weil, Adeles and Algebraic Groups, Birkhäuser, Boston, 1982.

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNI-VERSITY, 69978 TEL AVIV, ISRAEL

E-mail address: borovoi@math.tau.ac.il