# ON THE CAPACITY REGION OF MULTIPLE ACCESS ADDER CHANNEL

SHAOHUI ZHAI

ABSTRACT. We compute the capacity region of the *s*-user Multiple Access Adder Channel and obtain an explicit description of this region as only one polyhedron, instead of a convex hull of infinitely many polyhedrons. We obtain this by proving a conjecture raised by Lindström in 1975, which in turn follows from certain convexity results, both continuous and discrete.

### 1. Preliminaries

### 1.1. Multiple Access Channel.

**Definition 1.** An s-user Multiple Access Channel (MAC) consists of s input alphabets,  $\mathcal{X}_1, \mathcal{X}_2, \ldots, \mathcal{X}_s$ , an output alphabet  $\mathcal{Y}$ , and a probability transition matrix  $P(y \mid x_1, x_2, \ldots, x_s)$ , which defines the probability of receiving the output symbol y given input symbols  $x_1, x_2, \ldots, x_s$ .

**Definition 2.** An  $(M_1, M_2, \ldots, M_s; n)$  code for the s-user MAC consists of s encoding functions  $e_1, \ldots, e_s$  which encode each message  $m_i \in [M_i] = \{1, 2, \ldots, M_i\}$  into a string of length n in the input alphabet  $\mathcal{X}_i$ :

$$e_i: [M_i] \to \mathcal{X}_i^n m_i \mapsto e_i(m_i) = (e_i^{(1)}(m_i), e_i^{(2)}(m_i), \dots, e_i^{(n)}(m_i));$$

and a decoding function:

$$d: \qquad \mathcal{Y}^n \quad \to \quad [M_1] \times [M_2] \times \cdots \times [M_s] (y_1, \dots, y_n) \quad \mapsto \quad (\hat{m}_1, \dots, \hat{m}_s).$$

The model of communication is as follows (see Fig. 1). Each user *i* chooses a label  $m_i$  uniformly from its message set  $[M_i]$  and sends the codeword  $e_i(m_i)$ . All the users send their codewords simultaneously, symbol by symbol. After *n* transmissions, a sequence of *n* output symbols  $(y_1, y_2, \ldots, y_n)$  is received according to the transition probability matrix. This is then decoded by the decoding function *d* into a set of messages  $(\hat{m}_1, \ldots, \hat{m}_s)$ . If the decoded messages are identical to the messages sent, the communication has succeeded. Otherwise, we say an *error* has occurred.



Fig. 1. Multiple Access Channel

**Definition 3.** The average probability of error for an  $(M_1, M_2, \ldots, M_s; n)$  code is

$$P_e^{(n)} = \frac{1}{M_1 M_2 \cdots M_s} \sum_{\substack{(m_1, \dots, m_s) \\ \in [M_1] \times \dots \times [M_s]}} P_r(d(Y^n) \neq (m_1, \dots, m_s) \mid (m_1, \dots, m_s) \text{ sent}).$$

We use *rate* to measure the efficiency of an  $(M_1, \ldots, M_s; n)$  code.

**Definition 4.** The rate tuple  $(R_1, \ldots, R_s)$  of an  $(M_1, \ldots, M_s; n)$  code is defined as

$$R_i = \frac{\log_2 M_i}{n}$$
 bits per transmission in the *i*-th sender.

**Definition 5.** A rate tuple  $(R_1, R_2, \ldots, R_s)$  is achievable if there exists a sequence of  $(2^{nR_1}, 2^{nR_2}, \ldots, 2^{nR_s}; n)$  codes with  $P_e^{(n)} \to 0$  as  $n \to \infty$ .

**Definition 6.** The capacity region of the MAC is the closure of the set of all the achievable rate tuples.

**Definition 7.** A Multiple Access Channel is called Multiple Access Adder Channel (MAAC) when the input alphabets are  $\mathcal{X}_1 = \mathcal{X}_2 = \ldots = \mathcal{X}_s = \{0, 1\}$  and the channel output is the sum of the s input symbols:  $Y = X_1 + X_2 + \cdots + X_s$ , so that the output alphabet is  $\mathcal{Y} = \{0, 1, 2, \ldots, s\}$ .

This is a deterministic channel in that a given s-tuple of inputs always yields the same output. 1.2. Capacity Region. A simple characterization of the capacity region for the two-user MAC was first presented by Ahlswede [1] in 1973. Ulrey [2] generalized it to the MAC with more than two users.

Let  $A \subseteq [s] = \{1, 2, ..., s\}$ ,  $A^c$  denote the complement of A in [s],  $R(A) \stackrel{\Delta}{=} \sum_{i \in A} R_i, X(A) \stackrel{\Delta}{=} \{X_i : i \in A\}$ , and I(X; Y|Z) the conditional mutual information of random variable X and Y given random variable Z.

**Theorem 1** (Ulrey). The capacity region of the s-user Multiple Access Channel is the closure of the convex hull of the rate tuples  $(R_1, R_2, \ldots, R_s)$  satisfying

 $R(A) \leq I(X(A); Y|X(A^c))$  for all the subsets  $A \subseteq [s]$ 

for some product distribution  $p_1(x_1)p_2(x_2)\dots p_s(x_s)$ .

## 2. Results

For the s-user MAAC, we now present a finite description of its capacity region. Let  $\mathcal{B}(n,p)$  be the binomial distribution with n independent trials and probability p of a success in each trial, i.e., if  $X \sim \mathcal{B}(n,p)$ , then  $Pr(X = k) = {n \choose k} p^k (1-p)^{n-k}$ . Let  $H_n(p)$  denote the Shannon entropy of  $\mathcal{B}(n,p)$ , i.e.,

$$H_n(p) \stackrel{\Delta}{=} -\sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k} \cdot \log\left(\binom{n}{k} p^k (1-p)^{n-k}\right).$$

Now we state our main theorem.

**Theorem 2.** The capacity region for the s-user Multiple Access Adder Channel is the set

$$\{(R_1, R_2, \dots, R_s): R(A) \le H_{|A|}(\frac{1}{2}) \text{ for all } A \subseteq [s]\}.$$

This region is only one polyhedron. For example, the capacity region for the 2-user adder channel is

$$\{(R_1, R_2): 0 \le R_1 \le 1; 0 \le R_2 \le 1; 0 \le R_1 + R_2 \le 1/2\};$$

the capacity region for the 3-user adder channel is

$$\{ (R_1, R_2, R_3) : R_1 \le 1; R_2 \le 1; R_3 \le 1; R_1 + R_2 \le 1/2; R_1 + R_3 \le 1/2; R_2 + R_3 \le 1/2; R_1 + R_2 + R_3 \le (3 - \frac{3}{4} \log 3) \}.$$

A key part of our proof of Theorem 2 is the following result, conjectured by Lindström [3]:

**Theorem 3.** Let  $X_1, X_2, \ldots, X_n$  be independent random variables = 0 or 1 with  $Pr(X_i = 1) = p_i$  and  $Pr(X_i = 0) = 1 - p_i$ , where  $0 \le p_i \le 1$  for  $i = 1, 2, \ldots, n$ .

The Shannon entropy of the random variable  $X_1 + X_2 + \cdots + X_n$  is then maximal when  $p_i = 1/2$  for all  $i = 1, 2, \ldots, n$ , i.e.,

$$\max_{0 \le p_1, p_2, \dots, p_n \le 1} H(\sum_{i=1}^n X_i) = H_n(1/2)$$

**Remark.** The result in Theorem 2 has apparently obtained by Liao [4] in his Ph.D. dissertation in 1972, but his proof is never published. Our proof will include proving Theorem 3, conjectured in 1976.

### 3. Proof of Results

3.1. Theorem 3 implies Theorem 2. To compute the capacity region for the *s*-user adder channel, by Theorem 1, we need to maximize

$$I(X(A); Y|X(A^c))$$

for any  $A \subseteq [s]$ .

Since  $Y = X_1 + X_2 + \cdots + X_s$  is determined by X(A) and  $X(A^c)$  for all  $A \subseteq [s]$ , the conditional Shannon entropy of Y given X(A) and  $X(A^c)$  is zero, i.e.,  $H(Y|X(A), X(A^c)) = 0$ , so

$$\begin{split} I(X(A);Y|X(A^{c})) &= H\left(Y|X(A^{c})\right) - H\left(Y|X(A), X(A^{c})\right) \\ &= H\left(Y|X(A^{c})\right) \\ &= \sum_{x} H\left(Y|X(A^{c})=x\right) \cdot P_{r}\left(X(A^{c})=x\right) \\ &= \sum_{x} H\left(\sum_{i \in A} X_{i} + \sum_{i \in A^{c}} X_{i} \mid X(A^{c})=x\right) \cdot P_{r}\left(X(A^{c})=x\right) \\ &= \sum_{x} H\left(\sum_{i \in A} X_{i}\right) \cdot P_{r}\left(X(A^{c})=x\right) \\ &= H\left(\sum_{i \in A} X_{i}\right) \cdot \sum_{x} P_{r}\left(X(A^{c})=x\right) \\ &= H\left(\sum_{i \in A} X_{i}\right). \end{split}$$

So,

$$\max_{0 \le p_1, p_2, \dots, p_s \le 1} I(X(A); Y | X(A^c)) = \max_{\{i \in A: \ 0 \le p_i \le 1\}} H(\sum_{i \in A} X_i).$$

Thus, if Theorem 3 is true, then  $H\left(\sum_{i\in A} X_i\right)$  is maximized when  $p_i = 1/2, i \in A$ for all  $A \subseteq [s]$ . But when  $p_i = 1/2, i \in A$ , the random variable  $\sum_{i\in A} X_i$  has the binomial distribution with parameter (|A|, 1/2), i.e.,

$$\max_{0 \le p_1, p_2, \dots, p_s \le 1} I(X(A); Y | X(A^c)) = \max_{\{i \in A: \ 0 \le p_i \le 1\}} H(\sum_{i \in A} X_i) = H_{|A|}(1/2)$$

for all  $A \subseteq [s]$ . This, together with Theorem 1, implies Theorem 2.

3.2. **Proof of Theorem 3 (Lindström's conjecture).** Let  $p_i = P_r(X_i = 1), 0 \le p_i \le 1$ , and  $q_i = 1 - p_i$ , for i = 1, 2, ..., n. Let  $\mathbf{p} = (p_1, p_2, ..., p_n)$ ,  $\mathbf{p}_i = (p_1, ..., p_{i-1}, p_{i+1}, ..., p_n)$ , and more generally, for  $0 \le r \le n$ ,  $\mathbf{p}_{\hat{i}_1, \hat{i}_2, ..., \hat{i}_r} = (p_{j_1}, p_{j_2}, ..., p_{j_{n-r}})$ , where  $1 \le j_1 < j_2 < \cdots < j_{n-r} \le n$  and  $\{j_1, j_2, ..., j_{n-r}\} = [n] \setminus \{i_1, i_2, ..., i_r\}$ .

We need the following result of Mateev [5].

**Lemma 1** (Mateev). The Shannon entropy of the binomial distribution  $\mathcal{B}(n, p)$  reaches its maximum when p = 1/2, that is,

$$\max_{0 \le p \le 1} H_n(p) = H_n(1/2).$$

Let  $Z_{\mathbf{p}} = \sum_{i=1}^{n} X_i$ . We want to show that the Shannon entropy  $H(Z_{\mathbf{p}})$  of  $Z_{\mathbf{p}}$  is parimized when n = 1/2 for all  $i = 1, 2, \dots, n$  i.e.

maximized when  $p_i = 1/2$  for all i = 1, 2, ..., n, i.e.,

$$\max_{\leq p_1, p_2, \dots, p_n \leq 1} H(Z_{\mathbf{p}}) = H_n(1/2).$$

Notice that, when  $p_1 = p_2 = \cdots = p_n = p$ ,  $Z_{\mathbf{p}}$  has the binomial distribution with parameter (n, p), and  $H(Z_{\mathbf{p}}) = H_n(p)$ . By Lemma 1,  $\max_{0 \le p \le 1} H_n(p) = H_n(1/2)$ . Therefore, to prove Theorem 3, we only need to show that

(1) 
$$\max_{0 \le p_1, p_2, \dots, p_n \le 1} H(Z_{\mathbf{p}}) = \max_{0 \le p_1 = p_2 = \dots = p_n \le 1} H(Z_{\mathbf{p}})$$

We introduce the following lemma,

**Lemma 2.** Let  $\mathbf{p}' = (p'_1, \ldots, p'_n)$  be any point in  $\mathbf{R}^n$  with  $0 \le p'_1, \ldots, p'_n \le 1$ . Let  $1 \le l < m \le n$ , and  $\mathbf{\vec{u}} = (0, \ldots, \overset{l}{1}, 0, \ldots, \overset{m}{-1}, 0, \ldots, 0)$ . Then the function  $H(Z_{\mathbf{p}})$  restricted to the line  $\mathbf{p} = \mathbf{p}' + t \cdot \mathbf{\vec{u}}$ , where t is any real number such that  $\mathbf{0} \le \mathbf{p} \le \mathbf{1}$ , has a unique maximum at  $p_l = p_m$ .

Theorem 3 follows immediately from this lemma (which we will prove in the next paragraph). Indeed, let  $\mathbf{p}^* = (p_1^*, p_2^*, \ldots, p_n^*)$  be any maximal point of  $H(Z_{\mathbf{p}})$ . Then for any  $1 \leq l < m \leq n$ ,  $\mathbf{p}^*$  is also the maximal point of the function  $H(Z_{\mathbf{p}})$  restricted to the line  $\mathbf{p} = \mathbf{p}^* + t \cdot \vec{\mathbf{u}}$ . By Lemma 2,  $p_l^* = p_m^*$ , for any l, m, which proves (1).

**Proof of Lemma 2.** We first notice that  $H(Z_{\mathbf{p}})$  is symmetric in  $p_1, p_2$ ,

...,  $p_n$ , so the function  $H(Z_{\mathbf{p}})$  restricted to the line  $\mathbf{p} = \mathbf{p}' + t \cdot \mathbf{\vec{u}}$  is symmetric in  $p_l$ and  $p_m$ . We will prove that the function  $H(Z_{\mathbf{p}})$  restricted to the line  $\mathbf{p} = \mathbf{p}' + t \cdot \mathbf{\vec{u}}$ is concave by showing that the second directional derivative of  $H(Z_{\mathbf{p}})$  in the direction  $\mathbf{\vec{u}}$  is strictly negative. The concavity guarantees a unique maximum; by the symmetry, this maximum occurs at  $p_l = p_m$ .

Thus, to prove Lemma 2, and hence Theorem 3 and Theorem 2, we must show:

$$D_{\vec{\mathbf{u}}}^2 H(Z_{\mathbf{p}}) < 0$$
  
  $\forall \mathbf{p} = (p_1, p_2, \dots, p_n) \text{ with } 0 \le p_1, p_2, \dots, p_n \le 1,$ 

where  $D_{\vec{u}} = \frac{\partial}{\partial p_l} - \frac{\partial}{\partial p_m}$ . The probability distribution of  $Z_{\mathbf{p}}$  is described by

$$b(k,\mathbf{p}) \stackrel{\Delta}{=} \Pr(Z_{\mathbf{p}} = k) = \begin{cases} \sum_{A \subseteq [n] \ |A| = k} \prod_{i \in A} p_i \prod_{j \in A^c} q_j & \text{for } 0 \le k \le n \\ 0 & \text{otherwise,} \end{cases}$$

where  $A^c$  is the complement of A in [n]. Particularly,

$$b(0, \mathbf{p}) = \prod_{j=1}^{n} q_j, \qquad b(n, \mathbf{p}) = \prod_{i=1}^{n} p_i.$$

In general, the probability function  $Pr(Z_{\mathbf{p}_{\hat{i}_1,\hat{i}_2,\ldots,\hat{i}_r}} = k)$  of  $Z_{\mathbf{p}_{\hat{i}_1,\hat{i}_2,\ldots,\hat{i}_r}} \stackrel{(\Delta)}{=}$  $\sum_{\substack{1\leq i\leq n\\ i\neq i_1,\ldots,i_r}} X_i \big)$  is

$$b(k, \mathbf{p}_{\hat{i}_1, \hat{i}_2, \dots, \hat{i}_r}) = \begin{cases} \sum_{\substack{A \subseteq [n] \setminus \{i_1, \dots, i_r\} \\ |A| = k}} \prod_{i \in A} p_i \prod_{j \in A^c} q_j & \text{for } 0 \le k \le (n-r) \\ 0 & \text{otherwise,} \end{cases}$$

where  $A^c$  is the complement of A in  $[n] \setminus \{i_1, i_2, \ldots, i_r\}$ .

The Shannon entropy of  $Z_{\mathbf{p}}$  is

$$H(Z_{\mathbf{p}}) = -\sum_{k=0}^{n} b(k, \mathbf{p}) \log b(k, \mathbf{p}).$$

To compute the derivatives of  $H(Z_{\mathbf{p}})$ , we will repeatedly use the following formulas:  $\mathfrak{A}L(1)$ --- )

$$\frac{\partial b(k, \mathbf{p})}{\partial p_l} = \sum_{\substack{l \in A \subseteq [n] \\ |A|=k}} \prod_{\substack{i \neq l \\ i \in A}} p_i \prod_{\substack{j \in A^c}} q_j - \sum_{\substack{l \notin B \subseteq [n] \\ |B|=k}} \prod_{i \in B} p_i \prod_{\substack{j \neq l \\ |B|=k}} q_j$$

$$= \sum_{\substack{A' \subseteq [n] \setminus \{l\} \\ |A'|=k-1}} \prod_{i \in A'} p_i \prod_{\substack{j \in (A')^c \\ j \in (A')^c}} q_j - \sum_{\substack{B' \subseteq [n] \setminus \{l\} \\ |B'|=k}} \prod_{i \in B'} p_i \prod_{\substack{j \in (B')^c \\ j \in (B')^c}} q_j$$

$$= b(k-1, \mathbf{p}_{\hat{l}}) - b(k, \mathbf{p}_{\hat{l}})$$

where  $\Delta_k f(k) = f(k) - f(k-1)$ , the difference operator. On the boundaries k = 0, k = n, the above formula still holds:

$$\frac{\partial b(0,\mathbf{p})}{\partial p_l} = -\prod_{\substack{1 \le j \le n \\ j \ne l}} q_j = -b(0,\mathbf{p}_{\hat{l}}) = b(-1,\mathbf{p}_{\hat{l}}) - b(0,\mathbf{p}_{\hat{l}}) = -\Delta_k b(0,\mathbf{p}_{\hat{l}})$$
$$\frac{\partial b(n,\mathbf{p})}{\partial p_l} = \prod_{\substack{1 \le i \le n \\ i \ne l}} p_i = b(n-1,\mathbf{p}_{\hat{l}}) = b(n-1,\mathbf{p}_{\hat{l}}) - b(n,\mathbf{p}_{\hat{l}}) = -\Delta_k b(n,\mathbf{p}_{\hat{l}}).$$

Let us compute the second derivatives of  $b(k, \mathbf{p})$ :

$$\begin{array}{lll} \frac{\partial^2 b(k,\mathbf{p})}{\partial p_l \partial p_m} &=& \frac{\partial^2 b(k,\mathbf{p})}{\partial p_m \partial p_l} = \frac{\partial}{\partial p_m} ( \ b(k-1,\mathbf{p}_{\hat{l}}) - b(k,\mathbf{p}_{\hat{l}}) \ ) \\ &=& b(k-2,\mathbf{p}_{\hat{l},\hat{m}}) - b(k-1,\mathbf{p}_{\hat{l},\hat{m}}) - b(k-1,\mathbf{p}_{\hat{l},\hat{m}}) + b(k,\mathbf{p}_{\hat{l},\hat{m}}) \\ &=& \triangle_k^2 b(k,\mathbf{p}_{\hat{l},\hat{m}}), \end{array}$$

$$\frac{\partial^2 b(k, \mathbf{p})}{\partial p_l \partial p_l} = \frac{\partial}{\partial p_l} (b(k-1, \mathbf{p}_{\hat{l}}) - b(k, \mathbf{p}_{\hat{l}})) = 0.$$

When k = 0 and k = n, we have

$$\begin{split} \frac{\partial^2 b(0,\mathbf{p})}{\partial p_m \partial p_l} &= \frac{\partial}{\partial p_m} (\ b(-1,\mathbf{p}_{\hat{l}}) - b(0,\mathbf{p}_{\hat{l}}) \ ) = \frac{\partial}{\partial p_m} (\ 0 - \prod_{\substack{1 \le j \le n \\ j \ne l}} q_j) \\ &= \prod_{\substack{1 \le j \le n \\ j \ne l,m}} q_j = b(0,\mathbf{p}_{\hat{l},\hat{m}}) \\ &= b(-2,\mathbf{p}_{\hat{l},\hat{m}}) - 2b(-1,\mathbf{p}_{\hat{l},\hat{m}}) + b(0,\mathbf{p}_{\hat{l},\hat{m}}) = \triangle_k^2 b(0,\mathbf{p}_{\hat{l},\hat{m}}), \\ \frac{\partial^2 b(n,\mathbf{p})}{\partial p_m \partial p_l} &= \frac{\partial}{\partial p_m} (\ b(n-1,\mathbf{p}_{\hat{l}}) - b(n,\mathbf{p}_{\hat{l}}) \ ) = \frac{\partial}{\partial p_m} (\ \prod_{\substack{1 \le i \le n \\ i \ne l,m}} p_i = 0 \ ) \\ &= \prod_{\substack{1 \le i \le n \\ i \ne l,m}} p_i = b(n-2,\mathbf{p}_{\hat{l},\hat{m}}) \\ &= b(n-2,\mathbf{p}_{\hat{l},\hat{m}}) - 2b(n-1,\mathbf{p}_{\hat{l},\hat{m}}) + b(n,\mathbf{p}_{\hat{l},\hat{m}}) = \Delta_k^2 b(n,\mathbf{p}_{\hat{l},\hat{m}}). \end{split}$$

Then let us compute the derivatives of  $H(\mathbb{Z}_p)$ :

$$\begin{aligned} \frac{\partial H(Z_{\mathbf{p}})}{\partial p_l} &= -\sum_{k=0}^n \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \log b(k, \mathbf{p}) - \sum_{k=0}^n b(k, \mathbf{p}) \cdot \frac{1}{b(k, \mathbf{p})} \cdot \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \frac{1}{\ln 2} \\ &= -\sum_{k=0}^n \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \log b(k, \mathbf{p}) - \frac{1}{\ln 2} \frac{\partial}{\partial p_l} \sum_{k=0}^n b(k, \mathbf{p}) \\ &= -\sum_{k=0}^n \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \log b(k, \mathbf{p}) - \frac{1}{\ln 2} \frac{\partial}{\partial p_l} (1) \\ &= -\sum_{k=0}^n \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \log b(k, \mathbf{p}) \\ &= \sum_{k=0}^n \Delta_k b(k, \mathbf{p}_l) \cdot \log b(k, \mathbf{p}); \end{aligned}$$

and the second derivatives of  $H(\mathbb{Z}_{\mathbf{p}})$ :

$$\begin{aligned} \frac{\partial^2 H(Z_{\mathbf{p}})}{\partial p_l \partial p_m} &= -\sum_{k=0}^n \frac{\partial^2 b(k, \mathbf{p})}{\partial p_l \partial p_m} \cdot \log b(k, \mathbf{p}) - \sum_{k=0}^n \frac{\partial b(k, \mathbf{p})}{\partial p_l} \cdot \frac{1}{\ln 2 \cdot b(k, \mathbf{p})} \cdot \frac{\partial b(k, \mathbf{p})}{\partial p_m} \\ &= -\sum_{k=0}^n \triangle_k^2 b(k, \mathbf{p}_{\hat{l}, \hat{m}}) \cdot \log b(k, \mathbf{p}) - \sum_{k=0}^n \frac{\triangle_k b(k, \mathbf{p}_{\hat{l}}) \cdot \triangle_k b(k, \mathbf{p}_{\hat{m}})}{\ln 2 \cdot b(k, \mathbf{p})}, \\ \frac{\partial^2 H(Z_{\mathbf{p}})}{\partial p_l \partial p_l} &= -\sum_{k=0}^n \frac{(\triangle_k b(k, \mathbf{p}_{\hat{l}}))^2}{\ln 2 \cdot b(k, \mathbf{p})}. \end{aligned}$$

The second directional derivative of  $H(Z_{\mathbf{p}})$  in the direction  $\vec{\mathbf{u}}$  is

$$\begin{split} D_{\mathbf{t}}^{2}H(Z_{\mathbf{p}}) &= \frac{\partial^{2}H(Z_{\mathbf{p}})}{\partial p_{l}^{2}} - 2\frac{\partial^{2}H(Z_{\mathbf{p}})}{\partial p_{l}\partial p_{m}} + \frac{\partial^{2}H(Z_{\mathbf{p}})}{\partial p_{m}^{2}} \\ &= -\sum_{k=0}^{n} \frac{(\bigtriangleup_{k}b(k,\mathbf{p}_{\hat{l}}))^{2}}{\ln 2 \cdot b(k,\mathbf{p})} + 2\sum_{k=0}^{n} \bigtriangleup_{k}^{2}b(k,\mathbf{p}_{\hat{l},\hat{m}}) \cdot \log b(k,\mathbf{p}) \\ &+ 2\sum_{k=0}^{n} \frac{\bigtriangleup_{k}b(k,\mathbf{p}_{\hat{l}}) \cdot \bigtriangleup_{k}b(k,\mathbf{p}_{\hat{m}})}{\ln 2 \cdot b(k,\mathbf{p})} - \sum_{k=0}^{n} \frac{(\bigtriangleup_{k}b(k,\mathbf{p}_{\hat{m}}))^{2}}{\ln 2 \cdot b(k,\mathbf{p})} \\ &= 2\sum_{k=0}^{n} \bigtriangleup_{k}^{2}b(k,\mathbf{p}_{\hat{l},\hat{m}}) \cdot \log b(k,\mathbf{p}) - \sum_{k=0}^{n} \frac{(\bigtriangleup_{k}b(k,\mathbf{p}_{\hat{l}}) - \bigtriangleup_{k}b(k,\mathbf{p}_{\hat{m}}))^{2}}{\ln 2 \cdot b(k,\mathbf{p})}. \end{split}$$

The second term of  $D_{\vec{u}}^2$  is clearly negative. We will prove that the first term is strictly negative, therefore  $D_{\vec{u}}^2 H(Z_p) < 0$  will be proved. To evaluate the first term, we use "summation by parts":

$$\sum_{k=0}^{n} (\Delta_k f(k)) \cdot g(k) = -\sum_{k=0}^{n} f(k-1) \cdot (\Delta_k g(k)) \text{ provided } f(n) = f(-1) = 0.$$

Noticing that

$$\begin{split} b(k,\mathbf{p}_{\hat{l},\hat{m}}) &= 0 \quad \text{when } k < 0 \text{ or } k > n-2\\ \triangle_k b(k,\mathbf{p}_{\hat{l},\hat{m}}) &= 0 \quad \text{when } k < 0 \text{ or } k > n-1, \end{split}$$

we can apply summation by parts twice to the first term:

$$\sum_{k=0}^{n} \triangle_{k}^{2} b(k, \mathbf{p}_{\hat{l}, \hat{m}}) \cdot \log b(k, \mathbf{p})$$

$$= -\sum_{k=0}^{n} \triangle_{k} b(k-1, \mathbf{p}_{\hat{l}, \hat{m}}) \cdot \triangle_{k} (\log b(k, \mathbf{p}))$$

$$= \sum_{k=0}^{n} b(k-2, \mathbf{p}_{\hat{l}, \hat{m}}) \cdot \triangle_{k}^{2} (\log b(k, \mathbf{p}))$$

Since  $b(k, \mathbf{p}_{\hat{l},\hat{m}})$ 's are probabilities, they are non-negative, and at least one of them is positive. Finally, we have reduced our proof to the following lemma:

**Lemma 3.**  $b(k, \mathbf{p})$  is log-concave as a function of k, that is,

$$\Delta_k^2(\log b(k, \mathbf{p})) < 0, \quad \text{for any} \ \mathbf{0} \le \mathbf{p} \le \mathbf{1}.$$

Proof of Lemma 3. Since

$$\Delta_k^2(\log b(k, \mathbf{p})) = \log b(k, \mathbf{p}) - 2\log b(k-1, \mathbf{p}) + \log b(k-2, \mathbf{p})$$
$$= \log \frac{b(k-2, \mathbf{p}) \cdot b(k, \mathbf{p})}{(b(k-1, \mathbf{p}))^2},$$

we only need to show that

$$b(k-2,\mathbf{p}) \cdot b(k,\mathbf{p}) < (b(k-1,\mathbf{p}))^2,$$

or equivalently,

$$b(k-1,\mathbf{p}) \cdot b(k+1,\mathbf{p}) < (b(k,\mathbf{p}))^2.$$

Let us expand the left-hand side of the inequality:

$$\begin{split} b(k-1,\mathbf{p}) \cdot b(k+1,\mathbf{p}) \\ &= \sum_{\substack{A \subseteq [n] \\ |A|=k-1}} \prod_{a \in A} p_a \prod_{a' \in A^c} q_{a'} \cdot \sum_{\substack{B \subseteq [n] \\ |B|=k+1}} \prod_{b \in B} p_b \prod_{b' \in B^c} q_{b'} \\ &= \sum_{\substack{A,B \subseteq [n] \\ |A|=k-1,|B|=k+1}} \prod_{a \in A} p_a \prod_{b \in B} p_b \prod_{a' \in A^c} q_{a'} \prod_{b' \in B^c} q_{b'} \\ &= \sum_{\substack{A,B \subseteq [n] \\ |A|=k-1,|B|=k+1}} \prod_{A \cap B} p_i^2 \prod_{(A \cup B) \setminus (A \cap B)} p_{i'} \prod_{(A \cup B)^c} q_j^2 \prod_{(A \cap B)^c \setminus (A \cup B)^c} q_{j'} \\ &\stackrel{(1)}{=} \sum_{w=0}^{k-1} \sum_{\substack{I \subseteq [n] \\ |I|=w}} \sum_{\substack{J \subseteq [n] \setminus I \\ |J|=2k-2w}} (2k-2w) \prod_{I} p_i^2 \prod_{J} p_{i'} \prod_{(I \cup J)^c} q_j^2 \prod_{I^c \setminus (I \cup J)^c} q_{j'}. \end{split}$$

Equality (1) may be illustrated by Figure 1.



Figure 1.

The right-hand side of the inequality can be simplified as:

$$= \sum_{\substack{C \subseteq [n] \\ |C|=k}}^{C \subseteq [n]} \prod_{c \in C} p_{c} \prod_{c' \in C^{c}} q_{c'} \cdot \sum_{\substack{D \subseteq [n] \\ |D|=k}} \prod_{d \in D} p_{d} \prod_{d' \in D^{c}} q_{d'}$$

$$= \sum_{\substack{C,D \subseteq [n] \\ |C|=|D|=k}}^{C,D \subseteq [n]} \prod_{c \in C} p_{c} \prod_{d \in D} p_{d} \prod_{c' \in C^{c}} q_{c'} \prod_{d' \in D^{c}} q_{d'}$$

$$= \sum_{\substack{C=D \subseteq [n] \\ |C|=k}}^{C=D \subseteq [n]} \prod_{c \in C} p_{i}^{2} \prod_{c' \in C} q_{j}^{2} + \sum_{\substack{C \neq D \subseteq [n] \\ |C|=|D=k}}^{C=D \subseteq [n]} \prod_{c' \in C} p_{c'} \prod_{d \in D} p_{d} \prod_{c' \in C^{c}} q_{c'} \prod_{d' \in D^{c}} q_{d'}$$

$$\geq \sum_{\substack{C \neq D \subseteq [n] \\ |C|=k}}^{C\neq D \subseteq [n]} \prod_{c} p_{c} \prod_{D} p_{d} \prod_{c' \in C} q_{c'} \prod_{c' \in D^{c}} q_{d'}$$

$$\stackrel{(2)}{=} \sum_{w=0}^{k-1} \sum_{\substack{I \subseteq [n] \\ |I|=w}}^{I \subseteq [n]} \sum_{\substack{J \subseteq [n] \setminus I \\ |J|=2k-2w}}^{J \subseteq [n] \setminus I} (\frac{2k-2w}{k-w}) \prod_{I} p_{i}^{2} \prod_{J} p_{i'} \prod_{(I \cup J)^{c}} q_{j}^{2} \prod_{c \setminus (I \cup J)^{c}} q_{j'}$$

$$\stackrel{(3)}{=} \sum_{w=0}^{k-1} \sum_{\substack{I \subseteq [n] \\ |I|=w}}^{I \subseteq [n] \setminus I} \sum_{\substack{J \subseteq [n] \setminus I \\ |J|=2k-2w}}^{J \subseteq [n] \setminus I} (\frac{2k-2w}{k-w-1}) \prod_{I} p_{i}^{2} \prod_{J} p_{i'} \prod_{(I \cup J)^{c}} q_{j}^{2} \prod_{c \setminus (I \cup J)^{c}} q_{j'}$$

Equality (2) can be illustrated by Figure 2., and inequality (3) is true since  $\binom{2n}{n} > \binom{2n}{n-1}$ . Therefore Lemma 3 is proved.



Figure 2.

### References

- R. Ahlswede. "Multi-way Communication Channels". Proc. 2nd Int. Symp. Information Theory, Tsahkadsor, Armenian S.S.R., 1971, pp. 23-52. Publishing House of the Hungarian Academy of Sciences, 1973.
- [2] M. L. Ulrey. "The Capacity Region of a Channel with s senders and r receivers." Inform. Contr. vol. 29, pp. 185-203, 1975.
- [3] B. Lindström. "Determining Subsets by Unramified Experiments." J. N. Srivastava, ed., A survey of Statistical Design and Linear Models, North-Holland Publishing Company, 1975.
- [4] H. J. Liao. "Multiple-Access Channels". Ph.D. dissertation, Dept. of Elect. Eng. University of Hawaii, 1972.
- [5] P. S. Mateev. "On Entropy of a Polynomial Distribution." Theory Probab. and Appl. 23, 1, 196-198.

Home Address: 806 CHERRY LANE, APT. 208, EAST LANSING, MI 48823, USA *E-mail address*: szhai@lynx.neu.edu