RC-GRAPHS AND A GENERALIZED LITTLEWOOD-RICHARDSON RULE.

MIKHAIL KOGAN

ABSTRACT. Using a generalization of the Schensted insertion algorithm to rcgraphs, we provide a Littlewood-Richardson rule for multiplying certain Schubert polynomials by Schur polynomials.

1. INTRODUCTION

RC-graphs were originally introduced by Fomin and Kirillov in [4] in connection with Yang-Baxter equation and Schubert calculus. Combinatorial properties of rcgraphs have been later studied by Bergeron and Billey in [1], where rc-graphs were further applied to Schubert calculus, in particular, the Monk's rule was proved using a generalization of the Schensted insertion algorithm to rc-graphs. In this paper we use this generalized algorithm to provide a Littlewood-Richardson rule for multiplying certain Schubert polynomials by Schur polynomials.

Let us introduce some notation. An rc-graph R will be a collection of tuples $\{(i,k)|i,k \leq n\}$, which satisfy some additional properties (see Section 2 for precise definition). Graphically every rc-graph is given by a table of intersecting and non-intersecting strands, so that it represents a planar history of the permutation w_R (this permutation will permute all integers, which are not greater than n, such that there exists N with w(i) = i for every $i \leq N$.)

Define x^R to be the product of x_k 's, with one x_k for each $(i, k) \in R$. Then we can define a Schubert polynomial of w to be

$$S_w = \sum_{w_R = w} x^R.$$

(The standard definition of Schubert polynomials uses divided differences operators, but it was shown in [5] and [3] that the above formula holds.)

Let $\mu = (\mu_1, ..., \mu_n)$ be a partition with $\mu_1 \ge ... \ge \mu_n$. In Section 2 we will associate to each μ a permutation $w(\mu)$. The Schubert polynomial $S_{\mu} = S_{w(\mu)}$ is the Schur polynomial of μ . Since Schubert polynomials form a basis for the ring of all polynomials, we can write

$$S_w S_\mu = \sum_u c^u_{w,\mu} S_u,$$

where the sum is taken over all permutations u. The coefficients $c_{w,\mu}^{u}$ are known to be positive and are called the generalized Littlewood-Richardson coefficients.

Our goal is to provide a rule for computing Littlewood-Richardson coefficients in the case when w satisfies the following property:

$$w(i) > w(i-1)$$
if $i \le 0.$

(Note that this property will imply that if the permutation $w_R = w$ satisfies the above property, then k > 0 for each $(i, k) \in R$).

This rule will use the generalization of the Schensted insertion algorithm to the case of rc-graphs given in [1]. We describe this algorithm in detail in Section 3 and denote by $R \leftarrow k$ the result of the insertion of a number $1 \leq k \leq n$ into an rc-graph R, and by $R \leftarrow Y$ the result of the insertion of a Young tableau Y into an rc-graph R.

The key fact, which makes the generalized Littlewood-Richardson rule possible to prove is the following lemma, which generalizes the row bumping lemma (see [6]) in the case of the classical Schensted algorithm. We will give precise definitions of the paths of insertions in Section 3.3. Roughly speaking these paths are the parts of rc-graphs, which are changed during the insertion algorithms. Let us emphasize the fact that the following lemma does not hold for the general insertion algorithm, but works only in the special case we consider.

Lemma 3.2. If $x \leq y$, then the path of x is weakly to the left of the path of y in $R \leftarrow xy$.

If x > y then the path of x is weakly to the right of the path of y in $R \leftarrow xy$.

This lemma plays a pivotal role in the proof of the following theorem which gives the Littlewood-Richardson rule mentioned above.

Theorem 4.1. Let w be a permutation, which satisfies w(i) > w(i-1) for each $i \leq 0$ and let μ be any partition. Choose any rc-graph U and set $w_U = u$. Then $c_{w,\mu}^u$ is equal to the number of pairs (R, Y) of an rc-graph R and a Young tableau Y with w(R) = w and $\mu(Y) = \mu$, such that $R \leftarrow Y = U$.

Remark 1.1. It is not difficult to see that the insertion algorithm of [1] does not work in a setting more general than in Theorem 4.1. In particular, the Pieri's formula (see [2], [10], [12]) cannot be proved using this algorithm. But a modified insertion algorithm for rc-graphs, which proves the Pieri's formula is constructed in [7].

The paper is organized as follows. In Section 2 we recall basic definitions and properties of rc-graphs and Young tableaux. Section 3 describes the insertion algorithm together with the proof of Lemma 3.2. Section 4 outlines the proof of Theorem 4.1. Finally, Section 5 gives the technical details needed to prove Theorem 4.1.

Acknowledgments. I would like to thank my advisor Victor Guillemin. The results of this paper are a part of my Ph.D. thesis written under his supervision. I also thank Sara Billey and Allen Knutson for many helpful discussions and suggestions.

2. RC-graphs and Young Tableaux.

In this section we define rc-graphs, recall basic facts about Young tableaux and explain why rc-graphs are just generalizations of Young tableaux. At the end of the Section we talk about Schubert and Schur polynomials.

We start with a definition of rc-graphs (our conventions will differ from those of [4] and [1]). Let R be a finite set of pairs of integers $R = \{(i, k) | i \leq n, k \leq n\}$ (both i, k can be negative). We will think of R as a table of intersecting and

nonintersecting strands. Strands intersect for each $(i,k) \in R$ and do not intersect otherwise. The examples are provided on Figure 1, where we have three tables of strands $R_1 = \{(2,1), (1,1), (-1,2)\}, R_2 = \{(3,1), (2,3), (1,2)\}$ and $R_3 = \{(3,2), (2,2), (1,2), (2,3)\}.$



Figure 1: Examples of rc-graphs.

R is called an rc-graph if no two strands intersect twice. We can think of each rc-graph as a planar history of a permutation w_R , which is defined as follows. If we label each strand by the row where it starts from, then $w_R(i)$ is given by the column, where the i^{th} strand ends. Each w_R permutes all the integers, which are less than or equal to n. Moreover, there always exists some negative N such that $w_R(i) = i$ for i < N.

For example, for the rc-graphs from Figure 1, the corresponding permutations are given by $w_{R_1}(2, 1, 0, -1, -2) = (2, -2, 1, 0, -1)$ with $w_{R_1}(i) = i$ for i < -2, $w_{R_2}(3, 2, 1, 0, -1) = (3, 1, -1, 2, 0)$ with $w_{R_2}(i) = i$ for every i < -1 and finally $w_{R_3}(3, 2, 1, 0, -1) = (3, -1, 1, 2, 0)$ with $w_{R_3}(i) = i$ for every i < -1.

Let us show that R also provides a reduced expression for w_R , in other words we can write w_R as a composition of minimal number of simple transpositions (this minimal number is called the length $\ell(w)$ of a permutation w). Denote by s_i the simple transposition, which permutes i and i + 1 (i might be negative). Then to produce the reduced expression for w_R , read each row of the rc-graph from right to left, from the top row to the bottom one and multiply out simple transpositions $s_{i+k-n-1}$ for each (i, k) $\in R$. It is easy to see we get a reduced expression for w_R .

Let us recall some properties of rc-graphs, which were proved in [1]:

- Let R be an rc-graph, such that (i, k), (i 1, k) and $(i 1, k \ell)$ are not in R for some $i, k \leq n$ and positive ℓ , but all other (i', k') with $k \geq k' \geq k + \ell$ and $i \geq i' \geq i + 1$ are in R. Then we can substitute $(i, k - \ell)$ by (i - 1, k) without changing w_R . (We also can go backwards.) These operations are called *ladder* moves of size ℓ at the place (i, k). Examples of ladder moves of sizes 1 and 2 are shown on Figure 2
- For every permutation w there exist a unique rc-graph R_w (which we will call a top rc-graph of w), such that every other rc-graph R with $w_R = w$ could be constructed from R_w by a sequence of ladder moves, which change $(i, k - \ell)$ to (i - 1, k) (but not the other way).
- For a top rc-graph R_w , if $(i, k) \in R_w$ and i < n, then $(i+1, k) \in R_w$. In other words, all intersecting strands of R_w are concentrated to the left in each row of R_w .



Figure 2: Examples of ladder moves on rc-graphs of sizes 1 and 2.

From now on we will only work with those permutations w for which w(i) > w(i-1) for each $i \leq 0$. Equivalently, every rc-graph R with $w_R = w$ can be defined by the following property:

• *R* has no two nonpositive intersecting strands.

In particular, if R satisfies the above property it lies above the 0th row, that is if $(i,k) \in R$ then $k \ge 1$. Let us emphasize that starting from this point every rc-graph mentioned in this text has to satisfy the above property. In particular, the property is implicitly assumed in all the statements of theorems and lemmas stated below.

Let us now define Young diagrams and tableaux. A Young diagram will be given by a partition $\mu = (\mu_1, ..., \mu_n)$, where $\mu_1 \ge \mu_2 \ge ... \ge \mu_n > 0$. Graphically it will be given by μ_i boxes in i^{th} row, as shown on Figure 3, where Young diagrams correspond to partitions (3), (2, 1) and (3, 1, 1) respectively.



Figure 3: Examples of Young diagrams.

A Young tableaux Y is a filling of a Young diagram with numbers 1, ..., n which satisfies the following properties. If a and b are two boxes of the Young diagram, which lie in the same row, and a is to the left of b then the number in a is not greater than the number in b. If a and b are in the same column and a is on top of b, then the number in a should be strictly less than the number in b. We denote by $\mu(Y)$ the partition, which corresponds to the Young tableaux Y. Examples of Young tableaux are shown on Figure 4.



Figure 4: Examples of Young tableux.

Given a partition μ , we construct an rc-graph $R(\mu)$ as follows. Let $R(\mu) = \{(i,k)|1 \leq k \leq n, n \geq i \geq n - \mu_k + 1\}$. Then, set $w(\mu) = w_{R(\mu)}$. (Note $R(\mu)$ is the top rc-graph of $w(\mu)$.) Every such permutation has a unique ascent at 0, that is w(1) < w(0) but w(i) > w(i-1) if $i \neq 1$. In particular every permutation $w(\mu)$

satisfies w(i) > w(i-1) if $i \le 0$. The following lemma shows why we can think about rc-graphs as about generalizations of Young diagrams (similar results were obtained by Winkel in [13] and pointed out in [1]).

Lemma 2.1. *RC*-graphs *R* with $w_R = w(\mu)$ are in one to one correspondence with Young tableaux *Y* with $\mu(Y) = \mu$.

Proof. It is easy to see that we can apply only ladder moves of size 1 to any R with $w_R = w(\mu)$. Start with the top rc-graph $R(\mu)$ and the Young diagram, which is given by filling the i^{th} row of the Young diagram with entries i. Associate to each ladder move of size 1 an increase by 1 of the corresponding box in the Young tableaux. This obviously constructs a one to one correspondence between rc-graphs with permutation $w(\mu)$ and Young tableaux with partition μ .

Denote by R(Y) the rc-graph, which is constructed out of the Young tableau Y. As an illustration to the above lemma let us mention that the first Young tableau Y_1 on Figure 4 correspond to the first rc-graph R_1 on Figure 1. At the same time $w(\mu(Y_2)) = w_{R_2}$ but $R(Y_2) \neq R_2$.

Call any finite sequence of numbers 1, ..., n a *word*. For each Young tableau Y, associate a word v(Y), which is given by reading the entries of the tableau from left to right in each row, starting from the bottom row and going to the top one. For example, the words of Young diagrams from Figure 4 are $v(Y_1) = 112, v(Y_2) = 313$ and $v(Y_3) = 42122$.

On the set of all words we define Knuth moves (originally they appeared in [8]). These Knuth moves allow the following changes to a word:

$$...yxz... \Leftrightarrow ...yzx...$$
 if $x < y \leq z$

and

$$\dots xzy\dots \Leftrightarrow \dots zxy\dots$$
 if $x \leq y < z$

We say that two words v_1 and v_2 are Knuth equivalent if we can go from one of them to another by applying a sequence of Knuth moves.

The following theorem is the key fact in the Littlewood-Richardson rule for multiplying Schur polynomials and is very useful to us. The proof of it can be found in [6].

Theorem 2.2. If Y_1 and Y_2 are two distinct Young tableaux then $v(Y_1)$ and $v(Y_2)$ are not Knuth equivalent. Moreover, each word v is Knuth equivalent to exactly one word v(Y).

Let us now recall the definitions of Schur and Schubert polynomials. Each Young tableaux Y defines a monomial x^Y , which is equal to the product of x_i 's with one x_i for each entry *i* in the tableaux. Each partition μ defines a Schur polynomial

$$S_{\mu} = \sum_{\mu(Y)=\mu} x^{Y}.$$

It is well known that Schur polynomials are symmetric and that they form a basis for the ring of symmetric polynomials in n variables.

Similarly, given an rc-graph R we define x^R to be the product of x_k 's with one x_k for each $(i, k) \in R$. Then the Schubert polynomial for the permutation w is given by

$$S_w = \sum_{w_R = w} x^R.$$

MIKHAIL KOGAN

(Let us recall again that the standard definition of rc-graphs uses divided differences operators.) Polynomials S_w form a basis for the ring of all polynomials in n variables.

Since $x^{R(Y)} = x^Y$, Proposition 2.1 implies that $S_{w(\mu)} = S_{\mu}$, in other words we can think of Schubert polynomials as generalizations of Schur polynomials.

3. INSERTION ALGORITHM

The key tool in the classical Littlewood-Richardson rule for multiplying Schur polynomials is the Schensted insertion algorithm. This algorithm was generalized to the case of rc-graphs in [1] and used to prove Monk's formula. We will use a special case of this generalized algorithm to provide the Littlewood-Richardson rule for multiplying some Schubert polynomials by Schur polynomials. This section defines the algorithm and discusses its basic properties.

Let R be an rc-graph. We would like to provide an algorithm for inserting a number $1 \le k \le n$ into R.

Let us call a pair (i, j) an *open space*, if $(i, j) \notin R$ (two strands at position (i, j) do not intersect) and the bottom strand of the intersection is labeled by a nonpositive number, while the top strand is labeled by a positive number. (See Figure 5.)



Figure 5: An example of an open space: $a > 0 \ge b$.

Start at the row number $k_1 = k$ and find the smallest i_1 such that the space (i_1, k_1) is open (sometimes we will write $(i_1(k), k_1(k))$) to indicate the dependence on k). Insert (i_1, k_1) into R, in other words make the strands intersect at (i_1, k_1) . If a, b are the two labels of the strands going through the place (i_1, k_1) we set $a_1(k) = a$ and $b_1(k) = b$. If we constructed an rc-graph we stop, otherwise, it can be shown that the two stands which now intersect at the place (i_1, k_1) must also intersect at some other place (ℓ_2, k_2) with $k_2 > k_1$. We remove (ℓ_2, k_2) from R and find the smallest $i_2 > \ell_2$ such that (i_2, k_2) is open. We insert (i_2, k_2) into R, set $a_2(k)$ and $b_2(k)$ to be the labels of the strands passing through (i_2, k_2) and continue the process until it stops. For notational convenience set $k_{j+1}(k) = n + 1$, if the last intersection we inserted was $(i_j(k), k_j(k))$.

It was shown in [1] that the above algorithm stops at some point and produces a new rc-graph, which we denote by $R \leftarrow k$. Note, $R \leftarrow k$ and R have the same number of crossings in each row, except for the row k, where $R \leftarrow k$ has an additional crossing. Hence

$$x^R x_k = x^{R \leftarrow k}.$$

If v is a word, we denote by $R \leftarrow v$ the rc-graph we get after inserting one by one the letters of v. If Y is a Young tableau, we say $R \leftarrow Y = R \leftarrow v(Y)$. Obviously we have:

$$x^R x^Y = x^{R \leftarrow Y}$$

The above algorithm is a generalization of the Schensted row insertion algorithm (see [11] or [6]). To prove this we just have to translate what this algorithm means in the language of Young tableaux, in the case when R = R(Y) is constructed from

 $\mathbf{6}$

some young tableau Y as in Lemma 2.1. We omit the simple technical details of this proof, but recall a very important fact about this algorithm (see [6]). For two Young tableaux Y_1 and Y_2 we have

(1)
$$v(Y_1 \leftarrow Y_2)$$
 is Knuth equivalent to $v(Y_1)v(Y_2)$.

where $v(Y_1)v(Y_2)$ is just the concatenation of the two words $v(Y_1)$ and $v(Y_2)$.

Let us introduce new notations, which will be used later. During the insertion algorithm of k into R, each place (i_j, k_j) was connected to (ℓ_{j+1}, k_{j+1}) by two pieces of strands s_j and s^j . (Let us emphasize that s_j and s^j are just pieces of strands, which are between rows k_j and k_{j+1} , and their labels change during the insertion.) We say that the left strand s_j is a part of the left path $\ell(k)$ of the insertion while the right strand s^j is a part of the right path r(k) of the insertion. Both $\ell(k)$ and r(k) are collections of pieces of stands. The labeling of each piece s_j in $\ell(k)$ changes from being positive a_j to nonpositive b_j , while the labeling of s^j in r(k)change from b_j to a_j .

Figure 6 contains an example of inserting 1 into an rc-graph. The path of insertion is shown on the resulting rc-graph.



Figure 6: Insertion algorithm and path of insertion.

Recall that we are considering only those rc-graphs, for which no two nonpositive strands intersect. The following lemma shows that the insertion algorithm preserves this property.

Lemma 3.1. If no two nonpositive strands intersect in R, then no two nonpositive strands intersect in $R \leftarrow k$.

Proof. During the insertion algorithm the only possibility for introducing new intersections of nonpositive strands is when a strand s_j from $\ell(k)$ becomes nonpositive.

Let us show by contradiction that s_j cannot intersect any nonpositive strand. Assume that s_j (labeled by $b_j \leq 0$) in $R \leftarrow k$ is intersected by some nonpositive strand s in row $k_j < k' \leq k_{j+1}$. Look at the whole strand s' in $R \leftarrow k$, which is labeled by a_j . s' starts above zero, while s starts below zero, at the same time s is to the left of s' in the row k', hence these two strands must intersect in $R \leftarrow k$ below the row k'. The strand s' below the row k' consists of two parts: one of them is s^j , which was labeled by b_j in R, and the other one is the rest of the strand below the row k_j , which is labeled by a_j in both R and $R \leftarrow k$.

s cannot intersect s^j , since no two nonpositive strands intersect in R. At the same time s cannot intersect the rest of s', since it already intersects the strand

labeled by a_j in R once at row k', and cannot intersect it for the second time. So we found a contradiction and this lemma is proved.

The lemma immediately leads to the following property

• All strands passing between r(k) and $\ell(k)$ are positive.

Indeed, each strand between r(k) and $\ell(k)$ has to cross at least one strand from r(k) or $\ell(k)$, but, since r(k) is nonpositive in R and $\ell(k)$ is nonpositive in $R \leftarrow k$, and no two nonpositive strand can intersect, the above property holds.

Here is a very important lemma, which does not hold if we do not assume that no two nonpositive strands intersect in R (see [1] for a counterexample).

Lemma 3.2. If $x \leq y$, then the path of x is weakly to the left of the path of y in $R \leftarrow xy$.

If x > y then the path of x is weakly to the right of the path of y in $R \leftarrow xy$.

Remark 3.3. When we say that the path of x is to the left (right) of the path of y, we imply that the right path of x is to the left of the left path of y (respectively, the left path of x is to right of the right path of y). The word *weakly* stands for the fact that r(x) and $\ell(y)$ (respectively $\ell(x)$ and r(y)) might have some common parts.

Proof. For the case $x \leq y$ the right path r(x) of x in $R \leftarrow x$ contains strands which are all greater than zero after the insertion. Thus when we start inserting y into $R \leftarrow x$ each row $k \geq x$ should contain an open space to the right of the right path of x (since the right path of x is positive). Hence the left path of y is going to stay strictly to the right of the right path of x, until at some point it might happen that left path of y is the same as the right path of x.

It can occur only when an open space $(i,k) = (i_j(y), k_j(y))$ in $R \leftarrow x$ contains strands $s_j(y)$ and $s^j(y)$, such that part of $s_j(y)$ is a part of r(x). In other words, $s_j(y)$ and $s^{j'}(x)$ have a common part. If the insertion algorithm stops at this point there is nothing to prove. Otherwise, there should be a place $(i_{j+1}(y), k_{j+1}(y))$ where strands $s_j(y)$ and $s^j(y)$ intersect again. We would like to show

(2)
$$k_{j+1}(y) \ge k_{j'+1}(x)$$

This will be enough to prove the first part of the lemma. Indeed if r(x) and $\ell(y)$ coincide at the row k, they have to separate at the row $k_{j'+1}(x)$ by (2), so that r(x) moves to the left of $\ell(y)$. If they coincide again at some higher row, we can repeat the argument and show that they have to separate again.

To prove (2), note $s^{j'}(x)$ was nonpositive in R, hence it cannot intersect s^{j} , which was also nonpositive in R. Thus if $s_{j}(y)$ and $s^{j}(y)$ intersect in R, it should happen above the row where $s^{j'}(x)$ ends, in other words, above the row of intersection of $s_{j'}(x)$ and $s^{j'}(x)$, but this row is exactly $k_{j'+1}(x)$. Therefore (2) holds and the first part of the lemma is proved.

In the case x > y, the right path of y gets changed from being a set of nonpositive strands to positive strands. The left path of x in $R \leftarrow x$ contains only nonpositive strands, so these two paths cannot intersect (but some parts of them can coincide), since no two nonpositive strands can intersect.

Let's now argue by contradiction that r(y) is weakly to the left of $\ell(x)$ using the fact that r(y) cannot intersect $\ell(x)$. Pick the smallest k, such that the right path of y is to the right of the left path of x. This could not happen because of an intersection of r(y) and $\ell(x)$. Thus in the row k the insertion of x into R we had to remove some $(\ell_j(x), k_j(x)) = (\ell_j(x), k)$ from R and add some $(i_j(k), k)$ to R, moving $\ell(x)$ to the left. But only nonpositive strands pass in row k between $\ell_j(x)$ and $i_j(k)$ (otherwise, we would get an open space there, which is impossible), hence r(y) cannot pass between $\ell_j(x)$ and $i_j(x)$ and it must coincide with $\ell(x)$ in the row k-1. Moreover, the strand passing the row k directly to the left of $(\ell_j(x), k)$ is nonpositive in $R \leftarrow x$. At the same time, the strands between right and left paths of y are always positive, so the strand passing the row k directly to the left of $(\ell_j(x), k)$ is positive in $R \leftarrow x$. We found a contradiction, which means that the second part of the lemma is proved.

This Lemma immediately proves that if $0 < x < y \le z \le n$ then

Indeed in $R \leftarrow yx$ we know that $\ell(y)$ is weakly to the right of r(x), so r(y) is unchanged when we insert x into $R \leftarrow y$. At the same time when we insert z in $R \leftarrow y$ the left path $\ell(z)$ is weakly to the right of r(y). So, paths of x and z are separated by the path of y and, in particular, do not have any common strands. Hence multiplication of $R \leftarrow y$ by x commutes with multiplication by z, which proves (3).

Thus if v_1 and v_2 are two Knuth equivalent words, which can be gotten from one another using only Knuth moves of the first type, we have

$$R \leftarrow v_1 = R \leftarrow v_2$$

Let us talk about how the permutation of R changes after the insertion. Notice that after each step of the algorithm the permutation w_R does not change except for the last step. At the end we make two nonintersecting stands labeled by c and d intersect, which means that

$$w_{R\leftarrow x} = w_R s_{c,d}$$

where $s_{c,d}$ is the transposition (with $c > 0 \ge d$), which interchanges the elements in position c and d, when it acts on a permutation from the right. Moreover,

$$l(w_R s_{c,d}) = l(w_R) + 1$$

Conversely, given an rc-graph R' with $w_{R'} = w_R s_{c,d}$, such that $l(w_R s_{c,d}) = l(w_R)+1$ and $c > 0 \ge d$ we can traverse the above algorithm backwards starting by finding the unique intersection of strands labeled by c and d, making them nonintersecting and then proceeding in the opposite order. For more details about the inverse of the insertion algorithm see [1], where Monk's formula was proved using this inverse insertion algorithm.

4. LITTLEWOOD-RICHARDSON RULE FOR MULTIPLICATION SCHUBERT POLYNOMIALS BY SCHUR POLYNOMIALS.

Given a Schur polynomial S_{μ} and a Schubert polynomials S_w their product can be uniquely written as a sum of Schubert polynomials:

$$S_w S_\mu = \sum_u c^u_{w,\mu} S_u,$$

where the sum is taken over all the permutations u. The coefficients $c_{w,\mu}^u$ are called Littlewood-Richardson coefficients and are known to be positive. The following theorem provides a rule for computing these coefficients: **Theorem 4.1.** Let w be a permutation, which satisfies w(i) > w(i-1) for each $i \leq 0$ and let μ be any partition. Choose any rc-graph U and set $w_U = u$. Then $c_{w,\mu}^u$ is equal to the number of pairs (R, Y) of an rc-graph R and a Young tableau Y with w(R) = w and $\mu(Y) = \mu$, such that $R \leftarrow Y = U$.

The next three lemmas will lead to the proof of the above theorem. Let us define a Young diagram ν_m to be just one row of *m* boxes, so that the corresponding partition is given by one number *m*.

Lemma 4.2. Theorem 4.1 holds when $\mu = \nu_m$.

Remark 4.3. The above lemma is just a special case of the Pieri formula. Since Lemma 3.2 does not hold in general, Pieri formula was conjectured but was not proved in [1]. It was later proved by other methods in [2], [10], [12]. In [7] the Pieri formula is proved using a generalization of the insertion algorithm for rc-graphs.

Lemma 4.4. The polynomials S_{ν_m} generate the ring of symmetric polynomials in n variables. So that each symmetric polynomial S can be written as

(4)
$$S = \sum_{(m_1,\dots,m_k)\in M_+} S_{\nu_{m_1}}\cdots S_{\nu_{m_k}} - \sum_{(m_1,\dots,m_k)\in M_-} S_{\nu_{m_1}}\cdots S_{\nu_{m_k}}$$

where M_+ and M_- are two sets of sequences of positive numbers.

Lemma 4.5. Let R be an rc-graph then

$$R \leftarrow yxz = R \leftarrow yzx \quad if \quad 0 < x < y \le z \le n$$

and

$$R \leftarrow xzy = R \leftarrow zxy \quad if \quad 0 < x < y < z < n.$$

Corollary 4.6. If R is an rc-graph and Y_1 and Y_2 are two Young tableaux then

$$R \leftarrow (Y_1 \leftarrow Y_2) = (R \leftarrow Y_1) \leftarrow Y_2.$$

We postpone the proofs of the above three lemmas until the next Section. Let us just note that Corollary 4.6 follows easily from Lemma 4.5 and Fact (1).

Let us show how Theorem 4.1 can be proved using the above three lemmas. We define the sets \mathcal{R}_w and \mathcal{Y}_{μ} to be

$$\mathcal{R}_w = \bigcup_{w_R=w} R \text{ and } \mathcal{Y}_\mu = \bigcup_{\mu(Y)=\mu} Y$$

If

$$\mathcal{R} = \mathcal{R}_w \cdot \mathcal{Y}_\mu = \bigcup_{R \in \mathcal{R}_w, Y \in \mathcal{Y}_\mu} R \leftarrow Y,$$

we would like to show that

(5)
$$\mathcal{R} = \bigcup_{u} \mathcal{R}_{u}$$

This implies that each \mathcal{R}_u is taken $c_{w,\mu}^u$ times in the above union, since there is a unique way of writing $S_w S_\mu$ as a sum of Schubert polynomials. Hence (5) will prove the theorem. Use Lemma 4.4 to write

$$S_{\mu} = \sum_{(m_1, \dots, m_k) \in M_+} S_{\nu_{m_1}} \cdots S_{\nu_{m_k}} - \sum_{(m_1, \dots, m_k) \in M_-} S_{\nu_{m_1}} \cdots S_{\nu_{m_k}}$$

this immediately implies that

(6)
$$\mathcal{Y}_{\mu} = \bigcup_{(m_1,\dots,m_k)\in M_+} \mathcal{Y}_{\nu_{m_1}}\cdots\mathcal{Y}_{\nu_{m_k}} - \bigcup_{(m_1,\dots,m_k)\in M_-} \mathcal{Y}_{\nu_{m_1}}\cdots\mathcal{Y}_{\nu_{m_k}}$$

where the minus stands for the set theoretic difference of the two sets and where $\mathcal{Y}_{\mu_1} \cdot \mathcal{Y}_{\mu_2} = \bigcup_{\mu(Y_1)=\mu_1, \mu(Y_2)=\mu_2} Y_1 \leftarrow Y_2$. The reason why we can take the set theoretic difference in the above formula

The reason why we can take the set theoretic difference in the above formula is the following. By Lemma 4.2 both first and second sets in (6) could be broken up into unions of $\mathcal{Y}_{\mu'}$ (since any insertion into a Young tableaux produces a Young tableaux). But since S_{μ} cannot be written as a nontrivial linear expression of $S_{\mu'}$'s the set theoretical difference above is well defined.

Thus we can conclude:

$$\mathcal{R} = \mathcal{R}_w \cdot \big(\bigcup_{(m_1, \dots, m_k) \in M_+} \mathcal{Y}_{\nu_{m_1}} \cdots \mathcal{Y}_{\nu_{m_k}}\big) - \mathcal{R}_w \cdot \big(\bigcup_{(m_1, \dots, m_k) \in M_-} \mathcal{Y}_{\nu_{m_1}} \cdots \mathcal{Y}_{\nu_{m_k}}\big)$$

Using Corollary 4.6 we can immediately see that the set theoretic difference is welldefined in the above formula. On the other hand, this formula and Lemma 4.2 shows that \mathcal{R} can be written in the form (5), since by Lemma 4.2 each $\mathcal{R}_w \cdot \mathcal{Y}_{\nu_{m_1}} \cdots \mathcal{Y}_{\nu_{m_k}}$ is a union $\bigcup_u \mathcal{R}_u$. This finishes the proof of the Theorem 4.1.

5. TECHNICAL DETAILS IN THE PROOF OF THEOREM 4.1.

Proof. (of Lemma 4.2) The proof of this Lemma will just be a combination of Monk's rule and Lemma 3.2.

Let $Y = (1 \le a_1 \le a_2 \le \dots \le a_m \le n)$ be a filling of the Young diagram ν_m . We can easily see from the insertion algorithm that

$$w_{R\leftarrow Y} = w_{R\leftarrow a_1\dots a_m} = w_R s_{c_1,d_1}\dots s_{c_m,d_m}$$

where $c_i > 0 \ge d_i$, $d_1 > d_2 > ... > d_m$ and $l(w_R s_{c_1, d_1} ... s_{c_{m'}, d_{m'}}) = l(w_R) + m'$ for every $m' \le m$.

Conversely, assume we are given an rc-graph R' with $w_{R'} = w_R s_{c_1,d_1} \dots s_{c_m,d_m}$ with $c_i > 0 \ge d_i$, $d_1 > d_2 > \dots > d_m$ and $l(w_R s_{c_1,d_1} \dots s_{c_{m',d_{m'}}}) = l(w_R) + m'$ for every $m' \le m$. Then we can go through the inverse insertion algorithm and delete one by one intersections of strands c_i and d_i . We will get m numbers a_1, \dots, a_m .

Moreover, by Lemma 3.2, $a_i \leq a_{i+1}$ (otherwise we would not have $d_i < d_{i+1}$). Thus we have even proved a slightly better version of the Lemma:

$$S_w S_{\nu_m} = \sum S_{w s_{c_1, d_1} \dots s_{c_m, d_m}}$$

where $c_i > 0 \ge d_i$, $d_1 > d_2 > ... > d_m$ and $l(w_R s_{c_1, d_1} ... s_{c_{m', d_{m'}}}) = l(w_R) + m'$ for every $m' \le m$.

Proof. (of Lemma 4.4) This is an immediate corollary of the Jacobi-Trudi identity (see [9]). We thank Sara Billey for pointing this out to us. \Box

11

MIKHAIL KOGAN

The rest of this Section will be concerned with the proof of Lemma 4.5.

Recall that the first part of Lemma 4.5 followed from Lemma 3.2. So, we just have to prove the second part of it:

(7)
$$R \leftarrow xzy = R \leftarrow zxy$$
 for any R and $0 < x \le y < z \le n$

The path of x in $R \leftarrow xz$ is weakly to the left of the path of z. If it is strictly to the left of the path of z (in other words the right path r(x) of x has no common parts with the left path $\ell(z)$ of z), then clearly $R \leftarrow xz = R \leftarrow zx$ and (7) holds. An example for this situation would be x = 1, y = 2, z = 3 and $R = R_3$ (the third rc-graph from Figure 1).

Hence we just have to look at the case when right path of x partially coincides with the left path of z. Let's assume that the bottom row where this happens is k. Then by above argument, $R \leftarrow xzy = R \leftarrow zxy$ for all rows, which are below the row k.

Assume that during the insertion of x into R an intersection $(i_j(x), k_j(x))$ was inserted into R, such that $k_j(x) < k$ but $k_{j+1}(x) > k$. Denote by s_1 and s_2 the two pieces of strands, which connect $(i_j(x), k_j(x))$ with $(\ell_{j+1}(x), k_{j+1}(x))$. Set $a = a_j(x) > 0 \ge b = b_j(x)$. So that during the insertion of x into R, the labeling of s_1 changed from a to b, while the labeling of s_2 changed from b to a.

Assume that during the insertion of z into $R \leftarrow x$ we insert an intersection of strands at the place $(i, k) = (i_{j'}(z), k_{j'}(z))$, so that one of the strands at (i, k) is s_2 . Denote by s'_2 the piece of this strand, which connects (i, k) with $(\ell_{j'+1}(z), k_{j'+1}(z))$. $(s_2 \text{ and } s'_2 \text{ have a common piece between the rows k and <math>k_{j+1}(x)$.) Take the other strand coming out of (i, k) and denote the piece of this strand, which connects (i, k) with $(i_{j'+1}(z), k_{j'+1}(z))$, by s_3 . Clearly, $a_{j'}(z) = a$ and we set $b_{j'}(z) = c$.

Since the path of y has to sit between the right path of x and the left path of z below row k, the strand s_2 has to become a part of the left path of the insertion of y into $R \leftarrow xz$. Assume it happened at some place $(i_1, k_1) = (i_{j''}(y), k_{j''}(y))$. We claim that $(i, k) = (\ell_{j''+1}(y), k_{j''+1}(y))$. In other words, the strands, which pass through (i_1, k_1) in R have to pass through (i, k) in R. Indeed, if this claim does not hold, then the strand labeled by $c \leq 0$ has to pass between the left and right paths of y, which is impossible. Denote by s'_3 the right path, which connects (i_1, k_1) with (i, k), so that s'_3 and s_3 are two pieces of the same strand in R.

Hence during the insertion algorithm of y into $R \leftarrow xz$ we had to remove intersection (i, k) and find an open space to the left of it, call it $(\bar{i}, k) = (i_{j''+1}(y), k_{j''+1}(y))$. We have two cases:

Case 1. (\bar{i}, k) is to the left of the strand s_1 .

Case 2. (\bar{i}, k) is to the right of the strand s_1 .

Before going through the proofs for both cases, let us give two examples. Case 1 happens when we take x = y = 2, z = 3 and $R = R_3$ from Figure 1. For Case 2 take n = 2 and $R = \{(2, 2)\}$ then x = y = 1 and z = 2 will produce Case 2.

Proof of Case 1. First of all let us note that (\bar{i}, k) is to the left of s_1 if and only if the stand s_1 passes exactly to the left of strand s_2 in the row k, that is there are no other strands between s_1 and s_2 in the row k. Indeed, if we had other strands between them they had to be positive in R (since they lie between right and left paths of x), but then (i - 1, k) would be an open space, so that $\bar{i} = i - 1$, which contradicts the fact that (\bar{i}, k) is to the left of the strand s_1 . This argument also proves that (\bar{i}, k) is to the right of s_1 if and only if $\bar{i} = i - 1$, which will be used in the proof of the second case. Denote by p_1^x the path of the insertion of x into R above the row k, by p_1^z the path of the insertion of z into $R \leftarrow x$ above the row k and by p_1^y the path of the insertion of y into $R \leftarrow xz$ above the row k. Notice that since (\overline{i}, k) is to the left of s_1 we can conclude that p_1^z is weakly to the right of p_1^x , while p_1^y is weakly to the left of p_1^x .

Let us think how $R \leftarrow zxy$ looks like in this case. When we insert z into R, the open space $(i_{j'}(z), k_{j'}(z))$ in the row k is no longer (i, k), but it is now (i - 1, k). Indeed, s_1 is labeled by a > 0 while s_2 is labeled by $b \le 0$ in R, moreover, s_1 passes through the space (i - 1, k) and together with s_2 creates an open space. So we insert (i - 1, k) into R and denote by p_2^z the path of z in R above the row k. Notice that the paths p_2^z and p_1^x are identical. When we insert x into $R \leftarrow z$, at the row k we have to remove $(i - 1, k) = (\ell_{j+1}(x), k_{j+1}(x))$, since s_1 and s_2 intersect at (i - 1, k) in $R \leftarrow z$. So, we remove (i - 1, k) and insert $(\bar{i}, k) = (i_{j+1}(x), k_{j+1}(x))$ into $R \leftarrow z$. Denote by p_2^x the path of x in $R \leftarrow z$ above the row k. Notice that p_2^x is identical with p_1^y . At the same time, the path p_2^y of the insertion of y into $R \leftarrow zx$ above the row k will be identical with p_1^z .

To summarize, we have $R \leftarrow xzy = R \leftarrow zxy$ below the row k. Above the row k we first insert into R along the path $p_2^z = p_1^x$ in both cases. Then we insert along p_1^z and along p_1^y for $R \leftarrow xzy$ and along p_2^x and along p_2^y for $R \leftarrow xzy$. But since p_1^z, p_2^y are weakly to the right of $p_2^z = p_1^x$ while p_2^x, p_1^y are weakly to the left of $p_2^z = p_1^x$, we can apply Lemma 3.2 to show that paths $p_1^z = p_2^y$ and $p_2^x = p_1^y$ are separated by $p_2^z = p_1^x$ and hence it does not matter along which path above the row k we insert first. This proves $R \leftarrow xzy = R \leftarrow zxy$ above the row k and finishes the proof of Case 1.

Proof of Case 2. Case 2 is just slightly more difficult than Case 1.

Denote by k the row where the path of the insertion of y into $R \leftarrow xz$ moves to the left of s_1 (this has to happen above the row k, but below the row $k_{j+1}(x)$). As in Case 1, it can be shown that in the row \tilde{k} the strand s_1 has to pass directly to the left of s_2 . We define paths p_i^x , p_i^y and p_i^z for i = 1, 2, which lie above the row \tilde{k} , the same way we have done it in the first case. We also denote by p the path of the insertion of y between the rows k and \tilde{k} .

Then applying the same arguments as in Case 1, we can show that $R \leftarrow xzy = R \leftarrow zxy$ below the row k. We can also see that above the row k, p_2^z will be the path p between k and \tilde{k} and p_1^x above \tilde{k} . Similarly to Case 1, we apply Lemma 3.2 to show that p_2^x coincides with p_1^y and p_2^y is the same as p_2^z above the row \tilde{k} . Hence paths of insertion in both $R \leftarrow xzy$ and $R \leftarrow zxy$ above the row k are the same. This finishes the proof of Lemma 4.5 in the second case.

References

- N. Bergeron, S. Billey. RC-graphs and Schubert polynomials. Experiment. Math. 2 (1993), no. 4, 257–269.
- [2] N. Bergeron, F. Sotille. A Pieri-type formula for isotropic flag manifolds. MSRI Preprint 1998-050, arxiv:math.CO/9810025
- [3] S. Billey, W. Jockusch, R. Stanley. Some combinatorial properties of Schubert polynomials. J. Algebraic Combin. 2 (1993), no. 4, 345–374.
- [4] S. Fomin, A. Kirillov. The Yang-Baxter equation, symmetric functions, and Schubert polynomials. Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993). Discrete Math. 153 (1996), no. 1-3, 123–143.
- [5] S. Fomin, R. Stanley. Schubert polynomials and the nil-Coxeter algebra. Adv. Math. 103 (1994), no. 2, 196–207.

MIKHAIL KOGAN

- [6] W. Fulton. Young tableaux. With applications to representation theory and geometry. London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [7] M. Kogan, A. Kumar. A proof of Pieri's formula using generalized Schensted insertion algorithm for rc-graphs,. preprint
- [8] D. Knuth Permutations, matrices, and generalized Young tableaux. Pacific J. Math. 34 1970 709-727.
- [9] I. G. Macdonald *Notes on Schubert Polynomials* Montreal, Canada : Departement de mathematiques et d'informatique, Universite du Quebec a Montreal, 1991
- [10] A. Postnikov On a Quantum Version of Pieri's Formula, preprint March 23, 1997.
- [11] C. Schensted. Longest increasing and decreasing subsequences. Canad. J. Math. 13 1961 179–191.
- [12] F. Sottile. Pieri's formula for flag manifolds and Schubert polynomials, Annales de l'Institut Fourier 46 (1996), 89-110.
- [13] R. Winkel. A combinatorial bijection between Standard Young Tableaux and reduced words of Grassmannian permutations. Sem. Loth. Comb. B36h (1996).

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115 *E-mail address*: misha@research.neu.edu

14