

ON TWISTED REPRESENTATIONS OF VERTEX ALGEBRAS

MICHAEL ROITMAN

ABSTRACT. In this paper we develop a formalism for working with twisted realizations of vertex and conformal algebras. As an example, we study realizations of conformal algebras by twisted formal power series. The main application of our technique is the construction of a very large family of representations for the vertex superalgebra \mathfrak{V}_Λ corresponding to an integer lattice Λ . For an automorphism $\hat{\sigma} : \mathfrak{V}_\Lambda \rightarrow \mathfrak{V}_\Lambda$ coming from a finite order automorphism $\sigma : \Lambda \rightarrow \Lambda$ we define a category $\mathcal{O}_{\hat{\sigma}}$ of twisted representations of \mathfrak{V}_Λ and show that this category is semisimple with finitely many isomorphism classes of simple objects.

INTRODUCTION

One of the most fascinating features of vertex algebras is their “sensitivity” to finite order automorphisms. This is best illustrated by the construction of Moonshine representation of the Monster simple group [3, 12].

In general, vertex algebras are represented by formal power series, also called fields, of the form

$$\alpha(z) = \sum_{n \in \mathbb{Z}} \alpha(n) z^{-n-1}, \quad \alpha(n) \in \mathfrak{gl}(V),$$

with coefficients in the algebra of linear operators on some space V . In this case V is a module over the vertex algebra. Having a finite order automorphism on a vertex algebra \mathfrak{A} means that there is a grading $\mathfrak{A} = \bigoplus_{\lambda \in \Gamma} \mathfrak{A}^\lambda$ on \mathfrak{A} by a finite cyclic group Γ . Assume that $\Gamma \subset \mathbb{C}/\mathbb{Z}$, then it is tempting to represent the vertex algebra \mathfrak{A} by the so-called *twisted* or *generalized* fields, involving non-integer powers of z , so that an element $\alpha \in \mathfrak{A}^\lambda$ is represented by series of the form

$$\alpha(z) = \sum_{n \equiv \lambda \bmod \mathbb{Z}} \alpha(n) z^{-n-1}, \quad \alpha(n) \in \mathfrak{gl}(V).$$

This indeed could be done, and then V becomes a *twisted module* over \mathfrak{A} . In fact this construction works for any group $\Gamma \subset \mathbb{C}/\mathbb{Z}$.

The idea of twisted realizations of vertex algebras goes back to the work of Lepowsky and Wilson [19]. Twisted modules were systematically studied in the book [12], see also [6, 8, 17, 18, 21]. Closely related is the theory of orbifolds — the invariant subalgebras of vertex algebras under an action of a finite group of automorphisms. Twisted representations of the vertex algebra yield ordinary (non-twisted) representations of its orbifold, see e.g. [1, 4].

In this paper we develop a formalism for working with twisted fields and use it to construct twisted representations of conformal and vertex algebras in a very general setting. Then we consider two applications of this techniques.

First we study realizations of conformal algebras by twisted formal series. Conformal algebras were introduced by Kac in the book [15], see also [16, 25]. They proved to be a valuable tool in studying vertex algebras, the relation between the former and the later is somewhat like the relation between Lie and associative algebras.

The second application is the construction of generalized representations of vertex (super)algebra \mathfrak{V}_Λ corresponding to an integer lattice Λ . These algebras were extensively studied in e.g. [5, 6, 7, 8, 12, 23]. Lattice vertex algebras play a very important role in different areas of mathematics and physics, in particular the Moonshine vertex algebra V^\natural , mentioned above, is closely related to the lattice vertex algebra of certain even unimodular lattice of rank 24, called the Leech lattice. In fact

Date: December 2, 2024.

This research was conducted by the author for the Clay Mathematics Institute.

one needs to consider a twisted representation of the vertex algebra of the Leech lattice in order to construct the Moonshine vertex algebra.

This paper is organized as follows. We start with giving a formal definitions of conformal and vertex algebras. For more details the reader can consult the books [7, 12, 15]. Then, in §1.2–§1.7 we discuss some properties of these algebras in the context of generalized formal series. In §1.3 we derive a nice formula for the products of generalized series. In §1.5 we prove that conformal and vertex algebras are exactly the algebraic structures formed by generalized series with coefficients in a Lie algebra and generalized fields respectively. For vertex algebras this was proved by Li [21] in a slightly less general setting and using different methods. For the non-twisted case this result is well known [15, 22].

In §2 we study the twisted realizations of a conformal algebra \mathfrak{L} by generalized formal series. As in the non-twisted case, there is a universal realization with coefficients in a certain Lie algebra $\text{Coeff}_\Gamma \mathfrak{L}$, which can be constructed explicitly from \mathfrak{L} . In §2.2 we illustrate this by the example of an affine conformal algebra. Similar ideas appeared also in [16].

In §3 we study generalized representations of vertex (super)algebra \mathfrak{V}_Λ corresponding to an integral lattice Λ . After some preliminary information on representations of Heisenberg algebras (§3.1) and Fock spaces (§3.2) we define in §3.3 the twisted vertex operators, first introduced by Lepowsky [18]. In §3.4 we show that these operators generate a representation of the lattice vertex algebra. A similar result was obtained by Dong and Lepowsky [8], but we use different techniques and get a slightly more general construction.

In §3.6 we show that we have in fact constructed all reasonable generalized representations of lattice vertex algebras. The argument uses an idea of Lepowsky and Wilson [20], which was also used in [5, 6, 12, 23].

Finally, in §3.7–§3.8 we study the twisted representations of \mathfrak{V}_Λ , i.e. homogeneous with respect to a gradation induced by a finite order automorphism of Λ . In §3.8 we define a natural category of twisted representations of \mathfrak{V}_Λ , and show that this category is semisimple with finitely many isomorphism classes of simple objects. This result has been also obtained recently by Bakalov, Kac and Todorov [1]. Some special cases were known before, see e.g. [6, 10, 17].

The key idea is to show that the category of twisted representation of \mathfrak{V}_Λ is equivalent to the category of graded representations of certain graded associative algebra A , which turns out to be graded semisimple. This idea is similar to the idea of Zhu algebra, introduced by Zhu [27] and then generalized for the twisted case by Dong, Li and Mason [9]. However our algebra A is quite different from the Zhu algebra of \mathfrak{V}_Λ .

Acknowledgements. I would like to thank Bojko Bakalov, Chongying Dong and Jim Lepowsky for the inspiring discussions. I am very grateful to Bojko Bakalov for communicating the unpublished manuscript [1].

1. VERTEX ALGEBRA

1.1. Definitions of conformal and vertex algebras. We start with the algebraic definitions of the main objects of this paper. All algebras and spaces are over a ground field \mathbb{k} of characteristic 0.

A *Conformal algebra* [15, 16, 25] is a vector space \mathfrak{L} equipped with a sequence of bilinear products $\boxed{n} : \mathfrak{L} \otimes \mathfrak{L} \rightarrow \mathfrak{L}$, $n \in \mathbb{Z}_+$, and a linear operator $D : \mathfrak{L} \rightarrow \mathfrak{L}$, such that the following axioms hold for all $a, b, c \in \mathfrak{L}$ and $n \in \mathbb{Z}_+$:

- C1. (Locality) $a \boxed{n} b = 0$ for $n \gg 0$.
- C2. $D(a \boxed{n} b) = (Da) \boxed{n} b + a \boxed{n} (Db) = -na \boxed{n-1} b + a \boxed{n} (Db)$.
- C3. (Quasisymmetry)

$$a \boxed{n} b = - \sum_{i \geq 0} (-1)^{n+i} \frac{1}{i!} D^i (b \boxed{n+i} a).$$

- C4. (Conformal Jacoby identity)

$$(a \boxed{n} b) \boxed{m} c = \sum_{i=0}^n (-1)^i \binom{n}{i} \left(a \boxed{n-i} (b \boxed{m+i} c) - b \boxed{m+i} (a \boxed{n-i} c) \right).$$

Example: The Virasoro conformal algebra \mathfrak{Vir} is generated over $\mathbb{k}[D]$ by elements v and c , such that $Dc = 0$ (and therefore all products with c are 0 due to C2) and the products of v with itself are

$$v \boxed{0} v = Dv, \quad v \boxed{1} v = 2v, \quad v \boxed{3} v = c. \quad (1)$$

The rest of the products are 0.

A *vertex algebra* can be defined axiomatically as follows [3, 11, 12, 15]. Let \mathfrak{A} be a linear space endowed with a sequence of bilinear operations $\boxed{n} : \mathfrak{A} \otimes \mathfrak{A} \rightarrow \mathfrak{A}$, $n \in \mathbb{Z}$, and a distinguished element $\mathbf{1} \in \mathfrak{A}$. Let $D : \mathfrak{A} \rightarrow \mathfrak{A}$ be a linear map given by $Da = a \boxed{-2} \mathbf{1}$. Then \mathfrak{A} is a vertex algebra if it satisfies the following conditions for any $a, b, c \in \mathfrak{A}$ and $m, n \in \mathbb{Z}$:

V1. (Locality) $a \boxed{n} b = 0$ for $n \gg 0$.

$$V2. \text{ (Identity) } \mathbf{1} \boxed{n} a = \delta_{n,-1} a, \quad a \boxed{n} \mathbf{1} = \begin{cases} 0 & \text{if } n \geq 0, \\ \frac{1}{(-n-1)!} D^{-n-1} a & \text{if } n < 0. \end{cases}$$

V3. (Associativity)

$$(a \boxed{n} b) \boxed{m} c = \sum_{s \leq n} (-1)^{s+n} \binom{n}{n-s} a \boxed{s} (b \boxed{m+n-s} c) - \sum_{s \geq 0} (-1)^{s+n} \binom{n}{s} b \boxed{m+n-s} (a \boxed{s} c).$$

V4. (Commutation)

$$a \boxed{m} (b \boxed{n} c) - b \boxed{n} (a \boxed{m} c) = \sum_{s \geq 0} \binom{m}{s} (a \boxed{s} b) \boxed{m+n-s} c.$$

This is not the shortest possible list of axioms. See the references cited above for other equivalent definitions.

For $n, m \geq 0$, the associativity V3 is exactly the conformal Jacobi identity C4. Among other properties of vertex algebras are the formulas C2 as well as the quasisymmetry identity C3, which holds for all integer n . So vertex algebras are a special case of conformal algebras.

There is an additional axiom which is often imposed on vertex algebras, see [7, 11, 12, 22]. A vertex algebra \mathfrak{A} is called a *vertex operator algebra* if

V5. $\mathfrak{A} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{A}_n$ is graded so that $\mathbf{1} \in \mathfrak{A}_0$ and $\mathfrak{A}_i \boxed{n} \mathfrak{A}_j \subset \mathfrak{A}_{i+j-n-1}$.

V6. There exists an element $v \in \mathfrak{A}_2$ generating the Virasoro conformal algebra $\mathfrak{Vir} \subset \mathfrak{A}$ (i.e. the relations (1) hold for $c = c\mathbf{1}$, $c \in \mathbb{k}$) such that $v \boxed{0} a = Da$, $v \boxed{1} a = (\deg a) a$ for all homogeneous $a \in \mathfrak{A}$.

The number c is called the *conformal charge* of \mathfrak{A} .

In the next section we show that conformal and vertex algebras are precisely the algebraic structures formed by certain formal infinite series.

1.2. Formal series and locality. Let L be a Lie algebra. Denote by $L\{z\}$ the \mathbb{k} -linear span of all series of the form

$$\sum_{n \in \lambda + \mathbb{Z}} \alpha(n) z^{-n-1}, \quad \alpha(n) \in L, \quad \lambda \in \mathbb{k}.$$

For a linear space V , let $F\{V\} \subset \mathfrak{gl} V\{z\}$ be the space of all such series with coefficients in the Lie algebra $\mathfrak{gl} V$ with the property that $\alpha(n)v = 0$ for $n \gg 0$ for any fixed $v \in V$. We call $F\{V\}$ the space of *generalized fields*. It contains the space $F(V)$ of ordinary fields, that involve only integer powers of z . Denote by $\mathbf{1} \in F\{V\}$ the field with the only non-zero coefficient being $\mathbf{1}(-1) = \text{id}$.

The space $L\{z\} = \bigoplus_{[\lambda] \in \mathbb{k}/\mathbb{Z}} L\{z\}^{[\lambda]}$ is graded by the group \mathbb{k}/\mathbb{Z} so that $L\{z\}^{[\lambda]}$ is the space of all series of the form $z^{-\lambda} \alpha_\lambda(z)$, $\alpha_\lambda(z) \in L[[z^{\pm 1}]]$. We have $L[[z^{\pm 1}]] = L\{z\}^{[0]}$. The space of fields $F\{V\}$ is a homogeneous subspace of $\mathfrak{gl} V\{z\}$ and $F(V) = F\{V\}^{[0]}$.

A pair of series $\alpha, \beta \in L\{z\}$ are said to be *local of order* $N \in \mathbb{Z}_+$ [7, 15] if

$$(z-w)^N [\alpha(w), \beta(z)] = 0,$$

or, equivalently,

$$\sum_{s=0}^N (-1)^s \binom{N}{s} [\alpha(n-s), \beta(m+s)] = 0 \quad \forall m, n \in \mathbb{k}.$$

The same applies for $\alpha, \beta \in F\{V\}$. It is easy to see that if series $\alpha = \sum_{\lambda} z^{-\lambda} \alpha_{\lambda}$, $\beta = \sum_{\lambda} z^{-\lambda} \beta_{\lambda} \in L\{z\}$ are local of order N , then any two homogeneous components $\alpha_{\lambda}, \beta_{\mu} \in L[[z^{\pm 1}]]$ are local of the same order.

1.3. The products of formal series. Recall that for two ordinary fields $\alpha, \beta \in F(V)$, one can define products $\alpha \boxed{n} \beta \in F(V)$ in the following way. Let $\iota_{w,z} : \mathbb{k}(w, z) \rightarrow \mathbb{k}((w^{-1}, z))$ and $\iota_{z,w} : \mathbb{k}(w, z) \rightarrow \mathbb{k}((w, z^{-1}))$ be the expansions of a rational function into Laurent series at $(w, z) = (\infty, 0)$ and $(w, z) = (0, \infty)$ respectively, so that

$$\begin{aligned} \iota_{w,z}(w-z)^n &= \sum_{i \geq 0} (-1)^{n+i} \binom{n}{i} w^{n-i} z^i, \\ \iota_{z,w}(w-z)^n &= \sum_{i \geq 0} (-1)^i \binom{n}{i} w^i z^{n-i}. \end{aligned} \quad (2)$$

Of course, if $n \geq 0$ then $\iota_{w,z}(w-z)^n = \iota_{z,w}(w-z)^n$. We define

$$(\alpha \boxed{n} \beta)(z) = \text{Res}_w \left(\alpha(w) \beta(z) \iota_{w,z}(w-z)^n - \beta(z) \alpha(w) \iota_{z,w}(w-z)^n \right). \quad (3)$$

The m -th coefficient of $\alpha \boxed{n} \beta$ is given by

$$(\alpha \boxed{n} \beta)(m) = \sum_{s \leq n} (-1)^{s+n} \binom{n}{n-s} \alpha(s) \beta(m+n-s) - \sum_{s \geq 0} (-1)^{s+n} \binom{n}{s} \beta(m+n-s) \alpha(s). \quad (4)$$

If $n \geq 0$ then the products $\alpha \boxed{n} \beta$ make sense for formal series $\alpha, \beta \in L[[z^{\pm 1}]]$ as well. In this case (4) simplifies to

$$(\alpha \boxed{n} \beta)(m) = \sum_s (-1)^s \binom{n}{s} [\alpha(n-s), \beta(m+s)].$$

One can solve these equations with respect to the commutators and thus recover the bracket in L from the products in $L[[z^{\pm 1}]]$:

$$[\alpha(m), \beta(n)] = \sum_s \binom{m}{s} (\alpha \boxed{s} \beta)(m+n-s) \quad (5)$$

for every $m \in \mathbb{Z}_+$, $n \in \mathbb{Z}$. If α and β are local, then (5) holds for all $m, n \in \mathbb{Z}$.

Now we show how to expand these products to $F\{V\}$ and $L\{z\}$. For homogeneous $\alpha = z^{\lambda} \alpha_1$ and $\beta = z^{\mu} \beta_1$, where $\lambda, \mu \in \mathbb{k}$ and α_1, β_1 involve only integer powers of z , we set

$$\alpha \boxed{n} \beta = \sum_{j \geq 0} \binom{\lambda}{j} (\alpha_1 \boxed{n+j} \beta_1) z^{\mu+\lambda-j}. \quad (6)$$

Here

$$\binom{\lambda}{j} = \frac{\lambda(\lambda-1) \cdots (\lambda-j+1)}{j!}.$$

Note that the summation in (6) is finite due to the locality of α and β , and hence of α_1 and β_1 . One can easily check that the products (3) satisfy the identity (6) if we substitute $\alpha = z^{\lambda} \alpha_1$, $\beta = z^{\mu} \beta_1$ for $\lambda, \mu \in \mathbb{Z}$. We extend the definition of the products \boxed{n} by linearity to arbitrary $\alpha, \beta \in F\{V\}$ and $\alpha, \beta \in L\{z\}$ when $n \geq 0$.

It is clear that if α and β are local of order N then $\alpha \boxed{n} \beta = 0$ for $n \geq N$.

It could be shown that (5) remains valid for generalized series as well if $m, n \in \mathbb{k}$ and $\deg \alpha = [m]$, $\deg \beta = [n] \in \mathbb{k}/\mathbb{Z}$.

We now write explicitly the formula for the products of twisted fields $\alpha, \beta \in F\{V\}$, analogous to (3). Let N be the order of locality of α and β . Denote

$$\Delta(w, z) = \sum_{\lambda} \sum_{j=0}^{N-n-1} \binom{-\lambda}{j} w^{\lambda} z^{-\lambda-j} (w-z)^j,$$

where λ runs over the set of degrees that appear in α .

Then

$$\begin{aligned} (\alpha \boxed{n} \beta)(z) &= \sum_{\lambda} (z^{-\lambda} \alpha_{\lambda}) \boxed{n} \beta = \sum_{\lambda} \sum_{j \geq 0} \binom{-\lambda}{j} (\alpha \boxed{n+j} \beta) z^{-\lambda-j} \\ &= \text{Rez}_w \left(\left(\alpha(w) \beta(z) \iota_{w,z}(w-z)^n - \beta(z) \alpha(w) \iota_{z,w}(w-z)^n \right) \Delta(w, z) \right) \end{aligned} \quad (7)$$

1.4. The case of finite grading. Of a particular interest is the case when all the degrees of α are rational numbers with common denominator p , i.e.

$$\alpha(z) = \sum_{q=0}^{p-1} z^{-q/p} \alpha_q(z), \quad \alpha_q \in F(V).$$

In this case we can rewrite (7) in a different way. Some calculations show that

$$\Delta(w, z) = \sum_{q=0}^{p-1} \sum_{j=0}^m \binom{-q/p}{j} w^{\frac{q}{p}} z^{-\frac{q}{p}-j} (w-z)^j = \left(\frac{w-z}{w^{\frac{1}{p}} - z^{\frac{1}{p}}} \right)^{m+1} F_p(m),$$

where

$$F_p(m) = \sum_{l=1-p}^{1-p+m} \left(\sum_{q=0}^{p-1} \sum_{k \geq 0} (-1)^{l+q+kp} \binom{-q/p+k}{m} \binom{m+1}{l+q-kp} \right) w^{\frac{m-l+1}{p}-1} z^{\frac{l}{p}-m}.$$

Using this we get

$$\begin{aligned} (\alpha \boxed{n} \beta)(z) &= \text{Rez}_w \left(\left(\alpha(w) \beta(z) \iota_{w,z}(w-z)^n - \beta(z) \alpha(w) \iota_{z,w}(w-z)^n \right) \right. \\ &\quad \left. \times \left(\frac{w-z}{w^{\frac{1}{p}} - z^{\frac{1}{p}}} \right)^{N-n} F_p(N-n-1) \right). \end{aligned} \quad (8)$$

We also remark that the polynomial $F_p(m)$ has the following property:

$$F_p(m) \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} = p^{-m} z^{\frac{(m+1)(1-p)}{p}}. \quad (9)$$

1.5. Algebras of formal series. It is well known [15, 22] that a subspace $\mathfrak{V} \subset F(V)$ of ordinary fields such that $\mathbf{1} \in \mathfrak{V}$ and $\mathfrak{V} \boxed{n} \mathfrak{V} \subset \mathfrak{V}$ for all $n \in \mathbb{Z}$ is a vertex algebra. Similarly, a subspace $\mathfrak{L} \subset L[[z^{\pm 1}]]$ such that $\mathfrak{L} \boxed{n} \mathfrak{L} \subset \mathfrak{L}$ is a conformal algebra. Moreover, all vertex and conformal algebras are obtained in that way. Now we generalize this result to the case of twisted fields and series.

Theorem 1. a. Let $\mathfrak{A} \subset F\{V\}$ be a subspace of pairwise local generalized fields containing $\mathbf{1}$, closed with respect to the products \boxed{n} , $n \in \mathbb{Z}$. Then \mathfrak{A} is a vertex algebra.
b. Let $\mathfrak{L} \subset L\{z\}$ be a subspace of pairwise local twisted series with coefficients in a Lie algebra L , closed with respect to the products \boxed{n} , $n \in \mathbb{Z}_+$. Then \mathfrak{L} is a conformal algebra.

Proof. We will prove (a), the statement (b) is proved in the same way. We have to show that \mathfrak{A} satisfies the identities V2-V4, since V1 is true by assumption. All these identities are linear combinations of “vertex monomials” of the form $a_1 \boxed{n_1} \dots \boxed{n_{l-1}} a_l$ (with some order of parentheses) where some of a_i ’s could be equal to $\mathbf{1}$. Note that these identities are multilinear.

Let $\mathfrak{A}^{\text{gr}} \subset F\{V\}$ be the graded closure of \mathfrak{A} , i.e. the minimal graded subspace of $F\{V\}$ containing \mathfrak{A} . By the above remark, \mathfrak{A}^{gr} satisfies all the assumptions of the theorem, so we can assume that $\mathfrak{A} = \mathfrak{A}^{\text{gr}}$ is graded.

Consider all the fields $\beta \in F(V)$ such that $z^\lambda \beta \in \mathfrak{A}$ for some $\lambda \in \mathbb{k}$. Every two such fields are local, so since the theorem is known to be true in the non-twisted case, these fields generate a vertex algebra $\mathfrak{B} \subset F(V)$.

Let $R(a_1, \dots, a_l) = 0$ be an identity which we have to check. Since R is multilinear it is enough to check it for homogeneous $a_i = z^{-\lambda_i} b_i$, where $b_i \in \mathfrak{B}$, $\lambda_i \in \mathbb{k}$, $1 \leq i \leq l$. Sometimes we must set

$b_i = \mathbb{1}$ and $\lambda_i = 0$. When we substitute these expressions for a_i into $R(a_1, \dots, a_k) = 0$ and apply (6), we get a linear combination of conformal monomials of the form

$$P(\lambda_1, \dots, \lambda_l) z^{-\lambda_1 - \dots - \lambda_l} z^m (b_{\kappa(1)} \boxed{n_1} \dots \boxed{n_{l-1}} b_{\kappa(l)})$$

where $P(\lambda_1, \dots, \lambda_l)$ is a polynomial in $\lambda_1, \dots, \lambda_l$, $m \in \mathbb{Z}$, κ is a permutation of $\{1, 2, \dots, l\}$ and the products n_i 's are applied according to some order of parentheses. We can cancel the common factor $z^{-\lambda_1 - \dots - \lambda_l}$. Now we observe that the only remaining factor in these monomials that depends on λ_i 's is P , therefore for fixed b_1, \dots, b_l the map $(\lambda_1, \dots, \lambda_l) \mapsto R(z^{-\lambda_1} b_1, \dots, z^{-\lambda_l} b_l)$ is a polynomial map from \mathbb{k}^l to $F\{V\}$. But when all $\lambda_i \in \mathbb{Z}$ this map is equal to 0, because then $a_i \in F(V)$ generate a vertex algebra which satisfies the identity R , therefore, since $|\mathbb{k}| = \infty$, R must be identically 0. \square

Remark. The products of the twisted fields were first introduced by Li in [21]. However he deals only with generating functions of these products, formulas (7) and (8) seem to be new. Li also proves Theorem 1(a) using more straightforward techniques. Realizations of conformal algebras by generalized series were mentioned by Kac in [16].

1.6. Differentiation of fields. Assume that we are given a pair (V, D) consisting of a linear space V and a linear map $D : V \rightarrow V$. Let $\Delta : F\{V\} \rightarrow F\{V\}$ be a linear operator defined by $\Delta\phi = \frac{d}{dz}\phi - [D, \phi]$. A field $\phi \in F\{V\}$ is called to be of weight $\lambda \in \mathbb{k}$ if $\Delta\phi = \lambda z^{-1}\phi$. Denote by $F\{V\}_\lambda$ the space of all fields of weight λ . It is easy to see that if $\phi \in F\{V\}_\lambda$ then $z^\mu\phi \in F\{V\}_{\lambda+\mu}$.

In general it is not true that any field can be represented as a sum of homogeneous fields. However the following is true.

Proposition 1. *Let $S \subset F\{V\}$ be a set of pairwise local fields, and let $\mathfrak{A} \subset F\{V\}$ be the vertex algebra generated by S . If $S \subset F\{V\}_0$ then also $\mathfrak{A} \subset F\{V\}_0$ and if $S \subset \bigoplus_\lambda F\{V\}_\lambda$ then also $\mathfrak{A} \subset \bigoplus_\lambda F\{V\}_\lambda$.*

Proof. First of all we note that Δ is a derivation of all products: $\Delta(\alpha \boxed{n} \beta) = (\Delta\alpha) \boxed{n} \beta + \alpha \boxed{n} (\Delta\beta)$, because so are both $\text{ad } D$ and $\frac{d}{dz}$. Therefore if $\Delta\alpha = \Delta\beta = 0$ then $\Delta(\alpha \boxed{n} \beta) = 0$ and the first statement follows. For the second statement, it is enough to assume that all the generators from S are homogeneous. A pair $\alpha, \beta \in S$ can be written as $\alpha = z^\lambda \alpha_0$, $\beta = z^\mu \beta_0$ for some $\lambda, \mu \in \mathbb{k}$, where $\alpha_0, \beta_0 \in F\{V\}_0$. Now the statement follows from the formula (6). \square

1.7. Modules over vertex algebras. Now we give several definitions of modules for vertex algebras. We will call a vector space V a *module* over a vertex algebra \mathfrak{A} if there is a vertex algebra homomorphism $\pi : \mathfrak{A} \rightarrow F(V)$. In other words, for any $a, b \in \mathfrak{A}$, the fields $\pi(a)$ and $\pi(b)$ are local and $\pi(a \boxed{n} b) = \pi(a) \boxed{n} \pi(b)$, $\pi(\mathbb{1}) = \mathbb{1}$. We remark that sometimes what we call a module is called a weak module.

If instead of a homomorphism $\pi : \mathfrak{A} \rightarrow F(V)$ we have a homomorphism $\pi : \mathfrak{A} \rightarrow F\{V\}$ of \mathfrak{A} into the space of generalized fields over V , then V is called a *generalized module*.

Assume now that a vertex algebra $\mathfrak{A} = \bigoplus_{[\lambda] \in \Gamma} \mathfrak{A}^{[\lambda]}$ is graded by a group $\Gamma \subset \mathbb{k}/\mathbb{Z}$. Then a generalized module V is called *twisted* if the homomorphism $\pi : \mathfrak{A} \rightarrow F\{V\}$ is homogeneous, that is, $\pi(\mathfrak{A}^{[\lambda]}) \subset F\{V\}^{[\lambda]}$. This definition is due to Li [21]. Equivalently, twisted modules can be defined using the so-called twisted Jacobi identity, see e.g. [7, 9].

Assume that an \mathfrak{A} -module V (generalized or not) has a linear map $D : V \rightarrow V$. Then the module V is called *strong* if $\pi(\mathfrak{A}) \subset F\{V\}_0$, see §1.6. We note that if \mathfrak{A} contains an element v such that $Y(v)(0) = D$ (as it is the case if \mathfrak{A} is a vertex operator algebra), and V is a module over \mathfrak{A} such that $\pi(v) \in F\{V\}^{[0]}$, then V is a strong module. Indeed, by (5) we have $[v(0), a(n)] = (v \boxed{n} a)(n) = (Da)(n)$ for any $a \in \mathfrak{A}^{[n]}$.

Let again V be a module (generalized or not) over \mathfrak{A} and let $\pi : \mathfrak{A} \rightarrow F\{V\}$ be the representation map. Assume \mathfrak{A} is a vertex operator algebra and let $v \in \mathfrak{A}$ be the Virasoro element. We say that V is a module over the vertex *operator* algebra if $V = \bigoplus_{n \in \mathbb{k}} V_n$ is graded, $\pi(v) \in F\{V\}^{[0]}$ and $\pi(v)(1)|_{V_n} = n$. As it was mentioned above, in this case V is necessary a strong module over \mathfrak{A} .

Remark. Let \mathfrak{A} be a vertex algebra and let V be a strong twisted module over \mathfrak{A} . Then the semidirect product $\mathfrak{A} \ltimes V$ has a structure of *generalized vertex algebra*, introduced by Dong and Lepowsky [7]. The products in a generalized vertex algebra are indexed not necessarily by integers. If $a \in \mathfrak{A}$ and

$v \in V$ then $a \square v = a(n)v$ and the products $v \square a$ are defined using the quasismmetry identity C3, see §1.1.

2. TWISTED REALIZATIONS OF CONFORMAL ALGEBRAS

2.1. The coefficient algebra. Let \mathfrak{L} be a conformal algebra. Assume that $\mathfrak{L} = \bigoplus_{[\lambda] \in \Gamma} \mathfrak{L}^{[\lambda]}$ is graded by a group $\Gamma \subset \mathbb{K}/\mathbb{Z}$ so that $\mathfrak{L}^{[\lambda]} \square \mathfrak{L}^{[\mu]} \subseteq \mathfrak{L}^{[\lambda+\mu]}$, $D\mathfrak{L}^{[\lambda]} \subseteq \mathfrak{L}^{[\lambda]}$.

Define a Lie algebra $\text{Coeff}_\Gamma \mathfrak{L}$ in the following way. The underlying linear space of $\text{Coeff}_\Gamma \mathfrak{L}$ is spanned by the symbols $a(n)$ for all homogeneous $a \in \mathfrak{L}^{[n]}$ subject to the linear relations $(Da)(n) = -na(n-1)$. The brackets in $\text{Coeff}_\Gamma \mathfrak{L}$ are defined by the formula (5) for $a = \alpha \in \mathfrak{L}^{[m]}$ and $b = \beta \in \mathfrak{L}^{[n]}$. For $a \in \mathfrak{L}$ denote $a(n) = a^{[n]}(n)$, where $a^{[n]}$ is the projection of a onto the space $\mathfrak{L}^{[n]}$.

This construction generalizes the construction of usual coefficient algebra $\text{Coeff } \mathfrak{L} = \text{Coeff}_{[0]} \mathfrak{L}$ done in [15, 16, 26]. If \mathfrak{L} is a vertex algebra, then $\text{Coeff}_\Gamma \mathfrak{L}$ was considered in [9], at least when Γ is a finite cyclic group.

There is a map $\phi : \mathfrak{L} \rightarrow \text{Coeff}_\Gamma \mathfrak{L}\{z\}$ given by $\phi(a) = \sum_{n \in \Gamma + \mathbb{Z}} a(n) z^{-n-1}$. It is easy to see that ϕ is a homomorphism of \mathfrak{L} into a conformal subalgebra of $\text{Coeff}_\Gamma \mathfrak{L}\{z\}$ such that $\phi(\mathfrak{L}^{[\lambda]}) \subseteq \text{Coeff}_\Gamma \mathfrak{L}\{z\}^{[\lambda]}$. Moreover, this map is universal in the following sense: if $\rho : \mathfrak{L} \rightarrow K\{z\}$ is another homogeneous homomorphism of \mathfrak{L} into a conformal subalgebra of series with coefficients in some Lie algebra K , then there use unique Lie algebra homomorphism $\pi : \text{Coeff}_\Gamma \mathfrak{L} \rightarrow K$ making the following diagram commutative:

$$\begin{array}{ccc} \text{Coeff}_\Gamma \mathfrak{L}\{z\} & \xrightarrow{\pi} & K\{z\} \\ \phi \swarrow & & \nearrow \rho \\ & \mathfrak{L} & \end{array} \quad (10)$$

The proof of the fact that $\text{Coeff}_\Gamma \mathfrak{L}$ is indeed a Lie algebra and of the above universality property is done in the same way as in the non-twisted case.

We also give another construction of $\text{Coeff}_\Gamma \mathfrak{L}$. Let $L = \text{Coeff } \mathfrak{L}$ be the ordinary coefficient Lie algebra of \mathfrak{L} . Let $\mathfrak{L}_\Gamma \subset L\{z\}$ be the conformal algebra generated by all the series of the form $z^n a$ for $a \in \mathfrak{L}^{[n]}$. The algebra \mathfrak{L}_Γ is closed under the multiplication by $z^{\pm 1}$, in fact we have $z^{-\lambda} \mathfrak{L}_\Gamma^{[\lambda]} = \mathbb{K}[z^{\pm 1}] \mathfrak{L}^{[\lambda]}$.

Let $L_\Gamma \subset \text{Coeff } \mathfrak{L}_\Gamma$ be the subalgebra consisting of all elements of the form $a(0)$ for $a \in \mathfrak{L}_\Gamma$. In other words, $L_\Gamma = \text{Ker } D$, where $D : \text{Coeff } \mathfrak{L}_\Gamma \rightarrow \text{Coeff } \mathfrak{L}_\Gamma$ is the derivation of $\text{Coeff } \mathfrak{L}_\Gamma$ given by $a(n) \mapsto -na(n-1)$. There is a Lie algebra homomorphism $\text{Coeff } \mathfrak{L}_\Gamma \rightarrow L_\Gamma$ given by $a(n) \mapsto (z^n a)(0)$ for $a \in \mathfrak{L}^{[n]}$, which induces the conformal algebra homomorphism $\eta : \mathfrak{L}_\Gamma \rightarrow L_\Gamma[[w^{\pm 1}]]$ such that $\eta(z^k a) = w^k \eta(a)$ for $k \in \mathbb{Z}$. It could be shown that η is the universal among all realizations of \mathfrak{L}_Γ by integral formal series with this property.

We construct a homomorphism $\rho : \mathfrak{L} \rightarrow L_\Gamma\{w\}$ in the following way. Let $a \in \mathfrak{L}^{[\lambda]}$, define $\rho(a) = \eta(z^\lambda a) w^{-\lambda} \in L_\Gamma\{w\}$. Clearly, ρ is a linear map and it does not depend on the choice of representative $\lambda \in [\lambda]$. Now we show that ρ is indeed a homomorphism.

Let $a_1, \dots, a_k \in \mathfrak{L}$ be homogeneous elements of \mathfrak{L} satisfying a conformal identity $R(a_1, \dots, a_k) = 0$. Here R is a linear combination of conformal monomials in a_i 's with coefficients in $\mathbb{K}[D]$. We have to show that $R(\rho(a_1), \dots, \rho(a_k)) = 0$ in $L_\Gamma\{w\}$. Without loss of generality we can assume that R is homogeneous with respect to the gradation by Γ . Let $[\lambda_i] = \deg a_i$. Substitute $a_i = z^{-\lambda_i} b_i$ in R , apply the formula (6) and then cancel the common factor $z^{-\lambda_1 - \dots - \lambda_k}$. We get an identity $R_1(b_1, \dots, b_k) = 0$, which holds in $\mathfrak{L}_\Gamma \subset L\{z\}$, where R_1 is a combination of conformal monomials in b_1, \dots, b_k with coefficients in $\mathbb{K}[D, z^{\pm 1}]$. Since $\eta : \mathfrak{L}_\Gamma \rightarrow L_\Gamma[[w^{\pm 1}]]$ is a $\mathbb{K}[D, z^{\pm 1}]$ -module homomorphism, we have $R_1(\eta(b_1), \dots, \eta(b_k)) = 0$ in $L_\Gamma[[w^{\pm 1}]]$. Substitute now $\eta(b_i) = w^{\lambda_i} \rho(a_i)$ and apply (6) again. After dividing by $w^{\lambda_1 + \dots + \lambda_k}$, we get $R(\rho(a_1), \dots, \rho(a_k)) = 0$.

By the universality of $\text{Coeff}_\Gamma \mathfrak{L}$ we get a map $\pi : \text{Coeff}_\Gamma \mathfrak{L} \rightarrow L_\Gamma$ such that the diagram (10) commutes for $K = L_\Gamma$.

Proposition 2. *The map $\pi : \text{Coeff}_\Gamma \mathfrak{L} \rightarrow L_\Gamma$ is an isomorphism.*

Proof. We will show that the homomorphism $\rho : \mathfrak{L} \rightarrow L_\Gamma\{w\}$ constructed above has the same universality property as $\text{Coeff}_\Gamma \mathfrak{L}$. Then the fact that $\pi : \text{Coeff}_\Gamma \mathfrak{L} \rightarrow L_\Gamma$ is an isomorphism follows from uniqueness of $\text{Coeff}_\Gamma \mathfrak{L}$.

Let $\psi : \mathfrak{L} \rightarrow K\{w\}$ be a homogeneous homomorphism of \mathfrak{L} into a space of generalized series with coefficients in some Lie algebra K . This induces a homomorphism $\theta : \mathfrak{L}_\Gamma \rightarrow K[[w^{\pm 1}]]$ by $\theta(z^n a) = w^n \psi(a)$ for any $a \in \mathfrak{L}^{[\lambda]}$ and $n \in \lambda + \mathbb{Z}$. It is easy to see that $\theta(z^n a) = w^n \theta(a)$ for every $a \in \mathfrak{L}_\Gamma$ and $n \in \mathbb{Z}$. Hence there is a homomorphism $L_\Gamma \rightarrow K$ such that all the corresponding diagrams commute. \square

2.2. Twisted affine algebras. We illustrate the construction of §2.1 by the example of affine conformal algebras.

Let \mathfrak{g} be a Lie algebra with an invariant bilinear form $(\cdot | \cdot)$, i.e. such that $([a, b] | c) = (a | [b, c])$. Assume that the algebra $\mathfrak{g} = \bigoplus_{[\lambda] \in \Gamma} \mathfrak{g}^{[\lambda]}$ is graded by a group $\Gamma \subset \mathbb{k}/\mathbb{Z}$. Assume further that the gradation on \mathfrak{g} agrees with the form $(\cdot | \cdot)$ in the following way:

$$(\mathfrak{g}^{[\lambda]} | \mathfrak{g}^{[\mu]}) = 0 \quad \text{unless} \quad \lambda + \mu = 0. \quad (11)$$

Consider the Lie algebra $\widehat{L}_\Gamma = \mathfrak{g} \otimes \mathbb{k}[\Gamma + \mathbb{Z}] \oplus \mathbb{k}c$, where $\mathbb{k}[\Gamma + \mathbb{Z}]$ is the group algebra of $\Gamma + \mathbb{Z}$. We will write $a(n) = a \otimes n$ for $a \in \mathfrak{g}$, $n \in \Gamma + \mathbb{Z}$. The brackets in \widehat{L}_Γ are defined by

$$[a(m), b(n)] = [a, b](m+n) + \delta_{m+n,0} m (a^{[m]} | b^{[n]}) c, \quad c \in Z(\widehat{L}_\Gamma),$$

where $a^{[m]}$ and $b^{[n]}$ are projections of a and b onto $\mathfrak{g}^{[m]}$ and $\mathfrak{g}^{[n]}$ respectively. The twisted affine Lie algebra $L_\Gamma \subset \widehat{L}_\Gamma$ is the subalgebra of \widehat{L}_Γ spanned by c and all elements of the form $a(n)$ for $a \in \mathfrak{g}^{[n]}$. The grading on \mathfrak{g} induces a grading on L_Γ by setting $\deg a(n) = \deg a$, $\deg c = 0$. From now on if $a \otimes n \in \widehat{L}_\Gamma \setminus L_\Gamma$, we set $a(n) = 0$.

For any $a \in \mathfrak{g}$ consider series $\widetilde{a} = \sum_{n \in \Gamma + \mathbb{Z}} a(n) z^{-n-1} \in L_\Gamma\{z\}$. These series are local of order 2 and together with the series $c = c z^0$ they generate a conformal algebra $\mathfrak{L} \subset L_\Gamma\{z\}$. The products between \widetilde{a} 's are

$$\widetilde{a} \square \widetilde{b} = \widetilde{[a, b]}, \quad \widetilde{a} \sqcap \widetilde{b} = (a|b) c. \quad (12)$$

The affine conformal algebra \mathfrak{L} is a homogeneous subalgebra of $L_\Gamma\{z\}$ so that for $a \in \mathfrak{g}^{[\lambda]}$ we have $\widetilde{a} \in L_\Gamma\{z\}^{[\lambda]}$. It is independent on the Γ -grading of \mathfrak{g} . It is easy to see that $L_\Gamma = \text{Coeff}_\Gamma \mathfrak{L}$.

For each equivalence class $[\lambda] \in \Gamma \subset \mathbb{k}/\mathbb{Z}$ choose a representative $\lambda \in \mathbb{k}$ such that the representative of \mathbb{Z} is 0 and if $[\lambda] + [\mu] = 0$ then either $\lambda = \mu = 0$ or $\lambda + \mu = 1$. For example, if $\mathbb{k} \subseteq \mathbb{R}$, then we can take $0 \leq \lambda < 1$. A very important special case is when Γ is a finite cyclic group of order p , then one can take the set of representatives of Γ in \mathbb{k} to be $\{0, \frac{1}{p}, \frac{2}{p}, \dots, \frac{p-1}{p}\}$.

For a homogeneous element $a \in \mathfrak{g}^{[\lambda]}$ consider the series $\tau_a = z^\lambda \widetilde{a} \in L_\Gamma[[z^{\pm 1}]]$. Clearly, the series τ_a are pairwise local of order 2, so together with c they generate a conformal algebra $\mathfrak{L}_\Gamma \subset L_\Gamma[[z^{\pm 1}]]$, called a twisted affine conformal algebra. We calculate the non-zero products of these series, using (6) and (12). Here $a, b \in \mathfrak{g}$, $\deg a = [\lambda]$, $\deg b = [\mu]$.

$$\begin{aligned} \tau_a \square \tau_b &= z^{\lambda+\mu} \widetilde{[a, b]} + \lambda z^{\lambda+\mu-1} (a|b) c = \begin{cases} z \tau_{[a, b]} + \lambda (a|b) c & \text{if } \lambda + \mu \geq 1, \\ \tau_{[a, b]} & \text{if } \lambda + \mu < 1, \end{cases} \\ \tau_a \sqcap \tau_b &= z^{\lambda+\mu} (a|b) c = \begin{cases} (a|b) c & \text{if } \lambda = \mu = 0, \\ z (a|b) c & \text{if } \lambda + \mu = 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

It is not difficult to see that L_Γ is isomorphic to the subalgebra of $\text{Coeff } \mathfrak{L}_\Gamma$ consisting of all elements of the form $a(0)$, i.e., the embedding $\mathfrak{L}_\Gamma \rightarrow L_\Gamma[[z^{\pm 1}]]$ is the universal one among realizations of \mathfrak{L}_Γ by formal series that agree with the multiplication by z , see §2.1.

Remark. While in general we cannot guarantee that the twisted representation map $\rho : \mathfrak{L} \rightarrow L_\Gamma\{w\}$ is injective, in the particular case when \mathfrak{L} is an affine conformal algebra we do know that ρ is an isomorphism.

Remark. In principal, one can use the construction of §2.1 to get twisted representations of a vertex algebra. Namely, let $\mathfrak{A} = \bigoplus_{[\lambda] \in \Gamma} \mathfrak{A}^{[\lambda]}$ be a graded vertex algebra, and let $\pi : \mathfrak{A} \rightarrow F(V)$ be its representation. Let $\mathfrak{A}_\Gamma \subset F\{V\}$ be the vertex algebra generated by the generalized fields $z^n \pi(a)$

for $a \in \mathfrak{A}^{[n]}$. Then \mathfrak{A}_Γ will be closed under the multiplication by $z^{\pm 1}$. If we have a representation $\eta : \mathfrak{A}_\Gamma \rightarrow F(U)$ such that $\eta(z^k a) = w^k \eta(a)$ for $k \in \mathbb{Z}$, then the map $a \mapsto \eta(z^\lambda a) w^{-\lambda}$ for $a \in \mathfrak{A}^{[\lambda]}$ defines a twisted representation of \mathfrak{A} on U . However, in contrast with the conformal case, it is not clear how to construct such a representation η of \mathfrak{A}_Γ . On the other hand, if \mathfrak{A} is generated by a conformal algebra $\mathfrak{L} \subset \mathfrak{A}$, then applying the construction of §2.1 to \mathfrak{L} we can get a twisted representation of possibly some other enveloping vertex algebra $\mathfrak{B} \supset \mathfrak{L}$.

3. LATTICE VERTEX ALGEBRAS

In this section we assume that the ground field is \mathbb{C} and the group Γ is the cyclic group of order $p \in \mathbb{Z}_+$. We identify Γ with a subgroup of \mathbb{C}/\mathbb{Z} by $q \mapsto \frac{q}{p} + \mathbb{Z}$, $0 \leq q < p$. We remark that some of the constructions below can be done for a more general group $\Gamma \subset \mathbb{C}/\mathbb{Z}$.

3.1. Representation theory of Heisenberg algebras. Let \mathfrak{h} be a vector space of a finite dimension $\dim \mathfrak{h} = l$ equipped with a non-degenerate bilinear form $(\cdot | \cdot)$. Assume that $\mathfrak{h} = \bigoplus_{\lambda \in \Gamma} \mathfrak{h}^{[\lambda]}$ is graded by Γ and the grading agrees with the form in sense of (11). Recall that the grading induces an automorphism $\sigma : \mathfrak{h} \rightarrow \mathfrak{h}$ such that $\sigma|_{\mathfrak{h}^{[\lambda]}} = \exp(2\pi i \lambda)$. In fact the existence of an automorphism $\sigma : \mathfrak{h} \rightarrow \mathfrak{h}$ of order p is equivalent to the existence of the above grading of \mathfrak{h} by the group $\Gamma = \mathbb{Z}/p\mathbb{Z}$. Note that σ preserves the norm on \mathfrak{h} . For $h \in \mathfrak{h}$ we denote by $h^{[\lambda]}$ the projection of h onto $\mathfrak{h}^{[\lambda]}$.

View \mathfrak{h} as an abelian Lie algebra, and let $H = H_\Gamma = \text{Span}\{a(n) \mid a \in \mathfrak{h}^{[n]}\} \oplus \mathbb{C}c$ be the corresponding (twisted, unless $p = 1$) affine Lie algebra, see §2.2. It is usually called a Heisenberg Lie algebra. As in §2.2, for an element $a \in \mathfrak{h}$ consider formal series $\tilde{a} = \sum_{n \in \Gamma + \mathbb{Z}} a(n) z^{-n-1} \in H\{z\}$. These series, together with c , span over $\mathbb{C}[D]$ a copy of conformal Heisenberg algebra $\mathfrak{H} \subset H\{z\}$, so that $H = \text{Coeff}_\Gamma \mathfrak{H}$. The grading on \mathfrak{h} lifts to a grading on H and \mathfrak{H} and the automorphism σ lifts to automorphisms of H and \mathfrak{H} . Recall also that H is graded by setting $\deg a(n) = -n \in \frac{1}{p}\mathbb{Z}$.

We note that $\mathfrak{h}^{[0]} \subset Z(H)$ so that $H = H' \oplus \mathfrak{h}^{[0]}$, where $H' = \text{Span}\{a(n) \mid a \in \mathfrak{h}, n \in \mathbb{C}\} \cup \{c\}$. Denote $H_\pm = \text{Span}\{a(n) \mid n \gtrless 0\} \subset H'$. We have $H' = H_- \oplus \mathbb{C}c \oplus H_+$.

Now let M be a restricted H -module, i.e. such that for any $u \in M$ we have $h(n)u = 0$ for $n \gg 0$. Assume that c acts on M by the identity. Then the fields $\tilde{h} \in F\{M\}$ generate a vertex algebra $\mathfrak{V}_0 \subset F\{M\}$. It is called an *enveloping vertex algebra* of the conformal Heisenberg algebra \mathfrak{H} . The algebra \mathfrak{V}_0 is a module over the Heisenberg algebra H by $h(n)x = \tilde{h} \square x$. It is well known that \mathfrak{V}_0 is the unique enveloping vertex algebra of \mathfrak{H} such that $c = \mathbf{1}$. As a module over H , the vertex algebra \mathfrak{V}_0 is isomorphic to the so-called canonical relations representation $M(1) = U(H_-)\mathbf{1}$, which is generated by a single element $\mathbf{1}$ such that $H_+\mathbf{1} = 0$.

The vertex algebra \mathfrak{V}_0 is in fact a vertex operator algebra: it is graded so that $\deg \tilde{h} = 1$ for $h \in \mathfrak{h}$ and it contains a Virasoro element $v = \frac{1}{2} \sum_{i=1}^{\ell} \tilde{\alpha}_i \square \tilde{\beta}_i$, where $(\alpha_1, \dots, \alpha_\ell)$ and $(\beta_1, \dots, \beta_\ell)$ are dual bases of \mathfrak{h} , such that $v \square u = Du$ for all $v \in \mathfrak{V}_0$, $v \square u = (\deg u)u$ for all homogeneous $u \in \mathfrak{V}_0$, $v \square v = 0$ and $v \square v = \frac{1}{2} \mathbf{1}$.

An H -module V is called $\mathfrak{h}^{[0]}$ -diagonalizable if it can be decomposed into a direct sum of subspaces $V = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} V_\xi$, so that for $h \in \mathfrak{h}^{[0]}$ one has $h|_{V_\xi} = \xi(h)$. Recall [14] that an H -module V belongs to the category \mathcal{O} if V is $\mathfrak{h}^{[0]}$ -diagonalizable and for any $v \in V$ there is $n \in \frac{1}{p}\mathbb{Z}$ such that for any $x \in U(H)$ of $\deg x \geq n$ we have $xv = 0$. Clearly, any module from the category \mathcal{O} is restricted.

For the future reference we cite here a structural result from the representation theory of Heisenberg algebras.

Lemma 1. *Let V be a module over the Heisenberg Lie algebra H . Let $\Omega = \{v \in V \mid H_+v = 0\} \subset V$ be the vacuum subspace of V . Then the following conditions are equivalent:*

- (i) $V \cong M(1) \otimes \Omega$ and Ω is $\mathfrak{h}^{[0]}$ -diagonalizable;
- (ii) $V = U(H)\Omega$ and Ω is $\mathfrak{h}^{[0]}$ -diagonalizable;
- (iii) $V \in \mathcal{O}$ and V is completely reducible over H ;

- (iv) $V \in \mathcal{O}$ and there is a grading $V = \bigoplus_{n \in \frac{1}{p}\mathbb{Z}} V_n$ on V such that $\deg a(n) = -n$ and $v(1)|_{V_n} = n$, where

$$v(1) = \frac{1}{2} \sum_{i=1}^l \left(\sum_{s < 0} \alpha_i(s) \beta_i(-s) + \sum_{s \geq 0} \beta_i(-s) \alpha_i(s) \right) \in \overline{U(H)}$$

is the first coefficient of the Virasoro element $v \in \mathfrak{V}_0$.

Clearly, the vacuum space $\Omega \subset V$ is stable under the action of $\mathfrak{h}^{[0]}$. The condition (iv) means that V is a module over the vertex operator algebra \mathfrak{V}_0 , see §1.7. We remark that if in (iv) we assume that the grading on V is bounded from below, that is, there is $n_0 \in \frac{1}{p}\mathbb{Z}$ such that $V_n = 0$ for $n < n_0$, then the condition $V \in \mathcal{O}$ becomes obsolete.

3.2. Fock spaces. Let $\Lambda \subset \mathfrak{h}$ be a lattice in \mathfrak{h} of rank l . Assume that Λ agrees with the Γ -grading on \mathfrak{h} in the sense that $\text{rk } \Lambda^{[\lambda]} = \dim \mathfrak{h}^{[\lambda]}$ for $[\lambda] \in \Gamma$. Consider a central extension

$$1 \longrightarrow Z \longrightarrow \widehat{\Lambda} \longrightarrow \Lambda \longrightarrow 1$$

of Λ by the multiplicative group $Z \cong \mathbb{C}^\times = \mathbb{C} \setminus \{0\}$. Let $e : \Lambda \rightarrow \widehat{\Lambda}$ be a section and $\varepsilon : \Lambda \times \Lambda \rightarrow Z$ be the corresponding 2-cocycle, so that $e(\alpha)e(\beta) = \varepsilon(\alpha, \beta) e(\alpha + \beta)$ for $\alpha, \beta \in \Lambda$. Denote by R the quotient of the group algebra $\mathbb{C}[\widehat{\Lambda}]$ obtained by the identification of Z with $\mathbb{C}^\times \subset \mathbb{C}$.

The key object of the construction is a vector space V , which is a module over the associative algebra R and a restricted module over the twisted Heisenberg algebra H such that

$$[h(n), e(\alpha)] = \delta_{n,0}(h|\alpha) e(\alpha), \quad \text{for } h \in \mathfrak{h}, \quad n \in \Gamma + \mathbb{Z} \quad \text{and} \quad \mathbf{c} = \text{id}. \quad (13)$$

The formulas (13) mean that H acts on R by derivations, so we can form a skew tensor product algebra $U(H) \otimes R$ such that V is a $U(H) \otimes R$ -module.

We assume that $V = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} V_\xi$ is $\mathfrak{h}^{[0]}$ -diagonalizable. The bilinear form on Λ induces a homomorphism $\nu : \Lambda \rightarrow (\mathfrak{h}^{[0]})^*$ by $\nu(\alpha)\beta = (\alpha|\beta)$ for $\alpha \in \Lambda$, $\beta \in \Lambda^{[0]}$. The relation $[\beta(0), e(\alpha)] = (\alpha|\beta)e(\alpha)$ is equivalent to the fact that $e(\alpha)V_\xi \subseteq V_{\xi+\nu(\alpha)}$.

Such module V is sometimes referred to as a *Fock space*. In §3.4 we will make V to be a module over certain vertex algebra, under some additional assumptions.

A standard way to construct a Fock space V is as follows. Let $\Omega = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} \Omega_\xi$ be a $\mathfrak{h}^{[0]}$ -diagonalizable module over R and $\mathfrak{h}^{[0]}$ such that $[h, e(\alpha)] = (h|\alpha)e(\alpha)$ for $h \in \mathfrak{h}^{[0]}$. In other words, Ω is a module over a certain skew tensor product algebra $U(\mathfrak{h}^{[0]}) \otimes R$. As before, we have $e(\alpha)\Omega_\xi \subseteq \Omega_{\xi+\nu(\alpha)}$. Take now a restricted H -module M such that $\mathfrak{h}^{[0]}$ acts on M by 0. Let $V = M \otimes \Omega$. Then $V = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} V_\xi$, $V_\xi = M \otimes \Omega_\xi$, is also a restricted $\mathfrak{h}^{[0]}$ -diagonalizable H -module and also it is a module over R such that the relations (13) hold. The action is given by

$$a(n)(v \otimes u) = (a(n)v) \otimes u, \quad a(0)(v \otimes u) = v \otimes (a(0)u), \quad x(v \otimes u) = v \otimes x(u)$$

for $a \in \mathfrak{h}$, $0 \neq n \in \Gamma + \mathbb{Z}$, $x \in R$, $v \in M$, $u \in \Omega$. The central element \mathbf{c} acts on V by the identity.

For example, assume that V satisfies the conditions of Lemma 1 as a module over H . Then $V = M(1) \otimes \Omega$ can be obtained by the above construction for $M = M(1)$. The vacuum space $\Omega \cong \mathbf{1} \otimes \Omega \subset V$ becomes a module over $U(\mathfrak{h}^{[0]}) \otimes R$.

3.3. Twisted vertex operators. Now we are ready to define the main ingredient of this construction — the vertex operator $X_\alpha \in \text{F}\{V\}$ for $\alpha \in \Lambda$. Denote $h' = h - h^{[0]}$ for $h \in \mathfrak{h}$. We set

$$X_\alpha(z) = e(\alpha) E_-(\alpha, z) E_+(\alpha, z) z^{\alpha(0)} z^{-(\alpha'|\alpha')/2}, \quad (14)$$

where

$$E_\pm(\alpha, z) = \exp \sum_{n \in \Gamma + \mathbb{Z}, n \geq 0} -\frac{\alpha(n)}{n} z^{-n}.$$

Note that $\alpha(0) = \alpha^{[0]}$ is the projection of α onto $\mathfrak{h}^{[0]}$, so that $z^{\alpha(0)}|_{V_\xi} = z^{\xi(\alpha^{[0]})}$.

Remark. In fact we need that V is $\mathfrak{h}^{[0]}$ -diagonalizable only for the expression $z^{\alpha(0)}$ to make sense. We can instead interpret $z^{\alpha(0)} = \exp(\alpha(0) \log z)$, and then this expression is well defined under much weaker assumptions, for example it is enough to require that $\alpha(0)$ acts locally finite dimensionally.

Proposition 3. [18] *Let $h \in \mathfrak{h}$, $n \in \Gamma + \mathbb{Z}$, $\alpha, \beta \in \Lambda$. We have*

- a. $[h(n), X_\alpha(z)] = (\alpha|h) z^n X_\alpha(z);$
- b. $\tilde{h} \boxed{n} X_\alpha = 0$ if $1 \leq n \in \mathbb{Z}_+$ and $\tilde{h} \boxed{0} X_\alpha = (\alpha|h) X_\alpha;$
- c. $DX_\alpha = \tilde{\alpha} \boxed{-1} X_\alpha;$
- d. $v \boxed{0} X_\alpha = DX_\alpha, \quad v \boxed{1} X_\alpha = \frac{1}{2}(\alpha|\alpha) X_\alpha;$
- e. $X_\alpha(w)X_\beta(z) = \varepsilon(\alpha, \beta) X_{\alpha+\beta}(w, z) \iota_{w,z} \prod_{s=0}^{p-1} (w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}})^{(\sigma^{-s}\alpha|\beta)},$

where $\omega = \exp \frac{2\pi i}{p}$ is the primitive p -th root of unity and

$$X_{\alpha,\beta}(w, z) = e(\alpha + \beta) E_-(\alpha, w) E_-(\beta, z) E_+(\alpha, w) E_+(\beta, z) w^{\alpha(0)} z^{\beta(0)} w^{-(\alpha'|\alpha')/2} z^{-(\beta'|\beta')/2}.$$

We have $X_{\alpha,\beta}(w, z) = X_{\beta,\alpha}(z, w)$ and $X_{\alpha,\beta}(z, z) = X_{\alpha+\beta}(z) z^{(\alpha'|\beta')}$. The notation $\iota_{w,z}$ in (e) is a short for $\iota_{w^{\frac{1}{p}}, z^{\frac{1}{p}}}$, see (2).

Proof. (a). Let $0 \neq n \in \Gamma + \mathbb{Z}$. Then

$$\left[h(n), \exp \left(-\frac{\alpha(m)}{m} z^{-m} \right) \right] = \delta_{n,-m} \exp \left(-\frac{\alpha(m)}{m} z^{-m} \right) (\alpha|h) z^n,$$

and $h(n)$ commutes with all the rest of the factors in (14). Also $h(0)$ commutes with all the factors in (14) except $e(\alpha)$, whose commutators are given by (13), so (a) follows.

(b). It follows that $[h(n), X_\alpha(m)] = X_\alpha(m+n)$ for every $m, n \in \Gamma + \mathbb{Z}$. Hence we have for $n \in \mathbb{Z}_+$

$$\begin{aligned} (\tilde{h} \boxed{n} X_\alpha)(m) &= \sum_s (-1)^s \binom{n}{s} [h(n-s), X_\alpha(m+s)] \\ &= (\alpha|h) \sum_s (-1)^s \binom{n}{s} X_\alpha(m+n) = \begin{cases} 0 & \text{if } n > 0, \\ (\alpha|h) X_\alpha(m) & \text{if } n = 0. \end{cases} \end{aligned}$$

(c). Using that $(\alpha'|\alpha')/2 = \sum_{\lambda \in \Gamma} \lambda (\alpha^{[\lambda]}|\alpha)$, we get

$$\begin{aligned} DX_\alpha(z) &= \sum_{n < 0} \alpha(n) z^{-n-1} X_\alpha(z) + X_\alpha(z) \sum_{n > 0} \alpha(n) z^{-n-1} + X_\alpha(z) \alpha(0) z^{-1} \\ &\quad - \sum_{\lambda \in \Gamma} \lambda (\alpha^{[\lambda]}|\alpha) z^{-1} X_\alpha(z) \\ &= : \tilde{\alpha} X_\alpha : - \sum_{\lambda \in \Gamma} \lambda (\alpha^{[\lambda]}|\alpha) z^{-1} X_\alpha(z). \end{aligned}$$

On the other hand, set $\tilde{\alpha}(z) = \sum_\lambda z^{-\lambda} \tilde{\alpha}^{[\lambda]}(z)$. The locality of $\tilde{\alpha}$ and X_α is 1, hence, using (6),

$$\begin{aligned} \tilde{\alpha} \boxed{-1} X_\alpha &= \sum_\lambda (z^{-\lambda} \tilde{\alpha}^{[\lambda]}) \boxed{-1} X_\alpha \\ &= \sum_\lambda \left(z^{-\lambda} (\tilde{\alpha}^{[\lambda]} \boxed{-1} X_\alpha) - \lambda z^{-\lambda-1} (\tilde{\alpha}^{[\lambda]} \boxed{0} X_\alpha) \right) \\ &= : \tilde{\alpha} X_\alpha : - \sum_{\lambda \in \Gamma} \lambda (\alpha^{[\lambda]}|\alpha) z^{-1} X_\alpha(z). \end{aligned}$$

(d). Let us calculate $v \square X_\alpha$ by the associativity formula V3. We have, using (b) and (c):

$$\begin{aligned} v \square X_\alpha &= \frac{1}{2} \sum_i \left(\sum_{s < 0} \alpha_i(s) \beta_i(-s-1) + \sum_{s \geq 0} \beta_i(-s-1) \alpha_i(s) \right) X_\alpha \\ &= \frac{1}{2} \sum_i \left(\alpha_i(-1) (\beta_i | \alpha) + \beta_i(-1) (\alpha_i | \alpha) \right) X_\alpha \\ &= \alpha(-1) X_\alpha = D X_\alpha. \end{aligned}$$

The other relation is proved in the same way.

(e). Let us first calculate $S = E_+(\alpha, w) E_-(\beta, z) E_+^{-1}(\alpha, w)$. Since

$$\begin{aligned} \exp \left(-\frac{\alpha(n)}{n} w^{-n} \right) E_-(\beta, z) \exp \left(\frac{\alpha(n)}{n} w^{-n} \right) &= \exp \left(\frac{w^{-n}}{n} \text{ad } \alpha(n) \right) E_-(\beta, z) \\ &= (\alpha^{[n]} | \beta) \frac{z^n w^{-n}}{n} E_-(\beta, z), \end{aligned}$$

we have

$$S = \exp \sum_{0 < n \in \Gamma + \mathbb{Z}} -(\alpha^{[n]} | \beta) \frac{z^n w^{-n}}{n} = \exp \sum_{\lambda \in \{\frac{1}{p}, \dots, \frac{p-1}{p}, 1\}} -(\alpha^{[\lambda]} | \beta) \sum_{n \in \lambda + \mathbb{Z}_+} \frac{z^n w^{-n}}{n}.$$

Denote $y = (z/w)^{\frac{1}{p}}$ and let $\lambda = \frac{q}{p}$, $1 \leq q \leq p$. Then

$$\sum_{n \in \lambda + \mathbb{Z}_+} \frac{z^n w^{-n}}{n} = \int \frac{(z/w)^{\lambda-1}}{1 - (z/w)} d(z/w) = p \int \frac{y^{q-1}}{1 - y^p} dy.$$

Using the elementary fraction decomposition

$$\frac{y^r}{1 - y^p} = \frac{1}{p} \sum_{s=0}^{p-1} \frac{\omega^{-sr}}{1 - \omega^s y},$$

we get that

$$\sum_{n \in \lambda + \mathbb{Z}_+} \frac{z^n w^{-n}}{n} = - \sum_{s=0}^{p-1} \ln(1 - \omega^s y)^{\omega^{-sq}},$$

so that

$$S = \prod_{s,q=0}^{p-1} (1 - \omega^s y)^{\omega^{-sq} (\alpha^{(q/p)} | \beta)} = \prod_{s=0}^{p-1} (1 - \omega^s y)^{\sum_q \omega^{-sq} (\alpha^{(q/p)} | \beta)} = \prod_{s=0}^{p-1} (1 - \omega^s y)^{(\sigma^{-s} \alpha | \beta)}.$$

The rest of non-trivial commutation relations between the factors in (14) are

$$e(\alpha) \varepsilon(\beta) = \varepsilon(\alpha, \beta) e(\alpha + \beta), \quad w^{\alpha(0)} e(\beta) = e(\beta) w^{\alpha(0)} w^{(\alpha^{[0]} | \beta)},$$

so we finally get, using that $\alpha^{[0]} = \frac{1}{p} \sum_{s=0}^{p-1} \sigma^s \alpha$,

$$\begin{aligned} X_\alpha(w) X_\beta(z) &= \varepsilon(\alpha, \beta) X_{\alpha, \beta}(w, z) w^{(\alpha^{[0]} | \beta)} S \\ &= \varepsilon(\alpha, \beta) X_{\alpha, \beta}(w, z) \iota_{w, z} \prod_{s=0}^{p-1} (w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}})^{(\sigma^{-s} \alpha | \beta)}. \end{aligned}$$

□

The statement (b) implies that X_α generates a highest weight module over the Heisenberg Lie algebra H , where the action is given as usual by $h(n)u = \tilde{h} \square u$. This module is irreducible and is in fact a module over the Heisenberg vertex algebra $\mathfrak{V}_0 = U(H)\mathbb{1}$.

The Virasoro element $v \in \mathfrak{V}_0$ gives the operator

$$D = v(0) = \sum_i \left(\sum_{s < 0} \alpha_i(s) \beta_i(-s-1) + \sum_{s \geq 0} \beta_i(-s-1) \alpha_i(s) \right) \in \mathfrak{gl} V. \quad (15)$$

By (d) the vertex operators X_α are of weight 0 with respect to this D , see §1.6.

3.4. Lattice vertex algebras and their generalized representations. In this section we construct the vertex superalgebra \mathfrak{V}_Λ , known as the lattice vertex superalgebra. See the introduction for the discussion of its importance and references.

We start with the remark that only minimal modification is needed to transfer all the definitions and results of §1 and §2 to the realm of superalgebras. All commutators must be interpreted as supercommutators, the formula (3) must change to

$$(\alpha \square \beta)(z) = \text{Res}_w \left(\alpha(w) \beta(z) \iota_{w,z}(w-z)^n - (-1)^{p(\alpha)p(\beta)} \beta(z) \alpha(w) \iota_{z,w}(w-z)^n \right),$$

etc. Notably, in the definition of vertex operator superalgebra the eigenvalues of the grading derivation $v(1)$ are allowed to be half-integer, and the vertex algebra $\mathfrak{A} = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}} \mathfrak{A}_n$ is graded by $\frac{1}{2}\mathbb{Z}$ so that the even and odd parts of \mathfrak{A} are respectively $\mathfrak{A}^{\bar{0}} = \bigoplus_{n \in \mathbb{Z}} \mathfrak{A}_n$, $\mathfrak{A}^{\bar{1}} = \bigoplus_{n \in \mathbb{Z} + \frac{1}{2}} \mathfrak{A}_n$. A reader who is averse to supermathematics can assume that the lattice Λ is even, i.e. $(\alpha|\alpha) \in 2\mathbb{Z}$ for all $\alpha \in \Lambda$, see (i) below.

Here we make the following assumptions on the data introduced in §3.3.

- (i) For any $\alpha, \beta \in \Lambda$ the numbers $m_s = (\sigma^{-s}\alpha|\beta)$, $0 \leq s \leq p-1$, are integers, in other words $\Lambda \subset \mathfrak{h}$ is an integer lattice and the automorphism $\sigma : \mathfrak{h} \rightarrow \mathfrak{h}$ preserves the dual lattice $\Lambda' = \{\alpha \in \mathfrak{h} \mid (\alpha|\Lambda) \subset \mathbb{Z}\} \supseteq \Lambda$.
- (ii) The cocycle $\varepsilon : \Lambda \times \Lambda \rightarrow \mathbb{C}^\times$ is such that the corresponding commutator map is

$$C(\alpha, \beta) = \varepsilon(\alpha, \beta) \varepsilon(\beta, \alpha)^{-1} = (-1)^{(\alpha|\alpha)(\beta|\beta)} (-1)^{p(\alpha^{[0]}|\beta^{[0]})} \omega^{-\sum_{s=1}^{p-1} s m_s}. \quad (16)$$

Note that $p(\alpha^{[0]}|\beta^{[0]}) = \sum_{s=0}^{p-1} m_s \in \mathbb{Z}$.

The cocycle ε satisfying (16) can be easily constructed in the following way. Let $\alpha_1, \dots, \alpha_l$ be a \mathbb{Z} -basis of Λ . Define ε first on the elements of this basis such that (16) holds for any $\alpha, \beta \in \{\alpha_1, \dots, \alpha_l\}$. This is possible since $C(\alpha, \beta) = C(\beta, \alpha)^{-1}$ and $C(\alpha, \alpha) = 1$. Then, since $C : \Lambda \times \Lambda \rightarrow \mathbb{C}^\times$ is bimultiplicative, the identity (16) will continue to hold for the bimultiplicative extension of ε to the whole Λ . Note that for $p = 1$ or 2 we have $C(\alpha, \beta) = (-1)^{(\alpha|\alpha)(\beta|\beta)} (-1)^{(\alpha|\beta)}$.

Denote by $\mathcal{P}(m) = \{ \mathbf{r} = (r_1, r_2, \dots) \mid r_i \geq 0, \sum_{i \geq 1} i r_i = m \}$ the set of partitions of $m \in \mathbb{Z}$.

Theorem 2. *Under the assumptions (i) and (ii) above the vertex operators $X_\alpha, X_\beta \in \mathbf{F}\{V\}$ are local of order $N(\alpha, \beta) = \max \{-m_s \mid m_s < 0, 0 \leq s \leq p-1\} \cup \{0\}$. They generate the lattice vertex superalgebra $\mathfrak{V} = \mathfrak{V}_\Lambda \subset \mathbf{F}\{V\}$, which does not depend on the Γ -grading of \mathfrak{h} . The products of the generators are given by*

$$X_\alpha \square X_\beta = \varkappa(\alpha, \beta) \sum_{\mathbf{r} \in \mathcal{P}(-(\alpha|\beta)-n-1)} \prod_{j \geq 1} \left(\frac{\alpha(-j)}{j!} \right)^{r_j} X_{\alpha+\beta}, \quad (17)$$

for $n < -(\alpha|\beta)$, and $X_\alpha \square X_\beta = 0$ if $n \geq -(\alpha|\beta)$. Here

$$\varkappa(\alpha, \beta) = \varepsilon(\alpha, \beta) p^{-(\alpha|\beta)} \prod_{s=1}^{p-1} (1 - \omega^s)^{m_s}. \quad (18)$$

In particular,

$$X_\alpha \square_{-(\alpha|\beta)-1} X_\beta = \varkappa(\alpha, \beta) X_{\alpha+\beta}, \quad X_\alpha \square_{-(\alpha|\alpha)-2} X_{-\alpha} = \varkappa(\alpha, \alpha)^{-1} \tilde{\alpha}.$$

In the case when $p = 1$ this is just the usual construction of lattice vertex algebras, see e.g. [12].

The lattice vertex algebra $\mathfrak{V} = \bigoplus_{\alpha \in \Lambda} \mathfrak{V}_\alpha$ is graded by the lattice Λ . The subspace \mathfrak{V}_0 is the Heisenberg vertex subalgebra of \mathfrak{V} , and the rest of \mathfrak{V}_α 's are irreducible modules over \mathfrak{V}_0 .

The even and odd parts of \mathfrak{V} are

$$\mathfrak{V}^{\bar{0}} = \bigoplus_{\substack{\alpha \in \Lambda: \\ (\alpha|\alpha) \in 2\mathbb{Z}}} \mathfrak{V}_{\alpha}, \quad \mathfrak{V}^{\bar{1}} = \bigoplus_{\substack{\alpha \in \Lambda: \\ (\alpha|\alpha) \in 2\mathbb{Z}+1}} \mathfrak{V}_{\alpha}.$$

We also note that \mathfrak{V} is a simple vertex algebra.

Let us calculate the commutator map of the cocycle $\varkappa : \Lambda \times \Lambda \rightarrow \mathbb{C}^{\times}$ given by (18):

$$\begin{aligned} \varkappa(\alpha, \beta) \varkappa(\beta, \alpha)^{-1} &= C(\alpha, \beta) \prod_{s=1}^{p-1} (1 - \omega^s)^{m_s - m_{p-s}} \\ &= (-1)^{(\alpha|\alpha)(\beta|\beta)} (-1)^{(\alpha|\beta)} \prod_{s=1}^{p-1} (-\omega^{-s})^{m_s} \prod_{s=1}^{p-1} \left(\frac{1 - \omega^s}{1 - \omega^{-s}} \right)^{m_s} \\ &= (-1)^{(\alpha|\alpha)(\beta|\beta)} (-1)^{(\alpha|\beta)}. \end{aligned}$$

It follows that for different automorphisms σ all resulting cocycles \varkappa are cohomological and define certain class $\varkappa \in H^2(\Lambda, \mathbb{C}^{\times})$. Therefore, the vertex algebra $\mathfrak{V}_{\Lambda} \subset F\{V\}$ generated by X_{α} 's is indeed independent on σ .

Remark. If $(\alpha|\beta)$ is not an integer, then the vertex operators X_{α} and X_{β} are not local. However, they are local in a generalized sense, and they generate a generalized vertex algebra [7], mentioned at the end of §1.7. One can define products $X_{\alpha} \boxed{n} X_{\beta}$ for $n \equiv -(\alpha|\beta) \pmod{\mathbb{Z}}$.

3.5. Proof of Theorem 2. The assertion about locality follows from Proposition 3(e) and the formula

$$(w - z) = \prod_{s=0}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right).$$

Consider the operator $\delta = \iota_{w,z} - \iota_{z,w}$ equal to the difference between the two expansions of a rational function in w and z into a power series given by (2). It is easy to see that $\delta(g) = 0$ if and only if $g \in \mathbb{C}[w^{\pm 1}, z^{\pm 1}]$. In this case g commutes with δ : $\delta(gf) = g\delta(f)$. We will also make use of the following formula. For any formal power series $f(w, z)$ in the variables w, z one has

$$\text{Res}_w \left(f(w, z) \delta(w - z)^{-k-1} \right) = \frac{1}{k!} \frac{\partial^k}{\partial w^k} \Big|_{w=z} f(w, z)$$

whenever both sides make sense.

Let $n = -(\alpha|\beta) - k - 1$. We calculate the product $X_{\alpha} \boxed{n} X_{\beta}$ using (8) and Proposition 3(e):

$$\begin{aligned} (X_{\alpha} \boxed{n} X_{\beta})(z) &= \text{Res}_w \left(\left(X_{\alpha}(w) X_{\beta}(z) \iota_{w,z}(w - z)^n - (-1)^{(\alpha|\alpha)(\beta|\beta)} X_{\beta}(z) X_{\alpha}(w) \iota_{z,w}(w - z)^n \right) \times \right. \\ &\quad \left. \times F(N + m + k) \prod_{s=1}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{N-n} \right) \\ &= \varepsilon(\alpha, \beta) \text{Res}_w \left(X_{\alpha, \beta}(w, z) F(N + m + k) \prod_{s=1}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{N-n} \delta \prod_{s=0}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{m_s + n} \right) \\ &= \varepsilon(\alpha, \beta) \text{Res}_{w^{\frac{1}{p}}} \left(X_{\alpha, \beta}(w, z) w^{\frac{p-1}{p}} F(N + m + k) \prod_{s=1}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{m_s + N} \delta \left(w^{\frac{1}{p}} - z^{\frac{1}{p}} \right)^{-k-1} \right) \\ &= \varepsilon(\alpha, \beta) \frac{1}{k!} \frac{\partial^k}{\left(\partial w^{\frac{1}{p}} \right)^k} \Big|_{w^{\frac{1}{p}} = z^{\frac{1}{p}}} \left(X_{\alpha, \beta}(w, z) w^{\frac{p-1}{p}} F(N + m + k) \prod_{s=1}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{m_s + N} \right). \end{aligned} \tag{19}$$

We use here that $\text{Res}_w = \text{Res}_{w^{\frac{1}{p}}} w^{\frac{p-1}{p}}$. Note that $N + m_s \geq 0$ for all $0 \leq s \leq p-1$.

Recall that there is an operator $D : V \rightarrow V$ given by (15) such that the weights of all vertex operators X_α are 0, see §1.6. Denote

$$B = w^{\frac{p-1}{p}} F(N+m+k) \prod_{s=1}^{p-1} (w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}})^{m_s+N}.$$

Using (9) and the formula $\prod_{s=1}^{p-1} (1 - \omega^s) = p$, we calculate

$$B \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} = p^{-m-k} \prod_{s=1}^{p-1} (1 - \omega^s)^{m_s} z^{-(\alpha'|\beta')} z^{k\frac{1-p}{p}},$$

hence $\text{wt } B \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} = -(\alpha'|\beta') - k\frac{p-1}{p}$. Since B is a Laurent polynomial in $w^{\frac{1}{p}}$ and $z^{\frac{1}{p}}$, we get that

$$\text{wt } \frac{\partial^i}{(\partial w^{\frac{1}{p}})^i} \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} B = -(\alpha'|\beta') - k\frac{p-1}{p} - \frac{i}{p}.$$

Let us now calculate the derivative of $X_{\alpha,\beta}$, as in the proof of Proposition 3(c):

$$\begin{aligned} \frac{\partial}{\partial w^{\frac{1}{p}}} X_{\alpha,\beta}(w, z) &= p w^{\frac{p-1}{p}} \frac{\partial}{\partial w} F_{\alpha,\beta}(w, z) \\ &= p w^{\frac{p-1}{p}} \left(: \tilde{\alpha} X_{\alpha,\beta} : - \frac{1}{2} (\alpha'|\alpha') w^{-1} X_{\alpha,\beta} \right) \\ &= p w^{\frac{p-1}{p}} \left(\tilde{\alpha} \square X_{\alpha,\beta} + \frac{1}{2} (\alpha'|\beta') w^{-1} X_{\alpha,\beta} \right). \end{aligned}$$

Iterating this formula and using the fact that $X_{\alpha,\beta}(z, z) = X_{\alpha+\beta}(z) z^{(\alpha'|\beta')}$, we get

$$\begin{aligned} \frac{1}{i!} \frac{\partial^i}{(\partial w^{\frac{1}{p}})^i} \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} X_{\alpha,\beta} &= p^i z^{(\alpha'|\beta')} z^{i\frac{p-1}{p}} \sum_{(r_1, r_2, \dots) \in \mathcal{P}(i)} \prod_{j \geq 1} \left(\frac{\alpha(-j)}{j!} \right)^{r_j} X_{\alpha+\beta} \\ &+ \text{a sum of terms of the weight less than } i\frac{p-1}{p} + (\alpha'|\beta'). \end{aligned} \quad (20)$$

Now expand (19) using the Leibniz rule and (20), and note that the only term of weight 0 in this expansion is the biggest weight term in $\varepsilon(\alpha, \beta) \left(\frac{1}{i!} \partial^i / (\partial w^{\frac{1}{p}})^i \Big|_{w^{\frac{1}{p}}=z^{\frac{1}{p}}} X_{\alpha,\beta} \right) B$, which is precisely the right-hand side of the formula (17). This finishes the proof of Theorem 2. \square

Remark. One could have used the theorem of Li and Xu [23] on characterization of lattice vertex algebras to show that the vertex algebra generated by the vertex operators $X_\alpha \in \mathcal{F}\{V\}$ is isomorphic to the lattice vertex algebra. They consider only the case of an even lattice, but one can easily generalize their result to the case of a non-trivial odd part. It is easy then to deduce the product formula (17) but this proof would not give the explicit formula (18) for the cocycle \varkappa .

Remark. Dong and Lepowsky [8] have constructed all twisted modules of \mathfrak{V}_Λ for the case when Λ is even, see §3.7. They used, however, completely different techniques, in particular, they didn't get the cocycle \varkappa explicitly. Some special cases were also considered in [6, 10].

Corollary 1. *Let $U \subset V$ be a $U(H) \otimes R$ -submodule of V . Then U is a submodule over the vertex algebra \mathfrak{V}_Λ .*

3.6. Do we get all generalized modules of \mathfrak{V}_Λ ? In this section we show that in fact the construction of Theorem 2 exhausts all reasonable modules of \mathfrak{V}_Λ . A similar argument was also used in [5, 6, 20, 23].

Let V be a generalized module over the lattice vertex algebra \mathfrak{V}_Λ such that for any homogeneous $h \in \mathfrak{h}^{[\lambda]}$ the corresponding field $\tilde{h} \in \mathcal{F}\{V\}^{[\lambda]}$ is also homogeneous, in other words, V is a twisted module over \mathfrak{V}_0 . Then the space V becomes a module over the twisted Heisenberg algebra H , such that c acts as the identity.

We assume that $V = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} V_\xi$ is $\mathfrak{h}^{[0]}$ -diagonalizable, though as in the remark, preceding Proposition 3, we note that this assumption can be relaxed. We will show that the \mathfrak{V}_Λ -module V can be obtained by the construction of §3.4.

For $\alpha \in \Lambda$ define

$$e(\alpha) = E_-(\alpha, z)^{-1} X_\alpha(z) E_+(\alpha, z)^{-1} z^{-\alpha(0)} z^{(\alpha'|\alpha')/2}.$$

Using the same calculations as in the proof of (a) and (c) of Proposition 3 we see that $\frac{d}{dz}e(\alpha) = 0$ and $[h(n), e(\alpha)] = \delta_{n,0} (h|\alpha) e(\alpha)$, so that the relations (13) hold. Now we want to show that $e(\alpha) : V \rightarrow V$ generate an action of the central extension $\widehat{\Lambda}$ of the lattice Λ corresponding to a cocycle $\varepsilon : \Lambda \times \Lambda \rightarrow \mathbb{C}^\times$ satisfying (16). It will follow that V is a module over the skew tensor product $U(H) \otimes R$ introduced in §3.2.

Modifying slightly the proof of Proposition 3(e) we get that

$$X_\alpha(w) X_\beta(z) = e(\alpha) e(\beta) X'_{\alpha,\beta}(w, z) \prod_{s=0}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{(\sigma^{-s}\alpha|\beta)},$$

where

$$X'_{\alpha,\beta}(w, z) = E_-(\alpha, w) E_-(\beta, z) E_+(\alpha, w) E_+(\beta, z) w^{\alpha(0)} z^{\beta(0)} w^{-(\alpha'|\alpha')/2} z^{-(\beta'|\beta')/2}$$

is symmetric, $X'_{\alpha,\beta}(w, z) = X'_{\beta,\alpha}(z, w)$ and also $e(\alpha + \beta) X'_{\alpha,\beta}(z, z) = X_{\alpha+\beta}(z) z^{(\alpha'|\beta')}$.

Take N sufficiently large. Then, since X_α and X_β must be local, we get

$$\begin{aligned} 0 &= \left(X_\alpha(w) X_\beta(z) - (-1)^{(\alpha|\alpha)(\beta|\beta)} X_\beta(z) X_\alpha(w) \right) (w - z)^N \\ &= \left(e(\alpha) e(\beta) - C(\alpha, \beta) e(\beta) e(\alpha) \right) X'_{\alpha,\beta}(w, z) \prod_{s=0}^{p-1} \left(w^{\frac{1}{p}} - \omega^s z^{\frac{1}{p}} \right)^{n_s} \end{aligned}$$

for some $n_s \in \mathbb{Z}_+$ and $C(\alpha, \beta)$ as in (16). But this can only happen if

$$e(\alpha) e(\beta) = C(\alpha, \beta) e(\beta) e(\alpha),$$

because otherwise we would have $X'_{\alpha,\beta}(w, z) (w - z)^n = 0$ for $n = \max_s n_s$ and this is impossible since $X'_{\alpha,\beta}$ is regular at $w = z$.

Now we can apply the calculations (19) to the product $X_\alpha \square X_\beta$ for $n = -(\alpha|\beta) - 1$. Similar to (19) we get that

$$e(\alpha + \beta) X_\alpha \square X_\beta = e(\alpha) e(\beta) p^{-(\alpha|\beta)} \prod_{s=1}^{p-1} (1 - \omega^s)^{(\sigma^{-s}\alpha|\beta)} X_{\alpha+\beta}.$$

But we know that $X_\alpha \square X_\beta = \varkappa(\alpha, \beta) X_{\alpha+\beta}$ in \mathfrak{V}_Λ , therefore $e(\alpha) e(\beta) X_{\alpha+\beta} = \varepsilon(\alpha, \beta) e(\alpha + \beta) X_{\alpha+\beta}$, hence $e(\alpha) e(\beta) = \varepsilon(\alpha, \beta) e(\alpha + \beta)$.

So we have proved the following

Theorem 3. *Let V be a generalized module over the lattice vertex superalgebra \mathfrak{V}_Λ such that $\tilde{h} \in F\{V\}^{[\Lambda]}$ for $h \in \mathfrak{h}^{[\Lambda]}$. Assume that V is $\mathfrak{h}^{[0]}$ -diagonalizable. Then there is a unique action of R on V such that the highest weights vectors $X_\alpha \in \mathfrak{V}_\alpha$ for $\alpha \in \Lambda$ act on V by the vertex operators (14).*

It follows from Theorem 3 that a \mathfrak{V}_Λ -submodule $U \subset V$ is a submodule over $U(H) \otimes R$. Combining this with Corollary 1, we get the following statement.

Corollary 2. *Assume V is an irreducible module over $U(H) \otimes R$, satisfying the assumptions of Theorem 2. Then V is an irreducible module over the vertex algebra \mathfrak{V}_Λ . Conversely, let V be a module over \mathfrak{V}_Λ , satisfying the assumptions of Theorem 3. Then V is an irreducible $U(H) \otimes R$ -module.*

3.7. Twisted modules over lattice vertex superalgebras. In this section we study twisted modules over the lattice vertex algebra $\mathfrak{V} = \mathfrak{V}_\lambda$, see §1.7 for the definition. As in §3.3, we are fixing a p -th primitive root of unity $\omega = \exp\left(\frac{2\pi i}{p}\right)$.

Assume that the automorphism $\sigma : \mathfrak{h} \rightarrow \mathfrak{h}$ preserves the lattice Λ . Let $\hat{\sigma} : \mathfrak{V} \rightarrow \mathfrak{V}$ be an extension of the automorphism $\sigma : \mathfrak{V}_0 \rightarrow \mathfrak{V}_0$ to the whole vertex algebra \mathfrak{V} . It is easy to see that $\hat{\sigma}X_\alpha = \varphi(\alpha)X_{\sigma\alpha}$ for some 1-cocycle $\varphi : \Lambda \rightarrow \mathbb{C}^\times$ such that

$$d\varphi(\alpha, \beta) = \frac{\varphi(\alpha + \beta)}{\varphi(\alpha)\varphi(\beta)} = \frac{\varkappa(\sigma\alpha, \sigma\beta)}{\varkappa(\alpha, \beta)}.$$

Since σ preserves the norm $(\cdot | \cdot)$ on \mathfrak{h} and $\varkappa(\alpha, \beta)\varkappa(\beta, \alpha)^{-1} = (-1)^{(\alpha|\alpha)(\beta|\beta)}(-1)^{(\alpha|\beta)}$, we see that the cocycle $\frac{\varkappa(\sigma\alpha, \sigma\beta)}{\varkappa(\alpha, \beta)}$ is indeed symmetric and therefore equal to $d\varphi_0$ for

$$\varphi_0(\alpha) = \left(\frac{\varkappa(\sigma\alpha, \sigma\alpha)}{\varkappa(\alpha, \alpha)} \right)^{\frac{1}{2}}.$$

The formula (18) implies that

$$\frac{\varkappa(\sigma\alpha, \sigma\beta)}{\varkappa(\alpha, \beta)} = \frac{\varepsilon(\sigma\alpha, \sigma\beta)}{\varepsilon(\alpha, \beta)},$$

therefore the map $e(\alpha) \mapsto \varphi(\alpha)e(\sigma\alpha)$ defines an automorphism of the algebra R . And vice versa, any such automorphism of R defines an automorphism of \mathfrak{V} .

Remark. In general, the order of $\hat{\sigma} : \mathfrak{V} \rightarrow \mathfrak{V}$ may not be equal to p , in fact it may not be finite at all. However, the extension corresponding to φ_0 is of order p . Indeed,

$$\begin{aligned} \hat{\sigma}^p X_\alpha &= \varphi_0(\alpha)\varphi_0(\sigma\alpha) \cdots \varphi_0(\sigma^{p-1}\alpha)X_\alpha \\ &= \left(\frac{\varkappa(\sigma\alpha, \sigma\alpha)\varkappa(\sigma^2\alpha, \sigma^2\alpha) \cdots \varkappa(\alpha, \alpha)}{\varkappa(\alpha, \alpha)\varkappa(\sigma\alpha, \sigma\alpha) \cdots \varkappa(\sigma^{p-1}\alpha, \sigma^{p-1}\alpha)} \right)^{\frac{1}{2}} X_\alpha = X_\alpha. \end{aligned}$$

Now we show how the vertex algebra \mathfrak{V} is decomposed into a sum of the root spaces of $\hat{\sigma}$. Consider the action of $\hat{\sigma}$ on a linear span $E = \text{Span}\{X_\alpha, X_{\sigma\alpha}, \dots, X_{\sigma^{p-1}\alpha}\}$ of a $\hat{\sigma}$ -orbit of X_α . It is easy to see that the eigenvalues of $\hat{\sigma}$ on E are all the different roots $\mu = \sqrt[p]{\varphi(\alpha)\varphi(\sigma\alpha) \cdots \varphi(\sigma^{p-1}\alpha)}$. Fix such a root μ . The eigenvector corresponding to the eigenvalue $\mu\omega^j$, $0 \leq j \leq p-1$, is

$$Y_j = \sum_{s=0}^{p-1} \omega^{-js} k_s X_{\sigma^s \alpha}, \quad k_s = \mu^{-s} \varphi(\alpha)\varphi(\sigma\alpha) \cdots \varphi(\sigma^{s-1}\alpha) \text{ for } s > 0, \quad k_0 = 1. \quad (21)$$

So we deduce that $\mathfrak{V} = \bigoplus_{\mu \in \mathbb{C}^\times} \mathfrak{V}_\mu$ is decomposed into a direct sum of the root spaces of $\hat{\sigma}$ such that $\hat{\sigma}|_{\mathfrak{V}_\mu} = \mu$.

Let $V = \bigoplus_{\xi \in (\mathfrak{h}^{[0]})^*} V_\xi$ be a generalized $\mathfrak{h}^{[0]}$ -diagonalizable \mathfrak{V} -module. Denote by $\Xi = \{\xi \in (\mathfrak{h}^{[0]})^* \mid V_\xi \neq 0\}$ the set of weights of V . By Theorem 3 there are operators $e(\alpha) : V \rightarrow V$, $\alpha \in \Lambda$, that define an action of R on V , satisfying the condition (13).

Theorem 4. *The module V is twisted with respect to the grading induced by $\hat{\sigma}$ if and only if for every $\alpha \in \Lambda$ there is a root $\mu_\alpha = \sqrt[p]{\varphi(\alpha)\varphi(\sigma\alpha) \cdots \varphi(\sigma^{p-1}\alpha)}$ such that*

- (i) $e(\sigma^s \alpha) = k_s^{-1} e(\alpha)$ for every $0 \leq s \leq p-1$ and
- (ii) $\xi(\alpha(0)) \equiv \frac{(\alpha'|\alpha')}{2} - \lambda \pmod{\mathbb{Z}}$ for every $\xi \in \Xi$,

where p is the length of σ -orbit of α , k_s are given by (21) and $\lambda = \frac{1}{2\pi i} \ln \mu_\alpha$.

The condition (i) means that $e(\sigma\alpha) = \mu_\alpha \varphi(\alpha)^{-1} e(\alpha)$ for every $\alpha \in \Lambda$, where the root μ_α is the same for every vector in the σ -orbit of α . We also note that it is enough to impose (i) and (ii) only for α running over an integer basis of Λ .

Proof. Fix a root μ as above and let $\lambda = \frac{1}{2\pi i} \ln \mu$. Let $\pi : \mathfrak{V} \rightarrow F\{V\}$ be the representation map. Then V is twisted with respect to the above grading of \mathfrak{V} if $\pi(\mathfrak{V}_\mu) \subset F\{V\}^{[\lambda]}$. It is enough to require that $Y_j \in F\{V\}^{[j/p + \lambda]}$ where Y_j is (the image of) the eigenvector of $\hat{\sigma}$ given by (21).

Moreover, it is enough to require this only for a finite set of generators of \mathfrak{V} , for example for all the Y_j 's corresponding to the σ -orbits of an integer basis of Λ .

Denote $\mathring{X}_\alpha(z) = E_-(\alpha, z)E_+(\alpha, z) \in F\{V\}$ so that $X_\alpha = e(\alpha)\mathring{X}_\alpha z^{\alpha(0)} z^{-\frac{(\alpha'|\alpha')}{2}}$, see §3.3. Denote also $\mathring{Y}_j = \sum_{s=0}^{p-1} \omega^{-js} k_s e(\sigma^s \alpha) X_{\sigma^s(\alpha)}$ and then $Y_j = \mathring{Y}_j z^{\alpha(0)} z^{-\frac{(\alpha'|\alpha')}{2}}$. The field $Y_j \in F\{V\}$ is homogeneous if and only if \mathring{Y}_j is homogeneous and the values $\xi(\alpha(0))$ are the same modulo \mathbb{Z} for all $\xi \in \Xi$.

Assume that \mathring{Y}_j is homogeneous. Since $\mathring{X}_\alpha(z) \in \bigoplus_{q=0}^{p-1} F\{V\}^{[q/p]}$, we have $\mathring{Y}_j \in F\{V\}^{[(q+j)/p]}$ for some $0 \leq q \leq p-1$. Let τ be the automorphism of $\bigoplus_{q=0}^{p-1} F\{V\}^{[q/p]}$ such that $\tau|_{F\{V\}^{[q/p]}} = \omega^q$. It is easy to see that $\mathring{X}_{\sigma\alpha} = \tau \mathring{X}_\alpha$. Let $m \equiv -\frac{r}{p} \pmod{\mathbb{Z}}$. Denote by x_s the coefficient of z^m in $\mathring{X}_{\sigma^s \alpha}$. It follows that $x_s = \omega^{rs} x_0$. If $r \not\equiv q+j \pmod{p}$, then the coefficient of z^m in \mathring{Y}_j is equal to 0. This gives us the following system of linear equations:

$$\sum_{s=0}^{p-1} \omega^{(r-j)s} k_s e(\sigma^s \alpha) = 0 \quad \text{for } 0 \leq r \leq p-1, \quad r \not\equiv q+j \pmod{p}.$$

Since $k_0=1$, the solution of this system is $e(\sigma^s \alpha) = \omega^{qs} k_s^{-1} e(\alpha)$. Taking $\mu \omega^q$ instead of μ we get exactly (i).

The condition (ii) follows from the fact that in order to have $\deg Y_j = \frac{j}{p} + \lambda$ we must have $\xi(\alpha(0)) \equiv \frac{(\alpha'|\alpha')}{2} - \lambda - \frac{q}{p} \equiv \frac{(\alpha'|\alpha')}{2} - \frac{1}{2\pi i} \ln(\mu \omega^q) \pmod{\mathbb{Z}}$ for all $\xi \in \Xi$. \square

Corollary 3. *If for some $\alpha \in \Lambda$ and $0 \leq j \leq p-1$ we have $C(\alpha, \sigma^j \alpha) \neq 1$, then there are no non-trivial twisted modules of \mathfrak{V}_Λ corresponding to either of the continuations $\hat{\sigma} : \mathfrak{V}_\Lambda \rightarrow \mathfrak{V}_\Lambda$ of σ .*

Proof. Set $\beta = \sigma^j \alpha$. If $C(\alpha, \beta) \neq 1$, then $e(\alpha)e(\beta) \neq e(\beta)e(\alpha)$. Yet by Theorem 4, we have $e(\alpha)e(\beta)^{-1} \in \mathbb{C}^\times$. Therefore, R acts by 0 on every twisted \mathfrak{V}_Λ -module, hence by Theorem 3 so does \mathfrak{V}_Λ . \square

3.8. Semisimplicity of twisted representations. Consider the category $\mathcal{O}_{\hat{\sigma}}$ of $\mathfrak{h}^{[0]}$ -diagonalizable twisted with respect to $\hat{\sigma}$ modules V over \mathfrak{V}_Λ such that V satisfies the conditions of Lemma 1 as a module over the Heisenberg algebra H . In particular V must be a module over the vertex operator algebra \mathfrak{V}_Λ . Note that if $V = \bigoplus_n V_n$ is a $\hat{\sigma}$ -twisted module over the vertex operator algebra \mathfrak{V}_Λ such that $V_n = 0$ for $n \ll 0$, then $V \in \mathcal{O}_{\hat{\sigma}}$. The latter modules appear in the representation theory of vertex algebras quite often, in particular in connection with the Zhu theory [9, 27].

Note that by Corollary 3, the category $\mathcal{O}_{\hat{\sigma}}$ can sometimes be trivial.

Theorem 5. *The category $\mathcal{O}_{\hat{\sigma}}$ is semisimple with finitely many isomorphism classes of simple objects.*

In order to prove this theorem we have to study the quotient algebra A of R modulo the ideal generated by the condition (i) of Theorem 4. More precisely, let $\Pi \subset \Lambda$ be a finite set of vectors, closed under σ and spanning Λ over \mathbb{Z} . For $x \in R$ let \bar{x} be its image in A . The algebra A is generated by the set $\{\overline{e(\alpha)} \mid \alpha \in \Pi\}$ subject to the relations $\overline{e(\sigma\alpha)} = \mu_\alpha \phi(\alpha)^{-1} \overline{e(\alpha)}$ for all $\alpha \in \Pi$.

Let $\Pi = \bigsqcup_{j=1}^n S_j$ be the decomposition of Π into a disjoint union of σ -orbits. Choose some $\alpha_j \in S_j$ for all $1 \leq j \leq n$. Let $x_j = \overline{e(\alpha_j)} \in A$. Then the set $x_1^{\pm 1}, \dots, x_n^{\pm 1}$ is a set of generators of A . We note that $x_i x_j = c_{ij} x_j x_i$ for $c_{ij} = C(\alpha_i, \alpha_j) \in \mathbb{C}^\times$, see (16). The algebra A depends on the cocycle $\varphi : \Lambda \rightarrow \mathbb{C}^\times$ and also on the choice of a root $\mu_j = \sqrt[p]{\varphi(\alpha_j) \cdots \varphi(\sigma^{p-1} \alpha_j)}$ for $1 \leq j \leq n$.

Let $\nu : \Lambda \rightarrow (\mathfrak{h}^{[0]})^*$ be the map given by $\nu(\alpha)h = (\alpha|h)$. Then $\nu(\Lambda)$ is a sublattice of the dual lattice to $\Lambda^{[0]}$. The algebra R is graded by $\nu(\Lambda)$ by setting $\deg e(\alpha) = \nu(\alpha)$. We observe that the relations (i) of Theorem 4 are homogeneous with respect to this grading, because $\nu(\alpha) = \nu(\sigma\alpha)$ for all $\alpha \in \Lambda$. Hence A is graded by $\nu(\Lambda)$ as well.

Recall that a graded algebra is called semisimple if it is semisimple as a left graded module over itself. Like in a non-graded case, a graded algebra is semisimple if and only if it is decomposed into a direct product of graded simple algebras, the latter are isomorphic to matrix algebras over

graded division algebras, by a graded version of the classical Wedderburn theorem. If an algebra A is graded semisimple, then the category of graded R -modules is semisimple, i.e. every module is completely reducible. The simple graded A -modules are the simple homogeneous ideals of A with possible shift of degrees. For the non-graded case this is a classical theory presented in most of the graduate algebra textbooks, like Jacobson's *Basic Algebra* [13]. For the graded case the references are much more scarce, see, however, [24].

We will use the following rather obvious fact: A homomorphic image of a graded semisimple algebra modulo a homogeneous ideal is again a graded semisimple algebra.

Proposition 4. *The algebra A is $\nu(\Lambda)$ -graded semisimple.*

Proof. Let $\xi_j = \deg \alpha_j = \nu(\alpha_j) \in \nu(\Lambda)$. It is easy to see that $\xi_j = 0$ if and only if $\sum_{\alpha \in S_j} \alpha = 0$. We can assume that $\xi_1, \dots, \xi_m \neq 0$ for some $m \leq n$ and $\xi_{m+1} = \dots = \xi_n = 0$. Clearly, ξ_j 's span $\nu(\Lambda)$ over \mathbb{Z} .

Assume also that our choice of Π yields the minimal possible value of m . We claim that in this case ξ_1, \dots, ξ_m is a \mathbb{Z} -basis of $\nu(\Lambda)$. Indeed, otherwise there is an invertible matrix $M \in SL(m, \mathbb{Z})$ such that the m -th column of $(\xi_1 \dots \xi_m) M$ is 0. Let α'_j be the j -th column of $(\alpha_1 \dots \alpha_m) M$ for $1 \leq j \leq m$, and let Π' be the closure of $\{\alpha'_1, \dots, \alpha'_m\} \cup S_{m+1} \cup \dots \cup S_n$ with respect to the action of σ . Then Π' is a generating set of Λ , closed under σ and, since $\nu(\alpha'_m) = 0$ the number of σ -orbits in Π' with non-zero degree is at most $m - 1$.

Suppose that Π is not an integer basis of Λ . Then there is a linear relation in Λ of the form

$$\sum_{j=1}^n \sum_{s=0}^{p-1} r_{js} \sigma^s \alpha_j = 0, \quad r_{js} \in \mathbb{Z}.$$

In R this relation becomes $\prod_{j,s} e(\sigma^s \alpha_j)^{r_{js}} = \theta'$, where $\theta' \in \mathbb{C}^\times$ is a product of some values of the cocycle ε . In A this relation becomes

$$\prod_{j=1}^n x_j^{r_j} = \theta, \quad r_j = \sum_{s=0}^{p-1} r_{js},$$

for some other constant $\theta \in \mathbb{C}^\times$.

For $m+1 \leq j \leq n$ we have $\sum_s \sigma^s \alpha_j = 0$, hence in A we get the relation $x_j^p = \theta_j \in \mathbb{C}^\times$. By the change of variables $x_j \mapsto \theta_j^{-1/p} x_j$ we can make $\theta_j = 1$.

A look at the formula (16) for the cocycle C shows that if $\sum_{s=0}^p \sigma^s \alpha = 0$ for some $\alpha \in \Lambda$, then $C(\alpha, \beta)^p = 1$ for any other $\beta \in \Lambda$.

Summing up, we get that A is a homomorphic image of the algebra

$$B = \left\langle x_1^{\pm 1}, \dots, x_n^{\pm 1} \left| \begin{array}{l} x_i x_j = c_{ij} x_j x_i \text{ for } 1 \leq i, j \leq n \\ x_j^p = 1 \text{ for } m+1 \leq j \leq n \end{array} \right. \right\rangle, \quad (22)$$

where $c_{ij} = c_{ji}^{-1} \in \mathbb{C}^\times$ for $1 \leq i, j \leq n$ and $c_{ij}^p = 1$ if either $i > m$ or $j > m$.

Therefore, it is enough to show that B is $\nu(\Lambda)$ -graded semisimple. The proposition now follows from the following lemma. \square

Lemma 2. *Let B be the associative algebra defined by (22).*

- B is $\nu(\Lambda)$ -graded semisimple.
- A basis of B is given by all monomials $x_1^{r_1} x_2^{r_2} \dots x_n^{r_n}$ where $r_j \in \mathbb{Z}$ and $0 \leq r_j \leq p-1$ for $m+1 \leq j \leq n$.

Proof. Let us prove (a) by induction on $n - m$. If $n = m$, then B is a twisted Laurent polynomial algebra in x_1, \dots, x_m . Recall that the degrees $\xi_j = \deg x_j$ for $1 \leq j \leq m$ form a basis of the grading lattice $\nu(\Lambda)$, hence B in this case is a $\nu(\Lambda)$ -graded division algebra. Assume that $n > m$, and let $B' \subset B$ be the subalgebra generated by $x_1^{\pm 1}, \dots, x_{n-1}^{\pm 1}$. By induction, B' is graded semisimple, hence it is decomposed into a direct product $B' = \prod_r E_r$ of simple homogeneous ideals such that

$E_r E_s = \delta_{rs} E_r$. Note that a left ideal is also right, since for any $x, y \in B$ there is $c \in \mathbb{C}^\times$ such that $xy = c yx$. Let

$$e_j = \frac{1}{\sqrt{p}} \sum_{s=0}^{p-1} \omega^{js} x_n^s \in \mathbb{C}[x_n] / \langle x_n^p - 1 \rangle \subset B, \quad 0 \leq j \leq p-1,$$

be the idempotents of the subalgebra generated by x_n , so that $e_i e_j = \delta_{ij} e_i$. Then for each r and j the space $E_r e_j$ is a simple ideal of B , and we have $(E_r e_j)(E_s e_i) = \delta_{rs} \delta_{ij} E_r e_j$ and $B = \prod_{r,j} E_r e_j$. This proves (a).

To prove (b) we note that the defining relations of B in (22) form a Gröbner basis of the ideal of relations, and the statement can be easily verified using Diamond lemma techniques, see e.g. [2]. \square

Proof of Theorem 5. Let $V \in \mathcal{O}_{\hat{\sigma}}$. By (i) of Lemma 1 and Theorem 4 we have $V = M(1) \otimes \Omega$, where Ω is $\nu(\Lambda)$ -graded module over A . Hence it follows from Proposition 4 that Ω is decomposed into a direct sum of irreducible A -modules, and Corollary 2 implies that \mathfrak{V} is decomposed into a direct sum of irreducible \mathfrak{V}_Λ -modules.

A module $V = M(1) \otimes \Omega \in \mathcal{O}_{\hat{\sigma}}$ is simple if and only if Ω is a simple $\nu(\Lambda)$ -graded A -module. Such Ω must be isomorphic to a simple homogeneous ideal of A up to a shift of weights. The weights of Ω are restricted by (ii) of Theorem 4. We claim that a simple object of $\mathcal{O}_{\hat{\sigma}}$ is determined up to an isomorphism by a choice of a simple homogeneous ideal of A , an equivalence class $\eta + \nu(\Lambda) \in (\Lambda^{[0]})' / \nu(\Lambda)$ and a root $\mu_\alpha = \sqrt[p]{\varphi(\alpha) \cdots \varphi(\sigma^{p-1} \alpha)}$ for each generating σ -orbit of Λ .

Indeed, let $\xi \in (\mathfrak{h}^{[0]})^*$ be such that $\xi(\alpha(0)) = \frac{1}{2}(\alpha' | \alpha') - \frac{1}{2\pi i} \ln \mu_\alpha$. Let $I \subset A$ be a homogeneous ideal. Pick a homogeneous element $a \in I$. Then by Theorem 4 (ii), $\text{wt } a \equiv \xi \pmod{(\Lambda^{[0]})'}$. Automorphisms of I can vary $\text{wt } a$ by $\nu(\Lambda)$, so the weights of the graded A -module I are defined uniquely by a class in $(\Lambda^{[0]})' / \nu(\Lambda)$.

It follows that there are at most finitely many isomorphism classes of simple objects in $\mathcal{O}_{\hat{\sigma}}$. \square

Example 1. If $\sigma = \text{Id}$, then $\nu(\Lambda) \cong \Lambda$, since the form is non-degenerate, and $A = R$ is a Λ -graded division algebra. The automorphism $\hat{\text{id}} : \mathfrak{V}_\Lambda \rightarrow \mathfrak{V}_\Lambda$ is defined by choosing an arbitrary values $\mu_j = \varphi(\alpha_j) \in \mathbb{C}^\times$ for a basis $\alpha_1, \dots, \alpha_l$ of Λ . The simple objects of category $\mathcal{O}_{\hat{\text{id}}}$ are parametrized by Λ' / Λ , which agrees with the result of Dong [5]. Let $\xi \in \mathfrak{h}^*$ be the functional defined by $\xi(\alpha_j) = \frac{1}{2\pi i} \ln \mu_j$. Then the simple object of category $\mathcal{O}_{\hat{\text{id}}}$ corresponding to an equivalence class $\eta + \Lambda \in \Lambda^* / \Lambda$ is $M(1) \otimes \mathbb{C}[\Lambda]$, where $\mathbb{C}[\Lambda]$ is graded such that $\text{wt } e^0 = \eta + \xi$.

Example 2. A more interesting case is when $\sigma = -1$. Then $\Pi = \{\pm \alpha_1, \dots, \pm \alpha_l\}$ and the grading is trivial. One can choose the cocycle ε such that $\varepsilon(\alpha_j, \alpha_j) = 1$ and then $\varphi(-\alpha_j) = \varphi(\alpha_j)^{-1}$, hence $\mu_j = \pm 1$ for all $1 \leq j \leq l$. So the algebra A in this case is generated by x_1, \dots, x_l modulo the relations $x_i x_j = c_{ij} x_j x_i$, where $c_{ij} = (-1)^{(\alpha_i | \alpha_i)(\alpha_j | \alpha_j)} (-1)^{(\alpha_i | \alpha_j)}$, and $x_j^2 = \mu_j \phi(\alpha_j)$. By Lemma 2, A is a semisimple algebra of dimension 2^l for any choice of φ and μ_j . A simple module $V \in \mathcal{O}_{\hat{\sigma}}$ is decomposed $V = M(1) \otimes \Omega$, where Ω isomorphic to a simple ideal of A . This case was studied in [6, 10, 12].

We remark that -1 is an isomorphism of any lattice Λ . When Λ is the Leech lattice, twisted modules over \mathfrak{V}_Λ with respect to this isomorphism are used to construct the Moonshine vertex algebra V^\natural , see [12].

Remark. The case when Λ is a simply-laced root lattice and σ is an element of the Weyl group of corresponding affine Kac-Moody algebra was studied by Kac and Peterson [17].

REFERENCES

- [1] B. Bakalov, V. G. Kac, and I. T. Todorov. Lattice orbifolds. Preprint.
- [2] G. M. Bergman. The Diamond lemma for ring theory. *Adv. in Math.*, 29(2):178–218, 1978.
- [3] R. E. Borcherds. Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. U.S.A.*, 83(10):3068–3071, 1986.
- [4] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde. The operator algebra of orbifold models. *Comm. Math. Phys.*, 123(3):485–526, 1989.
- [5] C. Dong. Vertex algebras associated with even lattices. *J. Algebra*, 161(1):245–265, 1993.
- [6] C. Dong. Twisted modules for vertex algebras associated with even lattices. *J. Algebra*, 165(1):91–112, 1994.

- [7] C. Dong and J. Lepowsky. *Generalized vertex algebras and relative vertex operators*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [8] C. Dong and J. Lepowsky. The algebraic structure of relative twisted vertex operators. *J. Pure Appl. Algebra*, 110(3):259–295, 1996.
- [9] C. Dong, H. Li, and G. Mason. Twisted representations of vertex operator algebras. *Math. Ann.*, 310(3):571–600, 1998.
- [10] C. Dong and K. Nagatomo. Automorphism groups and twisted modules for lattice vertex operator algebras. In *Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998)*, pages 117–133. Amer. Math. Soc., Providence, RI, 1999.
- [11] I. B. Frenkel, Y.-Z. Huang, and J. Lepowsky. On axiomatic approaches to vertex operator algebras and modules. *Mem. Amer. Math. Soc.*, 104(494), 1993.
- [12] I. B. Frenkel, J. Lepowsky, and A. Meurman. *Vertex Operator Algebras and the Monster*. Academic Press, Boston, MA, 1988.
- [13] N. Jacobson. *Basic algebra. II*. W. H. Freeman and Company, New York, second edition, 1989.
- [14] V. G. Kac. *Infinite-Dimensional Lie Algebras*. Cambridge University Press, Cambridge, third edition, 1990.
- [15] V. G. Kac. *Vertex Algebras for Beginners*, volume 10 of *University Lecture Series*. AMS, Providence, RI, second edition, 1998.
- [16] V. G. Kac. Formal distribution algebras and conformal algebras. In *XIIth International Congress of Mathematical Physics (ICMP '97) (Brisbane)*, pages 80–97. Internat. Press, Cambridge, MA, 1999.
- [17] V. G. Kac and D. H. Peterson. 112 constructions of the basic representation of the loop group of E_8 . In *Symposium on anomalies, geometry, topology (Chicago, Ill., 1985)*, pages 276–298. World Sci. Publishing, Singapore, 1985.
- [18] J. Lepowsky. Calculus of twisted vertex operators. *Proc. Nat. Acad. Sci. U.S.A.*, 82(24):8295–8299, 1985.
- [19] J. Lepowsky and R. L. Wilson. Construction of the affine Lie algebra $A_1^{(1)}$. *Comm. Math. Phys.*, 62(1):43–53, 1978.
- [20] J. Lepowsky and R. L. Wilson. A new family of algebras underlying the Rogers-Ramanujan identities and generalizations. *Proc. Nat. Acad. Sci. U.S.A.*, 78(12, part 1):7254–7258, 1981.
- [21] H. Li. Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules. In *Moonshine, the Monster, and related topics (South Hadley, MA, 1994)*, pages 203–236. Amer. Math. Soc., Providence, RI, 1996.
- [22] H. Li. Local systems of vertex operators, vertex superalgebras and modules. *J. Pure Appl. Algebra*, 109(2):143–195, 1996.
- [23] H. Li and X. Xu. A characterization of vertex algebras associated to even lattices. *J. Algebra*, 173(2):253–270, 1995.
- [24] C. Năstăsescu and F. van Oystaeyen. *Graded ring theory*. North-Holland Publishing Co., Amsterdam, 1982.
- [25] M. Primc. Vertex algebras generated by Lie algebras. *J. Pure Appl. Algebra*, 135(3):253–293, 1999.
- [26] M. Roitman. On free conformal and vertex algebras. *J. Algebra*, 217(2):496–527, 1999.
- [27] Y. Zhu. Modular invariance of characters of vertex operator algebras. *Journal of the AMS*, 9(1):237–302, 1996.

M.S.R.I., 1000 CENTENNIAL DR., BERKELEY, CA 94720

E-mail address: roitman@msri.org