

ON CERTAIN REPRESENTATIONS OF AUTOMORPHISM GROUPS OF AN ALGEBRAICALLY CLOSED FIELD

M.ROVINSKY

1. INTRODUCTION

Let k be an algebraically closed field of characteristic zero, F its algebraically closed extension of transcendence degree $n < \infty$, and $G = G_{F/k}$ be the group of automorphisms over k of the field F . Let the set of subgroups $U_L := \text{Aut}(F/L)$ for all subfields L finitely generated over k be the basis of neighborhoods of the unity in G .

The main purpose of this note (cf. Theorem 2.6 and Corollary 2.9) is to show that any non-exact continuous representation of G factors through a discrete quotient of G . In Section 3 the smooth G -modules F/k and F^\times/k^\times are studied. It is shown that they are irreducible as modules over the subgroup G° of G generated by its compact subgroups, and the induced representations of the algebra \mathbf{D} of the compactly supported measures on G are exact. This shows that the radical of \mathbf{D} is trivial. However, it is shown in the Section 4 that the annihilator of the *admissible* irreducible G -modules, is non-trivial.

There is an evident link between G -modules and some geometric objects. Namely, for a variety X over k there is a natural smooth G -action on the group of cycles on $X_F := X \times_k F$. In particular, if X is irreducible of dimension $\leq n$ over k and $k(X)$ is embedded into F then the \mathbb{Q} -space of “generic” 0-cycles on X_F is isomorphic to the representation $\text{ind}(G, G_{F/k(X)}, \mathbb{Q})$ of G compactly induced by the trivial one-dimensional representation \mathbb{Q} of $G_{F/k(X)}$. More particularly, if X is smooth and proper, then the \mathbb{Q} -space $A^{\dim X}(X_F)$ of 0-cycles on X_F modulo an “adequate” equivalence relation is a quotient of the G -module $\text{ind}(G, G_{F/k(X)}, \mathbb{Q})$.

In some cases one can identify the groups of morphisms between geometric objects with the groups of morphisms between corresponding G -modules (cf. Proposition 5.4 and Corollary 5.5).

It is shown in Proposition 5.9 that each “primitive geometric representation of level n ” carries a non-degenerate symmetric G -equivariant form with values in a one-dimensional G -module, which is definite at least if $n \leq 2$.

1.1. Notations and terminology. For a field F_1 and its subfield F_2 we denote by G_{F_1/F_2} the group of automorphisms of the field F_1 over F_2 . Throughout the note k is an algebraically closed field of characteristic zero, F its algebraically closed extension of transcendence degree $1 \leq n < \infty$ and $G = G_{F/k}$. If K is a subfield of F then \overline{K} denotes its algebraic closure in F .

For a topological group H we denote by H° its subgroup generated by the compact subgroups.

A representation W of a topological group is called *smooth*, if stabilizers of all vectors in W are open. A smooth representation W is called *admissible* if fixed vectors of each open subgroup form a finite-dimensional subspace in W .

$\mathbb{Q}(\chi)$ is the quotient of the free abelian group generated by the set of compact open subgroups in G° by the relations $[U] = [U : U'] \cdot [U']$ for all $U' \subset U$. It is a one-dimensional \mathbb{Q} -vector space oriented by $[U] > 0$ for any U . The group G acts on it by the conjugations.

$\mathbf{D} = \lim_{\leftarrow U: \text{open}} \mathbb{Q}[G/U]$ is the algebra of compactly supported \mathbb{Q} -valued measures on G .

For each compact open subgroup U in G we denote by \mathcal{H}_U the Hecke subalgebra $h_U * \mathbf{D} * h_U$ in \mathbf{D} of U -invariant measures. Here h_U is the Haar measure on U , which is the identity in \mathcal{H}_U .

2. NORMAL CLOSED SUBGROUPS IN G°

We consider a topology on G with the basis of neighborhood of an automorphism $\sigma : F \xrightarrow{\sim} F$ over k given by the cosets of the form σU_L for all subfields L of F finitely generated over k , where $U_L = \text{Aut}(F/L)$. It is shown in [IIII-III], Lemma 1, Section 3 that the map

$$\{\text{subfields in } F \text{ over } k\} \longrightarrow \{\text{closed subgroups in } G\} \quad \text{given by} \quad K \longmapsto \text{Aut}(F/K)$$

is injective and induces one-to-one correspondences

- $\{\text{subfields } K \text{ of } F \text{ with } k \subseteq K \text{ and } F = \overline{K}\} \leftrightarrow \{\text{compact subgroups of } G\};$
- $\left\{ \begin{array}{l} \text{subfields } K \text{ of } F \text{ finitely} \\ \text{generated over } k \text{ with } F = \overline{K} \end{array} \right\} \leftrightarrow \{\text{compact open subgroups of } G\}.$

The inverse correspondences are given by $G \supset H \longmapsto F^H \subseteq F$.

One checks that the group G is Hausdorff, locally compact, and totally disconnected.

Denote by G° the subgroup of G generated by the compact subgroups. Obviously, G° is an open normal subgroup in G .

Lemma 2.1. *The common normalizer in G of all subgroups of index ≤ 2 in U_L coincides with U_L .*

Proof. For any extension L' of L any element τ in the normalizer of $U_{L'}$ satisfies $\tau(L') = L'$. Fix a smooth projective model X of L . Then τ induces a rational map $X \xrightarrow{\tau} X$. Suppose that $\tau|_L \neq \text{id}$. Then there is a point $x \in X$ outside the ramification such that $x \neq \tau(x)$. Let D and D' be linear equivalent very ample effective reduced divisors on X outside the branch locus passing through x , but not through $\tau(x)$, and $(f) = D - D'$. Then $D \ni x$, $\tau^{-1}(D) \not\ni x$ and all multiplicities are equal to 1, so $\tau f/f \notin L^{\times 2}$, and therefore, $f \in L^\times - \{g \in L^\times : \tau g \in L^{\times 2} g\}$, so by Kummer theory, $\tau(L(f^{1/2})) \neq L(f^{1/2})$. \square

Corollary 2.2. *For any $\xi \in G - \{1\}$ and an open subgroup $U \subseteq G$ there is $\sigma \in U$ such that $[\sigma, \xi] \in U - \{1\}$.*

Proof. Let $L \subset F$ be finitely generated over k with $\overline{L} = F$ and $U_L \subseteq U$. Set $L' = L\xi(L)$. Then $\xi^{-1}(\sigma\xi\sigma^{-1})|_L = \text{id}$ for any $\sigma \in U_{L'}$, so there is $\sigma \in U_{L'}$ with $\xi^{-1}\sigma\xi\sigma^{-1} \in U_L - \{1\}$. \square

Lemma 2.3. *Let H be a normal closed subgroup in G° . Then for any algebraically closed extension k' of k in F , over which F is of transcendence degree one, there is such $x \in F - k'$ that H contains $\text{Gal}(F/k'(x))$.*

Proof. By Corollary 2.2, for any $\xi \in H - \{1\}$ and any compact open subgroup U there is an element $\tau \in U$ such that $[\xi, \tau] \in U - \{1\}$, and therefore, $H \cap U \neq \{1\}$. For any element of $(H \cap U) - \{1\}$ there its power σ fixing a transcendence basis of k' over k , and thus, inducing

an automorphism of k' . By a well-known result of E. Artin, $\sigma \neq 1$. Then there is an element $x \in F - k'$ and an integer $N \geq 2$ such that $\sigma x^N = x^N$ and $\sigma x \neq x$.

Let L be a finite Galois extension of $k'(x)$. It is unramified over a finite set S of points on a smooth projective model of $k'(x)$ over k' . Then there is such an element $\bar{\alpha}_L \in \text{Aut}(k'(x)/k')$ that the set $\bar{\alpha}_L^{-1}\sigma\bar{\alpha}_L(S)$ does not intersect S , and therefore, for an extension $\alpha_L \in G_{F/k'}^\circ$ of $\bar{\alpha}_L$ to F , the field $L \cap \alpha_L^{-1}\sigma\alpha_L(L)$ is unramified over a smooth projective model of $k'(x)$, so $L \cap \alpha_L^{-1}\sigma\alpha_L(L) = k'(x)$.

Let $\beta \in G_{F/k'}^\circ$ be given on $k'(x)$ by $\bar{\alpha}_L$, and somehow extended to the field $\alpha_L^{-1}\sigma\alpha_L(L)$. Then $\beta^{-1} \circ \sigma^{-1} \circ \beta \circ \alpha_L^{-1} \circ \sigma \circ \alpha_L$ is the identity on $k'(x)$, and therefore, induces an automorphism of L . Since L and $\alpha_L^{-1}\sigma\alpha_L(L)$ are Galois extensions of $L \cap \alpha_L^{-1}\sigma\alpha_L(L) = k'(x)$, there is such an extension of β to F that for its restriction to L the composition $\beta^{-1} \circ \sigma^{-1} \circ \beta \circ \alpha_L^{-1} \circ \sigma \circ \alpha_L$ is any given automorphism of L over $k'(x)$. This means that the natural projection $H \cap \text{Gal}(F/k'(x)) \rightarrow \text{Gal}(L/k'(x))$ is surjective for any Galois extension L of $k'(x)$, i.e., that $H \cap \text{Gal}(F/k'(x))$ is dense in $\text{Gal}(F/k'(x))$. As $H \cap \text{Gal}(F/k'(x))$ is closed, we have $H \supseteq \text{Gal}(F/k'(x))$. \square

Lemma 2.4. *If $n = 1$ then there are no proper normal open subgroups in G° .*

Proof. Let H be a normal open subgroup in G° . Then for a subfield $L \subset F$ finitely generated over k one has $U_L \subseteq H$. For any purely transcendental extension $L' \subset F$ of k with $\bar{L}' = F$ one also has $U_{LL'} \subseteq H$, as well as $H \supseteq \langle \sigma U_{LL'} \sigma^{-1} \mid \sigma \in N_{G^\circ} U_{L'} \rangle \subseteq U_{L'}$.

The group in the middle is open, so it is closed, and therefore, it is the absolute Galois group of the field $\bigcap_{\sigma \in N_{G^\circ} U_{L'}} \sigma(LL')$, that coincides with L' . Indeed, LL' is ramified only over a divisor on \mathbb{P}_k^n , but the group $\text{PGL}_2 k$ does not preserve this divisor, so the intersection $\bigcap_{\sigma} \sigma(LL')$ consists of unramified elements, i.e., of the elements of L' .

This proves that $H \supseteq U_{L'}$ for any purely transcendental extension $L' \subset F$ of k with $\bar{L}' = F$, and therefore, H contains all compact subgroups in G , so finally, $H \supseteq G^\circ$. \square

Lemma 2.5. *Suppose that a closed subgroup H in G° contains subgroups $G_{F/k'}^\circ$ for all algebraically closed extensions k' of k in F of transcendence degree $n - 1$. Then $H = G^\circ$.*

Proof. Any element σ of a compact subgroup in G° can be presented as the limit of a compatible collection (σ_L) of embeddings σ_L of finitely generated extensions L of k in F , over which F is algebraic, into F . For L sufficiently large, F is algebraic over $L \cap F^{(\sigma)} =: L_0$.

Consider a finite extension L' of L , which is normal over L_0 . There is a $\text{Gal}(L'/L_0)$ -equivariant projective model X of L' over k . There is a fibration of X by $\text{Gal}(L'/L_0)$ -invariant irreducible curves. This fibration descends to a fibration of a model of L_0 over k with the same base. Then the function field of the base is a subfield k'' in L_0 , over which the transcendence degree of F is one, and which is algebraically closed in L' . Then there is an extension of σ_L , or even of $\sigma_{L'}$, to an element of $G_{F/\bar{k}''}^\circ$. \square

Theorem 2.6. *Any closed proper normal subgroup in G° is trivial.*

Proof. Let H be a non-trivial closed normal subgroup in G° . By Lemma 2.3, the group $H \cap G_{F/k'}^\circ$ is an open normal subgroup in $G_{F/k'}^\circ$, and thus, by Lemma 2.4, $H \cap G_{F/k'}^\circ = G_{F/k'}^\circ$. Then Lemma 2.5 implies that $H = G^\circ$. \square

Corollary 2.7. *There are no proper closed subgroups of finite index in G° .*

Proof. If H is a closed subgroup of finite index in G° then $\mathbb{Q}[G^\circ/H]$ is a finite-dimensional G° -representation. The kernel of such representation is non-trivial, since $\text{GL}(\mathbb{Q}[G^\circ/H])$ is countable, but G° is not. As the kernel of the G° -representation $\mathbb{Q}[G^\circ/H]$ is a closed and normal subgroup in G° , it coincides with G° , and thus, $H = G^\circ$. \square

Corollary 2.8. *Any finite-dimensional smooth representation of G° is trivial.*

(Proof. This is clear, since there are no proper open normal subgroups in G° .) \square

Corollary 2.9. *For any subgroup H of G containing G° and any continuous representation π of H either π is exact, or the restriction of π to G° is trivial.*

Proof. If π is not exact then, by Corollary 2.2, its kernel has a non-trivial intersection with G° . Then, by Theorem 2.6, the kernel of π contains G° . \square

3. THE G -MODULES F/k AND F^\times/k^\times

Lemma 3.1. *If $n = 1$ then for any element $\sigma \neq 1$ of G closure of its G° -orbit $\{\tau\sigma\tau^{-1} : \tau \in G^\circ\}$ in G is not compact. In particular, there are no normal compact subsets in G except $\{1\}$.*

Proof. Suppose that for an element $\sigma \neq 1$ of G the closure of its G° -orbit is compact. Then the G° -orbit of σ is a subset in a disjoint union $\bigcup_{j=1}^N \sigma_j U_L$ for some field $L \neq k$ finitely generated over k and elements $\sigma_j \in G$. We may assume that $\sigma_j \notin U_L$ for all $2 \leq j \leq N$ and the G° -orbit of σ is not a subset in a proper sub-union. This implies that for any $\tau \in G^\circ$ and each j intersection of $\sigma_j U_L$ with $\tau\sigma_s U_L \tau^{-1}$ is non-empty for at least one s .

Consider the k -subspace $V_j = \{t \in L \mid \sigma_j t - t \in k\}$ in L . If $V_j \neq V_j^{(\sigma_j)}$ then for any $t \in V_j - V_j^{(\sigma_j)}$ there is some $a \in k^\times$ with $\sigma_j t^2 - t^2 = 2at + a^2 \notin k$, so $V_j \neq L$. This shows that $L \not\subseteq \bigcup_{j=1}^N V_j$, so we can choose such an element $x \in L$ that $\sigma_j x - x \notin k$ for each $j \geq 2$.

Let τ be such an element of G° that $\tau x = x + 1$. Then for each pair (j, s) with intersecting $\sigma_j U_L$ and $\tau\sigma_s U_L \tau^{-1}$ action of an element in the intersection $\sigma_j U_L \cap \tau\sigma_s U_L \tau^{-1}$ on τx gives $\sigma_j x + 1 = \tau\sigma_s x$. As $\sigma_j U_L \cap U_{\tau(L)} = \emptyset$ for all $2 \leq j \leq N$, since otherwise $\sigma_j \in U_{\tau(L)} U_L \subseteq U_{k(x)}$, and thus, $\sigma_j x = x$, this implies that for some j with $\sigma_j \notin U_L$, $M > 0$ one has $\sigma_j x = \tau^M \sigma_j x - M$. Thus, $\sigma_j x - x = \tau^M(\sigma_j x - x)$, so τ belongs to a compact subgroup, contradicting to $\tau x = x + 1$. \square

Corollary 3.2. *The G° -modules F/k and F^\times/k^\times are irreducible.*

Proof. Let A be the additive subgroup of F generated by the G° -orbit of x . For any $y \in A - k$ one has $\frac{2}{y^2-1} = \frac{1}{y-1} - \frac{1}{y+1}$. As $\frac{1}{y-1}$ and $\frac{1}{y+1}$ are in the G° -orbit of y , this implies that $y^2 \in A$. As for any $y, z \in A$ one has $yz = \frac{1}{4}((y+z)^2 - (y-z)^2)$, the group A is a subring of F .

Let M be the multiplicative subgroup of F^\times generated by the G° -orbit some of $x \in F - k$. Then for any $y, z \in M$ one has $y + z = z(y/z + 1)$, so if $y/z \notin k$ then $y + z \in M$, and thus, $M \cup \{0\}$ is a G° -invariant subring of F .

Suppose that the G° -orbit of an element $x \in F - k$ generates a proper subring, and thus, a k -subalgebra of F . Since the G° -orbit of an element x contains all elements of $F - k(x)$, if $n \geq 2$ then each element of F is the sum of a pair of elements in the orbit. If $n = 1$ then $\text{Gal}(F/\mathbb{Q}(G^\circ x)) \subset G^\circ$ is a compact subgroup normalized by G° . Then by Lemma 3.1 we

have $\text{Gal}(F/\mathbb{Q}(G^\circ x)) = \{1\}$. As any element of $\mathbb{Q}(G^\circ x)$ is the fraction of a pair of elements in $\mathbb{Z}[G^\circ x]$ and for any $y \in F - k$ the element $1/y$ belongs to the G° -orbit of y , one has $\mathbb{Z}[G^\circ x] = F$. \square

Proposition 3.3. *The annihilators in \mathbf{D} of the smooth G -modules F/k and F^\times/k^\times are trivial.*

Proof. Let $\alpha \in \mathbf{D}$ annihilates F/k . Let $\sum_{j=1}^N a_j \cdot \sigma_j$ be the image of α in $\mathbb{Q}[G/U_L]$ for a compact open subgroup U_L , an integer $N \geq 1$, some $a_j \in \mathbb{Q}$ and some pairwise distinct embeddings $\sigma_j : L \hookrightarrow F$.

For any $x \in L$ one has $\sum_{j=1}^N a_j \cdot \sigma_j x \in k$, so for some $\tau \in G$ such that the embeddings $\sigma_1, \dots, \sigma_N, \tau\sigma_1, \dots, \tau\sigma_N$ are pairwise distinct one has $\sum_{j=1}^N a_j \cdot \sigma_j x + \sum_{j=1}^N (-a_j) \cdot \tau\sigma_j x = 0$ for all $x \in L$. Then, by Artin's theorem on independence of characters, $a_1 = \dots = a_N = 0$.

Similarly, let $\alpha \in \mathbf{D}$ annihilates F^\times/k^\times . Let $\sum_{j=1}^N a_j \cdot \sigma_j$ be the image of α in $\mathbb{Q}[G/U_L]$ for a compact open subgroup U_L , some $a_j \in \mathbb{Q}$ and some pairwise distinct embeddings $\sigma_j : L \hookrightarrow F$. We may suppose that $a_j \in \mathbb{Z}$.

For any $x \in L^\times$ one has $\prod_{j=1}^N (\sigma_j x)^{a_j} \in k^\times$, so for some $\tau \in G$ such that the embeddings $\sigma_1, \dots, \sigma_N, \tau\sigma_1, \dots, \tau\sigma_N : L \hookrightarrow F$ are pairwise distinct one has $\prod_{j=1}^{2N} (\sigma_j x)^{a_j} = 1$ for all $x \in L^\times$, where $\sigma_j = \tau\sigma_{j-N}$ and $a_j = -a_{j-N}$ for $N < j \leq 2N$.

Let τ_1, \dots, τ_M be a collection of pairwise distinct embeddings of L into F and b_1, \dots, b_M be such a collection of non-zero integers that $\sum_{j=1}^M |b_j|$ is minimal among those for which $\prod_{j=1}^M (\tau_j x)^{b_j} = 1$ for all $x \in L^\times$. Then $\prod_{j=1}^M (\tau_j x + 1)^{b_j} - \prod_{j=1}^M (\tau_j x)^{b_j} = 0$, which is equivalent to $\prod_{j:b_j > 0} (\tau_j x + 1)^{b_j} \prod_{j:b_j < 0} (\tau_j x)^{-b_j} - \prod_{j:b_j > 0} (\tau_j x)^{b_j} \prod_{j:b_j < 0} (\tau_j x + 1)^{-b_j} = 0$. We rewrite this as $\sum_{c_1, \dots, c_M} A_{c_1, \dots, c_M} \prod_{j=1}^M (\tau_j x)^{c_j} = 0$, where $0 \leq c_j \leq |b_j|$ and $\sum_{j=1}^M c_j < \sum_{j=1}^M |b_j|$. Then A_{c_1, \dots, c_M} are integers which are all non-zero if either $c_j = b_j$ for all j with $b_j > 0$, or $c_j = -b_j$ for all j with $b_j < 0$. By Artin's theorem on independence of characters, $\prod_{j=1}^M (\tau_j x)^{c_j} = \prod_{j=1}^M (\tau_j x)^{c'_j}$ for a pair of distinct collections (c_j) and (c'_j) as before, with $|c_j - c'_j| < |b_j|$ for some j , and for all $x \in L$. But then $\sum_{j=1}^M |c_j - c'_j| < \sum_{j=1}^M |b_j|$, contradicting our assumptions, so $a_1 = \dots = a_N = 0$. \square

4. REMARKS ON THE ADMISSIBLE RADICAL

As we have seen in the previous section, the common annihilator of all irreducible smooth representations of G is trivial. For each "integer" $1 \leq j \leq \infty$ we define \mathfrak{Rad}_j as the common annihilator of all admissible representations of G over \mathbb{Q} of length $\leq j$.

Clearly, $\bigcap_{j \geq 1} \mathfrak{Rad}_j \supseteq \mathfrak{Rad}_\infty$. We call $\mathfrak{Rad} := \mathfrak{Rad}_1$ the *admissible radical* of \mathbf{D} .

Let \mathbf{H} be the two-sided ideal in \mathbf{D} of locally translation-invariant measures on G . It is the union of the Hecke algebras \mathcal{H}_U for all open compact subgroups U in G .

Lemma 4.1. $\mathfrak{Rad}_i \cdot \mathfrak{Rad}_j \subseteq \mathfrak{Rad}_{i+j}$. *The restriction to \mathfrak{Rad} of the homomorphism $\mathbf{D} \xrightarrow{f_{\ker \chi}} \mathbb{Q}[\mathbb{Q}_+^\times]$, induced by the character χ of the representation $\mathbb{Q}(\chi)$, is zero.*

Proof. Any G -module W of length $\leq i + j$ contains a G -submodule W_1 of length $\leq i$ with the quotient $W/W_1 = W_2$ of length $\leq j$.

Let $\alpha \in \mathfrak{Rad}_i$ and $\beta \in \mathfrak{Rad}_j$. As $\beta(W_2) = 0$, one has $\beta(W) \subseteq W_1$, and thus, $\alpha\beta(W) = 0$.

For any $\beta \in \mathbb{Q}[\mathbb{Q}_+^\times] - \{0\}$ there is a homomorphism $\mathbb{Q}_+^\times \xrightarrow{\rho} \mathbb{Q}^\times$ inducing a ring homomorphism $\mathbb{Q}[\mathbb{Q}_+^\times] \xrightarrow{\rho} \mathbb{Q}$ such that $\rho(\beta) \neq 0$. If $\int_{\ker \chi}(\alpha) \neq 0$ then α acts on $\mathbb{Q}(\rho(\chi))$ non-trivially, contradicting the definition of \mathfrak{Rad} . \square

REMARK. This implies that there are admissible modules annihilated by \mathfrak{Rad} which are not semi-simple. Namely, let W be a finite-dimensional vector space over \mathbb{Q} and G acts via a composition $G \xrightarrow{\chi} \mathbb{Q}_+^\times \xrightarrow{\rho} \mathbb{Z} \xrightarrow{\lambda} \mathrm{GL}(W)$, where ρ is surjective and $\lambda(1)$ is a unipotent operator in $\mathrm{GL}(W)$. Then, by the above, the image of \mathfrak{Rad} in $\mathfrak{gl}(W)$ is zero.

Proposition 4.2. \mathfrak{Rad}_j are closed two-sided ideals in \mathbf{D} for all integer $j \geq 1$ and $j = \infty$.

$\mathfrak{Rad}_\infty \not\subseteq \mathbf{H}$. For any open compact subgroup U in G one has $\mathfrak{Rad}_\infty \cap \mathcal{H}_U \neq 0$, the algebra $\mathcal{H}_U/\mathfrak{Rad}$ is not semi-simple and it is infinite-dimensional.

Proof. It is straightforward that the common annihilator of any collection of G -modules is a two-sided ideal. The annihilator of a smooth vector w in any G -module is an open ideal, since it contains the left ideal generated by $\sigma - 1$ for all σ in the stabilizer of w . This implies that the common annihilator of any collection of smooth vectors in any collection of G -modules is a closed left ideal.

Let $L \subset F$ be a field with $\bar{L} = F$, h_L be the Haar measure on $\mathrm{Gal}(F/L) \subset G$, and suppose that $\sigma \in G$ induces an automorphism of L of finite order. Then $h_L\sigma = \sigma h_L = h_L\sigma h_L \in \mathbf{D}$. Let (W, π) be a smooth G -module. Since h_L is an idempotent, one has a decomposition $W = \ker(h_L) \oplus \ker(1 - h_L)$. Then $\pi(h_L\sigma)$ induces the zero map on $\ker(h_L)$, and it acts via an automorphism of finite order on $\ker(1 - h_L)$, which is the image of h_L . Suppose that σ is divisible as an element of $\mathrm{Aut}(L/k)$ and L is finitely generated over k . Let W be an admissible G -module such that $h_L(W) \neq 0$. Then, as the image of h_L coincides with $W^{U_L} \cong \mathbb{Q}^N$ for some integer $N \geq 1$, and $\pi(h_L\sigma)$ is a divisible torsion element of $\mathrm{GL}_N\mathbb{Q}$, one has $\pi(h_L\sigma) = \pi(h_L)$.

In particular, if U is a compact open subgroup in G and L is a purely transcendental extension of k in F^U , then, as $\mathrm{PGL}_{n+1}k$ is generated by divisible torsion elements, $h_L\sigma - h_L \in \mathfrak{Rad}_\infty \cap \mathcal{H}_U$ for any σ , inducing an automorphism of L in the subgroup $\mathrm{PGL}_{n+1}k \subseteq \mathrm{Aut}(L/k)$.

To construct an element in $\mathfrak{Rad}_\infty - \mathbf{H} \neq 0$, consider an abelian variety A of dimension n over k with the function field L_0 embedded into F . Each morphism $A \xrightarrow{N \times} A$ determines an extension L_N of L_0 in F of degree N^{2n} . Let L be the union of all these extensions. Let W be an admissible G -module. Since $L_N \cong L_0$, the subgroups U_{L_N} and U_{L_0} are conjugated in G , so $h_{L_0}(W)$ and $h_{L_N}(W)$ have the same dimension. As $h_{L_0}(W) \subseteq h_{L_N}(W)$, one has $h_{L_0}(W) = h_{L_N}(W)$. Let $e \in h_L(W)$ and $U_{L'} = \mathrm{Stab}_e \cap U_{L_0}$. As L' is a subfield in L finitely generated over k , one has $L' \subseteq L_N$ for some $N \geq 1$, so $e \in h_{L_N}(W)$, and thus, $h_{L_0}(W) = h_L(W)$. As h_{L_0} and h_L are projectors, this means that $h_{L_0} - h_L \in \mathfrak{Rad}_\infty$.

By Lemma 4.1, the surjection $\mathcal{H}_U \xrightarrow{\int_{\ker \chi}} \mathbb{Q}[\mathbb{Q}_+^\times]$ factors through $\mathcal{H}_U/\mathfrak{Rad}$, so for any open compact subgroup U in G the algebra $\mathcal{H}_U/\mathfrak{Rad}$ is infinite-dimensional and is not semi-simple. \square

Lemma 4.3. Let for a smooth proper curve X over k the canonical map $X \rightarrow \mathbb{P}_k^M$ be an embedding, and let $N \geq 2$ be such an integer that each projective $(N - 2)$ -subspace \mathbb{P} in \mathbb{P}_F^M intersects X_F at most at $N - 1$ point. Then the classes of pairwise distinct generic points $\sigma_1, \dots, \sigma_N$ of X are linearly independent in the group $\mathrm{Pic}(k(X) \otimes_k F)$.

Proof. Let $\{\omega_0, \dots, \omega_M\}$ be a basis of $\Gamma(X, \Omega_{X/k}^1)$. The canonical map $X_F \rightarrow \mathbb{P}_F^M$ is given by $\sigma \mapsto (\sigma\omega_0 : \dots : \sigma\omega_M)$ for any generic F -point σ . If for some m_j one has $\sum_{j=1}^N m_j \sigma_j = 0$ in the group $\text{Pic}(k(X) \otimes_k F)$ then $(\sum_{j=1}^N m_j \sigma_j \omega_0, \dots, \sum_{j=1}^N m_j \sigma_j \omega_M) = 0$, which means that the points $(\sigma_j \omega_0 : \dots : \sigma_j \omega_M)$ for $1 \leq j \leq N$ span at most an $(N-2)$ -dimensional projective subspace in \mathbb{P}_F^M , contradicting our assumption. \square

Proposition 4.4. *Support of any non-zero measure in \mathfrak{Rad} is infinite, i.e., $\mathfrak{Rad} \cap \mathbb{Q}[G] = 0$.*

Proof. By Proposition 5.3, the G -module $\text{Pic}(k(X) \otimes_k F)$ is admissible and semi-simple, and thus, it is annihilated by any element of \mathfrak{Rad} . It is enough to show that for any measure $\nu = \sum_{j=1}^N m_j \sigma_j$ with pairwise distinct σ_j and $m_j \in \mathbb{Q}^\times$ there is some smooth proper curve X with a non-zero action of ν on the corresponding $\text{Pic}(k(X) \otimes_k F)$.

There is such a smooth proper curve X of genus g that the minimal degree of maps from X to \mathbb{P}_k^1 is greater than N . Suppose that intersection of an $(N-2)$ -subspace $\mathbb{P} \subset \mathbb{P}^{g-1}$ with X in the canonical embedding contains a divisor D on X of degree N . Then under above assumptions $h^1(\Omega_X^1(-D)) = h^0(\mathcal{O}_X(D)) = 1$, and therefore, by Riemann–Roch formula, $\dim |\Omega_X^1(-D)| = g - N - 1$. On the other hand, the linear system of hyperplanes containing \mathbb{P} is $(g-N)$ -dimensional, leading to contradiction.

Let $L_{st} = F^{(\sigma_s = \sigma_t)} \subsetneq F$ for each pair $s \neq t$. Since any vector space over k is not a finite union of proper subspaces, the union $\bigcup_{1 \leq s < t \leq N} L_{st}$ never coincides with F , and we can embed the field $k(X)$ into F in such a way that $k(X)$ is not in the union $\bigcup_{1 \leq s < t \leq N} L_{st}$, so the points $\sigma_1|_{k(X)}, \dots, \sigma_N|_{k(X)}$ are pairwise distinct. Then, by Lemma 4.3, the classes of $\sigma_j|_{k(X)}$ are linearly independent in $\text{Pic}(k(X) \otimes_k F)$. \square

5. SOME GEOMETRIC REPRESENTATIONS

Now we are going to construct a supply of semi-simple admissible representations of G . Recall ([BZ]), that for each smooth G -module W and each open compact subgroup U of G the Hecke algebra \mathcal{H}_U acts on the space W^U .

Proposition 5.1 (=Proposition 2.10 of [BZ]). *1. A smooth \mathbb{Q} -representation W of G is irreducible if and only if for each subgroup U the \mathcal{H}_U -module W^U is irreducible.*
2. Let W_j for $j = 1, 2$ be smooth irreducible \mathbb{Q} -representations of G and $W_1^U \neq 0$ for some U . Then W_1 is equivalent to W_2 if and only if W_1^U is equivalent to W_2^U .
3. For each irreducible \mathbb{Q} -representation τ of the algebra \mathcal{H}_U there is a smooth irreducible representation W of G with $\tau \cong W^U$. \square

Lemma 5.2. *Let W be a smooth G -module. Suppose that for each compact open subgroup $U \subset G$ the \mathcal{H}_U -submodule W^U is semi-simple. Then W is semi-simple.*

Proof. Let W_0 be the sum of all irreducible G -submodules in W . Suppose $W \neq W_0$ and $e \in W - W_0$. Let V be the G -submodule in W generated by e , and $V_0 = V \cap W_0$. Let V_2 be a maximal G -submodule in $V - \{e\}$ containing V_0 . Then the G -module $V_1 = V/V_2$ is irreducible. Let U be such an open compact subgroup in G that $V_1^U \neq 0$. By Proposition 5.1(1), the \mathcal{H}_U -module V_1^U is irreducible. As the \mathcal{H}_U -module V^U is semi-simple and V^U/V_2^U is isomorphic to V_1^U , there is an \mathcal{H}_U -submodule N_U in V^U isomorphic to V_1^U and intersecting V_2 trivially.

For each open subgroup U' of U the $\mathcal{H}_{U'}$ -submodule $N_{U'}$ of U' -invariants of the G -submodule in V generated by N_U splits into a direct sum $\bigoplus_j A_j^{m_j}$ of irreducible $\mathcal{H}_{U'}$ -modules, $m_j \leq \infty$.

As $(N_{U'}^U)^U = h_U \mathbf{D} N_U = h_U \mathbf{D} h_U N_U = N_U$, there is exactly one index j such that $A_j^U \neq 0$. For this index one has $m_j = 1$. By Proposition 5.1(3), there are smooth irreducible representations W_1 and W_2 of G such that $W_1^{U'} \cong A_j$ and $W_2^{U'} \cong V_1^{U'}$. Since $W_1^U \cong W_2^U$, by Proposition 5.1(2), one has $W_1 \cong W_2$, so $A_j \cong V_1^{U'}$. Set $N_{U'} = A_j$.

Then $N := \bigcup_{U' \subset U} N_{U'}$ is a G -submodule in V intersecting V_0 trivially. (The action of $\sigma \in G$ on N_L for some finite extension L of F^U is given by the composition of the embedding $N_L \subseteq N_{L\sigma(L)}$ with $h_{L\sigma(L)} * \sigma * h_{L\sigma(L)} \in \text{End} N_{L\sigma(L)}$. Here h_L is the identity in \mathcal{H}_L , i.e., the Haar measure on U_L , and we have used $\sigma : W^{U_L} \rightarrow W^{U_{\sigma(L)}} \subseteq W^{U_{L\sigma(L)}}$.) By Proposition 5.1(1), the G -module N is irreducible. This contradicts the definition of W_0 . \square

Let $A^q(Y)$ be the \mathbb{Q} -space $Z^q(Y)$ of cycles on a smooth proper variety Y over k of codimension q modulo an adequate equivalence relation \sim .

Let X be an inverse limit of varieties of a given dimension over k : $X = \varprojlim_{j \in I} X_j$ for a set of indices I . Let $\alpha_j : X \rightarrow X_j$ be the canonical projections. Suppose that each X_j admits an open embedding into a smooth irreducible proper scheme $\overline{X_j}$ over k .

Let $A^q(X)$ be the cokernel of the map $\bigoplus_{j \in I} Z^q(\overline{X_j}) \xrightarrow{\sum_j \alpha_j^*} Z^q(X)$. This is independent of the choice of varieties $\overline{X_j}$. Set $B^q(X) = A^q(X)$ for \sim =numerical equivalence.

For a commutative k -algebra R we set $A^q(R) := A^q(\mathbf{Spec}(R))$.

Proposition 5.3. *The G -module $B_X^q = B^q(X \times_k F)$ is admissible and semi-simple for any smooth proper k -variety X and any $q \geq 0$. If $q \in \{0, 1, \dim X\}$ then B_X^q is of finite length.*

Proof. $B^q(X \times_k F)$ is admissible, since we may suppose that $k \subseteq \mathbb{C}$, and thus, for any smooth proper k -variety Y with $k(Y) \subset F$ the space of $U_{k(Y)}$ -invariants is a quotient of the finite-dimensional space $Z^q(X \times_k Y) / \sim_{\text{hom}} \subseteq H^{2q}((X \times_k Y)(\mathbb{C}), \mathbb{Q}(q))$.

For each Y with $k(Y) \subset F$ and $\overline{k(Y)} = F$ the kernel of $A^q(X \times_k Y) \rightarrow A^q(X \times_k k(Y))$ is a $A^n(Y \times_k Y)$ -submodule in $A^q(X \times_k Y)$, since for its arbitrary element α and for any element $\beta \in A^n(Y \times_k Y)$ one has $\alpha \circ \beta = \text{pr}_{13*}(\text{pr}_{12}^* \alpha \cdot \text{pr}_{23}^* \beta)$, so the projection to Y of the support of $\alpha \circ \beta$ is contained in $\text{pr}_2((D \times_k Y) \cap \text{supp}(\beta))$ for a divisor D on Y , which is of dimension $n - 1$, so cannot dominate Y . This implies that $A^q(X \times_k k(Y))$ has a natural structure of a $A^n(Y \times_k Y)$ -module.

By [J], the algebra $B^n(Y \times_k Y)$ is semi-simple, so the $B^n(Y \times_k Y)$ -module $B^q(X \times_k k(Y))$ is also semi-simple. Then by Lemma 5.2 the G -module $B^q(X \times_k F)$ is semi-simple.

As $B_X^{\dim X}$ is semi-simple, its length is does not exceed $\dim_{\mathbb{Q}} \text{End}_G(B_X^{\dim X})$. It follows from Proposition 5.4 that the latter space is finite-dimensional, and thus, $B_X^{\dim X}$ is of finite length. B_X^1 is a subquotient of B_C^1 for a curve C , so it is also of finite length. \square

Proposition 5.4. *Let Y be a smooth irreducible proper variety over k and $\dim Y \leq n$. Let X be a smooth variety over k . For each $q \geq 0$ there are canonical isomorphisms*

$$A^q(X_{k(Y)}) \xrightarrow{\sim} \text{Hom}_G(A^{\dim Y}(Y_F), A^q(X_F)) \xrightarrow{\sim} \text{Hom}_G(Z^{\dim Y}(k(Y) \otimes_k F), A^q(X_F)).$$

For each pair of reduced irreducible group schemes \mathcal{A} and \mathcal{B} over k there is a natural bijection $\text{Hom}_{\text{group schemes}/k}(\mathcal{A}, \mathcal{B}) \xrightarrow{\sim} \text{Hom}_G(\mathcal{A}(F), \mathcal{B}(F))$.

Proof. For each G -module M and each embedding $\sigma : k(Y) \hookrightarrow F$ one has an isomorphism $M^{U_{\sigma(k(Y))}} \longrightarrow \text{Hom}_G(Z^{\dim Y}(k(Y) \otimes_k F), M)$ given by $m \mapsto [\tau\sigma \mapsto \tau m]$.

Let $\varphi \in \text{Hom}_G(Z^{\dim Y}(k(Y) \otimes_k F), A^q(X_F))$. Fix an embedding $\sigma : k(Y) \hookrightarrow F$. For any $\xi \in G$ one has $\varphi(\xi\sigma) = \xi\varphi(\sigma)$, in particular, if $\xi|_{\sigma(k(Y))} = \text{id}$ then $\varphi(\sigma) \in A^q(X_F)^{U_{\sigma(k(Y))}} = A^q(X_{\sigma(k(Y))})$. From this and the fact that the pairing $A^q(X \times_k Y) \otimes A^{\dim Y}(Y_F) \longrightarrow A^q(X_F)$ factors through $A^q(X_{k(Y)}) \otimes A^{\dim Y}(Y_F) \longrightarrow A^q(X_F)$, one deduces that there is a canonical homomorphism $A^q(X_{k(Y)}) \longrightarrow \text{Hom}_G(A^{\dim Y}(Y_F), A^q(X_F))$ sending a cycle α to the action of its arbitrary lifting to a correspondence on $X \times_k Y$, and it is surjective. On the other hand, it has the inverse given by $\varphi \mapsto \sigma_*^{-1}\varphi(\sigma)$, where $\sigma_* : A^q(X_{k(Y)}) \xrightarrow{\sim} A^q(X_{\sigma(k(Y))})$.

The second map is clearly injective. Let $\varphi \in \text{Hom}_G(\mathcal{A}(F), \mathcal{B}(F))$. Fix an absolutely irreducible curve $C \subset \mathcal{A}$ over k generating \mathcal{A} as an algebraic group, i.e., with the dominant multiplication map $C^N \dashrightarrow \mathcal{A}$ for any $N \geq \dim \mathcal{A}$. Fix a generic point $x \in C(F)$. Since $\varphi(\tau x) = \tau\varphi(x)$, the element $\varphi(x)$ is fixed by any element τ fixing x , so any coordinate of $\varphi(x)$ is a rational function over k in coordinates of x , and therefore, this gives rise to a rational k -map $h : C \dashrightarrow \mathcal{B}$. Consider the rational k -map $C^N \dashrightarrow \mathcal{B}$ given by $(x_1, \dots, x_N) \mapsto h(x_1) \cdots h(x_N)$. On the set of F -points out of the union of “vertical” divisors, (i.e., on $\mathbf{Max}(\underbrace{k(C) \otimes_k \cdots \otimes_k k(C)}_{N \text{ copies}} \otimes_k F)$) this map coincides with one given by $(x_1, \dots, x_N) \mapsto$

$$\varphi(x_1) \cdots \varphi(x_N) = \varphi(x_1 \cdots x_N).$$

As the multiplication map $\mathbf{Max}(\underbrace{k(C) \otimes_k \cdots \otimes_k k(C)}_{N \text{ copies}} \otimes_k F) \longrightarrow \mathcal{A}(F)$ is surjective for any

integer $N \geq 2 \dim \mathcal{A}$, the rational map $C^N \dashrightarrow \mathcal{B}$ factors through $C^N \dashrightarrow \mathcal{A} \xrightarrow{\hat{h}} \mathcal{B}$. Since φ is a homomorphism, \hat{h} should also be a homomorphism, and in particular, regular. \square

Corollary 5.5. *For any pair (L, L') of fields finitely generated over k with $\dim_{L'} \Omega_{L'/k}^1 = m \leq n$ there is a canonical isomorphism*

$$A^q(L \otimes_k L') \xrightarrow{\sim} \text{Hom}_G(A^m(L' \otimes_k F), A^q(L \otimes_k F)). \quad \square$$

Corollary 5.6. *For any smooth irreducible variety X of dimension $\leq n + 1$ over k and any $q \in \{0, 1, 2, \dim X\}$ there is a unique G -submodule in $B_X^q = B^q(X \times_k F)$ isomorphic to $B^q(k(X) \otimes_k F)$.*

Proof. For any $q \geq 0$ and any adequate relation \sim one has the short exact sequence $\bigoplus_{D \in X^1} A^{q-1}(\tilde{D}_F) \longrightarrow A^q(X_F) \longrightarrow A^q(k(X) \otimes_k F) \longrightarrow 0$. As B_X^q is semi-simple, there is a G -submodule in B_X^q isomorphic to $B^q(k(X) \otimes_k F)$.

Such a submodule is unique, since, by Proposition 5.4, $\text{Hom}_G(A^{q-1}(\tilde{D}_F), A^q(k(X) \otimes_k F)) \xrightarrow{\sim} A^q(k(X) \otimes_k k(D)) = 0$ for $q = \dim X$; $\text{Hom}_G(A^{q-1}(\tilde{D}_F), A^q(k(X) \otimes_k F)) = 0$ for $q \in \{0, 1\}$; and $A^1(\tilde{D}_F)$ is isomorphic to a subquotient of $A^1(C_F) \oplus \mathbb{Q}^N$ for a smooth proper curve C , so $\text{Hom}_G(A^1(\tilde{D}_F), A^2(k(X) \otimes_k F)) \subseteq A^2(k(X) \otimes_k k(C)) = 0$. \square

For a G -module M define $N_j M$ as the subgroup generated by invariants $M^{G_{F/F_j}}$ for all subfields $F_j \subseteq F$ of transcendence degree j over k . Clearly, $N_j M$ is the G -submodule of M generated by $M^{G_{F/F_j}}$ for some algebraically closed F_j , $N_j M \subseteq N_{j+1} M$, and restriction to $N_j M_1$ of each G -homomorphism $M_1 \longrightarrow M_2$ factors through $N_j M_2$.

EXAMPLE. For any field L finitely generated over k one has $N_{q-1}A^q(L \otimes_k F) = 0$. If L is of transcendence degree q over k , or $q = 0$, or $q = 1$ then $N_q A^q(L \otimes_k F) = A^q(L \otimes_k F)$.

This implies that for any smooth irreducible variety Y over k one has $\text{Hom}_G(A^r(k(Y) \otimes_k F), A^q(k(X) \otimes_k F)) = 0$ if $r < q$ and either $r \geq \dim Y$, or $r \in \{0, 1\}$. This is because the restriction to $N_r A^r(k(Y) \otimes_k F) = A^r(k(Y) \otimes_k F)$ of each G -homomorphism $A^r(k(Y) \otimes_k F) \rightarrow A^q(k(X) \otimes_k F)$ factors through $N_r A^q(k(X) \otimes_k F) = 0$.

Lemma 5.7. *For any pair of k -varieties X, Y of dimension n , a k -variety Z of dimension $\leq n$ and integers $p, q \geq 0$ one has $\text{Hom}_{\mathcal{H}_U}(B^q(Z \times_k Y), B^p(k(X) \otimes_k k(Y))) = 0$, where either $q = \dim Z < p$, or $q = n$ and $\dim Z < p$, or $q > n$ and $p + q > \dim Z + n$, or $q < p$ and $q \in \{0, 1\}$.*

Proof.

- For each open compact subgroup $U \subset G$ and a smooth irreducible variety Y over k with $k(Y) = F^U$ we define a semi-simple G -module (of finite length) $B_{Z,Y}^q$ as the minimal one such that the \mathcal{H}_U -module $(B_{Z,Y}^q)^U$ is isomorphic to $B^q(Z \times_k Y)$. By Proposition 5.1, it exists and it is unique. There is a canonical morphism $B_{Z,Y}^q \rightarrow B_Z^q$. It is surjective if and only if B_Z^q is of finite type and U fixes a set of its generators.

Let $\dim Z = d < n$. Note, that $B^d(Z \times_k Y) = (B_{Z,Y}^d)^U$ and $W_1 = B_{Z,Y}^d$ is a quotient of the module $Z^d(k(Z) \otimes_k F) = N_d Z^d(k(Z) \otimes_k F)$ (since $W_1^U = W^U$ for $W = Z^d(k(Z) \otimes_k F) / \{G\text{-envelope of } \ker[Z^d(k(Z) \otimes_k k(Y)) \rightarrow B^d(Z \times_k Y)]\}$).

As $\text{Hom}_G(Z^d(k(Z) \otimes_k F), B^n(k(X) \otimes_k F)) = 0$, this gives $\text{Hom}_G(W_1, B^n(k(X) \otimes_k F)) = 0$. We need to show that $\text{Hom}_{\mathcal{H}_U}(W^U, B^n(k(X) \otimes_k k(Y))) = 0$.

By Proposition 5.1, for any pair (W_1, W_2) of semi-simple G -modules the natural homomorphism $\text{Hom}_G(W_1, W_2) \rightarrow \text{Hom}_{\mathcal{H}_U}(W_1^U, W_2^U)$ is surjective. Then, as its source is zero when $W_2 = B^n(k(X) \otimes_k F)$, we get that $\text{Hom}_{\mathcal{H}_U}((B_{Z,Y}^d)^U, B^n(k(X) \otimes_k k(Y))) = 0$.

- Let $c = n - \dim Z$, so the variety $Z \times \mathbb{P}^c$ is n -dimensional. As there is an embedding of \mathcal{H}_U -modules $B^n(Z \times_k Y) \hookrightarrow B^n(Z \times_k \mathbb{P}^c \times_k Y)$, it is enough to show the vanishing of $\text{Hom}_{\mathcal{H}_U}(B^n(Z \times \mathbb{P}^c \times_k Y), B^p(k(X) \otimes_k k(Y)))$ for $p > \dim Z$.

By semi-simplicity, the latter is a subgroup in $\text{Hom}_G(B_{Z \times \mathbb{P}^c, Y}^n, B^p(k(X) \otimes_k F)) \subseteq \text{Hom}_G(Z^n(k(Z \times \mathbb{P}^c) \otimes_k F), B^p(k(X) \otimes_k F))$. By Proposition 5.4, the latter coincides with $B^p(k(X) \otimes_k k(Z \times \mathbb{P}^c))$, which is dominated by $B^p(\mathbb{A}_{k(X) \otimes_k k(Z)}^c) \xrightarrow{\sim} B^p(k(X) \otimes_k k(Z)) = 0$, and thus, $\text{Hom}_{\mathcal{H}_U}(B^n(Z \times_k Y), B^p(k(X) \otimes_k k(Y))) = 0$, if $p > \dim Z$.

- As the \mathcal{H}_U -module $\bigoplus_{D \in Z^{q-n}} B^n(\tilde{D} \times_k Y)$ surjects onto \mathcal{H}_U -module $B^q(Z \times_k Y)$ when $q > n$, the vanishing of $\text{Hom}_{\mathcal{H}_U}(B^q(Z \times_k Y), B^p(k(X) \otimes_k k(Y)))$ follows from the vanishing of the group $\text{Hom}_{\mathcal{H}_U}(B^n(\tilde{D} \times_k Y), B^p(k(X) \otimes_k k(Y)))$ for each subvariety D of codimension $q - n$ on Z .
- If $q = 1$ then $B^1(Z \times_k Y)$ is a subquotient of $B^1(C \times_k Y)$ for a smooth curve C , so we are reduced to the case $q = \dim Z < p$. The case $q = 0$ is trivial. \square

Proposition 5.8. *Let X and Y be smooth irreducible varieties over k , and either $q \in \{0, 1, 2\}$, or $q = \dim X = \dim Y$. Then there is a unique $B^{\dim X}(X \times_k X) \times B^{\dim Y}(Y \times_k Y)$ -submodule in $B^q(X \times_k Y)$ isomorphic to its quotient $B^q(k(X) \otimes_k k(Y))$.*

Proof. The existence of such submodule follows from the semi-simplicity ([J]). As the kernel of the projection $B^q(X \times_k Y) \rightarrow B^q(k(X) \otimes_k k(Y))$ is generated by the images of $B^{q-1}(X \times_k \tilde{E})$ and $B^{q-1}(\tilde{D} \times_k Y)$ for all divisors D on X and all divisors E on Y , the

uniqueness would follow from $\text{Hom}_{B^{\dim Y}(Y \times_k Y)}(B^{q-1}(\tilde{D} \times_k Y), B^q(k(X) \otimes_k k(Y))) = 0$. But this is Lemma 5.7. \square

Proposition 5.9. *For any irreducible k -variety X of dimension n there is a non-degenerate G -equivariant pairing $W \otimes W \xrightarrow{\langle \cdot, \cdot \rangle} \mathbb{Q}(\chi)$, where $W = B^n(k(X) \otimes_k F) \neq 0$.*

If for $(n-1)$ -cycles on $2n$ -dimensional complex varieties the numerical equivalence coincides with the homological one, then $\langle \cdot, \cdot \rangle$ is $(-1)^n$ -definite. In particular, this holds for $n \leq 2$.

Proof. For a pair $\beta_1, \beta_2 \in W$ fixed by a compact open subgroup $U \subset G$ we define $\langle \alpha, \beta \rangle \in \mathbb{Q}(\chi)$ by $\langle \hat{\beta}_1 \cdot \hat{\beta}_2 \rangle \cdot [U]$, where $\hat{\beta}_1, \hat{\beta}_2$ are the images of $\beta_1, \beta_2 \in B^n(k(X) \otimes_k k(Y))$ in $B^n(X \times_k Y)$ in the sense of Proposition 5.8. Here Y is a smooth proper variety over k with the function field $k(Y)$ identified with F^U .

For a triple of smooth proper varieties X_1, X_2, X_3 , a triple of integers $a, b, c \geq 0$ with $a+b+c = \dim(X_1 \times X_2 \times X_3)$ and a triple $\alpha \in A^a(X_1 \times X_2), \beta \in A^b(X_2 \times X_3), \gamma \in A^c(X_1 \times X_3)$ one has $\langle \alpha \circ \beta, \gamma \rangle = \langle \beta, {}^t\alpha \circ \gamma \rangle = \langle \alpha, \gamma \circ {}^t\beta \rangle$. For any $\alpha \in W - \{0\}$ fixed by U there is $\beta \in B^n(X \times_k Y)$ such that $\langle \hat{\alpha}, \beta \rangle \neq 0$. Then, as $\hat{\alpha} = \Delta_{k(X)} \circ \hat{\alpha} \circ \Delta_{k(Y)}$, where $\Delta_{k(X)}$ is the identity element in $B^n(k(X) \otimes_k k(X))$ considered as an element of $B^n(X \times_k X)$, and similarly for $\Delta_{k(Y)}$, one has $\langle \hat{\alpha}, \beta \rangle = \langle \hat{\alpha}, \Delta_{k(X)} \circ \beta \circ \Delta_{k(Y)} \rangle \neq 0$. As $\Delta_{k(X)} \circ \beta \circ \Delta_{k(Y)} \in W$, this shows that $\langle \cdot, \cdot \rangle$ is non-degenerate.

By Lemma 5.7, $\text{Hom}_{\mathcal{H}_U}(B^{n+1}(X \times_k Y), B^n(k(X) \otimes_k k(Y))) = 0$, and therefore, for any $L \in \text{NS}(X)$ one has $\text{Hom}_{\mathcal{H}_U}(L \cdot B^n(X \times_k Y), B^n(k(X) \otimes_k k(Y))) = 0$. By the semi-simplicity, this means that the composition $B^n(k(X) \otimes_k k(Y)) \hookrightarrow B^n(X \times_k Y) \xrightarrow{[L]} B^{n+1}(X \times_k Y)$ is zero for any $L \in \text{NS}(X)$. Interchanging the roles of X and Y , we see that the image of $B^n(k(X) \otimes_k k(Y))$ in $B^n(X \times_k Y)$ is annihilated by any $L \in \text{NS}(X) \oplus \text{NS}(Y) \subseteq \text{NS}(X \times_k Y)$, i.e., it consists of primitive elements. Then, by the Hodge index theorem, if for $(n-1)$ -cycles on $2n$ -dimensional complex varieties the numerical equivalence coincides with the homological one, then the pairing $W \otimes W \rightarrow \mathbb{Q}(\chi)$ is $(-1)^n$ -definite. \square

REFERENCES

- [BZ] I.N.Bernstein, A.V.Zelevinsky, *Representations of the group $GL(n, F)$, where F is a local non-archimedean field*, Uspehi Matem.Nauk **31** (1976), no.3 (189), 5–70.
- [J] U.Jannsen, *Motives, numerical equivalence, and semi-simplicity*, Invent.Math. **107** (1992), no.3, 447–452.
- [III-III] I.I.Piatetskii-Shapiro, I.R.Shafarevich, *Galois theory of transcendental extensions and uniformization*, Izv.AN SSSR, ser. matem. **30** (1966), 671–704.