

Little q -Legendre polynomials and irrationality of certain Lambert series

Walter Van Assche *

Katholieke Universiteit Leuven and Georgia Institute of Technology

October 26, 2018

Abstract

Certain q -analogs $h_p(1)$ of the harmonic series, with $p = 1/q$ an integer greater than one, were shown to be irrational by Erdős [9]. In 1991–1992 Peter Borwein [4] [5] used Padé approximation and complex analysis to prove the irrationality of these q -harmonic series and of q -analogs $\ln_p(2)$ of the natural logarithm of 2. Recently Amdeberhan and Zeilberger [1] used the qEKHAD symbolic package to find q -WZ pairs that provide a proof of irrationality similar to Apéry's proof of irrationality of $\zeta(2)$ and $\zeta(3)$. They also obtain an upper bound for the measure of irrationality, but better upper bounds were earlier given by Bundschuh and Väänänen [8] and recently also by Matala-aho and Väänänen [14] (for $\ln_p(2)$). In this paper we show how one can obtain rational approximants for $h_p(1)$ and $\ln_p(2)$ (and many other similar quantities) by Padé approximation using little q -Legendre polynomials and we show that properties of these orthogonal polynomials indeed prove the irrationality, with an upper bound of the measure of irrationality which is as sharp as the upper bound given by Bundschuh and Väänänen for $h_p(1)$ and a better upper bound as the one given by Matala-aho and Väänänen for $\ln_p(2)$.

1 Introduction

Most important special functions, in particular hypergeometric functions, have q -extensions, usually obtained by replacing Pochhammer symbols $(a)_n = a(a+1)\cdots(a+n-1)$ by their q -analog $(a; q)_n = (1-a)(1-aq)(1-aq^2)\cdots(1-aq^{n-1})$. Since

$$\lim_{q \rightarrow 1} \frac{(q^a; q)_n}{(1-q)^n} = (a)_n,$$

one usually retrieves the original special function from its q -extension by letting $q \rightarrow 1$. A good source for q -extensions of hypergeometric series (basic hypergeometric series) is the book [10] by Gasper and Rahman. Our interest in this paper is the following q -extension of the harmonic series

$$h_p(1) = \sum_{k=1}^{\infty} \frac{1}{p^k - 1} = \sum_{k=1}^{\infty} \frac{q^k}{1 - q^k}, \quad 0 < q = \frac{1}{p} < 1, \quad (1)$$

*WVA is a Research Director of the Belgian Fund for Scientific Research (FWO).

and of the natural logarithm of 2

$$\ln_p(2) = \sum_{k=1}^{\infty} \frac{(-1)^k}{p^k - 1} = \sum_{k=1}^{\infty} \frac{(-q)^k}{1 - q^k}, \quad 0 < q = \frac{1}{p} < 1. \quad (2)$$

In 1948, Paul Erdős [9] proved that $h_2(1)$ is irrational. Peter Borwein [4] [5] showed that $h_p(1)$ (and other similar numbers) are irrational for every integer $p > 1$ and also proved the irrationality of $\ln_p(2)$ for every integer $p > 1$. If we denote $E_p(z) = (qz; q)_{\infty}$, with $q = 1/p$, then Bézivin [3] had earlier shown that $E_p(\alpha), E'_p(\alpha), \dots, E_p^{(k)}(\alpha)$ are linearly independent over \mathbb{Q} for every $k \in \mathbb{N}$ and $\alpha \in \mathbb{Q}$ with $\alpha p^j \neq -1$ for every integer j . The case $k = 1$ corresponds to irrationality of

$$\sum_{j=1}^{\infty} \frac{1}{\alpha + p^j}.$$

Borwein used Padé approximation techniques and complex analysis to obtain good rational approximants to $h_p(1)$ and $\ln_p(2)$. Indeed, one can use the following lemma to prove irrationality [16, Lemma 5.1]:

Lemma 1 *Let x be a real number, and suppose there exist integers a_n, b_n ($n \in \mathbb{N}$) such that*

1. $x \neq a_n/b_n$ for every $n \in \mathbb{N}$,
2. $\lim_{n \rightarrow \infty} (b_n x - a_n) = 0$,

then x will be irrational.

This lemma expresses the fact that the order of approximation of a rational number by rational numbers is one and not higher [12, Theorem 186]. Furthermore, if $|x - a_n/b_n| = \mathcal{O}(1/b_n^{1+s})$ with $0 < s < 1$ and $b_n < b_{n+1} < b_n^{1+o(1)}$, then the measure of irrationality $r(x)$ (Liouville-Roth number, order of approximation)

$$r(x) = \inf\{r \in \mathbb{R} : |x - a/b| < 1/b^r \text{ has at most finitely many integer solutions } (a, b)\}$$

satisfies $2 \leq r(x) \leq 1 + 1/s$ (see, e.g., [6, exercise 3 on p. 376] for the upper bound; the lower bound follows since every irrational number is approximable to order 2 [12, Theorem 187]). Recently Amdeberhan and Zeilberger [1] found q -WZ pairs to obtain rational approximants for $h_p(1)$ and $\ln_p(2)$, improving the upper bound for the measure of irrationality to $4.8 = 24/5$. However, four years earlier Bundschuh and Väänänen [8, p. 178] had established better upper bounds: $r(h_p(1)) \leq 1 + (\pi^2 + 2)/(\pi^2 - 2) = 2.50828 \dots$ and $r(\ln_p(2)) \leq 1 + (2\pi^2 + 3)/(\pi^2 - 3) = 4.310119 \dots$. The upper bound for $r(\ln_p(2))$ was improved by Matala-aho and Väänänen [14] to 3.9461.

In this paper we show that one can find rational approximants which are related to little q -Legendre polynomials and hence return to Padé approximation. We can then use some results for little q -Legendre polynomials to prove the irrationality once more, and with the aid of some elementary number theory we obtain the same bound for the measure of irrationality as the one obtained by Bundschuh and Väänänen for $h_p(1)$ and a better upper bound as the one given by Matala-aho and Väänänen for $\ln_p(2)$. The connection with little q -Legendre polynomials opens the way for proving the irrationality of q -extensions of $\zeta(2)$ and $\zeta(3)$ in Apéry's spirit [2], using multiple orthogonal q -polynomials [16].

2 Little q -Legendre polynomials

The little q -Legendre polynomials are defined by

$$\begin{aligned} P_n(x|q) &= {}_2\phi_1 \left(\begin{matrix} q^{-n}, q^{n+1} \\ q \end{matrix} \middle| q; qx \right) \\ &= \sum_{k=0}^n \frac{(q^{-n}; q)_k (q^{n+1}; q)_k}{(q; q)_k} \frac{q^k x^k}{(q; q)_k}, \quad 0 < q < 1, \end{aligned} \quad (3)$$

and they are orthogonal polynomials on the exponential lattice $\{q^k, k = 0, 1, 2, \dots\}$:

$$\sum_{k=0}^{\infty} q^k P_m(q^k|q) P_n(q^k|q) = \frac{q^n}{1 - q^{2n+1}} \delta_{n,m}. \quad (4)$$

If we use the q -binomial coefficients

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}, \quad (5)$$

and the formulas

$$(q^{-n}; q)_k = (-1)^k q^{-nk+k(k-1)/2} \frac{(q; q)_n}{(q; q)_{n-k}}, \quad (q^{n+1}; q)_k = \frac{(q; q)_{n+k}}{(q; q)_n}, \quad (6)$$

then (3) reduces to

$$P_n(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \begin{bmatrix} n+k \\ k \end{bmatrix}_q q^{-nk+k(k+1)/2} (-x)^k, \quad (7)$$

and since

$$\lim_{q \uparrow 1} \begin{bmatrix} n \\ k \end{bmatrix}_q = \binom{n}{k},$$

we see that we indeed find the Legendre polynomials on $[0, 1]$ by letting q tend to 1

$$\lim_{q \uparrow 1} P_n(x|q) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} (-x)^k = P_n(x).$$

Let $p = 1/q$ so that $p > 1$ whenever $0 < q < 1$, then

$$(q; q)_k = (-1)^k p^{-k(k+1)/2} (p; p)_k, \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = p^{-k(n-k)} \begin{bmatrix} n \\ k \end{bmatrix}_p, \quad (8)$$

and we can rewrite the little q -Legendre polynomial as

$$P_n(x|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p p^{-kn+k(k-1)/2} (-x)^k. \quad (9)$$

There is a Rodrigues formula for these little q -Legendre polynomials in terms of the q -difference operator D_q for which

$$D_q f(z) = \begin{cases} \frac{f(z) - f(qz)}{(1-q)z} & \text{if } z \neq 0, \\ f'(0) & \text{if } z = 0, \end{cases} \quad (10)$$

namely

$$P_n(x|q) = \frac{q^{n(n-1)/2}(1-q)^n}{(q; q)_n} D_p^n[(qx; q)_n x^n], \quad (11)$$

which will be useful later. We refer to [13] for more information and references for little q -Legendre polynomials.

Equation (9) expresses the little q -Legendre polynomials in the basis $\{1, x, x^2, \dots, x^n\}$ of monomials. Sometimes it is more convenient to use another basis of polynomials, and for orthogonal polynomials on $\{q^k, k = 0, 1, 2, \dots\}$ a convenient set of basis functions is $\{(qx; q)_k, k = 0, 1, 2, \dots, n\}$. We will need to use some q -series for this purpose. Recall the q -analog of Newton's binomium formula

$$(x; q)_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} (-x)^k, \quad (12)$$

and its dual

$$x^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-1)^k q^{-nk+k(k+1)/2} (x; q)_k. \quad (13)$$

A more general result is the q -binomial series

$$\sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} x^n = \frac{(ax; q)_{\infty}}{(x; q)_{\infty}}, \quad |q| < 1, |x| < 1. \quad (14)$$

Use (13) with argument $q^{n+1}x$ in the Rodrigues formula (11), then we have

$$P_n(x|q) = \frac{p^{n(n+1)}(1-p)^n}{(p; p)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-1)^k q^{-nk+k(k+1)/2} D_p^n(qx; q)_{n+k}.$$

One finds easily that

$$D_p^k(qx; q)_n = \frac{(q; q)_n}{(q; q)_{n-k}(1-p)^k} (qx; q)_{n-k}, \quad (15)$$

so that, using (8) we find the required expansion

$$P_n(x|q) = (-1)^n \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} (qx; q)_k. \quad (16)$$

3 The q -harmonic series

Orthogonal polynomials naturally arise in Padé approximation of a Stieltjes function

$$f(z) = \int \frac{d\mu(x)}{z-x}, \quad z \notin \text{supp}(\mu). \quad (17)$$

Suppose μ is a positive measure on the real line with infinite support and for which all the moments exist. If $P_n(x)$ ($n = 0, 1, 2, \dots$) are orthogonal polynomials for μ , i.e., P_n is of degree n and

$$\int P_n(x)P_m(x) d\mu(x) = 0, \quad m \neq n,$$

and if Q_n are the polynomials of degree $n-1$ given by

$$Q_n(z) = \int \frac{P_n(z) - P_n(x)}{z-x} d\mu(x), \quad (18)$$

then it is easy to see that

$$P_n(z)f(z) - Q_n(z) = \int \frac{P_n(x)}{z-x} d\mu(x), \quad z \notin \text{supp}(\mu). \quad (19)$$

If we expand $1/(z-x)$ around $z = \infty$ as

$$\frac{1}{z-x} = \sum_{k=0}^{n-1} \frac{x^k}{z^{k+1}} + \frac{x^n}{z^n} \frac{1}{z-x},$$

then by orthogonality we have

$$P_n(z)f(z) - Q_n(z) = \frac{1}{z^n} \int \frac{P_n(x)x^n}{z-x} d\mu(x) = \mathcal{O}(1/z^{n+1}).$$

This is precisely the (linearized) form of the interpolation conditions near $z = \infty$ for Padé approximation so that $Q_n(z)/P_n(z)$ is the $[\frac{n-1}{n}]$ Padé approximant for $f(z)$ near $z = \infty$.

For little q -Legendre polynomials the measure μ is supported on $\{q^k, k = 0, 1, 2, \dots\}$, which is a bounded set in $[0, 1]$ with one accumulation point at 0. The measure is given by

$$\int g(x) d\mu(x) = \sum_{k=0}^{\infty} g(q^k)q^k.$$

The Stieltjes function for this measure is

$$f(z) = \sum_{k=0}^{\infty} \frac{q^k}{z - q^k} = \sum_{k=0}^{\infty} \frac{1}{zp^k - 1}. \quad (20)$$

We will need this function at p^n , where it gives

$$f(p^n) = \sum_{k=0}^{\infty} \frac{1}{p^{n+k} - 1} = h_p(1) - \sum_{k=1}^{n-1} \frac{1}{p^k - 1}. \quad (21)$$

Hence if $p > 1$ is an integer, then $f(p^n)$ gives $h_p(1)$ up to $\sum_{k=1}^{n-1} 1/(p^k - 1)$, which is a rational number. Now use (19) for little q -Legendre polynomials at $z = p^n$ to find

$$P_n(p^n|q) \left(h_p(1) - \sum_{k=1}^{n-1} \frac{1}{p^k - 1} \right) - Q_n(p^n|q) = \sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{p^n - q^k} q^k. \quad (22)$$

Observe that (9) gives

$$P_n(p^n|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ n \end{bmatrix}_p (-1)^k p^{k(k-1)/2}, \quad (23)$$

which is nearly the b_n found in [1, p. 277] (their b_n corresponds to $P_n(p^{n+1}|q)$). Observe also that Borwein's construction [5, Lemma 2] uses $P_{n-1}(cp^{n+1}|q)$. The p -binomial numbers $\begin{bmatrix} n \\ k \end{bmatrix}_p$ are polynomials in p with integer coefficients, which follows easily from the q -version of Pascal's triangle identities

$$\begin{bmatrix} n \\ k \end{bmatrix}_p = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_p + p^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_p = \begin{bmatrix} n-1 \\ k \end{bmatrix}_p + p^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_p,$$

hence if $p > 1$ is an integer then $\begin{bmatrix} n \\ k \end{bmatrix}_p$ and $\begin{bmatrix} n+k \\ k \end{bmatrix}_p$ are integers. This means that (23) implies $P_n(p^n|q)$ to be an integer. Furthermore, since $p^n > 1$ and all the zeros of $P_n(x|q)$ are in $[0, 1]$, we also may conclude that $(-1)^n P_n(p^n|q)$ is positive for all n .

The second important quantity in (22) is $Q_n(p^n|q)$. This associated little q -Legendre polynomial can be computed explicitly using (18) and is given by

$$Q_n(x|q) = \sum_{j=0}^{\infty} \frac{P_n(x|q) - P_n(q^j|q)}{x - q^j} q^j.$$

Use (16) to write this as

$$Q_n(x|q) = (-1)^n \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \sum_{j=0}^{\infty} \frac{(qx; q)_k - (q^{j+1}; q)_k}{x - q^j} q^j.$$

Now use

$$\frac{(qx; q)_k - (qy; q)_k}{x - y} = - \sum_{\ell=1}^k q^\ell (qy; q)_{\ell-1} (q^{\ell+1}x; q)_{k-\ell},$$

which one can easily prove by induction, then this gives

$$\begin{aligned} Q_n(x|q) &= (-1)^{n+1} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \\ &\quad \times \sum_{\ell=1}^k q^\ell (q^{\ell+1}x; q)_{k-\ell} \sum_{j=0}^{\infty} q^j (q^{j+1}; q)_{\ell-1}. \end{aligned}$$

Using the q -binomial series (14) we can calculate the modified moments

$$\begin{aligned} \sum_{j=0}^{\infty} q^j (q^{j+1}; q)_{\ell-1} &= (q; q)_{\ell-1} \sum_{j=0}^{\infty} q^j \frac{(q^{\ell}; q)_j}{(q; q)_j} \\ &= \frac{(q; q)_{\ell-1} (q^{\ell+1}; q)_{\infty}}{(q; q)_{\infty}} \\ &= \frac{1}{1 - q^{\ell}}, \end{aligned}$$

so that

$$Q_n(x|q) = (-1)^{n+1} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \sum_{\ell=1}^k \frac{(q^{\ell+1}x; q)_{k-\ell}}{p^{\ell} - 1}. \quad (24)$$

Evaluating at $x = p^n$, and using

$$(q^{\ell+1}p^n; q)_{k-\ell} = (p^{n-k}; p)_{k-\ell},$$

then gives

$$Q_n(p^n|q) = (-1)^{n+1} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \sum_{\ell=1}^k \frac{(p^{n-k}; p)_{k-\ell}}{p^{\ell} - 1}. \quad (25)$$

All the terms in the sum for $Q_n(p^n|q)$ are now integers, except for the $p^{\ell} - 1$ in the denominators. In order to obtain an integer we therefore need to multiply everything by a multiple of all $p^{\ell} - 1$ for $\ell = 1, 2, \dots, n$. We will choose

$$d_n(p) = \prod_{k=1}^n \Phi_k(p), \quad (26)$$

where

$$\Phi_n(x) = \prod_{\substack{k=1 \\ \gcd(k,n)=1}}^n (x - \omega_n^k), \quad \omega_n = e^{2\pi i/n}, \quad (27)$$

are the cyclotomic polynomials [15, §4.8]. Each cyclotomic polynomial is monic, has integer coefficients, and the degree of Φ_n is $\phi(n)$ (Euler's totient function). It is known that

$$x^n - 1 = \prod_{d|n} \Phi_d(x), \quad (28)$$

and that every cyclotomic polynomial is irreducible over $\mathbb{Q}[x]$. Hence $d_n(p)$ is a multiple of all $p^{\ell} - 1$ for $\ell = 1, 2, \dots, n$. The growth of this sequence is given by the following lemma, which was essentially given by A. O. Gel'fond, who obtained the upper bound in [11, Equation (7)]. We give a proof to make this paper self-contained.

Lemma 2 Suppose p is an integer greater than one and let $d_n(p)$ be given by (26). Then

$$\lim_{n \rightarrow \infty} d_n(p)^{1/n^2} = p^{3/\pi^2}. \quad (29)$$

Proof: The degree of $d_n(p)$ as a monic polynomial in p is $\sum_{k=1}^n \varphi(k)$ and an old result of Mertens (1874) shows that for $n \rightarrow \infty$ this grows like [12, Theorem 330]

$$\sum_{k=1}^n \varphi(k) = \frac{3}{\pi^2} n^2 + \mathcal{O}(n \log n). \quad (30)$$

We will adapt the classical proof of (30) to prove our lemma. For this we use Möbius inversion of (28) to find the representation

$$\Phi_n(x) = \prod_{d|n} (x^d - 1)^{\mu(n/d)},$$

where μ is the Möbius function. Taking logarithms in (26) gives

$$\log d_n(p) = \sum_{k=1}^n \sum_{d|k} \mu(k/d) \log(p^d - 1).$$

Changing the order of summation gives

$$\log d_n(p) = \sum_{\ell=1}^n \mu(\ell) \sum_{d=1}^{\lfloor n/\ell \rfloor} \log(p^d - 1).$$

Now

$$\sum_{d=1}^m \log(p^d - 1) = \log p^{m(m+1)/2} (q; q)_m = \frac{m(m+1)}{2} \log p + \log(q; q)_m,$$

so that

$$\begin{aligned} \log d_n(p) &= \frac{n^2}{2} \log p \sum_{\ell=1}^n \frac{\mu(\ell)}{\ell^2} + \mathcal{O}\left(n \sum_{\ell=1}^n \frac{1}{\ell}\right) \\ &= \frac{n^2 \log p}{2\zeta(2)} + \mathcal{O}(n \log n). \end{aligned}$$

The lemma now follows by using $\zeta(2) = \pi^2/6$. \square

So far we have found that the numbers

$$b_n = d_n(p) P_n(p^n|q), \quad (31)$$

$$a_n = d_n(p) Q_n(p^n|q) + b_n \sum_{k=1}^{n-1} \frac{1}{p^k - 1}. \quad (32)$$

are integers and

$$b_n h_p(1) - a_n = d_n(p) \sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{p^n - q^k} q^k. \quad (33)$$

We now want to show that $b_n h_p(1) - a_n \neq 0$ for all n and

$$\lim_{n \rightarrow \infty} (b_n h_p(1) - a_n) = 0,$$

so that Lemma 1 implies the irrationality of $h_p(1)$. First observe that

$$\sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{p^n - q^k} q^k = \frac{1}{P_n(p^n|q)} \sum_{k=0}^{\infty} \frac{P_n(q^k|q) P_n(p^n|q)}{p^n - q^k} q^k.$$

If we add and subtract $P_n(q^k|q)$ in the sum, then we have

$$\sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{p^n - q^k} q^k = \frac{1}{P_n(p^n|q)} \sum_{k=0}^{\infty} P_n(q^k|q) \frac{P_n(p^n|q) - P_n(q^k|q)}{p^n - q^k} q^k + \frac{1}{P_n(p^n|q)} \sum_{k=0}^{\infty} \frac{P_n^2(q^k|q)}{p^n - q^k} q^k.$$

The first sum on the right hand side vanishes because of the orthogonality, so that

$$b_n h_p(1) - a_n = \frac{d_n(p)}{P_n(p^n|q)} \sum_{k=0}^{\infty} \frac{P_n^2(q^k|q)}{p^n - q^k} q^k. \quad (34)$$

All the terms in the sum are now positive, and $(-1)^n P_n(p^n|q)$ is positive for all n , hence we may conclude that

$$(-1)^n (b_n h_p(1) - a_n) > 0, \quad n = 1, 2, \dots$$

Next we show that this quantity converges to zero. Clearly $p^n - 1 \leq p^n - q^k \leq p^n$ so that

$$\frac{1}{p^n} \sum_{k=0}^{\infty} P_n^2(q^k|q) q^k \leq \sum_{k=0}^{\infty} \frac{P_n^2(q^k|q)}{p^n - q^k} q^k \leq \frac{1}{p^n - 1} \sum_{k=0}^{\infty} P_n^2(q^k|q) q^k.$$

Now we can use the norm of the little q -Legendre polynomial (4) to find

$$\frac{p}{p^{2n+1} - 1} \leq \sum_{k=0}^{\infty} \frac{P_n^2(q^k|q)}{p^n - q^k} q^k \leq \frac{p^{n+1}}{(p^n - 1)(p^{2n+1} - 1)}. \quad (35)$$

What remains is to find the asymptotic behavior of $P_n(p^n|q)$ as $n \rightarrow \infty$. For this we can use a very general theorem for sequences of polynomials with uniformly bounded zeros.

Lemma 3 *Suppose P_n ($n \in \mathbb{N}$) is a sequence of monic polynomials of degree n and that the zeros $x_{j,n}$ ($1 \leq j \leq n$) of P_n are such that $|x_{j,n}| \leq M$, with M independent of n . Then we have for $|x| > 1$ and every $c \in \mathbb{C}$*

$$\lim_{n \rightarrow \infty} |P_n(cx^n)|^{1/n^2} = |x|. \quad (36)$$

Proof: Factoring the polynomial P_n gives

$$|P_n(x)| = \prod_{j=1}^n |x - x_{j,n}|.$$

We have the obvious bounds

$$|x| - M \leq |x - x_{j,n}| \leq |x| + M,$$

hence when $|cx^n| > M$

$$(|cx^n| - M)^n \leq |P_n(cx^n)| \leq (|cx^n| + M)^n.$$

For n large enough this easily gives

$$|x| \left(|c| - \frac{M}{|x|^n} \right)^{1/n} \leq |P_n(cx^n)|^{1/n^2} \leq |x| \left(|c| + \frac{M}{|x|^n} \right)^{1/n},$$

which gives the desired result. \square

Observe that we may allow M to grow with n subexponentially. For little q -Legendre polynomials the zeros are all in $[0, 1]$ so that we can use the Lemma with $M = 1$. The leading coefficient κ_n of $P_n(x|q)$ is, by (9), equal to $\kappa_n = (-1)^n \begin{bmatrix} 2n \\ n \end{bmatrix}_p p^{-n(n+1)/2}$, giving

$$\lim_{n \rightarrow \infty} |\kappa_n|^{1/n^2} = \sqrt{p},$$

and hence Lemma 3 gives for $|x| > 1$ and $c \in \mathbb{C}$

$$\lim_{n \rightarrow \infty} |P_n(cx^n|q)|^{1/n^2} = \sqrt{p} |x|. \quad (37)$$

Theorem 1 *Suppose $p > 1$ is an integer. Let a_n and b_n be given by (31)–(32), then $a_n \in \mathbb{Z}$, $(-1)^n b_n \in \mathbb{N}$, and $(-1)^n (b_n h_p(1) - a_n) > 0$ for $n > 1$. Furthermore*

$$\lim_{n \rightarrow \infty} |b_n h_p(1) - a_n|^{1/n^2} = p^{-\frac{3(\pi^2-2)}{2\pi^2}} < 1,$$

which implies that $h_p(1)$ is irrational with measure of irrationality $r \leq \frac{2\pi^2}{\pi^2-2} = 2.50828 \dots$

Proof: If we take $x = p$ and $c = 1$ in (37) then we have

$$\lim_{n \rightarrow \infty} |P_n(p^n|q)|^{1/n^2} = p^{3/2},$$

and combining with (29) we have for the integer b_n in (31)

$$\lim_{n \rightarrow \infty} |b_n|^{1/n^2} = p^{\frac{3(\pi^2+2)}{2\pi^2}}.$$

Observe that b_n has the same sign as $P_n(p^n|q)$ which is $(-1)^n$. Furthermore (34) and (35) show that

$$\lim_{n \rightarrow \infty} |b_n h_p(1) - a_n|^{1/n^2} = \frac{\lim_{n \rightarrow \infty} d_n(p)^{1/n^2}}{\lim_{n \rightarrow \infty} |P_n(p^n|q)|^{1/n^2}} = p^{-\frac{3(\pi^2-2)}{2\pi^2}} < 1.$$

The irrationality now follows from Lemma 1. Observe that this gives rational approximants a_n/b_n for $h_p(1)$ satisfying

$$|h_p(1) - \frac{a_n}{b_n}| = \mathcal{O} \left(\frac{p^{(\frac{-3(\pi^2-2)}{2\pi^2} + \epsilon)n^2}}{b_n} \right)$$

for every $\epsilon > 0$. Now $b_n = p^{3n^2(\pi^2+2)/(2\pi^2)+o(n^2)}$, hence

$$|h_p(1) - \frac{a_n}{b_n}| = \mathcal{O} \left(\frac{1}{b_n^{1+\frac{\pi^2-2}{\pi^2+2}-\epsilon}} \right)$$

for every $\epsilon > 0$, which gives for the measure of irrationality the bound $r \leq 1 + \frac{\pi^2+2}{\pi^2-2} = \frac{2\pi^2}{\pi^2-2}$.

□

The upper bound for the measure of irrationality is better than the upper bound 4.8 obtained in [1], but the same as the (earlier) upper bound of Bundschuh and Väänänen [8].

4 The q -analog of the logarithm of 2

Next we show that a very similar analysis also proves the irrationality of $\ln_p(2)$ for every integer $p > 1$. First of all we rewrite $\ln_p(2)$ using the geometric series

$$\ln_p(2) = \sum_{k=1}^{\infty} (-1)^k \frac{q^k}{1 - q^k} = \sum_{k=1}^{\infty} (-1)^k q^k \sum_{j=0}^{\infty} q^{jk}.$$

Fubini's theorem allows us to change the order of the sums whenever $0 < q < 1$ and this gives

$$\begin{aligned} \ln_p(2) &= \sum_{j=0}^{\infty} \sum_{k=1}^{\infty} (-1)^k q^{k(j+1)} \\ &= - \sum_{j=0}^{\infty} \frac{q^{j+1}}{1 + q^{j+1}} \\ &= - \sum_{k=1}^{\infty} \frac{1}{p^k + 1}. \end{aligned}$$

Hence if we evaluate the Stieltjes function (20) at $z = -p^n$ then we find

$$f(-p^n) = - \sum_{k=0}^{\infty} \frac{1}{p^{n+k} + 1} = \ln_p(2) + \sum_{k=1}^{n-1} \frac{1}{p^k + 1},$$

so that $f(-p^n)$ gives the required $\ln_p(2)$ up to $\sum_{k=1}^{n-1} 1/(p^k + 1)$, which is a rational number. We can now proceed as in the previous section and evaluate (19) for little q -Legendre polynomials at $z = -p^n$ to find

$$P_n(-p^n|q) \left(\ln_p(2) + \sum_{k=1}^{n-1} \frac{1}{p^k + 1} \right) - Q_n(-p^n|q) = - \sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{p^n + q^k} q^k.$$

Here we can use (9) to see that

$$P_n(-p^n|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p p^{k(k-1)/2},$$

is a positive integer, and if we use (24) and $(-p^n q^{\ell+1}; q)_{k-\ell} = (-p^{n-k}; p)_{k-\ell}$ then

$$Q_n(-p^n|q) = (-1)^{n+1} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \sum_{\ell=1}^k \frac{(-p^{n-k}; p)_{k-\ell}}{p^\ell - 1}.$$

This quantity has the numbers $p^\ell - 1$ ($\ell = 1, 2, \dots, n$) in the denominator, so that we need to multiply it by a multiple of all $p^\ell - 1$ with $1 \leq \ell \leq n$. Now we have an additional quantity

$$\sum_{k=1}^{n-1} \frac{1}{p^k + 1}$$

and in order to make this an integer we need to multiply it by a multiple of all $p^k + 1$ for $k = 1, 2, \dots, n-1$. If we choose

$$\hat{d}_n(p) = \prod_{k=1}^n \Phi_k(p^2),$$

then because of $p^{2k} - 1 = (p^k - 1)(p^k + 1)$ we see that $\hat{d}_n(p)$ is a multiple of all $p^k + 1$ and all $p^k - 1$ for $k = 1, \dots, n$. Note that $\hat{d}_n(p) = d_n(p^2)$, where d_n is given by (26), so that Lemma 2 gives the growth

$$\lim_{n \rightarrow \infty} d_n(p^2)^{1/n^2} = p^{6/\pi^2}. \quad (38)$$

So if we choose

$$b_n = d_n(p^2) P_n(-p^n|q), \quad (39)$$

$$a_n = d_n(p^2) Q_n(-p^n|q) - b_n \sum_{k=1}^{n-1} \frac{1}{p^k + 1}, \quad (40)$$

then a_n and b_n are integers and

$$b_n \ln_p(2) - a_n = d_n(p^2) \sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{-p^n - q^j} q^k. \quad (41)$$

Theorem 2 Suppose $p > 1$ is an integer. Let a_n and b_n be given by (39)–(40), then $a_n \in \mathbb{Z}$, $b_n \in \mathbb{N}$, and $b_n \ln_p(2) - a_n < 0$. Furthermore

$$\lim_{n \rightarrow \infty} |b_n \ln_p(2) - a_n|^{1/n^2} = p^{-\frac{3(\pi^2-4)}{2\pi^2}} < 1,$$

which implies that $\ln_p(2)$ is irrational. Its measure of irrationality satisfies $r \leq \frac{2\pi^2}{\pi^2-4} = 3.36295\dots$

Proof: Use $c = -1$ and $x = p$ in (37), then together with (38) we find

$$\lim_{n \rightarrow \infty} b_n^{1/n^2} = p^{\frac{3(\pi^2+4)}{2\pi^2}n^2}.$$

Furthermore, we have

$$\sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{-p^n - q^j} q^k = \frac{1}{P_n(-p^n|q)} \sum_{k=0}^{\infty} \frac{P_n^2(q^k|q)}{-p^n - q^j} q^k$$

and $p^n \leq p^n + q^j \leq p^n + 1$ for every j , combined with (41) implies

$$\frac{d_n(p^2)}{P_n(-p^n|q)} \frac{p^{n+1}}{(p^n + 1)(p^{2n+1} - 1)} \leq -(b_n \ln_p(2) - a_n) \leq \frac{d_n(p^2)}{P_n(-p^n|q)} \frac{p}{p^{2n+1} - 1}.$$

From this one easily finds the required asymptotics and the irrationality then follows from Lemma 1. Observe that $b_n = p^{3n^2(\pi^2+4)/(2\pi^2)+o(n^2)}$ and

$$|\ln_p(2) - \frac{a_n}{b_n}| = \mathcal{O}\left(\frac{1}{b_n^{1+\frac{\pi^2-4}{\pi^2+4}-\epsilon}}\right)$$

for every $\epsilon > 0$, which gives for the measure of irrationality the bound $r \leq 1 + \frac{\pi^2+4}{\pi^2-4} = \frac{2\pi^2}{\pi^2-4}$.

□

The upper bound is better than the upper bounds 4.8 (obtained in [1]), 4.311 (obtained in [8]), and 3.9461 (obtained in [14]).

5 Extensions

The construction of rational approximants for $h_p(1)$ and $\ln_p(2)$ can be extended with little effort to series of the form

$$L = \sum_{k=1}^{\infty} \frac{1}{cp^k - 1},$$

where $c = a/b$ is a rational number and $cp^k \neq 1$ for every $k \geq 1$. Indeed, these series can be obtained by evaluating the Stieltjes function f in (20) at cp^n , giving

$$f(cp^n) = \sum_{k=1}^{\infty} \frac{1}{cp^k - 1} - \sum_{k=1}^{n-1} \frac{1}{cp^k - 1}.$$

We then get

$$P_n(cp^n|q) \left(L - \sum_{k=1}^{n-1} \frac{1}{cp^k - 1} \right) - Q_n(cp^n|q) = \sum_{k=0}^{\infty} \frac{P_n(q^k|q)}{cp^n - q^k} q^k,$$

where

$$P_n(cp^n|q) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p p^{k(k-1)/2} (-c)^k,$$

and

$$Q_n(cp^n|q) = (-1)^{n+1} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_p \begin{bmatrix} n+k \\ k \end{bmatrix}_p (-1)^k p^{(n-k)(n-k+1)/2} \sum_{\ell=1}^k \frac{(cp^{n-k}; p)_{k-\ell}}{p^\ell - 1}.$$

In order to have integers, the quantities $P_n(cp^n|q)$ and $Q(cp^n|q)$ now need to be multiplied by b^n and by a multiple of all $p^\ell - 1$ for $1 \leq \ell \leq n$ and of all $ap^k - b$ for $1 \leq k \leq n-1$. A possible factor is $b^{2n}d_n(p)(c; p)_n$. This factor grows like

$$\lim_{n \rightarrow \infty} |b^{2n}d_n(p)(c; p)_n|^{1/n^2} = p^{\frac{3}{\pi^2} + \frac{1}{2}}.$$

In a way similar to the proof of Theorems 1 and 2 we can then prove:

Theorem 3 *Suppose $p > 1$ is an integer and $c = a/b$ is rational but $cp^k \neq 1$ for every $k = 1, 2, \dots$. Let a_n and b_n be given by*

$$b_n = b^{2n}d_n(p)(c; p)_n P_n(cp^n|q), \quad (42)$$

$$a_n = b^{2n}d_n(p)(c; p)_n Q_n(cp^n|q) + b_n \sum_{k=1}^{n-1} \frac{1}{cp^k - 1}. \quad (43)$$

Then $a_n, b_n \in \mathbb{Z}$, and

$$\lim_{n \rightarrow \infty} |b_n L - a_n|^{1/n^2} = p^{-\frac{\pi^2-3}{\pi^2}} < 1,$$

which implies that the infinite sum L is irrational. Its measure of irrationality satisfies $r \leq \frac{3\pi^2}{\pi^2-3} = 4.310119\dots$

The upper bound for the measure of irrationality corresponds to the upper bound given by Bundschuh and Väänänen [8, p. 178]. For the cases $c = 1$ and $c = -1$, which we handled in Theorems 1 and 2, one can find better upper bounds. Note that the results in [8] and [14] are also valid for p and c in other number fields. Our main purpose in this paper, however, was to emphasise the use of little q -Legendre polynomials in the construction of rational approximants for certain important Lambert series.

Acknowledgments

This research was carried out while visiting Georgia Institute of Technology. The author wishes to thank the School of Mathematics for its hospitality. Also thanks to a referee for pointing out references [3], [8], [11], and [14], and to Roberto Costas for a useful conversation that led to Lemma 2. This research is partially funded by FWO research project G.0278.97 and INTAS 2000-272.

References

- [1] T. Amdeberhan, D. Zeilberger, *q-Apéry irrationality proofs by q-WZ pairs*, Adv. Appl. Math. **20** (1998), 275–283.

- [2] R. Apéry, *Irrationalité de $\zeta(2)$ et $\zeta(3)$* , Astérisque **61** (1979), 11–13.
- [3] J-P. Bézivin, *Indépendance linéaire des valeurs des solutions transcendentes de certaines équations fonctionnelles*, Manuscripta Math. **61** (1988), 103–129.
- [4] P. Borwein, *On the irrationality of $\sum \frac{1}{q^n+r}$* , J. Number Theory **37** (1991), 253–259.
- [5] P. Borwein, *On the irrationality of certain series*, Proc. Cambridge Philos. Soc. **112** (1992), 141–146.
- [6] J. M. Borwein, P. B. Borwein, *Pi and the AGM — A Study in Analytic Number Theory and Computational Complexity*, Wiley, New York, 1987.
- [7] P. Borwein, T. Erdélyi, *Polynomials and polynomial inequalities*, Graduate Texts in Mathematics **161**, Springer-Verlag, New York, 1995.
- [8] P. Bundschuh, K. Väänänen, *Arithmetical investigations of a certain infinite product*, Compositio Math. **91** (1994), 175–199.
- [9] P. Erdős, *On arithmetical properties of Lambert series*, J. Indiana Math. Soc. **12** (1948), 63–66.
- [10] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and its Applications **35**, Cambridge University Press, Cambridge, 1990.
- [11] A. O. Gel'fond, *Functions which take on integral values*, Mat. Zam. **1** No. 5 (1967), 509–513; translated in Math. Notes **1** (1967), 337–340.
- [12] G. H. Hardy, E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford University Press, 1938 (Fifth Edition, 1979).
- [13] R. Koekoek, R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue*, Faculty of Technical Mathematics and Informatics Report 98-17, Technical University Delft, 1998. Available at <ftp://ftp.twi.tudelft.nl/TWI/publications/tech-reports/1998/DUT-TWI-98-17.ps.gz>
- [14] T. Matala-aho, K. Väänänen, *On approximation measures of q -logarithms*, Bull. Australian Math. Soc. **58** (1998), 15–31.
- [15] J. Stillwell, *Elements of Algebra*, Undergraduate Texts in Mathematics, Springer, Berlin, 1994.
- [16] W. Van Assche, *Multiple orthogonal polynomials, irrationality and transcendence*, Contemporary Mathematics
- [17] A. van der Poorten, *A proof that Euler missed . . . , Apéry's proof of the irrationality of $\zeta(3)$* , Math. Intell. **1** (1979), 195–203.

Department of Mathematics
 Katholieke Universiteit Leuven
 Celestijnenlaan 200 B
 B-3001 Leuven
 BELGIUM
 walter@wis.kuleuven.ac.be