
P-TH POWERS IN MOD P COHOMOLOGY OF FIBERS

by Luc Menichi

ABSTRACT. — Let $F \hookrightarrow E \twoheadrightarrow B$ be a fibration whose base space B is a finite simply-connected CW-complex of dimension $\leq p$ and whose total space E is a path-connected CW-complex of dimension $\leq p-1$. If $\alpha \in H^+(F; \mathbb{F}_p)$ then $\alpha^p = 0$.

RÉSUMÉ (*Les p-ièmes puissances dans la cohomologie modulo p de fibres*)

Considérons une fibration $F \hookrightarrow E \twoheadrightarrow B$ dont la base B est un CW-complexe fini simplement connexe de dimension $\leq p$, et dont l'espace total E est un CW-complexe fini connexe par arcs de dimension $\leq p-1$. Si $\alpha \in H^+(F; \mathbb{F}_p)$ alors $\alpha^p = 0$.

We work over the prime field \mathbb{F}_p with p an odd or even prime. The homology and cohomology of spaces are considered with coefficients in \mathbb{F}_p .

In [1], Anick proved using algebraic models:

THEOREM [1, 9.1]. — *Let r be a non-negative integer. Let B be a simply-connected space with a finite type homology concentrated in degrees $i \in [r+1, rp]$. Then all p -th powers vanish in $H^+(\Omega B)$.*

This result was suggested by McGibbon and Wilkerson [6, p. 699]. The aim of this note is to give two different generalisations of Anick Theorem: Theorem A and Theorem C below.

LUC MENICHI, Université d'Angers, Faculté des Sciences, 2 Boulevard Lavoisier, 49045 Angers, FRANCE, Telephone: 02-41-73-50-25, Fax: 02-41-73-54-54
E-mail : Luc.Menichi@univ-angers.fr

2000 Mathematics Subject Classification. — 57T35, 55T20, 55R20.

Key words and phrases. — Eilenberg-Moore spectral sequence, fiber space, loop space, free loop space, p -th powers.

Topologie/Topology.

The first one, whose proof is inspired by a result of Lannes and Schwartz [4, Prop 0.6], uses the (vertical) Steenrod operations in the Eilenberg-Moore spectral sequence:

THEOREM A. — *Let r and k be two non-negative integers. Consider a fiber product of spaces*

$$\begin{array}{ccc} E \times_B X & \longrightarrow & E \\ \downarrow & & \downarrow \pi \\ X & \longrightarrow & B \end{array}$$

where

- π is a Serre fibration and
- $H^*(E)$, $H^*(X)$ and $H^*(B)$ are of finite type.

If B is simply-connected with homology concentrated in degrees $i \in [r+1, rp^k]$ and the product space $E \times X$ is path connected with homology $H_*(E \times X)$ concentrated in degrees $i \in [r, rp^k-1]$, then all p^k -th powers vanish in $H^+(E \times_B X)$.

Proof. — We suppose that p is an odd prime. The case $p = 2$ is similar. Let \mathcal{A} denote the mod p Steenrod algebra. The degree of an element α is denoted $|\alpha|$. Recall from [8, 11, 10], that the Eilenberg-Moore spectral sequence is a strongly convergent second quadrant cohomological spectral sequence of \mathcal{A} -modules

$$E_2^{-s,*} \cong \mathrm{Tor}_{H^*(B)}^{-s,*}(H^*(E), H^*(X)) \Rightarrow H^*(E \times_B X).$$

More precisely, there exists a convergent filtration of \mathcal{A} -modules on $H^*(E \times_B X)$:

$$H^*(E \times_B X) \supset \dots F_s \supset F_{s-1} \dots F_1 \supset F_0 \supset F_{-1} = \{0\}$$

such that $\Sigma^{-s} F_s / F_{s-1} \cong E_\infty^{-s,*}$, $s \geq 0$. Here Σ^{-s} denotes the s -th desuspension of an \mathcal{A} -module.

Let $\alpha \in F_s$ such that the class $[\alpha] \in F_s / F_{s-1}$ is non zero. We want to prove that $\alpha^{p^k} = 0$. As an \mathcal{A} -module, $\mathrm{Tor}_{H^*(B)}^{-s,*}(H^*(E), H^*(X))$ is the s -th homology group of a complex of \mathcal{A} -modules, namely the Bar construction, whose s -th term is $H^*(E) \otimes H^+(B)^{\otimes s} \otimes H^*(X)$.

The element $\Sigma^{-s}[\alpha] \in E_\infty^{-s,*}$ is represented by a cycle of the form $e[b_1] \dots [b_s]x$, where $e \in H^*(E)$, $(b_i)_{1 \leq i \leq s} \in H^+(B)$ and $x \in H^*(X)$. So $rs \leq |\alpha| \leq (rp^k - 1)(s+1)$.

Case 1: $|e| + |x| \geq r$. Then $|\alpha| \geq r(s+1)$. Therefore, by a degree argument, the element α^{p^k} of F_s is zero.

Case 2: $e = x = 1$. Since the Cartan formula applies, $\Sigma^{-s}[\alpha^p] = P^{|\alpha|/2} \Sigma^{-s}[\alpha]$ is represented by the element of the Bar construction,

$$\sum_{i_1 + \dots + i_s = |\alpha|/2} [P^{i_1} b_1 | \dots | P^{i_s} b_s].$$

So $\Sigma^{-s}[\alpha^{p^k}]$ is zero for degree reasons. Therefore α^{p^k} belongs to F_{s-1} which is concentrated in degrees $\leq (rp^k - 1)s$, thus $\alpha^{p^k} = 0$. \square

Let B be a space. The *free loop space* on B , denoted B^{S^1} , is the set of continuous (unpointed) maps from the circle S^1 to B . It can be defined as a fibre product:

$$\begin{array}{ccc} B^{S^1} & \xrightarrow{\quad} & B^{[0,1]} \\ \downarrow ev & & \downarrow \pi \\ B & \xrightarrow[\Delta]{} & B \times B \end{array}$$

In this particular case, we can improve Theorem A.

THEOREM B. — *Let r and k be two non-negative integers. If B is a simply-connected space with finite type homology concentrated in degrees $i \in [r+1, rp^k]$ then all p^k -th powers vanish in $H^+(B^{S^1})$.*

Proof. — The Eilenberg-Moore spectral sequence for the previous fiber product satisfies

$$E_2^{-s,*} \cong \mathrm{HH}_s(H^*(B)) \Rightarrow H^*(B^{S^1}).$$

Here HH_* denotes the Hochschild homology. As an \mathcal{A} -module, $\mathrm{HH}_s(H^*(B))$ is the s -th homology group of a complex of \mathcal{A} -modules, namely the Hochschild complex, whose s -th term is $H^*(B) \otimes H^+(B)^{\otimes s}$.

The same arguments as in the proof of Theorem A allow us to conclude except in Case 2 for $s = 1$. If $\alpha \in F_1 \subset H^*(B^{S^1})$, we can only affirm that $\alpha^{p^k} \in F_0$. The evaluation map $ev : B^{S^1} \rightarrow B$ admits a section σ . So $H^*(ev) : H^*(B) \hookrightarrow (B^{S^1})^*$ admits $H^*(\sigma)$ as retract. The edge homomorphism

$$H^*(B) = E_2^{0,*} \rightarrow E_3^{0,*} \cdots \rightarrow E_\infty^{0,*} = F_0 \subset H^*(B^{S^1})$$

correspond to $H^*(ev)$. Since $\alpha^{p^k} \in F_0 = H^*(B)$, $\alpha^{p^k} = [H^*(\sigma)(\alpha)]^{p^k}$. For degree reason, all p^k -th powers are zero in $H^+(B)$. So $\alpha^{p^k} = 0$. \square

In [3], Félix, Halperin and Thomas give a slightly more complicated proof of Anick Theorem. Their proof uses the vertical and horizontal Steenrod operations in the Serre spectral sequence:

THEOREM [3, 2.9(i)]. — *Let r and k be two non-negative integers. If B is a simply-connected space with a finite type homology concentrated in degrees $i \in [r+1, rp^k]$ then all p^k -th powers vanish in $H^+(\Omega B)$.*

This result generalizes in:

THEOREM C (Compare with [7, 10.8]). — *Let r and k be two non-negative integers. Let $F \xrightarrow{j} E \xrightarrow{\pi} B$ be a Serre fibration with E path connected. If B is a simply-connected space with finite type homology concentrated in degrees $i \in [r+1, rp^k]$ then, for any $\alpha \in H^*(F)$, $\alpha^{p^k} \in \text{Im } H^*(j)$.*

Proof. — The proof follows the lines of [3, 2.9]. Since $H^{\leq r}(B) = 0$, $\alpha \in E_2^{0,*}$ survives till $E_{r+1}^{0,*}$. Therefore by a Theorem of Araki [2] and Vázquez [12] (See also [9, Prop 2.5 Case 2]), $\alpha^{p^k} \in E_2^{0,*}$ survives till $E_{rp^k+1}^{0,*}$. Since $H^{>rp^k}(B) = 0$,

$$E_{rp^k+1}^{0,*} = E_{\infty}^{0,*} = \text{Im } H^*(j).$$

□

In order to see that the hypothesis in Félix-Halperin-Thomas Theorem (and in Theorem B) cannot be improved, consider $B = \Sigma \mathbb{CP}^{p^k}$, the suspension of the p^k -dimension complex projective space.

Observe also that in Theorem C, α^{p^k} is not zero in general. Indeed, take π to be the fibration associated to the suspension of the Hopf map from S^{2p^k-1} to \mathbb{CP}^{p^k} [7, Remark 9.9].

Finally, remark that the following question of McGibbon and Wilkerson remains unsolved.

QUESTION [6, p. 699] (See also [5, section 9.Question 3])

Let B be a finite simply-connected CW-complex and p a prime large enough. Do all the Steenrod operations act trivially on $H^(\Omega B)$?*

I wish to thank Professor Katsuhiko Kuribayashi, for his precious help with the Eilenberg-Moore spectral sequence.

References

- [1] D. J. ANICK – Hopf algebras up to homotopy, *J. Amer. Math. Soc.* **2** (1989), no. 3, p. 417–453.
- [2] S. ARAKI – Steenrod reduced powers in the spectral sequences associated with a fibering. I, II, *Mem. Fac. Sci. Kyushu Univ. Ser. A* **11** (1957), p. 15–64, 81–97.

- [3] Y. FÉLIX, S. HALPERIN & J.-C. THOMAS – The Serre spectral sequence of a multiplicative fibration, Prépublications, Université d’Angers, 1998, <http://sciences.univ-angers.fr/math/preprint.html>.
- [4] J. LANNES & L. SCHWARTZ – A propos de conjectures de Serre et Sullivan, *Invent. Math.* **83** (1986), p. 593–603.
- [5] C. A. MCGIBBON – Phantom maps, Handbook of Algebraic Topology (I. M. James, ed.), North-Holland, Amsterdam, 1995, p. 1209–1257.
- [6] C. A. MCGIBBON & C. W. WILKERSON – Loop spaces of finite complexes at large primes, *Proc. Amer. Math. Soc.* **96** (1986), p. 698–702.
- [7] L. MENICHI – On the cohomology algebra of a fiber, Preprint Hopf Topology Archive, 1999.
- [8] D. L. RECTOR – Steenrod operations in the Eilenberg-Moore spectral sequence, *Comment. Math. Helv.* **45** (1970), p. 540–552.
- [9] J. SAWKA – Odd primary Steenrod operations in first-quadrant spectral sequences, *Trans. Am. Math. Soc.* **273** (1982), p. 737–752.
- [10] L. SCHWARTZ – Unstable modules and unstable algebras. Applications to homotopy theory, Lectures Notes of the Grenoble Summer School, <http://www-fourier.ujf-grenoble.fr>, 1997.
- [11] L. SMITH – On the Künneth theorem. I: The Eilenberg-Moore spectral sequence, *Math. Z.* **116** (1970), p. 94–140.
- [12] R. VÁZQUEZ GARCIA – Note on Steenrod squares in the spectral sequence of a fibre space, *Bol. Soc. Math Mexicana* **2** (1957), p. 1–8.