

Quaternionic linear algebra and plurisubharmonic functions of quaternionic variables.

S.Alesker

Department of Mathematics, Tel Aviv University, Ramat Aviv
69978 Tel Aviv, Israel

Abstract

We remind known and establish new properties of the Dieudonné and Moore determinants of quaternionic matrices. Using these linear algebraic results we develop a basic theory of plurisubharmonic functions of quaternionic variables.

0 Introduction.

The main point of this paper is that in quaternionic algebra and analysis there exist structures which have analogues over the fields of real and complex numbers, but should reflect different phenomena.

The algebraic part is discussed in Section 1. There we remind the notions of the Moore and Dieudonné determinants of quaternionic matrices. It turns out that (under appropriate normalization) the Dieudonné determinant behaves exactly like the absolute value of the usual determinant of real or complex matrices from all points of view (algebraic and analytic). Let us state some of its properties discussed in more details in Subsection 1.2. Let us denote by $M_n(\mathbb{H})$ the set of all quaternionic $n \times n$ -matrices. The Dieudonné determinant D is defined on this set and takes values in non-negative real number:

$$D : M_n(\mathbb{H}) \longrightarrow \mathbb{R}_{\geq 0}$$

(see Definition 1.2.2). Then one has the following (known) results (see Theorems 1.2.3 and 1.2.4 below and references given at the beginning of Section 1):

Theorem.

(i) For any complex $n \times n$ - matrix X considered as quaternionic matrix the Dieudonné determinant $D(X)$ is equal to the absolute value of the usual determinant of X .

(ii) For any quaternionic matrix X

$$D(X) = D(X^t) = D(X^*),$$

where X^t and X^* denote the transposed and quaternionic conjugate matrices respectively.

(iii) $D(X \cdot Y) = D(X)D(Y)$.

Theorem 1.2.5. Let $A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \dots & \dots & \dots \\ a_{n1} & \dots & a_{nn} \end{bmatrix}$ be a quaternionic matrix.

Then

$$D(A) \leq \sum_{i=1}^n |a_{1i}| D(M_{1i}).$$

Similar inequalities hold for any other row or column.

(In this theorem $|a|$ denotes the absolute value of a quaternion a , and M_{pq} denotes the minor of the matrix A obtained from it by erasing the p -th row and q -th column).

In a sence, the Dieudonné determinant provides the theory of *absolute value* of determinant. However it is not always sufficient and we loose most of the algebraic properties of the usual determinant. The notion of Moore determinant provides such a theory, but only on the class of quaternionic *hyperhermitian* matrices. Remind that a square quaternionic matrix A is called hyperhermitian if its quaternionic conjugate $A^* = A$. The Moore determinant denoted by *det* is defined on the class of all hyperhermitian matrices and takes real values. (The Moore determinant is defined in Subsection 1.1. after Theorem 1.1.8). The important advantage of it with respect to the Dieudonné determinant is that it depends polynomially on the entries of a matrix; it has already all the algebraic and analytic properties of the usual determinant of real symmetric and complex hermitian matrices. Let us state some of them referring for the details to Subsection 1.1 (again, the references are given at the beginning of Section 1).

Theorem 1.1.9.

- (i) *The Moore determinant of any complex hermitian matrix considered as quaternionic hyperhermitian matrix is equal to its usual determinant.*
- (ii) *For any hyperhermitian matrix A and any quaternionic matrix C*

$$\det(C^*AC) = \det A \cdot \det(C^*C).$$

Example.

- (a) Let $A = \text{diag}(\lambda_1, \dots, \lambda_n)$ be a diagonal matrix with real λ_i 's. Then A is hyperhermitian and the Moore determinant $\det A = \prod_i \lambda_i$.
- (b) A general hyperhermitian 2×2 matrix A has the form

$$A = \begin{bmatrix} a & q \\ \bar{q} & b \end{bmatrix},$$

where $a, b \in \mathbb{R}$, $q \in \mathbb{H}$. Then its Moore determinant is equal to $\det A = ab - q\bar{q}$.

Next, in terms of the Moore determinant one can prove the generalization of the classical Sylvester criterion of positive definiteness of hyperhermitian matrices (Theorem 1.1.13). In terms of the Moore determinant one can introduce the notion of the mixed discriminant and to prove the analogues of Alexandrov's inequalities for mixed discriminants (Theorem 1.1.15 and Corollary 1.1.16).

The (well known) relation between the Dieudonné and Moore determinants is as follows: for any hyperhermitian matrix X

$$D(X) = |\det X|.$$

In Section 1 we prove some additional properties of the Dieudonné and Moore determinants; they are used in Section 2.

Note that the Dieudonné determinant was introduced originally by J. Dieudonné in [9] (see also [2] for his theory). It can be defined for arbitrary (non-commutative) field. On more modern language this result can be formulated as a computation of the K_1 -group of a non-commutative field (see e.g. [25]).

Section 2 of this paper develops the basic theory of plurisubharmonic functions of quaternionic variables on \mathbb{H}^n . It uses in essential way the linear algebraic results of Section 1. This theory is parallel to the classical theories of convex functions on \mathbb{R}^n and plurisubharmonic functions on \mathbb{C}^n .

The formal definition is as follows (for more discussion see Subsection 2.1).

Definition. A continuous real valued function

$$f : \mathbb{H}^n \longrightarrow \mathbb{R}$$

is called quaternionic plurisubharmonic if its restriction to any quaternionic line is subharmonic (i.e. its Laplacian is non-negative).

This definition is different but equivalent (by Proposition 2.1.6 below) to the original author's definition. In this form it was suggested by G. Henkin [15]. Let us describe the main results on plurisubharmonic functions we prove. We will write a quaternion q in the usual form

$$q = t + x \cdot i + y \cdot j + z \cdot k,$$

where t, x, y, z are real numbers, and i, j, k satisfy the usual relations

$$i^2 = j^2 = k^2 = -1, ij = -ji = k, jk = -kj = i, ki = -ik = j.$$

Let us introduce the differential operators $\frac{\partial}{\partial \bar{q}}$ and $\frac{\partial}{\partial q}$ as follows:

$$\frac{\partial}{\partial \bar{q}} f := \frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}, \text{ and}$$

$$\frac{\partial}{\partial q} f := \overline{\frac{\partial}{\partial \bar{q}} \bar{f}} = \frac{\partial f}{\partial t} - \frac{\partial f}{\partial x} i - \frac{\partial f}{\partial y} j - \frac{\partial f}{\partial z} k.$$

The operator $\frac{\partial}{\partial \bar{q}}$ is called the Cauchy-Riemann-Fueter operator since it was introduced by Fueter [12], [13] to define the notion of quaternionic analyticity. For further results on quaternionic analyticity we refer e.g. to [21], [22], [26]. First one has a simple

Proposition 2.1.6. *A real valued twice continuously differentiable function f on the domain $\Omega \subset \mathbb{H}^n$ is quaternionic plurisubharmonic if and only if at every point $q \in \Omega$ the matrix $(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})(q)$ is non-negative definite.*

Note that the matrix in the statement of proposition is quaternionic hyperhermitian. The more important thing is that in analogy to the real and complex cases one can define for any continuous quaternionic plurisubharmonic function f a non-negative measure $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})(q)$, where \det denotes

the Moore determinant (this measure is obviously defined for smooth f). We prove the following continuity result.

Theorem 2.1.11. *Let $\{f_N\}$ be sequence of continuous quaternionic plurisubharmonic function in a domain $\Omega \subset \mathbb{H}^n$. Assume that this sequence converges uniformly on compact subsets to a function f . Then f is continuous quaternionic plurisubharmonic function. Moreover the sequence of measures $\det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})$ weakly converges to the measure $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})$.*

The proofs of analogous results in real and complex cases can be found in [4], where the exposition of this topic follows the approach of Rauch- Taylor [24]. We adopt this method to the quaternionic situation. The difficulties come mostly from linear algebra since the technique of working with the Moore determinant is not sufficiently developed. For instance there is no formula of decomposition of the Moore determinant in row or column, and thus one should use some more tricky manipulations.

Next we would like to state a result on existence of solution of the Dirichlet problem for quaternionic Monge- Ampère equation (to be defined). The proof of this result is not presented in the paper, but it can be obtained by generalizing to the quaternionic case the method of Bedford and Taylor [6].

Definition. An open bounded domain $\Omega \subset \mathbb{H}^n$ is called strictly pseudoconvex if there exists a smooth strictly plurisubharmonic function h on \mathbb{H}^n such that $\Omega = \{h < 0\}$ and $\nabla h|_{\partial\Omega} \neq 0$.

The next result is quaternionic analogue of the results on Dirichlet problem of real and complex Monge- Ampere equations (see e.g. [4] and [6]).

Theorem. *Let Ω be a strictly pseudoconvex bounded domain in \mathbb{H}^n . Let ϕ be a continuous real valued function on the boundary $\partial\Omega$. Let f be a continuous function on the closure $\bar{\Omega}$, $f \geq 0$. Then there exists a unique continuous on $\bar{\Omega}$ plurisubharmonic function u such that*

$$\det(\frac{\partial^2 u}{\partial q_i \partial \bar{q}_j}) = f \text{ and}$$

$$u = \phi \text{ on } \Omega.$$

In appendix to this paper we prove the injectivity of Radon transform over quaternionic subspaces in the affine space \mathbb{H}^n . Probably this result is not new. It is included here since it was used in the proof of Lemma 2.1.7, and we could not find a reference.

Acknowledgements. We express our gratitude to J. Bernstein, P. Biran, G. Henkin, D. Kazhdan, L. Polterovich, and M. Sodin for useful discussions. We thank also E. Shustin who has shown us the reference [3].

1 Quaternionic linear algebra.

In this section we remind the construction and basic properties of the Dieudonné and Moore determinants and investigate further their properties. Part of them will be used in the next sections of this paper. For a survey of quaternionic determinants and references we refer to [3].

First of all remind that over any noncommutative field there exist usual notions of vector spaces over the field (however one should distinguish between left and right ones), their dimension, basis etc. (see e.g. [2]). However there is no construction of quaternionic determinant which would have all the properties of the determinant over commutative field. We are going to discuss this problem in this section. We will discuss only right vector spaces. The case of left ones can be considered similarly. Many results of Section 1 are a folklore. Theorems 1.1.8, 1.1.9, 1.1.4 are not new. We refer for the proofs to [5], [7],[8], [10],[11], [16], [17], [18], [19], [20], [23], [27], [28].

1.1 Hyperhermitian forms and the Moore determinant.

Let V be a right vector space over quaternions.

1.1.1. Definition. A *hyperhermitian semilinear form* on V is a map $a : V \times V \longrightarrow \mathbb{H}$ satisfying the following properties:

- (a) a is additive with respect to each argument;
- (b) $a(x, y \cdot q) = a(x, y) \cdot q$ for any $x, y \in V$ and any $q \in \mathbb{H}$;
- (c) $a(x, y) = \overline{a(y, x)}$.

1.1.2. Example. Let $V = \mathbb{H}^n$ be the standard coordinate space considered as right vector space over \mathbb{H} . Fix a *hyperhermitian* $n \times n$ matrix $(a_{ij})_{i,j=1}^n$, i.e. $a_{ij} = \overline{a_{ji}}$, where \bar{a} denotes the usual quaternionic conjugation. For $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$ define

$$A(x, y) = \sum_{i,j} \bar{x}_i a_{ij} y_j$$

(note the order of the terms!). Then A defines hyperhermitian semilinear form on V .

In general one has the following standard claims.

1.1.3. Claim. *Fix a basis in a finite dimensional right quaternionic vector space V . Then there is a natural bijection between hyperhermitian semilinear forms on V and $n \times n$ hyperhermitian matrices.*

This bijection is in fact described in previous Example 1.1.2.

1.1.4. Claim. *Let A be the matrix of the given hyperhermitian form in the given basis. Let C be transition matrix from this basis to another one. Then the matrix A' of the given form in the new basis is equal*

$$A' = C^* A C.$$

1.1.5. Remark. Note that for any hyperhermitian matrix A and for any matrix C the matrix $C^* A C$ is also hyperhermitian. In particular the matrix $C^* C$ is always hyperhermitian.

1.1.6. Definition. A hyperhermitian semilinear form a is called *positive definite* if $a(x, x) > 0$ for any non-zero vector x .

Let us fix on our quaternionic right vector space V a positive definite hyperhermitian form (\cdot, \cdot) . The space with a fixed such a form will be called *hyperhermitian space*.

For any quaternionic linear operator $\phi : V \longrightarrow V$ in hyperhermitian space one can define the adjoint operator $\phi^* : V \longrightarrow V$ in the usual way, i.e. $(\phi x, y) = (x, \phi^* y)$ for any $x, y \in V$. Then if one fixes an orthonormal basis in the space V then the operator ϕ is selfadjoint if and only if its matrix in this basis is hyperhermitian.

1.1.7. Claim. *For any selfadjoint operator in a hyperhermitian space there exists an orthonormal basis such that its matrix in this basis is diagonal and real. Moreover any operator with hyperhermitian matrix in an orthonormal basis is selfadjoint.*

The proof is standard. Now we are going to define the Moore determinant of a hyperhermitian matrices. The definition below is different from the original one [20] but equivalent to it.

First note that every hyperhermitian $n \times n$ - matrix A defines a hyperhermitian semilinear form on the coordinate space \mathbb{H}^n . It also can be considered as a *symmetric* bilinear form on \mathbb{R}^{4n} (which is the realisation of \mathbb{H}^n). Let us denote its $4n \times 4n$ - matrix by $\mathbb{R}A$. Let us consider the entries of A as formal variables (each quaternionic entry corresponds to four commuting real variables). Then $\det(\mathbb{R}A)$ is a homogeneous polynomial of degree $4n$ in $n(2n-1)$ variables. Let us denote by Id the identity matrix. One has the following result.

1.1.8. Theorem. *There exists a polynomial P defined on the space of all hyperhermitian $n \times n$ - matrices such that for any hyperhermitian $n \times n$ - matrix A one has $\det(\mathbb{R}A) = P^4(A)$ and $P(Id) = 1$. P is defined uniquely by these two properties. Furthermore P is homogeneous of degree n and has integer coefficients.*

Thus for any hyperhermitian matrix A the value $P(A)$ is a real number, and it is called the *Moore determinant* of the matrix A . The explicit formula for the Moore determinant was given by Moore [20] (see also [3]). From now on the Moore determinant will be denoted by $\det A$. This notation should not cause any confusion with the usual determinant of real or complex matrices due to part (i) of the next theorem.

1.1.9. Theorem. (i) *The Moore determinant of any complex hermitian matrix considered as quaternionic hyperhermitian matrix is equal to its usual determinant.*

(ii) *For any hyperhermitian matrix A and any matrix C*

$$\det(C^*AC) = \det A \cdot \det(C^*C).$$

1.1.10. Example. (a) Let $A = \text{diag}(\lambda_1, \dots, \lambda_n)$ be a diagonal matrix with real λ_i 's. Then A is hyperhermitian and the Moore determinant $\det A = \prod_i \lambda_i$.

(b) A general hyperhermitian 2×2 matrix A has the form

$$A = \begin{bmatrix} a & q \\ \bar{q} & b \end{bmatrix},$$

where $a, b \in \mathbb{R}$, $q \in \mathbb{H}$. Then $\det A = ab - q\bar{q}$.

Let us introduce more notation. Let A be any hyperhermitian $n \times n$ -matrix. For any non-empty subset $I \subset \{1, \dots, n\}$ the minor $M_I(A)$ of A which is obtained by erasing the rows and columns with indexes from the set I , is clearly hyperhermitien. For $I = \{1, \dots, n\}$ let $\det M_{\{1, \dots, n\}} = 1$.

1.1.11. Proposition. *For any hyperhermitian $n \times n$ - matrix A and any*

$$\text{diagonal real matrix } T = \begin{bmatrix} t_1 & & 0 \\ & \ddots & \\ 0 & & t_n \end{bmatrix}$$

$$\det(A + T) = \sum_{I \subset \{1, \dots, n\}} \left(\prod_{i \in I} t_i \right) \cdot \det M_I(A).$$

In particular

$$\det(A + t \cdot Id) = \sum_{I \subset \{1, \dots, n\}} t^{|I|} \cdot \det M_I(A),$$

where $|I|$ denotes the cardinality of the set I .

Remark. Clearly this formula is true for *arbitrary* matrix A over *commutative* field.

Proof. Fix a hyperhermitien matrix A . It is clear that $\det(A + T)$ is a polynomial in t_1, \dots, t_n of degree n . Since

$$A + \begin{bmatrix} t_1 & & 0 \\ & t_2 & \\ & & \ddots \\ 0 & & & t_n \end{bmatrix} = A + \begin{bmatrix} 0 & & 0 \\ & t_2 & \\ & & \ddots \\ 0 & & & t_n \end{bmatrix} + \begin{bmatrix} t_1 & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix}$$

one can apply induction in n to show that if $A = \left[\begin{array}{c|c} a_{11} & * \\ \hline * & B \end{array} \right]$, where $a_{11} \in \mathbb{R}$, and B is a hyperhermitian $(n-1) \times (n-1)$ - matrix. Set

$$f(t) := \det \left(A + \begin{bmatrix} t & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} \right).$$

It is sufficient to show that $f(t) = \det A + t \cdot \det B$. Clearly $f(0) = \det A$. Let k denote the degree of the polynomial f . Using Theorem 1.1.9(ii) one gets $f(t) =$

$$t^k \cdot \det \left(\begin{bmatrix} t^{-k/2} & & 0 \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{bmatrix} \left(A + \begin{bmatrix} t & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix} \right) \begin{bmatrix} t^{-k/2} & & 0 \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{bmatrix} \right) =$$

$$t^k \det \left[\begin{array}{c|ccc} a_{11}t^{-k} + t^{-k+1} & a_{12}t^{-k/2} & \dots & a_{1n}t^{-k/2} \\ \hline a_{21}t^{-k/2} & & & \\ \vdots & & B & \\ a_{n1}t^{-k/2} & & & \end{array} \right].$$

If $k > 1$ then

$$\frac{f(t)}{t^k} \longrightarrow \det \left[\begin{array}{c|c} 0 & 0 \\ \hline 0 & B \end{array} \right] = 0$$

when $t \longrightarrow \infty$. Hence $k = 1$ and

$$\frac{f(t)}{t} \longrightarrow \det \left[\begin{array}{c|c} 1 & 0 \\ \hline 0 & B \end{array} \right] = \det B.$$

Q.E.D.

1.1.12. Lemma. *Let A be a non-negative (resp. positive) definite hyperhermitian matrix. Then $\det A \geq 0$ (resp. $\det A > 0$).*

Proof. Let us prove it under the assumption that A is positive definite. By Claim 1.1.7 there exists a matrix $C \in Sp(n)$ (i.e. $C^*C = Id$) such that

$$A = C^* \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} C \text{ with } \lambda_i \in \mathbb{R}. \text{ Since } A \text{ is positive definite } \lambda_i > 0$$

for all i . By Theorem 1.1.9(ii) $\det A = \det(C^*C) \prod \lambda_i = \prod \lambda_i > 0$. Q.E.D.

The following theorem is a quaternionic generalization of the standard Sylvester criterion.

1.1.13. Theorem (Sylvester criterion). *A hyperhermitian $n \times n$ - matrix A is positive definite if and only if $M_{\{i+1, \dots, n\}}(A) > 0$ for any $0 \leq i \leq n-1$.*

Proof. The necessity follows from Lemma 1.1.12. Let us prove sufficiency by induction in n . For $n = 1$ the statement is trivial. Assume $n > 1$. Let

$$A = \left[\begin{array}{c|ccc} a_{11} & a_{12} & \dots & a_{1n} \\ \hline a_{21} & & & \\ \vdots & & B & \\ a_{n1} & & & \end{array} \right].$$

Consider the matrix

$$U = \left[\begin{array}{ccccc} 1 & -a_{12}/a_{11} & -a_{13}/a_{11} & \dots & -a_{1n}/a_{11} \\ 0 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{array} \right].$$

Then the matrix $A' := U^*AU$ has the form

$$A' = \left[\begin{array}{cccc} a_{11} & 0 & \dots & 0 \\ 0 & & & \\ \vdots & & B' & \\ 0 & & & \end{array} \right],$$

where B' is a hyperhermitian matrix. Moreover for any $1 \leq i \leq n-1$ one has

$$\det M_{\{i+1, \dots, n\}}(A') = \det M_{\{i+1, \dots, n\}}(A).$$

Indeed let us check it for $i = n$ (for $i < n$ the proof will be the same since the matrix U is triangular). Namely let us show that $\det A' = \det A$. By Theorem 1.1.9 (ii) $\det A' = \det A \cdot \det(U^*U)$. However using Theorem 1.1.8 and unipotence of U it is easy to see that $\det(U^*U) = 1$. Hence the matrix B' is positive definite by the induction assumption. Then A' is positive definite, and hence A is as well. Q.E.D.

Let us define now the mixed discriminant of hyperhermitian matrices in analogy with the case of real symmetric matrices [1].

1.1.14. Definition. Let A_1, \dots, A_n be hyperhermitian $n \times n$ -matrices. Consider the homogeneous polynomial in real variables $\lambda_1, \dots, \lambda_n$ of degree n equal to $\det(\lambda_1 A_1 + \dots + \lambda_n A_n)$. The coefficient of the monomial $\lambda_1 \dots \lambda_n$ divided by $n!$ is called the *mixed discriminant* of the matrices A_1, \dots, A_n , and it is denoted by $\det(A_1, \dots, A_n)$.

Note that the mixed discriminant is symmetric with respect to all variables, and linear with respect to each of them, i.e.

$$\det(\lambda A'_1 + \mu A_1'', A_2, \dots, A_n) = \lambda \cdot \det(A'_1, A_2, \dots, A_n) + \mu \cdot \det(A_1'', A_2, \dots, A_n)$$

for any real λ, μ . Note also that $\det(A, \dots, A) = \det A$. We will prove the following generalization of Alexandrov's inequalities for mixed discriminants [1] (though the proof will be very close to the original one).

1.1.15. Theorem. (i) *The mixed discriminant of positive (resp. non-negative) definite matrices is positive (resp. non-negative) definite.*

(ii) *Fix positive definite hyperhermitian $n \times n$ - matrices A_1, \dots, A_{n-2} . On the real linear space of hyperhermitian $n \times n$ - matrices consider the bilinear form*

$$B(X, Y) := \det(X, Y, A_1, \dots, A_{n-2}).$$

Then B is non-degenerate quadratic form, and its signature has one plus and the rest are minuses.

1.1.16. Corollary. *Let A_1, \dots, A_{n-1} be positive definite hyperhermitian $n \times n$ - matrices. Then for any hyperhermitian matrix X*

$$\det(A_1, \dots, A_{n-1}, X)^2 \geq \det(A_1, \dots, A_{n-1}, A_{n-1}) \cdot \det(A_1, \dots, A_{n-2}, X, X), \quad (1)$$

and the equality is satisfied if and only if the matrix X is proportional to A_{n-1} .

Proof of Corollary 1.1.16. By Theorem 1.1.15 (i) we get $\det(A_1, \dots, A_{n-1}, A_{n-1}) > 0$. Let

$$\lambda = \frac{\det(A_1, \dots, A_{n-1}, X)}{\det(A_1, \dots, A_{n-1}, A_{n-1})}.$$

Let $X' = X - \lambda A_{n-1}$. Then clearly $\det(A_1, \dots, A_{n-1}, X') = 0$. In the notation of Theorem 1.1.15 it means that $B(A_{n-1}, A_{n-1}) > 0$ and $B(A_{n-1}, X') = 0$. But the form B has just one plus. Hence $B(X', X') \leq 0$, and the equality is satisfied if and only if $X' = 0$. Developing $B(X', X')$ one gets inequality (1). The equality case follows as well. Q.E.D.

Proof of Theorem 1.1.15. (1) Let us prove the first part using induction in n . The case $n = 1$ is trivial. Assume that $n > 1$. Let A_1, \dots, A_n be positive definite hyperhermitian matrices. By Claim 1.1.7 and Theorem 1.1.9 (ii) we

can assume that the matrix A_n is diagonal, i.e. $A_n = \begin{bmatrix} t_1 & & 0 \\ & \ddots & \\ 0 & & t_n \end{bmatrix}$, and t_i 's are positive. By Proposition 1.1.11

$$\det(\lambda_1 A_1 + \cdots + \lambda_{n-1} A_{n-1} + \lambda_n A_n) = \sum_{I \subset \{1, \dots, n\}} \left(\prod_{i \in I} \lambda_n t_i \right) \cdot \det M_I(\lambda_1 A_1 + \cdots + \lambda_{n-1} A_{n-1}).$$

Since all the diagonal minors of positive definite matrix are positive definite and since $t_i > 0$ the assumption of induction implies the statement.

(2) Let us prove the second part of the theorem, i.e. that B is non-degenerate. First let us prove it for $n = 2$. Assume X_0 belongs to the kernel of B , i.e. $B(X, X_0) = 0$ for every X . One can assume that X_0 is diagonal: $X_0 = \begin{bmatrix} t_1 & 0 \\ 0 & t_2 \end{bmatrix}$. For any $X = \begin{bmatrix} x_1 & 0 \\ 0 & x_2 \end{bmatrix}$ with real x_1, x_2 one has $2\det(X, X_0) = t_1 x_2 + t_2 x_1 = 0$. Hence $t_1 = t_2 = 0$. Thus the form B is non-degenerate. Now, clearly $B(Id, Id) = 1 > 0$. Assume that $X \neq 0$ is orthogonal to Id with respect to B , i.e. $B(X, Id) = 0$. It remains to show that $B(X, X) < 0$. By Claim 1.1.7 we can assume that X is diagonal, $X = \begin{bmatrix} x_1 & 0 \\ 0 & x_2 \end{bmatrix}$. Then $2B(X, Id) = x_1 + x_2 = 0$. But $B(X, X) = x_1 x_2 = -x_1^2 < 0$.

Let us assume that $n > 2$. Assume also that the theorem is true for matrices of size at most $n - 1$. Let us prove first that the form B is non-degenerate. Assume that X_0 belongs to the kernel of B . Since A_{n-2} is positive definite, by Claim 1.1.7 one can assume that the matrix A_{n-2} is equal to Id and X_0 is diagonal. For $1 \leq i \leq n$ and for $n \times n$ -matrices C_1, \dots, C_{n-1} let us denote by $\det(C_1, \dots, C_{n-1})_i$ the mixed discriminant of $(n-1) \times (n-1)$ -matrices obtained from C_j 's by erasing the i 's row and the i 's column. Let

$T = \begin{bmatrix} t_1 & & 0 \\ & \ddots & \\ 0 & & t_n \end{bmatrix}$. Using Proposition 1.1.11 one can easily see that

$$c \cdot \det(A_1, \dots, A_{n-2}, T, X_0) = \sum_{i=1}^n t_i \det(A_1, \dots, A_{n-2}, X_0)_i, \quad (2)$$

where c is a normalizing constant. Hence $\det(A_1, \dots, A_{n-2}, X_0)_i = 0$ for all i . By the induction assumption and Corollary 1.1.16 (which is also satisfied for matrices of size $n - 1$)

$$\det(A_1, \dots, A_{n-3}, X_0, X_0)_i \leq 0 \text{ for } i = 1, \dots, n \quad (3)$$

with equalities if and only if the matrix X_0 vanishes. Since $A_{n-2} = Id$ and X_0 belongs to the kernel of B the equality analogous to (2) implies that

$$0 = c \cdot \det(A_1, \dots, A_{n-3}, A_{n-2}, X_0, X_0) = \sum_{i=1}^n \det(A_1, \dots, A_{n-3}, X_0, X_0)_i.$$

By inequalities (3) one gets that $\det(A_1, \dots, A_{n-3}, X_0, X_0)_i = 0$ for all i . Hence X_0 vanishes. This proves that the form B is non-degenerate.

It remains to compute the signature of B . Remind that B depends on positive definite matrices A_1, \dots, A_{n-2} . The space of positive definite matrices is connected (indeed if A and B are positive definite then $tA + (1 - t)B$ is positive definite for $0 \leq t \leq 1$). The signature of a family of non-degenerate quadratic forms cannot jump. Hence it is constant. Thus we can assume that $A_1 = \dots = A_{n-2} = Id$. As in the case $n = 2$ it is sufficient to check that if $X \neq 0$ satisfies $B(X, Id) = 0$ then $B(X, X) < 0$. Again we can assume that X is diagonal,

$$X = \begin{bmatrix} x_1 & & 0 \\ & \ddots & \\ 0 & & x_n \end{bmatrix}.$$

The condition $B(X, Id) = 0$ means that $\sum_{i=1}^n x_i = 0$. Also it is easy to see that

$$\kappa \cdot B(X, X) = 2 \sum_{i < j} x_i x_j,$$

where κ is a positive normalization constant. But

$$2 \sum_{i < j} x_i x_j = \left(\sum_i x_i \right)^2 - \sum_i x_i^2 = - \sum_i x_i^2 < 0.$$

The theorem is proved. Q.E.D.

1.2 Dieudonné determinant.

We will remind the construction of the Dieudonné determinant referring for the details and proofs to [2]. Also we will prove some properties of it which will be used in the subsequent sections of the paper. Intuitively the Dieudonné determinant of an arbitrary quaternionic matrix has the same algebraic and analytic properties as the *absolute value* of the usual determinant of real or complex matrices. First let us discuss purely algebraic construction.

Let F be an infinite field, not necessarily commutative. Let $M_n(F)$ denote the ring of $n \times n$ - matrices with coefficients in F . Let $GL_n(F)$ denote the group of invertible $n \times n$ - matrices. By an *elementary* matrix one calls a matrix which has units on the diagonal and at most one non-zero element out of the diagonal. Let E denote the subgroup of $GL_n(F)$ generated by all elementary matrices. Set also $F_{ab}^* := F^*/[F^*, F^*]$ the abelinization of the multiplicative group of F (here F^* denotes the multiplicative group of F , and $[F^*, F^*]$ denotes its commutator subgroup).

1.2.1. Theorem (Dieudonné). *Let $n \geq 2$. The group E is normal subgroup of $GL_n(F)$. For the quotient- group $GL_n(F)/E$ there exists a natural isomorphism $D : GL_n(F)/E \longrightarrow F_{ab}^*$.*

This isomorphism D is uniquely defined by the property that for any non-degenerate diagonal matrix $X = \begin{bmatrix} x_1 & & 0 \\ & \ddots & \\ 0 & & x_n \end{bmatrix}$, $D(X) = \prod_i x_i \text{ mod } [F^, F^*]$.*

1.2.2. Definition (Dieudonné determinant). The Dieudonné determinant is a map

$$D : M_n(F) \longrightarrow F_{ab}^* \cup \{0\}$$

defined as follows: if X is an invertible matrix then $D(X)$ is as in Theorem 1.2.1; if X is not invertible then $D(X) := 0$.

Note also that it is convenient to define the Dieudonné determinant of elements of F , i.e. 1×1 - matrices, as $D(0) = 0$ and for $x \neq 0$ as $D(x) := x \text{ mod } [F^*, F^*]$.

Let us state some basic general properties of the Dieudonné determinant. For the proofs we again refer to [2].

1.2.3. Theorem. (i) $D(Id) = 1$.

(ii) For $X, Y \in M_n(F)$

$$D(XY) = D(X)D(Y).$$

(iii) For any block- matrix $A = \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}$ with X, Y being square matrices

$$D(A) = D(X)D(Y).$$

(iv) If one interchanges two rows or two columns of the matrix then the Dieudonné determinant is multiplied by $-1 \bmod k_{ab}^*$.

Now let us consider in more details the case of quaternionic field $F = \mathbb{H}$. The commutator subgroup $[\mathbb{H}^*, \mathbb{H}^*]$ coincides with the subgroup of quaternions of absolute value 1. Thus we can identify \mathbb{H}_{ab}^* with the multiplicative group $\mathbb{R}_{>0}$ by

$$q \bmod [\mathbb{H}^*, \mathbb{H}^*] \leftrightarrow |q| := \sqrt{q\bar{q}}.$$

So in the quaternionic case the Dieudonné determinant maps

$$D : M_n(\mathbb{H}) \longrightarrow \mathbb{R}_{\geq 0}.$$

In the rest of the paper we will denote by $D(X)$ the Dieudonné determinant of quaternionic matrix X , and by $\det(X)$ the Moore determinant of hyperhermitian matrix X .

1.2.4. Theorem. (i) For any complex $n \times n$ - matrix X considered as quaternionic matrix the Dieudonné determinant $D(X)$ is equal to the absolute value of the usual determinant of X .

(ii) Let X be a quaternionic hyperhermitian $n \times n$ - matrix. Then its Dieudonné determinant $D(X)$ is equal to the absolute value of its Moore determinant $|\det(X)|$.

(iii) For any X

$$D(X) = D(X^t) = D(X^*),$$

where X^t and X^* denote the transposed and quaternionic conjugate matrices respectively.

For any $n \times n$ - matrix X and any subsets $I, J \subset \{1, \dots, n\}$ let us denote by $M_{I,J}(X)$ the matrix obtained from X by erasing the rows with indexes not in I and columns with indexes not in J . The following result is a weakened version of usual formula of the decomposition of the determinant with respect to row. Note that this result is satisfied for the absolute value of complex matrices.

1.2.5. Theorem. Let $A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \dots & \dots & \dots \\ a_{n1} & \dots & a_{nn} \end{bmatrix}$ be a quaternionic matrix.

Then

$$D(A) \leq \sum_{i=1}^n |a_{1i}| D(M_{1i}).$$

Similar inequalities hold for any other row or column.

Proof. From Theorem 1.2.3 it follows that

$$D \left(\left[\begin{array}{c|ccc} a & 0 & \dots & 0 \\ \hline * & & & \\ \vdots & & B & \\ * & & & \end{array} \right] \right) = |a| D(B).$$

Hence to prove the statement it is sufficient to show that the Dieudonné determinant is subadditive with respect to the first row; namely if the matrices A, A', A'' are such that the first row of A is the sum of first rows of A' and A'' and all the other rows are the same, then $D(A) \leq D(A') + D(A'')$. But the Dieudonné determinant has the following property over arbitrary (non-commutative) field k ([2], Thm. 4.5):

$$D(A) \subset D(A') + D(A''),$$

where the inclusion and addition are understood in the sense of conjugacy classes modulo $[k_{ab}^*, k_{ab}^*]$. But under our identification of H_{ab}^* with $\mathbb{R}_{>0}$ the last inclusion implies the desired inequality $D(A) \leq D(A') + D(A'')$. Q.E.D.

The next two propositions will be used in the sequel. It will be convenient to introduce the following notation. Set $M'_{IJ}(A) := M_{\{1, \dots, n\} - I, \{1, \dots, n\} - J}(A)$, i.e. it denotes the minor which stays on the intersection of the rows with indexes from I and columns with indexes from J .

1.2.6. Proposition. Let A be hyperhermitian non-negative definite $n \times n$ -matrix. Fix an integer k , $1 \leq k \leq n$ and two subsets $I, J \subset \{1, \dots, n\}$ of cardinality k . Then

$$2D(M'_{IJ}(A)) \leq D(M'_{II}(A)) + D(M'_{JJ}(A)).$$

Proof. For simplicity of the notation and without loss of generality we may assume that $I \cup J = \{1, \dots, n\}$, $I = \{1, \dots, k\}$, and $J = \{n - k + 1, \dots, n\}$.

First let us reduce to the case $I \cap J = \emptyset$. We have

$$A = \left[\begin{array}{c|c|c} * & * & * \\ \hline * & M'_{I \cap J, I \cap J}(A) & * \\ \hline * & * & * \end{array} \right].$$

For generic matrix A the (hyperhermitian) minor $M'_{I \cap J, I \cap J}(A)$ is invertible. Then by Claim 1.1.7 that one can choose an invertible matrix U_0 such that

$$U_0^* M'_{I \cap J, I \cap J}(A) U_0 = Id. \text{ Let } U = \left[\begin{array}{c|c|c} Id & 0 & 0 \\ \hline 0 & U_0 & 0 \\ \hline 0 & 0 & Id \end{array} \right]. \text{ Consider matrix } A_1 :=$$

$U^* A U$. Clearly $D(M'_{II}(A_1)) = D(M'_{II}(A)) D(U)^2$, and similarly for M'_{IJ} and M'_{JJ} . Hence replacing A by A_1 we may assume that $M'_{I \cap J, I \cap J}(A) = Id$. Thus A has the form

$$A = \left[\begin{array}{c|c|c} * & X & * \\ \hline X^* & Id & Y^* \\ \hline * & Y & * \end{array} \right].$$

$$\text{Set } V = \left[\begin{array}{c|c|c} Id & -X & 0 \\ \hline 0 & Id & 0 \\ \hline 0 & -Y & Id \end{array} \right]. \text{ Consider}$$

$$A_2 := V A V^* = \left[\begin{array}{c|c|c} P & 0 & R \\ \hline 0 & Id & 0 \\ \hline R^* & 0 & Q \end{array} \right].$$

Here P and Q are hyperhermitian matrices. Then A_2 has the same Dieudonné determinants of the minors M'_{II} , M'_{IJ} , M'_{JJ} as A . Hence we may replace A by

$$A_2, \text{ and we will denote it by the same letter } A. \text{ Then } M'_{II}(A) = \begin{bmatrix} P & 0 \\ 0 & Id \end{bmatrix},$$

$$M'_{IJ}(A) = \begin{bmatrix} 0 & R \\ Id & 0 \end{bmatrix}, M'_{JJ}(A) = \begin{bmatrix} Id & 0 \\ 0 & Q \end{bmatrix}. \text{ So one has to show that}$$

$$2D(R) \leq D(P) + D(Q).$$

This inequality is the statement of the proposition for the matrix $\tilde{A} := \begin{bmatrix} P & R \\ R^* & Q \end{bmatrix}$ which is also hyperhermitian and positive definite since A is.

Replacing \tilde{A} by the matrix $\begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix} \tilde{A} \begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix}^*$ with $U_1, U_2 \in Sp(k)$ one can assume that the matrices P and Q are diagonal.

Fix now some $U, V \in Sp(k)$ (the choice of them will be clear later). Let $T := \begin{bmatrix} P^{1/2}UP^{-1/2} & 0 \\ 0 & Q^{1/2}VQ^{-1/2} \end{bmatrix}$. Then

$$T\tilde{A}T^* = \begin{bmatrix} P & R_1 \\ R_1^* & Q \end{bmatrix},$$

where $R_1 = P^{1/2}U(P^{-1/2}RQ^{-1/2})V^*Q^{1/2}$. Note that $D(R_1) = D(R)$. Since P and Q are diagonal, by a choice of $U, V \in Sp(k)$ one can make the matrix R_1 diagonal.

Finally we are reduced to the hyperhermitian non-negative definite matrix

$$A \text{ of the form } A = \left[\begin{array}{cc|cc} \lambda_1 & 0 & \nu_1 & 0 \\ & \ddots & & \ddots \\ 0 & \lambda_k & 0 & \nu_k \\ \hline \bar{\nu}_1 & 0 & \mu_1 & 0 \\ & \ddots & & \ddots \\ 0 & \bar{\nu}_k & 0 & \mu_k \end{array} \right]. \text{ We have to show that}$$

$$2 \prod_{i=1}^k |\nu_i| \leq \prod_{i=1}^k |\lambda_i| + \prod_{i=1}^k |\mu_i|.$$

Consider the 2×2 - matrix $\begin{bmatrix} \lambda_i & \nu_i \\ \bar{\nu}_i & \mu_i \end{bmatrix}$ which is clearly non-negative definite.

Take a vector $\begin{pmatrix} 1 \\ t \cdot q \end{pmatrix}$ for any $t \in \mathbb{R}$ and any quaternion q of norm 1. Applying that matrix to this vector we get

$$\lambda_i + t^2 \mu_i + 2t \operatorname{Re}(\nu_i q) \geq 0.$$

Hence $|\nu_i| \leq \sqrt{\lambda_i \mu_i}$. Then

$$2 \prod_i |\nu_i| \leq 2 \sqrt{\prod_i |\lambda_i| \cdot \prod_i |\mu_i|} \leq \prod_i |\lambda_i| + \prod_i |\mu_i|.$$

Q.E.D.

1.2.7. Proposition. *Let $A = (a_{ij}), B$ be $n \times n$ - hyperhermitian matrices. Then the mixed discriminant satisfies*

$$|\det(A, \underbrace{B, \dots, B}_{n-1 \text{ times}})| \leq c_n \cdot \max |a_{ij}| \cdot \left(\sum_{|I|, |J|=n-1} D(B_{IJ}) \right),$$

where c_n is a constant depending on n only.

Proof. Since $\det(A, B, \dots, B)$ is linear in A it is sufficient to prove the inequality in the following two cases.

$$1) A = \begin{bmatrix} 1 & & 0 \\ & 0 & \\ & & \ddots \\ 0 & & & 0 \end{bmatrix};$$

$$2) A = \left[\begin{array}{cc|c} 0 & q & 0 \\ \bar{q} & 0 & \\ \hline & & 0 \\ & & \ddots \\ 0 & & & 0 \end{array} \right]. \text{ The first case follows from Proposition}$$

1.1.11. Let us consider the second case. Replacing A by the matrix

$$\begin{bmatrix} \frac{\bar{q}}{|q|} & & 0 \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{bmatrix} A \begin{bmatrix} \frac{q}{|q|} & & 0 \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{bmatrix} = \left[\begin{array}{cc|c} 0 & |q| & 0 \\ |q| & 0 & \\ \hline & & 0 \\ & & \ddots \\ 0 & & & 0 \end{array} \right]$$

$$\text{we can assume that } A = \left[\begin{array}{cc|c} 0 & 1 & 0 \\ 1 & 0 & \\ \hline & & 0 \\ & & \ddots \\ 0 & & & 0 \end{array} \right]. \text{ Let } B = \left[\begin{array}{c|c} P & R \\ \hline R^* & Q \end{array} \right]. \text{ Here}$$

P and Q are hyperhermitian matrices of sizes 2×2 and $(n-2) \times (n-2)$ respectively.

Claim.

$$\det(A, B, \dots, B) \leq M'_{\{2,3,\dots,n\},\{1,3,\dots,n\}}(B) + M'_{\{1,3,\dots,n\},\{2,3,\dots,n\}}(B).$$

We may also assume that Q is invertible. Set $S := \left[\begin{array}{c|c} Id & Id - RQ^{-1} \\ \hline 0 & Id \end{array} \right]$. Consider

$$B_1 := SBS^* = \begin{bmatrix} * & 0 \\ 0 & * \end{bmatrix}.$$

Note also that $S^*AS = A$. It is easy to see that

$$M'_{\{2,3,\dots,n\},\{1,3,\dots,n\}}(B_1) = M'_{\{2,3,\dots,n\},\{1,3,\dots,n\}}(B) \text{ and}$$

$$M'_{\{1,3,\dots,n\},\{2,3,\dots,n\}}(B_1) = M'_{\{1,3,\dots,n\},\{2,3,\dots,n\}}(B).$$

Hence it is sufficient to prove the claim under assumption $R = 0$, i.e. $B = \begin{bmatrix} P & 0 \\ 0 & Q \end{bmatrix}$. Then clearly

$$\det(A, B, \dots, B) = \det\left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, P\right) \cdot \det Q.$$

If $P = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$ then

$$|\det\left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, P\right)| = \operatorname{Re}(b_{12}) \leq |b_{11}| + |b_{22}|,$$

where the last inequality follows from Proposition 1.2.6. Proposition 1.2.7 is proved. Q.E.D.

From Propositions 1.2.6 and 1.2.7 one can easily deduce

1.2.8. Proposition. *Let $A = (a_{ij})$ be a hyperhermitian matrix and B_1, \dots, B_{n-1} be non-negative definite hyperhermitian matrices. Then*

$$|\det(A, B_1, \dots, B_{n-1})| \leq c_n \cdot \max_{i,j} |a_{ij}| \cdot \sum_{1 \leq i_1, \dots, i_{n-1} \leq n-1} \det(B_{i_1}, \dots, B_{i_{n-1}}),$$

where c_n is a constant depending on n only.

2 Plurisubharmonic functions of quaternionic variables.

In this part we will develop a basic theory of plurisubharmonic functions of quaternionic variables.

First let us remind few standard notions. Below Ω will denote an open domain. As usual we will denote by $C^k(\Omega)$ the class of k times continuously differentiable functions on Ω , and by $C_0^k(\Omega)$ the class of k times continuously differentiable functions on Ω with compact support.

2.1.1. Definition. A real valued continuous function $f : \Omega \subset \mathbb{R}^m \longrightarrow \mathbb{R}$ is called *subharmonic* if $\Delta f \geq 0$, where $\Delta = \sum_{i=1}^m \frac{\partial^2}{\partial x_i^2}$ is the Laplacian.

Note that if f is not smooth then Δf is understood in the sence of distributions.

2.1.2. Definition. A real valued continuous function

$$f : \Omega \subset \mathbb{R}^n \longrightarrow \mathbb{R}$$

is called *convex* if its restriction to any (real) line is subharmonic.

2.1.3. Definition. A real valued continuous function

$$f : \Omega \subset \mathbb{C}^n \longrightarrow \mathbb{R}$$

is called *plurisubharmonic* if its restriction to any *complex* line is subharmonic.

Now let us introduce a new definition.

2.1.4. Definition. A real valued continuous function

$$f : \Omega \subset \mathbb{H}^n \longrightarrow \mathbb{R}$$

is called *quaternionic plurisubharmonic* if its restriction to any *quaternionic* line is subharmonic.

2.1.5. Example. 1) Any convex function on \mathbb{H}^n is quaternionic plurisubharmonic .

2) Fix on \mathbb{H}^n one of the complex structures compatible with the quaternionic structure; say, let us fix i . Let f be a continuous function which is plurisubharmonic with respect to this complex structure in the sence of Definition 2.1.3. It is easy to see that f is plurisubharmonic in the quaternionic sence.

Let q be a quaternionic coordinate,

$$q = t + ix + jy + kz,$$

where t, x, y, z are real numbers. Consider the following operators defined on the class of smooth \mathbb{H} -valued functions of the variable $q \in \mathbb{H}$:

$$\frac{\partial}{\partial \bar{q}} f := \frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}, \text{ and}$$

$$\frac{\partial}{\partial q} f := \overline{\frac{\partial}{\partial \bar{q}} \bar{f}} = \frac{\partial f}{\partial t} - \frac{\partial f}{\partial x} i - \frac{\partial f}{\partial y} j - \frac{\partial f}{\partial z} k.$$

Note that $\frac{\partial}{\partial \bar{q}}$ is called Cauchy-Riemann-Fueter operator. It is easy to see that $\frac{\partial}{\partial \bar{q}}$ and $\frac{\partial}{\partial q}$ commute, and if f is a *real valued* function then

$$\frac{\partial}{\partial \bar{q}} \frac{\partial}{\partial q} f = \Delta f = \left(\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) f.$$

For any real valued C^2 -smooth function f the matrix $(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})_{i,j=1}^n$ is obviously hyperhermitian. For brevity we will use the following notation:

$$\det(f_1, \dots, f_n) := \det \left(\left(\frac{\partial^2 f_1}{\partial q_i \partial \bar{q}_j} \right), \dots, \left(\frac{\partial^2 f_n}{\partial q_i \partial \bar{q}_j} \right) \right),$$

where \det denotes the mixed discriminant of hyperhermitian matrices (see Definition 1.1.14). One can easily check the following identities.

Claim. (i) Let $f : \mathbb{H}^n \rightarrow \mathbb{H}$ be a smooth function. Then for any \mathbb{H} -linear transformation A of \mathbb{H}^n (as right \mathbb{H} -vector space) one has the identities

$$\left(\frac{\partial^2 f(Aq)}{\partial \bar{q}_i \partial q_j} \right) = A^* \left(\frac{\partial^2 f}{\partial \bar{q}_i \partial q_j}(AQ) \right) A.$$

(ii) If, in addition, f is real valued then for any \mathbb{H} -linear transformation A of \mathbb{H}^n and any quaternion a with $|a| = 1$

$$\left(\frac{\partial^2 f(A(q \cdot a))}{\partial \bar{q}_i \partial q_j} \right) = A^* \left(\frac{\partial^2 f}{\partial \bar{q}_i \partial q_j}(AQ) \right) A.$$

2.1.6. Proposition. *A real valued twice continuously differentiable function f on the domain $\Omega \subset \mathbb{H}^n$ is quaternionic plurisubharmonic if and only if at every point $q \in \Omega$ the matrix $(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})(q)$ is non-negative definite.*

The proof of this proposition is streightforward. The following lemma will be useful in the sequel.

2.1.7. Lemma. *Let f_0, f_1, \dots, f_n be real valued compactly supported sufficiently smooth functions on \mathbb{H}^n . The $(n+1)$ - linear functional*

$$L(f_0, f_1, \dots, f_n) := \int_{\mathbb{H}^n} f_0(q) \cdot \det(f_1, \dots, f_n)(q) dq$$

is symmetric with respect to all f_0, f_1, \dots, f_n .

Proof. Note that L is symmetric with respect to the last n arguments. Thus it is sufficient to check that

$$L(f_0, f_1, f_2, \dots, f_n) = L(f_1, f_0, f_2, \dots, f_n) \quad (4)$$

for any smooth compactly supported functions f_0, f_1, \dots, f_n . The last expression makes sense if f_1 is a generalized function. Since linear combinations of delta- functions of points δ_q are dense in the space of all the generalized functions it is sufficient to prove (4) for $f_0 = \delta_0$, namely

$$(\det(f_1, \dots, f_n))|_{q=0} = \int_{\mathbb{H}^n} f_1(q) \det(\delta_0, f_2, \dots, f_n). \quad (5)$$

Clearly the right hand side in equation (5) depends only on derivatives at 0 of f_1, \dots, f_n up to order 2. Consider the terms of the Taylor series of f_1 at 0:

$$f_1(q) = g(q) + h(q) + O(|q|^2),$$

where g is a polynomial of degree one, and h is a quadratic term. So it is sufficient to prove the following two statements:

Case 1.

$$L(h, \delta_0, f_2, \dots, f_n) = \det(h, f_2, \dots, f_n)|_{q=0} \quad (6)$$

for any smooth compactly supported function h which is equal to a homogeneous polynomial of degree 2 in a neighbourhood of 0, and for any smooth compactly supported functions f_2, \dots, f_n .

Case 2.

$$L(g, \delta_0, f_2, \dots, f_n) = 0 \quad (7)$$

for any smooth compactly supported function g which is equal to a polynomial of degree 1 in a neighbourhood of 0, and for any smooth compactly supported functions f_2, \dots, f_n .

Let us consider Case 1. If we write down the formula for $L(h, \delta_0, f_2, \dots, f_n)$ as a polynomial in $\frac{\partial^2 f_k}{\partial t_i \partial t_j}$, $\frac{\partial^2 f_k}{\partial t_i \partial x_j}$ etc. and in $\frac{\partial^2 \delta_0}{\partial t_i \partial t_j}$, $\frac{\partial^2 \delta_0}{\partial t_i \partial x_j}$ etc. then we see that the derivatives of δ_0 enter at each monomial only once because of linearity of L with respect to each argument. For example consider a monomial containing $\frac{\partial^2 \delta_0}{\partial t_i \partial t_j}$. Let it be $\int_{\mathbb{H}^n} h \cdot \frac{\partial^2 \delta_0}{\partial t_i \partial t_j} \cdot \partial^2 f_2 \cdot \dots \cdot \partial^2 f_n$, where $\partial^2 f_k$ denotes certain partial derivative of order 2 of f_k . But

$$\begin{aligned} \int_{\mathbb{H}^n} h \cdot \frac{\partial^2 \delta_0}{\partial t_i \partial t_j} \cdot \partial^2 f_2 \cdot \dots \cdot \partial^2 f_n &= \frac{\partial^2}{\partial t_i \partial t_j} (h \cdot \partial^2 f_2 \cdot \dots \cdot \partial^2 f_n)|_{q=0} = \\ &= \frac{\partial^2 h}{\partial t_i \partial t_j}(0) \cdot \partial^2 f_2(0) \cdot \dots \cdot \partial^2 f_n(0), \end{aligned}$$

where the last equality is satisfied since the first derivatives of h at 0 vanish. Thus in each monomial the term $h \cdot \frac{\partial^2 \delta_0}{\partial t_i \partial t_j}$ is just replaced by $\frac{\partial^2 h}{\partial t_i \partial t_j}(0)$. Hence the final expression is $\det(h, f_2, \dots, f_n)|_{q=0}$. This proves the first case.

Let us prove Case 2. It is convenient to prove a more general statement.

Claim. *Let U be a fixed neighbourhood of the origin 0. Let g be any smooth compactly supported function which is equal to a polynomial of degree 1 inside U . Let f_1 be a generalized function with support contained in U . Let f_2, \dots, f_n be smooth compactly supported functions.*

Then

$$\int_{\mathbb{H}^n} g \det(f_1, f_2, \dots, f_n) = 0.$$

The proof of the claim will be by induction in n . If $n = 1$ then using selfadjointness of the Laplacian one gets:

$$\int_{\mathbb{H}} g \Delta f_1 = \int_{\mathbb{H}} \Delta g \cdot f_1 = \int_U \Delta g \cdot f_1 = 0.$$

Assume that $n > 1$. It is well known (see Appendix) that the linear combinations of delta- functions of quaternionic hyperplanes are dense in the

space of all generalized functions (this fact is equivalent to the injectivity of the Radon transform with respect to quaternionic hyperplanes). Hence it is sufficient to prove the claim for $f_1 = \delta_L$, where L is the hyperplane $\{q_1 = 0\}$.

Since δ_L is invariant with respect to translations in directions q_2, \dots, q_n then $\frac{\partial^2 \delta_L}{\partial q_i \partial \bar{q}_j} = 0$ unless $i = j = 1$. Thus

$$\frac{\partial^2 \delta_L}{\partial q_i \partial \bar{q}_j} = \left[\begin{array}{c|ccc} \Delta_1 \delta_L & 0 & \dots & 0 \\ \hline 0 & & & \\ \vdots & & 0 & \\ 0 & & & \end{array} \right],$$

where Δ_1 denotes the Laplacian with respect to the first coordinate: $\Delta_1 = \frac{\partial^2}{\partial t_1^2} + \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial y_1^2} + \frac{\partial^2}{\partial z_1^2}$. Using Proposition 1.1.11 it is easy to see that

$$c \cdot \det(\delta_L, f_2, \dots, f_n) = \Delta_1 \delta_L \cdot \det(B_2, \dots, B_n),$$

where c is a positive normalizing constant, and B_k denotes the $(n-1) \times (n-1)$ -matrix $(\det \frac{\partial^2 f_k}{\partial q_i \partial \bar{q}_j})_{i,j=2}^n$. Then

$$\int_{\mathbb{H}^n} g \cdot \det(\delta_L, f_2, \dots, f_n) = \int_{\mathbb{H}^n} g \cdot \Delta_1 \delta_L \cdot \det(B_2, \dots, B_n).$$

Clearly the last expression depends only on the 2-jets of g, f_2, \dots, f_n in the direction q_1 . Thus we may assume that the functions f_k are of the form

$$f_k(q_1, q_2, \dots, q_n) = p_k(q_1) \cdot f'_k(q_2, \dots, q_n),$$

where $p_k(q_1)$ are polynomials (of degree at most 2) depending only on t_1, x_1, y_1, z_1 , and f'_k are smooth compactly supported functions depending only on q_2, \dots, q_n .

Since $\deg g \leq 1$ we may assume (by linearity) that either $g(q_1, q_2, \dots, q_n) = g(q_1)$ or $g(q_1, q_2, \dots, q_n) = g(q_2, \dots, q_n)$. In the first case

$$\begin{aligned} \int_{\mathbb{H}^n} g \cdot \Delta_1 \delta_L \cdot \det(B_2, \dots, B_n) = \\ \Delta_1 (g(q_1) \cdot p_2(q_1) \cdot \dots \cdot p_n(q_1))|_{q_1=0} \cdot \int_L \det(g, B'_2, \dots, B'_n), \end{aligned}$$

where B'_k denotes the matrix $(\frac{\partial^2 f'_k}{\partial q_i \partial \bar{q}_j})_{i,j=2}^n$. The last integral vanishes by the induction assumption.

Now consider the second case $g(q_1, q_2, \dots, q_n) = g(q_2, \dots, q_n)$. We have

$$\int_{\mathbb{H}^n} g \cdot \Delta_1 \delta_L \cdot \det(B_2, \dots, B_k) = \Delta_1(p_2 \dots p_n)|_{q_1=0} \int_L \det(g, B_2, \dots, B_n).$$

Again the last expression vanishes by the induction assumption. Thus our claim, and hence Proposition 2.1.6, are proved. Q.E.D.

The next result is again a quaternionic analogue of the corresponding property of convex functions and complex plurisubharmonic functions. We adopt the arguments of Rauch and Taylor [24] (see also [4]).

2.1.8. Proposition. *Let $\Omega \subset \mathbb{H}^n$ be an open domain. Assume that a sequence $\{f_N\}$ of twice continuously differentiable quaternionic plurisubharmonic functions converges uniformly on compact subsets of Ω to a twice continuously differentiable function f . Then f is also quaternionic plurisubharmonic, and for every function $\phi \in C_0^\infty(\Omega)$*

$$\int_{\Omega} \phi \cdot \det\left(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j}\right) \text{ tends to } \int_{\Omega} \phi \cdot \det\left(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j}\right) \text{ as } N \longrightarrow \infty.$$

We will need a lemma. But first let us introduce a notation. For subsets $I, J \subset \{1, \dots, n\}$ and a function g let us denote by $M'_{IJ}(g)$ the matrix which stays on the intersection of rows with indexes from I and columns with indexes from J in the matrix $\det(\frac{\partial^2 g}{\partial q_i \partial \bar{q}_j})_{i,j=1}^n$. Also for a set U and a function g on it let us denote by $\|g\|_{C^0(U)} := \sup_{q \in U} |g(q)|$, and by $\|g\|_{C^k(U)}$ the maximum of $C^0(U)$ - norms of all partial derivatives of g up to order k . Below we will denote for brevity $\det(\frac{\partial^2 g}{\partial q_i \partial \bar{q}_j})_{i,j=1}^n$ by $\det(g)$.

2.1.9. Lemma. *Let I, J be subsets of $\{1, \dots, n\}$ of cardinality k . Let f be a continuous function on Ω , and let g be a twice continuously differentiable quaternionic plurisubharmonic function on a domain $\Omega \subset \mathbb{H}^n$. Let K be a compact subset of Ω , and let U be a neighbourhood of K in Ω . Then*

$$\left| \int_K f \cdot D(M'_{IJ}(g)) \right| \leq C(U) \|f\|_{C^0(U)} \|g\|_{C^0(U)}^k,$$

where $C(U)$ is a constant depending on U only.

Proof. Since g is plurisubharmonic, Proposition 1.2.6 implies the estimate $D(M'_{IJ}(g)) \leq D(M'_{II}(g)) + D(M'_{JJ}(g))$. Hence

$$\left| \int_K f \cdot D(M'_{IJ}(g)) \right| \leq \|f\|_{C^0(K)} \cdot \int_K (D(M'_{II}(g)) + D(M'_{JJ}(g))).$$

It remains to prove that for any subset $I \subset \{1, \dots, n\}$ of cardinality k

$$\int_K \det(M'_{II}(g)) \leq C(U) \|g\|_{C^0(U)}^k.$$

Let us prove this inequality by induction in k . For $k = 0$ the statement is trivial. Assume that $k > 0$. Let us fix a non-negative function $\gamma \in C_0^\infty(\Omega)$ such that $\gamma|_K \equiv 1$ and γ vanishes on $\Omega - U$. Then using Lemma 2.1.7

$$\int_K \det M'_{II}(g) \leq \int_{\mathbb{H}^n} \gamma \cdot \det M'_{II}(g) = \int_{\mathbb{H}^n} g \cdot \det_I(\gamma, g, \dots, g),$$

where \det_I denotes the mixed discriminant of matrices of order $|I|$. By Proposition 1.2.7 the last expression is at most

$$\|g\|_{C^0(U)} \cdot \|\gamma\|_{C^2(U)} \sum_{S,T} \int_U D_{ST}(g),$$

where the sum extends over all subsets S, T of I of cardinality $k - 1$. Again by Proposition 1.2.6

$$\int_U D_{ST}(g) \leq \int_U (\det_S(g) + \det_T(g)).$$

Now the estimate follows by the assumption of induction. Q.E.D.

Now let us prove Proposition 2.1.8. First let us show that the limit f is plurisubharmonic. Indeed the restriction of f to any quaternionic line is subharmonic since the uniform limit of subharmonic functions is subharmonic.

Let us prove the second part of Proposition 2.1.8. Fix $\varepsilon > 0$, and a compact neighborhood U of K . Let us choose a function $\psi \in C_0^\infty(\Omega)$ such that $\|\phi - \psi\|_{C^0(U)} \leq \varepsilon$. We have

$$\begin{aligned} & \left| \int_K \psi(\det(f_N) - \det(f)) - \int_K \phi(\det(f_N) - \det(f)) \right| \leq \\ & \left| \int_K (\psi - \phi) \det(f_N) \right| + \left| \int_K (\psi - \phi) \det(f) \right| \leq \\ & C(U) (\|f_N\|_{C^0(U)}^n + \|f\|_{C^0(U)}^n) \cdot \varepsilon, \end{aligned}$$

where the last inequality follows from Lemma 2.1.9. The last expression can be estimated for large N by $3C(U) \|f\|_{C^0(U)}^n \cdot \varepsilon$. Thus it is sufficient to prove that

$$\int_\Omega \psi \cdot \det(f_N) \text{ tends to } \int_\Omega \psi \cdot \det(f) \text{ as } N \longrightarrow \infty.$$

We have

$$\begin{aligned}
\left| \int_{\Omega} \psi \cdot (\det(f_N) - \det(f)) \right| &= \left| \int_{\Omega} \sum_{i=0}^{n-1} \psi \cdot \det(\underbrace{f_N, \dots, f_N}_i \text{ times}, f_N - f, \underbrace{f, \dots, f}_{n-i-1} \text{ times}) \right| = \\
&= \left| \sum_{i=0}^{n-1} \int_{\Omega} (f_N - f) \det(\underbrace{f_N, \dots, f_N}_i, \psi, \underbrace{f, \dots, f}_{n-i-1}) \right| \leq \\
&\leq C \cdot \|f_N - f\|_{C^0(\Omega)} \|\psi\|_{C^2(\Omega)} \sum_{i=0}^{n-1} \int_{\text{supp } \psi} \det_{n-1}(\underbrace{f_N, \dots, f_N}_i, \underbrace{f, \dots, f}_{n-i-1}) \quad (8)
\end{aligned}$$

by Proposition 1.2.8 (here we have used the fact that the functions f_N and f are plurisubharmonic). Now let us estimate the last expression.

$$\begin{aligned}
\int_{\text{supp } \psi} \det_{n-1}(f_N, \dots, f_N, f, \dots, f) &\leq \int_{\text{supp } \psi} \det_{n-1}(\underbrace{f + f_N, \dots, f + f_N}_{n-1}) \leq \\
&\leq C' \|f + f_N\|_{C^0}^{n-1},
\end{aligned}$$

where the last inequality holds by Lemma 2.1.9. Hence the expression (8) tends to 0 as $N \rightarrow \infty$. This proves Proposition 2.1.8. Q.E.D.

Now let us study *continuous* quaternionic plurisubharmonic functions which are not necessarily smooth. For every continuous plurisubharmonic function f we will define a non-negative measure such that if f is smooth it coincides with $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})$. To do it let us observe first of all that any continuous plurisubharmonic function f on a domain $\Omega \subset \mathbb{H}^n$ can be approximated by C^∞ -smooth plurisubharmonic functions uniformly on compact subsets of Ω . (To see it consider the convolution of f with a delta-sequence of non-negative C^∞ -smooth functions. Each such convolution is infinitely smooth and plurisubharmonic). The next theorem is first main result of this section; it provides the definition of the measure $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})$ for any continuous plurisubharmonic function f .

2.1.10. Theorem. *Let f be a continuous quaternionic plurisubharmonic function on a domain Ω . Let $\{f_N\}$ be a sequence of twice continuously differentiable plurisubharmonic functions converging to f uniformly on compact subsets of Ω . Then $\det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})$ weakly converges to a non-negative measure on Ω . This measure depends only on f and not on the choice of an approximating sequence $\{f_N\}$.*

This measure will be denoted by $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})$.

Proof. By Lemma 2.1.9 one sees that for any compact subset $K \subset \Omega$ the sequence of measures $\det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})|_K$ is bounded. Thus it is sufficient to show that for any continuous compactly supported function ϕ the sequence $\int_{\Omega} \phi \cdot \det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})$ is a Cauchy sequence. Let us fix $\varepsilon > 0$, and a function $\psi \in C_0^\infty(\Omega)$ such that $\|\phi - \psi\|_{C^0(\Omega)} < \varepsilon$. Let us also fix an arbitrary compact subset $K \subset \Omega$ and a compact neighbourhood U of K in Ω . As in the proof of Proposition 2.1.8 we have

$$\begin{aligned} & \left| \int_K \psi(\det(f_N) - \det(f_M)) - \int_K \phi(\det(f_N) - \det(f_M)) \right| \leq \\ & \left| \int_K (\psi - \phi)\det(f_N) \right| + \left| \int_K (\psi - \phi)\det(f_M) \right| \leq \\ & C(U)(\|f_N\|_{C^0(U)}^n + \|f_M\|_{C^0(U)}^n) \cdot \varepsilon, \end{aligned}$$

where the last inequality follows from Lemma 2.1.9. For large M and N the last expression can be estimated from above by $3C(U)\|f\|_{C^0(U)}^n \cdot \varepsilon$. Hence it is sufficient to prove that for any function $\psi \in C_0^\infty(\Omega)$ the sequence $\int_{\Omega} \psi \cdot \det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})$ is a Cauchy sequence. We have the following estimate exactly as in the inequality (8) (with f_M instead of f):

$$\begin{aligned} & \left| \int_{\Omega} \psi \cdot (\det(f_N) - \det(f_M)) \right| \leq \\ & C \cdot \|f_N - f_M\|_{C^0(\text{supp } \psi)} \|\psi\|_{C^2(\Omega)} \sum_{i=0}^{n-1} \int_{\text{supp } \psi} \det_{n-1}(\underbrace{f_N, \dots, f_N}_i, \underbrace{f_M, \dots, f_M}_{n-i-1}). \end{aligned}$$

and again as in the proof of Proposition 2.1.8 we get

$$\int_{\text{supp } \psi} \det_{n-1}(\underbrace{f_N, \dots, f_N}_i, \underbrace{f_M, \dots, f_M}_{n-i-1}) \leq C\|f_N + f_M\|_{C^0(\text{supp } \psi)}^{n-1} < C'.$$

This proves Theorem 2.1.10. Q.E.D.

The second main result of this section is as follows.

2.1.11. Theorem. *Let $\{f_N\}$ be sequence of continuous quaternionic plurisubharmonic function in a domain $\Omega \subset \mathbb{H}^n$. Assume that this sequence converges uniformly on compact subsets to a function f . Then f is continuous quaternionic plurisubharmonic function. Moreover the sequence of measures $\det(\frac{\partial^2 f_N}{\partial q_i \partial \bar{q}_j})$ weakly converges to the measure $\det(\frac{\partial^2 f}{\partial q_i \partial \bar{q}_j})$.*

Proof. The limit f is a plurisubharmonic function. Indeed the restriction of f to any quaternionic line is subharmonic as a uniform limit of subharmonic functions.

Let us prove the second part of the statement. The functions f_N can be approximated uniformly on compact subsets as good as we wish by smooth plurisubharmonic functions g_N such that the sequence g_N will converge uniformly on compact subsets to f . Then the result follows from previous Theorem 2.1.10. Q.E.D.

Appendix.

In this appendix we prove that the linear combinations of delta-functions of quaternionic hyperplanes in \mathbb{H}^n are dense in the space of distributions (this fact was needed in the proof of Lemma 2.1.7). By the Hahn- Banach theorem it is equivalent to the injectivity of the Radon transform over quaternionic hyperplanes. We beleave that the injectivity of quaternionic Radon transform is a well known fact, but we include the proof for completeness, since we could not find the reference.

Let us fix hyperhermitian metric on \mathbb{H}^n , i.e. a Euclidean metric such that for any two vectors $x, y \in \mathbb{H}^n$ and any quaternion a with $|a| = 1$

$$(x \cdot a, y \cdot a) = (x, y).$$

Let f be any smooth compactly supported function on \mathbb{H}^n . The quaternionic Radon transform of f is a function on the manifold of all affine quaternionic hyperplanes defined as

$$Rf(E) = \int_E f(q) dq,$$

where the integration is with respect the volume form on E defined by the metric.

Proposition. *The quaternionic Radon transform is injective.*

Proof. We will just present the inversion formula completely analogous to the complex Radon transform (see [14]). Let us fix the origin $0 \in \mathbb{H}^n$ for convenience. Let us denote by \mathcal{A} the manifold of affine quaternionic hyperplanes in \mathbb{H}^n . For any point $q \in \mathbb{H}^n$ let \mathcal{P}_q denotes the manifold of quaternionic hyperplanes passing through q . For $E \in \mathcal{A}$ let us denote by E^\perp the quaternionic line orthogonal to E and passing through the origin.

Let us define the operator

$$\mathcal{D} : C^\infty(\mathcal{A}) \longrightarrow C^\infty(\mathbb{H}^n)$$

as follows. Let $g \in C^\infty(\mathcal{A})$. Set

$$\mathcal{D}g(q) := \int_{E \in \mathcal{P}_q} (\Delta_{E^\perp})^{2(n-1)} g(E + w),$$

where Δ_{E^\perp} denotes the (4- dimensional) Laplacian with respect to $w \in E^\perp$, and the first integration is with respect the Haar measure on \mathcal{P}_q .

Claim. *For any smooth rapidly decreasing function f of \mathbb{H}^n*

$$\mathcal{D}(Rf) = c \cdot f,$$

where c is a non-zero constant.

It is sufficient to check this claim pointwise, say at 0. The operators R and \mathcal{D} commute with translations and the action of the group Sp_n . Then $\mathcal{D}(Rf)(0)$ defines a distribution invariant with respect to the action of Sp_n . Moreover it is easy to check that this distribution is homogeneous of degree $-4n$ (exactly as the delta- function at 0). It is easy to see that there is at most one dimensional space of Sp_n - invariant distributions homogeneous of degree $-4n$. Hence they must be proportional to the delta- function at 0. Thus $\mathcal{D}(Rf) = c \cdot f$ for some constant c . So see that $c \neq 0$ it is sufficient to check it by an explicite computation for the function $f(q) = \exp(-|q|^2/2)$. Q.E.D.

References

- [1] Alexandrov, A.D.; Die gemischte Diskriminanten und die gemischte Volumina. Math. Sbornik 3(1938), pp.227-251.
- [2] Artin, Emil; Geometric algebra. Interscience Publishers, Inc., New York-London, 1957.
- [3] Aslaksen, Helmer; Quaternionic determinants. Math. Intelligencer 18 (1996), no. 3, 57–65.
- [4] Aubin, Thierry; Nonlinear analysis on manifolds. Monge-Ampère equations. Grundlehren der Mathematischen Wissenschaften [Fundamental

- Principles of Mathematical Sciences], 252. Springer-Verlag, New York, 1982.
- [5] Barnard, R. W., Moore, E. H.; General Analysis, Part 1. Memoirs of the American Philosophical Society, 1935.
 - [6] Bedford, Eric; Taylor, B. A.; The Dirichlet problem for a complex Monge-Ampère equation. *Invent. Math.* 37 (1976), no. 1, 1–44.
 - [7] Chen, Long Xuan; Definition of determinant and Cramer solutions over the quaternion field. A Chinese summary appears in *Acta Math. Sinica* 35 (1992), no. 3, 432. *Acta Math. Sinica (N.S.)* 7 (1991), no. 2, 171–180.
 - [8] Chen, Long Xuan; Inverse matrix and properties of double determinant over quaternion field. *Sci. China Ser. A* 34 (1991), no. 5, 528–540.
 - [9] Dieudonné, Jean; Les déterminants sur un corps non commutatif. (French) *Bull. Soc. Math. France* 71, (1943). 27–45.
 - [10] Dyson, Freeman J.; Correlations between eigenvalues of a random matrix. *Comm. Math. Phys.* 19 1970 235–250.
 - [11] Dyson, Freeman J.; Quaternionic determinants. *Helv. Phys. Acta* 45, (1972), 289–302.
 - [12] Fueter, R.; Die Funktionentheorie der Differentialgleichungen $\Delta u = 0$ und $\Delta \Delta u = 0$ mit vier reellen Variablen. *Comment. Math. Helv.* 7 (1935), 307–330.
 - [13] Fueter, R.; Über die analytische Darstellung der regulären Functionen einer Quaternionenvariablen. *Comment. Math. Helv.* 8 (1936), 371–378.
 - [14] Gelfand, I. M.; Graev, M. I.; Vilenkin, N. Ya.; Generalized functions. Vol. 5. Integral geometry and representation theory. Translated from the Russian by Eugene Saletan. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1966 [1977].
 - [15] Henkin, Gennadi; Private communication.
 - [16] Jacobson, N.; Normal semi-linear transformations. *Amer. J. Math.* 61 (1939), 45–58.

- [17] Jacobson, N.; An application of E. H. Moore's determinant of a Hermitian matrix. Bull. Amer. Math. Soc. 45, (1939). 745–748.
- [18] Mehta, Madan Lal; Determinants of quaternionic matrices. J. Math. Phys. Sci. 8 (1974), 559–570.
- [19] Mehta, Madan Lal; Elements of matrix theory. Dehli Hindustan Pub. Corp., 1977.
- [20] Moore, E. H.; On the determinant of an hermitian matrix of quaternionic elements. Bull. Amer. Math. Soc. 28 (1922), 161–162.
- [21] Palamodov, Victor P.; Holomorphic synthesis of monogenic functions of several quaternionic variables. J. Anal. Math. 78 (1999), 177–204.
- [22] Pertici, Donato; Quaternion regular functions and domains of regularity. Boll. Un. Mat. Ital. B (7) 7 (1993), no. 4, 973–988.
- [23] Piccinni, Paolo; Dieudonné determinant and invariant real polynomials on $gl(n, \mathbb{H})$. Rend. Mat. (7) 2 (1982), no. 1, 31–45.
- [24] Rauch, Jeffrey; Taylor, B. A. The Dirichlet problem for the multidimensional Monge-Ampère equation. Rocky Mountain J. Math. 7 (1977), no. 2, 345–364.
- [25] Rosenberg, Jonathan; Algebraic K -theory and its applications. Graduate Texts in Mathematics, 147. Springer-Verlag, New York, 1994.
- [26] Sudbery, A.; Quaternionic analysis. Math. Proc. Cambridge Philos. Soc. 85 (1979), no. 2, 199–224.
- [27] Van Praag, Paul; Sur les déterminants des matrices quaternioniennes. (French) [On the determinants of quaternion matrices] Helv. Phys. Acta 62 (1989), no. 1, 42–46.
- [28] Van Praag, Paul; Sur la norme réduite du déterminant de Dieudonné des matrices quaternioniennes. (French) [On the reduced norm of the Dieudonné determinant of quaternion matrices] J. Algebra 136 (1991), no. 2, 265–274.