

Product formulas on a unitary group in three variables

LEI YANG

1. Introduction

In the present paper, we consider the eigenvalue problems which concern a differential equation

$$(1.1) \quad L_k f(z_1, z_2) = \lambda f(z_1, z_2).$$

Here f is a function on the Siegel domain of type II

$$(1.2) \quad \mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} - z_2 \overline{z_2} > 0\},$$

and L_k is a differential operator given by

$$(1.3) \quad \begin{aligned} L_k = & (z_1 + \overline{z_1} - z_2 \overline{z_2}) \\ & \times \left[(z_1 + \overline{z_1}) \frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2} \frac{\partial^2}{\partial z_1 \partial \overline{z_2}} - k \left(\frac{\partial}{\partial z_1} - \frac{\partial}{\partial \overline{z_1}} \right) \right], \end{aligned}$$

where k is an integer. Let $\Gamma \subset U(2, 1)$ be an arithmetic subgroup or a convex cocompact subgroup, where

$$(1.4) \quad U(2, 1) = \{g \in GL(3, \mathbb{C}) : g^* J g = J\}, \quad J = \begin{pmatrix} & & -1 \\ & 1 & \\ -1 & & \end{pmatrix},$$

and assume that f is a Γ -automorphic form of weight k in the sense that it is invariant under

$$(1.5) \quad f \mapsto f(\gamma(z_1, z_2)) (a_1 z_1 + a_2 z_2 + a_3)^{-k} \overline{(a_1 z_1 + a_2 z_2 + a_3)}^k$$

for every $\gamma = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix} \in \Gamma$. In fact, L_k commutes with (1.5).

Now, let us recall some basic facts about complex hyperbolic geometry (see [Ap] and [Go]). Geometry of complex hyperbolic space $\mathbf{H}_{\mathbb{C}}^n$ is the geometry of the unit ball $\mathbb{B}_{\mathbb{C}}^n$

in \mathbb{C}^n with the Kähler structure given by the Bergman metric whose automorphisms are biholomorphic automorphisms of the ball, i.e., elements of $PU(n, 1)$. Any complex hyperbolic manifold can be represented as the quotient $M = \mathbf{H}_{\mathbb{C}}^n/\Gamma$ by a discrete torsion free isometric action of the fundamental group of M , $\pi_1(M) \cong \Gamma \subset PU(n, 1)$, its boundary at infinity $\partial_{\infty} M$ is naturally identified as the quotient $\Omega(\Gamma)/\Gamma$ of the discontinuity set of Γ at infinity. Here the discontinuity set $\Omega(\Gamma)$ is the maximal subset of $\partial \mathbf{H}_{\mathbb{C}}^n$ where Γ acts discretely, its complement $\Lambda(\Gamma) = \partial \mathbf{H}_{\mathbb{C}}^n \setminus \Omega(\Gamma)$ is the limit set of Γ , $\Lambda(\Gamma) = \overline{\Gamma(x)} \cap \partial \mathbf{H}_{\mathbb{C}}^n$ for any $x \in \mathbf{H}_{\mathbb{C}}^n$.

In general, let G be a connected, linear, real simple Lie group of rank one, $G = KAN$ be an Iwasawa decomposition of G , $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$ be the corresponding Iwasawa decomposition of the Lie algebra \mathfrak{g} , and $P = MAN$ be a minimal parabolic subgroup. The group G acts isometrically on the rank-one symmetric space $X = G/K$. Let $\partial X = G/P = K/M$ be its geodesic boundary. We regard $\overline{X} := X \cup \partial X$ as a compact manifold with boundary.

By the classification of symmetric spaces with strictly negative sectional curvature, we know that X is one of the following spaces: a real hyperbolic space $\mathbf{H}_{\mathbb{R}}^n$ ($n \geq 1$), a complex hyperbolic space $\mathbf{H}_{\mathbb{C}}^n$ ($n \geq 2$), a quaternionic hyperbolic space $\mathbf{H}_{\mathbb{H}}^n$ ($n \geq 2$) or the Cayley hyperbolic plane $\mathbf{H}_{\mathbb{O}}^2$, and G is a linear group finitely covering the orientation-preserving isometric group of X .

Following [BO], we consider a torsion-free discrete subgroup $\Gamma \subset G$ such that ∂X admits a Γ -invariant partition $\partial X = \Omega(\Gamma) \cup \Lambda(\Gamma)$, where $\Omega(\Gamma) \neq \emptyset$ is open and Γ acts freely and cocompactly on $X \cup \Omega(\Gamma)$. The closed subset $\Lambda(\Gamma)$ is called the limit set of Γ . The locally symmetric space $Y := \Gamma \backslash X$ is a complete Riemannian manifold of infinite volume without cusps. It can be compactified by adjoining the geodesic boundary $B := \Gamma \backslash \Omega(\Gamma)$.

A subgroup Γ satisfying this assumption is called convex cocompact or geometrically cocompact since it acts cocompactly on the convex hull of the limit set. The quotient Y is called a Kleinian manifold.

For $H \in \mathfrak{a}$, we define $\rho \in \mathfrak{a}^*$ by $\rho(H) = \frac{1}{2}\text{tr}(\text{ad}(H)|_{\mathfrak{n}})$. Then $\rho(\mathbf{H}_{\mathbb{R}}^n) = \frac{n-1}{2}$ and $\rho(\mathbf{H}_{\mathbb{C}}^n) = n$.

Now, we need the following definition (see [Co]):

Definition. For any discrete group $\Gamma \subset G$, the critical exponent $\delta(\Gamma)$ is the infimum of all s such that

$$\sum_{\gamma \in \Gamma} \exp(-sd(x, \gamma y))$$

converges for some (or any) $x, y \in X$, where $d(x, y)$ is the Riemannian distance from x to y .

The critical exponent $\delta(\Gamma)$ has been extensively studied (see [Pa1], [Su], [Co] and [CoI]). It is known that $\delta(\Gamma) \in [0, 2\rho]$ if Γ is nontrivial. In fact, if $\Omega(\Gamma) \neq \emptyset$, then $\delta(\Gamma)$ is equal to the Hausdorff dimension of the limit set with respect to the natural class of sub-Riemannian metrics on ∂X .

In the case of $U(2, 1)$, $\delta(\Gamma) \in [0, 4]$ for $X = \mathbf{H}_{\mathbb{C}}^2$.

The theory of Eisenstein series and the spectral theory for Kleinian groups on real hyperbolic spaces $\mathbf{H}_{\mathbb{R}}^n$ has been extensively studied, in particular, by Mandouvalos [Ma1], [Ma2], Patterson [Pa3], and Perry [Pe1], [Pe2], [FHP]. A Kleinian group Γ is a discrete subgroup acting on $\mathbf{H}_{\mathbb{R}}^n$, which acts discontinuously on $\mathbf{H}_{\mathbb{R}}^n$ and has a fundamental domain in $\mathbf{H}_{\mathbb{R}}^n$ of infinite hyperbolic volume. This in particular implies that it is a non-arithmetic subgroup. Their theory extended previous work of Roelcke [Ro], Elstrodt [El], Patterson [Pa2] and Fay [Fa] on the spectral theory and Eisenstein series for Fuchsian groups of the second kind.

Now, we give the corresponding results on the complex hyperbolic space $\mathbf{H}_{\mathbb{C}}^2$. We give the Eisenstein series for the Picard modular group, which is an arithmetic subgroup of $U(2, 1)$. We also give the Eisenstein series for any discrete subgroup (either arithmetic or non-arithmetic). In particular, for a convex cocompact subgroup Γ which satisfies that $\delta(\Gamma) < 1$, we give the product formulas. Now, we state the results of this paper.

The Poisson kernel is given by

$$(1.6) \quad P(Z, W) = \frac{\rho(Z)}{|\rho(Z, W)|^2}, \quad \text{for } Z = (z_1, z_2) \in \mathfrak{S}_2, W = (w_1, w_2) \in \partial\mathfrak{S}_2,$$

where

$$(1.7) \quad \rho(Z, W) := \overline{z_1} + w_1 - \overline{z_2}w_2, \quad \rho(Z) = \rho(Z, Z).$$

Let

$$(1.8) \quad \sigma(Z, Z') := \frac{|\rho(Z, Z')|^2}{\rho(Z)\rho(Z')}.$$

It is a point-pair invariant under the action of $U(2, 1)$. The critical exponent is given by

$$(1.9) \quad \delta(\Gamma) := \inf\{s > 0 : \sum_{\gamma \in \Gamma} \sigma(Z, \gamma(Z'))^{-s} < \infty\}.$$

For a discrete subgroup $\Gamma \subset U(2, 1)$, the Eisenstein series is defined by

$$(1.10) \quad E(Z, W; s) := \sum_{\gamma \in \Gamma} P(\gamma(Z), W)^s, \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

The automorphic Green function is given by

$$(1.11) \quad G(Z, Z'; s) = \sum_{\gamma \in \Gamma} r(Z, \gamma(Z'); s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma),$$

with

$$(1.12) \quad r(Z, Z'; s) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s - \frac{1}{2})} (\frac{1}{4})^{s-1} u^{-s} F(s, s-1; 2s-1; -u^{-1}).$$

Here

$$(1.13) \quad u = u(Z, Z') = \sigma(Z, Z') - 1.$$

Now, the first main result of this paper is stated as follows:

THEOREM 1.1. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following product formula holds:*

$$(1.14) \quad \begin{aligned} & \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) E(Z', W; 2-s) dm(W) \\ &= -\frac{\pi}{s-1} c(s) c(2-s) [G(Z, Z'; s) - G(Z, Z'; 2-s)], \end{aligned}$$

where $\delta(\Gamma) < \operatorname{Re}(s) < 2 - \delta(\Gamma)$ and $c(s) = \sqrt{\pi}\Gamma(s - \frac{1}{2})\Gamma(s)^{-1}$.

For $\gamma = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix} \in U(2, 1)$, set

$$(1.15) \quad j(\gamma, Z) = a_1 z_1 + a_2 z_2 + a_3 z_3.$$

The S-matrix is given by

$$(1.16) \quad S(W, W'; s) := \sum_{\gamma \in \Gamma} |\rho(\gamma(W), W')|^{-2s} |j(\gamma, W)|^{-2s}$$

for $W, W' \in \Omega(\Gamma)$ and $\operatorname{Re}(s) > \delta(\Gamma)$. By the S-matrix, we give the functional equation of Eisenstein series:

THEOREM 1.2. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following functional equation for Eisenstein series holds:*

$$(1.17) \quad \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) S(W, W'; 2-s) dm(W) = \frac{4^{s-1} \pi}{s-1} \frac{\sqrt{\pi}\Gamma(s - \frac{1}{2})}{\Gamma(s)} E(Z, W'; 2-s),$$

for $1 < \operatorname{Re}(s) < 2 - \delta(\Gamma)$.

The Poisson kernel of weight k is given by

$$(1.18) \quad P_k(Z, W; s) = \rho(Z)^s \rho(Z, W)^{k-s} \rho(\overline{Z}, \overline{W})^{-k-s},$$

where $Z \in \mathfrak{S}_2$, $W \in \partial\mathfrak{S}_2$. The Eisenstein series of weight k is defined by

$$(1.19) \quad E(Z, W; k, s) = \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k P_k(\gamma(Z), W; s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

Set

$$(1.20) \quad \begin{aligned} K(Z, W; k, s) &= \frac{\pi^{\frac{3}{2}} \Gamma(\frac{3}{2} - s)}{(|k| + 1 - s) \Gamma(2 - s)} \left(\frac{1}{4}\right)^{s-1} \rho(Z, W)^k \rho(\bar{Z}, \bar{W})^{-k} \\ &\times \sigma^{-|k|} (\sigma - 1)^{|k|-s} F(s - |k|, s - 1 - |k|; 2s - 1; -\frac{1}{\sigma - 1}). \end{aligned}$$

The automorphic Green function of weight k is given by

$$(1.21) \quad G(Z, Z'; k, s) = \sum_{\gamma \in \Gamma} K(Z, \gamma(Z'); k, s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

Now, we state the main theorem of this paper:

THEOREM 1.3 (MAIN THEOREM). *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following product formula holds:*

$$(1.22) \quad \begin{aligned} &\int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; k, s) E(Z', W; -k, 2 - s) dm(W) \\ &= G(Z, Z'; k, s) + G(Z, Z'; k, 2 - s) \\ &= G(Z', Z; -k, s) + G(Z', Z; -k, 2 - s), \end{aligned}$$

for $\delta(\Gamma) < \operatorname{Re}(s) < 2 - \delta(\Gamma)$ and $k = 0$ or $k = \pm 1$.

The S-matrix of weight k is defined by

$$(1.23) \quad S(W, W'; k, s) := \sum_{\gamma \in \Gamma} j(\gamma, W)^{k-s} j(\bar{\gamma}, \bar{W})^{-k-s} \rho(\gamma(W), W')^{-k-s} \rho(\bar{\gamma}(W), \bar{W}')^{k-s}$$

for $W, W' \in \Omega(\Gamma)$ and $\operatorname{Re}(s) > \delta(\Gamma)$. By the S-matrix of weight k , we obtain the functional equation of Eisenstein series of weight k .

THEOREM 1.4. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following functional equation for Eisenstein series of weight k holds:*

$$(1.24) \quad \begin{aligned} &\int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; k, s) S(W, W'; -k, 2 - s) dm(W) \\ &= \frac{\pi^{\frac{3}{2}} \Gamma(s - \frac{1}{2})}{(|k| + s - 1) \Gamma(s)} 4^{s-1} E(Z, W'; k, 2 - s), \end{aligned}$$

for $1 < \operatorname{Re}(s) < 2 - \delta(\Gamma)$ and $k = 0, \pm 1$.

In fact, when $k = 0$, Theorem 1.3 and Theorem 1.4 reduces to Theorem 1.1 and Theorem 1.2.

In the appendix, we give the product formulas on $SL(2, \mathbb{R})$.

THEOREM 1.5. *Assume that Γ is convex cocompact and $\delta(\Gamma) < \frac{1}{2}$. Then the following product formula on $SL(2, \mathbb{R})$ holds:*

$$(1.25) \quad \int_{\Gamma \backslash \Omega(\Gamma)} E(z, \zeta; k, s) E(z', \zeta; -k, 1-s) dm(\zeta) = G(z, z'; k, s) + G(z, z'; k, 1-s),$$

for $\delta(\Gamma) < \operatorname{Re}(s) < 1 - \delta(\Gamma)$ and $k = 0, \pm 1$.

2. Epstein zeta function and Eisenstein series on Picard modular groups

Let \mathfrak{S}_2 be the Siegel domain

$$(2.1) \quad \mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} > |z_2|^2\}.$$

The imaginary quadratic field $K = \mathbb{Q}(\sqrt{-3})$ is called the field of Eisenstein numbers. Its ring of integers

$$(2.2) \quad \mathcal{O} = \mathcal{O}_K = \{a/2 + b\sqrt{-3}/2 : a, b \in \mathbb{Z}, a \equiv b \pmod{2}\} = \mathbb{Z} + \mathbb{Z}[\omega]$$

with $\omega = \frac{-1+\sqrt{-3}}{2}$ is called the ring of Eisenstein integers.

In their paper [KW], Korányi and Wolf studied the general Cayley transform which carries the bounded domain of the Harish-Chandra realization into a generalized half-plane. In [KR], Korányi and Reimann gave the Cayley transform on $U(2, 1)$. Now, we define the Cayley transform $C : \mathbb{B}^2 := \{(w_1, w_2) \in \mathbb{C}^2 : |w_1|^2 + |w_2|^2 < 1\} \rightarrow \mathfrak{S}_2$, where

$$(2.3) \quad C = \begin{pmatrix} -\bar{\omega} & -\omega \\ -1 & 1 \end{pmatrix} \in GL(3, \mathcal{O}).$$

It is known that

$$(2.4) \quad U(2, 1) = \{g \in GL(3, \mathbb{C}) : g I_{2,1} g^* = I_{2,1}\},$$

where $I_{2,1} = \operatorname{diag}\{1, 1, -1\}$.

Let $E_{i,j}$ be the 3×3 matrix with a 1 at the i, j -th component, zeros elsewhere. Using Kronecker's delta, we have

$$[E_{p,q}, E_{r,s}] = \delta_{q,r} E_{p,s} - \delta_{s,p} E_{r,q}.$$

A basis of $\mathfrak{g} = u(2, 1)$ is as follows:

$$\begin{aligned} X_1 &= E_{2,3} + E_{3,2}, \quad X_2 = E_{1,3} + E_{3,1}, \quad X_3 = E_{1,2} - E_{2,1}, \\ X_4 &= iE_{2,3} - iE_{3,2}, \quad X_5 = iE_{1,3} - iE_{3,1}, \quad X_6 = iE_{1,2} + iE_{2,1}, \\ X_7 &= iE_{1,1}, \quad X_8 = iE_{2,2}, \quad X_9 = iE_{3,3}. \end{aligned}$$

Thus,

$$\begin{aligned} E_{2,3} &= \frac{1}{2}(X_1 - iX_4), \quad E_{3,2} = \frac{1}{2}(X_1 + iX_4), \\ E_{1,3} &= \frac{1}{2}(X_2 - iX_5), \quad E_{3,1} = \frac{1}{2}(X_2 + iX_5), \\ E_{1,2} &= \frac{1}{2}(X_3 - iX_6), \quad E_{2,1} = -\frac{1}{2}(X_3 + iX_6), \\ E_{1,1} &= -iX_7, \quad E_{2,2} = -iX_8, \quad E_{3,3} = -iX_9. \end{aligned}$$

The action of $U(2, 1)$ on \mathbb{B}^2 is as follows:

$$\begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} (w_1, w_2) = \left(\frac{a_1 w_1 + a_2 w_2 + a_3}{c_1 w_1 + c_2 w_2 + c_3}, \frac{b_1 w_1 + b_2 w_2 + b_3}{c_1 w_1 + c_2 w_2 + c_3} \right).$$

We have

$$\begin{aligned} X_3 f(w_1, w_2, \overline{w_1}, \overline{w_2}) &= \frac{d}{dt} [f(\exp(t \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix})(w_1, w_2)) + f(\exp(t \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix})(\overline{w_1}, \overline{w_2}))]|_{t=0} \\ &= \frac{d}{dt} [f(\begin{pmatrix} \cos t & \sin t & 0 \\ -\sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix}(w_1, w_2)) + f(\begin{pmatrix} \cos t & \sin t & 0 \\ -\sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix}(\overline{w_1}, \overline{w_2}))]|_{t=0} \\ &= \frac{d}{dt} f(w_1 \cos t + w_2 \sin t, -w_1 \sin t + w_2 \cos t)|_{t=0} \\ &\quad + \frac{d}{dt} f(\overline{w_1} \cos t + \overline{w_2} \sin t, -\overline{w_1} \sin t + \overline{w_2} \cos t)|_{t=0} \\ &= w_2 \frac{\partial f}{\partial w_1} - w_1 \frac{\partial f}{\partial w_2} + \overline{w_2} \frac{\partial f}{\partial \overline{w_1}} - \overline{w_1} \frac{\partial f}{\partial \overline{w_2}}. \end{aligned}$$

So, the corresponding operator is

$$X_3 = w_2 \frac{\partial}{\partial w_1} - w_1 \frac{\partial}{\partial w_2} + \overline{w_2} \frac{\partial}{\partial \overline{w_1}} - \overline{w_1} \frac{\partial}{\partial \overline{w_2}}.$$

Similarly, we have

$$\begin{aligned}
X_2 &= (1 - w_1^2) \frac{\partial}{\partial w_1} - w_1 w_2 \frac{\partial}{\partial w_2} + (1 - \overline{w_1}^2) \frac{\partial}{\partial \overline{w}_1} - \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_2}, \\
X_1 &= -w_1 w_2 \frac{\partial}{\partial w_1} + (1 - w_2^2) \frac{\partial}{\partial w_2} - \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_1} + (1 - \overline{w_2}^2) \frac{\partial}{\partial \overline{w}_2}, \\
X_6 &= i w_2 \frac{\partial}{\partial w_1} + i w_1 \frac{\partial}{\partial w_2} - i \overline{w_2} \frac{\partial}{\partial \overline{w}_1} - i \overline{w_1} \frac{\partial}{\partial \overline{w}_2}, \\
X_5 &= i(1 + w_1^2) \frac{\partial}{\partial w_1} + i w_1 w_2 \frac{\partial}{\partial w_2} - i(1 + \overline{w_1}^2) \frac{\partial}{\partial \overline{w}_1} - i \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_2}, \\
X_4 &= i w_1 w_2 \frac{\partial}{\partial w_1} + i(1 + w_2^2) \frac{\partial}{\partial w_2} - i \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_1} - i(1 + \overline{w_2}^2) \frac{\partial}{\partial \overline{w}_2}, \\
X_7 &= i w_1 \frac{\partial}{\partial w_1} - i \overline{w_1} \frac{\partial}{\partial \overline{w}_1}, \\
X_8 &= i w_2 \frac{\partial}{\partial w_2} - i \overline{w_2} \frac{\partial}{\partial \overline{w}_2}, \\
X_9 &= -i w_1 \frac{\partial}{\partial w_1} - i w_2 \frac{\partial}{\partial w_2} + i \overline{w_1} \frac{\partial}{\partial \overline{w}_1} + i \overline{w_2} \frac{\partial}{\partial \overline{w}_2}.
\end{aligned}$$

We denote X_{ij} as the differential operator corresponding to $E_{i,j}$, then

$$\begin{aligned}
X_{23} &= \frac{1}{2}(X_1 - iX_4) = \frac{\partial}{\partial w_2} - \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_1} - \overline{w_2}^2 \frac{\partial}{\partial \overline{w}_2}, \quad X_{32} = \frac{1}{2}(X_1 + iX_4) = \overline{X_{23}}, \\
X_{13} &= \frac{1}{2}(X_2 - iX_5) = \frac{\partial}{\partial w_1} - \overline{w_1}^2 \frac{\partial}{\partial \overline{w}_1} - \overline{w_1 w_2} \frac{\partial}{\partial \overline{w}_2}, \quad X_{31} = \frac{1}{2}(X_2 + iX_5) = \overline{X_{13}}, \\
X_{12} &= \frac{1}{2}(X_3 - iX_6) = w_2 \frac{\partial}{\partial w_1} - \overline{w_1} \frac{\partial}{\partial \overline{w}_2}, \quad X_{21} = -\frac{1}{2}(X_3 + iX_6) = -\overline{X_{12}}, \\
X_{11} &= -iX_7 = w_1 \frac{\partial}{\partial w_1} - \overline{w_1} \frac{\partial}{\partial \overline{w}_1}, \\
X_{22} &= -iX_8 = w_2 \frac{\partial}{\partial w_2} - \overline{w_2} \frac{\partial}{\partial \overline{w}_2}, \\
X_{33} &= -iX_9 = -w_1 \frac{\partial}{\partial w_1} - w_2 \frac{\partial}{\partial w_2} + \overline{w_1} \frac{\partial}{\partial \overline{w}_1} + \overline{w_2} \frac{\partial}{\partial \overline{w}_2}.
\end{aligned}$$

Now, if n is any positive integer, we claim that

$$\sum_{i_1=1}^3 \sum_{i_2=1}^3 \cdots \sum_{i_n=1}^3 X_{i_1 i_2} X_{i_2 i_3} \cdots X_{i_n i_1}$$

lies in the center of $U(\mathfrak{g})$. To prove this, without loss of generality, let us assume $n = 2$. Then

$$[X_{pq}, \sum_{i_1} \sum_{i_2} X_{i_1 i_2} X_{i_2 i_1}] = \sum_{i_1} \sum_{i_2} ([X_{pq}, X_{i_1 i_2}] X_{i_2 i_1} + X_{i_1 i_2} [X_{pq}, X_{i_2 i_1}]).$$

By $[X_{pq}, X_{rs}] = \delta_{qr} X_{ps} - \delta_{sp} X_{rq}$, this equals

$$\begin{aligned} & \sum_{i_1} \sum_{i_2} (\delta_{qi_1} X_{pi_2} X_{i_2 i_1} - \delta_{i_2 p} X_{i_1 q} X_{i_2 i_1} + X_{i_1 i_2} \delta_{qi_2} X_{pi_1} - X_{i_1 i_2} \delta_{i_1 p} X_{i_2 q}) \\ &= \sum_{i_2} X_{pi_2} X_{i_2 q} - \sum_{i_1} X_{i_1 q} X_{pi_1} + \sum_{i_1} X_{i_1 q} X_{pi_1} - \sum_{i_2} X_{pi_2} X_{i_2 q} = 0. \end{aligned}$$

Thus the above sum commutes with the generators X_i of $U(\mathfrak{g})$, and lies in the center of $U(\mathfrak{g})$. The proof is similar if $n \neq 2$.

Set

$$\begin{aligned} D_1 &= \sum_{i_1=1}^3 \sum_{i_2=1}^3 E_{i_1 i_2} E_{i_2 i_1}, \\ D_2 &= \sum_{i_1=1}^3 \sum_{i_2=1}^3 \sum_{i_3=1}^3 E_{i_1 i_2} E_{i_2 i_3} E_{i_3 i_1}. \end{aligned}$$

In the category of differential operators,

$$[X_{pq}, X_{rs}] = \delta_{sp} X_{rq} - \delta_{qr} X_{ps}.$$

Therefore, we have

$$D_1 = 2(X_{11}^2 + X_{22}^2 + X_{11}X_{22} + X_{12}X_{21} + X_{23}X_{32} + X_{31}X_{13}) = 2\Delta,$$

$$\begin{aligned} D_2 = & 3[X_{11}(X_{12}X_{21} - X_{32}X_{23} - X_{22}^2) + X_{22}(X_{21}X_{12} - X_{31}X_{13} - X_{11}^2) \\ & + X_{23}X_{31}X_{12} + X_{13}X_{32}X_{21} - X_{31}X_{13} - X_{32}X_{23} + X_{11} + X_{22}] = -3\Delta, \end{aligned}$$

where

$$\begin{aligned} (2.5) \quad \Delta &= (1 - |w_1|^2 - |w_2|^2) \\ &\times \left[(1 - |w_1|^2) \frac{\partial^2}{\partial w_1 \partial \bar{w}_1} + (1 - |w_2|^2) \frac{\partial^2}{\partial w_2 \partial \bar{w}_2} - w_1 \bar{w}_2 \frac{\partial^2}{\partial w_1 \partial \bar{w}_2} - \bar{w}_1 w_2 \frac{\partial^2}{\partial \bar{w}_1 \partial w_2} \right] \end{aligned}$$

is the Laplace-Beltrami operator of $U(2, 1)$ on \mathbb{B}^2 .

Let $G = CU(2, 1)C^{-1}$, $K = C(U(2) \times U(1))C^{-1}$ and $G(\mathcal{O}) = CU(2, 1; \mathcal{O})C^{-1}$, then $G(\mathcal{O})$ is an arithmetic subgroup of G and $S := G(\mathcal{O})\backslash G/K$ is called a Picard modular surface (see [Pic1], [Pic2], [Gir], [Ho1], [Ho2], [LR] and [BB] for the details).

The complex hyperbolic Gauss-Bonnet formula states that (see [HP] and [P])

$$\text{Vol}(\mathbb{H}_{\mathbb{C}}^2/\Gamma) = \frac{8\pi^2}{3}\chi(\mathbb{H}_{\mathbb{C}}^2/\Gamma),$$

where $\chi(\mathbb{H}_{\mathbb{C}}^2/\Gamma)$ is the Euler number of $\mathbb{H}_{\mathbb{C}}^2/\Gamma$. Holzapfel showed that

$$\chi(\mathbb{B}^2/PSU(2, 1; \mathcal{O})) = \frac{1}{24}.$$

As $PSU(2, 1; \mathcal{O})$ has index 3 in $PU(2, 1; \mathcal{O})$, we see that $\chi(\mathbb{B}^2/PU(2, 1; \mathcal{O})) = \frac{1}{72}$, and so

$$\text{Vol}_{\mathbb{B}^2}(\mathbb{B}^2/PU(2, 1; \mathcal{O})) = \frac{8\pi^2}{3} \cdot \frac{1}{72} = \frac{\pi^2}{27}.$$

Denote $J = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$, then $CI_{2,1}C^* = J$. It follows that

$$(2.6) \quad G = \{g \in GL(3, \mathbb{C}) : gJg^* = J\}.$$

This is the other realization of $U(2, 1)$. For simplicity, from now on, we use the same symbol $U(2, 1)$ to denote as G . In fact, $G = \{g \in GL(3, \mathbb{C}) : g^*Jg = J\}$.

Let P be the parabolic subgroup of $U(2, 1)$ which consists of upper triangular matrices. This group P has the Langlands decomposition $P = NAM$, where

$$\begin{aligned} N &= \left\{ n = [z, t] = \begin{pmatrix} 1 & z & \frac{1}{2}|z|^2 + it \\ 0 & 1 & \bar{z} \\ 0 & 0 & 1 \end{pmatrix} : z \in \mathbb{C}, t \in \mathbb{R} \right\}. \\ A &= \left\{ a = \begin{pmatrix} \rho^{-1} & & \\ & 1 & \\ & & \rho \end{pmatrix} : \rho > 0 \right\}, \\ M &= \left\{ m = \begin{pmatrix} \beta & & \\ & \beta^{-2} & \\ & & \beta \end{pmatrix} : \beta \in \mathbb{C}, |\beta|^2 = 1 \right\}. \end{aligned}$$

Then $G = NAMK = NAK$ is the Iwasawa decomposition. Set $\Gamma = G(\mathcal{O})$, then

$$(2.7) \quad \Gamma \cap N = \left\{ [\alpha, \beta] = \begin{pmatrix} 1 & \alpha & \beta \\ & 1 & \bar{\alpha} \\ & & 1 \end{pmatrix} : |\alpha|^2 = \beta + \bar{\beta}, \alpha, \beta \in \mathcal{O} \right\}.$$

If $k \in K$, then

$$k = C \begin{pmatrix} \alpha & \beta & 0 \\ \gamma & \delta & 0 \\ 0 & 0 & e^{i\theta} \end{pmatrix} C^{-1},$$

where $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in U(2)$ and $\theta \in [0, 2\pi)$. Write explicitly,

$$k = \begin{pmatrix} -\bar{\omega}\alpha - \omega e^{i\theta} & -\bar{\omega}\beta & -\alpha + e^{i\theta} \\ \gamma & \delta & \omega\gamma \\ -\alpha + e^{i\theta} & -\beta & -\omega\alpha - \bar{\omega}e^{i\theta} \end{pmatrix}.$$

We have $k(-\omega, 0) = (-\omega, 0)$. Note that $\rho(-\omega, 0) = 1$. Therefore, $g(-\omega, 0) = (-\omega, 0)$ if and only if $g \in K$. Define $\pi : G \rightarrow \mathfrak{S}_2$ by $\pi(g) = g(-\omega, 0)$. It induces a bijection $G/K \rightarrow \mathfrak{S}_2$. For $g = nak$,

$$\pi(g) = nak(-\omega, 0) = na(-\omega, 0) = (-\omega\rho^{-2} + \frac{1}{2}|z|^2 + it, \bar{z}).$$

Thus, if we set $(z_1, z_2) = \pi(g)$, then $z_1 + \bar{z}_1 - z_2\bar{z}_2 = \rho^{-2}$.

Following [He], let $\Delta = \Delta(\Gamma) = \left\{ \begin{pmatrix} 1 & ix \\ & 1 \\ & & 1 \end{pmatrix} : x \in \mathbb{R} \right\} \cap \Gamma$ and $q = q(\Gamma)$ be the positive integer such that $\begin{pmatrix} 1 & q\sqrt{-3} \\ & 1 \\ & & 1 \end{pmatrix}$ generates Δ . For $[\alpha, \beta], [\alpha', \beta'] \in \Gamma \cap N$,

$$[\alpha', \beta'][\alpha, \beta]([\alpha, \beta][\alpha', \beta'])^{-1} = [0, 2\sqrt{-3}\operatorname{Im}(\alpha'\bar{\alpha})] \in \Delta,$$

where $\operatorname{Im}\alpha := b$ for $\alpha = a + b\sqrt{-3}$, $a, b \in \mathbb{Z}$. Hence, $2q^{-1}\operatorname{Im}(\alpha'\bar{\alpha}) \in \mathbb{Z}$.

By Cayley transform, we have

THEOREM 2.1. *The Laplace-Beltrami operator of $U(2, 1)$ on \mathfrak{S}_2 is*

$$(2.8) \quad L = (z_1 + \bar{z}_1 - |z_2|^2) \left[(z_1 + \bar{z}_1) \frac{\partial^2}{\partial z_1 \partial \bar{z}_1} + \frac{\partial^2}{\partial z_2 \partial \bar{z}_2} + z_2 \frac{\partial^2}{\partial \bar{z}_1 \partial z_2} + \bar{z}_2 \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \right].$$

If $g = \begin{pmatrix} 1 & \alpha & \beta \\ & 1 & \gamma \\ & & 1 \end{pmatrix} \in \Gamma$, then $\alpha = \bar{\gamma}$, $|\alpha|^2 = \beta + \bar{\beta}$. Denote $Z = (z_1, z_2)$, set $\rho(Z) = z_1 + \bar{z}_1 - |z_2|^2$, we have

$$\rho(g, Z) := \rho(g(Z)) = (z_1 + \alpha z_2 + \beta) + (\bar{z}_1 + \bar{\alpha} \bar{z}_2 + \bar{\beta}) - (z_2 + \gamma)(\bar{z}_2 + \bar{\gamma}) = \rho(Z).$$

If $g = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix} \in U(2, 1)$, then $a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}$. By a straightforward calculation, we have

$$(2.9) \quad \rho(g, Z) = \frac{\rho(Z)}{|j(\gamma, Z)|^2}, \quad \text{with } j(\gamma, Z) := a_1z_1 + a_2z_2 + a_3.$$

For the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$, the Eisenstein series is defined as

$$(2.10) \quad E(Z; s) = \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} \rho(\gamma, Z)^s, \quad \text{for } \operatorname{Re}(s) > 2.$$

The unit group of \mathcal{O} is $\{\pm 1, \pm \omega, \pm \overline{\omega}\}$, its order is 6.

$$E(Z; s) = \frac{1}{6} \sum_{\substack{a_1, a_2, a_3 \in \mathcal{O}, (a_1, a_2, a_3) = \mathcal{O}, \\ a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}}} \frac{\rho^s}{|a_1z_1 + a_2z_2 + a_3|^{2s}},$$

where (a_1, a_2, a_3) denotes the ideal generated by a_1, a_2 and a_3 .

THEOREM 2.2. *The Eisenstein series satisfies the following equation:*

$$(2.11) \quad LE(Z; s) = s(s-2)E(Z; s).$$

Proof. For $j(\gamma, Z) = a_1z_1 + a_2z_2 + a_3$,

$$L\rho(\gamma, Z)^s = s(s-2)\rho(\gamma, Z)^s + (-\overline{a_1}a_3 - a_1\overline{a_3} + a_2\overline{a_2})s^2 \frac{\rho(Z)^s}{|j(\gamma, Z)|^{2(s+1)}}.$$

□

Let

$$\mathcal{SP}_3(\mathbb{C}) = \{Y \in M_3(\mathbb{C}) : Y \text{ positive definite, } Y^* = Y, \det Y = 1\},$$

$\mathcal{H}_3(\mathbb{C}) := \mathcal{SP}_3(\mathbb{C}) \cap U(2, 1)$ and denote $Y[a] = a^*Ya$, for $Y \in \mathcal{SP}_3(\mathbb{C})$.

For $a \in A, n \in N$ as above,

$$a[n] := n^*an = \begin{pmatrix} \frac{1}{\rho} & \frac{z}{\rho} & \frac{\frac{1}{2}|z|^2+it}{\rho} \\ \frac{\overline{z}}{\rho} & \frac{|z|^2}{\rho} + 1 & \frac{\overline{z}(\frac{1}{2}|z|^2+it)}{\rho} + 1 \\ \frac{\frac{1}{2}|z|^2-it}{\rho} & z(\frac{\frac{1}{2}|z|^2-it}{\rho} + 1) & \frac{\frac{1}{4}|z|^4+t^2}{\rho} + |z|^2 + \rho \end{pmatrix}.$$

We can identify \mathfrak{S}_2 with $\mathcal{H}_3(\mathbb{C})$ by $(z_1, z_2) \mapsto W_{(\rho, t, z)} = a[n]$, where

$$z_1 = \rho + \frac{1}{2}|z|^2 - it, z_2 = -\bar{z}, \rho = \frac{1}{2}(z_1 + \bar{z}_1 - |z_2|^2).$$

LEMMA 2.3. *If $a = (a_3, -a_2, a_1) \in \mathcal{O}^3$, where $a_1\bar{a}_3 + \bar{a}_1a_3 = a_2\bar{a}_2$, then*

$$(2.12) \quad W_{(\rho, t, z)}[a^*] = \rho^{-1}|a_1z_1 + a_2z_2 + a_3|^2.$$

Proof. In fact,

$$\rho^{-1}|a_1z_1 + a_2z_2 + a_3|^2 - W_{(\rho, t, z)}[a^*] = a_1\bar{a}_3 + \bar{a}_1a_3 - a_2\bar{a}_2 = 0.$$

□

Definition 2.4. For the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$ the Epstein zeta function is given by

$$(2.13) \quad Z(Y, s) := \frac{1}{6} \sum_{a \in \mathcal{O}^3 - \{0\}, a_1\bar{a}_3 + \bar{a}_1a_3 = a_2\bar{a}_2} Y[a]^{-s},$$

where $\text{Re}(s) > 2$.

LEMMA 2.5. $Z(W_{(\rho, t, z)}, s) = \zeta_K(2s)E(z_1, z_2; s)$, where $\zeta_K(s)$ is the Dedekind zeta function of the number field $K = \mathbb{Q}(\sqrt{-3})$.

Because $E(z_1, z_2; s) = Z(W_{(\rho, t, z)}, s)/\zeta_K(2s)$, any zero of $\zeta_K(2s)$ will give a pole of $E(z_1, z_2; s)$ unless $Z(W, s)$ vanishes at that value of s . The trivial zeros of $Z(W, s)$ and $\zeta_K(2s)$ are both $s = -1, -2, -3, \dots$, and both have order 1. So the trivial zeros cancel out in $E(z_1, z_2; s)$.

Suppose $a_1, a_2, a_3 \in \mathcal{O}$ and $\bar{a}_1a_3 + a_1\bar{a}_3 = a_2\bar{a}_2$, then $J[(a_1, a_2, a_3)^*] = 0$ and $J[(a_3, -a_2, a_1)^*] = 0$.

For $n, a \in U(2, 1)$ as above, then $n^* \in U(2, 1)$, thus $W \in U(2, 1)$. If $g \in \mathcal{H}_3(\mathbb{C})$, $gJg = J$, thus $g^{-1} = JgJ$. Suppose $a = (a_1, a_2, a_3) \in \mathcal{O}^3 - \{0\}$, we have the following lemma:

LEMMA 2.6. *For $g \in \mathcal{H}_3(\mathbb{C})$, $g[(a_3, -a_2, a_1)^*] = g^{-1}[(a_1, a_2, a_3)^*]$.*

Proof. In fact,

$$g^{-1}[(a_1, a_2, a_3)^*] = JgJ[(a_1, a_2, a_3)^*] = (-a_3, a_2, -a_1)g \begin{pmatrix} -\bar{a}_3 \\ \bar{a}_2 \\ -\bar{a}_1 \end{pmatrix} = g[(a_3, -a_2, a_1)^*].$$

□

By Lemma 2.6, it follows that

THEOREM 2.7. *The functional equation of Epstein zeta functions:*

$$(2.14) \quad Z(Y^{-1}, s) = Z(Y, s)$$

if $Y \in \mathcal{H}_3(\mathbb{C})$.

Following [Ho2], we define the Picard modular forms on \mathfrak{S}_2 as follows:

Definition 2.8. A holomorphic function $f : \mathfrak{S}_2 \rightarrow \mathbb{C}$ is a Picard modular form of $K = \mathbb{Q}(\sqrt{-3})$ and of weight k , if it satisfies:

$$(2.15) \quad f(\gamma(Z)) = \text{Jac}(\gamma, Z)^{-k} f(Z) \quad \text{for all } \gamma \in \Gamma,$$

where $\text{Jac}(\gamma, Z)$ is the Jacobi determinant of $\gamma : \mathfrak{S}_2 \rightarrow \mathfrak{S}_2$ at $Z = (z_1, z_2)$.

Now, for the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$, we define the Eisenstein series of weight k :

$$(2.16) \quad E_k(Z) = \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} \text{Jac}(\gamma, Z)^k, \quad \text{for } k > 1.$$

In fact, in [BB] and [Sh1], the authors studied the Poincaré-Eisenstein series.

The following proposition is important.

PROPOSITION 2.9. $j(g, Z)$ satisfies the following identity:

$$(2.17) \quad j(g_1 g_2, Z) = j(g_1, g_2(Z)) j(g_2, Z).$$

Proof. For $g_1 = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2, 1)$ and $g_2 = \begin{pmatrix} a'_1 & a'_2 & a'_3 \\ b'_1 & b'_2 & b'_3 \\ c'_1 & c'_2 & c'_3 \end{pmatrix} \in U(2, 1)$, we have

$$g_1 g_2 = \begin{pmatrix} * & * & * \\ * & * & * \\ a'_1 c_1 + b'_1 c_2 + c'_1 c_3 & a'_2 c_1 + b'_2 c_2 + c'_2 c_3 & a'_3 c_1 + b'_3 c_2 + c'_3 c_3 \end{pmatrix}.$$

Thus,

$$j(g_1 g_2, Z) = (a'_1 c_1 + b'_1 c_2 + c'_1 c_3) z_1 + (a'_2 c_1 + b'_2 c_2 + c'_2 c_3) z_2 + (a'_3 c_1 + b'_3 c_2 + c'_3 c_3).$$

On the other hand.

$$g_2(Z) = \left(\frac{a'_1 z_1 + a'_2 z_2 + a'_3}{c'_1 z_1 + c'_2 z_2 + c'_3}, \frac{b'_1 z_1 + b'_2 z_2 + b'_3}{c'_1 z_1 + c'_2 z_2 + c'_3} \right).$$

$$\begin{aligned} j(g_1, g_2(Z)) &= c_1 \frac{a'_1 z_1 + a'_2 z_2 + a'_3}{c'_1 z_1 + c'_2 z_2 + c'_3} + c_2 \frac{b'_1 z_1 + b'_2 z_2 + b'_3}{c'_1 z_1 + c'_2 z_2 + c'_3} + c_3. \\ j(g_2, Z) &= c'_1 z_1 + c'_2 z_2 + c'_3. \end{aligned}$$

So,

$$j(g_1, g_2(Z))j(g_2, Z) = j(g_1 g_2, Z).$$

□

3. Product formulas on $U(2, 1)$

In his paper [Ko], the Poisson kernel $P : B \times D \rightarrow \mathbb{R}$ of D is defined by

$$P(u, z) = \frac{|S(u, z)|^2}{S(z, z)}$$

where $D = \{(z_1, z_2) | \operatorname{Im} z_1 - \Phi(z_2, z_1) \in \Omega\}$ and B is the distinguished boundary of D :

$$B = \{(z_1, z_2) | \operatorname{Im} z_1 - \Phi(z_2, z_1) = 0\}.$$

The Szegö kernel function S is given by

$$S(z, w) = \frac{1}{(2\pi)^{n_1}} \int_{\Omega'} e^{-<\alpha, \rho(z, w)>} L(\alpha)^{-1} d\alpha,$$

where

$$\rho(z, w) = \frac{z_1 - \overline{w_1}}{i} - 2\Phi(z_2, w_2)$$

for $z = (z_1, z_2)$, $w = (w_1, w_2)$, and

$$L(\alpha) = \left(\frac{i}{2}\right)^{n_2} \int_{V_2} e^{-2<\alpha, \Phi(z_2, z_2)>} dz_2 d\overline{z}_2.$$

Now, we define the Poisson kernel as follows:

$$(3.1) \quad P(Z, W) = \frac{\rho(Z)}{|\rho(Z, W)|^2}, \quad \text{for } Z = (z_1, z_2) \in \mathfrak{S}_2, W = (w_1, w_2) \in \partial \mathfrak{S}_2,$$

where

$$\rho(Z, W) := \overline{z_1} + w_1 - \overline{z_2} w_2.$$

In fact, for $Z = (\frac{\rho+|z|^2}{2} + it, z)$ and $Z' = (\frac{\rho'+|z'|^2}{2} + it', z')$, where $\rho > 0$, $\rho' > 0$, $t, t' \in \mathbb{R}$ and $z, z' \in \mathbb{C}$, we have

$$|\rho(Z, Z')|^2 = \frac{1}{4}(\rho + \rho' + |z - z'|^2)^2 + (t' - t + \operatorname{Im} z \overline{z'})^2 \geq 0.$$

$\rho(Z, Z') = 0$ if and only if $Z = Z' \in \partial\mathfrak{S}_2$.

In their papers [Ko], [KW], Korányi and Wolf studied the realization of Siegel domain of type II and the Poisson integral. Our definition has a little difference from theirs. We have

$$(3.2) \quad LP(Z, W)^s = s(s-2)P(Z, W)^s - s^2\rho(W)P(Z, W)^{s+1} = s(s-2)P(Z, W)^s.$$

In fact, by Helgason's conjecture, which was proved by Kashiwara et al. in [K], that the eigenfunctions on Riemannian symmetric spaces can be represented as Poisson integrals of their hyperfunction boundary values.

The boundary of \mathfrak{S}_2 :

$$(3.3) \quad \partial\mathfrak{S}_2 = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 + \overline{z_1} = z_2\overline{z_2}\}.$$

In the homogeneous coordinates, one has

$$\overline{\partial\mathfrak{S}_2} = \{(z_1, z_2, z_3) \in \mathbb{C}^3 : z_1\overline{z_3} + \overline{z_1}z_3 = z_2\overline{z_2}\}.$$

In particular,

$$\overline{\partial\mathfrak{S}_2}(\mathcal{O}) = \{(a_1, a_2, a_3) \in \mathcal{O}^3 : a_1\overline{a_3} + \overline{a_1}a_3 = a_2\overline{a_2}\}.$$

The Green function associated to $Lf = 0$ on \mathfrak{S}_2 is defined as

$$(3.4) \quad G(Z, W) = \log|\rho(Z, W)| \quad \text{for } Z, W \in \mathfrak{S}_2.$$

It satisfies that $LG(Z, W) = 0$.

If $g = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2, 1)$, then $g^*Jg = J$, i.e.,

$$\begin{cases} -\overline{c_1}a_1 + \overline{b_1}b_1 - \overline{a_1}c_1 = 0, \\ -\overline{c_2}a_1 + \overline{b_2}b_1 - \overline{a_2}c_1 = 0, \\ -\overline{c_2}a_2 + \overline{b_2}b_2 - \overline{a_2}c_2 = 1, \\ -\overline{c_3}a_1 + \overline{b_3}b_1 - \overline{a_3}c_1 = -1, \\ -\overline{c_3}a_2 + \overline{b_3}b_2 - \overline{a_3}c_2 = 0, \\ -\overline{c_3}a_3 + \overline{b_3}b_3 - \overline{a_3}c_3 = 0. \end{cases}$$

By a straightforward calculation, we have

$$\rho(g(Z), g(W)) = \frac{\overline{z_1} + w_1 - \overline{z_2}w_2}{(c_1z_1 + c_2z_2 + c_3)(c_1w_1 + c_2w_2 + c_3)}.$$

Hence,

$$(3.5) \quad \rho(g(Z), g(W))\overline{j(g, Z)}j(g, W) = \rho(Z, W), \quad \text{for } g \in G.$$

Therefore,

$$(3.6) \quad P(g(Z), g(W)) = |j(g, W)|^2 P(Z, W).$$

By Proposition 2.9, we have

$$(3.7) \quad P(g(Z), W) = |j(g, g^{-1}(W))|^2 P(Z, g^{-1}(W)) = |j(g^{-1}, W)|^{-2} P(Z, g^{-1}(W)).$$

The point-pair invariant on \mathfrak{S}_2 is defined by

$$(3.8) \quad \sigma(Z, Z') := \frac{|\rho(Z, Z')|^2}{\rho(Z)\rho(Z')},$$

which is invariant under the action of $U(2, 1)$. The critical exponent is given by

$$(3.9) \quad \delta(\Gamma) = \inf\{s > 0 : \sum_{\gamma \in \Gamma} \sigma(Z, \gamma(Z'))^{-s} < \infty\}.$$

The number $\delta(\Gamma)$ does not depend on the choice of Z, Z' . It is an invariant of the group Γ .

The Eisenstein series is given by

$$(3.10) \quad E(Z, W; s) := \sum_{\gamma \in \Gamma} P(\gamma(Z), W)^s, \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma),$$

where Γ is a discrete subgroup of $U(2, 1)$. We have the following lemma:

LEMMA 3.1. *The properties of Eisenstein series:*

- (1) $LE(Z, W; s) = s(s - 2)E(Z, W; s)$.
- (2) $E(\gamma(Z), W; s) = E(Z, W; s)$.
- (3) $E(Z, \gamma(W); s) = |j(\gamma, W)|^{2s}E(Z, W; s)$ for $\gamma \in \Gamma$.

We have the following lemma:

LEMMA 3.2. *Let $f = f(u)$ with $u = u(Z, Z') = \sigma(Z, Z') - 1$, then*

$$(3.11) \quad Lf = u(u + 1)f''(u) + (3u + 2)f'(u).$$

By $Lf = s(s-2)f$, we have

$$(3.12) \quad u(u+1)f''(u) + (3u+2)f'(u) + s(2-s)f(u) = 0.$$

A solution is

$$(3.13) \quad \varphi_s(u(Z, Z')) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})} \left(\frac{1}{4}\right)^{s-1} u^{-s} F(s, s-1; 2s-1; -u^{-1}).$$

Here $F = {}_2F_1$ is the Gauss hypergeometric function.

Let

$$(3.14) \quad G(Z, Z'; s) = \sum_{\gamma \in \Gamma} r(Z, \gamma(Z'); s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma),$$

where $r(Z, Z'; s) = \varphi_s(u(Z, Z'))$. By the integral representation of hypergeometric functions, we can only consider the series

$$\sum_{\gamma \in \Gamma} \frac{1}{[1 + u(Z, \gamma(Z'))]^s} = \sum_{\gamma \in \Gamma} \sigma(Z, \gamma(Z'))^{-s},$$

which is convergent for $\operatorname{Re}(s) > \delta(\Gamma)$. Thus, the right hand side of (3.14) is convergent when $\operatorname{Re}(s) > \delta(\Gamma)$. We have

$$\lim_{\rho(Z) \rightarrow 0} \rho(Z)^{-s} r(Z, Z'; s) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})} \left(\frac{1}{4}\right)^{s-1} P(Z', W)^s,$$

where $Z \rightarrow W \in \partial \mathfrak{S}_2$ as $\rho(Z) \rightarrow 0$. Consequently,

$$\lim_{\rho(Z) \rightarrow 0} \rho(Z)^{-s} G(Z, Z'; s) = \frac{\Gamma(s)}{\sqrt{\pi}\Gamma(s-\frac{1}{2})} \left(\frac{1}{4}\right)^{s-1} E(Z', W; s).$$

In the real hyperbolic geometry, the double transitivity of the action of the isometric groups on the real hyperbolic space has been studied extensively (see [ElGM] for three dimensional hyperbolic space). Now, we give the corresponding results on the complex hyperbolic space.

PROPOSITION 3.3. *The group $U(2, 1)$ acts in the following sense doubly transitively on \mathfrak{S}_2 : For all $P, P', Q, Q' \in \mathfrak{S}_2$, such that $d(P, P') = d(Q, Q')$, there exists an element $g \in U(2, 1)$ such that $g(P) = Q$, $g(P') = Q'$.*

Proof. Set $P = (z_1, z_2)$ with $z_1 = -\omega\rho + \frac{|z|^2}{2} + it$, $z_2 = z$, where $\rho > 0$, $t \in \mathbb{R}$ and $z \in \mathbb{C}$. Then $\rho(z_1, z_2) = \rho$.

$$T_1 := \begin{pmatrix} \frac{1}{\sqrt{\rho}} & & \\ & 1 & \\ & & \sqrt{\rho} \end{pmatrix} \begin{pmatrix} 1 & -\bar{z} & \frac{|z|^2}{2} - it \\ & 1 & -z \\ & & 1 \end{pmatrix} \in U(2, 1).$$

Moreover, $T_1(z_1, z_2) = (-\omega, 0)$, i.e. T_1 maps P onto $(-\omega, 0)$. Set $(z'_1, z'_2) = (-\omega\rho' + \frac{|z'|^2}{2} + it', z')$, then

$$T_1(P') = \left(-\omega \frac{\rho'}{\rho} + \frac{1}{2} \left(\frac{|z' - z|}{\sqrt{\rho}} \right)^2 + i \frac{t' - t + \operatorname{Im}(z\bar{z}')}{\rho}, \frac{z' - z}{\sqrt{\rho}} \right),$$

i.e., $T_1(P')$ has the form of $(-\omega\rho + \frac{|z|^2}{2} + it, z)$.

If $k(-\omega\rho + \frac{|z|^2}{2} + it, z) = (-\omega\rho', 0)$, where $k \in K$, then we obtain two complex equations about $\alpha, \beta, \gamma, \delta$ and θ . In fact, the number of real linear equations is four. While, the real dimension of K is five. Hence, the existence of k is proved. Applying a suitable element T_2 of the stabilizer K of $(-\omega, 0)$ in $U(2, 1)$ to $T_1(P')$ we get a transform $T = T_2 T_1$ such that $T(P) = (-\omega, 0)$, $T(P') = (-\lambda\omega, 0)$ with $\lambda \geq 1$. Since the isometric group of \mathfrak{S}_2 is $U(2, 1)$, we have

$$d(P, P') = d(T(P), T(P')) = d((- \omega, 0), (-\lambda\omega, 0)) = \int_1^\lambda \frac{dr}{r} = \log \lambda.$$

So $\lambda = \exp(d(P, P'))$. Similarly, there exists an element $S \in U(2, 1)$ such that $S(Q) = (-\omega, 0)$, $S(Q') = (-\lambda\omega, 0)$ with the same λ , since $d(P, P') = d(Q, Q')$. Thus, $g := S^{-1}T$ has the desired properties. □

In [ElGM], the concept of real hyperbolic distance was introduced and the basic properties was studied. Now, we give the complex hyperbolic distance and obtain the triangle inequality for our function δ .

PROPOSITION 3.4. *The complex hyperbolic distance $d(P, P')$ is given by*

$$(3.15) \quad \cosh d(P, P') = \delta(P, P'),$$

where δ is defined by

$$(3.16) \quad \delta(P, P') = \delta(Z, Z') = \frac{1}{2}[\sigma(Z, Z') + 1],$$

and $P = Z$, $P' = Z'$.

Proof. If $P = (-\omega, 0)$ and $P' = (-\lambda\omega, 0)$ ($\lambda \geq 1$), we have

$$d((- \omega, 0), (-\lambda\omega, 0)) = \int_1^\lambda \frac{dr}{r} = \log \lambda.$$

$$\delta((- \omega, 0), (-\lambda\omega, 0)) = \frac{1}{2}(\lambda + \frac{1}{\lambda}) = \cosh d((- \omega, 0), (-\lambda\omega, 0)).$$

Thus the proposition is true in the special case $P = (-\omega, 0)$, $P' = (-\lambda\omega, 0)$. Note that δ is a point-pair invariant. Since $U(2, 1)$ acts doubly transitively on \mathfrak{S}_2 , there exists for all $P, P' \in \mathfrak{S}_2$ an element $g \in U(2, 1)$, such that $g(P) = (-\omega, 0)$, $g(P') = (-\lambda\omega, 0)$, $\lambda = \exp(d(P, P'))$. Therefore, by point-pair invariance, we have

$$\begin{aligned} \cosh d(P, P') &= \cosh d((- \omega, 0), (-\lambda\omega, 0)) = \delta((- \omega, 0), (-\lambda\omega, 0)) = \delta(g(P), g(P')) \\ &= \delta(P, P'). \end{aligned}$$

□

PROPOSITION 3.5. *For $g = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2, 1)$, the following formulas hold:*

(1)

$$(3.17) \quad \begin{aligned} &\delta(P, g(P)) + \delta(Q, g(Q)) + \delta(P, g(Q)) + \delta(Q, g(P)) \\ &= \frac{5}{2} + 2(|a_1|^2 + |a_3|^2 + |c_1|^2 + |c_3|^2) + (|a_2|^2 + |c_2|^2 + |b_1|^2 + |b_3|^2) + \frac{1}{2}|b_2|^2, \end{aligned}$$

where $P = (-\omega, 0)$ and $Q = (-\bar{\omega}, 0)$.

(2)

$$(3.18) \quad \delta((1, 0), g(1, 0)) = \frac{5}{8} + \frac{1}{8} \sum_{j=1}^3 (|a_j|^2 + |b_j|^2 + |c_j|^2).$$

Proof. It is obtained by a straightforward calculation. □

PROPOSITION 3.6. *The triangle inequality for the metric function δ on the complex hyperbolic space: For all $P, Q, R \in \mathfrak{S}_2$, the following inequality holds:*

$$(3.19) \quad \frac{1}{72} \frac{\delta(Q, R)}{\delta(P, Q)} \leq \delta(P, R) \leq 72\delta(P, Q)\delta(Q, R).$$

Proof. Since δ is a point-pair invariant, it is sufficient to prove (3.19) only in the special case $Q = (\frac{1}{2}, 0)$. Set $P = (\frac{\rho_1 + |z_1|^2}{2} + it_1, z_1)$ and $R = (\frac{\rho_2 + |z_2|^2}{2} + it_2, z_2)$. Then

$$\delta(P, Q) = \frac{\frac{1}{4}(1 + \rho_1 + |z_1|^2)^2 + t_1^2 + \rho_1}{2\rho_1}, \quad \delta(Q, R) = \frac{\frac{1}{4}(1 + \rho_2 + |z_2|^2)^2 + t_2^2 + \rho_2}{2\rho_2},$$

and

$$\delta(P, R) = \frac{1}{8\rho_1\rho_2} [(\rho_1 + \rho_2 + |z_1 - z_2|^2)^2 + 4(t_2 - t_1 + \text{Im}z_1\bar{z}_2)^2 + 4\rho_1\rho_2].$$

Note that $|z_1 - z_2|^2 \leq 2(|z_1|^2 + |z_2|^2)$ and $|\text{Im}z_1\overline{z_2}| \leq |z_1||z_2| \leq \frac{1}{2}(|z_1|^2 + |z_2|^2)$, we have

$$8\rho_1\rho_2\delta(P, R) \leq (\rho_1 + \rho_2 + 2|z_1|^2 + 2|z_2|^2)^2 + (2|t_1| + 2|t_2| + |z_1|^2 + |z_2|^2)^2 + 4\rho_1\rho_2.$$

By $2|t_1||t_2| \leq t_1^2 + t_2^2$ and $2|t_i||z_j|^2 \leq t_i^2 + |z_j|^4$, $i, j = 1, 2$, we have

$$\begin{aligned} & 8\rho_1\rho_2\delta(P, R) \\ & \leq \rho_1^2 + \rho_2^2 + 9|z_1|^4 + 9|z_2|^4 + 6\rho_1\rho_2 + 4(\rho_1 + \rho_2)(|z_1|^2 + |z_2|^2) + 10|z_1|^2|z_2|^2 \\ & \quad + 12t_1^2 + 12t_2^2 \\ & \leq 9(1 + \rho_1^2 + |z_1|^4 + 6\rho_1 + 2|z_1|^2 + 2\rho_1|z_1|^2 + 4t_1^2) \\ & \quad \times (1 + \rho_2^2 + |z_2|^4 + 6\rho_2 + 2|z_2|^2 + 2\rho_2|z_2|^2 + 4t_2^2) \\ & = 576\delta(P, Q)\delta(Q, R). \end{aligned}$$

Hence, $\delta(P, R) \leq 72\delta(P, Q)\delta(Q, R)$. Interchanging P and Q , we obtain the other part of the inequality. \square

THEOREM 3.7. *Let*

$$(3.20) \quad \phi(Z, Z'; s) = \int_{\partial\mathfrak{S}_2} P(Z, W)^s P(Z', W)^{2-s} dm(W).$$

Then one has the following formula:

$$(3.21) \quad \phi(Z, Z'; s) = -\frac{\pi}{s-1} \frac{\sqrt{\pi}\Gamma(\frac{3}{2}-s)}{\Gamma(2-s)} \left(\frac{1}{4}\right)^{s-1} u^{-s} F(s, s-1; 2s-1; -u^{-1}) + (s \mapsto 2-s),$$

where $dm(W)$ is the Lebesgue measure on $\partial\mathfrak{S}_2$, $u = u(Z, Z')$ is the point-pair invariant, and $(s \mapsto 2-s)$ is equal to the first term on the right hand side with s replaced by $2-s$.

Proof. $\phi(Z, Z'; s)$ is a point-pair invariant. Thus, it suffices to calculate it for $Z = (z, t, \rho) = (0, 0, \rho)$ and $Z' = (z', t', \rho') = (0, 0, \rho')$, i.e., $Z = (\frac{1}{2}\rho, 0)$ and $Z' = (\frac{1}{2}\rho', 0)$. Let ρ and ρ' stand for $(\rho, 0)$ and $(\rho', 0)$, respectively. Now, $\rho(Z, Z') = \frac{1}{2}(\rho + \rho')$ and

$$\sigma(Z, Z') = \frac{(\rho + \rho')^2}{4\rho\rho'} = \frac{(\lambda + 1)^2}{4\lambda}, \quad u = \frac{(\lambda - 1)^2}{4\lambda},$$

where $\lambda = \rho/\rho'$. For $W \in \partial\mathfrak{S}_2$, $w_1 = \frac{1}{2}|w|^2 + iv$ and $w_2 = w$, where $w \in \mathbb{C}$, $v \in \mathbb{R}$. $\rho(Z, W) = \frac{1}{2}(\rho + |w|^2) + iv$ and $\rho(Z', W) = \frac{1}{2}(\rho' + |w|^2) + iv$. Hence,

$$\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; s\right) = \int_{\mathbb{C}} \int_{\mathbb{R}} \frac{\rho^s \rho'^{2-s}}{|\frac{1}{2}(\rho + |w|^2) + iv|^{2s} |\frac{1}{2}(\rho' + |w|^2) + iv|^{2(2-s)}} dm(w) dv.$$

By transform $w = re^{i\theta}$ with $r \geq 0, \theta \in [0, 2\pi)$ and $u = r^2$, we have

$$\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; s\right) = 16\pi\rho^s\rho'^{2-s} \int_0^\infty \int_0^\infty \frac{dudv}{[(\rho+u)^2+v^2]^s[(\rho'+u)^2+v^2]^{2-s}}.$$

Set $v = (\rho' + u)\sqrt{w}$, then

$$\int_0^\infty \frac{dv}{[(\rho+u)^2+v^2]^s[(\rho'+u)^2+v^2]^{2-s}} = \frac{1}{2}(\rho'+u)^{-3} \int_0^\infty w^{-\frac{1}{2}}(1+w)^{s-2}(l^2+w)^{-s}dw,$$

where $l = l(u) = \frac{\rho+u}{\rho'+u}$.

By (see [Er], p.115, (5))

$$F(a, b; c; 1-z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^\infty s^{b-1}(1+s)^{a-c}(1+sz)^{-a}ds$$

for $\operatorname{Re}(c) > \operatorname{Re}(b) > 0$, $|\arg(z)| < \pi$, we have

$$\int_0^\infty w^{-\frac{1}{2}}(1+w)^{s-2}(1+wl^{-2})^{-s}dw = \frac{\pi}{2}F(s, \frac{1}{2}; 2; 1-l^{-2}).$$

Now, we need the following two formulas:

$$(1) z = 1 - l^{-2}, \frac{z}{z-1} = 1 - l^2.$$

By (see [Er], p.105, 2.9.(3))

$$F(a, b; c; z) = (1-z)^{-a}F(a, c-b; c; \frac{z}{z-1}),$$

one has

$$F(s, \frac{1}{2}; 2; 1-l^{-2}) = l^{2s}F(s, \frac{3}{2}; 2; 1-l^2).$$

$$(2) z = 1 - l^2, 1 - z = l^2.$$

By (see [Er], p.108, 2.10.(1))

$$\begin{aligned} F(a, b; c; z) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}F(a, b; a+b-c+1; 1-z) \\ &\quad + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}(1-z)^{c-a-b}F(c-a, c-b; c-a-b+1; 1-z), \end{aligned}$$

where $|\arg(1-z)| < \pi$, one has

$$\begin{aligned} F(s, \frac{3}{2}; 2; 1-l^2) &= \frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}F(s, \frac{3}{2}; s+\frac{1}{2}; l^2) \\ &\quad + \frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}l^{1-2s}F(2-s, \frac{1}{2}; \frac{3}{2}-s; l^2). \end{aligned}$$

Hence, $\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; s) = \phi_I + \phi_{II}$, where

$$\phi_I = 4\pi^2 \lambda^s \rho'^2 \frac{\Gamma(\frac{1}{2} - s)}{\sqrt{\pi} \Gamma(2 - s)} \int_0^\infty (\rho' + u)^{-3} F(s, \frac{3}{2}; s + \frac{1}{2}; l^2) du,$$

$$\phi_{II} = 4\pi^2 \lambda^s \rho'^2 \frac{2\Gamma(s - \frac{1}{2})}{\sqrt{\pi} \Gamma(s)} \int_0^\infty (\rho' + u)^{-3} l^{1-2s} F(2 - s, \frac{1}{2}; \frac{3}{2} - s; l^2) du.$$

$l = \frac{\rho+u}{\rho'+u}$ implies that $u = \frac{\rho-\rho'l}{l-1}$. Thus, $du = \frac{\rho'-\rho}{(l-1)^2} dl$ and $\rho' + u = \frac{\rho-\rho'}{l-1}$. Without loss of generality, we can assume that $\rho < \rho'$. Then

$$\begin{aligned} \phi_I &= -\frac{4\pi^2 \lambda^s}{(1-\lambda)^2} \frac{\Gamma(\frac{1}{2} - s)}{\sqrt{\pi} \Gamma(2 - s)} \int_\lambda^1 (l-1) F(s, \frac{3}{2}; s + \frac{1}{2}; l^2) dl, \\ \phi_{II} &= -\frac{4\pi^2 \lambda^s}{(1-\lambda)^2} \frac{2\Gamma(s - \frac{1}{2})}{\sqrt{\pi} \Gamma(s)} \int_\lambda^1 (l-1) l^{1-2s} F(2 - s, \frac{1}{2}; \frac{3}{2} - s; l^2) dl. \end{aligned}$$

We have

$$\begin{aligned} &\int_\lambda^1 (l-1) F(s, \frac{3}{2}; s + \frac{1}{2}; l^2) dl \\ &= \sum_{k=0}^{\infty} \frac{(s)_k (\frac{3}{2})_k}{k! (s + \frac{1}{2})_k} \left(\frac{\lambda^{2k+1}}{2k+1} - \frac{\lambda^{2k+2}}{2k+2} - \frac{1}{(2k+1)(2k+2)} \right) \\ &= \lambda F(s, \frac{1}{2}; s + \frac{1}{2}; \lambda^2) - \frac{2s-1}{2s-2} F(s-1, \frac{1}{2}; s - \frac{1}{2}; \lambda^2) + \frac{1}{s-1} \frac{\Gamma(s + \frac{1}{2})}{\sqrt{\pi} \Gamma(s)}. \end{aligned}$$

Here we use the following formula:

$$F(s-1, -\frac{1}{2}; s - \frac{1}{2}; 1) = \frac{\Gamma(s - \frac{1}{2})}{\sqrt{\pi} \Gamma(s)}, \quad \text{for } \operatorname{Re}(s) > 0.$$

Similarly,

$$\begin{aligned} &\int_\lambda^1 (l-1) l^{1-2s} F(2 - s, \frac{1}{2}; \frac{3}{2} - s; l^2) dl \\ &= \frac{\lambda^{2-2s}}{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2} - s; \lambda^2) - \frac{\lambda^{3-2s}}{3-2s} F(2-s, \frac{1}{2}; \frac{5}{2} - s; \lambda^2) - \frac{1}{2-2s} \frac{\Gamma(\frac{3}{2} - s)}{\sqrt{\pi} \Gamma(2 - s)}, \end{aligned}$$

for $\operatorname{Re}(s) < 2$.

Set $z = -\lambda$, then $\frac{4z}{(1+z)^2} = -u^{-1}$. By (see [Er], p.111, 2.11.(5))

$$F(a, b; 2b; \frac{4z}{(1+z)^2}) = (1+z)^{2a} F(a, a + \frac{1}{2} - b; b + \frac{1}{2}; z^2),$$

we have

$$\begin{aligned} F(s, \frac{1}{2}; s + \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2s} F(s, s; 2s; -u^{-1}), \\ F(s-1, \frac{1}{2}; s - \frac{1}{2}; \lambda^2) &= (1-\lambda)^{2-2s} F(s-1, s-1; 2s-2; -u^{-1}), \\ F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) &= (1-\lambda)^{2s-2} F(1-s, 1-s; 2-2s; -u^{-1}), \\ F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2) &= (1-\lambda)^{2s-4} F(2-s, 2-s; 4-2s; -u^{-1}). \end{aligned}$$

Thus,

$$\begin{aligned} \phi_I &= -4\pi^2 \frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} (4u)^{-s-1} F(s, s; 2s; -u^{-1}) \\ &\quad - \frac{4\pi^2}{s-1} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} (4u)^{-s} F(s-1, s-1; 2s-2; -u^{-1}) \\ &\quad - \frac{4\pi}{s-1} \frac{\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2})}{\Gamma(s)\Gamma(2-s)} \lambda^s (1-\lambda)^{-2}, \end{aligned}$$

and

$$\begin{aligned} \phi_{II} &= -\frac{4\pi^2}{1-s} \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} (4u)^{s-2} F(1-s, 1-s; 2-2s; -u^{-1}) \\ &\quad - 4\pi^2 \frac{\Gamma(s-\frac{3}{2})}{\sqrt{\pi}\Gamma(s)} (4u)^{s-3} F(2-s, 2-s; 4-2s; -u^{-1}) \\ &\quad - \frac{4\pi}{s-1} \frac{\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)}{\Gamma(s)\Gamma(2-s)} \lambda^s (1-\lambda)^{-2}. \end{aligned}$$

Note that

$$\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2}) + \Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s) = (s-\frac{1}{2})\Gamma(\frac{1}{2}-s)\Gamma(s-\frac{1}{2}) + (\frac{1}{2}-s)\Gamma(s-\frac{1}{2})\Gamma(\frac{1}{2}-s) = 0.$$

Moreover, we need the following lemma.

LEMMA 3.8. *The following formula about hypergeometric functions holds:*

$$\gamma F - \beta z F(\beta+1, \gamma+1) - \gamma F(\alpha-1) = 0,$$

where F denotes $F(\alpha, \beta; \gamma; z)$, $F(\beta+1, \gamma+1)$ stands for $F(\alpha, \beta+1; \gamma+1; z)$ and $F(\alpha-1)$ stands for $F(\alpha-1, \beta; \gamma; z)$.

Proof. It is obtained by the integral representation of hypergeometric functions. \square

By Lemma 3.8 and the following formula (see [Er], p.103, (35)):

$$(\gamma - 1)F(\gamma - 1) - \alpha F(\alpha + 1) - (\gamma - \alpha - 1)F = 0.$$

we get

$$(2s - 1)F(s, s - 1; 2s - 1; z) - (s - 1)zF(s, s; 2s; z) - (2s - 1)F(s - 1, s - 1; 2s - 1; z) = 0,$$

$$(2s - 2)F(s - 1, s - 1; 2s - 2; z) - (s - 1)F(s, s - 1; 2s - 1; z) - (s - 1)F(s - 1, s - 1; 2s - 1; z) = 0.$$

Thus, we have

$$\frac{4s - 2}{s - 1}F(s, s - 1; 2s - 1; z) = zF(s, s; 2s; z) + \frac{4s - 2}{s - 1}F(s - 1, s - 1; 2s - 2; z).$$

Substituting this identity to the expression of $\phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; s)$, We get the desired results. \square

Consequently, we have

COROLLARY 3.9. *The following formula holds:*

$$(3.22) \quad \phi(Z, Z'; s) = -\frac{\pi}{s - 1}c(s)c(2 - s)[r(Z, Z'; s) - r(Z, Z'; 2 - s)],$$

where

$$c(s) = \frac{\sqrt{\pi}\Gamma(s - \frac{1}{2})}{\Gamma(s)}.$$

COROLLARY 3.10. *The functional equation for the trivial group on \mathfrak{S}_2 : If $\text{Re}(s) > 1$, then the following identity holds:*

$$(3.23) \quad \int_{\partial\mathfrak{S}_2} P(Z, W)^s |\rho(W, W')|^{-2(2-s)} dm(W) = \frac{\pi}{s - 1} \frac{\sqrt{\pi}\Gamma(s - \frac{1}{2})}{\Gamma(s)} 4^{s-1} P(Z, W')^{2-s},$$

where $W' \in \partial\mathfrak{S}_2$.

Proof. The integral on the left-hand side is absolutely convergent. We set

$$P(Z', W) = \frac{\rho(Z')}{|\rho(Z', W)|^2}$$

in the formula of Theorem 3.7. Next, we multiply both sides of the formula by $\rho(Z')^{s-2}$ and take the limit as $Z' \rightarrow W' \in \partial\mathfrak{S}_2$. Thus we get the desired results. \square

PROPOSITION 3.11. *The integral formula for the Poisson kernel:*

$$(3.24) \quad \int_{\partial\mathfrak{S}_2} P(Z, W)^s dm(W) = \frac{\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} 4^{s-1} \rho^{2-s}, \quad \text{for } \operatorname{Re}(s) > 1.$$

Proof. Denote the left hand side of (3.24) as $f(Z)$, then

$$f(Z) = \rho^s \int_{\partial\mathfrak{S}_2} |\overline{z_1} + w_1 - \overline{z_2}w_2|^{-2s} dm(W).$$

Set $z_1 = \frac{\rho+|z|^2}{2} + ir$, $z_2 = z$, $w_1 = \frac{|w|^2}{2} + it$ and $w_2 = w$, where $z, w \in \mathbb{C}$, $t, r \in \mathbb{R}$, we have

$$\begin{aligned} f(Z) &= \rho^s \int_{\mathbb{C}} \int_{\mathbb{R}} \left| \frac{\rho + |w - z|^2}{2} + i(t - r + \operatorname{Im}(z\bar{w})) \right|^{-2s} dm(w) dt \\ &= \rho^s \int_{\mathbb{C}} \int_{\mathbb{R}} \left| \frac{\rho + |w|^2}{2} + it \right|^{-2s} dm(w) dt \\ &= 2\pi\rho^s \int_0^\infty \int_{-\infty}^\infty \frac{rdrdt}{[\frac{1}{4}(\rho+r^2)^2+t^2]^s} \\ &= 2^{2s-1}\pi\rho^s \int_0^\infty \int_{-\infty}^\infty \frac{drdu}{[(\rho+r)^2+u^2]^s}. \end{aligned}$$

By the well known formula (see [Ku], p.15):

$$\int_{-\infty}^\infty \frac{dt}{(1+t^2)^s} = \sqrt{\pi} \frac{\Gamma(s-\frac{1}{2})}{\Gamma(s)}, \quad \text{for } \operatorname{Re}(s) > 1,$$

we have

$$f(Z) = \frac{\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} 4^{s-1} \rho^{2-s}.$$

□

We define the S-matrix as follows:

$$(3.25) \quad S(W, W'; s) := \sum_{\gamma \in \Gamma} |\rho(\gamma(W), W')|^{-2s} |j(\gamma, W)|^{-2s}$$

for $W, W' \in \Omega(\Gamma)$ and $\operatorname{Re}(s) > \delta(\Gamma)$.

THEOREM 3.12. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following functional equation for Eisenstein series holds:*

$$(3.26) \quad \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) S(W, W'; 2-s) dm(W) = \frac{4^{s-1}\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} E(Z, W'; 2-s),$$

for $1 < \operatorname{Re}(s) < 2 - \delta(\Gamma)$.

Proof. By Corollary 3.10,

$$(3.27) \quad \int_{\partial\mathfrak{S}_2} E(Z, W; s) |\rho(W, W')|^{-2(2-s)} dm(W) = \frac{4^{s-1}\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} E(Z, W'; 2-s).$$

On the other hand,

$$\begin{aligned} \int_{\partial\mathfrak{S}_2} f(W) dW &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} f(\gamma(W)) dm(\gamma(W)) \\ &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |J(\gamma(W))| f(\gamma(W)) dm(W). \end{aligned}$$

Hence, the left hand side of (3.27):

$$\begin{aligned} LHS &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |J(\gamma(W))| |E(Z, \gamma(W); s)| |\rho(\gamma(W), W')|^{-2(2-s)} dm(W) \\ &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |j(\gamma, W)|^{-4} |j(\gamma, W)|^{2s} E(Z, W; s) |\rho(\gamma(W), W')|^{-2(2-s)} dm(W) \\ &= \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) S(W, W'; 2-s) dm(W). \end{aligned}$$

PROPOSITION 3.13. Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following product formula holds: \square

$$(3.28) \quad \begin{aligned} &\int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) E(Z', W; 2-s) dm(W) \\ &= -\frac{\pi}{s-1} c(s) c(2-s) [G(Z, Z'; s) - G(Z, Z'; 2-s)], \end{aligned}$$

where $\delta(\Gamma) < \operatorname{Re}(s) < 2 - \delta(\Gamma)$.

For $s \in \mathbb{C}$, similar as in [Ma1] and [Ma2], we set

$$L_s^2(\Gamma) := \{f(s, \cdot) : f(s, \cdot) \in C^\infty(\Omega(\Gamma)), f(s, \gamma(W)) j(\gamma, W)^{2-s} = f(s, W), \gamma \in \Gamma\}.$$

If $f(s, \cdot) \in L_s^2(\Gamma)$, we define the Eisenstein integral:

$$E(Z, \cdot, s) : L_s^2(\Gamma) \rightarrow C^\infty(\Gamma \setminus \mathfrak{S}_2),$$

$$(3.29) \quad E(Z, f, s) = \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; s) f(s, W) dm(W), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma),$$

and the scattering operator:

$$(3.30) \quad S_s(f)(z, t) := \lim_{Z \rightarrow (z, t)} \rho(Z)^{-s} [E(Z, f, s) - \frac{4^{s-1}\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} f(s, (z, t)) \rho(Z)^{2-s}],$$

where $\delta(\Gamma) < \operatorname{Re}(s) < 2$.

Now, the functional equation for Eisenstein series can be written in terms of Eisenstein integrals as

$$(3.31) \quad E(Z, S_{2-s}(f), s) = \frac{4^{s-1}\pi}{s-1} \frac{\sqrt{\pi}\Gamma(s-\frac{1}{2})}{\Gamma(s)} E(Z, f, 2-s),$$

and in terms involving the scattering operator as:

$$(3.32) \quad S_s S_{2-s} = -\frac{\pi^3}{(s-1)^2} \frac{\Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)}{\Gamma(s)\Gamma(2-s)} I.$$

4. Product formulas of weight k on $U(2, 1)$

In the case of $SL(2, \mathbb{R})$, many people, especially Maass [M], Selberg [Se], Roelcke [Ro], Fay [Fa], Elstrodt [El], Hejhal [H], and Shimura [Sh2] studied the eigenvalue problems which concern a homogeneous equation

$$(\Delta_k - \lambda)f(z) = 0.$$

Here f is a function on the complex upper plane

$$\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im}(z) > 0\},$$

and L_k is a differential operator given by

$$\Delta_k = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) - iky \frac{\partial}{\partial x},$$

where $z = x + iy$ and k is an integer. Let Γ be a congruence subgroup of $SL(2, \mathbb{Z})$ and assume that f is a Γ -automorphic form of weight k in the sense that it is invariant under

$$f \mapsto f(\gamma(z))(cz+d)^{-k} |cz+d|^k$$

for any $\gamma = \begin{pmatrix} * & * \\ c & d \end{pmatrix} \in \Gamma$. In fact, L_k commutes with the above map. One of the most remarkable facts about such λ and f is the existence of Selberg zeta function (see [Se]), whose set of zeros coincides essentially with the set $S(\Gamma, k)$ of $s \in \mathbb{C}$ such that $\lambda = s(1 - s)$ occurs as an eigenvalue of L_k as in the above equation with a cusp form f (see [Sh2]).

Now, we study the corresponding problems in the case of $U(2, 1)$.

PROPOSITION 4.1. *Let*

$$(4.1) \quad L_{(\alpha, \beta)} = (z_1 + \overline{z_1} - z_2 \overline{z_2}) \\ \times \left[(z_1 + \overline{z_1}) \frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2} \frac{\partial^2}{\partial z_1 \partial \overline{z_2}} + \beta \frac{\partial}{\partial z_1} + \alpha \frac{\partial}{\partial \overline{z_1}} \right]$$

and

$$(4.2) \quad E(Z, \overline{Z}; \alpha, \beta) = \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta},$$

where $\Gamma = U(2, 1; \mathcal{O})$ is the Picard modular group. Then the following formula holds:

$$(4.3) \quad L_{(\alpha, \beta)} E(Z, \overline{Z}; \alpha, \beta) = 0.$$

Proof. We have

$$\begin{aligned} & \left[(z_1 + \overline{z_1}) \frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2} \frac{\partial^2}{\partial z_1 \partial \overline{z_2}} \right] [j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta}] \\ &= \alpha \beta [a_1 j(\gamma, Z)^{-\alpha-1} \overline{j(\gamma, Z)}^{-\beta} + \overline{a_1} j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta-1}] \\ &= - \left(\beta \frac{\partial}{\partial z_1} + \alpha \frac{\partial}{\partial \overline{z_1}} \right) [j(\gamma, Z)^{-\alpha} \overline{j(\gamma, Z)}^{-\beta}]. \end{aligned}$$

Hence, $L_{(\alpha, \beta)} E(Z, \overline{Z}; \alpha, \beta) = 0$.

□

Set

$$(4.4) \quad E_{3k}(Z, \overline{Z}; \beta) = \rho(Z)^\beta E(Z, \overline{Z}; \alpha, \beta) = \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} \frac{\rho(Z)^\beta}{j(\gamma, Z)^{3k} |j(\gamma, Z)|^{2\beta}}$$

with $3k = \alpha - \beta$.

In particular, when $\beta = -k$ and $\alpha = k$, we have

$$\begin{aligned} L_k &= (z_1 + \overline{z_1} - z_2 \overline{z_2}) \\ &\times \left[(z_1 + \overline{z_1}) \frac{\partial^2}{\partial z_1 \partial \overline{z_1}} + \frac{\partial^2}{\partial z_2 \partial \overline{z_2}} + z_2 \frac{\partial^2}{\partial \overline{z_1} \partial z_2} + \overline{z_2} \frac{\partial^2}{\partial z_1 \partial \overline{z_2}} - k \left(\frac{\partial}{\partial z_1} - \frac{\partial}{\partial \overline{z_1}} \right) \right]. \end{aligned}$$

PROPOSITION 4.2. *The operator L_k has the following automorphic property:*

$$(4.5) \quad L_k[f(\gamma(Z))j(\gamma, Z)^{-k} \overline{j(\gamma, Z)}^k] = j(\gamma, Z)^{-k} \overline{j(\gamma, Z)}^k (L_k f)(\gamma(Z)),$$

for $\gamma \in U(2, 1)$ and $Z = (z_1, z_2) \in \mathbb{C}^2$, where $f = f(Z, \overline{Z})$ is a real analytic function. In other words, L_k commutes with the following map

$$(4.6) \quad f(Z) \mapsto f(\gamma(Z))j(\gamma, Z)^{-k} \overline{j(\gamma, Z)}^k.$$

Proof. For $\gamma = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix} \in U(2, 1)$, one has $\gamma J \gamma^* = J$, i.e.,

$$\begin{cases} -\overline{a_1}a_3 + a_2\overline{a_2} - a_1\overline{a_3} = 0, \\ -a_3\overline{b_1} + a_2\overline{b_2} - a_1\overline{b_3} = 0, \\ -a_3\overline{c_1} + a_2\overline{c_2} - a_1\overline{c_3} = -1, \\ -b_3\overline{b_1} + b_2\overline{b_2} - b_1\overline{b_3} = 1, \\ -b_3\overline{c_1} + b_2\overline{c_2} - b_1\overline{c_3} = 0, \\ -c_3\overline{c_1} + c_2\overline{c_2} - c_1\overline{c_3} = 0. \end{cases}$$

Denote

$$\gamma(z_1, z_2) = (u(z_1, z_2), v(z_1, z_2)) = \left(\frac{a_1 z_1 + a_2 z_2 + a_3}{c_1 z_1 + c_2 z_2 + c_3}, \frac{b_1 z_1 + b_2 z_2 + b_3}{c_1 z_1 + c_2 z_2 + c_3} \right).$$

Set

$$A_{1,1} := \frac{\partial}{\partial z_1} \left(\frac{a_1 z_1 + a_2 z_2 + a_3}{c_1 z_1 + c_2 z_2 + c_3} \right) = [(a_1 c_2 - a_2 c_1) z_2 + (a_1 c_3 - a_3 c_1)] j(\gamma, Z)^{-2}.$$

$$A_{1,2} := \frac{\partial}{\partial z_2} \left(\frac{a_1 z_1 + a_2 z_2 + a_3}{c_1 z_1 + c_2 z_2 + c_3} \right) = [(a_2 c_1 - a_1 c_2) z_1 + (a_2 c_3 - a_3 c_2)] j(\gamma, Z)^{-2}.$$

$$A_{2,1} := \frac{\partial}{\partial z_1} \left(\frac{b_1 z_1 + b_2 z_2 + b_3}{c_1 z_1 + c_2 z_2 + c_3} \right) = [(b_1 c_2 - b_2 c_1) z_2 + (b_1 c_3 - b_3 c_1)] j(\gamma, Z)^{-2}.$$

$$A_{2,2} := \frac{\partial}{\partial z_2} \left(\frac{b_1 z_1 + b_2 z_2 + b_3}{c_1 z_1 + c_2 z_2 + c_3} \right) = [(b_2 c_1 - b_1 c_2) z_1 + (b_2 c_3 - b_3 c_2)] j(\gamma, Z)^{-2}.$$

We have

$$\begin{aligned} L_k(\gamma(Z)) &= (u + \bar{u} - v\bar{v}) \\ &\quad \times \left[(u + \bar{u}) \frac{\partial^2}{\partial u \partial \bar{u}} + \frac{\partial^2}{\partial v \partial \bar{v}} + v \frac{\partial^2}{\partial \bar{u} \partial v} + \bar{v} \frac{\partial^2}{\partial u \partial \bar{v}} - k \left(\frac{\partial}{\partial u} - \frac{\partial}{\partial \bar{u}} \right) \right], \end{aligned}$$

while $u + \bar{u} - v\bar{v} = (z_1 + \bar{z}_1 - z_2 \bar{z}_2) |j(\gamma, Z)|^{-2}$.

On the other hand,

$$\begin{aligned} &L_k[f(\gamma(Z)) j(\gamma, Z)^{-k} \overline{j(\gamma, Z)}^k] \\ &= j(\gamma, Z)^{-k} \overline{j(\gamma, Z)}^k (z_1 + \bar{z}_1 - z_2 \bar{z}_2) [F_{1,1} \frac{\partial^2 f}{\partial u \partial \bar{u}} + F_{1,2} \frac{\partial^2 f}{\partial u \partial \bar{v}} + F_{2,1} \frac{\partial^2 f}{\partial \bar{u} \partial v} \\ &\quad + F_{2,2} \frac{\partial^2 f}{\partial v \partial \bar{v}} + H_1 \frac{\partial f}{\partial u} + H_2 \frac{\partial f}{\partial v} + G_1 \frac{\partial f}{\partial \bar{u}} + G_2 \frac{\partial f}{\partial \bar{v}}](\gamma(Z)), \end{aligned}$$

where

$$\begin{cases} F_{1,1} = (z_1 + \bar{z}_1) A_{1,1} \overline{A_{1,1}} + A_{1,2} \overline{A_{1,2}} + z_2 A_{1,2} \overline{A_{1,1}} + \bar{z}_2 A_{1,1} \overline{A_{1,2}}, \\ F_{1,2} = (z_1 + \bar{z}_1) A_{1,1} \overline{A_{2,1}} + A_{1,2} \overline{A_{2,2}} + z_2 A_{1,2} \overline{A_{2,1}} + \bar{z}_2 A_{1,1} \overline{A_{2,2}}, \\ F_{2,2} = (z_1 + \bar{z}_1) A_{2,1} \overline{A_{2,1}} + A_{2,2} \overline{A_{2,2}} + z_2 A_{2,2} \overline{A_{2,1}} + \bar{z}_2 A_{2,1} \overline{A_{2,2}}, \\ H_1 = -k A_{1,1} + k[(z_1 + \bar{z}_1) \overline{c_1} A_{1,1} + \overline{c_2} A_{1,2} + z_2 \overline{c_1} A_{1,2} + \bar{z}_2 \overline{c_2} A_{1,1}] j(\bar{\gamma}, \bar{Z})^{-1}, \\ H_2 = -k A_{2,1} + k[(z_1 + \bar{z}_1) \overline{c_1} A_{2,1} + \overline{c_2} A_{2,2} + z_2 \overline{c_1} A_{2,2} + \bar{z}_2 \overline{c_2} A_{2,1}] j(\bar{\gamma}, \bar{Z})^{-1}, \\ F_{2,1} = \overline{F_{1,2}}, \quad G_1 = -\overline{H_1}, \quad G_2 = -\overline{H_2}. \end{cases}$$

Now, we will prove that $F_{1,1} = (u + \bar{u}) |j(\gamma, Z)|^{-2}$. Note that

$$\begin{aligned} &F_{1,1} |j(\gamma, Z)|^4 \\ &= (z_1 + \bar{z}_1) [(a_1 c_2 - a_2 c_1) z_2 + (a_1 c_3 - a_3 c_1)] [(\overline{a_1 c_2} - \overline{a_2 c_1}) \bar{z}_2 + (\overline{a_1 c_3} - \overline{a_3 c_1})] \\ &\quad + [(a_2 c_1 - a_1 c_2) z_1 + (a_2 c_3 - a_3 c_2)] [(\overline{a_2 c_1} - \overline{a_1 c_2}) \bar{z}_1 + (\overline{a_2 c_3} - \overline{a_3 c_2})] \\ &\quad + z_2 [(a_2 c_1 - a_1 c_2) z_1 + (a_2 c_3 - a_3 c_2)] [(\overline{a_1 c_2} - \overline{a_2 c_1}) \bar{z}_2 + (\overline{a_1 c_3} - \overline{a_3 c_1})] \\ &\quad + \bar{z}_2 [(a_1 c_2 - a_2 c_1) z_2 + (a_1 c_3 - a_3 c_1)] [(\overline{a_2 c_1} - \overline{a_1 c_2}) \bar{z}_1 + (\overline{a_2 c_3} - \overline{a_3 c_2})]. \end{aligned}$$

On the other hand,

$$u + \bar{u} = [(a_1 z_1 + a_2 z_2 + a_3)(\overline{c_1 z_1} + \overline{c_2 z_2} + \overline{c_3}) + (\overline{a_1 z_1} + \overline{a_2 z_2} + \overline{a_3})(c_1 z_1 + c_2 z_2 + c_3)] |j(\gamma, Z)|^{-2}.$$

Now, we need the following six identities:

$$\begin{aligned}
& (a_2c_1 - a_1c_2)(\overline{a_2c_1} - \overline{a_1c_2}) \\
&= (\overline{a_1}a_3 + a_1\overline{a_3})c_1\overline{c_1} - a_1\overline{a_2c_1}c_2 - \overline{a_1}a_2c_1\overline{c_2} + a_1\overline{a_1}(c_1\overline{c_3} + c_3\overline{c_1}) \\
&= \overline{a_1}c_1(a_3\overline{c_1} - a_2\overline{c_2} + a_1\overline{c_3}) + a_1\overline{c_1}(\overline{a_3}c_1 - \overline{a_2}c_2 + \overline{a_1}c_3) \\
&= \overline{a_1}c_1 + a_1\overline{c_1}. \\
\\
& (a_1c_2 - a_2c_1)(\overline{a_1c_3} - \overline{a_3c_1}) \\
&= \overline{a_1}c_2(1 - a_3\overline{c_1} + a_2\overline{c_2}) + a_2\overline{c_1}(1 + \overline{a_2}c_2 - \overline{a_1}c_3) - \overline{a_1}a_2c_1\overline{c_3} - a_1\overline{a_3}c_1c_2 \\
&= \overline{a_1}c_2 + a_2\overline{c_1} + \overline{c_1}c_2(-\overline{a_1}a_3 + a_2\overline{a_2} - a_1\overline{a_3}) + \overline{a_1}a_2(c_2\overline{c_2} - \overline{c_1}c_3 - c_1\overline{c_3}) \\
&= \overline{a_1}c_2 + a_2\overline{c_1}. \\
\\
& (a_2c_3 - a_3c_2)(\overline{a_1c_2} - \overline{a_2c_1}) + (a_1c_2 - a_2c_1)(\overline{a_2c_3} - \overline{a_3c_2}) \\
&= a_2\overline{c_2}(\overline{a_1}c_3 + \overline{a_3}c_1) + \overline{a_2}c_2(a_3\overline{c_1} + a_1\overline{c_3}) - \overline{a_1}a_3c_2\overline{c_2} - a_2\overline{a_2}c_1\overline{c_3} - a_1\overline{a_3}c_2\overline{c_2} \\
&= a_2\overline{c_2} + \overline{a_2}c_2 + c_2\overline{c_2}(a_2\overline{a_2} - \overline{a_1}a_3 - a_1\overline{a_3}) + a_2\overline{a_2}(c_2\overline{c_2} - \overline{c_1}c_3 - c_1\overline{c_3}) \\
&= a_2\overline{c_2} + \overline{a_2}c_2. \\
\\
& (a_1c_3 - a_3c_1)(\overline{a_1c_3} - \overline{a_3c_1}) + (a_2c_1 - a_1c_2)(\overline{a_2c_3} - \overline{a_3c_2}) \\
&= a_1\overline{c_3}(\overline{a_1}c_3 - \overline{a_2}c_2) + \overline{a_3}c_1(a_3\overline{c_1} - a_2\overline{c_2}) - \overline{a_1}a_3c_1\overline{c_3} - a_1\overline{a_3}c_1c_3 + a_2\overline{a_2}c_1\overline{c_3} + a_1\overline{a_3}c_2\overline{c_2} \\
&= a_1\overline{c_3} + \overline{a_3}c_1 - a_1\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) - c_1\overline{c_3}(a_1\overline{a_3} + \overline{a_1}a_3 - a_2\overline{a_2}) \\
&= a_1\overline{c_3} + \overline{a_3}c_1. \\
\\
& (a_2c_3 - a_3c_2)(\overline{a_1c_3} - \overline{a_3c_1}) \\
&= a_2\overline{c_3}(1 - \overline{a_3}c_1 + \overline{a_2}c_2) + \overline{a_3}c_2(1 + a_2\overline{c_2} - a_1\overline{c_3}) - \overline{a_1}a_3c_2\overline{c_3} - a_2\overline{a_3}c_1c_3 \\
&= a_2\overline{c_3} + \overline{a_3}c_2 - a_2\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) + c_2\overline{c_3}(a_2\overline{a_2} - a_1\overline{a_3} - \overline{a_1}a_3) \\
&= a_2\overline{c_3} + \overline{a_3}c_2. \\
\\
& (a_2c_3 - a_3c_2)(\overline{a_2c_3} - \overline{a_3c_2}) \\
&= a_3\overline{c_3}(1 - \overline{a_3}c_1 - \overline{a_1}c_3) + \overline{a_3}c_3(1 - a_3\overline{c_1} - a_1\overline{c_3}) + a_2\overline{a_2}c_3\overline{c_3} + a_3\overline{a_3}c_2\overline{c_2} \\
&= a_3\overline{c_3} + \overline{a_3}c_3 - a_3\overline{a_3}(c_1\overline{c_3} + \overline{c_1}c_3 - c_2\overline{c_2}) - c_3\overline{c_3}(\overline{a_1}a_3 + a_1\overline{a_3} - a_2\overline{a_2}) \\
&= a_3\overline{c_3} + \overline{a_3}c_3.
\end{aligned}$$

By these identities, we get the desired formula.

Similarly, we have

$$F_{1,2} = \overline{v}|j(\gamma, Z)|^{-2}, \quad F_{2,1} = v|j(\gamma, Z)|^{-2}, \quad F_{2,2} = |j(\gamma, Z)|^{-2},$$

and

$$H_1 = -k|j(\gamma, Z)|^{-2}, \quad H_2 = 0, \quad G_1 = k|j(\gamma, Z)|^{-2}, \quad G_2 = 0.$$

Therefore,

$$\begin{aligned}
& L_k(f(\gamma(Z))j(\gamma, Z)^{-k}j(\bar{\gamma}, \bar{Z})^k) \\
&= j(\gamma, Z)^{-k}j(\bar{\gamma}, \bar{Z})^k(z_1 + \bar{z}_1 - z_2\bar{z}_2)|j(\gamma, Z)|^{-2} \\
&\quad \times \left[(u + \bar{u})\frac{\partial^2}{\partial u \partial \bar{u}} + \frac{\partial^2}{\partial v \partial \bar{v}} + v\frac{\partial^2}{\partial \bar{u} \partial v} + \bar{v}\frac{\partial^2}{\partial u \partial \bar{v}} - k\left(\frac{\partial}{\partial u} - \frac{\partial}{\partial \bar{u}}\right)\right] f(\gamma(Z)) \\
&= j(\gamma, Z)^{-k}j(\bar{\gamma}, \bar{Z})^k(L_k f)(\gamma(Z)).
\end{aligned}$$

□

For the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$, the Eisenstein series of weight k is defined by

$$(4.7) \quad E(Z; k, s) = \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} j(\bar{\gamma}, \bar{Z})^k j(\gamma, Z)^{-k} \rho(\gamma, Z)^s, \quad \text{for } \operatorname{Re}(s) > 2, k \in \mathbb{Z}.$$

THEOREM 4.3. *The Eisenstein series satisfies the following equation:*

$$(4.8) \quad L_k E(Z; k, s) = s(s-2)E(Z; k, s).$$

Proof. First, we note that

$$L_k \rho^s = L\rho^s + \rho \cdot k \left(\frac{\partial}{\partial z_1} - \frac{\partial}{\partial \bar{z}_1} \right) \rho^s = s(s-2)\rho^s.$$

By Proposition 4.2, we have

$$\begin{aligned}
L_k E(Z; k, s) &= \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} L_k [\rho(\gamma(Z))^s j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k] \\
&= \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k (L_k \rho^s)(\gamma(Z)) \\
&= s(s-2) \sum_{\gamma \in \Gamma \cap N \setminus \Gamma} j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k \rho^s(\gamma(Z)) \\
&= s(s-2)E(Z; k, s).
\end{aligned}$$

□

Set

$$(4.9) \quad J(\gamma, Z) := \frac{j(\gamma, Z)}{|j(\gamma, Z)|}.$$

Then $J(\gamma, Z) = e^{i \arg j(\gamma, Z)}$.

A generalized automorphic form on the Picard modular group $\Gamma = U(2, 1; \mathcal{O})$ of weight $3k$ is given by

$$(4.10) \quad f(\gamma(Z)) = \frac{j(\gamma, Z)^{3k}}{|j(\gamma, Z)|^{3k}} f(Z) \quad \text{for } \gamma \in \Gamma.$$

Put

$$(4.11) \quad H_k(Z, W) = \frac{\rho(Z, W)^{2k}}{|\rho(Z, W)|^{2k}} \quad \text{for } Z, W \in \mathfrak{S}_2.$$

Definition 4.4. For any function $\Phi \in C^\infty(\mathbb{R})$, the associated real point-pair invariant is given by

$$(4.12) \quad k(Z, W) = \Phi \left[\frac{|\rho(Z, W)|^2}{\rho(Z)\rho(W)} \right],$$

where $Z, W \in \mathfrak{S}_2$.

PROPOSITION 4.5. *The following formulas hold:*

- (1) $k(Z, W) = k(g(Z), g(W))$, for any $g \in G$.
- (2) $k(Z, W) = k(W, Z)$.

Definition 4.6. Let $\Phi \in C^\infty(\mathbb{R})$ be a real-valued function. The complex point-pair invariant is given by

$$(4.13) \quad K_k(Z, W) = H_k(Z, W) \Phi \left[\frac{|\rho(Z, W)|^2}{\rho(Z)\rho(W)} \right],$$

where $Z, W \in \mathfrak{S}_2$.

PROPOSITION 4.7. *The following formulas hold:*

- (1) $H_k(g(Z), g(W)) = J(g, Z)^{2k} H_k(Z, W) J(g, W)^{-2k}$, for $g \in G$.
- (2) $K_k(g(W), g(W)) = J(g, Z)^{2k} K_k(Z, W) J(g, W)^{-2k}$, for $g \in G$.
- (3) $|H_k(Z, W)| = 1$, and $H_k(Z, W) = \overline{H_k(W, Z)}$.
- (4) $K_k(Z, W) = \overline{K_k(W, Z)}$.

Proof. Since $\rho(g(Z), g(W)) \overline{j(g, Z)} j(g, W) = \rho(Z, W)$ for $g \in G$. We have

$$H_k(g(Z), g(W)) = \frac{|j(g, Z)|^{2k}}{|j(g, Z)|^{2k}} H_k(Z, W) \frac{|j(g, W)|^{2k}}{|j(g, W)|^{2k}} = J(g, Z)^{2k} H_k(Z, W) J(g, W)^{-2k}.$$

The others are trivial. □

LEMMA 4.8. *The following formula holds:*

$$(4.14) \quad L_k K_k(Z, W) = H_k(Z, W) \left[(\sigma^2 - \sigma)\Phi''(\sigma) + (3\sigma - 1)\Phi'(\sigma) + \frac{k^2}{\sigma}\Phi(\sigma) \right],$$

where $\sigma(Z, W) = \frac{|\rho(Z, W)|^2}{\rho(Z)\rho(W)}$.

Proof. It is obtained by a straightforward calculation. □

By $L_k K_k(Z, W) = s(s-2)K_k(Z, W)$, one has

$$(4.15) \quad \Phi''(\sigma) + \left(\frac{1}{\sigma} + \frac{2}{\sigma-1} \right) \Phi'(\sigma) + \frac{1}{\sigma(\sigma-1)} \left[\frac{k^2}{\sigma} - s(s-2) \right] \Phi(\sigma) = 0.$$

It is known that the Fuchsian equation with three regular singularities a, b, c is given by

$$\begin{aligned} & \frac{d^2w}{dz^2} + \left[\frac{1-\alpha_1-\alpha_2}{z-a} + \frac{1-\beta_1-\beta_2}{z-b} + \frac{1-\gamma_1-\gamma_2}{z-c} \right] \frac{dw}{dz} + \frac{1}{(z-a)(z-b)(z-c)} \\ & \times \left[\frac{\alpha_1\alpha_2(a-b)(a-c)}{z-a} + \frac{\beta_1\beta_2(b-c)(b-a)}{z-b} + \frac{\gamma_1\gamma_2(c-a)(c-b)}{z-c} \right] w = 0, \end{aligned}$$

where $(\alpha_1, \alpha_2), (\beta_1, \beta_2), (\gamma_1, \gamma_2)$ are exponents belonging to a, b, c , respectively, and, they satisfy that $\alpha_1 + \alpha_2 + \beta_1 + \beta_2 + \gamma_1 + \gamma_2 = 1$. The solution of this equation is given in Riemann P-notation by

$$w(z) = P \left\{ \begin{matrix} a & b & c \\ \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \end{matrix}; z \right\}.$$

When $c = \infty$, it reduces to

$$\begin{aligned} & \frac{d^2w}{dz^2} + \left[\frac{1-\alpha_1-\alpha_2}{z-a} + \frac{1-\beta_1-\beta_2}{z-b} \right] \frac{dw}{dz} + \frac{1}{(z-a)(z-b)} \\ & \times \left[\frac{\alpha_1\alpha_2(a-b)}{z-a} + \frac{\beta_1\beta_2(b-a)}{z-b} + \gamma_1\gamma_2 \right] w = 0. \end{aligned}$$

Now, in our case,

$$a = 0, b = 1, c = \infty, \alpha_1 = -|k|, \alpha_2 = |k|, \beta_1 = 0, \beta_2 = -1, \gamma_1 = s, \gamma_2 = 2-s,$$

and

$$\begin{aligned}
\Phi(\sigma) &= \Phi_{|k|}(\sigma) \\
&= P \left\{ \begin{array}{cccc} 0 & 1 & \infty \\ -|k| & 0 & s \\ |k| & -1 & 2-s \end{array}; \sigma \right\} \\
&= \sigma^{-|k|} P \left\{ \begin{array}{cccc} 0 & 1 & \infty \\ 0 & 0 & s-|k| \\ 2|k| & -1 & 2-s-|k| \end{array}; \sigma \right\} \\
&= \sigma^{-|k|} F(s-|k|, 2-s-|k|; 1-2|k|; \sigma) \\
&= \sigma^{-|k|} (1-\sigma)^{|k|-s} F(s-|k|, s-1-|k|; 1-2|k|; \frac{\sigma}{\sigma-1}) \\
&= \sigma^{-|k|} (1-\sigma)^{|k|-s} F(s-|k|, s-1-|k|; 2s-1; -\frac{1}{\sigma-1}).
\end{aligned}$$

Now, we set

$$\begin{aligned}
(4.16) \quad K(Z, W; k, s) &= \frac{\pi^{\frac{3}{2}} \Gamma(\frac{3}{2} - s)}{(|k| + 1 - s) \Gamma(2 - s)} \left(\frac{1}{4} \right)^{s-1} \rho(Z, W)^k \rho(\overline{Z}, \overline{W})^{-k} \\
&\quad \times \sigma^{-|k|} (\sigma - 1)^{|k|-s} F(s-|k|, s-1-|k|; 2s-1; -\frac{1}{\sigma-1}).
\end{aligned}$$

It has the following property:

$$(4.17) \quad K(\gamma(Z), \gamma(Z'); k, s) = j(\gamma, Z)^k j(\overline{\gamma}, \overline{Z})^{-k} j(\gamma, Z')^{-k} j(\overline{\gamma}, \overline{Z'})^k K(Z, Z'; k, s).$$

Moreover, note that $\rho(Z', Z) = \rho(\overline{Z}, \overline{Z'})$, we have

$$K(Z', Z; k, s) = K(Z, Z'; -k, s).$$

The automorphic Green function of weight k is given by

$$(4.18) \quad G(Z, Z'; k, s) = \sum_{\gamma \in \Gamma} K(Z, \gamma(Z'); k, s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

By the same method, we can prove that the right hand side of (4.18) is convergent if $\operatorname{Re}(s) > \delta(\Gamma)$.

The automorphic Green function of weight k has the following properties:

$$G(Z, Z'; k, s) = G(Z', Z; -k, s), \quad L_k(Z) G(Z, Z'; k, s) = s(s-2) G(Z, Z'; k, s),$$

Furthermore,

$$\begin{aligned}
L_{-k}(Z') G(Z, Z'; k, s) &= L_{-k}(Z') G(Z', Z; -k, s) \\
&= s(s-2) G(Z', Z; -k, s) = s(s-2) G(Z, Z'; k, s).
\end{aligned}$$

The Poisson kernel of weight k is given by

$$(4.19) \quad P_k(Z, W; s) = \rho(Z)^s \rho(Z, W)^{k-s} \rho(\overline{Z}, \overline{W})^{-k-s},$$

where $Z \in \mathfrak{S}_2$, $W \in \partial\mathfrak{S}_2$. By

$$\rho(\gamma(Z), \gamma(W)) \overline{j(\gamma, Z)} j(\gamma, W) = \rho(Z, W), \quad j(\gamma, \gamma^{-1}(W)) = j(\gamma^{-1}, W)^{-1},$$

we have

$$P_k(\gamma(Z), W; s) = P_k(Z, \gamma^{-1}(W); s) j(\gamma, Z)^k j(\overline{\gamma}, \overline{Z})^{-k} j(\gamma^{-1}, W)^{k-s} j(\overline{\gamma}^{-1}, \overline{W})^{-k-s}.$$

The Eisenstein series with respect to $\partial\mathfrak{S}_2$ of weight k is defined by

$$(4.20) \quad E(Z, W; k, s) = \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\overline{\gamma}, \overline{Z})^k P_k(\gamma(Z), W; s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

Here Γ is a discrete subgroup of $U(2, 1)$. Now, we have the following formula:

$$\lim_{\rho(Z) \rightarrow 0} \rho(Z)^{-s} K_k(Z, Z') = \frac{\pi^{\frac{3}{2}} \Gamma(\frac{3}{2} - s)}{(|k| + 1 - s) \Gamma(2 - s)} \left(\frac{1}{4}\right)^{s-1} P_k(Z', W; s),$$

where $Z \rightarrow W \in \partial\mathfrak{S}_2$ as $\rho(Z) \rightarrow 0$. Consequently,

$$\lim_{\rho(Z) \rightarrow 0} \rho(Z)^{-s} G(Z, Z'; k, s) = \frac{\pi^{\frac{3}{2}} \Gamma(\frac{3}{2} - s)}{(|k| + 1 - s) \Gamma(2 - s)} \left(\frac{1}{4}\right)^{s-1} E(Z', W; k, s).$$

LEMMA 4.9. *The Eisenstein series of weight k satisfies the following three equations:*

- (1) $L_k E(Z, W; k, s) = s(s-2) E(Z, W; k, s)$.
- (2) $E(\gamma(Z), W; k, s) = E(Z, W; k, s)$, for $\gamma \in \Gamma$.
- (3) $E(Z, g(W); k, s) = j(g, W)^{s-k} j(\overline{g}, \overline{W})^{s+k} E(Z, W; k, s)$, where $g \in \Gamma$.

Proof. In fact,

$$L_k P_k(Z, W; s) = s(s-2) P_k(Z, W; s) + (k^2 - s^2) \rho(W) P_k(Z, W; s+1).$$

Since $\rho(W) = 0$, $L_k P_k(Z, W; s) = s(s-2) P_k(Z, W; s)$. By Proposition 4.2,

$$\begin{aligned} L_k E(Z, W; k, s) &= \sum_{\gamma \in \Gamma} L_k [P_k(\gamma(Z), W; s) j(\gamma, Z)^{-k} j(\overline{\gamma}, \overline{Z})^k] \\ &= \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\overline{\gamma}, \overline{Z})^k (L_k P_k)(\gamma(Z), W; s) \\ &= s(s-2) \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\overline{\gamma}, \overline{Z})^k P_k(\gamma(Z), W; s) \\ &= s(s-2) E(Z, W; k, s). \end{aligned}$$

To prove the third identity, we note that

$$P_k(\gamma(Z), \gamma(W); s) = j(\gamma, Z)^k j(\bar{\gamma}, \bar{Z})^{-k} j(\gamma, W)^{s-k} j(\bar{\gamma}, \bar{W})^{s+k} P_k(Z, W; s).$$

By Proposition 2.9, we have

$$\begin{aligned} E(Z, g(W); k, s) &= \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k P_k(\gamma(Z), g(W); s) \\ &= \sum_{\gamma \in \Gamma} j(g, \gamma(Z))^{-k} j(\bar{g}, \bar{\gamma}(\bar{Z}))^k j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k P_k(g(\gamma(Z)), g(W); s) \\ &= \sum_{\gamma \in \Gamma} j(\gamma, Z)^{-k} j(\bar{\gamma}, \bar{Z})^k P_k(\gamma(Z), W; s) j(g, W)^{s-k} j(\bar{g}, \bar{W})^{s+k} \\ &= j(g, W)^{s-k} j(\bar{g}, \bar{W})^{s+k} E(Z, W; k, s). \end{aligned}$$

□

THEOREM 4.10. Set

$$(4.21) \quad \phi(Z, Z'; k, s) = \int_{\partial \mathfrak{S}_2} P_k(Z, W; s) P_{-k}(Z', W; 2-s) dm(W).$$

Then the following formula holds:

$$\begin{aligned} (4.22) \quad \phi(Z, Z'; k, s) &= K(Z, Z'; k, s) + K(Z, Z'; k, 2-s) \\ &= \phi(Z', Z; -k, 2-s) = K(Z', Z; -k, s) + K(Z', Z; -k, 2-s), \end{aligned}$$

where $k = 0$ or $k = \pm 1$.

Proof. In fact, if $k = 0$, this is Theorem 3.7. Now, we only consider the nontrivial case $k = \pm 1$.

For $\gamma \in G$,

$$\begin{aligned} \phi(\gamma(Z), \gamma(Z'); k, s) &= \int_{\partial \mathfrak{S}_2} P_k(\gamma(Z), W; s) P_{-k}(\gamma(Z'), W; 2-s) dm(W) \\ &= j(\gamma, Z)^k j(\bar{\gamma}, \bar{Z})^{-k} j(\gamma, Z')^{-k} j(\bar{\gamma}, \bar{Z}')^k \int_{\partial \mathfrak{S}_2} P_k(Z, \gamma^{-1}(W); s) \\ &\quad \times P_{-k}(Z', \gamma^{-1}(W); 2-s) j(\gamma^{-1}, W)^{-2} j(\bar{\gamma}^{-1}, \bar{W})^{-2} dm(W). \end{aligned}$$

Since $dm(\gamma^{-1}(W)) = |j(\gamma^{-1}, W)|^{-4} dm(W)$, we have

$$(4.23) \quad \phi(\gamma(Z), \gamma(Z'); k, s) = j(\gamma, Z)^k j(\bar{\gamma}, \bar{Z})^{-k} j(\gamma, Z')^{-k} j(\bar{\gamma}, \bar{Z}')^k \phi(Z, Z'; k, s).$$

Thus, $\phi(Z, Z'; k, s)$ is a point-pair invariant covariant with respect to the weight k .

By the double transitivity of $U(2, 1)$ on \mathfrak{S}_2 , there exists a $\gamma \in U(2, 1)$, such that

$$Z = \gamma\left(\frac{1}{2}\rho, 0\right), \quad Z' = \gamma\left(\frac{1}{2}\rho', 0\right).$$

Let ρ and ρ' stand for $(\rho, 0)$ and $(\rho', 0)$, respectively. Set $\gamma = \begin{pmatrix} * & * & * \\ * & * & * \\ a_1 & a_2 & a_3 \end{pmatrix}$, then

$$|j(\gamma, \frac{1}{2}\rho)|^2 = \left|\frac{1}{2}\rho a_1 + a_3\right|^2 = \frac{1}{4}\rho^2|a_1|^2 + \frac{1}{2}\rho|a_2|^2 + |a_3|^2.$$

Denote

$$(4.24) \quad \mu = \frac{\frac{1}{2}(\rho + \rho')}{\frac{1}{4}\rho^2|a_1|^2 + \frac{1}{2}\rho|a_2|^2 + |a_3|^2},$$

then $\mu > 0$.

Since

$$\begin{aligned} \phi(Z, Z'; k, s) &= \phi\left(\gamma\left(\frac{1}{2}\rho\right), \gamma\left(\frac{1}{2}\rho'\right); k, s\right) \\ &= j(\gamma, \frac{1}{2}\rho)^k j(\bar{\gamma}, \frac{1}{2}\rho)^{-k} j(\gamma, \frac{1}{2}\rho')^{-k} j(\bar{\gamma}, \frac{1}{2}\rho')^k \phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right). \end{aligned}$$

By (4.24), we have

$$\rho(Z, Z') = \rho\left(\gamma\left(\frac{1}{2}\rho\right), \gamma\left(\frac{1}{2}\rho'\right)\right) = \frac{\frac{1}{2}(\rho + \rho')}{j(\bar{\gamma}, \frac{1}{2}\rho)j(\gamma, \frac{1}{2}\rho')} = \mu \frac{j(\gamma, \frac{1}{2}\rho)}{j(\gamma, \frac{1}{2}\rho')}.$$

Hence,

$$(4.25) \quad \begin{aligned} \phi(Z, Z'; k, s) &= [\mu^{-1}\rho(Z, Z')]^k [\mu^{-1}\rho(\bar{Z}, \bar{Z}')^{-k}] \phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) \\ &= \rho(Z, Z')^k \rho(\bar{Z}, \bar{Z}')^{-k} \phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right). \end{aligned}$$

For $W \in \partial\mathfrak{S}_2$, i.e., $w_1 = \frac{|w|^2}{2} + iv$, $w_2 = w$ with $w \in \mathbb{C}$, $v \in \mathbb{R}$.

$$\rho(Z, W) = \frac{\rho + |w|^2}{2} + iv, \quad \rho(Z', W) = \frac{\rho' + |w|^2}{2} + iv.$$

Therefore,

$$\begin{aligned} \phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 16\rho^s\rho'^{2-s} \int_{\mathbb{R}} \int_{\mathbb{C}} \frac{1}{|\rho + |w|^2 + 2iv|^{2s}} \frac{1}{|\rho' + |w|^2 + 2iv|^{4-2s}} \\ &\quad \times \left(\frac{\rho + |w|^2 + 2iv}{\rho + |w|^2 - 2iv}\right)^k \left(\frac{\rho' + |w|^2 + 2iv}{\rho' + |w|^2 - 2iv}\right)^{-k} dw d\bar{w} dv. \end{aligned}$$

Set $w = re^{i\theta}$ and $u = r^2$, then

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 16\pi\rho^s\rho'^{2-s} \int_{-\infty}^{\infty} \int_0^{\infty} \frac{1}{|\rho + u + 2iv|^{2s}} \frac{1}{|\rho' + u + 2iv|^{4-2s}} \\ &\quad \times \left(\frac{\rho + u + 2iv}{\rho + u - 2iv}\right)^k \left(\frac{\rho' + u + 2iv}{\rho' + u - 2iv}\right)^{-k} dudv \\ &= 8\pi\rho^s\rho'^{2-s} \int_{-\infty}^{\infty} \int_0^{\infty} \frac{1}{|\rho + u + iv|^{2s}} \frac{1}{|\rho' + u + iv|^{2(2-s)}} \\ &\quad \times \exp(2ki(\arctan \frac{v}{\rho + u} - \arctan \frac{v}{\rho' + u})) dudv.\end{aligned}$$

By

$$\arctan \frac{v}{\rho + u} - \arctan \frac{v}{\rho' + u} = \arctan \frac{(\rho' - \rho)v}{(\rho + u)(\rho' + u) + v^2},$$

we have

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 8\pi\rho^s\rho'^{2-s} \int_{-\infty}^{\infty} \int_0^{\infty} \frac{1}{[(\rho + u)^2 + v^2]^s} \frac{1}{[(\rho' + u)^2 + v^2]^{2-s}} \\ &\quad \times \exp(2ki \arctan \frac{(\rho' - \rho)v}{(\rho + u)(\rho' + u) + v^2}) dudv \\ &= 16\pi\rho^s\rho'^{2-s} \int_0^{\infty} \int_0^{\infty} \frac{1}{[(\rho + u)^2 + v^2]^s} \frac{1}{[(\rho' + u)^2 + v^2]^{2-s}} \\ &\quad \times \cos 2k \arctan \frac{(\rho' - \rho)v}{(\rho + u)(\rho' + u) + v^2} dudv.\end{aligned}$$

Assume that $\rho < \rho'$, set $\lambda = \frac{\rho}{\rho'}$, $v = (\rho' - \rho)w$ and $l = \frac{\rho+u}{\rho'+u}$, then $u = \frac{l\rho'-\rho}{1-l}$, $du = \frac{\rho'-\rho}{(1-l)^2}dl$. We have

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 16\pi\lambda^s(1-\lambda)^{-2} \int_0^{\infty} \int_{\lambda}^1 \left[w^2 + \frac{l^2}{(1-l)^2}\right]^{-s} \left[w^2 + \frac{1}{(1-l)^2}\right]^{s-2} \\ &\quad \times \cos 2k \arctan \left(\frac{w}{w^2 + \frac{l}{(1-l)^2}}\right) (1-l)^{-2} dl dw.\end{aligned}$$

Set $w = \frac{\sqrt{v}}{1-l}$, then

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 8\pi\lambda^s(1-\lambda)^{-2} \int_{\lambda}^1 (1-l)dl \int_0^{\infty} v^{-\frac{1}{2}} (1+v)^{s-2} (l^2 + v)^{-s} \\ &\quad \times \cos 2k \arctan \left[\frac{(1-l)\sqrt{v}}{v+l}\right] dv.\end{aligned}$$

Let $x = \arctan \frac{(1-l)\sqrt{v}}{v+l}$, then

$$\cos 2x = \frac{1 - \tan^2 x}{1 + \tan^2 x} = \frac{(v+l)^2 - (1-l)^2 v}{(v+l)^2 + (1-l)^2 v}.$$

We have

$$\cos 2kx = \cos \left(k \arccos \frac{(v+l)^2 - (1-l)^2 v}{(v+l)^2 + (1-l)^2 v} \right).$$

It is known that the Tchebichef polynomial

$$T_n(x) = \cos(n \arccos x) = F(-n, n; \frac{1}{2}; \frac{1-x}{2}).$$

Hence

$$\cos 2kx = F(-k, k; \frac{1}{2}; \frac{(1-l)^2 v}{v^2 + (1+l^2)v + l^2}).$$

Therefore,

$$\begin{aligned} \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) &= 8\pi\lambda^s(1-\lambda)^{-2} \int_{\lambda}^1 (1-l)dl \int_0^{\infty} v^{-\frac{1}{2}}(1+v)^{s-2}(l^2+v)^{-s} \\ &\quad \times F(-k, k; \frac{1}{2}; \frac{(1-l)^2 v}{(1+v)(l^2+v)})dv. \end{aligned}$$

We know that

$$F(-k, k; \frac{1}{2}; x) = \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n! (\frac{1}{2})_n} x^n.$$

Thus

$$\begin{aligned} \phi(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s) &= 8\pi\lambda^s(1-\lambda)^{-2} \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n! (\frac{1}{2})_n} \int_{\lambda}^1 (1-l)^{2n+1} l^{-2(s+n)} dl \\ &\quad \times \int_0^{\infty} v^{n-\frac{1}{2}}(1+v)^{s-n-2}(1+vl^{-2})^{-s-n} dv. \end{aligned}$$

By the following formula

$$F(a, b; c; 1-z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^{\infty} s^{b-1} (1+s)^{a-c} (1+sz)^{-a} ds$$

for $\operatorname{Re}(c) > \operatorname{Re}(b) > 0$, $|\arg(z)| < \pi$, we have

$$\begin{aligned} &\int_0^{\infty} v^{n-\frac{1}{2}}(1+v)^{s-n-2}(1+vl^{-2})^{-s-n} dv \\ &= \frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)} F(s+n, \frac{1}{2}+n; 2n+2; 1-l^{-2}). \end{aligned}$$

Hence,

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 8\pi\lambda^s(1-\lambda)^{-2} \sum_{n=0}^{|k|} \frac{(-k)_n(k)_n}{n!(\frac{1}{2})_n} \frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)} \\ &\quad \times \int_{\lambda}^1 (1-l)^{2n+1} l^{-2(s+n)} F(s+n, \frac{1}{2}+n; 2n+2; 1-l^{-2}) dl.\end{aligned}$$

Note that

$$\frac{1}{n!(\frac{1}{2})_n} \frac{\Gamma(n+\frac{1}{2})\Gamma(n+\frac{3}{2})}{\Gamma(2n+2)} = \frac{\pi}{2^{2n+1}(n!)^2}.$$

We have

$$\begin{aligned}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) &= 4\pi^2\lambda^s(1-\lambda)^{-2} \sum_{n=0}^{|k|} \frac{(-k)_n(k)_n}{2^{2n}(n!)^2} \int_{\lambda}^1 (1-l)^{2n+1} l^{-2(s+n)} \\ &\quad \times F(s+n, \frac{1}{2}+n; 2n+2; 1-l^{-2}) dl.\end{aligned}$$

$$(1) \ z = 1 - l^{-2}, \ \frac{z}{z-1} = 1 - l^2.$$

By $F(a, b; c; z) = (1-z)^{-a}F(a, c-b; c; \frac{z}{z-1})$, we have

$$F(s+n, \frac{1}{2}+n; 2n+2; 1-l^{-2}) = l^{2(s+n)} F(s+n, n+\frac{3}{2}; 2n+2; 1-l^2).$$

$$(2) \ z = 1 - l^2, \ 1 - z = l^2.$$

By

$$\begin{aligned}F(a, b; c; z) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F(a, b; a+b-c+1; 1-z) \\ &\quad + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} (1-z)^{c-a-b} F(c-a, c-b; c-a-b+1; 1-z),\end{aligned}$$

where $|\arg(1-z)| < \pi$, we have

$$\begin{aligned}&F(s+n, n+\frac{3}{2}; 2n+2; 1-l^2) \\ &= \frac{\Gamma(2n+2)\Gamma(\frac{1}{2}-s)}{\Gamma(n-s+2)\Gamma(n+\frac{1}{2})} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) \\ &\quad + \frac{\Gamma(2n+2)\Gamma(s-\frac{1}{2})}{\Gamma(s+n)\Gamma(n+\frac{3}{2})} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2).\end{aligned}$$

Therefore,

$$\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; k, s\right) = 4\pi^2\lambda^s(1-\lambda)^{-2} \sum_{n=0}^{|k|} \frac{(-k)_n(k)_n}{2^{2n}(n!)^2} (F_1 + F_2),$$

where

$$\begin{aligned} F_1 &= \frac{(2n+1)!\Gamma(\frac{1}{2}-s)}{\Gamma(n-s+2)\Gamma(n+\frac{1}{2})} \int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl, \\ F_2 &= \frac{(2n+1)!\Gamma(s-\frac{1}{2})}{\Gamma(s+n)\Gamma(n+\frac{3}{2})} \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl. \end{aligned}$$

It is known that the incomplete beta function is defined by (see [Er], p.87)

$$B_x(p, q) = \int_0^x t^{p-1} (1-t)^{q-1} dt.$$

Set $I_x(p, q) = \frac{B_x(p, q)}{B_1(p, q)}$. Note that $B_1(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$.

Hence,

$$\begin{aligned} &\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl \\ &= \sum_{m=0}^{\infty} \frac{(s+n)_m(n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m} [B_1(2m+1, 2n+2) - B_{\lambda}(2m+1, 2n+2)]. \end{aligned}$$

Similarly,

$$\begin{aligned} &\int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl \\ &= \sum_{m=0}^{\infty} \frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} [B_1(2m-2s+2, 2n+2) - B_{\lambda}(2m-2s+2, 2n+2)]. \end{aligned}$$

We have

$$\begin{aligned} (4.26) \quad &\sum_{m=0}^{\infty} \frac{(s+n)_m(n+\frac{3}{2})_m}{m!(s+\frac{1}{2})_m} B_1(2m+1, 2n+2) = \sum_{m=0}^{\infty} \frac{(s+n)_m}{(s+\frac{1}{2})_m} \frac{(2m-1)!!(2n)!!}{(2m+2n+2)!!} \\ &= \frac{1}{2n+2} \sum_{m=0}^{\infty} \frac{(s+n)_m(\frac{1}{2})_m}{(s+\frac{1}{2})_m(n+2)_m}. \end{aligned}$$

By

$$(n+2)_m = \frac{(n+m+1)!}{(n+1)!}, \quad (\frac{1}{2})_m = \frac{(-\frac{1}{2}-n)_{m+n+1}}{(-\frac{1}{2}-n)_{n+1}},$$

$$(s+\frac{1}{2})_m = \frac{(s-n-\frac{1}{2})_{m+n+1}}{(s-n-\frac{1}{2})_{n+1}}, \quad (s+n)_m = \frac{(s-1)_{m+n+1}}{(s-1)_{n+1}},$$

(4.26) is equal to

$$\begin{aligned} & \frac{1}{2n+2} \frac{(n+1)!(s-n-\frac{1}{2})_{n+1}}{(s-1)_{n+1}(-\frac{1}{2}-n)_{n+1}} \sum_{m=0}^{\infty} \frac{(s-1)_{m+n+1}(-\frac{1}{2}-n)_{m+n+1}}{(s-n-\frac{1}{2})_{m+n+1}(m+n+1)!} \\ (4.27) \quad &= \frac{1}{2n+2} \frac{(n+1)!(s-n-\frac{1}{2})_{n+1}}{(s-1)_{n+1}(-\frac{1}{2}-n)_{n+1}} [F(s-1, -\frac{1}{2}-n; s-n-\frac{1}{2}; 1) \\ & \quad - \sum_{r=0}^n \frac{(s-1)_r (-\frac{1}{2}-n)_r}{(s-n-\frac{1}{2})_r r!}]. \end{aligned}$$

Similarly,

$$\begin{aligned} & \sum_{m=0}^{\infty} \frac{(2-s+n)_m (n+\frac{1}{2})_m}{m! (\frac{3}{2}-s)_m} B_1(2m-2s+2, 2n+2) \\ (4.28) \quad &= \frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)} \sum_{m=0}^{\infty} \frac{(1-s)_m (n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} \\ &= \frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)} F(1-s, n+\frac{1}{2}; \frac{5}{2}-s+n; 1). \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{m=0}^{\infty} \frac{(s+n)_m (n+\frac{3}{2})_m}{m! (s+\frac{1}{2})_m} B_{\lambda}(2m+1, 2n+2) \\ &= \sum_{m=0}^{\infty} \frac{(s+n)_m (n+\frac{3}{2})_m}{m! (s+\frac{1}{2})_m} B_1(2m+1, 2n+2) I_{\lambda}(2m+1, 2n+2) \\ &= \frac{1}{2n+2} \sum_{m=0}^{\infty} \frac{(s+n)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (n+2)_m} I_{\lambda}(2m+1, 2n+2). \end{aligned}$$

Now, we need the following lemma:

LEMMA 4.11. *The explicit expression of $I_x(p, q)$: For $p \in \mathbb{C}$, $\operatorname{Re}(p) > 0$ and $q \in \mathbb{Z}$ with $q > 0$, the following formula holds:*

$$(4.29) \quad I_x(p, q+1) = (p)_{q+1} \sum_{r=0}^q \frac{(-1)^r}{r!(q-r)!} \frac{x^{p+r}}{p+r}.$$

Proof.

$$\begin{aligned}
I_x(p, q+1) &= \frac{(p)_{q+1}}{q!} B(p, q+1) I_x(p, q+1) \\
&= \frac{(p)_{q+1}}{q!} B_x(p, q+1) = \frac{(p)_{q+1}}{q!} \int_0^x t^{p-1} (1-t)^q dt \\
&= \frac{(p)_{q+1}}{q!} \sum_{r=0}^q (-1)^r C_q^r \frac{x^{p+r}}{p+r} = (p)_{q+1} \sum_{r=0}^q \frac{(-1)^r}{r!(q-r)!} \frac{x^{p+r}}{p+r}.
\end{aligned}$$

□

By Lemma 4.11, we have

$$\begin{aligned}
&\frac{1}{2n+2} \sum_{m=0}^{\infty} \frac{(s+n)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (n+2)_m} I_{\lambda}(2m+1, 2n+2) \\
&= \frac{1}{2n+2} \sum_{m=0}^{\infty} \frac{(s+n)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (n+2)_m} (2m+1)_{2n+2} \sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \frac{\lambda^{2m+r+1}}{2m+r+1} \\
&= \frac{1}{2n+2} \sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \sum_{m=0}^{\infty} \frac{(s+n)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (n+2)_m} \\
&\quad \times \frac{(2m+1)(2m+2)\cdots(2m+2n+2)}{2m+r+1} \lambda^{2m+r+1}.
\end{aligned}$$

If $n = 0$, then $r = 0, 1$.

(1) $r = 0$,

$$\sum_{m=0}^{\infty} \frac{(s)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (2)_m} (2m+2) \lambda^{2m+1} = 2\lambda F(s, \frac{1}{2}; s+\frac{1}{2}; \lambda^2).$$

(2) $r = 1$,

$$\sum_{m=0}^{\infty} \frac{(s)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (2)_m} (2m+1) \lambda^{2m+2} = \frac{2s-1}{s-1} [F(s-1, \frac{1}{2}; s-\frac{1}{2}; \lambda^2) - 1].$$

On the other hand, when $n = 0$, (4.27) is equal to

$$\frac{2s-1}{2s-2} [1 - F(s-1, -\frac{1}{2}; s-\frac{1}{2}; 1)].$$

Hence,

$$\begin{aligned}
&\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl|_{n=0} \\
&= -\lambda F(s, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) + \frac{2s-1}{2s-2} F(s-1, \frac{1}{2}; s-\frac{1}{2}; \lambda^2) - \frac{2s-1}{2s-2} F(s-1, -\frac{1}{2}; s-\frac{1}{2}; 1).
\end{aligned}$$

If $n = 1$, then $r = 0, 1, 2, 3$.

(1) $r = 0$,

$$\begin{aligned} & \sum_{m=0}^{\infty} \frac{(s+1)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (3)_m} (2m+2)(2m+3)(2m+4) \lambda^{2m+1} \\ &= 8\lambda F(s+1, \frac{3}{2}; s+\frac{1}{2}; \lambda^2) + 16\lambda F(s+1, \frac{1}{2}; s+\frac{1}{2}; \lambda^2). \end{aligned}$$

(2) $r = 1$,

$$\sum_{m=0}^{\infty} \frac{(s+1)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (3)_m} (2m+1)(2m+3)(2m+4) \lambda^{2m+2} = 8\frac{s-\frac{1}{2}}{s} [F(s, \frac{3}{2}; s-\frac{1}{2}; \lambda^2) - 1].$$

(3) $r = 2$,

$$\sum_{m=0}^{\infty} \frac{(s+1)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (3)_m} (2m+1)(2m+2)(2m+4) \lambda^{2m+3} = 8\lambda^3 F(s+1, \frac{3}{2}; s+\frac{1}{2}; \lambda^2).$$

(4) $r = 3$,

$$\begin{aligned} & \sum_{m=0}^{\infty} \frac{(s+1)_m (\frac{1}{2})_m}{(s+\frac{1}{2})_m (3)_m} (2m+1)(2m+2)(2m+3) \lambda^{2m+4} \\ &= 8\frac{s-\frac{1}{2}}{s} \lambda^2 [F(s, \frac{3}{2}; s-\frac{1}{2}; \lambda^2) - 1] - 16\frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} [F(s-1, \frac{1}{2}; s-\frac{3}{2}; \lambda^2) \\ & \quad - \frac{s-1}{2s-3} \lambda^2 - 1]. \end{aligned}$$

On the other hand, when $n = 1$, (4.27) is equal to

$$\frac{2}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} F(s-1, -\frac{3}{2}; s-\frac{3}{2}; 1) - \frac{2}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} + \frac{s-\frac{1}{2}}{s}.$$

Therefore,

$$\begin{aligned} & \int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl_{n=1} \\ &= \frac{2}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} F(s-1, -\frac{3}{2}; s-\frac{3}{2}; 1) - \frac{2}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} F(s-1, \frac{1}{2}; s-\frac{3}{2}; \lambda^2) \\ & \quad - \frac{2}{3} \lambda F(s+1, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) + (\frac{1}{3}\lambda^2 + 1) \frac{s-\frac{1}{2}}{s} F(s, \frac{3}{2}; s-\frac{1}{2}; \lambda^2) \\ & \quad - (\lambda^3 + \frac{1}{3}\lambda) F(s+1, \frac{3}{2}; s+\frac{1}{2}; \lambda^2). \end{aligned}$$

Now, we need the following relation of Gauss between contiguous functions (see [Er], p.103, (36)):

$$(c-a-b)F - (c-a)F(a-1) + b(1-z)F(b+1) = 0,$$

where F denotes $F(a, b; c; z)$ and $F(a \pm 1)$, $F(b \pm 1)$, and $F(c \pm 1)$ stands for $F(a \pm 1, b; c; z)$, $F(a, b \pm 1; c; z)$, and $F(a, b; c \pm 1; z)$ respectively.

We have

$$\begin{aligned} F(s-1, \frac{1}{2}; s-\frac{3}{2}; \lambda^2) &= \frac{1}{2}F(s-2, \frac{1}{2}; s-\frac{3}{2}; \lambda^2) + \frac{1}{2}(1-\lambda^2)F(s-1, \frac{3}{2}; s-\frac{3}{2}; \lambda^2), \\ F(s+1, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) &= \frac{1}{2}F(s, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) + \frac{1}{2}(1-\lambda^2)F(s+1, \frac{3}{2}; s+\frac{1}{2}; \lambda^2). \end{aligned}$$

By [Er], p.104, (46),

$$F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad c \neq 0, -1, -2, \dots, \operatorname{Re}(c) > \operatorname{Re}(a+b),$$

we have

$$F(s-1, -\frac{3}{2}; s-\frac{3}{2}; 1) = \frac{\Gamma(s-\frac{3}{2})}{-2\sqrt{\pi}\Gamma(s)}, \quad F(s-1, -\frac{1}{2}; s-\frac{1}{2}; 1) = \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)}.$$

Therefore,

$$\begin{aligned} &\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl_{n=0} \\ &= -\lambda F(s, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) + \frac{s-\frac{1}{2}}{s-1} F(s-1, \frac{1}{2}; s-\frac{1}{2}; \lambda^2) - \frac{1}{\sqrt{\pi}} \frac{\Gamma(s+\frac{1}{2})}{(s-1)\Gamma(s)}, \end{aligned}$$

and

$$\begin{aligned} &\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl_{n=1} \\ &= -\frac{1}{3} \frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}(s-1)\Gamma(s+1)} - \frac{1}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} F(s-2, \frac{1}{2}; s-\frac{3}{2}; \lambda^2) \\ &\quad - \frac{1}{3} \frac{(s-\frac{1}{2})(s-\frac{3}{2})}{s(s-1)} (1-\lambda^2) F(s-1, \frac{3}{2}; s-\frac{3}{2}; \lambda^2) + (\frac{1}{3}\lambda^2 + 1) \frac{s-\frac{1}{2}}{s} F(s, \frac{3}{2}; s-\frac{1}{2}; \lambda^2) \\ &\quad - \frac{1}{3} \lambda F(s, \frac{1}{2}; s+\frac{1}{2}; \lambda^2) - \frac{2}{3} \lambda (\lambda^2 + 1) F(s+1, \frac{3}{2}; s+\frac{1}{2}; \lambda^2). \end{aligned}$$

By [Er], p.111, (5),

$$F(a, b; 2b; \frac{4z}{(1+z)^2}) = (1+z)^{2a} F(a, a + \frac{1}{2} - b; b + \frac{1}{2}; z^2),$$

we have

$$\begin{aligned} F(s, \frac{1}{2}; s + \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2s} F(s, s; 2s; -u^{-1}), \\ F(s-1, \frac{1}{2}; s - \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2(s-1)} F(s-1, s-1; 2s-2; -u^{-1}), \\ F(s-2, \frac{1}{2}; s - \frac{3}{2}; \lambda^2) &= (1-\lambda)^{-2(s-2)} F(s-2, s-2; 2s-4; -u^{-1}), \\ F(s-1, \frac{3}{2}; s - \frac{3}{2}; \lambda^2) &= (1-\lambda)^{-2(s-1)} F(s-1, s-2; 2s-4; -u^{-1}), \\ F(s, \frac{3}{2}; s - \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2s} F(s, s-1; 2s-2; -u^{-1}), \\ F(s, \frac{1}{2}; s + \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2s} F(s, s; 2s; -u^{-1}), \\ F(s+1, \frac{3}{2}; s + \frac{1}{2}; \lambda^2) &= (1-\lambda)^{-2(s+1)} F(s+1, s; 2s; -u^{-1}). \end{aligned}$$

where $u = \frac{(\lambda-1)^2}{4\lambda}$. Hence,

$$\begin{aligned} &\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n + \frac{3}{2}; s + \frac{1}{2}; l^2) dl_{n=0} = -\lambda (1-\lambda)^{-2s} F(s, s; 2s; -u^{-1}) \\ &+ \frac{s - \frac{1}{2}}{s-1} (1-\lambda)^{-2(s-1)} F(s-1, s-1; 2s-2; -u^{-1}) - \frac{1}{\sqrt{\pi}} \frac{1}{s-1} \frac{\Gamma(s + \frac{1}{2})}{\Gamma(s)}. \\ &\int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n + \frac{3}{2}; s + \frac{1}{2}; l^2) dl_{n=1} = -\frac{1}{3} \frac{1}{\sqrt{\pi}} \frac{1}{s-1} \frac{\Gamma(s + \frac{1}{2})}{\Gamma(s+1)} \\ &- \frac{1}{3} \frac{(s - \frac{1}{2})(s - \frac{3}{2})}{s(s-1)} (1-\lambda)^{-2(s-2)} F(s-2, s-2; 2s-4; -u^{-1}) \\ &- \frac{1}{3} \frac{(s - \frac{1}{2})(s - \frac{3}{2})}{s(s-1)} (1-\lambda^2)(1-\lambda)^{-2(s-1)} F(s-1, s-2; 2s-4; -u^{-1}) \\ &+ (\frac{1}{3}\lambda^2 + 1) \frac{s - \frac{1}{2}}{s} (1-\lambda)^{-2s} F(s, s-1; 2s-2; -u^{-1}) \\ &- \frac{1}{3} \lambda (1-\lambda)^{-2s} F(s, s; 2s; -u^{-1}) - \frac{2}{3} \lambda (\lambda^2 + 1) (1-\lambda)^{-2(s+1)} F(s+1, s; 2s; -u^{-1}). \end{aligned}$$

Similarly,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} B_{\lambda}(2m+2-2s, 2n+2) \\
&= \sum_{m=0}^{\infty} \frac{(2-s+n)_m(n+\frac{1}{2})_m}{m!(\frac{3}{2}-s)_m} B_1(2m+2-2s, 2n+2) I_{\lambda}(2m+2-2s, 2n+2) \\
&= \frac{(2n+1)!\Gamma(2-2s)}{\Gamma(4-2s+2n)} \sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} I_{\lambda}(2m+2-2s, 2n+2).
\end{aligned}$$

By Lemma 4.11, we have

$$I_{\lambda}(2m+2-2s, 2n+2) = (2m+2-2s)_{2n+2} \sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \frac{\lambda^{2m+2-2s+r}}{2m+2-2s+r}.$$

Thus,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} I_{\lambda}(2m+2-2s, 2n+2) \\
&= \sum_{r=0}^{2n+1} \frac{(-1)^r}{r!(2n+1-r)!} \sum_{m=0}^{\infty} \frac{(1-s)_m(n+\frac{1}{2})_m}{(\frac{5}{2}-s+n)_m m!} \\
&\quad \times \frac{(2m+2-2s)(2m+3-2s)\cdots(2m+2n+3-2s)}{2m+2-2s+r} \lambda^{2m+2-2s+r}.
\end{aligned}$$

If $n = 0$, then $r = 0, 1$.

(1) $r = 0$,

$$\sum_{m=0}^{\infty} \frac{(1-s)_m(\frac{1}{2})_m}{(\frac{5}{2}-s)_m m!} (2m+3-2s) \lambda^{2m+2-2s} = (3-2s) \lambda^{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2).$$

(2) $r = 1$,

$$\sum_{m=0}^{\infty} \frac{(1-s)_m(\frac{1}{2})_m}{(\frac{5}{2}-s)_m m!} (2m+2-2s) \lambda^{2m+3-2s} = 2(1-s) \lambda^{3-2s} F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2).$$

On the other hand, when $n = 0$, (4.28) is equal to

$$\frac{1}{(3-2s)(2-2s)} F(1-s, \frac{1}{2}; \frac{5}{2}-s; 1).$$

Thus,

$$\begin{aligned}
& \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=0} \\
&= \frac{1}{(3-2s)(2-2s)} F(1-s, \frac{1}{2}; \frac{5}{2}-s; 1) - \frac{\lambda^{2-2s}}{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) \\
&\quad + \frac{\lambda^{3-2s}}{3-2s} F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2).
\end{aligned}$$

If $n = 1$, then $r = 0, 1, 2, 3$.

(1) $r = 0$,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(1-s)_m (\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!} (2m+3-2s)(2m+4-2s)(2m+5-2s) \lambda^{2m+2-2s} \\
&= 8(1-s)(\frac{3}{2}-s)(\frac{5}{2}-s) \lambda^{2-2s} F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) \\
&\quad + 8(\frac{3}{2}-s)(\frac{5}{2}-s) \lambda^{2-2s} F(1-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2).
\end{aligned}$$

(2) $r = 1$,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(1-s)_m (\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!} (2m+2-2s)(2m+4-2s)(2m+5-2s) \lambda^{2m+3-2s} \\
&= 8(1-s)(2-s)(\frac{5}{2}-s) \lambda^{3-2s} F(3-s, \frac{3}{2}; \frac{5}{2}-s; \lambda^2).
\end{aligned}$$

(3) $r = 2$,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(1-s)_m (\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!} (2m+2-2s)(2m+3-2s)(2m+5-2s) \lambda^{2m+4-2s} \\
&= 8(1-s)(\frac{3}{2}-s)(\frac{5}{2}-s) \lambda^{4-2s} F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2).
\end{aligned}$$

(3) $r = 3$,

$$\begin{aligned}
& \sum_{m=0}^{\infty} \frac{(1-s)_m (\frac{3}{2})_m}{(\frac{7}{2}-s)_m m!} (2m+2-2s)(2m+3-2s)(2m+4-2s) \lambda^{2m+5-2s} \\
&= 8(1-s)(2-s)(\frac{5}{2}-s) \lambda^{5-2s} F(3-s, \frac{3}{2}; \frac{5}{2}-s; \lambda^2) \\
&\quad - 8(1-s)(2-s) \lambda^{5-2s} F(3-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2).
\end{aligned}$$

On the other hand, when $n = 1$, (4.28) is equal to

$$\frac{6\Gamma(2-2s)}{\Gamma(6-2s)}F(1-s, \frac{3}{2}; \frac{7}{2}-s; 1).$$

Thus,

$$\begin{aligned} & \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=1} \\ &= \frac{6\Gamma(2-2s)}{\Gamma(6-2s)} F(1-s, \frac{3}{2}; \frac{7}{2}-s; 1) - \frac{1}{2} \frac{1}{(2-s)} \lambda^{2-2s} (1+3\lambda^2) F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) \\ &+ \frac{1}{2} \frac{1}{(\frac{3}{2}-s)} \lambda^{3-2s} (3+\lambda^2) F(3-s, \frac{3}{2}; \frac{5}{2}-s; \lambda^2) - \frac{1}{2} \frac{1}{(1-s)(2-s)} \lambda^{2-2s} \\ &\times F(1-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) - \frac{1}{2} \frac{1}{(\frac{3}{2}-s)(\frac{5}{2}-s)} \lambda^{5-2s} F(3-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2). \end{aligned}$$

Now, we need the following relation of Gauss between contiguous functions (see [Er], p.103, (33)):

$$(c-a-b)F + a(1-z)F(a+1) - (c-b)F(b-1) = 0.$$

We have

$$F(1-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) = (1-s)(1-\lambda^2)F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) + sF(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2).$$

$$F(3-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2) = (3-s)(1-\lambda^2)F(4-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2) - (2-s)F(3-s, \frac{1}{2}; \frac{7}{2}-s; \lambda^2).$$

By [Er], p.104, (46),

$$F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad c \neq 0, -1, -2, \dots, \quad \operatorname{Re}(c) > \operatorname{Re}(a+b),$$

we have

$$F(1-s, \frac{1}{2}; \frac{5}{2}-s; 1) = 2 \frac{\Gamma(\frac{5}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}, \quad F(1-s, \frac{3}{2}; \frac{7}{2}-s; 1) = \frac{4}{3} \frac{\Gamma(\frac{7}{2}-s)}{\sqrt{\pi}\Gamma(2-s)}.$$

Therefore,

$$\begin{aligned} & \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=0} \\ &= \frac{1}{2-2s} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} - \frac{\lambda^{2-2s}}{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) + \frac{\lambda^{3-2s}}{3-2s} F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2), \end{aligned}$$

and

$$\begin{aligned}
& \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=1} \\
&= \frac{1}{2-2s} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} - \frac{1}{2-s} (1+\lambda^2) \lambda^{2-2s} F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) \\
&\quad + \frac{1}{2} \frac{1}{(\frac{3}{2}-s)} (3+\lambda^2) \lambda^{3-2s} F(3-s, \frac{3}{2}; \frac{5}{2}-s; \lambda^2) \\
&\quad - \frac{1}{2} \frac{s}{(1-s)(2-s)} \lambda^{2-2s} F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) \\
&\quad - \frac{1}{2} \frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)} (1-\lambda^2) \lambda^{5-2s} F(4-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2) \\
&\quad + \frac{1}{2} \frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)} \lambda^{5-2s} F(3-s, \frac{1}{2}; \frac{7}{2}-s; \lambda^2).
\end{aligned}$$

By [Er], p.111, (5),

$$F(a, b; 2b; \frac{4z}{(1+z)^2}) = (1+z)^{2a} F(a, a+\frac{1}{2}-b; b+\frac{1}{2}; z^2),$$

we have

$$\begin{aligned}
F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) &= (1-\lambda)^{-2(1-s)} F(1-s, 1-s; 2-2s; -u^{-1}), \\
F(2-s, \frac{1}{2}; \frac{5}{2}-s; \lambda^2) &= (1-\lambda)^{-2(2-s)} F(2-s, 2-s; 4-2s; -u^{-1}), \\
F(2-s, \frac{3}{2}; \frac{3}{2}-s; \lambda^2) &= (1-\lambda)^{-2(2-s)} F(2-s, 1-s; 2-2s; -u^{-1}), \\
F(3-s, \frac{3}{2}; \frac{5}{2}-s; \lambda^2) &= (1-\lambda)^{-2(3-s)} F(3-s, 2-s; 4-2s; -u^{-1}), \\
F(1-s, \frac{1}{2}; \frac{3}{2}-s; \lambda^2) &= (1-\lambda)^{-2(1-s)} F(1-s, 1-s; 2-2s; -u^{-1}), \\
F(4-s, \frac{3}{2}; \frac{7}{2}-s; \lambda^2) &= (1-\lambda)^{-2(4-s)} F(4-s, 3-s; 6-2s; -u^{-1}), \\
F(3-s, \frac{1}{2}; \frac{7}{2}-s; \lambda^2) &= (1-\lambda)^{-2(3-s)} F(3-s, 3-s; 6-2s; -u^{-1}).
\end{aligned}$$

Therefore,

$$\begin{aligned}
& \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=0} \\
&= \frac{1}{2-2s} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} - \frac{1}{2-2s} \lambda^{2-2s} (1-\lambda)^{-2(1-s)} F(1-s, 1-s; 2-2s; -u^{-1}) \\
&\quad + \frac{1}{3-2s} \lambda^{3-2s} (1-\lambda)^{-2(2-s)} F(2-s, 2-s; 4-2s; -u^{-1}).
\end{aligned}$$

$$\begin{aligned}
& \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=1} \\
&= \frac{1}{2-2s} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} - \frac{1}{2-s} (1+\lambda^2) \lambda^{2-2s} (1-\lambda)^{-2(2-s)} F(2-s, 1-s; 2-2s; -u^{-1}) \\
&\quad + \frac{1}{2} \frac{1}{(\frac{3}{2}-s)} (3+\lambda^2) \lambda^{3-2s} (1-\lambda)^{-2(3-s)} F(3-s, 2-s; 4-2s; -u^{-1}) \\
&\quad - \frac{1}{2} \frac{s}{(1-s)(2-s)} \lambda^{2-2s} (1-\lambda)^{-2(1-s)} F(1-s, 1-s; 2-2s; -u^{-1}) \\
&\quad - \frac{1}{2} \frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)} (1-\lambda^2) \lambda^{5-2s} (1-\lambda)^{-2(4-s)} F(4-s, 3-s; 6-2s; -u^{-1}) \\
&\quad + \frac{1}{2} \frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)} \lambda^{5-2s} (1-\lambda)^{-2(3-s)} F(3-s, 3-s; 6-2s; -u^{-1}).
\end{aligned}$$

Since $u = \frac{(\lambda-1)^2}{4\lambda}$, we have

$$\begin{aligned}
& \frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} \lambda^s (1-\lambda)^{-2} \int_{\lambda}^1 (1-l)^{2n+1} F(s+n, n+\frac{3}{2}; s+\frac{1}{2}; l^2) dl_{n=0} \\
&= \frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} [-(4u)^{-(s+1)} F(s, s; 2s; -u^{-1}) + \frac{s-\frac{1}{2}}{s-1} (4u)^{-s} \\
&\quad \times F(s-1, s-1; 2s-2; -u^{-1}) - \frac{1}{s-1} \frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} \lambda^s (1-\lambda)^{-2}],
\end{aligned}$$

and

$$\begin{aligned}
& \frac{2\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} \lambda^s (1-\lambda)^{-2} \int_{\lambda}^1 (1-l)^{2n+1} l^{1-2s} F(n-s+2, n+\frac{1}{2}; \frac{3}{2}-s; l^2) dl_{n=0} \\
&= \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} [\frac{1}{1-s} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} \lambda^s (1-\lambda)^{-2} - \frac{1}{1-s} (4u)^{-(2-s)} \\
&\quad \times F(1-s, 1-s; 2-2s; -u^{-1}) + \frac{1}{\frac{3}{2}-s} (4u)^{-(3-s)} F(2-s, 2-s; 4-2s; -u^{-1})].
\end{aligned}$$

Note that

$$\begin{aligned}
& \lambda^s (1-\lambda)^{-2} \left[\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} \frac{(-1)}{\sqrt{\pi}} \frac{1}{s-1} \frac{\Gamma(s+\frac{1}{2})}{\Gamma(s)} + \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} \frac{1}{\sqrt{\pi}} \frac{1}{1-s} \frac{\Gamma(\frac{3}{2}-s)}{\Gamma(2-s)} \right] \\
&= \frac{\lambda^s (1-\lambda)^{-2}}{\pi(1-s)\Gamma(s)\Gamma(2-s)} [\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2}) + \Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s)] \\
&= 0.
\end{aligned}$$

Now, we need the following formulas:

$$\gamma F - \beta z F(\beta + 1, \gamma + 1) - \gamma F(\alpha - 1) = 0.$$

$$(\gamma - 1)F(\gamma - 1) - \alpha F(\alpha + 1) - (\gamma - \alpha - 1)F = 0.$$

We have

$$(2s - 1)F(s, s - 1; 2s - 1; z) - (s - 1)zF(s, s; 2s; z) - (2s - 1)F(s - 1, s - 1; 2s - 1; z) = 0,$$

and

$$(2s - 2)F(s - 1, s - 1; 2s - 2; z) - (s - 1)F(s, s - 1; 2s - 1; z) - (s - 1)F(s - 1, s - 1; 2s - 1; z) = 0.$$

Thus,

$$\frac{4s - 2}{s - 1}F(s, s - 1; 2s - 1; z) = zF(s, s; 2s; z) + \frac{4s - 2}{s - 1}F(s - 1, s - 1; 2s - 2; z).$$

Hence, we have

$$\begin{aligned} & \frac{s - \frac{1}{2}}{s - 1}F(s, s - 1; 2s - 1; -u^{-1}) \\ &= -(4u)^{-1}F(s, s; 2s; -u^{-1}) + \frac{s - \frac{1}{2}}{s - 1}F(s - 1, s - 1; 2s - 2; -u^{-1}). \end{aligned}$$

Similarly,

$$\begin{aligned} & \frac{\frac{3}{2} - s}{1 - s}F(2 - s, 1 - s; 3 - 2s; -u^{-1}) \\ &= -(4u)^{-1}F(2 - s, 2 - s; 4 - 2s; -u^{-1}) + \frac{\frac{3}{2} - s}{1 - s}F(1 - s, 1 - s; 2 - 2s; -u^{-1}). \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{1}{4\pi^2}\phi\left(\frac{1}{2}\rho, \frac{1}{2}\rho'; 0, s\right) &= \frac{1}{1 - s} \frac{\Gamma(\frac{3}{2} - s)}{\sqrt{\pi}\Gamma(2 - s)}(4u)^{-s}F(s, s - 1; 2s - 1; -u^{-1}) \\ &\quad + \frac{1}{s - 1} \frac{\Gamma(s - \frac{1}{2})}{\sqrt{\pi}\Gamma(s)}(4u)^{s-2}F(2 - s, 1 - s; 3 - 2s; -u^{-1}). \end{aligned}$$

For $n = 1$, note that

$$\frac{3!\Gamma(\frac{1}{2} - s)}{\Gamma(3 - s)\Gamma(\frac{3}{2})} = \frac{12\Gamma(\frac{1}{2} - s)}{\sqrt{\pi}\Gamma(3 - s)},$$

we have

$$\begin{aligned}
& \frac{12\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\lambda^s(1-\lambda)^{-2}\int_{\lambda}^1(1-l)^{2n+1}F(s+n,n+\frac{3}{2};s+\frac{1}{2};l^2)dl_{n=1} \\
& = -4\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\frac{1}{s-1}\frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\lambda^s(1-\lambda)^{-2} \\
& \quad -4\frac{\Gamma(\frac{5}{2}-s)}{s(s-1)\sqrt{\pi}\Gamma(3-s)}\lambda(4u)^{-(s-1)}F(s-2,s-2;2s-4;-u^{-1}) \\
& \quad -4\frac{\Gamma(\frac{5}{2}-s)}{s(s-1)\sqrt{\pi}\Gamma(3-s)}(1-\lambda^2)(4u)^{-s}F(s-1,s-2;2s-4;-u^{-1}) \\
& \quad -4\frac{\Gamma(\frac{3}{2}-s)}{s\sqrt{\pi}\Gamma(3-s)}(\lambda+\frac{3}{\lambda})(4u)^{-(s+1)}F(s,s-1;2s-2;-u^{-1}) \\
& \quad -4\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}(4u)^{-(s+1)}F(s,s;2s;-u^{-1}) \\
& \quad -8\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}(\lambda+\frac{1}{\lambda})(4u)^{-(s+2)}F(s+1,s;2s;-u^{-1}).
\end{aligned}$$

Similarly,

$$\frac{3!\Gamma(s-\frac{1}{2})}{\Gamma(s+1)\Gamma(\frac{5}{2})}=\frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)},$$

and

$$\begin{aligned}
& \frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\lambda^s(1-\lambda)^{-2}\int_{\lambda}^1(1-l)^{2n+1}l^{1-2s}F(n-s+2,n+\frac{1}{2};\frac{3}{2}-s;l^2)dl_{n=1} \\
& = \frac{8\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)}\times[\frac{1}{2}\frac{1}{(1-s)}\frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)}\lambda^s(1-\lambda)^{-2} \\
& \quad -\frac{1}{2-s}(\lambda+\frac{1}{\lambda})(4u)^{-(3-s)}F(2-s,1-s;2-2s;-u^{-1}) \\
& \quad +\frac{1}{2}(\frac{1}{(\frac{3}{2}-s)})^2(\lambda+\frac{3}{\lambda})(4u)^{-(4-s)}F(3-s,2-s;4-2s;-u^{-1}) \\
& \quad -\frac{1}{2}\frac{s}{(1-s)(2-s)}(4u)^{-(2-s)}F(1-s,1-s;2-2s;-u^{-1}) \\
& \quad -\frac{1}{2}\frac{(3-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}(1-\lambda^2)(4u)^{-(5-s)}F(4-s,3-s;6-2s;-u^{-1}) \\
& \quad +\frac{1}{2}\frac{(2-s)}{(\frac{3}{2}-s)(\frac{5}{2}-s)}\lambda(4u)^{-(4-s)}F(3-s,3-s;6-2s;-u^{-1})].
\end{aligned}$$

Note that

$$\begin{aligned}
& \left[\frac{-4\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} \frac{1}{s-1} \frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)} + \frac{-4\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)} \frac{1}{s-1} \frac{\Gamma(\frac{3}{2}-s)}{\sqrt{\pi}\Gamma(3-s)} \right] \lambda^s (1-\lambda)^{-2} \\
&= \frac{-4\lambda^s (1-\lambda)^{-2}}{\pi(s-1)\Gamma(s+1)\Gamma(3-s)} \left[\Gamma(\frac{1}{2}-s)\Gamma(s+\frac{1}{2}) + \Gamma(s-\frac{1}{2})\Gamma(\frac{3}{2}-s) \right] \\
&= 0.
\end{aligned}$$

For $k = 1$, if $n = 0$, $\frac{(-1)_n(1)_n}{2^{2n}(n!)^2} = 1$. If $n = 1$, $\frac{(-1)_n(1)_n}{2^{2n}(n!)^2} = -\frac{1}{4}$. Let us add the terms of $n = 0$ and $n = 1$, we have

$$\frac{\Gamma(\frac{1}{2}-s)}{\sqrt{\pi}\Gamma(2-s)} (4u)^{-s} S_I + \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} (4u)^{s-3} S_{II},$$

where

$$\begin{aligned}
S_I &= \frac{\frac{1}{2}-s}{1-s} F(s, s-1; 2s-1; -u^{-1}) + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} 4u\lambda F(s-2, s-2; 2s-4; -u^{-1}) \\
&\quad + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} (1-\lambda^2) F(s-1, s-2; 2s-4; -u^{-1}) \\
&\quad + \frac{\frac{1}{2}-s}{s(2-s)} (\lambda + \frac{3}{\lambda}) \frac{1}{4u} F(s, s-1; 2s-2; -u^{-1}) \\
&\quad + \frac{1}{2-s} \frac{1}{4u} F(s, s; 2s; -u^{-1}) + \frac{1}{2-s} 2(\lambda + \frac{1}{\lambda}) \frac{1}{(4u)^2} F(s+1, s; 2s; -u^{-1}),
\end{aligned}$$

and

$$\begin{aligned}
S_{II} &= \frac{1}{s-1} 4u F(2-s, 1-s; 3-2s; -u^{-1}) \\
&\quad + \frac{2}{s(2-s)} (\lambda + \frac{1}{\lambda}) F(2-s, 1-s; 2-2s; -u^{-1}) \\
&\quad - \frac{1}{s(\frac{3}{2}-s)} (\lambda + \frac{3}{\lambda}) (4u)^{-1} F(3-s, 2-s; 4-2s; -u^{-1}) \\
&\quad + \frac{1}{(1-s)(2-s)} 4u F(1-s, 1-s; 2-2s; -u^{-1}) \\
&\quad + \frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)} (1-\lambda^2) (4u)^{-2} F(4-s, 3-s; 6-2s; -u^{-1}) \\
&\quad - \frac{(2-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)} \lambda (4u)^{-1} F(3-s, 3-s; 6-2s; -u^{-1}).
\end{aligned}$$

Since $u = \frac{(\lambda-1)^2}{4\lambda}$, we have

$$\lambda = \frac{1}{(\sqrt{u+1} + \sqrt{u})^2}.$$

Hence,

$$\begin{aligned}\lambda + \frac{1}{\lambda} &= 4u + 2, \quad \lambda + \frac{3}{\lambda} = 4(2u + 1 + \sqrt{u+1}\sqrt{u}), \\ 1 - \lambda^2 &= \lambda\left(\frac{1}{\lambda} - \lambda\right) = \frac{4\sqrt{u}\sqrt{u+1}}{(\sqrt{u+1} + \sqrt{u})^2}.\end{aligned}$$

Thus, we have

$$\begin{aligned}S_I &= \frac{s - \frac{1}{2}}{s-1} F(s, s-1; 2s-1; -u^{-1}) \\ &\quad + \frac{\left(\frac{3}{2}-s\right)\left(\frac{1}{2}-s\right)}{s(s-1)(2-s)} \frac{4}{(\sqrt{1+\frac{1}{u}}+1)^2} F(s-2, s-2; 2s-4; -u^{-1}) \\ &\quad + \frac{\left(\frac{3}{2}-s\right)\left(\frac{1}{2}-s\right)}{s(s-1)(2-s)} \frac{4\sqrt{1+\frac{1}{u}}}{(\sqrt{1+\frac{1}{u}}+1)^2} F(s-1, s-2; 2s-4; -u^{-1}) \\ &\quad + \frac{\left(\frac{1}{2}-s\right)}{s(2-s)} \left(2 + \frac{1}{u} + \sqrt{1+\frac{1}{u}}\right) F(s, s-1; 2s-2; -u^{-1}) \\ &\quad + \frac{1}{2-s} \frac{1}{4u} F(s, s; 2s; -u^{-1}) + \frac{1}{2-s} \left(\frac{1}{2u} + \frac{1}{4u^2}\right) F(s+1, s; 2s; -u^{-1}). \\ S_{II} &= \frac{1}{s-1} 4u F(2-s, 1-s; 3-2s; -u^{-1}) \\ &\quad + \frac{1}{s(2-s)} 4(2u+1) F(2-s, 1-s; 2-2s; -u^{-1}) \\ &\quad - \frac{1}{s\left(\frac{3}{2}-s\right)} \left(2 + \frac{1}{u} + \sqrt{1+\frac{1}{u}}\right) F(3-s, 2-s; 4-2s; -u^{-1}) \\ &\quad + \frac{1}{(1-s)(2-s)} 4u F(1-s, 1-s; 2-2s; -u^{-1}) \\ &\quad + \frac{(3-s)}{s\left(\frac{3}{2}-s\right)\left(\frac{5}{2}-s\right)} \frac{\sqrt{1+\frac{1}{u}}}{(1+\sqrt{1+\frac{1}{u}})^2} \frac{1}{4u^2} F(4-s, 3-s; 6-2s; -u^{-1}) \\ &\quad - \frac{(2-s)}{s\left(\frac{3}{2}-s\right)\left(\frac{5}{2}-s\right)} \frac{1}{(1+\sqrt{1+\frac{1}{u}})^2} \frac{1}{4u^2} F(3-s, 3-s; 6-2s; -u^{-1}).\end{aligned}$$

Set $z = -u^{-1}$ and $x = \sqrt{1-z}$, then

$$\frac{u}{u+1} = \frac{1}{x^2}, \quad \frac{u^2}{u+1} = \frac{1}{x^2(x^2-1)}.$$

$$\begin{aligned} S_I &= \frac{s-\frac{1}{2}}{s-1} F(s, s-1; 2s-1; 1-x^2) \\ &\quad + \frac{\left(\frac{3}{2}-s\right)\left(\frac{1}{2}-s\right)}{s(s-1)(2-s)} \frac{4}{(x+1)^2} F(s-2, s-2; 2s-4; 1-x^2) \\ &\quad + \frac{\left(\frac{3}{2}-s\right)\left(\frac{1}{2}-s\right)}{s(s-1)(2-s)} \frac{4x}{(x+1)^2} F(s-1, s-2; 2s-4; 1-x^2) \\ &\quad + \frac{\left(\frac{1}{2}-s\right)}{s(2-s)} (x^2+x+1) F(s, s-1; 2s-2; 1-x^2) \\ &\quad + \frac{1}{2-s} \frac{1}{4} (x^2-1) F(s, s; 2s; 1-x^2) + \frac{1}{2-s} \frac{1}{4} (x^4-1) F(s+1, s; 2s; 1-x^2). \\ S_{II} &= \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &\quad + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &\quad - \frac{1}{s\left(\frac{3}{2}-s\right)} (x^2+x+1) F(3-s, 2-s; 4-2s; 1-x^2) \\ &\quad + \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &\quad + \frac{(3-s)}{s\left(\frac{3}{2}-s\right)\left(\frac{5}{2}-s\right)} \frac{1}{4} x(x-1)^2 F(4-s, 3-s; 6-2s; 1-x^2) \\ &\quad - \frac{(2-s)}{s\left(\frac{3}{2}-s\right)\left(\frac{5}{2}-s\right)} \frac{1}{4} (x-1)^2 F(3-s, 3-s; 6-2s; 1-x^2). \end{aligned}$$

At first, we calculate S_I . It is known that (see [Er], p.103, (29))

$$(c-b)F(b-1) + (2b-c-bz+az)F + b(z-1)F(b+1) = 0.$$

Set $a = s$, $b = s$, $c = 2s$ and $z = 1-x^2$, then

$$F(s, s+1; 2s; 1-x^2) = \frac{1}{x^2} F(s, s-1; 2s; 1-x^2),$$

i.e.,

$$F(s+1, s; 2s; 1-x^2) = \frac{1}{x^2} F(s, s-1; 2s; 1-x^2).$$

Set $a = s - 1$, $b = s - 1$, $c = 2s - 2$ and $z = 1 - x^2$, then

$$F(s - 1, s; 2s - 2; 1 - x^2) = \frac{1}{x^2} F(s - 1, s - 2; 2s - 2; 1 - x^2),$$

i.e.,

$$F(s, s - 1; 2s - 2; 1 - x^2) = \frac{1}{x^2} F(s - 1, s - 2; 2s - 2; 1 - x^2).$$

In the formula (see [Er], p.103, (34))

$$c[a - (c - b)z]F - ac(1 - z)F(a + 1) + (c - a)(c - b)zF(c + 1) = 0,$$

set $a = s - 2$, $b = s - 2$, $c = 2s - 4$ and $z = 1 - x^2$, then

$$\begin{aligned} & F(s - 2, s - 2; 2s - 4; 1 - x^2) \\ &= F(s - 1, s - 2; 2s - 4; 1 - x^2) + \frac{x^2 - 1}{2x^2} F(s - 2, s - 2; 2s - 3; 1 - x^2). \end{aligned}$$

Now, we have

$$\begin{aligned} S_I &= \frac{s - \frac{1}{2}}{s - 1} F(s, s - 1; 2s - 1; 1 - x^2) \\ &+ \frac{\left(\frac{3}{2} - s\right)\left(\frac{1}{2} - s\right)}{s(s - 1)(2 - s)} \frac{4}{x + 1} F(s - 1, s - 2; 2s - 4; 1 - x^2) \\ &+ \frac{\left(\frac{3}{2} - s\right)\left(\frac{1}{2} - s\right)}{s(s - 1)(2 - s)} \frac{2(x - 1)}{x^2(x + 1)} F(s - 2, s - 2; 2s - 3; 1 - x^2) \\ &+ \frac{\left(\frac{1}{2} - s\right)}{s(2 - s)} \frac{x^2 + x + 1}{x^2} F(s - 1, s - 2; 2s - 2; 1 - x^2) \\ &+ \frac{1}{2 - s} \frac{1}{4} (x^2 - 1) F(s, s; 2s; 1 - x^2) \\ &+ \frac{1}{2 - s} \frac{1}{4} \frac{x^4 - 1}{x^2} F(s, s - 1; 2s; 1 - x^2). \end{aligned}$$

It is known that

$$\gamma F - \beta zF(\beta + 1, \gamma + 1) - \gamma F(\alpha - 1) = 0.$$

Set $\alpha = s$, $\beta = s - 1$, $\gamma = 2s - 1$ and $z = 1 - x^2$, then

$$F(s, s; 2s; 1 - x^2) = \frac{1}{1 - x^2} \frac{2s - 1}{s - 1} [F(s, s - 1; 2s - 1; 1 - x^2) - F(s - 1, s - 1; 2s - 1; 1 - x^2)].$$

Set $\alpha = s - 1$, $\beta = s - 1$, $\gamma = 2s - 1$ and $z = 1 - x^2$, then

$$\begin{aligned} F(s, s - 1; 2s; 1 - x^2) &= F(s - 1, s; 2s; 1 - x^2) \\ &= \frac{1}{1 - x^2} \frac{2s - 1}{s - 1} [F(s - 1, s - 1; 2s - 1; 1 - x^2) - F(s - 2, s - 1; 2s - 1; 1 - x^2)]. \end{aligned}$$

Thus, we have

$$\begin{aligned} S_I &= \frac{\left(\frac{1}{2} - s\right)\left(s - \frac{3}{2}\right)}{(2 - s)(s - 1)} F(s, s - 1; 2s - 1; 1 - x^2) \\ &\quad + \frac{\left(\frac{3}{2} - s\right)\left(\frac{1}{2} - s\right)}{s(s - 1)(2 - s)} \frac{4}{x + 1} F(s - 1, s - 2; 2s - 4; 1 - x^2) \\ &\quad + \frac{\left(\frac{3}{2} - s\right)\left(\frac{1}{2} - s\right)}{s(s - 1)(2 - s)} \frac{2(x - 1)}{x^2(x + 1)} F(s - 2, s - 2; 2s - 3; 1 - x^2) \\ &\quad + \frac{\left(\frac{1}{2} - s\right)}{s(2 - s)} \frac{x^2 + x + 1}{x^2} F(s - 1, s - 2; 2s - 2; 1 - x^2) \\ &\quad + \frac{\left(\frac{1}{2} - s\right)}{(2 - s)(s - 1)} \frac{1}{2x^2} F(s - 1, s - 1; 2s - 1; 1 - x^2) \\ &\quad - \frac{\left(\frac{1}{2} - s\right)}{(2 - s)(s - 1)} \frac{x^2 + 1}{2x^2} F(s - 1, s - 2; 2s - 1; 1 - x^2). \end{aligned}$$

It is known that (see [Er], p.103, (37))

$$(b - a)(1 - z)F - (c - a)F(a - 1) + (c - b)F(b - 1) = 0.$$

Set $a = s$, $b = s - 1$, $c = 2s - 1$ and $z = 1 - x^2$, then

$$F(s, s - 1; 2s - 1; 1 - x^2) = \frac{1 - s}{x^2} F(s - 1, s - 1; 2s - 1; 1 - x^2) + \frac{s}{x^2} F(s, s - 2; 2s - 1; 1 - x^2).$$

By [Er], p.103, (30),

$$c(c - 1)(z - 1)F(c - 1) + c[c - 1 - (2c - a - b - 1)z]F + (c - a)(c - b)zF(c + 1) = 0,$$

set $a = s - 1$, $b = s - 2$, $c = 2s - 3$ and $z = 1 - x^2$, then

$$\begin{aligned} F(s - 1, s - 2; 2s - 4; 1 - x^2) &= F(s - 1, s - 2; 2s - 3; 1 - x^2) + \frac{(s - 1)}{4(s - \frac{3}{2})} \frac{1 - x^2}{x^2} F(s - 1, s - 2; 2s - 2; 1 - x^2). \end{aligned}$$

Hence,

$$\begin{aligned}
S_I = & \frac{s(\frac{1}{2}-s)(\frac{3}{2}-s)}{(2-s)(1-s)} \frac{1}{x^2} F(s, s-2; 2s-1; 1-x^2) \\
& + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} \frac{4}{x+1} F(s-1, s-2; 2s-3; 1-x^2) \\
& + \frac{(\frac{3}{2}-s)(\frac{1}{2}-s)}{s(s-1)(2-s)} \frac{2(x-1)}{x^2(x+1)} F(s-2, s-2; 2s-3; 1-x^2) \\
& + \frac{(\frac{1}{2}-s)}{s(2-s)} \frac{x+2}{x} F(s-1, s-2; 2s-2; 1-x^2) \\
& - \frac{(\frac{1}{2}-s)^2}{s-1} \frac{1}{x^2} F(s-1, s-1; 2s-1; 1-x^2) \\
& - \frac{(\frac{1}{2}-s)}{(2-s)(s-1)} \frac{x^2+1}{2x^2} F(s-1, s-2; 2s-1; 1-x^2).
\end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = s-2$, $b = s-1$, $c = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(s-1, s-1; 2s-1; 1-x^2) \\
& = \frac{2(s-1)}{s-2} F(s-2, s-1; 2s-2; 1-x^2) - \frac{s}{s-2} F(s-2, s-1; 2s-1; 1-x^2).
\end{aligned}$$

Set $a = s-2$, $b = s-2$, $c = 2s-2$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(s-2, s-2; 2s-3; 1-x^2) \\
& = \frac{(s-1)}{2(s-\frac{3}{2})} F(s-2, s-2; 2s-2; 1-x^2) + \frac{(s-2)}{2(s-\frac{3}{2})} F(s-1, s-2; 2s-2; 1-x^2).
\end{aligned}$$

Set $a = s-1$, $b = s-2$, $c = 2s-1$ and $z = 1-x^2$, then

$$F(s, s-2; 2s-1; 1-x^2) = 2F(s-1, s-2; 2s-2; 1-x^2) - F(s-1, s-2; 2s-1; 1-x^2).$$

Hence,

$$\begin{aligned}
S_I = & \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{x^2+1}{2x^2} \right] F(s-1, s-2; 2s-1; 1-x^2) \\
& + \left[\frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{x^2(x+1)} - \frac{(\frac{1}{2}-s)}{s(s-2)} \frac{x+2}{x} \right] \\
& \times F(s-1, s-2; 2s-2; 1-x^2) \\
& + \frac{(\frac{1}{2}-s)(\frac{3}{2}-s)}{s(s-1)(2-s)} \frac{4}{x+1} F(s-1, s-2; 2s-3; 1-x^2) \\
& + \frac{(\frac{1}{2}-s)}{s(s-2)} \frac{(x-1)}{x^2(x+1)} F(s-2, s-2; 2s-2; 1-x^2).
\end{aligned}$$

We know that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = s-2$, $b = s-2$, $c = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(s-2, s-2; 2s-2; 1-x^2) \\
& = \frac{s}{2(s-1)} F(s-2, s-2; 2s-1; 1-x^2) + \frac{s-2}{2(s-1)} F(s-1, s-2; 2s-1; 1-x^2).
\end{aligned}$$

For $F(s-1, s-2; 2s-3; 1-x^2) = F(s-2, s-1; 2s-3; 1-x^2)$, set $a = s-2$, $b = s-1$, $c = 2s-2$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(s-1, s-2; 2s-3; 1-x^2) \\
& = \frac{s-1}{2(s-\frac{3}{2})} F(s-1, s-2; 2s-2; 1-x^2) + \frac{s-2}{2(s-\frac{3}{2})} F(s-1, s-1; 2s-2; 1-x^2).
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
S_I = & \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{x^2+1}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{2x^2(x+1)} \right] \\
& \times F(s-1, s-2; 2s-1; 1-x^2) \\
& + \left[\frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{x^2(x+1)} - \frac{(\frac{1}{2}-s)}{s(s-2)} \frac{(x+2)}{x} + \frac{(\frac{1}{2}-s)}{s(s-2)} \frac{2}{x+1} \right] \\
& \times F(s-1, s-2; 2s-2; 1-x^2) \\
& + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{2}{x+1} F(s-1, s-1; 2s-2; 1-x^2) \\
& + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)} F(s-2, s-2; 2s-1; 1-x^2).
\end{aligned}$$

Since (see [Er], p.103, (38))

$$c(1-z)F - cF(a-1) + (c-b)zF(c+1) = 0.$$

Set $a = s-1$, $b = s-1$, $c = 2s-2$ and $z = 1-x^2$, then

$$\begin{aligned} & F(s-1, s-1; 2s-2; 1-x^2) \\ &= \frac{1}{x^2} F(s-1, s-2; 2s-2; 1-x^2) + \frac{x^2-1}{2x^2} F(s-1, s-1; 2s-1; 1-x^2). \end{aligned}$$

Thus,

$$\begin{aligned} S_I = & \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x^2+1)}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{2x^2(x+1)} \right] \\ & \times F(s-1, s-2; 2s-1; 1-x^2) \\ & + \frac{(\frac{1}{2}-s)}{s(s-2)} \frac{-(x-1)(x^2+2x+2)}{x^2(x+1)} F(s-1, s-2; 2s-2; 1-x^2) \\ & + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{x^2} F(s-1, s-1; 2s-1; 1-x^2) \\ & + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)} F(s-2, s-2; 2s-1; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = s-2$, $b = s-1$, $c = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned} & F(s-1, s-2; 2s-2; 1-x^2) \\ &= \frac{s}{2(s-1)} F(s-1, s-2; 2s-1; 1-x^2) + \frac{(s-2)}{2(s-1)} F(s-1, s-1; 2s-1; 1-x^2). \end{aligned}$$

Hence,

$$\begin{aligned} S_I = & \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x^2+1)}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{2x^2(x+1)} \right. \\ & \left. + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{-(x-1)(x^2+2x+2)}{2x^2(x+1)} \right] F(s-1, s-2; 2s-1; 1-x^2) \\ (4.30) \quad & + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{-(x-1)}{2(x+1)} F(s-1, s-1; 2s-1; 1-x^2) \\ & + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)} F(s-2, s-2; 2s-1; 1-x^2). \end{aligned}$$

In the following formula (see [Er], p.103, (36))

$$(c-a-b)F - (c-a)F(a-1) + b(1-z)F(b+1) = 0,$$

set $a = s-1$, $b = s-2$, $c = 2s-1$ and $z = 1-x^2$, then

$$\begin{aligned} & 2F(s-1, s-2; 2s-1; 1-x^2) \\ &= -(s-2)x^2F(s-1, s-1, 2s-1; 1-x^2) + sF(s-2, s-2; 2s-1; 1-x^2). \end{aligned}$$

Now, the sum of the latter two terms in (4.30) is equal to

$$\begin{aligned} & \frac{(\frac{1}{2}-s)}{s(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)} [-(s-2)x^2F(s-1, s-1, 2s-1; 1-x^2) \\ & + sF(s-2, s-2; 2s-1; 1-x^2)] \\ &= \frac{(\frac{1}{2}-s)}{s(s-1)(s-2)} \frac{(x-1)}{x^2(x+1)} F(s-1, s-2; 2s-1; 1-x^2). \end{aligned}$$

Thus,

$$\begin{aligned} S_I = & \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x^2+1)}{2x^2} + \frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{2x^2(x+1)} \right. \\ & + \left. \frac{(\frac{1}{2}-s)}{s(s-1)(s-2)} \frac{(x-1)}{x^2(x+1)} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{-(x-1)(x^2+2x+2)}{2x^2(x+1)} \right] \\ & \times F(s-1, s-2; 2s-1; 1-x^2). \end{aligned}$$

Note that

$$\frac{x^2+1}{2x^2} - \frac{(x-1)(x^2+2x+2)}{2x^2(x+1)} = \frac{x+3}{2x^2(x+1)},$$

and

$$\frac{(\frac{1}{2}-s)}{s(s-1)} \frac{(x-1)}{2x^2(x+1)} + \frac{(\frac{1}{2}-s)}{s(s-1)(s-2)} \frac{(x-1)}{x^2(x+1)} = \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)}.$$

We have

$$\begin{aligned} S_I &= \left[\frac{-s(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{1}{x^2} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x+3)}{2x^2(x+1)} + \frac{(\frac{1}{2}-s)}{(s-1)(s-2)} \frac{(x-1)}{2x^2(x+1)} \right] \\ &\quad \times F(s-1, s-2; 2s-1; 1-x^2) \\ &= \frac{\frac{1}{2}-s}{2-s} \frac{1}{x^2} F(s-1, s-2; 2s-1; 1-x^2) \\ &= \frac{\frac{1}{2}-s}{2-s} \frac{u}{u+1} F(s-1, s-2; 2s-1; -u^{-1}). \end{aligned}$$

Therefore, we have

$$\frac{\Gamma(\frac{1}{2} - s)}{\sqrt{\pi}\Gamma(2-s)}(4u)^{-s}S_I = \frac{\Gamma(\frac{3}{2} - s)}{\sqrt{\pi}\Gamma(3-s)}4^{-s}(u+1)^{-1}u^{-(s-1)}F(s-1, s-2; 2s-1; -u^{-1}).$$

Now, we compute S_{II} . It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1) + (2a-c-az+bz)F + a(z-1)F(a+1) = 0.$$

Set $a = 3-s$, $b = 3-s$, $c = 6-2s$ and $z = 1-x^2$, then

$$F(4-s, 3-s; 6-2s; 1-x^2) = \frac{1}{x^2}F(2-s, 3-s; 6-2s; 1-x^2).$$

By [Er], p.103, (39),

$$[a-1-(c-b-1)z]F + (c-a)F(a-1) - (c-1)(1-z)F(c-1) = 0,$$

set $a = 3-s$, $b = 3-s$, $c = 6-2s$ and $z = 1-x^2$, then

$$\begin{aligned} & F(3-s, 3-s; 6-2s; 1-x^2) \\ &= \frac{2(\frac{5}{2}-s)}{2-s}F(3-s, 3-s; 5-2s; 1-x^2) - \frac{3-s}{2-s}\frac{1}{x^2}F(2-s, 3-s; 6-2s; 1-x^2). \end{aligned}$$

Thus,

$$\begin{aligned} S_{II} &= \frac{1}{s-1}\frac{4}{x^2-1}F(2-s, 1-s; 3-2s; 1-x^2) \\ &+ \frac{1}{s(2-s)}\frac{4(x^2+1)}{x^2-1}F(2-s, 1-s; 2-2s; 1-x^2) \\ &- \frac{1}{s(\frac{3}{2}-s)}(x^2+x+1)F(3-s, 2-s; 4-2s; 1-x^2) \\ &+ \frac{1}{(1-s)(2-s)}\frac{4}{x^2-1}F(1-s, 1-s; 2-2s; 1-x^2) \\ &+ \frac{(3-s)}{s(\frac{3}{2}-s)(\frac{5}{2}-s)}\frac{(x-1)^2(x+1)}{4x^2}F(3-s, 2-s; 6-2s; 1-x^2) \\ &- \frac{1}{s(\frac{3}{2}-s)}\frac{1}{2}(x-1)^2F(3-s, 3-s; 5-2s; 1-x^2). \end{aligned}$$

We know that (see [Er], p.103, (30))

$$c(c-1)(z-1)F(c-1) + c[c-1-(2c-a-b-1)z]F + (c-a)(c-b)zF(c+1) = 0.$$

Set $a = 3 - s$, $b = 2 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned} F(3-s, 2-s; 6-2s; 1-x^2) &= \frac{-4(\frac{5}{2}-s)}{3-s} \frac{x^2}{x^2-1} F(3-s, 2-s; 4-2s; 1-x^2) \\ &\quad + \frac{4(\frac{5}{2}-s)}{3-s} \frac{x^2}{x^2-1} F(3-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = 2 - s$, $b = 3 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$F(3-s, 3-s; 5-2s; 1-x^2) = 2F(3-s, 2-s; 4-2s; 1-x^2) - F(3-s, 2-s; 5-2s; 1-x^2).$$

Hence,

$$\begin{aligned} S_{II} &= \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &\quad + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &\quad - \frac{1}{s(\frac{3}{2}-s)} (2x^2+1) F(3-s, 2-s; 4-2s; 1-x^2) \\ &\quad + \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &\quad + \frac{1}{s(\frac{3}{2}-s)} \frac{1}{2} (x^2-1) F(3-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

By [Er], p.103, (39)

$$[a-1-(c-b-1)z]F + (c-a)F(a-1) - (c-1)(1-z)F(c-1) = 0,$$

set $a = 3 - s$, $b = 2 - s$, $c = 5 - 2s$ and $z = 1 - x^2$, then

$$F(3-s, 2-s; 5-2s; 1-x^2) = -\frac{1}{x^2} F(2-s, 2-s; 5-2s; 1-x^2) + 2F(3-s, 2-s; 4-2s; 1-x^2).$$

We have

$$\begin{aligned} S_{II} &= \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &\quad + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &\quad - \frac{1}{s(\frac{3}{2}-s)} (x^2+2) F(3-s, 2-s; 4-2s; 1-x^2) \\ &\quad + \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &\quad - \frac{1}{s(\frac{3}{2}-s)} \frac{x^2-1}{2x^2} F(2-s, 2-s; 5-2s; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1) + (2a-c-az+bz)F + a(z-1)F(a+1) = 0.$$

Set $a = 2-s$, $b = 2-s$, $c = 4-2s$ and $z = 1-x^2$, then

$$F(3-s, 2-s; 4-2s; 1-x^2) = \frac{1}{x^2} F(2-s, 1-s; 4-2s; 1-x^2).$$

By

$$\gamma F - \beta z F(\beta+1, \gamma+1) - \gamma F(\alpha-1) = 0,$$

set $\alpha = 2-s$, $\beta = 1-s$, $\gamma = 4-2s$ and $z = 1-x^2$, then

$$\begin{aligned} & F(2-s, 2-s; 5-2s; 1-x^2) \\ &= \frac{2(2-s)}{1-s} \frac{1}{x^2-1} [F(1-s, 1-s; 4-2s; 1-x^2) - F(2-s, 1-s; 4-2s; 1-x^2)]. \end{aligned}$$

So,

$$\begin{aligned} S_{II} &= \frac{1}{s-1} \frac{4}{x^2-1} F(2-s, 1-s; 3-2s; 1-x^2) \\ &+ \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2-1} F(2-s, 1-s; 2-2s; 1-x^2) \\ &+ \left[\frac{1}{(1-s)(\frac{3}{2}-s)} \frac{1}{x^2} - \frac{1}{s(\frac{3}{2}-s)} \right] F(2-s, 1-s; 4-2s; 1-x^2) \\ &+ \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} F(1-s, 1-s; 2-2s; 1-x^2) \\ &- \frac{(2-s)}{s(1-s)(\frac{3}{2}-s)} \frac{1}{x^2} F(1-s, 1-s; 4-2s; 1-x^2). \end{aligned}$$

We know that (see [Er], p.103, (38))

$$c(1-z)F - cF(a-1) + (c-b)zF(c+1) = 0.$$

Set $a = 2-s$, $b = 1-s$, $c = 2-2s$ and $z = 1-x^2$, then

$$\begin{aligned} & F(2-s, 1-s; 2-2s; 1-x^2) \\ &= \frac{1}{x^2} F(1-s, 1-s; 2-2s; 1-x^2) + \frac{x^2-1}{2x^2} F(2-s, 1-s; 3-2s; 1-x^2). \end{aligned}$$

Thus,

$$\begin{aligned}
S_{II} = & \left[\frac{1}{s-1} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{2(x^2+1)}{x^2} \right] F(2-s, 1-s; 3-2s; 1-x^2) \\
& + \left[\frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2(x^2-1)} \right] F(1-s, 1-s; 2-2s; 1-x^2) \\
& + \left[\frac{1}{(1-s)(\frac{3}{2}-s)} \frac{1}{x^2} - \frac{1}{s(\frac{3}{2}-s)} \right] F(2-s, 1-s; 4-2s; 1-x^2) \\
& - \frac{(2-s)}{s(1-s)(\frac{3}{2}-s)} \frac{1}{x^2} F(1-s, 1-s; 4-2s; 1-x^2).
\end{aligned}$$

It is known that

$$\gamma F - \beta z F(\beta+1, \gamma+1) - \gamma F(\alpha-1) = 0.$$

Set $\alpha = 1-s$, $\beta = -s$, $\gamma = 3-2s$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(1-s, 1-s; 4-2s; 1-x^2) \\
& = \frac{2(\frac{3}{2}-s)}{s} \frac{1}{x^2-1} [F(1-s, -s; 3-2s; 1-x^2) - F(-s, -s; 3-2s; 1-x^2)].
\end{aligned}$$

Set $\alpha = 2-s$, $\beta = -s$, $\gamma = 3-2s$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(2-s, 1-s; 4-2s; 1-x^2) \\
& = \frac{2(\frac{3}{2}-s)}{s} \frac{1}{x^2-1} [F(2-s, -s; 3-2s; 1-x^2) - F(1-s, -s; 3-2s; 1-x^2)].
\end{aligned}$$

Therefore,

$$\begin{aligned}
S_{II} = & \left[\frac{1}{s-1} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{2(x^2+1)}{x^2} \right] F(2-s, 1-s; 3-2s; 1-x^2) \\
& + \left[\frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2(x^2-1)} \right] F(1-s, 1-s; 2-2s; 1-x^2) \\
& + \left[\frac{1}{s(1-s)} \frac{2}{x^2(x^2-1)} - \frac{1}{s^2} \frac{2}{x^2-1} \right] F(2-s, -s; 3-2s; 1-x^2) \\
& + \frac{(2-s)}{s^2(1-s)} \frac{2}{x^2(x^2-1)} F(-s, -s; 3-2s; 1-x^2) \\
& + \left[-\frac{1}{s^2(1-s)} \frac{4}{x^2(x^2-1)} + \frac{1}{s^2} \frac{2}{x^2-1} \right] F(1-s, -s; 3-2s; 1-x^2).
\end{aligned}$$

It is known that (see [Er], p.103, (28))

$$(c-a)F(a-1) + (2a-c-az+bz)F + a(z-1)F(a+1) = 0.$$

Set $a = 1-s$, $b = 1-s$, $c = 3-2s$ and $z = 1-x^2$, then

$$\begin{aligned} & F(2-s, 1-s; 3-2s; 1-x^2) \\ &= \frac{2-s}{1-s} \frac{1}{x^2} F(1-s, -s; 3-2s; 1-x^2) - \frac{1}{1-s} \frac{1}{x^2} F(1-s, 1-s; 3-2s; 1-x^2). \end{aligned}$$

Set $a = 1-s$, $b = -s$, $c = 3-2s$ and $z = 1-x^2$, then

$$\begin{aligned} & F(2-s, -s; 3-2s; 1-x^2) \\ &= \frac{2-s}{1-s} \frac{1}{x^2} F(-s, -s; 3-2s; 1-x^2) + \frac{1}{1-s} \frac{x^2-2}{x^2} F(1-s, -s; 3-2s; 1-x^2). \end{aligned}$$

Note that

$$\begin{aligned} & \frac{1}{(1-s)(2-s)} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{4(x^2+1)}{x^2(x^2-1)} \\ &= \frac{1}{s(1-s)(2-s)} \frac{4}{x^2-1} + \frac{1}{s(2-s)} \frac{4}{x^2(x^2-1)}, \end{aligned}$$

and

$$-\frac{1}{s^2(1-s)} \frac{4}{x^2(x^2-1)} + \frac{1}{s^2} \frac{2}{x^2-1} - \frac{1}{s^2(1-s)} \frac{2(x^2-2)}{x^2(x^2-1)} = \frac{-1}{s(1-s)} \frac{2}{x^2-1}.$$

We have

$$\begin{aligned} S_{II} = & \left[-\frac{1}{s(1-s)} \frac{2}{x^2-1} - \frac{(2-s)}{(1-s)^2} \frac{4}{x^2(x^2-1)} + \frac{1}{s(1-s)} \frac{2(x^2+1)}{x^4} \right. \\ & + \left. \frac{1}{s(1-s)^2} \frac{2(x^2-2)}{x^4(x^2-1)} \right] F(1-s, -s; 3-2s; 1-x^2) \\ & + \left[\frac{1}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{s(1-s)(2-s)} \frac{2(x^2+1)}{x^4} \right] F(1-s, 1-s; 3-2s; 1-x^2) \\ & + \left[\frac{1}{s(2-s)} \frac{4}{x^2(x^2-1)} + \frac{1}{s(1-s)(2-s)} \frac{4}{x^2-1} \right] F(1-s, 1-s; 2-2s; 1-x^2) \\ & + \frac{(2-s)}{s(1-s)^2} \frac{2}{x^4(x^2-1)} F(-s, -s; 3-2s; 1-x^2). \end{aligned}$$

It is known that (see [Er], p.103, (38))

$$c(1-z)F - cF(a-1) + (c-b)zF(c+1) = 0.$$

Set $a = 1 - s$, $b = 1 - s$, $c = 2 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned} & F(1-s, 1-s; 2-2s; 1-x^2) \\ &= \frac{1}{x^2} F(-s, 1-s; 2-2s; 1-x^2) + \frac{x^2-1}{2x^2} F(1-s, 1-s; 3-2s; 1-x^2). \end{aligned}$$

Note that

$$\begin{aligned} & \frac{1}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{s(1-s)(2-s)} \frac{2(x^2+1)}{x^4} + \frac{1}{s(2-s)} \frac{2}{x^4} + \frac{1}{s(1-s)(2-s)} \frac{2}{x^2} \\ &= \frac{1}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)} \frac{2}{x^4}. \end{aligned}$$

Thus,

$$\begin{aligned} S_{II} = & \left[-\frac{1}{s(1-s)} \frac{2}{x^2-1} - \frac{(2-s)}{(1-s)^2} \frac{4}{x^2(x^2-1)} + \frac{1}{s(1-s)} \frac{2(x^2+1)}{x^4} \right. \\ & + \left. \frac{1}{s(1-s)^2} \frac{2(x^2-2)}{x^4(x^2-1)} \right] F(1-s, -s; 3-2s; 1-x^2) \\ & + \left[\frac{1}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)} \frac{2}{x^4} \right] F(1-s, 1-s; 3-2s; 1-x^2) \\ & + \left[\frac{1}{s(2-s)} \frac{4}{x^4(x^2-1)} + \frac{1}{s(1-s)(2-s)} \frac{4}{x^2(x^2-1)} \right] F(-s, 1-s; 2-2s; 1-x^2) \\ & + \left. \frac{(2-s)}{s(1-s)^2} \frac{2}{x^4(x^2-1)} F(-s, -s; 3-2s; 1-x^2) \right]. \end{aligned}$$

We know that (see [Er], p.103, (35))

$$(c-a-1)F + aF(a+1) - (c-1)F(c-1) = 0.$$

Set $a = -s$, $b = 1 - s$, $c = 3 - 2s$ and $z = 1 - x^2$, then

$$\begin{aligned} & F(-s, 1-s; 2-2s; 1-x^2) \\ &= \frac{2-s}{2(1-s)} F(-s, 1-s; 3-2s; 1-x^2) - \frac{s}{2(1-s)} F(1-s, 1-s; 3-2s; 1-x^2). \end{aligned}$$

Note that

$$\begin{aligned} & \frac{1}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{(1-s)(2-s)} \frac{2}{x^4} - \frac{1}{(1-s)(2-s)} \frac{2}{x^4(x^2-1)} \\ & - \frac{1}{(1-s)^2(2-s)} \frac{2}{x^2(x^2-1)} = \frac{1}{(1-s)^2} \frac{2}{x^2(x^2-1)}. \end{aligned}$$

Therefore,

$$\begin{aligned}
S_{II} = & \left[-\frac{1}{s(1-s)} \frac{2}{x^2-1} - \frac{(2-s)}{(1-s)^2} \frac{4}{x^2(x^2-1)} + \frac{1}{s(1-s)} \frac{2(x^2+1)}{x^4} \right. \\
& + \frac{1}{s(1-s)^2} \frac{2(x^2-2)}{x^4(x^2-1)} + \frac{1}{s(1-s)} \frac{2}{x^4(x^2-1)} + \frac{1}{s(1-s)^2} \frac{2}{x^2(x^2-1)} \Big] \\
& \times F(1-s, -s; 3-2s; 1-x^2) \\
& + \frac{1}{(1-s)^2} \frac{2}{x^2(x^2-1)} F(1-s, 1-s; 3-2s; 1-x^2) \\
& + \frac{(2-s)}{s(1-s)^2} \frac{2}{x^4(x^2-1)} F(-s, -s; 3-2s; 1-x^2)
\end{aligned}$$

Note that

$$-\frac{1}{s(1-s)} \frac{2}{x^2-1} + \frac{1}{s(1-s)} \frac{2}{x^4(x^2-1)} + \frac{1}{s(1-s)} \frac{2(x^2+1)}{x^4} = 0,$$

and

$$\begin{aligned}
& \frac{1}{s(1-s)^2} \frac{2(x^2-2)}{x^4(x^2-1)} + \frac{1}{s(1-s)^2} \frac{2}{x^2(x^2-1)} \\
& = \frac{-1}{s(1-s)^2} \frac{4}{x^4(x^2-1)} + \frac{1}{s(1-s)^2} \frac{4}{x^2(x^2-1)}.
\end{aligned}$$

We have

$$\begin{aligned}
S_{II} = & \left[-\frac{(2-s)}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{s(1-s)^2} \frac{4}{x^4(x^2-1)} + \frac{1}{s(1-s)^2} \frac{4}{x^2(x^2-1)} \right] \\
& \times F(1-s, -s; 3-2s; 1-x^2) \\
& + \frac{1}{(1-s)^2} \frac{2}{x^2(x^2-1)} F(1-s, 1-s; 3-2s; 1-x^2) \\
& + \frac{2-s}{s(1-s)^2} \frac{2}{x^4(x^2-1)} F(-s, -s; 3-2s; 1-x^2).
\end{aligned}$$

In the formula (see [Er], p.103, (36))

$$(c-a-b)F - (c-a)F(a-1) + b(1-z)F(b+1) = 0,$$

set $a = 1-s$, $b = -s$, $c = 3-2s$ and $z = 1-x^2$, then

$$\begin{aligned}
& F(1-s, 1-s; 3-2s; 1-x^2) + \frac{2-s}{s} \frac{1}{x^2} F(-s, -s; 3-2s; 1-x^2) \\
& = \frac{1}{s} \frac{2}{x^2} F(1-s, -s; 3-2s; 1-x^2).
\end{aligned}$$

Now, we have

$$\begin{aligned}
& \frac{1}{(1-s)^2} \frac{2}{x^2(x^2-1)} F(1-s, 1-s; 3-2s; 1-x^2) \\
& + \frac{(2-s)}{s(1-s)^2} \frac{2}{x^4(x^2-1)} F(-s, -s; 3-2s; 1-x^2) \\
& = \frac{1}{(1-s)^2} \frac{2}{x^2(x^2-1)} \frac{1}{s} \frac{2}{x^2} F(1-s, -s; 3-2s; 1-x^2) \\
& = \frac{1}{s(1-s)^2} \frac{4}{x^4(x^2-1)} F(1-s, -s; 3-2s; 1-x^2).
\end{aligned}$$

Hence,

$$\begin{aligned}
S_{II} &= \left[-\frac{(2-s)}{(1-s)^2} \frac{4}{x^2(x^2-1)} - \frac{1}{s(1-s)^2} \frac{4}{x^4(x^2-1)} + \frac{1}{s(1-s)^2} \frac{4}{x^2(x^2-1)} \right. \\
&\quad \left. + \frac{1}{s(1-s)^2} \frac{4}{x^4(x^2-1)} \right] F(1-s, -s; 3-2s; 1-x^2) \\
&= \frac{1}{s} \frac{4}{x^2(x^2-1)} F(1-s, -s; 3-2s; 1-x^2) \\
&= \frac{4}{s} \frac{u^2}{u+1} F(1-s, -s; 3-2s; -u^{-1}).
\end{aligned}$$

Therefore, we have

$$\frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} (4u)^{s-3} S_{II} = \frac{\Gamma(s-\frac{1}{2})}{\sqrt{\pi}\Gamma(s+1)} 4^{s-2} u^{s-1} (u+1)^{-1} F(1-s, -s; 3-2s; -u^{-1}).$$

Consequently,

$$\phi(Z, Z'; k, s) = K(Z, Z'; k, s) + K(Z, Z'; k, 2-s),$$

where $k = \pm 1$.

COROLLARY 4.12. *The functional equation of Eisenstein series of weight k for the trivial group on \mathfrak{S}_2 : If $\operatorname{Re}(s) > 1$, then the following identity holds:* □

$$\begin{aligned}
& \int_{\partial\mathfrak{S}_2} P_k(Z, W; s) \rho(W, W')^{s+k-2} \rho(\overline{W}, \overline{W'})^{s-k-2} dm(W) \\
(4.31) \quad &= \frac{\pi^{\frac{3}{2}} \Gamma(s-\frac{1}{2})}{(|k|+s-1)\Gamma(s)} 4^{s-1} P_k(Z, W'; 2-s),
\end{aligned}$$

for $k = 0, \pm 1$, where $W' \in \partial\mathfrak{S}_2$.

Proof. The integral on the left-hand side is absolutely convergent. We set

$$P_{-k}(Z', W; 2-s) = \left[\frac{\rho(Z', W)}{\rho(\overline{Z'}, \overline{W})} \right]^{-k} \frac{\rho(Z')^{2-s}}{|\rho(Z', W)|^{2(2-s)}}$$

in Theorem 4.10. Next, we multiply both sides of the formula by $\rho(Z')^{s-2}$ and take the limit as $Z' \rightarrow W' \in \partial\mathfrak{S}_2$. This completes the proof. \square

We define the S-matrix of weight k as follows:

$$(4.32) \quad S(W, W'; k, s) := \sum_{\gamma \in \Gamma} j(\gamma, W)^{k-s} j(\overline{\gamma}, \overline{W})^{-k-s} \rho(\gamma(W), W')^{-k-s} \rho(\overline{\gamma(W)}, \overline{W'})^{k-s}$$

for $W, W' \in \Omega(\Gamma)$ and $\text{Re}(s) > \delta(\Gamma)$.

THEOREM 4.13. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following functional equation for Eisenstein series of weight k holds:*

$$(4.33) \quad \begin{aligned} & \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; k, s) S(W, W'; -k, 2-s) dm(W) \\ &= \frac{\pi^{\frac{3}{2}} \Gamma(s - \frac{1}{2})}{(|k| + s - 1) \Gamma(s)} 4^{s-1} E(Z, W'; k, 2-s), \end{aligned}$$

for $1 < \text{Re}(s) < 2 - \delta(\Gamma)$ and $k = 0, \pm 1$.

Proof. By Corollary 4.12,

$$(4.34) \quad \begin{aligned} & \int_{\partial\mathfrak{S}_2} E(Z, W; k, s) \rho(W, W')^{s+k-2} \rho(\overline{W}, \overline{W'})^{s-k-2} dm(W) \\ &= \frac{\pi^{\frac{3}{2}} \Gamma(s - \frac{1}{2})}{(|k| + s - 1) \Gamma(s)} 4^{s-1} E(Z, W'; k, 2-s). \end{aligned}$$

On the other hand,

$$\begin{aligned} \int_{\partial\mathfrak{S}_2} f(W) dW &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} f(\gamma(W)) dm(\gamma(W)) \\ &= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |J(\gamma(W))| f(\gamma(W)) dm(W). \end{aligned}$$

By Lemma 4.9, (3), the left hand side of (4.34) is equal to

$$\begin{aligned}
& \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |J(\gamma(W))| E(Z, \gamma(W); k, s) \rho(\gamma(W), W')^{s+k-2} \rho(\overline{\gamma(W)}, \overline{W'})^{s-k-2} dm(W) \\
&= \int_{\Gamma \setminus \Omega(\Gamma)} \sum_{\gamma \in \Gamma} |j(\gamma, W)|^{-4} j(\gamma, W)^{s-k} j(\overline{\gamma}, \overline{W})^{s+k} E(Z, W; k, s) \\
&\quad \times \rho(\gamma(W), W')^{s+k-2} \rho(\overline{\gamma(W)}, \overline{W'})^{s-k-2} dm(W) \\
&= \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; k, s) S(W, W'; -k, 2-s) dm(W).
\end{aligned}$$

□

By Theorem 4.10, we get the main theorem of this paper.

THEOREM 4.14. *Assume that Γ is convex cocompact and $\delta(\Gamma) < 1$. Then the following product formula holds:*

$$\begin{aligned}
(4.35) \quad & \int_{\Gamma \setminus \Omega(\Gamma)} E(Z, W; k, s) E(Z', W; -k, 2-s) dm(W) \\
&= G(Z, Z'; k, s) + G(Z, Z'; k, 2-s) \\
&= G(Z', Z; -k, s) + G(Z', Z; -k, 2-s),
\end{aligned}$$

for $\delta(\Gamma) < \operatorname{Re}(s) < 2 - \delta(\Gamma)$ and $k = 0$ or $k = \pm 1$.

Appendix. Product formulas on $SL(2, \mathbb{R})$

In this appendix, by the same method as in the above argument we give the product formulas on $SL(2, \mathbb{R})$. For simplicity, we omit some details.

Let

$$\Delta_k = -(z - \bar{z})^2 \frac{\partial^2}{\partial z \partial \bar{z}} - k(z - \bar{z}) \left(\frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \right).$$

Set $H_k(z, w) = (\bar{z} - w)^k (z - \bar{w})^{-k}$ and $P(z, \zeta) = \frac{\operatorname{Im}(z)}{|\bar{z} - \zeta|^2}$, where $z, w \in \mathbb{H}$ and $\zeta \in \partial \mathbb{H} = \mathbb{R}$. The Poisson kernel of weight k is defined as follows:

$$P_k(z, \zeta; s) = (\bar{z} - \zeta)^k (z - \zeta)^{-k} P(z, \zeta)^s.$$

$$E(z, \zeta; k, s) := \sum_{\gamma \in \Gamma} P_k(\gamma(z), \zeta; k, s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

Here Γ is a discrete subgroup of $SL(2, \mathbb{R})$. The point-pair invariant

$$u(z, z') := \frac{|z - z'|^2}{4\operatorname{Im}(z)\operatorname{Im}(z')}, \quad \sigma(z, z') := \frac{|\bar{z} - z'|^2}{4\operatorname{Im}(z)\operatorname{Im}(z')}.$$

In fact, $u = \sigma - 1$.

If $f = H_k(z, w)\Phi(\sigma(z, w))$, then

$$\Delta_k f = H_k(z, w) \left[\sigma(\sigma - 1)\Phi''(\sigma) + (2\sigma - 1)\Phi'(\sigma) + \frac{k^2}{\sigma}\Phi(\sigma) \right].$$

By $\Delta_k f = s(s - 1)f$, we have

$$\sigma(\sigma - 1)\Phi''(\sigma) + (2\sigma - 1)\Phi'(\sigma) + \left[\frac{k^2}{\sigma} - s(s - 1) \right] \Phi(\sigma) = 0.$$

A solution is

$$\begin{aligned} \Phi(\sigma) &= P \left\{ \begin{matrix} 0 & 1 & \infty \\ -|k| & 0 & s \\ |k| & 0 & 1-s \end{matrix}; \sigma \right\} \\ &= \sigma^{-|k|} (1 - \sigma)^{|k|-s} F(s - |k|, s - |k|; 2s; -\frac{1}{\sigma - 1}). \end{aligned}$$

Set

$$\begin{aligned} K(z, z'; k, s) &= \frac{\sqrt{\pi}\Gamma(\frac{1}{2} - s)}{\Gamma(1 - s)} 4^{-s} \frac{s}{s - |k|} H_k(z, z') \\ &\quad \times u^{|k|-s} (1 + u)^{-|k|} F(s - |k|, s - |k|; 2s; -u^{-1}), \end{aligned}$$

where $u = u(z, z')$. The automorphic Green function of weight k is given by

$$G(z, z'; k, s) = \sum_{\gamma \in \Gamma} K(z, \gamma(z'); k, s), \quad \text{for } \operatorname{Re}(s) > \delta(\Gamma).$$

Let

$$\phi(z, z'; k, s) = \int_{\mathbb{R}} P_k(z, \zeta; s) P_{-k}(z', \zeta; 1 - s) d\zeta.$$

Then ϕ is a point-pair invariant covariant with respect to the weight k .

Put $z = iy$, $z' = iy'$, $y > 0$, $y' > 0$. Then

$$\begin{aligned} \phi(iy, iy'; k, s) &= y^s y'^{1-s} \int_{-\infty}^{\infty} \left(\frac{\zeta + iy}{\zeta - iy} \right)^k \left(\frac{\zeta + iy'}{\zeta - iy'} \right)^{-k} |\zeta - iy|^{-2s} |\zeta - iy'|^{-2(1-s)} d\zeta \\ &= y^s y'^{1-s} \int_{-\infty}^{\infty} \exp(2ki \arctan \frac{y}{\zeta} - 2ki \arctan \frac{y'}{\zeta}) \frac{1}{(\zeta^2 + y^2)^s (y'^2 + \zeta^2)^{1-s}} d\zeta. \end{aligned}$$

By $\tan(\arctan \frac{y}{\zeta} - \arctan \frac{y'}{\zeta}) = \frac{\zeta(y-y')}{\zeta^2 + yy'}$,

$$\cos(2 \arctan \left[\frac{(y-y')\zeta}{yy'+\zeta^2} \right]) = \frac{(yy'+\zeta^2)^2 - (y-y')^2\zeta^2}{(yy'+\zeta^2)^2 + (y-y')^2\zeta^2},$$

and

$$\begin{aligned} & \cos(2k \arctan \left[\frac{(y-y')\zeta}{yy'+\zeta^2} \right]) \\ &= \cos(k \arccos \frac{(yy'+\zeta^2)^2 - (y-y')^2\zeta^2}{(yy'+\zeta^2)^2 + (y-y')^2\zeta^2}) \\ &= F(-k, k; \frac{1}{2}; \frac{1}{2} \left[1 - \frac{(yy'+\zeta^2)^2 - (y-y')^2\zeta^2}{(yy'+\zeta^2)^2 + (y-y')^2\zeta^2} \right]) \\ &= F(-k, k; \frac{1}{2}; \frac{(y-y')^2\zeta^2}{(y^2+\zeta^2)(y'^2+\zeta^2)}), \end{aligned}$$

we have

$$\begin{aligned} \phi(iy, iy'; k, s) &= 2y^s y'^{1-s} \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n! (\frac{1}{2})_n} (y-y')^{2n} \\ &\quad \times \int_0^\infty \zeta^{2n} (\zeta^2 + y^2)^{-s-n} (\zeta^2 + y'^2)^{s-n-1} d\zeta. \end{aligned}$$

Set $l = \frac{y'}{y}$, then $u = \frac{(l-1)^2}{4l}$. By the same method as the above argument for $GU(2, 1)$, we have

$$\begin{aligned} & \phi(iy, iy'; k, s) \\ &= \sum_{n=0}^{|k|} \frac{(-k)_n (k)_n}{n! (\frac{1}{2})_n} \Gamma(n + \frac{1}{2}) [\frac{\Gamma(\frac{1}{2} - s)}{\Gamma(n+1-s)} (4u)^{-s} F(n+s, s; 2s; -u^{-1}) \\ &\quad + \frac{\Gamma(s - \frac{1}{2})}{\Gamma(n+s)} (4u)^{-(1-s)} F(n+1-s, 1-s; 2-2s; -u^{-1})]. \end{aligned}$$

(1) $k = 0$,

$$\phi(iy, iy'; k, s) = \frac{\sqrt{\pi} \Gamma(\frac{1}{2} - s)}{\Gamma(1-s)} (4u)^{-s} F(s, s; 2s; -u^{-1}) + (s \mapsto 1-s).$$

(2) $k = \pm 1$,

$$S_I := \frac{\sqrt{\pi} \Gamma(\frac{1}{2} - s)}{\Gamma(1-s)} (4u)^{-s} [F(s, s; 2s; -u^{-1}) + \frac{1}{s-1} F(s+1, s; 2s; -u^{-1})].$$

By [Er], p.103, (28),

$$(c-a)F(a-1) + (2a-c-az+bz)F + a(z-1)F(a+1) = 0,$$

set $a = s$, $b = s$, $c = 2s$ and $z = -u^{-1}$, then

$$F(s+1, s; 2s; -u^{-1}) = \frac{u}{u+1} F(s-1, s; 2s; -u^{-1}).$$

By [Er], p.103, (36),

$$(c-a-b)F - (c-a)F(a-1) + b(1-z)F(b+1) = 0,$$

set $a = s$, $b = s-1$, $c = 2s$ and $z = -u^{-1}$, then

$$\begin{aligned} & \frac{1}{s-1} \frac{u}{u+1} F(s, s-1; 2s; -u^{-1}) + F(s, s; 2s; -u^{-1}) \\ &= \frac{s}{s-1} \frac{u}{u+1} F(s-1, s-1; 2s; -u^{-1}). \end{aligned}$$

Thus,

$$S_I = \frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)} (4u)^{-s} \frac{s}{s-1} \frac{u}{u+1} F(s-1, s-1; 2s; -u^{-1}).$$

Consequently,

$$\phi(iy, iy'; k, s) = \frac{\sqrt{\pi}\Gamma(\frac{1}{2}-s)}{\Gamma(1-s)} 4^{-s} \frac{s}{s-1} u^{1-s} (u+1)^{-1} F(s-1, s-1; 2s; -u^{-1}) + (s \mapsto 1-s).$$

Thus, we have the following theorem:

THEOREM. *Assume that Γ is convex cocompact and $\delta(\Gamma) < \frac{1}{2}$. Then the following product formula on $SL(2, \mathbb{R})$ holds:*

$$\int_{\Gamma \backslash \Omega(\Gamma)} E(z, \zeta; k, s) E(z', \zeta; -k, 1-s) dm(\zeta) = G(z, z'; k, s) + G(z, z'; k, 1-s),$$

for $\delta(\Gamma) < Re(s) < 1 - \delta(\Gamma)$ and $k = 0, \pm 1$.

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY
 BEIJING 100871, P. R. CHINA
E-mail address: yanglei@sxx0.math.pku.edu.cn

REFERENCES

- [Ap] B. APANASOV, Deformations and stability in complex hyperbolic geometry, MSRI preprint, No. 1997-111.
- [BB] W. L. BAILY AND A. BOREL, Compactification of arithmetic quotients of bounded symmetric domains, Ann. of Math. **84** (1966), 442-528.
- [BO] U. BUNKE AND M. OLBRICH, The spectrum of Kleinian manifolds, math.DG/9810146.
- [Co] K. CORLETTE, Hausdorff dimensions of limit sets I, Invent. Math. **102** (1990), 521-542.
- [CoI] K. CORLETTE AND A. IOZZI, Limit sets of discrete groups of isometries of exotic hyperbolic spaces, Trans. AMS **351** (1999), 1507-1530.
- [El] J. ELSTRODT, Die Resolvente zum Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene, I, Math. Ann. **203** (1973), 295-330; II, Math. Z. **132** (1973), 99-134; III, Math. Ann. **208** (1974), 99-132.
- [ElGM] J. ELSTRODT, F. GRUNEWALD AND J. MENNICKE, *Groups Acting on Hyperbolic Space, Harmonic Analysis and Number Theory*, Springer-Verlag, 1998.
- [Er] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER AND F. G. TRICOMI, *Higher Transcendental Functions*, Vol. I, McGraw-Hill Book Company, Inc., 1953.
- [Fa] J. D. FAY, Fourier coefficients of the resolvent for a Fuchsian group, J. Reine Angew. Math. **294** (1977), 143-203.
- [FHP] R. FROESE, P. HISLOP AND P. PERRY, The Laplace operator on hyperbolic three manifolds with cusps of non-maximal rank, Invent. Math. **106** (1991), 295-333.
- [Gir] G. GIRAUD, Sur certaines fonctions automorphes de deux variables, Ann. Ec. Norm., (3), **38** (1921), 43-164.
- [Go] W. M. GOLDMAN, *Complex Hyperbolic Geometry*, Oxford Mathematical Monographs, Oxford, 1999.
- [H] D. HEJHAL, *The Selberg Trace Formula for $PSL(2, \mathbb{R})$* , Vol. 1, Lecture Notes in Math. **548**, Springer-Verlag, 1976; Vol. 2, Lecture Notes in Math. **1001**, Springer-Verlag, 1983.
- [He] J. C. HEMPERLY, The parabolic contribution to the number of linearly independent automorphic forms on a certain bounded domain, Amer. J. Math. **94** (1972), 1078-1100.
- [HP] S. HERSONSKY AND F. PAULIN, On the volumes of complex hyperbolic manifolds, Duke Math. J. **84** (1996), 719-737.
- [Ho1] R. P. HOLZAPFEL, *Geometry and Arithmetic Around Euler Partial Differential Equations*, Dt. Verl. d. Wiss., Berlin/Reidel, Dordrecht 1986.
- [Ho2] R. P. HOLZAPFEL, *The Ball and Some Hilbert Problems*, Lectures in Mathematics, ETH Zürich, Birkhäuser, 1995.
- [K] M. KASHIWARA, A. KOWATA, K. MINEMURA, K. OKAMOTO, T. OSHIMA AND M. TANAKA, Eigenfunctions of invariant differential operators on a symmetric space, Ann. of Math. 107(1978), 1-39.
- [Ko] A. KORÁNYI, The Poisson integral for generalized half-planes and bounded symmetric domains, Ann. of Math. **82** (1965), 332-350.
- [KR] A. KORÁNYI AND H. M. REIMANN, Quasiconformal mappings on the Heisenberg group, Invent. Math. **80** (1985), 309-338.
- [KW] A. KORÁNYI AND J. A. WOLF, Realization of hermitian symmetric spaces as generalized half-planes, Ann. of Math. **81** (1965), 265-288.
- [Ku] T. KUBOTA, *Elementary Theory of Eisenstein Series*. Wiley, New York, 1973.
- [LR] R. P. LANGLANDS AND D. RAMAKRISHNAN, *The Zeta Functions of Picard Modular Surfaces*, Les Publications CRM, Université de Montréal, 1992.
- [M] H. MAASS, Die Differentialgleichungen in der Theorie der elliptischen Modulfunktionen, Math. Ann. **125** (1953), 235-263.
- [Ma1] N. MANDOUVALOS, Spectral theory and Eisenstein series for Kleinian groups, Proc. London Math. Soc. (3) **57** (1988), 209-238.

- [Ma2] N. MANDOUVALOS, *Scattering Operator, Eisenstein Series, Inner Product Formula and “Maass-Selberg” Relations for Kleinian Groups*, Memoirs of A.M.S. **78**, No. **400** (1989).
- [P] J. R. PARKER, On the volumes of cusped, complex hyperbolic manifolds and orbifolds, Duke Math. J. **94** (1998), 433-464.
- [Pa1] S. J. PATTERSON, The limit set of a Fuchsian group, Acta Math. **136** (1976), 241-273.
- [Pa2] S. J. PATTERSON, The Laplacian operator on a Riemann surface, I, Comp. Math. **31** (1975), 83-107; II, Comp. Math. **32** (1976), 71-119, III, Comp. Math. **33** (1976), 227-259.
- [Pa3] S. J. PATTERSON, The Selberg zeta function of a Kleinian group. In *Number Theory, Trace Formulas, and Discrete Groups*: Symposium in honor of Atle Selberg, Oslo, Norway (1987), 409-442.
- [Pe1] P. A. PERRY, The Laplace operator on a hyperbolic manifold. II. Eisenstein series and the scattering matrix, J. Reine Angew. Math. **398** (1989), 67-91.
- [Pe2] P. A. PERRY, The Selberg zeta function and a local trace formula for Kleinian groups, J. Reine Angew. Math. **410** (1990), 116-152.
- [Pic1] E. PICARD, Sur une extension aux fonctions de deux variables du problème de Riemann relatif aux fonctions hypergéométriques séries hypergéométriques, Ann. Ec. Norm. **10** (1881), 305-322.
- [Pic2] E. PICARD, Sur les fonctions hyperfuchsiennes provenant des séries hypergéométriques de deux variables, Ann. Ec. Norm. **III 2** (1885), 357-384.
- [Ro] W. ROELCKE, Das Eigenwertproblem der automorphen Formen in der hyperbolischen Ebene, I, Math. Ann. **167** (1966), 292-337; II, Math. Ann. **168** (1967), 261-324.
- [Se] A. SELBERG, Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series, J. Indian Math. Soc. B. **20** (1956), 47-87.
- [Sh1] G. SHIMURA, The arithmetic of automorphic forms with respect to a unitary group, Ann. of Math. **107** (1978), 569-605.
- [Sh2] G. SHIMURA, L-functions and eigenvalue problems. In *Algebraic Analysis, Geometry, and Number Theory*, (J. I. Igusa, ed.), Johns Hopkins University Press, Baltimore, 1989, 341-396.
- [Su] D. SULLIVAN, The density at infinity of a discrete group of hyperbolic motions, Publ. Math. I.H.E.S. **50** (1979), 171-209.