# Integral Transform and Segal-Bargmann Representation associated to q-Charlier Polynomials

Nobuhiro ASAI International Institute for Advanced Studies Kyoto, 619-0225, Japan. asai@iias.or.jp

April 24, 2001

#### Abstract

Let  $\mu_p^{(q)}$  be the q-deformed Poisson measure in the sense of Saitoh-Yoshida [24] and  $\nu_p$  be the measure given by Equation (3.6). In this short paper, we introduce the q-deformed analogue of the Segal-Bargmann transform associated with  $\mu_p^{(q)}$ . We prove that our Segal-Bargmann transform is a unitary map of  $L^2(\mu_p^{(q)})$  onto the Hardy space  $\mathcal{H}^2(\nu_q)$ . Moreover, we give the Segal-Bargmann representation of the multiplication operator by x in  $L^2(\mu_p^{(q)})$ , which is a linear combination of the q-creation, q-annihilation, q-number, and scalar operators.

#### 1 Introduction

The classical Segal-Bargmann transform in Gaussian analysis yields a unitary map of  $L^2$  space of the Gaussian measure on  $\mathbb{R}$  onto the space of  $L^2$  holomorphic functions of the Gaussian measure on  $\mathbb{C}$ , see papers [7, 8, 15, 16, 19, 25, 26].

Recently, Accardi-Bożejko [1] showed the existence of a unitary operator between a one-mode interacting Fock space and  $L^2$  space of a probability measure on  $\mathbb{R}$  by making use of the basic properties of classical orthogonal polynomials and associated recurrence formulas [14, 27]. Inspired by this work, the author [3] has recently extended the Segal-Bargmann transform to non-Gaussian cases. The crucial point is to introduce a coherent state vector as a kernel function in such a way that a transformed function, which is a holomorphic function on a certain domain in general, becomes a power series expression. Along this line, Asai-Kubo-Kuo [6] have considered the case of the Poisson measure compared with the case of the Gaussian measure. However, the case of  $L^2$  space of Wigner's semi-circle distributions in free probability theory [28] is beyond their scope.

On the other hand, Leeuwen-Maassen [22] considered a transform associated with q deformation of the Gaussian measure [10, 11, 12] and showed that for a given real number  $q \in [0,1)$  it is a unitary map of  $L^2$  space of q-defomed Gaussian measure onto Hardy space  $\mathcal{H}^2(\nu_q)$  where  $\nu_q$  is given in (3.6). Biane

[9] examined the case of q=0 (Free case). Roughly speaking, their methods are valid only if kernel functions for their transforms are known explicitly. As observed in Section 3 and Appendix A, our approach does not require to possess the explicit forms of kernel functions.

In this paper, we shall consider the q-deformed version of the Segal-Bargmann transform  $S_{\mu_p^{(q)}}$  associated with q-deformed Poisson measure, denoted by  $\mu_p^{(q)}$ , in the sense of Saitoh-Yoshida [24]. As a main result, we shall provide Proposition 4.1, which claims that  $S_{\mu_p^{(q)}}$  is a unitary map of  $L^2(\mu_p^{(q)})$  onto  $\mathcal{H}^2(\nu_q)$ . Moreover, in Theorem 4.3 we shall give the representation in  $\mathcal{H}^2(\nu_q)$  of the multiplication operator by x in  $L^2(\mu_p^{(q)})$ , which is a linear combination of the q-creation, q-annihilation, q-number, and scalar operators. We remark that our representation is compatible with that on the q-Fock space by Saitoh-Yoshida [23] and can be viewed as the q-analogue of the Hudson-Parthasarathy [17] decomposition of the usual Poisson random variable on the standard Boson Fock space (q=1). Ito-Kubo [18] also studied a similar decomposition in details from the point of white noise calculus [20, 21] (For more recent formulation, see papers [4, 5, 13]).

The present article serves a good example to papers by Accardi-Bożejko [1] and Asai [3]. The present paper is organized as follows. In Section 2, we recall the recurrence formula for q-Charlier polynomials. In Section 3, we introduce a q-deformed coherent state vector and a Segal-Bargmann transform associated to a q-deformed Poisson measure. In addition, we quickly define the Hardy space as the Segal-Bargmann representation space. In Section 4, our main results are given. In Appendix A, we give some remarks on known results [7, 9, 22, 25, 26] related to q-Hermite polynomials.

#### Notation.

Let us recall standard notation from q-analysis [2]. We put for  $n \in \mathbb{N}_0$ ,

$$[n]_q := 1 + q + \dots + q^{n-1}$$
 with  $[0]_q = 0$ .

Then q-factorial is naturally defined as

$$[n]_q! := [1]_q \cdots [n]_q$$
 with  $[0]_q! = 0$ .

The q-exponential is given by

$$\exp_q(x) := \sum_{n=0}^{\infty} \frac{x^n}{[n]_q!}.$$

whose radious of convergence is 1/(1-q). In addition, another symbol used is the q-analogue of the Pochhammer symbol,

$$(a;q)_n = \prod_{j=0}^{n-1} (1 - aq^j)$$
 and  $(a;q)_\infty = \prod_{j=0}^\infty (1 - aq^j)$ 

with the convention  $(a;q)_0 = 1$ .

## 2 q-deformed Charlier Polynomials

From now on, we always assume that  $q \in [0,1)$  is fixed. Recently, Saitoh-Yoshida [24] calculated the explicit form of the q-deformed Poisson measure with a parameter  $\beta > 0$  for  $q \in [0,1)$ . We denote it by  $\mu_p^{(q)}$ . The orthogonal polynomials associated to  $\mu_p^{(q)}$  are the q-Charlier polynomials  $\{C_n^{(q)}(x)\}$  with the Szegö-Jacobi parameters  $\alpha_n = [n]_q + \beta - 1, \omega_n = \beta[n]_q$ . See papers [23, 24]. We also refer the book [14] for the standard Charlier polynomials case (q = 1). Let  $\lambda = \{\beta^n[n]_q!\}_{n=0}^{\infty}$ . The following relations hold for each  $n \geq 1$ :

$$(x - [n]_q - \beta)C_n^{(q)}(x) = C_{n+1}^{(q)}(x) + \beta[n]_q C_{n-1}^{(q)}(x)$$
(2.1)

where  $C_0^{(q)} = 1$ ,  $C_1^{(q)} = x - \beta$ . Then, for any  $L^2$ -convergent decompositions  $f(x) = \sum a_n C_n^{(q)}(x)$  and  $g(x) = \sum b_n C_n^{(q)}(x)$ , the inner product  $\langle \cdot, \cdot \rangle_{L^2(\mu_p^{(q)})}$  is given by the form

$$\langle f, g \rangle_{L^2(\mu_p^{(q)})} = \sum_n \beta^n n! \overline{a}_n b_n.$$
 (2.2)

Moreover, let  $||f||_{L^2(\mu_p^{(q)})}^2 = \sum_{n=0}^{\infty} \beta^n [n]_q! |a_n|^2$ .

### 3 Segal-Bargmann Transform and Hardy Space

Let us define the q-deformed coherent state vector for  $\{C_n^{(q)}(x)\}$  by

$$E_{\lambda,p}^{(q)}(x,z) = \sum_{n=0}^{\infty} \frac{C_n^{(q)}(x)}{\beta^n [n]_q!} z^n, \quad z \in \Omega_q^{\beta} := \left\{ z \in \mathbb{C} : |z|^2 < \frac{\beta}{1-q} \right\}.$$
 (3.1)

It is easy to see that  $E_p^{(q)}(x,z) \in L^2(\mu_p^{(q)})$  due to

$$||E_{\lambda,p}^{(q)}(x,z)||_{L^2(\mu_p^{(q)})}^2 = \exp_q(\beta^{-1}|z|^2) < \infty.$$
 (3.2)

Moreover, it can be shown that  $\{E_{\lambda,p}^{(q)}(x,y)\}$  is linearly independent and total in  $L^2(\mu_p)$ .

Now we are in a position to introduce our key tool in this paper. Let us consider the q-analogue of the Segal-Bargmann transform  $S_{\mu_p^{(q)}}$  associated to  $\{C_n^{(q)}(x)\}$  given by

$$(S_{\mu_p^{(q)}}f)(z) = \langle E_{\lambda,p}^{(q)}(x,\overline{z}), f(x) \rangle_{L^2(\mu_p^{(q)})} \text{ for any } f \in L^2(\mu_p^{(q)}).$$
 (3.3)

**Lemma 3.1.** Let  $f \in L^2(\mu_p^{(q)})$ . Then  $(S_{\mu_p^{(q)}}f)(z)$  converges absolutely for all  $z \in \Omega_q^{\beta}$ .

*Proof.* For  $f(x) = \sum_{n=0}^{\infty} a_n C_n^{(q)}(x)$ , it is quite easy to see

$$(S_{\mu_p^{(q)}}f)(z) = \sum_{n=0}^{\infty} a_n z^n.$$
(3.4)

By the Schwartz inequality, we get the inequality

$$\sum_{n=0}^{\infty} |a_n z^n| \le \|f\|_{L^2(\mu_p^{(q)})} \exp_q\left(\frac{|z|^2}{\beta}\right).$$

This shows that the  $(S_{\mu_p^{(q)}}f)(z)$  converges absolutely for all  $z\in\Omega_q^\beta$ .

The completion of the space of holomorphic functions F,G on  $\Omega_q^\beta$  with respect to the inner product,

$$(F,G)_{\mathcal{H}_q^2} = \frac{1}{2\pi} \int_0^{2\pi} F\left(\sqrt{\frac{\beta}{1-q}} e^{i\theta}\right) G\left(\sqrt{\frac{\beta}{1-q}} e^{i\theta}\right) d\theta \tag{3.5}$$

is nothing but the Hardy space  $\mathcal{H}^2(\nu_q)$ . Here  $\nu_q$  means that

$$\nu_q(dz) = (q;q)_{\infty} \sum_{j=0}^{\infty} \frac{q^j}{(q;q)_j} \lambda_{r_j}^{\beta}(dz), \quad q \in [0,1), \ r_j = q^{\frac{j}{2}} \sqrt{\frac{\beta}{1-q}}$$
 (3.6)

where  $\lambda_{r_j}^{\beta}(dz)$  is the Lebesgue measure on the circle of radious  $r_j$ .

**Lemma 3.2.**  $\{z^n\}_{n=0}^{\infty}$  forms an orthogonal basis of  $\mathcal{H}^2(\nu_q)$ .

*Proof.* We adopt the same idea as in the proof [22].

$$\begin{split} (z^{n}, z^{m})_{\mathcal{H}_{q}^{2}} &= \int_{\Omega_{q}^{\beta}} \overline{z}^{n} z^{m} \nu_{q}(dz) \\ &= (q; q)_{\infty} \sum_{j=0}^{\infty} \frac{q^{j}}{(q; q)_{j}} \int_{\Omega_{q}^{\beta}} \overline{z}^{n} z^{m} \lambda_{r_{j}}^{\beta}(dz) \\ &= \frac{(q; q)_{\infty}}{2\pi} \sum_{j=0}^{\infty} \frac{q^{j}}{(q; q)_{j}} r_{j}^{n+m} \int_{0}^{2\pi} e^{i(m-n)\theta} d\theta \\ &= \delta_{n,m} \frac{(q; q)_{\infty}}{(1-q)^{n}} \sum_{j=0}^{\infty} \frac{q^{(n+1)j}}{(q; q)_{j}} \\ &= \delta_{n,m} [n]_{q}!. \end{split}$$

Note that we have used the q-Gamma function [2, 22],

$$\Gamma_q(n+1) := \frac{(q;q)_{\infty}}{(1-q)^n} \sum_{j=0}^{\infty} \frac{q^{(n+1)j}}{(q;q)_j} = [n]_q! \text{ for any } n \in \mathbb{N}.$$

Hence, for any  $F = \sum_{n=0}^{\infty} a_n z^n$ ,  $G = \sum_{n=0}^{\infty} b_n z^n \in \mathcal{H}^2(\nu_q)$ , the inner product  $(\cdot, \cdot)_{\mathcal{H}^2_q}$  is written as

$$(F,G)_{\mathcal{H}_q^2} = \sum_{n=0}^{\infty} \beta^n [n]_q! \overline{a}_n b_n \tag{3.7}$$

and the corresponding norm of F is

$$||F||_{\mathcal{H}_q^2}^2 = \sum_{n=0}^{\infty} \beta^n [n]_q! |a_n|^2.$$
 (3.8)

#### 4 Main Results

**Proposition 4.1.**  $S_{\mu_p^{(q)}}$  is a unitary map of  $L^2(\mu_p^{(q)})$  onto  $\mathcal{H}^2(\nu_q)$ .

Proof. As we have seen in Lemma 3.1,

$$(S_{\mu_n^{(q)}}C_n^{(q)})(z) = z^n. (4.1)$$

In addition, we derive by Lemma 3.2,

$$\|C_n^{(q)}\|_{L^2(\mu_p^{(q)})}^2 = \|z^n\|_{\mathcal{H}_q^2}^2 = \beta^n.$$

Therefore, we finish the proof.

Let us define operators  $Z_q$  and  $D_q$  in  $\mathcal{H}^2(\nu_q)$  satisfying

$$Z_q F(z) = z F(z). (4.2)$$

and

$$D_{q,\beta}F(z) := 0 \ (z = 0), \quad D_{q,\beta}F(z) := \frac{\beta(F(z) - F(qz))}{z(1 - q)}. \ (z \neq 0)$$
 (4.3)

Operators  $Z_q$  and  $D_{q,\beta}$  play the roles of the q-creation operator and q-annihilation operator respectively and satisfy the q-deformed commutation relation  $D_{q,\beta}Z_q - qZ_qD_{q,\beta} = I$ . The q-number operator acting on  $\mathcal{H}^2(\nu_q)$  is defined by

$$\widetilde{N}_q F(z) = [n]_q F(z), \ n \ge 0. \tag{4.4}$$

In addition, the operator  $\tilde{\alpha}_{N_q}$  acting on  $\mathcal{H}^2(\nu_q)$  is defined by

$$\widetilde{\alpha}_{N_q} F(z) = ([n]_q + \beta) F(z), \quad n \ge 0. \tag{4.5}$$

Remark that  $\widetilde{\alpha}_{N_q}F(z)=0$  for q-Gaussian case, see Appendix A. By the direct calculation, we have

**Lemma 4.2.** (1)  $S_{\mu_n^{(q)}} 1 = 1$ 

- $(2) D_{q,\beta}z^n = \beta[n]_q z^{n-1}$
- $(3) Z_q z^n = z^{n+1}$

The transformation of the multiplication operator  $Q_p^{(q)}$  by x in  $L^2(\mu_p^{(q)})$  satisfies the following relation.

Theorem 4.3. 
$$S_{\mu_{q}^{(q)}}Q_{p}^{(q)}=(D_{q,\beta}+Z_{q}+\widetilde{\alpha}_{N_{q}})S_{\mu_{q}^{(q)}}$$

Proof. By the recurrence formula (2.1), Equation (4.1) and Lemma 4.2, we derive

$$\begin{split} (S_{\mu_p^{(q)}}Q_p^{(q)}C_n^{(q)})(z) &= \big\langle E_{\lambda,p}^{(q)}(x,\overline{z}), C_{n+1}^{(q)} + \beta[n]_q C_{n-1}^{(q)} + ([n]_q + \beta) C_n^{(q)} \big\rangle_{L^2(\mu_p^{(q)})} \\ &= (D_{q,\beta} + Z_q + \widetilde{\alpha}_{N_q}) (S_{\mu^{(q)}}C_n^{(q)})(z). \end{split}$$

Moreover we obtain

Corollary 4.4. The operators  $Z_q, D_q, \widetilde{N}_q, \widetilde{\alpha}_{N_q}$  have the following properties:

(1) 
$$\widetilde{\alpha}_{N_q} = \frac{1}{\beta} Z_q D_{q,\beta} + \beta I = \widetilde{N}_q + \beta I$$

(2) 
$$Z_q + D_{q,\beta} + \widetilde{\alpha}_{N_q} = \left(\frac{1}{\sqrt{\beta}}Z_q + \sqrt{\beta}I\right)\left(\frac{1}{\sqrt{\beta}}D_{q,\beta} + \sqrt{\beta}I\right).$$

*Proof.* The proof is done by Equations (4.2), (4.3), (4.4), (4.5), and Lemma 4.2.

Therefore, by Theorem 4.3 and Corollary 4.4, the multiplication operator  $Q_p^{(q)}$  by x in  $L^2(\mu_p^{(q)})$  is represented as a linear combination of the q-creation, q-annihilation, q-number, and scalar operators in the Hardy space  $\mathcal{H}^2(\nu_q)$ . In the q-deformed Gaussian case, due to Equation (A.4) in Appendix A, the multiplication operator  $Q_g^{(q)}$  in  $L^2(\mu_g^{(q)})$  is represented as a linear combination of q-creation and q-annihilation operators in the "same" Hardy space  $\mathcal{H}^2(\nu_q)$ .

Remark. (1) The analogous results in this paper to q=1 have been considered by Asai, et al. [6]. (2) In general, if variances of two given measures  $\mu_1$  and  $\mu_2$  are the same, then the actions of creation and annihilation operators are the same. In addition, the transformed function spaces by  $S_{\mu_1}$  and  $S_{\mu_2}$  are also the same. However, if one of the measure  $\mu_1$  is non-symmetric and the other  $\mu_2$  is symmetric, then the representation of multiplication in  $L^2(\mu_1)$  is different from that in  $L^2(\mu_2)$ .

# Acknowledgments

The author expresses sincere thanks to Professors T. Hida and K. Saitô for their kind arrangement of the conference. He is grateful for research support by International Institute of Advanced Studies, Kyoto, Japan.

# A Appendix. On q-Hermite Polynomials

First of all, we refer to the papers [12, 22] and references cited therein for the detailed description of q-Hermite polynomials. For q = 1, see books [14, 21, 27].

Let  $\mu_g^{(q)}$  be the q-deformed Gaussian measure with mean zero and variance  $\beta > 0$ . It is well-known that an associated orthogonal polynomial to  $\mu_g^{(q)}$  is the q-Hermite polynomial  $\{H_n^{(q)}(x)\}$  with the Szegö-Jacobi parameters  $\alpha_n = 0, \omega_n = \beta[n]_q$ . In this case the following relations hold for each  $n \geq 1$ :

$$xH_n^{(q)}(x) = H_{n+1}^{(q)}(x) + \beta[n]_q H_{n-1}^{(q)}(x)$$
(A.1)

where  $H_0^{(q)}=1$  and  $H_1^{(q)}=x$ . Then for any  $f(x)=\sum a_n H_n^{(q)}(x)$  and  $g(x)=\sum b_n H_n^{(q)}(x)$  in  $L^2(\mu_g^{(q)})$ , the inner product  $\langle\cdot,\cdot\rangle_{L^2(\mu_g^{(q)})}$  is given by the form

$$\langle f, g \rangle_{L^2(\mu_g^{(q)})} = \sum_n \beta^n [n]_q! \overline{a}_n b_n.$$

Moreover, let  $||f||_{L^{2}(\mu_{\alpha}^{(q)})}^{2} = \sum_{n=0}^{\infty} \beta^{n}[n]_{q}! |a_{n}|^{2}$ .

A q-deformed coherent state vector  $E_{\lambda,q}^{(q)}(x,z)$  for  $\{H_n^{(q)}(x)\}$  is defined by

$$E_{\lambda,g}^{(q)}(x,z) = \sum_{n=0}^{\infty} \frac{H_n^{(q)}(x)}{\beta^n n!} z^n, \quad z \in \Omega_q^{\beta}.$$
 (A.2)

It can be shown that the set  $\{E_{\lambda,g}^{(q)}(x,z):z\in\Omega_q^\beta\}$  is linearly independent and total in  $L^2(\mu_g^{(q)})$ . The q-analogue of the Segal-Bargmann transform  $S_{\mu_g^{(q)}}$  associated to  $\{H_n^{(q)}(x)\}$  is given by

$$(S_{\mu_g^{(q)}}f)(z) = \langle E_{\lambda,g}^{(q)}(x,\overline{z}), f(x) \rangle_{L^2(\mu_g^{(q)})} \text{ for any } f \in L^2(\mu_g^{(q)}).$$
 (A.3)

With this transform, we can reproduce the same results as for  $q \in [0,1)$  Theorem III.4 in Leeuwen-Maassen [22] and for q=0 Proposition 1 in Biane [9]. That is,  $S_{\mu_g^{(q)}}$ -transform yields a unitary isomorphism between  $L^2(\mu_g^{(q)})$  and  $\mathcal{H}^2(\nu_q)$  and

$$S_{\mu_{\alpha}^{(q)}}Q_g^{(q)} = (D_{q,\beta} + Z_q)S_{\mu_{\alpha}^{(q)}}$$
 (A.4)

where  $Q_g^{(q)}$  is the multiplication operator by x in  $L^2(\mu_g^{(q)})$ .

### References

- [1] L. Accardi and M. Bożejko, *Interacting Fock space and Gaussianization of probability measures*. Infinite Dimensional Analysis, Quantum Probability and Related Topics, **1** (1998), 663–670.
- [2] G. E. Andrew, R. Askey and R. Roy, Special Functions. Cambridge, 1999.
- [3] N. Asai: Analytic characterization of one-mode interacting Fock space. Preprint (2001)
- [4] N. Asai, I. Kubo, and H.-H. Kuo, Roles of Log-concavity, log-convexity, and growth order in white noise analysis. to appear in Infinite Dimensional Analysis, Quantum Probability and Related Topics, 4 (2001). Universidade da Madeira CCM Preprint 37 (1999)
- [5] N. Asai, I. Kubo, and H.-H. Kuo, General characterization theorems and intrinsic topologies in white noise analysis. to appear in Hiroshima Math. J., 31 (2001). Louisiana State University Preprint (1999)
- [6] N. Asai, I. Kubo, and H.-H. Kuo, One-mode interacting Fock spaces associated to Gaussian and Poisson measures. In Preparation (2001).
- [7] V. Bargmann, On a Hilbert space of analytic functions and an associated integral transform, I. Comm. Pure Appl. Math., 14 (1961), 187–214.
- [8] V. Bargmann, On a Hilbert space of analytic functions and an associated integral transform, II. Comm. Pure Appl. Math., 20 (1967), 1–101.

- [9] P. Biane, Segal-Bargmann transform, functional calculus on matrix spaces and the theory of semi-circular and circular systems. J. Funct. Anal., 144 (1997), 232–286.
- [10] M. Bożejko and R. Speicher, An example of a generalized Brownian motion I. Comm. Math. Phys., 137 (1991), 519–531.
- [11] M. Bożejko and R. Speicher, An example of a generalized Brownian motion II. in Quantum Probability and Related Topics VII, L. Accardi (ed.) World Scientific, 1992, pp. 67–77.
- [12] M. Bożejko, B. Kümmerer and R. Speicher, q-Gaussian processes: Noncommutative and classical aspects. Comm. Math. Phys., 185 (1997), 129– 154.
- [13] W. G. Cochran, H.-H. Kuo, and A. Sengupta, A new class of white noise generalized functions. Infinite Dimensional Analysis, Quantum Probability and Related Topics, 1 (1998), 43–67.
- [14] T. S. Chihara, An Introduction to Orthogonal Polynomials. Gordon and Breach, 1978.
- [15] T. Dwyer, Partial differential equations in Fischer-Fock spaces for Hilbert-Schmidt holomorphy type. Bull. Amer. Math. Soc. 77 (1971), 725–730
- [16] L. Gross P. and Malliavin, Hall's transform and the Segal-Bargmann map. in: Itô's Stochastic Calculus and Probability Theory, M. Fukushima et al. (eds.) Springer-Verlag, 1996, pp. 73–116.
- [17] R. L. Hudson and K. R. Parthasarathy, Quantum Ito's formula and stochastic evolutions. Comm. Math. Phys., 93 (1984), 301–323.
- [18] Y. Ito and I. Kubo, Calculus on Gaussian and Poisson white noises. Nagoya Math. J., 111 (1988), 41–84.
- [19] I. Kubo and H.-H. Kuo, Finite dimensional Hida distributions. J. Funct. Anal., 128 (1995), 1–47.
- [20] Kubo, I. and Takenaka, S.: Calculus on Gaussian white noise I, II, III, IV; Proc. Japan Acad. 56A (1980) 376–380, 56A (1980) 411–416, 57A (1981) 433–437, 58A (1982) 186–189.
- [21] H.-H. Kuo, White Noise Distribution Theory. CRC Press, 1996.
- [22] H. van Leeuwen and H. Maassen, A q deformation of the Gauss distribution. J. Math. Phys., 36 (1995), 4743–4756.
- [23] N. Saitoh and H. Yoshida, The q-deformed Poisson random variables on the q-Fock space. J. Math. Phys., 41 (2000), 5767–5772.
- [24] N. Saitoh and H. Yoshida, The q-deformed Poisson distribution based on orthogonal polynomials. J. Phys. A: Gen., 33 (2000), 1435–1444.
- [25] I. E. Segal, Mathematical characterization of the physical vacuum for a linear Bose-Einstein field. Illinois J. Math., 6 (1962), 500–523.

- [26] I. E. Segal, The complex wave representation of the free Boson field. in: Essays Dedicated to M. G. Krein on the Occassion of His 70th Birthday, Advances in Math.: Supplementary Studies Vol.3, I. Goldberg and M. Kac (eds.) Academic, 1978, pp. 321–344.
- [27] M. Szegö, Orthogonal Polynomials. Coll. Publ. 23, Amer. Math. Soc., 1975.
- [28] D. Voiculescu, K. J. Dykema and A. Nica, *Free Rondom Variables*. CRM Monograph Ser. 1, Amer. Math. Soc., 1992.