## A RATIONAL NONCOMMUTATIVE INVARIANT OF BOUNDARY LINKS

STAVROS GAROUFALIDIS AND ANDREW KRICKER

A bstract. We construct an invariant of boundary links which is at least as strong as the K ontsevich integral, determ ines the S-equivalence class of a boundary link and which takes values in a space of trivalent graphs whose edges are decorated by rational functions in noncom muting variables. Our invariant is axiom atically characterized by a universal property closely related to the H om ology Surgery view of boundary links, and comes equipped with a diagram matic integration theory, in the spirit of perturbative quantum eld theory.

#### 1. Introduction

1.1. H istory. It is well-known that the problem of H om obgy Surgery and its application to the classi cation of boundary links, up to cobordism, is closely related to the set of H em it ian matrices over Z F ] (the group-ring of the free group), invertible over Z. C appell-Shaneson formulated the problem of boundary link cobordism (in high dimensions) in terms of -groups (see [CS1, CS2]) and Vogel identied these -groups in terms of surgery L-groups of a suitable localization (the Cohn localization) of Z F ], [Vo]. Later on, Vogel and Farber identied the Cohn localization of Z F ] with the ring of rational functions in noncommuting variables, FV]. In addition, Farber constructed an invariant of boundary links with values in that ring, Fa].

In a completely di erent direction, K ontævich de ned a graph-valued invariant of links (in particular, boundary links) that is closely related to perturbative quantum eld theory, [K o].

12. A brief sum m ary. The paper introduces a graph-valued invariant Z<sup>rat</sup> (pronounced: the rational Z -invariant) of boundary links (that is, links whose components bound pairw ise disjoint in bedded surfaces) with the following features:

 $Z^{rat}$  is de ned for boundary links (and their cousins, F -links) L in integral hom ology 3-spheres M. It takes values in a space of trivalent graphs, whose edges are decorated by rational functions in noncommuting variables ft<sub>1</sub>;:::;t<sub>g</sub>g, see Theorem 1.3. For good boundary links, i.e., for those whose free cover is acyclic, the edges of the graphs are decorated by polynom ials in noncommuting variables, see Remark 1.6. A word of caution: the word \rational" in the name of Z<sup>rat</sup> and in the title of the paper refers to rational functions and not to the edd of rational numbers.

The degree 0 part of  $Z^{rat}$  is a matrix-valued invariant that determ ines the B lanch eld pairing of (M; L)(thus also all classical concordance invariants of boundary links), and the analytic invariant of Farber [Fa] and the of F -links of [G L2], see P roposition 12.

Z<sup>rat</sup> determ ines the LM O -A arhus extension of the K ontsevich integral, see Theorem 1.3.

 $Z^{rat}$  com es equipped with a diagram m atic integration theory  $n^{rat}$ , in the spirit of perturbative quantum eld theory, see Sections 3-5.

 $Z^{rat}$  can be axiom atically characterized by a universal property, namely that it is the universal nite type invariant of F-links with respect to the loop ltration, after Cohn-localization, see Theorem 1.12. For knots, the existence of  $Z^{rat}$  gives a new proof the M M R Conjecture, originally proven in [BG] as well as a strong form of Rozansky's Rationality Conjecture for the Kontsevich integral, [R], previously unknown, see Remark 1.8.

Date: This edition: A pril 18, 2001 First edition: A pril 18, 2001.

S.G. was partially supported by an NSF grant D.M.S-98-00703 and by an Israel-U.S.B.SF grant. A.K. was partially supported by a JSPS fellow ship.

This and related preprints can also be obtained at http://www.math.gatech.edu/ stavros .

<sup>1991</sup> M athem atics C lassi cation. P rim ary 57N 10. Secondary 57M 25.

K ey words and phrases: Boundary links, K ontsevich integral, C ohn localization.

An application of the invariant  $Z^{rat}$  is a formula for the LM O invariant of cyclic branched covers of knots K in integral hom ology 3-spheres M in terms of residues of  $Z^{rat}$  and the signature function of (M; K), see [GK].

A nother application of the degree 2 part of  $Z^{rat}$  and its properties under surgery is a result of Teichner and the rst author on knots with trivial A lexander polynom ial, [G T].

Question: W hy consider boundary links and not all links, or at least all links with linking numbers zero? Answer: There are two reasons. The Z<sup>rat</sup>-invariant is closely related to the surgery view of boundary links (and their cousins, the F -links), see Section 1.4. At present a surgery view of links with linking number zero is unknown. An additional reason is that the comparison between the Z<sup>rat</sup> and Z invariants requires links with vanishing -invariants, see Remark 7.2. Examples of such links are the boundary links.

A re nem ent of boundary links (abbreviated (e-links) are F-links, i.e., a triple (M; L; ) of a link L in an integral hom ology 3-sphere M and an onto map  $: _1 (M L) F$ , where F is the free group on a set T (in 1-1 correspondence with the components of L) such that the ith meridian is sent to the ith generator. Two F-links (M; L; ) and  $(M^0; L^0; {}^0)$  are equivalent i (M; L) and  $(M^0; L^0)$  are isotopic and the maps and  ${}^0$  di er by an inner autom orphism of F. The underlying link of an F-link is a (e-link).

K nots K in integral hom ology 3-spheres M are examples of (-links) (in fact, F -links) with one component whose canonical F-structure is given by the map  $_1(M \times ) ! H_1(M \times ) = Z$ .

1.3. Statem ent of the results. As a warm-up, recall that the A lexander polynom ial of a knot m easures the order of the rst hom ology of the free abelian cover of its complement. Given an F-link (M;L; ), we can consider the free cover X<sup>!</sup> of its complement, and its hom ology H<sub>1</sub> (X<sup>!</sup>), which is a module over the group-ring = Z F] of the free group. Since the ring is not Noetherian (when L is not a knot), the notion of a torsion module does not make sense. However, in [GL2], Levine and the rst author de ned an invariant of F-links with values in  $^{-}$  (cyclic), the group of form alpower series (with rational coe cients) in noncommutative variables, module cyclic permutation. The invariant is a reform ulation of an invariant de ned earlier by Farber, Fa], and in the case of knots equals, up to a multiple, to the logarithm of the A lexander polynom ial. Thus, can be thought as a noncommutative torsion invariant of F-links.

Our not goal is to de ne a matrix-valued generalization of using the surgery view of F-links. The surgery view considers the set N (O) of nullhom otopic links L with Z-invertible linking matrix in the complement of a standard trivial unlink O in S<sup>3</sup>. It turns out that surgery on L transforms (S<sup>3</sup>;O) to an an F-link. A lterning L by a kirby move (i.e., by adding a parallel of a link component to another, or adding/rem oving a unit-fram ed unlink away from L) preserves surgery and does not change the corresponding F-link. Let N (O)=h i denote the set of equivalence classes of N (O) under the K irby equivalence relation. The next theorem explains what we mean by the surgery view of F-links.

Theorem 1.1. (Proof in Sections 2.1-2.2) The surgery map gives a 1-1 onto correspondence

N (O)=h i ! F links:

The equivariant linking matrix of L 2 N (O) is a Herm itian matrix (that is, a matrix A over is that satis es A = A<sup>?</sup> where A<sup>?</sup> is the transpose of A followed by the natural involution of ), de ned up to an indeterm inacy that is a relation of the K inby moves; compare also [G L1]. Indeed, recall that two Herm itian matrices A; B are simply stably congruent i A  $S_1 = P(B - S_2)P^2$  for some diagonal matrices  $S_1; S_2$  with

1 entries and some elementary matrix P, i.e., one which is either invertible over or a diagonal matrix with entries in . There is a natural map of rings ! Z that sends F to 1. Let B (! Z) denote the set of simple stable congruence classes of H erm itian matrices over , invertible over Z<sup>1</sup>. The K index equivalence class of the equivariant linking matrix of L as above lies in B (! Z).

Let be the  $\hat{}=(cyclic)$ -valued invariant on the set Herm (! Z) of Herm itian matrices W over , nonsingular over Z, de ned by

$$(W) = \operatorname{trlog}^{0}(W) := \frac{X^{2}}{n} \frac{(1)^{n}}{n} \operatorname{tr} (W (W)^{1} I)^{n}$$

where : ! Z.W ith the above notation we have the following

 $<sup>^{1}</sup>$ By invertible over Z, we mean that the image of a matrix which is dened over under the map ? Z is invertible over Z.

P roposition 1.2. (Proof in Section 2.3) (a) There exist an invariant W of F-links with values in B (! Z). (b)  $H_1(X^{!})$  is a -m odule with presentation W. (c) The Blanch eld pairing [Dv]

H<sub>1</sub>(X<sup>!</sup>;Z) H<sub>1</sub>(X<sup>!</sup>;Z)! <sub>loc</sub>=

is given by W  $^{1}$  mod , where  $_{loc}$  is the Cohn localization of , FV]. (d) = .

We now introduce a graph-valued deform ation of the matrix-valued invariant W of F - and @-links. Recall that the Kontsevich integral (or rather, its LMO-A arhus extension) Z (M;L) of a link L in an integral hom ology 3-sphere M takes values in in a completed vector space A ( $_{\rm H}$ ) over Q of diagram swhose skeleton is a disjoint union of solid circles [ $_{\rm i}$   $_{\rm h_i}$ , one circle labeled by a variable  $h_{\rm i}$  for each component of L, [LMO, A]. The Z<sup>rat</sup> invariant takes values in a set

$$A^{\circ}(_{loc}) = B(_{loc} ! Z) A(_{loc})$$

where A ( $_{\rm bc}$ ) is the completed vector space of vertex and edge oriented, trivalent diagram s whose edges are decorated by elements of the Cohn localization  $_{\rm bc}$  of , [BR], or equivalently, by the ring of rational functions in noncommuting variables, [FV]. The diagram s in A ( $_{\rm bc}$ ) are considered modulo som e algebraic analogs of geom etric relations on links, explained in detail in Section 3), nam ely

the well-known AS and IHX relations,

relations on the labels of the edges of diagrams (that rst appeared in [GL1]), see Figures 1 and 2. There are some additional! and CA relations on  $A^{0}(_{bc})$ , (see Sections 3 and 5.4). Moreover, there is a map Hair :  $A^{0}(_{bc})=h!$ ; CAi! A (<sub>H</sub>) that replaces a generator  $t_{i}$  of the free group F by an exponential of i-colored legs, labeled by  $h_{i}$ , explained in Section 7.1 and in Figure 4.

Question: W hy consider the Cohn localization loc of ?

Answer: For two independent reasons, see Remark 5.2.

Theorem 1.3. (Proof in Sections 5.5, 6.5, 7.4) There exist invariants Z<sup>rat</sup> of F-links and Q-links that t in a commutative diagram

W e will call Z<sup>rat</sup> the rational Z -invariant of Q-links. The above theorem states that Z<sup>rat</sup> is a lift of the LM O -A arhus-K ontsevich invariant of Q-links.

Som e com m ents m ay help the reader fam iliarize with the rational invariant Z  $^{\rm rat}$  :

Rem ark 1.4. First a rem ark about the empty set. For the unlink O of g components, we have that Z<sup>rat</sup> (O) = (empty matrix; 1) 2 A<sup>0</sup> ( $_{bc}$ )=h!; CAi. On the other hand, the K ontsevich intregalofO is given by Z (O) = (h<sub>1</sub>)t t<sub>g</sub>)(f(where (h<sub>i</sub>) is the K ontsevich integral of the unknot, whose diagram s are colored by h<sub>1</sub>), and

Hair  $(em pty m atrix; 1) = (h_1) t t_g)$ 

This is how Z  $^{\rm rat}$  (O) determ ines Z (O), as stated in the above theorem .

Rem ark 1.5. In case L = ; is the empty link, then Z<sup>rat</sup> (M;;) = (empty matrix; Z (M;;)). Using the map ': A ( $_{\rm loc}$ ) ! A (Q) = A () induced by the canonical ring hom on orphism :  $_{\rm loc}$  ! Q, it follows that ' Z<sup>rat</sup> (M;L) = Z<sup>rat</sup> (M;;). In other words, evaluating the rational invariant at  $t_i = 1$  has the e ect of rem oving the components of the  $\ell$ -link.

Rem ark 1.6. The Z<sup>rat</sup> invariant of F-links takes values in a rather strange quotient A<sup>0</sup> ( $_{loc}$ )=h! i of a set of pairs of matrices and vectors A<sup>0</sup>. If we restrict Z<sup>rat</sup> j to a class S of F-links with the same B lanch eld pairing (or the same S-equivalence class), then Z<sup>rat</sup> j is equivalent to an invariant

that takes values in a graded vector space, where the degree of a graph is the number of trivalent vertices. In addition, the edges of the trivalent graphs are decorated by elements of a subring of  $_{\rm loc}$  that depends on S. In particular, for good boundary links (i.e., for links whose free cover is acyclic, or equivalently, whose B lanch eld pairing is trivial) Z<sup>rat</sup> is equivalent to an invariant with values in the more manageable graded vector space A ()=(!;CA); see Section 5.5.

Rem ark 1.7. For g = 1, the sets of 1-component F -links and Q-links coincide with the set of knots in integral hom ology 3-spheres, and the relations ! and CA<sub>1</sub> are trivial, see Lemmas 3.8 and 6.7. In this case, Z<sup>rat</sup> takes values in B ( $_{bc}$ ! Z) A ( $_{bc}$ ), and the matrix-free part of Z<sup>rat</sup> takes values in the graded algebra A ( $_{bc}$ ), where the multiplication in A ( $_{bc}$ ) is given by the disjoint union of graphs. It turns out that the matrix-free part of Z<sup>rat</sup> is a group-like element in A ( $_{bc}$ ), i.e., it equals to the exponential of a series of connected graphs.

For g = 2 (i.e., for two component  $\ell$ -links) the relation  $CA_g$  is trivial, see Lemma 6.7 and the set of 2-component F-links coincides with the one of  $\ell$ -links.

Remark 1.8. How unique is the lift  $Z^{rat}$  of Z? A weak form of the RC Conjecture for e-links states that the image of Z lies in the image of the map Hair :  $A^0$  ( $_{loc}$ )=h!; CAi! A ( $_H$ ). A strong form of the RC Conjecture for knots states that the Kontsevich integral lifts. The map Hair is not 1-1 even for knots, since in that case  $A^0$  ( $_{loc}$ )=h!; CAi = B ( $_{loc}$ ! Z) A ( $_{loc}$ ) and the map Hair restricted on B ( $_{loc}$ ! Z) depends on the determ inant of the stable equivalence class of a Herm itian matrix and its size. The weak form of RC Conjecture for knots was form ulated by Rozansky in [R] and was proven by the second nam ed author in [K r1]. The strong form of the RC Conjecture for F-links was form ulated by the rst nam ed author in [G a] and is proven in Theorem 1.3 above.

Question 1.9. In the case of knots, is the map Hair : A  $(_{\rm loc})$  ! A  $(_{\rm H})$  1-1?

Remark 1.10. In a sense, Z is a graph-valued  $-invariant of @-links, where <math>^{-}$  is the completion of Q with respect to the augmentation ideal whereas Z<sup>rat</sup> is a graph-valued  $_{loc}$ -invariant. It often happens that  $^{-}$ -invariants can be lifted to  $_{loc}$ -invariants, see for example the discussion of Farber-Ranicki, [FRa], on a problem of N ovikov about circle-valued M orse theory. It would be nice if an extension of this circle-valued M orse theory to in nite dimensions (using the circle-valued Chem-Sim ons action on the space of gauge equivalence classes of connections on a principal bundle over a 3-m anifold) would lead to an explanation for the existence of the Z <sup>rat</sup>-invariant.

Rem ark 1.11. As will become apparent from the construction of the Z<sup>rat</sup> invariant, it is closely related to equivariant linking matrices of nullhom otopic links in the complement of an F-link, see Proposition 8.2. It is an interesting question to ask whether the Z<sup>rat</sup> invariant of an F-link can be dened in terms of an equivariant construction on the free cover of the F-link.

The next theorem characterizes Z<sup>rat</sup> by a universal property. Using the term inology of [G a], recall that a bop move on an F-link (M;L;) can be described by surgery on a clover G in M L whose leaves lie in the kerner of , [G a]. The result (M;L;)<sub>G</sub> of surgery is another F-link. As usual, one can de ne a notion of nite type invariants on the set of F-links using bop m oves. The num ber of invariants of degree n can be identified with a quotient vector space  $G_n^{\text{loop}}$ . In [G a], it was shown that  $G_n^{\text{loop}}$  is a module over n, and that after localization, (i.e., after tensoring with bc),  $G_{n;bc}^{\text{loop}} = 0$  for odd n and is a quotient of  $A_n^0$  (bc)=h! i for even n. The next theorem constructs enough nite type invariants to characterize Z<sup>rat</sup> by a universal property.

Theorem 1.12. (Proof in Section 8) (a) The matrix-part of Z<sup>rat</sup> is at least as strong as a universal nite type invariant.

(b) The matrix-free part of Z<sup>rat</sup> is the universal A ( $_{loc}$ )=(!)-valued nite type invariant of F -links with respect to the loop limit on. In particular, we have that A<sub>2n</sub> ( $_{loc}$ )=(!) =  $G_{2n;loc}^{loop}$  for n 1.

1.4. The main ideas of the proof. The three main ideas of the paper are sum marized as follows:



The invariant Z<sup>rat</sup> is closely related to the surgery view of @-links and their cousins, the F-links.

Section 2, which consists entirely of classical geom etric topology, identi es the set of F -links (resp. (e-links) with the quotient (i.e., the set of equivalence classes) N (O)=h i of N (O) under a suitable K irby-type (resp. and in addition, a  $CA_g$ -type) equivalence relation . Using this, and arguments of surgery theory, we give a proof of Proposition 1.2. The rest of the section identies the set N (O) it terms of a set of tangles D (D<sub>g</sub>)=hr; ;! i of N (O), namely a a set of sliced crossed links in a solid handlebody, m odulo certain equivalence relations that include regular isotopy, r basing, and wrapping !.

The road to de ne the invariant Z<sup>rat</sup> that appears in Theorem 1.3 is long, and given in Sections 3-7.

Section 3 presents the various sets of diagram s and their relations, where ultim ately all the invariants will take value. The algebraic relations on diagram s are motivated by the geometric relations of Section 2.

In Section 4 we will de ne an invariant D (D<sub>g</sub>) ! A <sup>gp</sup> (?<sub>X</sub>; ) and show that it maps the various geom etric relations on the set D (D<sub>g</sub>) to the corresponding algebraic relations on diagram s, thus giving rise to a map N (O) ! A <sup>gp</sup> (~<sub>X</sub>; )=h! <sup>gp</sup> i.

Unfortunately, the map N (O) !  $A^{qp}$  (~ x; )=h!  $^{qp}$  is not invariant under the K inby moves . To accommodate for that, in Section 5 we construct an integration theory  $^{R_{rat}}$  in the spirit of the A arhus Integral [A] with the property that the above map, composed with integration (and a slight normalization), is invariant under the K inby moves. This is our desired invariant that appears in Theorem 1.3. The  $^{R_{rat}}$ -integration is a formal diagram matic analogue of G aussian Integration, expressed in terms of gluing legs of diagrams to each other and preserves the basing, wrapping and K inby equivalence relations. Some identities of this theory of integration motivate the algebraic form of the wrapping relations, as well as what the value of the invariant Z <sup>rat</sup> in degree 0, see Section 5.4 and Section B. Section 5.5 ends by de ning the invariants of F -links and Q-links that appear in Theorem 1.3.

The comparison between the  $Z^{rat}$  invariant and the Kontsevich integral Z of boundary links is done in Section 7, using a magic form ula for the Kontsevich integral of the Long HopfLink, conjectured in [A 0] (in conjunction with the W heels and W heeling Conjectures) and proven in [BLT]. Section 7.4 com pletes the proof of Theorem 1.3.

Section 8 is devoted to the proof of Theorem 1.12, namely the universal property of  $Z^{rat}$ . This follows from the general principle of locality of the invariant Z<sup>rat</sup> (i.e., with the behavior of Z<sup>rat</sup> under sublinks) together with the identi cation of the covariance of Z<sup>rat</sup> with an equivariant linking function.

The paper ends with som e appendices on diagram m atic integration and its various identities.

1.5. A cknow ledgem ent. The results of this work were rst announced by the rst named author in a meeting on \In uence of Physics on Topology" on August 2000, in the sunny San Diego. We thank the organizers, M.Freedm an and P.Teichner for arranging such a fertile meeting. We also wish to express our gratitude to J.Levine who shared with the rst author his expertise on topology, and to J.Hillm an and T. O htsuki for encouragem ent and support of the second author.

Contents

Introduction
 H istory

12. A brief sum mary	1
13. Statem ent of the results	2
1.4. The main ideas of the proof	5
1.5. A cknow ledgem ent	5
2. A surgery view of F-links	6
2.1. A surgery description of F-links	6
22. A relative version of Fenn-Rourke's theorem	7
2.3. A matrix-valued invariant of F-links	8
2.4. A tangle description of F-links and Q-links	10
3. An assortm ent of diagram s and their relations	12
3.1. A review of the Cohn localization	14
32. The in nitesimal wrapping relations	15
3.3. Group-like relations	15
4. An invariant of links in the complement of an unlink	17
5. A rational version of the Aarhus integral	20
5.1. The de nition of Trat	20
52. respects the basing relations	21
5.3. Dealing with the Kirby moves	22
5.4. Dealing with the wrapping relations	24
5.5. The de nition of the rational invariant Z <sup>rat</sup> of F-links	24
6. The CA action and invariants of boundary links	25
6.1. A surgery description of boundary links	25
62. The action of CA on diagram s	27
6.3. $Z_{rat}^{rat}$ commutes with the CA action	29
6.4. $\operatorname{com} m$ utes with the CA <sub>g</sub> action	29
6.5. The de nition of the rational invariant Z <sup>rat</sup> of Q-links	30
7. A comparison between Z and Z <sup>rat</sup>	31
7.1. The Hairmap	31
7.2. A comparison between $Z_{rat}^{rat}$ and $Z_{rat}$	32
7.3. A comparison between <sup>Trat</sup> and	33
7.4. End of the proof of Theorem 1.3	33
8. A universal property of Z <sup>rat</sup>	34
Appendix A. A brief review of diagram matic calculus	36
A.1. Som e useful identities	36
A 2. Iterated Integration	37
A.3. Integration by parts	38
Appendix B. The W heels Identity	40
Appendix C. In nitesimal versus group-like w rapping relations	41
Appendix D. In nitesimal basing relations	45
Appendix E. A useful calculation	46
References	47

## 2. A surgery view of F-links

2.1. A surgery description of F-links. The purpose of this section is to give a link-surgery description of the set of F-links and @-links. This will be achieved by adapting classical surgery theory arguments to the case of 3-m anifolds with boundary. This section consists entirely of classical geometric topology.

Before we proceed, a rem ark on notation:

Remark 2.1. Throughout the paper, given an equivalence relation on a set X, we will denote by X = h is the set of equivalence classes. Occasionally, an equivalence relation on X will be defined by one of the following ways:

E ither by the action of a group G on X , in which case X =h i coincides with the set of orbits of G on X .

Or by a m ove on X, i.e., by a subset of X = X, in which case the equivalence relation is the sm allest one that contains the m ove.

Proof.(of Theorem 1.1) F is once and for all a based unlink 0 of g components in  $S^3$ . Let N (0) denote the set of nullhom otopic links L with Z-invertible linking matrix in the complement of 0.

Surgery on an element C of N (O) transform s (S<sup>3</sup>;O) to a F -link (M ;L). Indeed, since C is nullhom otopic the natural map  $_1$  (S<sup>3</sup> O)! F gives rise to a map  $_1$  (M L)! F. A lternatively, one can construct disjoint Seifert surfaces for each component of L by tubing the disjoint disks that O bounds, which is possible, since each component of C is nullhom otopic. Since the linking matrix of C is invertible over Z, M is an integral hom ology 3-sphere. Let denote the equivalence relation on N (O) generated by the moves of handle-slide  $_1$  (i.e., adding a parallel of a link component to another component) and stabilization  $_2$  (i.e., adding to a

link an unknot away from the link with fram ing 1). It is well-known that -equivalence preserves surgery, thus we get a surgery map

Now, we will show that the surgery map is onto. Given a F -link (M ;L) choose disjoint Seifert surfaces for its components and choose a fram ed link C<sup>0</sup> in M such that  $M_c = S^3$ . C might intersect , how ever by a small isotopy (which does not change the property that  $M_c = S^3$ ) we may assume that C is disjoint from . Consider the image <sup>0</sup> of in S<sup>3</sup> = M<sub>c</sub>. By doing crossing changes among the bands of <sup>0</sup>, translated in terms of surgery on a unit-fram ed link C S<sup>3</sup> <sup>0</sup>, we can assume that each component of <sup>0</sup> lies in a ball disjoint from the other components and further that the boundary of <sup>0</sup> is an unlink. The link C [ C<sup>0</sup> is nullhom otopic in the complement of the unlink (<sup>0</sup> since it is disjoint from <sup>0</sup> and the linking matrix of C [ C<sup>0</sup> is invertible over Z since S<sup>3</sup><sub>c [ C<sup>0</sup></sub> = M , an integral hom ology 3-sphere. By moving the basing around, if necessary, we can assume that the basing of (<sup>0</sup> is standard. By construction, C [ C<sup>0</sup> 2 N (O ) maps to (M ;L).

Next, we need to show that the surgery map is 1-1; we will do this in the next section.

2.2. A relative version of Fenn-Rourke's theorem . The goal of this independent section is to show a relative version of Fenn-Rourke's theorem [FR, Theorem 6]. We begin with some notation. Given a 3-m anifold M (possibly with nonempty boundary) and a framed link L in its interior, surgery produces a 3-m anifold M  $_{\rm L}$  whose boundary is canonically identied with that of M and a cobordism W  $_{\rm L}$  with boundary M [ (@M I) [ M  $_{\rm L}$ .

Theorem 2.2. If M is a 3-m anifold with boundary,  $L_1$ ,  $L_2$  two framed links in its interior, then  $L_1$  and  $L_2$  are -equivalent i there exist

- (a)  $h: M_{L_1} ! M_{L_2}$ , a di eo rel. boundary.
- (b) An isomorphism  $:_1 (W_{L_1}) ! _1 (W_{L_2}).$
- (c) A commutative diagram ()



such that () =  $0.2 \text{ H}_4$ ().

Here, () 2 H<sub>4</sub>() is the hom ology class of (W), where :W ! K (;1), W = W<sub>L1</sub> [ $_{0}$  W<sub>L2</sub> (where the boundaries are identied via the identity on M, @M I and h on the top piece) and =  $_{1}$  (W<sub>L1</sub>) =  $_{1}$  (W<sub>L2</sub>). Note that W is a closed 4-m anifold.

The proof of Theorem 2.2 follows mainly from Fenn-Rourke [FR], with some simplications due to Roberts, [Rb]. We begin with the following lemma from [FR], which reformulates the vanishing of the condition () = 0.

Lem m a 2.3. [FR, Lem m a 9] Assume that we are in the situation of Theorem 2.2 and that there exists h; making commute. Then () = 0 i W #  $_{r}CP^{2}$  #  $_{s}CP^{2}$  = @ <sup>5</sup> for some r; s and closed 5-m anifold such that the diagram



commutes and  $(j_1)$ ;  $(j_2)$  are split in jections.

Proof. (of Theorem 2.2) If  $L_1$  and  $L_2$  are -equivalent, then it follows from the \only if" statement of Lemma 2.3 that () = 0 since the induced homeomorphisms on  $_1$  do not change.

So, assume that the algebraic conditions of Theorem 2.2 are satised for  $L_1$ ,  $L_2$ . We will show that  $L_1$  is -equivalent to  $L_2$ .

The rst step is to arrange via -m oves that  $W_{L_1}$  and  $W_{L_2}$  dieom orphic. The proof of [FR, p.9] (where they assume @M = ) using surgery works here, too.

The next step is to use Cerf theory. Namely, consider M orse functions  $f_i$  realizing the cobordism s W  $_{L_1}$  for i = 1;2. Since W  $_{L_1}$  is di com orphic to W  $_{L_2}$ ,  $f_1$  and  $f_2$  are related by a one parameter family of sm ooth functions  $f_t$  for 1 t 2. The arguments of Roberts Rb, Stage 2, p.3] in this case imply that  $L_1$  is -equivalent to  $L_2$ .

Remark 2.4. If L is a nullhom otopic link in M, then W<sub>L</sub> is hom otopic to a one point union of M and some number of 2-spheres S<sup>2</sup>. In particular,  $_1 (M)'_1 (W_L)$ .

End of the Proof of Theorem 1.1. We will apply Theorem 2.2 to  $M = S^3 = 0$ . Consider (L<sub>i</sub>; <sub>i</sub>) two surgery presentations of the same F -link for i = 1; 2. Then, we have a hom otopy commutative diagram



where  $_i: M_{L_i}! M$  are splitting maps. It follows by Remark 2.4 that a map exists such that the following diagram () commutes



The obstruction () lies in H<sub>4</sub>(F). However, K(F;1) =  $\_S^1$ , thus H<sub>4</sub>(F) = H<sub>4</sub>( $\_S^1$ ) = 0. Thus, Theorem 2.2 in plies that L<sub>1</sub> is -equivalent to L<sub>2</sub>.

2.3. A m atrix-valued invariant of F-links. The goal of this section is to prove P roposition 1.2. We begin by dening them ap W. G iven a nullhom otopic link L in the complement of O, consider its equivariant linking matrix, i.e., the equivariant linking numbers of the lifts of L to the universal cover of S<sup>3</sup> O. The equivariant linking matrix is Hermitian and well-dened modulo congruence by diagonal matrices with elements in . If L is in N (O), then its equivariant linking matrix is invertible over Z, and if we modify L by K inby moves, the class of its equivariant linking matrix in B (! Z) remains unchanged. Thus, we have well-dened map

which, combined with Theorem 1.1, gives rise to a map

Recall the -invariant of Section 1.3.

P roposition 2.5. (a) For Herm itian matrices A; B over  $_{loc}$ , nonsingular over Z, we have in  $\hat{}=$  (cyclic) that

(AB) = (A) + (B) and (A B) = (A) + (B):

(b) descends to a  $\hat{}=$  (cyclic)-valued invariant of the set B (  $_{loc}$  ! Z). (c) For A as above, (A) = (A A<sup>1</sup>) where : ! Z.

Proof. (b) follows from (a). For (a), let R denote the ring of square matrices with entries in  $\hat{}$ , and [R; R] denote the subgroup of it generated by matrices of the form AB BA where A; B 2 R. We claim that if A; B are Herm it is over  $_{loc}$ , with augmentations A<sub>0</sub>; B<sub>0</sub> invertible over Z, then

(1) 
$$\log^{0}(AB) = \log^{0}(A) + \log^{0}(B) \mod [R;R]$$

Indeed, if  $A_0 = B_0 = I$ , then this follows from the Baker-C am bell-H ausdor form ula, since  $\log^0(AB) = \log(AB)$ . For the general case, we have modulo [R; R], that

$$\log^{0}(AB) = \log(AB(A_{0}B_{0})^{-1}) = \log(ABB_{0}^{-1}A^{-1}AA_{0}^{-1})$$
  
= 
$$\log(ABB_{0}^{-1}A^{-1}) + \log(AA_{0}^{-1})$$
  
= 
$$A \log(BB_{0}^{-1})A^{-1} + \log(AA_{0}^{-1}) = \log^{0}(B) + \log^{0}(A):$$

Taking traces in  $\hat{}=$  (cyclic), Equation (1) in plies (a). (c) follows immediately from the denition of and the fact that (A A<sup>1</sup>) = I.

We will also denote by the ^= (cyclic)-valued invariant of F -links de ned by

Proof. (of Proposition 1.2) Consider C 2 N (O) and let (M;  $L_b$ ) the corresponding F-link. Let W denote the equivariant linking matrix of C, and X<sup>!</sup> denote the free cover of M  $L_b$  given by .

Parts (a) and (b) follow from classical surgery arguments presented in the case of knots by Levine in [Le1] and adapted without di culty in the case of F -links; see also [GL3, Lem m a 2.4].

For part (c), observe that the and the invariant of [GL2] are de ned, respectively, via the Seifert surface and the surgery view of boundary links. The two views can be related using the so-called Y-view of knots presented in [GGP, Section 6.4] and extended without di culty to the case of boundary links. Consider a disjoint union of Seifert surfaces in S<sup>3</sup> which we think of as a disjoint union of in bedded disks with pairs of bands attached in an alternating way along each disk.

Consider an additional link  $L^0$  in  $S^3$  such that its linking matrix C satis esdet (C) = 1 and such that the linking number between the cores of the bands and  $L^0$  vanishes. W ith respect to a choice of orientation of 1-cycles corresponding to the cores of the bands, a Seifert matrix of is given by

$$A = \begin{array}{cc} L^{xx} & L^{xy} + I \\ L^{yx} & L^{yy} \end{array}$$

Start from a surgery presentation of  $(S^3; )$  in terms of clovers with three leaves, as was explained in [GGP, Section 6.4] and sum marized in the following gure:



Surgery on a clover with three leaves can be described in terms of surgery on a six component link  $L^{\infty}$ . It was observed by the second author in [K r2, F igure 3.1] that  $L^{\infty}$  can be simplied via K index moves to a four

component link  $L^{00}$  shown below:



The equivariant linking matrix of (a based representative of)  $L^{00}$  [  $L^{0}$  is given by W C where W is the following matrix in  $fx_i; y_i; x_i^0; y_i^0$  basis:

$$W = \begin{bmatrix} 0 & 1 \\ B & L^{xx} & L^{xy} & I & 0 \\ B & L^{xy} & L^{yy} & 0 & I \\ 0 & I & 0 & 0 & xI^{A} \\ 0 & I & xI & 0 \end{bmatrix}$$

where x = t 1 and  $x = t^{1}$  1. Thus

WW(1)<sup>1</sup> =  $\begin{array}{cccc} L & I & 0 & I \\ I & B & I & L \end{array}$  =  $\begin{array}{cccc} I & 0 \\ B & I & BL \end{array}$ :

and (W) = (I BL).

It is easy to see that every F -link has a surgery presentation  $L^{00}$  [  $L^{0}$  as above by adapting the onto part of the proof of Theorem 1.1.

U sing the notation of [GL2], let  $Z = A (A A^0)^{-1}$ . Since  $(A A^0)^{-1} = \begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}$  it follows that

$$L + \begin{array}{ccc} 0 & I \\ 0 & 0 \end{array} = A = Z \begin{array}{ccc} 0 & I \\ I & 0 \end{array}$$
 thus  $L = Z \begin{array}{ccc} 0 & I \\ I & 0 \end{array} \begin{array}{ccc} 0 & I \\ 0 & 0 \end{array}$ :

Substituting, we obtain that

$$I BL = \begin{cases} 1 & 0 & x & 0 & z & 0 & I \\ 0 & t^{1} & 0 & x & I & 0 & 0 & t^{1} \\ 0 & t^{1} & 0 & x & I & 0 & 0 & x & I & 0 \\ \end{cases}$$
$$= \begin{cases} 0 & 1 \\ t^{1} & 0 & (I + XZ) & I & 0 & I \\ 0 & I & 0 & I & 0 & 1 \\ \end{bmatrix}$$

Proposition 2.5 in plies that  $(I F L) = \begin{pmatrix} 0 & 1 \\ t^1 & 0 \end{pmatrix} + (I + X Z) = (A) as required. \Box$ 

2.4. A tangle description of F-links and Q-links. In this section we give a tangle diagram description of F-links and Q-links which is useful in de ning invariants. We begin by giving a tangle description of the set N (O). Consider the following surfaces in  $\mathbb{R}^2$ 



together with a distinguished part of their boundary (called a gluing site) marked by an arrow in the gure above. By a tangle on a surface  $_x$  (for x = a;b;c) we mean an oriented, framed, smooth, proper immersions of an oriented 1-manifold into the surface, up to isotopies rel boundary of the surface, double points being equipped with crossing information, with the boundary points of the tangle lying on standard points at t = 0and t = 1.

A crossed tangle on a surface x is a tangle diagram on x possibly with some crosses marking those boundary points of the diagram that lie either on a t = 0 edge of x for x = a; b or on a t = 1 edge of c.

The following is an example of a crossed tangle diagram on b:



The cross notation evokes a small pair of scissors  $\searrow_{0}^{0}$  that will be sites where the skeleton of the tangle will be cut. A sliced crossed link on D<sub>g</sub> is a sequence of crossed tangles,  $fT_1$ ;:::; $T_k$ g such that

the boundaries of the surfaces m atch up, that is, their shape, distribution of endpoints, and crosses.

For example, if  $T_i$  is of the form



the top and bottom boundaries of  $T_1$  and  $T_k$  look like:



A fler stacking the tangles  $fT_i$ ;::; $T_kg$  from top to bottom, we obtain a crossed link in  $D_g$  I, where  $D_g$  is a disk with g holes. Note that the holes in  $D_g$  are ordered from the top one to the bottom one and that  $D_g$  is equipped with g gluing sites.

C learly, every crossed link in  $D_g$  I can be obtained this way. Here is an example of a sliced crossed link in  $D_1$  whose corresponding link in N (O) is a surgery presentation for the Figure 8 knot:



Let D (D  $_{g}$ ) denote the set of sliced crossed links on D  $_{g}$  such that each component of the associated link is nullhom otopic in D  $_{g}$  I and m arked with precisely one cross.

W e now introduce som e equivalence relations on D (D $_g$ ) that are in portant in comparing D (D $_g$ ) to N (O). Regular isotopy <sup>r</sup> of sliced crossed links is the equivalence relation generated by regular isotopy of individual tangles and the following m oves:



where the glued edge of  $T_e$  must have no crosses marking it.

Basing relation of sliced crossed links is the equivalence relation generated by moving a cross of each component of the associated link in some other admissible position in the sliced crossed link. Notice that moving the cross of a sliced crossed link can be obtained (in an equivalent way) by the local moves  $_1$  and  $_2$ , where  $_1$  (resp.  $_2$ ) moves the cross across consecutive tangles in such a way that a gluing site is not

(resp. is) crossed. Here is an example of a 2 move:



The wrapping relation <sup>!</sup> of sliced crossed links is generated by the following move



Recall that Theorem 1.1 identi es the set of F -links with a quotient of N (O). Since we are interested to give a description of the set of F -links and N (O) with a quotient of D (D<sub>g</sub>), we need to introduce analogs of the -relation on D (D<sub>g</sub>).

The handleslide move is generated by the move:

f:::;
$$T_{i-1}$$
; $T_{i}$ ; $x_i$ ; $x_j$ ; $T_{i+2}$ ;::: $g$  ! f:::; $T_{i-1}^0$ ; $x_i$ ; $x_j$ ; $x_i$ ; $x_j$ ; $T_{i+2}^0$ ;::: $g$ :

where each tangle in  $fT_k^0$  is obtained from the corresponding tangle in  $fT_k$  by taking a fram ed parallel of every component contributing to the component  $x_i$ ;

Proposition 2.6. There is a 1-1 correspondence

$$D(D_q)=hr;$$
;!i! N(O) and  $D(D_q)=hr;$ ;!; i! F links

Proof. The rst identication is well-known without crosses, see for example [AMR] and [Tu, Figure 7]. The idea is that it follows from general position arguments analogous to those that prove the su ciency of the Reidem aster moves for planar knot projections. The addition of crosses presents no extra di culty. The second identication follows easily from the rst, together with the following observation that takes care of inner automorphisms: the group Inn (F) of inner automorphisms acts trivially on the set of F -links (by de nition) and it also acts trivially on the set D (D<sub>g</sub>)=hr; ;! i. The proof of this observation is a pleasant exercise, left to the reader.

### 3. An assortment of diagrams and their relations

In this section we explain what are the spaces of diagram s in which our invariants will take values.

We begin with some notation, which generalizes the notion of the A-groups introduced in [GL1]: given a ring R with a distinguished group of units U, and (possibly empty sets) X;Y;T, D ("<sub>X</sub>;?<sub>Y</sub>;~<sub>T</sub>;R;U) is the completed graded vector space over Q generated by

Uni-trivalent diagram s with skeleton 1/2, with symmetric univalent vertices labeled by Y [ T.

The diagram s have oriented edges and skeleton and each edge is labeled by an element of R, such that the edges that are part of the skeleton are labeled only by U. M oreover, the product of the labels along each component of the skeleton is 1. Labels on edges or part of the skeleton will be called beads.

The degree of a diagram is the num ber of trivalent vertices, and the completion is with respect to this degree. Empty sets will be omitted from the notation, and so will U, the selected group of units of R. For example, A (?<sub>Y</sub>;R), A (R) and A () stands for A (";?<sub>Y</sub>;~;R;U), A (";?;~;R;U) and A (";?;~;Z;1) respectively. Univalent vertices of diagram s will often be called legs. D iagram s will sometimes be referred to as graphs. Special diagram s, called struts, labeled by a; c with bead b are drawn as follows

oriented from bottom to top. Multiplication of diagram  $s D_1$  and  $D_2$ , unless otherwise mentioned, means their disjoint union and will be denoted by  $D_1 D_2$  and occasionally by  $D_1 t D_2$ .

We will only be interested in the rings = Q [F] (the group ring of the free group F on some xed set  $T = ft_1; :::; t_g g$  of generators), its completion  $\hat{}$  (with respect to the powers of the augmentation ideal) and its C ohn localization  $_{loc}$  (i.e., the localization with respect to the set of matrices over that are invertible

over Z) all with selected group of units F.; and  $_{loc}$  are rings with involution induced by  $g = g^{-1}$  for  $g \ge F$ , with augmentation over Q, and with a commutative diagram



where !  $is given by the m odern version of the M agnus expansion t_i ! e^{h_i} . can be indentied with the ring of noncommuting variables fh_1;:::;h_g g and ______ with the ring of rational functions in noncommuting variables ft_1;:::;t_g, [FV, BR].$ 

W e now collect various relations on spaces of diagram s. W e de ne

$$A(''_X;?_Y;R;U) = D(''_X;?_Y;R;U) = (AS;IHX;Multilinear;Push)$$

where AS; HX and Multilinear are the relations shown in Figure 1 and Push is the relation shown in Figure 2. Note that the AS and IHX are well-known relations in the theory of nite type invariants of links and 3-m anifolds, whereas the Multilinear and Push are more recent, rst appearing in work of [GL1] in relation to nite type invariants of hom ology surgery and in [GR]. Note also that the above relations on are hom ogenous with respect to the grading, thus A is a completed graded vector space, too.



Figure 1. The AS, HX (for arbitrary orientations of the edges) and multilinear relations. Here r! r is the involution of R, r; s 2 R.



Figure 2. Pushing a unit g 2 U past a trivalent vertex.

The following is a companion to Remark 2.1:

Remark 3.1. Throughout the paper, given a subspace W of a vector space V, we will denote by V = (W) the quotient of V modulo W. O ccasionally, W will be dened by one of the following ways:

Either by a linear action of a group G on V, in which case  $W = fgv \quad vg$  and V = (W) coincides with the space of coinvariants of G on V.

Orby a subset of V, in which case W is the subspace spanned by that subset.

We now introduce on A algebraic analogs of the geometric wrapping relations of sliced crossed links in  $D_g$ . We begin by dening the in nitesimal wrapping relations ! on A. This will require an enlargement of the vector space A by a set of matrices as stated in Theorem 1.3, as well as some analysis of the notion of localization. We begin with the latter.

3.1. A review of the C ohn localization. In this self-contained section we review several facts about the C ohn localization that are used in the text. Consider the free group  $F_g$  on generators  $ft_1$ ;:::; $t_gg$ . The C ohn localization  $f_{bc}$  of its group-ring = Q  $F_g$ ] is characterized by a universal property,



namely that for every -inverting ring hom om orphism : ! R there exists a unique ring hom om orphism :  $l_{\text{loc}}$  ! R that m akes the above diagram commute, [C].

Farber and Vogel identied loc with the ring of rational functions in noncommuting variables, [FV, BR]. An example of such a rational function is

$$(3 t_1^2 t_2^{-1} (t_3 + 1)t_1^{-1})^{-1} t_2 5$$
:

There is a close relation between rational functions in noncommuting variables and nite state automata, sederibed in [BR, FV]. Farber-Vogel showed that every elements 2  $_{loc}$  can be represented as a solution to a system of equations M = D where M is a matrix over , invertible over Z, and D is a column vector over , FV, Proposition 2.1]. More precisely, we have that

## $s = (1;0;...;0)M^{-1}b;$

which we will call a matrix presentation of s 2  $_{loc}$ .

In the text, we often use substitutions of the form  $'_{t_i! t_i e^h}$ , or  $'_{t_i! e^h t_i}$  or  $'_{t_i! e^h t_i e^h}$ , where h does not commute with  $F_g$ . In order to make sense of these substitutions, we need to enlarge our ring  $_{bc}$ . This can be achieved in the following way.  $F_g$  can be included in the free group F with one additional generator t. Let L denote the group-ring of F,  $L_{bc}$  denote its C ohn localization and  $f_{bc}^l$  denote the completion of  $L_{bc}$  with respect to the the ideal generated by t 1. There is an identication of  $f_{bc}^l$  with the following ring R whose elements consist of form all sums of the form  $\int_{m,f}^{1} \int_{i=1}^{1} h^{m_i} f_i$  where  $f_i 2 \int_{bc}^{1} h^{m_i} f_i$  and

 $m = (m_1; m_2; \dots) : N ! N$  is eventually 0 and  $f = (f_1; f_2; \dots) : N ! loc$  is eventually 1.

In the above sum, for each xed k, there are nitely many sequences m with  $jn j = m_i k$ .

It is easy to see that R is a ring with involution (in fact a subring of the completion of L with respect to the augmentation ideal) such that  $t_i$  and  $e^h$  are units, where  $h = \log t$ . In particular,  $\hat{B}_{loc}$  is an h-graded ring. Let  $\deg_h^n : \hat{B}_{loc} ! \hat{B}_{loc}^n$  denote the projection on the h-degree n part of  $\hat{B}_{loc}$ . There is a map 'i: !  $\hat{B}_{loc}$  given by substituting  $t_i$  for  $e^h t_i e^h$  for 1 i g. It extends to a map 'i:  $b_c$  !  $\hat{B}_{loc}$ . Indeed, using the dening property of the Cohn localization, it suces to show that 'i is -inverting, [C, FV]. In other words, we need to show that if a matrix W over is invertible over Z, then 'i(W) is invertible over  $\hat{B}_{loc}$ . However,  $\deg_h^0$  ('i(W)) = W, invertible over  $b_{loc}$  and thus also over  $L_{loc}$ . The result follows.

$$: loc ! \mathbf{P}_{loc}$$

denote the h-degree 1 part of ' $_{i}$ . The following lemma gives an axiomatic de nition of  $_{i}$  by properties analogous to a derivation. Compare also with the derivations of Fox di erential calculus, [FV].

Lem m a 3.2.  $_{\rm i}$  is characterized by the following properties:

$$i(t_j) = i_j[t_i;h] = i_j(t_ih h t_i)$$
  
 $i(ab) = i(a)b + a_i(b)$   
 $i(a + b) = i(a) + i(b):$ 

Proof. It is easy to see that  $_{i}$  satis es the above properties, by looking at the h-degree 0 and 1 part in  $'_{i}$  (a). This determ ines  $_{i}$  uniquely on . By the universal property, we know that  $_{i}$  extends over  $_{bc}$ ; a priori the extension need not be unique. Since however,  $_{i}$  extends uniquely to matrices over , and since every element of  $_{bc}$  has a matrix presentation (see the discussion in the beginning of the section), it follows that  $_{i}$  is characterized by the derivation properties.

3.2. The in nitesim alw rapping relations. We have all the required prelim inaries to introduce an innitesim alw rapping relation on  $A^0 = B(_{bc} ! Z) A \cdot G$  iven a vector v 2 A (?<sub>X</sub>;  $_{bc}$ ) and a diagram D 2 A (?<sub>X</sub>;  $_{bc}$ ) with a single special leg colored by @h and M 2 Herm ( $_{bc} ! Z$ ), the in nitesim alw rapping relation  $!_i$  at the ith place in Herm ( $_{bc} ! Z$ ) A (?<sub>X</sub>;  $_{bc}$ ) is generated by the move

$$(M;v)^{!_{i}} M;v + con_{fhg} i (D) \frac{1}{2} D t (fr(M^{-1};M))$$

It is easy to see that the above de nition depends only on the image of M in B (  $_{\rm loc}$  ! Z). We de ne

$$A^{0}(?_{x}; b_{c})=h! i = A^{0}(?_{x}; b_{c})=h!_{1};...;!_{q}i:$$

The following example is a special case of the  $!_i$  relation for a diagram labeled whose edges are monomials  $t_i$ , with M the empty matrix, and where all the edges labeled by  $t_i$  are shown:



Note that in the ! relation, we allow for D to be a diagram of a circle (with a bead and with a special  $\theta$  leg). In addition, we allow the  $\theta$  heg to have a bead in  $_{\rm loc}$ .

3.3. G roup-like relations. Although the invariant  $Z^{rat}$  that appears in Theorem 1.3 takes values in  $A^0$ , we will actually construct a re ned invariant =  $Z^{rat;gp}$  which takes values in a set of group-like elements, modulo appropriate equivalence relations.

Let A <sup>gp</sup> denote the set of group-like elements of A i.e., elements of the form exp(a) where a is a sum of connected diagram s.

We now de ne an analog of the basing relations on  $A^{gp}$ . Let  $\mathfrak{m}_{z}^{x,y} : A^{gp}("_{fx,yg} [ "_{x}; _{bc}) ! A^{gp}("_{fzg} [ "_{x}; _{bc}) )$  be the map that glues the end of the x-skeleton to the beginning of the y-skeleton and then relabeling the skeleton by z, for x; y; z  $\mathfrak{I}$  X. Given  $s_1; s_2 \ 2 \ A^{gp}(?_{x}; _{bc})$ , we say that

 $s_1^{(1)}$  s<sub>2</sub>, if there exists an element s 2 A <sup>gp</sup> ("<sub>fxg</sub>"<sub>fx<sup>0</sup>g</sub>"<sub>x fxg</sub>; <sub>loc</sub>) with the property that

(2) 
$$(\mathfrak{m}_{x}^{x,x^{0}}(s)) = s_{1} \text{ and } (\mathfrak{m}_{x}^{x^{0},x}(s)) = s_{2}$$

 $s_2 = s_2 = s_2 = s_2$  is obtained from  $s_1$  by pushing an element of F on each of the X -labeled legs of  $s_1$ . For example, if all  $x_1$  legs are shown below, we have:

$D_1$	$D_2$		$D_1$	$D_2$
+		2	$f \bullet \bullet f$ +	$f \bullet$
$x_1 x_1 x_2 x_2$	$x_1 x_2 x_2 x_2$		$x_1 x_1 x_2 x_2$	$x_1 x_2 x_2 x_2$

Remark 3.3. When we push an element of the free group f in the legs of a diagram, depending on the orientation of the leg, we add f or f to the leg. See also Figure 2 for a convention for all choices of orientations. Similarly, when we glue two legs of diagrams together, the label on the bead depends on the orientations of the legs. For example,

$$a \stackrel{\frown}{\bullet} b \stackrel{\frown}{\bullet} a_{ab} \stackrel{\frown}{\bullet} b \stackrel{\frown}{\bullet} a_{ab} \stackrel{\frown}{\bullet} b \stackrel{\frown}{\bullet} a_{\overline{b}} \stackrel{\frown}{\bullet} b \stackrel{\bullet}{\bullet} a_{\overline{b}} \stackrel{\bullet}{\bullet} \stackrel{\bullet}{\bullet} a$$

Wede ne

$$A^{gp} (\sim_X ; bc) = A^{gp} (?_X ; bc) = h_1^{gp} ; 2^{gp} i$$

We mention a useful reform ulation of the  $\frac{gp}{1}$  relation for diagram s in A (?<sub>x</sub>; <sub>loc</sub>), which makes obvious the form alsim ilarity between the  $\frac{gp}{1}$  and  $\frac{gp}{2}$  relations. In order to do so, it is convenient to introduce a contraction m ap con: A ( $x_1$  ( $x_2$ ;  $x_1$ ,  $x_2$ )! A (:::;  $x_1$ ,  $x_2$ ), de ned as follows. For s 2 A ( $x_1$  ( $x_2$ ;  $x_2$ ,  $x_2$ ),

$$\operatorname{con}_X(s) = (\operatorname{div}_X)_{X \to 0}$$
:

In words, contraction glues all ways of pairing all legs labeled from @X to all legs labeled from X. For  $s_1; s_2 \ge A^{gp}$  (?x;  $l_{oc}$ ), we say that

$$s_1 = s_2$$
 i there exists  $s_{12} \ge A^{gp}$  (?<sub>X [@h</sub>; <sub>bc</sub>) such that

$$s_1 = con_{fhg}(s_{12})$$
 and  $s_2 = con_{fhg}(s_{12}) i_{X ! X e^h}$ 

where  $s_{12} \, \dot{x}_{\,! X \, e^h}$  is by de nition the result of pushing  $e^h$  to each X -labeled leg of  $s_{12}$ .

Lem m a 3.4. The equivalence relations  $\frac{gp}{1}^{\circ}$  and  $\frac{gp}{1}$  are equal on A  $^{gp}$  (?x;  $_{loc}$ ).

Proof. Suppose that  $s_1 \stackrel{\text{gp}}{\stackrel{1}{1}} s_2$  and consider the corresponding element s from Equation (2). We start by presenting s as a result of a contraction. Let  $= \iint_{x} e^h e^h f_{xg}(s) = \int_{x^+ e^h} e^h e^h ds$ . Let  $m = \int_{x^+}^{y} e^h e^h f_{xg}(s) = \int_{x^+ e^h} e^h ds$ .

$$s = con_{fhg} (s(@h));$$
  $s_1 = con_{fhg} \times m_x^{x,x^0} ();$   $s_2 = con_{fhg} \times m_x^{x^0,x} ():$ 

Now the term  $m_x^{x,x^0}$  () is obtained from the term  $m_x^{x^0,x}$  () by conjugating by an e<sup>h</sup>. Indeed:



C on jugating with  $e^h$  com m utes with the sym m etrization m ap  $_X$  (E xercise!). Letting

$$s_{12} = \operatorname{con}_{fhg} \quad x \quad \operatorname{re}_{x}^{x,x^{\circ}}() \quad j_{eh!} \quad e_{h+e_{h^{\circ}}} \quad 2 \quad A^{gp} \quad (?_{x \ [e_{h^{\circ}}; \ b_{c}});$$

it follows that

 $s_1 = con_{fh^0g}(s_{12})$  and  $s_2 = con_{fh^0g}(s_{12}j_{k! xe^{h^0}})$ 

as needed. The converse can be obtained by reversing the steps in the above proof.

Now, we de ne a group-like wrapping relation on A  $^{\rm gp}$ . This enlargement m ight sound articial, however there are several good reasons for doing so. For example, the bop move on the set of F-links and its associated ltration predicts that in degree 0, the universal invariant is given by the S-equivalence class (of F-links), or equivalently, by the set of B lanch eld pairings of F-links. A  $^{\rm gp}$ , 0 (  $_{\rm bc}$ ), de ned below, m aps onto B (  $_{\rm bc}$  ! Z) which in turn m aps onto the set of B lanch eld pairings of F-links.

In addition, the A-groups, studied in relation to Homology Surgery in [GL1], are secondary obstructions to the vanishing of a W itt-type obstruction which lies in B (Z [ ]! Z). In the case of F-links, this motivates the fact that the set B ( $_{loc}$ ! Z) should be taken together with A ( $_{loc}$ ). W ith these motivations, we will consider the set

$$A^{gp;0}(?_X; bc) = B(bc!Z) A^{gp}(?_X; bc):$$

De nition 3.5. Two pairs (M;  $s_m$ ) 2 Herm ( $_{loc}$ ! Z) A<sup>gp</sup> (?<sub>X</sub>;  $_{loc}$ ) m = 1;2 are related by a wrapping move at the jth gluing site (for some j such that 1 j g) if there exists an s 2 A<sup>gp</sup> (?<sub>X;@h</sub>;  $_{loc}$ ) such that:

$$s_m = con_{fhg} \prime_m (s) t exp_t \frac{1}{2} h (M^{-1} \prime_m M)$$

for m = 1;2 where '<sub>1</sub> is the identity and '<sub>2</sub> is the substitution '<sub>t<sub>i</sub>! e h t<sub>i</sub>eh, and</sub>

$$h(A) = \bigcirc trlog(A) 2 A (?_h; loc):$$

Our not task is to make sure that the above de nition makes sense. Using the h-graded ring with involution  $\mathbf{E}_{\text{loc}}^{l}$  from Section 3.1, it follows that 'm (s<sub>12</sub>) 2 A <sup>gp</sup> (?<sub>X</sub>;  $\mathbf{E}_{\text{loc}}^{l}$ ). Furthermore, h M <sup>1</sup>'m M) can be thought of either as a circle with a bead in  $\mathbf{E}_{\text{loc}}^{l}$  (cyclic) that has vanishing h-degree 0 part, or as a sum of wheels with h-colored legs and beads in  $\mathbf{E}_{\text{loc}}^{l}$ . Thus, the contraction  $\operatorname{con}_{\text{fhg}}$  in the above de nition results in an element of A <sup>gp</sup> (?<sub>X</sub>;  $\mathbf{E}_{\text{loc}}^{l}$ ). It is easy to see that h satisfies the properties in Proposition 2.5; thus h depends only on the image of M in B ( $\mathbf{E}_{\text{loc}}^{l}$  Z) and the above de nition makes sense in A <sup>gp;0</sup>.

Remark 3.6. In the above de nition 3.5 of the wrapping relation, we could have chosen equivalently that  $'_1$  and  $'_2$  denote the substitutions  $'_{t_i! t_i e^h}$  and  $'_{t_i! e^h t_i}$ .

Remark 3.7. In the above de nition, the @h colored legs are alloed to have beads in  $_{loc}$ . A lso, we allow the case of the empty matrix M, with the understanding that the  $_{h}$  term is absent.

Lem m a 3.8. For g = 1, the wrapping relation  $!^{gp}$  on  $A^{gp;0}$  (?<sub>X</sub>;  $_{loc}$ ) is trivial.

Proof. It follows from a swapping argument analogous to the isotopy of the following gure



W e end this section with a comparison between A<sup>0</sup> and A<sup>gp;0</sup>. There is an obvious inclusion

 $A^{gp;0}$  (?<sub>x</sub>; loc) !  $A^{0}$  (?<sub>x</sub>; loc):

The next proposition states that

Proposition 3.9. The inclusion  $A^{gp;0}$  (?<sub>x</sub>; <sub>loc</sub>) !  $A^{0}$  (?<sub>x</sub>; <sub>loc</sub>) maps the !<sup>gp</sup>-relations to !-relations, and thus induces a map

$$A^{gp;0}(?_X; b_c)=h!^{gp}i ! A^0(?_X; b_c)=h!i:$$

Proof. See Appendix C.

4. An invariant of links in the complement of an unlink

The goal of this section is to de ne an invariant  $Z^{rat} : D_X (D_g) ! A^{gp} (~_X; )$  of sliced crossed links whose components are in 1-1 correspondence with a set X.

G iven a nite set X, an X -indexed sliced crossed link on D $_g$  is a sliced crossed link on D $_g$  equipped with bijection of the set of components of the presented link with X. All the de nitions above have obvious adaptations for X -indexed objects, indicated by an appropriate X subscript when required.

Theorem 4.1. There exists an invariant

that m aps the topological basing relations to group-like basing relations and the topological wrapping relations to wrapping relations. In particular, it induces a map

$$Z^{rat}:N_X(O) ! A^{gp}(\sim_X;):$$

Proof. Our rst task is to de ne Z<sup>rat</sup>. In order to do so, we need to introduce som em ore notation. G iven a word w in the letters " and #, let

", denote the corresponding skeleton.

I 2 A (", ; ) denote the identity element.

 $G_{i,w}$  denote the gluing word at the ith gluing site, that is,  $I_w$  with a label of  $t_i$  on each component, in the sense described by the following example: if w = "##, then  $G_{2;"##} = [t_2]_{i} t_2 = t_2$ .

Given a word w in the symbols ";#;" and #, let w<sup>0</sup> denote w with the crosses forgotten. Let  $I_w$  be  $I_w^{\circ}$  with every crossed strand broken. For example,

$$\mathbb{I}^{\mathbb{X} \# \mathbb{X}} = \left( \begin{array}{c} \\ \\ \\ \\ \end{array} \right) \left( \begin{array}{c} \\ \\ \\ \\ \end{array} \right) \left( \begin{array}{c} \\ \\ \\ \\ \\ \end{array} \right)$$

D e nition 4.2. Take an X -indexed element D =  $fT_1$ ;:::; $T_kg 2 D_X (D_g)$ .

- 1. Select q-tangle lifts of each tangle in the sequence,  $fT_1^0$ ; ...;  $T_k^0$ g, such that
  - (a) the strands of each q-tangle which lie on boundary lines of the following form



are bracketed  $(w_1)(w_2)$ , where  $w_1$  (resp.  $w_2$ ) corresponds to the strands going to the left (resp. right) of the rem oved disc, and these bracketings m atch up,

(b) boundary words at boundaries of the form



are given the canonical left bracketing.

- 2. Compose the (usual) K ontsevich invariants of these q-tangles in the following way:
  - (a) if two consecutive tangles are as follows,

with a word w in the symbols "; #; X; # describing their m atching boundary word, then they are to be composed

(b) if, on the other hand, two consecutive tangles m eet as follows,

f:::; 
$$\begin{bmatrix} T_i \\ \\ \\ \\ \end{bmatrix}$$
 ;  $\begin{bmatrix} T_{i+1} \\ \\ \\ \\ \\ \end{bmatrix}$  :::g

with their matching boundary word at the jth gluing site factored as  $(w_1)(w_2)$ , then they are to be composed

(I) (I) (I, G<sub>1</sub>, G<sub>1</sub>, C) Z (I<sup>0</sup><sub>i+1</sub>) ::::

This completes the denition of Z<sup>rat</sup> (D). A normalized version Z<sup>rat</sup> (D) (useful for invariance under K irby m oves) is dened by connect-sum m ing a copy of into each component of the crossed link associated to D. O bserve that the result does not depend on the choice of a place to connect-sum to.

The proof of Theorem 4.1 follows from Lemmas 4.4 and 4.5 below.

The next lem ma, observed by the second author in [Krl] follows from a \sweeping" argum ent.

Lem m a 4.3. Let s denote an element of A ( $"_X$ ;Q) A ( $"_X$ ; ), (i.e., some diagram with labels 1 at each edge), where the top boundary word of  $"_X$  is  $w_1$ , and the bottom boundary word of  $"_X$  is  $w_2$ . Then

$$G_{W_1}$$
  $S = S$   $G_{W_2}$ :

Lem m a 4.4. The map  $Z^{rat}$  is well de ned on  $D_X$  ( $D_q$ ) and moreover descends to an invariant of  $D_X$  ( $D_q$ )=hri.

Proof. For the rst part we only need to show that the function does not depend on the choice of bracketings. This follows because of the following, where  $_1$ ;  $_2$  are some expressions built from the associator.

$$\stackrel{\text{(I)}}{=} (\mathbf{T} (1 2) (\mathbf{I}_{1} \mathbf{G}_{j;W_{2}}) (1^{1} 2^{1}) \mathbf{Z} (\mathbf{I}_{i+1}^{0}) ::::;$$

$$= \stackrel{\text{(I)}}{=} (\mathbf{T} (\mathbf{I}_{i_{1}} \mathbf{G}_{j;W_{2}}) (1^{1} 2) (1^{1} 2^{1}) \mathbf{Z} (\mathbf{I}_{i+1}^{0}) ::::;$$

$$= \stackrel{\text{(I)}}{=} (\mathbf{T} (\mathbf{I}_{i_{1}} \mathbf{G}_{j;W_{2}}) \mathbf{Z} (\mathbf{I}_{i+1}^{0}) :::::$$

where the rst step follows from Lemma 4.3. The invariance of Z<sup>rat</sup> under the r equivalence relation follows the same way.

Lem m a 4.5. The map

$$Z^{rat}:D_X(D_q)=hr; i! A^{gp}(?_X;)=h^{gp}i$$

is well-de ned.

Proof. We need to show invariance of  $Z^{rat}$  (or equivalently,  $Z^{rat}$ ) under the  $_1$  and  $_2$  moves. Enlarge  $D_X (D_g)$  to  $D_X (D_g)^0$ , the set of X-indexed sliced crossed links on  $D_g$  whose presented links in  $D_g$ . I have at least one cross on every component, thus  $D(D_g) = D(D_g)^0$ . The function  $Z^{rat}$  immediately extends to this generality. For two diagram  $sD_1; D_2 2 D_X (D_g)$  such that  $D_1 = D_2$  let  $D_{12}$  denote the diagram in  $D(D_g)^0$  such that  $D_1$  is obtained by forgetting some cross, and  $D_2$  is obtained by forgetting the other cross on that component. Nootice that in this case  $Z^{rat}(D_{12}) 2 A^{gp}("fxg"fx^0g"X fxg;)$ : The key point is the fact that the product of labels around each component of the skeleton equals to 1; this is true since the locations of the two crosses cut the component into two arcs, both of which have trivial intersection word with the gluing sites. It follows by construction that

$$Z^{\text{rat}}(\mathbb{D}_{1}) = \mathfrak{m}_{x}^{x;x^{\circ}}(Z^{\text{rat}}(\mathbb{D}_{12}));$$

$$Z^{\text{rat}}(\mathbb{D}_{2}) = \mathfrak{m}_{x}^{x^{\circ};x}(Z^{\text{rat}}(\mathbb{D}_{12}));$$

$$Z^{\text{rat}}(\mathbb{D}_{12}) = 2 \quad A^{\text{gp}}(\mathbf{"}_{fxg}\mathbf{"}_{fx^{\circ}g}\mathbf{"}_{x} fxg; loc):$$

Thus,  $(\mathbb{Z}^{\operatorname{rat}}(\mathbb{D}_1))^{\operatorname{gp}}$   $(\mathbb{Z}^{\operatorname{rat}}(\mathbb{D}_2)):$ 

If D  $_1$  <sup>2</sup> D  $_2$ , then for some appropriate expression built from the associator, we have that:

$$(Z^{rat}(f:::;) ) = ( _{W} I (I_{1} G_{i_{W_{2}}}) ^{1} :::)$$

$$(Z^{rat}(f:::;) ) = ( _{W} I (I_{1} G_{i_{W_{2}}}) ^{1} :::)$$

$$( _{W_{1}} (IG_{i_{W_{2}}}) I_{W} ^{1} :::)$$

$$( _{W_{1}} (IG_{i_{W_{2}}}) I_{W} ^{1} :::)$$

$$= (Z^{rat}(f:::;) ) ( _{W_{1}} (IG_{i_{W_{2}}}) I_{W} ^{1} :::))$$

Lem m a 4.6. The following m ap is well-de ned:

Let.

Z<sup>rat</sup> :D (D<sub>g</sub>)=h!i! A <sup>gp</sup> (?<sub>X</sub> ; )=h! <sup>gp</sup>i

P roof. Consider two sliced crossed links  $D_1; D_2 2 D (D_g)$  related by a topological wrapping m ove at the jth gluing site as in Section 2.4. In other words,  $D_1$  (shown on the left) and  $D_2$  (shown on the right) are given by:



Now, we can compute, for some element built from an associator:

In the above,  $\frac{w_0}{v_0}$  means the comultiplication of the diagram x with pattern given by the gluing word  $w_2$ .

Theorem 4.1 now follows, using the identication of N<sub>X</sub> (O) with the set  $D_X$  ( $D_g$ )=hr; ;! i given by Proposition 2.6.

Rem ark 4.7. An application of the Z<sup>rat</sup> invariant is a formula for the LMO invariant of cyclic branched covers of knots, [GK]. This application requires a renormalized version Z<sup>rat</sup>[] where =  $(_1; :::;_g)$  and  $_i 2 A^{gp}$  (?<sub>k</sub>). Z<sup>rat</sup>[] is de ned using D e nition 4.2 and replacing the formula in 2(b) by

$$(I)$$
 ( $I$  ( $I_{w_1}$  G  $_{j,w_2}$  ( $_{j}$ )) Z ( $I_{i+1}^0$ ) ::::

where (j) places a copy of j in each of the skeleton segments of the jth gluing site. The renormalized invariant Z<sup>rat</sup> [] still satisfies the properties of Theorem 4.1.

# 5. A rational version of the Aarhus integral

The main purpose of this section is to introduce a notion  $R_{rat}$  of Rational Form al Gaussian Integration  $R_{rat}$  which respects the K index moves the basing and the winding relations. The reader who is not fam iliar with the combinatorics of gluing legs of diagram s together, is urged to look at the appendix.

5.1. The densition of  $R_{rat}$ . Recall that Herm ( $_{loc}$ ! Z) denotes the set of Herm itian matrices over  $_{loc}$ , invertible over Z. Let X denote a nite set and let X<sup>0</sup> = fx<sub>i</sub>g denote an arbitrary subset of X. A diagram D 2 A (?<sub>X</sub>;  $_{loc}$ ) is called X<sup>0</sup>-substantial if there are no struct components (i.e., components comprising of a single edge) both of whose univalent vertices are labeled from the subset X<sup>0</sup>. An element of A (?<sub>X</sub>;  $_{loc}$ ) is a series of X<sup>0</sup>-substantial diagram s.

An element s 2 A ( $?_X$ ;  $_{loc}$ ) is called integrable with respect to X  $^0$  if there exists an Herm itian matrix M 2 Herm ( $_{loc}$ ! Z) such that

(3) 
$$s = \exp^{\frac{0}{2} \frac{1}{2} \sum_{i,j=x_i}^{x_j} M_{ij}^A t R;}$$

with R an X  $^{0}$ -substantial element. Notice that M, the covariance matrix cov(s) of s, and R, the X  $^{0}$ -substantial part of s, are uniquely determined by s.

Let  $Int_X \circ A$  (?<sub>X</sub>;  $l_{bc}$ ) A (?<sub>X</sub>;  $l_{bc}$ ) denote the subset of elements that are integrable with respect to X<sup>0</sup>. O bserve that although  $Int_X \circ A$  (?<sub>X</sub>;  $l_{bc}$ ) has non-empty intersection with A<sup>gp</sup> (?<sub>X</sub>;  $l_{bc}$ ), it is not a subset of it. Further, observe that if two elements of A<sup>gp</sup> (?<sub>X</sub>;  $l_{bc}$ ) are related by a basing move and one of them lies in  $Int_X \circ A$  (?<sub>X</sub>;  $l_{bc}$ ), so does the other. Thus the following de nition makes sense:

$$\operatorname{Int}_{X} \circ A^{\operatorname{gp}} (\sim_{X} ; _{\operatorname{bc}}) = (A^{\operatorname{gp}} (?_{X} ; _{\operatorname{bc}}) \setminus \operatorname{Int}_{X} \circ A (?_{X} ; _{\operatorname{bc}})) = h_{1}^{\operatorname{gp}} ; _{2}^{\operatorname{gp}} i:$$

For the next de nition, recall that every matrix W 2 Herm ( loc! Z) is invertible over loc.

De nition 5.1. De neamap (Rational Formal Gaussian Integration)

Z rat  
dX 
$$^{0}$$
 : Int<sub>X</sub>  $\circ$ A (?<sub>X</sub> ;  $_{loc}$ ) ! A (?<sub>X X</sub>  $\circ$ ;  $_{loc}$ )

as follows. If s 2 Int<sub>x</sub> A (?<sub>x</sub>; loc) has associated decomposition given in Equation (3), then

The gluing of the legs of diagram s occurs according to Remark 3.3.

Remark 52. The denition of  $R_{rat}$  for elements swhose covariance matrix M is dened over, and invertible over Z, requires beads labeled by the inverse matrix M  $^1$ . This necessitates the need to replace by an appropriate ring  $R_{rat}$  that makes all such matrices M invertible. It follows by denition that the universal choice of  $R_{rat}$  is the C ohn localization of .

A complementary reason to consider the Cohn localization, from the (dual) point of view of Itrations comes from the the study of the loop Itration, given in [Ga].

5.2.  $R_{rat}$  respects the basing relations. The next theorem shows that  $R_{rat}$  is compatible with the basing relations.

Theorem 5.3.  $R_{rat}$  descends to a map:  $Z_{rat}$  $dX^{0}: Int_{X^{0}}A^{gp}(?_{X}; b_{c})=h_{X}^{gp}i! A^{gp}(?_{X X^{0}}; b_{c})=h_{X}^{gp}_{X^{0}}i!$ 

 $\begin{array}{c} \text{Proof. Using Lemma 3.4, we need to show that given $s_1;s_2$ 2 $Int_X \circ A$ (?_X; $loc$) \ A$ $^{gp}$ (?_X; $loc$) such that $s_1^{$gp^0$}$ $s_2$ or $s_1^{$gp^0$}$ $s_2$ or $s_1^{$gp^0$}$ $s_2$ , based on some element $x_k$ 2 X $^0$, then $s_1$ then $s_1$ $s_2$ or $s_1^{$gp^0$}$ $s_2$ or $s_1^{$gp^0$}$ $s_2$ or $s_1$ $s_2$ or $s_2$ or $s_1$ $s_2$ or $s_2$ or $s_1$ $s_2$ or $s_2$ or $s_1$ $s_2$ or $s_2$ or $s_1$ $s_2$ or $s_1$ $s_2$ or $s_1$ $s_2$ or $s_2$ or $s_2$ or $s_2$ or $s_2$ or $s_1$ $s_2$ or $s$ 

$$S_2 \text{ or } S_1$$
  $S_2$ , based on some element  $X_k 2 X^\circ$ , then  
 $Z_{rat}$   $Z_{rat}$   
 $dX^0(S_1) = dX^0(S_2)$ :

The case of  $\frac{gp}{2}$ . Consider the decomposition of  $s_1$ 

Ζ

$$s_1 = \exp \frac{1}{2} X \qquad \bigcup_{\substack{x_j \\ x_i}}^{x_j} M_{ij} tR:$$

Suppose that an element f 2  $F_g$  is pushed onto the legs labeled by  $x_k$  2 X<sup>0</sup>, to form  $s_2$ . Now, let D<sub>f,k</sub> be diag(1;1;:::;f;:::;1), where f is in the kth entry, and let R<sup>0</sup> denote R with an f pushed onto every leg marked  $x_k$ . Then,

$$\begin{array}{rcl} {}^{\mathrm{rat}} & & {}^{\mathrm{Z}} & {}^{\mathrm{rat}} & & {}^{\mathrm{rat}} & & {}^{\mathrm{rat}} & {}^{\mathrm{rat}} & {}^{\mathrm{rat}} & {}^{\mathrm{rat}} & {}^{\mathrm{rat}} & {}^{\mathrm{rat}} & {}^{\mathrm{g}} & {}^{\mathrm{g}$$

because all the beads labeled f and f  $^{1}$  m atch up and cancel.

The case of  $\frac{gp}{1}$ . Consider s 2 A gp (?<sub>X [@h</sub>; bc) such that

$$s_1 = con_{fhg}(s)$$
 and  $s_2 = con_{fhg}(s_{x_k}! x_k e^h)$ :

Notice that if  $s_1$ ;  $s_2$  are X<sup>0</sup>-integrable, so is s. Furthermore,  $con_{fhg}$  commutes with  $R_{rat}$ . The result follows from Lemma 5.4 below.

Proof. The idea of this lemma is the same as in the case  $\frac{gp}{2}$ ", but instead of pushing an element of the free group onto legs labeled by some element, we push an exponential of hair.

The substitution  $x ! xe^h$  de nes a map

with the property that s  $s_{1, xe^{h}}$  is X<sup>0</sup>-substantial for all s 2 A ( $?_{X}$ ;  $_{bc}$ ).

A ssum e that the canonical decom position of s is given by Equation (3). Let  $D_{f,k}$  be diag(1;1;:::;e<sup>h</sup>;:::;1), where e<sup>h</sup> is in the kth entry, and let  $R^0$  denote R with an e<sup>h</sup> pushed onto every leg m arked  $x_k$ . Then, Theorem E.1 from Appendix E im plies that

$$Z_{rat} = \exp \frac{1}{2} h M D_{e^{h};k} M^{-1} D_{e^{h};k} t \exp \frac{1}{2} X_{i} X_{ij}^{j} M_{ij}^{-1}; R$$

$$= \exp \frac{1}{2} X_{ij}^{j} M_{ij}^{-1}; R$$

where the last line follows from Proposition 2.5.

Remark 5.5. An alternative proof of Theorem 5.3 can be obtained by adapting the proof of [A, Part II, Proposition 5.6] to the rational integration  $R_{rat}^{rat}$ .

5.3. Dealing with the K inby m oves. Recall from Section 2.4 the handleslide m ove on the set D (D<sub>g</sub>). In a similar fashion, we de ne an X <sup>0</sup>-handleslide m ove on A ( $?_X$ ;  $_{loc}$ ), as follows: For two elements  $s_1$ ;  $s_2$  2 A ( $?_X$ ;  $_{loc}$ ), say that  $s_1 \xrightarrow{x^0} s_2$ , if for some  $x_1 \ge X$ ; and  $x_1 \ge X^0$ ; if j,

$$s_{2} = f_{x_{i}g} \quad m_{x_{i}}^{y,x_{i}} \quad f_{y,x_{i}g} \quad (s_{1} (x_{1}; \dots; x_{j-1}; x_{j} + y; x_{j+1}; \dots)) :$$

Notice that if  $s_1$ ;  $s_2 \ 2 \ A^{gp}$  (?<sub>X</sub>;  $_{bc}$ ),  $s_1 \xrightarrow{x^0} s_2$  and one of  $s_1$  lies in  $Int_X \circ A$  (?<sub>X</sub>;  $_{bc}$ ), so does the other. This is because the covariance matrix of  $s_2$  is obtained from the covariance matrix of  $s_1$  by a unimodular congruence.

Theorem 5.6. The map Z<sup>rat</sup> descends to a map:

$$Z^{rat}:D_X(D_q)=hr;$$
;  $X^{oi}!$  Int $X^{oA}$  (? $X^{gp}$ (? $X^{gp}$ )=h  $gp$ ;  $X^{oi}$ :

P roof. The argument of [LM M O, P roposition 1], adapts to this setting without di culty. There are two possible subtleties that must be observed.

1. The proof in [LM M O] is based on taking a parallel of the parts of a component (here, the component  $x_j$ ) away from the site of the cross. In our setting, the paralleling operation of the involved components of tangles behaves as usual (sum s over lifts of legs). At a gluing word, one simply labels both of the two components arising from the parallel operation with a bead labeled by the appropriate letter. Then, when

 $x_{i}$ , y is applied, these two identical beads are pushed onto the legs sitting on those components side-by-side. This may alternatively be done by rst pushing the original bead up onto the legs, and then taking the parallel, as is required in this situation.

2. We have de ned the bracketing at boundary lines with crosses to be the standard left-bracketing, while the move in [LM M O] is given for a presentation where the two involved strands are bracketed together. This is treated by preceding and following the  $x \circ w$  it a  $_1$  move, to account for the change in bracketing, exactly as in the proof of the  $_2$ -case of Lem m a 4.5.

Theorem 5.7. The map  $R_{rat} dX^0$  descends to a map:

Z rat  
dX <sup>0</sup>: Int<sub>x</sub> 
$$\circ$$
 A <sup>gp</sup> (?<sub>x</sub>; <sub>bc</sub>)=h x  $\circ$  i! A <sup>gp</sup> (?<sub>x x</sub>  $\circ$ ; <sub>bc</sub>):

This is proved by the following lemma.

Lem m a 5.8. Let  $s_1 \ 2 \ \text{Int}_X \circ A$  (?x ;  $_{\text{loc}}$ ) and let

$$s_2 = m_{x_i}^{y,x_i} (s_1 (x_1; \dots; x_{j-1}; x_j + y; x_{j+1}; \dots)):$$

If  $x_1 2 X^0$  then

$$\begin{array}{ccc} Z & & & Z \\ & & & \\ & & dX \\ \end{array} \begin{array}{c} & & Z \\ & & \\$$

Proof. We prove this in two steps, which usually go by the names of the \lucky" and \unlucky" cases. The lucky case:  $s_1; s_2 2$   $Int_{fx_1g}A$  (?x;  $b_c$ ). In this case, we can use the Iterated Integration Lemma A 2.  $\rm Z_{rat}$ Z <sub>rat</sub>  $Z_{rat}$  $dX^{0}(s_{2}) =$ d(X<sup>0</sup> dfx<sub>i</sub>g(s<sub>2</sub>) ; fxjg)  $Z_{rat} = Z_{rat}$   $d(X^{0} fx_{j}g) = dfx_{j}g_{fx_{i}g} m_{x_{i}}^{y;x_{i}} fy;x_{i}g (s_{1}(x_{1};...;x_{j-1};x_{j}+y;x_{j+1};...));$ Z<sub>rat</sub> d(X<sup>0</sup> fx<sub>j</sub>g)  $\begin{array}{cccc} & & & & & & & \\ & & & & & \\ fx_ig & & fx_{i} & & fy_{i}x_{i}g & & & & dfx_jg \mbox{ (s_1 (x_1; :::; x_{j-1}; x_j + y_{i}x_{j+1}; :::)) } \end{array} \ ,$ Z <sub>rat</sub>  $d(X ^{0} fx_{j}g) fx_{ig} m_{x_{i}}^{y,x_{i}} fy_{x_{i}g}$ 0 0 11 y because  $\operatorname{div}_{fx_jg}^{0} \exp^{(0)} \int_{j}^{j} 1^{A_i} A = 1$ : The unlucky case:  $s_1; s_2 \neq Int_{fx_jg} A$  (?x;  $_{1^{loc}}$ ). This pathology is treated by deform ation. Namely, we \deform "s<sub>1</sub> by multiplying it by  $\exp^{(0)}$ "  $_{j}^{1A}$ ; for a \small real," commuting with  $_{loc}$ . Formally, we Let be a variable and consider an integration theory based on Q ()  $\mathbb{E}_{\mathbb{C}}$  and its C ohn localization  $\mathbb{E}_{\mathbb{C}}$ . We identify  $Q^0() = Q()$ , the subring of rational functions in non-singular at zero, and the corresponding Cohn localization  $^{,0}_{\rm loc}$ . So, letting  $\begin{array}{rcl} 0 & 1 \\ & & & \\ s_{1} & = & \exp{(\theta)} & & \\ & & & \\ & & & \\$  $s_2 = m_{x_ig} m_{x_i}^{y_i x_i} f_{y_i x_ig} (s_1 (x_1; \dots; x_{j-1}; x_j + y_i x_{j+1}; \dots));$ it follows from the argument of the \lucky case" that  $Z_{rat} \qquad Z_{rat} \qquad Z_{rat} \qquad dX^{0}(s_{2}) = \qquad dX^{0}(s_{1}):$ 

Now, there is a ring hom om orphism :  $i_{\text{loc}}^{;0}$  !  $i_{\text{loc}}$ ; obtained by \setting to zero". Observing that, although of course  $r_{x_{ij}}^{\text{R}_{\text{rat}}} df_{x_{j}g}(s_{i}) \neq A (?_{x}; i_{\text{loc}}^{;0})$  (though it is now well-de ned),

$$dX^{0}(s_{i}) 2 A (?_{X}; c_{loc}^{0}); and dX^{0}(s_{i}) = dX^{0}(s_{i});$$

the proof follows.

Remark 5.9. These last lines are best understood from the realization of  $_{\rm loc}$  and  $_{\rm loc}$  as spaces of form al power series. The elements  ${}^{\rm R}{}_{\rm rat} dX^{0} (s_{\rm i})$  lie in A (?<sub>x</sub>;  ${}^{i0}_{\rm loc}$ ) because the determ inant of the augmentation of the covariance m atrix of  $s_{\rm i}$  is invertible in Q<sup>0</sup>().

5.4. Dealing with the w rapping relations. In this section we show that  $^{R_{rat}}$ -integration respects the group-like w rapping relations ! <sup>gp</sup> that were introduced in D e nition 3.5. Since the w rapping relations involve pairs ofm atrices and graphs, we begin by extending the de nition of  $^{R_{rat}}$  to pairs (M ;s) by setting

$$dX^{0}(M;s) = (M \cos(s); dX^{0}(s));$$

where  $\cos(s)$  is the covariance matrix of s. The motivation for this comes from the next theorem which states that preserves the wrapping relations.

Theorem 5.10. The map  $R_{rat}^{R_{rat}} dX^{0}$  descends to a map  $Z_{rat}^{Z_{rat}} dX^{0}$ : Int<sub>X 0</sub>A <sup>gp;0</sup> (?<sub>X</sub>; <sub>bc</sub>)=h! <sup>gp</sup>i! A <sup>gp;0</sup> (?<sub>X X 0</sub>; <sub>bc</sub>)=h! <sup>gp</sup>i

Proof. Consider two pairs (M ;s<sub>1</sub>)  $^{!^{gp}}$  (M ;s<sub>2</sub>) related by a wrapping m ove.

Using Theorem E 1 in Appendix E, it follows that if W is the common covariance matrix of  $s_1; s_2$  and  $s_{12}$ , then

$$\begin{array}{rcl} & & & & & & & & & \\ & & & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & &$$

Rem ark 5.11. The group-like w rapping relation is the m inim al relation that we need to impose on diagram s so that  $Z^{at}$  is invariant under the topological w rapping relation, as follows from the proof of Theorem 5.10.

5.5. The de nition of the rational invariant Z<sup>rat</sup> of F-links. Theorem s 4.1, 5.3, 5.7, and 5.10 in ply that there is a map

de ned as follows: for a F-link (M;L) of g components, choose a sliced crossed link C in D (D<sub>g</sub>) whose components are in 1-1 correspondence with a set X, and de ne

$$Z^{\text{rat};\text{gp}}(M;L) = dX \quad Z^{\text{rat}}(C) \quad c^{+(B)}c \quad (B)^{2} A^{\text{gp};0}(_{\text{loc}}) = h!^{\text{gp}}i$$

where  $c = {}^{K} dUZ (S^{3};U)$  are some universal constants of the unit-fram ed unknot U and where B is the linking matrix of C and (B) is the number of positive (resp. negative) eigenvalues of the symmetric matrix B. The slight renormalization given by multiplication by c is needed to deal with the stabilization K index move on C, and is justified as follows: recall that  $A^{gp}()$  is a group, which acts on the set  $A^{gp;0}(_{loc})$ by a (M;s) = (M;ats). Since the beads of the graphs in  $A^{gp}()$  are rational numbers independent of the  $t_i$ , it follows easily that the above action of  $A^{gp}()$  on  $A^{gp;0}(_{loc})$  induces an action of  $A^{gp}()$  on the quotient set  $A^{gp;0}(_{loc})=h!$  i. This is exactly what appears in the above de nition of  $Z^{rat}$ .

Composing further with the map  $A^{gp;0}$  ( $_{loc}$ )=h!  $^{gp}$ ; CAi !  $A^{0}$ =h!; CAi of Proposition 3.9, de nes the rational invariant Z<sup>rat</sup> of F-links that appears in Theorem 1.3.

Proof. (of R em ark 1.6) R ecall that the m atrix part W of Z<sup>rat</sup> is an element of B ( $_{loc}$ ! Z) and determines the S-equivalence class of the F-link, as follows from Proposition 1.2. R ecall also that the w rapping relations ! <sup>gp</sup> and ! depend only on W . Thus, it follows that S-equivalent F-links have equal matrix-part, and that the matrix-free part of Z<sup>rat</sup> m akes sense as an element of the graded vector space A ( $_{loc}$ )=(!).

Proposition 1.2 implies that (M; L) is a good boundary link, (i.e.,  $H_1(X^{!}; Z) = 0$ ) i its associated matrix W is invertible over . In that case, the construction of the Z<sup>rat</sup> invariant implies that Z<sup>rat</sup> (M; L) 2  $A^0() = h!$ ; CAi.

### 6. The CA action and invariants of boundary links

The goal of this section is to use the already constructed invariant of F -links in order to de ne an invariant of boundary links. The key observation is that the set of boundary links can be de ned with the set of orbits of an action

#### CAF Links! F Links

for a suitable group CA. Starting from this observation, we re ne the results of Sections 2-5 to take into account the action of the group CA.

61. A surgery description of boundary links. We begin by giving a surgery description of boundary links. Recall that the set of  $\ell$ -links can be identi ed as a quotient of the set of F-links modulo an action of a group of autom orphism scAg of the free group, K, CS2]. The group CAg in question is the subgroup of the autom orphism group of Fg that consists of all autom orphism s such that  $(t_i) = w_i^{-1} t_i w_i$  for som  $e w_i 2 F_g$  and for all generators  $t_1$ ; :::; $t_g$  of Fg. In particular, CAg contains (and for g = 1; 2 coincides with) the group of inner autom orphism sof Fg. It is easy to see that CAg is generated by the special autom orphism s  $_{ij}$  (for  $1 \quad i \in i \quad g$ ) acting on the generators of Fg by  $_{ij}(t_k) = t_k$  (resp.  $t_j^{-1}t_it_j$ ) for  $k \in i$  (resp. k = i), see K, Lem m a 2.4]. The group CAg acts on the set of F-links by changing the basing of the F-link, in other words by sending (M;  $L_b$ ; ) to for  $2 CA_g$ . An alternative and m ore geometric way to describe this action is as follows: identify the set of F-links with the set of Seifert surfaces modulo isotopy and tube equivalence, see [G L2]. There is an action of CAg on the set of Seifert surfaces, described on generators  $_{ij}$  by cocooning the ith surface to the jth surface using suitable arcs to a base point, well-de ned modulo isotopy and tube equivalence.

Since the set N (O)=h i is in 1-1 correspondence with the set of F -links and since CA acts on the latter set, it follows that it acts on the form er set, too. We now describe this action of  $CA_g$  (or rather, of its generating set f  $_{ij}$ g) on the set N (O)=h i.

Consider a link L 2 N (O). The action of  $_{ij}$  on such a link is the following: \the ith unlink component is isotoped through the jth unlink component, and then put back where it started".

Let us exam ine what happens when we try to implement such a principle. To do this, we take a parallel copy of the ith unlink component and drag it so that, at the end, the initial link (resp. the link that results from the move) is recovered by forgetting the parallel copy (resp. forgetting the original copy).



Figure 3. D ragging the ith unlink component through the jth unlink component, and then put back where it started.

It is easy to see that the  $_{ij}$  m ove does not depend on the choice of arc, m odulo K index m oves, and that the above discussion together with Theorem 1.1 in ply that

Theorem 6.1. There exists an action

which induces a 1-1 onto correspondence

In order to give a tangle description of the set of  $(P-links, we need to introduce a geometric action of the set CA<sub>g</sub> on the set D (D<sub>g</sub>). Recalling the action of CA<sub>g</sub> on the set N (O), we see that this action can be specified by a framed arc, and that as this proceeds, then all the strands going between the two copies are parallel to each other (that is, dragging the copy away from the original \lines-up" all the strands in between); see also Figure 3. This will be the idea of our action of CA<sub>g</sub> on D (D<sub>g</sub>). To form alize this, we (brie y) enlarge our notion of sliced crossed links, to allow for such an arc. That is, we will consider the set D <sup>arc</sup> (D<sub>g</sub>) of arc-decorated sliced crossed links, represented by sequences of tangles such that, in addition to all closed components, there is an arc which starts and nishes at som e meridional disc:$ 



For an element of this set, we will say that the result of a drag along the arc is the sliced crossed link that arises by making the following replacements, accompanied by taking a parallel of the arc a according to the appropriate word (in the example below, according to  $w_{b}\overline{w_{b}}$ ).



Given two sliced crossed links  $L_1$ ;  $L_2 2 D (D_g)$ , we say that  $L_2$  is obtained from  $L_1$  by an  $_{ij}$  m ove if there exists an arc-decorated sliced crossed link  $L_{12} 2 D^{arc} (D_{g+1})$  with the following properties:

- 1. there are no intersections of link components with the (i+1)st m eridianal disc,
- 2. gluing back in the (i + 1)st removed disc (and forgetting the arc) gives a diagram which presents  $L_{1}$ ,
- 3. the arc travels from the ith m eridianal disc, intersects only the jth m eridianal disc, and only once, and then ends at the ith m eridianal disc.
- 4. D ragging L<sub>12</sub> along the arc and then gluing back the ith rem oved unlink component recovers L<sub>2</sub>.

Remark 62. The above de nition implements the procedure described in Figure 3. Actually, this  $_{ij}$  move does not depend on the choice of arc (this is an instructive exercise). In addition, this move translated to the set of F-links coincides with the action of CA<sub>g</sub> by cocooning.

Proposition 2.6 and Theorem 6.1 im ply that

Proposition 6.3. There is an action

```
CA D(D<sub>g</sub>)=hr; ;!; i ! D(D<sub>g</sub>)=hr; ;!; i
```

which induces a 1-1 correspondence

D (D<sub>g</sub>)=hr; ;!; ;CAi ! @ links:

62. The action of CA on diagram s. We now de ne an algebraic action of CA<sub>g</sub> on diagram s. Recall the group CA<sub>g</sub> that acts on the free group F<sub>g</sub> (see Section 6.1), thus also on . The action of CA<sub>g</sub> on extends to the localization  $_{\rm loc}$ , since for 2 CA<sub>g</sub>, the induced ring hom om orphism !  $_{\rm loc}$  is -inverting, and thus extends to a ring hom om orphism  $_{\rm loc}$ !  $_{\rm loc}$ . This induces an action

$$CA_{q} = A^{0}(?_{X}; b_{c}) ! A^{0}(?_{X}; b_{c})$$

given by acting on the beads of diagram s and the entries of matrices by special autom orphism s of the free group. Following Remark 3.1, we denote by  $A^0$  (?<sub>x</sub>;  $l_{bc}$ )=hCAi the quotient space.

Sim ilarly, there is an action

$$CA_{q} = A^{qp;0}(?_{X}; _{loc}) ! A^{qp;0}(?_{X}; _{loc});$$

which results in a quotient space  $A^{gp;0}$  (?<sub>x</sub>;  $_{loc}$ )=hCAi, too.

The next proposition show s that the action of CA<sub>g</sub> descends to one on the quotient set A  $g^{p;0}$  (?<sub>x</sub>;  $l_{oc}$ )=h  $g^{p;0}$ ;  $g^{p;0}$ :

Theorem 6.4. The  $CA_g$  action on  $A^{gp;0}$  (?<sub>x</sub>;  $_{bc}$ ) descends to an action on  $A^{gp;0}$  (?<sub>x</sub>;  $_{bc}$ )=h  $^{gp;1}$ .

Proof. It is an exercise in the denitions to see that if  $(M; s_1); (M; s_2) \ge A^{gp;0}(?_X; _{bc})$  and and  $(M; s_1)$   $(M; s_2)$ , then  $_{ij}:(M; s_1) \xrightarrow{g_P} _{ij}:(M; s_2)$ . However, this is not so clear for the case of the wrapping move  $!^{gp}$ . Consider pairs  $(M; s_1) \xrightarrow{l_{k}^{gp}} _{ij}:(M; s_2)$  by a wrapping move at the kth site. We need to show that  $_{ij}:(M; s_1) \xrightarrow{g_P} _{ij}:(M; s_2)$ .

By de nition, there exists an s 2 A  $^{\rm gp}$  (?<sub>X [f@hg</sub>; loc) such that

$$s_1 = con_{fhg}$$
 (s) and  $s_2 = con_{fhg} = t_k ! e^{h} t_k e^{h}$  (s)  $t exp_t = \frac{1}{2} h (M^{-1} t_k ! e^{h} t_k e^{h} M)$ ;

Then, we have that

$$i_j s_1 = con_{fhg} (i_j (s))$$
 and  $i_j s_2 = con_{fhg} i_j t_k ! e^h t_k e^h (s) t exp_t \frac{1}{2} h (M^{-1} t_k ! e^h t_k e^h M)$ :

If  $k \in i$ ; j then  $_{ij}$  commutes with  $_{t_k$   $r e^h t_k e^h}$ , thus the result follows. It remains to consider the cases of k = i or k = j.

The case of k = i. Observe that ij equals to the substitution  $t_i T t_i^{-1} t_i t_i$ , thus

We will rst compare  $t_i \mathbf{\tilde{r}} e^{h} t_j^{-1} t_i t_j e^{h}$  and  $t_i \mathbf{\tilde{r}} t_j^{-1} e^{h} t_i e^{h} t_j$  using the Push Identity, and then  $t_i \mathbf{\tilde{r}} t_j^{-1} e^{h} t_i e^{h} t_j$  with the identity using the wrapping move to reach the desired conclusion. Explicitly, the Push Identity of Figure 2 in plies that

if all appearances of t and all legs (h are drawn. Let s((h) denote s with the special leg colored explicitly by (h, and use the following abbreviation

$$(N) = \exp_t \frac{1}{2} h(N)$$
 :

The above discussion implies that:

$$\begin{array}{rcl} \text{ij}\,(M\,\,;s_2) &=& \text{ij}M\,\,;\text{con}_{\text{fhg}} & {}_{t_i}\,^{\gamma}\,e^{\,h}\,t_j^{\,1}\,t_it_je^{\,h}}\,(s\,(e\,h\,))\,t &(\,\,ij\,M^{-1}\,\colon_{t_i}\,^{\gamma}\,e^{\,h}\,t_j^{\,1}\,t_it_je^{\,h}}\,(M\,\,)) \\ & \stackrel{\text{Push}}{=} & ijM\,\,;\text{con}_{\text{fhg}} & {}_{t_i}\,^{\gamma}\,t_j^{\,1}\,e^{\,h}\,t_ie^{\,h}\,t_j}\,s\,(t_j\,e\,h\,)\,t & ijM^{-1}\,\colon_{t_i}\,^{\gamma}\,t_j^{\,1}\,e^{\,h}\,t_ie^{\,h}\,t_j}\,M \\ & \stackrel{\text{I}_{i}^{\text{gp}}}{=} & ijM\,\,;\text{con}_{\text{fhg}} & {}_{t_i}\,^{\gamma}\,t_j^{\,1}\,t_it_j}\,s\,(t_j\,e\,h\,)\,t & ijM^{-1}\,\colon_{t_i}\,^{\gamma}\,t_j^{\,1}\,t_it_j}\,M \\ & = & (\,\,ijM\,\,;\,(\,\,ijs\,(t_j\,e\,h\,)\,j_{e\,h\,\gamma\,0}\,)) \\ & = & ijM\,\,;\text{con}_{\text{fhg}}\,(\,\,ij\,s\,) \\ & = & ijM\,\,;\text{con}_{\text{fhg}}\,(\,\,ij\,s\,) \\ & = & ijM\,\,;\text{con}_{\text{fhg}}\,(\,\,ij\,s\,) \end{array}$$

The case of k = j. Observe that

First, we will compare the substitutions  $^{(0)}$  with  $^{(4)}$  (dened below) using a wrapping move !  $_{j}^{gp}$ , the following --identity

 $(A^{1}B)t (B^{1}C) = (A^{1}C)$ 

(which is a consequence of Proposition 2.5) and Equation (13) of Lemma A.1.

Then, we will compare  ${}^{(4)}$  with  ${}^{(8)}$  using the wrapping move  $!\,{}^{\rm gp}_{i}$  .

Finally, we will compare (8) with the identity using the case of k = i above. Here, (a) denote the substitutions:

(0) =	tj 7 e <sup>h</sup> tje <sup>h</sup> ;	t <sub>i</sub> 7 t <sub>j</sub> <sup>1</sup> t <sub>i</sub> t <sub>j</sub> ;
(2) =	$t_j 7 e^{h} e^{h^0} t_j e^{h^0} e^h$ ;	$t_i 7 e^{h^0} t_j^{-1} e^{-h^0} t_i e^{h^0} t_j e^{-h^0};$
(3) =	$t_j 7 e^{h^0} t_j e^{h^0};$	
(4) =	t <sub>j</sub> 7 t <sub>j</sub> ;	t <sub>i</sub> 7 e <sup>h</sup> t <sub>j</sub> <sup>1</sup> e <sup>h</sup> t <sub>i</sub> e <sup>h</sup> t <sub>j</sub> e <sup>h</sup> ;
(6) =	t <sub>j</sub> 7 t <sub>j</sub> ;	$t_i $ ? $e^h t_j^{1} e^h e^{h^0} t_i e^{h^0} e^h t_j e^h$ ;
(7) =	$t_i$ 7 $e^{h^0}t_i e^{h^0}$ ;	
(8) =	tj 7 tj;	t <sub>i</sub> 7 e <sup>h</sup> t <sub>j</sub> <sup>1</sup> t <sub>i</sub> t <sub>j</sub> e <sup>h</sup> :

Then, we have that

6.3. Z  $^{\rm rat}$  com m utes with the CA action. Recall the invariant Z  $^{\rm rat}$  of sliced crossed links of Section 4.

Theorem 6.5. The map

$$Z^{\text{rat}}:D_X(D_q)! A^{\text{gp}}(?_X;)=h!^{\text{gp}}i$$

is CAg-equivariant, and induces a well-de ned map

$$Z^{\text{rat}}:D_X (D_g)=hCA_gi! A^{gp}(?_X;)=h!^{gp};CA_gi:$$

Proof. W e will show that for D  $2 D_X$  (D<sub>g</sub>), we have

$$ij Z^{rat}(D) = Z^{rat}(ij D):$$

G iven sliced crossed links  $L_1; L_2$  such that  $L_2 = _{ij} L_1$ , consider an accompanying link  $L_{12}$  as in the previous section.  $L_{12}$  may be presented so that all the strands going through the ith meridianal disc proceed, parallel to each other, through the jth meridianal disc, and then onto the (i + 1)st meridianal disc. That is, we can nd a sequence of tangles, containing some arc 'x' (not exactly the arc 'a' of Section 2.4), such that the link  $L_{12}$  is presented by taking a parallel the arc 'x' according to som e word (in the example at hand, w<sub>b</sub>). Making the following de nitions  $T_p; T_q; T_r; T_s; T_t; T_u$  respectively



and continuing to work with the example of the above paragraph, then the sequence will bok as follows (where the jth, the ith and then the (i + 1)st m eridianal discs are displayed):

$$f:::; \quad \stackrel{w_{b}}{x} (T_{p}); \quad \stackrel{w_{b}}{x} (T_{q});:::; \quad \stackrel{w_{b}}{x} (T_{r}); T_{s};:::; T_{t}; \quad \stackrel{w_{b}}{x} (T_{u});:::g$$

Now, the link  $L_1$  (resp.  $L_2$ ) is presented by the diagram s recovered by gluing back in the (i + 1)st (resp. ith) rem oved disc.

So, on the one hand (and the reader is advised to follow the following compositions on paper):

$$Z^{\text{rat}}(L_{1}) = \begin{pmatrix} w_{\text{b}}(Z(T_{\text{p}})) & (I_{\text{t}_{d}} & G_{j;w_{\text{b}}w_{e}}) & w_{\text{b}}(Z(T_{\text{q}})) & ::: \\ & w_{\text{b}}(Z(T_{\text{r}})) & (I_{\text{t}_{a}} & G_{i;w_{\text{b}}}) & Z(T_{\text{s}}) & tZ(T_{\text{s}}) & tZ(T_{\text{s}}) & tZ(T_{\text{s}}) & tZ(T_{\text{s}}) & \vdots:: \\ & w_{\text{b}}(Z(T_{\text{p}})) & (I_{\text{t}_{d}} & G_{j;w_{\text{b}}w_{e}}) & (I_{\text{t}_{d}} & G_{i;w_{\text{b}}} & I_{w_{e}}) & w_{\text{b}}(Z(T_{\text{q}})) & ::: \\ & w_{\text{b}}(Z(T_{\text{r}})) & Z(T_{\text{s}}) & tZ(T_{\text{s}}) & tZ(T_{\text{s}}) & \vdots: \end{pmatrix}$$

A  $!^{gp}$ -m ove has been used here to move the gluing word G<sub>i,w</sub> up to (just underneath) the gluing word arising from the jth meridional slice. This is legitim at because the meaning of the  $!^{gp}$ -m ove is that such an element (if it involves every appearance of  $t_i$ ) can be slid along a paralleled arc (here arising from the component x).

0 n the other hand:

$$Z^{\text{rat}}(L_{2}) = \begin{cases} w^{\text{b}}_{\text{x}}(Z(T_{p})) & (I_{w_{d}} G_{j_{jw_{b}w_{e}}}) & w^{\text{b}}_{\text{x}}(Z(T_{q})) & ::: \\ & w^{\text{b}}_{\text{x}}(Z(T_{r})) & Z(T_{s}) & tZ(T(I_{w_{d}}^{W_{b}} G_{j_{jw_{b}}}) & w^{\text{b}}_{\text{x}}(Z(T_{u})) & ::: \\ & w^{\text{b}}_{\text{x}}(Z(T_{p})) & (I_{w_{d}} G_{j_{jw_{b}}} I_{w_{e}}) & (I_{w_{d}} G_{j_{jw_{b}w_{e}}}) & w^{\text{b}}_{\text{x}}(Z(T_{q})) & ::: \\ & w^{\text{b}}_{\text{x}}(Z(T_{r})) & Z(T_{s}) & tZ(T^{w_{b}}_{\text{x}}(Z(T_{u})) & ::: \end{cases}$$

Observe that this line results from the line ending the previous paragraph of equations by the algebraic action of  $_{ij}$ , as is required for the proof.

6.4.  $R_{rat}$  com mutes with the CA<sub>g</sub> action. The following theorem says that the action of CA<sub>g</sub> on diagram s com mutes with  $R_{rat}$ -integration.

Theorem 6.6. The map

Z<sub>rat</sub> dX<sup>0</sup>: Int<sub>x °</sub>A<sup>gp;0</sup> (?<sub>x</sub>;  $_{loc}$ ) ! A<sup>gp;0</sup> (?<sub>x x °</sub>;  $_{loc}$ ) is CA<sub>q</sub> equivariant and induces a well-de ned map:

Z rat  
dX <sup>0</sup>: Int<sub>x 
$$\circ$$</sub> A <sup>gp;0</sup> (?<sub>x</sub>; <sub>bc</sub>)=hCAi! A <sup>gp;0</sup> (?<sub>x x  $\circ$ ; <sub>bc</sub>)=hCAi</sub>

Proof. Let the canonical decomposition of s 2 Int<sub>X</sub>  $_{\circ}A^{gp}$  (?<sub>X</sub> ;  $_{bc}$ ) with respect to X  $^{0}$  be

$$s = \exp \frac{1}{2} \frac{X}{\underset{k_{1} \\ k_{1} \\ x_{1}}} M_{k_{1}} t R;$$

for some matrix M 2 Herm (  $_{loc}$  ! Z). Then, the canonical decomposition of  $_{ij}$  s with respect to X  $^0$  is

Observe that ( $_{ij}M$ ) <sup>1</sup> =  $_{ij}M$  <sup>1</sup> 2 Herm ( $_{loc}$ ! Z) which implies that  $_{ij}s$  is X <sup>0</sup>-integrable. Furthermore, we have that

$$Z_{rat} = X_{rat} = X_{r$$

Note that we could have used Theorem E.1 to deduce the above equality.

W e end this section with the following

Lem m a 6.7. (a) For all g, the identity m ap A ( $?_X$ ;  $_{loc}$ ) ! A ( $?_X$ ;  $_{loc}$ ) m aps the  $Inn_g$ -relation to the basing relation.

(b) For g = 1;2, the identity map A ( $?_X$ ;  $_{loc}$ ) ! A ( $?_X$ ;  $_{loc}$ ) maps the  $CA_g$ -relation to the basing relation.

 $P \, \text{roof.} R$  ecall that  $\text{Inn}_g$  is the subgroup of CA  $_g$  that consists of inner autom orphism s of the free group F .

For the rst part, an inner autom orphism by an element  $g \ 2 \ F$  m aps a bead  $b \ 2 \ _{loc}$  to  $g \ ^{1}bg \ 2 \ _{loc}$ . The relations of Figure 1 cancel the g at every trivalent vertex of a diagram. The remaining vertices change a diagram by a basing move  $\frac{gp}{2}$ .

The second part follows from the fact that  $Inn_q = CA_q$  for g = 1;2, see [CS2, K].

6.5. The de nition of the rational invariant Z rat of Q-links. .

Theorem s 61, 63, 65 and 6.6 im ply that there is a map

 $Z^{rat;gp}: @ links! A^{gp;0}(_{bc})=h!^{gp}i$ 

de ned as follows: for a (-link (M; L)) of g components, choose a sliced crossed link C in D (D<sub>g</sub>) whose components are in 1-1 correspondence with a set X, and de ne

$$Z^{\text{rat};\text{gp}}(M;L) = dX \quad Z^{\text{rat}}(C) \quad c^{+(B)}c^{-(B)}c^{-(B)}2A^{\text{gp};0}(_{\text{loc}})=h!^{\text{gp}};CAi$$

where  $c = {}^{R} dUZ (S^{3};U)$  are some universal constants of the unit-fram ed unknot U and where B is the linking matrix of C and (B) is the number of positive (resp. negative) eigenvalues of B. The above de nition makes sense and uses the fact that as in Section 5.5, the set  $A^{gp;0}(_{bc})=h! g^{gp}$  is a well-de ned action of the group  $A^{gp}()$ , which commutes with the CA action (where CA acts trivially on  $A^{gp}()$ ).

C on posing the Z<sup>rat,gp</sup> invariant with the projection  $A^{gp}$ ,<sup>0</sup> ( $_{bc}$ )=h! <sup>gp</sup>; CAi!  $A^{0}$ =h!; CAi of P roposition 3.9, de nes the rational invariant Z<sup>rat</sup> of  $\ell$ -links that appears in Theorem 1.3.

#### 7. A comparison between Z and Z<sup>rat</sup>

The ultim ate goal of this section is to compare the invariants Z<sup>rat</sup> and Z of F - inks and to nigh the proof of Theorem 1.3. This will be obtained by comparing the construction of the A arhus integral with that of  $R^{rat}$ .

7.1. The Hairm ap. First, we need to de neam ap Hair from the space of diagram s that Z <sup>rat</sup> takes values to the space of diagram s that Z takes values. G iven generators  $ft_1; :::;t_g g$  of the free group, let  $fh_1; :::;h_g g$  be noncommuting variables such that  $t_i = e^{h_i}$  and let Hair: A (?<sub>X</sub>; )! A (?<sub>X [H</sub>) be de ned as follows: replace a variable by the exponential of hair, as in Figure 4. This extends to a map Hair: A (?<sub>X</sub>; `)! A (?<sub>X [H</sub>) be de ned as follows: replace (?<sub>X [H</sub>)) (and, after composition with A (?<sub>X ; bc</sub>)! A (?<sub>X ; `</sub>)), also to a map

whose in age lies in the span of all diagram s that do not contain a H -labeled tree as a connected component.

"j t<sub>i</sub> ! 
$$\frac{X^{i}}{n=0} \frac{1}{n!}$$
 n h<sub>i</sub>-labeled legs

Figure 4. From beads to legs.

Recall the extension  $A^{gp;0}$  of  $A^{gp}$ . G iven a pair (M;s) where M = 2 Herm ( $_{loc} ! Z$ )  $A^{gp}$ , we can de ne an extended m ap Hair:  $A^{gp;0}$  (?<sub>X</sub>;  $_{loc}$ ) !  $A^{gp}$  (?<sub>X [H</sub>) by

$$Hair(M;s) = exp_t \frac{1}{2} h(M) t Hair(s):$$

Note that the  $exp_{t}$  term in the above expression is a disjoint union of wheels.

Lem m a 7.1. There exist m aps that t in a commutative diagram

$$A^{gp;0} (\sim_{X}; _{bc})=h!^{gp}; CA_{gi} \longrightarrow A^{0} (\sim_{X}; _{bc})=h!^{gp}; CA_{gi}$$

$$Hair$$

$$A (\sim_{X[H]})$$

Proof. Notice that  $_{h}$  (M) depends only on the equivalence class of M 2 B ( $_{loc}$  ! Z); see Proposition 2.5. It is easy to show that the X – avored group-like and in tesim albasing relations (de ned in Section D) are mapped by the Hairm ap to X – avored in nitesim albasing relations. It is also easy to see that the ! <sup>gp</sup> and CA<sub>g</sub> relations are mapped to H – avored in nitesim albasing relations.

Remark 7.2. The invariant Z<sup>rat</sup> takes values in A  $^{\rm gp;0}$  ( $_{\rm bc}$ )=h!  $^{\rm gp}$ i, the invariant Z takes values in A ( $_{\rm H}$ ) and the image of the map Hair : A  $^{\rm gp;0}$  ( $_{\rm bc}$ )=h!  $^{\rm gp}$ i ! A ( $_{\rm H}$ ) lies in the subspace of A ( $_{\rm H}$ ) spanned by diagram s without tree components. If we expect Z  $^{\rm rat}$  to determ ine Z, we should restrict attention to the class of links whose K ontsevich integral has vanishing \tree part". Since the vanishing of the tree-part is equivalent to the vanishing of all M ilnor's invariants, we should only consider links with vanishing M ilnor's invariants. Such links include, and perhaps coincide with, all sublinks of hom obgy boundary links, see for example [Le2]. For simplicity, we will not deal with hom obgy boundary links, but rather with boundary links.

We will need a slightly renorm alized version H air of the H air map, obtained by considering the disjoint union of the map H air and  $(h_1)$  t  $t_g)$  (hw here  $= Z (S^3; U)$ .

72. A comparison between  $Z^{rat}$  and Z. Consider the set  $L_X$  (O) of links L [O in  $S^3$  where O is an unlink whose components are in 1-1 correspondence with the set  $H = fh_1; :::;h_gg$  and L is a link whose components are in 1-1 correspondence with a set X. De ne a map

that m aps the associated link L of a sliced crossed link in  $D_g$  I to the link L [O S<sup>3</sup> according to a standard inclusion of  $D_g$  I in S<sup>3</sup>, by adding a g component unlink O, one component for each gluing site of  $D_g$  as follows:

$$\begin{array}{c} & & \\ & &$$

Consider in addition the Kontsevich integral

where A (~  $_{X [H}$ ) denotes the space of unitrivalent graphs whose legs are colored by X [ H, modulo the AS; IHX relations and the in nitesimal basing relations discussed in Appendix C. The purpose of this section is to show that

Theorem 7.3. The following diagram commutes

The proof will utilize the following magic formula for the Kontzevich integral of the Long Hopf Link, conjectured in [A 0] (in conjunction with the so-called W heels and W heeling conjectures) and proven in [B LT].

Theorem 7.4. [BLT]

$$f_{x,yg} \quad Z \quad \stackrel{x}{\underset{|}{\bigcirc}} \begin{array}{c} y \\ y \end{array} = \exp \left| \begin{array}{c} x \\ y \end{array} \right|_{y} t \quad (y) \ 2 \ A \ (?_{fxg} \sim fyg):$$

Proof of Theorem 7.3. We will show rst that the following diagram commutes:

$$\begin{array}{c} D_{X} (O) \xrightarrow{Z^{rat}} A^{gp} ("_{X}; ) \\ \uparrow [ \\ L_{X} (O) \xrightarrow{Z} A (\sim_{X [H]}) \end{array}$$

(4)

Given a word w in the symbols " and #, let

LongH op  $f_{j,w}$  denote the long H opf link cabled according to the word w, with the closed component labeled  $h_j$ . For example:

LongH opf<sub>j;#""</sub> = 
$$\begin{pmatrix} & & & \\ & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & & \\ &$$

 $G_{inv}^{h}$  denote the jth hairy gluing word corresponding to w. This is de nition by example:

$$G_{j;\#""}^{h} = \vec{\mu} e^{h_{j}} \vec{j} e^{h_{j}} \vec{j} e^{h_{j}}$$

W e use the following corollary to Theorem 7.4.

Corollary 7.5.

$$fh_{jg}$$
 Z (LongH opf<sub>jw</sub>) =  $G_{jw}^{h}$  t (h<sub>j</sub>) 2 A ("<sub>w</sub> ~  $fh_{jg}$ ):

The commutativity of diagram (4) now follows by the denition of Z<sup>rat</sup> Corollary 7.5. Theorem 7.3 follows from the commutativity of diagram (4), after considering the quotient N<sub>X</sub> (O) of D<sub>X</sub> (O) (given by Proposition 2.6) and symmetrizing (i.e., composing with  $:A^{gp}("_X;)! A^{gp}(\sim_X;)$ ) together with the fact that Hair commutes with composition.

7.3. A comparison between  $\frac{R_{rat}}{R}$  and  $\frac{R}{R}$ . Recall that the A arhus integral respects the group-like basing relations as shown in A, Part II, Proposition 5.6] and also the in nitesim albasing relations as was observed by D. Thurston, see BL, Proposition 2.2]. In other words, there is a well-de ned m ap

$$dX^{0}$$
: Int<sub>x</sub>  $\circ$ A (~ <sub>X [H</sub>) ! Int<sub>x</sub>  $\circ$ A (~ <sub>(X X  $\circ$ ) [H</sub>);

where  $H = fh_1; \dots; h_q g$ . The purpose of this section is to show that

Theorem 7.6. The following diagram commutes

$$\begin{array}{cccc} \operatorname{Int}_{X} \circ A \overset{\operatorname{gp};0}{\operatorname{gp}} (?_{X} ; _{\operatorname{loc}}) & \xrightarrow{\operatorname{R}_{\operatorname{rat}} dX ^{\circ}} A \overset{\operatorname{gp};0}{\longrightarrow} (?_{X} _{X} \circ ; _{\operatorname{loc}}) \\ & & & & & \\ & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & \\ & & & & & \\ & & & & & \\ & & & & &$$

Proof. It follows from Remark E 2 of Appendix E.

7.4. End of the proof of Theorem 1.3. In this section we show that the diagram of Theorem 1.3 commutes. We begin by observing that the following diagram commutes, by the de nition of the invariant  $Z^{rat;gp}$ :

F links 
$$\xrightarrow{Z^{rat;qp}} A^{gp;0} (_{bc})=h!^{gp}i$$
  
 $\downarrow$   $\downarrow$   $\downarrow$   
 $\emptyset$  links  $\xrightarrow{Z^{rat;qp}} A^{gp;0} (_{bc})=h!^{gp};CAi$ 

We now claim that the following diagram commutes:

$$\begin{array}{cccc} 0 & links \xrightarrow{Z \text{ rat;gp}} A \text{ gp;}^0 (_{\text{loc}})=h! \text{ gp;}CA i \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ H \text{ air} \\ 0 & links \xrightarrow{Z} A (\sim_H) \end{array}$$

(5)

The above two diagrams, together with Lemma 7.1 in ply the commutativity of the diagrams of Theorem 1.3. In order to show the commutativity of (5), we will assem ble several commutative diagrams in one with two layers. The top layer consists of the rational invariants Z<sup>rat</sup> and its integration theory  $R_{rat}$ ; the bottom layer consists of the LM O -A arhus-K ontsevich integral Z and its integration theory .



Theorem s 4.1, 7.3 and 7.6 in ply that all faces except of the above cube except the side right one commute. Since the map  $N_X$  (O) !  $L_X$  (O)=h ;CAi is onto, it follows that the remaining face of the cube commutes. U sing the identications A ( \_ H ) = A (~ H ) (see Proposition D 1), and

$$L_x$$
 (O)= $h_x$ ; CAi=  $N_x$  (O)= $h_x$ ; CAi= @ links;

(see Proposition 2.6) it follows that diagram (5) commutes. This concludes the proof of Theorem 1.3.

#### 8. A universal property of Z<sup>rat</sup>

The goal of this section is to show that the rational invariant Z<sup>rat</sup> of F-links of Theorem 1.3 has a universal property, namely that it is the universal  $l_{loc}$ -valued nite type invariant of F-links with respect to the loop

Itration, [G a]. We begin by recalling the notion of nite type invariants on the set of F-links introduced at [G a]. Given an F-link (M;L; ), where :  $_1$  (M L)! F, we will call a link C 2 M L good relative to (M;L; ) if each component of C is mapped to 1 under . Notice that surgery under a good link preserves the underlying F-structure. Given a good link C relative to an F-link (M;L), choose a sliced crossed link L<sup>0</sup> [C<sup>0</sup> 2 D (D<sub>g</sub>) such that after surgery on L<sup>0</sup> S<sup>3</sup> O it gives rise to the F-link (M;L) with C<sup>0</sup> S<sup>3</sup> O transform ed to a good link C relative to (M;L). We de ne

$$Z^{\text{rat};\text{gp}}(\mathcal{M};L);C) = dX (Z^{\text{rat}}(L^{0}[C^{0}]) = (\text{norm alization}) 2 A^{\text{gp};0}(\sim_{Y}; _{\text{loc}}) = h!^{\text{gp};1}$$

$$Z^{\text{rat};\text{gp}}(\mathcal{M};L)_{C} = dY (Z^{\text{rat}}(\mathcal{M};L);C)) = (\text{norm alization}) 2 A^{\text{gp};0}(\sim; _{\text{loc}}) = h!^{\text{gp};1}$$

where X (resp. Y) is in 1-1 correspondence with the components of L (resp. C), and where (norm alization) refers to factors inserted to deal with the K inby move of adding a distant union of unit-fram ed unlinks, as in Section 5.5. Theorem 1.3 has the following relative version (as is revealed by its proof in Section 5.5):

Theorem 8.1. There exists an invariant  $Z^{rat;gp}$ : F links [good C !  $A^{gp;0}$  (~  $_{C}$ ;  $_{loc}$ )=h! i which is local (i.e., compatible with respect to considering sublinks of C) and ts in a commutative diagram

F links [good C<sup>Z</sup> 
$$\xrightarrow{\text{rat},\text{gp}}$$
 A  $\xrightarrow{\text{gp};0}$  (~  $_{\text{Y}}$  ;  $_{\text{loc}}$ )=h!  $\xrightarrow{\text{gp}}$  i  
c surgery  
F links  $\xrightarrow{Z^{\text{rat},\text{gp}}}$  A  $\xrightarrow{\text{gp};0}$  ( $_{\text{loc}}$ )=h!  $\xrightarrow{\text{gp}}$  i

As in Section 5.5, we denote by Z<sup>rat</sup> the composition of Z<sup>rat;gp</sup> with the map  $A^{gp;0}$  (~<sub>Y</sub>; <sub>loc</sub>)=h! <sup>gp</sup>i!  $A^{0}$  (~<sub>Y</sub>; <sub>loc</sub>)=h! i.

Given a good link C relative to (M;L), one can de ne an equivariant linking matrix  $\mathbb{k}_{(M;L)}(C)$ , by considering the linking numbers of the components of the lift  $\mathcal{C}$  of C in the free cover of M L.  $\mathbb{k}_{(M;L)}(C)$  is a H erm itian matrix over  $_{loc}$ , well de ned up to conjugation by a diagonal matrix with elements in F. The details of the de nition, together with the following proposition will appear in a subsequent publication, [G a2]:

Proposition 8.2. For C and M;L) as above, the covariance matrix cov of Z rat satis es

$$\operatorname{cov}(\mathbb{Z}^{\operatorname{rat}}(\mathbb{M};\mathbb{L});\mathbb{C})) = \mathbb{k}_{\mathbb{M};\mathbb{L}}(\mathbb{C}):$$

Proof. (of Theorem 1.12) Recall rst that we consider the A  $^0$ -valued invariant Z  $^{rat}$  rather than the A  $^{gp;0}$ -valued invariant Z  $^{rat;gp}$ .

For part (a), the matrix-part of Z<sup>rat</sup> lies in the set B ( $_{loc}$ ! Z). In [G a] it was shown that two F -links are 0-equivalent (i.e., equivalent under a sequence of loop moves) i they have isom orphic B lanch eld pairings. Proposition 1.2 implies that the B lanch eld pairing is determined by the matrix part of Z<sup>rat</sup>. The result follow s.

For part (b), the proof is similar to the case of knots, given in [GR, Section 3]. For the convenience of the reader, we will give the proof in a self-contained manner, using occasional notation of [GR].

First, we explain why it makes sense to separate the matrix-free part of  $Z^{rat}$ . Surgery on a good Y-graph G relative to an F-link (M;L) gives an S-equivalent F-link, as was shown in [Ga]. In this case, Remark 1.6

in plies that we can x the matrix-part of Z<sup>rat</sup> and regard its matrix-free part as an element of the graded vector space A ( $_{loc}$ )=(!). In the rest of the proof, Z<sup>rat</sup> refers to this invariant.

Next, we need to recall some properties of surgery on good Y -graphs. Consider a good Y -graph G relative to an F -link (M ;L). Recall that surgery on G corresponds to surgery on a six component link partitioned in three blocks  $A_i$  of two components each, for i = 1;2;3. Each  $A_i$  consists of an edge and a leaf. A fler choosing appropriate sliced crossed links, it follows that the covariance m atrix of G relative to (M ;L), and its inverse are given by

0 I and W I I W I 0

respectively, where  ${\tt W}_{-}$  is the covariance  ${\tt m}$  atrix of the leaves of {\tt G} .

We now claim that the degree 2n part of Z<sup>rat</sup> is an invariant of F-links of loop type 2n. Fix an F-link (M; L), and let G be a good Y-graph of degree 1. A key feature of surgery on G is the fact that surgery on any proper subset of the set of arms does not alter (M; L). In other words, alternating (M; L) with respect to surgery on G equals to alternating (M; L) with respect to surgery on all nine subsets of the set  $fA_1; A_2; A_3g$ . Due to the locality property of the rational invariant  $Z^{rat}$  (i.e., the fact that  $Z^{rat}((M; L); C)$ ) determ ines  $Z^{rat}((M; L); C^{(0)})$  for  $C^{(0)}$  a sublink of C), the nontrivial contributions to  $Z^{rat}((M; L); G) = Z^{rat}((M; L); A_1; A_2; A_3g)$  come from the part of  $Z^{rat}((M; L); A_1; A_2; A_3g)$  that consists of diagram swith legs on  $A_1$  and on  $A_2$  and on  $A_3$ . Here, the [; ] notation, borrowed from the theory of nite type invariants, means that

$$[M;B] = \begin{bmatrix} X \\ B^{\circ} & B \end{bmatrix} (1)^{jB^{\circ}} M_{B^{\circ}}:$$

Now, let G be a disjoint of  $2n + 1 \mod Y$ -graphs, each of degree 1, and let A denote its set of arm s. W hen we compute Z<sup>rat</sup> ([M;L);G]) = Z<sup>rat</sup> ([M;L);A]), we need to concentrate on all diagram s D (every connected component of which has at least one trivalent vertex) with legs on each arm, and glue pairwise the legs on the arm s. Thus, the diagram s D with nonzero contribution should have at least 3(2n + 1) + 1 = 6n + 4 A-colored legs, to be glued pairwise. Thus, at most three A-colored legs meet at a trivalent vertex, and after gluing we will obtain a trivalent graph D<sup>0</sup> with at least (6n + 4)=3 = 2n + 4=3 trivalent vertices, in other words of degree at least 2n + 2. This im plies that the degree 2n part of Z<sup>rat</sup> is a loop-type 2n invariant with values on A<sub>2n</sub> ( $_{loc}$ )=(!). Note that in [GR, Section 3], this was called counting above the critical degree.

We next compute  $Z_{2n}^{rat}([M;L);G]) \ge A(_{loc})=(!)$  for good Y-graphs G of degree 2n; this was called counting on the critical degree in [GR]. The contributions to  $Z_{2n}^{rat}$  come from contractions of a disjoint union of diagrams of the form Y glued using the negative inverse covariance matrix of G. Due to the form of that matrix, as discussed above, and just as in [GR], the only contributions to  $Z_{2n}^{rat}$  come from a disjoint union  $= [_{2n} Y \text{ of diagram s in } 1-1 \text{ correspondence with the components } G_i \text{ of } G$ , such that the legs of each component of are labeled by the three edges of the corresponding component of G, and such that the legs of are glued using the covariance matrix of G.

We now recall the notion of a universal nite type invariant on the set of F -links with respect to the loop move, following [Ga]. The following de nition is most suitable for our purposes:

De nition 8.3.

U:F links! A 
$$(_{loc})=(!)$$

is a universal nite type invariant i for every n 1 and every F -link (M;L) and every G good with respect to (M;L) of degree 2n, we have that

$$U_{2n} ([(M;L);G]) = \exp \frac{1}{2} \cos (M;L) (Leaves(G)) ; G^{0} 2 A_{2n} (L_{bc}) = (!)$$

where  $G^0$  is G the unitrivalent graph G m inus the leaves of G. In other words, the sym  $bolU_{2n}$  ([M;L);G]) of U is given by the sum of contractions of the legs of G using the negative equivariant linking m atrix of the leaves of G.

Rem ark 8.4. Note the resemblance with perturbative quantum eld theory, where the vertex is the graph G and the propagator is  $cov_{(M;L)}$  (Leaves(G)).

It follows from the above discussion and de nition that  $Z_{2n}^{rat}$  is a universal nite type invariant for n 1.

#### Appendix A. A brief review of diagrammatic calculus

A 1. Some useful identities. In this section we establish some notations and collect some lemm as. Let X denote some nite set. Given a sum of diagram ss with legs colored by a set X, and a subset X<sup>0</sup> =  $fx_1; :::;x_rg$  of X, we will often denote s by  $s(x_1; :::;x_r)$  or simply by s(x). This notation allows us to extend linearly to diagram swhose legs are labeled by form al linear combinations of elements of X. For example, for a; b2 Q, v; w 2 X,  $s(:::;x_{i-1};av + bw; x_{i+1};:::)$  denotes the element obtained from s by replacing each diagram appearing in s by the sum of all diagram s obtained by relabeling each leg labeled  $x_i$  by either v (with a multiplicative factor of b). Given diagram s A (x); B (x), we let

 $hA(@x);B(x)i_x$  and  $A(@x)[_XB(x)$ 

denote respectively the sum of all ways of pairing all X -colored legs of A with all (resp. som e) X -colored legs of B.

Furtherm ore, for some q 2  $_{bc}$ , s(:::;x<sub>i</sub> 1;x<sub>i</sub>q;x<sub>i+1</sub>;:::) denotes the element obtained by pushing a bead labeled by q onto each leg labeled x<sub>i</sub> in the sense speci ed by the following diagram :

Taken together, these conventions give meaning to such expressions as s(xM), for M = 2 Herm  $(_{bc} ! Z)$ .

The following kmmma collects a number of useful identities, which the reader would be well-advised to understand.

Lem m a A .1. (a) Let A 2 A ( $?_X$ ;  $_{bc}$ ); B 2 A ( $?_{X[Y}$ ;  $_{bc}$ ) (for Y a bijective copy of X), and let X  $^0$  and X  $^{00}$  denote bijective copies of X.

(6) hA (@x); B (x; y) 
$$\dot{\mathbf{x}} = \mathbf{x} \mathbf{i}_{\mathbf{x}} = \mathbf{h} \mathbf{A} (\mathbf{e} \mathbf{x}^{0} + \mathbf{e} \mathbf{x}^{0}); \mathbf{B} (\mathbf{x}^{0}; \mathbf{x}^{0}) \dot{\mathbf{x}}_{\mathbf{x}} \mathbf{i}_{\mathbf{x}} \mathbf{i}_{\mathbf{x}} \mathbf{i}_{\mathbf{x}}$$

(b) Let 
$$A_1; A_2; B 2 A (?_X; bc)$$
.

(7) 
$$hA_1(@x)A_2(@x);B(x)i_x = hA_2(@x);A_1(@x)[_x B(x)i_x :$$

(c) Let A; B 2 A ( $?_X$ ; loc); and let Y be a bijective copy of X.

(8) 
$$A (@x) [_X B (x) = hA (@y); B (x + y)i_Y$$

(d) Let  $A = \exp(a) 2 A$  (?<sub>X</sub>;  $_{loc}$ ) and  $B = \exp(b) 2 A$  (?<sub>X</sub>;  $_{loc}$ ). Then

(9) 
$$\exp(a) \left[ x \circ \exp(b) = \exp(c) \right];$$

where c is the sum of all connected diagrams that arise by pairing all legs labeled from X  $^0$  of som e diagram (not necessarily connected) appearing in A to som e legs of som e diagram (not necessarily connected) appearing in B.

(e) Let A 2 A (?x ;  $_{\rm loc}$ ), let Y be a bijective copy of X , and let M 2 H erm (  $_{\rm loc}$  ! Z).

(10) 
$$\overset{\text{(ex)}}{\underset{y_{i}}{\overset{\text{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}}{\overset{(ex)}{\overset{(ex)}}{\overset{($$

(f) In particular,

(11) 
$$\exp \begin{vmatrix} e^{x} \\ e^{x} \end{vmatrix} = B(y)$$
  
(12) 
$$\exp \begin{vmatrix} e^{x} \\ e^{x} \\ e^{x} \end{vmatrix} = \exp \begin{pmatrix} X \\ e^{x} \\ e^{x} \end{vmatrix} = \exp \begin{pmatrix} X \\ e^{x} \\ e^{x}$$

(g) In addition, if  $s (@z) 2 A (?_{X [ fh^{\circ};h^{\circ\circ};@zg}; _{bc})$  then (13)  $con_{fh^{\circ};h^{\circ\circ}g} (s (@h^{\circ} + @h^{\circ\circ})) = con_{fhg} (s (@h) j_{h^{\circ};h^{\circ\circ}g} h);$
A 2. Iterated Integration. This section proves an Iterated Integration property of the  $R_{rat}$  integral. Fix labeling sets X  $^{0}$  X  $^{0}$  X.

Lem m a A .2 (Iterated integration). Given s 2 Int<sub>x</sub>  $\circ$ A (?<sub>x</sub> ;  $_{\rm loc}$ ) \ Int<sub>x</sub>  $\circ$ A (?<sub>x</sub> ;  $_{\rm loc}$ ), then, Z  $_{\rm rat}$ 

P roof. D enote the components of X  $^{0}$  and X  $^{00}$  by  $a_{i}$  and  $b_{j}$  respectively. Consider the covariance m atrix of s with respect to X  $^{0}$ , presented with respect to this basis:

$$M = \begin{array}{cc} A & C \\ C & B \end{array}$$

where  $A = A^{?}$  and  $B = B^{?}$ , where here and below  $A^{?}$  denotes the transpose of C, followed by the conjugation  $t_{i} ! t_{i}^{1}$ . If the canonical decomposition of s with respect to  $X^{0}$  is then,

$$\exp^{0} \frac{1}{2} \sum_{a_{i}}^{a_{j}} A_{ij} + \frac{1}{2} \sum_{b_{i}}^{b_{j}} B_{ij} + \sum_{b_{i}}^{a_{j}} C_{ij}^{A} t R;$$

(suppressing the sum m ations) then the canonical decomposition of s with respect to X  $^{(0)}$  is

$$\exp \frac{1}{2} \int_{a_{i}}^{a_{j}} A_{ij} t^{0} \exp \frac{1}{2} \int_{b_{i}}^{b_{j}} B_{ij} + \int_{b_{i}}^{a_{j}} C_{ij}^{A} t R^{A}:$$

Using the identities of Lemma A.1, we have

$$Z_{rat} = \exp^{Q} \frac{1}{2} \begin{bmatrix} e_{a_{j}} \\ i_{j} \\ e_{a_{1}} \end{bmatrix}^{A} i_{j} \\ A_{ij} \\ A_{$$

It is easy to see that if M and A are invertible over Z, then so is B  $C^{2}A^{-1}C$  over Z, see [A, Part II, Proposition 2.13]. Thus, the rst part of the theorem follows. Continuing,

The identication of the term on the left of the pairing with  $\exp \begin{pmatrix} 0 & x_j \\ 1 & y \end{pmatrix} M_{ij}^{-1} A$  completes the proof.  $\Box$ 

0

1

Remark A 3. W ith the notation of the previous lemma, the matrices A (B C<sup>2</sup>A<sup>1</sup>C) and M are congruent by a product of elementary matrices, as follows from the following identity

 $\begin{array}{cccc} A & 0 \\ 0 & B & C^{?}A^{-1}C \end{array} = \begin{array}{cccc} I & AC^{?} & ? & A & C & I & AC^{?} \\ 0 & I & C^{?}B & 0 & I \end{array}$ 

and from the fact that the conjugating matrix is a product of elementary matrices. It follows from this, that the Iterated Integration Lemma A 2 is valid on A  $^{gp;0}$  (?<sub>x</sub>;  $_{loc}$ ).

A 3. Integration by parts. Fix labeling sets X and X<sup>0</sup> X. Recall from Section 5.1 the notion of X<sup>0-</sup> substantial diagram s, that is diagram s that do not contain a strut labeled from X<sup>0</sup> [ @X <sup>0</sup>. The divergence of an X <sup>0-</sup> substantial diagram (extended by linearity to sum s of diagram s) with respect to X <sup>0-</sup> X is de ned to be the sum of diagram s obtained by pairing all legs labeled from  $@_{X^0} w$  it homes are legs labeled from X <sup>0-</sup>.

Lem m a A .4 (Integration by Parts). Let s be an X  $^{0}$ -substantial diagram and t 2 Int<sub>x  $^{0}$ </sub>A (?<sub>x</sub>;  $_{loc}$ ). Then

$$s \begin{bmatrix} x t & 2 & \text{Int}_{X} \circ A (?_{X}; _{bc}) \\ Z_{rat} & Z_{rat} \\ dX^{0}(s \begin{bmatrix} x t \end{pmatrix}) = dX^{0}(div_{X} \circ (s) \begin{bmatrix} x & x \circ t \end{pmatrix}):$$

Proof. For the rst statement, assume, for convenience, that  $X^0 = ff_i g$  and  $X = fg_j g$ , and rename  $X^0$  and  $X = X^0$  by F and G respectively. Applying Lemma A 1.(8), introducing a labeling set F (resp. G), bijective with F (resp. G) (and associated sets  $e_r$  and  $e_G$ ), gives the expression in question as:

If the canonical decomposition of twith respect to F is

$$t(f;g) = \exp^{\frac{0}{2}} \frac{1}{2} \int_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f;g);$$

then the previous expression is equal to

Now, because s is an X  $^0\!$  -substantial operator, this gives the required decomposition of s [\_F [\_G t with respect to F .

For the second statem ent, we com pute

Z <sub>rat</sub>

$$dF (s [_{F[G} t) \\ * 0 \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{1} A_{i} s (f;g;0f;0g);exp^{0} \bigcup_{f_{i}}^{f_{j}} M_{ij} + \frac{1}{2} \bigcup_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{1} A_{i} s (f;g;0f;0g);exp^{0} \bigcup_{f_{i}}^{f_{j}} M_{ij} + \frac{1}{2} \bigcup_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{1} A_{i} s (f;g;0f;0g);exp^{0} \bigcup_{f_{i}}^{f_{j}} M_{ij} + \frac{1}{2} \bigcup_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{1} A_{i} s (f;g;0f;0g);exp^{0} \bigcup_{f_{i}}^{f_{j}} M_{ij} + \frac{1}{2} \bigcup_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{1} A_{i} s (f;g;0f;0g);exp^{0} \bigcup_{f_{i}}^{f_{j}} M_{ij} + \frac{1}{2} \bigcup_{f_{i}}^{f_{j}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{A} t \prod_{f_{i}}^{f_{i}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{A} t \prod_{f_{i}}^{f_{i}} M_{ij}^{A} t T (f + f;g + g) \\ = exp^{0} \frac{1}{2} \bigcup_{i=1}^{0} M_{ij}^{A} t \prod_{f_{i}}^{f_{i}} M_{ij}^{A} t T (f + f;g + g)$$

To proceed, we must adjust this calculation so that it can distinguish struts, appearing in the left of the pairing, according to which factors, appearing in the right of the pairing, they are glued to. This adjustment is made with Lemma A 1.(6), and leads to the following (introducing sets F<sup>a</sup>, F<sup>b</sup> and F<sup>c</sup>, bijective copies of F):

$$\begin{array}{c} \star \quad 0 \\ \exp{(\theta - \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{a} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta - f_{j}^{b} \\ 0 \end{pmatrix}_{ij}^{1} + \frac{1}{2} \begin{pmatrix} \theta -$$

The calculation proceeds by removing factors appearing in left hand side of the pairing, at the expense of operating with them on the right hand side, follow Lemma A 1.(7)  $\frac{1}{1}$ 

The rst term to be treated in this way will be  $\exp^{(0)} \left[\frac{1}{2} \prod_{j=1}^{p} M_{ij}\right]^{1} A$ ; giving (following an application of  $e_{f_{1}^{b}}$ 

The next term to be so treated will be exp  $\begin{pmatrix} 0 & f_j^c \\ & j & M \\ & j & M \\ & 0 & f_j^b \end{pmatrix}^{-1} A$ ; which gives:

$$exp^{(l)} \begin{bmatrix} 0 & 0 & 1 & 1 & 0 & 1 \\ gf_{j}^{c} & & f_{j} & & f_{j} \\ gf_{j}^{c} & & f_{j} & & f_{j} \\ gf_{j}^{b} & & f_{j} & & f_{j} \\ gf_{j}^{b} & & & f_{j} \\ gf_{j}^{b$$

The result of these operations is

\* 0  
exp 
$$\begin{pmatrix} 1 \\ 2 \\ gf_{1}^{a} \\ gf_{1}^{a} \\ gf_{1}^{a} \\ gf_{1}^{a} \\ gf_{1}^{a} \\ gf_{1}^{c} \\ ff_{1}^{c} \\ gf_{1}^{c} \\ gf_{1}^{c}$$

Perform ing the pairing with respect to the variables F, leads to the replacement of the the innerh; i with

$$s(f^{a};g;f^{b}M;@g);T(f^{c};g+g)_{G}:$$

$$0 \qquad ef_{j}^{b} \qquad 1$$
Trading the factor exp 
$$(i = f_{i}^{a})^{i}M_{ij} = A;gives:$$

$$ef_{i}^{a} \qquad 1$$

$$exp \qquad ef_{j}^{b} \qquad 1$$

$$exp \qquad if_{j}^{b}M_{ij} = A_{i}[F^{b}S(f^{a};g;f^{b}M;@g) = div_{F^{a}} = S(f^{a};g;f^{b}M + @f^{a};@g)_{@g!@g}:$$

Perform ing the pairing with respect to  $F^{b}$  sets  $f^{b}$  to zero, and thus we arrive at the following expression.

$$\exp^{\mathbb{Q}} \left( \frac{1}{2} \int_{\mathbb{Q}_{f_{i}}^{a}}^{\mathbb{Q}_{f_{i}}^{a}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{a}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{c}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{c}}}^{\mathbb{Q}_{f_{i}}^{c}}} \int_{\mathbb{Q}_{f_{i}}^{$$

The proof follows by inspection.

#### Appendix B. The W heels Identity

The goal of this section is to show the following W heels Identity:

Theorem B.1. For every M 2 Herm ( $_{loc}$  ! Z), we have that

7

dX exp 
$$\frac{1}{2} X \overset{x_j}{\underset{x_i}{\overset{y}{\xrightarrow{y_j}}}} M_{ij}(e^{h_1}; ...; e^{h_g}) = exp_t \frac{1}{2} (M);$$

where we think of (M) as taking values in  $\hat{}=(cyclic)$ , which is idential ed via the map  $t_i ! e^{h_i}$  with the set Q hh<sub>1</sub>;:::;h<sub>g</sub>ii=(cyclic) of form all power series in noncommuting variables modulo cyclic permutations.

Proof. Choose generators  $ft_1$ ;:::; $t_g$ g for the free group F and identify  $\hat{}$  with the ring of noncommutative power series  $fh_1$ ;:::; $h_n$ g. Recall the map !  $\hat{}$  given by  $t_i$  !  $e^{h_i}$ .

The A arhus integral is calculated by splitting the quadratic part from the rest of the integrand as follows:

W ith the following notation for  $\Red$  and  $\G$  reen" struts,

\*

it follows that the desired integral equals to the following form alseries (which makes sense, since nitely many terms appear in each degree)

$$\exp \quad \frac{1}{2} \, \overset{"}{j}_{R}^{R} ; \exp \quad \frac{1}{2} \, \overset{"}{j}_{J}^{G} = \overset{"}{\underset{n=0}{\overset{"}{x}}} (1)^{n} \, (\frac{1}{2})^{n} \, (\frac{1}{2})^{n} \frac{1}{n!n!} \overset{"}{\underset{n!}{n!}} R ; \overset{"}{\underset{j}{f}} G :$$

Speci cally, the answer is recovered if each \jpin" is accompanied by a K ronecker delta. For example:

$$\overset{"}{j}_{G}^{R} = \begin{array}{c} X & \overset{x_{k}}{j}_{M jk} (1; \dots; 1)^{1} & X & \overset{x_{k}}{j}_{M jk} (1; \dots; 1)^{1} \\ \overset{"}{j}_{G} & \overset{i}{j}_{i;j;k} & \overset{j}{j}_{M ij} (e^{h_{1}}; \dots; e^{h_{g}}) & \overset{i}{j}_{i;j;k} & \overset{j}{j}_{M ij} (1; \dots; 1) \end{array} :$$

We focus on a particular sum m and. It will help in the m anagement of the combinatorics if we label each of the arcs that appear in the struts:

Now we introduce some de nitions to assist in the enumeration of the polygons that arise from this pairing. An RG-shape of degree n is a collection of even-sided polygons whose edges are alternately colored red and green. The total number of edges is 2n.

A labeled RG-shape of degree n is an RG-shape with an extra bivalent vertex in each edge, such that:

1. the resulting set of red (resp. green) edges is equipped with a bijection to  $fr_{1;1};r_{1;2};\ldots;r_{n;1};r_{n;2}g$  (resp.  $fg_{1;1};g_{1;2};\ldots;g_{n;1};g_{n;2}g$ ),

2. if an edge is labeled, for example  $r_{i;1}$  (resp.  $g_{i;1}$ ), then its neighboring red (resp. green) edge is labeled  $r_{i;2}$  (resp.  $g_{i;2}$ ).

Let the constant C , for x = x = x = x = x some RG -shape, denote the number of labeled RG -shapes which recover when their labels and extra bivalent vertices are forgotten.

$$\begin{bmatrix} n & i & n \\ j & i \end{bmatrix} G = \begin{bmatrix} X & X \\ i & j \end{bmatrix} G = \begin{bmatrix} X & C \\ i & i \end{bmatrix} C = \begin{bmatrix} X & C \\ i & i \end{bmatrix} C$$
 a labeled RG -shape of deg n an RG -shape of deg n

To calculate this constant describe an RG-shape of degree n, , by a sequence  $(\underline{i}\,;\ldots;\underline{i}_n\,)$  where  $\underline{i}_j$  counts the number of polygons with 2j edges.

LemmaB.2.

$$C = \frac{n \, 2^n n \, 2^n}{((2) \, (1))^{i_1} \, i_1 \, ! \, ((2) \, (2))^{i_2} \, i_2 \, ! \, : : : ((2) \, (n))^{i_n} \, i_n \, !};$$

Proof. Fix a copy of with its set of edges ordered (any old how, so as to distinguish them from each other). There are precisely  $2^n n 2^n n! w$  ays of labeling this so as to respect the conditions of the above de nition.

But this overcounts by a multiplicative factor of the order of the sym metry group of (that is, with the ordering of the edges forgotten). That sym metry group is precisely the following group (that is, swaps of identical polygons, and rotations and ips of individual polygons):

$$(i_1 n (D_1 :::D_1)) = i_n (n (D_n :::D_n));$$

where D  $_{\rm n}$  represents the nth dihedral group.

1 -

Thusweconclude:

$$C = \frac{2^{n} n 2^{n} n!}{\# f(_{i_{1}} n (D_{1} _{1})) _{i_{n}}(n (D_{n} _{n}))g}$$

If i denotes a connected polygon with 2i edges, it follows that

Substituting the de nitions of the  $\Red$ " and  $\G$  reen" struts into  $_{p}$  yields

 $p = tr (M (1;:::;1) {}^{1} (M (e^{h_{1}};:::;e^{h_{g}}) M (1;:::;1))^{p} = tr (M (1;:::;1) {}^{1} M (e^{h_{1}};:::;e^{h_{g}}) I)^{p} : from which the result follow s.$ 

# Appendix C. Infinitesimal versus group-like wrapping relations

In this section we show that the in nitesim alw rapping relations ! in ply the group-like w rapping relation  $!^{gp}$ . In order to achieve this, it will be useful to introduce the variant  $!_i^0$  of  $!_i$  for i = 1; :::; g where we allow the diagram D 2 A ( $?_{X [f@hg}; _{bc}$ ) to have several legs labeled by @h. In other words,  $!_i^0$  is generated by the move  $(M; D_1)$   $(M; D_2)$  where

$$D_{m} = \operatorname{con}_{fhg} '_{m,i}(D) t \exp_{t} \frac{1}{2} h (M^{-1} '_{m,i}M)$$

for m = 1;2 and  $'_{1;i}$  is the identity and  $'_{2;i}$  (abbreviated by  $'_i$  below) denotes the substitution  $t_i! e^{h} t_i e^{h}$  on beads.

It is obvious that the  $!_i^0$  relation includes the  $!_i$  relation. Our goal for the rest of the section is to show that they coincide (see P roposition C.7) and that the  $!^{gp}$ -relations are implied by the !-relations (see P roposition 3.9).

Recall the m aps  $_{i: bc}$ !  $E_{bc}$  of Section 3.1 that keep track of the h-degree 1 part of the substitutions  $'_{t_i! e^{h} t_i e^{h}}$  (a) of a 2  $_{bc}$ . We will introduce an extension  $b_i$  of  $_i$  which will allow us to keep track of the h-degree n part of  $'_{t_i! e^{h} t_i e^{h}}$  (a), for all n. The following kemma (which the reader should compare with Lemma 3.2) summarizes the properties of  $b_i$ .

Lem m a C .1. For every i = 1;:::;g, there exists a unique

$$b_i: P_{loc}! P_{loc}$$

that satis es the following properties:

$$\begin{array}{rcl} b_{i}(t_{j}) &=& _{ij}[t_{i};h] = & _{ij}(t_{i}h & ht_{i}) \\ b_{i}(h) &=& 0 \\ b_{i}(ab) &=& b_{i}(a)b + & ab_{i}(b) \\ b_{i}(a + b) &=& b_{i}(a) + & b_{i}(b) \end{array}$$

In particular, the restriction of  $b_i$  to  $\ _{\rm loc}$   $\ {\rm I}\!{\rm d}_{\rm loc}$  equals to  $\ _i.$ 

Proof. First, we need to show the existence of such a  $b_i$ . A coordinate approach is to de ne $b_i: R \ ! \ R$  on the ring R of Section 3.1 which is isomorphic to  $B_{\rm loc}$ . G iven the description of R, it is easy to see that  $b_i$  exists and is uniquely determined by the above properties.

An alternative, coordinate-free de nition of  $b_i$  is the following. Let  $L_{loc}$  denote the Cohn localization of the group-ring Q F ? Z ? Z], followed by a completion with respect to t 1 and t<sup>0</sup> 1 (where t and t<sup>0</sup> are generators of Z ? Z). Let  $h = \log t$  and  $h^0 = \log t^0$ . The map  $t_{i,!} e^{h} t_{i,e^{h}} : Q F ? Z ? Z]! Q F ? Z ? Z]$  is -inverting, thus it extends to a map of the localization, and further to a map  $L_{loc}! L_{loc}$  of the completion.  $L_{loc}$  is a  $h^0$ -graded ring, and we can de ne

$$b_i: P_{bc}! P_{bc}$$

by

$$b_i = \prime_{h^0! h} deg_{0} \prime_{t_i! e^{h} t_i e^{h}}$$

It is easy to see that this de nition of  $b_i$  satis es the properties of the km m a, and that the properties characterize  $b_i$ .

F in ally, Lemma 3.2 in plies that the restriction of  $b_i$  on  $b_{ic}$   $B_{ic}$  equals to i.

Lem m a C .2. For all n and i we have that

$$\frac{1}{n!}b_i^n = deg_h^n \quad \prime_{t_i! \ e^h \ t_i e^h}:$$

considered as maps !  $P_{loc}$ .

Proof. In view of Lemma 32, it su ces to show that the maps agree on generators  $t_j$  of  $F_g$ . For  $j \notin i$ , both maps vanish. For j = i, the following identity

$$e^{h}t_{i}e^{h} = \exp(ad_{h})(t_{i}) = \frac{x^{h}}{n!} \frac{1}{n!} [[t_{i};h];h]; \dots; h]$$

and Lemma 3.2 show that both maps agree. The result follows.

Next, we extend the above lem m a for m atrices.

Lem m a C .3. For all n and sites i we have that

$$\frac{1}{n!}b_{i}^{n} = deg_{h}^{n} \quad \prime_{t_{i}! e^{h} t_{i}e^{h}}:$$

considered as maps M at( ! Z) ! M at( ${\rm E}^{l}_{\rm loc}$  ! Z).

Proof. Observe the identity

(14)

$$b_i$$
 (AB) =  $b_i$  (A)B + A $b_i$  (B)

for matrices A, B (not necessarily square) that can be multiplied. It im plies that

$$0 = \frac{1}{n!} b_{i}^{n} (I) = \frac{1}{n!} b_{i}^{n} (M M^{-1}) = \sum_{k=0}^{X^{n}} \frac{1}{k!} b_{i}^{k} (M ) b_{i}^{n-k} (M^{-1}):$$

The lemma follows by induction on n and the above identity which solves  ${}^n_i M$  ) in terms of quantities known by induction.

P roposition C .4. For all n and sites i we have that

$$\frac{1}{n!}b_i^n = deg_h^n \quad \prime_{t_i! \ e^h \ t_i e^h}:$$

considered as maps  $loc ! Pl_{loc}$ .

Proof. Consider a matrix presentation of s 2 loc. I.e.,

$$s = (1;0;...;0)M^{-1}b$$

for M 2 M at ( ! Z) and b a column vector over . Equation (14) im plies that

$$\frac{1}{n!} b_{1}^{n} (s) = \sum_{k=0}^{X^{n}} \frac{1}{k!} b_{1}^{k} (M^{-1}) b_{1}^{n-k} (5):$$

The proposition now follows from Lemmas C 2 and C 3.

Proposition C.5. For all diagram s D 2 A ( $?_{X;@h}$ ;  $_{loc}$ ), M 2 Herm ( $_{loc}$  ! Z), and i = 1;:::;g, we have that

$$\frac{1}{n!} b_{i}^{n} (D) = \deg_{h}^{n} '_{t_{i}! e^{h} t_{i} e^{h}} (D) t \exp_{t} \frac{1}{2} h (M^{-1} '_{t_{i}! e^{h} t_{i} e^{h}} (M)) :$$

Proof. The proof follows from a careful combinatorial counting similar to that of the W heels Identity of Theorem B.1. Let us precede the proof with an inform alexam ination of what takes place when we repeatedly apply  $b_i$  to the empty diagram, D = . The rst application results in a term :

$$\frac{1}{2} \sum_{k}^{X} (\mathbf{b}_{1} (\mathbf{b}_{1} (\mathbf{M})))_{kk}$$

The second application requires the result of applying  $b_i$  to M<sup>1</sup>. This can be done using the following special case of Equation (14):

(15) 
$$b_i (M^{-1}) = M^{-1} (b_i (M)) M^{-1}$$
:

So, the second application results in the following term s:

$$\frac{1}{2} X \longrightarrow M^{-1} (b_{i} (M))_{kl} + \frac{1}{2} X \longrightarrow M^{-1} (b_{i} (M))_{kl} + \frac{1}{2} X \longrightarrow M^{-1} (b_{i})^{2} (M))_{kk} + \frac{(\frac{1}{2})^{2}}{2!} X \longrightarrow M^{-1} (b_{i} (M))_{kk} + \frac{(\frac{1}{2})^{2}}{2!} X \longrightarrow M^{-1} (b_{i}$$

W e see that the term s arising from the application of  $b_i$  fall into three categories.

(16) 
$$b_{i}() = \frac{1}{2} \bigwedge_{k} M^{-1}(b_{i}(M))_{kk}$$

(17) 
$$(b_{1} (M^{-1}))_{kl} = (M^{-1})_{ml} (b_{1} (M^{-1})_{ml})_{km}$$

(18) 
$$(b_{i}(b_{i})^{p}(M))_{kl} = ((b_{i})^{p+1}(M))_{kl}$$

Thus, we see that  $b_i^n$  () equals to a linear combination of diagram s, each consisting of a number of wheel components, each of which is of the following form (this can be (over-)determined by a sequence (1;:::; ) of positive integers):

$$\begin{array}{c} X \\ k_{1}, \dots, k_{k} \end{array} \begin{pmatrix} M^{-1} ( (b_{1}) (M^{-1}))_{k-1} \\ & \vdots \\ M^{-1} ( (b_{1})^{-1} (M^{-1}))_{k-k_{1}} \end{pmatrix}$$

The question, then, is to calculate an expression in term s of such diagram s, counting  $\allways"$  that they arise from the application of moves (16)-(18). We will generate a sequence of such moves with the following object, an edge-list on a shape.

The shape determ ined by a set of positive integers ( $_1$ ;:::; ) is a set of oriented polygons, one with  $_1$  edges, and so on, such that each has a distinguished bivalent vertex, its basepoint. A labeled shape is a shape with its edges labeled by positive integers. The diagram D determ ined by some labeled shape is the diagram that arises by replacing each polygon by a series of wheels, in the follow ing manner (to distinguish them from beads, the bivalent vertices of the polygons will be drawn by small horizontal segments, and the base-point by an x).

$$\begin{array}{c} \begin{array}{c} & X \\ & & \end{array} \\ \end{array} \begin{array}{c} & X \\ & & \end{array} \\ \begin{array}{c} & M \end{array} \begin{array}{c} M \end{array} \begin{pmatrix} M \\ ( \begin{array}{c} (b_{1}) \\ ( \end{array}{c} (b_{1}) \\ ( \begin{array}{c} (b_{1}) \\ ( \end{array}{c} (b_{1}) \\$$

An edge-list on a shape is a nite sequence of the edges of the shape such that each edge is represented at least once. (It is convenient to enum erate the edges along the orientation, starting from the basepoint.) The labeled shape associated to some edge-list on that shape is that shape with each edge labeled by the num ber of times that edge appears in the edge-list.

An edge-list (1;:::; r) on a shape will determ ine a sequence of moves ((16)-(18)) as follows.

- 1. If some  $_{i}$  is the rst appearance in the edge-list of an edge from some polygon with edges, then that term determ ines a move (16), where for the purposes of discussion the introduced wheel is oriented with beads, with one such distinguished as the \ rst", and with the label M<sup>-1</sup> ( $b_{i}$  M) sitting on bead number  $_{i}$ , and with all other beads labeled 1.
- 2. If some  $_{i}$  is the rst appearance in the edge-list of som e edge of a polygon, som e other edge of which has already appeared in the edge-list, then that term determ ines a move (17) applied to the factor of M<sup>-1</sup> that immediately follows the edge  $_{i}$  on the (oriented) wheel. (That is, the factor of M<sup>-1</sup> ( $b_{i}$  M)) that is introduced slides" back around the polygon from there to the bead number  $_{i}$ .)
- 3. If some edge  $_{i}$  has already appeared in the path (say, p times) then that step determ ines the application of a move (18) to the factor  $(b_{i})^{p}$  (M) appearing on the bead number  $_{i}$ .

Rem ark C.6. O beerve that the result of these m oves will be the diagram determ ined by the labeled shape associated to the edge-list.

O berve also that the num ber of di erent edge-lists on a shape which give rise to a given associated labeled shape, e.g.

Two edge-lists determ ine the same sequence of moves if and only if they are related by symmetries of the shape:

1. rotating individual polygons (i.e. shifting the \base-point"),

2. permuting a subset of polygons which all have the same num ber of edges.

We now de ne a function Sym from the set of shapes to the positive integers as follows. If a shape is specified by a sequence  $(_1; _2; :::;_k)$ , where  $_1$  counts the number of polygons with one edge,  $_2$  counts the number of polygons with 2 edges, and so on, then

Sym () = 
$$(_1!_2!:::_k!)(1^12^2:::k^k):$$

We are now ready to prove Proposition C 5. First consider the case where the given diagram D = i is empty.

$$\begin{array}{rcl}
X & \underbrace{(b_{i})^{n}}{n!}(\textbf{;}) &= & X & \underbrace{1}{\text{Sym}(.)} & X & \underbrace{1}{(1,1!\cdots 1, 1!)\cdots (1,1!\cdots 1, 1!)} D \\
& & \text{a shape} & \underbrace{1}{\text{Sym}(.)} & \text{: a labeling of} & \underbrace{(1,1!\cdots 1, 1!)\cdots (1,1!\cdots 1, 1!)}_{(1,1!\cdots 1, 1!)\cdots (1,1!\cdots 1, 1!)} D \\
& & = & \exp & \frac{1}{2}\text{Tr} & \underbrace{X^{i}}_{m=1} & \underbrace{1}{m} & M^{-1} & b_{i}(M) + \underbrace{(b_{i})^{2}}{2!}(M) + \underbrace{(b_{i})^{3}}{3!}(M) + \cdots \\
& & = & \exp_{t} & \frac{1}{2} h & M^{-1} & t_{i}! e^{h} t_{i}e^{h}(M) & \textbf{;}
\end{array}$$

as needed. In case D is not empty, assume for convenience that there are k beads on D, and explicitly record their labels D  $(q_1; ::: q_k)$ . Now,

$$\begin{array}{ccc} X & \underbrace{(b_{1})^{n}}{n!} (D) = & X \\ & & \\ n & \underbrace{(b_{1})^{n_{1}}}{n!} (D) = & \underbrace{(b_{1})^{n_{1}}}{n_{1}!} (D) \\ & & \\ & & \\ \end{array}$$

and the P roposition follows from the above case of D = ;.

Proposition C.7. The ! and  $!^{0}$  relations on A (?<sub>x</sub> ;  $_{loc}$ ) coincide.

Proof. Consider two pairs  $(M; s_1)$  and  $(M; s_2)$  related by a wrapping move  $!_i^0$  as in de nition 3.5. In other words, for some s 2 A gp ( $?_{X;@h}; _{loc}$ ) we have that

$$s_m = con_{fhg} \prime_{m;i}(s) t exp_t \frac{1}{2} h (M^{-1} \prime_{m;i} M)$$

for m = 1;2 where '<sub>1;i</sub> is the identity and '<sub>2;i</sub> =  $t_{i!} e^{h} t_{i}e^{h}$ . Consider the case that s has a diagram with n legs labeled by @h. Proposition C.5 in plies that

$$s_2 = \frac{1}{n!} con_{fhg} (b_i^n (s)):$$

Let  $s^0$  denote the result of gluing all but one of the n  $\theta$ h legs of  $b_i^{n-1}$  (s) and renam ing the resulting one  $\theta h^0$ . In other words,  $s^0 = con_{fhg} (b_i^{n-1} (s) (\theta + \theta h^0))$ . Let  $b_i^0$  denote the action of  $b_i$  for  $\theta h^0$ -labeled legs. If s has at least one  $\theta h$  leg, it follows that

$$s_2 = con_{fh^0q} (b_i^0 (s^0)) = 0 \mod !_i$$

by the in nitesimal wrapping relations, and it follows that  $s_1 = 0$  by de nition. If s has no  $(h \log, then s_1 = s_2$ . Thus, in all cases,  $(M; s_1)^{!_1} (M; s_2)$ . The converse is obvious; thus the result follows.

Proof. (of Proposition 3.9) It is obvious that the  $!^{0}$  relation in plies the group-like  $!^{gp}$  relation of Section 5.4. Since  $!^{0} = !$ , the result follows.

### Appendix D. Infinitesimal basing relations

This section, although it is not used anywhere in the paper, is added for completeness. The reader may have noticed that in Section 3, we mentioned in nitesimal versions of the wrapping and the CA relations, but not of the basing relations. The reason is that we did not need them; in addition <sup>R</sup><sub>rat</sub> is not known to preserve them. Nevertheless, we introduce them here.

Wedene

$$A (\sim_X ; b_c) = A (?_X ; b_c) = (1; 2);$$

where  $_{1:X}$  is the subspace of X - avored link relations that appeared in A, Part II, Section 5.2]:

and  $_{2:X}$  denote subspace generated by an action

$$A(?_{X}; _{loc}) ! A(?_{X}; _{loc})$$

described by pushing some element of f 2  $F_g$  onto all X -labeled legs of a diagram . We now compare the in nitesim al and group-like basing relations as follows.

F

Proposition D.1. (a) We have an isom orphism

$$A (:::; \sim_X ; _{bc}) = A (:::;?_X ; _{bc}) = h_X i! A (:::; _X ; _{bc}):$$

,

(b) The inclusion  $A^{gp;0}$  (:::;?<sub>X</sub>;  $_{loc}$ ) !  $A^{0}$  (:::;?<sub>X</sub>;  $_{loc}$ ) maps the  $^{gp}$ -relations to -relations, and induces a map

$$A^{gp; 0}$$
 (:::;~x; bc) !  $A^{0}$  (:::;?x; bc):

Proof. The rst part follows from the proof of [A, Part II, Theorem 3] without essential changes. The second part is standard.

# Appendix E.A useful calculation

In this section, we quote a useful and often appearing computation that uses the identities of Lemma A 1. Let ':  $A^{gp}$  (?<sub>X</sub>; <sub>bc</sub>) !  $A^{gp}$  (?<sub>X [fhg</sub>; <sub>bc</sub>) be a map that commutes with respect to gluing X -colored legs.

Theorem E.1. For X<sup>0</sup> X and s 2 Int<sub>x</sub> 
$$\circ$$
 A<sup>gp</sup> (~ <sub>x</sub>; <sub>loc</sub>), decomposed as in Equation (3), we have that  
 $Z$  and  $Z$  and

$$\begin{array}{ccc} {}^{rat} & {}^{Z} {}^{rat} \\ {}^{dX {}^{0}}(' {}^{\prime}(s)) = & {}^{dX {}^{0}} \exp \left( \frac{1}{2} X \right) \left( \frac{x_{j}}{2} X \right) \left($$

If in addition '(s) s is X -substantial for all s 2 A  $^{\rm gp}$  (?<sub>X</sub> ;  $_{\rm loc}$ ), then Z  $_{\rm rat}$ 

$$dX^{0}('(s)) = exp = \frac{1}{2} M^{1}(M^{-1}(M)) + dX^{0}(s) :$$

Proof. Consider the decomposition of s given by Equation (3). We begin with a  $\$  -function tryck" of Equation (11) of A 1:

$$s = \exp^{0} \frac{1}{2} \frac{X}{j} \frac{X_{j}}{j} M_{ij}^{A} t R = R (0 y); \exp \frac{1}{2} \frac{X}{j} \frac{X_{j}}{j} M_{ij} + \frac{y}{j} : +$$

Thus,

$$Z_{rat} = (R (@y)); \quad dX^{0} exp = \frac{1}{2} X_{ij} M_{ij} + U_{j} H_{ij} + U_{ij} H_{ij} + U_$$

Since div<sub>X</sub>  $\circ \exp e^{0} = \frac{1}{\int_{Y_{i}}^{e_{X_{j}}} M_{ij}^{1}A} = 1$ , the integration by parts Lemma A.4 in plies that

Equation (12) of Lemma A 1 implies that the above equals to:

$$\exp \frac{1}{2} X y_{j} Y_{i} X_{ij} X_{i$$

Returning to the expression in question:

$$Z_{rat} = \frac{Z_{rat}}{dX^{0}('(s))} = \frac{Z_{rat}}{dX^{0} \exp \frac{1}{2} X_{x_{i}}^{x_{j}} M_{ij} t' (R(@y)); \exp \frac{1}{2} X_{y_{i}}^{y_{j}} M_{ij}^{1} :$$

The second factor equals to  $'^{R_{rat}} dX^{0}(s)$  , and concludes the rst part of the theorem .

In order to compute the rst factor, we need to separate its X  $^0$ -substantial part. W ith the added assum p-tion on ', we can compute the rst rst factor in a manner analogous to the W heels Identity of Theorem B.1

Remark E 2. The proof of Theorem E 1 can be applied without changes to the case of the map ' = Hair<sub>H</sub> : A (?<sub>X</sub>;  $_{loc}$ ) ! A (?<sub>X [H</sub>) in order to show that for s 2 Int<sub>X 0</sub>A <sup>gp</sup> (?<sub>X</sub>;  $_{loc}$ ), we have that

$$dX^{0}(Hair_{H}(s)) = exp = \frac{1}{2} h(M^{-1}Hair_{H}M) Hair_{H}^{2} dX^{0}(s) :$$

### References

- [AMR] J.Andersen and J.M attes and N.Reshetikhin, The Poisson structure on the moduli space of at connections and chord diagram s, Topology 35 (1996) 1069{1083.
- [A 0] D.Bar-Natan, S.Garoufalidis, L.Rozansky and D.Thurston, W heels, wheeling and the Kontsevich integral of the unknot, Israel J.M ath. 119 (2000) 217{238.
- [A] D.Bar-Natan, S.Garoufalidis, L.Rozansky and D.Thurston, The Aarhus integral of rational homology 3-spheres I-II, Selecta M ath. in press.

[BL] D.Bar-Natan and R.Law rence, A rational surgery form ula for the LMO invariant, preprint 2000 math.GT/0007045.

[BLT] D.Bar-Natan, T.T.Q.Le and D.Thurston.Two applications of elementary knot theory to Lie algebras and Vassiliev invariants, preprint.

- [BG] D.Bar-Natan, S.Garoufalidis, On the Melvin-Morton-Rozansky conjecture, Inventiones Math. 125 (1996) 103(133.
- [BR] J.Berstel and C.Reutenauer, Rational Series and their Languages, Springer-Verlag 1984.
- [CS1] S.Cappell and J.Shaneson, The codimension two placement problem and homology equivalent manifolds, Annals M ath.99 (1974) 277{348.
- [CS2] \_\_\_\_\_, Link cobordism, CommentariiMath.Helv.55 (1980) 20{49.
- [C] PM.Cohn, Free rings and their relations, A cadem ic Press, 1985.
- [Dv] J.Duval, Form e de Blanch eld et cobordism e d'enterlacs bords, Commentarii Math. Helv. 61 (1986) 617(635.
- FV] M.Farber and P.Vogel, The Cohn localization of the free group ring, M ath.Proc.Cam bridge Phil.Soc.111 (1992)
  432{443.

[Fa] , Noncommutative rational functions and boundary links, M ath. Annalen 293 (1992) 543 [568.

- [FRa] \_\_\_\_\_and A.Ranicki, The Morse-Novikov theory of circled-valued functions and noncommutative localization, Tr. M ath.Inst.Steklova 225 (1999) 381{388, and also math.DG/9812122.
- [FR] R.Fenn and C.Rourke, On Kirby's calculus of links, Topology 18 (1979) 1{15.
- [GR] S.G aroufalidis and L.Rozansky, The loop expansion of the Kontsevich integral, abelian invariants of knots and S-equivalence, preprint 2000, math.GT/0003187.
- [GL1] \_\_\_\_\_and J. Levine, Hom obgy surgery and invariants of 3-m anifolds, preprint 2000, math.GT/0005280.
- [GL2] \_\_\_\_\_\_ and \_\_\_\_\_, A nalytic invariants of boundary links, preprint 2000, math.GT/0102103.
- [GL3] \_\_\_\_\_\_ and \_\_\_\_\_, Concordance and 1-bop covers, preprint 2001, math.GT/0102102.
- [GGP] \_\_\_\_\_, M.Goussarov and M.Polyak, Calculus of clovers and nite type invariants of 3-m anifolds, Geometry and Topology, 5 (2001) 75{108.
- [Ga] \_\_\_\_\_, Boundary links, localization and the loop move, preprint 2001.
- [G a2] \_\_\_\_, to appear.
- [GK] \_\_\_\_\_and A.K ricker, Finite type invariants of cyclic branched covers, preprint 2001.

- [GT] \_\_\_\_\_and P. Teichner, in preparation.
- [Gu] M.Gutierrez, Boundary links and an unlinking theorem, Trans. Amer. Math. Soc. 171 (1972) 491 (499.
- [Ko] M.Kontsevich, Vassiliev's knot invariants, Adv. in Sov. Math., 16(2) (1993), 137{150.
- [K] K.H.Ko, Seifert matrices and boundary link cobordisms, Trans. Amer. Math. Soc. 299 (1987) 657 [681.
- [Kr1] A. Kricker, The lines of the Kontsevich integral and Rozansky's Rationality Conjecture, preprint 2000, math/GT.0005284.
- [Kr2] \_\_\_\_\_, C overing spaces over claspered knots, preprint 1999, math.GT/9901029.
- [Le1] J.Levine, Knotmodules I, Trans. AM S 229 (1977) 1{50.
- [Le2] \_\_\_\_\_, Link concordance and algebraic closure, II, Inventiones M ath.96 (1989) 571{592.
- [LM] T.T.Q. Le, J. M urakami, Parallel version of the universal Vassiliev-Kontsevich invariant, J. Pure and Applied Algebra, 121 (1997) 271(291.
- [LM M O] \_\_\_\_\_, H.M urakam i, J.M urakam i and T.Ohtsuki, A three-m anifold invariant via the Kontsevich integral, O saka J.M ath 36 (1999) 365{395.
- [LMO] T.T.Q.Le, J.M. urakami, T.Ohtsuki, A universal quantum invariant of 3-m anifolds, Topology 37 (1998) 539 (574.
- [R] L.Rozansky, Notes from lectures delivered in Grenoble, 1999.
- [Rb] J.Roberts, K irby calculus in manifolds with boundary, Turkish J.M ath.21 (1997) 111{117.
- [Sm] N.Smythe, Boundary links, Topology Sem inar (Wisconsin 1965) Princeton Univ. Press, Princeton, NJ 1966, 69{72.
- [Tu] V.Turaev, Conway and Kau man modules of a solid torus, J.Soviet M ath.52 (1990) 2799(1805.
- [Vo] P.Vogel, The Cohn boalization of the free group ring, M ath. Proc. C am bridge Phil. Soc. 111 (1992) 433{443.

School of M athematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA. E-m ailaddress: stavros@math.gatech.edu

E -m ailaddress: kricker@math.gatech.edu