

A RATIONAL NONCOMMUTATIVE INVARIANT OF BOUNDARY LINKS

STAVROS GAROUFALIDIS AND ANDREW KRICKER

ABSTRACT. We construct an invariant of boundary links which is at least as strong as the Kontsevich integral, determines the S -equivalence class of a boundary link and which takes values in a space of trivalent graphs whose edges are decorated by rational functions in noncommuting variables. Our invariant is axiomatically characterized by a universal property closely related to the Homology Surgery view of boundary links, and comes equipped with a diagrammatic integration theory, in the spirit of perturbative quantum field theory.

1. INTRODUCTION

1.1. Introduction. The Kontsevich integral of a link is a powerful invariant that takes values in a completed vector space of graphs. The graphs in question have univalent and trivalent vertices only (so-called *unitrivalent graphs*), and their univalent vertices are labeled by the components of the link. In addition, the graphs are considered modulo some well-known relations, that include a 3-term relation (the IHX relation) depicted as follows:

$$\begin{array}{c} \diagup \quad \diagdown \\ | \\ \hline \end{array} = \begin{array}{c} | \quad | \\ | \quad | \\ \hline \end{array} - \begin{array}{c} \diagdown \quad \diagup \\ | \\ \hline \end{array}$$

Every unitrivalent graph G is the union of a trivalent graph G^t together with a number of unitrivalent trees attached on each edge of the trivalent graph. This is true provided that no component of G is a tree, and provided that we allow G^t to be a circle in case a connected G has one loop (i.e., G has betti number 1). Using the IHX relation, it follows that every unitrivalent graph with no tree components is a sum of trivalent graphs with *hair* attached on their edges (by 'hair' we mean the tree with one edge and no trivalent vertices).

We will be interested in links whose Kontsevich integral has no tree part. For such links, the Kontsevich integral takes values in a completed vector space generated by trivalent graphs with hair. The hair are colored by the components of the link, and we can record this information by placing monomials in variables h_i (one variable per link component) on the edges of a trivalent graph as follows:

$$\begin{array}{c} L_i \quad L_j \quad L_k \\ | \quad | \quad | \\ \hline \end{array} \longrightarrow \begin{array}{c} h_i \quad h_j \quad h_k \\ \hline \end{array}$$

The variables h_i do not commute with each other. In the case of a knot, it was conjectured by Rozansky [R2] that there is a rational lift Z^{rat} of the Kontsevich integral Z that takes values in a completed vector space of trivalent graphs whose edges are labeled by certain rational functions in a variable $t = e^h$. It is a purpose of our paper to provide a proof of this Rationality Conjecture for the Kontsevich integral of a knot.

The rational lift Z^{rat} determines the Kontsevich integral Z and also determines the S -equivalence class of a knot. Although the appropriate (Hair) map on diagrams is not 1-1, [P], it is still conceivable that Z^{rat} carries the same information as Z . At any rate, one can ask in what sense is Z^{rat} a better invariant than Z ? The power of the Z^{rat} invariant is that it packages whole power series of Vassiliev invariants into rational functions. These rational functions even though they are no longer Vassiliev invariants, they are nonetheless finite type invariants with respect to a new move. This so called *null move* is described in terms of surgery on a null homologous clasper in the knot complement, and was introduced and studied extensively by [GR]. As a result, we can give *realization theorems* for these rational functions in much the same spirit as was done

Date: This edition: April 18, 2002 First edition: April 18, 2001.

S.G. was partially supported by an NSF grant DMS-98-00703 and by an Israel-US BSF grant. A.K. was partially supported by a JSPS fellowship.

This and related preprints can also be obtained at <http://www.math.gatech.edu/~stavros> .

1991 *Mathematics Classification.* Primary 57N10. Secondary 57M25.

Key words and phrases: Boundary links, Kontsevich integral, Cohn localization.

decades ago for the Alexander polynomial, [Le1]. A sample realization is the computation of the symbol of Z^{rat} given in Section 8, and is used in [GT] to settle a question of low-dimensional topology (namely, to separate minimal rank knots from Alexander polynomial 1 knots), and to fix an error in earlier work of M. Freedman. This is perhaps the first application of finite type invariants in a purely 3-dimensional question, which was made possible by using the 2-loop part of the Z^{rat} invariant (the so-called Q function). Another application of the 2-loop part is a formula for the Casson-Walker invariant of cyclic branched coverings of a knot in terms of the signature of the knot and residues of the Q function, obtained in joint work [GK1].

From the above mentioned applications, it seems that the rational form Z^{rat} of the Kontsevich integral is a powerful new invariant of knots, which is closely related to classical topology.

For simplicity, we restricted the above discussion to knots. We mentioned earlier that we will focus on links whose Kontsevich integral has vanishing tree part. Using [HM], this condition is equivalent to the vanishing of all Milnor invariants of a link, which is a well-defined statement. This class of links contains (and perhaps coincides with) the class of sublinks of homology boundary links, [Le2]. For simplicity, we will focus on *boundary links*, namely those each component of which bounds a surface, such that the surfaces are pairwise disjoint.

It turns out that there is a rational form Z^{rat} of the Kontsevich integral of a boundary link, which takes values in a completed vector space spanned by trivalent graphs with beads. The beads in question are *rational functions in noncommuting variables*, and they lie in a ring Λ_{loc} which is the noncommutative (Cohn) localization of the group-ring $\Lambda = \mathbb{Z}[F]$ of the free group F . Here F is the free group in generators t_1, \dots, t_g , corresponding to the components of the boundary link.

Thus, our rational form Z^{rat} of boundary links is intimately related to the noncommutative localization Λ_{loc} of Λ . Two previous relations among boundary links and noncommutative localization are already known through the work of Cappell-Shaneson and Farber-Vogel. Let us comment briefly on this topic. It is well-known that the problem of Homology Surgery and its application to the classification of boundary links, up to cobordism, is closely related to the set of Hermitian matrices over Λ , invertible over \mathbb{Z} . Cappell-Shaneson formulated the problem of boundary link cobordism (in high dimensions) in terms of Γ -groups (see [CS1, CS2]) and Vogel identified these Γ -groups in terms of surgery L -groups of a suitable localization of Λ , [Vo]. Later on, Vogel and Farber identified this localization with the ring of rational functions in noncommuting variables, [FV]. In addition, Farber constructed an invariant of boundary links with values in Λ_{loc} , [Fa].

Our invariant Z^{rat} is a graph-valued generalization of the L -theory invariants of Cappell-Shaneson and the invariant of Farber. It is rather surprising to reinvent the localization of Λ by looking at the Kontsevich integral of a boundary link.

One more comment about links. Rozansky has recently introduced a Rationality Conjecture for a class of *algebraically connected* links, that is links with nonvanishing Alexander polynomial, [R3]. This class is disjoint from ours, since a boundary link with more than one component has vanishing Alexander polynomial. It is an interesting question to extend our results in this setting of Rozansky. We plan to address this issue in a later publication.

1.2. Statement of the results. A refinement of *boundary links* (abbreviated ∂ -links) are *F-links*, i.e., a triple (M, L, ϕ) of a link L in an integral homology 3-sphere M and an onto map $\phi : \pi_1(M - L) \twoheadrightarrow F$, where F is the free group on a set T (in 1-1 correspondence with the components of L) such that the i th meridian is sent to the i th generator. Two F -links (M, L, ϕ) and (M', L', ϕ') are equivalent iff (M, L) and (M', L') are isotopic and the maps ϕ and ϕ' differ by an inner automorphism of F . The underlying link of an F -link is a ∂ -link. There is an action

$$\text{String}_g \times F - \text{links} \longrightarrow F - \text{links}$$

whose orbits can be identified with the set of boundary links. Here String_g is the group of group of motions of an unlink in 3-space, and can be described by the automorphisms of the free group that map generators to conjugates of themselves. The action is given by composition with the map ϕ , as is explained in Section 6.1.

Our results are closely related to the *surgery view of boundary links* which we introduced at [GK2]. By this we mean that every F -link can be obtained from the standard unlink (S^3, \mathcal{O}, ϕ) via surgery on a framed nullhomotopic link $C \subset S^3 - \mathcal{O}$ with \mathbb{Z} -invertible linking matrix. C is not unique but its equivalence class

(under a sequence of handle slides and stabilization) is. This was shown in [GK2], where we also constructed a map:

$$W : F\text{-links} \longrightarrow \mathcal{B}(\Lambda \rightarrow \mathbb{Z})$$

where $\mathcal{B}(\Lambda \rightarrow \mathbb{Z})$ is the set of simple stably congruence class of Hermitian matrices A , invertible over \mathbb{Z} . Here, two Hermitian matrices A, B are *simply stably congruent* iff $A \oplus S_1 = P(B \oplus S_2)P^*$ for some diagonal matrices S_1, S_2 with ± 1 entries and some *elementary* matrix P (i.e., one which differs from the identity matrix on a single non-diagonal entry) or a diagonal matrix with entries in $\pm F \subset \Lambda$. The map W sends an F -link to the equivariant linking matrix of \tilde{C} , a lift of a surgery presentation C , to the free cover of $S^3 - \mathcal{O}$. It was shown in [GK2] that the map W determines the Blanchfield form of the F -link, as well as a noncommutative version of the Alexander polynomial defined by Farber [Fa]. Our rational invariant Z^{rat} is a vast generalization of this classical invariant.

Recall that the Kontsevich integral (or rather, its LMO-Aarhus extension) $Z(M, L)$ of a link L in an integral homology 3-sphere M takes values in a completed vector space $\mathcal{A}(\circlearrowright_H)$ over \mathbb{Q} of diagrams whose skeleton is a disjoint union of solid circles $\cup_i \circlearrowright_{h_i}$, one circle labeled by a variable h_i for each component of L , [LMO, A]. The Z^{rat} invariant takes values in a set

$$\mathcal{A}^0(\Lambda_{\text{loc}}) = \mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}(\Lambda_{\text{loc}})$$

where $\mathcal{A}(\Lambda_{\text{loc}})$ is the completed vector space of vertex and edge oriented, trivalent diagrams whose edges are decorated by elements of the *Cohn localization* Λ_{loc} of Λ , [BR], or equivalently, by the ring of *rational functions in noncommuting variables*, [FV]. The diagrams in $\mathcal{A}(\Lambda_{\text{loc}})$ are considered modulo the AS, IHX, Orientation Reversal, Linearity and Holonomy Relations shown in Figure 1.




Figure 1. The AS, IHX (for arbitrary orientations of the edges), Orientation Reversal, Linearity and Holonomy Relations.

$\mathcal{A}(\Lambda_{\text{loc}})$ is a graded vector space, where the degree of a diagram is the number of trivalent vertices, and the completion refers to the above grading. There are some additional ω and String relations on $\mathcal{A}^0(\Lambda_{\text{loc}})$ described in Sections 3 and 5.4).

We now introduce an important map that maps diagrams with beads to power series of diagrams with hair.

Definition 1.1. The Hair $^\nu$ map

$$\text{Hair}^\nu : \mathcal{A}^0(\Lambda_{\text{loc}}) / \langle \omega, \text{String} \rangle \rightarrow \mathcal{A}(\circlearrowright_H)$$

is defined as follows. For $s \in \mathcal{A}(\Lambda_{\text{loc}})$, $\text{Hair}(s)$ replaces a generator t_i of the free group F by an exponential of i -colored legs, labeled by h_i :

$$\uparrow t_i \rightarrow \sum_{n=0}^{\infty} \frac{1}{n!} \begin{array}{c} \uparrow \\ \rightarrow \\ \rightarrow \\ \rightarrow \end{array} n \text{ } h_i\text{-labeled legs}$$

Given a pair $(M, s) \in \mathcal{B}(\Lambda \rightarrow \mathbb{Z}) \times \mathcal{A}^{\text{gp}}$, we define

$$\text{Hair}^\nu(M, s) = \exp_{\sqcup} \left(-\frac{1}{2} \chi(M) \right) \sqcup \text{Hair}(s) \sqcup \nu(h_1) \cdots \sqcup \nu(h_g),$$

where $\nu(h) = Z(S^3, \text{unknot})$ and

$$\chi(M) = \text{tr} \log'(M) := \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{tr}((M(\epsilon M)^{-1} - I)^n)$$

where $\epsilon : \Lambda \rightarrow \mathbb{Z}$. Note that the \exp_{\sqcup} term in the above expression is a disjoint union of *wheels*, and that χ was shown in [GK2] to coincide with an invariant defined by Farber [Fa].

We now have all the ingredients to formulate our first result:

Theorem 1.2. (*Proof in Sections 5.5, 6.5, 7.4*) *There exist an invariant of F-links*

$$Z^{\text{rat}} : F\text{-links} \longrightarrow \mathcal{A}^0(\Lambda_{\text{loc}})/\langle \omega \rangle$$

which descends to and invariant of boundary links

$$Z^{\text{rat}} : \partial\text{-links} \longrightarrow \mathcal{A}^0(\Lambda_{\text{loc}})/\langle \omega, \text{String} \rangle$$

and determines the Kontsevich integral Z of a boundary link

$$Z = \text{Hair}^{\nu} \circ Z^{\text{rat}}.$$

as well as the Blanchfield pairing of a boundary link.

We will call Z^{rat} the *rational Z -invariant* of ∂ -links. The above theorem states that Z^{rat} is a lift of the LMO-Aarhus-Kontsevich invariant of ∂ -links. Let us comment on the above result.

Remark 1.3. First a remark about the empty set. For the unlink \mathcal{O} of g components, we have that $Z^{\text{rat}}(\mathcal{O}) = (\text{empty matrix}, 1) \in \mathcal{A}^0(\Lambda_{\text{loc}})/\langle \omega, \text{String} \rangle$. On the other hand, the Kontsevich intregal of \mathcal{O} is given by $Z(\mathcal{O}) = \nu(h_1) \sqcup \cdots \sqcup \nu(h_g)$ (where $\nu(h_i)$ is the Kontsevich integral of the unknot, whose diagrams are colored by h_i), and

$$\text{Hair}^{\nu}(\text{empty matrix}, 1) = \nu(h_1) \sqcup \cdots \sqcup \nu(h_g).$$

This is how $Z^{\text{rat}}(\mathcal{O})$ determines $Z(\mathcal{O})$, as stated in the above theorem.

Remark 1.4. Erasing a component L_i of a boundary link corresponds to substituting $t_i = 1$. In other words,

$$Z^{\text{rat}}(M, L - L_i) = Z^{\text{rat}}(M, L)|_{t_i=1}.$$

Remark 1.5. The Z^{rat} invariant of F -links takes values in a rather strange quotient $\mathcal{A}^0(\Lambda_{\text{loc}})/\langle \omega \rangle$ of a set of pairs of matrices and vectors \mathcal{A}^0 . If we restrict $Z^{\text{rat}}|_{\mathcal{S}}$ to a class \mathcal{S} of F -links with the same Blanchfield pairing (or the same S -equivalence class), then $Z^{\text{rat}}|_{\mathcal{S}}$ is equivalent to an invariant

$$\mathcal{S} \longrightarrow \mathcal{A}(\Lambda_{\text{loc}})/\langle \omega \rangle$$

that takes values in a *graded vector space*, where the degree of a graph is the number of trivalent vertices. In addition, the edges of the trivalent graphs are decorated by elements of a subring of Λ_{loc} that depends on \mathcal{S} . In particular, for good boundary links (i.e., for links whose free cover is acyclic, or equivalently, whose Blanchfield pairing is trivial) Z^{rat} is equivalent to an invariant with values in the more manageable graded vector space $\mathcal{A}(\Lambda)/\langle \omega, \text{String} \rangle$; see Section 5.5.

Remark 1.6. For $g = 1$, the sets of 1-component F -links and ∂ -links coincide with the set of knots in integral homology 3-spheres, and the relations ω and String_1 are trivial, see Lemmas 3.11 and 6.8. In this case, Z^{rat} takes values in $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}(\Lambda_{\text{loc}})$, and the matrix-free part of Z^{rat} takes values in the *graded algebra* $\mathcal{A}(\Lambda_{\text{loc}})$, where the multiplication in $\mathcal{A}(\Lambda_{\text{loc}})$ is given by the disjoint union of graphs. It turns out that the matrix-free part of Z^{rat} is a group-like element in $\mathcal{A}(\Lambda_{\text{loc}})$, i.e., it equals to the exponential of a series of *connected* graphs.

For $g = 2$ (i.e., for two component ∂ -links) the relation String_g is trivial, see Lemma 6.8 and the set of 2-component F -links coincides with the one of ∂ -links.

Remark 1.7. [P] has shown that the Hair map is not 1-1 (even in the case of knots), thus the lift Z^{rat} of Z is not unique. A *weak form* of the RC Conjecture for ∂ -links states that the image of Z lies in the image of the map $\text{Hair}^{\nu} : \mathcal{A}^0(\Lambda_{\text{loc}})/\langle \omega, \text{String} \rangle \rightarrow \mathcal{A}(\odot_H)$. A *strong form* of the RC Conjecture for knots states that the Kontsevich integral lifts. The weak form of RC Conjecture for knots was formulated by Rozansky in [R2] and was proven by the second named author in [Kr1]. The strong form of the RC Conjecture for F -links was formulated by the first named author in [Ga] and is proven in Theorem 1.2 above.

Remark 1.8. The reader may wonder why in the construction of Z^{rat} we needed to introduce diagrams with beads in Λ_{loc} . For two independent reasons, see Remark 5.2.

Remark 1.9. In a sense, Z is a graph-valued $\hat{\Lambda}$ -invariant of ∂ -links, where $\hat{\Lambda}$ is the completion of $\Lambda \otimes \mathbb{Q}$ with respect to the augmentation ideal whereas Z^{rat} is a graph-valued Λ_{loc} -invariant. It often happens that $\hat{\Lambda}$ -invariants can be lifted to Λ_{loc} -invariants, see for example the discussion of Farber-Ranicki, [FRa], on a problem of Novikov about circle-valued Morse theory. It would be nice if an extension of this circle-valued Morse theory to infinite dimensions (using the circle-valued Chern-Simons action on the space of gauge equivalence classes of connections on a principal bundle over a 3-manifold) would lead to an explanation for the existence of the Z^{rat} -invariant.

Remark 1.10. As will become apparent from the construction of the Z^{rat} invariant, it is closely related to equivariant linking matrices of nullhomotopic links in the complement of an F -link, see Proposition 8.2. It is an interesting question to ask whether the Z^{rat} invariant of an F -link can be defined in terms of an equivariant construction on the free cover of the F -link.

The next theorem characterizes Z^{rat} by a universal property. Using the terminology of [Ga], recall that a *null move* on an F -link (M, L, ϕ) can be described by surgery on a clover G in $M - L$ whose leaves lie in the kernel of ϕ , [Ga]. The result $(M, L, \phi)_G$ of surgery is another F -link. As usual, one can define a notion of finite type invariants on the set of F -links using null moves; we will call the corresponding filtration the *Euler filtration*. The number of invariants of degree n can be identified with a quotient vector space $\mathcal{G}_n^{\text{null}}$. For simplicity, we will confine ourselves to the case of *acyclic F -links*, that is F -links whose free cover X^ω satisfies $\bar{H}_*(X^\omega, \mathbb{Z}) = 0$. In [Ga], it was shown that there is an onto map

$$\mathcal{A}(\Lambda)/\langle \omega \rangle \longrightarrow \mathcal{G}^{\text{null}}$$

which preserves degrees. The next theorem constructs an inverse to this map and characterizes Z^{rat} by a universal property.

Theorem 1.11. *(Proof in Section 8) Z^{rat} is a universal $\mathcal{A}(\Lambda)/\langle \omega \rangle$ -valued finite type invariant of acyclic F -links with respect to the Euler filtration. In particular, we have that $\mathcal{A}(\Lambda) \cong \mathcal{G}^{\text{null}}$.*

1.3. The main ideas of the proof. The construction of the rational invariant Z^{rat} is obtained in the following steps:

- We begin by identifying the set of F -links with the set $\mathcal{N}(\mathcal{O})$ of nullhomotopic links C in the complement of an unlink, modulo Kirby moves. The linking matrix of C is invertible over \mathbb{Z} . This is the so-called surgery view of F -links.
- In Section 4 we define an invariant of links C as above. It takes values in a vector space of univalent graphs with beads. The beads are elements of Λ , and record the winding of a tangle representative of C in $S^3 - \mathcal{O}$. The legs of the graphs are labeled by the components of C . The strut part of this invariant records the equivariant linking matrix of C .
- Unfortunately, this invariant of links C is not independent under Kirby moves of C . To accommodate for that, in Section 5 we construct an integration theory \int^{rat} in the spirit of the Aarhus Integral [A]. In that integration theory, we separate the strut part of a diagram with beads, invert their matrix (it is here that Λ_{loc} is needed) to construct new struts, and glue the legs of these new struts to the rest of the diagrams. The result is a formal linear combination of trivalent graphs whose beads are elements of Λ_{loc} . The main property of this integration theory, is that the resulting invariant is independent under Kirby moves and gives an invariant of F -links.
- The comparison between the Z^{rat} invariant and the Kontsevich integral Z of boundary links is done in Section 7, using a *magic formula* for the Kontsevich integral of the Long Hopf Link, conjectured in [A0] (in conjunction with the Wheels and Wheeling Conjectures) and proven in [BLT]. Section 7.4 completes the proof of Theorem 1.2.
- Section 8 is devoted to the proof of Theorem 1.11, namely the universal property of Z^{rat} . This follows from the general principle of locality of the invariant Z^{rat} (i.e., with the behavior of Z^{rat} under sublinks) together with the identification of the covariance of Z^{rat} with an equivariant linking function.
- The paper ends with some appendices on diagrammatic integration and its various identities.

1.4. Acknowledgement. The results of this work were first announced by the first named author in a meeting on “Influence of Physics on Topology” on August 2000, in the sunny San Diego. We thank the organizers, M. Freedman and P. Teichner for arranging such a fertile meeting. We also wish to express our gratitude to J. Levine who shared with the first author his expertise on topology, and to J. Hillman and T. Ohtsuki for encouragement and support of the second author.

CONTENTS

1. Introduction	1
1.1. Introduction	1
1.2. Statement of the results	2
1.3. The main ideas of the proof	5
1.4. Acknowledgement	6
2. A surgery view of F -links	7
2.1. A surgery description of F -links	7
2.2. A tangle description of F -links	7
3. An assortment of diagrams and their relations	9
3.1. A review of the Cohn localization	11
3.2. The infinitesimal wrapping relations	12
3.3. Group-like relations	13
4. An invariant of links in the complement of an unlink	16
5. A rational version of the Aarhus integral	18
5.1. The definition of \int^{rat}	19
5.2. \int^{rat} respects the basing relations	19
5.3. Dealing with the Kirby moves	20
5.4. Dealing with the wrapping relations	22
5.5. The definition of the rational invariant Z^{rat} of F -links	23
6. The String action and invariants of boundary links	23
6.1. A surgery description of boundary links	24
6.2. The action of String on diagrams	25
6.3. Z^{rat} commutes with the String action	27
6.4. \int^{rat} commutes with the String_g action	28
6.5. The definition of the rational invariant Z^{rat} of ∂ -links	29
7. A comparison between Z and Z^{rat}	29
7.1. The Hair map	29
7.2. A comparison between Z^{rat} and Z	30
7.3. A comparison between \int^{rat} and \int	31
7.4. End of the proof of Theorem 1.2	32
8. A universal property of Z^{rat}	32
8.1. A relative version of the Z^{rat} invariant	33
8.2. Surgery on clasps	33
8.3. Counting above the critical degree	34
8.4. Counting on the critical degree	34
Appendix A. A brief review of diagrammatic calculus	35
A.1. Some useful identities	35
A.2. Iterated Integration	36
A.3. Integration by parts	37
Appendix B. The Wheels Identity	39
Appendix C. Infinitesimal versus group-like wrapping relations	41
Appendix D. Infinitesimal basing relations	45
Appendix E. A useful calculation	46
References	47

2. A SURGERY VIEW OF F -LINKS

2.1. A surgery description of F -links. In this section we recall the surgery view of links, introduced in [GK2], which is a key ingredient in the construction of the rational invariant Z^{rat} . Let us recall some important notions. Fix once and for all a based unlink \mathcal{O} of g components in S^3 .

Definition 2.1. Let $\mathcal{N}(\mathcal{O})$ denote the set of *nullhomotopic links C with \mathbb{Z} -invertible linking matrix* in the complement of \mathcal{O} .

Surgery on an element C of $\mathcal{N}(\mathcal{O})$ transforms $(S^3, \mathcal{O} \cup C)$ to a F -link (M, L, ϕ) . Indeed, since C is nullhomotopic the natural map $\pi_1(S^3 - \mathcal{O}) \rightarrow F$ gives rise to a map $\pi_1(M - L) \rightarrow F$. Alternatively, one can construct disjoint Seifert surfaces for each component of L by tubing the disjoint disks that \mathcal{O} bounds, which is possible, since each component of C is nullhomotopic.

Since the linking matrix of C is invertible over \mathbb{Z} , M is an integral homology 3-sphere. Let \approx denote the equivalence relation on $\mathcal{N}(\mathcal{O})$ generated by the moves of *handle-slide* κ_1 (i.e., adding a parallel of a link component to another component) and *stabilization* κ_2 (i.e., adding to a link an unknot away from the link with framing ± 1). It is well-known that κ -equivalence preserves surgery. A main result of [GK2] is the following:

Theorem 2.2. ([GK2]) *The surgery map gives a 1-1 and onto correspondence*


$$\mathcal{N}(\mathcal{O}) / \langle \kappa \rangle \rightarrow F\text{-links.}$$

2.2. A tangle description of F -links. In this section we give a tangle diagram description of F -links and ∂ -links which is useful in defining invariants. We begin by giving a tangle description of the set $\mathcal{N}(\mathcal{O})$. Before we proceed, a remark on notation:

Remark 2.3. Throughout the paper, given an equivalence relation ρ on a set X , we will denote by $X / \langle \rho \rangle$ the set of *equivalence classes*. Occasionally, an equivalence relation on X will be defined by one of the following ways:

- Either by the *action* of a group G on X , in which case $X / \langle \rho \rangle$ coincides with the set of orbits of G on X .
- Or by a *move* on X , i.e., by a subset of $X \times X$, in which case the equivalence relation is the smallest one that contains the move.

Consider the following surfaces in \mathbb{R}^2



together with a distinguished part of their boundary (called a *gluing site*) marked by an arrow in the figure above. By a *tangle on a surface Σ_x* (for $x = a, b, c$) we mean an oriented, framed, smooth, proper immersion of an oriented 1-manifold into the surface, up to isotopies rel boundary of the surface, double points being equipped with crossing information, with the boundary points of the tangle lying on standard points at $t = 0$ and $t = 1$.

A *crossed tangle on a surface Σ_x* is a tangle diagram on Σ_x possibly with some crosses marking those boundary points of the diagram that lie either on a $t = 0$ edge of Σ_x for $x = a, b$ or on a $t = 1$ edge of Σ_c . The following is an example of a crossed tangle diagram on Σ_b :



The cross notation evokes a small pair of scissors \times that will be sites where the skeleton of the tangle will be cut. A *sliced crossed link on D_g* is a sequence of crossed tangles, $\{T_1, \dots, T_k\}$ such that

- the boundaries of the surfaces match up, that is, their shape, distribution of endpoints, and crosses.


For example, if T_i is of the form  then T_{i+1} is of the form 

- the top and bottom boundaries of T_1 and T_k look like:



- After stacking the tangles $\{T_1, \dots, T_k\}$ from top to bottom, we obtain a crossed link in $D_g \times I$, where D_g is a disk with g holes. Note that the holes in D_g are ordered from the top one to the bottom one and that D_g is equipped with g gluing sites.


Clearly, every crossed link in $D_g \times I$ can be obtained this way. Here is an example of a sliced crossed link in D_1 whose corresponding link in $\mathcal{N}(\mathcal{O})$ is a surgery presentation for the Figure 8 knot:



Let $\mathcal{D}(D_g)$ denote the set of sliced crossed links on D_g such that each component of the associated link is nullhomotopic in $D_g \times I$ and marked with precisely one cross.


We now introduce some equivalence relations on $\mathcal{D}(D_g)$ that are important in comparing $\mathcal{D}(D_g)$ to $\mathcal{N}(\mathcal{O})$.

Regular isotopy \sim^r of sliced crossed links is the equivalence relation generated by regular isotopy of individual tangles and the following moves:




where the glued edge of T_e must have no crosses marking it.

Basing relation \sim^β of sliced crossed links is the equivalence relation generated by moving a cross of each component of the associated link in some other admissible position in the sliced crossed link. Notice that moving the cross of a sliced crossed link can be obtained (in an equivalent way) by the local moves β_1 and β_2 , where β_1 (resp. β_2) moves the cross across consecutive tangles in such a way that a gluing site is not (resp. is) crossed. Here is an example of a β_2 move:



The *wrapping relation* \sim^ω of sliced crossed links is generated by the following move



Recall that Theorem 2.2 identifies the set of F -links with a quotient of $\mathcal{N}(\mathcal{O})$. Since we are interested to give a description of the set of F -links and $\mathcal{N}(\mathcal{O})$ with a quotient of $\mathcal{D}(D_g)$, we need to introduce analogs of the κ -relation on $\mathcal{D}(D_g)$.

The *handleslide move* $\overset{\kappa}{\sim}$ is generated by the move:

$$\{\dots, T_{i-1}, T_i, \begin{array}{|c|c|} \hline \times & \times \\ \hline x_i & x_j \\ \hline \end{array}, T_{i+2}, \dots\} \longleftrightarrow \{\dots, T'_{i-1}, T'_i, \begin{array}{|c|c|} \hline \times & \\ \hline x_i & x_j \\ \hline \times & \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \times \\ \hline x_i & x_j \\ \hline \end{array}, T'_{i+2}, \dots\}.$$

where each tangle in $\{T'_k\}$ is obtained from the corresponding tangle in $\{T_k\}$ by taking a framed parallel of every component contributing to the component x_j ,

Proposition 2.4. *There is a 1-1 correspondence*

$$\mathcal{D}(D_g)/\langle r, \beta, \omega \rangle \longleftrightarrow \mathcal{N}(\mathcal{O}) \quad \text{and} \quad \mathcal{D}(D_g)/\langle r, \beta, \omega, \kappa \rangle \longleftrightarrow F\text{-links}.$$

Proof. The first identification is well-known without crosses, see for example [AMR] and [Tu, Figure 7]. The idea is that it follows from general position arguments analogous to those that prove the sufficiency of the Reidemaster moves for planar knot projections. The addition of crosses presents no extra difficulty. The second identification follows easily from the first, together with the following observation that takes care of inner automorphisms: the group $\text{Inn}(F)$ of inner automorphisms acts trivially on the set of F -links (by definition) and it also acts trivially on the set $\mathcal{D}(D_g)/\langle r, \beta, \omega \rangle$. The proof of this observation is a pleasant exercise, left to the reader. \square

3. AN ASSORTMENT OF DIAGRAMS AND THEIR RELATIONS

In this section we explain what are the spaces of diagrams in which our invariants will take values.

We begin with some notation, which generalizes the notion of the \mathcal{A} -groups introduced in [GL1]: given a ring R with a distinguished group of units U , and (possibly empty sets) X, Y, T , $\mathcal{D}(\uparrow_X, \star_Y, \otimes_T, R, U)$ is the completed graded vector space over \mathbb{Q} generated by

- Uni-trivalent diagrams with skeleton \uparrow_X , with symmetric univalent vertices labeled by $Y \cup T$.
- The diagrams have oriented edges and skeleton and each edge is labeled by an element of R , such that the edges that are part of the skeleton are labeled *only* by U . Moreover, the product of the labels along each component of the skeleton is 1. Labels on edges or part of the skeleton will be called *beads*.

The *degree* of a diagram is the number of trivalent vertices, and the completion is with respect to this degree. Empty sets will be omitted from the notation, and so will U , the selected group of units of R . For example, $\mathcal{A}(\star_Y, R)$, $\mathcal{A}(R)$ and $\mathcal{A}(\phi)$ stands for $\mathcal{A}(\uparrow_\phi, \star_Y, \otimes_\phi, R, U)$, $\mathcal{A}(\uparrow_\phi, \star_\phi, \otimes_\phi, R, U)$ and $\mathcal{A}(\uparrow_\phi, \star_\phi, \otimes_\phi, \mathbb{Z}, 1)$ respectively. Univalent vertices of diagrams will often be called *legs*. Diagrams will sometimes be referred to as *graphs*. Special diagrams, called *struts*, labeled by a, c with bead b are drawn as follows

$$\begin{array}{c} a \\ \uparrow \\ \bullet \\ \downarrow \\ c \end{array} b.$$

oriented from bottom to top. Multiplication of diagrams D_1 and D_2 , unless otherwise mentioned, means their disjoint union and will be denoted by $D_1 D_2$ and occasionally by $D_1 \sqcup D_2$.

We will only be interested in the rings $\Lambda = \mathbb{Q}[F]$ (the group ring of the free group F on some fixed set $T = \{t_1, \dots, t_g\}$ of generators), its completion $\hat{\Lambda}$ (with respect to the powers of the augmentation ideal) and its *Cohn localization* Λ_{loc} (i.e., the localization with respect to the set of matrices over Λ that are invertible over \mathbb{Z}) all with selected group of units F . $\Lambda, \hat{\Lambda}$ and Λ_{loc} are rings with involution induced by $\bar{g} = g^{-1}$ for $g \in F$, with augmentation over \mathbb{Q} , and with a commutative diagram

$$\begin{array}{ccc} \Lambda & \xrightarrow{\quad} & \Lambda_{\text{loc}} \\ & \searrow & \swarrow \\ & \hat{\Lambda} & \\ & 9 & \end{array}$$

where $\Lambda \rightarrow \hat{\Lambda}$ is given by the modern version of the Magnus expansion $t_i \rightarrow e^{h_i}$. $\hat{\Lambda}$ can be identified with the ring of noncommuting variables $\{h_1, \dots, h_g\}$ and Λ_{loc} with the ring of rational functions in noncommuting variables $\{t_1, \dots, t_g\}$, [FV, BR].

We now collect various relations on spaces of diagrams. We define

$$\mathcal{A}(\uparrow_X, \star_Y, R, U) = \mathcal{D}(\uparrow_X, \star_Y, R, U) / (\text{AS, IHX, Multilinear, Holonomy})$$

where AS, IHX and Multilinear are the relations shown in Figure 2 and Holonomy is the relation shown in Figure 3. Note that the AS and IHX are well-known relations in the theory of finite type invariants of links and 3-manifolds, whereas the Multilinear and Holonomy are more recent, first appearing in work of [GL1] in relation to finite type invariants of homology surgery and in [GR]. Note also that the above relations on Δ are homogenous with respect to the grading, thus \mathcal{A} is a completed graded vector space, too.




Figure 2. The AS, IHX (for arbitrary orientations of the edges), Orientation Reversal and Linearity relations. Here $r \rightarrow \bar{r}$ is the involution of R , $r, s \in R$.




Figure 3. The Holonomy relation that pushes a unit $g \in U$ past a trivalent vertex.

The following is a companion to Remark 2.3:

Remark 3.1. Throughout the paper, given a subspace W of a vector space V , we will denote by $V/(W)$ the quotient of V modulo W . Occasionally, W will be defined by one of the following ways:

- Either by a *linear action* of a group G on V , in which case $W = \{gv - v\}$ and $V/(W)$ coincides with the space of *coinvariants* of G on V .
- Or by a subset of V , in which case W is the subspace spanned by that subset.

We now introduce on \mathcal{A} algebraic analogs of the geometric wrapping relations of sliced crossed links in D_g . We begin by defining the infinitesimal wrapping relations ω on \mathcal{A} . This will require an enlargement of the vector space \mathcal{A} by a set of matrices as stated in Theorem 1.2, as well as some analysis of the notion of localization. We begin with the latter.

3.1. A review of the Cohn localization. In this self-contained section we review several facts about the Cohn localization that are used in the text. Consider the free group F_g on generators $\{t_1, \dots, t_g\}$. The Cohn localization Λ_{loc} of its group-ring $\Lambda = \mathbb{Q}[F_g]$ is characterized by a universal property,

$$\begin{array}{ccc} \Lambda & \longrightarrow & \Lambda_{\text{loc}} \\ & \searrow \alpha & \downarrow \bar{\alpha} \\ & & R \end{array}$$

namely that for every Σ -inverting ring homomorphism $\alpha : \Lambda \rightarrow R$ there exists a unique ring homomorphism $\bar{\alpha} : \Lambda_{\text{loc}} \rightarrow R$ that makes the above diagram commute, [C].

Farber and Vogel identified Λ_{loc} with the ring of *rational functions in noncommuting variables*, [FV, BR]. An example of such a rational function is

$$(3 - t_1^2 t_2^{-1} (t_3 + 1) t_1^{-1})^{-1} t_2 - 5.$$

There is a close relation between rational functions in noncommuting variables and *finite state automata*, sedcribed in [BR, FV]. Farber-Vogel showed that every element $s \in \Lambda_{\text{loc}}$ can be represented as a solution to a system of equations $M\vec{x} = \vec{b}$ where M is a matrix over Λ , invertible over \mathbb{Z} , and \vec{b} is a column vector over Λ , [FV, Proposition 2.1]. More precisely, we have that

$$s = (1, 0, \dots, 0)M^{-1}\vec{b},$$

which we will call a *matrix presentation* of $s \in \Lambda_{\text{loc}}$.

In the text, we often use substitutions of the form $\varphi_{t_i \rightarrow t_i e^h}$, or $\varphi_{t_i \rightarrow e^h t_i}$ or $\varphi_{t_i \rightarrow e^{-h} t_i e^h}$, where h does not commute with F_g . In order to make sense of these substitutions, we need to enlarge our ring Λ_{loc} . This can be achieved in the following way. F_g can be included in the free group F with one additional generator t . Let \mathcal{L} denote the group-ring of F , \mathcal{L}_{loc} denote its Cohn localization and $\widehat{\mathcal{L}}_{\text{loc}}$ denote the completion of \mathcal{L}_{loc} with respect to the ideal generated by $t - 1$. There is an identification of $\widehat{\mathcal{L}}_{\text{loc}}$ with the following ring \mathcal{R} whose elements consist of formal sums of the form $\sum_{\mathbf{m}, \mathbf{f}} \prod_{i=1}^{\infty} h^{m_i} f_i$ where $f_i \in \Lambda_{\text{loc}}$ and

- $\mathbf{m} = (m_1, m_2, \dots) : \mathbb{N} \rightarrow \mathbb{N}$ is eventually 0 and $\mathbf{f} = (f_1, f_2, \dots) : \mathbb{N} \rightarrow \Lambda_{\text{loc}}$ is eventually 1.
- In the above sum, for each fixed k , there are finitely many sequences \mathbf{m} with $|\mathbf{m}| = \sum_i m_i \leq k$.

It is easy to see that \mathcal{R} is a ring with involution (in fact a subring of the completion of \mathcal{L} with respect to the augmentation ideal) such that t_i and e^h are units, where $h = \log t$. In particular, $\widehat{\mathcal{L}}_{\text{loc}}$ is an h -graded ring. Let $\text{deg}_h^n : \widehat{\mathcal{L}}_{\text{loc}} \rightarrow \widehat{\mathcal{L}}_{\text{loc}}^n$ denote the projection on the h -degree n part of $\widehat{\mathcal{L}}_{\text{loc}}$. There is a map $\varphi_i : \Lambda \rightarrow \widehat{\mathcal{L}}_{\text{loc}}$ given by substituting t_i for $e^{-h} t_i e^h$ for $1 \leq i \leq g$. It extends to a map $\varphi_i : \Lambda_{\text{loc}} \rightarrow \widehat{\mathcal{L}}_{\text{loc}}$. Indeed, using the defining property of the Cohn localization, it suffices to show that φ_i is Σ -inverting, [C, FV]. In other words, we need to show that if a matrix W over Λ is invertible over \mathbb{Z} , then $\varphi_i(W)$ is invertible over $\widehat{\mathcal{L}}_{\text{loc}}$. Since $\widehat{\mathcal{L}}_{\text{loc}}$ is a completion of \mathcal{L}_{loc} , it suffices to show that $\text{deg}_h^0(\varphi_i(W))$ is invertible over \mathcal{L}_{loc} . However, $\text{deg}_h^0(\varphi_i(W)) = W$, invertible over Λ_{loc} and thus also over \mathcal{L}_{loc} . The result follows.

Let

$$\eta_i : \Lambda_{\text{loc}} \rightarrow \widehat{\mathcal{L}}_{\text{loc}}^1$$

denote the h -degree 1 part of φ_i . The following lemma gives an axiomatic definition of η_i by properties analogous to a *derivation*. Compare also with the derivations of Fox differential calculus, [FV].

Lemma 3.2. η_i is characterized by the following properties:

$$\begin{aligned} \eta_i(t_j) &= \delta_{ij}[t_i, h] = \delta_{ij}(t_i h - h t_i) \\ \eta_i(ab) &= \eta_i(a)b + a\eta_i(b) \\ \eta_i(a + b) &= \eta_i(a) + \eta_i(b). \end{aligned}$$

Proof. It is easy to see that η_i satisfies the above properties, by looking at the h -degree 0 and 1 part in $\varphi_i(a)$. This determines η_i uniquely on Λ . By the universal property, we know that η_i extends over Λ_{loc} ; a priori the extension need not be unique. Since however, η_i extends uniquely to matrices over Λ , and since every

element of Λ_{loc} has a matrix presentation (see the discussion in the beginning of the section), it follows that η_i is characterized by the derivation properties. \square

3.2. The infinitesimal wrapping relations. We have all the required preliminaries to introduce an infinitesimal wrapping relation on $\mathcal{A}^0 = \mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}$. Given a vector $v \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ and a diagram $D \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ with a *single* special leg colored by ∂h and $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$, the *infinitesimal wrapping relation* ω_i at the i th place in $\text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ is generated by the move

$$(M, v) \stackrel{\omega_i}{\sim} \left(M, v + \text{con}_{\{h\}} \left(\eta_i(D) - \frac{1}{2} D \sqcup \begin{array}{c} \curvearrowright \\ \bullet \end{array} \text{tr}(M^{-1} \eta_i(M)) \right) \right).$$

It is easy to see that the above definition depends only on the image of M in $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$. We define

$$\mathcal{A}^0(\star_X, \Lambda_{\text{loc}}) / \langle \omega \rangle = \mathcal{A}^0(\star_X, \Lambda_{\text{loc}}) / \langle \omega_1, \dots, \omega_g \rangle.$$

We give two examples of infinitesimal wrapping relations:

Example 3.3. In this example, the beads of a diagram are monomials t_j , M is the empty matrix, and *all* the edges labeled by t_i are shown:

Note that in the ω relation, we allow for D to be a diagram of a circle (with a bead and with a special ∂h leg). In addition, we allow the ∂h leg to have a bead in Λ_{loc} .

The next example illustrates the infinitesimal wrapping relation in its full complication.

Example 3.4. Consider the matrix

$$M = \begin{bmatrix} 1 & t_3 - t_2 t_1^{-1} \\ t_3^{-1} - t_1 t_2^{-1} & 1 \end{bmatrix}$$

which is Hermitian over Λ , and invertible over \mathbb{Z} . Lemma 3.2 and the following property of η_i ,

$$\eta_i(AB) = \eta_i(A)B + A\eta_i(B)$$

valid for not necessarily square matrices A, B of appropriate size, implies that


$$\begin{aligned} \eta_1(M^{-1}) &= M^{-1}(t_2 t_1^{-1}, 0)' h(0, 1) M^{-1} + M^{-1}(-t_2, 0)' h(0, t_1^{-1}) M^{-1} \\ &+ M^{-1}(0, t_1)' h(t_2^{-1}, 0) M^{-1} + M^{-1}(0, -1)' h(t_1 t_2^{-1}, 0) M^{-1}. \end{aligned}$$

Further, observe that


$$\begin{aligned} -1/2 \text{tr}(M^{-1} \eta_1(M)) &= (1/2, 0) M^{-1}(0, t_1)' h t_2^{-1} + (1/2, 0) M^{-1}(0, -1)' h t_1 t_2^{-1} \\ &+ (0, 1/2) M^{-1}(t_2 t_1^{-1}, 0)' h + (0, 1/2) M^{-1}(-t_2, 0)' h t_1^{-1}. \end{aligned}$$

Now, consider the infinitesimal wrapping relation arising from the diagram


That, is we have $(M, v) \stackrel{\omega_1}{\sim} (M, v + x)$, where $x = x_1 + x_2$, and x_1 is given by gluing the $(h, \partial h)$ legs in each of the following diagrams,



as well as these diagrams



and summing the result up, and x_2 is given by gluing the $(h, \partial h)$ legs of the disjoint union of D with the following sum:



Notice that x is a sum of diagrams of degree 2 of the form Θ and $\bigcirc\bigcirc$. It is an interesting question to understand the infinitesimal wrapping relation in degree 2.

3.3. Group-like relations. Although the invariant Z^{rat} that appears in Theorem 1.2 takes values in \mathcal{A}^0 , we will actually construct a refined invariant $= Z^{\text{rat,sp}}$ which takes values in a set of group-like elements, modulo appropriate equivalence relations.


Let \mathcal{A}^{sp} denote the set of group-like elements of \mathcal{A} i.e., elements of the form $\exp(a)$ where a is a sum of connected diagrams.

We now define an analog of the basing relations on \mathcal{A}^{sp} . Let $\vec{m}_z^{x,y} : \mathcal{A}^{\text{sp}}(\uparrow_{\{x,y\}} \cup \uparrow_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^{\text{sp}}(\uparrow_{\{z\}} \cup \uparrow_X, \Lambda_{\text{loc}})$ be the map that glues the end of the x -skeleton to the beginning of the y -skeleton and then relabeling the skeleton by z , for $x, y, z \notin X$. Given $s_1, s_2 \in \mathcal{A}^{\text{sp}}(\star_X, \Lambda_{\text{loc}})$, we say that


- $s_1 \stackrel{\beta_1^{\text{sp}}}{\sim} s_2$, if there exists an element $s \in \mathcal{A}^{\text{sp}}(\uparrow_{\{x\}} \uparrow_{\{x'\}} \uparrow_{X-\{x\}}, \Lambda_{\text{loc}})$ with the property that

$$(1) \quad \sigma(\vec{m}_x^{x,x'}(s)) = s_1 \quad \text{and} \quad \sigma(\vec{m}_x^{x',x}(s)) = s_2.$$

- $s_1 \stackrel{\beta_2^{\text{sp}}}{\sim} s_2$ iff s_2 is obtained from s_1 by pushing an element of F on each of the X -labeled legs of s_1 . For example, if all x_1 legs are shown below, we have:



Remark 3.5. When we push an element of the free group f in the legs of a diagram, depending on the orientation of the leg, we add f or \bar{f} to the leg. See also Figure 3 for a convention for all choices of orientations. Similarly, when we glue two legs of diagrams together, the label on the bead depends on the orientations of the legs. For example,



We define

$$\mathcal{A}^{\text{gp}}(\otimes_X, \Lambda_{\text{loc}}) = \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}}) / \langle \beta_1^{\text{gp}}, \beta_2^{\text{gp}} \rangle.$$

We mention a useful reformulation of the β_1^{gp} relation for diagrams in $\mathcal{A}(\star_X, \Lambda_{\text{loc}})$, which makes obvious the formal similarity between the β_1^{gp} and β_2^{gp} relations. In order to do so, it is convenient to introduce a *contraction map* $\text{con} : \mathcal{A}(\cdots \star_{X \cup \partial X}, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\cdots, \Lambda_{\text{loc}})$, defined as follows. For $s \in \mathcal{A}(\cdots \star_{X \cup \partial X}, \Lambda_{\text{loc}})$,

$$\text{con}_X(s) = (\text{div}_X)|_{X \rightarrow 0}.$$

In words, contraction glues all ways of pairing *all* legs labeled from ∂X to *all* legs labeled from X . For $s_1, s_2 \in \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$, we say that

- $s_1 \stackrel{\beta_1^{\text{gp}'}}{\sim} s_2$ iff there exists $s_{12} \in \mathcal{A}^{\text{gp}}(\star_{X \cup \partial h}, \Lambda_{\text{loc}})$ such that

$$s_1 = \text{con}_{\{h\}}(s_{12}) \quad \text{and} \quad s_2 = \text{con}_{\{h\}}(s_{12}|_{X \rightarrow X e^h})$$


where $s_{12}|_{X \rightarrow X e^h}$ is by definition the result of pushing e^h to each X -labeled leg of s_{12} .

Lemma 3.6. *The equivalence relations $\beta_1^{\text{gp}'}$ and β_1^{gp} are equal on $\mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$.*

Proof. Suppose that $s_1 \stackrel{\beta_1^{\text{gp}'}}{\sim} s_2$ and consider the corresponding element s from Equation (1). We start by presenting s as a result of a contraction. Let $\alpha = \uparrow_x e^h (\sigma_{\{x\}}(s))|_{x \rightarrow \partial h}$. Lemma A.1 implies that

$$s = \text{con}_{\{h\}}(s(\partial h)), \quad s_1 = (\text{con}_{\{h\}} \circ \sigma_X) \left(\vec{m}_x^{x, x'}(\alpha) \right), \quad s_2 = (\text{con}_{\{h\}} \circ \sigma_X) \left(\vec{m}_x^{x', x}(\alpha) \right).$$

Now the term $\vec{m}_x^{x, x'}(\alpha)$ is obtained from the term $\vec{m}_x^{x', x}(\alpha)$ by conjugating by an e^h . Indeed:



Conjugating with e^h commutes with the symmetrization map σ_X (Exercise!). Letting

$$s_{12} = \text{con}_{\{h\}} \left(\sigma_X \left(\vec{m}_x^{x, x'}(\alpha) \right) |_{\partial h \rightarrow \partial h + \partial h'} \right) \in \mathcal{A}^{\text{gp}}(\star_{X \cup \partial h'}, \Lambda_{\text{loc}}),$$

it follows that

$$s_1 = \text{con}_{\{h'\}}(s_{12}) \quad \text{and} \quad s_2 = \text{con}_{\{h'\}}(s_{12}|_{x \rightarrow x e^{h'}})$$

as needed. The converse can be obtained by reversing the steps in the above proof. \square

Now, we define a group-like wrapping relation on \mathcal{A}^{gp} . This enlargement might sound artificial, however there are several good reasons for doing so. For example, the null move on the set of F -links and its associated filtration predicts that in degree 0, the universal invariant is given by the S-equivalence class (of F -links), or equivalently, by the set of Blanchfield pairings of F -links. $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}})$, defined below, maps onto $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ which in turn maps onto the set of Blanchfield pairings of F -links.

In addition, the \mathcal{A} -groups, studied in relation to Homology Surgery in [GL1], are secondary obstructions to the vanishing of a Witt-type obstruction which lies in $\mathcal{B}(\mathbb{Z}[\pi] \rightarrow \mathbb{Z})$. In the case of F -links, this motivates the fact that the set $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ should be taken together with $\mathcal{A}(\Lambda_{\text{loc}})$. With these motivations, we will consider the set

$$\mathcal{A}^{\text{gp}, 0}(\star_X, \Lambda_{\text{loc}}) = \mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}}).$$

Definition 3.7. Two pairs $(M, s_m) \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$ $m = 1, 2$ are related by a *wrapping move* at the j th gluing site (for some j such that $1 \leq j \leq g$) if there exists an $s \in \mathcal{A}^{\text{gp}}(\star_X, \partial h, \Lambda_{\text{loc}})$ such that:

$$s_m = \text{con}_{\{h\}} \left(\varphi_m(s) \sqcup \left(\exp_{\sqcup} \left(-\frac{1}{2} \chi_h(M^{-1} \varphi_m M) \right) \right) \right)$$

for $m = 1, 2$ where φ_1 is the identity and φ_2 is the substitution $\varphi_{t_j \rightarrow e^{-h} t_j e^h}$, and

$$\chi_h(A) = \text{tr} \log(A) \in \mathcal{A}(\star_h, \Lambda_{\text{loc}}).$$

Our first task is to make sure that the above definition makes sense. Using the h -graded ring with involution $\widehat{\mathcal{L}}_{\text{loc}}$ from Section 3.1, it follows that $\varphi_m(s_{12}) \in \mathcal{A}^{\text{gp}}(\star_X, \widehat{\mathcal{L}}_{\text{loc}})$. Furthermore, $\chi_h(M^{-1} \varphi_m M)$ can be thought of either as a circle with a bead in $\widehat{\mathcal{L}}_{\text{loc}}/(\text{cyclic})$ that has vanishing h -degree 0 part, or as a sum of wheels with h -colored legs and beads in Λ_{loc} . Thus, the contraction $\text{con}_{\{h\}}$ in the above definition results in an element of $\mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$. It is easy to see that χ_h satisfies the following properties (compare with [GK2, Proposition 2.3])

Proposition 3.8. ([GK2]) (a) For Hermitian matrices A, B over Λ_{loc} , nonsingular over \mathbb{Z} , we have in $\widehat{\Lambda}/(\text{cyclic})$ that

$$\chi_h(AB) = \chi_h(A) + \chi_h(B) \quad \text{and} \quad \chi_h(A \oplus B) = \chi_h(A) + \chi_h(B).$$

(b) χ_h descends to a $\widehat{\Lambda}/(\text{cyclic})$ -valued invariant of the set $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$.

(c) For A as above, $\chi_h(A) = \chi_h(A \epsilon A^{-1})$ where $\epsilon : \Lambda \rightarrow \mathbb{Z}$.


From this, it follows that χ_h depends only on the image of M in $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ and the above definition makes sense in $\mathcal{A}^{\text{gp},0}$.

Remark 3.9. In the above definition 3.7 of the wrapping relation, we could have chosen equivalently that φ_1 and φ_2 denote the substitutions $\varphi_{t_j \rightarrow t_j e^h}$ and $\varphi_{t_j \rightarrow e^h t_j}$.

Remark 3.10. In the above definition, the ∂h colored legs are allowed to have beads in Λ_{loc} . Also, we allow the case of the empty matrix M , with the understanding that the χ_h term is absent.

Lemma 3.11. For $g = 1$, the wrapping relation ω^{gp} on $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})$ is trivial.

Proof. It follows from a swapping argument analogous to the isotopy of the following figure



□

We end this section with a comparison between \mathcal{A}^0 and $\mathcal{A}^{\text{gp},0}$. There is an obvious inclusion

$$\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) \longrightarrow \mathcal{A}^0(\star_X, \Lambda_{\text{loc}}).$$

The next proposition states that

Proposition 3.12. The inclusion $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) \longrightarrow \mathcal{A}^0(\star_X, \Lambda_{\text{loc}})$ maps the ω^{gp} -relations to ω -relations, and thus induces a map

$$\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \longrightarrow \mathcal{A}^0(\star_X, \Lambda_{\text{loc}}) / \langle \omega \rangle.$$

Proof. See Appendix C. □

4. AN INVARIANT OF LINKS IN THE COMPLEMENT OF AN UNLINK

The goal of this section is to define an invariant $Z^{\text{rat}} : \mathcal{D}_X(D_g) \rightarrow \mathcal{A}^{\text{gp}}(\otimes_X, \Lambda)$ of sliced crossed links whose components are in 1-1 correspondence with a set X .

Given a finite set X , an X -indexed sliced crossed link on D_g is a sliced crossed link on D_g equipped with bijection of the set of components of the presented link with X . All the definitions above have obvious adaptations for X -indexed objects, indicated by an appropriate X subscript when required.

Theorem 4.1. *There exists an invariant*

$$\check{Z}^{\text{rat}} : \mathcal{D}_X(D_g)/\langle r \rangle \longrightarrow \mathcal{A}^{\text{gp}}(\uparrow_X, \Lambda)$$

that maps the topological basing relations to group-like basing relations and the topological wrapping relations to wrapping relations. In particular, it induces a map

$$\sigma \circ \check{Z}^{\text{rat}} : \mathcal{N}_X(\mathcal{O}) \longrightarrow \mathcal{A}^{\text{gp}}(\otimes_X, \Lambda).$$

Proof. Our first task is to define \check{Z}^{rat} . In order to do so, we need to introduce some more notation. Given a word w in the letters \uparrow and \downarrow , let


- \uparrow_w denote the corresponding skeleton.
- $I_w \in \mathcal{A}(\uparrow_w, \Lambda)$ denote the identity element.
- $G_{i,w}$ denote the gluing word at the i th gluing site, that is, I_w with a label of t_i on each component, in the sense described by the following example: if $w = \uparrow\downarrow\downarrow$, then $G_{2,\uparrow\downarrow\downarrow} = \uparrow t_2 \downarrow t_2^{-1} \downarrow t_2^{-1}$.

Given a word w in the symbols $\uparrow, \downarrow, \bowtie$ and \bowtie , let w' denote w with the crosses forgotten. Let I_w^\times be $I_{w'}$ with every crossed strand broken. For example,

$$I_{\bowtie\downarrow\bowtie}^\times = \begin{array}{c} \uparrow \quad | \quad \downarrow \\ \uparrow \quad | \quad \downarrow \end{array}$$


Definition 4.2. Take an X -indexed element $D = \{T_1, \dots, T_k\} \in \mathcal{D}_X(D_g)$.

- (1) Select q-tangle lifts of each tangle in the sequence, $\{T'_1, \dots, T'_k\}$, such that
 - (a) the strands of each q-tangle which lie on boundary lines of the following form




are bracketed $(w_1)(w_2)$, where w_1 (resp. w_2) corresponds to the strands going to the left (resp. right) of the removed disc, and these bracketings match up,

- (b) boundary words at boundaries of the form



are given the canonical left bracketing.

- (2) Compose the (usual) Kontsevich invariants of these q-tangles in the following way:
 - (a) if two consecutive tangles are as follows,



with a word w in the symbols $\uparrow, \downarrow, \bowtie, \bowtie$ describing their matching boundary word, then they are to be composed

$$\dots \circ Z(T'_i) \circ I_w^\times \circ Z(T'_{i+1}) \circ \dots$$

(b) if, on the other hand, two consecutive tangles meet as follows,



with their matching boundary word at the j th gluing site factored as $(w_1)(w_2)$, then they are to be composed

$$\cdots \circ Z(T'_i) \circ (I_{w_1} \otimes G_{j,w_2}) \circ Z(T'_{i+1}) \circ \cdots$$

This completes the definition of $Z^{\text{rat}}(D)$. A normalized version $\check{Z}^{\text{rat}}(D)$ (useful for invariance under Kirby moves) is defined by connect-summing a copy of ν into each component of the crossed link associated to D . Observe that the result does not depend on the choice of a place to connect-sum to.

The proof of Theorem 4.1 follows from Lemmas 4.4 and 4.5 below. \square

The next lemma, observed by the second author in [Kr1] follows from a “sweeping” argument.

Lemma 4.3. *Let s denote an element of $\mathcal{A}(\uparrow_X, \mathbb{Q}) \subset \mathcal{A}(\uparrow_X, \Lambda)$, (i.e., some diagram with labels 1 at each edge), where the top boundary word of \uparrow_X is w_1 , and the bottom boundary word of \uparrow_X is w_2 . Then*

$$G_{w_1} \circ s = s \circ G_{w_2}.$$

Lemma 4.4. *The map \check{Z}^{rat} is well defined on $\mathcal{D}_X(D_g)$ and moreover descends to an invariant of $\mathcal{D}_X(D_g)/\langle r \rangle$.*

Proof. For the first part we only need to show that the function does not depend on the choice of bracketings. This follows because of the following, where ψ_1, ψ_2 are some expressions built from the associator.

$$\begin{aligned} & \cdots \circ Z(T'_i) \circ (\psi_1 \otimes \psi_2) \circ (I_{w_1} \otimes G_{j,w_2}) \circ (\psi_1^{-1} \otimes \psi_2^{-1}) \circ Z(T'_{i+1}) \circ \cdots, \\ = & \cdots \circ Z(T'_i) \circ (I_{w_1} \otimes G_{j,w_2}) \circ (\psi_1 \otimes \psi_2) \circ (\psi_1^{-1} \otimes \psi_2^{-1}) \circ Z(T'_{i+1}) \circ \cdots, \\ = & \cdots \circ Z(T'_i) \circ (I_{w_1} \otimes G_{j,w_2}) \circ Z(T'_{i+1}) \circ \cdots \end{aligned}$$

where the first step follows from Lemma 4.3. The invariance of \check{Z}^{rat} under the \sim^r equivalence relation follows the same way. \square

Lemma 4.5. *The map*

$$\sigma \circ \check{Z}^{\text{rat}} : \mathcal{D}_X(D_g)/\langle r, \beta \rangle \rightarrow \mathcal{A}^{\text{gp}}(\star_X, \Lambda)/\langle \beta^{\text{gp}} \rangle$$

is well-defined.

Proof. We need to show invariance of $\sigma \circ \check{Z}^{\text{rat}}$ (or equivalently, $\sigma \circ Z^{\text{rat}}$) under the β_1 and β_2 moves. Enlarge $\mathcal{D}_X(D_g)$ to $\mathcal{D}_X(D_g)'$, the set of X -indexed sliced crossed links on D_g whose presented links in $D_g \times I$ have at least one cross on every component, thus $\mathcal{D}(D_g) \subset \mathcal{D}(D_g)'$. The function Z^{rat} immediately extends to this generality. For two diagrams $D_1, D_2 \in \mathcal{D}_X(D_g)$ such that $D_1 \stackrel{\beta^{\text{gp}}}{\sim} D_2$ let D_{12} denote the diagram in $\mathcal{D}(D_g)'$ such that D_1 is obtained by forgetting some cross, and D_2 is obtained by forgetting the other cross on that component. Notice that in this case $Z^{\text{rat}}(D_{12}) \in \mathcal{A}^{\text{gp}}(\uparrow_{\{x\}} \uparrow_{\{x'\}} \uparrow_{X-\{x\}}, \Lambda)$. The key point is the fact that the product of labels around each component of the skeleton equals to 1; this is true since the locations of the two crosses cut the component into two arcs, both of which have trivial intersection word with the gluing sites. It follows by construction that

$$\begin{aligned} Z^{\text{rat}}(D_1) &= \vec{m}_x^{x'}(Z^{\text{rat}}(D_{12})), \\ Z^{\text{rat}}(D_2) &= \vec{m}_x^{x'}(Z^{\text{rat}}(D_{12})), \\ Z^{\text{rat}}(D_{12}) &\in \mathcal{A}^{\text{gp}}(\uparrow_{\{x\}} \uparrow_{\{x'\}} \uparrow_{X-\{x\}}, \Lambda_{\text{loc}}). \end{aligned}$$

Thus, $\sigma(Z^{\text{rat}}(D_1)) \stackrel{\beta_1^{\text{gp}}}{\sim} \sigma(Z^{\text{rat}}(D_2))$.

If $D_1 \stackrel{\beta_2}{\sim} D_2$, then for some appropriate expression ψ built from the associator, we have that:

$$\begin{aligned}
\sigma(Z^{\text{rat}}(\{\dots, \text{diagram}_1, \text{diagram}_2, \dots\})) &= \sigma(\dots \circ I_w^\times \circ \psi \circ (I_{w_1} \otimes G_{i,w_2}) \circ \psi^{-1} \circ \dots) \\
&\stackrel{\beta_1^{\text{gp}}}{\sim} \sigma(\dots \circ \psi \circ I_w^\times \circ (I_{w_1} \otimes G_{i,w_2}) \circ \psi^{-1} \circ \dots) \\
&\stackrel{\beta_2^{\text{gp}}}{\sim} \sigma(\dots \circ \psi \circ (I_{w_1} \otimes G_{i,w_2}) \circ I_w^\times \circ \psi^{-1} \circ \dots) \\
&\stackrel{\beta_1^{\text{gp}}}{\sim} \sigma(\dots \circ \psi \circ (I_{w_1} \otimes G_{i,w_2}) \circ \psi^{-1} \circ I_w^\times \circ \dots) \\
&= \sigma(Z^{\text{rat}}(\{\dots, \text{diagram}_2, \text{diagram}_1, \dots\})).
\end{aligned}$$

□

Lemma 4.6. *The following map is well-defined:*

$$\sigma \circ \check{Z}^{\text{rat}} : \mathcal{D}(D_g) / \langle \omega \rangle \rightarrow \mathcal{A}^{\text{gp}}(\star_X, \Lambda) / \langle \omega^{\text{gp}} \rangle$$

Proof. Consider two sliced crossed links $D_1, D_2 \in \mathcal{D}(D_g)$ related by a topological wrapping move at the j th gluing site as in Section 2.2. In other words, D_1 (shown on the left) and D_2 (shown on the right) are given by:

$$\{\dots, \text{diagram}_1, \text{diagram}_2, \dots\} \longleftrightarrow \{\dots, \text{diagram}_2, \text{diagram}_1, \dots\}$$

Let

$$z_1 = Z^{\text{rat}} \left(\text{diagram}_1 \right), \quad z'_1 = \uparrow_{x'} e^h \cup (\sigma_x(z_1))_{x \rightarrow \partial h}, \quad z_2 = Z^{\text{rat}} \left(\text{diagram}_2 \right)$$

Now, we can compute, for some element ψ built from an associator:

$$\begin{aligned}
(\sigma \circ Z^{\text{rat}})(D_2) &= \sigma \circ (\text{con}_{\{h\}}(\dots \circ \psi \circ \Delta_{x'}^{w_2}(z'_1) \circ (I_{w_1} \otimes G_{j,w_2}) \circ \Delta_x^{w_2}(z_2) \circ \psi^{-1} \circ \dots)) \\
&\stackrel{\omega^{\text{gp}}}{\sim} \sigma \circ (\text{con}_{\{h\}}(\dots \circ \psi \circ (I_{w_1} \otimes G_{j,w_2}) \circ \Delta_{x'}^{w_2}(z'_1) \circ \Delta_x^{w_2}(z_2) \circ \psi^{-1} \circ \dots)), \\
&= (\sigma \circ Z^{\text{rat}})(D_1).
\end{aligned}$$

In the above, $\Delta_{x'}^{w_2}$ means the comultiplication of the diagram x with pattern given by the gluing word w_2 . □

Theorem 4.1 now follows, using the identification of $\mathcal{N}_X(\mathcal{O})$ with the set $\mathcal{D}_X(D_g) / \langle r, \beta, \omega \rangle$ given by Proposition 2.4.

Remark 4.7. An application of the Z^{rat} invariant is a formula for the LMO invariant of cyclic branched covers of knots, [GK1]. This application requires a *renormalized version* $Z^{\text{rat}}[\alpha]$ where $\alpha = (\alpha_1, \dots, \alpha_g)$ and $\alpha_i \in \mathcal{A}^{\text{gp}}(\star_k)$. $Z^{\text{rat}}[\alpha]$ is defined using Definition 4.2 and replacing the formula in 2(b) by

$$\dots \circ Z(T'_i) \circ (I_{w_1} \otimes G_{j,w_2} \Delta(\alpha_j)) \circ Z(T'_{i+1}) \circ \dots$$

where $\Delta(\alpha_j)$ places a copy of α_j in each of the skeleton segments of the j th gluing site. The renormalized invariant $Z^{\text{rat}}[\alpha]$ still satisfies the properties of Theorem 4.1.

5. A RATIONAL VERSION OF THE AARHUS INTEGRAL

The main purpose of this section is to introduce a notion \int^{rat} of Rational Formal Gaussian Integration \int^{rat} which respects the Kirby moves the basing and the winding relations. The reader who is not familiar with the combinatorics of gluing legs of diagrams together, is urged to look at the appendix.

5.1. **The definition of \int^{rat} .** Recall that $\text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ denotes the set of Hermitian matrices over Λ_{loc} , invertible over \mathbb{Z} . Let X denote a finite set and let $X' = \{x_i\}$ denote an arbitrary subset of X . A diagram $D \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ is called X' -*substantial* if there are no *strut* components (i.e., components comprising of a single edge) both of whose univalent vertices are labeled from the subset X' . An element of $\mathcal{A}(\star_X, \Lambda_{\text{loc}})$ is X' -substantial if it is a series of X' -substantial diagrams.

An element $s \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ is called *integrable with respect to X'* if there exists an Hermitian matrix $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ such that

$$(2) \quad s = \exp \left(\frac{1}{2} \sum_{i,j} \begin{array}{c} x_j \\ \uparrow \\ \bullet \\ \downarrow \\ x_i \end{array} M_{ij} \right) \sqcup R,$$

with R an X' -substantial element. Notice that M , the *covariance matrix* $\text{cov}(s)$ of s , and R , the X' -substantial part of s , are uniquely determined by s .

Let $\text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \subset \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ denote the subset of elements that are integrable with respect to X' . Observe that although $\text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ has non-empty intersection with $\mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$, it is not a subset of it. Further, observe that if two elements of $\mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$ are related by a basing move and one of them lies in $\text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, so does the other. Thus the following definition makes sense:

$$\text{Int}_{X'} \mathcal{A}^{\text{gp}}(\otimes_X, \Lambda_{\text{loc}}) = (\mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}}) \cap \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})) / \langle \beta_1^{\text{gp}}, \beta_2^{\text{gp}} \rangle.$$

For the next definition, recall that every matrix $W \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ is invertible over Λ_{loc} .

Definition 5.1. Define a map (*Rational Formal Gaussian Integration*)

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_{X-X'}, \Lambda_{\text{loc}})$$

as follows. If $s \in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ has associated decomposition given in Equation (2), then

$$\int^{\text{rat}} dX'(s) = \left\langle \exp_{\sqcup} \left(-\frac{1}{2} \sum_{i,j} \begin{array}{c} \partial x_j \\ \uparrow \\ \bullet \\ \downarrow \\ \partial x_i \end{array} M_{ij}^{-1} \right), R \right\rangle_{X'}.$$

The gluing of the legs of diagrams occurs according to Remark 3.5.

Remark 5.2. The definition of \int^{rat} for elements s whose covariance matrix M is defined over Λ , and invertible over \mathbb{Z} , requires beads labeled by the inverse matrix M^{-1} . This necessitates the need to replace Λ by an appropriate ring Λ_{loc} that makes all such matrices M invertible. It follows by definition that the universal choice of Λ_{loc} is the *Cohn localization* of Λ .

A complementary reason to consider the Cohn localization, from the (dual) point of view of filtrations comes from the the study of the *null filtration*, given in [Ga].

5.2. \int^{rat} **respects the basing relations.** The next theorem shows that \int^{rat} is compatible with the basing relations.

Theorem 5.3. \int^{rat} *descends to a map:*

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}}) / \langle \beta_X^{\text{gp}} \rangle \rightarrow \mathcal{A}^{\text{gp}}(\star_{X-X'}, \Lambda_{\text{loc}}) / \langle \beta_{X-X'}^{\text{gp}} \rangle$$

Proof. Using Lemma 3.6, we need to show that given $s_1, s_2 \in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \cap \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$ such that $s_1 \stackrel{\beta_1^{\text{gp}}}{\sim} s_2$ or $s_1 \stackrel{\beta_2^{\text{gp}}}{\sim} s_2$, based on some element $x_k \in X'$, then

$$\int^{\text{rat}} dX'(s_1) = \int^{\text{rat}} dX'(s_2).$$

The case of β_2^{gp} . Consider the decomposition of s_1

$$s_1 = \exp \left(\frac{1}{2} \sum_{i,j} \begin{array}{c} x_j \\ \uparrow \\ \bullet \\ \downarrow \\ x_i \end{array} M_{ij} \right) \sqcup R.$$

Suppose that an element $f \in F_g$ is pushed onto the legs labeled by $x_k \in X'$, to form s_2 . Now, let $D_{f,k}$ be $\text{diag}(1, 1, \dots, f, \dots, 1)$, where f is in the k th entry, and let R' denote R with an f pushed onto every leg marked x_k . Then,

$$\begin{aligned} \int^{\text{rat}} dX'(s_2) &= \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \uparrow (D_{f,k} M D_{f^{-1},k})_{ij} \right) \sqcup R' \right) \\ &= \left\langle \exp \left(-\frac{1}{2} \sum_{x_i}^{x_j} \uparrow (D_{f,k} M^{-1} D_{f^{-1},k})_{ij} \right), R' \right\rangle \\ &= \left\langle \exp \left(-\frac{1}{2} \sum_{x_i}^{x_j} \uparrow (M^{-1})_{ij} \right), R \right\rangle \\ &= \int^{\text{rat}} dX'(s_1) \end{aligned}$$

because all the beads labeled f and f^{-1} match up and cancel.

The case of $\beta_1^{\text{gp}'}$. Consider $s \in \mathcal{A}^{\text{gp}}(\star_{X \cup \partial h}, \Lambda_{\text{loc}})$ such that

$$s_1 = \text{con}_{\{h\}}(s) \quad \text{and} \quad s_2 = \text{con}_{\{h\}}(s|_{x_k \rightarrow x_k e^h}).$$

Notice that if s_1, s_2 are X' -integrable, so is s . Furthermore, $\text{con}_{\{h\}}$ commutes with \int^{rat} . The result follows from Lemma 5.4 below. \square

Lemma 5.4. *Let $s \in \text{Int}_{X'} \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$, and let $x \in X' \subset X$.*

$$\int^{\text{rat}} dX'(s|_{x \rightarrow x e^h}) = \int^{\text{rat}} dX'(s) \in \mathcal{A}^{\text{gp}}(\star_{X \cup \{h\}}, \Lambda_{\text{loc}}).$$

Proof. The idea of this lemma is the same as “in the case $\beta_2^{\text{gp}''}$ ”, but instead of pushing an element of the free group onto legs labeled by some element, we push an exponential of hair.

The substitution $x \rightarrow x e^h$ defines a map

$$\mathcal{A}(\star_X, \Lambda_{\text{loc}}) \longrightarrow \mathcal{A}(\star_{X \cup \{h\}}, \Lambda_{\text{loc}})$$

with the property that $s - s|_{x \rightarrow x e^h}$ is X' -substantial for all $s \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$.

Assume that the canonical decomposition of s is given by Equation (2). Let $D_{f,k}$ be $\text{diag}(1, 1, \dots, e^h, \dots, 1)$, where e^h is in the k th entry, and let R' denote R with an e^h pushed onto every leg marked x_k . Then, Theorem E.1 from Appendix E implies that

$$\begin{aligned} \int^{\text{rat}} dX'(s|_{x \rightarrow x e^h}) &= \exp \left(-\frac{1}{2} \chi_h (M D_{e^h, k} M^{-1} D_{e^{-h}, k}) \right) \sqcup \left\langle \exp \left(-\frac{1}{2} \sum_{x_i}^{x_j} \uparrow M_{ij}^{-1} \right), R \right\rangle \\ &= \left\langle \exp \left(-\frac{1}{2} \sum_{x_i}^{x_j} \uparrow M_{ij}^{-1} \right), R \right\rangle \end{aligned}$$

where the last line follows from Proposition 3.8. \square

Remark 5.5. An alternative proof of Theorem 5.3 can be obtained by adapting the proof of [A, Part II, Proposition 5.6] to the rational integration \int^{rat} .

5.3. Dealing with the Kirby moves. Recall from Section 2.2 the κ handleslide move on the set $\mathcal{D}(D_g)$. In a similar fashion, we define an X' -handleslide move on $\mathcal{A}(\star_X, \Lambda_{\text{loc}})$, as follows: For two elements $s_1, s_2 \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, say that $s_1 \stackrel{\kappa_{X'}}{\sim} s_2$, if for some $x_i \in X$, and $x_j \in X'$, $i \neq j$,

$$s_2 = (\sigma_{\{x_i\}} \circ \bar{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) (s_1(x_1, \dots, x_{j-1}, x_j + y, x_{j+1}, \dots)).$$

Notice that if $s_1, s_2 \in \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$, $s_1 \stackrel{\kappa_{X'}}{\sim} s_2$ and one of s_i lies in $\text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, so does the other. This is because the covariance matrix of s_2 is obtained from the covariance matrix of s_1 by a unimodular congruence.

Theorem 5.6. *The map \tilde{Z}^{rat} descends to a map:*

$$\tilde{Z}^{\text{rat}} : \mathcal{D}_X(D_g)/\langle r, \beta, \kappa_{X'} \rangle \rightarrow \text{Int}_{X'} \mathcal{A}^{\text{gp}}(\star_X, \Lambda)/\langle \beta^{\text{gp}}, \kappa_{X'} \rangle.$$

Proof. The argument of [LMMO, Proposition 1], adapts to this setting without difficulty. There are two possible subtleties that must be observed.

1. The proof in [LMMO] is based on taking a parallel of the parts of a component (here, the component x_j) away from the site of the cross. In our setting, the paralleling operation of the involved components of tangles behaves as usual (sums over lifts of legs). At a gluing word, one simply labels both of the two components arising from the parallel operation with a bead labeled by the appropriate letter. Then, when $\sigma_{x_i, y}$ is applied, these two identical beads are pushed onto the legs sitting on those components side-by-side. This may alternatively be done by first pushing the original bead up onto the legs, *and then* taking the parallel, as is required in this situation.
2. We have defined the bracketing at boundary lines with crosses to be the standard left-bracketing, while the move in [LMMO] is given for a presentation where the two involved strands are bracketed together. This is treated by preceding and following the $\kappa_{X'}$ with a β_1 move, to account for the change in bracketing, exactly as in the proof of the β_2 -case of Lemma 4.5. \square

Theorem 5.7. *The map $\int^{\text{rat}} dX'$ descends to a map:*

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})/\langle \kappa_{X'} \rangle \rightarrow \mathcal{A}^{\text{gp}}(\star_{X-X'}, \Lambda_{\text{loc}}).$$

This is proved by the following lemma.

Lemma 5.8. *Let $s_1 \in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ and let*

$$s_2 = (\sigma_{\{x_i\}} \circ \vec{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) (s_1(x_1, \dots, x_{j-1}, x_j + y, x_{j+1}, \dots)).$$

If $x_j \in X'$ then

$$\int^{\text{rat}} dX'(s_1) = \int^{\text{rat}} dX'(s_2).$$

Proof. We prove this in two steps, which usually go by the names of the “lucky” and “unlucky” cases.

The lucky case: $s_1, s_2 \in \text{Int}_{\{x_j\}} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$. In this case, we can use the Iterated Integration Lemma A.2.

$$\begin{aligned} \int^{\text{rat}} dX'(s_2) &= \int^{\text{rat}} d(X' - \{x_j\}) \left(\int^{\text{rat}} d\{x_j\}(s_2) \right), \\ &= \int^{\text{rat}} d(X' - \{x_j\}) \left(\int^{\text{rat}} d\{x_j\} (\sigma_{\{x_i\}} \circ \vec{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) (s_1(x_1, \dots, x_{j-1}, x_j + y, x_{j+1}, \dots)) \right), \\ &= \int^{\text{rat}} d(X' - \{x_j\}) \\ &\quad \left((\sigma_{\{x_i\}} \circ \vec{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) \left(\int^{\text{rat}} d\{x_j\} (s_1(x_1, \dots, x_{j-1}, x_j + y, x_{j+1}, \dots)) \right) \right), \\ &= \int^{\text{rat}} d(X' - \{x_j\}) \left((\sigma_{\{x_i\}} \circ \vec{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) \right. \\ &\quad \left. \left(\int^{\text{rat}} d\{x_j\} \left(\exp \left(\begin{array}{c} y \\ \uparrow \\ \bullet \\ \partial x_j \end{array} \right) 1 \right) b_{\{x_j\}} s_1(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots) \right) \right) \end{aligned}$$

Now, using the Integration by Parts Lemma A.4, it follows that

$$\begin{aligned} \int^{\text{rat}} d\{x_j\} \left(\exp \left(\begin{array}{c} y \\ \uparrow \\ \bullet \\ \partial x_j \end{array} \right) 1 \right) b_{\{x_j\}} s_1(x_1, \dots) &= \int^{\text{rat}} d\{x_j\} \left(\text{div}_{\{x_j\}} \left(\exp \left(\begin{array}{c} y \\ \uparrow \\ \bullet \\ \partial x_j \end{array} \right) 1 \right) b_{\{x_j\}} s_1(x_1, \dots) \right) \\ &= \int^{\text{rat}} d\{x_j\} (s_1(x_1, \dots)), \end{aligned}$$

because $\text{div}_{\{x_j\}} \left(\exp \left(\begin{array}{c} y \\ \uparrow \\ \epsilon \uparrow 1 \\ \partial x_j \end{array} \right) \right) = 1$.

The unlucky case: $s_1, s_2 \notin \text{Int}_{\{x_j\}} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$. This pathology is treated by deformation. Namely, we “deform” s_1 by multiplying it by $\exp \left(\begin{array}{c} x_j \\ \uparrow \\ \epsilon \uparrow 1 \\ x_j \end{array} \right)$, for ϵ a “small real,” commuting with Λ_{loc} . Formally, we let ϵ be a variable and consider an integration theory based on $\mathbb{Q}(\epsilon)[F_g]$ and its Cohn localization $\Lambda_{\text{loc}}^\epsilon$. We identify $\mathbb{Q}^0(\epsilon) \subset \mathbb{Q}(\epsilon)$, the subring of rational functions in ϵ non-singular at zero, and the corresponding Cohn localization $\Lambda_{\text{loc}}^{\epsilon,0}$.

So, letting

$$\begin{aligned} s_1^\epsilon &= \exp \left(\begin{array}{c} x_j \\ \uparrow \\ \epsilon \uparrow 1 \\ x_j \end{array} \right) \sqcup s_1 \in \mathcal{A}(\star_X, \Lambda_{\text{loc}}^\epsilon), \\ s_2^\epsilon &= (\sigma_{\{x_i\}} \circ \vec{m}_{x_i}^{y, x_i} \circ \chi_{\{y, x_i\}}) (s_1^\epsilon(x_1, \dots, x_{j-1}, x_j + y, x_{j+1}, \dots)), \end{aligned}$$

it follows from the argument of the “lucky case” that

$$\int^{\text{rat}} dX'(s_2^\epsilon) = \int^{\text{rat}} dX'(s_1^\epsilon).$$

Now, there is a ring homomorphism $\phi : \Lambda_{\text{loc}}^{\epsilon,0} \rightarrow \Lambda_{\text{loc}}$, obtained by “setting ϵ to zero”. Observing that, although of course $\int^{\text{rat}} d\{x_j\}(s_i^\epsilon) \notin \mathcal{A}(\star_X, \Lambda_{\text{loc}}^{\epsilon,0})$ (though it is now well-defined),

$$\int^{\text{rat}} dX'(s_i^\epsilon) \in \mathcal{A}(\star_X, \Lambda_{\text{loc}}^{\epsilon,0}), \quad \text{and} \quad \phi \left(\int^{\text{rat}} dX'(s_i^\epsilon) \right) = \int^{\text{rat}} dX'(s_i),$$

the proof follows. □

Remark 5.9. These last lines are best understood from the realization of Λ_{loc} and $\Lambda_{\text{loc}}^\epsilon$ as spaces of formal power series. The elements $\int^{\text{rat}} dX'(s_i^\epsilon)$ lie in $\mathcal{A}(\star_X, \Lambda_{\text{loc}}^{\epsilon,0})$ because the determinant of the augmentation of the covariance matrix of s_i is invertible in $\mathbb{Q}^0(\epsilon)$.

5.4. Dealing with the wrapping relations. In this section we show that \int^{rat} -integration respects the group-like wrapping relations ω^{gp} that were introduced in Definition 3.7. Since the wrapping relations involve pairs of matrices and graphs, we begin by extending the definition of \int^{rat} to pairs (M, s) by setting

$$\int^{\text{rat}} dX'(M, s) = (M \oplus \text{cov}(s), \int^{\text{rat}} dX'(s)),$$

where $\text{cov}(s)$ is the covariance matrix of s . The motivation for this comes from the next theorem which states that \int^{rat} preserves the wrapping relations.

Theorem 5.10. *The map $\int^{\text{rat}} dX'$ descends to a map*

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \rightarrow \mathcal{A}^{\text{gp},0}(\star_{X-X'}, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle$$

Proof. Consider two pairs $(M, s_1) \stackrel{\omega^{\text{gp}}}{\sim} (M, s_2)$ related by a wrapping move.

Using Theorem E.1 in Appendix E, it follows that if W is the common covariance matrix of s_1, s_2 and s_{12} , then

$$\begin{aligned}
\int^{\text{rat}} dX'(M, s_m) &= \left(M \oplus W, \text{con}_{\{h\}} \left(\int^{\text{rat}} dX'(\varphi_m(s_{12})) \sqcup \left(\exp \left(-\frac{1}{2} \chi_h(M^{-1} \varphi_m M) \right) \right) \right) \right) \\
&= \left(M \oplus W, \text{con}_{\{h\}} \left(\varphi_m \left(\int^{\text{rat}} dX'(s_{12}) \right) \sqcup \left(\exp \left(-\frac{1}{2} \chi_h(W^{-1} \varphi_m W) \right) \right) \right) \right) \\
&\quad \sqcup \left(\exp \left(-\frac{1}{2} \chi_h(M^{-1} \varphi_m M) \right) \right) \\
&= \left(M \oplus W, \text{con}_{\{h\}} \left(\varphi_m \left(\int^{\text{rat}} dX'(s_{12}) \right) \sqcup \left(\exp \left(-\frac{1}{2} \chi_h((W \oplus M)^{-1} \varphi_m (W \oplus M)) \right) \right) \right) \right) \\
&\stackrel{\omega^{\text{gp}}}{\sim} \left(M, \int^{\text{rat}} dX'(M, s_{12}) \right).
\end{aligned}$$

□

Remark 5.11. The group-like wrapping relation is the minimal relation that we need to impose on diagrams so that $\int^{\text{rat}} \circ \sigma \circ Z^{\text{rat}}$ is invariant under the topological wrapping relation, as follows from the proof of Theorem 5.10.

5.5. The definition of the rational invariant Z^{rat} of F -links. Theorems 4.1, 5.3, 5.7, and 5.10 imply that there is a map

$$Z^{\text{rat}, \text{gp}} : F - \text{links} \longrightarrow \mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle$$

defined as follows: for a F -link (M, L) of g components, choose a sliced crossed link C in $\mathcal{D}(D_g)$ whose components are in 1-1 correspondence with a set X , and define

$$Z^{\text{rat}, \text{gp}}(M, L) = \left(\int^{\text{rat}} dX(\sigma \circ \check{Z}^{\text{rat}}(C)) \right) \cdot c_+^{-\sigma_+(B)} c_-^{-\sigma_-(B)} \in \mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle$$

where $c_{\pm} = \int dU \check{Z}(S^3, U_{\pm})$ are some universal constants of the unit-framed unknot U_{\pm} and where B is the linking matrix of C and $\sigma_{\pm}(B)$ is the number of positive (resp. negative) eigenvalues of the symmetric matrix B . The slight renormalization given by multiplication by c_{\pm} is needed to deal with the stabilization Kirby move on C , and is justified as follows: recall that $\mathcal{A}^{\text{gp}}(\phi)$ is a group, which acts on the set $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}})$ by $a \cdot (M, s) = (M, a \sqcup s)$. Since the beads of the graphs in $\mathcal{A}^{\text{gp}}(\phi)$ are rational numbers independent of the t_i , it follows easily that the above action of $\mathcal{A}^{\text{gp}}(\phi)$ on $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}})$ induces an action of $\mathcal{A}^{\text{gp}}(\phi)$ on the quotient set $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega \rangle$. This is exactly what appears in the above definition of Z^{rat} .

Composing further with the map $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}}, \text{String} \rangle \rightarrow \mathcal{A}^0 / \langle \omega, \text{String} \rangle$ of Proposition 3.12, defines the rational invariant Z^{rat} of F -links that appears in Theorem 1.2.

Proof. (of Remark 1.5) Recall that the matrix part W of Z^{rat} is an element of $\mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ and determines the S -equivalence class of the F -link, as follows from [GK2]. Recall also that the wrapping relations ω^{gp} and ω depend only on W . Thus, it follows that S -equivalent F -links have equal matrix-part, and that the matrix-free part of Z^{rat} makes sense as an element of the graded vector space $\mathcal{A}(\Lambda_{\text{loc}}) / \langle \omega \rangle$.

[GK2, Theorem 3] implies that (M, L) is a good boundary link, (i.e., $H_1(X^{\omega}, \mathbb{Z}) = 0$) iff its associated matrix W is invertible over Λ . In that case, the construction of the Z^{rat} invariant implies that $Z^{\text{rat}}(M, L) \in \mathcal{A}^0(\Lambda) / \langle \omega, \text{String} \rangle$. □

6. THE String ACTION AND INVARIANTS OF BOUNDARY LINKS

The goal of this section is to use the already constructed invariant of F -links in order to define an invariant of boundary links. The key observation is that the set of boundary links can be defined with the set of orbits of an action

$$\text{String} \times F - \text{links} \rightarrow F - \text{links}$$

for a suitable group String . Starting from this observation, we refine the results of Sections 2-5 to take into account the action of the group String .

6.1. A surgery description of boundary links. We begin by giving a surgery description of boundary links. Recall that the set of ∂ -links can be identified as a quotient of the set of F -links modulo an action of a group of automorphisms String_g of the free group, [K, CS2]. The group String_g in question is the subgroup of the automorphism group of F_g that consists of all automorphisms α such that $\alpha(t_i) = w_i^{-1}t_iw_i$ for some $w_i \in F_g$ and for all generators t_1, \dots, t_g of F_g . In particular, String_g contains (and for $g = 1, 2$ coincides with) the group of inner automorphisms of F_g . It is easy to see that String_g is generated by the special automorphisms α_{ij} (for $1 \leq i \neq j \leq g$) acting on the generators of F_g by $\alpha_{ij}(t_k) = t_k$ (resp. $t_j^{-1}t_it_j$) for $k \neq i$ (resp. $k = i$), see [K, Lemma 2.4]. The group String_g acts on the set of F -links by changing the basing of the F -link, in other words by sending (M, L_b, ϕ) to $\alpha \circ \phi$ for $\alpha \in \text{String}_g$. An alternative and more geometric way to describe this action is as follows: identify the set of F -links with the set of Seifert surfaces modulo isotopy and *tube equivalence*, see [GL2]. There is an action of String_g on the set of Seifert surfaces, described on generators α_{ij} by *cocooning* the i th surface to the j th surface using suitable arcs to a base point, well-defined modulo isotopy and tube equivalence.

Since the set $\mathcal{N}(\mathcal{O})/\langle \kappa \rangle$ is in 1-1 correspondence with the set of F -links and since String acts on the latter set, it follows that it acts on the former set, too. We now describe this action of String_g (or rather, of its generating set $\{\alpha_{ij}\}$) on the set $\mathcal{N}(\mathcal{O})/\langle \kappa \rangle$.

Consider a link $L \in \mathcal{N}(\mathcal{O})$. The action of α_{ij} on such a link is the following: “the i th unlink component is isotoped through the j th unlink component, and then put back where it started”.

Let us examine what happens when we try to implement such a principle. To do this, we take a parallel copy of the i th unlink component and drag it so that, at the end, the initial link (resp. the link that results from the move) is recovered by forgetting the parallel copy (resp. forgetting the original copy).




Figure 4. Dragging the i th unlink component through the j th unlink component, and then put back where it started.

It is easy to see that the α_{ij} move does not depend on the choice of arc, modulo Kirby moves, and that the above discussion together with Theorem 2.2 imply that

Theorem 6.1. *There exists an action*

$$\text{String} \times (\mathcal{N}(\mathcal{O})/\langle \kappa \rangle) \longrightarrow \mathcal{N}(\mathcal{O})/\langle \kappa \rangle$$

which induces a 1-1 onto correspondence


$$\mathcal{N}(\mathcal{O})/\langle \kappa, \text{String} \rangle \longleftrightarrow \partial - \text{links}.$$

Remark 6.2. It is instructive to compare the above theorem with a result of Roberts [Rb], which states that two 3-manifolds N_L and $N_{L'}$ are diffeomorphic relative boundary iff L and L' are related by a sequence of handle-slides (κ_1), stabilization (κ_2), and insertion/deletion of an arbitrary knot in N together with a 0-framed meridian of it (κ_3). In case the knot participating in the κ_3 move is nullhomotopic, the move is equivalent to a composition of κ_1 and κ_2 moves, as was observed by Kirby and Fenn-Rourke. In general, the set $\mathcal{N}(\mathcal{O})$ is not closed under the moves κ_3 , however we can consider the set $\widehat{\mathcal{N}}(\mathcal{O})$ that consists of all links in $S^3 - \mathcal{O}$ that are obtained from nullhomotopic links by $\kappa_{1,2,3}$ moves. Roberts' theorem implies that there is a 1-1 onto correspondence


$$\widehat{\mathcal{N}}(\mathcal{O})/\langle \kappa_1, \kappa_2, \kappa_3 \rangle \longleftrightarrow \partial - \text{links}.$$

It is an amusing exercise to show that the links of Figure 4 are related by a sequence of $\kappa_{1,2,3}$ moves. \square

In order to give a tangle description of the set of ∂ -links, we need to introduce a geometric action of the set String_g on the set $\mathcal{D}(D_g)$. Recalling the action of String_g on the set $\mathcal{N}(\mathcal{O})$, we see that this action can be specified by a framed arc, and that as this proceeds, then all the strands going between the two copies are parallel to each other (that is, dragging the copy away from the original “lines-up” all the strands in between); see also Figure 4. This will be the idea of our action of String_g on $\mathcal{D}(D_g)$. To formalize this, we (briefly) enlarge our notion of sliced crossed links, to allow for such an arc. That is, we will consider the set $\mathcal{D}^{\text{arc}}(D_g)$ of *arc-decorated sliced crossed links*, represented by sequences of tangles such that, in addition to all closed components, there is an arc which starts and finishes at some meridional disc:



For an element of this set, we will say that the result of a *drag* along the arc is the sliced crossed link that arises by making the following replacements, accompanied by taking a parallel of the arc a according to the appropriate word (in the example below, according to $w_b\bar{w}_b$).



Given two sliced crossed links $L_1, L_2 \in \mathcal{D}(D_g)$, we say that L_2 is obtained from L_1 by an α_{ij} move if there exists an arc-decorated sliced crossed link $L_{12} \in \mathcal{D}^{\text{arc}}(D_{g+1})$ with the following properties:

- (1) there are no intersections of link components with the $(i + 1)$ st meridional disc,
- (2) gluing back in the $(i + 1)$ st removed disc (and forgetting the arc) gives a diagram which presents L_1 ,
- (3) the arc travels from the i th meridional disc, intersects only the j th meridional disc, and only once, and then ends at the i th meridional disc.
- (4) Dragging L_{12} along the arc and then gluing back the i th removed unlink component recovers L_2 .

Remark 6.3. The above definition implements the procedure described in Figure 4. Actually, this α_{ij} move does not depend on the choice of arc (this is an instructive exercise). In addition, this move translated to the set of F -links coincides with the action of String_g by cocooning.

Proposition 2.4 and Theorem 6.1 imply that

Proposition 6.4. *There is an action*

$$\text{String} \times \mathcal{D}(D_g) / \langle r, \beta, \omega, \kappa \rangle \longleftrightarrow \mathcal{D}(D_g) / \langle r, \beta, \omega, \kappa \rangle$$

which induces a 1-1 correspondence

$$\mathcal{D}(D_g) / \langle r, \beta, \omega, \kappa, \text{String} \rangle \longleftrightarrow \partial - \text{links.}$$

6.2. The action of String on diagrams. We now define an algebraic action of String_g on diagrams. Recall the group String_g that acts on the free group F_g (see Section 6.1), thus also on Λ . The action of String_g on Λ extends to the localization Λ_{loc} , since for $\alpha \in \text{String}_g$, the induced ring homomorphism $\Lambda \rightarrow \Lambda \subset \Lambda_{\text{loc}}$ is Σ -inverting, and thus extends to a ring homomorphism $\Lambda_{\text{loc}} \rightarrow \Lambda_{\text{loc}}$. This induces an action

$$\text{String}_g \times \mathcal{A}^0(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^0(\star_X, \Lambda_{\text{loc}})$$

given by acting on the beads of diagrams and the entries of matrices by special automorphisms of the free group. Following Remark 3.1, we denote by $\mathcal{A}^0(\star_X, \Lambda_{\text{loc}})/\langle \text{String} \rangle$ the quotient space.

Similarly, there is an action

$$\text{String}_g \times \mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}),$$

which results in a quotient space $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})/\langle \text{String} \rangle$, too.

The next theorem shows that the action of String_g descends to one on the quotient set $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})/\langle \beta^{\text{gp}}, \omega^{\text{gp}} \rangle$.

Theorem 6.5. *The String_g action on $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})$ descends to an action on $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})/\langle \beta^{\text{gp}}, \omega^{\text{gp}} \rangle$.*

Proof. It is an exercise in the definitions to see that if $(M, s_1), (M, s_2) \in \mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})$ and $(M, s_1) \stackrel{\beta^{\text{gp}}}{\sim} (M, s_2)$, then $\alpha_{ij} \cdot (M, s_1) \stackrel{\beta^{\text{gp}}}{\sim} \alpha_{ij} \cdot (M, s_2)$. However, this is not so clear for the case of the wrapping move ω^{gp} . Consider pairs $(M, s_1) \stackrel{\omega^{\text{gp}}}{\sim} (M, s_2)$ by a wrapping move at the k th site. We need to show that $\alpha_{ij} \cdot (M, s_1) \stackrel{\beta^{\text{gp}}}{\sim} \alpha_{ij} \cdot (M, s_2)$.

By definition, there exists an $s \in \mathcal{A}^{\text{gp}}(\star_{X \cup \{\partial h\}}, \Lambda_{\text{loc}})$ such that

$$s_1 = \text{con}_{\{h\}}(s) \quad \text{and} \quad s_2 = \text{con}_{\{h\}} \left(\phi_{t_k \rightarrow e^{-h} t_k e^h}(s) \sqcup \exp_{\square} \left(-\frac{1}{2} \chi_h(M^{-1} \phi_{t_k \rightarrow e^{-h} t_k e^h} M) \right) \right),$$

Then, we have that

$$\alpha_{ij} s_1 = \text{con}_{\{h\}}(\alpha_{ij}(s)) \quad \text{and} \quad \alpha_{ij} s_2 = \text{con}_{\{h\}} \left(\alpha_{ij} \phi_{t_k \rightarrow e^{-h} t_k e^h}(s) \sqcup \left(\exp_{\square} \left(-\frac{1}{2} \chi_h(M^{-1} \phi_{t_k \rightarrow e^{-h} t_k e^h} M) \right) \right) \right).$$

If $k \neq i, j$ then α_{ij} commutes with $\phi_{t_k \rightarrow e^{-h} t_k e^h}$, thus the result follows. It remains to consider the cases of $k = i$ or $k = j$.

The case of $k = i$. Observe that α_{ij} equals to the substitution $\phi_{t_i \rightarrow t_j^{-1} t_i t_j}$, thus

$$\alpha_{ij} \circ \phi_{t_i \rightarrow e^{-h} t_i e^h} = \phi_{t_i \rightarrow e^{-h} t_j^{-1} t_i t_j e^h}$$

We will first compare $\phi_{t_i \rightarrow e^{-h} t_j^{-1} t_i t_j e^h}$ and $\phi_{t_i \rightarrow t_j^{-1} e^{-h} t_i e^h t_j}$ using a Holonomy Relation, and then $\phi_{t_i \rightarrow t_j^{-1} e^{-h} t_i e^h t_j}$ with the identity using the wrapping move to reach the desired conclusion. Explicitly, the Holonomy Relation of Figure 3 implies that

$$\text{con}_{\{h\}} \left(\begin{array}{c} e^h \uparrow e^h \uparrow e^h \uparrow \partial h \text{ ---} \\ t \bullet t \bullet t \bullet \hat{\partial} h \text{ ---} \\ e^{-h} \uparrow e^{-h} \uparrow e^{-h} \uparrow \partial h \text{ ---} \end{array} \right) = \text{con}_{\{h\}} \left(\begin{array}{c} \uparrow \uparrow \uparrow \partial h \text{ ---} \\ t \bullet t \bullet t \bullet \hat{\partial} h \text{ ---} \\ \uparrow \uparrow \uparrow \partial h \text{ ---} \end{array} \right)$$

if all appearances of t and all legs ∂h are drawn. Let $s(\partial h)$ denote s with the special leg colored explicitly by ∂h , and use the following abbreviation

$$\Psi(N) = \left(\exp_{\square} \left(-\frac{1}{2} \chi_h(N) \right) \right).$$

The above discussion implies that:

$$\begin{aligned} \alpha_{ij}(M, s_2) &= \left(\alpha_{ij} M, \text{con}_{\{h\}} \left(\phi_{t_i \rightarrow e^{-h} t_j^{-1} t_i t_j e^h}(s(\partial h)) \sqcup \Psi(\alpha_{ij} M^{-1} \cdot \phi_{t_i \rightarrow e^{-h} t_j^{-1} t_i t_j e^h}(M)) \right) \right) \\ &\stackrel{\text{Holonomy}}{=} \left(\alpha_{ij} M, \text{con}_{\{h\}} \left(\phi_{t_i \rightarrow t_j^{-1} e^{-h} t_i e^h t_j}(s(t_j \partial h)) \sqcup \Psi \left(\alpha_{ij} M^{-1} \cdot \phi_{t_i \rightarrow t_j^{-1} e^{-h} t_i e^h t_j}(M) \right) \right) \right) \\ &\stackrel{\omega^{\text{gp}}}{\sim} \left(\alpha_{ij} M, \text{con}_{\{h\}} \left(\phi_{t_i \rightarrow t_j^{-1} t_i t_j}(s(t_j \partial h)) \sqcup \Psi \left(\alpha_{ij} M^{-1} \cdot \phi_{t_i \rightarrow t_j^{-1} t_i t_j}(M) \right) \right) \right) \\ &= (\alpha_{ij} M, (\alpha_{ij} s(t_j \partial h) |_{\partial h \rightarrow 0})) \\ &= (\alpha_{ij} M, \text{con}_{\{h\}}(\alpha_{ij} s)) \\ &= \alpha_{ij}(M, s_1). \end{aligned}$$

The case of $k = j$. Observe that

$$\alpha_{ij} \circ \phi_{t_j \rightarrow e^{-h} t_j e^h} = \phi_{t_j \rightarrow e^{-h} t_j e^h, t_i \rightarrow t_j^{-1} t_i t_j}$$

First, we will compare the substitutions $\phi^{(0)}$ with $\phi^{(4)}$ (defined below) using a wrapping move ω_j^{gp} , the following Ψ -identity

$$\Psi(A^{-1}B) \sqcup \Psi(B^{-1}C) = \Psi(A^{-1}C)$$

(which is a consequence of Proposition 3.8) and Equation (14) of Lemma A.1.

Then, we will compare $\phi^{(4)}$ with $\phi^{(8)}$ using the wrapping move ω_i^{gp} .

Finally, we will compare $\phi^{(8)}$ with the identity using the case of $k = i$ above.

Here, $\phi^{(a)}$ denote the substitutions:

$$\begin{aligned} \phi^{(0)} &= & t_j &\mapsto e^{-h}t_j e^h, & t_i &\mapsto t_j^{-1}t_i t_j, \\ \phi^{(2)} &= & t_j &\mapsto e^{-h}e^{h'}t_j e^{-h'}e^h, & t_i &\mapsto e^{h'}t_j^{-1}e^{-h'}t_i e^{h'}t_j e^{-h'}, \\ \phi^{(3)} &= & t_j &\mapsto e^{h'}t_j e^{-h'}, & & \\ \phi^{(4)} &= & t_j &\mapsto t_j, & t_i &\mapsto e^h t_j^{-1} e^{-h} t_i e^h t_j e^{-h}, \\ \phi^{(6)} &= & t_j &\mapsto t_j, & t_i &\mapsto e^h t_j^{-1} e^{-h} e^{h'} t_i e^{-h'} e^h t_j e^{-h}, \\ \phi^{(7)} &= & t_i &\mapsto e^{h'} t_i e^{-h'}, & & \\ \phi^{(8)} &= & t_j &\mapsto t_j, & t_i &\mapsto e^h t_j^{-1} t_i t_j e^{-h}. \end{aligned}$$

Then, we have that

$$\begin{aligned} \alpha_{ij}(M, s_2) &= & & \left(\alpha_{ij}M, \text{con}_{\{h\}} \left(\phi^{(0)} s(\partial h) \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(0)}M) \right) \right) \\ &\stackrel{\omega_j^{\text{gp}}}{\sim} & & \left(\alpha_{ij}M, \text{con}_{\{h, h'\}} \left(\phi^{(2)} s(\partial h + \partial h') \sqcup \Psi(\phi^{(3)}\alpha_{ij}M^{-1}\phi^{(2)}M) \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(3)}\alpha_{ij}M) \right) \right) \\ &\stackrel{\Psi\text{-identity}}{=} & & \left(\alpha_{ij}M, \text{con}_{\{h, h'\}} \left(\phi^{(2)} s(\partial h + \partial h') \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(2)}M) \right) \right) \\ &\stackrel{\text{Equation (14)}}{=} & & \left(\alpha_{ij}M, \text{con}_{\{h\}} \left(\phi^{(4)} s(\partial h) \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(4)}M) \right) \right) \\ &\stackrel{\omega_i^{\text{gp}}}{\sim} & & \left(\alpha_{ij}M, \text{con}_{\{h, h'\}} \left(\phi^{(6)} s(\partial h + \partial h') \sqcup \Psi(\phi^{(7)}\alpha_{ij}M^{-1}\phi^{(6)}M) \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(7)}\alpha_{ij}M) \right) \right) \\ &\stackrel{\Psi\text{-identity}}{=} & & \left(\alpha_{ij}M, \text{con}_{\{h, h'\}} \left(\phi^{(6)} s(\partial h + \partial h') \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(6)}M) \right) \right) \\ &\stackrel{\text{Equation (14)}}{=} & & \left(\alpha_{ij}M, \text{con}_{\{h\}} \left(\phi^{(8)} s(\partial h) \sqcup \Psi(\alpha_{ij}M^{-1}\phi^{(8)}M) \right) \right) \\ &\stackrel{\text{Case of } k=i}{\sim} & & \left(\alpha_{ij}M, \text{con}_{\{h\}} (\alpha_{ij}s(\partial h)) \right) \\ &= & & \alpha_{ij}(M, s_1). \end{aligned}$$

□

6.3. Z^{rat} commutes with the String action. Recall the invariant Z^{rat} of sliced crossed links of Section 4.

Theorem 6.6. *The map*

$$\sigma \circ Z^{\text{rat}} : \mathcal{D}_X(D_g) \rightarrow \mathcal{A}^{\text{gp}}(\star_X, \Lambda) / \langle \omega^{\text{gp}} \rangle$$


is String_g -equivariant, and induces a well-defined map

$$\sigma \circ Z^{\text{rat}} : \mathcal{D}_X(D_g) / \langle \text{String}_g \rangle \rightarrow \mathcal{A}^{\text{gp}}(\star_X, \Lambda) / \langle \omega^{\text{gp}}, \text{String}_g \rangle.$$

Proof. We will show that for $D \in \mathcal{D}_X(D_g)$, we have

$$\alpha_{ij}.Z^{\text{rat}}(D) = Z^{\text{rat}}(\alpha_{ij}.D).$$

Given sliced crossed links L_1, L_2 such that $L_2 = \alpha_{ij}.L_1$, consider an accompanying link L_{12} as in the previous section. L_{12} may be presented so that all the strands going through the i th meridional disc proceed, parallel to each other, through the j th meridional disc, and then onto the $(i+1)$ st meridional disc. That is, we can find a sequence of tangles, containing some arc 'x' (not exactly the arc 'a' of Section 2.2), such that the link L_{12} is presented by taking a parallel the arc 'x' according to some word (in the example at hand, w_b). Making the following definitions $T_p, T_q, T_r, T_s, T_t, T_u$ respectively



and continuing to work with the example of the above paragraph, then the sequence will look as follows (where the j th, the i th and then the $(i + 1)$ st meridional discs are displayed):

$$\{\dots, \Delta_x^{w_b}(T_p), \Delta_x^{w_b}(T_q), \dots, \Delta_x^{w_b}(T_r), T_s, \dots, T_t, \Delta_x^{w_b}(T_u), \dots\}$$

Now, the link L_1 (resp. L_2) is presented by the diagrams recovered by gluing back in the $(i + 1)$ st (resp. i th) removed disc.

So, on the one hand (and the reader is advised to follow the following compositions on paper):

$$\begin{aligned} Z^{\text{rat}}(L_1) &= \dots \circ \Delta_x^{w_b}(Z(T_p)) \circ (I_{w_d} \otimes G_{j,w_b,w_e}) \circ \Delta_x^{w_b}(Z(T_q)) \circ \dots \\ &\quad \dots \circ \Delta_x^{w_b}(Z(T_r)) \circ (I_{w_a} \otimes G_{i,w_b}) \circ Z(T_s) \circ \dots \circ Z(T_t) \circ \Delta_x^{w_b}(Z(T_u)) \circ \dots \\ &\stackrel{\omega^{\text{SP}}}{\sim} \dots \circ \Delta_x^{w_b}(Z(T_p)) \circ (I_{w_d} \otimes G_{j,w_b,w_e}) \circ (I_{w_d} \otimes G_{i,w_b} \otimes I_{w_e}) \circ \Delta_x^{w_b}(Z(T_q)) \circ \dots \\ &\quad \dots \circ \Delta_x^{w_b}(Z(T_r)) \circ Z(T_s) \circ \dots \circ Z(T_t) \circ \Delta_x^{w_b}(Z(T_u)) \circ \dots \end{aligned}$$

A ω^{SP} -move has been used here to move the gluing word $G_{i,w}$ up to (just underneath) the gluing word arising from the j th meridional slice. This is legitimate because the meaning of the ω^{SP} -move is that such an element (if it involves *every* appearance of t_i) can be slid along a *paralleled* arc (here arising from the component x).

On the other hand:

$$\begin{aligned} Z^{\text{rat}}(L_2) &= \dots \circ \Delta_x^{w_b}(Z(T_p)) \circ (I_{w_d} \otimes G_{j,w_b,w_e}) \circ \Delta_x^{w_b}(Z(T_q)) \circ \dots \\ &\quad \dots \circ \Delta_x^{w_b}(Z(T_r)) \circ Z(T_s) \circ \dots \circ Z(T_t) \circ (I_{w_c \overline{w_b}} \otimes G_{i,w_b}) \circ \Delta_x^{w_b}(Z(T_u)) \circ \dots \\ &\stackrel{\omega^{\text{SP}}}{\sim} \dots \circ \Delta_x^{w_b}(Z(T_p)) \circ (I_{w_d} \otimes G_{i,w_b} \otimes I_{w_e}) \circ (I_{w_d} \otimes G_{j,w_b,w_e}) \circ \Delta_x^{w_b}(Z(T_q)) \circ \dots \\ &\quad \dots \circ \Delta_x^{w_b}(Z(T_r)) \circ Z(T_s) \circ \dots \circ Z(T_t) \circ \Delta_x^{w_b}(Z(T_u)) \circ \dots \end{aligned}$$

Observe that this line results from the line ending the previous paragraph of equations by the algebraic action of α_{ij} , as is required for the proof. \square

6.4. \int^{rat} **commutes with the String $_g$ action.** The following theorem says that the action of String $_g$ on diagrams commutes with \int^{rat} -integration.

Theorem 6.7. *The map*

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}^{\text{SP},0}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^{\text{SP},0}(\star_{X-X'}, \Lambda_{\text{loc}})$$

is String $_g$ equivariant and induces a well-defined map:

$$\int^{\text{rat}} dX' : \text{Int}_{X'} \mathcal{A}^{\text{SP},0}(\star_X, \Lambda_{\text{loc}}) / \langle \text{String} \rangle \rightarrow \mathcal{A}^{\text{SP},0}(\star_{X-X'}, \Lambda_{\text{loc}}) / \langle \text{String} \rangle.$$

Proof. Let the canonical decomposition of $s \in \text{Int}_{X'} \mathcal{A}^{\text{SP}}(\star_X, \Lambda_{\text{loc}})$ with respect to X' be

$$s = \exp \left(\frac{1}{2} \sum_{kl} \begin{array}{c} x_k \\ \uparrow \\ M_{kl} \\ \downarrow \\ x_l \end{array} \right) \sqcup R,$$

for some matrix $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$. Then, the canonical decomposition of $\alpha_{ij}.s$ with respect to X' is

$$\alpha_{ij}.s = \exp \left(\frac{1}{2} \sum_{kl} \begin{array}{c} x_k \\ \uparrow \\ \alpha_{ij}.M_{kl} \\ \downarrow \\ x_l \end{array} \right) \sqcup \alpha_{ij}.R.$$

Observe that $(\alpha_{ij}.M)^{-1} = \alpha_{ij}.M^{-1} \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$ which implies that $\alpha_{ij}.s$ is X' -integrable. Furthermore, we have that

$$\begin{aligned}
\int^{\text{rat}} dX'(\alpha_{ij}.s) &= \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{kl} \overset{x_k}{\uparrow} \alpha_{ij}.M_{kl} \right) \sqcup (\alpha_{ij}.R) \right) \\
&= \left\langle \exp \left(-\frac{1}{2} \sum_{kl} \overset{x_k}{\uparrow} \alpha_{ij}.(M_{kl}^{-1}) \right), \alpha_{ij}.R \right\rangle \\
&= \alpha_{ij}. \int^{\text{rat}} dX'(s).
\end{aligned}$$

Note that we could have used Theorem E.1 to deduce the above equality. \square

We end this section with the following

Lemma 6.8. (a) For all g , the identity map $\mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ maps the Inn_g -relation to the basing relation.

(b) For $g = 1, 2$, the identity map $\mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ maps the String_g -relation to the basing relation.

Proof. Recall that Inn_g is the subgroup of String_g that consists of inner automorphisms of the free group F .

For the first part, an inner automorphism by an element $g \in F$ maps a bead $b \in \Lambda_{\text{loc}}$ to $g^{-1}bg \in \Lambda_{\text{loc}}$. The relations of Figure 2 cancel the g at every trivalent vertex of a diagram. The remaining vertices change a diagram by a basing move β_2^{SP} .

The second part follows from the fact that $\text{Inn}_g = \text{String}_g$ for $g = 1, 2$, see [CS2, K]. \square

6.5. The definition of the rational invariant Z^{rat} of ∂ -links. .

Theorems 6.1, 6.4, 6.6 and 6.7 imply that there is a map

$$Z^{\text{rat}, \text{gp}} : \partial\text{-links} \rightarrow \mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}}, \text{String} \rangle$$

defined as follows: for a ∂ -link (M, L) of g components, choose a sliced crossed link C in $\mathcal{D}(D_g)$ whose components are in 1-1 correspondence with a set X , and define

$$Z^{\text{rat}, \text{gp}}(M, L) = \left(\int^{\text{rat}} dX (\sigma \circ \check{Z}^{\text{rat}}(C)) \right) \cdot c_+^{\sigma_+(B)} c_-^{\sigma_-(B)} \in \mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}}, \text{String} \rangle$$

where $c_{\pm} = \int dU \check{Z}(S^3, U_{\pm})$ are some universal constants of the unit-framed unknot U_{\pm} and where B is the linking matrix of C and $\sigma_{\pm}(B)$ is the number of positive (resp. negative) eigenvalues of B . The above definition makes sense and uses the fact that as in Section 5.5, the set $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle$ has a well-defined action of the group $\mathcal{A}^{\text{gp}}(\phi)$, which commutes with the String action (where String acts trivially on $\mathcal{A}^{\text{gp}}(\phi)$).

Composing the $Z^{\text{rat}, \text{gp}}$ invariant with the projection $\mathcal{A}^{\text{gp}, 0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}}, \text{String} \rangle \rightarrow \mathcal{A}^0 / \langle \omega, \text{String} \rangle$ of Proposition 3.12, defines the rational invariant Z^{rat} of ∂ -links that appears in Theorem 1.2.

7. A COMPARISON BETWEEN Z AND Z^{rat}

The goal of this section is to compare the invariants Z^{rat} and Z of F -links and to finish the proof of Theorem 1.2. This will be obtained by comparing the construction of the Aarhus integral \int with that of \int^{rat} .

7.1. The Hair map. First, we need to define a map Hair from the space of diagrams in which Z^{rat} takes values to the space of diagrams in which Z takes values. Given generators $\{t_1, \dots, t_g\}$ of the free group, let $\{h_1, \dots, h_g\}$ be noncommuting variables such that $t_i = e^{h_i}$ and let $\text{Hair} : \mathcal{A}(\star_X, \Lambda) \rightarrow \mathcal{A}(\star_{X \cup H})$ be defined as follows: replace a variable by the exponential of hair, as in Figure 5. This extends to a map $\text{Hair} : \mathcal{A}(\star_X, \hat{\Lambda}) \rightarrow \mathcal{A}(\star_{X \cup H})$ (and, after composition with $\mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_X, \hat{\Lambda})$), also to a map

$$\text{Hair} : \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_{X \cup H})$$

whose image lies in the span of all diagrams that do not contain a H -labeled tree as a connected component.

$$\uparrow t_i \rightarrow \sum_{n=0}^{\infty} \frac{1}{n!} \begin{array}{c} \uparrow \\ \rightarrow \\ \rightarrow \\ \rightarrow \end{array} n \text{ } h_i\text{-labeled legs}$$

Figure 5. From beads to legs.

Recall the extension $\mathcal{A}^{\text{gp},0}$ of \mathcal{A}^{gp} . Given a pair $(M, s) \in \mathcal{B}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z}) \times \mathcal{A}^{\text{gp}}$, we can define an extended map $\text{Hair} : \mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^{\text{gp}}(\star_{X \cup H})$ by

$$\text{Hair}(M, s) = \exp_{\sqcup} \left(-\frac{1}{2} \chi(M) \right) \sqcup \text{Hair}(s).$$

Note that the \exp_{\sqcup} term in the above expression is a disjoint union of *wheels*.

Lemma 7.1. *There exist maps that fit in a commutative diagram*

$$\begin{array}{ccc} \mathcal{A}^{\text{gp},0}(\otimes_X, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle & \xrightarrow{\quad\quad\quad} & \mathcal{A}^0(\otimes_X, \Lambda_{\text{loc}}) / \langle \omega \rangle \\ \downarrow & & \downarrow \\ \mathcal{A}^{\text{gp},0}(\otimes_X, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}}, \text{String}_g \rangle & \xrightarrow{\quad\quad\quad} & \mathcal{A}^0(\otimes_X, \Lambda_{\text{loc}}) / \langle \omega, \text{String}_g \rangle \\ & \searrow \text{Hair} & \swarrow \text{Hair} \\ & \mathcal{A}(\otimes_{X \cup H}) & \end{array}$$

Proof. Proposition D.1 implies that the horizontal maps are well-defined. It is easy to show that the X -flavored group-like and infinitesimal basing relations (defined in Section D) are mapped by the Hair-map to X -flavored infinitesimal basing relations. It is also easy to see that the ω^{gp} and String_g relations are mapped to H -flavored basing relations. \square

Remark 7.2. The invariant Z^{rat} takes values in $\mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle$, the invariant Z takes values in $\mathcal{A}(\otimes_H)$ and the image of the map $\text{Hair} : \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \rightarrow \mathcal{A}(\otimes_H)$ lies in the subspace of $\mathcal{A}(\otimes_H)$ spanned by diagrams without tree components. If we expect Z^{rat} to determine Z , we should restrict attention to the class of links whose Kontsevich integral has vanishing “tree part”. Since the vanishing of the tree-part is equivalent to the vanishing of all Milnor’s invariants, we should only consider links with vanishing Milnor’s invariants. Such links include, and perhaps coincide with, all *sublinks of homology boundary links*, see for example [Le2]. For simplicity, we will not deal with homology boundary links, but rather with boundary links.

We will need a slightly renormalized version Hair' of the Hair map, obtained by considering the disjoint union of the map Hair and $\nu(h_1) \sqcup \cdots \sqcup \nu(h_g)$, where $\nu = Z(S^3, U)$.

7.2. A comparison between Z^{rat} and Z . Consider the set $\mathcal{L}_X(\mathcal{O})$ of links $L \cup \mathcal{O}$ in S^3 where \mathcal{O} is an unlink whose components are in 1-1 correspondence with the set $H = \{h_1, \dots, h_g\}$ and L is a link whose components are in 1-1 correspondence with a set X . Define a map

$$\varphi_{\sqcup} : \mathcal{D}_X(\mathcal{O}) \rightarrow \mathcal{L}_X(\mathcal{O})$$

that maps the associated link L of a sliced crossed link in $D_g \times I$ to the link $L \cup \mathcal{O} \subset S^3$ according to a standard inclusion of $D_g \times I$ in S^3 , by adding a g component unlink \mathcal{O} , one component for each gluing site of D_g as follows:

$$\varphi_{\sqcup} \left(\left\{ \dots, \begin{array}{|c|} \hline T_i \\ \hline \end{array}, \begin{array}{|c|} \hline T_{i+1} \\ \hline \end{array}, \dots \right\} \right) \rightarrow \begin{array}{|c|} \hline T_i \\ \hline \end{array} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array}$$

Consider in addition the Kontsevich integral

$$\sigma \circ Z : \mathcal{L}_X(\mathcal{O}) \rightarrow \mathcal{A}(\otimes_{X \cup H})$$

where $\mathcal{A}(\otimes_{X \cup H})$ denotes the space of univalent graphs whose legs are colored by $X \cup H$, modulo the AS, IHX relations and the *infinitesimal basing relations* discussed in Appendix C. The purpose of this section is to show that

Theorem 7.3. *The following diagram commutes*

$$\begin{array}{ccc} \mathcal{N}_X(\mathcal{O}) & \xrightarrow{\sigma \circ Z^{\text{rat}}} & \mathcal{A}^{\text{SP}}(\otimes_X, \Lambda) / \langle \omega^{\text{SP}} \rangle \\ \varphi_{\cup} \downarrow & & \downarrow \text{Hair}^{\nu} \\ \mathcal{L}_X(\mathcal{O}) & \xrightarrow{\sigma \circ Z} & \mathcal{A}(\otimes_{X \cup H}) \end{array}$$

The proof will utilize the following *magic formula* for the Kontsevich integral of the Long Hopf Link, conjectured in [A0] (in conjunction with the so-called Wheels and Wheeling conjectures) and proven in [BLT].

Theorem 7.4. [BLT]

$$(\sigma_{\{x,y\}} \circ Z) \left(\begin{array}{c} \uparrow x \\ \text{---} \\ \downarrow y \end{array} \right) = \exp \left(\begin{array}{c} x \\ | \\ y \end{array} \right) \sqcup \nu(y) \in \mathcal{A}(\star_{\{x\}} \otimes \{y\}).$$

Proof of Theorem 7.3. We will show first that the following diagram commutes:

$$(3) \quad \begin{array}{ccc} \mathcal{D}_X(\mathcal{O}) & \xrightarrow{\sigma \circ Z^{\text{rat}}} & \mathcal{A}^{\text{SP}}(\otimes_X, \Lambda) \\ \varphi_{\cup} \downarrow & & \downarrow \text{Hair}^{\nu} \\ \mathcal{L}_X(\mathcal{O}) & \xrightarrow{\sigma \circ Z} & \mathcal{A}(\otimes_{X \cup H}) \end{array}$$

Given a word w in the symbols \uparrow and \downarrow , let

- $\text{LongHopf}_{j,w}$ denote the long Hopf link cabled according to the word w , with the closed component labeled h_j . For example:

$$\text{LongHopf}_{j,\downarrow\uparrow\uparrow} = \begin{array}{c} \uparrow \\ \uparrow \\ \uparrow \\ \text{---} \\ \downarrow \\ \downarrow \\ \downarrow \end{array} h_j$$

- $G_{j,w}^h$ denote the j th hairy gluing word corresponding to w . This is definition by example:

$$G_{j,\downarrow\uparrow\uparrow}^h = \downarrow e^{-h_j} \uparrow e^{h_j} \uparrow e^{h_j}$$

We use the following corollary to Theorem 7.4.

Corollary 7.5.

$$(\sigma_{\{h_j\}} \circ Z) (\text{LongHopf}_{j,w}) = G_{j,w}^h \sqcup \nu(h_j) \in \mathcal{A}(\uparrow_w \otimes \{h_j\}).$$

The commutativity of diagram (3) now follows by the definition of Z^{rat} . Theorem 7.3 follows from the commutativity of diagram (3), after considering the quotient $\mathcal{N}_X(\mathcal{O})$ of $\mathcal{D}_X(\mathcal{O})$ (given by Proposition 2.4) and symmetrizing (i.e., composing with $\sigma : \mathcal{A}^{\text{SP}}(\uparrow_X, \Lambda) \rightarrow \mathcal{A}^{\text{SP}}(\otimes_X, \Lambda)$) together with the fact that Hair^{ν} commutes with symmetrization. □

7.3. A comparison between \int^{rat} and \int . Recall that the Aarhus integral respects the group-like basing relations as shown in [A, Part II, Proposition 5.6] and also the infinitesimal basing relations as was observed by D. Thurston, see [BL, Proposition 2.2]. In other words, there is a well-defined map

$$\int dX' : \text{Int}_{X'} \mathcal{A}(\otimes_{X \cup H}) \rightarrow \mathcal{A}(\otimes_{(X-X') \cup H}),$$

where $H = \{h_1, \dots, h_g\}$. The purpose of this section is to show that

Theorem 7.6. *The following diagram commutes*

$$\begin{array}{ccc}
\text{Int}_{X'} \mathcal{A}^{\text{gp},0}(\otimes_X, \Lambda_{\text{loc}}) & \xrightarrow{f^{\text{rat}} dX'} & \mathcal{A}^{\text{gp},0}(\otimes_{X-X'}, \Lambda_{\text{loc}}) \\
\text{Hair}' \downarrow & & \downarrow \text{Hair}' \\
\text{Int}_{X'} \mathcal{A}(\otimes_{X \cup H}) & \xrightarrow{f dX'} & \mathcal{A}(\otimes_{(X-X') \cup H})
\end{array}$$

Proof. It follows from Remark E.2 of Appendix E. \square

7.4. End of the proof of Theorem 1.2. We begin by observing that the following diagram commutes, by the definition of the invariant $Z^{\text{rat},\text{gp}}$, using the work of Section 6:

$$\begin{array}{ccc}
F\text{-links} & \xrightarrow{Z^{\text{rat},\text{gp}}} & \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}})/\langle \omega^{\text{gp}} \rangle \\
\downarrow & & \downarrow \\
\partial\text{-links} & \xrightarrow{Z^{\text{rat},\text{gp}}} & \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}})/\langle \omega^{\text{gp}}, \text{String} \rangle
\end{array}$$

We now claim that the following diagram commutes:

$$(4) \quad \begin{array}{ccc}
\partial\text{-links} & \xrightarrow{Z^{\text{rat},\text{gp}}} & \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}})/\langle \omega^{\text{gp}}, \text{String} \rangle \\
\parallel & & \downarrow \text{Hair}' \\
\partial\text{-links} & \xrightarrow{Z} & \mathcal{A}(\otimes_H)
\end{array}$$

The above two diagrams, together with Lemma 7.1 Theorem 1.2. In order to show the commutativity of (4), we will assemble several commutative diagrams in one with two layers. The top layer consists of the rational invariants Z^{rat} and its integration theory f^{rat} ; the bottom layer consists of the LMO-Aarhus-Kontsevich integral Z and its integration theory f .

$$\begin{array}{ccccc}
& & \mathcal{N}_X(\mathcal{O}) & \xrightarrow{\quad\quad\quad} & \mathcal{N}_X(\mathcal{O})/\langle \kappa_X, \text{String} \rangle \\
& \swarrow & \vdots & & \swarrow Z^{\text{rat},\text{gp}} \\
\mathcal{A}^{\text{gp}}(\otimes_X, \Lambda)/\langle \omega^{\text{gp}} \rangle & \xrightarrow{f^{\text{rat}} dX} & \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}})/\langle \omega^{\text{gp}}, \text{String} \rangle & & \\
\vdots & & \vdots & & \vdots \\
& \swarrow & \mathcal{L}_X(\mathcal{O}) & \xrightarrow{\quad\quad\quad} & \partial\text{-links} \\
& \swarrow & \vdots & & \swarrow Z \\
\mathcal{A}(\otimes_{X \cup H}) & \xrightarrow{f dX} & \mathcal{A}(\otimes_H) & &
\end{array}$$

Theorems 4.1, 7.3 and 7.6 imply that all faces of the above cube except the side right one commute. Since the map $\mathcal{N}_X(\mathcal{O}) \rightarrow \mathcal{L}_X(\mathcal{O})/\langle \kappa, \text{String} \rangle$ is onto, it follows that the remaining face of the cube commutes.

Using the identification $\mathcal{A}(\cup_H) \cong \mathcal{A}(\otimes_H)$ (see Proposition D.1) it follows that diagram (4) commutes. This concludes the proof of Theorem 1.2. \square

8. A UNIVERSAL PROPERTY OF Z^{rat}

In this section we will show that the rational invariant Z^{rat} , evaluated on acyclic F -links, is characterized by a universal property. Namely, it is the universal finite type invariant of acyclic F -links with respect to the null filtration, [Ga].

The arguments in this section are minor modifications of arguments given in [GR] for the case of knots. For self-completeness, though, we will include them here.

8.1. A relative version of the Z^{rat} invariant. We already discussed in the introduction (before the statement of Theorem 1.11) the notion of finite type invariants of F -links, based on the null move. The null move is described in terms of surgery on a clasper whose leaves lie in the kernel of the basing map ϕ of an F -link.

Our task here is to give a *relative version* of the Z^{rat} invariant for null links in the complement of an F -link. Given an F -link (M, L, ϕ) , where $\phi : \pi_1(M - L) \rightarrow F$, we will call a link $C \in M - L$ *null relative to* (M, L, ϕ) if each component of C is mapped to 1 under ϕ . Notice that surgery under a null link preserves the underlying F -structure. In what follows, we will omit ϕ from the notation, for typographical reasons.

Given a null link C relative to an F -link (M, L) , choose a sliced crossed link $L' \cup C' \in \mathcal{D}(D_g)$ such that after surgery on $L' \subset S^3 - \mathcal{O}$ it gives rise to the F -link (M, L) with $C' \subset S^3 - \mathcal{O}$ transformed to a null link C relative to (M, L) . We define

$$\begin{aligned} Z^{\text{rat,gp}}((M, L), C) &= \int^{\text{rat}} dX(\sigma \circ \check{Z}^{\text{rat}}(L' \cup C')) / (\text{normalization}) \in \mathcal{A}^{\text{gp},0}(\otimes_Y, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \\ Z^{\text{rat,gp}}((M, L)_C) &= \int^{\text{rat}} dY(Z^{\text{rat}}((M, L), C)) / (\text{normalization}) \in \mathcal{A}^{\text{gp},0}(\otimes, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \end{aligned}$$

where X (resp. Y) is in 1-1 correspondence with the components of L (resp. C), and where (normalization) refers to factors inserted to deal with the Kirby move of adding a distant union of unit-framed unlinks, as in Section 5.5. Theorem 1.2 has the following relative version (as is revealed by its proof in Section 5.5):

Theorem 8.1. *There exists an invariant*

$$Z^{\text{rat,gp}} : F\text{-links} \cup \text{null } C \rightarrow \mathcal{A}^{\text{gp},0}(\otimes_C, \Lambda_{\text{loc}}) / \langle \omega \rangle$$

which is local (i.e., compatible with respect to considering sublinks of C) and fits in a commutative diagram

$$\begin{array}{ccc} F\text{-links} \cup \text{null } C & \xrightarrow{Z^{\text{rat,gp}}} & \mathcal{A}^{\text{gp},0}(\otimes_Y, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \\ \downarrow C\text{-surgery} & & \downarrow \int^{\text{rat}} dY \\ F\text{-links} & \xrightarrow{Z^{\text{rat,gp}}} & \mathcal{A}^{\text{gp},0}(\Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \end{array}$$

As in Section 5.5, we denote by Z^{rat} the composition of $Z^{\text{rat,gp}}$ with the map $\mathcal{A}^{\text{gp},0}(\otimes_Y, \Lambda_{\text{loc}}) / \langle \omega^{\text{gp}} \rangle \rightarrow \mathcal{A}^0(\otimes_Y, \Lambda_{\text{loc}}) / \langle \omega \rangle$.

Given a null link C relative to (M, L) , one can define an *equivariant linking matrix* $\text{lk}_{(M,L)}(C)$, by considering the linking numbers of the components of the lift \tilde{C} of C in the free cover of $M - L$. $\text{lk}_{(M,L)}(C)$ is a Hermitian matrix over Λ_{loc} , well defined up to conjugation by a diagonal matrix with elements in F . The details of the definition, together with the following proposition will appear in a subsequent publication, [Ga2]:

Proposition 8.2. *For C and (M, L) as above, the covariance matrix cov of Z^{rat} satisfies*

$$\text{cov}(Z^{\text{rat}}((M, L), C)) = \text{lk}_{(M,L)}(C).$$

The discussion in this section simplifies considerably when the F -links in question are acyclic. In this case, there is no need to localize, and the wrapping relation becomes trivial. In other words, we get an invariant

$$Z^{\text{rat}} : \text{Acyclic } F\text{-links} \cup \text{null links} \rightarrow \mathcal{A}(\otimes_Y, \Lambda) / \langle \omega \rangle.$$

8.2. Surgery on claspers. Let us review in brief some elementary facts about surgery on claspers. For a more detailed discussion, we refer the reader to [GGP] and also [GR, Section 3].

Consider a null clasper G of degree 1 in $S^3 - \mathcal{O}$ (a so-called Y -graph). Surgery on G can be described by surgery on a six component link $E \cup L$ associated to G , where E (resp. L) is the three component link that consists of the edges (resp. leaves) of G .

Lemma 8.3. *The equivariant linking matrix of $E \cup L$ and its negative inverse are given as follows:*

$$(5) \quad \begin{pmatrix} 0 & I \\ I & \text{lk}_{(S^3, \mathcal{O})}(L_i, L_j) \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \text{lk}_{(S^3, \mathcal{O})}(L_i, L_j) & -I \\ -I & 0 \end{pmatrix}.$$

Proof. The 0 block follows from the fact that $\{E_i, E_j, \mathcal{O}\}$ is an unlink. The I block follows from the fact that E_i is a meridian of L_i , using the formula for the Kontsevich integral of the Long Hopf Link, together with the fact that the linking number between L_i and \mathcal{O} is zero. \square

The six component link $E \cup L$ is partitioned in three blocks of two component links $A_i = \{E_i, L_i\}$ each for $i = 1, 2, 3$, the *arms* of G . A key feature of surgery on G is the fact that surgery on any proper subset of the set of arms does not alter M . In other words, alternating (S^3, \mathcal{O}) with respect to surgery on G equals to alternating (S^3, \mathcal{O}) with respect to surgery on all nine subsets of the set of arms $A = \{A_1, A_2, A_3\}$. That is,

$$(6) \quad Z^{\text{rat}}([(S^3, \mathcal{O}), G]) = Z^{\text{rat}}([(S^3, \mathcal{O}), A]).$$

Due to the locality property of the Z^{rat} invariant (i.e., Theorem 8.1) the nontrivial contributions to the right hand side of Equation (6) come from the *strutless part* $Z^t((S^3, \mathcal{O}), A_1 \cup A_2 \cup A_3)$ of $Z^t((S^3, \mathcal{O}), A_1 \cup A_2 \cup A_3)$ that consists of graphs with legs *touch* (i.e., are colored by) all three arms of G . The above discussion generalizes to the case of an arbitrary disjoint union of clasps.

8.3. Counting above the critical degree.

Proposition 8.4. *The Euler degree n part of Z^{rat} is a type n invariant of acyclic F -links with values in $\mathcal{A}_n(\Lambda)$.*

Proof. Since $\mathcal{A}_{\text{odd}}(\Lambda) = 0$, it suffices to consider the case of even n . Suppose that $G = \{G_1, \dots, G_m\}$ (for $m \geq 2n + 1$) is a collection of null clasps in $S^3 - \mathcal{O}$ each of degree 1, and let A denote the set of arms of G . Equation (6) and its following discussion implies that

$$Z^{\text{rat}}([(S^3, \mathcal{O}), G]) = Z^{\text{rat}}([(S^3, \mathcal{O}), A])$$

and that the nonzero contribution to the right hand side come from diagrams in $Z^{\text{rat},t}((S^3, \mathcal{O}), A)$ that touch all arms. Thus, contributing diagrams have at least $3(2n + 1) + 1 = 6n + 4$ A -colored legs, to be glued pairwise.

Notice that the diagrams in $Z^{\text{rat},t}((S^3, \mathcal{O}), A)$ contain no struts. Thus, at most three A -colored legs meet at a vertex, and after gluing the A -colored legs we obtain trivalent graphs with at least $(6n + 4)/3 = 2n + 4/3$ trivalent vertices, in other words of Euler degree at least $2n + 2$. Thus, $Z_{2n}^{\text{rat}}([(S^3, \mathcal{O}), G]) = 0 \in \mathcal{A}_{2n}(\Lambda)$, which implies that Z_{2n}^{rat} is a invariant of acyclic F -links of type $2n$ with values in $\mathcal{A}_{2n}(\Lambda)$. \square

Sometimes the above vanishing statement is called *counting above the critical degree*.

8.4. Counting on the critical degree. Our next statement can be considered as *counting on the critical degree*. We need a preliminary definition.

Definition 8.5. Consider a null clasper G in $S^3 - \mathcal{O}$ of degree $2n$, and let $G^{\text{break}} = \{G_1, \dots, G_{2n}\}$ denote the collection of degree 1 clasps G_i which are obtained by inserting a Hopf link in the edges of G . Let $G^{\text{nl}} = \{G_1^{\text{nl}}, \dots, G_{2n}^{\text{nl}}\}$ denote the collection of abstract univalent graph obtained by removing the leaves of the G_i (and leaving one leg, or univalent vertex, for each leave behind). Choose arcs from a fixed base point to the trivalent vertex of each G_i^{nl} , which allows us to define the equivariant linking numbers of the leaves of G^{break} . Then the *complete contraction* $\langle G \rangle \in \mathcal{A}(\Lambda)$ of G is defined to be the sum over all ways of gluing pairwise the legs of G^{nl} , and placing as a bead the equivariant linking number of the corresponding leaves.

The result of a complete contraction of a null clasper G is a well-defined element of $\mathcal{A}(\Lambda)$. Changing the arcs is taken care by the holonomy relations in $\mathcal{A}(\Lambda)$. The next proposition computes the *symbol* of Z^{rat} :

Proposition 8.6. *If G is a null clasper of degree $2n$ in $S^3 - \mathcal{O}$, then*

$$Z_{2n}^{\text{rat}}([(S^3, \mathcal{O}), G]) = \langle G \rangle \in \mathcal{A}_{2n}(\Lambda) / \langle \omega \rangle$$

Proof. It suffices to consider a collection $G = \{G_1, \dots, G_{2n}\}$ of clasps in S^3 each of degree 1. Let A denote the set of arms of G . The counting argument of the above Proposition shows that the contributions to $Z_{2n}^{\text{rat}}([(S^3, \mathcal{O}), G]) = Z_{2n}^{\text{rat}}([(S^3, \mathcal{O}), A])$ come from complete contractions of a disjoint union $D = Y_1 \cup \dots \cup Y_{2n}$ of $2n$ *vortices*. A vortex is the diagram Y , the next simplest univalent graph after the strut. Furthermore,

the $6n$ legs of D should touch all $6n$ arms of G . In other words, there is a 1-1 correspondence between the legs of such D and the arms of G .

Consider a leg l of D that touches an arm $A_l = \{E_l, L_l\}$ of G . If l touches L_l , then due to the restriction of the negative inverse linking matrix of G (see Lemma 8.3), it needs to be contracted to another leg of D that touches E_l . But this is impossible, since the legs of D are in 1-1 correspondence with the arms of G .

Thus, each leg of D touches precisely one edge of G . In particular, each leg of D is colored by three edges of G .

Consider a vortex colored by three edges of G as part of the Z^{rat} invariant. Since the three edges of G is an unlink in a ball disjoint from \mathcal{O} , it follows that the beads on the edges of the vortex are 1, if the three edges belong to the same G_i , and 0 otherwise.

Thus, the diagrams D that contribute are a disjoint union of $2n$ vortices $Y = \{Y_1, \dots, Y_{2n}\}$ and these vortices are in 1-1 correspondence with the set of claspers $\{G_1, \dots, G_{2n}\}$, in such a way that the legs of each vortex Y_i are colored by the edges of a unique clasper G_j .

After we glue the legs of such Y using the negative inverse linking matrix of G , the result follows. \square

Let us mention that the discussion of Propositions 8.4 and 8.6 really applied to the unnormalized rational Aarhus integral; however since we are counting above the critical degree, we need only use the degree 0 part of the normalization which equals to 1; in other words we can forget about the normalization.

The above proposition is useful in realization properties of the Z_{2n} invariant, but also in proving the following *Universal Property*:

Proposition 8.7. *For all n , the composite map of $[\text{Ga}]$ with that of Proposition 8.6*

$$\mathcal{A}_{2n}(\Lambda)/\langle\omega\rangle \longrightarrow \mathcal{G}_{2n}^{\text{null}} \xrightarrow{Z_{2n}^{\text{rat}}} \mathcal{A}_{2n}(\Lambda)/\langle\omega\rangle$$

is the identity. Since the map on the left is onto, it follows that the above maps are isomorphism.

Remark 8.8. In the above propositions 8.4-8.7, S^3 can be replaced by any integral homology 3-sphere (or even a rational homology 3-spheres) M . \square

Clearly, Propositions 8.4-8.7 imply Theorem 1.11.

APPENDIX A. A BRIEF REVIEW OF DIAGRAMMATIC CALCULUS

A.1. Some useful identities. In this section we establish some notations and collect some lemmas. Let X denote some finite set. Given a sum of diagrams s with legs colored by a set X , and a subset $X' = \{x_1, \dots, x_r\}$ of X , we will often denote s by $s(x_1, \dots, x_r)$ or simply by $s(x)$. This notation allows us to extend linearly to diagrams whose legs are labeled by formal linear combinations of elements of X . For example, for $a, b \in \mathbb{Q}$, $v, w \in X$, $s(\dots, x_{i-1}, av + bw, x_{i+1}, \dots)$ denotes the element obtained from s by replacing each diagram appearing in s by the sum of all diagrams obtained by relabeling each leg labeled x_i by either v (with a multiplicative factor of a) or w (with a multiplicative factor of b). Given diagrams $A(x), B(x)$, we let

$$\langle A(\partial x), B(x) \rangle_X \quad \text{and} \quad A(\partial x) \flat_X B(x)$$

denote respectively the sum of all ways of pairing all X -colored legs of A with all (resp. some) X -colored legs of B .

Furthermore, for some $q \in \Lambda_{\text{loc}}$, $s(\dots, x_{i-1}, x_i q, x_{i+1}, \dots)$ denotes the element obtained by pushing a bead labeled by q onto each leg labeled x_i in the sense specified by the following diagram:

$$\begin{array}{ccc} \uparrow 1 & \rightarrow & \uparrow q \\ x_i & & x_i \end{array}$$

Taken together, these conventions give meaning to such expressions as $s(xM)$, for $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$.

The following lemma collects a number of useful identities, which the reader would be well-advised to understand.

Lemma A.1. (a) *Let $A \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, $B \in \mathcal{A}(\star_{X \cup Y}, \Lambda_{\text{loc}})$ (for Y a bijective copy of X), and let X' and X'' denote bijective copies of X .*

$$(7) \quad \langle A(\partial x), B(x, y)|_{x=y} \rangle_X = \langle A(\partial x' + \partial x''), B(x', x'') \rangle_{X' \cup X''}$$

(b) Let $A_1, A_2, B \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$.

$$(8) \quad \langle A_1(\partial x)A_2(\partial x), B(x) \rangle_X = \langle A_2(\partial X), A_1(\partial x) \flat_X B(x) \rangle_X.$$

(c) Let $A, B \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, and let Y be a bijective copy of X .

$$(9) \quad A(\partial x) \flat_X B(x) = \langle A(\partial y), B(x+y) \rangle_Y$$

(d) Let $A = \exp(a) \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$ and $B = \exp(b) \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$. Then

$$(10) \quad \exp(a) \flat_{X'} \exp(b) = \exp(c),$$

where c is the sum of all connected diagrams that arise by pairing all legs labeled from X' of some diagram (not necessarily connected) appearing in A to some legs of some diagram (not necessarily connected) appearing in B .

(e) Let $A \in \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, let Y be a bijective copy of X , and let $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$.

$$(11) \quad \left(\exp \left(\begin{array}{c} \partial x_j \\ \uparrow \\ \bullet \\ y_i \end{array} M_{ij} \right) \right) \flat_X A(x) = A(yM).$$

(f) In particular,

$$(12) \quad \left\langle \exp \left(\begin{array}{c} \partial x \\ | \\ y \end{array} \right), B(x) \right\rangle_x = B(y)$$

$$(13) \quad \exp \left(\begin{array}{c} \partial x \\ | \\ y \end{array} \right) \flat_x \exp(B(x)) = \exp \left(\sum \text{all ways of replacing some of the } x\text{-legs of } B \text{ by } y\text{-legs} \right)$$

(g) In addition, if $s(\partial z) \in \mathcal{A}(\star_{X \cup \{h', h'', \partial z\}}, \Lambda_{\text{loc}})$ then

$$(14) \quad \text{con}_{\{h', h''\}}(s(\partial h' + \partial h'')) = \text{con}_{\{h\}}(s(\partial h))|_{h', h'' \mapsto h}.$$

A.2. Iterated Integration. This section proves an Iterated Integration property of the \int^{rat} integral. Fix labeling sets $X'' \subset X' \subset X$.

Lemma A.2 (Iterated integration). *Given $s \in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \cap \text{Int}_{X''} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$, then,*

$$\int^{\text{rat}} dX''(s) \in \text{Int}_{X' - X''} \mathcal{A}(\star_{X - X''}, \Lambda_{\text{loc}})$$

$$\int^{\text{rat}} d(X' - X'') \left(\int^{\text{rat}} dX''(s) \right) = \int^{\text{rat}} dX'(s).$$

Proof. Denote the components of X' and X'' by a_i and b_j respectively. Consider the covariance matrix of s with respect to X' , presented with respect to this basis:

$$M = \begin{bmatrix} A & C \\ C^* & B \end{bmatrix},$$

where $A = A^*$ and $B = B^*$, where here and below A^* denotes the transpose of C , followed by the conjugation $t_i \rightarrow t_i^{-1}$. If the canonical decomposition of s with respect to X' is then,

$$\exp \left(\begin{array}{c} a_j \\ \uparrow \\ \frac{1}{2} \bullet A_{ij} \\ a_i \end{array} + \begin{array}{c} b_j \\ \uparrow \\ \frac{1}{2} \bullet B_{ij} \\ b_i \end{array} + \begin{array}{c} b_j \\ \uparrow \\ \bullet C_{ij} \\ a_i \end{array} \right) \sqcup R,$$

(suppressing the summations) then the canonical decomposition of s with respect to X'' is

$$\exp \left(\begin{array}{c} a_j \\ \uparrow \\ \frac{1}{2} \bullet A_{ij} \\ a_i \end{array} \right) \sqcup \left(\exp \left(\begin{array}{c} b_j \\ \uparrow \\ \frac{1}{2} \bullet B_{ij} \\ b_i \end{array} + \begin{array}{c} b_j \\ \uparrow \\ \bullet C_{ij} \\ a_i \end{array} \right) \sqcup R \right).$$

Using the identities of Lemma A.1, we have

$$\begin{aligned}
\int^{\text{rat}} dX''(s) &= \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial a_i}^{\partial a_j} A_{ij}^{-1} \right), \exp \left(\frac{1}{2} \uparrow_{b_i}^{b_j} B_{ij} + \uparrow_{a_i}^{b_j} C_{ij} \right) \sqcup R \right\rangle_{X''} \\
&= \left\langle \left(\exp \left(\uparrow_{\partial a_i}^{b_j} C_{ij} \right) \flat_{X'} \exp \left(-\frac{1}{2} \uparrow_{a_i}^{a_j} A_{ij}^{-1} \right) \right) \Big|_{a_i \rightarrow \partial a_i}, \exp \left(\frac{1}{2} \uparrow_{b_i}^{b_j} B_{ij} \right) \sqcup R \right\rangle_{X''} \\
&= \exp \left(\frac{1}{2} \uparrow_{b_i}^{b_j} (B - C^* A^{-1} C)_{ij} \right) \sqcup \left(\left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial a_i}^{\partial a_j} A_{ij}^{-1} - \uparrow_{\partial a_i}^{b_j} (A^{-1} C)_{ij} \right), R \right\rangle_{X''} \right)
\end{aligned}$$

It is easy to see that if M and A are invertible over \mathbb{Z} , then so is $B - C^* A^{-1} C$ over \mathbb{Z} , see [A, Part II, Proposition 2.13]. Thus, the first part of the theorem follows. Continuing,

$$\begin{aligned}
&\int^{\text{rat}} d(X'' - X') \left(\int^{\text{rat}} dX'(s) \right) \\
&= \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial b_i}^{\partial b_j} (B - C^* A^{-1} C)_{ij}^{-1} \right), \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial a_i}^{\partial a_j} A_{ij}^{-1} - \uparrow_{\partial a_i}^{b_j} (A^{-1} C)_{ij} \right), R \right\rangle_{X''} \right\rangle_{X' - X''} \\
&= \left\langle \left(\exp \left(-\uparrow_{\partial a_i}^{\partial b_j} (A^{-1} C)_{ij} \right) \flat_{X' - X''} \exp \left(-\frac{1}{2} \uparrow_{b_i}^{b_j} (B - C^* A^{-1} C)_{ij}^{-1} \right) \right) \Big|_{b_i \rightarrow \partial b_i}, \exp \left(-\frac{1}{2} \uparrow_{\partial a_i}^{\partial a_j} A_{ij}^{-1} \right), R \right\rangle_{X'} \\
&= \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial a_i}^{\partial a_j} (A^{-1} + A^{-1} C (B - C^* A^{-1} C)^{-1} C^* A^{-1})_{ij} \right. \right. \\
&\quad \left. \left. + \uparrow_{\partial a_i}^{\partial b_j} (A^{-1} C (B - C^* A^{-1} C)^{-1})_{ij} - \frac{1}{2} \uparrow_{\partial b_i}^{\partial b_j} (B - C^* A^{-1} C)_{ij}^{-1} \right), R \right\rangle_{X'}
\end{aligned}$$

The identification of the term on the left of the pairing with $\exp \left(-\frac{1}{2} \uparrow_{\partial x_i}^{\partial x_j} M_{ij}^{-1} \right)$ completes the proof. \square

Remark A.3. With the notation of the previous lemma, the matrices $A \oplus (B - C^* A^{-1} C)$ and M are congruent by a product of elementary matrices, as follows from the following identity

$$\begin{bmatrix} A & 0 \\ 0 & B - C^* A^{-1} C \end{bmatrix} = \begin{bmatrix} I & -AC^* \\ 0 & I \end{bmatrix}^* \begin{bmatrix} A & C \\ C^* & B \end{bmatrix} \begin{bmatrix} I & -AC^* \\ 0 & I \end{bmatrix}.$$

and from the fact that the conjugating matrix is a product of elementary matrices. It follows from this, that the Iterated Integration Lemma A.2 is valid on $\mathcal{A}^{\text{gp},0}(\star_X, \Lambda_{\text{loc}})$.

A.3. Integration by parts. Fix labeling sets X and $X' \subset X$. Recall from Section 5.1 the notion of X' -substantial diagrams, that is diagrams that do not contain a strut labeled from $X' \cup \partial X'$. The *divergence* of an X' -substantial diagram (extended by linearity to sums of diagrams) with respect to $X' \subset X$ is defined to be the sum of diagrams obtained by pairing *all* legs labeled from $\partial X'$ with *some* legs labeled from X' .

Lemma A.4 (Integration by Parts). *Let s be an X' -substantial diagram and $t \in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}})$. Then*

$$\begin{aligned}
s \flat_{X'} t &\in \text{Int}_{X'} \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \\
\int^{\text{rat}} dX'(s \flat_{X'} t) &= \int^{\text{rat}} dX'(\text{div}_{X'}(s) \flat_{X - X'} t).
\end{aligned}$$

Proof. For the first statement, assume, for convenience, that $X' = \{f_i\}$ and $X - X' = \{g_j\}$, and rename X' and $X - X'$ by F and G respectively. Applying Lemma A.1.(9), introducing a labeling set \tilde{F} (resp. \tilde{G}), bijective with F (resp. G) (and associated sets $\partial_{\tilde{F}}$ and $\partial_{\tilde{G}}$), gives the expression in question as:

$$s(f, g, \partial f, \partial g) \flat_{F \cup G} t(f, g) = \left\langle s(f, g, \partial \tilde{f}, \partial \tilde{g}), t(f + \tilde{f}, g + \tilde{g}) \right\rangle_{\tilde{F} \cup \tilde{G}}$$

If the canonical decomposition of t with respect to F is

$$t(f, g) = \exp \left(\frac{1}{2} \uparrow_{f_i}^{f_j} M_{ij} \right) \sqcup T(f, g),$$

then the previous expression is equal to

$$\exp \left(\frac{1}{2} \uparrow_{f_i}^{f_j} M_{ij} \right) \sqcup \left\langle s(f, g, \partial \tilde{f}, \partial \tilde{g}), \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i} M_{ij}} + \frac{1}{2} \uparrow_{\tilde{f}_i}^{\tilde{f}_j} M_{ij} \right) \sqcup T(f + \tilde{f}, g + \tilde{g}) \right\rangle_{\tilde{F} \cup \tilde{G}}.$$

Now, because s is an X' -substantial operator, this gives the required decomposition of $s \flat_{F \cup G} t$ with respect to F .

For the second statement, we compute

$$\begin{aligned} & \int^{\text{rat}} dF(s \flat_{F \cup G} t) \\ &= \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial f_i}^{\partial f_j} M_{ij}^{-1} \right), \left\langle s(f, g, \partial \tilde{f}, \partial \tilde{g}), \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i} M_{ij}} + \frac{1}{2} \uparrow_{\tilde{f}_i}^{\tilde{f}_j} M_{ij} \right) \sqcup T(f + \tilde{f}, g + \tilde{g}) \right\rangle_{\tilde{F} \cup \tilde{G}} \right\rangle_F. \end{aligned}$$

To proceed, we must adjust this calculation so that it can distinguish struts, appearing in the left of the pairing, according to which factors, appearing in the right of the pairing, they are glued to. This adjustment is made with Lemma A.1.(7), and leads to the following (introducing sets F^a , F^b and F^c , bijective copies of F):

$$\begin{aligned} & \left\langle \exp \left(-\frac{1}{2} \uparrow_{\partial f_i^a}^{\partial f_j^a} M_{ij}^{-1} \quad -\frac{1}{2} \uparrow_{\partial f_i^b}^{\partial f_j^b} M_{ij}^{-1} \quad -\frac{1}{2} \uparrow_{\partial f_i^c}^{\partial f_j^c} M_{ij}^{-1} \quad -\uparrow_{\partial f_i^b}^{\partial f_j^a} M_{ij}^{-1} \quad -\uparrow_{\partial f_i^c}^{\partial f_j^a} M_{ij}^{-1} \quad -\uparrow_{\partial f_i^c}^{\partial f_j^b} M_{ij}^{-1} \right), \right. \\ & \left. \left\langle s(f^a, g, \partial \tilde{f}, \partial \tilde{g}), \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i^b} M_{ij}} + \frac{1}{2} \uparrow_{\tilde{f}_i}^{\tilde{f}_j} M_{ij} \right) \sqcup T(f^c + \tilde{f}, g + \tilde{g}) \right\rangle_{\tilde{F} \cup \tilde{G}} \right\rangle_{F^a \cup F^b \cup F^c}. \end{aligned}$$

The calculation proceeds by removing factors appearing in left hand side of the pairing, at the expense of operating with them on the right hand side, follow Lemma A.1.(8).

The first term to be treated in this way will be $\exp \left(-\frac{1}{2} \uparrow_{\partial f_i^b}^{\partial f_j^b} M_{ij}^{-1} \right)$, giving (following an application of Lemma A.1.(10)):

$$\exp \left(-\frac{1}{2} \uparrow_{\partial f_i^b}^{\partial f_j^b} M_{ij}^{-1} \right) \flat_{F^b} \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i^b} M_{ij}} \right) = \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i^b} M_{ij}} \right) \sqcup \exp \left(-\frac{1}{2} \uparrow_{\tilde{f}_i}^{\tilde{f}_j} M_{ij} \right).$$

The next term to be so treated will be $\exp \left(-\uparrow_{\partial f_i^c}^{\partial f_j^a} M_{ij}^{-1} \right)$, which gives:

$$\exp \left(-\uparrow_{\partial f_i^c}^{\partial f_j^a} M_{ij}^{-1} \right) \flat_{F^b \cup F^c} \left(\exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i^b} M_{ij}} \right) \sqcup T(f^c + \tilde{f}, g + \tilde{g}) \right) = \exp \left(\frac{\tilde{f}_j}{\uparrow_{f_i^b} M_{ij}} \right) \sqcup T((f^c - \tilde{f}) + \tilde{f}, g + \tilde{g}).$$

The result of these operations is

$$\left\langle \exp \left(-\frac{1}{2} \begin{array}{c} \partial f_j^a \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^a \end{array} - \frac{1}{2} \begin{array}{c} \partial f_j^c \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^c \end{array} - \begin{array}{c} \partial f_j^a \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^b \end{array} - \begin{array}{c} \partial f_j^a \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^c \end{array} \right), \\ \left\langle s(f^a, g, \partial \tilde{f}, \partial \tilde{g}), \exp \left(\begin{array}{c} \tilde{f}_j \\ \uparrow \\ M_{ij} \\ f_i^b \end{array} \right) \sqcup T(f^c, g + \tilde{g}) \right\rangle_{\tilde{F} \cup \tilde{G}} \Big|_{F^a \cup F^b \cup F^c} \right\rangle.$$

Performing the pairing with respect to the variables \tilde{F} , leads to the replacement of the the inner $\langle \cdot, \cdot \rangle$ with

$$\langle s(f^a, g, f^b M, \partial \tilde{g}), T(f^c, g + \tilde{g}) \rangle_{\tilde{G}}.$$

Trading the factor $\exp \left(-\begin{array}{c} \partial f_j^b \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^a \end{array} \right)$, gives:

$$\exp \left(-\begin{array}{c} \partial f_j^b \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^a \end{array} \right) \Big|_{F^a \cup F^b} s(f^a, g, f^b M, \partial \tilde{g}) = \operatorname{div}_{F^a} (s(f^a, g, f^b M + \partial f^a, \partial g)) \Big|_{\partial g \rightarrow \partial \tilde{g}}.$$

Performing the pairing with respect to F^b sets f^b to zero, and thus we arrive at the following expression.

$$\left\langle \exp \left(-\frac{1}{2} \begin{array}{c} \partial f_j^a \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^a \end{array} - \frac{1}{2} \begin{array}{c} \partial f_j^c \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^c \end{array} - \begin{array}{c} \partial f_j^a \\ \uparrow \\ M_{ij}^{-1} \\ \partial f_i^c \end{array} \right), \left\langle \operatorname{div}_F (s(f^a, g, \partial f^a, \partial g)) \Big|_{\partial g \rightarrow \partial \tilde{g}}, T(f^c, g + \tilde{g}) \right\rangle_{\tilde{G}} \right\rangle_{F^a \cup F^c}.$$

The proof follows by inspection. \square

APPENDIX B. THE WHEELS IDENTITY

The goal of this section is to show the following *Wheels Identity*:

Theorem B.1. *For every $M \in \operatorname{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$, we have that*

$$\int dX \left(\exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (e^{h_1}, \dots, e^{h_g}) \right) \right) = \exp_{\square} \left(-\frac{1}{2} \begin{array}{c} \circlearrowleft \\ \bullet \\ \chi(M) \end{array} \right),$$

where we think of $\chi(M)$ as taking values in $\hat{\Lambda}/(\text{cyclic})$, which is identified via the map $t_i \rightarrow e^{h_i}$ with the set $\mathbb{Q}\langle\langle h_1, \dots, h_g \rangle\rangle/(\text{cyclic})$ of formal power series in noncommuting variables modulo cyclic permutations.

Proof. Choose generators $\{t_1, \dots, t_g\}$ for the free group F and identify $\hat{\Lambda}$ with the ring of noncommutative power series $\{h_1, \dots, h_n\}$. Recall the map $\Lambda \rightarrow \hat{\Lambda}$ given by $t_i \rightarrow e^{h_i}$.

The Aarhus integral is calculated by splitting the quadratic part from the rest of the integrand as follows:

$$\begin{aligned} & \int dX \left(\exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (e^{h_1}, \dots, e^{h_g}) \right) \right) \\ &= \int dX \left(\exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (1, \dots, 1) \right) \sqcup \exp \left(\frac{1}{2} \sum_{x_i} \left(\begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (e^{h_1}, \dots, e^{h_g}) - \begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (1, \dots, 1) \right) \right) \right) \\ &= \left\langle \exp \left(-\frac{1}{2} \sum_{\partial x_i} \begin{array}{c} \partial x_j \\ \uparrow \\ M_{ij} \\ \uparrow \end{array} (1, \dots, 1)^{-1} \right), \exp \left(\frac{1}{2} \sum_{x_i} \left(\begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (e^{h_1}, \dots, e^{h_g}) - \begin{array}{c} \uparrow \\ M_{ij} \\ \uparrow \end{array} (1, \dots, 1) \right) \right) \right\rangle_X. \end{aligned}$$

With the following notation for ‘‘Red’’ and ‘‘Green’’ struts,

$$\begin{aligned}\uparrow R &= \sum_{\frac{\partial x_j}{\partial x_i}} \uparrow M_{ij}(1, \dots, 1)^{-1}, \\ \uparrow G &= \sum_{x_i} \left(\uparrow M_{ij}(e^{h_1}, \dots, e^{h_g}) - \uparrow M_{ij}(1, \dots, 1) \right).\end{aligned}$$

it follows that the desired integral equals to the following formal series (which makes sense, since finitely many terms appear in each degree)

$$\left\langle \exp \left(-\frac{1}{2} \uparrow R \right), \exp \left(\frac{1}{2} \uparrow G \right) \right\rangle = \sum_{n=0}^{\infty} (-1)^n \left(\frac{1}{2} \right)^n \left(\frac{1}{2} \right)^n \frac{1}{n!} \frac{1}{n!} \left\langle \left(\uparrow R \right)^n, \left(\uparrow G \right)^n \right\rangle.$$

Specifically, the answer is recovered if each ‘‘join’’ is accompanied by a Kronecker delta. For example:

$$\begin{aligned}\uparrow R \\ \uparrow G\end{aligned} = \sum_{i,j,k} \begin{aligned} \uparrow \\ \uparrow \\ \uparrow \end{aligned} \begin{aligned} M_{jk}(1, \dots, 1)^{-1} \\ M_{ij}(e^{h_1}, \dots, e^{h_g}) \\ M_{ij}(1, \dots, 1) \end{aligned} - \sum_{i,j,k} \begin{aligned} \uparrow \\ \uparrow \\ \uparrow \end{aligned} \begin{aligned} M_{jk}(1, \dots, 1)^{-1} \\ M_{ij}(1, \dots, 1) \\ M_{ij}(1, \dots, 1) \end{aligned}.$$

We focus on a particular summand. It will help in the management of the combinatorics if we label each of the arcs that appear in the struts:

$$\left\langle \left(\uparrow R \right)^n, \left(\uparrow G \right)^n \right\rangle \rightarrow \left\langle \begin{matrix} r_{1,1} & r_{n,1} & g_{1,1} & g_{n,1} \\ \uparrow & \dots & \uparrow & \dots & \uparrow \\ r_{1,2} & r_{n,2} & g_{1,2} & g_{n,2} \end{matrix} \right\rangle$$

Now we introduce some definitions to assist in the enumeration of the polygons that arise from this pairing.

- An *RG-shape of degree n* is a collection of even-sided polygons whose edges are alternately colored red and green. The total number of edges is $2n$.
- A *labeled RG-shape of degree n* is an RG-shape with an extra bivalent vertex in each edge, such that:
 - (1) the resulting set of red (resp. green) edges is equipped with a bijection to $\{r_{1,1}, r_{1,2}, \dots, r_{n,1}, r_{n,2}\}$ (resp. $\{g_{1,1}, g_{1,2}, \dots, g_{n,1}, g_{n,2}\}$),
 - (2) if an edge is labeled, for example $r_{i,1}$ (resp. $g_{i,1}$), then its neighboring red (resp. green) edge is labeled $r_{i,2}$ (resp. $g_{i,2}$).
- Let the constant C_τ , for τ some RG-shape, denote the number of labeled RG-shapes which recover τ when their labels and extra bivalent vertices are forgotten.

Observe that:

$$\left\langle \left(\uparrow R \right)^n, \left(\uparrow G \right)^n \right\rangle = \sum_{\tau \text{ a labeled RG-shape of deg } n} \tau = \sum_{\tau \text{ an RG-shape of deg } n} C_\tau \tau.$$

To calculate this constant describe an RG-shape of degree n , τ , by a sequence (i_1, \dots, i_n) where i_j counts the number of polygons with $2j$ edges.

Lemma B.2.

$$C_\tau = \frac{n! 2^n n! 2^n}{((2)(1))^{i_1} i_1! ((2)(2))^{i_2} i_2! \dots ((2)(n))^{i_n} i_n!}.$$

Proof. Fix a copy of τ with its set of edges ordered (any old how, so as to distinguish them from each other). There are precisely $2^n n! 2^n n!$ ways of labeling this so as to respect the conditions of the above definition.

But this overcounts by a multiplicative factor of the order of the symmetry group of τ (that is, with the ordering of the edges forgotten). That symmetry group is precisely the following group (that is, swaps of identical polygons, and rotations and flips of individual polygons):

$$(\Sigma_{i_1} \times (D_1 \times \dots \times D_1)) \times \dots \times (\Sigma_{i_n} \times (D_n \times \dots \times D_n)),$$

where D_n represents the n th dihedral group.

Thus we conclude:

$$C_\tau = \frac{2^n n! 2^{2n} n!}{\#\{(\Sigma_{i_1} \times (D_1 \times \cdots \times D_1)) \times \cdots \times (\Sigma_{i_n} \times (D_n \times \cdots \times D_n))\}}$$

□

If τ_i denotes a connected polygon with $2i$ edges, it follows that

$$\begin{aligned} \left\langle \exp\left(-\frac{1}{2} \uparrow R\right), \exp\left(\frac{1}{2} \uparrow G\right) \right\rangle &= \sum_{n=0}^{\infty} (-1)^n \left(\frac{1}{2}\right)^n \left(\frac{1}{2}\right)^n \frac{1}{n!} \frac{1}{n!} \left\langle \left(\uparrow R\right)^n, \left(\uparrow G\right)^n \right\rangle, \\ &= \sum_n \sum_{i_1, \dots, i_n} \left(\prod_{p=1}^n \left(\frac{(-1)^p}{(2p)} \right)^{i_p} \frac{1}{i_p!} \right) \tau_1^{i_1} \sqcup \cdots \sqcup \tau_n^{i_n}, \\ &= \exp\left(\frac{1}{2} \sum_{p=1}^{\infty} \frac{(-1)^p}{p} \tau_p\right). \end{aligned}$$

Substituting the definitions of the ‘‘Red’’ and ‘‘Green’’ struts into τ_p yields

$$\tau_p = \text{tr} \left((M(1, \dots, 1))^{-1} (M(e^{h_1}, \dots, e^{h_g}) - M(1, \dots, 1))^p \right) = \text{tr} \left((M(1, \dots, 1))^{-1} M(e^{h_1}, \dots, e^{h_g}) - I \right)^p.$$

from which the result follows. □

APPENDIX C. INFINITESIMAL VERSUS GROUP-LIKE WRAPPING RELATIONS

In this section we show that the infinitesimal wrapping relations ω imply the group-like wrapping relation ω^{SP} . In order to achieve this, it will be useful to introduce the variant ω'_i of ω_i for $i = 1, \dots, g$ where we allow the diagram $D \in \mathcal{A}(\star_{X \cup \{\partial h\}}, \Lambda_{\text{loc}})$ to have *several* legs labeled by ∂h . In other words, ω'_i is generated by the move $(M, D_1) \sim (M, D_2)$ where

$$D_m = \text{con}_{\{h\}} \left(\varphi_{m,i}(D) \sqcup \left(\exp_{\sqcup} \left(-\frac{1}{2} \chi_h (M^{-1} \varphi_{m,i} M) \right) \right) \right)$$

for $m = 1, 2$ and $\varphi_{1,i}$ is the identity and $\varphi_{2,i}$ (abbreviated by φ_i below) denotes the substitution $\phi_{t_i \rightarrow e^h t_i e^{-h}}$ on beads.

It is obvious that the ω'_i relation includes the ω_i relation. Our goal for the rest of the section is to show that they coincide (see Proposition C.7) and that the ω^{SP} -relations are implied by the ω -relations (see Proposition 3.12).

Recall the maps $\eta_i : \Lambda_{\text{loc}} \rightarrow \widehat{\mathcal{L}}_{\text{loc}}$ of Section 3.1 that keep track of the h -degree 1 part of the substitutions $\varphi_{t_i \rightarrow e^{-h} t_i e^h}(a)$ of $a \in \Lambda_{\text{loc}}$. We will introduce an extension $\widehat{\eta}_i$ of η_i which will allow us to keep track of the h -degree n part of $\varphi_{t_i \rightarrow e^{-h} t_i e^h}(a)$, for all n . The following lemma (which the reader should compare with Lemma 3.2) summarizes the properties of $\widehat{\eta}_i$.

Lemma C.1. *For every $i = 1, \dots, g$, there exists a unique*

$$\widehat{\eta}_i : \widehat{\mathcal{L}}_{\text{loc}} \rightarrow \widehat{\mathcal{L}}_{\text{loc}}$$

that satisfies the following properties:

$$\begin{aligned} \widehat{\eta}_i(t_j) &= \delta_{ij}[t_i, h] = \delta_{ij}(t_i h - h t_i) \\ \widehat{\eta}_i(h) &= 0 \\ \widehat{\eta}_i(ab) &= \widehat{\eta}_i(a)b + a\widehat{\eta}_i(b) \\ \widehat{\eta}_i(a+b) &= \widehat{\eta}_i(a) + \widehat{\eta}_i(b). \end{aligned}$$

In particular, the restriction of $\widehat{\eta}_i$ to $\Lambda_{\text{loc}} \subset \widehat{\mathcal{L}}_{\text{loc}}$ equals to η_i .

Proof. First, we need to show the existence of such a $\widehat{\eta}_i$. A coordinate approach is to define $\widehat{\eta}_i : \mathcal{R} \rightarrow \mathcal{R}$ on the ring \mathcal{R} of Section 3.1 which is isomorphic to $\widehat{\mathcal{L}}_{\text{loc}}$. Given the description of \mathcal{R} , it is easy to see that $\widehat{\eta}_i$ exists and is uniquely determined by the above properties.

An alternative, coordinate-free definition of $\widehat{\eta}_i$ is the following. Let $\widehat{\mathcal{L}\mathcal{L}_{\text{loc}}}$ denote the Cohn localization of the group-ring $\mathbb{Q}[F \star \mathbb{Z} \star \mathbb{Z}]$, followed by a completion with respect to $t - 1$ and $t' - 1$ (where t and t' are generators of $\mathbb{Z} \star \mathbb{Z}$). Let $h = \log t$ and $h' = \log t'$. The map $\varphi_{t_i \rightarrow e^{-h} t_i e^h} : \mathbb{Q}[F \star \mathbb{Z} \star \mathbb{Z}] \rightarrow \widehat{\mathcal{L}\mathcal{L}_{\text{loc}}}$ is Σ -inverting, thus it extends to a map of the localization, and further to a map $\widehat{\mathcal{L}\mathcal{L}_{\text{loc}}} \rightarrow \widehat{\mathcal{L}\mathcal{L}_{\text{loc}}}$ of the completion. $\widehat{\mathcal{L}\mathcal{L}_{\text{loc}}}$ is a h' -graded ring, and we can define

$$\widehat{\eta}_i : \widehat{\mathcal{L}_{\text{loc}}} \rightarrow \widehat{\mathcal{L}_{\text{loc}}}$$

by

$$\widehat{\eta}_i = \varphi_{h' \rightarrow h} \circ \text{deg}_{h'}^1 \circ \varphi_{t_i \rightarrow e^{-h} t_i e^h}.$$

It is easy to see that this definition of $\widehat{\eta}_i$ satisfies the properties of the lemma, and that the properties characterize $\widehat{\eta}_i$.

Finally, Lemma 3.2 implies that the restriction of $\widehat{\eta}_i$ on $\Lambda_{\text{loc}} \subset \widehat{\mathcal{L}_{\text{loc}}}$ equals to η_i . \square

Lemma C.2. *For all n and i we have that*

$$\frac{1}{n!} \widehat{\eta}_i^n = \text{deg}_h^n \circ \varphi_{t_i \rightarrow e^{-h} t_i e^h}.$$

considered as maps $\Lambda \rightarrow \widehat{\mathcal{L}_{\text{loc}}}$.

Proof. In view of Lemma 3.2, it suffices to show that the maps agree on generators t_j of F_g . For $j \neq i$, both maps vanish. For $j = i$, the following identity

$$e^{-h} t_i e^h = \exp(-\text{ad}_h)(t_i) = \sum_{n=0}^{\infty} \frac{1}{n!} [[t_i, h], h], \dots, h]$$

and Lemma 3.2 show that both maps agree. The result follows. \square

Next, we extend the above lemma for matrices.

Lemma C.3. *For all n and sites i we have that*

$$\frac{1}{n!} \widehat{\eta}_i^n = \text{deg}_h^n \circ \varphi_{t_i \rightarrow e^{-h} t_i e^h}.$$

considered as maps $\text{Mat}(\Lambda \rightarrow \mathbb{Z}) \rightarrow \text{Mat}(\widehat{\mathcal{L}_{\text{loc}}} \rightarrow \mathbb{Z})$.

Proof. Observe the identity

$$(15) \quad \widehat{\eta}_i(AB) = \widehat{\eta}_i(A)B + A\widehat{\eta}_i(B)$$

for matrices A, B (not necessarily square) that can be multiplied. It implies that

$$0 = \frac{1}{n!} \widehat{\eta}_i^n(I) = \frac{1}{n!} \widehat{\eta}_i^n(MM^{-1}) = \sum_{k=0}^n \frac{1}{k!} \widehat{\eta}_i^k(M) \widehat{\eta}_i^{n-k}(M^{-1}).$$

The lemma follows by induction on n and the above identity which solves $\widehat{\eta}_i^n(M)$ in terms of quantities known by induction. \square

Proposition C.4. *For all n and sites i we have that*

$$\frac{1}{n!} \widehat{\eta}_i^n = \text{deg}_h^n \circ \varphi_{t_i \rightarrow e^{-h} t_i e^h}.$$

considered as maps $\Lambda_{\text{loc}} \rightarrow \widehat{\mathcal{L}_{\text{loc}}}$.

Proof. Consider a matrix presentation of $s \in \Lambda_{\text{loc}}$. I.e.,

$$s = (1, 0, \dots, 0)M^{-1}\vec{b}$$

for $M \in \text{Mat}(\Lambda \rightarrow \mathbb{Z})$ and \vec{b} a column vector over Λ . Equation (15) implies that

$$\frac{1}{n!} \widehat{\eta}_i^n(s) = \sum_{k=0}^n \frac{1}{k!} \widehat{\eta}_i^k(M^{-1}) \widehat{\eta}_i^{n-k}(\vec{b}).$$

The proposition now follows from Lemmas C.2 and C.3. \square

Proposition C.5. For all diagrams $D \in \mathcal{A}(\star_{X,\partial h}, \Lambda_{\text{loc}})$, $M \in \text{Herm}(\Lambda_{\text{loc}} \rightarrow \mathbb{Z})$, and $i = 1, \dots, g$, we have that

$$\frac{1}{n!} \widehat{\eta}_i^n(D) = \text{deg}_h^n \left(\varphi_{t_i \rightarrow e^{-h} t_i e^h}(D) \sqcup \exp_{\sqcup} \left(-\frac{1}{2} \chi_h(M^{-1} \varphi_{t_i \rightarrow e^{-h} t_i e^h}(M)) \right) \right).$$

Proof. The proof follows from a careful combinatorial counting similar to that of the Wheels Identity of Theorem B.1. Let us precede the proof with an informal examination of what takes place when we repeatedly apply $\widehat{\eta}_i$ to the empty diagram, $D = \phi$. The first application results in a term:

$$\frac{1}{2} \sum_k \circlearrowleft (M^{-1}(-\widehat{\eta}_i(M)))_{kk}$$

The second application requires the result of applying $\widehat{\eta}_i$ to M^{-1} . This can be done using the following special case of Equation (15):

$$(16) \quad \widehat{\eta}_i(M^{-1}) = M^{-1}(-\widehat{\eta}_i(M))M^{-1}.$$

So, the second application results in the following terms:

$$\begin{aligned} & \frac{1}{2} \sum_{k,l} \left(\begin{array}{c} \bullet \\ \circlearrowleft \\ \bullet \end{array} \begin{array}{c} (M^{-1}(-\widehat{\eta}_i(M)))_{lk} \\ (M^{-1}(-\widehat{\eta}_i(M)))_{kl} \end{array} \right) + \frac{1}{2} \sum_k \circlearrowleft \left(M^{-1} \frac{(-\widehat{\eta}_i)^2}{2!}(M) \right)_{kk} \\ & + \frac{(\frac{1}{2})^2}{2!} \sum_{k,l} \circlearrowleft \left(M^{-1}(-\widehat{\eta}_i(M)) \right)_{kk} \sqcup \circlearrowleft \left(M^{-1}(-\widehat{\eta}_i(M)) \right)_{ll} \end{aligned}$$

We see that the terms arising from the application of $\widehat{\eta}_i$ fall into three categories.

$$(17) \quad \widehat{\eta}_i(\phi) = \frac{1}{2} \sum_k \circlearrowleft (M^{-1}(-\widehat{\eta}_i(M)))_{kk}$$

$$(18) \quad \begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} (\widehat{\eta}_i(M^{-1}))_{kl} = \begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} \begin{array}{c} (M^{-1})_{ml} \\ (M^{-1}(-\widehat{\eta}_i(M)))_{km} \end{array}$$

$$(19) \quad \begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} (\widehat{\eta}_i((\widehat{\eta}_i)^p(M)))_{kl} = \begin{array}{c} \bullet \\ \downarrow \\ \bullet \end{array} ((\widehat{\eta}_i)^{p+1}(M))_{kl}$$

Thus, we see that $\widehat{\eta}_i^n(\phi)$ equals to a linear combination of diagrams, each consisting of a number of wheel components, each of which is of the following form (this can be (over-)determined by a sequence $(\kappa_1, \dots, \kappa_\mu)$ of positive integers):

$$\sum_{k_1, \dots, k_\mu} \left(\begin{array}{c} \bullet \\ \circlearrowleft \\ \bullet \end{array} \begin{array}{c} (M^{-1}(-\widehat{\eta}_i)^{\kappa_\mu}(M))_{k_{\mu-1}k_\mu} \\ \dots \\ (M^{-1}(-\widehat{\eta}_i)^{\kappa_1}(M))_{k_\mu k_1} \end{array} \right)$$

The question, then, is to calculate an expression in terms of such diagrams, counting “all ways” that they arise from the application of moves (17)-(19). We will generate a sequence of such moves with the following object, an *edge-list* on a *shape*.

The *shape* determined by a set of positive integers (μ_1, \dots, μ_τ) is a set of oriented polygons, one with μ_1 edges, and so on, such that each has a distinguished bivalent vertex, its *basepoint*. A *labeled shape* is a shape with its edges labeled by positive integers. The diagram D_κ determined by some labeled shape κ is the diagram that arises by replacing each polygon by a series of wheels, in the following manner (to distinguish them from beads, the bivalent vertices of the polygons will be drawn by small horizontal segments, and the base-point by an x).

$$\left(\begin{array}{c} \kappa_\mu \\ \dots \\ \kappa_1 \end{array} \right) \times \rightarrow \sum_{k_1, \dots, k_\mu} \left(\begin{array}{c} \bullet \\ \circlearrowleft \\ \bullet \end{array} \begin{array}{c} (M^{-1}(-\widehat{\eta}_i)^{\kappa_\mu}(M))_{k_{\mu-1}k_\mu} \dots \\ (M^{-1}(-\widehat{\eta}_i)^{\kappa_1}(M))_{k_\mu k_1} \end{array} \right)$$

An *edge-list* on a shape is a finite sequence of the edges of the shape such that each edge is represented at least once. (It is convenient to enumerate the edges along the orientation, starting from the basepoint.)

Proof. Consider two pairs (M, s_1) and (M, s_2) related by a wrapping move ω'_i as in definition 3.7. In other words, for some $s \in \mathcal{A}^{\text{gp}}(\star_X, \partial h, \Lambda_{\text{loc}})$ we have that

$$s_m = \text{con}_{\{h\}} \left(\varphi_{m,i}(s) \sqcup \left(\exp_{\sqcup} \left(-\frac{1}{2} \chi_h(M^{-1} \varphi_{m,i} M) \right) \right) \right)$$

for $m = 1, 2$ where $\varphi_{1,i}$ is the identity and $\varphi_{2,i} = \phi_{t_i \rightarrow e^{-h} t_i e^h}$. Consider the case that s has a diagram with n legs labeled by ∂h . Proposition C.5 implies that

$$s_2 = \frac{1}{n!} \text{con}_{\{h\}} (\widehat{\eta}_i^n(s)).$$

Let s' denote the result of gluing all but one of the n ∂h legs of $\widehat{\eta}_i^{n-1}(s)$ and renaming the resulting one $\partial h'$. In other words, $s' = \text{con}_{\{h\}} (\widehat{\eta}_i^{n-1}(s)(\partial h + \partial h'))$. Let $\widehat{\eta}'_i$ denote the action of $\widehat{\eta}_i$ for $\partial h'$ -labeled legs. If s has at least one ∂h leg, it follows that

$$s_2 = \text{con}_{\{h'\}} (\widehat{\eta}'_i(s')) = 0 \text{ mod } \omega_i$$

by the infinitesimal wrapping relations, and it follows that $s_1 = 0$ by definition. If s has no ∂h leg, then $s_1 = s_2$. Thus, in all cases, $(M, s_1) \stackrel{\omega_i}{\sim} (M, s_2)$. The converse is obvious; thus the result follows. \square

Proof. (of Proposition 3.12) It is obvious that the ω' relation implies the group-like ω^{gp} relation of Section 5.4. Since $\omega' = \omega$, the result follows. \square


APPENDIX D. INFINITESIMAL BASING RELATIONS

This section, although it is not used anywhere in the paper, is added for completeness. The reader may have noticed that in Section 3, we mentioned infinitesimal versions of the wrapping and the String relations, but not of the basing relations. The reason is that we did not need them; in addition \int^{rat} is not known to preserve them. Nevertheless, we introduce them here.

We define

$$\mathcal{A}(\otimes_X, \Lambda_{\text{loc}}) = \mathcal{A}(\star_X, \Lambda_{\text{loc}}) / \langle \beta_1, \beta_2 \rangle,$$

where $\beta_{1,X}$ is the subspace of X -flavored link relations that appeared in [A, Part II, Section 5.2]:



and $\beta_{2,X}$ denote subspace generated by an action

$$F \times \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_X, \Lambda_{\text{loc}})$$

described by pushing some element of $f \in F_g$ onto all X -labeled legs of a diagram.

We now compare the infinitesimal and group-like basing relations as follows.

Proposition D.1. (a) *We have an isomorphism*

$$\mathcal{A}(\dots, \otimes_X, \Lambda_{\text{loc}}) = \mathcal{A}(\dots, \star_X, \Lambda_{\text{loc}}) / \langle \beta_X \rangle \xrightarrow{\cong} \mathcal{A}(\dots, \cup_X, \Lambda_{\text{loc}}).$$

(b) *The inclusion $\mathcal{A}^{\text{gp},0}(\dots, \star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^0(\dots, \star_X, \Lambda_{\text{loc}})$ maps the β^{gp} -relations to β -relations, and induces a map*

$$\mathcal{A}^{\text{gp},0}(\dots, \otimes_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^0(\dots, \star_X, \Lambda_{\text{loc}}).$$

Proof. The first part follows from the proof of [A, Part II, Theorem 3] without essential changes. The second part is standard. \square

APPENDIX E. A USEFUL CALCULATION

In this section, we quote a useful and often appearing computation that uses the identities of Lemma A.1.

Let $\varphi : \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}^{\text{gp}}(\star_{X \cup \{h\}}, \Lambda_{\text{loc}})$ be a map that commutes with respect to gluing X -colored legs.

Theorem E.1. *For $X' \subset X$ and $s \in \text{Int}_{X'} \mathcal{A}^{\text{gp}}(\otimes_X, \Lambda_{\text{loc}})$, decomposed as in Equation (2), we have that*

$$\int^{\text{rat}} dX'(\varphi(s)) = \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} \right) \right) \varphi \left(\int^{\text{rat}} dX'(s) \right).$$

If in addition $\varphi(s) - s$ is X -substantial for all $s \in \mathcal{A}^{\text{gp}}(\star_X, \Lambda_{\text{loc}})$, then

$$\int^{\text{rat}} dX'(\varphi(s)) = \exp \left(-\frac{1}{2} \chi_h(M^{-1} \varphi M) \right) \varphi \left(\int^{\text{rat}} dX'(s) \right).$$

Proof. Consider the decomposition of s given by Equation (2). We begin with a “ δ -function tryck” of Equation (12) of A.1:

$$s = \exp \left(\frac{1}{2} \sum_{i,j} \begin{matrix} x_j \\ \uparrow \\ x_i \end{matrix} M_{ij} \right) \sqcup R = \left\langle R(\partial y), \exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} M_{ij} + \begin{matrix} y_j \\ \uparrow \\ x_i \end{matrix} \right) \right\rangle_Y.$$

Thus,

$$\int^{\text{rat}} dX'(\varphi(s)) = \left\langle \varphi(R(\partial y)), \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} + \begin{matrix} y_j \\ \uparrow \\ x_i \end{matrix} \right) \right) \right\rangle_Y.$$

Since $\text{div}_{X'} \exp \left(-\sum_{\substack{\partial x_j \\ \uparrow \\ y_i}} \varphi M_{ij}^{-1} \right) = 1$, the integration by parts Lemma A.4 implies that

$$\begin{aligned} \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} + \begin{matrix} y_j \\ \uparrow \\ x_i \end{matrix} \right) \right) = \\ \int^{\text{rat}} dX' \left(\left(\exp \left(-\sum_{\substack{\partial x_j \\ \uparrow \\ y_i}} \varphi M_{ij}^{-1} \right) \right) \flat_{X'} \exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} + \begin{matrix} y_j \\ \uparrow \\ x_i \end{matrix} \right) \right). \end{aligned}$$

Equation (13) of Lemma A.1 implies that the above equals to:

$$\exp \left(-\frac{1}{2} \sum_{\substack{y_j \\ \uparrow \\ y_i}} \varphi M_{ij}^{-1} \right) \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} \right) \right).$$

Returning to the expression in question:

$$\int^{\text{rat}} dX'(\varphi(s)) = \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{\substack{x_j \\ \uparrow \\ x_i}} \varphi M_{ij} \right) \right) \sqcup \left\langle \varphi(R(\partial y)), \exp \left(-\frac{1}{2} \sum_{\substack{y_j \\ \uparrow \\ y_i}} \varphi M_{ij}^{-1} \right) \right\rangle_Y.$$

The second factor equals to $\varphi \left(\int^{\text{rat}} dX'(s) \right)$, and concludes the first part of the theorem.

In order to compute the first factor, we need to separate its X' -substantial part. With the added assumption on φ , we can compute the first factor in a manner analogous to the Wheels Identity of Theorem B.1

$$\begin{aligned}
\int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \uparrow \varphi M_{ij} \right) \right) &= \int^{\text{rat}} dX' \left(\exp \left(\frac{1}{2} \sum \left(\uparrow_{x_i}^{x_j} M_{ij} + \uparrow_{x_i}^{x_j} \varphi M_{ij} - \uparrow_{x_i}^{x_j} M_{ij} \right) \right) \right) \\
&= \left\langle \exp \left(-\frac{1}{2} \sum_{\partial x_i}^{\partial x_j} \uparrow M_{ij}^{-1} \right), \exp \left(\frac{1}{2} \sum_{x_i}^{x_j} \uparrow \varphi M_{ij} - \uparrow_{x_i}^{x_j} M_{ij} \right) \right\rangle_{X'} \\
&= \exp \left(-\frac{1}{2} \chi_h(M^{-1} \varphi M) \right).
\end{aligned}$$

□

Remark E.2. The proof of Theorem E.1 can be applied without changes to the case of the map $\varphi = \text{Hair}_H : \mathcal{A}(\star_X, \Lambda_{\text{loc}}) \rightarrow \mathcal{A}(\star_{X \cup H})$ in order to show that for $s \in \text{Int}_{X'} \mathcal{A}^{\text{SP}}(\star_X, \Lambda_{\text{loc}})$, we have that

$$\int dX'(\text{Hair}_H(s)) = \exp \left(-\frac{1}{2} \chi_h(M^{-1} \text{Hair}_H M) \right) \text{Hair}_H \left(\int^{\text{rat}} dX'(s) \right).$$

REFERENCES

- [AMR] J. Andersen and J. Mattes and N. Reshetikhin, *The Poisson structure on the moduli space of flat connections and chord diagrams*, *Topology* **35** (1996) 1069–1083.
- [A0] D. Bar-Natan, S. Garoufalidis, L. Rozansky and D. Thurston, *Wheels, wheeling and the Kontsevich integral of the unknot*, *Israel J. Math.* **119** (2000) 217–238.
- [A] D. Bar-Natan, S. Garoufalidis, L. Rozansky and D. Thurston, *The Aarhus integral of rational homology 3-spheres I-II*, *Selecta Math.* in press.
- [BL] D. Bar-Natan and R. Lawrence, *A rational surgery formula for the LMO invariant*, preprint 2000 [math.GT/0007045](#).
- [BLT] D. Bar-Natan, T.T.Q. Le and D. Thurston, *Two applications of elementary knot theory to Lie algebras and Vassiliev invariants*, preprint 2002 [math.GT/0204311](#).
- [BG] D. Bar-Natan, S. Garoufalidis, *On the Melvin-Morton-Rozansky conjecture*, *Inventiones Math.* **125** (1996) 103–133.
- [BR] J. Berstel and C. Reutenauer, *Rational Series and their Languages*, Springer-Verlag 1984.
- [CS1] S. Cappell and J. Shaneson, *The codimension two placement problem and homology equivalent manifolds*, *Annals Math.* **99** (1974) 277–348.
- [CS2] ———, *Link cobordism*, *Commentarii Math. Helv.* **55** (1980) 20–49.
- [C] P.M. Cohn, *Free rings and their relations*, Academic Press, 1985.
- [Dv] J. Duval, *Forme de Blanchfield et cobordisme d'entrelacs bords*, *Commentarii Math. Helv.* **61** (1986) 617–635.
- [FV] M. Farber and P. Vogel, *The Cohn localization of the free group ring*, *Math. Proc. Cambridge Phil. Soc.* **111** (1992) 432–443.
- [Fa] ———, *Noncommutative rational functions and boundary links*, *Math. Annalen* **293** (1992) 543–568.
- [FRa] ——— and A. Ranicki, *The Morse-Novikov theory of circled-valued functions and noncommutative localization*, *Tr. Math. Inst. Steklova* **225** (1999) 381–388, and also [math.DG/9812122](#).
- [FR] R. Fenn and C. Rourke, *On Kirby's calculus of links*, *Topology* **18** (1979) 1–15.
- [GR] S. Garoufalidis and L. Rozansky, *The loop expansion of the Kontsevich integral, abelian invariants of knots and S-equivalence*, preprint 2000, [math.GT/0003187](#), to appear in *Topology*.
- [GL1] ——— and J. Levine, *Homology surgery and invariants of 3-manifolds*, *Geometry and Topology*, **5** (2001) 551–578.
- [GL2] ——— and ———, *Analytic invariants of boundary links*, preprint 2000, [math.GT/0102103](#), to appear in *J. Knot Theory and its Rami*.
- [GL3] ——— and ———, *Concordance and 1-loop clovers*, *Algebraic and Geometric Topology*, **1** (2001) 687–697.
- [GGP] ———, M. Goussarov and M. Polyak, *Calculus of clovers and finite type invariants of 3-manifolds*, *Geometry and Topology*, **5** (2001) 75–108.
- [Ga] ———, *Boundary links, localization and the loop move*, preprint 2001.
- [Ga2] ———, to appear.
- [GK1] ——— and A. Kricker, *Finite type invariants of cyclic branched covers*, preprint 2001, [math.GT/0107220](#).
- [GK2] ——— and ———, *A surgery view of boundary links*, preprint 2002 [math.GT/0205328](#).
- [GT] ——— and P. Teichner, *On knots with trivial Alexander polynomial*, preprint 2002, [math.GT/02XXXXX](#).
- [Gu] M. Gutierrez, *Boundary links and an unlinking theorem*, *Trans. Amer. Math. Soc.* **171** (1972) 491–499.
- [HM] N. Habegger, G. Masbaum, *The Kontsevich Integral and Milnor's Invariants*, *Topology* **39** (2000) 1253–1289.
- [Ko] M. Kontsevich, *Vassiliev's knot invariants*, *Adv. in Sov. Math.*, **16(2)** (1993), 137–150.
- [K] K.H. Ko, *Seifert matrices and boundary link cobordisms*, *Trans. Amer. Math. Soc.* **299** (1987) 657–681.
- [Kr1] A. Kricker, *The lines of the Kontsevich integral and Rozansky's Rationality Conjecture*, preprint 2000, [math/GT.0005284](#).
- [Kr2] ———, *Covering spaces over claspered knots*, preprint 1999, [math.GT/9901029](#).

- [Le1] J. Levine, *Knot modules I*, Trans. AMS **229** (1977) 1–50.
- [Le2] ———, *Link concordance and algebraic closure, II*, Inventiones Math. **96** (1989) 571–592.
- [LM] T.T.Q. Le, J. Murakami, *Parallel version of the universal Vassiliev-Kontsevich invariant*, J. Pure and Applied Algebra, **121** (1997) 271–291.
- [LMMO] ———, H. Murakami, J. Murakami and T. Ohtsuki, *A three-manifold invariant via the Kontsevich integral*, Osaka J. Math **36** (1999) 365–395.
- [LMO] T.T.Q. Le, J. Murakami, T. Ohtsuki, *A universal quantum invariant of 3-manifolds*, Topology **37** (1998) 539–574.
- [P] B. Patureau-Mirand, *Non-Injectivity of the “hair” map*, preprint 2002, [math.GT/0202065](#).
- [R1] L. Rozansky, *A contribution of a $U(1)$ -reducible connection to quantum invariants of links I: R-matrix and Bureau representation*, [math.QA/9806004](#).
- [R2] ———, *A rationality conjecture about Kontsevich integral of knots and its implications to the structure of the colored Jones polynomial*, preprint [math.GT/0106097](#).
- [R3] ———, *A universal $U(1)$ -RCC invariant of links and rationality conjecture*, preprint 2002 [math.GT/0201139](#).
- [Rb] J. Roberts, *Kirby calculus in manifolds with boundary*, Turkish J. Math. **21** (1997) 111–117.
- [Sm] N. Smythe, *Boundary links*, Topology Seminar (Wisconsin 1965) Princeton Univ. Press, Princeton, NJ 1966, 69–72.
- [Tu] V. Turaev, *Conway and Kauffman modules of a solid torus*, J. Soviet Math. **52** (1990) 2799–1805.
- [Vo] P. Vogel, *The Cohn localization of the free group ring*, Math. Proc. Cambridge Phil. Soc. **111** (1992) 433–443.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA.
E-mail address: stavros@math.gatech.edu

E-mail address: kriker@math.gatech.edu