

ON KAN FIBRATIONS FOR MALTSEV ALGEBRAS

MAMUKA JIBLADZE AND TEIMURAZ PIRASHVILI

It is well known that simplicial groups are Kan complexes and more generally any surjective homomorphism of simplicial groups is a Kan fibration. From results in [1] among other things the following statements follow:

1. **Theorem.** *Any simplicial model of a Maltsev theory is a Kan complex.*
2. **Theorem.** *If \mathbb{T} is an algebraic theory, such that any simplicial \mathbb{T} -model is a Kan complex, then \mathbb{T} is a Maltsev theory.*

The aim of this note is twofold, firstly to give direct proofs of these facts without using category theory machinery and secondly to get sharper results. In particular we prove that any surjective homomorphism of Maltsev algebras is a Kan fibration.

A *Maltsev operation* in an algebraic theory \mathbb{T} is a ternary operation $[-, -, -]$ in \mathbb{T} satisfying the identities

$$[a, a, b] = b \text{ and } [a, b, b] = a.$$

An algebraic theory is called Maltsev if it possesses a Maltsev operation. Clearly the theory of groups is a Maltsev theory by taking $[a, b, c] = ab^{-1}c$. More generally theory of loops is a Maltsev theory; this last fact has already been used in homotopy theory – see [3]. Another sort of example of Maltsev theory is the theory of Heyting algebras.

We start by proving a more strong version of Theorem 2. We let

$$S^1 = \Delta^1 / \partial \Delta^1$$

be the smallest simplicial model of the circle. Moreover, we let $S_{\mathbb{T}}^1$ be the simplicial \mathbb{T} -model, which is obtained by applying the free \mathbb{T} -model functor degreewise to S^1 .

3. **Proposition.** *Let \mathbb{T} be an algebraic theory such that $S_{\mathbb{T}}^1$ satisfies the (1, 2)-th Kan condition in dimension 2. That is, for any 1-simplices x_1, x_2 with $d_1 x_1 = d_1 x_2$ there is a 2-simplex x with $d_1 x = x_1$ and $d_2 x = x_2$. Then \mathbb{T} is a Maltsev theory.*

Proof. Denote the unique nondegenerate 1-simplex of S^1 by σ and the unique vertex $d_0 \sigma = d_1 \sigma$ by $*$. So $S_{\mathbb{T}}^1$ in dimension zero is the free \mathbb{T} -model generated by $*$. Similarly $S_{\mathbb{T}}^1$ in dimension one is the free \mathbb{T} -model generated by $s_0 *$ and σ , and in dimension two it is the free \mathbb{T} -model generated by $s_1 s_0 *$, $s_0 \sigma$, and $s_1 \sigma$. Since $d_1 s_0 * = d_1 \sigma = *$, the (1, 2)-th Kan condition implies existence of a 2-simplex x of $S_{\mathbb{T}}^1$ with $d_1 x = s_0 *$ and $d_2 x = \sigma$. This means there is an element $x(s_1 s_0 *, s_0 \sigma, s_1 \sigma)$ in the free \mathbb{T} -model with three generators $s_1 s_0 *$, $s_0 \sigma$, $s_1 \sigma$ such that the equalities $x(d_1 s_1 s_0 *, d_1 s_0 \sigma, d_1 s_1 \sigma) = s_0 *$ and $x(d_2 s_1 s_0 *, d_2 s_0 \sigma, d_2 s_1 \sigma) = \sigma$ hold in the free \mathbb{T} -algebra with two generators $s_0 *$, σ . Applying standard simplicial identities we see that this means $x(s_0 *, \sigma, \sigma) = s_0 *$ and $x(s_0 *, s_0 *, \sigma) = \sigma$, i. e. that x is a Maltsev operation. \square

The following theorem shows that if \mathbb{T} is a Maltsev theory then all surjective homomorphisms of simplicial \mathbb{T} -models are Kan fibrations which obviously implies Theorem 1. Our proof uses exactly the same inductive argument as the one given in [2] for simplicial groups (see page 130 in [2]), except that we modify slightly the initial input.

4. Theorem. *Any surjective homomorphism $f : X \rightarrow Y$ of simplicial models of a Maltsev theory is a Kan fibration.*

Proof. For $n > 0$ and $0 \leq k \leq n$, given $y \in Y_n$ with $d_i y = f(x_i)$ for $i \neq k$, $0 \leq i \leq n$, where x_i are elements of X_{n-1} with matching faces, we have to find an $x \in X_n$ with $f(x) = y$ and $d_i x = x_i$ for $i \neq k$. Take $w_{-1} \in f^{-1}(y)$ and then put

$$w_j = [w_{j-1}, s_j d_j w_{j-1}, s_j x_j]$$

for $0 \leq j < k$; moreover put $w_{n+1} = w_{k-1}$ and

$$w_j = [w_{j+1}, s_{j-1} d_j w_{j+1}, s_{j-1} x_j]$$

for $n \geq j > k$.

We then have by induction for $0 \leq j < k$

$$f(w_j) = [f(w_{j-1}), s_j d_j f(w_{j-1}), s_j f(x_j)] = [y, s_j d_j y, s_j d_j y] = y$$

and for $n \geq j > k$

$$f(w_j) = [f(w_{j+1}), s_{j-1} d_j f(w_{j+1}), s_{j-1} f(x_j)] = [y, s_{j-1} d_j y, s_{j-1} d_j y] = y.$$

Furthermore for $0 \leq j < k$

$$\begin{aligned} d_i w_j &= [d_i w_{j-1}, d_i s_j d_j w_{j-1}, d_i s_j x_j] \\ &= \begin{cases} [x_i, s_{j-1} d_{j-1} d_i w_{j-1}, s_{j-1} d_i x_j] = \\ [x_i, s_{j-1} d_{j-1} x_i, s_{j-1} d_{j-1} x_i] = x_i, & 0 \leq i < j, \\ [d_j w_{j-1}, d_j w_{j-1}, x_j] = x_j, & i = j \end{cases} \end{aligned}$$

and then for $n \geq j > k$,

$$\begin{aligned} d_i w_j &= [d_i w_{j+1}, d_i s_{j-1} d_j w_{j+1}, d_i s_{j-1} x_j] \\ &= \begin{cases} [x_i, s_{j-2} d_{j-1} d_i w_{j+1}, s_{j-2} d_i x_j] = \\ [x_i, s_{j-2} d_{j-1} x_i, s_{j-2} d_{j-1} x_i] = x_i, & 0 \leq i < k, \\ [x_i, s_{j-1} d_j d_i w_{j+1}, s_{j-1} d_{i-1} x_j] = \\ [x_i, s_{j-1} d_j x_i, s_{j-1} d_j x_i] = x_i, & j < i \leq n, \\ [d_j w_{j+1}, d_j w_{j+1}, x_j] = x_j, & i = j \end{cases} \end{aligned}$$

Thus with $x = w_{k+1}$ one obtains $f(x) = y$ and $d_i x = x_i$ for $i \neq k$, $0 \leq i \leq n$, as desired. \square

REFERENCES

- [1] A. CARBONI, G. M. KELLY and M. C. PEDICCHIO. *Some remarks on Maltsev and Goursat categories*. Appl. Categ. Structures 1(1993), no. 4, 385–421.
- [2] E. B. CURTIS. *Simplicial homotopy theory*. Advan. in Math. 6 1971 107–209 (1971).
- [3] S. KLAUS. *On simplicial loops and H-spaces*. Topology Appl. 112 (2001), no. 3, 337–348.

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, DÉPARTEMENT DE MATHÉMATIQUE, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM

A. RAZMADZE MATHEMATICAL INSTITUTE, M. ALEXIDZE STR. 1, TBILISI 380093, REPUBLIC OF GEORGIA