

Projective bundles over hyperkähler manifolds and stability of Fourier-Mukai transform

Misha Verbitsky,¹

verbit@thelema.dnttm.ru, verbit@mccme.ru

Abstract

Let M be a K3 surface, B a stable bundle on M , and X the coarse moduli of stable deformations of B . The Fourier-Mukai transform is a functor $FM : D_b(M) \rightarrow D_b(X)$, where $D_b(M)$, $D_b(X)$ denotes the derived category of coherent sheaves on M , X .

Consider a stable bundle B_1 on M , and let $FM(B_1)$ be the corresponding complex of coherent sheaves on X . Assume X is compact. It was conjectured that the cohomology of the complex $FM(B_1)$ are polystable sheaves on X . This conjecture is known when the bundles B , B_1 have zero degree. We prove it for arbitrary stable bundles B , B_1 .

Contents

1	Introduction	2
1.1	Fourier-Mukai transform and stability	2
1.2	Fourier-Mukai transform and Yang-Mills geometry	5
1.3	Projectively hyperholomorphic bundles	8
1.4	Quaternionic Dolbeault complex	9
1.5	Contents	10
2	Hyperkähler manifolds	12
3	Hyperkähler manifolds and stable bundles	14
3.1	Holomorphic and hyperholomorphic connections	14
3.2	Hyperholomorphic bundles and Yang-Mills connections.	16
4	Projective bundles and projectively hyperholomorphic bundles	17
4.1	Projective bundles and $PGL(n)$ -bundles	18
4.2	Projective hyperholomorphic bundles: the definition	19
4.3	Moduli of projectively hyperholomorphic bundles	24

¹The author is partially supported by CRDF grant RM1-2087 and EPSRC grant GR/R05215.

5	Hyperholomorphic sheaves	25
5.1	Stable sheaves and Yang-Mills connections	25
5.2	Stable hyperholomorphic sheaves over hyperkähler manifolds	27
5.3	Weakly hyperholomorphic sheaves	28
6	Deformation spaces and universal bundles	28
6.1	Coarse moduli spaces	28
6.2	Existence of the universal bundle	29
7	Fourier-Mukai transform	31
8	Quaternionic Dolbeault complex	34
8.1	Quaternionic Dolbeault complex: the definition	34
8.2	Connections in modules over DG-algebras	37
8.3	Hodge decomposition for the quaternionic Dolbeault complex	40
8.4	The Dolbeault bicomplex and quaternionic Dolbeault complex	41
9	qD-modules and coherent sheaves	44
9.1	qD-modules: the definition	45
10	qD-Laplace operator and the extended qD-complex	47
10.1	Extended quaternionic Dolbeault bicomplex	47
10.2	qD-Laplace operator	48
10.3	qD-Laplace operator and the quaternionic Kodaira relations	50
11	Quaternionic Dolbeault complex of a product	51
11.1	Bidegree: the definition	51
11.2	Fiberwise connection operators on a product of two hyperkähler manifolds	52
12	Hyperholomorphic structure on a pushforward of a hyperholomor- phic bundle	54
12.1	Quaternionic Dolbeault complex and a C^∞ -pushforward of a bundle	55
12.2	Quaternionic Dolbeault complex and the derived direct image functor	58

1 Introduction

1.1 Fourier-Mukai transform and stability

Fourier-Mukai transform was discovered by S. Mukai in 1981 ([Mu1]). Mukai worked in the following situation. Let T be a compact torus or abelian variety, and $\hat{T}$ the dual torus, which is by definition a moduli space of line bundles of degree 0 on T . The **Poincaré bundle** P is a line bundle of degree zero on the product $T \times \hat{T}$, defined in such a way that for all $t \in \hat{T}$, the

restriction of P to $T \times \{t\}$ is isomorphic to the line bundle corresponding to the point $t \in \hat{T}$. This bundle is also called **the universal bundle**.

Given a coherent sheaf (or a complex of sheaves) F on T , let $FM(F)$ be the total derived direct image

$$FM(F) := R^\bullet(\pi_2)_*(P \otimes \pi_1^*F),$$

where $\pi_1, \pi_2 : T \times \hat{T} \rightarrow T, \hat{T}$ are the natural projection maps.

Clearly, $F \rightarrow FM(F)$ defines a functor of derived categories of coherent sheaves

$$FM : D_b(T) \rightarrow D_b(\hat{T}).$$

Consider the dual torus $\hat{\hat{T}}$ to $\hat{T}$. Clearly, $\hat{\hat{T}}$ is naturally isomorphic to T . Applying Fourier-Mukai functor to the torus $\hat{T}$, we obtain the transform

$$F\hat{M} : D_b(\hat{T}) \rightarrow D_b(T).$$

It is easy to check that the composition functor

$$FM \circ F\hat{M} : D_b(T) \rightarrow D_b(T)$$

is equivalent to identity.

The usual Fourier transform has a similar interpretation in terms of D -moduli. There is also a version of Fourier transform for perverse sheaves over $\mathbb{A}_{\mathbb{F}_q}$, where $\mathbb{A}_{\mathbb{F}_q}$ denotes an affine space over a field of char p .

This reveals the Fourier-Mukai transform as one of the most fundamental operations in algebraic geometry.

M. Kontsevich proposed an interpretation of the Mirror Conjecture in terms of the derived category of coherent sheaf on one of the mirror manifolds ([Kon]). It turns out that the derived category of coherent sheaves on a complex manifold M contains a great wealth of information about the geometry of M . In fact, if M is projective and the canonical or anti-canonical bundle of M is ample, the manifold M can be reconstructed from $D_b(M)$ ([BO], [P]). This puts a spotlight on the cases when such a reconstruction is **not** possible.

For instance, the Fourier-Mukai transform implies the existence of an equivalence between $D_b(T)$ and $D_b(\hat{T})$ ([Mu1]).

In 1990-ies, a number of similar construction appeared (see e.g. [BBR1]-[BBR3], [Ma]). Let M be a K3 surface, B a stable bundle, and X the moduli of stable deformations of B . S. Mukai [Mu2] investigated this situation in

great detail. It was shown that X is smooth and hyperkähler. Under some additional assumptions, X is also compact, and there exists a universal bundle $\mathcal{B}$ on $M \times X$. In more special cases, X is a K3 surface with periods prescribed by S. Mukai [Mu5].

This situation is analogous to the usual duality between the abelian varieties, with the role of a dual manifold played by X . In such a case, the Fourier-Mukai transform is also defined, in the same way it was defined for a torus (see Section 7 for details). It is also known that FM is (under additional assumptions) invertible ([Ma]). This establishes an equivalence of the derived categories of coherent sheaves ([Br], [BrM], [O]).

Given a complex of coherent sheaves $F^\bullet \in D_b(M)$, we may speak of its cohomology sheaves, which are coherent sheaves on M . It was conjectured that Fourier-Mukai transform (under some additional assumptions) preserves the Mumford stability of coherent sheaves ([Mu3]). We prove this statement when the base is a K3 surface.

Theorem 1.1: Let M be a K3 surface, B a stable bundle, and X the moduli of stable deformations of B . Assume that X is compact, and the universal bundle $\tilde{B}$ is well defined on $M \times X$. Consider the functor of Fourier-Mukai transform $FM : D_b(M) \longrightarrow D_b(X)$,

$$FM(B_1) = R^\bullet(\pi_2)_*(\tilde{B} \otimes \pi_1^* B_1).$$

Let B_1 be a stable bundle on M , and denote by $FM^i(B_1)$ the cohomology sheaves of the complex $FM(B_1)$. Then $FM^i(B_1)$ is polystable, up to torsion.¹

Proof: See Section 7. ■

Let M be a Kähler manifold, with $\omega \in H^2(M)$ a Kähler class, and $c \in H^2(M)$ an arbitrary cohomology class. Consider the number

$$\deg(c) := \int_M c \wedge \omega^{\dim_{\mathbb{C}} M - 1}$$

This number is called **the degree of c** .

¹That is, the reflexive hull $FM^i(B_1)^{**}$ is polystable. For a definition of reflexive sheaves and reflexive hull, see Definition 5.1.

In [BBR2] (see also [BBR3]), Theorem 1.1 was proven for $\deg c_1(B) = \deg c_1(B_1) = 0$, and $\dim_{\mathbb{C}} X = 2$. We prove Theorem 1.1 in full generality (Section 7).

In [BBR3], there was a sketch of the proof of Theorem 1.1 for arbitrary dimension of X (and $\deg c_1(B) = \deg c_1(B_1) = 0$), under an additional assumption that the cohomology sheaves $H^i(FM(B_1))$ are smooth. In the following Subsection, we extend the arguments of [BBR3] to deal with possible singularities.

A version of Theorem 1.1 is true when M is a compact 2-dimensional torus and $\deg c_1(B) = \deg c_1(B_1) = 0$ (see [DK], [FL], [BBR3]).

However, for arbitrary $\deg c_1(B)$, $\deg c_1(B_1)$ and M a torus, the Theorem 1.1 is false, as established by Kota Yoshioka ([Y1]). It is also false when M is a K3 surface and B is not locally free; the counterexample can be constructed using [Y3].

In [Y2], [Y4], the Fourier-Mukai transform was used at great length to obtain information about the geometry of the moduli of sheaves on a K3 surface. Under some additional assumptions, it is shown that the moduli of sheaves on a K3 surface are deformation equivalent to the Hilbert scheme of points on a K3 surface.

1.2 Fourier-Mukai transform and Yang-Mills geometry

In this Subsection, we state the proof of Theorem 1.1, for $\deg c_1(B) = \deg c_1(B_1) = 0$. This proof is an immediate application of Yang-Mills theory for hyperkähler manifolds.

Further on in this paper, we give an independent proof of Theorem 1.1 in full generality.

The present subsection is not used anywhere in this paper; however, the proof of a simple special case of Theorem 1.1 might be instructive for the reader.

A hyperkähler manifold (Definition 2.1) is a Riemannian manifold with an action of a quaternion algebra $\mathbb{H}$ in its tangent bundle TM , which is compatible with the metrics and the Levi-Civita connection. The group of unitary quaternions is isomorphic to $SU(2)$. This gives an action of $SU(2)$ on the tangent bundle of a hyperkähler manifold. By multiplicativity, we may extend this action to an action on differential forms.

The Calabi-Yau theorem (Theorem 2.4) provides any K3 surface, and any 2-dimensional compact complex torus, with a hyperkähler structure. Consider the corresponding action of $SU(2)$ on the cohomology of M .

Lemma 1.2: Let M be a hyperkähler manifold, $\dim H^{2,0}(M) = 1$, and $\eta \in H^{1,1}(M)$ a 1, 1-class in cohomology. Then η is $SU(2)$ -invariant if and only if $\deg \eta = 0$.

Proof: Follows from an elementary linear algebra argument (see, e.g. [V1]). ■

Clearly, Lemma 1.2 can be applied to a K3 surface.

Corollary 1.3: Let M be a K3 surface, and B a vector bundle on M . The Chern classes of B are $SU(2)$ -invariant if and only if $\deg c_1(B) = 0$.

Proof: Follows immediately from Lemma 1.2. ■

In [V1] we studied the geometry of stable bundles with $SU(2)$ -invariant Chern classes.

Definition 1.4: Let M be a compact hyperkähler manifold, I an induced complex structure, B a holomorphic bundle on (M, I) and ∇ a Hermitian connection on B . Then ∇ is called hyperholomorphic if its curvature $\Theta \in \Lambda^2(M) \otimes \text{End}(B)$ is a $SU(2)$ -invariant 2-form.

From the Gauss-Bonnet formula, it follows that the Chern classes of a hyperholomorphic bundle are $SU(2)$ -invariant.

In [V1] (see also Section 3) it was shown that a connection is hyperholomorphic if and only if it induces a holomorphic structure on B for any induced complex structure L on M .

Hyperholomorphic bundles were studied in [V1] in great detail. The following results were obtained.

Theorem 1.5: Let M be a compact hyperkähler manifold, I an induced complex structure and B a holomorphic vector bundle on (M, I) . Then B admits a hyperholomorphic connection if and only if B is a direct sum of stable bundles with $SU(2)$ -invariant Chern classes c_1, c_2 . In this case, the hyperholomorphic connection is unique.

Proof: [V1]; see also Section 3. ■

Holomorphic bundles admitting hyperholomorphic connection are called **hyperholomorphic**.

Theorem 1.6: Let M be a compact hyperkähler manifold, I an induced complex structure and B a hyperholomorphic vector bundle on (M, I) . Denote by X the coarse moduli space of stable deformations of B . Then X is singular hyperkähler,² and its normalization is a hyperkähler manifold.

Proof: [V1], [V3] (see also Subsection 4.3). ■

Theorem 1.7: In assumptions of Theorem 1.6, consider the universal bundle $\tilde{B}$, which is defined locally in X . Consider the natural (Peterson-Weil) metrics and a connection on ∇ in B . Then ∇ is hyperholomorphic.

Proof: Follows from an explicit description of the connection in $\tilde{B}$ given in [V1]. ■

Return to the assumptions of Theorem 1.1. Assume also that

$$\deg c_1(B) = \deg c_1(B_1) = 0.$$

By Corollary 1.3, then, the bundles B, B_1 are hyperholomorphic. Then X is hyperkähler, and the bundle $\pi^* B_1 \otimes \tilde{B}$ is hyperholomorphic by multiplicativity of the curvature.

Therefore, Theorem 1.1 for $\deg c_1(B) = \deg c_1(B_1) = 0$ is implied by the following result.

Theorem 1.8: Let $M = M_1 \times M_2$ be a product of compact hyperkähler manifolds, I an induced complex structure, and V a holomorphic vector bundle. Consider the natural projection $\pi_2 : M \rightarrow M_2$ and the derived direct image $R^i \pi_{2*} V$, which is a coherent sheaf on M_2 . Assume that V is hyperholomorphic. Then the reflexive hull $(R^i \pi_{2*} V)^{**}$ is polystable of degree 0.³

Proof: The proof of Theorem 1.8 for $R^i \pi_{2*} V$ a bundle is essentially contained in [BBR3]. Here we give a sketch of this proof.

²For a definition of a singular hyperkähler variety, see, e.g., [V3].

³A sheaf is called polystable of degree 0 if it is a direct sum of stable sheaves of degree 0; see Definition 3.4.

Consider the spinor bundle S on (M, I) . Since the canonical class of (M, I) is trivial, the bundle S is isomorphic to $\oplus_p \Lambda^{p,0}(M, I)$. Consider the sub-bundle of the spinor bundle generated by the vector tangent to the fibers of π_2 ; this bundle is isomorphic to $\pi_1^* \left(\oplus_p \Lambda^{p,0}(M_1, I) \right)$, where $\pi_1 : M \rightarrow M_1$ is the standard projection.

Let $D_1 : S_1 \rightarrow S_1$ be the composition of the Dirac operator and the orthogonal projection. Clearly, the cohomology of D_1 are identified with $R := \oplus_i R^i \pi_{2*} V$. Outside of singularities, we may consider R as a bundle on M_2 . Clearly, the Weil-Peterson metric and the connection on R are independent from the choice of induced complex structure I . Therefore, R is holomorphic with respect to any of the induced complex structures. This means that R is hyperholomorphic.

A hyperholomorphic bundle is polystable (Theorem 1.5). Therefore, R is polystable, if it is smooth.

If we apply this argument to the case when R might be singular, we obtain that R is equipped, outside of its singularities, with a hyperholomorphic connection. In [V2] we studied such sheaves at a great length. In [V4] the following theorem was proven.

Theorem 1.9: [V4] Let M be a compact hyperkähler manifold, I an induced complex structure, and F a reflexive coherent sheaf on (M, I) . Assume that F , outside of its singularities, is equipped with a hyperholomorphic connection. Then F is polystable of degree 0.

■

This finishes the proof of Theorem 1.8, when R is reflexive.

1.3 Projectively hyperholomorphic bundles

Let M be a compact hyperkähler manifold, I an induced complex structure, and B a holomorphic bundle on (M, I) . The bundle B is called **projectively hyperholomorphic** if $\text{End } B$ is hyperholomorphic. Deformation theory of projectively hyperholomorphic bundles was developed in [V1], and it is essentially not different from that of hyperholomorphic bundles.

In particular, it was shown that the deformation space of projectively hyperholomorphic bundles is (possibly singular) hyperkähler, and its normalization is smooth ([V1], [V3]).

The main idea of this paper is to expand on the similarity between hyperholomorphic and projectively hyperholomorphic bundles, using it to

prove the stability of Fourier-Mukai transform.

The first step is the following trivial claim ([V1]).

Claim 1.10: Let M be a K3 surface or a compact 2-dimensional torus, and B a stable bundle on M . Then B is projectively hyperholomorphic.

Proof: Clearly, $\text{End } B$ is polystable and $c_1(\text{End } B) = 0$. By Corollary 1.3, this implies that $\text{End } B$ is hyperholomorphic. ■

Let M be a hyperkähler manifold, I an induced complex structure, B a projectively hyperholomorphic bundle on (M, I) and X the space of stable deformations of B . If X is non-smooth, we replace it by its normalization, which is smooth by [V3].

If it exists, the universal bundle $\tilde{B}$ is projectively hyperholomorphic (see Theorem 7.3). The argument of Subsection 1.2 can be repeated word-by-word, until we come across an analogue of Theorem 1.8.

Theorem 1.11: Let $M = M_1 \times M_2$ be a product of compact hyperkähler manifolds, I an induced complex structure, and V a holomorphic vector bundle. Consider the natural projection $\pi_2 : M \rightarrow M_2$ and the derived direct image $R^i\pi_{2*}V$, which is a coherent sheaf on M_2 . Assume that V is projectively hyperholomorphic, and either $H^1(M_1)$ or $H^1(M_2)$ vanishes. Then the sheaf $R^i\pi_{2*}V$ is polystable of degree 0.

Proof: Follows from Theorem 7.5. ■

The proof of Theorem 1.11 takes a technique which is radically different from those employed to prove Theorem 1.8. Most of the present paper is dedicated to developing the machinery of quaternionic Dolbeault complex, which is used to prove Theorem 1.11.

1.4 Quaternionic Dolbeault complex

The main idea of the proof of Theorem 1.11 comes from the theory of D -modules and the study of quaternionic Dolbeault complex by S. Salamon and others ([S], [CS], [B], [L], [Ka]). Let M be a smooth manifold, and B a vector bundle. It is well known that to define a flat connection on B it means to define a map

$$\Lambda^0(M) \otimes B \xrightarrow{\nabla} \Lambda^1(M) \otimes B \xrightarrow{\nabla} \Lambda^2(M) \otimes B \xrightarrow{\nabla} \dots$$

which satisfies the Leibniz identity

$$\nabla(a \otimes b) = a \wedge \nabla(b) + da \otimes b, \quad (1.1)$$

where d denotes de Rham differential, $a \in \Lambda^*(M)$, $b \in B$.

A similar interpretation exists for holomorphic vector bundles (see Example 8.8). Given a differential graded algebra (A^*, d) we define a **bundle with connection over (A^*, d)** (Definition 8.4) as an A^0 -module equipped with a map

$$B \xrightarrow{\nabla} B \otimes_{A^0} A^1 \xrightarrow{\nabla} B \otimes_{A^0} A^2 \xrightarrow{\nabla} B \otimes_{A^0} A^3 \xrightarrow{\nabla} \dots$$

satisfying the Leibniz relation (1.1) (see Definition 8.4 for details).

For any hyperkähler manifold, one defines a canonical DG-algebra called **quaternionic Dolbeault complex** (Section 8). We show that a bundle with connection over a quaternionic Dolbeault complex is the same as a bundle with hyperholomorphic connection.

We compute the quaternionic Dolbeault complex explicitly in terms of $(p, 0)$ -differential forms (see (8.18)). This allows to express the holomorphic cohomology of a bundle in terms of the quaternionic Dolbeault complex (Subsection 12.2).

Comparing this with the decomposition-type results for quaternionic Dolbeault complex of a product (Section 11), we conclude that the direct image of a projectively hyperholomorphic bundle is projectively hyperholomorphic (Subsection 12.2).

1.5 Contents

- The present Introduction is independent from the rest of this paper. We give an abbreviated history of the problem, and explain the way it is solved.
- Section 2 is a compendium of basic results of hyperkähler geometry: Calabi-Yau theorem and $SU(2)$ -action on the differential forms and cohomology.
- Section 3 deals with Yang-Mills theory and hyperholomorphic connections on stable bundles over a hyperkähler manifold. We follow [V1].
- In Section 4 we give some expository treatment of projective bundles and projectively hyperholomorphic bundles. The main ideas of Section 4 are contained in [V1], but the approach of Section 4 is more

fundamental, connecting projectively hyperholomorphic bundles with the general theory of holomorphic G -bundles.

- Section 5 gives an abbreviated exposition of some of results in [V2] and [V4]. We define hyperholomorphic sheaves and list some of their properties.
- Section 6 deals with the coarse moduli of G -bundles and vector bundles. All the results Section 6 of are well known. Given a stable projective bundle P on M , we show that the universal bundle exists on the product of M and the moduli of P .
- In Section 7, we define the Fourier-Mukai transform, following [Mu1], [Ma] and others. We state our main result, Theorem 7.2, which says that Fourier-Mukai transform preserves stability. We reduce Theorem 7.2 to another theorem, stating that a derived push-forward of a projectively hyperholomorphic bundle is projectively hyperholomorphic, outside of singularities.

Section 7 is based on preliminary Sections 3-6, where we state the necessary results and definitions. The material of Sections 5-6 is not used anywhere except in Section 7.

The original results of this paper start from Section 7.

- In Section 8, we define (following S. Salamon) the quaternionic Dolbeault complex and relate it to hyperholomorphic and projectively hyperholomorphic bundles. We describe the quaternionic Dolbeault complex explicitly in terms of the usual Dolbeault complex of a manifold. Section 8 depends on the results and definitions of Sections 2-4.
- In Section 9, we deal with the singularities of the sheaves equipped with quaternionic Dolbeault connections. We construct a functor from the category of such sheaves to the category of quasicoherent sheaves on (M, I) . We show that this functor is compatible with the pushforward under a map $j : (M \setminus Z) \rightarrow M$, for $\text{codim}_{\mathbb{C}} Z \geq 2$. This is used to relate the quaternionic Dolbeault action and the reflexization of coherent sheaves. Together with the results of [V4], this is used further on to show that a pushforward of a projectively hyperholomorphic sheaf from $M_1 \times M_2$ to M_2 is projectively hyperholomorphic, if it is projectively hyperholomorphic outside of singularities and either $H^1(M_1) = 0$ or $H^2(M_2) = 0$. Section 9 is based on Section 8.

- In Section 10, we study the natural Laplace operator on the quaternionic Dolbeault complex. Using the hyperkähler version of the Kodaira relations ([V1]), we relate this Laplacian and the standard Laplacian associated with the Dolbeault differential. Section 10 is based on Section 8 and the results of [V1].
- In Section 11, we study the natural decomposition of a quaternionic Dolbeault complex of a product of hyperkähler manifolds. We use the explicit computation of the quaternionic Dolbeault complex. Section 11 is based on Section 8 and Sections 2-4.
- In Section 12, we relate the cohomology of quaternionic Dolbeault complex with coefficients in a bundle and the holomorphic cohomology of a bundle. Together with the decomposition obtained in Section 11 and the explicit computation of quaternionic Dolbeault complex (Section 8), this implies that the derived direct image of a projectively hyperholomorphic bundle is projectively hyperholomorphic outside of singularities.

2 Hyperkähler manifolds

This Section contains a compression of the basic and best known results and definitions from hyperkähler geometry, found, for instance, in [Bes], [Be1] and [V1].

Definition 2.1: ([Bes]) A **hyperkähler manifold** is a Riemannian manifold M endowed with three complex structures I , J and K , such that the following holds.

- (i) the metric on M is Kähler with respect to these complex structures and
- (ii) I , J and K , considered as endomorphisms of a real tangent bundle, satisfy the relation $I \circ J = -J \circ I = K$.

The notion of a hyperkähler manifold was introduced by E. Calabi ([C]).

Clearly, a hyperkähler manifold has a natural action of the quaternion algebra $\mathbb{H}$ in its real tangent bundle TM . Therefore its complex dimension

is even. For each quaternion $L \in \mathbb{H}$, $L^2 = -1$, the corresponding automorphism of TM is an almost complex structure. It is easy to check that this almost complex structure is integrable ([Bes]).

Definition 2.2: Let M be a hyperkähler manifold, and L a quaternion satisfying $L^2 = -1$. The corresponding complex structure on M is called **an induced complex structure**. The M , considered as a Kähler manifold, is denoted by (M, L) . In this case, the hyperkähler structure is called **compatible with the complex structure L** .

Definition 2.3: Let M be a complex manifold and Θ a closed holomorphic 2-form over M such that $\Theta^n = \Theta \wedge \Theta \wedge \dots$, is a nowhere degenerate section of a canonical class of M ($2n = \dim_{\mathbb{C}}(M)$). Then M is called **holomorphically symplectic**.

Let M be a hyperkähler manifold; denote the Riemannian form on M by $\langle \cdot, \cdot \rangle$. Let the form $\omega_I := \langle I(\cdot), \cdot \rangle$ be the usual Kähler form which is closed and parallel (with respect to the Levi-Civita connection). Analogously defined forms ω_J and ω_K are also closed and parallel.

A simple linear algebraic consideration ([Bes]) shows that the form $\Theta := \omega_J + \sqrt{-1}\omega_K$ is of type $(2, 0)$ and, being closed, this form is also holomorphic. Also, the form Θ is nowhere degenerate, as another linear algebraic argument shows. It is called **the canonical holomorphic symplectic form of a manifold M** . Thus, for each hyperkähler manifold M , and an induced complex structure L , the underlying complex manifold (M, L) is holomorphically symplectic. The converse assertion is also true:

Theorem 2.4: ([Be1], [Bes]) Let M be a compact holomorphically symplectic Kähler manifold with the holomorphic symplectic form Θ , a Kähler class $[\omega] \in H^{1,1}(M)$ and a complex structure I . Let $n = \dim_{\mathbb{C}} M$. Assume that $\int_M \omega^n = \int_M (Re\Theta)^n$. Then there is a unique hyperkähler structure $(I, J, K, (\cdot, \cdot))$ over M such that the cohomology class of the symplectic form $\omega_I = (\cdot, I\cdot)$ is equal to $[\omega]$ and the canonical symplectic form $\omega_J + \sqrt{-1}\omega_K$ is equal to Θ .

Theorem 2.4 follows from the conjecture of Calabi, proven by S.-T. Yau ([Yau]). ■

Let M be a hyperkähler manifold. We identify the group $SU(2)$ with the group of unitary quaternions. This gives a canonical action of $SU(2)$

on the tangent bundle, and all its tensor powers. In particular, we obtain a natural action of $SU(2)$ on the bundle of differential forms.

The following lemma is clear.

Lemma 2.5: The action of $SU(2)$ on differential forms commutes with the Laplacian.

Proof: This is Proposition 1.1 of [V0]. ■

Thus, for compact M , we may speak of the natural action of $SU(2)$ in cohomology.

Further in this article, we use the following statement.

Lemma 2.6: Let ω be a differential form over a hyperkähler manifold M . The form η is $SU(2)$ -invariant if and only if it is of Hodge type (p, p) with respect to all induced complex structures on M .

Proof: Let I be an induced complex structure, and $\rho_I : U(1) \rightarrow SU(2)$ the corresponding embedding, induced by the map $\mathbb{R} = \mathfrak{u}(1) \rightarrow \mathfrak{su}(2)$, $1 \rightarrow I$. The Hodge decomposition on $\Lambda^*(M)$ coincides with the weight decomposition of the $U(1)$ -action ρ_I . An $SU(2)$ -invariant form is also invariant with respect to ρ_I , and therefore, has Hodge type (p, p) . Conversely, if a η is invariant with respect to ρ_I , for all induced complex structures I , then η is invariant with respect to the Lie group G generated by these $U(1)$ -subgroups of $SU(2)$. A trivial linear-algebraic argument ensures that the group G coincides with the whole $SU(2)$. This proves Lemma 2.6. ■

3 Hyperkähler manifolds and stable bundles

This section contains several versions of a definition of hyperholomorphic connection in a complex vector bundle over a hyperkähler manifold. We follow [V1].

3.1 Holomorphic and hyperholomorphic connections

Let B be a holomorphic vector bundle over a complex manifold M , ∇ a connection in B and $\Theta \in \Lambda^2 \otimes \text{End}(B)$ be its curvature. This connection is called **compatible with a holomorphic structure** if $\nabla_X(\zeta) = 0$ for any holomorphic section ζ and any antiholomorphic tangent vector field

$X \in T^{0,1}(M)$. If there exists a holomorphic structure compatible with the given Hermitian connection then this connection is called **integrable**.

One can define a **Hodge decomposition** in the space of differential forms with coefficients in any complex bundle, in particular, $\text{End}(B)$.

Theorem 3.1: Let ∇ be a Hermitian connection in a complex vector bundle B over a complex manifold. Then ∇ is integrable if and only if $\Theta \in \Lambda^{1,1}(M, \text{End}(B))$, where $\Lambda^{1,1}(M, \text{End}(B))$ denotes the forms of Hodge type $(1,1)$. Also, the holomorphic structure compatible with ∇ is unique.

Proof: This is Proposition 4.17 of [Kob], Chapter I. ■

This proposition is a version of Newlander-Nirenberg theorem. For vector bundles, it was proven by M. Atiyah and R. Bott.

Definition 3.2: Let B be a Hermitian vector bundle with a connection ∇ over a hyperkähler manifold M . Then ∇ is called **hyperholomorphic** if ∇ is integrable with respect to each of the complex structures induced by the hyperkähler structure.

This definition can be word-by-word extended to principal G -bundles, with G a compact form of a complex Lie group.

As follows from Theorem 3.1, ∇ is hyperholomorphic if and only if its curvature Θ is of Hodge type $(1,1)$ with respect to any of the complex structures induced by a hyperkähler structure.

As follows from Lemma 2.6, ∇ is hyperholomorphic if and only if Θ is an $SU(2)$ -invariant differential form.

Example 3.3: (Examples of hyperholomorphic bundles)

- (i) Let M be a hyperkähler manifold, and TM be its tangent bundle equipped with the Levi-Civita connection ∇ . Consider a complex structure on TM induced from the quaternion action. Then ∇ is a Hermitian connection which is integrable with respect to each induced complex structure, and hence, is hyperholomorphic.
- (ii) For B a hyperholomorphic bundle, all its tensor powers are hyperholomorphic.
- (iii) Thus, the bundles of differential forms on a hyperkähler manifold are also hyperholomorphic.

3.2 Hyperholomorphic bundles and Yang-Mills connections.

Definition 3.4: Let F be a coherent sheaf over an n -dimensional compact Kähler manifold M . We define **the degree** $\deg(F)$ (sometimes the degree is also denoted by $\deg c_1(F)$) as

$$\deg(F) = \int_M \frac{c_1(F) \wedge \omega^{n-1}}{\text{vol}(M)}$$

and $\text{slope}(F)$ as

$$\text{slope}(F) = \frac{1}{\text{rank}(F)} \cdot \deg(F).$$

The number $\text{slope}(F)$ depends only on a cohomology class of $c_1(F)$.

Let F be a coherent sheaf on M and $F' \subset F$ its proper subsheaf. Then F' is called **destabilizing subsheaf** if $\text{slope}(F') \geq \text{slope}(F)$

A coherent sheaf F is called **stable**¹ if it has no destabilizing subsheaves. A coherent sheaf F is called **polystable** if it is a direct sum of stable sheaves of the same slope.

Let M be a Kähler manifold with a Kähler form ω . For differential forms with coefficients in any vector bundle there is a Hodge operator $L : \eta \rightarrow \omega \wedge \eta$. There is also a fiberwise-adjoint Hodge operator Λ (see [GH]).

Definition 3.5: Let B be a holomorphic bundle over a Kähler manifold M with a holomorphic Hermitian connection ∇ and a curvature $\Theta \in \Lambda^{1,1} \otimes \text{End}(B)$. The Hermitian metric on B and the connection ∇ defined by this metric are called **Yang-Mills** if

$$\Lambda(\Theta) = \text{constant} \cdot \text{Id} \Big|_B,$$

where Λ is a Hodge operator and $\text{Id} \Big|_B$ is the identity endomorphism which is a section of $\text{End}(B)$.

A holomorphic bundle is called **indecomposable** if it cannot be decomposed onto a direct sum of two or more holomorphic bundles.

¹In the sense of Mumford-Takemoto

The following fundamental theorem provides examples of Yang-Mills bundles.

Theorem 3.6: (Uhlenbeck-Yau) Let B be an indecomposable holomorphic bundle over a compact Kähler manifold. Then B admits a Hermitian Yang-Mills connection if and only if it is stable. Moreover, the Yang-Mills connection is unique, if it exists.

Proof: [UY]. ■

Proposition 3.7: Let M be a hyperkähler manifold, L an induced complex structure and B be a complex vector bundle over (M, L) . Then every hyperholomorphic connection ∇ in B is Yang-Mills and satisfies $\Lambda(\Theta) = 0$, where Θ is a curvature of ∇ .

Proof: We use the definition of a hyperholomorphic connection as one with $SU(2)$ -invariant curvature. Then Proposition 3.7 follows from the

Lemma 3.8: Let $\Theta \in \Lambda^2(M)$ be a $SU(2)$ -invariant differential 2-form on M . Then $\Lambda_L(\Theta) = 0$ for each induced complex structure L .²

Proof: This is Lemma 2.1 of [V1]. ■

Let M be a compact hyperkähler manifold, I an induced complex structure. For any stable holomorphic bundle on (M, I) there exists a unique Hermitian Yang-Mills connection which, for some bundles, turns out to be hyperholomorphic. It is possible to tell exactly when this happens.

Theorem 3.9: Let B be a stable holomorphic bundle over (M, I) , where M is a hyperkähler manifold and I is an induced complex structure over M . Then B admits a compatible hyperholomorphic connection if and only if the first two Chern classes $c_1(B)$ and $c_2(B)$ are $SU(2)$ -invariant.³

Proof: This is Theorem 2.5 of [V1]. ■

4 Projective bundles and projectively hyperholomorphic bundles

²By Λ_L we understand the Hodge operator Λ associated with the Kähler complex structure L .

³We use Lemma 2.5 to speak of action of $SU(2)$ in cohomology of M .

4.1 Projective bundles and $PGL(n)$ -bundles

Let M be a complex manifold, and B a holomorphic vector bundle, $\dim B = n$. Consider the principal $GL(n)$ -bundle G_B of linearly independent n -tuples of vectors in B . The bundle B can be reconstructed from G_B as follows

$$B = G_B \times_{GL(n)} V_n,$$

where $V_n = \mathbb{C}^n$ denotes the fundamental representation of $GL(n)$. This is well known.

A similar construction exists for principal $PGL(n)$ -bundles. Given a principal $PGL(n)$ -bundle G_P , we can consider the associated fibration

$$P := G_P \times_{PGL(n)} \mathbb{P}, \quad (4.1)$$

where $\mathbb{P}$ is $\mathbb{C}P^n$, equipped with the natural action of $PGL(n)$.

Definition 4.1: A projective vector bundle over M is a holomorphic fibration P over M , obtained from a principal $PGL(n)$ -bundle as in (4.1).

Remark 4.2: Any automorphism of $\mathbb{C}P^n$ is projective linear. Given a locally trivial holomorphic fibration P with the fibers isomorphic to $\mathbb{C}P^n$, it is very easy to reconstruct the corresponding principal $PGL(n+1)$ -bundle G_P . For any point $x \in M$, the fiber of G_P in x is the set of linearly independent n -tuples $p_1, \dots, p_n$ in the space $V_x = \mathbb{C}^{n+1}$ corresponding to $P|_x \cong \mathbb{C}P^n$, up to a complex multiplier. The group $PGL(P) = \text{Aut}_M(P)$ acts on $G_P|_x$ freely and transitively, in such a way that G_P forms a principal $PGL(n+1)$ -bundle.

Clearly, then, the bundle P satisfies (4.1). In other words, projective vector bundles are in one-to-one correspondence with the locally trivial holomorphic fibrations with the fibers isomorphic to $\mathbb{C}P^n$.

Definition 4.3: Let P be a projective vector bundle, and G_P the corresponding $PGL(n)$ -bundle. Assume that P is equipped with a smooth Hermitian metric h such that the bundle of Hermitian automorphisms of P is isomorphic to $PSU(n)$. Then P is called a Hermitian projective vector bundle.

Remark 4.4: In the above assumptions, the bundle $\text{Aut}_M(P)$ is equipped with a unique anticomplex involution $t : G_P \rightarrow G_P$ which fixes a $PSU(n)$ -subbundle defined by the Hermitian structure on P . This involution defines

a Hermitian metrics on $\text{Aut}(P)$ as follows:

$$h(a, b) := \text{Tr}(a \cdot t(b))$$

Since G_P is a principal $\text{Aut}(P)$ -bundle, a Hermitian structure on $\text{Aut}(P)$ induces a Hermitian structure h on G_P . Using the Hermitian metrics h and the holomorphic structure, one defines the connection on G_P in the usual way. This connection is called **the Hermitian connection**.

Remark 4.5: Let B be a Hermitian holomorphic vector bundle on M , and $\mathbb{P}B$ is projectivization. Clearly, $\mathbb{P}B$ is a Hermitian holomorphic projective vector bundle. The natural Hermitian connection obtained this way on $PGL(B)$ coincides with that constructed in Remark 4.4.

4.2 Projective hyperholomorphic bundles: the definition

Now assume that M is a hyperkähler manifold, and I an induced complex structure. Consider (M, I) as a Kähler manifold.

Definition 4.6: Let P be a projective bundle over (M, I) , equipped with a Hermitian structure. We say that P is hyperholomorphic if the Hermitian connection in the corresponding principal $PGL(n)$ -bundle is hyperholomorphic (Definition 3.2).

Remark 4.7: Given a hyperholomorphic projective vector bundle P over (M, I) , we obtain that P is a holomorphic projective bundle over (M, J) for any induced complex structure J . This is implied by the G -bundle version of the Newlander-Nirenberg theorem.

Let M be a Kähler manifold, B a Hermitian holomorphic vector bundle, and Λ, L the standard Hodge operators on differential forms (Subsection 3.2). We denote by Θ the curvature of B . Let $\text{Tr}(\Theta) \in \Lambda^{1,1}(M)$ be the trace of the curvature,

$$\text{Tr}(\Theta)(x, y) = \text{Tr}(\Theta(x, y)); \quad x, y \in TM$$

(we interpret $\Theta(x, y)$ as an endomorphism of B and take its trace).

Proposition 4.8: Let M be a connected hyperkähler manifold of real dimension $4n$, I an induced complex structure, and B a holomorphic Hermitian vector bundle on M . Denote the curvature of B by

$$\Theta \in \Lambda^{1,1}(M, \text{End}(B)).$$

Then the following conditions are equivalent

- (i) The bundle $\text{End}(B)$ is hyperholomorphic
- (ii) The traceless part of the curvature

$$\Theta_{tr} := \Theta - \frac{1}{\dim B} \text{Tr}(\Theta)$$

is $SU(2)$ -invariant, with respect to the natural action of $SU(2)$ on the differential forms.

- (iii) The projectivization $\mathbb{P}B$ is a hyperholomorphic projective bundle on M .

Proof: The implication (ii) $\Rightarrow$ (i) is quite easy. The curvature $\Theta' \in \text{End}(\text{End}(B))$ of $\text{End } B$ is expressed via Θ as follows

$$\Theta' = l(\Theta) - r(\Theta), \quad (4.2)$$

where $l : \text{End}(B) \longrightarrow \text{End}(\text{End}(B))$ is an operation of left multiplication by the sections of $\text{End}(B)$, and r is an operation of right multiplication. Now, the trace of Θ belongs to the center of $\text{End}(B)$, and therefore the traceless part of the curvature satisfies

$$\Theta' = l(\Theta) - r(\Theta) = l(\Theta_{tr}) - r(\Theta_{tr}) \quad (4.3)$$

Since the 2-form Θ_{tr} is $SU(2)$ -invariant, Θ' is also $SU(2)$ -invariant. By definition, this implies that $\text{End}(B)$ is hyperholomorphic. This proves an implication (ii) $\Rightarrow$ (i). The converse implication is obvious. Clearly, the map

$$LR : \text{End}(B) \longrightarrow \text{End}(\text{End}(B)), \quad LR(a) = l(a) - r(a)$$

is injective on the traceless part of $\text{End}(B)$. Therefore, $\text{End}(B)$ is hyperholomorphic if and only if the traceless part of Θ is $SU(2)$ -invariant. This proves the equivalence (i) $\Leftrightarrow$ (ii).

To prove (i) $\Rightarrow$ (iii), we notice that $\mathfrak{pgl}(B)$ is a quotient of $\mathfrak{gl}(B)$, which is hyperholomorphic by assumptions of (i). This means that $\mathbb{P}B$ is also hyperholomorphic. The converse implication is also clear. The Lie algebra bundle $\mathfrak{pgl}(B)$ corresponding to $PGL(B)$ is the bundle $\text{End}_{tr}(B)$ of traceless automorphisms of B , and it is hyperholomorphic by the assumptions of (iii). On the other hand, $\text{End}(B) = \text{End}_{tr}(B) \oplus \mathbb{C}$ where $\mathbb{C}$ is the trivial

holomorphic Hermitian bundle. Therefore, $\text{End}(B)$ is hyperholomorphic if and only if $\text{End}_{tr}(B)$ is hyperholomorphic. We proved the equivalence (i) $\Leftrightarrow$ (iii). ■

Definition 4.9: [V1] Let M be a hyperkähler manifold, I an induced complex structure, and B a Hermitian holomorphic vector bundle on (M, I) . We say that B is projectively hyperholomorphic if any of the conditions of Proposition 4.8 hold.

The following claim is trivial.

Claim 4.10: [V1] Let M be a compact hyperkähler manifold of real dimension 4, I an induced complex structure, and B a holomorphic vector bundle on (M, I) . Assume that B is stable. Then B admits a unique projectively hyperholomorphic Yang-Mills connection.

Proof: Let ∇ be the Yang-Mills connection on B , which exists and is unique by Theorem 3.6. Denote the corresponding connection on $\text{End}(B)$ by ∇_1 . The curvature of $\text{End}(B)$ can be algebraically expressed through the curvature of B . Using this expression, it is easy to show that $\text{End}(B)$ is also Yang-Mills (see (4.3)). By Theorem 3.9, to show that the Yang-Mills connection in $\text{End}(B)$ is hyperholomorphic, one needs to prove that the Chern classes $c_1(\text{End}(B))$ and $c_2(\text{End}(B))$ are $SU(2)$ -invariant. The first Chern class of $\text{End}(B)$ is zero, hence $SU(2)$ -invariant. The second Chern class belongs to the one-dimensional space $H^4(M)$, and it is clear that the standard action of $SU(2)$ on $H^4(M)$ is trivial. This proves that $c_1(\text{End}(B))$ and $c_2(\text{End}(B))$ are $SU(2)$ -invariant, and $\text{End}(B)$ is hyperholomorphic. By Proposition 4.8 (i) this implies that B is projectively hyperholomorphic. ■

Claim 4.11: Let M be a compact hyperkähler manifold, I an induced complex structure, and B a holomorphic vector bundle on (M, I) . Assume that B admits a projectively hyperholomorphic connection ∇ . Then B is polystable¹. Moreover, the Yang-Mills connection on B is projectively hyperholomorphic.

Proof: By Proposition 4.8 (i), the bundle $\text{End}(B)$ is hyperholomorphic; in particular, it is Yang-Mills and polystable. Given a destabilizing subsheaf

¹A polystable bundle is a direct sum of several stable bundles of the same slope

$F \subset B$, we obtain that the sheaf $\mathrm{Hom}(B, F) \subset \mathrm{End}(B)$ is also destabilising.

Denote by F' the quotient sheaf $F' := B/F$. By definition, V is isomorphic to $B^* \otimes F'$. We obtain a decomposition

$$\mathrm{End} B = B \otimes B^* = F \otimes B^* \oplus F' \otimes B^*. \quad (4.4)$$

To prove that B is polystable, we need to show that the extension

$$0 \longrightarrow F \longrightarrow B \longrightarrow F' \longrightarrow 0 \quad (4.5)$$

splits. If we tensor this exact sequence by B , it will be split by (4.4). Now, let us tensor (4.5) by $B \otimes B^*$. Thus obtained exact sequence splits because (4.5) tensored by B splits. On the other hand, the trivial sheaf $\mathcal{O}_M$ is a direct summand of $B \otimes B^*$, hence the extension (4.5) is a direct summand of the extension

$$0 \longrightarrow F \otimes B \otimes B^* \longrightarrow B \otimes B \otimes B^* \longrightarrow F' \otimes B \otimes B^* \longrightarrow 0.$$

We have shown that this extension splits; therefore, its direct summand (4.5) also splits. We have shown that B is polystable.

Let ∇ be the Yang-Mills connection on B . Then ∇ induces a Yang-Mills connection ∇_e on $\mathrm{End}(B)$. On the other hand, the Yang-Mills connection is unique. Therefore, ∇_e coincides with the original hyperholomorphic connection on $\mathrm{End}(B)$. We obtained that ∇ is projectively hyperholomorphic. ■

Remark 4.12: It is possible for a connection in bundle to be projectively hyperholomorphic, in the sense of Definition 4.9, and not Yang-Mills. For instance, any line bundle is obviously projectively hyperholomorphic. From Claim 4.11 it follows that the distinction between “projectively hyperholomorphic” and “projectively hyperholomorphic Yang-Mills” is not very important.

In [V1], we used a different definition of “projectively hyperholomorphic”. Namely, a projectively hyperholomorphic bundle, in the sense of [V1], is a holomorphic vector bundle with a Yang-Mills Hermitian metric satisfying conditions of Definition 4.9. In the present paper these bundles are called **projectively hyperholomorphic Yang-Mills**, sometimes abbreviated for **phYM-bundles**.

The following trivial lemma is used further on in the study of derived direct images.

Lemma 4.13: Let B be a projectively hyperholomorphic Yang-Mills bundle, Θ its curvature, and $Tr(\Theta) \in \Lambda^{1,1}(M)$ its trace. Then $Tr(\Theta)$ is harmonic.²

Proof: Clearly, $Tr(\Theta)$ is the curvature of the line bundle $\Lambda^q B$, $q = \dim B$ (the top exterior power of B). Since $\Lambda^q B$ is a tensor power of B , it is also a Yang-Mills bundle. We reduced Lemma 4.13 to the case when B is a line bundle. In this case, $\Theta \in \Lambda^{1,1}(M, \text{End } B) = \Lambda^{1,1}(M)$ is a closed 1,1-form which satisfies $\Lambda\Theta = \text{const}$. Therefore,

$$\partial\Lambda\Theta = 0. \quad (4.6)$$

By Kodaira's formulae, $[\Lambda\partial] = \sqrt{-1} \bar{\partial}^*$. Therefore, (4.6) implies $\bar{\partial}^*\Theta = 0$. Since Θ is of type (1,1) and closed, we also have $\bar{\partial}\Theta = 0$. Therefore, Θ is harmonic. ■

Remark 4.14: Yang-Mills bundles can be characterized as bundles which have harmonic curvature. This is shown by the same argument that proves Lemma 4.13.

The following Corollary gives a characterization of projectively hyperholomorphic Yang-Mills bundles in terms of curvature.

Corollary 4.15: Let M be a hyperkaehler manifold, I an induced complex structure, and B a holomorphic Hermitian vector bundle over (M, I) . Denote by

$$\Theta = \Theta_+ + \Theta_{SU(2)}$$

the isotypical decomposition of its curvature, with $\Theta_{SU(2)}$ being $SU(2)$ -invariant, and Θ_+ of weight 2 with respect to $SU(2)$. Then the following conditions are equivalent

- (i) The form $\Theta_+ \in \Lambda^{1,1}(M) \otimes \text{End } B$ is equal to $\omega \times Id_B$, with $Id_B \in \text{End } B$ the identity endomorphism of B and $\omega \in \Lambda^{1,1}(M)$ a harmonic form.
- (ii) The bundle B is projectively hyperholomorphic Yang-Mills.

²In fact, $Tr(\Theta)$ is harmonic for any Yang-Mills bundle; the same proof works.

Proof: The implication (ii) $\Rightarrow$ (i) is clear by Lemma 4.13. Indeed, since B is projectively hyperholomorphic, we have $\Theta_+ \omega \times Id_B$ (the non- $SU(2)$ -invariant part of the curvature is proportional to Id_B). Then Θ_+ is a non- $SU(2)$ -invariant part of the form $Tr(\Theta)$ which is harmonic by Lemma 4.13. Since the action of $SU(2)$ commutes with the Laplacian, the form Θ_+ is also harmonic.

The converse implication is clear by Lemma 3.8. If the bundle B satisfies the assumption (i), then it is by definition projectively hyperholomorphic; we need only to show it is Yang-Mills, that is, $\Lambda\Theta = const \cdot Id_B$. Given an $SU(2)$ -invariant 2-form η , we have $\Lambda\eta = 0$ (Lemma 3.8). Therefore, $\Lambda(\Theta) = \Lambda(\Theta_+)$. On the other hand, the operator Λ commutes with the Laplacian. Since, Θ_+ is harmonic, $\Lambda(\Theta_+)$ is also harmonic. A harmonic 0-form is constant. Therefore, $\Lambda(\Theta_+) = const \cdot Id_B$. We proved the implication (i) $\Rightarrow$ (ii). Corollary 4.15 is proven. ■

4.3 Moduli of projectively hyperholomorphic bundles

Let M be a compact hyperkähler manifold, I an induced complex structure, and B a stable holomorphic vector bundle on (M, I) . Assume that the Chern classes of $\text{End}(B)$ are $SU(2)$ -invariant, or, what is the same, B is projectively hyperholomorphic.

Consider the coarse moduli space of deformations of B (see Definition 6.1 for details).

In [V1] (see [V3] for details and a more rigorous definition), we defined singular hyperkähler varieties. A singular hyperkähler variety is a real analytic variety S , equipped with a real analytic metrics and a quaternion action in its sheaf of Kähler differentials, such that for every embedding $\mathbb{C} \hookrightarrow \mathbb{H}$ the corresponding almost complex structure on S is integrable and Kähler outside of singularities.

It was shown that the moduli space of projectively hyperholomorphic bundles is singular hyperkähler ([V1]).

In [V3], a desingularization theorem was proven for singular hyperkähler varieties. Given a singular hyperkähler variety X , and an induced complex structure I , consider a normalization $\tilde{X}$ of the complex variety (X, I) . It was shown that $\tilde{X}$ is smooth, hyperkähler, and does not depend on the choice of I . Moreover, locally in a neighbourhood of a singular point, the complex variety (X, I) is isomorphic to a union of several planes in a vector space.

To avoid the complicated notation, further on we shall speak of the moduli of projectively hyperholomorphic bundles as of a smooth hyperkähler manifold; if it is non-smooth, we replace it by its desingularization.

5 Hyperholomorphic sheaves

In [BS], S. Bando and Y.-T. Siu developed machinery allowing one to apply the methods of Yang-Mills theory to torsion-free coherent sheaves. In [V2], their work was applied to generalise the results of [V1] (see Section 3) to sheaves.

5.1 Stable sheaves and Yang-Mills connections

In this subsection, we repeat the basic definitions and results from [BS] and [OSS].

Definition 5.1: Let X be a complex manifold, and F a coherent sheaf on X . Consider the sheaf $F^* := \mathcal{H}om_{\mathcal{O}_X}(F, \mathcal{O}_X)$. There is a natural functorial map $\rho_F : F \longrightarrow F^{**}$. The sheaf F^{**} is called a **reflexive hull**, or **reflex-ization** of F . The sheaf F is called **reflexive** if the map $\rho_F : F \longrightarrow F^{**}$ is an isomorphism.

Remark 5.2: For all coherent sheaves F , the map $\rho_{F^*} : F^* \longrightarrow F^{***}$ is an isomorphism ([OSS], Ch. II, the proof of Lemma 1.1.12). Therefore, a reflexive hull of a sheaf is always reflexive.

Reflexive hull can be obtained by restricting to an open subset and taking the pushforward.

Lemma 5.3: Let X be a complex manifold, F a coherent sheaf on X , Z a closed analytic subvariety, $\text{codim } Z \geq 2$, and $j : (X \setminus Z) \hookrightarrow X$ the natural embedding. Assume that the pullback j^*F is reflexive on $(X \setminus Z)$. Then the pushforward j_*j^*F is also reflexive.

Proof: This is [OSS], Ch. II, Lemma 1.1.12. ■

Lemma 5.4: Let F be a reflexive sheaf on M , and X its singular set. Then $\text{codim}_M X \geq 3$

Proof: This is [OSS], Ch. II, 1.1.10. ■

Claim 5.5: Let X be a Kähler manifold, and F a torsion-free coherent sheaf over X . Then F (semi)stable if and only if F^{**} is (semi)stable.

Proof: This is [OSS], Ch. II, Lemma 1.2.4. ■

The admissible Hermitian metrics, introduced by Bando and Siu in [BS], play the role of the ordinary Hermitian metrics for vector bundles.

Let X be a Kähler manifold. In Hodge theory, one considers the operator $\Lambda : \Lambda^{p,q}(X) \longrightarrow \Lambda^{p-1,q-1}(X)$ acting on differential forms on X , which is adjoint to the multiplication by the Kähler form. This operator is defined on differential forms with coefficient in every bundle. Considering the curvature Θ of a bundle B as a 2-form with coefficients in $\text{End}(B)$, we shall speak of $\Lambda\Theta$ which is a section of $\text{End}(B)$.

Definition 5.6: Let X be a Kähler manifold, and F a reflexive coherent sheaf over X . Denote by $U \subset X$ the set of all points at which F is locally trivial. By definition, the restriction $F|_U$ of F to U is a bundle. An **admissible metric** on F is a Hermitian metric h on the bundle $F|_U$ which satisfies the following assumptions

- (i) the curvature Θ of (F, h) is square integrable, and
- (ii) the corresponding section $\Lambda\Theta \in \text{End}(F|_U)$ is uniformly bounded.

Definition 5.7: Let X be a Kähler manifold, F a reflexive coherent sheaf over X , and h an admissible metric on F . Consider the corresponding Hermitian connection ∇ on $F|_U$. The metric h and the Hermitian connection ∇ are called **hyperholomorphic** if its curvature satisfies

$$\Lambda\Theta \in \text{End}(F|_U) = c \cdot \text{id}$$

where c is a constant and id the unit section $\text{id} \in \text{End}(F|_U)$.

One of the main results of [BS] is the following analogue of the Uhlenbeck-Yau theorem (Theorem 3.6).

Theorem 5.8: Let M be a compact Kähler manifold, and F a coherent sheaf without torsion. Then F admits an admissible Yang–Mills metric if and only if F is polystable. Moreover, if F is stable, then this metric is unique, up to a constant multiplier.

Proof: [BS], Theorem 3. ■

5.2 Stable hyperholomorphic sheaves over hyperkähler manifolds

Let M be a compact hyperkähler manifold, I an induced complex structure, F a torsion-free coherent sheaf over (M, I) and F^{**} its reflexization. Recall that the cohomology of M are equipped with a natural $SU(2)$ -action (Lemma 2.5). The motivation for the following definition is Theorem 3.9 and Theorem 5.8.

Definition 5.9: Assume that the first two Chern classes of the sheaves F , F^{**} are $SU(2)$ -invariant. Then F is called **stable (polystable) hyperholomorphic** if F is stable (polystable), and **stable (polystable) projectively hyperholomorphic**, if $\text{End}(F)$ is stable (polystable) hyperholomorphic.

Remark 5.10: The slope of a hyperholomorphic sheaf is zero, because a degree of an $SU(2)$ -invariant 2-form is zero (Lemma 3.8).

Let M be a hyperkähler manifold, I an induced complex structure, and F a torsion-free sheaf over (M, I) . Consider the natural $SU(2)$ -action in the bundle $\Lambda^i(M, B)$ of the differential i -forms with coefficients in a vector bundle B . Let $\Lambda_{inv}^i(M, B) \subset \Lambda^i(M, B)$ be the bundle of $SU(2)$ -invariant i -forms.

Definition 5.11: Let $X \subset (M, I)$ be a complex subvariety of codimension at least 2, such that $F|_{M \setminus X}$ is a bundle, h be a metric on $F|_{M \setminus X}$ and ∇ the associated connection. Then ∇ is called **hyperholomorphic** if its curvature

$$\Theta_{\nabla} = \nabla^2 \in \Lambda^2 \left(M, \text{End} \left(F|_{M \setminus X} \right) \right)$$

is $SU(2)$ -invariant, i. e. belongs to $\Lambda_{inv}^2 \left(M, \text{End} \left(F|_{M \setminus X} \right) \right)$. The connection ∇ is called **admissible hyperholomorphic** if ∇ is hyperholomorphic and the metric h is admissible, in the sense of Definition 5.6.

Theorem 5.12: Let M be a compact hyperkähler manifold, I an induced complex structure and F a reflexive sheaf on (M, I) . Then F admits an admissible hyperholomorphic connection if and only if F is polystable hyperholomorphic.

Proof: This is [V2], Theorem 3.19. ■

5.3 Weakly hyperholomorphic sheaves

In most situations in this paper, we operate with a weaker notion of hyperholomorphic connection - there is no way to obtain the admissibility of the metrics. However, for the present purposes, the results of [V4] will suffice.

Definition 5.13: Let M be a hyperkähler manifold, I an induced complex structure, and F a torsion-free coherent sheaf on M . Assume that outside of a closed complex analytic set $Z \subset (M, I)$, $\text{codim}_{\mathbb{C}} Z \geq 3$, the sheaf F is smooth and equipped with a connection ∇ . Assume, moreover, that ∇ is hyperholomorphic. Then F is called weakly hyperholomorphic.

Theorem 5.14: Let M be a compact hyperkähler manifold, I an induced complex structure, and F a reflexive coherent sheaf on M which is indecomposable (that is, cannot be represented as a direct sum of non-trivial coherent sheaves). Assume that F is weakly hyperholomorphic. Then F is stable.

Proof: This is the main result of [V4]. ■

By Theorem 5.12, F admits a unique admissible hyperholomorphic Yang-Mills connection ∇_1 . However, we do not know whether $\nabla = \nabla_1$ or not.

Conjecture 5.15: In assumptions of Theorem 5.14, the connection ∇ is admissible.

Clearly, Theorem 5.14 is implied by Theorem 5.12 and Conjecture 5.15.

6 Deformation spaces and universal bundles

6.1 Coarse moduli spaces

Let M be a compact complex manifold, and G a complex Lie group. In this section, we study the deformations of vector bundles, or G -bundles, over M . We work in a rather abstract setting; in the applications, we deal with vector bundles, or $PGL(n)$ -bundles satisfying some stability conditions.

Let F be a vector bundle or a G -bundle, and $\mathcal{C}$ the category of vector bundles or G -bundles over M containing F . Denote by $\mathcal{C}(M)$ the corresponding category of bundles over M . Let $\mathcal{T}$ be some condition (such as stability) that might be true or false for objects of $\mathcal{C}(M)$.

Assume that all objects of $\mathcal{C}$ satisfying $\mathcal{T}$ are **simple**, that is, they have the smallest automorphism group that the geometry of $\mathcal{C}$ allows (for vector bundles, this condition translates to the usual notion of simplicity of a holomorphic vector bundle; for projective bundles, it means that the corresponding $\mathfrak{pgl}(n) = \mathfrak{sl}(n)$ -bundle does not have Lie algebra automorphisms).

Let X be a complex analytic space, with the marked point x_0 . Given the above data, a variation of F over a base X is an object $\mathcal{F}$ of $\mathcal{C}$ (a vector bundle or a G -bundle) over the product $M \times X$, such that for all $x \in X$ the restriction $\mathcal{F}|_{M \times \{x\}}$ of $\mathcal{F}$ to $M \times \{x\}$ satisfies $\mathcal{T}$, and $\mathcal{F}|_{M \times \{x_0\}}$ is isomorphic to F .

Definition 6.1: In the above assumptions, let S be a complex analytic space with the marked point denoted by $[F]$. Then S is called the **coarse moduli space of deformations of F , satisfying $\mathcal{T}$** , if the following conditions hold

- (i) There is a bundle $F' \in \mathcal{C}$ over M associated to each point $[F'] \in S$. This bundle always satisfies $\mathcal{T}$. The bundle F is associated with the marked point $[F] \in S$.
- (ii) For any variation $(X, \mathcal{F})$ of F with a connected base X there is a unique map $f : X \rightarrow S$, mapping $[F]$ to the marked point x_0 , such that for each point $x \in X$ the restriction of $\mathcal{F}$ on $\{x\} \times M$ is isomorphic to the bundle associated with the point $f(x) \in S$ by (i).
- (iii) For any point $[F'] \in S$ and a sufficiently small open neighbourhood U of $[F']$, there exists a bundle $\mathcal{F} \in \mathcal{C}$ over $M \times U$ such that the restriction of $\mathcal{F}$ to $\{[F']\} \times M$ is isomorphic to F' .

This definition is usually applied when $\mathcal{T}$ is some stability condition. In this case, S is called **the coarse moduli of stable deformations of F** .

6.2 Existence of the universal bundle

Definition 6.2: In assumptions of Definition 6.1, let $\mathcal{U}$ be a bundle from $\mathcal{C}$ over $M \times S$. We say that $\mathcal{U}$ is **universal** if for any point $[F'] \in S$, the restriction of $\mathcal{U}$ to $\{[F']\} \times M$ is isomorphic to F' .

The existence of the universal bundle is known for projective vector bundles. Recall that the projective vector bundles are in one-to-one correspondence with principal holomorphic $PGL(n)$ -bundles (Remark 4.2).

Proposition 6.3: Let M be a compact complex manifold, P a holomorphic projective vector bundle, $\mathcal{T}$ a true-or-false condition for holomorphic projective vector bundles, satisfying assumptions of Subsection 6.1, and S the coarse moduli of deformations of P satisfying $\mathcal{T}$ (see Definition 6.1 for details).¹ Then the universal bundle $\mathcal{U}$ exists over $M \times S$.

Proof: Locally in a neighbourhood of a point $[P'] \in S$, the universal bundle exists by Definition 6.1 (iii), and is, by definition, unique up to an isomorphism. To prove its existence globally, we need to glue together these local versions of the universal bundle. Generally speaking, this might be non-trivial, because the bundle P , which we are trying to deform, might have automorphisms. Then, gluing together the universal bundle from its local versions will have obstructions. These obstructions lie in the first cohomology $H^1(S, \text{Aut}(\mathcal{F}))$ of the sheaf of local automorphisms of the (locally defined) universal bundle of P . However, the universal bundle exists when the deformations of P have no automorphisms. We assumed that the condition $\mathcal{T}$ implies simplicity; for a projective bundle, this means that the corresponding $\mathfrak{sl}(n)$ -bundle does not have Lie algebra automorphisms. Therefore, the next lemma finishes the proof of Proposition 6.3.

Lemma 6.4: Let M be a compact complex manifold, P a projective bundle, and G_P the corresponding principal $PGL(n)$ -bundle. Consider the Lie algebra bundle $\mathfrak{g}_P$ associated with G_P . Assume that $\mathfrak{g}_P$ has no Lie algebra automorphisms. Then the fibration P has no automorphisms over M .

Proof: Let $\sigma : P \rightarrow P$ be a non-trivial automorphism. The group $PGL(n)$ has zero center. Therefore, σ acts non-trivially on the Lie algebra of infinitesimal automorphisms of P . This contradicts the simplicity

¹Please notice, however, that the existence of the coarse moduli is itself quite non-trivial.

assumption (Subsection 6.1).

We proved Proposition 6.3. ■

Let B be a vector bundle over a compact complex manifold M , and S its deformation space. The universal bundle $\mathcal{U}$ does not always exist. However, the universal bundle, corresponding to the projectivization of B , exists by Proposition 6.3. Denote by $\text{End}_{tr}(\mathcal{U})$ the Lie algebra of the corresponding $PGL(n)$ -bundle.

Definition 6.5: The bundle $\text{End}_{tr}(\mathcal{U})$ is called **the bundle of traceless endomorphisms of the universal bundle**.

Notice that $\text{End}_{tr}(\mathcal{U})$ is well defined even when $\mathcal{U}$ is not defined.

7 Fourier-Mukai transform

Let M be a compact complex manifold, B a vector bundle, and S the coarse deformation space of B . If the universal bundle exists, we denote it by $\mathcal{U}$. The bundle $\text{End}_{tr}(\mathcal{U})$ of traceless endomorphisms of the universal bundle (Definition 6.5) is well-defined, regardless of the existence of $\mathcal{U}$. Denote the projections from $M \times S$ to M , S by π_1 , π_2

Definition 7.1: In the above assumptions, let $D_b(M)$, $D_b(S)$ be the derived category of bounded complexes of coherent sheaves on M , S . **Fourier-Mukai transform** is a functor from $D_b(M)$ to $D_b(S)$, which is defined as follows. Given an object A of $D_b(M)$, we take its pullback π_1^*A to $M \times S$, tensor it with $\mathcal{U}$ and take the pushforward to S :

$$FM(A) = R^\bullet(\pi_2)_* \left(\pi_1^*A \otimes \mathcal{U} \right)$$

However, it is not defined when $\mathcal{U}$ does not exist. It is convenient to define **the modified Fourier-Mukai transform**:

$$FM_1(A) = R^\bullet(\pi_2)_* \left(\pi_1^*A \otimes \text{End}_{tr} \mathcal{U} \right)$$

which exists always.

Given a coherent sheaf A , the Fourier-Mukai transform of A is no longer a sheaf, but a complex of sheaves. However, its cohomology are coherent

sheaves on S . Thus, the Fourier-Mukai functor gives a rise to a series of functors on the corresponding categories of coherent sheaves:

$$A \longrightarrow H^i(FM(A)).$$

The main result of this paper is the following theorem (stability¹ of Mukai transform).

Theorem 7.2: Let M be a K3 surface, and B a stable bundle on M . Denote by S the coarse moduli of stable deformations of B , equipped with the natural (Weil-Peterson) Kähler metrics. Let A be an arbitrary stable bundle on M . Consider the modified Fourier-Mukai transform $FM_1(A)$, and the corresponding cohomology sheaves $H^i(FM_1(A))$. Assume that S is compact. Then the cohomology sheaf $H^i(FM_1(A))$ is (up to torsion) polystable², for all i . Moreover, the cohomology sheaves of $FM(A)$ are also polystable, assuming that the universal bundle $\mathcal{U}$ and the Fourier-Mukai transform is defined.

Proof: The proof of Theorem 7.2 is done in two major steps. First, we show that the universal bundle is projectively hyperholomorphic. Second, we prove that a pushforward of a projectively hyperholomorphic bundle is a complex of sheaves with polystable cohomology.

The K3 surface is hyperkähler. Fix a hyperkähler structure which is compatible with the Kähler structure on M (Definition 2.2). Then the bundles A and B are equipped with a unique projectively hyperholomorphic Yang-Mills connection (Claim 4.10). The following result is implied by the general machinery of hyperholomorphic bundles.

Theorem 7.3: Let M be a compact hyperkähler manifold, I an induced complex structure, and B a projectively hyperholomorphic bundle on (M, I) . Consider the coarse moduli S of the stable deformations of B . By [V1], Theorem 11.3 (see also Subsection 4.3), the variety S has a natural hyperkähler structure. Let $\text{End}_{tr}(\mathcal{U})$ be the sheaf of traceless endomorphisms of the universal bundle. Then $\text{End}_{tr}(\mathcal{U})$ is a hyperholomorphic bundle over a hyperkähler manifold $M \times S$.

¹Throughout this paper, stability is understood in the sense of Mumford-Takemoto, see Definition 3.4.

²“Polystable” means “a direct sum of stable sheaves of the same slope”, see Definition 3.4.

Proof: Consider the projective bundle $P = \mathbb{P}B$, obtained as a projectivization of B . Let G_P be the corresponding principal $PGL(n)$ -bundle. Using an appropriate definition of stability, one can define the coarse moduli S_P of stable deformations of G_P . There is a natural map

$$S \xrightarrow{\varphi} S_P$$

corresponding to taking a projectivization of a vector bundle. Using results of [V1] it is easy to check that S_P is hyperkähler, and, moreover, the map φ is compatible with the hyperkähler structure.

Let $\mathcal{U}_P$ be the universal bundle on $M \times S_P$, which exists by Proposition 6.3, and $\mathfrak{pgl}(\mathcal{U}_P)$ the corresponding Lie algebra bundle. From the definition of $\mathrm{End}_{tr} \mathcal{U}$ it follows that the sheaf $\mathrm{End}_{tr} \mathcal{U}$ is canonically isomorphic to the pullback $\varphi^* \mathfrak{pgl}(\mathcal{U}_P)$. To show that $\mathrm{End}_{tr} \mathcal{U}$ is hyperholomorphic, we need only to prove that $\mathfrak{pgl}(\mathcal{U}_P)$ is hyperholomorphic. This is a consequence of the following general statement.

Proposition 7.4: Let M be a compact hyperkähler manifold, I an induced complex structure, G a compact Lie group, and P a hyperholomorphic G -bundle. Denote the coarse moduli space of P by S . Assume that the universal bundle $\mathcal{U}$ exists. Then $\mathcal{U}$ is hyperholomorphic.

Proof: Let $s \in S$ be a point. Locally in a neighbourhood S_0 of s , there exists a universal bundle $\mathcal{U}_0$ on $M \times S_0$, which is constructed explicitly using Hodge theory ([V1], Section 7). Consider the natural (Peterson-Weil) metrics on $\mathcal{U}_0$. From the explicit construction of $\mathcal{U}_0$ it immediately follows that $\mathcal{U}_0$ is hyperholomorphic (in fact, $\mathcal{U}_0$ can be constructed without mentioning the induced complex structure I at all; hence, $\mathcal{U}_0$ is holomorphic with respect to all induced complex structures on $M \times S_0$; then, the bundle $\mathcal{U}_0$ is holomorphic in each of these complex structures, and therefore hyperholomorphic).

This proves Proposition 7.4 and Theorem 7.3. ■

Return to the proof of Theorem 7.2. By definition of Fourier-Mukai transform,

$$H^i(FM(A)) = R^i(\pi_2)_*(\pi_1^* A \otimes \mathcal{U})$$

and

$$H^i(FM_1(A)) = R^i(\pi_2)_*(\pi_1^* A \otimes \mathrm{End}_{tr} \mathcal{U}).$$

Now, $\pi_1^* A$ projectively hyperholomorphic as a pullback of a projectively hyperholomorphic bundle, and $\mathcal{U}$ is projectively hyperholomorphic by Theorem 7.3. The tensor product of projectively hyperholomorphic sheaves is projectively hyperholomorphic. Therefore, the sheaf $H^i(FM(A))$ is a pushforward of a projectively hyperholomorphic sheaf; the same is true for $H^i(FM_1(A))$. To prove Theorem 7.2 it remains to show that a pushforward of a projectively hyperholomorphic sheaf is polystable. By Theorem 5.14 for this purpose it is sufficient to prove that a higher direct image of a projectively hyperholomorphic sheaf is projectively hyperholomorphic, outside of singularities.

The following theorem concludes the proof of Theorem 7.2.

Theorem 7.5: Let $M \times S$ be a product of compact hyperkähler manifolds, I an induced complex structure, and B a projectively hyperholomorphic bundle on $(M \times S, I)$, which is also Yang-Mills. Denote by π the projection of $M \times S$ to S . Assume that either $H^1(S) = 0$ or $H^1(M) = 0$. Let $R := R^i \pi_* B$ be the higher direct image of B , $\mathcal{T} \subset R$ its torsion component, and $R/\mathcal{T}$ the torsion-free part. Then the sheaf $R/\mathcal{T}$ is polystable.

Proof: See Subsection 12.2. ■

8 Quaternionic Dolbeault complex

The quaternionic cohomology is a well known subject, introduced by M. Capria and S. Salamon ([CS], [S], [B], [L]). Here we give an exposition of quaternionic cohomology and quaternionic Dolbeault complex for hyperkähler manifolds. However, the generalization of all these subjects to quaternionic and quaternionic-Kähler manifolds is transparent.

8.1 Quaternionic Dolbeault complex: the definition

Let M be a hyperkähler manifold, and

$$\Lambda^0 M \xrightarrow{d} \Lambda^1 M \xrightarrow{d} \Lambda^2 M \xrightarrow{d} \dots$$

its de Rham complex. Consider the natural action of $SU(2)$ on $\Lambda^* M$ (see Section 2). Clearly, $SU(2)$ acts on $\Lambda^i M$, $i \leq \frac{1}{2} \dim_{\mathbb{R}} M$ with weights $i, i-2, i-4, \dots$

We denote by Λ_+^i the maximal $SU(2)$ -subspace of Λ^i , on which $SU(2)$ acts with weight i .

The following linear algebraic lemma allows one to compute Λ_+^i explicitly

Lemma 8.1: In the above assumptions, let I be an induced complex structure, and $\mathbb{H}_I$ the quaternion space, considered as a 2-dimensional complex vector space with the complex structure induced by I . Denote by $\Lambda_I^{p,0}(M)$ the space of $(p,0)$ -form on M , with respect to the Hodge decomposition associated with the complex structure I . The space $\mathbb{H}_I$ is equipped with a natural action of $SU(2)$. Consider $\Lambda_I^{p,0}(M)$ as a representation of $SU(2)$, with trivial group action. Then, there is a canonical isomorphism

$$\Lambda_+^p(M) \cong S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M), \quad (8.1)$$

where $S_{\mathbb{C}}^p \mathbb{H}_I$ denotes a p -th symmetric power of $\mathbb{H}_I$. Moreover, the $SU(2)$ -action on $\Lambda_+^p(M)$ is compatible with the isomorphism (8.1).

Proof: Fix a standard basis $1, I, J, K$ in $\mathbb{H}$. Let $h_1, h_2 \in \mathbb{H}_I$ be the corresponding basis in $\mathbb{H}_I$, $h_1 = 1 + \sqrt{-1}I$, $h_2 = J - \sqrt{-1}K$. Consider the $SU(2)$ -invariant homomorphism

$$\mathbb{H}_I \otimes \Lambda_I^{1,0}(M) \longrightarrow \Lambda^1(M) \quad (8.2)$$

mapping $h_1 \otimes \eta$ to η and $h_2 \otimes \eta$ to $J(\eta)$, where J denotes an endomorphism of $\Lambda^1(M)$ induced by J . The equation (8.1) is obvious for $p = 1$:

$$\Lambda^1(M) = \Lambda_+^1(M) = \mathbb{H}_I \otimes \Lambda_I^{1,0}(M) \quad (8.3)$$

This isomorphism is by construction $SU(2)$ -invariant. Given two vector spaces A and B , we have a natural map

$$S^i A \otimes \Lambda^i B \longrightarrow \Lambda^i(A \otimes B). \quad (8.4)$$

From (8.4) and (8.3), we obtain a natural $SU(2)$ -invariant homomorphism

$$S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M) \longrightarrow \Lambda_+^p(M). \quad (8.5)$$

Since $S_{\mathbb{C}}^p \mathbb{H}_I$ has weight p , the arrow (8.5) maps $S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M)$ to $\Lambda_+^p(M)$. We have constructed a map

$$S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M) \xrightarrow{\Psi} \Lambda_+^p(M). \quad (8.6)$$

It remains to show that it is an isomorphism. Let $adI : \Lambda^* M \longrightarrow \Lambda^* M$ act on (p,q) -forms $ad(\eta) = (p - q)\sqrt{-1} \eta$. Clearly, $-\sqrt{-1} adI$ is a root

of the Lie algebra $SU(2)$. It is well known that a representation of a Lie algebra is generated by the highest vector. For the Lie algebra $\mathfrak{su}(2)$, this means that $\Lambda_+^p(M)$ is a subspace of $\Lambda^p(M)$ generated by $SU(2)$ from the subspace $W \subset \Lambda_+^p(M)$ consisting of all vectors on which $-\sqrt{-1} adI$ acts as a multiplication by p . On the other hand, W coincides with $\Lambda_I^{p,0}(M)$. We obtained the following

$$\begin{aligned} \text{The space } \Lambda_+^p(M) \text{ is generated by } SU(2) \text{ from its subspace} \\ \Lambda_I^{p,0}(M). \end{aligned} \quad (8.7)$$

The image of

$$\Psi : S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M) \longrightarrow \Lambda_+^p(M)$$

is an $SU(2)$ -invariant subspace of $\Lambda^p(M)$ containing $\Lambda_I^{p,0}(M)$. By (8.7), this means that Ψ is surjective.

Let $R \subset S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M)$ be the kernel of Ψ . By construction, R is $SU(2)$ -invariant, of weight p . By the same arguments as above, R is generated by its subspace of highest weight, that is, the vectors of type $h_1^p \eta$, where $\eta \in \Lambda_I^{p,0}(M)$ (see (8.2)). On the other hand, on the subspace

$$h_1^p \cdot \Lambda_I^{p,0}(M) \subset S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M)$$

the map Ψ is by construction injective. Therefore, the intersection $h_1^p \cdot \Lambda_I^{p,0}(M) \cap R$ is empty. We proved that Ψ is an isomorphism. Lemma 8.1 is proven. ■

Consider an $SU(2)$ -invariant decomposition

$$\Lambda^p(M) = \Lambda_+^p(M) \oplus V^p, \quad (8.8)$$

where V^p is the sum of all $SU(2)$ -subspaces of $\Lambda^p(M)$ of weight less than p . Since $SU(2)$ -action is multiplicative, the subspace $\oplus_p V^p \subset \Lambda^p(M)$ is an ideal. Therefore, the quotient

$$\Lambda_+^*(M) = \Lambda^*(M) / \oplus_p V^p$$

is an algebra. Using the decomposition (8.8), we define the quaternionic Dolbeault differential $d_+ : \Lambda_+^*(M) \longrightarrow \Lambda_+^*(M)$ as the composition of de Rham differential and projection of to $\Lambda_+^*(M) \subset \Lambda^*(M)$. Since de Rham differential cannot increase the $SU(2)$ -weight of a form more than by 1,

d preserves the subspace $V^* \subset \Lambda^*(M)$. Therefore, d_+ is a differential in $\Lambda_+^*(M)$.

Definition 8.2: Let

$$\Lambda^0 M \xrightarrow{d_+} \Lambda^1 M \xrightarrow{d_+} \Lambda_+^2 M \xrightarrow{d_+} \Lambda_+^3 M \xrightarrow{d_+} \dots$$

be the differential graded algebra constructed above¹. It is called **the quaternionic Dolbeault complex**, or **Salamon complex**.

Remark 8.3: The isomorphism (8.1) is clearly multiplicative.

8.2 Connections in modules over DG-algebras

Using the quaternionic Dolbeault complex, it is possible to give an alternative definition for hyperholomorphic bundles, in the same way as one defines the holomorphic bundles using the usual Dolbeault complex (Theorem 3.1).

We state the definition of a connection and flat connection for a general differential graded algebra; flat bundles, holomorphic bundles and hyperholomorphic bundles become special cases of this algebraic definition.

Throughout this paper, the differential graded algebras are always assumed to be supercommutative.

Definition 8.4: Let

$$A^0 \xrightarrow{d} A^1 \xrightarrow{d} A^2 \xrightarrow{d} \dots$$

be a differential graded supercommutative algebra, and B an A^0 -module. Consider an operator

$$\nabla : B \longrightarrow B \otimes_{A^0} A^1$$

satisfying the Leibniz identity:

$$\nabla(ab) = da \otimes b + a\nabla b, \tag{8.9}$$

($a \in A^0$, $b \in B$). Then ∇ is called (A^*, d) -connection.

¹We identify $\Lambda^0 M$ and $\Lambda_+^0 M$, $\Lambda^1 M$ and $\Lambda_+^1 M$.

The following claim is trivial

Claim 8.5: Let A^* be a differential graded algebra, which is multiplicatively generated by A_0, A_1 , and (B, ∇) an A^0 -module with a connection over A^* . Then ∇ can be uniquely extended to a sequence of operators

$$B \xrightarrow{\nabla} B \otimes_{A^0} A^1 \xrightarrow{\nabla} B \otimes_{A^0} A^2 \xrightarrow{\nabla} B \otimes_{A^0} A^3 \xrightarrow{\nabla} \dots \quad (8.10)$$

which satisfies the Leibniz identity.

Proof: Clear. ■

A sequence (8.10) is called **an extended connection** over A^* . Further on, we shall always work with the differential graded algebras, which are multiplicatively generated by A_0, A_1 (such as the Dolbeault complex, de Rham complex, quaternionic Dolbeault complex, etc.) For such algebras, we do not always make a distinction between a connection and an extended connection.

Definition 8.6: Let A^*, d be a differential graded algebra, B an A_0 -module, and

$$B \xrightarrow{\nabla} B \otimes_{A^0} A^1 \xrightarrow{\nabla} B \otimes_{A^0} A^2 \xrightarrow{\nabla} B \otimes_{A^0} A^3 \xrightarrow{\nabla} \dots \quad (8.11)$$

an extended connection on B . We say that ∇ is **flat** if $\nabla^2 = 0$, where

$$\nabla^2 : B \otimes_{A^0} A^i \longrightarrow B \otimes_{A^0} A^{i+2}$$

is a composition of two successive arrows in (8.11). We say that ∇ is **projectively flat** if for all $b \in B$, we have

$$\nabla^2 B = ab,$$

where $a \in A^2$.

Example 8.7: Let (A^*, d) be de Rham algebra over M , and B a C^∞ -vector bundle. There is a natural action of A^0 on B . Clearly, an A^* -connection on B is simply a connection on B . An A^* -connection is flat (projectively flat) if and only if B is flat (resp. projectively flat).

Example 8.8: Let (A^*, d) be the Dolbeault complex:

$$(A^*, D) = (\Lambda^{0,*}, \bar{\partial}),$$

and B a complex vector C^∞ -bundle. The Newlander-Nierenberg theorem (Theorem 3.1) states that flat (A^*, d) -connections are in one to one correspondence with the holomorphic structures on B .

The projective flatness of a connection can be dealt with in the same fashion as we dealt with the projectively hyperholomorphic bundles in Proposition 4.8.

Claim 8.9: Let (A^*, d) be a differential graded algebra, and B an A_0 -module with a connection ∇ . Assume that B is projective and finite generated² over A_0 . Then the connection in B is projectively flat over A^* if and only if the associated connection in $\text{End}_{A_0} B$ is flat over A^* .

Proof: Claim 8.9 is a straightforward generalization of Proposition 4.8, and its proof is essentially the same. Let $\Theta : A^* \otimes_{A_0} B \longrightarrow A^{*+2} \otimes_{A_0} B$ be the curvature of ∇ . From the Leibniz identity it immediately follows that Θ is A^0 -linear, that is, satisfies

$$\Theta(ab) = a\Theta(b)$$

for $a \in A^*$, $b \in B$. Therefore, Θ is determined by the value it takes on B . We may interpret Θ as an A_0 -linear homomorphism

$$\Theta : B \longrightarrow A^2 \otimes_{A_0} B.$$

In other words, Θ can be considered as an element in $A^2 \otimes_{A_0} \text{End}(B)$. Clearly, B is projectively flat if and only if Θ belongs to $A^2 \subset A^2 \otimes_{A_0} \text{End}(B)$, where $A^2 \hookrightarrow A^2 \otimes_{A_0} \text{End}(B)$ is the canonical embedding corresponding to the identity section of $\text{End} B$.

Now, the curvature Θ' of $\text{End} B$ is expressed through Θ by the relation (4.2). Using this relation just as in the proof of Proposition 4.8, we obtain that Θ belongs to $A^2 \subset A^2 \otimes_{A_0} \text{End}(B)$ if and only if Θ' is zero. This proves Claim 8.9. ■

Our main working example comes from the hyperkähler geometry. The following claim is clear.

²This assumption is clear when A_0 is $C^\infty(M)$ and B a (finite-dimensional) bundle over M , for M any smooth manifold.

Claim 8.10: Let M be a hyperkähler manifold and B a Hermitian vector bundle with a connection

$$B \xrightarrow{\nabla} B \otimes \Lambda^1 M.$$

Using the identification $\Lambda^1(M) \cong \Lambda_+^1(M)$, we may consider ∇ as a (Λ_+^*, d_+) -connection, where (Λ_+^*, d_+) is the quaternionic Dolbeault complex. Then ∇ is flat (resp. projectively flat) over (Λ_+^*, d_+) if and only if ∇ is hyperholomorphic (resp. projectively hyperholomorphic).

Proof: Claim 8.10 is obvious from the definitions. To avoid misunderstanding, we give its proof, but for most readers it will be easier to invent a proof of their own.

Extend ∇ to a sequence

$$B \xrightarrow{\nabla} B \otimes \Lambda_+^1 \xrightarrow{\nabla} B \otimes \Lambda_+^2 \xrightarrow{\nabla} B \otimes \Lambda_+^3 \xrightarrow{\nabla} \dots \quad (8.12)$$

(Claim 8.5). Then ∇^2 maps a form η to $\Pi(\eta \wedge \Theta)$, where $\Theta \in \Lambda^2(M) \otimes \text{End}(B)$ is the curvature of the connection $\nabla : B \rightarrow B \otimes \Lambda^1 M$, and Π a standard projection of $\Lambda^2(M)$ to $\Lambda_+^2(M)$.

Consider the standard action of $SU(2)$ on differential forms. We say that a form κ is pure of weight p if κ belongs to an $SU(2)$ -representation of weight p .

If Θ is $SU(2)$ -invariant, and η is pure of weight p , then $\eta \wedge \Theta$ also has weight p and belongs to the kernel of Π ; this means that ∇ is flat over $\Lambda_+^*(M)$. Conversely, if ∇ is flat over $\Lambda_+^*(M)$, then Θ is a sum of forms which are of weight less than 2; however, Θ is a 2-form, and 2-forms can only have weights 2 and 0. Therefore, the curvature Θ is pure of weight 0, that is, $SU(2)$ -invariant.

Applying this argument to the bundle $\text{End } B$, we obtain that B is projectively hyperholomorphic if and only if $\text{End } B$ is flat over $\Lambda_+^*(M)$. By Claim 8.9, this is equivalent to B being projectively flat over $\Lambda_+^*(M)$.

Claim 8.10 is proven. ■

8.3 Hodge decomposition for the quaternionic Dolbeault complex

Let M be a hyperkähler manifold, and I an induced complex structure. As usually, we define an operator $adI : \Lambda^*(M) \rightarrow \Lambda^*(M)$ mapping a (p, q) -form η to $\sqrt{-1}(p - q)\eta$. By definition, adI belongs to the Lie algebra $\mathfrak{su}(2)$

acting on $\Lambda^*(M)$ in the standard way. Therefore, adI preserves the subspace $\Lambda_+^*(M) \subset \Lambda^*(M)$. We obtain the Hodge decomposition

$$\Lambda_+^*(M) = \oplus_{p,q} \Lambda_{+,I}^{p,q}(M).$$

Definition 8.11: The decomposition

$$\Lambda_+^*(M) = \oplus_{p,q} \Lambda_{+,I}^{p,q}(M)$$

is called **the Hodge decomposition for the quaternionic Dolbeault complex**.

The following claim is trivial

Claim 8.12: Given a hyperkähler manifold M and an induced complex structure I , the following subspaces of $\Lambda^*(M)$ coincide:

$$\Lambda_{+,I}^{p,0}(M) = \Lambda_I^{p,0}(M),$$

where $\Lambda_I^{p,0}(M)$ denotes the space of all $(p, 0)$ -forms.

Proof: Immediately follows from Lemma 8.1. ■

8.4 The Dolbeault bicomplex and quaternionic Dolbeault complex

Let M be a hyperkähler manifold, I an induced complex structure, and $I, J, K \in \mathbb{H}$ the standard triple of induced complex structures. Clearly, J acts on the complexified co tangent space $\Lambda^1 M \otimes \mathbb{C}$ mapping $\Lambda_I^{0,1}(M)$ to $\Lambda_I^{1,0}(M)$. Consider a differential operator

$$\partial_J : C^\infty(M) \longrightarrow \Lambda_I^{1,0}(M),$$

mapping f to $J(\bar{\partial}f)$, where $\bar{\partial} : C^\infty(M) \longrightarrow \Lambda_I^{0,1}(M)$ is the standard Dolbeault differential on a Kähler manifold (M, I) . We extend ∂_J to a differential

$$\partial_J : \Lambda_I^{p,0}(M) \longrightarrow \Lambda_I^{p+1,0}(M),$$

using the Leibniz rule.

Let B be a bundle with connection. In the same fashion as above, we define the operator

$$\nabla'_J : \Lambda_I^{p,0}(M) \otimes B \longrightarrow \Lambda_I^{p+1,0}(M) \otimes B, \quad \nabla'_J = J \circ \nabla'' \circ J^{-1}; \quad (8.13)$$

this operator is equal to ∂_J when B is a trivial bundle.

Proposition 8.13: Let M be a hyperkähler manifold, I an induced complex structure, I, J, K the standard basis in quaternion algebra, and

$$\Lambda_+^*(M) = \oplus_{p,q} \Lambda_{I,+}^{p,q}(M)$$

the Hodge decomposition of the quaternionic Dolbeault complex (Subsection 8.3). Then there exists a canonical isomorphism

$$\Lambda_{I,+}^{p,q}(M) \cong \Lambda_I^{p+q,0}(M). \quad (8.14)$$

Under this identification, the quaternionic Dolbeault differential

$$d_+ : \Lambda_{I,+}^{p,q}(M) \longrightarrow \Lambda_{I,+}^{p+1,q}(M) \oplus \Lambda_{I,+}^{p,q+1}(M)$$

corresponds to a sum

$$\partial \oplus \partial_J : \Lambda_I^{p+q,0}(M) \longrightarrow \Lambda_I^{p+q+1,0}(M) \oplus \Lambda_I^{p+q+1,0}(M).$$

Proof: Consider the isomorphism

$$\Lambda_+^p \cong S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M). \quad (8.15)$$

The Hodge decomposition of (8.15) is induced by the $SU(2)$ -action, as follows. Let $\rho_I : U(1) \longrightarrow SU(2)$ be a group map corresponding to the Lie algebra embedding $\mathfrak{u}(1) = \mathbb{R} \hookrightarrow \mathfrak{su}(2)$, $1 \longrightarrow I$. From the definition of the $SU(2)$ -action it follows that the Hodge decomposition of $\Lambda^*(M)$ coincides with the weight decomposition under the action of $\rho_I : U(1) \longrightarrow \text{End}(\Lambda^*(M))$.

The $SU(2)$ -action on $S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M)$ is trivial on the second component. Consider the weight decomposition

$$S_{\mathbb{C}}^i \mathbb{H}_I \cong \bigoplus_{p+q=i} S_{\mathbb{C}}^{p,q} \mathbb{H}_I$$

associated with ρ_I . Then (8.15) translates to the isomorphism

$$\Lambda_{I,+}^{p,q}(M) \cong S_{\mathbb{C}}^{p,q} \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p+q,0}(M).$$

Let h_1, h_2 be a basis in $\mathbb{H}_I$ defined as in the proof of Lemma 8.1. An elementary calculation shows that h_1 has weight $(1,0)$, and h_2 has weight $(0,1)$. Therefore, the space $S_{\mathbb{C}}^{p,q}\mathbb{H}_I$ is 1-dimensional and generated by $h_1^p h_2^q$. We obtained an isomorphism

$$\Lambda_{I,+}^{p,q}(M) \cong h_1^p \cdot h_2^q \cdot \Lambda_I^{p+q,0}(M). \quad (8.16)$$

This proves (8.14).

The isomorphism (8.16) is clearly multiplicative. Consider the differential

$$\hat{d}_+ = h_1 \partial + h_2 \partial_J : S_{\mathbb{C}}^p \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p,0}(M) \longrightarrow S_{\mathbb{C}}^{p+1} \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{p+1,0}(M)$$

To prove Proposition 8.13, we need to show that the quaternionic Dolbeault differential d_+ coincides with $\hat{d}_+$ under the identification (8.16). The isomorphism (8.16) is by construction multiplicative, and the differentials d_+ and $\hat{d}_+$ both satisfy the Leibniz rule. Therefore, it is sufficient to show that

$$d_+ = \hat{d}_+ \quad (8.17)$$

on $C^\infty(M) = \Lambda_+^0(M)$.

On functions, the equality (8.17) is immediately implied by the definition of the isomorphism

$$\Lambda_+^1(M) \cong \mathbb{H}_I \otimes_{\mathbb{C}} \Lambda_I^{1,0}(M).$$

Proposition 8.13 is proven. ■

Remark 8.14: A similar result is true for bundles with connection. Given a vector bundle B with a connection ∇ , the $\Lambda_+^*(M)$ -connection operator

$$\nabla_+ : \Lambda_{I,+}^{p,q}(M) \otimes B \longrightarrow \Lambda_{I,+}^{p+1,q}(M) \otimes B \oplus \Lambda_{I,+}^{p,q+1}(M) \otimes B$$

(Claim 8.10) corresponds to a sum

$$\nabla' \oplus \nabla'_J : \Lambda_I^{p+q,0}(M) \otimes B \longrightarrow \Lambda_I^{p+q+1,0}(M) \otimes B \oplus \Lambda_I^{p+q+1,0}(M) \otimes B.$$

where ∇'_J is defined in the same way as ∂_J (8.13). The proof of this statement is the same as the proof of Proposition 8.13.

The statement of Proposition 8.13 can be represented by the following diagram

$$\begin{array}{ccc}
\Lambda_+^0(M) & & \Lambda_I^{0,0}(M) \\
\swarrow d'_+ \quad \searrow d''_+ & & \swarrow h_1 \partial \quad \searrow h_2 \partial_J \\
\Lambda_+^{1,0}(M) \quad \Lambda_+^{0,1}(M) & \cong & h_1 \Lambda_I^{1,0}(M) \quad h_2 \Lambda_I^{1,0}(M) \\
\swarrow d'_+ \quad \searrow d''_+ \quad \swarrow d'_+ \quad \searrow d''_+ & & \swarrow h_1 \partial \quad \searrow h_2 \partial_J \quad \swarrow h_1 \partial \quad \searrow h_2 \partial_J \\
\Lambda_+^{2,0}(M) \quad \Lambda_+^{1,1}(M) \quad \Lambda_+^{0,2}(M) & & h_1^2 \Lambda_I^{2,0}(M) \quad h_1 h_2 \Lambda_I^{2,0}(M) \quad h_2^2 \Lambda_I^{2,0}(M)
\end{array} \tag{8.18}$$

where $d_+ = d'_+ + d''_+$ is the Hodge decomposition of the quaternionic Dolbeault differential.

Definition 8.15: In the above assumptions, the bicomplex (8.18) is called **the quaternionic Dolbeault bicomplex**.

Applying the above argument to the quaternionic Dolbeault complex with coefficients in a bundle, we obtain the following claim.

Claim 8.16: In assumptions of Proposition 8.13, let B be a bundle with connection over M , and

$$\nabla_+ : \Lambda_+^*(M) \otimes B \longrightarrow \Lambda_+^{*+1}(M) \otimes B$$

the corresponding connection over $\Lambda_+^*(M)$. Consider the operators

$$\nabla', \quad \nabla'_J : \Lambda_+^{*,0}(M) \otimes B \longrightarrow \Lambda_+^{*+1,0}(M) \otimes B.$$

(Remark 8.14). Then B is hyperholomorphic if and only if ∇' commutes with ∇'_J

Proof: Follows directly from Remark 8.14, the isomorphism (8.18) and Claim 8.10. ■

We do not use Claim 8.16 anywhere in this paper.

9 qD-modules and coherent sheaves

9.1 qD-modules: the definition

The following generalization of a notion of a hyperholomorphic bundle is useful in the study of coherent sheaves over hyperkähler manifolds.

Definition 9.1: Let M be a hyperkähler manifold, F a $C^\infty(M)$ -module (not necessarily finitely generated) and

$$F \xrightarrow{\nabla} F \otimes_{C^\infty(M)} \Lambda_+^1(M) \xrightarrow{\nabla} F \otimes_{C^\infty(M)} \Lambda_+^2(M) \dots$$

a quaternionic Dolbeault connection. Then F is called a **quaternionic Dolbeault module**, or **qD-module** for short, if ∇ is flat.

Let I be an induced complex structure. Consider the operator $L : \Lambda_+^i(M) \rightarrow \Lambda_+^{i+2}(M)$, mapping η to $\eta \wedge \omega$, where ω is the Kaehler form on (M, I) . By $\Lambda : \Lambda_+^i(M) \rightarrow \Lambda_+^{i-2}(M)$ we denote the Hermitian adjoint operator. These operators can be obtained from the usual Hodge operators L, Λ by taking an orthogonal projection to $\Lambda_+^*(M)$.

Definition 9.2: Let I be an induced complex structure, and F a $\Lambda_+^*(M)$ -module with a connection ∇ . Assume that the curvature of ∇ is of Hodge type $(1, 1)$. If ∇ is projectively flat, and its curvature Θ satisfies

$$\Lambda(\Theta) = cId,$$

then F is called a **projective quaternionic Dolbeault Yang-Mills module**, pqDYM-module for short.

This term is coined in analogy to the usual D-modules; in fact, any D-module is automatically a qD-module. The converse is not true, of course.

Definition 9.3: Let M be a hyperkähler manifold, I an induced complex structure, and F a qD-module or a pqDYM-module. Consider the $(0, 1)$ -part ∇'' of the connection ∇ :

$$\nabla'' : F \rightarrow F \otimes_{C^\infty(M)} \Lambda_I^{0,1}(M).$$

Let $\mathcal{H}_I(F)$ be the sheaf of all sections $f \in F$ satisfying $\nabla'' f = 0$.

Clearly, $\mathcal{H}_I(F)$ is a $\mathcal{O}_{(M, I)}$ -module. We call $\mathcal{H}_I(F)$ a **canonical quasi-coherent sheaf on (M, I) associated with F** . When F is a hyperholomorphic bundle, $\mathcal{H}_I(F)$ is the corresponding holomorphic bundle on (M, I) .

Definition 9.4: Let M be a hyperkähler manifold, I an induced complex structure, and $\mathcal{F}$ a quasicoherent sheaf on (M, I) . A **qD- or pqDYM-structure on $\mathcal{F}$** is a qD- or pqDYM-module F equipped with an isomorphism $\mathcal{H}_I(F) \cong \mathcal{F}$.

The main result of this section is the following proposition

Proposition 9.5: Let M be a hyperkähler manifold, I an induced complex structure, and $\mathcal{F}$ a coherent sheaf on (M, I) equipped with a qD- or pqDYM-structure. Then the following claims are true

- (i) The torsion subsheaf $\mathcal{T} \subset \mathcal{F}$, and the quotient $\mathcal{F}/\mathcal{T}$ is also equipped with a qD- (respectively, pqDYM-) structure.
- (ii) Let $Z \subset (M, I)$, $\text{codim}_{\mathbb{C}} \geq 2$ be a closed complex subvariety of (M, I) , $U := M \setminus Z$ the corresponding open submanifold, and $j : U \hookrightarrow M$ the natural open embedding. Consider the quasicoherent sheaf $j_*j^*\mathcal{F}$. Then the sheaf $j_*j^*\mathcal{F}$ is also equipped with a qD- (respectively, pqDYM-) structure.

Proof: Let F be a qD- or pqDYM-module such that $\mathcal{H}_I(F) = \mathcal{F}$, and $F_0 \subset F$ the torsion subsheaf of F , that is, the union of all sections of F which vanish on a dense open subset of M . Clearly, $\mathcal{H}_I(F_0) = \mathcal{T}$. Therefore, $\mathcal{T} \subset \mathcal{F}$ is compatible with the qD- or pqDYM-structure on $\mathcal{F}$. This proves Proposition 9.5 (i).

To prove Proposition 9.5 (ii), consider the sheaf j_*j^*F . By definition, for any open subset $W \subset M$, we have

$$\Gamma_W(j_*j^*F) := \Gamma_{W \cap U} F.$$

Clearly, j_*j^*F is a qD- or pqDYM-module. Moreover, $\mathcal{H}_I(j_*j^*F) = j_*j^*\mathcal{F}$. This proves Proposition 9.5 (ii). ■

Corollary 9.6: Let M be a hyperkähler manifold, I an induced complex structure, and $\mathcal{F}$ a torsion-free coherent sheaf on (M, I) equipped with a qD- or pqDYM-structure. Then the reflexive hull $\mathcal{F}^{**}$ of $\mathcal{F}$ is also equipped with a qD- (respectively, pqDYM-) structure.

Proof: In assumptions of Proposition 9.5, let Z be the singular set of $\mathcal{F}$. Then $j_*j^*\mathcal{F} \cong \mathcal{F}^{**}$ (Lemma 5.3). Corollary 9.6 is implied immediately by Proposition 9.5. ■

Remark 9.7: Let M, M' be compact hyperkähler manifolds, $H^1(M') = 0$, I an induced complex structure and $\pi : M \times M' \rightarrow M$ the standard projection. Consider a hyperholomorphic or projectively hyperholomorphic bundle B on $(M \times M', I)$. In Section 12, we prove that the higher direct image $\mathcal{F} := R^i \pi_* B$ is equipped with a qD- or pqDYM-structure. Let $\mathcal{T}$ be the torsion part of $\mathcal{F}$. By Proposition 9.5 (i), the quotient $\mathcal{F}/\mathcal{T}$ is equipped with a qD- or pqDYM-structure. By Corollary 9.6, the reflexive hull F of $\mathcal{F}/\mathcal{T}$ is also equipped with a qD- or pqDYM-structure. We arrive in the situation described by Theorem 5.14: $\text{End}(F)$ is a reflexive sheaf equipped with a hyperholomorphic connection outside of its singularities. By Theorem 5.14, $\text{End}(F)$ is polystable. It is easy to check that the polystability of $\text{End}(F)$ implies polystability of F .

10 qD-Laplace operator and the extended qD-complex

10.1 Extended quaternionic Dolbeault bicomplex

To define the quaternionic Dolbeault Laplace operator, we need to extend the qD-bicomplex (8.18) as follows.

Definition 10.1: Let M be a hyperkähler manifold and I an induced complex structure. Consider the bicomplex

$$\bigoplus_{p,q \in \mathbb{Z}} \left(\tilde{\Lambda}_{I,+}^{p,q}(M) \right) := \bigoplus_{p,q \in \mathbb{Z}} \left(h_1^p h_2^q \Lambda_I^{p+q,0}(M) \right)$$

with the differential $d_+ := h_1 \partial + h_2 \partial_J$ (Proposition 8.13.) The bicomplex $(\tilde{\Lambda}_+^*(M), d_+)$ is called **the extended quaternionic Dolbeault bicomplex**. Clearly, the subspace

$$(\Lambda_+^*(M), d_+) \subset (\tilde{\Lambda}_+^*(M), d_+)$$

is preserved by d_+ .

Remark 10.2: In most respects, the extended qD-bicomplex can play the role of the usual quaternionic Dolbeault complex just as nicely. For instance, given a C^∞ -bundle B with a $(\Lambda_+^*(M), d_+)$ -connection ∇ , we can immediately obtain a $(\tilde{\Lambda}_+^*(M), d_+)$ -connection $\tilde{\nabla}$ on the bundle B ; the converse

is also true (Claim 8.16). Moreover, the connection $\tilde{\nabla}$ is flat (projectively flat) if and only if ∇ is flat (projectively flat).

Remark 10.3: The usual qD-complex is embedded to $\tilde{\Lambda}_+^*(M), d_+$ as follows

$$\begin{aligned} \Lambda_+^*(M) &= \bigoplus_{p,q \geq 0} \left(h_1^p h_2^q \Lambda_I^{p+q,0}(M) \right) \\ &\subset \bigoplus_{p,q \in \mathbb{Z}} \left(h_1^p h_2^q \Lambda_I^{p+q,0}(M) = \tilde{\Lambda}_+^*(M) \right). \end{aligned}$$

10.2 qD-Laplace operator

Let M be a hyperkähler manifold, $\dim_{\mathbb{H}} M = n$, I, J, K the standard triple of induced complex structures, and $\Omega_I := \omega_J + \sqrt{-1} \omega_K$ the standard holomorphic symplectic form on (M, I) . Consider the corresponding section of the canonical class $\Omega_I^n \in \Lambda_I^{2n,0}(M)$. Let

$$* : \Lambda^{p,0} \longrightarrow \Lambda^{2n-p,2n}(M)$$

be the Hodge star operator,

$$\overline{K} : \Lambda^{p,0} \longrightarrow \Lambda^{p,2n}(M)$$

the multiplication by $\overline{\Omega}_I^n$, and

$$\overline{K}^* : \Lambda^{p,2n} \longrightarrow \Lambda^{p,0}(M)$$

the Hermitian adjoint operator. We define the hyperkähler star operator

$$\star : \Lambda^{p,0} \longrightarrow \Lambda^{2n-p,0}(M)$$

([V5]) as a composition of $*$ and $\overline{K}^*$. Clearly, $\partial^* = -\star \partial \star$ and $\partial_J^* = -\star \partial_J \star$, where

$$\partial, \partial_J : \Lambda_I^{p,0}(M) \longrightarrow \Lambda_I^{p+1,0}(M)$$

are the standard differentials (Proposition 8.13), and ∂^*, ∂_J^* their Hermitian adjoint operators, defined as

$$\partial^* = - * \partial *, \quad \partial_J^* = - * \partial_J *,$$

([V1]).

We consider $\star$ as an endomorphism of

$$\tilde{\Lambda}_+(M) = \oplus_{p,q \geq 0} h_1^p h_2^q \Lambda_I^{p+q,0}(M),$$

$$\star(h_1^p h_2^q \eta) := h_1^{n-p} h_2^{n-q} \star(\eta).$$

Clearly, $\star$ preserves $\Lambda_+(M) \subset \tilde{\Lambda}_+(M)$. Therefore, we may consider $\star$ as an endomorphism of the space of quaternionic Dolbeault forms. The operator $\star$ plays the same role as the usual Hodge star operator plays in Hodge theory.

Definition 10.4: In the above assumptions, the quaternionic Dolbeault Laplace operator is defined on $\tilde{\Lambda}_+(M)$ as

$$\Delta_+ := d_+ d_+^* + d_+^* d_+,$$

where $d_+^* := -\star d_+ \star$.

Given a bundle B with a connection over $\Lambda_+(M)$, we may define the qD-Laplace operator on $\tilde{\Lambda}_+(M) \otimes B$ as

$$\Delta_+ := \tilde{\nabla} \tilde{\nabla}^* + \tilde{\nabla}^* \tilde{\nabla},$$

where $\tilde{\nabla}$ is the corresponding connection over $\tilde{\Lambda}_+(M)$, d_+ , and

$$\tilde{\nabla}^* := -\star \tilde{\nabla} \star.$$

Remark 10.5: Given $\eta \in \Lambda_I^{p,0}(M)$, let $h_1^i h_2^j \eta$ be the corresponding vector in $\Lambda_+(M)$. Clearly, $\Delta_+(h_1^i h_2^j \eta)$ is independent from the choice of i, j . Therefore, we may consider Δ_+ as an operator on $\Lambda_I^{p,0}(M)$. By the hyperkaehler Kodaira relations ([V1]; see also (10.1)), the anticommutators $\{\partial, \partial_J^*\}$ and $\{\partial^*, \partial_J\}$ vanish, and we have

$$\Delta_+ = \partial \partial^* + \partial^* \partial + \partial_J \partial_J^* + \partial_J^* \partial_J$$

A similar interpretation exists for the quaternionic Dolbeault Laplacian with coefficients in a bundle.

10.3 qD-Laplace operator and the quaternionic Kodaira relations

Let M be a hyperkähler manifold, $\dim_{\mathbb{H}} M = n$, I, J, K the standard triple of induced complex structures, and $\Omega_I := \omega_J + \sqrt{-1} \omega_K$ the standard holomorphic symplectic form on (M, I) . Consider the operator of multiplication by Ω_I

$$L_\Omega : \Lambda_I^{p,0}(M) \longrightarrow \Lambda_I^{p+2,0}(M).$$

Let $\Lambda_\Omega := \star L_\Omega \star$ be the Hermitian adjoint operator.
Consider the differentials

$$\partial, \partial_J : \Lambda_I^{p,0}(M) \longrightarrow \Lambda_I^{p+1,0}(M)$$

(Proposition 8.13). In [V1], the following Kodaira-type relations were obtained.

$$\begin{aligned} &= -\sqrt{-1} \partial_J^*, \quad [\Lambda_\Omega, \partial_J] = -\sqrt{-1} \partial^*, \\ [L_\Omega, \partial^*] &= \sqrt{-1} \partial_J, \quad [L_\Omega, \partial_J^*] = \sqrt{-1} \partial. \end{aligned} \tag{10.1}$$

Let B be a bundle with a connection over $\Lambda_+^*(M)$, ∇' denote the $(1,0)$ -component of a connection, ∇'' the $(0,1)$ -component of a connection, and $\nabla'_J := -J \circ \nabla'' \circ J$ the operator obtained from ∇'' by conjugation with J .

The same relations are true for forms with coefficients in a bundle:

$$\begin{aligned} [\Lambda_\Omega, \nabla'] &= -\sqrt{-1} \nabla'^*_J, \quad [\Lambda_\Omega, \nabla'_J] = -\sqrt{-1} \nabla'^*, \\ [L_\Omega, \nabla'^*] &= \sqrt{-1} \nabla'_J, \quad [L_\Omega, \nabla'^*_J] = \sqrt{-1} \nabla'. \end{aligned}$$

From these relations, we obtain the following results about the Laplace operators ([V1]). Let

$$\Delta_\partial := \nabla' \nabla'^* + \nabla'^* \nabla'$$

be the standard Dolbeault Laplacian,

$$\Delta_{\partial_J} := \nabla'_J \nabla'^*_J + \nabla'^*_J \nabla'_J$$

the Laplacian corresponding to ∇'_J , and Δ_+ the qD-Laplace operator (Definition 10.4)

Using the quaternionic Dolbeault Kodaira relations in the standard way ([GH]), we obtain the following results about the Laplace operators ([V1]).

$$\Delta_+ = \Delta_\partial + \Delta_{\partial_J}, \quad \Delta_\partial - \Delta_{\partial_J} = -\sqrt{-1} [\Theta, \Lambda_\Omega], \quad (10.2)$$

where $\Theta := \nabla' \nabla'_J + \nabla'_J \nabla'$ is the curvature of ∇_+ , considered as an operator on differential forms.

11 Quaternionic Dolbeault complex of a product

In this section, we study the linear algebra of the quaternionic Dolbeault complex of a product of two hyperkähler manifolds.

11.1 Bidegree: the definition

Let $M = M_1 \times M_2$ be a product of two hyperkähler manifolds,

$$\pi_1 : M \longrightarrow M_1, \pi_2 : M \longrightarrow M_2$$

the standard projections, and I an induced complex structure. We have

$$\Lambda_+^i(M) = \Lambda_I^{i,0}(M) \otimes_{\mathbb{C}} S^i \mathbb{H}_I$$

(Lemma 8.1). Therefore, the natural multiplicative homomorphism

$$\pi_1^* \Lambda_+^*(M_1) \otimes_{C^\infty M} \pi_2^* \Lambda_+^*(M_2) \longrightarrow \Lambda_+^*(M) \quad (11.1)$$

is surjective. It is very easy to see that (11.1) not injective. Still, it is possible to speak of a bidegree of a form.

Definition 11.1: Let

$$\boxtimes : \pi_1^* \Lambda_+^*(M_1) \otimes_{C^\infty M} \pi_2^* \Lambda_+^*(M_2) \longrightarrow \Lambda_+^*(M)$$

be the natural surjective map defined above. We say that the form $\eta \in \Lambda_+^{p+q}(M)$ has bidegree (p, q) if η belongs to

$$\boxtimes \left(\pi_1^* \Lambda_+^p(M_1) \otimes_{C^\infty M} \pi_2^* \Lambda_+^q(M_2) \right) \subset \Lambda_+^*(M)$$

Let

$$\rho : \mathfrak{u}(1) \longrightarrow \text{End } \Lambda^*(M)$$

be a representation of $\mathfrak{u}(1)$ acting as $\sqrt{-1}(p-q)$ on forms of bidegree (p, q) . From Lemma 8.1, it follows that ρ is well defined. Clearly, ρ commutes with the natural $SU(2)$ -action on $\Lambda^*(M)$. Therefore, ρ preserves the subspace $\Lambda_+^* \subset \Lambda^*(M)$. We obtained the following claim

Claim 11.2: In the above assumptions, we have a direct sum decomposition

$$\Lambda_+^*(M) = \bigoplus_{p,q} \Lambda_+^*(M)_{p,q}, \quad (11.2)$$

where $\Lambda_+^*(M)_{p,q} \subset \Lambda_+^*(M)$ denotes the subset of bidegree (p, q) .

■

11.2 Fiberwise connection operators on a product of two hyperkähler manifolds

Let B be a vector bundle with connection over $M = M_1 \times M_2$.

A bidegree decomposition exists for forms with coefficients in a vector bundle:

$$\Lambda_+^*(M) \otimes B = \bigoplus_{p,q} \Lambda_+^*(M)_{p,q} \otimes B.$$

Let ∇ be a connection in B , and $\nabla = \nabla_1 + \nabla_2$ be the decomposition of ∇ with respect to bidegree, with ∇_1 being of bidegree $(1,0)$, and ∇_2 of bidegree $(0,1)$.

Clearly, the operator ∇_1 is $C^\infty(M_2)$ -linear, and ∇_2 is $C^\infty(M_1)$ -linear. One may think of ∇_1, ∇_2 as of connection operators acting along the fibers of the standard projections $\pi_1 : M \rightarrow M_1$, $\pi_2 : M \rightarrow M_2$. Indeed, we have

$$\nabla_1 : B \rightarrow B \otimes \pi_1^* \Lambda_+^* M_1, \quad \nabla_2 : B \rightarrow B \otimes \pi_1^* \Lambda_+^* M_2.$$

Let $M = M_1 \times M_2$ be a product of two hyperkähler manifolds, I an induced complex structure, and B a Hermitian holomorphic vector bundle over (M, I) . Assume that M is projectively hyperholomorphic and Yang-Mills. Let

$$\nabla : B \rightarrow \Lambda^1(M) \otimes B$$

be the Hermitian connection in B . We denote by the same letter the associated extended connection operator

$$\nabla : \Lambda_+^*(M) \otimes B \longrightarrow \Lambda_+^{*+1}(M) \otimes B.$$

Consider the decomposition $\nabla = \nabla_1 + \nabla_2$ constructed as above.

Proposition 11.3: Let $M = M_1 \times M_2$ be a product of two compact hyperkaehler manifolds, with $H^1(M_1) = 0$. Consider a projectively hyperholomorphic Yang-Mills bundle B on $M_1 \times M_2$. In the above notation, we have

$$\nabla_1 \nabla_2 = -\nabla_2 \nabla_1 \quad (11.3)$$

Proof: We have $\nabla = \nabla_1 + \nabla_2$, and the bidegree-(1,1) part of ∇^2 is equal to $\nabla_1 \nabla_2 + \nabla_2 \nabla_1$. On the other hand, B is Yang-Mills, and therefore, the curvature of B over $\Lambda_+^*(M)$ is equal to $\omega \otimes Id_B$, where $\omega \in \Lambda^{1,1}(M)$ is a harmonic form (Corollary 4.15). To prove Proposition 11.3, we have to show that the bidegree-(1,1) part of $\nabla^2 = \omega$ is zero. However, since $H^1(M_1) = 0$, by Künneth formula we have $H^2(M) = H^2(M_1) \oplus H^2(M_2)$. Therefore, any harmonic form on M is a sum of a harmonic form lifted from M_1 and a harmonic form lifted from M_2 . In other words, any harmonic form is a sum of a form of bidegree (2,0) (these are lifted from M_1) and a form of bidegree (0,2) (these are lifted from M_2). We have shown that the curvature of B has zero bidegree-(1,1) part. This proves (11.3). ■

Consider the extended qD-complex

$$\tilde{\Lambda}_+^*(M) \otimes B$$

(Definition 10.1). Clearly, the operators ∇_1, ∇_2 are extended naturally to $\tilde{\Lambda}_+^*(M) \otimes B$, and commute here by Proposition 11.3. Consider the hyperkähler star-operator

$$\star_1 : \tilde{\Lambda}_+^i(M_1) \longrightarrow \Lambda_+^{2 \dim_{\mathbb{H}} M_1 - i}(M_1)$$

(Subsection 10.2). Let $\star_1$ act as identity on $\tilde{\Lambda}_+^i(M_2)$. We extend $\star_1$ to $\tilde{\Lambda}_+^*(M) \otimes B$ multiplicatively via (11.1). Since ∇_2 is $\tilde{\Lambda}_+^i(M_1)$ -linear, this operator commutes with $\star_1$.

Consider the Laplace operator associated with $\nabla_1 : \tilde{\Lambda}_+^*(M) \longrightarrow \tilde{\Lambda}_+^*(M)$,

$$\Delta_1 := \nabla_1^* \nabla_1 + \nabla_1 \nabla_1^*,$$

where $\nabla_1^* := -\star_1 \nabla_1 \star_1$. From (11.3) it follows immediately that

$$[\Delta_1, \nabla_2] = 0. \quad (11.4)$$

Let Δ_{∂_1} be the Dolbeault Laplace operator on $\Lambda_+^*(M) \otimes B$,

$$\Delta_{\partial_1} := \nabla_1' \nabla_1'^* + \nabla_1'^* \nabla_1'.$$

By (10.2), we have

$$\Delta_1 - 2\Delta_{\partial_1} = [\Theta_1, \Lambda_1], \quad (11.5)$$

where $\Theta_1 = \nabla_1^2$ is a curvature operator, and Λ_1 is the standard Λ lifted from M_1 . On the other hand, Δ_1 and ∇_2 commute by (11.4). We obtained the following proposition

Proposition 11.4: Let $M = M_1 \times M_2$ be a product of compact hyperkähler manifolds, B a bundle on M which is projectively hyperholomorphic Yang-Mills, and $\Delta_{\partial_1}, \nabla_2$ the operators on $\Lambda_+^*(M) \otimes B$ defined above. Assume that $H^1(M_1) = 0$ or $H^1(M_2) = 0$. Then

$$[\Delta_{\partial_1}, \nabla_2] = 0.$$

Proof: By (11.4), ∇_2 commutes with Δ_1 . The equation (11.5) expresses Δ_{∂_1} via Δ_1 and $[\Theta_1, \Lambda]$. Therefore, to prove Proposition 11.4, it suffices to show that ∇_2 commutes with $[\Theta_1, \Lambda]$. However, $\Theta_1 = \nabla_1^2$, and by Proposition 11.3, ∇_2 commutes with ∇_1 . The operator Λ is trivial along M_2 . Representing a form as a product of a form lifted from M_1 and one lifted from M_2 , we conclude that ∇_2 commutes with Λ_1 (see the proof of Lemma 12.2 for a more detailed version of this argument). This proves Proposition 11.4. ■

12 Hyperholomorphic structure on a pushforward of a hyperholomorphic bundle

In this Section we use the results of Section 11 to study the pushforward of a projectively hyperholomorphic bundle.

12.1 Quaternionic Dolbeault complex and a C^∞ -pushforward of a bundle

Let $M = M_1 \times M_2$ be a product of two hyperkähler manifolds, and I an induced complex structure on M .

Lemma 12.1: In the above assumptions, consider the map

$$\boxtimes : \Lambda_I^{*,0}(M_1) \otimes_{\mathbb{C}} \Lambda_+^*(M_2) \longrightarrow \Lambda_+^*(M) \quad (12.1)$$

induced by (11.1). Then (12.1) is an embedding.

Proof: Follows immediately from Lemma 8.1. ■

Let B be a bundle with connection on M . We denote the corresponding extended connection

$$\Lambda_+^*(M) \otimes B \longrightarrow \Lambda_+^{*+1}(M) \otimes B$$

by the same letter ∇ . Consider the natural embedding

$$\left(\Lambda_I^{*,0}(M_1) \otimes_{\mathbb{C}} \Lambda_+^*(M_2) \right) \otimes_{C^\infty M} B \longrightarrow \Lambda_+^*(M) \otimes B \quad (12.2)$$

obtained from (12.1) by tensoring with B . As in Section 11, we consider a decomposition $\nabla = \nabla_1 + \nabla_2$, with ∇_1, ∇_2 being parts of the connection of bidegree (1,0) and (0,1).

Lemma 12.2: In the above assumptions, the operator

$$\nabla_2 : \Lambda_+^*(M) \otimes B \longrightarrow \Lambda_+^{*+1}(M) \otimes B$$

preserves the subspace

$$V := \left(\Lambda_I^{*,0}(M_1) \otimes_{\mathbb{C}} \Lambda_+^*(M_2) \right) \otimes_{C^\infty M} B \subset \Lambda_+^*(M) \otimes B$$

(12.2)

Proof: Let $\eta_1 \in \Lambda_I^{*,0}(M_1)$, $\eta_2 \in \Lambda_+^*(M_2)$, and $b \in B$. Consider the form $\eta := \pi_1^* \eta_1 \otimes \pi_2^* \eta_2 \otimes b$, where $\pi_i : M \rightarrow M_i$, $i = 1, 2$ are the standart projection maps. Such forms generate the whole space V . Therefore, to

prove Lemma 12.2 it suffices to show that $\nabla_2(\eta)$ belongs to the image V of (12.2).

By definition, the form η_1 is constant along M_1 . This implies

$$\nabla_2(\eta) = \pi_1^* \eta_1 \cdot \nabla_2(\pi_2^* \eta_2 \otimes b).$$

The form $\nabla_2(\pi_2^* \eta_2 \otimes b)$ has bidegree $(0, p)$, and $\pi_1^* \eta_1$ belongs to $\pi^*(\Lambda_I^{*,0}(M_1))$. We obtain that the $\nabla_2(\eta)$ is a product of a $(p, 0)$ -form of M_1 and a form bidegree $(0, p)$; this is exactly the definition of V . We proved Lemma 12.2. ■

Given $x \in M_2$, consider the space Γ_x of C^∞ -sections of the bundle

$$\left(\pi_2^* \Lambda_I^{*,0}(M_1) \otimes_{C^\infty} B \right) \Big|_{M_1 \times \{x\}},$$

$$\Gamma_x = \Gamma_{M_1 \times \{x\}} \left(\pi_2^* \Lambda_I^{*,0}(M_1) \otimes_{C^\infty} B \right),$$

where $\Gamma_Z(B)$ denotes the space of sections of a restriction $B|_Z$. Clearly, the correspondence $x \longrightarrow \Gamma_x$ gives an infinite-dimensional C^∞ -vector bundle over M_2 . Denote this bundle as Γ . By definition, the fibers of Γ can be also represented as

$$\pi_2^* \Lambda_I^{p,0}(M_1) \otimes_{C^\infty} B \cong \Lambda_{I,+}^{p,0}(M)_{p,0} \otimes B,$$

where $\Lambda_{I,+}^{p,q}(M)$ denotes the Hodge grading on the space $\Lambda_+^*(M)$ (Subsection 8.3) and $\Lambda_+^*(M)_{p,q}$ the space of bidegree- (p, q) forms (Definition 11.1).

Using Lemma 12.2, we may consider ∇_2 as an operator

$$\nabla_2 : \Gamma \otimes \Lambda_+^i(M_2) \longrightarrow \Gamma \otimes \Lambda_+^{i+1}(M_2) \quad (12.3)$$

Clearly, this gives a connection (in the sense of Definition 8.4) in an infinite-dimensional vector bundle Γ over a DG-algebra $(\Lambda_+^*(M_2), d_+)$.

Proposition 12.3: Let $M = M_1 \times M_2$ be a product of two hyperkähler manifolds, I an induced complex structure, and B a holomorphic bundle over (M, I) . Consider the connection

$$\nabla_\Gamma : \Gamma \otimes \Lambda_+^i(M_2) \longrightarrow \Gamma \otimes \Lambda_+^{i+1}(M_2) \quad (12.4)$$

defined above (see (12.3)). Assume that B is hyperhomomorphic (projectively hyperhomomorphic Yang-Mills). Then Γ is also hyperhomomorphic (resp., projectively hyperhomomorphic Yang-Mills)

Proof: Let γ be a section of Γ , which is an element of $\Lambda_{I,+}^{p,0}(M)_{p,0} \otimes B$. By construction,

$$\nabla_{\Gamma}(\gamma) = \nabla_2(\gamma),$$

where $\nabla_2(\gamma)$ is understood as an element of

$$\Gamma \otimes \Lambda_+^*(M_2) = V = \left(\Lambda_I^{*,0}(M_1) \otimes_{\mathbb{C}} \Lambda_+^*(M_2) \right) \otimes_{C^\infty M} B \subset \Lambda_+^*(M) \otimes B \quad (12.5)$$

Under the identification (12.5), the section $\nabla_{\Gamma}^2 \gamma$ is equal to $(\nabla_2)^2 \gamma$. By definition, $(\nabla_2)^2 \gamma$ is equal to $\Theta_{0,2}(\gamma)$, where $\Theta_{0,2}$ is the bidegree-(0,2) part of the curvature Θ of B over $\Lambda_+^*(M)$.

If B is hyperhomomorphic, then $\Theta = 0$, hence its bidegree-(0,2) part is also zero, and we have $\nabla_{\Gamma}^2 \gamma = 0$; this means that Γ is flat over $\Lambda_+^*(M_2)$, that is, hyperhomomorphic.

If B is projectively hyperhomomorphic Yang-Mills, then $\Theta(\gamma) = \omega \wedge \gamma$, where ω is a harmonic form of M . Therefore, a bidegree-(0,2) part $\Theta_{0,2}$ of Θ is equal to ω_2 , where ω_2 is bidegree (0,2)-part of ω . Using the Künneth formula, we obtain that ω_2 is harmonic. We have shown, then, that the $\Lambda_+^*(M)$ -curvature of Γ is a harmonic form. By Corollary 4.15, this means that Γ is projectively hyperhomomorphic Yang-Mills. We proved Proposition 12.3. ■

Consider the Dolbeault Laplace operator associated with ∇_1'

$$\Delta_{\partial_1} : \Lambda_*^+(M) \otimes B \longrightarrow \Lambda_*^+(M) \otimes B$$

(Proposition 11.4). Clearly, Δ_{∂_1} preserves the bidegree and the Hodge decomposition on $\Lambda_*^+(M)$. Therefore, $\Delta_{\partial_1}(\Gamma) \subset \Gamma$. Let $\Gamma_{\Delta_{\partial_1}} \subset \Gamma$ be the kernel of $\Delta_{\partial_1} : \Gamma \longrightarrow \Gamma$. If $H^1(M_1) = 0$ or $H^1(M_2) = 0$, the operators Δ_{∂_1} and ∇_2 commute (Proposition 11.4). Therefore, the restriction of ∇_2 to $V_{\Delta_{\partial_1}}$ establishes a $\Lambda_+^*(M_2)$ -connection on the subspace $\Gamma_{\Delta_{\partial_1}} \subset \Gamma$. We have proven the following result

Corollary 12.4: In the assumptions of Proposition 12.3, let $\Gamma_{\Delta_{\partial_1}}$ be the kernel of the Dolbeault Laplace operator defined above. Then the subsheaf $\Gamma_{\Delta_{\partial_1}} \subset \Gamma$ is invariant with respect to the $\Lambda_+^*(M_2)$ -connection constructed in Proposition 12.3. In particular, the sheaf $\Gamma_{\Delta_{\partial_1}}$ is a pqDYM-module, in the sense of Definition 9.1.

■

12.2 Quaternionic Dolbeault complex and the derived direct image functor

In this Subsection, we conclude the proof Theorem 7.5, summing up the arguments given earlier.

Let $M = M_1 \times M_2$ be a product of two hyperkähler manifolds, I an induced complex structure on M , and B a projectively hyperholomorphic Yang-Mills bundle on (M, I) . Denote by π_1, π_2 the standard projections to M_1, M_2 . As usually, we write the connection in B as

$$\nabla = \nabla_1 + \nabla_2 : \Lambda_+^*(M) \otimes B \longrightarrow \Lambda_+^{*+1}(M) \otimes B,$$

with ∇_1, ∇_2 being the bidegree $(1,0)$ and $(0,1)$ -components of ∇ (Definition 11.1). We shall think of ∇_1, ∇_2 as of parts of the connection acting along the fibers of π_2, π_1 respectively. This is to say, ∇_1 “acts along M_1 ”, ∇_2 “acts along M_2 ”. Using the Hodge decomposition, we can write

$$\nabla_1 = \nabla'_1 + \nabla''_1,$$

with ∇'_1, ∇''_1 being $(1,0)$ - and $(0,1)$ -parts of ∇_1 , with respect to the Hodge decomposition.

In the above assumptions, let

$$\Gamma := \bigoplus_p \pi_{2*} \Lambda^{p,0}(M)_{p,0} \otimes B = \bigoplus_p \Lambda_{I,+}^{p,0}(M)_{p,0} \otimes B \subset \Lambda_+^*(M) \otimes B.$$

be the bundle over M_2 defined in Subsection 12.1, and ∇_Γ the standard projectively hyperholomorphic Yang-Mills $\Lambda_+^*(M_2)$ -connection in Γ (Proposition 12.3). The fiber of Γ in $x \in M_2$ is identified with the space of sections of the bundle

$$\bigoplus_p \Lambda^{p,0}(M_1) \otimes_{C^\infty M} \left(B \Big|_{M_1 \times \{x\}} \right),$$

where $B|_{M_1 \times \{x\}}$ is a restriction of B from $M = M_1 \times M_2$ to $M_1 \times \{x\} \subset M$.

Let Δ_{∂_1} be the Laplace operator on the fibers of Γ , and $\Gamma_{\Delta_{\partial_1}}$ its kernel. In Corollary 12.4, we have established a structure of pqDYM-module on the sheaf $\Gamma_{\Delta_{\partial_1}}$. On the other hand, outside of its singularities, the sheaf $\Gamma_{\Delta_{\partial_1}}$ is naturally identified with the derived direct image of B . Let $\mathcal{F} := \mathcal{H}_I(\Gamma_{\Delta_{\partial_1}})$ be the corresponding quasicoherent sheaf (Definition 9.3), and $\mathcal{T}$ be the torsion component of R . By Remark 9.7, the reflexive hull of $\mathcal{F}/\mathcal{T}$ is polystable. This proves Theorem 7.5. ■

Acknowledgements: This paper appeared as a result of fruitful talks with D. Kaledin. I am grateful to M. Kontsevich, for encouraging me to think of quaternionic cohomology, and A. Losev, who suggested the close study of Lie superalgebras arising from the natural DG-algebras on manifolds. Many thanks S. Merkulov for his interest and encouragement, R. Bielawski for interesting discussions, J.-P. Demailly for answering my questions about positive forms, and T. Pantev for useful advice. After this paper was written, K. Yoshioka found a counterexample to Mukai conjecture on stability for abelian surfaces; he gave me some historical reference and also helped to find a serious error in the manuscript.

And last but not least, my gratitude to Glasgow University, where I visited while writing this paper, for warmth and hospitality.

References

- [BS] Bando, S., Siu, Y.-T., *Stable sheaves and Einstein-Hermitian metrics*, In: Geometry and Analysis on Complex Manifolds, Festschrift for Professor S. Kobayashi's 60th Birthday, ed. T. Mabuchi, J. Noguchi, T. Ochiai, World Scientific, 1994, pp. 39-50.
- [B] Baston, R. J. *Quaternionic complexes*. J. Geom. Phys. **8** (1992), no. 1-4, 29-52.
- [Be1] Beauville, A. *Varieties Kähleriennes dont la première classe de Chern est nulle*. J. Diff. Geom. **18**, pp. 755-782 (1983).
- [Be2] Beauville, A. *Quelques remarques sur la transformation de Fourier dans l'anneau de Chow d'une variété abélienne*, Lecture Notes in Math., **1016**, 238-260.
- [Bes] Besse, A., *Einstein Manifolds*, Springer-Verlag, New York (1987)

- [BBR1] Bartocci, C.; Bruzzo, U.; Hernandez Ruiperez, D. *A Fourier-Mukai transform for stable bundles on K3 surfaces*, J. Reine Angew. Math. **486** (1997), 1–16.
- [BBR2] Bartocci, C.; Bruzzo, U.; Hernandez Ruiperez, D. *Existence of stable vector bundles on K3 surfaces and the Fourier-Mukai transform*, Algebraic geometry (Catania, 1993/Barcelona, 1994), 245–257, Lecture Notes in Pure and Appl. Math., 200, Dekker, New York, 1998.
- [BBR3] Bartocci, C., Bruzzo, U., Hernandez Ruiperez, D. *A hyper-Kähler Fourier transform*, Differential Geom. Appl. **8** (1998), no. 3, 239–249.
- [BO] A. Bondal, D. Orlov, *Semiorthogonal decomposition for algebraic varieties*, alg-geom/9506012
- [Br] Bridgeland, T., *Equivalences of triangulated categories and Fourier-Mukai transforms*, Bull. London Math. Soc. **31** (1999), 25–34,
- [BrM] Bridgeland, T., Maciocia A., *Fourier-Mukai transforms for K3 and elliptic fibrations*, math.AG/9908022
- [C] Calabi, E., *Metriques kähleriennes et fibrès holomorphes*, Ann. Ecol. Norm. Sup. **12** (1979), 269–294.
- [CS] Capria, M. M., Salamon, S. M. *Yang-Mills fields on quaternionic spaces*, Nonlinearity **1** (1988), no. 4, 517–530.
- [DK] Donaldson, S., Kronheimer, P., *The geometry of four-manifolds*, Oxford University Press (1990).
- [FL] Fahlaoui, R., Laszlo, Y. *Transformée de Fourier et stabilité sur les surfaces abéliennes*, Compositio Math. **79** (1991), no. 3, 271–278.
- [GH] Griffiths, Ph., Harris, J., *Principles of Algebraic Geometry*, Wiley-Interscience, New York, 1978.
- [Ka] Kaledin, D., *Hyperkähler structures on total spaces of holomorphic cotangent bundles*, alg-geom/9710026; published in a book “Hyperkähler manifolds” (joint with M. Verbitsky), Int’l Press, Boston, 2001.
- [Kob] Kobayashi S., *Differential geometry of complex vector bundles*, // Princeton University Press, 1987.
- [Kon] Kontsevich, M., *Homological algebra of mirror symmetry*. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 120–139, Birkhäuser, Basel, 1995.
- [L] Leung N. C., Yi S., *Analytic Torsion for Quaternionic manifolds and related topics*, dg-ga/9710022
- [Ma] Maciocia, Antony, *Gieseker stability and the Fourier-Mukai transform for abelian surfaces* Quart. J. Math. Oxford Ser. (2) **47** (1996), no. 185, 87–100.

- [Mu1] Mukai, S., *Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves*, Nagoya Math. J. (1981), 153-175.
- [Mu2] Mukai, S., *On the moduli space of bundles on K3 surfaces. I*, Vector bundles on algebraic varieties (Bombay, 1984), 341-413, Tata Inst. Fund. Res. Stud. Math., 11, Tata Inst. Fund. Res., Bombay, 1987.
- [Mu3] Mukai, S., *Fourier functor and its application to moduli of bundles on an abelian variety*, Adv. in Pure Math. (1987), 515-550.
- [Mu4] Mukai, S., *Abelian variety and spin representation (in Japanese)*, Proceedings of symposium "Hodge theory and algebraic geometry (Sapporo, 1994)", 110-135: English translation, Univ. of Warwick preprint, 1998
- [Mu5] Mukai, S., *Duality of polarized K3 surfaces*, London Math. Soc. Lect. Notes Series 264, 311-326
- [OSS] Christian Okonek, Michael Schneider, Heinz Spindler, *Vector bundles on complex projective spaces*. Progress in mathematics, vol. 3, Birkhauser, 1980.
- [O] Orlov, D., *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. (NY), **84** (1997) 1361-1381
- [P] Positselski, L., *All strictly exceptional collections in $D_{coh}^b(P^m)$ consist of vector bundles*, alg-geom/9507014
- [S] Salamon, S., *Quaternionic Manifolds*, Comunicazione inviata all'Instituto nazionale di Alta Matematica Francesco Severi.
- [V0] Verbitsky M., *Hyperkähler embeddings and holomorphic symplectic geometry II*, alg-geom electronic preprint 9403006 (1994), 14 pages, LaTeX, also published in: GAFA **5** no. 1 (1995), 92-104.
- [V1] Verbitsky M., *Hyperholomorphic bundles over a hyperkähler manifold*, alg-geom electronic preprint 9307008 (1993), 43 pages, LaTeX, also published in: Journ. of Alg. Geom., **5** no. 4 (1996) pp. 633-669.
- [V2] Verbitsky M., *Hyperholomorphic sheaves and new examples of hyperkähler manifolds*, alg-geom 9712012 - 113 pages, LaTeX 2e; published in a book "Hyperkähler manifolds" (joint with D. Kaledin), Int'l Press, Boston, 2001.
- [V3] Verbitsky M., *Hypercomplex Varieties*, alg-geom/9703016 (1997); published in: Comm. Anal. Geom. **7** (1999), no. 2, 355-396.
- [V4] Verbitsky, M., *Hyperholomorphic connections on coherent sheaves and stability*, 40 pages, math.AG/0107182
- [V5] Verbitsky, M., *Mirror Symmetry for hyperkähler manifolds*, electronic preprint alg-geom 9512195 (published in the book "Mirror Symmetry III", International Press).

- [UY] Uhlenbeck K., Yau S. T., *On the existence of Hermitian Yang-Mills connections in stable vector bundles*, // Comm. on Pure and Appl. Math., **39**, p. S257-S293 (1986).
- [Yau] Yau, S. T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I*. // Comm. on Pure and Appl. Math. 31, 339-411 (1978).
- [Y1] Yoshioka, K., *A note on Fourier-Mukai transform*, 10 pages, math.AG/0112267
- [Y2] Yoshioka, K., *Twisted stability and Fourier-Mukai transform*, math.AG/0106118, 28 pages.
- [Y3] Yoshioka, K., *Some examples of Mukai's reflections on K3 surfaces*, J. reine angew. Math. **515** (1999), 97–123
- [Y4] Yoshioka, K., *An application of exceptional bundles to the moduli of stable sheaves on a K3 surface*, 12 pages, alg-geom/9705027