

NON-COMPLEX SYMPLECTIC 4-MANIFOLDS WITH $b_2^+ = 1$

JONGIL PARK

ABSTRACT. In this paper we give a criterion whether a given minimal symplectic 4-manifold with $b_2^+ = 1$ having a torsion-free canonical class is rational or ruled. As a corollary, we confirm that most of homotopy elliptic surfaces $\{E(1)_K \mid K \text{ is a fibered knot in } S^3\}$ constructed by R. Fintushel and R. Stern are minimal symplectic 4-manifolds with $b_2^+ = 1$ which do not admit a complex structure.

1. INTRODUCTION

As an application of Seiberg-Witten theory to symplectic 4-manifolds, C. Taubes proved that every minimal symplectic 4-manifold with $b_2^+ > 1$ satisfies $K^2 \geq 0$ ([T]). Here $K = -c_1(TX) \in H^2(X; \mathbf{Z})$ denotes the canonical class associated to a compatible almost complex structure on X . In the case when $b_2^+ = 1$, the situation is quite different, i.e. there are examples satisfying $K^2 > 0$, $K^2 = 0$ and $K^2 < 0$ respectively. Hence it is an interesting problem to classify minimal symplectic 4-manifolds with $b_2^+ = 1$. Note that most known minimal symplectic 4-manifolds with $b_2^+ = 1$ except Barlow surfaces and Dolgachev surfaces are rational or ruled, i.e. \mathbf{CP}^2 , $S^2 \times S^2$ or S^2 -bundles over a Riemann surface. Regarding this, Liu proved that if $K^2 < 0$, then it is an irrational ruled surface ([MS]). In this paper, when $K^2 \geq 0$, we give a criterion whether a given symplectic 4-manifold with $b_2^+ = 1$ having a torsion-free canonical class is rational or ruled. Explicitly, we prove

Theorem 1.1. *Suppose X is a closed minimal symplectic 4-manifold with $b_2^+ = 1$ such that its canonical class is a torsion-free class of non-negative square. Then X is rational or ruled if and only if its Seiberg-Witten invariant SW_X° vanishes.*

R. Gompf constructed infinitely many minimal symplectic 4-manifolds which do not admit a complex structure using a fiber sum technique ([G]). All those manifolds have $b_2^+ > 1$. In the case when $b_2^+ = 1$, it has not been much known about minimal symplectic 4-manifolds except complex surfaces. In Section 3 we compute the Seiberg-Witten invariant $SW_{E(1)_K}^\circ$, obtained by a small generic perturbation of Seiberg-Witten equations, of a family of homotopy elliptic surfaces $\{E(1)_K \mid K \text{ is a fibered knot in } S^3\}$ constructed by R. Fintushel and R. Stern in [FS] and, as a corollary, we conclude

Corollary 1.1. *If K is a fibered knot in S^3 whose Alexander polynomial is non-trivial and is different from that of any (p, q) -torus knot, then $E(1)_K$ is a simply connected minimal symplectic 4-manifold which do not admit a complex structure.*

Date: March 1, 2001.

1991 Mathematics Subject Classification. 57R17, 57R57.

Key words and phrases. Canonical class, rational, ruled, SW-invariant, symplectic.

This work was supported by grant No. 1999-2-102-002-3 from the interdisciplinary research program of the KOSEF.

2. THE SEIBERG-WITTEN INVARIANT

In this section we briefly review the Seiberg-Witten invariant of smooth 4-manifolds ([KM], [M] for details).

Let X be a closed, oriented smooth 4-manifold with $b_2^+ > 0$ and a fixed metric g , and let L be a characteristic line bundle on X , i.e. $c_1(L)$ is an integral lift of $w_2(X)$. This determines a $Spin^c$ -structure on X which induces a unique complex spinor bundle $W \cong W^+ \oplus W^-$, where W^\pm is the associated $U(2)$ -bundles on X such that $\det(W^\pm) \cong L$. Note that the Levi-Civita connection on TX together with a unitary connection A on L induces a connection $\nabla_A : \Gamma(W^+) \rightarrow \Gamma(T^*X \otimes W^+)$. This connection, followed by Clifford multiplication, induces a $Spin^c$ -Dirac operator $D_A : \Gamma(W^+) \rightarrow \Gamma(W^-)$. Then, for each self-dual 2-form $h \in \Omega_+^2(X; \mathbf{R})$, the following pair of equations for a unitary connection A on L and a section Ψ of $\Gamma(W^+)$ are called the *perturbed Seiberg-Witten equations*:

$$(1) \quad (SW_{g,h}) \begin{cases} D_A \Psi & = 0 \\ F_A^{+g} & = i(\Psi \otimes \Psi^*)_0 + ih. \end{cases}$$

Here F_A^{+g} is the self-dual part of the curvature of A with respect to a metric g on X and $(\Psi \otimes \Psi^*)_0$ is the trace-free part of $(\Psi \otimes \Psi^*)$ which is interpreted as an endomorphism of W^+ . The gauge group $\mathcal{G} := Aut(L) \cong Map(X, S^1)$ acts on the space $\mathcal{A}_X(L) \times \Gamma(W^+)$ by

$$g \cdot (A, \Psi) = (g \circ A \circ g^{-1}, g \cdot \Psi)$$

Since the set of solutions is invariant under the action, it induces an orbit space, called the *Seiberg-Witten moduli space*, denoted by $M_{X,g,h}(L)$, whose formal dimension is

$$\dim M_{X,g,h}(L) = \frac{1}{4}(c_1(L)^2 - 3\sigma(X) - 2e(X))$$

where $\sigma(X)$ is the signature of X and $e(X)$ is the Euler characteristic of X . Note that if $b_2^+(X) > 0$ and $M_{X,g,h}(L) \neq \emptyset$, then for a generic metric g and a generic self-dual 2-form h on X the moduli space $M_{X,g,h}(L)$ contains no reducible solutions, so that it is a compact, smooth manifold of the given dimension. Furthermore the moduli space $M_{X,g,h}(L)$ is orientable and its orientation is determined by a choice of orientation on $\det(H^0(X; \mathbf{R}) \oplus H^1(X; \mathbf{R}) \oplus H_+^2(X; \mathbf{R}))$.

Definition The *Seiberg-Witten invariant* (for brevity, *SW-invariant*) for a smooth 4-manifold X with $b_2^+ > 0$ is a function $SW_X : Spin^c(X) \rightarrow \mathbf{Z}$ defined by

$$(2) \quad SW_X(L) = \begin{cases} \int_{M_{X,g,h}} \mu^{d_L} & \text{if } \dim M_{X,g,h}(L) := 2d_L \geq 0 \text{ and even} \\ 0 & \text{otherwise.} \end{cases}$$

Here $\mu \in H^2(M_{X,g,h}(L); \mathbf{Z})$ is the first Chern class of the S^1 -bundle

$$\widetilde{M_{X,g,h}(L)} = \{\text{solutions}(A, \Psi)\} / Aut^0(L) \longrightarrow M_{X,g,h}(L)$$

where $Aut^0(L)$ consists of gauge transformations which are the identity on the fiber of L over a fixed base point in X . Note that if $b_2^+(X) > 1$, the Seiberg-Witten invariant, denoted by $SW_X = \sum SW_X(L) \cdot e^L$, is a diffeomorphic invariant, i.e. SW_X does not depend on the choice of a generic metric on X and a generic perturbation of Seiberg-Witten equations. Furthermore, only finitely many $Spin^c$ -structures on X have a non-zero Seiberg-Witten invariant. We say that the characteristic line bundle L , equivalently a cohomology class $c_1(L) \in H^2(X; \mathbf{Z})$, is a *SW-basic class* of X if $SW_X(L) \neq 0$.

When $b_2^+(X) = 1$, the SW-invariant $SW_X(L)$ defined in (2) above depends not only on a metric g but also on a self-dual 2-form h . Because of this fact, there are several

types of Seiberg-Witten invariants for a smooth 4-manifold with $b_2^+ = 1$ depending on how to perturb Seiberg-Witten equations. We introduce three types of SW-invariants and investigate how they are related. First we allow all metrics and self-dual 2-forms to perturb Seiberg-Witten equations. Then the SW-invariant $SW_X(L)$ defined in (2) above has generically two values which are determined by the sign of $(2\pi c_1(L) + [h]) \cdot [\omega_g]$, where ω_g is a unique g -self-dual harmonic 2-form of norm one lying in the positive component of $H_+^2(X; \mathbf{R})$. We denote the SW-invariant for a generic metric g and a self-dual 2-form h satisfying $(2\pi c_1(L) + [h]) \cdot [\omega_g] > 0$ by $SW_X^+(L)$ and denote the other one by $SW_X^-(L)$. Then the wall crossing formula tells us the relation between $SW_X^+(L)$ and $SW_X^-(L)$.

Theorem 2.1 (Wall crossing formula, [LL], [M]). *Suppose that X is a closed, oriented smooth 4-manifold with $b_1 = 0$ and $b_2^+ = 1$. Then for each characteristic line bundle L on X such that the formal dimension of the moduli space $M_{X,g,h}(L)$ is non-negative and even, say $2d_L$, we have*

$$SW_X^+(L) - SW_X^-(L) = -(-1)^{d_L}.$$

In general case ($b_1 \neq 0$), Li and Liu proved that the difference is also expressed explicitly by some numerical invariants of $c_1(L)$ and X ([LL]).

Second one may perturb the Seiberg-Witten equations by adding only a small generic self-dual 2-form $h \in \Omega_+^2(X; \mathbf{R})$, so that one can define the SW-invariants as in (2) above. In this case we denote the SW-invariant for a generic metric g satisfying $(2\pi c_1(L)) \cdot [\omega_g] > 0$ by $SW_X^{\circ,+}(L)$ and we denote the other one by $SW_X^{\circ,-}(L)$. Note that $SW_X^{\circ,\pm}(L) = SW_X^\pm(L)$. But it sometimes happens that the sign of $(2\pi c_1(L)) \cdot [\omega_g]$ is the same for all generic metrics, so that there exists only one SW-invariant obtained by a small generic perturbation of Seiberg-Witten equations. In such a case we define the SW-invariant of L on X by

$$SW_X^\circ(L) = \begin{cases} SW_X^{\circ,+}(L) & \text{if } 2\pi c_1(L) \cdot [\omega_g] > 0 \\ SW_X^{\circ,-}(L) & \text{if } 2\pi c_1(L) \cdot [\omega_g] < 0. \end{cases}$$

If $SW_X^\circ(L) \neq 0$, we also call the corresponding $c_1(L)$ (or L) a *SW-basic class* which will be a diffeomorphic invariant of X . Furthermore we can extend many results obtained for smooth 4-manifolds with $b_2^+ > 1$ to this case. For example, we have

Theorem 2.2. *Let X be a simply connected, closed, oriented smooth 4-manifold with $b_2^+ = 1$ and $b_2^- \leq 9$. Then*

- (i) *There are only finitely many characteristic line bundles L on X such that $SW_X^\circ(L) \neq 0$.*
- (ii) *If X admits a metric of positive scalar curvature, then the SW-invariant of X vanishes, that is, $SW_X^\circ(L) = 0$ for any characteristic line bundle L on X .*
- (iii) *It also satisfies the generalized adjunction inequality. That is, if Σ is a homologically non-trivial, smoothly embedded, oriented surface in X with $[\Sigma]^2 \geq 0$, then any characteristic line bundle L on X with $SW_X^\circ(L) \neq 0$ satisfies the following inequality:*

$$2 \cdot \text{genus}(\Sigma) - 2 \geq |c_1(L) \cdot [\Sigma]| + [\Sigma]^2.$$

Proof : Proofs of (i), (ii) and (iii) are exactly the same as the case $b_2^+ > 1$ as long as the SW-invariant SW_X° is well defined, i.e. it is independent of metrics on X . Let L be a characteristic line bundle on X such that the formal dimension, $\frac{1}{4}(c_1(L)^2 - 3\sigma(X) - 2e(X))$, of the moduli space is non-negative and even. The condition $b_2^+ = 1$ and $b_2^- \leq 9$ imply that $c_1(L)^2 \geq 3\sigma(X) + 2e(X) \geq 0$. Furthermore, since X is simply connected and $c_1(L)$ is characteristic, $c_1(L) \neq 0$. Thus, for any metric g on X , the light cone lemma (see Lemma 3.1) implies $c_1(L) \cdot [\omega_g] \neq 0$, so that the sign of $(2\pi c_1(L)) \cdot [\omega_g]$ is the same for all generic metrics. Hence the SW-invariant $SW_X^\circ(L)$ is well defined. \square

Finally we introduce one more type of Seiberg-Witten invariants for $b_2^+ = 1$ - Given a fixed cohomology class $[x] \in H^2(X; \mathbf{Z})$ with $[x] \cdot [x] \geq 0$, one may divide a set of metrics and self-dual 2-forms into two classes according to the sign of $proj_{+g}(2\pi c_1(L) + [h]) \cdot [x]$, where $proj_{+g}$ is the projection of $\Omega^2(X; \mathbf{R})$ into g -self-dual harmonic 2-form space $H_{+g}^2(X; \mathbf{R})$. In this case we denote the SW-invariant for a generic metric g and a generic self-dual 2-form h satisfying $proj_{+g}(2\pi c_1(L) + [h]) \cdot [x] > 0$ by $SW_X^{[x],+}(L)$ and we denote the other one by $SW_X^{[x],-}(L)$. R. Fintushel and R. Stern used this type ($[x] = [T]$) of SW-invariants for $b_2^+ = 1$ in [FS]. Note that if $[x] \cdot [\omega_g] > 0$, then $SW_X^{[x],\pm}(L) = SW_X^{\pm}(L)$ and if $[x] \cdot [\omega_g] < 0$, then $SW_X^{[x],\pm}(L) = SW_X^{\mp}(L)$.

We close this section by mentioning that C. Taubes' theorem on a symplectic 4-manifold with $b_2^+ > 1$ can be easily extended to $b_2^+ = 1$ case.

Theorem 2.3. *Suppose X is a closed symplectic 4-manifold with $b_2^+ = 1$ and a canonical class K_X . Then $SW_X^-(K_X) = \pm 1$.*

3. MINIMAL SYMPLECTIC 4-MANIFOLDS WITH $b_2^+ = 1$

In this section we investigate minimal symplectic 4-manifolds with $b_2^+ = 1$. First we compute the SW-invariant SW_X° , obtained by a small generic perturbation of Seiberg-Witten equations, of such symplectic 4-manifolds. And then we show that there exist infinitely many simply connected minimal symplectic 4-manifolds which cannot admit a complex structure. Let us begin by stating known results in the classification of symplectic 4-manifolds. As mentioned in Introduction, most known minimal symplectic 4-manifolds with $b_2^+ = 1$ are complex surfaces such as rational or ruled surfaces, which are recognized by Liu's and Ohta-Ono's results.

Theorem 3.1 (Liu, [MS]). *If X is a closed minimal symplectic 4-manifold with $b_2^+ = 1$ such that the canonical class K_X has $K_X^2 < 0$, then X is an irrational ruled surface.*

Furthermore, when $K_X^2 \geq 0$, they also proved the following fact by similar arguments used in the proof of Theorem 3.1 above.

Theorem 3.2 (Liu, Ohta-Ono, [MS]). *Let X be a closed minimal symplectic 4-manifold with $b_2^+ = 1$ and a canonical class K_X . Then the followings are equivalent.*

- (i) X admits a metric of positive scalar curvature.
- (ii) X admits a symplectic structure ω with $K_X \cdot [\omega] < 0$.
- (iii) X is either rational or ruled.

Hence, by combining two theorems above, we conclude that if X is a minimal symplectic 4-manifold with $b_2^+ = 1$ which is neither rational nor ruled, then it admits a symplectic structure ω satisfying $K_X^2 \geq 0$ and $K_X \cdot [\omega] \geq 0$. In this case we compute the Seiberg-Witten invariant $SW_X^{\circ}(K_X)$ of the canonical class K_X on X .

Before stating a main result, we introduce some terminology - Given a symplectic 4-manifold X with $b_2^+ = 1$ and a symplectic form ω , the positive cone $\{\alpha \in H^2(X; \mathbf{R}) \mid \alpha^2 > 0\}$ has two connected components in $H^2(X; \mathbf{R})$. We write P^+ for the forward positive cone, which is the component containing $[\omega]$, and P^- for the backward positive cone. Then the closure of P^+ in $H^2(X; \mathbf{R}) \setminus \{0\}$ is given by

$$\overline{P^+} = \{\alpha \in H^2(X; \mathbf{R}) \mid \alpha^2 \geq 0, \alpha \neq 0, \alpha \cdot [\omega] \geq 0\}.$$

Note that the next lemma, sometimes known as the light cone lemma, follows from Cauchy-Schwarz inequality.

Lemma 3.1 ([MS]). *If $\alpha, \beta \in \overline{P^+}$, then $\alpha \cdot \beta \geq 0$ with equality if and only if $\alpha = \lambda\beta$ for some $\lambda > 0$.*

Now, in the case when $K^2 \geq 0$, we get a criterion whether a given symplectic 4-manifold with $b_2^+ = 1$ having a torsion-free canonical class is rational or ruled. Explicitly, we have

Theorem 3.3. *Suppose X is a closed minimal symplectic 4-manifold with $b_2^+ = 1$ such that its canonical class K_X is a torsion-free class of non-negative square. Then X is rational or ruled if and only if its Seiberg-Witten invariant SW_X° vanishes.*

Proof : As stated above, if X is neither rational nor ruled, then it admits a symplectic structure ω satisfying $K_X^2 \geq 0$ and $K_X \cdot [\omega] \geq 0$. Furthermore, $K_X^2 = 3\sigma(X) + 2e(X) \geq 0$ implies $b_2^- \leq 9$ and the light cone lemma implies $K_X \cdot [\omega] > 0$. Note that, since $b_2^- \leq 9$ and $(-K_X) \cdot [\omega] < 0$, SW_X° is well defined and $SW_X^\circ(-K_X) = SW_X^{\circ,-}(-K_X) = SW_X^-(-K_X)$. Hence Theorem 2.3 implies $SW_X^\circ(-K_X) = \pm 1$, so that $SW_X^\circ \neq 0$. Conversely, if X is rational or ruled with $K_X^2 \geq 0$, Theorem 3.2 implies that X admits a metric of positive scalar curvature. Hence its SW-invariant SW_X° vanishes by Theorem 2.2. \square

Next, as an application of Theorem 3.3 above, we show that a family of homotopy elliptic surfaces $\{E(1)_K \mid K \text{ is a fibered knot in } S^3\}$ constructed by R. Fintushel and R. Stern in [FS] are minimal symplectic 4-manifolds which are neither rational nor ruled. First we briefly review their construction - Suppose K is a fibered knot in S^3 with a punctured surface Σ_g° of genus g as fiber. Let M_K be a 3-manifold obtained by performing 0-framed surgery on K , and let m be a meridional circle to K . Then the 3-manifold M_K can be considered as a fiber bundle over circle with a closed Riemann surface Σ_g as a fiber, and there is a smoothly embedded torus $T_m := m \times S^1$ of square 0 in $M_K \times S^1$. Thus $M_K \times S^1$ fibers over $S^1 \times S^1$ with Σ_g as fiber and with $T_m = m \times S^1$ as section. It is a Thurston's theorem that such a 4-manifold $M_K \times S^1$ has a symplectic structure with symplectic section T_m . Thus, if X is a symplectic 4-manifold with a symplectically embedded torus T of square 0, then the fiber sum 4-manifold $X_K := X \natural_{T=T_m}(M_K \times S^1)$, obtained by taking a fiber sum along $T = T_m$, is symplectic. R. Fintushel and R. Stern proved that X_K is homotopy equivalent to X under a mild condition on X and computed the SW-invariant of X_K (In the case when $b_2^+ = 1$, they computed the relative SW-invariant of X_K). For example, applying the construction above on an elliptic surface $E(1)$, they get a family of homotopy elliptic surfaces $\{E(1)_K \mid K \text{ is a fibered knot in } S^3\}$ and computed the relative SW-invariant $SW_{E(1)_K, T}^\pm$ of $E(1)_K$:

Theorem 3.4 ([FS]). *For each fibered knot K in S^3 , a homotopy elliptic surface $E(1)_K$ is a simply connected symplectic 4-manifold whose $[T]$ -relative SW-invariants are*

$$\begin{aligned} \sum_{L \cdot [T]=0} SW_{E(1)_K}^{[T], \pm}(L) \cdot e^L &= \sum_{L \cdot [T]=0} SW_{E(1)}^{[T], \pm}(L) \cdot e^L \cdot \Delta_K(e^{2[T]}) \\ &= \sum_{n=0}^{\infty} (\mp 1) \cdot e^{\mp(2n+1)[T]} \cdot \Delta_K(e^{2[T]}) \end{aligned}$$

where Δ_K is the Alexander polynomial of K and T is a symplectically embedded torus induced from a standard torus fiber lying in $E(1)$.

Now we are ready to compute the SW-invariant $SW_{E(1)_K}^\circ$ of $E(1)_K$, which is obtained by a small generic perturbation of Seiberg-Witten equations.

Theorem 3.5. *For each fibered knot K in S^3 , a homotopy elliptic surface $E(1)_K$ has a SW-invariant denoted by $SW_{E(1)_K}^\circ = P_{E(1)_K}^+ + P_{E(1)_K}^-$ such that $P_{E(1)_K}^\pm$ is the partial sum consisting of only positive (negative) multiples of $[T]$ in the exponent of the series $\mp \sum_{n=0}^\infty e^{\mp(2n+1)[T]} \cdot \Delta_K(e^{2[T]})$, where Δ_K is the Alexander polynomial of K and T is a symplectically embedded torus induced from a standard torus fiber lying in $E(1)$.*

Proof : Since $E(1)_K$ is symplectic, one may choose a metric g on $E(1)_K$ so that the symplectic form ω is g -self-dual. Then a symplectically embedded torus T of square 0 in $E(1)_K$, induced from a standard torus fiber lying in $E(1)$, satisfies $[T] \cdot [\omega] > 0$. Thus $SW_{E(1)_K}^{\circ, \pm} = SW_{E(1)_K}^\pm = SW_{E(1)_K}^{[T], \pm}$. Since $E(1)_K$ is also simply connected and $b_2^- = 9$, each characteristic line bundle L on $E(1)_K$ has at most one SW-invariant, either $SW_{E(1)_K}^+(L)$ or $SW_{E(1)_K}^-(L)$. Hence $SW_{E(1)_K}^\circ(L)$ is well defined, i.e.

$$(3) \quad SW_{E(1)_K}^\circ(L) = \begin{cases} SW_{E(1)_K}^{\circ, +}(L) = SW_{E(1)_K}^{[T], +}(L) & \text{if } 2\pi c_1(L) \cdot [\omega] > 0 \\ SW_{E(1)_K}^{\circ, -}(L) = SW_{E(1)_K}^{[T], -}(L) & \text{if } 2\pi c_1(L) \cdot [\omega] < 0. \end{cases}$$

Furthermore a generalized adjunction inequality (Theorem 2.2 - (iii)) implies that any characteristic line bundle L with $SW_{E(1)_K}^\circ(L) \neq 0$ satisfies $L \cdot [T] = 0$, so that it is of the form $L = \lambda[T]$, for some integer λ , due to the light cone lemma (see Lemma 3.1). On the other hand, Theorem 3.4 above implies that

$$(4) \quad \sum_{\lambda \in \mathbf{Z}} SW_{E(1)_K}^{[T], \pm}(\lambda[T]) = \mp \sum_{n=0}^\infty e^{\mp(2n+1)[T]} \cdot \Delta_K(e^{2[T]}).$$

Hence, combining Equations (3) and (4), we have

$$SW_{E(1)_K}^\circ := P_{E(1)_K}^+ + P_{E(1)_K}^-,$$

where $P_{E(1)_K}^\pm$ is the partial sum consisting of only positive (negative) multiples of $[T]$ in the exponent of the series $\mp \sum_{n=0}^\infty e^{\mp(2n+1)[T]} \cdot \Delta_K(e^{2[T]})$. \square

Corollary 3.1. *For each fibered knot K in S^3 , a homotopy elliptic surface $E(1)_K$ is rational or ruled if and only if the Alexander polynomial Δ_K of K is trivial.*

Proof : Note that the Alexander polynomial Δ_K is non-trivial (i.e. $\Delta_K \neq 1$) if and only if the highest power of $[T]$ in the series $-\sum_{n=0}^\infty e^{-(2n+1)[T]} \cdot \Delta_K(e^{2[T]})$ coming from $(-e^{-[T]}) \cdot (\text{the highest power of } \Delta_K(e^{2[T]}))$ is non-negative, equivalently, the SW-invariant $SW_{E(1)_K}^\circ$ of $E(1)_K$ is not zero. Thus our claim follows from Theorem 3.3. \square

Corollary 3.2. *For each fibered knot K with a non-trivial Alexander polynomial in S^3 , a homotopy elliptic surface $E(1)_K$ is a minimal symplectic 4-manifold which is neither rational nor ruled.*

Proof : Corollary 3.1 implies that $E(1)_K$ is not rational or ruled. Furthermore since the set of SW-basic classes of $E(1)_K$ is $\{\lambda_i[T] \mid \lambda_1, \dots, \lambda_n \text{ are some integers}\}$, we have $(\lambda_i[T] - \lambda_j[T])^2 = 0 \neq -4$. Hence $E(1)_K$ is minimal. \square

Examples. Let K be a (p, q) -torus knot with relatively prime integers $p, q \geq 2$ in S^3 . Then $E(1)_K$ is diffeomorphic to a Dolgachev surface $E(1; p, q)$, which is obtained by p - and q - logarithmic transforms on $E(1)$. Hence the SW-invariant $SW_{E(1; p, q)}^\circ$ of Dolgachev surfaces is easily computed by Theorem 3.5. Furthermore, it is an easy outcome of Corollary 3.2 that Dolgachev surfaces are neither rational nor ruled.

Corollary 3.3. *If K is a fibered knot in S^3 whose Alexander polynomial is non-trivial and is different from that of any (p, q) -torus knot, then $E(1)_K$ is a simply connected minimal symplectic 4-manifold which do not admit a complex structure.*

Proof : Note that simply connected complex surfaces satisfying $c_1^2 = 0$ and $b_2^+ = 1$ are classified as a blowing up of other complex surfaces and Dolgachev surfaces. Since the Seiberg-Witten invariant $SW_{E(1)_K}^\circ$ is different from that of any such complex surfaces, $E(1)_K$ is not diffeomorphic to any of them so that it cannot be a complex surface. The rest of proof follows from Corollary 3.2. \square

Acknowledgement. The author would like to thank Prof. Ron Fintushel for many helpful discussions including his explanation of Dolgachev surfaces.

REFERENCES

- [FS] R. Fintushel and R. Stern, Knots, links and 4-manifolds, *Invent. Math.* **134** (1998), 363-400
- [G] R. Gompf, A new construction of symplectic manifolds, *Annals of Math.* **142** no **3** (1995), 527-595
- [KM] P.B. Kronheimer and T.S. Mrowka, The genus of embedded surfaces in the projective plane, *Math. Res. Letters* **1** (1994), 709-808
- [LL] T.J. Li and A. Liu, General wall crossing formula, *Math. Res. Letters* **2** (1995), 797-810
- [M] J. Morgan, The Seiberg-Witten equations and applications to the topology of smooth four-manifolds, *Math. Notes* **44**, Princeton University Press, 1996
- [MS] D. McDuff and D. Salamon, A survey of symplectic 4-manifolds with $b_2^+ = 1$, *Turkish Jour. Math.* **20** (1996), 47-61
- [T] C.H. Taubes, The Seiberg-Witten invariants and symplectic forms, *Math. Res. Letters* **1** (1994), 809-822

DEPARTMENT OF MATHEMATICS, KONKUK UNIVERSITY, 1 HWAYANG-DONG, KWANGJIN-GU, SEOUL 143-701, KOREA

E-mail address: jipark@konkuk.ac.kr