

# Resolutions of $p$ -Modular TQFT's and Representations of Symmetric Groups<sup>1</sup>

Thomas Kerler

September 2001

## Contents

1. Introduction .....	1
2. Frohman-Nicas TQFT's over $\mathbb{Z}$ .....	6
3. Lefschetz Decompositions and Specht Modules .....	10
4. $F_p = \mathbb{Z}/p\mathbb{Z}$ -Reductions and the Sequences $C_{p,k}$ .....	12
5. Exactness of $C_{p,k}$ .....	15
6. Characters, Dimensions, and the Alexander Polynomial ....	20
7. Johnson-Morita Extensions .....	23
8. Open Questions.....	26

## 1. Introduction

In this article we construct and study several families of topological quantum field theories (TQFT's) over the finite number field  $F_p = \mathbb{Z}/p\mathbb{Z}$ . They are extracted in a series of steps from the integral TQFT's over lattices constructed by Frohman and Nicas [3] via the intersection homology of the Jacobians of surfaces. Their  $p$ -modular reductions have rather non-trivial modular structures, which reveal themselves to be closely related to those of  $p$ -modular representations of the symmetric groups. We obtain resolutions of the semisimple subquotients of these  $p$ -modular TQFT's that permit us to express the coefficients of the Alexander Polynomial of a 3-manifold evaluated at a  $p$ -th root of unity in terms of the respective traces over covering cobordisms.

Our original motivation to construct and study the TQFT's of this article is that they are likely to describe the constant order in the cyclotomic integer expansion of the Reshetikhin-Turaev TQFT's at a  $p$ -th root of unity  $\zeta_p$ . Remarkably, the latter TQFT's are obtained by entirely different methods using quantum groups [25]. The fact that the associated invariants of closed 3-manifolds live in the ring of cyclotomic integers  $\mathbb{Z}[\zeta_p]$  is established in [22, 18]. The

---

<sup>1</sup> 2000 Mathematics Subject Classification: Primary 57R56, 57M27; Secondary: 17B10, 17B37, 17B50, 20B30, 20C30, 20C20.

coefficients of the corresponding expansions are dominated in the case of homology 3-spheres by the universal Ohtsuki invariants [24], all of which are of finite type [15]. In the simplest case, Murakami [22] finds that the first order Ohtsuki invariant coincides with the Casson invariant. In [5] Glimmer proves from the integrality of the invariants for closed 3-manifolds that also the Reshetikhin-Turaev TQFT's for "grounded" cobordisms have presentations in the cyclotomic integers.

Yet, until now, the TQFT content of the Ohtsuki invariants that is entailed by the corresponding expansions is as mysterious as it is of obvious importance for structural insights into these invariants. Particularly, TQFT interpretations would allow us to find non-trivial extensions of the invariants to 3-manifolds with Bettinumbers  $\beta_1 > 0$ , gain greater flexibility in their computation, understand underlying representations of mapping class groups, and find useful geometrical interpretations.

Inevitably, any progress in this direction requires, before anything, a thorough analysis of the constant order contributions. In particular, these determine the ranks of the  $\mathbb{Z}[\frac{1}{p}]$ -modules associated to the surfaces, and the complete TQFT can be understood as a deformation of the constant order theory. Moreover, they compute the first non-vanishing Ohtsuki invariant for a given manifold, which is of higher than constant order for  $\beta_1 > 0$ . The argument that leads us to believe that the TQFT's considered here dominate the constant contribution of the Reshetikhin-Turaev theory is roughly as follows:

Assuming that a specific block structure can be preserved when choosing an integral basis (as it is the case for  $p = 5$  [13]) it follows that the mapping class group representations contain the commutators of the Torelli groups in their kernels. Hence, we generally expect the structure of the constant order TQFT to be described in terms of the invariant theory of the symplectic groups, or certain abelian extensions thereof, acting on homology. The Lefschetz components of the Frohman-Nicas theory provide a family of TQFT's which are characterized in precisely this way. For a fixed surface they generate all of the fundamental highest weight representations of the symplectic groups, and hence, by taking tensor products and decompositions, all possible representations, including the ones emerging from the Reshetikhin-Turaev theory. To be more precise, we will also take non-trivial null space subquotients of the  $p$ -reduced homological TQFT's, given that the Reshetikhin-Turaev spaces all have non-degenerate inner forms. In [13] we give concrete evidence for our conjecture by establishing an explicit isomorphism between the constant order quantum theory at  $r = 5$  and one of the TQFT's over  $\mathbb{F}_5$  constructed in this article.

From this point we are able to derive a number of pertinent topological results by combining the trace formulae for the Alexander Polynomial or Reidemeister-Milnor Torsion of a 3-manifold with identifications of the same TQFT's as constant orders of a quantum group generated TQFT. One of the more interesting consequences is that due to the functorial nature of the Reshetikhin-Turaev TQFT's it will turn out that specific parts of the Alexander Polynomial are independent of the choice of a 1-cocycle. In turn this imposes non-trivial constraints on a group  $G$  in order for it to be realized as  $G = \pi_1(M)$  the fundamental group of a closed, compact 3-manifold, see [13] and below.

More generally, we expect that TQFT identifications of the sort described above and proved in [13] yield important insights into the role of the Alexander Polynomial of a 3-manifold in the context of extending finite type invariants to manifolds with non-trivial homology. Investigations into the low order behavior thus far suggest that for 3-manifolds with  $\beta_1 > 0$  the first order Ohtsuki invariant is given by polynomials in torsion coefficients rather than the Casson invariant, while the constant order coefficient will be zero rather than one.

Further, from the general axiomatics it follows that TQFT interpretations will deliver formulae for the Ohtsuki invariants (mod  $p$ ) for 3-manifolds given in terms of Heegaard splittings. For example, in first non-trivial order, we infer that the Casson invariant for integral homology spheres is determined by the constant order TQFT, which is expectedly described by symplectic representation theory. In fact, Morita [20] has worked out such formulae for the Casson invariant in terms of  $\mathrm{Sp}(2g; \mathbb{Z})$ -invariant theory. In special cases we do indeed reproduce Morita's expressions from the expansions of the quantum theory [13].

Other topological applications, that will be further explained and developed in [6], include the fact that we can estimate the cut numbers of 3-manifolds by their quantum orders. Here the cut number of a 3-manifold  $M$  can be defined as the maximal rank of a (non-abelian) free group onto which  $\pi_1(M)$  can be mapped, and the quantum order is the order of the lowest non-vanishing Ohtsuki invariant.

Let us summarize next the main results of this paper along with an outline of the construction of the  $p$ -modular TQFT's and symmetric group representations:

We revisit the Frohman-Nicas theory  $V^{FN}$  in Section 2 as an integral TQFT acting on lattices associated to surfaces in lieu of vector spaces. All lattices are endowed with a non-degenerate inner form. We isolate in Lemma 3 integral representations of the symmetric groups  $S_n$  by restricting the  $\mathrm{Sp}(2g; \mathbb{Z})$  action on the lattices to the parts of the Weyl subgroup that stabilize the weight spaces. In Section 3 we recall the decomposition of  $V^{FN}$  according to the Lefschetz  $\mathrm{sl}_2$ -action from [12] yielding as summands the irreducible, integral TQFT's  $V_Z^{(k)}$ . The corresponding symmetric group representations on the weight sublattices are identified in Lemma 4 with standard Specht modules  $S_Z$  for two-row Young diagrams.

In Section 4 we consider the (rank preserving) reductions  $V_p^{(k)}$  to the number field  $F_p$ . They inherit a natural inner form with values in  $F_p$  from the one of  $V^{FN}$  by restriction to sublattices and subsequent  $p$ -reduction. This process typically introduces quite non-trivial degeneracies. Consequently, we obtain new TQFT's  $\overline{V}_p^{(k)}$  from the  $V_p^{(k)}$  by dividing out the null spaces of the induced inner forms over  $F_p$ . Although the resulting TQFT's now act on ordinary vector spaces over  $F_p$  they can no longer be consistently described as  $p$ -reductions of TQFT's over  $\mathbb{Z}$ . However, by definition, they have a non-degenerate inner form and will turn out to be irreducible.

By Howe duality we have in characteristic zero that the operators of the Lefschetz  $\mathrm{sl}_2$ -action map only isomorphic TQFT's and  $\mathrm{Sp}(2g; \mathbb{Z})$ -representations to each other. However, in the  $p$ -modular reduction we find, quite surprisingly, that powers of the  $E$ -operator define non-trivial maps between symmetric and symplectic group representations that were irreducible and inequivalent over  $\mathbb{Z}$ , see Corollary 6 and Corollary 7. In Section 4 we discover further that these maps can be arranged into a differential complex.

In Section 5 we prove one of the main results of this article, namely that the so constructed complexes are acyclic, i.e., have no homology. This applies also when we put the respective irreducible,  $p$ -modular TQFT's at the ends. Thus the exact sequences we found are really resolutions of the  $\overline{V}_p^{(j)}$ . The arrows between the TQFT's are natural transformations, meaning, collections of linear maps between the vector spaces that commute with the actions of the cobordisms.

**Theorem 1** For any prime  $p \geq 3$  and integer  $k$  with  $0 < k < p$  we have an exact sequence of natural transformations of TQFT's

$$\cdots \rightarrow V_p^{((i+1)p + k_{i+1})} \rightarrow V_p^{(ip + k_i)} \rightarrow \cdots \rightarrow V_p^{(2p - k)} \rightarrow V_p^{(k)} \rightarrow \overline{V}_p^{(k)} \rightarrow 0; \quad (1)$$

where we set  $k_i = k$  for even  $i$  and  $k_i = p - k$  for odd  $i$ .

In the proof we first view (1) as a direct sum of differential complexes of corresponding  $S_n$ -representations by restriction to the individual weight spaces. We thus obtain a sequence of maps between the  $p$ -reductions of Specht modules  $S_p$  ending in the standard irreducible subquotient  $D_p$ . The proof of Theorem 1 thus amounts to showing that we have resolutions of the  $D_p$  by standard Specht modules.

Such types of resolutions appear to be new in the theory of modular symmetric group representations and interesting by themselves. The precise statement is given in Theorem 10. The proof uses very recent results of Kleshchev and Sheth [16] on the precise modular structure of the  $S_p$ .

An obvious application of such resolutions is that we can write the irreducible characters for the  $D_p$  as alternating sums of the ordinary  $S_p$ -characters. They are given in Corollary 11 of Section 6. Theorem 1 implies an analogous relation for traces over cobordisms for the different TQFT's.

The need to prove such character formulae was indeed what initially triggered our efforts to find the TQFT resolutions from Theorem 1. Specially, they allow us, in combination with trace formulae from [3] and [12], to express the Alexander Polynomial of a 3-manifold evaluated at a  $p$ -th root of unity as a sum of traces over the covering cobordisms evaluated in the irreducible  $p$ -modular TQFT's. Assuming that the latter describe the Reshetikhin-Turaev theory in lowest order the precise formula that we give next thus provides the crucial link with torsion coefficients. See also (56).

**Theorem 2** Let  $(M) \in \mathbb{Z}[x; x^{-1}]$  be the Alexander Polynomial of a closed, compact, oriented 3-manifold  $M$  with respect to an epimorphism  $\pi: H_1(M) \twoheadrightarrow \mathbb{Z}$ . Choose a surface  $\Sigma$  dual to  $\pi$  and denote by  $C$  the cobordism from  $\Sigma$  to itself obtained by removing a neighborhood of  $\Sigma$  from  $M$ .

Let  $\overline{\pi}_p(M)$  be the reduction of  $(M)$  obtained by substituting  $x = \zeta_p$ , with  $\zeta_p$  a  $p$ -th root of unity, and taking the  $\mathbb{Z}$  coefficients modulo  $p$ . Then

$$\overline{\pi}_p(M) = \sum_{k=1}^{p-1} (-1)^{k-1} [k]_p \text{trace}(\overline{V}_p^{(k)}(C)) \in \mathbb{F}_p[p]:$$

In Section 6 we also deal with the closely related problem of computing the dimensions of the modules  $D_p$  and  $\overline{V}_p^{(k)}$  as alternating sums of the dimensions of the  $S_p$  and  $V_p^{(k)}$  respectively. As a serendipitous result we find in the case of  $p = 5$  a set of curious identities between Fibonacci numbers and Catalan numbers, which we could not find elsewhere in the literature. For the TQFT's the dimensions are more conveniently compared using the formalism of fusion algebras or Verlinde algebras.

All TQFT's constructed up to this point contain the Torelli groups  $I_g$  in their kernels. In Section 7, however, we show how to "glue" the  $\text{Sp}(2g; \mathbb{Z})$ -representations of  $V^{(j)}$  and  $V^{(j+3)}$  for some  $j$  together in order to obtain an extension which represents  $I_g$  non-trivially. Over  $\mathbb{Z}$  this yields faithful representations of  $Q_g = \Gamma_g / K_g$ , where  $K_g$  is the normal subgroup of the mapping class group  $\Gamma_g$  generated by Dehn twists along bounding cycles. We will prove that this extension over  $\mathbb{Z}$  is compatible with the  $p$ -modular resolutions above, and thus factors through the irreducible subquotients. In summary we find the following.

**Theorem 3** For  $0 < k < p - 3$  there are indecomposable representations  $\overline{U}_p^{(k)}(\Gamma_g)$  of the mapping class group  $\Gamma_g$  that factor through  $Q_g = \Gamma_g / K_g$  but represent the Torelli group  $I_g$  non-trivially. There is a short, non-split exact sequence as follows:

$$0 \rightarrow \overline{V}_p^{(k+3)}(\Gamma_g) \rightarrow \overline{U}_p^{(k)}(\Gamma_g) \rightarrow \overline{V}_p^{(k)}(\Gamma_g) \rightarrow 0: \quad (2)$$

We summarize the different TQFT's of this paper, the constructions connecting them, and the included modules of the symmetric groups  $S_n$  in the following table:

SL(2;R)-Lefschetz Decomposition	$V_Z^{FN}$	Jacobian TQFT [3]	-	Permutation Modules $M$
	$\frac{\mathbb{H}}{\mathbb{H}_5}$	Fully reducible over $\mathbb{Z}$		
p-Reduction	$V_Z^{(k)}$	Lefschetz Component, irreducible over $\mathbb{Z}$	-	Specht Modules $S_Z$
	$\frac{\mathbb{H}}{\mathbb{H}_5}$			
Null space quotient	$V_p^{(k)}$	Reducible over $F_p$ with inner form.	-	p-Specht Modules $S_p$
	$\frac{\mathbb{H}}{\mathbb{H}_5}$			
Johnson-Moriai Extension	$\overline{V}_p^{(k)}$	Irreducible over $F_p$ . Not $\mathbb{Z}$ -liftable, but resolutions in $V_p^{(j)}$ 's	-	Simple $S_n$ -modules $D_p$
	$\frac{\mathbb{H}}{\mathbb{H}_5}$			
	$\overline{U}_p^{(k)}$	Indecomposable with two factors $\overline{V}_p^{(k)}$ and $\overline{V}_p^{(k+3)}$		

We conclude with the precise statements of some of the results in [13] on the Reshetikhin Turaev TQFT at a 5-th root of unity  $\zeta_5$ , which we already alluded to in the beginning of the introduction, and which inspired the constructions carried out in this article. There  $V_5^{RT}$  is identified as a halfprojective TQFT in the sense of [10] over  $\mathbb{Z}[\zeta_5]$  with parameter  $x = \zeta_5^{-1}$ . Hence we can consider the constant order reduction  $V_{0,5}^I$  as a TQFT over  $F_5$ , also halfprojective but with parameter  $x = 0$ .

Theorem 4 ([13])  $\overline{V}_{0,5}^I(\mathfrak{g})$  and  $\overline{U}_5^{(1)}(\mathfrak{g})$  are equivalent as  $\mathfrak{g}$ -representations over  $F_5$ .

This identification is remarkable and by no means obvious since the Reshetikhin Turaev TQFT is obtained via the representation category of quantum  $\mathfrak{sl}_2$  at a 5-th root of unity, which bears little resemblance to the way we construct  $\overline{U}_5^{(1)}$ . Combining Theorem 4 with the character formula from Theorem 2 we derive the equation

$$2^{++}_{,5}(\mathfrak{M}) = \det(\ker(\cdot)) \quad xV_5^{RT}(\mathfrak{M}) + O(x^3) : \quad (3)$$

The interesting observation about this formula for  $\chi_1(\mathfrak{M}) > 1$  is that  $2^{++}_{,5}(\mathfrak{M})$  depends up to order  $x^3$  only on the Reshetikhin Turaev invariant and is thus independent of the choice of the cyclic covering determined by  $\mathfrak{M}$ . Now, the Alexander Polynomial is entirely determined by the 3-manifold group, that is,  $\chi_1(\mathfrak{M}) = \chi_1(G)$  with  $G = \pi_1(\mathfrak{M})$  and  $\chi_1 : G \rightarrow \mathbb{Z}$ . Hence, the fact that the  $x^2$ -coefficient is (up to a sign) independent of the choice of  $\mathfrak{M}$  translates into a condition on  $G$ . In separate work we will exploit this apparently new and non-trivial criterion for excluding families of algebraic groups  $G$  as 3-manifold groups.

Acknowledgements: I am much indebted to Alexander Kleshchev for pointing out his results in [16], and to Gordon James and Alain Reuter for checking the arguments in Section 5. I thank Ronald Solomon, Stephen Rallis, Adalbert Kerber, and Charles Frohman for comments

and interest. Also, I have benefited much from discussions with Pat Gilmer on related integral TQFT's and their applications. Finally, I would like to express my thanks to Tomotada Ohtsuki and Hitoshi Murakami for hospitality at the RIMS, Kyoto, where final parts of this paper were completed, and for the opportunity to present and discuss the results.

This paper was typeset in  $\text{\LaTeX}$  supported by `diagrams.sty` by Paul Taylor and a modification of `young.sty` by Jorg Knappen.

## 2. Frohman-Nicas TQFT's over $\mathbb{Z}$

In [3] Frohman and Nicas construct a TQFT that allows the interpretation of the Alexander Polynomial as a weighted Lefschetz trace. The vector spaces are the cohomology rings of the Jacobians of the surfaces. In [12] we extract a natural Lefschetz  $\text{SL}(2; \mathbb{R})$ -action on these spaces, with respect to which this TQFT is equivariant. Let us describe in this section the basic ingredients of the Frohman Nicas theory, reviewing also the conventions of [12].

We denote by  $\text{Cob}_3^{22\text{fr}}$  the category of evenly framed cobordisms between connected standard surfaces  $\Sigma_g$ . These are essentially the cobordisms obtained by an even number of surgeries, see Lemma 10, [12]. The Frohman Nicas TQFT is then given as a functor

$$V^{\text{FN}} : \text{Cob}_3^{22\text{fr}} \rightarrow \text{SL}(2; \mathbb{R}) \text{ mod } ; \quad (4)$$

for which

$$V^{\text{FN}}(\Sigma_g) = \bigvee H_1(\Sigma_g)$$

for the surface  $\Sigma_g$  of genus  $g$ . For each surface we pick a symplectic basis  $a_1, \dots, a_g; b_1, \dots, b_g$  for  $H_1(\Sigma_g)$  and introduce the inner form  $\langle \cdot, \cdot \rangle$ , for which this basis is orthonormal. We denote  $i : H_1(\Sigma_g) \rightarrow H_1(\Sigma_{g+1})$  the inclusion map compatible with the choice of bases and the construction of  $\Sigma_{g+1}$  from  $\Sigma_g$  by handle addition.

The bases naturally provide lattice bases for  $\bigvee H_1(\Sigma_g; \mathbb{Z})$ , given by monomials in the  $a_i$  and  $b_i$ . Moreover, the inner form extends to the exterior product to make the monomial basis an orthonormal one. We also denote by  $\omega_g = \bigwedge_{i=1}^g a_i \wedge b_i \in \bigwedge^2 \bigvee H_1(\Sigma_g; \mathbb{Z})$  the symplectic form. The  $\text{sl}(2; \mathbb{R})$ -action on  $\bigvee H_1(\Sigma_g)$  is given in terms of the standard generators  $E, F$ , and  $H$  as

$$H = (j - g) \cdot \bigwedge^2 \bigvee H_1(\Sigma_g); \quad E = \wedge^2 \omega_g; \quad F = E : \quad (5)$$

From the explicit actions of the generators in [12] we see that the lattices  $\bigvee H_1(\Sigma_g; \mathbb{Z})$  are preserved by the maps  $V^{\text{FN}}(M)$  for cobordisms  $M$  in  $\text{Cob}_3^{22\text{fr}}$ . Moreover, also the standard generators of  $\text{sl}(2; \mathbb{R})$  preserve the lattices. Hence also the universal enveloping algebra over  $\mathbb{Z}$  generated by the operators in (5), which we shall (abusively) denote by  $\text{sl}(2; \mathbb{Z})$ , acts on the lattices. We denote the respective equivariant, lattice TQFT functor by:

$$V_{\mathbb{Z}} : \text{Cob}_3^{22\text{fr}} \rightarrow \text{sl}(2; \mathbb{Z}) \text{ mod } ; \quad (6)$$

More specially, the extension of the mapping class group in  $\text{Cob}_3^{22\text{fr}}$  factors under  $V_{\mathbb{Z}}$  through a split  $F_2 = \mathbb{Z} \oplus \mathbb{Z}$  extension of  $\text{Sp}(2g; \mathbb{Z})$ , acting in the natural way on  $\bigvee H_1(\Sigma_g; \mathbb{Z})$ . We also denote by  $\text{sp}(2g; \mathbb{Z})$  the algebra over  $\mathbb{Z}$  generated by the standard  $\text{Sp}(2g; \mathbb{R})$  Lie algebra generators. It is not hard to show that in this representation it coincides with the group algebra  $\mathbb{Z}[\text{Sp}(2g; \mathbb{Z})]$ . Besides the mapping class groups the other generators of  $\text{Cob}_3^{22\text{fr}}$  are the handle attachment cobordisms  $H_g^+ : \Sigma_g \rightarrow \Sigma_{g+1}$  and  $H_g : \Sigma_{g+1} \rightarrow \Sigma_g$  with actions given by

$$V_{\mathbb{Z}}(H_g^+) : \bigwedge^i(\cdot) \wedge a_{g+1} \quad \text{and} \quad V_{\mathbb{Z}}(H_g) = V_{\mathbb{Z}}(H_g^+) ; \quad (7)$$

where we use the notation  $i$  also for the inclusion  $H_1(\mathfrak{g}) \hookrightarrow H_1(\mathfrak{g}+1)$  extended to the exterior powers.

The structure of  $V_Z$  and the dual  $\mathfrak{sl}(2; \mathbb{Z})$ -action can be better understood if we decompose the lattices  $V_Z(\mathfrak{g})$  according to  $\mathfrak{sp}(2g; \mathbb{Z})$ -weights. We write an  $\mathfrak{sp}(2g; \mathbb{Z})$ -weight in the standard basis  $= \sum_{i=1}^g \lambda_i \epsilon_i$  as given in [7]. Specifically, we have  $h\alpha_i = h_i; h\alpha_i$  and  $h\beta_i = -h_i; h\beta_i$  for  $h \in \mathbb{Z}$ , the diagonal matrices in  $\mathfrak{sp}(2g; \mathbb{Z})$ .

We denote by  $r_g = \{ \sum_{i=1}^g \lambda_i \epsilon_i : \lambda_i \in \mathbb{Z} \}$  the set of possible weights of vectors in  $H_1(\mathfrak{g})$ . This yields a decomposition into weight spaces denoted as follows.

$$V_Z(\mathfrak{g}) = \bigoplus_{\lambda \in r_g} W_Z(\lambda; \mathfrak{g}) \quad (8)$$

For a given weight  $\lambda \in r_g$  let  $N(\lambda) = \{i : \lambda_i = 0\}$  and  $n(\lambda) = |N(\lambda)|$ . A special vector  $w(\lambda) \in W_Z(\lambda; \mathfrak{g})$  is given by

$$w(\lambda) = w_1(\epsilon_1) \wedge \dots \wedge w_g(\epsilon_g) \in V_{g-n(\lambda)} H_1(\mathfrak{g}) \quad \text{where} \quad \begin{cases} w_i(\epsilon_1) = a_i \\ w_i(\epsilon_{-1}) = b_i \\ w_i(0) = 1 \end{cases}$$

Let  $e_i$  be generators of a 2-dimensional lattice  $\mathbb{Z}\epsilon_i$ , and  $L_Z^n = \mathbb{Z}\epsilon_i$  the lattice of rank  $2^n$ . Given  $N(\lambda) = \{j_1, \dots, j_{n(\lambda)}\}$  with  $j_1 < \dots < j_{n(\lambda)}$  we thus define maps

$$\begin{aligned} \phi : L_Z^{n(\lambda)} &\rightarrow W(\lambda; \mathfrak{g}) \\ \phi : e_1 \otimes \dots \otimes e_n &\mapsto w(\lambda) \wedge \phi_{j_1}(\epsilon_1) \wedge \dots \wedge \phi_{j_n}(\epsilon_n) \end{aligned} \quad (9)$$

where  $n = n(\lambda)$  and  $\begin{cases} \phi_j(\epsilon_j) = a_j \wedge b_j \\ \phi_j(\epsilon_i) = 1 \end{cases}$

The lattices  $L_Z^n$  also have a natural inner product for which the monomials  $e_1 \otimes \dots \otimes e_n$  form an orthonormal basis, and carry a natural  $\mathfrak{sl}(2; \mathbb{Z})$ -action given by  $E e_i = e_i, F e_i = -e_i, H e_i = e_i$ , and  $E e_{-i} = F e_{-i} = 0$ , for which also  $E = F$  and  $H = H$ .

**Lemma 1** The  $\phi$  are  $\mathfrak{sl}(2; \mathbb{Z})$ -equivariant isomorphisms of lattices with inner forms.

**Proof:** They are obviously isomorphisms of lattices as they map orthonormal bases to each other. It is also easy to see that the  $H$ -weight for a monomial is  $n(\lambda) + 2 \sum_{i=1}^n \lambda_i$  on both sides of (9). Now,  $E$  is multiplication with  $\epsilon_j = a_j \wedge b_j$ . Clearly,  $w(\lambda) \wedge a_j \wedge b_j = 0$  if  $j \notin N(\lambda)$  so we multiply only with  $\epsilon_j = a_j \wedge b_j = \phi_{j_1}(\epsilon_1) \wedge \dots \wedge \phi_{j_n}(\epsilon_n)$ . The  $E$ -equivariance then follows from  $\phi_j(\epsilon_j) \wedge \phi_j(\epsilon_j) = 0$  and  $\phi_j(\epsilon_i) \wedge \phi_j(\epsilon_i) = \phi_j(\epsilon_i)$ .  $F$ -equivariance follows from  $F = E$ . ■

Although not of immediate necessity for the main result let us record here also how the actions of morphisms  $V_Z(M)$  on the lattices  $L_Z^n$  look like. We give them in terms of generators of  $\mathfrak{sp}(2g; \mathbb{Z})$  and the handle attaching maps. We introduce  $\mathfrak{sl}(2; \mathbb{Z})$ -equivariant morphisms

$$\text{coev}_k = I_{k-1} \otimes \text{coev} : L_Z^n \rightarrow L_Z^{n+2} \quad (10)$$

$$\text{where } \text{coev} : 1 \mapsto (e_1 \otimes e_1 - e_2 \otimes e_2) \quad \text{and } I_n = \text{id}_{L_Z^n} :$$

$$\text{and } \text{ev}_k = (\text{coev}_k) : L_Z^{n+2} \rightarrow L_Z^n$$

The obey relation

$$ev_k \circ \text{coev} = I \quad \text{and} \quad ev_k \circ \text{coev} = 2 \cdot I :$$

For a map  $\phi : W_Z(\cdot; g) \rightarrow W_Z(\cdot^0; g^0)$  we denote  $\phi = \phi_0^{n(\cdot)} : L_Z^{n(\cdot)} \rightarrow L_Z^{n(\cdot^0)}$ . Moreover, as in [7], we denote the standard generators  $e_i$  and  $f_j$  of  $sp(2g; \mathbb{Z})$  for simple roots  $\alpha_i = \alpha_i - \alpha_{i+1}$  for  $i < g$  and  $\alpha_g = 2\alpha_g$ . We have  $e_i a_{i+1} = a_i$ ,  $e_i b_i = b_{i+1}$ , and  $e_i v = 0$  for all other basis vectors  $v$  if  $i < g$ . Also  $e_g b_g = e_g$  and  $e_g v = 0$  for all others. This further determines the action of the other generators with  $f_i = e_i$ .

Denote now by  $e_{i;}$  the restriction  $e_i : W_Z(\cdot; g) \rightarrow W_Z(\cdot + \alpha_i; g)$ , where we put  $e_{i;} = 0$  if  $\alpha_i + \alpha_{i+1} \notin r_g$ . Let us also denote the restriction of the handle attaching map  $H = V_Z(H_g^+) : W_Z(\cdot; g) \rightarrow W_Z(\cdot + \alpha_{g+1}; g+1)$ . The following is the result of a straight forward calculation.

Lemma 2 For  $2r_g$  and  $n = n(\cdot)$  we have, when  $i < g$ ,

$$e_{i;} = \begin{pmatrix} I_n & \text{for } (\alpha_i; \alpha_{i+1}) = (0; 1) \\ I_n & \text{for } (\alpha_i; \alpha_{i+1}) = (-1; 0) \\ \text{coev}_k & \text{for } (\alpha_i; \alpha_{i+1}) = (-1; 1) \text{ with } j_{i-1} < i < j_k \\ ev_k & \text{for } (\alpha_i; \alpha_{i+1}) = (0; 0) \text{ with } i = j_k; i+1 = j_{k+1} \\ 0 & \text{elsewise} \end{pmatrix} ;$$

$$e_{g;} = \begin{pmatrix} I_n & \text{for } \alpha_g = 1 \\ 0 & \text{elsewise} \end{pmatrix} ; \quad \text{and} \quad H = I_n :$$

Another prominent, and for our purposes more important, action on the lattices  $V_Z(\cdot_g)$  is that of subgroups of the Weyl group  $W_g = (F_2)^g \circ S_g$  of  $Sp(2g; \mathbb{Z})$ , where  $S_g$  denotes the symmetric group in  $g$  letters. The  $j$ -th  $F_2$ -generator of  $W_g$  acts on weights by changing the sign of  $\alpha_j$ . It is realized as a subgroup  $\tilde{W}_g \subset Sp(2g; \mathbb{Z})$  by an extension  $1 \rightarrow (\mathbb{Z} = 2\mathbb{Z}) \rightarrow \tilde{W}_g \rightarrow W_g \rightarrow 1$ , with  $\tilde{W}_g = (F_4)^g \circ S_g$ . The  $F_4$ -generators are given by the "S-matrices"  $S_j \in Sp(2g; \mathbb{Z})$ , see [12], defined by  $S_j a_j = b_j$ ,  $S_j b_j = a_j$  and  $S_j a_i = a_i$  and  $S_j b_i = b_i$  for  $i \neq j$ . We specify two relevant representations of  $\tilde{W}_k$ :

1.  $L_Z^k = (\mathbb{Z}^2)^k$ : This action factors through  $\tilde{W}_k \rightarrow S_k$  the symmetric group, which acts canonically on the lattice by permutation of factors.

2.  $M_Z^k = (\mathbb{Z}^2)^k$ : Here the  $j$ -th  $F_4$ -factor is represented by the matrices  $S_j = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  acting on the  $j$ -th factor of the tensor product. The action of  $S_k$  on  $M_Z^k$  is the canonical representation multiplied by the alternating representation, i.e.,  $(v_1 \otimes \dots \otimes v_k) = \text{sign}(\sigma) v_{\sigma^{-1}(1)} \otimes \dots \otimes v_{\sigma^{-1}(k)}$ .

We have natural subgroups  $\tilde{W}_{g-n} \subset \tilde{W}_n \subset \tilde{W}_g$ , for which the right coset

$$C_n^g = \tilde{W}_g \cdot \tilde{W}_{g-n} \cdot \tilde{W}_n = S_g \cdot S_{g-n} \cdot S_n ;$$

is identified with the set of subsets  $A \subset \{1, \dots, g\}$  of size  $|A| = n$ . We denote

$$W_Z(n; g) = \bigcup_{A \subset C_n^g} W_Z(A; g) = \bigcup_{\substack{M \\ 2r_{g-n}(\cdot) = n}} W_Z(\cdot; g) ; \quad (11)$$

$$\text{where } W_Z(A; g) = \bigcup_{\substack{M \\ 2r_g(N(\cdot)) = A}} W_Z(\cdot; g) : \quad (12)$$



Clearly, the summands of  $V_Z = \bigoplus_n W_Z(n; g)$  are invariant under the  $\mathbb{W}_g$ -action for each  $n$ . These subrepresentations are identified next as induced representations.

Lemma 3 For every  $n$  with  $0 \leq n \leq g$  there is a natural isomorphism of  $\mathbb{W}_g$ -modules

$$\text{Ind}_{\mathbb{W}_{g-n} \times \mathbb{W}_n}^{\mathbb{W}_g} M_Z^{g-n} \otimes L_Z^n \cong W_Z(n; g) :$$

This map is an  $\mathfrak{sl}(2; \mathbb{Z})$ -equivariant isometry.

Proof: For  $N_n = \{fg \mid n+1 \leq g \leq g\}$  the  $\mathbb{W}_{g-n} \times \mathbb{W}_n$ -module is readily identified via the isomorphism

$$M_Z^{g-n} \otimes L_Z^n \cong W_Z(N_n; g) : e_1 \otimes \dots \otimes e_{n-g} \otimes 1 \otimes [1, \dots, n-g, 0, \dots, 0] \quad (1)$$

with the submodule  $W_Z(N_n; g) \cong H_1(\mathfrak{g})$ , where the action is defined by restricting the action of  $\mathbb{W}_g$ . We next define a natural section

$$: C_n^g \rightarrow S_g \times \mathbb{W}_g : A \mapsto A \quad (13)$$

as follows. Let  $A = f_1, \dots, f_g$  with  $n = |A|$ . There is a unique permutation  $\sigma \in S_g$  such that  $A = \sigma(f_{g-n+1}, \dots, f_g)$ ,  $\sigma(1) < \sigma(2) < \dots < \sigma(g-n)$ , and  $\sigma(g-n+1) < \sigma(g-n+2) < \dots < \sigma(g)$ . Clearly, we have

$$A : W_Z(N_n; g) \cong W_Z(A; g) :$$

The induced representation by definition the space of all maps  $f : \mathbb{W}_g \rightarrow M_Z^{g-n} \otimes L_Z^n$  such that  $f(\gamma) = \gamma^{-1} \cdot f(1)$  for  $\gamma \in \mathbb{W}_{g-n} \times \mathbb{W}_n$ , equipped with the left regular  $\mathbb{W}_g$ -action  $(\gamma f)(\gamma^{-1}) = f(1)$ . Now, every  $\gamma \in \mathbb{W}_g$  has a unique decomposition  $\gamma = \gamma_{(N_n)}$ , with  $\gamma_{(N_n)} \in \mathbb{W}_{g-n} \times \mathbb{W}_n$ . Thus we may identify the induced representation with the space of maps  $\bar{f} : C_n^g \rightarrow M_Z^{g-n} \otimes L_Z^n$ , setting  $\bar{f}(A) = f(\sigma(A))$  and hence  $f(\gamma) = \gamma^{-1} \cdot \bar{f}(\sigma(N_n))$ . The isomorphism is now given by

$$\text{Ind} = \text{Map}(C_n^g; M_Z^{g-n} \otimes L_Z^n) \rightarrow W_Z(n; g) : \bar{f} \mapsto (f) = \sum_{A \in C_n^g} \bar{f}(A) \quad (14)$$

An inverse is obtained by mapping  $v \in W_Z(A; g)$  to  $\sigma_A^{-1}(v) \in f_{A,g}$ , where  $f_{A,g}(B) = 1$  for  $A = B$  and 0 otherwise. In order to show that it is equivariant let  $\bar{f}$  be an arbitrary map  $C_n^g \rightarrow W_Z(N_n; g)$ ,  $\gamma \in \mathbb{W}_g$ , and  $\bar{f}^0 = \gamma^{-1} \cdot \bar{f}$ . Let  $A \in C_n^g$  and define  $\sigma_A : W_{g-n} \times \mathbb{W}_n$  by  $\sigma_A^{-1} A = \sigma^{-1}(\sigma(A))$ . Hence we have  $\bar{f}^0(A) = f^0(\sigma(A)) = (\gamma f)(\sigma(A)) = f(\sigma^{-1}(\sigma(A))) = f(\sigma^{-1}(\sigma(A))) = \sigma_A^{-1}(\sigma(A)) = \sigma_A^{-1}(\sigma(A))$ . Thus

$$\begin{aligned} (\bar{f}^0) &= \sum_{A \in C_n^g} \bar{f}^0(A) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) \\ &= \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) \\ &= \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) = \sum_{A \in C_n^g} \sigma_A^{-1}(\sigma(A)) \end{aligned}$$

which is what we needed to show. Isometry of  $\bar{f}$  is with respect to the natural inner product on  $\text{Map}(C_n^g; M_Z^{g-n} \otimes L_Z^n)$  given by

$$\langle \bar{f}, \bar{g} \rangle = \sum_{A \in C_n^g} \langle \bar{f}(A), \bar{g}(A) \rangle \quad (15)$$

given the inner form on  $M_Z^{g,n} = L_Z^n$ . Also, as  $L_Z^n$  is an  $sl(2;Z)$ -module also  $\text{Map}(C_n^g; M_Z^{g,n} = L_Z^n)$  is. Both, equivariance and isometry, follows immediately from the form of the isomorphism in (14) and the fact that the  $\mathcal{A}_\alpha$  are isometric equivariant maps. ■

The relation of the isomorphisms in Lemma 3 to the ones in Lemma 1 is given by the restrictions of  $\chi^{-1}$  to the weight spaces

$$\chi_{N(\cdot)}^{-1} = \chi^{-1} : W(\cdot; g) \rightarrow W(\cdot; g) : W(N_n; g) \quad (16)$$

Here  $\chi$  denotes the "sign-content" of a weight  $\lambda$  defined by

$$\chi_{N(\cdot)}^{-1} = \chi_{N(\cdot)}^{-1} = \prod_{j=1}^{g \cdot X^n(\cdot)} \chi_{N(\cdot)(j)}^{-1} = 1 + \dots + g \cdot n(\cdot) :$$

### 3. Lefschetz Decompositions and Specht Modules

As in [12] we consider the decomposition of the Frohman Nicas TQFT according to  $SL(2;R)$ -representations:

$$V_Z^{FN} = \bigoplus_{j=1}^M V_j = V^{(j)} ;$$

where  $V_j$  is the  $j$ -dimensional irreducible representation of  $SL(2;R)$ . Note, that the convention we use here for the superscript in  $V^{(j)}$  is shifted by one from the one used in [12]. For weights we follow the notations of [7]. The sublattices of the irreducible TQFT components can be defined as the  $SL(2;R)$  lowest weight spaces

$$V_Z^{(j)}(\cdot) = \{v \in V_Z(\cdot) : Fv = 0 \text{ and } Hv = (j-1)v\} = V_{g \cdot j+1}^{(j)}(H_1(\cdot; Z) \setminus \ker(F)) :$$

The representation of  $Sp(2g;Z)$  on  $V_Z^{(j)}(\cdot)$  is irreducible of fundamental highest weight  $\$_{g \cdot j+1} = 1 + \dots + g \cdot j+1$  with height weight vector

$$w(\$_{g \cdot j+1}) = a_1 \wedge \dots \wedge a_{g \cdot j+1} = \$_{g \cdot j+1}(e^{j+1}) :$$

The possible weights in this representation are given by

$$r_g^{(j)} = \{ \lambda \in 2r_g : n(\lambda) = j-1 \text{ and } n(\lambda) = j-1 \pmod{2g} :$$

We obtain an analogous weight space decomposition

$$V_Z^{(j)}(\cdot) = \bigoplus_{2r_g^{(j)}}^M W_Z^{(j)}(\cdot; g) = \bigoplus_{2r_g^{(j)}}^M L_Z^{n(\cdot)} \setminus \ker(F) \setminus \ker(H + j-1) : \quad (17)$$

Lemma 4 The spaces  $W_Z^{(j)}(\cdot; g)$  are as  $S_{n(\cdot)}$ -modules isomorphic to the standard irreducible Specht modules  $S^{[a;b]}$  for the two-row Young-diagram

$$[a;b] = \frac{h_{n(\cdot)} + j-1}{2} ; \frac{n(\cdot) - j+1}{2} :$$

Proof: Although this may be standard we shall provide a proof to fix conventions. We largely follow here the definitions and notations of [8]. First we note that  $\ker(H + j - 1)$  is naturally isomorphic to the permutation module  $M^{[a,b]}$ , where  $n() = a + b$  and  $j = a - b + 1$ . The isomorphism  $M^{[a,b]} \cong \ker(H + j - 1) \setminus \mathbb{Z}^{n()}$  maps a tabloid  $ftg = \frac{i_1 \ i_2 \ \dots \ i_a}{j_1 \ \dots \ j_b}$  to the basis vector  $e_{ftg} = e_{i_1} \ e_{i_2} \ \dots \ e_{i_n}$  with  $i_k = +$  if  $k \in \{j_1, \dots, j_b\}$  and  $i_k = -$  if  $k \in \{i_1, \dots, i_a\}$ . It is obvious that the  $e_{ftg}$  indeed span  $\ker(H + j - 1)$ . Consider a tableau  $t = \begin{array}{|c|} \hline i_1 \\ \hline j_1 \\ \hline \end{array} \begin{array}{|c|c|} \hline j_b \ j_{b+1} \\ \hline j_b \\ \hline \end{array} \begin{array}{|c|} \hline i_a \\ \hline \end{array}$  of shape  $[a;b]$ . Let  $C_t$  be the column stabilizer group and

$$t = \sum_{\sigma \in C_t} \text{sign}(\sigma) = \sum_{k=1}^b (1 \ (i_k; j_k)) \quad (18)$$

the signed column sum. We set  $e_t = t e_{ftg}$ . The Specht module  $S^{[a,b]}$  is the space generated by these  $e_t$  and we want to show that it coincides with  $\ker(F)$ . The easy part is to show  $S^{[a,b]} \subseteq \ker(F)$ . Using that  $F$  commutes with  $t$  we compute  $F e_t = F t e_{ftg} = t F e_{ftg} = \sum_{k=1}^b t e_{ft_k g}$ , where  $ft_k g$  is the tabloid of shape  $[a+1; b-1]$ , in which we have removed  $j_k$  from the bottom row and added to the top row. As a result  $(1 \ (i_k; j_k)) ft_k g = 0$ , hence  $t ft_k g = 0$  so that  $F e_t = 0$ .

In order to prove  $\ker(F) \subseteq S^{[a,b]}$  we proceed by induction.  $\ker(F)$  on  $M^{[a,b]}$  is given by  $\ker(F^2)$  on  $M^{[a-1;b]}$ . We have a map

$$\ker(F^2) \setminus M^{[a-1;b]} \rightarrow \ker(F) \setminus M^{[a,b]} : x \mapsto e \quad x \mapsto Fx :$$

It is easy to see that this map is an isomorphism. We use first that  $Fx \in \ker(F) \setminus M^{[a,b-1]}$  and hence by induction  $Fx = \sum_{t \in T} b_t e_t$ . Here  $T$  is the set of tableaux of the form  $t =$

$$\begin{array}{|c|} \hline i_1 \\ \hline j_1 \\ \hline \end{array} \begin{array}{|c|c|} \hline j_{b-1} \ j_b \\ \hline j_{b-1} \\ \hline \end{array} \begin{array}{|c|} \hline i_a \\ \hline \end{array}$$

with numbers from  $\{2, 3, \dots, a+b\}$ . Now as  $b-1 < a$  we have that  $i_a$  is not permuted by  $C_t$ . Thus if we denote  $E_j = I^{j-1} E \ I^{N-j}$  we have that  $E_{i_a}$  commutes with  $t$ . Let  $h_t = t E_{i_a} e_{ftg}$ . Now  $F h_t = F E_{i_a} t e_{ftg} = [F; E_{i_a}] t e_{ftg} + E_{i_a} F t e_{ftg} = H_{i_a} t e_{ftg}$  since  $e_t \in \ker(F)$  as shown above. Now,  $H_{i_a}$  commutes with  $t$  and  $H_{i_a} e_{ftg} = e_{ftg}$  by construction. Hence  $F h_t = e_t$ . In other words  $S^{[a,b-1]} \subseteq \text{im}(F)$ . Consider now  $y = x - \sum_{t \in T} b_t h_t$ . We thus have  $y \in \ker(F) \setminus M^{[a-1;b]}$  so that by induction  $y = \sum_{s \in S} c_s e_s$ , where  $S$  denotes the tableau of the form  $s = \begin{array}{|c|} \hline k_1 \\ \hline l_1 \\ \hline \end{array} \begin{array}{|c|c|} \hline k_b \ k_{b+1} \\ \hline l_b \\ \hline \end{array} \begin{array}{|c|} \hline k_{a-1} \\ \hline \end{array}$  with all numbers in  $\{2, 3, \dots, a+b\}$ . Inserting everything we find

$$z = e \quad x \mapsto Fx = \sum_{s \in S} c_s e_s - \sum_{t \in T} b_t (e \quad h_t - e \quad e) :$$

Now, it is not hard to see that  $e_s = (e \quad e_s)$  with  $s = \begin{array}{|c|} \hline k_1 \\ \hline l_1 \\ \hline \end{array} \begin{array}{|c|c|} \hline k_b \ k_{b+1} \\ \hline l_b \\ \hline \end{array} \begin{array}{|c|c|} \hline k_{a-1} \ 1 \\ \hline \end{array}$  given that  $a-1 < b$  so that 1 is not permuted by  $C_s$ . Moreover,  $(e \quad h_t - e \quad e) = (1 \ (l_1; i_a)) (e \quad h_t) (1 \ (l_1; i_a)) t E_{i_a} e_{ftg} = \hat{t} e_{ftg} = e_{\hat{t}}$ , where  $\hat{t} = \begin{array}{|c|c|} \hline 1 \ i_1 \\ \hline i_a \ j_1 \\ \hline \end{array} \begin{array}{|c|c|} \hline j_{b-1} \ j_b \\ \hline j_{b-1} \\ \hline \end{array} \begin{array}{|c|c|} \hline i_{a-1} \ 1 \\ \hline \end{array}$ . Thus  $z = \sum_{s \in S} c_s e_s + \sum_{t \in T} b_t e_{\hat{t}}$  so that  $z \in S^{[a,b]}$ . ■

Combining Lemma 3 with Lemma 4 and using notation  $W_z^{(j)}(A; g)$  and  $W_z^{(j)}(n; g)$  with  $n \equiv j-1$  and  $n \equiv j-1 \pmod{2}$  analogous to (11) so that  $\bigvee_z^{(j)}(g) = \bigwedge_n W_z^{(j)}(n; g)$  we find the following structure.

Corollary 5 For every  $1 \leq j \leq n - g$  with  $n - j \equiv 1 \pmod{2}$  there is an isomorphism of  $\mathbb{W}_g$ -modules

$$W_Z^{(j)}(n; g) : \text{Ind}_{\mathbb{W}_{g-n} \times \mathbb{W}_n}^{\mathbb{W}_g} M_Z^{g-n} S^{[\frac{n+j-1}{2}, \frac{n-j+1}{2}]} = ! \quad W_Z^{(j)}(n; g) :$$

Let us also describe the  $\mathfrak{sp}(2g; \mathbb{Z})$ -generators on the vectors  $e_t$  spanning the Specht modules  $W_Z^{(j)}(n; g)$ . To this end it is convenient to use tableaux in which entries are taken from the set  $N(n)$  rather than  $1; \dots; n - g$ , related to the standard ones by application of  $N(n)$ . We denote the set of these tableaux by  $T^{(j)}(n)$ .

Lemma 5 The  $\mathfrak{sp}(2g; \mathbb{Z})$ -generators act on the tableau vectors of Specht modules in the  $V_Z^{(j)}$ -TQFT as follows. For the  $e_i$ , with  $1 \leq i \leq g - 1$  we have:

1. If  $(i; i+1) = (0; 1)$  then  $e_i, e_t = e_s$  where  $s \in T^{(j)}(n + i)$  is obtained from  $t \in T^{(j)}(n)$  by replacing the label  $i \in N(n)$  in  $t$  by the label  $i+1 \in N(n + i)$ .
2. If  $(i; i+1) = (-1; 0)$  then  $e_i, e_t = e_s$  where  $s \in T^{(j)}(n + i)$  is obtained from  $t \in T^{(j)}(n)$  by replacing the label  $i+1 \in N(n)$  in  $t$  by the label  $i \in N(n + i)$ .
3. If  $(i; i+1) = (-1; 1)$  we have  $f_i, i+1g = N(n + i) \setminus N(n)$  and  $e_i, e_t = e_s$  where  $s \in T^{(j)}(n + i)$  by adding a column  $\begin{bmatrix} i \\ i+1 \end{bmatrix}$  to  $t \in T^{(j)}(n)$ .
4. If  $(i; i+1) = (0; 0)$  so that  $f_i, i+1g = N(n) \setminus N(n + i)$  then  $e_i, e_t$ , with  $t \in T^{(j)}(n)$ , is

(a) 0 if the labels  $i$  and  $i+1$  occur in columns of height 1.

(b)  $2e_s$  if  $t$  is given by adding the column  $\begin{bmatrix} i \\ i+1 \end{bmatrix}$  to  $s$ .

(c)  $e_s$  if  $i$  and  $i+1$  occur in different columns of height 2.

Here  $s$  is obtained from  $t$  by replacing the double column  $\begin{bmatrix} k & 1 \\ i & i+1 \end{bmatrix}$  by  $\begin{bmatrix} 1 \\ k \end{bmatrix}$

(d)  $e_s$  if  $i$  is in column of height 2 and  $i+1$  in columns of height 1, where  $s$  is obtained from  $t$  by deleting the column  $\begin{bmatrix} i \\ k \end{bmatrix}$  and replacing  $\begin{bmatrix} i+1 \end{bmatrix}$  by  $\begin{bmatrix} k \end{bmatrix}$

(e)  $e_s$  if  $i+1$  is in column of height 2 and  $i$  in columns of height 1, where  $s$  is obtained from  $t$  by deleting the column  $\begin{bmatrix} k \\ i+1 \end{bmatrix}$  and replacing  $\begin{bmatrix} i \end{bmatrix}$  by  $\begin{bmatrix} k \end{bmatrix}$

All other cases of positions of  $i$  and  $i+1$  follow from the symmetry properties of the vectors  $e_t$  under permutations of columns or within columns. Since  $N(n + g) = N(n)$  if  $n + g \equiv 2 \pmod{g}$  we have that  $e_{-g}$  acts as identity also on the vectors  $e_t$ . Similarly,  $H$  acts as identity.

#### 4. $F_p = \mathbb{Z}/p\mathbb{Z}$ -Reductions and the Sequences $C_{p,k}$

For the remainder of this article let  $p$  be an odd prime number. Since the TQFT's  $V_Z^{(j)}$  are defined over free  $\mathbb{Z}$ -modules (lattices) we naturally obtain TQFT's  $V_p^{(j)}$  over the number field  $F_p$  by setting

$$V_p^{(j)}(g) = V_Z^{(j)}(g) \cdot pV_Z^{(j)}(g)$$

Now, each  $V_Z^{(j)}(\mathfrak{g})$  inherits a non-degenerate inner product as a sublattice of  $V_{H_1(\mathfrak{g};Z)}$ . This, however, will in general degenerate if we consider the  $p$ -reduction  $h; i_p : V_p^{(j)}(\mathfrak{g}) \rightarrow F_p$ . We denote the corresponding null space as follows.

$${}^4V_p^{(j)}(\mathfrak{g}) = \{v \in V_p^{(j)}(\mathfrak{g}) : hv; w i_p = 0 \ \forall w \in V_p^{(j)}(\mathfrak{g})\}$$

The elements are represented by vectors  $v \in V_Z^{(j)}(\mathfrak{g})$  for which  $hv; w i \in pZ$  for all  $w \in V_Z^{(j)}(\mathfrak{g})$  although  $v \notin pV_Z^{(j)}(\mathfrak{g})$ .

Lemma 6 There are well defined TQFT's  ${}^4V_p^{(j)}$  and  $\overline{\overline{V}}_p^{(j)}$  which assign to a surface  $\mathfrak{g}$  the  $F_p$ -vector spaces

$${}^4V_p^{(j)}(\mathfrak{g}) \quad \text{and} \quad \overline{\overline{V}}_p^{(j)}(\mathfrak{g}) = V_p^{(j)}(\mathfrak{g}) \cdot {}^4V_p^{(j)}(\mathfrak{g}) :$$

Proof: We note that since for a cobordism  $M$  the map  $V_Z(M)$  commutes with  $E$  we have that  $V_Z(M)$  commutes with  $F = E$  and hence also maps the  $V_Z^{(j)}(\mathfrak{g})$  to themselves. Thus if  $v_i \in V_Z^{(j)}(\mathfrak{g}_i)$  for  $i = 1, 2$ ,  $M$  is a cobordism from  $\mathfrak{g}_1$  to  $\mathfrak{g}_2$ , and  $v_1$  represents a vector in  ${}^4V_p^{(j)}(\mathfrak{g}_1)$ , then  $hv_2; V_Z^{(j)}(M)v_1 i = hv_2; V_Z^{(j)}(M)v_1 i \in pZ$  as  $V_Z^{(j)}(M)v_2 \in V_Z^{(j)}(\mathfrak{g}_2)$ . ■

We extend the previous notations to the weight spaces  $\overline{\overline{W}}_p^{(j)}(\mathfrak{g})$ ,  $\overline{\overline{W}}_p^{(j)}(\mathfrak{g})$ ,  $\overline{\overline{W}}_p^{(j)}(n; \mathfrak{g})$ , etc.. Since the weight spaces are all orthogonal to each other these subspaces can be defined also as the null spaces from the respective restriction of the inner forms. Now, also the Specht modules  $S$  for a diagram  $\mathfrak{g} = [a; b]$  inherit an inner form from the permutation module  $M$ , which is via the isometry compatible with the one on the weight spaces. As in the standard literature, e.g., [8], we set

$$D_p = S_p \cdot {}^4S_p \quad \text{where} \quad {}^4S_p = S_p \setminus S_p^?$$

and  $S_p$  is the  $p$ -reduction of  $S$ . They are related to irreducible TQFT's as follows.

Lemma 7 Let  $p \geq 3$  be a prime. The TQFT's  $\overline{\overline{V}}_p^{(j)}$  are irreducible over  $F_p$  and the weight spaces are identified as  $\overline{\overline{W}}_p^{(j)}$ -modules by equivariant isomorphisms:

$$\overline{\overline{V}}_p^{(j)} : \text{Ind}_{\overline{\overline{W}}_p^{(j)}(\mathfrak{g}_n)}^{\overline{\overline{W}}_p^{(j)}(\mathfrak{g})} M_p^{g, n} \xrightarrow{D_p^{[\frac{n+j-1}{2}, \frac{n-j+1}{2}]}} \overline{\overline{W}}_p^{(j)}(n; \mathfrak{g}) : \quad (19)$$

Proof: The isomorphism in (19) follows from the definitions and properties of  $\overline{\overline{V}}_p^{(j)}$ . We first show that the spaces  $\overline{\overline{W}}_p^{(j)}(n; \mathfrak{g})$  are irreducible with respect to the semidirect product  $X_g$  of the Cartan algebra  $Z[h_g]_p \cong \mathfrak{sl}(2g; Z)$  and the algebra of the Weyl group  $Z[\overline{\overline{W}}_g] \cong Z[S_p(2g; Z)]$ . For any vector  $v = \sum v_i \in \overline{\overline{W}}_p^{(j)}(n; \mathfrak{g})$  the action of  $h_g$  shows that each weight component  $v_i \in \overline{\overline{W}}_p^{(j)}(\mathfrak{g})$  has to lie in  $X_g v$ . Each  $\overline{\overline{W}}_p^{(j)}(n; \mathfrak{g})$  is a module of a symmetric group  $S_n \rtimes \overline{\overline{W}}_g \cong X_g$  which is equivalent to  $D_p^{[\frac{n+j-1}{2}, \frac{n-j+1}{2}]}$ . It now follows from Theorem 4.9 in [8] that these representations are irreducible. In fact, as  $p \geq 3$  any two-row diagram is  $p$ -regular so that these representations are never zero, see Theorem 11.1 in [8]. In particular, if  $v \neq 0$  then  $Z[S_n]v$  is the entire module. Hence, for  $v \neq 0$  we must have  $\overline{\overline{W}}_p^{(j)}(\mathfrak{g}) \subset X_g v$  for at least one  $\mathfrak{g} \in \mathfrak{r}_g$  with  $n(\mathfrak{g}) = n$ . Since  $\overline{\overline{W}}_g$  acts transitively on all of such weights and provides isomorphisms between the weight spaces we thus have  $\overline{\overline{W}}_p^{(j)}(n; \mathfrak{g}) = X_g v$ , which implies

irreducibility since  $v$  was arbitrary. A submodule of  $\overline{V}_p^{(j)}(g)$  must therefore be a direct sum of the  $\overline{W}_p^{(j)}(n;g)$ . Each of these contains a special vector  $w_n^g = s_{g-j+1}(e_{t(n;j)})$  with  $t(n;j) = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array} \quad \begin{array}{|c|} \hline m-1 \\ \hline m \\ \hline \end{array} \quad \begin{array}{|c|} \hline n \\ \hline \end{array}$  where  $m = n - j + 1$ . Using  $2_{t(n;j)} = 2^m$  for the antisymmetrizer (18), we find  $hw_n^g; w_n^g i_p = h e_{t(n;j)}; e_{t(n;j)} i_p = 2^m h e_{ft(n;j)g}; e_{ft(n;j)g} i_p = 2^{m-6} \pmod p$  for  $p \geq 3$  so that all of these vectors are non-zero in  $\overline{W}_p^{(j)}(n;g)$ . It can be computed from the rules 3. and 4.(b) in Lemma 5 that  $e_{g-n-1} S_{g-n-1} w_n^g = w_{n-2}^g$  and  $e_{g-n+1} w_n^g = 2w_{n+2}^g$ , where  $S_1 \in W_g$  maps  $a_j$  to  $b_j$  so that with  $2 \leq 6 \pmod p$  we have non-trivial maps between all of the irreducible  $X_g$ -components. Consequently, the  $\overline{V}_p^{(j)}(g)$  are irreducible as  $sp(2g;Z)$ -representations and any sub-TQFT must assign either this space or 0 to a surface  $g$ . As in [12] we easily check that the handle attachment maps are non-trivial between these spaces so that  $\overline{V}_p^{(j)}$  does in fact contain no proper sub-TQFT.  $\blacksquare$

We next construct a sequence of maps between the  $p$ -reductions of the Specht modules, using the  $sl(2;Z)$ -actions. As before we fix  $n \geq N$  and denote the Specht module

$$S_Z^{fcg} = L_Z^n \setminus \ker(F) \setminus \ker(H + c - 1) = S;$$

where  $c = a - b + 1$  with tableau  $\begin{array}{|c|c|} \hline a & b \\ \hline \end{array} = [a; b] = [\frac{n+c-1}{2}; \frac{n-c+1}{2}]$ .

Lemma 8 Let  $p \geq 3$ ,  $c \leq 0 \pmod p$ , and  $c \geq 1; \dots; p-1 \pmod p$  such that  $c \not\equiv 0 \pmod p$ . Then

$$E^{c_0}(S_Z^{fcg}) \subset S_Z^{fc-2c_0g} + pL_Z^n;$$

Moreover,

$$E^{c_0}(S_Z^{fcg}) \not\subset pL_Z^n \quad \text{and} \quad E^p(L_Z^n) \subset pL_Z^n;$$

Proof: Now, with notation as in the proof of Lemma 19,  $e_{t(n;c-1)} \in S_Z^{fcg}$  is a cyclic vector so it suffices to show that  $E^{c_0}(e_{t(n;c-1)}) \in (S_Z^{fc-2c_0g} + pL_Z^n) \cup pL_Z^n$ . Furthermore,  $e_{t(n;c-1)} = e_{t(m;0)} e^{c-1}$  and  $E e_{t(m;0)} = 0$  so that we really need to show that  $E^{c_0}(e^{c-1}) \in (\ker(F) + pL_Z^{c-1}) \cup pL_Z^{c-1}$ , where  $c = c_0 + kp$ . We do this by induction in  $c_0$ .

For  $c_0 = 1$  we have  $c - 1 = kp$  and, using  $sl_2$ -relations,  $FE(e^{kp}) = He^{kp} + EFE^{kp} = kpe^{kp}$ . Hence, if we set  $w = ke^{kp-1}$  we find  $Fw = e^{kp}$  so that  $E(e^{kp}) = kpw \in \ker(F)$ , meaning  $E(e^{kp}) \in \ker(F) + pL_Z^{c-1}$ . Next assume the assertion is true for  $c_0$  so that  $E^{c_0}(e^{c-1}) = y + pz$  with  $Fy = 0$  and  $Hy = (c - 2c_0 - 1)y$ . We have by the binomial formula and  $Ee = e_+$  and  $E^2e = 0$  that  $E^{c_0+1}(e^c) = E^{c_0+1}(e^{c-1}e) = (E^{c_0+1}e^{c-1})e + (c_0 + 1)E^{c_0}e^{c-1} = (Ey)e + (c_0 + 1)y = y + pz$ . Thus we need to show  $t = (Ey)e + (c_0 + 1)y \in \ker(F) \cup pL_Z^c$ . We compute  $Ft = (FEy)e + (c_0 + 1)ye = (Hy)e + (c_0 + 1)ye = (c - c_0)ye = kpy \in \ker(F)$ . Also  $F(y - e) = y - e$  so that  $t - kpy \in \ker(F)$  and hence  $t \in \ker(F) \cup pL_Z^c$ .

Finally, it is not hard to see that  $he_+^{c_0} e^{c-c_0-1}; E^{c_0}(e^{c-1})i = c_0! \not\equiv 0 \pmod p$  if  $c_0 < p$  so that  $E^{c_0}(e^{c-1}) \not\in pL_Z^{c-1}$  and hence  $E^{c_0}e_{t(n;c-1)} \not\in pL_Z^n$ . Also, the binomial formula yields  $E^p = \sum_{j=0}^p E^p E^{p-j} = E^j$  so that we can conclude  $E^p = 0$  by a similar induction argument.  $\blacksquare$

In particular Lemma 8 implies that we have well defined, non-zero maps  $E^{c_0} : S_p^{fcg} \rightarrow S_p^{fc-2c_0g}$  on the respective  $p$ -reductions  $S_p^{fcg} = S_Z^{fcg} \pmod p = S_Z^{fcg} \pmod p = S_Z^{fcg} \pmod p = pL_Z^n$  using that  $pS_Z^{fcg} = S_Z^{fcg} \setminus pL_Z^n$ .

Corollary 6 For  $p$  and  $n$  as above and  $k = 1; \dots; p-1$  with  $k \equiv n+1 \pmod{2}$  there is a sequence  $C_{p;k}$  of Specht modules over  $F_p$  as follows:

$$0 \rightarrow S_p^{fn+1-2lg} \rightarrow \dots \rightarrow S_p^{f2p+kg} \xrightarrow{E^k} S_p^{f2p-kg} \xrightarrow{E^{p-k}} S_p^{fkg} \rightarrow D_p^{fkg} \rightarrow 0 : \quad (20)$$

All maps (except the first and last one) are non-zero, and any two consecutive maps compose to zero.

More precisely, we have that the  $i$ -th component of this sequence is  $C_{p;k}^{(i)} = S_p^{fip+k_i g}$ , where  $k_i = k$  if  $i$  is even and  $k_i = p-k$  if  $i$  is odd. The maps are  $E^{k_i} : C_{p;k}^{(i)} \rightarrow C_{p;k}^{(i+1)}$  so that two consecutive maps compose as  $E^{k_i} E^{k_{i+1}} = E^p = 0$ .

We have that  $C_{p;k}^{(0)} = S_p^{fkg} \rightarrow C_{p;k}^{(1)} = D_p^{fkg}$  is the (non-zero) quotient map. Now, it is clear that  $\text{im}(E) \subseteq \ker(F)^2 \subseteq S_p^{4fkg}$  given that  $E = F$ . Hence also the composite  $C_{p;k}^{(1)} \rightarrow C_{p;k}^{(0)} \rightarrow C_{p;k}^{(1)}$  is zero. In order to characterize the last index write  $\frac{n+1+k}{2} = ph + q$  with  $h \geq 0$  and  $q = 0; \dots; p-1$ . We have

$$1 = \begin{cases} q-k & \text{if } q \geq k \\ q & \text{if } q < k \end{cases} : \quad (21)$$

The maps in Corollary 6 thus extend to a sequence of the  $p$ -reductions of the induced representations from Corollary 5 as well as the weight spaces  $W_p^{(c)}(n;g)$ . The respective maps on the vector spaces  $V_p^{(c)}(g)$  are, by equivariance of the  $\psi^{(j)}$ , given by the restriction and  $p$ -reductions of the maps  $E^c$  on  $H_1(g)$ . In particular, these maps commute with the TQFT-images of the cobordisms by equivariance of  $V_Z$ .

Corollary 7 For  $p, n$  and  $k$  as above there is a sequence of natural transformations of TQFT's

$$\dots \rightarrow V_p^{((i+1)p+k_{i+1})} \rightarrow V_p^{(ip+k_i)} \rightarrow \dots \rightarrow V_p^{(2p-k)} \rightarrow V_p^{(k)} \rightarrow \overline{V}_p^{(k)} \rightarrow 0 \quad (22)$$

with  $k_i = k; p-k$  as above and any two consecutive transformations compose to zero.

## 5. Exactness of $C_{p;k}$

Exactness of the sequence  $C_{p;k}$  defined in Corollary 6 follows from the modular structure of the involved Specht modules. The irreducible factors of the  $S_p$  for two row diagrams are determined in Theorem 24.15 of [8]. For our proof we will, however, have to make use also of the precise submodule structure, which turns out to be rather rigid. We use the recent result of Kleshchev and Sheth in [16] that describes this structure precisely. In order to state it we need to introduce some more conventions and definitions.

For a Young diagram  $\lambda = [a; b]$  we set  $c = a - b + 1$ , and consider the  $p$ -adic expansion

$$c = c_0 + c_1 p + c_2 p^2 + \dots + c_r p^r \quad \text{with } c_j \in \{0; 1; \dots; p-1\} : \quad (23)$$

As in [16] we denote by  $\hat{\Lambda}$  the family of sets of integers of the form

$$I = [i_1; i_2] [i_3; i_4] \dots [i_{2u-1}; i_{2t}] \quad (24)$$

such that  $i_1 < i_2 < \dots < i_{2u} ; \quad c_{i_{2j-1}} \neq 0 \quad \text{and} \quad c_{i_{2j}} \neq p-1 : \quad (25)$

For such a set  $I \in \mathcal{A}$  we define as in [16] the number

$$I = \prod_{i \in I} (p-1) q p^i + \prod_{j=1}^{X^1} p^{i_{2j}-1} = \prod_{j=1}^{X^1} [i_{2j-1}; i_{2j}] ; \quad (26)$$

where  $\prod_{i=u}^{X^1} (p-1) q p^i + p^u$ . Also, as in [16] we introduce the smaller set  $\hat{\mathcal{A}}$  given by

$$\hat{\mathcal{A}} = \{I \in \mathcal{A} : I \text{ is big}\}; \quad (27)$$

as well as for any  $I \in \hat{\mathcal{A}}$  the function  $\chi_I$  on Young diagrams  $\lambda = [a; b]$  defined as

$$\chi_I(\lambda) = [a + \chi_I; b - \chi_I]; \quad (28)$$

As before we denote by  $D_p$  the irreducible quotient of the Specht module  $S_p$  over  $F_p$  for any two row diagram  $\lambda$ . Moreover, denote by  $F(M)$  the set of irreducible factors that occur in a composition series of a representation  $M$ . Further, let  $M$  be the smallest submodule of  $S_p$  such that  $D_p \subset F(M)$ . The description of the submodule structure in [16] uses the partial order on  $F(S_p)$  defined as  $D_{p^1} \leq D_{p^2}$  if and only if  $M_1 \subset M_2$ , i.e., if and only if  $D_{p^1} \subset M_2$ .

Theorem 8 (Corollary 3.4 of [16]) For two-row Young diagrams

1. all multiplicities of  $D_p$  in  $S_p$  are zero or one.
2.  $F(S_p) = \{D_{p^I}(\lambda) : I \in \mathcal{A} \text{ big}\}$
3.  $D_{p^J}(\lambda) \leq D_{p^I}(\lambda)$  if and only if  $J \leq I$ . ■

We define next submodules of the  $S_p$  by choosing a special subset of  $\mathcal{A}$  and  $\hat{\mathcal{A}}$ . It is defined as

$$\hat{\mathcal{A}}^0 = \{I \in \hat{\mathcal{A}} : 0 \in I\} \quad \text{and} \quad \hat{\mathcal{A}}^+ = \hat{\mathcal{A}} \setminus \hat{\mathcal{A}}^0 = \{I \in \hat{\mathcal{A}} : 0 \notin I\} \quad (29)$$

as well as  $\mathcal{A}^0 = \mathcal{A} \setminus \mathcal{A}^+$  and  $\mathcal{A}^+ = \mathcal{A} \setminus \mathcal{A}^0$ . Let us also introduce the number  $k = \min \{j \geq 1 : c_j \neq 0\}$  with  $k = 1$  if  $c = c_0$ . Hence

$$c = c_0 + c_k p^k + \dots + c_r p^r \quad \text{and} \quad [0; k) \setminus I = \emptyset; \quad \text{for any } I \in \hat{\mathcal{A}}^+ : \quad (30)$$

The latter follows since if  $0 \notin I$  we must have  $i_1 > 0$ . But with  $c_{i_1} \neq 0$  we find  $i_1 \leq k$ . For the following we assume  $c \notin 0 \pmod{p}$ , i.e.,  $c \neq 0$ . Given this we introduce for a diagram  $\lambda = [a; b]$  with  $a - b = c - 1 - 2c$  (so that  $k < 1$ ) the notation

$$\lambda^0 = [a - c; b + c] \quad \text{so that} \quad \lambda = [0; k) + \lambda^0 \quad (31)$$

For  $\lambda^0$  we thus have  $c^0 = c - 2c$  and for the  $p$ -adic expansion of  $c^0 = \sum_{j=0}^p c_j^0 p^j$  we obtain

$$\begin{aligned} c_0^0 &= p - c & c_k^0 &= c_k - 1 \\ c_i^0 &= p - 1 & \text{for } 1 \leq i < k & \\ c_i^0 &= c_i & \text{for } i > k & \end{aligned} \quad (32)$$

From these equations it is clear that  $c_0^0 \neq 0$  and  $c_k^0 \neq p - 1$  so that  $[0; k)$  is an admissible interval for  $\hat{\mathcal{A}}^0$ . In fact, it is the unique minimal interval of the special subset

$$\hat{\mathcal{A}}_0^0 = \{I \in \hat{\mathcal{A}}^0 : [0; k) \subset I\} \quad \text{and} \quad [0; k) \subset \hat{\mathcal{A}}_0^0 : \quad (33)$$



This is obvious since we must have  $i_1 = 0$  for  $0 \leq i_1$  and then the next possible  $i_2$  is  $k$ . It is also easy to see that  $c_{0;k}^0 = c_0$ , confirming the relation in (31) between  $c$  and  $c^0$  via (28). We have the following simple but crucial observation.

**Lemma 9** With  $c$  and  $c^0$  as above we have a well defined bijection with inverse

$$: \hat{A}^0 \rightarrow \hat{A}^+ : I \mapsto I \cap [0; k) \quad \text{and} \quad : \hat{A}^+ \rightarrow \hat{A}^0 : J \mapsto J \cap [0; k)$$

**Proof:** We first show that  $\hat{A}^+$  is well defined. From (33) we know that every  $I \in \hat{A}^0$  contains the special interval and is thus of the form

$$I = [0; i_2) \cup [i_3; i_4) \cup \dots \cup [i_{2u-1}; i_{2u}) \quad \text{with } i_2 \leq k :$$

Clearly,  $0 \in I$ . We further distinguish the following two cases:

**Case  $i_2 = k$  :** In this situation  $I = [i_3; i_4) \cup \dots$ . Now  $I \in \hat{A}^+$  since by (32) the coefficients of  $c$  and  $c^0$  and hence the conditions on the  $i_j$  for  $j \geq 3$  are the same.

**Case  $i_2 > k$  :** In this situation  $I = [k; i_2) \cup R$  with  $R = [i_3; i_4) \cup \dots$ . We have  $c_k \neq 0$  by assumption, and  $c_{i_2}^0 \neq c_{i_2}^0 \neq 0$  as  $i_2 > k$ . Also all intervals in  $R$  are admissible. Hence  $I \in \hat{A}^+$  again.

In a similar fashion we show that  $^{-1}$  is well defined using the fact that any  $L \in \hat{A}^+$  is by (30) of the form  $L = [j_1; j_2) \cup [j_3; j_4) \cup \dots$  with  $j_1 \leq k$ . For  $j_1 > k$  we have  $^{-1}(L) = [0; k) \cup [j_1; j_2) \cup [j_3; j_4) \cup \dots$ . This is  $\hat{A}^0$  as  $c_0^0 \neq 0$ ,  $c_k^0 \neq 0$ , and  $c_j^0 = c_j$  for  $j \geq j_1$ . In case that  $j_1 = k$  we have  $^{-1}(L) = [0; j_2) \cup [j_3; j_4) \cup \dots$ , which is again in  $\hat{A}^0$  since  $c_j^0 = c_j$  for  $j \geq j_1$ . It is obvious that  $^{-1}$  and  $^1$  are inverses of each other given (30) and (33). ■

Next we consider how the differential from (26) changes under this bijection.

**Lemma 10** We have

$$I^0 = (I) + c_0$$

**Proof:** This is a computation done according to the same cases as in the proof of Lemma 9:

**Case  $i_2 = k$  :** Then  $I^0 = c_{0;k}^0 + \sum_{j=2}^P c_{[i_{2j-1}; i_{2j})}^0$ . Now, since  $c_i^0 = p^{-1}$  for  $i = 1, \dots, k-1$  we have  $c_{0;k}^0 = (p^{-1} - c_0) + p^0 = c_0$ . We also have  $c_{[i_{2j-1}; i_{2j})}^0 = c_{[i_{2j-1}; i_{2j})}$  since the  $c_i$  are all the same with  $i_j > k$ . But  $I \cap [0; k) = \sum_{j=2}^P c_{[i_{2j-1}; i_{2j})}$  which implies the assertion.

**Case  $i_2 > k$  :** As before we write  $I = [0; i_2) \cup [i_3; i_4) \cup \dots = [0; i_2) \cup R$  and  $(I) = [k; i_2) \cup R$

$$I^0 = \sum_{i=0}^{k-1} (p^{-1} - c_i) p^i + 1 + R^0 \quad (34)$$

$$= (p^{-1} - c_0) + 1 + (p^{-1} - c_k) p^k + \sum_{i>k} (p^{-1} - c_i) p^i + R \quad (35)$$

$$= c_0 + (p^{-1} - c_k) p^k + p^k + \sum_{i>k} (p^{-1} - c_i) p^i + R \quad (36)$$

$$= c_0 + \sum_{i=k}^{k-1} (p^{-1} - c_i) p^i + p^k + R \quad (37)$$

$$= c_0 + [k; i_2) + R = c_0 + [k; i_2) \cup R = c_0 + I \cap [0; k) \quad (38)$$

which proves the assertion. ■

Corollary 9 The map  $\pi$  restricts to a bijection  $\pi : A^0 \rightarrow A^+$  with the properties that it is monotonous with respect to inclusions and

$$\pi(I^0) = \pi(I) : \quad (39)$$

Proof: From (31) we have  $b^0 = b + c_0$ . Moreover, Lemma 9 and Lemma 10 imply that  $I \in A^0$  if and only if  $I \in \pi(I) + c_0 = b + c_0 + \pi(I) = b + \pi(I) \in A^+$ . The fact that  $\pi(A) \subseteq \pi(B) \Rightarrow A \subseteq B$  is obvious. Relation (39) is immediate from Lemma 10, (28) and (31). ■

Let us turn now to the analogous relations between the modules. The set  $A^0$  contains a minimal element, namely  $I^0 = [0; h)$ , which is the smallest interval with  $h > 0$  and  $c_h \notin p - 1$ . We denote the modules

$$S_p = M_{\pi(I^0)} \quad \text{and} \quad \check{S}_p = S_p \cdot S_p :$$

By 3. of Theorem 8 we know that the modules in the composition series of  $M_{\pi(I^0)}$  are of the form  $D_p^{J^0}$ , where  $I^0 \subseteq J$ , which is equivalent to  $J \in A^0$ . Hence

$$F(S_p) = \{D_p^{I^0} : I \in A^0\} \quad \text{and} \quad F(\check{S}_p) = \{D_p^{I^0} : I \in A^+\} : \quad (40)$$

With  $h_0 = k$  and  $I^0 = (;$  we thus have  $S_p^0 = M^0$ . The bijection from Corollary 9 allows us now to show that the composition series of  $S_p^0$  and  $\check{S}_p$  yield exactly the same sets of irreducibles via the identification

$$F(S_p^0) \cong F(\check{S}_p) : D_p^{I^0} \cong D_p^{\pi(I^0)} : \quad (41)$$

Lemma 11 Let  $\pi$  and  $\pi^0$  be as above. Suppose there is a non-zero map

$$\pi : S_p \rightarrow S_p^0 \quad \text{with} \quad S_p \not\subseteq \ker(\pi) :$$

Then we have

$$\text{im}(\pi) = S_p^0 \quad \text{and} \quad \ker(\pi) = S_p :$$

Proof: Consider the map  $\bar{\pi} : \check{S}_p \rightarrow S_p^0$  defined on the quotient by  $S_p$  as well as the composite  $\bar{\pi} : \check{S}_p \rightarrow S_p^0$  with the projection. Now, since by 1. of Theorem 8 we have a disjoint union  $F(S_p^0) = F(S_p^0) \sqcup F(\check{S}_p^0)$  so that by (41)  $F(\check{S}_p^0) \setminus F(\check{S}_p) = \emptyset$ . Hence  $\bar{\pi}$  is a map between modules with no common irreducibles in their composition series so that  $\bar{\pi} = 0$ . Thus the image of  $\pi$  or  $\bar{\pi}$  must lie in  $S_p^0$ . Consider the sequence of maps

$$\pi : M = S_p \rightarrow S_p^0 \rightarrow \check{S}_p \rightarrow S_p^0 = M^0 :$$

Since all multiplicities are one we have again a disjoint union  $F(M) = F(\ker(\pi)) \sqcup F(\text{im}(\pi))$  so that either  $D_p \in F(\ker(\pi))$  or  $D_p \in F(\text{im}(\pi))$ . In the first case this would mean that  $\ker(\pi) = M$  is a submodule that contains  $D_p$  in its composition series so that by minimality  $\ker(\pi) = M$ . We had, however, assumed that  $\pi \neq 0$ . Hence  $D_p$  must be contained in the

composition series of  $\text{im}(\bar{\cdot}) = M^0$ . Again it follows by minimality that  $\text{im}(\bar{\cdot}) = M^0$ . This also means that  $\bar{\cdot}$  is a surjective map. The fact that all multiplicities are one together with (41) implies that  $\dim(S_p^0) = \dim(S_p^0)$ . Hence  $\bar{\cdot}$  must be an isomorphism. ■

We are finally in the position to establish following resolution of irreducible modules by Specht modules.

**Theorem 10** The sequences (20) from Corollary 6 are exact.

**Proof:** We begin by deriving from Theorem 8 that the first term  $S_p^{fn+1-2lg}$  in the sequence is irreducible. Here we have  $b=1$  and  $c=n+1-2l=2ph+2q-k-2l$  so that by (21)

$$b = \begin{pmatrix} q & k \\ q & \end{pmatrix} \begin{matrix} \text{if } q < k \\ \text{if } q < k \end{matrix} \quad \text{and} \quad c_0 = \begin{pmatrix} k \\ p & k \end{pmatrix} \begin{matrix} \text{if } q < k \\ \text{if } q < k \end{matrix}$$

Now for a set  $I \subseteq A$  with  $I \neq \emptyset$ ; we have  $I \subseteq p \subseteq b$ . We also need  $I \subseteq b$  and hence  $p \subseteq b$ . For  $q < k$  this condition reduces to  $p \subseteq q$ , which is not possible, and for  $q < k$  we find  $q < k$ , a contradiction as well. Hence  $A = f;g$  so that  $S_p^{fn+1-2lg} = S_p^{fn+1-2lg} = D_p^{fn+1-2lg}$  and  $S_p^{fn+1-2lg} = 0$ .

Next, we observe that all maps  $E^{k_i} : S_q^{f(i+1)p+k_{i+1}g} \rightarrow S_q^{fip+k_ig}$  in the sequence are of the type of the ones in Lemma 11 with diagrams related by (31). More precisely, we have  $c_0 = k_{i+1}$  so that  $c^0 = c - 2q = (i+1)p + k_{i+1} - 2k_{i+1} = jp + (p - k_{i+1}) = ip + k_i$ . Corollary 6 also implies that these maps are non-zero and equivariant. It now follows by induction, going from large to small  $i$ , that

$$\ker(E^{k_i}) = S_q^{f(i+1)p+k_{i+1}g} \quad \text{and} \quad \text{im}(E^{k_i}) = S_q^{fip+k_ig} :$$

For the first two maps in the sequence this is clear by irreducibility of the first module and the fact that the next map is non-zero.

Once we have proved the relation for  $E^{k_{i+1}}$  we know from  $E^{k_i}E^{k_{i+1}} = 0$  that  $S_q^{f(i+1)p+k_{i+1}g} = \text{im}(E^{k_{i+1}}) \subseteq \ker(E^{k_i})$ . We can thus apply Lemma 11 to  $E^{k_i}$  and infer the statement for  $i$  from  $i+1$ . Hence exactness holds for the terms before  $S_p^{fkg}$ .

For this last Specht module we have  $c_0 = k$  and  $c_j = 0$  for  $j > 0$ . Thus  $A^+ = f;g$  and  $A^0 = f(0;j) : p^j \xrightarrow{\frac{n+k+1}{2}g}$  using that  $(0;j) = p^j - k$  and  $b = \frac{n+k+1}{2}$ . Particularly, we have that  $S_p^{fkg} = D_p^{fkg}$  is already irreducible. The kernel of the last map already contains  $S_p^{fkg}$ , the image of the previous map, in its kernel. If the kernel was bigger the map would thus have to be zero, which is not the case. Hence exactness also holds at  $S_p^{fkg}$ . Finally, the last map is by irreducibility onto so that exactness holds through out the sequence. ■

As a result we obtain that the sequence of TQFT's in (22) of Corollary 7 is exact and this yields a resolutions of the TQFT's  $\bar{V}_p^{(k)}$  with  $0 < k < p$ , thus proving Theorem 1. The kernels and cokernels of these sequences also define TQFT's, which we denote in analogy to the symmetric group representations by  $V_p^{(j)}$  and  $\check{V}_p^{(j)}$  so that we have short exact sequences

$$0 \rightarrow V_p^{(j)} \rightarrow V_p^{(j)} \rightarrow \check{V}_p^{(j)} \rightarrow 0 : \quad (42)$$

## 6. Characters, Dimensions, and the Alexander Polynomial

An obvious application of Theorem 10 is that we can express the characters  $\chi_p^k$  and dimensions of the  $D_p^k = D_p^{\text{fkg}}$  for diagram  $s_k = [\frac{n+k-1}{2}; \frac{n-k+1}{2}]$  with  $k \equiv n+1 \pmod{2}$  and  $0 < k < p$  in terms of the ordinary characters of Specht modules  $S^\lambda$ .

Corollary 11 For  $k \equiv n+1 \pmod{2}$  and  $0 < k < p$  we have the following identity of  $S_n$ -characters

$$\chi_p^k = \sum_{i=0}^X (-1)^i \chi_{j_i}; \quad (43)$$

where  $j_i = ip + k_i$  and  $k_i = k$  for  $i$  even and  $k_i = p - k$  for  $i$  odd.

As an example we consider the special case  $p = 5$ , where  $k = 1$  or  $3$  if  $n$  is even and  $k = 2$  or  $4$  if  $n$  odd. In [26] Ryba constructs a family of irreducible so called Fibonacci representations  $R_n$  and  $R_n^0$  of  $S_n$  over  $F_5$  with Brauer characters  $\chi_n$  and  $\chi_n^0$  respectively. It follows by straightforward computation from (43) and the formulae in Definition 2 of [26] that  $\chi_n^0 = \chi_{[r;r]}$  and  $\chi_n = \chi_{[r+1;r-1]}$  if  $n = 2r$  is even, and  $\chi_n^0 = \chi_{[r+2;r-1]}$  and  $\chi_n = \chi_{[r+1;r]}$  if  $n = 2r+1$  is odd.

Corollary 12 The Fibonacci representations from [26] are

$$R_n = \begin{cases} D_5^{[r+1;r-1]} & \text{if } n = 2r \\ D_5^{[r+1;r]} & \text{if } n = 2r+1 \end{cases} \quad \text{and} \quad R_n^0 = \begin{cases} D_5^{[r;r]} & \text{if } n = 2r \\ D_5^{[r+2;r-1]} & \text{if } n = 2r+1 \end{cases}$$

We expect similar relations with the generalizations of these representations obtained by Kleshchev in [14].

The dimensions of the Specht modules are naturally given by the Catalan numbers  $C(n; j) = \frac{n!}{j! (j+1)!}$ . More precisely,  $\dim(S^{[n-b;b]}) = C(n; b)$ . Particularly, we find for the components of the sequence in (20) that  $\dim(S^{\text{fc}_{2s}g}) = C(n; b-sp)$  and  $\dim(S^{\text{fc}_{2s+1}g}) = C(n; b+(s+1)p)$ , with  $b = \frac{n+1-k}{2}$ , where we also use that  $C(n; j) = C(n; n+1-j)$ . The alternating sum of the Specht module dimensions comes out to be

$$d_k^n = \dim(D_p^{[n-b;b]}) = \sum_{s \in \mathbb{Z}} C(n; b+sp) \quad \text{provided } 0 < k = n - 2b + 1 < p; \quad (44)$$

Note, that if we extend the above formula for  $d_p^n$  to the next indices we find

$$d_0^n = 0 \text{ for odd } n; \quad \text{and} \quad d_p^n = 0 \text{ for even } n; \quad (45)$$

In order to describe generating functions for these dimensions we introduce some notation. First we write  $[n]_x = \frac{x^n - x^{-n}}{x - x^{-1}} \in \mathbb{Z}[x; x^{-1}]$  for the usual quantum integers. Moreover, we consider now the ring of cyclotomic integers  $\mathbb{Z}[\zeta_p]$  obtained  $\mathbb{Z}[x; x^{-1}]$  by imposing the relation  $\sum_{j=0}^{p-1} x^j = 0$ . It is free as a  $\mathbb{Z}$ -module of rank  $p-1$ . We also denote by  $A[\zeta_p] \subset \mathbb{Z}[\zeta_p]$  the subring invariant under conjugation  $\zeta_p \mapsto \zeta_p^{-1}$ . This is a free  $\mathbb{Z}$ -module of rank  $\frac{p-1}{2}$ . It is not hard to see that the set of  $[k]_p$  restricted to either even  $k$  or to odd  $k$  yields a  $\mathbb{Z}$ -basis for  $A[\zeta_p]$ .

Lemma 12 We have the following identity in  $A[\zeta_p]$ :

$$[2]_p^n = \sum_{\substack{0 < k < p \\ k \equiv n+1 \pmod{2}}} d_k^n [k]_p \quad (46)$$

Proof: We have by Schur-Weyl duality that  $L_2^n = V_2^n = \bigoplus_{j \equiv n-1 \pmod{2}} V_j \otimes S^{[\frac{n+j-1}{2}, \frac{n-j+1}{2}]}$ . The operator  $x^H$  with  $H \in \mathfrak{sl}(2; \mathbb{Z})$  is well defined and has trace  $\text{tr}_{V_j}(x^H) = [j]_k$ . It thus follows that  $[2]_k^n = \sum_{l \equiv 1, n+1 \pmod{2}} [l]_k C(n; \frac{n-l+1}{2})$ . Now, any such  $l$  can be uniquely written in the form  $l = k + 2sp$  or  $l = k + 2(s+1)p$  with  $s \geq 0$  and  $k \equiv n+1 \pmod{2}$  and  $0 < k < p$ . Specializing to a root of unity  $x = \zeta_p$  we have then that  $[l]_p = [k]_p$  and  $[l]_p = [k]_p$  respectively. Also  $C(n; \frac{n-l+1}{2}) = C(n; \frac{n-k+1}{2} - sp)$  and  $C(n; \frac{n-l+1}{2}) = C(n; \frac{n-k+1}{2} + (s+1)p) = C(n; \frac{n-k+1}{2} + (s+1)p)$  respectively. Hence, the terms for a fixed  $k$  are given by  $[k]_p C(n; \frac{n-k+1}{2} - sp)$  and  $[k]_p C(n; \frac{n-k+1}{2} + (s+1)p)$  for  $s \geq 0$  and  $b = \frac{n+1-k}{2}$ , which with  $s \in \mathbb{Z}$  adds up to the expression in (44). ■

Using  $[2]_k [k]_k = [k-1]_k + [k+1]_k$  and the bases of  $A[p] \subset \mathbb{Z}[p]$  we readily derive from (46) the recursion  $d_k^{n+1} = d_{k-1}^n + d_{k+1}^n$ . This translates for  $0 \leq a \leq b \leq p-2$  to

$$\dim(D_p^{[a,b]}) = \begin{cases} \dim(D_p^{[a,a-1]}) & \text{if } a = b \\ \dim(D_p^{[a-1,b]}) + \dim(D_p^{[a,b-1]}) & \text{if } 0 < a < b < p-2 \\ \dim(D_p^{[a-1,b]}) & \text{if } a = b = p-2 \end{cases} \quad (47)$$

It is easy to see from this form that the dimensions are indeed given by the number of paths through the set of diagrams with  $a \leq b \leq p-2$  as described in in [19] for general diagrams.

In the case  $p = 5$  this recursion reduces to  $\dim(R_n) = \dim(R_{n-1}) + \dim(R_{n-1}^0)$  and  $\dim(R_n^0) = \dim(R_{n-1})$  so that the dimensions are given by Fibonacci numbers. More precisely, we have  $\dim(R_n) = f_n$  and  $\dim(R_n^0) = f_{n-1}$ , where  $f_n$  are the Fibonacci numbers defined by  $f_0 = 0, f_1 = 1$  and  $f_{n+1} = f_n + f_{n-1}$ . Note that together with (44) we find interesting presentations of Fibonacci numbers in terms of alternating, 5-periodic sums of Catalan numbers:

$$\begin{aligned} f_{2r} &= C(2r; r-1) - C(2r; r-3) + C(2r; r-6) - C(2r; r-8) + \dots \\ &= C(2r+1; r-1) - C(2r+1; r-2) + C(2r+1; r-6) - C(2r+1; r-7) + \dots \end{aligned} \quad (48)$$

and

$$\begin{aligned} f_{2r+1} &= C(2r+1; r) - C(2r+1; r-3) + C(2r+1; r-5) - C(2r+1; r-8) + \dots \\ &= C(2r+2; r+1) - C(2r+2; r-3) + C(2r+2; r-4) - C(2r+2; r-8) + \dots \end{aligned} \quad (49)$$

The reader is invited to check these identities independently via recursion relations such as  $C(n+1; j) = C(n; j) + C(n; j-1)$  or via the well known generating functions of Catalan and Fibonacci numbers.

Another useful tool in the determination of dimensions are fusion algebras or Verlinde algebras, see [2]. For the quantum group  $U_q(\mathfrak{sl}_k)$  at a  $p$ -th root of unity the ring of irreducible representations yields the fusion algebra  $\mathcal{F}_p$  generated by the irreducibles  $[1], [2], \dots, [p-1]$ . Let us denote by  $\text{mult}(k; R) \in \mathbb{N}$  the multiplicity of  $[k]$  in an element  $R \in \mathcal{F}_p$  so that  $R = \sum_j \text{mult}(k; R) [k]$ . We have relations  $[1] \cdot [k] = [k]$ ,  $[p-1] \cdot [k] = [p-k]$  and  $[2] \cdot [k] = [k+1] + [k-1]$  for  $1 < k < p-1$ . Comparing this to the recursion in (47) we find that

$$\dim(D_p^{[\frac{n+k-1}{2}, \frac{n-k+1}{2}]}) = \text{mult}(k; [2]^n) : \quad (50)$$

Now, as the  $D_p$  are isomorphic to the weight spaces  $\overline{W}_p^{(k)}(\cdot; g)$  with  $n = n(\cdot)$  we find for the dimensions  $\dim(\overline{V}_p^{(k)}(\cdot; g)) = \sum_{n=0}^p 2^g \sum_n \dim(D_p^{[\frac{n+k-1}{2}, \frac{n-k+1}{2}]})$  so that we have the following Verlinde type formula.

Lemma 13

$$\dim(\overline{V}_p^{(k)}(g)) = \text{mult}(k; \mathbb{F}_p^g) \quad \text{with } \mathbb{F}_p = 2\mathbb{H}_1 + \mathbb{H}_2 \otimes \mathbb{F}_p : \quad (51)$$

Compare this to the TQFT's  $V_p^{RT}$  and  $V_p^{RT}$  of Reshetikhin-Turaev for the quantum groups  $U_q(\mathfrak{sl}_2)$  and  $U_q(\mathfrak{so}_3)$  respectively at a  $p$ -th root of unity. ( $V_p^{RT}$  is really a factor  $P$ -TQFT obtained from  $V_p^{RT}$  by restricting to odd dimensional representations). With  $\mathbb{F}_p = \mathbb{H}_2 j + 1 i^2$  and  $\mathbb{F}_p = 2\mathbb{F}_p = \mathbb{H}_k k i^2$  we have

$$\dim(V_p^{RT}(g)) = \text{mult}(1; \mathbb{F}_p^g) = 2^g \text{mult}(1; \mathbb{F}_p^g) = 2^g \dim(V_p^{RT}(g)) : \quad (52)$$

In the case  $p = 5$  these formulae allow us to efficiently compute and compare dimensions.

Lemma 14 We have for the dimensions  $D_g^{(k)} = \dim(\overline{V}_5^{(k-1)}(g))$ .

1. For even  $g$ :

$$\begin{aligned} D_g^{(1)} &= \frac{1}{2} 5^{\frac{g}{2}} f_{g-1} + f_{2g+1} & D_g^{(2)} &= \frac{1}{2} 5^{\frac{g}{2}} f_g + f_{2g} \\ D_g^{(4)} &= \frac{1}{2} 5^{\frac{g}{2}} f_{g-1} - f_{2g+1} & D_g^{(3)} &= \frac{1}{2} 5^{\frac{g}{2}} f_g - f_{2g} \end{aligned}$$

2. For odd  $g$ :

$$\begin{aligned} D_g^{(1)} &= \frac{1}{2} 5^{\frac{g-1}{2}} (f_{g-2} + f_g) + f_{2g+1} & D_g^{(2)} &= \frac{1}{2} 5^{\frac{g-1}{2}} (f_{g-1} + f_{g+1}) + f_{2g} \\ D_g^{(4)} &= \frac{1}{2} 5^{\frac{g-1}{2}} (f_{g-2} + f_g) - f_{2g+1} & D_g^{(3)} &= \frac{1}{2} 5^{\frac{g-1}{2}} (f_{g-1} + f_{g+1}) - f_{2g} \end{aligned}$$

and

$$\dim(V_5^{RT}(g)) = D_g^{(1)} + D_g^{(4)} - 8g - 0 : \quad (53)$$

Proof: Let us use a more convenient notation  $1 = \mathbb{H}_1 i, \mathbb{H}_3 i, \mathbb{H}_4 i$ , and  $\mathbb{H}_2 i$  subject to relations  $\mathbb{H}_1 i = 1 + \mathbb{H}_3 i$  and  $\mathbb{H}_2 i^2 = 1$ . These relations imply  $f_n = f_{n-1} 1 + f_n$ ,  $\mathbb{H}_3 i = 2 + \mathbb{H}_4 i$  and  $\mathbb{H}_5 i = 2 + \mathbb{H}_2 i$ . We note now that  $\mathbb{F}_5^2 = (2 + \mathbb{H}_2 i)^2 = 4 + 4\mathbb{H}_2 i + \mathbb{H}_2 i^2 = 5(1 + \mathbb{H}_2 i) = 5\mathbb{F}_5^2$  so that  $\mathbb{F}_5^g = 5^{\frac{g}{2}} (f_{g-1} + f_g)$  if  $g$  is even. From there we compute directly that for odd  $g$  we have  $\mathbb{F}_5^g = 5^{\frac{g-1}{2}} ((f_{g-2} + f_g) + (f_{g-1} + f_{g+1}))$ . Consider also  $\mathbb{H}_2 i = 1 + \mathbb{H}_4 i$ . We have  $\mathbb{H}_2 i^2 = 1 - 2\mathbb{H}_4 i + \mathbb{H}_4 i^2 = 2 - 1 + \mathbb{H}_4 i$  and thus again  $f_n = f_{n-1} + f_n$ . We find  $(2 - \mathbb{H}_4 i)^g = 2^g = f_{2g-1} + f_{2g} = f_{2g+1} - f_{2g}$ . Now, we have  $(1 + \mathbb{H}_2 i)^g = (1 + \mathbb{H}_4 i)^g$  and thus  $(1 + \mathbb{H}_2 i)^g = (1 + \mathbb{H}_4 i)^g$ . Similarly, we find  $(1 - \mathbb{H}_2 i)^g = (1 - \mathbb{H}_4 i)^g$  so that  $\mathbb{F}_5^g = \frac{1}{2} (1 + \mathbb{H}_2 i)^g + \frac{1}{2} (1 - \mathbb{H}_2 i)^g = \frac{1}{2} (1 + \mathbb{H}_4 i)^g + \frac{1}{2} (1 - \mathbb{H}_4 i)^g$ . With the previous results on  $\mathbb{F}_5^g$  and  $(2 - \mathbb{H}_4 i)^g$  we thus find a formula for  $\mathbb{F}_5^g$ , which inserted into (51) yields the asserted formulae. ■

The formula in (53) reflects the fact that  $\overline{V}_5^{(1)} - \overline{V}_5^{(4)}$  is the sum of the irreducible constituents of the  $\mathbb{F}_5$ -reductions of  $V_5^{RT}$  as TQFT's, see [13]. For larger primes  $p$  it is, however, not possible that a  $\mathbb{F}_p$ -reduction  $V_p^{RT}$  has only the  $V_p^{(j)}$  as irreducible components. This is easily seen by looking at the large  $g$  asymptotics of the dimension expressions in (51) and (52), which grow like  $k\mathbb{F}_p k^g$  and  $k\mathbb{F}_p k^g$ . Here the Perron-Frobenius eigenvalues are given by  $k\mathbb{F}_p k = \frac{p}{4 \sin^2(\frac{\pi}{p})}$  and  $k\mathbb{F}_p k = 4 \cos^2(\frac{\pi}{2p})$  and we have  $k\mathbb{F}_p k > k\mathbb{F}_p k$  if  $p > 5$ . In Section 8 we outline how tensor products will possibly remedy the situation.

The analogs of the character expansions given in Corollary 11 and relations as in Lemma 12 in the context of the corresponding TQFT's attain a topological interpretation via the Alexander

Polynomial  $(M) \in \mathbb{Z}[x; x^{-1}]$  for a compact, oriented 3-manifold  $M$  with a selected epimorphism  $\pi: H_1(M; \mathbb{Z}) \twoheadrightarrow \mathbb{Z}$ . Up to  $S$ -equivalence the cocycle defines a two-sided embedded surface  $\Sigma \subset M$  and cobordism  $C: \Sigma \rightarrow \Sigma$  obtained by removing a neighborhood of  $\Sigma$  from  $M$ . The Alexander Polynomial is then given (up to a sign, which is determined by the additional framing structure on  $M$ ) by the following identity extracted in Section 11 of [12]:

$$(M) = \text{trace}(x^H V_Z(C)) = \sum_{j=1}^X [j]_x \text{trace}(V_Z^{(j)}(C)) \quad (54)$$

By inserting  $x = \omega_p$  or  $x = \omega_p^{-1}$ , a  $p$ -th root of unity, and reducing the integer coefficients to  $\mathbb{F}_p$  we consider the image of the Alexander Polynomial under the two so defined natural maps

$$\mathbb{Z}[x; x^{-1}] \rightarrow \mathbb{F}_p[x] : x \mapsto \omega_p : (M) \mapsto {}^{\omega_p}(M) : \quad (55)$$

Now, if we insert  $x = \omega_p$  into (54) and use  $[j]_p = (\omega_p)^{j-1} (1 - \omega_p^j) [k]_p$ , where  $j_i = ip + k_i$  as before, we infer from the resolutions of TQFT's in (22) the following identities

$${}^{\omega_p}(M) = \sum_{k=1}^{\frac{p-1}{X^2}} (\omega_p)^{k-1} [k]_p \text{trace}(\overline{V}_p^{(k)}(C)) + \text{trace}(\overline{V}_p^{(p-k)}(C)) ; \quad (56)$$

$${}^{\omega_p^{-1}}(M) = \sum_{k=1}^{\frac{p-1}{X^2}} [k]_p \text{trace}(\overline{V}_p^{(k)}(C)) - \text{trace}(\overline{V}_p^{(p-k)}(C)) : \quad (57)$$

This implies Theorem 2. At a 5-th root of unity  ${}^{\omega_p}(M)$  depends only on traces of  $C$  under  $V_5^{(1)} - V_5^{(4)}$  and  $V_5^{(2)} - V_5^{(3)}$ . The former is in [13] identified with the integral semisimple reduction of the Reshetikhin-Turaev theory  $V_5^{RT}$  and implies (3).

## 7. Johnson-Morita Extensions

As before we consider for  $H = H_1(\Sigma)$  a standard basis  $a_1, \dots, a_g; b_1, \dots, b_g$  that is symplectic with respect to the standard skew form  $(;)$  and orthonormal with respect to the inner form  $\langle \cdot, \cdot \rangle$ . Let  $J \in \text{Sp}(2g; \mathbb{Z})$  be the special element defined by  $Ja_i = b_i$  and  $Jb_i = -a_i$  so that  $\langle Jx, y \rangle = \langle x, Jy \rangle$  and  $Jg^{-1}J^{-1} = g$ . Also denote by  $! = \sum_{j=1}^g a_j \wedge b_j$  the standard invariant 2-form.

For any  $x \in \bigwedge^m H$  we can now define a degree- $m$  map  $\varphi(x) : \bigwedge^V H \rightarrow \bigwedge^{V+m} H$  by  $\varphi(x)y = x \wedge y$ . From this we define another map as  $\psi(x) = (Jx) : H \rightarrow \bigwedge^m H$  of degree  $-m$ .

Lemma 15 The maps  $\varphi$  and  $\psi$  have the following properties:

1. Covariance:  $g \varphi(x) g^{-1} = \varphi(gx)$  and  $g \psi(x) g^{-1} = \psi(gx)$  for all  $x \in \bigwedge^m H$  and  $g \in \text{Sp}(2g; \mathbb{Z})$ .
2. Homomorphism:  $\varphi(x \wedge y) = \varphi(x) \varphi(y)$  and  $\psi(x \wedge y) = \psi(y) \psi(x)$ .
3. Generators:  $\varphi(!) = E$  and  $\psi(!) = F$  so that  $[E; \varphi(x)] = [F; \psi(x)] = 0$  for all  $x \in \bigwedge^m H$ .
4. Anticommutator:  $\varphi(x) \varphi(y) + \varphi(y) \varphi(x) = \langle x, y \rangle I$  for all  $x, y \in \bigwedge^m H$ .
5. Commutators:  $[E; \varphi(x)] = \varphi(x)$  and  $[F; \psi(x)] = \psi(x)$  for  $x \in \bigwedge^m H$ .





Proof: In general if  $V, W$  and  $M$  are  $G$ -modules, and  $\alpha: M \rightarrow \text{Hom}(V; W)$  a covariant map, we construct a module  $V \oplus W$  of  $M \circ G$  by letting  $(m; g)$  act on the sum  $V \oplus W$  by the block matrix  $\begin{pmatrix} g & 0 \\ (m)g & g \end{pmatrix}$  so that  $W \oplus V \oplus W$  is a submodule with subquotient  $V$ . Thus the map from Lemma 16 defines such a module for  $JM_1(m; g)$  over  $Z$ . In the  $F_p$ -reduction  $\alpha$  is invertible so that  $\alpha$  can be extended to  $JM_a(m; g)$ .

By exactness of the sequences in (22) we have that  $V_p^{(j)}(g) = V_p^{(j)}(g) \setminus \text{im}(E^{p-k})$ . Thus by (58) of Lemma 16 we have that  $(x)(V_p^{(j)}(g)) = V_p^{(j+m)}(g) \setminus \text{im}(E^{p-k-m}) = V_p^{(j+m)}(g)$ . This implies, by construction, that  $V_p^{(j)}(g) \subset V_p^{(j+m)}(g)$  is indeed a submodule. ■

Note that for  $0 < k < p$  the factors in (60) are irreducible. We also write

$$\overline{U}_p^{(j)}(m; g) = \overline{V}_p^{(j)}(g) \oplus \overline{V}_p^{(j+m)}(g) : \quad (61)$$

The case that is topologically relevant is  $m = 3$ . For this case Morita constructed in [21] a homomorphism  $\tilde{K}: g \rightarrow JM_2(3; g)$  on the mapping class group  $g$ . Its kernel is the group  $K_g$  generated by bounding cycles and its restriction to the Torelli group coincides with the Johnson homomorphism  $\gamma_2: I_g \rightarrow \frac{V_3 H}{H}$  from [9].

Theorem 14 (Johnson [9], Morita [21]) There is a finite index subgroup  $Q_g \subset JM_2(3; g) = \frac{1}{2} \frac{V_3 H}{H} \circ Sp(2g; Z)$ , and homomorphisms  $\gamma_2$  and  $\tilde{K}$  such that the following diagram is commutative with all rows and the last column exact.

$$\begin{array}{ccccccc} & & & & 0 & & \\ & & & & \downarrow & & \\ & & & & V_3 H & & \\ 0 & \longrightarrow & K_g & \longrightarrow & I_g & \xrightarrow{\gamma_2} & \frac{V_3 H}{H} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & K_g & \longrightarrow & I_g & \xrightarrow{\tilde{K}} & Q_g \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & I_g & \longrightarrow & g & \longrightarrow & Sp(2g; Z) \longrightarrow 0 \\ & & & & & & \downarrow \\ & & & & & & 0 \end{array}$$

Combining this result with Proposition 13 now yields Theorem 3, where we denoted the special module  $\overline{U}_p^{(j)}(g) = \overline{U}_p^{(j)}(3; g)$ . Note, that we obtain from the sequences in (22) similar resolutions. Particularly, for  $g = 0$ ,  $p = 5$  a prime and  $0 < k < p - 3$  there is an exact sequence of maps of  $F_p$ -modules as follows.

$$\begin{aligned} \cdots \rightarrow U_p^{((i+1)p+k_{i+1})}(g) \rightarrow U_p^{(ip+k_i)}(g) \rightarrow \cdots \rightarrow U_p^{(2p+k)}(g) \rightarrow \\ \rightarrow U_p^{(2p-k-3)}(g) \rightarrow U_p^{(k)}(g) \rightarrow \overline{U}_p^{(k)}(g) \rightarrow 0; \end{aligned} \quad (62)$$

where now  $k_i = k$  for  $i$  is even, and  $k_i = p - i - 3$  if  $i$  is odd.

The maps are alternatingly given by  $E^k \rightarrow E^{k+3}$  and  $E^{p-k} \rightarrow E^{p-k-3}$ . Note, however, that the module extensions work alternatingly in opposite ways so that the maps cannot be  $Q_g$ -equivariant. It is true that by setting  $\tilde{R}(g) = R(\hat{J}g^{-1}\hat{J}^{-1})$ , where  $\hat{J} \in g$  is a representative of

the  $J \subset \text{Sp}(2g; \mathbb{Z})$  above we can reverse the exact sequences that define an extension. Yet, even by flipping every second extension in (62) still does not yield equivariant maps.

## 8. Open Questions

1. What is the precise submodule structure of the  $V_p^{(j)}$ ? To answer this question the results from [16] should be combined with the  $\text{sp}(2g; \mathbb{Z})$  and handle mappings between the Specht modules as worked out explicitly in Lemma 5 of Section 3.
2. Are there generalizations to the resolutions in Corollary 6 to general Young diagrams? Schur-Weyl duality suggests to use an analogous  $\text{sl}(n; \mathbb{Z})$  action, identify Specht modules as lowest weight spaces, and attempt to define differential operators between different modules from the Borel algebra generators  $E_i \in \text{sl}(n; \mathbb{Z})$ . Such maps and generalized resolutions should yield many more character formulae.
3. Is there an independent proof of exactness? One should note that we the property of the modular structure from [16] that we really used was that the heads of some representations are irreducible, implying maximal elements and cyclicity. It would be interesting to use this more directly, possibly by employing also the operators  $\frac{1}{k!} E^k$  that are well defined for all  $k \geq 0$  on the lattices, though they do not map  $\ker(F)$ 's to each other are likely to play a rôle. Note also that similar such resolutions are of interest in the representation theory of  $p$ -adic groups and quantum groups.
4. Are there TQFT extensions for all of the  $\overline{U}_p^{(j)}(\mathfrak{g})$  and  $U_Z^{(j)}(\mathfrak{g})$  modules? There are two principal approaches of constructing these. One is to attempt to enlarge the gauge group used by Frohman and Nicas from  $U(1)$  to a continuous group  $G$  for which  $G^0 = [G; G] \neq 0$  but  $[G^0; G] = 0$  or at least  $[G^0; G^0] = 0$  such as the Heisenberg group or the group of upper triangular  $3 \times 3$ -matrices. The representations of the surface groups  $\pi_1(\Sigma_g)$  will factor then through  $N = \overline{[G; G]}$ , which is the natural space for the Morita extension  $\mathcal{Q}_g$  to act on, see [21]. The obvious difficulty is that such groups are non-compact so that intersection homology is not well defined. The other line of approach is to consider deformations of the algebra  $N$  considered in [12] that would lead to non-trivial contributions of the Torelli groups.
5. What are the constant order contributions of the Reshetikhin-Turaev TQFT's for other roots of unity? We noted in Section 4 that the dimensions of these vector spaces grow as  $k\mathbb{F}_p k^g$  for large  $g$ , where  $k\mathbb{F}_p k = \frac{p}{4 \sin^2(\frac{\pi}{p})}$ , but the dimensions of the modular components only with  $k\mathbb{F}_p k^g$ , where  $k\mathbb{F}_p k = 4 \cos^2(\frac{\pi}{2p})$ . For example, for  $p = 7$  these are different numbers but they are related by  $k\mathbb{F}_7 k = 2k\mathbb{F}_7 k^2 - 7k\mathbb{F}_7 k + 7$ . A way of accounting for quadratic terms is to take tensor products of TQFT's. These will lead to  $\text{Sp}(2g; \mathbb{Z})$ -representations that are no longer of fundamental weight, and also will in general have more components in the modular than the in the integral case. Integer factor can be accounted for by tensoring with TQFT's,  $C_n$ , for the simple objects of the underlying category are all invertible forming the cyclic group  $\mathbb{Z}/n\mathbb{Z}$ . We thus expect that the constant order of the Reshetikhin-Turaev TQFT at a 7-th root of unity is described as a sub and/or quotient TQFT of some  $C_2 \otimes \overline{V}_7^{(\#)} \otimes \overline{V}_7^{(\#)}$ , where the  $\#$  indicate any linear combination of degrees. In general,  $k\mathbb{F}_p k$  is a degree  $\frac{p-3}{2}$  polynomial in  $k\mathbb{F}_p k$  with integer coefficients, which gives some indication as to how this generalizes to higher primes  $p$ .

6. How do the tensor product TQFT's decompose? This question is closely related to the previous one. Since the Frohman and Nicas theories yield  $Sp(2g; \mathbb{Z})$ -representations for all fundamental weights we expect the ring of TQFT's generated by them via tensoring, summing and decomposing to contain all possible modular irreducible  $Sp(2g; \mathbb{Z})$ -representations that come from highest weight representations.
7. How are lowest orders of the Reshetikhin Turaev Invariants related to Torsion coefficients? This considers generalizations of the formula in (3). Assuming that the constant order RT-TQFT's are obtained from tensor products from the  $p$ -modular Jacobian TQFT's we expect by taking traces to find corresponding polynomial expressions in the coefficients of the Alexander Polynomial.
8. Which modular parts of the Alexander Polynomial are independent of the choice of a covering, i.e., 1-cocycle, and why? The observation made in (3) derived from the independence of the Reshetikhin Turaev invariant is expected to have generalizations. Since such independence theorems are formulated without any reference to quantum topology it is a nearby task to try to find independent proof entirely within the language of classical algebraic topology. Interpretations in terms of cyclic covering spaces, generalized signature defects, etc. would be quite useful.
9. Which criteria for 3-manifold groups can be derived from that? As already noted in the introduction the Alexander Polynomial can be computed for any group  $G$  with 1-cocycle  $\gamma: G \rightarrow \mathbb{Z}$ . Thus independence of  $\gamma$  of the  $x^2$ -coefficient in (3) can be thought of as a condition on  $G$  in order for it to be 3-manifold group  $G = \pi_1(M)$ . Typically, this criterion allows us to exclude direct sums of groups with sufficiently different Alexander Polynomials as 3-manifold groups, but applies also to more subtle examples. Details will be worked out in separate work.
10. What is the relation with other TQFT constructions that relate to the Alexander Polynomial and Torsion? Among these are the generalizations of Frohman and Nicas to  $PSU(n)$ -theories [4], and constructions by Donaldson [1] that universally reduce to the Alexander Polynomial, or polynomial expressions in its coefficients despite rather intricate non-abelian Casson-type constructions. This is somewhat parallel to what we expect for the constant order expansion of the Reshetikhin Turaev invariants as outlined above.
11. What are the possible "deformations" of these modular TQFT's? Essentially what we mean is a classification of TQFT's over  $R_{p,1} = \mathbb{F}_p[x] = x^{1-p}$  for  $1 < p$  such that their reduction under  $R_{p,1} \rightarrow R_{p,0} = \mathbb{F}_p$  is one of the given modular TQFT's. The Reshetikhin Turaev theory is in this sense an example of a deformation under the reduction  $\mathbb{Z}[p] \rightarrow \mathbb{F}_p[p] = R_{p,p-2}$  mapping the root of unity to  $p-1+x$ . The rigidity of the  $Sp(2g; \mathbb{Z})$ -representation theory and its actions on the lower parts of the filtrations of the mapping class groups appear to impose string constraints on the space of possible deformations.
12. What are connections to the universal finite type TQFT's? The TQFT's discussed here represent the Torelli groups non-trivially and thus have some overlap with the lowest orders of the TQFT extensions of [23] to homologically trivial cobordisms the famous LMO invariant from [17]. Similarly, we can consider deformations of the modular TQFT's as described previously and compare them up to higher orders of the finite type theory. These connections should shed some light on possible modular reductions of the finite type theories that have a chance to be extended to a homologically non-trivial theory.

# Bibliography

- [1] Donaldson, S.K. : Topological field theories and formulae of Casson and Meng-Taubes. Proceedings of the Kirbyfest. Geom. Topol. 2 (1999) 87{102 (electronic).
- [2] Frohlich, J., Kerler, T. : Quantum groups, quantum categories and quantum field theory. Lecture Notes in Mathematics 1542 Springer-Verlag, Berlin, 1993.
- [3] Frohman, C., Nicas, A. : The Alexander Polynomial via topological quantum field theory. Differential Geometry, Global Analysis, and Topology, Canadian Math. Soc. Conf. Proc. Vol. 12, Amer. Math. Soc. Providence, RI, (1992) 27{40.
- [4] Frohman, C., Nicas, A. : An intersection homology invariant for knots in a rational homology 3-sphere. Topology 33 (1994) no. 1, 123-158.
- [5] Gilmer, P. : Integrality for TQFT's. Preprint 2001.math.QA/0105059.
- [6] Gilmer, P., Kerler, T. : Cut Numbers via Quantum Orders for  $p = 5$ . (in preparation)
- [7] Goodman, R., Wallach, N.R. : Representations and invariants of the classical groups. Encyclopedia of Mathematics and its Applications 68 Cambridge University Press, 1998.
- [8] James, G.D. : The representation theory of the symmetric groups. Lecture Notes in Mathematics 682 Springer, Berlin, 1978.
- [9] Johnson, D. : An abelian quotient of the mapping class group  $I_g$ . Math. Ann. 249 (1980), no. 3, 225{242.
- [10] Kerler, T. : On the connectivity of cobordisms and half-projective TQFT's. Commun. Math. Phys. 198 No. 3 (1998) 535-590.
- [11] Kerler, T., Lyubashenko, V.V. : Non-semisimple topological quantum field theories for 3-manifolds with corners. Lecture Notes in Mathematics 1765, xi+ 376 pp.. Springer Verlag 2001.
- [12] Kerler, T. : Homology TQFT's and the Alexander-Reidemeister Invariant of 3-Manifolds via Hopf Algebras and Skein Theory". 50 pages 38 figures.math.GT/0008204
- [13] Kerler, T. : On the Structure of Fibonacci TQFT's. (in preparation).
- [14] Kleshchev, A. : Completely splittable representations of symmetric groups. J. Algebra 181 (1996), no. 2, 584{592.
- [15] Krieger, A., Spence, B. : Ohtsuki's invariants are of finite type. J. Knot Theory Ramifications 6 (1997), no. 4, 583{597.

- [16] Kleshchev, A. S., Sheth, J.: On extensions of simple modules over symmetric and algebraic groups. *J. Algebra* 221 (1999), no. 2, 705{722.
- [17] Le, T. T. Q., Murakami, J., Ohtsuki, T.: On a universal perturbative invariant of 3-manifolds. *Topology* 37 (1998), no. 3, 539{574.
- [18] Masbaum, G.; Roberts, J. D.: A simple proof of integrality of quantum invariants at prime roots of unity. *Math. Proc. Cambridge Philos. Soc.* 121 (1997), no. 3, 443{454.
- [19] Mathieu, O.: On the dimension of some modular irreducible representations of the symmetric group. *Lett. Math. Phys.* 38 (1996), no. 1, 23{32.
- [20] Morita, S.: Casson's invariant for homology 3-spheres and characteristic classes of surface bundles. I. *Topology* 28 (1989), no. 3, 305{323.
- [21] Morita, S.: The extension of Johnson's homomorphism from the Torelli group to the mapping class group. *Invent. Math.* 111 (1993), no. 1, 197{224.
- [22] Murakami, H.: Quantum  $SO(3)$ -invariants dominate the  $SU(2)$ -invariant of Casson and Walker. *Math. Proc. Cambridge Philos. Soc.* 117 (1995), no. 2, 237{249.
- [23] Murakami, J., Ohtsuki, T.: Topological quantum field theory for the universal quantum invariant. *Comm. Math. Phys.* 188 (1997), no. 3, 501{520.
- [24] Ohtsuki, T.: A polynomial invariant of rational homology 3-spheres. *Invent. Math.* 123 (1996), no. 2, 241{257.
- [25] Reshetikhin, N. Yu., Turaev, V. G.: Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.* 103 (1991), 547{597.
- [26] Ryba, A. J. E.: Fibonacci representations of the symmetric groups. *J. Algebra* 170 (1994), no. 2, 678{686.

The Ohio State University,  
 Department of Mathematics,  
 231 West 18th Avenue,  
 Columbus, OH 43210, U.S.A.  
 E-mail: kerler@math.ohio-state.edu