

THE CONNES-KASPAROV CONJECTURE FOR ALMOST CONNECTED GROUPS

JÉRÔME CHABERT, SIEGFRIED ECHTERHOFF, AND RYSZARD NEST

ABSTRACT. Let G be a locally compact group with cocompact connected component. We prove that the assembly map from the topological K-theory of G to the K-theory of the reduced C^* -algebra of G is an isomorphism.

CONTENTS

1. Introduction and statement of results	1
2. Some preliminaries and first reductions	3
3. Baum-Connes for continuous fields of C^* -algebras	8
4. The semi-simple case	13
5. The general case	17
6. Relations to the K -theory of the maximal compact subgroup	25
References	28

1. INTRODUCTION AND STATEMENT OF RESULTS

In this paper we give a proof of the Connes-Kasparov conjecture for almost connected groups. To be more precise, we prove the following

Theorem 1.1. *Let G be a second countable almost connected group (i.e., G/G_0 is compact, where G_0 denotes the connected component of G). Then G satisfies the Baum-Connes conjecture with trivial coefficients \mathbb{C} , i.e., if $K_*^{\text{top}}(G)$ denotes the topological K-theory of G , then the Baum-Connes assembly map*

$$\mu : K_*^{\text{top}}(G) \rightarrow K_*(C_r^*(G))$$

is an isomorphism.

It was already shown by Kasparov in [23] that the theorem is true if G is amenable. In fact, by a more recent result of Higson and Kasparov, we know that the Baum-Connes conjecture with arbitrary coefficients holds for any amenable group. By work of A. Wassermann [38], we also know that the result is true for all connected reductive linear Lie groups. More recently, Lafforgue used quite different methods to give a proof of the conjecture for all connected semi-simple groups with finite center (which are not necessarily linear). The main idea of the proof of the general result of Theorem 1.1 is to use the Mackey-machine approach, as outlined in [10], in order to reduce to the reductive case. The strategy for doing this bases

This research has been supported by the Deutsche Forschungsgemeinschaft (SFB 478).

heavily on some ideas presented in Pukánszky's recent book [34] where he reports on his deep analysis of the representation theory of connected groups. In particular the methods of his proof that locally algebraic connected real Lie groups are type I, presented on the first four pages of his book, where most enlightening.

The above theorem is actually a special case of a more general result which we shall explain below. If G is a second countable locally compact group, then by a G -algebra A we shall always understand a C^* -algebra A equipped with a strongly continuous action of G by $*$ -automorphisms of A . Let $\mathcal{E}(G)$ denote a locally compact universal proper G -space in the sense of [25] (we refer to [11] for a discussion about the relation to the notion of universal proper G -space as introduced by Baum, Connes and Higson in [3]). If A is a G -algebra, the topological K -theory of G with coefficients in A is defined as

$$K_*^{\text{top}}(G, A) = \lim_X \text{KK}_*^G(C_0(X), A),$$

where X runs through the G -compact subspaces of $\mathcal{E}(G)$ (i.e., X/G is compact) ordered by inclusion, and $\text{KK}_*^G(C_0(X), A)$ denotes Kasparov's equivariant KK -theory. If $A = \mathbb{C}$, we simply write $K_*^{\text{top}}(G)$ for $K_*^{\text{top}}(G, \mathbb{C})$.

The construction of Baum, Connes and Higson presented in [3, §9] determines a homomorphism

$$\mu_A : K_*^{\text{top}}(G, A) \rightarrow K_*(A \rtimes_r G),$$

usually called the *assembly map*. We say that G satisfies BC for A (i.e., G satisfies the Baum-Connes conjecture for the coefficient algebra A), if μ_A is an isomorphism. Theorem 1.1 is then a special case of

Theorem 1.2. *Suppose that G is any second countable locally compact group such that G/G_0 satisfies BC for arbitrary coefficients, where G_0 denotes the connected component of G . (By the results of Higson and Kasparov [21] this is in particular true if G/G_0 is amenable or, more general, if G/G_0 satisfies the Haagerup property.) Then G satisfies BC for $\mathcal{K}(H)$, H a separable Hilbert space, with respect to any action of G on $\mathcal{K}(H)$.*

It is well known that in case of almost connected groups, the topological K -theory $K_*^{\text{top}}(G, A)$ has a very nice description in terms of the maximal compact subgroup L of G . In fact, under some mild extra conditions on G , the group $K_*^{\text{top}}(G, A)$ can be computed by means of the K -theory of the crossed product $A \rtimes L$. We give a brief discussion of these relations in §6 below. As was already pointed out in [35], our results have important applications to the study of square-integrable representations. In fact, combining our results with [35, Theorem 4.6] gives

Corollary 1.3 (cf [35, Corollary 4.7]). *Let G be a connected unimodular Lie group. Then all square-integrable factor representations of G are type I. Moreover, G has no square-integrable factor representations if $\dim(G/L)$ is odd, where L denotes the maximal compact subgroup of G .*

The paper is outlined as follows: In our preliminary section, §2, we recall the main results from [9, 10] on the permanence properties of the Baum-Connes conjecture which are needed in this work. We will also use these results to perform some first reductions of the problem. In §3 we prove a result on continuous fields of actions, showing under some mild conditions on

the group G and the base space X of the field, that G satisfies the Baum-Connes conjecture with coefficients in the algebra of C_0 -sections of the field if it satisfies the conjecture for all fibres. This result will be another basic tool for the proof of our main theorem.

In §4 we are concerned with the conjecture for reductive groups. Using (and slightly extending) some recent results of Lafforgue [29] on the conjecture for semi-simple groups with finite center, we will show that the results on continuous fields obtained in §3 imply that the conjecture (with trivial coefficients) holds for all reductive groups without any linearity conditions. In §5 we will then use Pukánzsky's methods in combination with an extensive use of the permanence properties for BC to give the final steps for the proof of Theorem 1.2. In §6 we shall then discuss the implications for almost connected groups as mentioned above. Note that §6 does not contain any new material, except the conclusion drawn out of our main theorem.

Acknowledgments. The authors profited quite a lot from valuable discussions with several friends and colleagues. We are most grateful to Alain Connes who suggested (after an Oberwolfach lecture on the permanence properties for BC) to use the results of [9, 10] to attack the Connes-Kasparov conjecture with trivial coefficients. We are also grateful to Bachir Bekka, Ludwig Bröcker, Guennadi Kasparov, Guido Kings, Hervé Oyono-Oyono, Jörg Schürmann and Peter Slodowy for some useful discussions or comments.

2. SOME PRELIMINARIES AND FIRST REDUCTIONS

Let us collect some general facts which were presented in [10] – for the definitions of twisted actions and twisted equivariant KK-theory we refer to [9]. Assume that G is a second countable group and let B be a G - C^* -algebra. We say that G satisfies BC with coefficients in B if the assembly map

$$\mu_B : K_*^{\text{top}}(G, B) \rightarrow K_*(B \rtimes_r G)$$

is an isomorphism. If N is a closed normal subgroup of G , there exists a twisted action of (G, N) on $B \rtimes_r N$ such that the twisted crossed product $(B \rtimes_r N) \rtimes_r (G, N)$ is canonically isomorphic to $B \rtimes_r G$. Moreover, we can use the twisted equivariant KK-theory of [9] to define the topological K-theory $K_*^{\text{top}}(G/N, B \rtimes_r N)$ with respect to the twisted action of (G, N) on $B \rtimes_r N$, and a twisted version of the assembly map

$$\mu_{B \rtimes_r N} : K_*^{\text{top}}(G/N, B \rtimes_r N) \rightarrow K_*((B \rtimes_r N) \rtimes_r (G, N)).$$

In [9] we constructed a partial assembly map

$$\mu_{N,B}^G : K_*^{\text{top}}(G, B) \rightarrow K_*^{\text{top}}(G/N, B \rtimes_r N)$$

such that the following diagram commutes

$$\begin{array}{ccc} K_*^{\text{top}}(G, B) & \xrightarrow{\mu_{N,B}^G} & K_*^{\text{top}}(G/N, B \rtimes_r N) \\ \mu_B \downarrow & & \downarrow \mu_{B \rtimes_r N} \\ K_*(B \rtimes_r G) & \xrightarrow{\cong} & K_*((B \rtimes_r N) \rtimes_r (G, N)). \end{array}$$

Using this, the first two authors were able to prove the following extension results:

Theorem 2.1. *Assume that B is a G -algebra and let N be a closed normal subgroup of G . Let $q : G \rightarrow G/N$ denote the quotient map and assume that one of the following conditions is satisfied*

- (i) G/N has a compact open subgroup \dot{K} and for any compact subgroup \dot{C} of G/N , the group $C = q^{-1}(\dot{C})$ satisfies BC for B .
- (ii) G has a γ -element $\gamma \in \text{KK}_0^G(\mathbb{C}, \mathbb{C})$ (which is automatically true if G is almost connected), G/N is almost connected and $K = q^{-1}(\dot{K})$ satisfies BC for B , where \dot{K} is a maximal compact subgroup of G/N .

Then the partial assembly map $\mu_{N,B}^G : \text{K}_*^{\text{top}}(G, B) \rightarrow \text{K}_*^{\text{top}}(G/N, B \rtimes_r N)$ is an isomorphism. In particular, G satisfies BC for B if and only if G/N satisfies BC for $B \rtimes_r N$.

Proof. See [10, Theorem 3.3 and Theorem 3.7]. □

In order to avoid the use of twisted actions we may use the version of the Packer-Raeburn stabilization trick as given in [32, 15]:

Proposition 2.2 (cf [32, Theorem 3.4] and [15, Corollary 1]). *Assume that G is a second countable group and let N be a closed normal subgroup of G . Let (α, τ) be a twisted action of (G, N) on the separable C^* -algebra A . Then there exists an ordinary action $\beta : G/N \rightarrow \text{Aut}(A \otimes \mathcal{K})$, $\mathcal{K} = \mathcal{K}(l^2(\mathbb{N}))$, such that β is stably exterior equivalent (and hence Morita equivalent) to (α, τ) .*

Note that BC is invariant under passing to Morita equivalent actions. Thus, in order to conclude that (G, N) satisfies BC for $B \rtimes_r N$, it is enough to show that G/N satisfies BC for $(B \rtimes_r N) \otimes \mathcal{K}$ with respect to an appropriate action of G/N on $(B \rtimes_r N) \otimes \mathcal{K}$. In particular, if G/N is amenable, it follows that $\mu_{B \rtimes_r N}^G : \text{K}_*^{\text{top}}(G/N, B \rtimes_r N) \rightarrow \text{K}_*((B \rtimes_r N) \rtimes_r (G, N))$ is always an isomorphism.

In what follows we need to study the following special situation: Assume that $\alpha : G \rightarrow \text{Aut}(\mathcal{K})$ is an action of G on $\mathcal{K} = \mathcal{K}(H)$ for some separable Hilbert space H . Since $\text{Aut}(\mathcal{K}) \cong \text{PU}(H) = U(H)/\mathbb{T}1$, we can choose a Borel map $V : G \rightarrow U(H)$ such that $\alpha_s = \text{Ad } V_s$ for all $s \in G$. Since α is a homomorphism, we see that there exists a Borel cocycle $\omega \in Z^2(G, \mathbb{T})$ such that

$$V_s V_t = \omega(s, t) V_{st} \quad \text{for all } s, t \in G.$$

The class $[\omega] \in H^2(G, \mathbb{T})$ is called the *Mackey obstruction* for α being unitary. Let

$$1 \mapsto \mathbb{T} \rightarrow G_\omega \rightarrow G \rightarrow 1$$

be the central extension of G by \mathbb{T} corresponding to ω , i.e., we have $G_\omega = G \times \mathbb{T}$ with multiplication given by

$$(g, z)(g', z') = (gg', \omega(g, g')zz'),$$

and the unique locally compact group topology which generates the product Borel structure on $G \times \mathbb{T}$ (see [30]). Then the following is true

Lemma 2.3. *For each $n \in \mathbb{Z}$ let $\chi_n : \mathbb{T} \rightarrow \mathbb{T}; \chi_n(z) = z^n$. Let $\alpha : G \rightarrow \text{Aut}(\mathcal{K})$ and G_ω be as above. Then α is Morita equivalent to the twisted action (id, χ_1) of (G_ω, \mathbb{T}) on \mathbb{C} .*

Proof. Let $V : G \rightarrow U(H)$ be as in the discussion above, i.e., $\alpha_s = \text{Ad } V_s$ and $V_s V_t = \omega(s, t) V_{st}$ for all $s, t \in G$. Then it is easy to check that $\tilde{V} : G_\omega \rightarrow U(H)$ defined by $\tilde{V}_{(s,z)} = z V_s$ is a homomorphism which implements the desired equivalence on the $\mathcal{K} - \mathbb{C}$ bimodule H (we refer to [15] for an extensive discussion of Morita equivalence for twisted actions). \square

Another important result is the continuity of the Baum-Connes conjecture with respect to inductive limits of the coefficients, at least if G is exact. For this we need

Lemma 2.4. *Assume that $(B_i)_{i \in I}$ is an inductive system of G -algebras and let $B = \lim_i B_i$ be the C^* -algebraic inductive limit. Assume further that one of the following conditions is satisfied:*

- (i) *All connecting maps $B_i \rightarrow B_j$, $i \leq j \in I$ are injective, or*
- (ii) *G is exact.*

Then $B \rtimes_r G = \lim_i (B_i \rtimes_r G)$ with respect to the obvious connecting homomorphisms.

Proof. If all connecting maps are injective, we may regard each B_i as a subalgebra of B . But this implies that we also have $\overline{B_i \rtimes_r G}$ as subalgebras of $B \rtimes_r G$, and hence the inductive limit $\lim_i (B_i \rtimes_r G) = \overline{\cup \{B_i \rtimes_r G : i \in I\}}$ sits inside $B \rtimes_r G$. But it is easy to check that $\cup \{C_c(G, B_i) : i \in I\} \subseteq \lim (B_i \rtimes_r G)$ is dense in $B \rtimes_r G$.

Suppose now that G is exact. In this situation we want to reduce the proof to situation (i). Consider the canonical homomorphisms $\phi_i : B_i \rightarrow B$. Let $I_i = \ker \phi_i$ and let $I_{ij} = \ker \phi_{ij}$, where the $\phi_{ij} : B_i \rightarrow B_j$ denote the connecting homomorphisms for $j \geq i$. Of course, if $i \leq j \leq j'$ then $I_{ij} \subseteq I_{ij'}$, so for each $i \in I$ the system $(I_{ij})_{j \geq i}$ is an inductive system with injective connecting maps. It follows directly from the definition of the inductive limit that $I_i = \overline{\cup \{I_{ij} : j \geq i\}} = \lim_{j \geq i} I_{ij}$, and hence it follows from (i) that $I_i \rtimes_r G = \lim_{j \geq i} (I_{ij} \rtimes_r G)$. By exactness of G it follows that $I_i \rtimes_r G$ is the kernel of $\phi_i \rtimes_r G : B_i \rtimes_r G \rightarrow B \rtimes_r G$. By the previous discussion it follows that $I_i \rtimes_r G = \lim_{j \geq i} (I_{ij} \rtimes_r G)$ is also the kernel of the canonical homomorphism $B_i \rtimes_r G \rightarrow \lim (B_j \rtimes_r G)$. Thus, dividing out the kernels, i.e., by considering the system $(B'_i)_{i \in I}$ with $B'_i = B_i / I_i$ we conclude from another use of (i) that

$$B \rtimes_r G = \lim (B'_i \rtimes_r G) = \lim (B_i \rtimes_r G).$$

\square

As a direct consequence we obtain

Proposition 2.5. *Assume that the G -algebra B is an inductive limit of the G -algebras B_i , $i \in I$, such that G satisfies BC for all B_i . Assume further that G is exact or that all connecting homomorphisms $B_i \rightarrow B_j$ are injective. Then G satisfies BC for B .*

Proof. It follows from Lemma 2.4 and the continuity of K -theory that $K_*(B \rtimes_r G) = \lim_i K_*(B_i \rtimes_r G)$. On the other side, it is shown in [10, Proposition 7.1] that $K_*^{\text{top}}(G, B) = \lim_i K_*^{\text{top}}(G, B_i)$. Since by assumption $K_*^{\text{top}}(G, B_i) \cong K_*(B_i \rtimes_r G)$ via the assembly map, and since the assembly map commutes with the K -theory maps induced by the G -equivariant homomorphism $B_i \rightarrow B_j$, the result follows. \square

As a first application we get

Proposition 2.6. *Let G be a separable locally compact group such that G/G_0 satisfies BC for arbitrary coefficients. Then the following are equivalent:*

- (1) *For every central extension $1 \rightarrow \mathbb{T} \rightarrow \bar{G} \rightarrow G \rightarrow 1$ the group \bar{G} satisfies BC for \mathbb{C} .*
- (2) *G satisfies BC with coefficients in the compact operators $\mathcal{K} \cong \mathcal{K}(H)$ for all separable Hilbert spaces H and with respect to all possible actions of G on \mathcal{K} .*

Proof. Assume that (1) holds. Let $\alpha : G \rightarrow \text{Aut}(\mathcal{K})$ be any action of G on \mathcal{K} and let $[\omega] \in H^2(G, \mathbb{T})$ denote the Mackey obstruction for this action. Let

$$1 \rightarrow \mathbb{T} \rightarrow G_\omega \rightarrow G \rightarrow 1$$

denote the central extension determined by ω . It follows from Lemma 2.3 that α is Morita equivalent to the twisted action (id, χ_1) of (G_ω, \mathbb{T}) on \mathbb{C} . By assumption, we know that G_ω satisfies BC for \mathbb{C} . It follows from Theorem 2.1 that (G_ω, \mathbb{T}) satisfies BC for $C^*(\mathbb{T}) \cong C_0(\mathbb{Z})$, or, equivalently, that G satisfies BC for $C_0(\mathbb{Z}, \mathcal{K})$ with respect to the appropriate action of G (use Proposition 2.2). Since our group G does not satisfy directly the assumptions of Theorem 2.1, let us briefly explain how it is used: first apply part (i) of Theorem 2.1 to $N = G_0$, which implies that G satisfies BC for $C_0(\mathbb{Z}, \mathcal{K})$ if and only if every compact extension C of G_0 in G satisfies BC for $C_0(\mathbb{Z}, \mathcal{K})$, and then apply part (ii) of Theorem 2.1 to the subgroup \mathbb{T} of $C_\omega \subseteq G_\omega$.

Writing $C_0(\mathbb{Z}) = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}$, the twisted action of (G_ω, \mathbb{T}) is given by the twisted action (id, χ_n) of (G_ω, \mathbb{T}) on the n 'th summand. Let $q_1 : C_0(\mathbb{Z}) \rightarrow \mathbb{C}$ be the projection on the summand corresponding to $1 \in \mathbb{Z}$. Consider the diagram

$$\begin{array}{ccc} K_*^{\text{top}}(G, C_0(\mathbb{Z})) & \xrightarrow{\mu_{C_0(\mathbb{Z})}} & K_*(C_0(\mathbb{Z}) \rtimes_r (G_\omega, \mathbb{T})) \\ q_{1,*} \downarrow & & \downarrow q_{1,*} \\ K_*^{\text{top}}(G, \mathbb{C}) & \xrightarrow{\mu_{\mathbb{C}}} & K_*(\mathbb{C} \rtimes_r (G_\omega, \mathbb{T})). \end{array}$$

(Here the topological K -theory $K_*^{\text{top}}(G, \mathbb{C})$ is computed with respect to the twisted action (id, χ_1) of $G \cong G_\omega/\mathbb{T}$ and $\mu_{\mathbb{C}}$ denotes the twisted assembly map!) Since the vertical arrows are split-surjective and the upper horizontal arrow is bijective, it follows that the lower horizontal arrow is also bijective. Thus we see that (G_ω, \mathbb{T}) satisfies BC for \mathbb{C} with respect to the twisted action (id, χ_1) . By Morita equivalence this implies that G satisfies BC for \mathcal{K} with respect to α .

For the opposite direction assume that (2) holds. Let $1 \rightarrow \mathbb{T} \rightarrow \bar{G} \rightarrow G \rightarrow 1$ be as in (1). As explained above it follows from Theorem 2.1 that \bar{G} satisfies BC for \mathbb{C} if (\bar{G}, \mathbb{T}) satisfies BC for $C^*(\mathbb{T}) = C_0(\mathbb{Z})$. Using the stabilization trick, the latter is true if G satisfies BC for $C_0(\mathbb{Z}, \mathcal{K})$ with respect to an appropriate action of G on $C_0(\mathbb{Z}, \mathcal{K})$ which fixes the base \mathbb{Z} . Using continuity of BC, this follows easily from the fact that G satisfies BC for arbitrary actions on \mathcal{K} . \square

We also need a result on induced algebras as obtained in [10]. For this recall that if H is a closed subgroup of G and A is an H -algebra, then the induced algebra $\text{Ind}_H^G A$ is defined as

$$\text{Ind}_H^G A = \{f \in C_b(G, A) : f(sh) = h^{-1}(f(s)) \text{ and } (sH \mapsto \|f(s)\|) \in C_0(G/H)\}.$$

Equipped with the pointwise operations and the supremum-norm, $\text{Ind}_H^G A$ becomes a C^* -algebra with G -action defined by

$$s \cdot f(t) = f(s^{-1}t).$$

The following result follows from [10, Theorem 2.2]:

Theorem 2.7. *Let G , H , A and $\text{Ind}_H^G A$ be as above. Then G satisfies BC for $\text{Ind}_H^G A$ if and only if H satisfies BC for A .*

The result becomes most valuable for us when combined with the following result of [14]:

Proposition 2.8. *Suppose that H is a closed subgroup of G and B is a G -algebra. Let \widehat{B} denote the set of equivalence classes of irreducible representations of B equipped with the usual G -action defined by $s \cdot \pi(b) = \pi(s^{-1} \cdot b)$. Then B is isomorphic (as a G -algebra) to $\text{Ind}_H^G A$ for some H -algebra A if and only if there exists a G -equivariant continuous map $\varphi : \widehat{B} \rightarrow G/H$. Moreover, if $\varphi : \widehat{B} \rightarrow G/H$ is such a map, then A can be chosen to be B/I , with $I = \cap \{\ker \pi : \varphi(\pi) = eH\}$ equipped with the obvious H -action.*

As a corollary of Theorem 2.7 and Proposition 2.8 we get in particular:

Corollary 2.9. *Suppose that G is a locally compact group and B is a G -algebra which is type I and such that G acts transitively on \widehat{B} . Let $\pi \in \widehat{B}$ and let G_π denote the stabilizer of π for the action of G on \widehat{B} . Then G satisfies BC for B if and only if G_π satisfies BC for $B/\ker \pi \cong \mathcal{K}(H_\pi)$, where H_π denotes the Hilbert space of π .*

Proof. Since there is only one orbit for the G -action on \widehat{B} , it follows from results of Glimm [19], that \widehat{B} is homeomorphic to G/G_π via $sG_\pi \mapsto s \cdot \pi$. In particular, it follows that \widehat{B} is Hausdorff, which implies that $B/\ker \pi \cong \pi(B) = \mathcal{K}(H_\pi)$. The inverse of the above map is clearly a continuous G -equivariant map of \widehat{B} to G/G_π , and Proposition 2.8 then implies that $B \cong \text{Ind}_{G_\pi}^G (B/\ker \pi)$. The result then follows from Theorem 2.7. \square

We now give a short outline of the proof of Theorem 1.1. The main work is required for proving the following proposition:

Proposition 2.10. *Assume that G is a Lie group with finitely many components and let $\alpha : G \rightarrow \text{Aut}(\mathcal{K})$ be an action of G on the compact operators on some separable Hilbert space H . Then G satisfies BC for \mathcal{K} .*

The body of this paper is devoted to give a proof of this proposition by using induction on the dimension of G . It is fairly easy to see that the above proposition implies Theorem 1.2. Indeed, using the first part of Theorem 2.1 we can directly reduce to the case where G is almost connected. Hence Theorem 1.2 follows from

Proposition 2.11. *Suppose that Proposition 2.10 holds. Let G be any almost connected group and let $\alpha : G \rightarrow \text{Aut}(\mathcal{K})$ be any action of G on the compact operators on some separable Hilbert space H . Then G satisfies BC with coefficients in \mathcal{K} .*

Proof. By the structure theory of almost connected groups (e.g. see [31]) we can find a compact normal subgroup $C \subseteq G$ such that G/C is a Lie group with finitely many components. Using Theorem 2.1 we see that G satisfies BC for \mathcal{K} if and only if G/C satisfies BC for

$\mathcal{K} \rtimes C$ (with respect to an appropriate twisted action). Since C is compact, it follows that $X := (\mathcal{K} \rtimes C)^\wedge$ is discrete, and (after stabilizing if necessary), $\mathcal{K} \rtimes C \cong C_0(X, \mathcal{K})$. Let $\tilde{G} := G/C$ and let X/\tilde{G} denote the space of \tilde{G} -orbits in X . Since X is discrete, the same is true for X/\tilde{G} , and we get a decomposition $C_0(X, \mathcal{K}) \cong \bigoplus_{\tilde{G}(x) \in X/\tilde{G}} C_0(\tilde{G}(x), \mathcal{K})$. By continuity of BC (see Proposition 2.5), we conclude that \tilde{G} satisfies BC for $C_0(X, \mathcal{K})$ if and only if \tilde{G} satisfies BC for $C_0(\tilde{G}(x), \mathcal{K})$ for all $x \in X$. Using Corollary 2.9, this will follow if all stabilizers $\tilde{G}_x \subseteq \tilde{G}$ satisfy BC for \mathcal{K} . But since X is discrete, it follows that each stabilizer \tilde{G}_x contains the connected component \tilde{G}_0 of \tilde{G} . Thus, each stabilizer is a Lie group with finitely many components and the result will follow from Proposition 2.10. \square

As mentioned above, the main idea for the proof of Proposition 2.10 is to use induction on the dimension $\dim(G)$ of the Lie group G . For this we were very much influenced by Pukánszky's proof of the fact that locally algebraic groups (i.e., Lie groups having the same Lie algebra as some real algebraic group) have type I group C^* -algebras as presented in his recent book [34]. We split the induction argument into two main parts, which deal with the cases whether G is semi-simple or not. Note that even in the semi-simple case the result does not follow directly from the existent results, since all known results only work for trivial coefficients and require that the groups have finite centers.

3. BAUM-CONNES FOR CONTINUOUS FIELDS OF C^* -ALGEBRAS

Let G be a separable locally compact group. Then G is called K -exact, if the functor $A \mapsto K_*(A \rtimes_r G)$ is half-exact, that is: whenever $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$ is a short exact sequence of G -algebras, then the sequence

$$K_*(I \rtimes_r G) \rightarrow K_*(A \rtimes_r G) \rightarrow K_*(A/I \rtimes_r G)$$

is exact in the middle term. Clearly, every exact group is K -exact. Note that every almost connected group is exact by [27, Corollary 6.9].

Recall also that an element $\gamma \in \text{KK}_0^G(\mathbb{C}, \mathbb{C})$ is called a γ -element for G if there exists a locally compact proper G -space Y , a C^* -algebra D equipped with a nondegenerate and G -equivariant $*$ -homomorphism $\phi : C_0(Y) \rightarrow ZM(D)$, the center of the multiplier algebra $M(D)$ of D , and (Dirac and dual-Dirac) elements

$$\alpha \in \text{KK}_0^G(D, \mathbb{C}) \quad \beta \in \text{KK}_0^G(\mathbb{C}, D)$$

such that

$$\gamma = \beta \otimes_D \alpha \quad \text{and} \quad p_Z^*(\gamma) = 1 \in \text{RKK}_0^G(Z; \mathbb{C}, \mathbb{C})$$

for all locally compact proper G -spaces Z , where $p_Z : Z \rightarrow \{pt\}$. It is a basic result of Kasparov [24, Theorem 5.7] that every almost connected group has a γ -element and it follows also from the work of Kasparov (but see also [37, §5]) that a γ -element of G is unique and that it is an idempotent with the remarkable property that for every G -algebra B the image $\mu_B(K_*^{\text{top}}(G; B))$ of the assembly map is equal to the γ -part

$$\gamma \cdot K_*(B \rtimes_r G) := \{x \otimes_{B \rtimes_r G} j_G(\sigma_B(\gamma)) : x \in K_*(B \rtimes_r G)\}.$$

Here and below, we denote by $j_G : \text{KK}_*^G(A, B) \rightarrow \text{KK}_*(A \rtimes_r G, B \rtimes_r G)$ the (reduced) descent homomorphism of Kasparov and we denote by $\sigma_B : \text{KK}_*^G(A, D) \rightarrow \text{KK}_*(B \otimes A, D \otimes B)$ the external tensor product homomorphism (see [24, Definition 2.5]). Note that it follows from

the above discussion that a group G with γ -element satisfies BC for a given G -algebra B if and only if γ (i.e., $j_G(\sigma_B(\gamma))$) acts as the identity on $K_*(B \rtimes_r G)$. We want to exploit these facts to prove the following basic result:

Proposition 3.1. *Suppose that X is a separable locally compact space which can be realized as the geometric realization of a (probably infinite) finite dimensional simplicial complex. Let A be the algebra of C_0 -sections of a continuous field of C^* -algebras $\{A_x : x \in X\}$, and let $\alpha : G \rightarrow \text{Aut}(A)$ be a $C_0(X)$ -linear action of G on A . Assume further that G is exact and has a γ -element $\gamma \in \text{KK}_0^G(\mathbb{C}, \mathbb{C})$. Then, if G satisfies BC with coefficients in each fibre A_x , G satisfies BC for A .*

For the general notion of continuous fields of C^* -algebras and their basic properties we refer to [18, 17, 5, 26].

The idea of the proof is to show first that it holds for any closed interval $I \subseteq \mathbb{R}$. Then a short induction argument will show that it holds for any cube in \mathbb{R}^n . Then the result will follow from a Mayer-Vietoris argument. For the proof we first need the following lemma.

Lemma 3.2. *Assume that G is a K -exact group with a γ -element $\gamma \in \text{KK}_0^G(\mathbb{C}, \mathbb{C})$. Let A be a G -algebra and let $I \subseteq A$ be a G -invariant closed ideal of A . Then there is a natural six-term exact sequence*

$$\begin{array}{ccccc} (1 - \gamma) \cdot K_0(I \rtimes_r G) & \longrightarrow & (1 - \gamma) \cdot K_0(A \rtimes_r G) & \longrightarrow & (1 - \gamma) \cdot K_0(A/I \rtimes_r G) \\ & & \uparrow & & \downarrow \\ (1 - \gamma) \cdot K_1(A/I \rtimes_r G) & \longleftarrow & (1 - \gamma) \cdot K_1(A \rtimes_r G) & \longleftarrow & (1 - \gamma) \cdot K_1(I \rtimes_r G). \end{array}$$

Proof. Since G is K -exact, it follows that $A \mapsto K_*(A \rtimes_r G)$ is a homotopy invariant and half-exact functor on the category of G - C^* -algebras which also satisfies Bott-periodicity (with respect to the trivial G -action on $C_0(\mathbb{R}^2)$). Then it follows from some general arguments (e.g., see [4, Chapter IX]) that there exists a six-term exact sequence

$$\begin{array}{ccccc} K_0(I \rtimes_r G) & \longrightarrow & K_0(A \rtimes_r G) & \longrightarrow & K_0(A/I \rtimes_r G) \\ & & \uparrow & & \downarrow \\ K_1(A/I \rtimes_r G) & \longleftarrow & K_1(A \rtimes_r G) & \longleftarrow & K_1(I \rtimes_r G). \end{array}$$

We want to show that all maps in the sequence commute with multiplication with the γ -element. By the construction of the connecting maps in the above sequence as given in [4, Chapters VIII and IX], it is enough to show that for any pair of G -algebras A and B and any $y \in \text{KK}_*^G(A, B)$

$$K_*(A \rtimes_r G) \rightarrow K_*(B \rtimes_r G); x \mapsto x \otimes_{A \rtimes_r G} j_G(y)$$

commutes with multiplication with γ . But for this it is enough to show that

$$j_G(y) \otimes_{B \rtimes_r G} j_G(\sigma_B(\gamma)) = j_G(\sigma_A(\gamma)) \otimes_{A \rtimes_r G} j_G(y).$$

This follows from the fact that the descent homomorphism j_G is compatible with Kasparov products and the fact that

$$y \otimes_B \sigma_B(\gamma) = y \otimes_{\mathbb{C}} \gamma = \gamma \otimes_{\mathbb{C}} y = \sigma_A(\gamma) \otimes_A y,$$

which follows from [24, Theorem 2.14].

It follows now that multiplication with $1 - \gamma$ also commutes with all maps in the above commutative diagram. Since $1 - \gamma$ is an idempotent, it is now easy to see that the full six-term exact sequence restricts to a six-term exact sequence on the $1 - \gamma$ -parts of the respective K-theory groups of the crossed products. \square

Remark 3.3. It is now a direct consequence of the above proposition that if G is a K-exact group possessing a γ -element, and if $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$ is a short exact sequence of G -algebras, then G satisfying BC for two of the algebras in this sequence implies that G satisfies BC for all three algebras in the sequence. The same result holds without the assumption on the γ -element (see [10, Proposition 4.2] – which was actually deduced as an easy consequence of a result of Kasparov and Skandalis in [25]).

We also need the following easy lemma.

Lemma 3.4. *Assume that X is a locally compact space and that A is the algebra of C_0 -sections of the continuous field $\{A_x : x \in X\}$ of C^* -algebras. Assume further that $z \in K_i(A)$, $i = 0, 1$, such that $q_{x,*}(z) = 0$ for some evaluation map $q_x : A \rightarrow A_x$. Then there exists a compact neighborhood C of x such that $q_{C,*}(z) = 0$ in $K_0(A|_C)$, where $A|_C$ denotes the restriction of A to C and $q_C : A \rightarrow A|_C$ denotes the quotient map.*

Proof. We may assume without loss of generality that X is compact. Using suspension, it is enough to give a proof for the case $i = 0$. In what follows, if B is any C^* -algebra, we denote by B^1 the algebra obtained from B by adjoining a unit (even if B is already unital). Then $\{A_x^1 : x \in X\}$ is a continuous field of C^* -algebras in a canonical way. The algebra \tilde{A} of sections can be written as the set of pairs $\{(a, f) : a \in A, f \in C_0(X)\}$ with multiplication given pointwise by the multiplication rule of the fibres A_x^1 . Moreover, we have an obvious unital embedding $A^1 \rightarrow \tilde{A}$.

Assume now that $z \in K_0(A)$ and $x \in X$ are as in the lemma. We represent z as a formal difference $[p - p']$ for some projections $p, p' \in M_l(A^1)$. Since $q_{x,*}(z) = 0$ we may assume (after increasing dimension if necessary) that there exists a unitary $u_x \in M_l(A_x^1)$ such that $u_x p u_x^* = p'_x$. After passing to $\begin{pmatrix} u & 0 \\ 0 & u^* \end{pmatrix}$ if necessary, we may further assume that u_x lies in the connected component of the identity of $U(M_l(A_x^1))$. Thus, there exists a unitary $u \in M_l(A^1)$ such that $q_x(u) = u_x$. Since u is a continuous section in \tilde{A} , it follows that there exists a compact neighborhood C of x such that $\|u_y p_y u_y^* - p'_y\| < 1$ for all $y \in C$, which implies that $[p_C] = [u_C p_C u_C^*] = [p'_C] \in K_0(A|_C)$, where p_C, u_C , and p'_C denote the restrictions of p, u, p' to C , respectively. But this shows that $q_{C,*}(z) = [p_C - p'_C] = 0$ in $K_0(A|_C)$. \square

Proof of Proposition 3.1. Since G is exact, it follows from [26, Theorem] that the crossed products $\{A_x \rtimes_r G : x \in X\}$ form a continuous bundle such that $A \rtimes_r G$ is the algebra of continuous sections of this bundle. We start with proving the result in the special case where $X = [0, 1] \subseteq \mathbb{R}$.

Recall from the above discussions that G satisfies BC for a given G -algebra B if and only if $(1 - \gamma) \cdot K_*(B \rtimes_r G) = \{0\}$. In particular, it follows from our assumptions that $(1 - \gamma) \cdot K_*(A_x \rtimes_r G) = \{0\}$ for all $x \in I$. Assume now that $z \in (1 - \gamma) \cdot K_i(A \rtimes_r G)$, $i = 0, 1$, and let $q_x : A \rtimes_r G \rightarrow A_x \rtimes_r G$ denote the evaluation maps for each $x \in X$. Then $q_{x,*}(z) \in (1 - \gamma) \cdot K_i(A_x \rtimes_r G) = \{0\}$ for all $x \in I$. Thus, using Lemma 3.4, we see that there exists a

partition $0 = x_0 < x_1 < \dots < x_l = 1$ such that $q_{[x_{j-1}, x_j], *}(z) = 0$ in $K_i(A|_{[x_{j-1}, x_j]} \rtimes_r G)$. Now let $O = [0, 1] \setminus \{x_0, \dots, x_l\}$ and let $A|_O = C_0(O) \cdot A \cong \bigoplus_{j=1}^l A|_{(x_{j-1}, x_j)}$. It follows from the exact sequence

$$(1 - \gamma) \cdot K_i(A_O \rtimes_r G) \rightarrow (1 - \gamma) \cdot K_i(A \rtimes_r G) \rightarrow \bigoplus_{j=0}^l (1 - \gamma) \cdot K_i(A_{x_j} \rtimes_r G) = \{0\}$$

that there exists a $z' \in (1 - \gamma) \cdot K_i(A_O \rtimes_r G)$ such that z is the image of z' under the inclusion. Since

$$(1 - \gamma) \cdot K_i(A_O \rtimes_r G) = \bigoplus_{j=1}^l (1 - \gamma) \cdot K_i(A|_{(x_{j-1}, x_j)} \rtimes_r G),$$

we may write $z' = \sum_{j=1}^l z'_j$ with $z'_j \in (1 - \gamma) \cdot K_i(A|_{(x_{j-1}, x_j)} \rtimes_r G)$ for each $1 \leq j \leq l$. Thus it is enough to show that $z'_j = 0$ for each $1 \leq j \leq l$. In what follows, we write $A_j = A|_{(x_{j-1}, x_j)}$ and $\bar{A}_j = A|_{[x_{j-1}, x_j]}$. Since $(1 - \gamma) \cdot K_i(A_{x_k} \rtimes_r G) = \{0\}$ for all $0 \leq k \leq l$ we obtain a six-term exact sequence

$$\begin{array}{ccccc} (1 - \gamma) \cdot K_0(A_j \rtimes_r G) & \longrightarrow & (1 - \gamma) \cdot K_0(\bar{A}_j \rtimes_r G) & \longrightarrow & 0 \\ \uparrow & & & & \downarrow \\ 0 & \longleftarrow & (1 - \gamma) \cdot K_1(\bar{A}_j \rtimes_r G) & \longleftarrow & (1 - \gamma) \cdot K_1(A_j \rtimes_r G). \end{array}$$

Since the image of z'_j in $K_i(\bar{A}_j \rtimes_r G)$ coincides with the image of z in $K_i(\bar{A}_j \rtimes_r G)$, we see that z'_j maps to 0 under the isomorphism $(1 - \gamma) \cdot K_0(A_j \rtimes_r G) \rightarrow (1 - \gamma) \cdot K_0(\bar{A}_j \rtimes_r G)$, so $z'_j = 0$.

We now show by induction on n that the result is true for $[0, 1]^n \subseteq \mathbb{R}^n$. For this assume that $\{A_x : x \in [0, 1]^n\}$ is a continuous field over the cube and A is the algebra of continuous sections of this field. We write $[0, 1]^n = \cup_{y \in [0, 1]} \{y\} \times [0, 1]^{n-1}$ and put $A_y = A|_{\{y\} \times [0, 1]^{n-1}}$. Then $\{A_y : y \in [0, 1]\}$ is a continuous field over $[0, 1]$ and A is also the section algebra of this bundle. If α is a $C([0, 1]^n)$ -linear action on A , it is also $C([0, 1])$ -linear with respect to the bundle structure of A over $[0, 1]$ coming from the above decomposition of the cube. Moreover, the actions on the fibres A_y are clearly $C([0, 1]^{n-1})$ -linear, so by the induction assumption we know that G satisfies BC with coefficients in A_y for all $y \in [0, 1]$. We now apply the above result to the bundle $\{A_y : y \in [0, 1]\}$ to conclude that G satisfies BC with coefficients in A .

In a next step we show that the result holds for the open cubes $(0, 1)^n \subseteq \mathbb{R}^n$. By similar arguments as given above it suffices to show that the result holds for open intervals. So assume that $\{A_x : x \in (0, 1)\}$ is a continuous field with section algebra A . Let $x_1 < x_2 \in (0, 1)$. Then it follows from the first part of the proof that G satisfies BC with coefficients in $A|_{[x_1, x_2]}$. Since, by assumption, G also satisfies BC for the fibres, a six-term sequence argument shows that it also satisfies BC with coefficients in (x_1, x_2) . Writing $A = \lim_{n \rightarrow \infty} A|_{(\frac{1}{n}, 1 - \frac{1}{n})}$ and using continuity of the BC conjecture, it follows that G satisfies BC for A .

Since the result of the proposition is clearly invariant under replacing the space X by a homeomorphic space Y , we now see that the result holds for all open or closed simplices. We now proof the general result for simplicial complexes via induction on the dimension of the complex. By continuity of the conjecture, the result is clear for zero-dimensional complexes. If X has dimension n , let W_n denote the interiors of all n -dimensional simplices in X . Then

W_n is homeomorphic to a disjoint union of open n -dimensional cubes, so the result holds for W_n . Since $X \setminus W_n$ is a simplicial complex of dimension $n - 1$, the result is true for $X \setminus W_n$ by the induction assumption. The result then follows from another easy application of the six-term sequence (see Remark 3.3). \square

Remark 3.5. Let G be a second countable locally compact group and let X be a second countable locally compact G -space. Following Glimm we say that the quotient space X/G is *countably separated*, if all orbits $G(x)$ are locally closed, i.e. $G(x)$ is open in its closure $\overline{G(x)}$. Glimm showed in [19, Theorem] that X/G being countably separated is equivalent to each of the following conditions:

- The canonical map $G/G_x \rightarrow G(x), gG_x \mapsto g \cdot x$ is a homeomorphism for each $x \in X$.
- There exists a sequence of G -invariant open subsets $\{U_\nu\}_\nu$ of X , where ν runs through the ordinal numbers such that
 - (i) $U_\nu \subseteq U_{\nu+1}$ for each ν and $(U_{\nu+1} \setminus U_\nu)/G$ is Hausdorff.
 - (ii) If ν is a limit ordinal, then $U_\nu = \cup_{\mu < \nu} U_\mu$.
 - (iii) There exists an ordinal number ν_0 such that $X = U_{\nu_0}$.

Unfortunately, in order to apply our bundle results to such actions, we need to know that the spaces $(U_{\nu+1} \setminus U_\nu)/G$ have a simplicial structure as required in Proposition 3.1. However, we can summarize the above results to prove the following theorem which turns out to be sufficient for our purposes:

Theorem 3.6. *Suppose that G is a second countable exact group which has a γ -element. Let X be a second countable locally compact G -space, let $\{A_x : x \in X\}$ be a continuous bundle of C^* -algebras over X with section algebra A and let $\alpha : G \rightarrow \text{Aut}(A)$ be an action of G on A which is compatible with the given action of G on X . Assume further that the following assumptions are satisfied:*

- (i) X/G is countably separated.
- (ii) *There exists an increasing sequence of open G -invariant subsets $\{U_\nu\}_\nu$ of X such that $U_\nu = \cup_{\mu < \nu} U_\mu$ if ν is a limit ordinal and $X = U_{\nu_0}$ for some ν_0 .*
- (iii) *For each ν there exists a finite dimensional simplicial complex Y_ν and a continuous and open surjection $q_\nu : U_{\nu+1} \setminus U_\nu \rightarrow Y_\nu$ such that for all $y \in Y_\nu$, $q_\nu^{-1}(\{y\})$ is a finite union of orbits in X .*
- (iv) *Each stabilizer G_x , $x \in X$, satisfies BC for A_x .*

Then G satisfies BC for A .

Proof. For each ordinal ν let $A_\nu := C_0(U_\nu)A$ denote the ideal corresponding to U_ν . We show by transfinite induction that G satisfies BC with coefficients in A_ν for each ν . Since $A = A_{\nu_0}$ for some ν_0 , the result will follow.

We start by showing that G satisfies BC with coefficients in A_1 . Since the map $q_1 : U_1 \rightarrow Y_1$ is open, we can regard A_1 as a section algebra of a continuous bundle over Y_1 with fibres isomorphic to $A|_{q_1^{-1}(y)}$. By Proposition 3.1 it is therefore sufficient to prove that G satisfies BC for $A|_{q_1^{-1}(y)}$ for all $y \in Y_1$. Fix $y \in Y_1$ and put $Z := q_1^{-1}(y)$. Since Z is a finite union of G -orbits, we find a finite sequence

$$Z = Z_0 \supseteq Z_1 \supseteq \cdots \supseteq Z_l = \emptyset$$

of open invariant subsets of Z such that $(Z_{i-1} \setminus Z_i)/G$ is a discrete finite set. To see this let C_1 be the union of all closed G -orbits in Z (such orbits must exist by the finiteness of $Z/G!$). Then C_1 is closed in Z and C_1/G is discrete. Put $Z_1 = Z \setminus C_1$ and then define the Z_i 's, $i > 1$, inductively by the same procedure. Using six-term sequences (e.g., see Remark 3.3), G satisfies BC for $A|_Z$ if G satisfies BC for all $A|_{Z_{i-1} \setminus Z_i}$, which in turn follows if G satisfies BC for $A|_{G(x)}$ for any G -orbit $G(x) \subseteq X$. Since $G(x)$ is homeomorphic to G/G_x via the canonical map (see Remark 3.5 above), it follows from Proposition 2.8, that $A|_{G(x)}$ is G -equivariantly isomorphic to $\text{Ind}_{G_x}^G A_x$. Since, by assumption, G_x satisfies BC with coefficients in A_x , it follows then from Theorem 2.7 that G satisfies BC with coefficients in $A|_{G(x)}$. This completes the proof for A_1 .

Assume now that ν is an ordinal number and that we have already shown that G satisfies BC for A_μ for all $\mu < \nu$. If $\nu = \mu + 1$ for some ordinal μ , it follows from the same reasoning as for the case $\nu = 1$ that G satisfies BC for $A|_{U_\nu \setminus U_\mu} \cong A_\nu/A_\mu$. Since G satisfies BC for A_μ by the induction assumption, it follows from Remark 3.3 that G satisfies BC for A_ν .

Assume now that ν is a limit ordinal and G satisfies BC for each $\mu < \nu$. Then $U_\nu = \cup_{\mu < \nu} U_\mu$ which implies that $A_\nu = \lim_{\mu < \nu} A_\mu$ is the inductive limit of the A_μ . Thus it follows from Proposition 2.5 that G satisfies BC for A_ν . \square

4. THE SEMI-SIMPLE CASE

In this section we want to show that Proposition 2.10 is true if G is semi-simple. For this we first have to obtain a slight extension of Lafforgue's results on the Baum-Connes conjecture for semi-simple groups with finite center.

Let us first recall the basic idea of Lafforgue's proof of the Baum-Connes conjecture for such groups. If G is a locally compact group we let $C_c(G)$ denote the convolution algebra of G consisting of continuous functions with compact supports. A norm $\| \cdot \|$ on $C_c(G)$ is called *good* if convolution is continuous with respect to this norm and if $\|f\|$ only depends on the absolute value of f for all $f \in C_c(G)$ (i.e., $\|f\| = \|\ |f|\ |$ for all $f \in C_c(G)$). A *good completion* $\mathcal{A}(G)$ of $C_c(G)$ is a completion with respect to a good norm on $C_c(G)$. Note that $L^1(G)$ is always a good completion of $C_c(G)$, but $C^*(G)$ and $C_r^*(G)$ are in general not good completions of $C_c(G)$.

If $\mathcal{A}(G)$ is a good completion of $C_c(G)$, then Lafforgue constructed an assembly map

$$\mu_{\mathcal{A}(G)} : K_*^{\text{top}}(G, \mathbb{C}) \rightarrow K_*(\mathcal{A}(G)).$$

Moreover, if the identity on $C_c(G)$ extends to a continuous embedding $\iota : \mathcal{A}(G) \rightarrow C_r^*(G)$, he also shows that the assembly map $\mu : K_*^{\text{top}}(G, \mathbb{C}) \rightarrow K_*(C_r^*(G))$ factors through $K_*(\mathcal{A}(G))$, i.e.,

$$\mu = \iota_* \circ \mu_{\mathcal{A}(G)}$$

(see [29, Proposition 1.7.6]). Thus, if we know that $\mu_{\mathcal{A}(G)}$ is an isomorphism for all good completions of $C_c(G)$, and if we further know that there exists a good completion $\mathcal{A}(G) \subseteq C_r^*(G)$ such that the inclusion $\iota_* : K_*(\mathcal{A}(G)) \rightarrow K_*(C_r^*(G))$ is an isomorphism, it follows that $\mu : K_*^{\text{top}}(G, \mathbb{C}) \rightarrow K_*(C_r^*(G))$ is an isomorphism. Note that $\iota_* : K_*(\mathcal{A}(G)) \rightarrow K_*(C_r^*(G))$ is an isomorphism whenever $\mathcal{A}(G)$ is closed under holomorphic functional calculus in $C_r^*(G)$. Now Lafforgue was able to prove the following deep results:

Theorem 4.1 (cf [29, Théorème 17.13 and Chapitre 3]). *Assume that G is a second countable locally compact group such that G acts isometrically and properly on a Riemannian manifold with nonpositive sectional curvature which is bounded below. Then $\mu_{\mathcal{A}(G)} : K_*^{\text{top}}(G, \mathbb{C}) \rightarrow K_*(\mathcal{A}(G))$ is an isomorphism for every good completion $\mathcal{A}(G)$ of $C_c(G)$.*

In fact, Lafforgue was even able to show that the above result holds with arbitrary C^* -algebra coefficients, but we do not need this more general result here. Note that if G is semi-simple with finite center, then the Riemannian manifold of the theorem can be chosen to be the symmetric space G/K , where K is the maximal compact subgroup of G .

In the second step for the proof of BC for semi-simple groups, Lafforgue constructed a *Schwartz-algebra* $\mathcal{S}(G) \subseteq C_r^*(G)$ which is a good completion of $C_c(G)$ which is closed under holomorphic functional calculus in $C_r^*(G)$. In fact, this construction followed a more general principle, which we are now going to describe in more detail.

Assume that G is a unimodular group and K is a compact subgroup of G . Then, following Lafforgue (see [29, Chapitre 4]), we say that the pair (G, K) satisfies property (HC) if the following conditions are satisfied

- (HC1) There exists a continuous function $d : G \rightarrow [0, \infty)$ such that $d(e) = 0$, $d(kgk') = d(g)$ for all $k, k' \in K, g \in G$ and $d(gg') \leq d(g) + d(g')$ for all $g, g' \in G$.
- (HC2) There exists a continuous function $\phi : G \rightarrow (0, 1]$ such that $\phi(e) = 1$, $\phi(g^{-1}) = \phi(g)$, $\phi(kgk') = \phi(g)$ for all $g \in G$ and $k, k' \in K$, and

$$\int_K \phi(gkg') dk = \phi(g)\phi(g') \quad \text{for all } g, g' \in G,$$

with respect to the normalized Haar measure on K .

- (HC3) There exists a $t_0 \in \mathbb{R}$ such that $(t \mapsto \phi(g)(1 + d(g))^{-t}) \in L^2(G)$ for all $t > t_0$.

If G is a connected semi-simple group with finite center, and if K is the maximal compact subgroup of G , then Lafforgue was able to show that (G, K) satisfies (HC). The following theorem then completes the proof of BC for connected semi-simple groups with finite center.

Theorem 4.2 (cf [29, Proposition 4.2.1]). *Assume that G is a unimodular group and K is a compact subgroup of G such that the pair (G, K) satisfies (HC). Then there exists a good completion $\mathcal{S}(G) \subseteq C_r^*(G)$ of $C_c(G)$ such that $\mathcal{S}(G)$ is closed under holomorphic functional calculus in $C_r^*(G)$.*

In order to extend Lafforgue's methods to extensions of semi-simple groups, we first show that property (HC) is closed under compact extensions.

Lemma 4.3. *Assume that (G, K) satisfies property (HC). Assume further that*

$$1 \rightarrow C \rightarrow \tilde{G} \xrightarrow{q} G \rightarrow 1$$

is a group extension with C compact. Let $\tilde{K} = q^{-1}(K) \subseteq \tilde{G}$. Then (\tilde{G}, \tilde{K}) satisfies (HC).

Proof. Since compact extensions of unimodular groups are unimodular, \tilde{G} is unimodular. Let (d, ϕ) be a pair of functions which satisfy HC1, HC2, and HC3 with respect to (G, K) . Define $\tilde{d}(g) = d(q(g))$ and $\tilde{\phi}(g) = \phi(q(g))$. Then a straightforward computation shows that $(\tilde{d}, \tilde{\phi})$ satisfies HC1, HC2, and HC3 with respect to (\tilde{G}, \tilde{K}) . \square

The second result is slightly more technical.

Lemma 4.4. *Assume that G is a unimodular Lie group with finitely many components and let K be a maximal compact subgroup of G . Let G_0 denote the connected component of G and let K_0 be a maximal compact subgroup of G_0 such that (G_0, K_0) satisfies (HC). Then (G, K) satisfies (HC), too.*

Proof. First note that we may assume that $K_0 = K \cap G_0$. To see this observe first that, since K_0 is a compact subgroup of G , we may assume without loss of generality that $K_0 \subseteq K \cap G_0$. But the maximality of K_0 then implies equality. It follows from this that K_0 is a normal subgroup of K and since G/K is connected, it follows that the inclusion $K \rightarrow G$ induces a group isomorphism $K/K_0 \cong G/G_0$.

Let (d_0, ϕ_0) be a pair of functions satisfying conditions HC1–HC3 for (G_0, K_0) . It follows from the above remarks that we can write every element of G as a product kg with $k \in K$, $g \in G_0$. We then define

$$d(kg) = \int_K d_0(lgl^{-1}) dl \quad \text{and} \quad \phi(kg) = \int_K \phi_0(lgl^{-1}) dl.$$

To see that d and ϕ are well defined assume that we have two factorizations $kg = k'g'$ with $k, k' \in K, g, g' \in G_0$. Then $g = k^{-1}k'g'$ with $h := k^{-1}k' \in K_0$. Since K normalizes K_0 , it follows from the left and right K_0 -invariance of d_0 that

$$\begin{aligned} d(kg) &= \int_K d_0(lgl^{-1}) dl = \int_K d_0(lhg't^{-1}) dl = \int_K d_0((lhl^{-1})(lg't^{-1})) dl \\ &= \int_K d_0(lg't^{-1}) dl = d(k'g'). \end{aligned}$$

A similar computation shows that ϕ is well defined.

We are now going to check properties HC1–HC3 for (d, ϕ) . It follows directly from the definition of d and ϕ that they are left invariant under the action of K . To see right invariance, we compute for $h \in K$:

$$d(kgh) = d(khh^{-1}gh) = d(h^{-1}gh) = \int_K d_0(lh^{-1}ghl^{-1}) dl \stackrel{h \rightarrow lh}{=} \int_K d_0(lhl^{-1}) dl = d(kg).$$

So d is also right invariant and a similar computation show that the same is true for ϕ . Since Haar measure on K is normalized, it follows that $d(e) = 0$ and $\phi(e) = 1$. Moreover, if $kg, hg' \in G$ with $k, h \in K, g, g' \in G_0$ we get

$$\begin{aligned} d(kghg') &= d(khh^{-1}ghg') = \int_K d_0(lh^{-1}ghg'l^{-1}) dl = \int_K d_0(lh^{-1}ghl^{-1}lg't^{-1}) dl \\ &\leq \int_K d_0(lh^{-1}ghl^{-1}) dl + \int_K d_0(lg't^{-1}) dl = d(kg) + d(hg'). \end{aligned}$$

In order to prove the multiplication rule for ϕ we use Weil's formula

$$\int_K \varphi(l) dl = \int_{K/K_0} \left(\int_{K_0} \varphi(lm) dm \right) dl = \int_K \left(\int_{K_0} \varphi(lm) dm \right) dl,$$

with respect to normalized Haar measures on K, K_0 , and K/K_0 to compute

$$\begin{aligned}
\int_K \phi(kglhg') dl &= \int_K \phi(glhg'h^{-1}) dl = \int_K \int_{K_0} \phi(glmhg'h^{-1}) dm dl \\
&\stackrel{(*)}{=} \int_K \int_{K_0} \phi(l^{-1}glmhg'h^{-1}) dm dl \\
&= \int_K \int_{K_0} \int_K \phi_0(nl^{-1}glmhg'h^{-1}n^{-1}) dn dm dl \\
&\stackrel{(**)}{=} \int_K \int_K \int_{K_0} \phi_0(nl^{-1}gln^{-1}mnhg'h^{-1}n^{-1}) dm dl dn \\
&\stackrel{(***)}{=} \int_K \int_K \phi_0(nl^{-1}gln^{-1}) \phi_0(nhg'h^{-1}n^{-1}) dl dn \\
&\stackrel{l \mapsto l^{-1}n}{=} \left(\int_K \phi_0(lgl^{-1}) dl \right) \left(\int_K \phi_0(ng'n^{-1}) dn \right) \\
&= \phi(kg)\phi(hg').
\end{aligned}$$

Here the equation (*) follows from the K -invariance of ϕ , (**) follows from Fubini together with the transformation $m \mapsto n^{-1}mn$, and (***) follows from property HC2 for ϕ_0 . This completes the proof of HC1 and HC2.

For the proof of HC3 we write $F_s(g) := \phi(g)(1+d(g))^{-s}$, $s > 0$. Since $G = KG_0$, it follows from the K -invariance of ϕ and d that the integrals of F_s^2 over the G_0 -cosets coincide. Thus, since G/G_0 is finite, it is enough to show that $F_s|_{G_0} \in L^2(G_0)$ for some $s > 0$. For this we first choose a set of representatives $t_1, \dots, t_n \in K$ for K/K_0 with $t_1 = e$. Then, for $g \in G_0$, we obtain the inequality

$$1 + d(g) = 1 + \frac{1}{n} \sum_{i=1}^n d_0(t_i g t_i^{-1}) \geq 1 + \frac{1}{n} d_0(g),$$

from which it follows that $(1 + d(g))^n \geq 1 + d_0(g)$ for all $g \in G_0$. Thus, if $t \in \mathbb{R}$ such that $(g \mapsto \phi_0(g)(1 + d_0(g))^{-t}) \in L^2(G_0)$, then we also have $(g \mapsto \phi_0(g)(1 + d(g))^{-nt}) \in L^2(G_0)$. So let $s = nt$ with t as above. Then we get

$$\begin{aligned}
F_s(g) &= \phi(g)(1 + d(g))^{-s} = \frac{1}{n} \sum_{i=1}^n \phi_0(t_i g t_i^{-1})(1 + d(g))^{-s} \\
&= \frac{1}{n} \sum_{i=1}^n \phi_0(t_i g t_i^{-1})(1 + d(t_i g t_i^{-1}))^{-s}.
\end{aligned}$$

Since G is unimodular, it follows that each summand $(g \mapsto \phi_0(t_i g t_i^{-1})(1 + d(t_i g t_i^{-1}))^{-s}) \in L^2(G_0)$, and hence $F_s|_{G_0} \in L^2(G_0)$. \square

We are now ready to combine the above results to get

Proposition 4.5. *Let G be a locally compact group with finitely many components. Assume further that G has a compact normal subgroup $C \subseteq G_0$ such that G_0/C is semi-simple with finite center. Then G satisfies BC with trivial coefficients.*

Proof. Let K_0 denote the maximal compact subgroup of G_0 . Then $G_0/K_0 \cong (G_0/C)/(K_0/C)$ is a symmetric space and therefore has nonpositive sectional curvature. Moreover, if K is

a maximal compact subgroup of G such that $K \cap G_0 = K_0$, we see that $G/K \cong G_0/K_0$ as a Riemannian manifold. Since G acts isometrically and properly on G/K , G satisfies the assumptions of Theorem 4.1. By Lafforgue's results we also know that $(G_0/C, K_0/C)$ satisfies (HC). Lemmas 4.3 and 4.4 then imply that (G, K) also satisfies property (HC). Thus, it follows from the combination of Theorem 4.1 with Theorem 4.2 that G satisfies BC with coefficients in \mathbb{C} . \square

Using the results on continuous fields of actions as presented in the previous section, we are now able to prove

Proposition 4.6. *Assume that G is a Lie group with finitely many components such that G_0 is reductive, i.e., the Lie algebra \mathfrak{g} of G is a direct sum of two ideals $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{z}$ with \mathfrak{s} semi-simple and \mathfrak{z} abelian. Then G satisfies BC for \mathbb{C} .*

Proof. Let $Z = Z(G_0)$ denote the center of G_0 . Using Theorem 2.1 it is enough to show that G/Z satisfies BC with coefficients in $C^*(Z) \otimes \mathcal{K} \cong C_0(\widehat{Z}, \mathcal{K})$, where the action of G/Z on the dual space \widehat{Z} of Z is given via conjugation. Since Z is central in G_0 , it follows that this action factors through an action of the finite group G/G_0 . Moreover, since \widehat{Z} is a manifold (since Z is a compactly generated abelian group), it follows that the quotients of the orbit-types in \widehat{Z} are manifolds. From this we easily obtain a finite decomposition sequence

$$\emptyset = U_0 \subseteq U_1 \subseteq \cdots \subseteq U_l = \widehat{Z}$$

of open G -invariant subsets of \widehat{Z} such that the quotients of the differences $U_j \setminus U_{j-1}$ are homeomorphic to geometric realizations of finite dimensional simplicial complexes. Moreover, since all stabilizers for the action of G/Z on \widehat{Z} contain G_0/Z , which is semi-simple with trivial center, it follows from a combination of Proposition 4.5 with Proposition 2.6 that all stabilizers satisfy BC for \mathcal{K} . The result then follows from Theorem 3.6. \square

Since any central extension of a semi-simple group is reductive, we now get the desired result for general semi-simple groups.

Corollary 4.7. *Let G be a semi-simple Lie group with finitely many components and let*

$$1 \rightarrow \mathbb{T} \rightarrow \bar{G} \rightarrow G \rightarrow 1$$

be a central extension of G by \mathbb{T} . Then \bar{G} satisfies BC for \mathbb{C} . As a consequence (using Proposition 2.6), G satisfies BC for \mathcal{K} with respect to arbitrary actions of G on \mathcal{K} .

5. THE GENERAL CASE

We now want to give a proof of Proposition 2.10. As indicated in the first section, we are going to use an induction argument on the dimension $n = \dim(G)$. Since any one-dimensional Lie group with finitely many components is amenable, and since amenable groups satisfy BC for arbitrary coefficients, the case $n = 1$ is clear. Assume now that G is an arbitrary Lie group with finitely many components. Let G_0 denote the connected component of G . Let N denote the nilradical of G and let \mathfrak{n} and \mathfrak{g} denote the Lie algebras of N and G , respectively. If $\mathfrak{n} = \{0\}$, then G is semi-simple and the result follows from the previous section. So we may assume that $\mathfrak{n} \neq \{0\}$.

It is shown in [34, Lemma 4 on p. 24] that the subgroup H of G_0 corresponding to the subalgebra $\mathfrak{h} = [\mathfrak{g}, \mathfrak{g}] + \mathfrak{n}$ of \mathfrak{g} is closed in G_0 . Further, if \mathfrak{s} is a Levi section in \mathfrak{g} , i.e., \mathfrak{s} is a maximal semi-simple subalgebra of \mathfrak{g} , then $\mathfrak{h} = \mathfrak{s} + \mathfrak{n}$ (e.g., see the discussion in the proof of [34, Sublemma on p. 24]). In particular, H/N is semi-simple. Clearly, H is a normal subgroup of G and G/H is a finite extension of a connected abelian Lie group. Let M denote the inverse image of the maximal compact subgroup of G/H in G . It follows then from Theorem 2.1 that G satisfies BC for \mathcal{K} if and only if M satisfies BC for \mathcal{K} . Note that the connected component of M/H is a compact connected abelian Lie group, hence a torus group.

Thus, replacing G by M , we may from now on assume that G has the following structure: There exist closed normal subgroups

$$(5.1) \quad N \subseteq H \subseteq G_0 \subseteq G$$

such that N is a non-trivial connected nilpotent Lie group, H/N is semi-simple, G_0/H is a torus group and G/G_0 is finite. Moreover, by induction we may assume that every almost connected Lie group with smaller dimension satisfies BC for \mathcal{K} with respect to arbitrary actions on \mathcal{K} , or, equivalently (by Proposition 2.6), every central extension by \mathbb{T} satisfies BC for \mathbb{C} . It is now useful to recall the following result of Chevalley (see [12, Proposition 5, p.324]):

Proposition 5.1. *Let $\mathfrak{g} \subseteq \mathfrak{gl}(V)$ be a Lie-algebra of endomorphisms of the finite dimensional real vector space V . Then \mathfrak{g} is algebraic (i.e. it corresponds to a real algebraic subgroup $G \subseteq \mathrm{GL}(V)$) if and only if there exist subalgebras $\mathfrak{s}, \mathfrak{a}, \mathfrak{n}$ of \mathfrak{g} with $\mathfrak{g} = \mathfrak{s} + \mathfrak{a} + \mathfrak{n}$, \mathfrak{s} is semi-simple, \mathfrak{n} is the largest ideal of \mathfrak{g} consisting of nilpotent endomorphisms, and \mathfrak{a} is an algebraic abelian subalgebra of $\mathfrak{gl}(V)$ consisting of semi-simple endomorphisms such that $[\mathfrak{s}, \mathfrak{a}] \subseteq \mathfrak{a}$.*

Using Ado's theorem (see [12, Théorème 5 on p. 333]) and Proposition 5.1, it follows that the group H considered above is locally algebraic, i.e., the Lie algebra \mathfrak{h} has a faithful representation as an algebraic Lie subalgebra into some $\mathfrak{gl}(V)$.

Using this structure, the main idea is to apply the Mackey machine to a suitable abelian subgroup S of N which is normal in G . The fact that G is very close to an algebraic group implies that the action of G on the dual \widehat{S} of S has very good topological properties, which is precisely what we need to make everything work. As a first hint that this approach is feasible we prove:

Lemma 5.2. *Assume that G is a Lie group with finitely many components. Let $H \subseteq G_0$ be a connected closed normal subgroup of G such that H is locally algebraic, and such that G/H is compact. Let N denote the nilpotent radical of H , and let $S \subseteq N$ be a connected abelian normal subgroup of G . Let \widehat{S} denote the character group of S and let G act on \widehat{S} via conjugation. Then the following assertions are true:*

- (i) *The orbit space \widehat{S}/G is countably separated, i.e., all G -orbits in \widehat{S} are locally closed.*
- (ii) *If G_χ is the stabilizer of some $\chi \in \widehat{S}$ for the action of G on \widehat{S} , then $G_\chi/(G_\chi)_0$ is amenable.*

Proof. We first show that it is sufficient to prove the result for the case $G = H$. Indeed, if we already know that \widehat{S}/H is countably separated, then we observe that \widehat{S}/H is a topological G/H -space such that $\widehat{S}/G \cong (\widehat{S}/H)/(G/H)$. But it is an easy exercise to prove that the quotient space of a countably separated space by a compact group action is countably separated.

Assume now that G_χ is the stabilizer of some $\chi \in \widehat{S}$ in G . Then $H_\chi = G_\chi \cap H$ is the stabilizer in H . It follows that H_χ is a normal subgroup of G_χ such that G_χ/H_χ is compact. If $H_\chi/(H_\chi)_0$ is amenable, it also follows that $G_\chi/(H_\chi)_0$, and hence also $G_\chi/(G_\chi)_0$ are amenable.

So, for the rest of the proof we assume that $G = H$. In the next step we reduce to the case where H is simply connected. For this let \tilde{H} denote the universal covering group of H . Then \tilde{H} has the same Lie algebra as H , and therefore it is locally algebraic. Let $q : \tilde{H} \rightarrow H$ denote the quotient map and let $C = \ker q$. Then C is a discrete central subgroup of \tilde{H} . Let $\mathfrak{r} \subseteq \mathfrak{h}$ denote the Lie algebra of S and let \tilde{S} denote the connected closed normal subgroup of \tilde{H} corresponding to \mathfrak{r} . Then \tilde{S} is a vector subgroup of the nilpotent radical \tilde{N} of \tilde{H} and the quotient map $\tilde{H} \rightarrow H$ maps \tilde{S} surjectively onto S , i.e., we have $S \cong \tilde{S}/(\tilde{S} \cap C)$. In particular, it follows that we may view \widehat{S} as a closed \tilde{H} -invariant subspace of $\widehat{\tilde{S}}$, and we have $\widehat{S}/H = \widehat{S}/\tilde{H}$ (since the central subgroup C acts trivially on \widehat{S}). Thus, if $\widehat{\tilde{S}}/\tilde{H}$ is countably separated, the same is true for \widehat{S}/H .

We now consider the stabilizers. It follows from the above considerations that if H_χ is the stabilizer of some $\chi \in \widehat{S}$, then $q^{-1}(H_\chi) \subseteq \tilde{H}$ is the stabilizer of χ in \tilde{H} . Thus it follows that $H_\chi = \tilde{H}_\chi/(C \cap \tilde{H}_\chi)$. Since the connected component of \tilde{H}_χ is mapped onto the connected component of H_χ under the quotient map, it follows that $H_\chi/(H_\chi)_0$ is a quotient of $\tilde{H}_\chi/(\tilde{H}_\chi)_0$. Thus, if the latter is amenable, the same is true for $H_\chi/(H_\chi)_0$.

Thus, in what follows we may assume without loss of generality that H is simply connected. We are then in precisely the same situation as in the proof of Case (A) of the proof of the Theorem on page 2 of [34], and from now on we can follow the line of arguments as given on pages 2 and 3 of [34] to see that \widehat{S}/H is countably separated. Moreover, the arguments presented in steps c) and d) on page 3 of Pukánszky's book imply that for each stabilizer H_χ the quotient $H_\chi/(H_\chi)_0$ is a finite extension of an abelian group, and hence is amenable. \square

Remark 5.3. We should point out that the result on the stabilizers in Lemma 5.2 is most satisfying: Indeed if we know that every almost connected Lie group with dimension $\dim(G) < n$ satisfies BC for \mathcal{K} , say, then, by an easy application of Theorem 2.1 the same is true for all Lie groups H with $\dim(H) < n$ and H/H_0 amenable!

Unfortunately, the result on the orbit space \widehat{S}/G is not sufficient for a direct application of Theorem 3.6. So we have to do some extra work to obtain more information on the structure of \widehat{S}/G . To do this we have to do two steps:

- (i) Reduce to cases where the action of G on \widehat{S} factors through an algebraic action of some real algebraic group G' (or a subgroup of finite index in G').
- (ii) Show that that the topological orbit-spaces of algebraic group actions on real affine varieties have nice stratifications as required by Theorem 3.6.

Note that Pukánszky does the first reduction for the cocompact subgroup H of G , which allowed us to draw the conclusions of the previous lemma. However, with a bit more work we

obtain similar conclusion for G . The following result is certainly well-known to the experts, but since we didn't find a direct reference we included the easy proof.

Lemma 5.4. *Assume that G is a Lie group with finitely many components such that G has a connected closed normal subgroup H with H semi-simple and G_0/H a torus group. Let V be a finite dimensional real vector space and let $\rho : G \rightarrow \mathrm{GL}(V)$ be any continuous homomorphism. Then the Zariski closure G' of $\rho(G)$ is a (reductive) real algebraic group which contains $\rho(G)$ as a subgroup of finite index.*

Proof. Let $R : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ denote the differential of ρ and let \mathfrak{h} denote the ideal of \mathfrak{g} corresponding to H . Then $R(\mathfrak{h})$ is semi-simple (or trivial). Since $\rho(H)$ is a semi-simple subgroup of $\mathrm{GL}(V)$ it is closed in $\mathrm{GL}(V)$. This follows from the fact that every semi-simple subalgebra of $\mathfrak{gl}(V)$ is algebraic (by Proposition 5.1), which implies that $\rho(H)$ is the connected component of some algebraic linear subgroup of $\mathrm{GL}(V)$. Since G/H is compact, it follows that $\rho(G)$ is a closed subgroup of $\mathrm{GL}(V)$, too.

To simplify notation we assume from now on that G itself is a closed subgroup of $\mathrm{GL}(V)$ and that ρ is the identity map. Let $\mathfrak{g} = \mathfrak{s} + \mathfrak{z}$ be a Levi decomposition of \mathfrak{g} . Since G_0/H is abelian and H is semi-simple, it follows that $\mathfrak{s} = [\mathfrak{g}, \mathfrak{g}] = \mathfrak{h}$ and $[\mathfrak{h}, \mathfrak{z}] = [\mathfrak{z}, \mathfrak{z}] = \{0\}$. Thus $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{z}$ is reductive.

We now show that \mathfrak{g} is an algebraic subalgebra of $\mathfrak{gl}(V)$. By Proposition 5.1 it suffices to show that \mathfrak{z} is algebraic and consists of semi-simple elements. But this will follow if we can show that $Z = \exp(\mathfrak{z}) \subseteq \mathrm{GL}(V)$ is compact, and hence a torus group. Since $Z \cap H$ is finite (since every linear semi-simple group has finite center), the restriction to Z of the quotient map $q : G \rightarrow G/H$ has finite kernel. Since $q(Z) = G_0/H$, $q(Z)$ is compact by assumption, and hence Z is compact, too.

It follows that the algebraic closure \tilde{G} of G_0 is a reductive algebraic group which contains G_0 as a subgroup of finite index. Since every element of G fixes the Lie algebra \mathfrak{g} via the adjoint action, it also normalizes \tilde{G} . Therefore, $G' = G\tilde{G}$ is a reductive algebraic group which contains G as a subgroup of finite index. \square

We now show that quotients of linear algebraic group actions on affine varieties have nice stratifications in the sense of Theorem 3.6. We are very grateful to Jörg Schürmann and Peter Slodowy for some valuable comments, which helped us to replace a previous version of the following result (which, as was pointed out to us by Jörg Schürmann, contained a gap) by

Proposition 5.5. *Suppose that G is a closed subgroup of finite index of a Zariski closed subgroup G' of $\mathrm{GL}(n, \mathbb{R})$ and that $V \subseteq \mathbb{R}^n$ is a G' -invariant Zariski closed subset of \mathbb{R}^n . Then there exists a stratification*

$$\emptyset = V_0 \subseteq V_1 \subseteq \cdots \subseteq V_l = V$$

of open G -invariant subsets V_i of V such that $(V_i \setminus V_{i-1})/G$ admits a continuous and open finite-to-one map onto a differentiable manifold.

Since every manifold has a triangulation, the above result really gives what we need to apply Theorem 3.6. For the proof we need the following lemma about certain decompositions of continuous semi-algebraic maps.

Lemma 5.6. *Let X, Y be semi-algebraic sets and let $f : X \rightarrow Y$ be a continuous semi-algebraic map (see [6] for the notations). Then there exists a stratification*

$$\emptyset = Z_0 \subseteq Z_1 \cdots \subseteq Z_l = f(X),$$

with each Z_i open in $f(X)$, $Z_i \setminus Z_{i-1}$ is a differentiable manifold and

$$f : f^{-1}(Z_i \setminus Z_{i-1}) \rightarrow Z_i \setminus Z_{i-1}$$

is open (in the euclidean topology) for all $1 \leq i \leq l$.

Proof. Since the image of a semi-algebraic set under a semi-algebraic map is semi-algebraic (see [6, Proposition 2.2.7]), we may assume without loss of generality that $Y = f(X)$. By [6, Corollary 9.3.3] there exists a closed semi-algebraic subset $Y_1 \subseteq Y$ with $\dim(Y_1) < \dim(Y)$, such that $Y \setminus Y_1$ is a finite disjoint union of connected components (combine with [6, Theorem 2.4.5]) and such that the restriction of f to the inverse image of each component is a projection, hence open. Thus $f : f^{-1}(Y \setminus Y_1) \rightarrow Y \setminus Y_1$ is open, too. Indeed, the construction (using [6, Proposition 9.18]) implies that $Y \setminus Y_1$ is homeomorphic to a submanifold of some \mathbb{R}^m . Put $Z_0 = Y \setminus Y_1$. Since $\dim(Y_1) < \dim(Y)$, the result follows by induction. \square

Remark 5.7. Let $G \subseteq \mathrm{GL}(n, \mathbb{R})$ be a real linear algebraic group, and let $G_{\mathbb{C}} \subseteq \mathrm{GL}(n, \mathbb{C})$ be its complexification. Then it follows from [7, Proposition 2.3] that each $G_{\mathbb{C}}$ -orbit in \mathbb{C}^n contains at most finitely many G -orbits in $\mathbb{R}^n \subseteq \mathbb{C}^n$.

Proof of Proposition 5.5. We first note that we may assume without loss of generality that $G = G'$. Indeed, since G has finite index in G' , every G' -orbit decomposes into finitely many G -orbits. Thus, if $\emptyset = V_0 \subseteq V_1 \subseteq \cdots \subseteq V_l = V$ is a stratification of V for the G' -action with the required properties, it is also a stratification for the G -action with the same properties. Thus we assume from now on that G is a Zariski closed subgroup of $\mathrm{GL}(V)$.

Let $V_{\mathbb{C}} \subseteq \mathbb{C}^n$ denote the complexification of V . Consider the diagram

$$\begin{array}{ccc} V & \longrightarrow & V_{\mathbb{C}} \\ \downarrow & & \downarrow \\ V/G & \longrightarrow & V_{\mathbb{C}}/G_{\mathbb{C}}. \end{array}$$

By the theorem of Rosenlicht ([36], but see also [28, Satz 2.2 on p. 23]), there exists a sequence

$$V_{\mathbb{C}} = W_0 \supseteq W_1 \supseteq W_2 \supseteq \cdots \supseteq W_r = \emptyset,$$

of Zariski-closed $G_{\mathbb{C}}$ -invariant subsets of strictly decreasing dimension such that $W_i \setminus W_{i+1}$ has closed $G_{\mathbb{C}}$ -orbits and the geometric quotient by $G_{\mathbb{C}}$ of $W_i \setminus W_{i+1}$ exists. This means that the quotient $(W_i \setminus W_{i+1})/G_{\mathbb{C}}$ can be realized as an algebraic set and the quotient map is also algebraic. Let \mathcal{O} be the first of the sets $W_i \setminus W_{i+1}$ which has nonempty intersection with V . Restricting the maps in the above diagram gives

$$\begin{array}{ccc} V \cap \mathcal{O} & \longrightarrow & \mathcal{O} \\ \downarrow & & \downarrow \\ (V \cap \mathcal{O})/G & \longrightarrow & \mathcal{O}/G_{\mathbb{C}}. \end{array}$$

The resulting map f from $V \cap \mathcal{O}$ to $\mathcal{O}/G_{\mathbb{C}}$ is an algebraic map, and hence it is a continuous semi-algebraic map. Thus it follows from Lemma 5.6 that, if Y denotes the image of $X := V \cap \mathcal{O}$ in $\mathcal{O}/G_{\mathbb{C}}$, then Y has a stratification

$$\emptyset = Z_0 \subseteq Z_1 \subseteq \cdots \subseteq Z_s = Y$$

such that $f : f^{-1}(Z_i \setminus Z_{i-1}) \rightarrow Z_i \setminus Z_{i-1}$ is open for all $1 \leq i \leq s$, each Z_i is open in Y , and the difference sets $Z_i \setminus Z_{i-1}$ are submanifolds of some \mathbb{R}^m . Put $V_i = f^{-1}(Z_i)$ for $0 \leq i \leq s$. Then $V_s = V \cap \mathcal{O}$. By Remark 5.7, if we pass through the lower left corner of the diagram, the corresponding maps $(V_i \setminus V_{i-1})/G \rightarrow Z_i \setminus Z_{i-1}$ are open, finite-to-one, onto the manifolds $Z_i \setminus Z_{i-1}$.

Now replace V by the invariant Zariski-closed subset $V \setminus \mathcal{O}$. Repeating the above arguments finitely many times gives the desired stratification (the procedure stops after finitely many steps, since any increasing sequence of Zariski open sets eventually stabilizes). \square

Using the above results, we are now able to prove

Proposition 5.8. *Suppose that G is a Lie group with finitely many components and with connected closed normal subgroups $N \subseteq H \subseteq G_0 \subseteq G$ as in (5.1), i.e., N is the nilradical of H , H/N is semi-simple and G_0/H is a torus group. Let $S \subseteq Z(N)$ be a connected closed subgroup which is normal in G , where $Z(N)$ denotes the center of N . Then \widehat{S} decomposes into a countable disjoint union of open G -invariant sets V_n such that each V_n has a stratification*

$$\emptyset = U_0 \subseteq U_1 \subseteq \cdots \subseteq U_l = V_n$$

(where l may depend on n) of open G -invariant subsets of V_n , and continuous open surjections

$$q_i : U_i \setminus U_{i-1} \rightarrow Y_i, \quad 1 \leq i \leq l,$$

such that each Y_i is a differentiable manifold and inverse images of points in Y_i are finite unions of G -orbits in V_n for all $1 \leq i \leq l$.

Proof. Let \mathfrak{s} denote the ideal of \mathfrak{g} corresponding to S . Then we may identify \widehat{S} with a closed G -stable subset of \mathfrak{s}^* of the form $R \times Z$ with R being a vector subgroup of \mathfrak{s}^* and Z a finitely generated free abelian group. Note that Z can be identified with the dual of the maximal compact subgroup in S , and therefore we can decompose Z into a disjoint union of G -orbits, which are all finite since G_0 acts trivially on Z . It then follows that \widehat{S} can be decomposed into a disjoint union of G -invariant sets of the form $R \times F$ with $F \subseteq Z$ finite.

The action of G on \widehat{S} is given via the coadjoint representation $\text{Ad}_{\mathfrak{s}}^* : G \rightarrow GL(\mathfrak{s}^*)$. Since $S \subseteq Z(N)$, it follows that this representation factors through a representation of G/N . Thus it follows from the general assumptions on G and Lemma 5.4 that the algebraic closure G' of $\text{Ad}_{\mathfrak{s}}^*(G)$ in $GL(\mathfrak{s}^*)$ is a reductive algebraic group which contains the image of $\text{Ad}_{\mathfrak{s}}^*(G)$ as a subgroup of finite index. Since the G -stable sets of the form $R \times F$ of the previous paragraph are closed algebraic subvarieties of \mathfrak{s}^* , it follows that these sets are also invariant under the action of the Zariski closure G' of $\text{Ad}_{\mathfrak{s}}^*(G)$. Thus it follows from Proposition 5.5 that for each such set we obtain a stratification

$$\emptyset = U_0 \subseteq U_1 \subseteq \cdots \subseteq U_l = R \times F$$

with the required properties. \square

We are now ready for the final step:

Proof of Proposition 2.10. By the discussion at the beginning of this section we may assume without loss of generality that G is as in (5.1), i.e., we have connected closed normal subgroups

$$N \subseteq H \subseteq G_0 \subseteq G$$

such that N is a nontrivial nilpotent group H/N is semi-simple and G_0/H is a torus group. For the induction step we have to show that every central extension

$$1 \rightarrow \mathbb{T} \rightarrow \bar{G} \rightarrow G \rightarrow 1$$

satisfies BC for \mathbb{C} . Let \bar{N} , \bar{H} and \bar{G}_0 denote the inverse images of N , H and G_0 in \bar{G} . Then the sequence of normal subgroups

$$\bar{N} \subseteq \bar{H} \subseteq \bar{G}_0 \subseteq \bar{G}$$

has the same general properties as the sequence $N \subseteq H \subseteq G_0 \subseteq G$, in particular, \bar{N} is the nilradical of \bar{H} and \bar{H} is locally algebraic. Let T denote the central copy of \mathbb{T} in \bar{G} coming from the given central extension. We now divide the proof into the following cases:

C(1) The center $S = Z(\bar{N})$ of \bar{N} has dimension greater or equal to two.

C(2) $Z(\bar{N}) = T$.

We start with Case C(1): By Theorem 2.1 (and the discussion following that theorem) it suffices to show that \bar{G}/S satisfies BC with coefficients in $C_0(\hat{S}, \mathcal{K})$ where the action of \bar{G}/S on \hat{S} is given by conjugation. By Theorem 3.6 it suffices to show that all stabilizers $(\bar{G}/S)_\chi = \bar{G}_\chi/S$ satisfy BC for \mathcal{K} and that \hat{S} has a nice stratification. While the latter follows from Proposition 5.8, the requirement on the stabilizers follows from Lemma 5.2, Remark 5.3, and the induction assumption since

$$\dim(\bar{G}_\chi/S) \leq \dim(\bar{G}) - 2 < \dim(G).$$

This finishes the proof in Case C(1).

For the proof of Case C(2) we have to do some more reduction steps in order to use the same line of arguments as in C(1). For this it is useful to consider the following two subcases:

(2)a If $\bar{Z}(N)$ denotes the inverse image of the center $Z(N)$ of N in \bar{G} , then $Z(\bar{Z}(N)) = T$.

(2)b $\dim(Z(\bar{Z}(N))) \geq 2$.

In Case (2)a we consider the normal subgroup $S = \bar{Z}(N)$ of G . Then S is a connected two-step nilpotent Lie group with one-dimensional center T , and therefore a Heisenberg group. It follows that $C_r^*(S) = C^*(S)$ can be written as the direct sum

$$C^*(S) = \bigoplus_{\chi \in \hat{T}} A_\chi$$

with

$$A_\chi \cong \mathcal{K}, \quad \text{if } \chi \neq 1, \quad \text{and} \quad A_1 = C_0(\widehat{S/T}).$$

Since \bar{G} acts trivially on \hat{T} , it follows that the decomposition action of \bar{G}/S on $C^*(S) \otimes \mathcal{K}$ induces an action on each fibre A_χ , and, by Theorem 2.1 together with Proposition 2.5, it follows that \bar{G} satisfies BC with coefficients in \mathbb{C} if \bar{G}/S satisfies BC with coefficients in $A_\chi \otimes \mathcal{K}$ for each $\chi \in \hat{T}$. If $\chi \neq 1$, we get $A_\chi \otimes \mathcal{K} \cong \mathcal{K}$, and the desired result follows from the induction assumption and the fact that $\dim(\bar{G}/S) < \dim(G)$.

So we only have to deal with the case $\chi = 1$, where we have to deal with the fibre $C_0(\widehat{S/T}, \mathcal{K}) = C_0(\widehat{Z(N)}, \mathcal{K})$. But here we are exactly in the same situation as in the proof of Case C(1), since the action of \bar{G} on $\widehat{Z(N)}$ factors through an action of G/N and all stabilizers of the characters of $Z(N)$ have dimension strictly smaller than $\dim(G)$.

We have to work a bit more for the Proof of Case (2)b. Here we put $S = Z(\bar{Z}(N))$. Then S is a connected abelian subgroup of \bar{N} and it follows from Lemma 5.2, Remark 5.3, the fact that $\dim(\bar{G}/S) < \dim(G)$ and the induction assumption that all stabilizers for the action of \bar{G}/S on \widehat{S} satisfy BC for \mathcal{K} .

Again we study the structure of the orbit space \widehat{S}/\bar{G} . For each $\chi \in \widehat{T}$ we define

$$\widehat{S}_\chi = \{\mu \in \widehat{S} : \mu|_T = \chi\}.$$

Since T is central in \bar{G} , it follows that \bar{G} acts trivially on \widehat{T} , and hence that \widehat{S}_χ is \bar{G} -invariant for all $\chi \in \widehat{T}$. Since \widehat{T} is discrete, we may write

$$C_0(\widehat{S}, \mathcal{K}) \cong \bigoplus_{\chi \in \widehat{T}} C_0(\widehat{S}_\chi, \mathcal{K})$$

with fiberwise action of \bar{G}/S . Thus by continuity of BC it suffices to deal with the single fibers. For $\chi = 1$ we are looking at the action of $\bar{G}/S \cong G/(S/T)$ on $\widehat{S}_1 \cong \widehat{S/T}$, and since S/T is a central subgroup of N we may again argue precisely as in the proof of Case C(1) to see that \bar{G}/S satisfies BC for $C_0(\widehat{S}_1, \mathcal{K})$.

In order to deal with the other fibers we are now going to show that \bar{G} acts transitively on \widehat{S}_χ for each nontrivial character $\chi \in \widehat{T}$. It follows then directly from Corollary 2.9 that \bar{G}/S satisfies BC for $C_0(\widehat{S}_\chi, \mathcal{K})$. In fact, Lemma 5.9 below shows that \bar{N} already acts transitively on \widehat{S}_χ for $\chi \neq 1$ and the result will follow from that lemma. \square

The following lemma is certainly well known to the experts on the representation theory of nilpotent groups. For the readers convenience we give the elementary proof.

Lemma 5.9. *Assume that N is a connected nilpotent Lie group with one-dimensional center $Z(N) = T$. Let S be a closed connected abelian normal subgroup of N such that $T \subseteq S$ and $S/T \subseteq Z(N/T)$. Let $1 \neq \chi \in \widehat{T}$ and let $\widehat{S}_\chi = \{\mu \in \widehat{S} : \mu|_T = \chi\}$. Then N acts transitively on \widehat{S}_χ by conjugation.*

Proof. We may assume without loss of generality that N is simply connected. In fact, if this is not the case, we pass to the universal covering group \tilde{N} of N and the universal covering $\tilde{S} \subseteq \tilde{N}$ of S and observe that there exists a discrete subgroup $D \subseteq \tilde{T} = Z(\tilde{N})$ such that $N = \tilde{N}/D$, $S = \tilde{S}/D$, $T = \tilde{T}/D$ and \widehat{S}_χ can then be (equivariantly) identified with $\widehat{\tilde{S}}_\chi$ for all $\chi \in \widehat{T} \subseteq \widehat{\tilde{T}}$.

Let \mathfrak{n} , \mathfrak{s} and \mathfrak{t} denote the Lie algebras of N , S and T , respectively. Since N is simply connected, we can write

$$N = \{\exp(X) : X \in \mathfrak{n}\}$$

with multiplication given by the Campbell-Hausdorff formula. In particular, if $Y \in \mathfrak{s}$, then

$$\exp(X)\exp(Y) = \exp(X + Y + [X, Y])$$

for all $X \in \mathfrak{n}$, since it follows from the assumption that $S/T \subseteq Z(N/T)$ that $[X, Y] \in \mathfrak{t} = \mathfrak{z}(\mathfrak{n})$ and all commutators with $[X, Y]$ vanish. In particular, if we conjugate $\exp(Y)$ by $\exp(X)$ we get the formula

$$(5.2) \quad \exp(X) \exp(Y) \exp(-X) = \exp(Y + [X, Y])$$

for all $Y \in \mathfrak{s}$.

Assume now that $\dim(\mathfrak{s}) = n + 1$ and let $0 \neq Z \in \mathfrak{t}$. We claim that we can find a basis $\{Y_1, \dots, Y_n, Z\}$ of \mathfrak{s} and elements $X_1, \dots, X_n \in \mathfrak{n}$ such that

$$(5.3) \quad [X_i, Y_i] = Z \quad \text{and} \quad [X_i, Y_j] = 0$$

for all $1 \leq i, j \leq n, i \neq j$. Indeed this follows from an easy Schmidt-orthogonalization procedure: Consider the bilinear form

$$(\cdot, \cdot) : \mathfrak{n} \times \mathfrak{s} \rightarrow \mathbb{R}; \quad (X, Y) = \lambda \Leftrightarrow [X, Y] = \lambda Z.$$

Choose $Y_1 \in \mathfrak{s} \setminus \mathfrak{t}$. Then there exists $X_1 \in \mathfrak{n}$ with $[X_1, Y_1] \neq 0$ (since Y_1 is not central), and hence we may assume that $[X_1, Y_1] = Z$. Assume now that we have chosen Y_1, \dots, Y_l and $X_1, \dots, X_l, l < n$, with the desired properties. Choose Y_{l+1} not in the span of $\{Y_1, \dots, Y_l, Z\}$ and $X' \in \mathfrak{n}$ with $[X', Y_{l+1}] = Z$. Then it is easy to check that $X_{l+1} = X' - \sum_{i=1}^l (X', Y_i) Y_i$ satisfies $[X_{l+1}, Y_{l+1}] = Z$ and $[X_{l+1}, Y_i] = 0$ for $1 \leq i \leq l$.

Now we identify S with \mathfrak{s} (via \exp) and \widehat{S} with \mathfrak{s}^* . The conjugation action of N on \widehat{S} is then transferred to the coadjoint action Ad^* . If $\{f_1, \dots, f_n, g\}$ is a dual basis for the basis $\{Y_1, \dots, Y_n, Z\}$ of \mathfrak{s} , the result will follow if we can show that

$$\text{Ad}^*(N)(\lambda g) = \text{span}\{f_1, \dots, f_n\} + \lambda g$$

for all $0 \neq \lambda \in \mathbb{R}$. By rescaling we may assume that $\lambda = 1$. But for $\lambda_1, \dots, \lambda_n \in \mathbb{R}$ we can compute

$$\left(\text{Ad}^*(\exp(\lambda_1 X_1 + \dots + \lambda_n X_n))(g) \right)(Y_i) = g(Y_i + \lambda_i [X_i, Y_i]) = \lambda_i g(Z) = \lambda_i.$$

Since Z is central in \mathfrak{n} it follows that $\left(\text{Ad}^*(\exp(X))(g) \right)(Z) = g(Z) = 1$ for all $X \in \mathfrak{n}$. Thus

$$\text{Ad}^*(\exp(\lambda_1 X_1 + \dots + \lambda_n X_n))(g) = \lambda_1 f_1 + \dots + \lambda_n f_n + g.$$

□

6. RELATIONS TO THE K -THEORY OF THE MAXIMAL COMPACT SUBGROUP

In this section we want to describe the relations between the K -theory of $C_r^*(G)$ and the K -theory of $C^*(L)$, where L denotes the maximal compact subgroup of the almost connected group G (we chose the letter L to avoid confusion). We should mention that all results presented here (accept the conclusions drawn out of our main theorem) are well known, but since they have important impact on our results, we found it useful to give at least a brief report. The main references for these results are [13, 22], and we refer especially to [13, §4] for a more geometric discussion of some the results presented in this section.

If G and L are as above, it follows from work of Abels (see [1]) that G/L is a universal proper G -space. Thus we have

$$K_*^{\text{top}}(G, A) \cong \text{KK}_*^G(C_0(G/L), A) \stackrel{\text{res}_L^G}{\cong} \text{KK}_*^L(C_0(G/L), A),$$

where the second isomorphism follows from [24, Corollary to Theorem 5.7]. Also by the work of Abels [1], G/L is a Riemannian manifold which is L -equivariantly diffeomorphic to the tangent space $V := T_{eL}$ equipped with the adjoint action of L on V . It follows then from Kasparov's work in [22] (see [9, Lemma 7.7] for a more extensive discussion) that tensoring with $C_0(V)$ gives a natural isomorphism

$$\sigma_{C_0(V)} : \mathrm{KK}_*^L(C_0(V), A) \rightarrow \mathrm{KK}_*^L(C_0(V) \otimes C_0(V), A \otimes C_0(V)),$$

and by Kasparov's Bott-periodicity theorem (see [22, Theorem 7]) we know that $C_0(V) \otimes C_0(V)$, equipped with the diagonal action, is KK^L -equivalent to \mathbb{C} (but see also the discussion below). Thus we obtain the following chain of isomorphisms

$$\begin{aligned} \mathrm{KK}_*^L(C_0(G/L), A) &\cong \mathrm{KK}_*^L(C_0(V), A) \stackrel{\sigma_{C_0(V)}}{\cong} \mathrm{KK}_*^L(C_0(V) \otimes C_0(V), A \otimes C_0(V)) \\ &\cong \mathrm{KK}_*^L(\mathbb{C}, A \otimes C_0(V)) = \mathrm{K}_*((A \otimes C_0(V)) \rtimes L), \end{aligned}$$

where the last isomorphism follows from the Green-Julg theorem. Hence, as a direct consequence of Theorem 1.2 we can deduce

Theorem 6.1. *Assume that G is an almost connected (second countable) group with maximal compact subgroup L . Let $\mathcal{K} = \mathcal{K}(H)$ be the algebra of compact operators on the separable Hilbert space H equipped with any action of G . Then $\mathrm{K}_*(\mathcal{K} \rtimes_r G)$ is naturally isomorphic to $\mathrm{K}_*((\mathcal{K} \otimes C_0(V)) \rtimes L)$.*

By Kasparov's Bott-periodicity theorem (see [22, Theorem 7]) it follows that $C_0(V)$ is KK^L -equivalent to the graded complex Clifford algebra $Cl(V)$ (with respect to a compatible inner product on V), equipped with the action of L induced by the given action on V . So we can replace $C_0(V)$ by the graded C^* -algebra $Cl(V)$, but then we have to use graded K -theory!

Let us look a bit closer to the implications of this Bott-periodicity theorem. Assume for the moment that V is even dimensional and that the action of L on V preserves a given orientation of V , i.e., the action factors through a homomorphism $\varphi : L \rightarrow SO(V)$. We have a central extension

$$0 \rightarrow \mathbb{T} \rightarrow \mathrm{Spin}^c(V) \rightarrow SO(V) \rightarrow 0$$

of $SO(V)$, where $\mathrm{Spin}^c(V) \subseteq Cl(V)$ denotes the group of complex spinors (e.g. see [2]). The corresponding action of L on $Cl(V)$ is given by the homomorphism

$$L \rightarrow SO(V) \cong \mathrm{Spin}^c(V)/\mathbb{T} = \mathrm{Ad}(\mathrm{Spin}^c(V)).$$

Now choose a fixed orthonormal base $\{e_1, \dots, e_n\}$ of V . Then the grading of $Cl(V)$ is given by conjugation with the symmetry $J = e_1 \cdot e_2 \cdots e_n \in Cl(V)$. One can show that, up to a sign, J does not depend on the choice of this basis, and the sign only depends on the orientation of the basis. In particular, J is invariant under conjugation with elements in $\mathrm{Spin}^c(V)$. From this it follows that the graded L -algebra $Cl(V)$ is L -equivariantly Morita equivalent to the trivially graded L -algebra $Cl(V)$ – a Morita equivalence is given by the module $Cl(V)$ with given L -action and grading automorphism given by left multiplication with J . Moreover, since $n = \dim(V)$ is even, $Cl(V)$ is isomorphic to the simple matrix algebra $M_{2^n}(\mathbb{C})$.

Assume now that $\dim(G/L)$ is odd. Then, replacing G by $G \times \mathbb{R}$ (with trivial action of \mathbb{R} on \mathcal{K}) we get

$$\mathrm{K}_*(\mathcal{K} \rtimes_r G) = \mathrm{K}_{*+1}(\mathcal{K} \rtimes_r (G \times \mathbb{R})).$$

Moreover, if the action of L on $V = T_{eL}$ is orientation preserving, the same is true for the resulting action of L on $V \times \mathbb{R}$, which we identify with the tangent space at eL in the group $G \times \mathbb{R}$. Hence, modulo a dimension shift, we can use the above considerations also for this case. Thus, as a consequence of Theorem 6.1 we obtain

Theorem 6.2. *Assume that G is an almost connected group with maximal compact subgroup L such that the adjoint action of L on $V = T_{eL}$ is orientation preserving. Then there are natural isomorphisms*

$$K_*(\mathcal{K} \rtimes_r G) \cong K_*((\mathcal{K} \otimes Cl(V)) \rtimes L)$$

if $\dim(G/L)$ is even and

$$K_{*+1}(\mathcal{K} \rtimes_r G) \cong K_*((\mathcal{K} \otimes Cl(V \times \mathbb{R})) \rtimes L)$$

if $\dim(G/L)$ is odd. Here all algebras are trivially graded!

Perhaps, the above result has its most satisfying formulation if translated into the language of twisted group algebras. For this let $\omega \in Z^2(G, \mathbb{T})$ denote a representative of the Mackey obstruction for the action of G on \mathcal{K} (see the discussion preceding Lemma 2.3). Then $\mathcal{K} \rtimes_r G$ is isomorphic to $C_r^*(G, \omega) \otimes \mathcal{K}$, where $C_r^*(G, \omega)$ denotes the reduced twisted group algebra $C_r^*(G, \omega)$ (e.g., see [20, Theorem 18]). Recall that $C_r^*(G, \omega)$ can be defined either as the reduced twisted crossed product $\mathbb{C} \rtimes_r (G_\omega, \mathbb{T})$ with respect to the twisted action (id, χ_1) (which, by Lemma 2.3, is Morita equivalent to the given action on \mathcal{K}), or as the completion of $L^1(G) \subseteq B(L^2(G))$, where $L^1(G)$ acts on $L^2(G)$ by the twisted convolution

$$f * \xi(s) = \int_G f(t)\omega(t, t^{-1}s)\xi(t^{-1}s) dt, \quad f \in L^1(G), \xi \in L^2(G).$$

Up to isomorphism, $C_r^*(G, \omega)$ only depends on the class $[\omega] \in H^2(G, \mathbb{T})$. Conversely, given any cocycle, the representation $\lambda_\omega : G \rightarrow U(L^2(G))$ given by

$$(\lambda_\omega(t)\xi)(s) = \omega(t, t^{-1}s)\xi(t^{-1}s)$$

determines an action of G on $\mathcal{K}(L^2(G))$ with Mackey obstruction represented by ω .

Note that the Mackey obstruction for the action of L on \mathcal{K} is given by the restriction of ω to L and the obstruction for the action of L on $Cl(V) \cong M_{2^n}(\mathbb{C})$ (if $\dim(V)$ is even) is given by the pull-back, say μ_L , to L of a cocycle representing the central extension

$$1 \rightarrow \mathbb{T} \rightarrow \text{Spin}^c(V) \rightarrow SO(V) \rightarrow 1.$$

Since $\text{Spin}^c(V) \cong (\mathbb{T} \times \text{Spin}(V))/\mathbb{Z}_2$ (diagonal action), where

$$1 \rightarrow \mathbb{Z}_2 \rightarrow \text{Spin}(V) \rightarrow SO(V) \rightarrow 1$$

is the real group of spinors, the cocycle μ_L can be chosen to take values in the subgroup $\mathbb{Z}_2 \subseteq \mathbb{T}$, and therefore $\mu_L^2 = 1$. Note that μ_L is trivial if and only if the homomorphism $\varphi : L \rightarrow SO(V)$ factorizes through $\text{Spin}^c(V)$ (i.e., if and only if G/L carries a G -invariant Spin^c -structure). If $\dim(V)$ is odd, we may define μ_L in the same way as above, noticing that this cocycle is equivalent to the pull back of (a cocycle representing) the extension

$$1 \rightarrow \mathbb{T} \rightarrow \text{Spin}^c(V \times \mathbb{R}) \rightarrow SO(V \times \mathbb{R}) \rightarrow 1,$$

which follows from the fact that L acts trivially on \mathbb{R} ! Since the Mackey obstruction of a tensor product of actions is the product of the Mackey obstructions of the factors, we obtain

Theorem 6.3. *Assume that G is an almost connected group with maximal compact subgroup L such that the adjoint action of L on $V = T_{eL}$ is orientation preserving. Let $n = \dim(G/L)$ and let $\omega \in Z^2(G, \mathbb{T})$ be any cocycle on G . Then*

$$K_* (C_r^*(G, \omega)) \cong K_{*+n} (C^*(L, \omega \cdot \mu_L)).$$

In particular, in the special case where ω is trivial, we obtain an isomorphism

$$K_* (C_r^*(G)) \cong K_{*+n} (C^*(L, \mu_L)).$$

Again, μ_L is trivial if and only if G/L carries a G -invariant Spin^c -structure. In general, since $C^*(L, \omega \cdot \mu_L)$ is the quotient of the central extension $L_{\omega \cdot \mu_L}$ of L by \mathbb{T} corresponding to the character χ_1 of \mathbb{T} , it follows that $C^*(L, \omega \cdot \mu_L)$ is a direct sum of (possibly infinitely many) matrix algebras. Thus as a direct corollary of the above result we obtain:

Corollary 6.4. *Assume that G , L and ω are as in Theorem 6.3. Then $K_{0+n} (C_r^*(G, \omega))$ is isomorphic to a free abelian group in at most countably many generators and $K_{1+n} (C_r^*(G, \omega)) = \{0\}$.*

This result has interesting consequence towards the question of existence of square integrable representations of connected unimodular Lie groups. In fact, combining the above corollary with [35, Theorem 4.6] gives:

Corollary 6.5 (cf [35, Corollary 4.7]). *Let G be a connected unimodular Lie group. Then all square-integrable factor representations of G are type I. Moreover, G has no square-integrable factor representations if $\dim(G/L)$ is odd, where L denotes the maximal compact subgroup of G .*

We refer to [35] for more detailed discussions on this kind of applications of the positive solution of the Connes-Kasparov conjecture. Note that Theorem 6.3 and Corollary 6.4 do not hold in general without the assumption that the action of L on $V = T_{eL}$ is orientation preserving. In fact an easy six-term-sequence argument shows that it cannot hold for the group $G = \mathbb{R} \rtimes \mathbb{Z}_2$, where \mathbb{Z}_2 acts on \mathbb{R} by reflection through 0.

REFERENCES

- [1] H. Abels. *Parallelizability of proper actions, global K-slices and maximal compact subgroups*, Math. Ann. **212**, 1–19 (1974).
- [2] M. Atiyah, R. Bott and A. Shapiro. *Clifford Modules*, Topology 3, Suppl. 1, 3-38 (1964).
- [3] P. Baum, A. Connes and N. Higson. *Classifying space for proper actions and K-theory of group C^* -algebras*, Contemp. Math. **167**, 241-291 (1994).
- [4] B. Blackadar. *K-theory for operator algebras*, MSRI pub., **5**, (1986), Springer Verlag.
- [5] E. Blanchard. *Deformations de C^* -algebres de Hopf*, Bull. Soc. Math. Fr. **124**, 141-215 (1996).
- [6] J. Bochnak, M. Coste, and M.-F. Roy. *Real algebraic geometry*. Ergebnisse der Mathematik und ihrer Grenzgebiete Vol. 36, Springer 1998.
- [7] A. Borel and Harish-Chandra. *Arithmetic subgroups of algebraic groups*. Annals of Math. **75** (1962), 485–535.
- [8] A. Borel. *Linear Algebraic Groups*, Springer Verlag **GTM 126** (1991).
- [9] J. Chabert and S. Echterhoff. *Twisted equivariant KK-theory and the Baum-Connes conjecture for group extensions*, To appear in *K-Theory*.
- [10] J. Chabert and S. Echterhoff. *Permanence properties of the Baum-Connes conjecture*, Doc. Math. **6**, 127–183 (2001).

- [11] J. Chabert, S. Echterhoff, and Ralf Meyer. *Deux remarques sur la conjecture de Baum-Connes*, C. R. Acad. Sci., Paris, Ser. I **332**, Série I, 607–610 (2001).
- [12] C. Chevalley. *Theorie des groupes de Lie. Groupes algebriques. Theoremes generaux sur les algebres de Lie*. 2ieme ed. Paris: Hermann & Cie. IX (1968).
- [13] A. Connes and H. Moscovici. *The L^2 -index theorem for homogeneous spaces of Lie groups*, Ann. Math., II. Ser. **115**, 291-330 (1982).
- [14] S. Echterhoff. *On induced covariant systems*. Proc. Amer. Math. Soc. **108** (1990), 703–708.
- [15] S. Echterhoff. *Morita equivalent actions and a new version of the Packer-Raeburn stabilization trick*. J. London Math. Soc. (2), **50** (1994), 170–186.
- [16] S. Echterhoff. *Crossed products with continuous trace*, Mem. Amer. Math. Soc. **123** (1996), no. 586, 1–134.
- [17] G. Elliott, T. Natsume, R. Nest. *The Heisenberg group and K-theory*, K-Theory **7**, 409-428 (1993).
- [18] J. Fell. *The structure of algebras of operator fields*, Acta Math. **106**, 233-280 (1961).
- [19] J. Glimm. *Locally compact transformation groups*, Trans. Am. Math. Soc. **101**, 124-138 (1961).
- [20] P. Green. *The local structure of twisted covariance algebras*. Acta. Math., **140** (1978), 191–250.
- [21] N. Higson and G. Kasparov, *E-theory and KK-theory for groups which act properly and isometrically on Hilbert space*, Invent. Math. **144**, 23–74, (2001).
- [22] G. Kasparov. *The operator K-functor and extensions of C^* -algebras*, Math. USSR Izvestija, Vol. **16**, No. 3 (1981) 513–572.
- [23] G. Kasparov. *K-theory, group C^* -algebras, higher signatures (Conspectus)*. in: Novikov conjectures, index theorems and rigidity. London Math. Soc, lecture notes series **226** (1995), 101–146.
- [24] G. Kasparov. *Equivariant KK-theory and the Novikov conjecture*, Invent. Math. **91**, (1988) 147-201.
- [25] G. Kasparov and G. Skandalis, *Groups acting properly on “bolic” spaces and the Novikov conjecture*, Preprint (1999).
- [26] E. Kirchberg and S. Wassermann. *Exact groups and continuous bundles of C^* -algebras*, Math. Ann. **315**, 169-203 (1999).
- [27] E. Kirchberg and S. Wassermann. *Permanence properties of C^* -exact groups*. Doc. Math. **5** (2000), 513–558.
- [28] H. Kraft, P. Slodowy, and T.A. Springer. *Algebraische Transformationsgruppen und Invariantentheorie*. DMV-Seminar Band 13, Birkhäuser 1989.
- [29] V. Lafforgue, *K-theory bivariente pour les algebres de Banach et conjecture de Baum-Connes*, PhD Dissertation, Universite Paris Sud, (1999).
- [30] G. Mackey. *Borel structure in groups and their duals*, Trans. Am. Math. Soc. **85**, 134-165 (1957).
- [31] D. Montgomery and L. Zippin. *Topological transformation groups*. Interscience Tracts in Pure and Applied Mathematics. New York: Interscience Publishers, Inc. XI (1955).
- [32] J. Packer and I. Raeburn. *Twisted crossed products of C^* -algebras*, Math. Proc. Camb. Philos. Soc. **106**, 293-311 (1989).
- [33] G.K. Pedersen. *C^* -Algebras and their Automorphism Groups*. Academic Press, London, 1979.
- [34] Lajos Pukánszky. *Characters of connected Lie groups*. Mathematical surveys and Monographs Vol 71. American Mathematical Society, Rhode Island 1999.
- [35] J. Rosenberg. *Group C^* -algebras and topological invariants*. In: Operator algebras and group representations, Proc. int. Conf., Neptun/Rom. 1980, Vol. II, Monogr. Stud. Math. **18**, 95-115 (1984).
- [36] M. Rosenlicht. *A remark on quotient spaces*. An. Acad. Brasil. Ciênc. **35** (1963), 487–489.
- [37] J.L. Tu. *La conjecture de Novikov pour les feuilletages hyperboliques*, K-theory, **16**, No.2, 129-184 (1999).
- [38] A. Wassermann. *Une demonstration de la conjecture of Connes-Kasparov pour les groupes de Lie lineaires connexes reductifs*, C. R. Acad. Sci., Paris, Ser. I **304**, 559-562 (1987).

JÉRÔME CHABERT: UNIVERSITÉ BLAISE PASCAL, BÂT. DE MATHÉMATIQUES, 63177, AUBIÈRE, FRANCE

WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, MATHEMATISCHES INSTITUT, EINSTEINSTR. 62 D-48149 MÜNSTER, GERMANY

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5, DK-2100
COPENHAGEN, DENMARK

E-mail address: `chabert@math.univ-bpclermont.fr`, `echters@math.uni-muenster.de`, `rnest@math.ku.dk`