Relative family Gromov-Witten invariants and symplectomorphisms

Olguța Bușe^{*}

November 6, 2018

Abstract

We study the symplectomorphism groups $G_{\lambda} = \operatorname{Symp}_{0}(M, \omega_{\lambda})$ of an arbitrary closed manifold M equipped with a 1-parameter family of symplectic forms ω_{λ} with variable cohomology class. We show that the existence of nontrivial elements in $\pi_{*}(\mathcal{A}, \mathcal{A}')$, where $(\mathcal{A}, \mathcal{A}')$ is a suitable pair of spaces of almost complex structures, implies the existence of families of nontrivial elements in $\pi_{*-i}G_{\lambda}$, for i = 1 or 2. Suitable parametric Gromov Witten invariants detect nontrivial elements in $\pi_{*}(\mathcal{A}, \mathcal{A}')$. By looking at certain resolutions of quotient singularities we investigate the situation $(M, \omega_{\lambda}) = (S^{2} \times S^{2} \times X, \sigma_{F} \oplus \lambda \sigma_{B} \oplus \omega_{st})$, with (X, ω_{st}) an arbitrary symplectic manifold. We find families of nontrivial elements in $\pi_{k}(G_{\lambda}^{X})$, for countably many k and different values of λ . In particular we show that the fragile elements w_{ℓ} found by Abreu-McDuff [3] in $\pi_{4\ell}(G_{\ell+1}^{\operatorname{pt}})$ do not disappear when we consider them in $S^{2} \times S^{2} \times X$.

1 Introduction

Consider (M^{2n}, ω) a 2*n* dimensional compact symplectic manifold. A basic invariant which distinguishes among different symplectic structures on M is the group of symplectomorphisms, $\operatorname{Symp}(M, \omega)$. This is an infinite dimensional group endowed with a natural C^{∞} topology.

Two natural questions arise in relation with $\text{Symp}(M, \omega)$ namely

- (1) What can be said about the topological type of $\text{Symp}(M, \omega)$?
- (2) How does the topological type change as ω varies?

Research has been done in this direction by various authors [Abreu [2], Le-Ono [9], Mc-Duff [10], Seidel [15]] by using information on *J*-holomorphic curves. We investigate these questions by defining relative parametric GW invariants, which are sensitive to the topology of appropriate spaces of almost complex structures. The connection between the spaces of

^{*}SUNY at Stony Brook, buse@math.sunysb.edu

almost complex structures and the symplectomorphism groups is achieved by means of the following fibration, introduced by Kronheimer [7] and used in McDuff [11]:

$$\operatorname{Symp}_{0}(M,\omega) \longrightarrow \operatorname{Diff}_{0}(M) \xrightarrow{\psi \to (\psi^{-1})^{*}\omega} S_{\omega}$$
(1)

where S_{ω} is the space of symplectic forms which can be joined to ω through a path of cohomologous symplectic forms, $\operatorname{Diff}_0(M)$ is the connected component of the identity inside the group of diffeomorphism and $\operatorname{Symp}_0(M,\omega) = \operatorname{Symp}(M,\omega) \cap \operatorname{Diff}_0(M)$. Now consider \mathcal{A}_{ω} the space of all almost complex structures tamed by some symplectic form ω' in \mathcal{S}_{ω} . By [11], \mathcal{A}_{ω} is homotopy equivalent to \mathcal{S}_{ω} . This yields the following homotopy fibration:

$$\operatorname{Symp}_{0}(M,\omega) \longrightarrow \operatorname{Diff}_{0}(M) \longrightarrow \mathcal{A}_{\omega}.$$
(2)

Our strategy will be to define suitable pairs $(\mathcal{A}, \mathcal{A}')$ of spaces of almost complex structures, such that information on nontrivial homotopy groups in $(\mathcal{A}, \mathcal{A}')$ extends to information on $\operatorname{Symp}_0(M, \omega)$. We develop a version of relative GW invariants in family which detects such nontrivial elements in $\pi_*(\mathcal{A}, \mathcal{A}')$.

Outline of the methods

In section **2** we will define the invariants as follows: Consider $D \in H_2(M, \mathbb{Z})$ and let $\mathcal{A}_{\omega,D}^c$ be the subspace of \mathcal{A}_{ω} consisting of those almost complex structures J which do not admit J-holomorphic stable maps in the class D. For I an interval in \mathbb{R} further define $(\mathcal{A}_I, \mathcal{A}_{I,D}^c) = \bigcup_{\omega_{\lambda} \in L} (\mathcal{A}_{\omega_{\lambda}}, \mathcal{A}_{\omega_{\lambda},D}^c)$, where the cohomology of symplectic form $[\omega_{\lambda}]$ is deformed along a line L inside a positive cone $\mathcal{K} \in H^2(M, \mathbb{R})$.

Consider a family of almost complex structures $(J_B, J_{\partial B})$ that represent an element in $\pi_*(\mathcal{A}_I, \mathcal{A}_{I,D}^c)$. We will define a homomorphism

$$PGW_{D,0,k}^{M,(J_B,J_{\partial B})}: \bigoplus_{i=1}^k H^{a_i}(M,\mathbb{Q})^k \to \mathbb{Q}$$
(3)

by counting J_b -holomorphic stable maps in class D, for all $b \in B$. This is well defined because the class D is never represented as a J_b -holomorphic stable maps if $b \in \partial B$. We have the following

Theorem 1.1 i) The invariants $PGW_{D,0,k}^{M,(J_B,J_{\partial B})}$ are symplectic deformation invariants and depend only on the relative homotopy class of the pair $(J_B, J_{\partial B})$.

ii) For a fixed choice of k, D and α_i the map $\Theta_{0,k,\alpha_1,\dots,\alpha_k} : \pi_*(\mathcal{A}_I, \mathcal{A}_{I,D}^c) \to \mathbb{Q}$, given by

$$\Theta_{k,\alpha_1,\ldots,\alpha_k}([(J_B, J_{\partial B})] = PGW_{D,0,k}^{M,(J_B,J_{\partial B})}(\alpha_1,\ldots,\alpha_k)$$

is a homomorphism.

The reason why (i) holds is that the class D is never represented for a J_b with $b \in \partial B$.

In section **3** we will exhibit some examples of nontrivial PGW. There we consider the case when (M, ω) is $S^2 \times S^2 \times X$, where X is an arbitrary symplectic manifold and $\omega = \omega_\lambda \oplus \omega_{st}$, with $\omega_\lambda = \sigma_F \oplus \lambda \sigma_B$. Here σ_F, σ_B are forms on the fiber and base respectively, of total area $1, \lambda \geq 1$, and ω_{st} is arbitrary symplectic form on X. The families $(J_B, J_{\partial B})$ of almost complex structures are provided for $S^2 \times S^2$ in [7] and then further investigated in [3]. One has to look at a quotient singularity, $\mathbb{C}^2/C_{2\ell}$, where $C_{2\ell}$ is the cyclic group of order 2ℓ acting diagonally by scalars on \mathbb{C}^2 . The deformation space for the canonical resolution of this singularity provides a $4\ell - 2$ family $(J_{B_\ell}, \partial J_{B_\ell}) \in (\mathcal{A}_{[\ell+\epsilon,\ell]}, \mathcal{A}_\ell)$ for which suitable PGW are nontrivial.

The link between these examples and the corresponding groups of symplectomorphisms will be explained in section **4**. It will be there where we explain the extent to which the known homotopy properties (see [3]) of $\operatorname{Symp}_0(S^2 \times S^2, \omega_\lambda)$ are reflected in the high homotopy groups of $G_\lambda^X := \operatorname{Symp}_0(S^2 \times S^2 \times X, \omega_\lambda \oplus \omega_{st})$. For every M, ω_λ a general symplectic manifold, we set the notation $G_\lambda := \operatorname{Symp}_0(M, \omega_\lambda)$.

To be able to give any answers related to the two questions posed in the beginning, one has to establish first a more precise language in which they make sense. One of the difficulties is that in general there is no direct map $G_{\lambda} \to G_{\lambda+\epsilon}$. In the particular situation $M = S^2 \times S^2 \times \text{pt}$ Abreu-McDuff in [3] and [11] find natural maps $G_{\lambda}^{\text{pt}} \to G_{\lambda+\epsilon}^{\text{pt}}$, well defined up to homotopy, and prove:

Theorem 1.2 (Abreu-McDuff) (i) The homotopy type of G_{λ}^{pt} is constant on all the intervals $(\ell - 1, \ell]$ with $\ell \geq 2$ a natural number. Moreover, as λ passes an integer $\ell, \ell \geq 2$ the groups $\pi_i(G_{\lambda}^{\text{pt}}), i \leq 4\ell - 5$, do not change.

(ii) There is an element $w_{\ell} \in \pi_{4\ell-4}(G_{\lambda}^{\text{pt}}) \times \mathbb{Q}$ when $\ell-1 < \lambda \leq \ell$ that vanishes for $\lambda > \ell$.

When we deal with a general manifold M, to get around the fact that there is no map $G_{\lambda} \to G_{\lambda+\epsilon}$ we show that for any compact $K \subset G_{\lambda}$, the inclusion $0 \times K \subset G_{\lambda}$ extends to a map h that fits into the following commuting diagram:

$$h: [-\epsilon, \epsilon] \times K \longrightarrow \mathcal{G} := \bigcup (G_{\lambda} \times \lambda) \subset \text{Diff} \times \mathbb{R}$$

$$\downarrow^{pr_1} \qquad \qquad \downarrow^{pr_2}$$

$$[-\epsilon, \epsilon] \xrightarrow{incl} \qquad (-\infty, \infty)$$

$$(4)$$

Moreover, for any two such maps h and h' which coincide on $0 \times K$, there is, for ϵ' small enough, a homotopy between them $H : [0,1] \times [-\epsilon',\epsilon'] \times K \to \mathcal{G}$ which also preserves the fibers of the natural projections. We therefore see that, for any cycle ρ in G_{λ} there are extensions ρ_{ϵ} in $G_{\lambda+\epsilon}$ which, for ϵ sufficiently small, are unique up to homotopy. Hence they give well defined elements in $\pi_*G_{\lambda+\epsilon}$.

It will therefore make sense to ask what will become of an element $\rho \in \pi_*G_\lambda$ inside $\pi_*G_{\lambda+\epsilon}$, for small ϵ . In this language we say that an element $\theta_\ell \in \pi_*G_\ell$ is *fragile* if any extension $\theta_{\ell+\epsilon}$ is null-homotopic in $\pi_*(G_{\ell+\epsilon})$ for $\epsilon > 0$. Also, we say that a family $\eta_{\ell+\epsilon} \in \pi_*G_{\ell+\epsilon}, 0 < \epsilon$ is *new* if there is no $\eta_\ell \in \pi_*G_\ell$ whose extension is $\eta_{\ell+\epsilon}$. We consider the space \mathcal{A}_{ℓ^+} roughly given by $\mathcal{A}_{\ell^+} := \bigcap_{0 < \epsilon < \epsilon_0} \mathcal{A}_{\ell+\epsilon}$.

We say that an element $\alpha \in \pi_*(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$ is *persistent* if it has nonzero image under the map $\pi_*(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell}) \to \pi_*(\mathcal{A}_{[\ell, \ell+\epsilon]}, \mathcal{A}_{\ell}).$

The content of our main theorem is the following:

Theorem 1.3 Assume that we have a persistent element $0 \neq \beta_{\ell} \in \pi_k(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$ Exactly one of the statements below holds.

A) There is an uniquely associated non-zero fragile element $\theta_{\ell} \in \pi_{k-2}G_{\ell}$, such that $i_*(\theta_{\ell}) = 0$ in $\pi_{k-2}\text{Diff}_0(M)$.

B) There exists an $\epsilon_{\ell} > 0$ and an uniquely associated family of **new** elements $0 \neq \eta_{\ell+\epsilon} \in \pi_{k-1}G_{\ell+\epsilon}, 0 < \epsilon < \epsilon_{\ell}$.

We should point out that our methods do not allow us to decide in general whether the image of $\eta_{\ell+\epsilon}$ in $\text{Diff}_0(M)$ is zero or not.

We show that the hypothesis of the theorem is verified when $M = S^2 \times S^2 \times X$. We consider $D = A - \ell F$. Since $[\sigma_F \oplus \lambda \sigma_B \oplus \omega_{st}](A - \ell F) = 0$ we get that $\mathcal{A}_{\ell} \subset \mathcal{A}^c_{[\ell,\ell+\epsilon],D}$. In this situation the $4\ell - 2$ dimensional elements $(B_{\ell}, \partial B_{\ell})$ obtained in section **3** are detected as nontrivial in $\pi_{4\ell-2}(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$ and are persistent. In fact in general PGW invariants detect persistent elements. By varying the value of the integer ℓ we obtain infinitely many values of λ for which higher order homotopy groups of G_{λ}^X are nontrivial and also make a more detailed discussion regarding the stability of the elements w_{ℓ} provided by theorem 1.2 inside G_{λ}^X . This is the content of the following:

Corollary 1.4 For any natural number $\ell \geq 1$, exactly one of the statements below holds.

A) There is a non-zero fragile element $w_{\ell}^X \in \pi_{4\ell-4}G_{\ell}^X$, such that $i_*(w_{\ell}^X) = 0$ in $\pi_*\text{Diff}_0(S^2 \times S^2 \times X)$. This element can be identified with $w_{\ell} \times id$.

B) There exists an $\epsilon_{\ell} > 0$ and a family of **new** elements $0 \neq \eta_{\ell+\epsilon}^X \in \pi_{4\ell-3}G_{\ell+\epsilon}^X, 0 < \epsilon < \epsilon_{\ell}$.

In particular this shows that the fragile elements obtained by Abreu-McDuff for $\ell > 1$ do not disappear when we consider them inside $S^2 \times S^2 \times X$. One possibility is that $0 \neq w_{\ell} \times id \in \pi_{4\ell-4}(G_{\ell}^X)$. If this is not the case, then we have the associated new $4\ell-3$ dimensional elements $0 \neq \eta_{\ell+\epsilon}^X$ in $\pi_{4\ell-3}G_{\ell+\epsilon}^X$ for small $\epsilon > 0$. For general X, when $\ell = 1$ it is known by work of Le-Ono that **B** takes place and $0 \neq i_*(\eta_{\ell+\epsilon}) \in \text{Diff}(S^2 \times S^2 \times X)$. Also, for X = ptand $\ell > 1$ from the work of Abreu-McDuff we know that **A** takes place.

We do not have examples when case **B** takes place and $i_*(\eta_{\ell+\epsilon}) \neq 0 \in \text{Diff}(M)$.

Our method had been inspired by P. Kronheimer's work, who uses parametric Seiberg-Witten invariants in dimension 4, as well as by the work of D. McDuff [10]. Similar work has been done in this direction by Le-Ono in[9]; by looking at related but slightly different parametric GW invariants they get results about $\pi_i(\text{Symp}_0(S^2 \times S^2 \times X, \omega_{\omega_1 \oplus \omega_{st}}))$ when i = 1, 3. In section **3** we could consider $\mathbb{C}^2/\mathbb{C}_{2\ell+1}$ instead and by carrying out similar arguments get the same type of results for $\mathbb{C}P^2 \#\mathbb{C}P^2 \times X$.

Acknowledgments This is part of the author's doctoral research at SUNY Stony Brook. The author would like to thank her advisor, Dusa McDuff, for her suggestions, advice and comments on earlier drafts.

2 Relative parametric GW invariants

2.1 Definition and properties

Consider B to be a compact manifold with boundary and a smooth map $i : (B, \partial B) \rightarrow (\mathcal{A}_I, \mathcal{A}_{I,D}^c)$. Although the invariants can be defined with this data regarding the parameter space, for the applications we have in mind we will consider B to be an n-ball such that i represents a relative homotopy class in $\pi_*(\mathcal{A}_I, \mathcal{A}_{I,D}^c)$. We will often write $J_b := i(b)$ and

 $J_B = \operatorname{im}(i)$, and refer to $\operatorname{im}B$ in \mathcal{A}_I as J_B . Consider also a smooth family of symplectic forms $(\omega_b)_{b\in B}$ where ω_b tames J_b . We point out that the ω_b need not be cohomologous, as the taming condition is an open condition. Our goal here is to show how we can define parametric GW invariants relative to the boundary ∂J_B , which count J_b holomorphic maps for some $b \in B$. These will not depend either on deformations of the family ω_B or on the relative homotopy class $(J_B, \partial J_B) \subset (\mathcal{A}_I, \mathcal{A}_{I,D}^c)$.

Consider $\widetilde{\mathcal{M}}_{0,k}^*(M, D, (J_B, \partial J_B))$ the space of tuples (b, f, x_1, \ldots, x_k) where $f: S^2 \to M$ is a simple ¹ J_b -holomorphic map in class D, for some $b \in B$ and x_i are pairwise distinct points on S^2 . We will consider

$$\mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B)) = \widetilde{\mathcal{M}}_{0,k}^*(M, D, (J_B, \partial J_B))/G$$

where $G = \text{PSL}(2, \mathbb{C})$ acts on the moduli space by reparametrizations of the domain. Denote the elements of $\mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B))$ by $[b, f, x_1, \ldots, x_k]$.

In the best scenario, for a good choice of $(J_B, \partial J_B)$, the following hold:

(1) $\mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B))$ is a manifold of dimension $2n + 2c_1(D) + 2k + \dim B$ and (2) $\mathcal{M}_{0,k}^* := \mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B))$ is compact. Then the image of the map

$$ev: \mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B)) \to M^k \tag{5}$$

with $ev([b, f, x_1, \ldots, x_k]) := (f(x_1), \ldots, f(x_k))$ will provide a cycle $ev_*(\mathcal{M}_{0,k}^*)$ in M^k which, by intersection with homology classes of complementary dimension in M^k , gives the parametric Gromov-Witten invariants.

As we will see in the regularity discussion below, (1) is always possible to accomplish by Sard-Smale theorem. However, (2) is seldom true; the compactification $\overline{\mathcal{M}}_{0,k}(M, D, (J_B, \partial J_B))$ of $\mathcal{M}_{0,k}^*(M, D, (J_B, \partial J_B))$ contains both *stable maps* and *nonsimple curves*, which we sometimes call multiple cover curves. We will spell out some of these notions later in this section; for more information, the reader can check [12], [8], [14], [4].

 $\overline{\mathcal{M}}_{0,k}(M, D, (J_B, \partial J_B))$ is a stratified space. The best we can hope is that the image of the evaluation map $ev : \mathcal{M}_{0,k}^* \to M^k$ a *pseudo-cycle*, or differently said, the boundary strata in the image will be codimension 2 or bigger. If this scenario works we can still define the PGW as the intersection between the image of ev and classes of complementary dimension in $H_*(M^k)$. This will be for instance the case when the class D is J_b indecomposable for any $b \in B$, that is, no J_b holomorphic map in class D can decompose into a connected union of J_b holomorphic spheres $C = C^1 \bigcup C^2 \bigcup \ldots \bigcup C^N$ such that each C^i represents the class D_i and $D = D_1 + \ldots + D_N$. In fact, in this situation the image of ev is a cycle. This hypothesis will be enough for the application we have in mind.

Definition 2.1 We will say that the hypothesis H_1 is satisfied if the class D is J_b indecomposable for any $b \in B$.

Parametric regularity

We begin by explaining what is *D*-parametric regularity and contrast it with the usual D-regularity for J (see [12]). For this we need to introduce the following facts.

¹We say that $f: \Sigma \to M$ is simple if it is not the composite of a holomorphic branched covering map $(\Sigma, j) \to (\Sigma', j')$ of degree greater than 1 with a J-holomorphic map $\Sigma' \to M$.

Let $\mathcal{X} = \operatorname{Map}(\Sigma, M; D)$ be the space of somewhere injective ² smooth maps $f: \Sigma \to M$ representing class D. This is an infinite dimensional manifold with $T_f \mathcal{X} = C^{\infty}(f^*TM)$. We will next consider the following generalized vector bundle $\mathcal{E} \longrightarrow B \times \mathcal{X}$, whose fiber at (b, f)is the space $\mathcal{E}_{b,f} = \Omega_{J_b}^{0,1}(f^*TM)$ of smooth J_b antilinear forms with values in f^*TM . In this vector bundle we consider a section $\Phi: B \times \mathcal{X} \longrightarrow \mathcal{E}$, given by

$$\Phi(b,f) = \frac{1}{2}(df + J_b \circ df \circ j) \tag{6}$$

The zeros of Φ are precisely J_b holomorphic maps and thus the moduli space

$$\mathcal{M}_{0,0}^*(M, D, (J_B, \partial J_B)) = \Phi^{-1}(0)$$

is the intersection of im Φ with the zero section of the bundle. Since we would like $\widetilde{\mathcal{M}}^*_{0,k}(M, D, (J_B, \partial J_B))$ to be a manifold we require that Φ is transversal to the zero section. This means that the image of $d\Phi(b, f)$ is complementary to the tangent space $T_b B \oplus T_f \mathcal{X}$ of the zero section. But for any f which is J_b holomorphic, $d\Phi$ is given by

$$d\Phi(b,f): T_bB \oplus C^{\infty}(f^*TM) \longrightarrow T_bB \oplus T_f\mathcal{X} \oplus \mathcal{E}_{b,f}$$

If we consider now the projection onto the vertical space of the bundle:

$$proj_2: T_bB \oplus T_f\mathcal{X} \oplus \mathcal{E}_{b,f} \longrightarrow \mathcal{E}_{b,f}$$

the above transversality translates into the fact that

$$d\Phi(b,f) \circ proj_2 : T_b B \oplus C^{\infty}(f^*TM) \longrightarrow \Omega^{0,1}_{J_b}(\Sigma, f^*TM)$$
(7)

is onto. We will make the notation $D\Phi(b, f) = d\Phi(b, f) \circ proj_2$. We then have:

Definition 2.2 We say that a J_b holomorphic map f is J_B parametric regular if $D\Phi(b, f)$ is onto.

Observation: The linearized operator is well defined if there is no pair (b, f) with $f J_b$ holomorphic and $b \in \partial B$. This is precisely the condition we imposed on $(J_B, \partial J_B)$ to give a relative cycle in $(\mathcal{A}_I, \mathcal{A}_{I,D}^c)$.

Definition 2.3 Consider (J_B, ω_B) as above. We say that $(J_B, J_{\partial B})$ is an D-parametric regular family of almost complex structures if any J_b holomorphic map in class D is parametric regular. We denote by $J_{preg}(D)$ the set of all D-parametric regular families $(J_B, \partial J_B) \in (\mathcal{A}_I, \mathcal{A}_{I,D}^c)$.

In order to apply the implicit function theorem and Sard-Smale theorem we must work on Banach manifolds and hence complete all spaces under suitable Sobolev norms. For example, one should to work on spaces consisting of almost complex structures of class C^l , on $\mathcal{X}^{k,p}$, with kp > 2, the space of maps whose k-th derivatives are of class L^p . Also, we should work on

$$\mathcal{E}_f^p = L^p(\Lambda^{0,1} \otimes_J f^*TM))$$

²We say that a map $f : \sigma \to M$ is somewhere injective if $df(z) \neq 0, f^{-1}(f(z) = z$ for some $z \in \Sigma$. A simple J-holomorphic map is somewhere injective (see [12].

rather that with $\Omega_J^{0,1}(\Sigma, f^*TM)$. There are standard arguments [12] to show that one can pass the following arguments from spaces of C^l objects (which are Banach manifolds) to spaces of C^{∞} objects (which are Frechet manifolds). For simplicity we will drop the superscripts l, k, p unless it will be relevant to specify them. We have the following:

Theorem 2.4 If $J_B \in J_{preg}(D)$, then the moduli space $\widetilde{\mathcal{M}}^*_{0,0}(M, D, (J_B, \partial J_B))$ is a smooth open manifold of dimension $2n + 2c_1(D) + \dim B$, with a natural orientation.

Moreover, if one considers $\widetilde{\mathcal{M}}_{0,0}^*(M, D, (J_B, \partial J_B)) \times (S^2)^k$ and takes away all the diagonals of the type $\widetilde{\mathcal{M}}_{0,0}^*(M, D, (J_B, \partial J_B)) \times \operatorname{diag}_{i,j}$, what we obtain is precisely $\widetilde{\mathcal{M}}_{0,k}^*(M, D, (J_B, \partial J_B))$ This will therefore be a manifold of dimension $2n + 2c_1(D) + \dim B + 2k$.

Let $\mathcal{M}_{0,0}^*(M, D, \mathcal{A}_I)$ be the *universal* moduli space consisting of pairs (f, J) where $J \in \mathcal{A}_I$ and f is J-holomorphic. It will be more relevant to the story to point out the following characterization of parametric regularity.

Proposition 2.5 Consider the diagram

$$\begin{array}{ccc}
\widetilde{\mathcal{M}}_{0,0}^{*}(M, D, \mathcal{A}_{I}) & (8) \\
& & & \downarrow \Pi \\
(B, \partial B) \xrightarrow{i} (\mathcal{A}_{I}, \mathcal{A}_{I,D}^{c})
\end{array}$$

Then $J_B \in J_{preg}(A)$ iff $i \pitchfork \Pi$.

Proof: For simplicity we will denote by $D_{f,b} = D\Phi(b, f)_{|C^{\infty}(f^*(TM))|}$. By (7) the surjectivity of $D\Phi(b, f)$ is then equivalent with the surjectivity of the following linear operator

$$D\phi_{|T_bB}: T_bB \to \operatorname{coker} D_{b,f}$$

We will denote i(b) = J. The tangent space $T_J \mathcal{A}_I$ to \mathcal{A}_I consists of all sections Y of the bundle $\operatorname{End}(TM, J)$ whose fiber at $p \in M$ is the space of linear maps $Y : T_pM \to T_pM$ such that YJ + JY = 0; we will consider the map

$$R: T_J \mathcal{A}_I \to \Omega_J^{0,1}(\Sigma, f^*TM)$$

given by $R(Y) = \frac{1}{2}Y \circ df \circ j$. The map

$$d\Pi: T_{f,J}\widetilde{\mathcal{M}}^*_{0,0}(M,D,\mathcal{A}_I) \to T_J\mathcal{A}_I$$

is given by $d\Pi(\xi, Y) = Y$, where the pair (ξ, Y) is in $T_{f,J}\widetilde{\mathcal{M}}^*_{0,0}(M, D, \mathcal{A}_I)$ if and only if

$$D_{f,b}(\xi) + R(Y) = 0 (9)$$

From this one can see that $\operatorname{im} D_{f,b} = R(\operatorname{im}(d\Pi))$. Since $D_{b,f}$ is elliptic and $\ker R \subset \operatorname{im} d\Pi$, it follows that $\operatorname{coker} d\Pi$ has finite dimension. If we consider the map $\mathcal{F} : \mathcal{X} \times \mathcal{A}_I \to \mathcal{E}$, given by $\mathcal{F}(f, J) = \overline{\partial}_J(f)$ then (see [12]) the linearization at a zero (f, J) with f simple is onto. That is

$$D\mathcal{F}(f,J)(\xi,Y) = D_f\xi + R(Y)$$

is onto. This implies that $\operatorname{coker} D_f$ is covered by R. We can show that there is an induced map

$$R: \operatorname{coker} d\Pi \to \operatorname{coker} D_{b,f}$$

which is isomorphism. The proof of the proposition then follows easily. $D\Phi_{|T_bB}(Y) = R \circ di$, so we have $i \pitchfork \Pi \Leftrightarrow di \to \operatorname{coker} d\Pi$ onto $\Leftrightarrow \widetilde{R} \circ di \to \operatorname{coker} D_{b,f}$ onto.

There are few key points to be noticed here. Notice that parametric regularity is a generalization of the usual regularity. Indeed, if we consider $J_b = J$ to be constant for b in a neighborhood around b_0 then the regularity of an almost complex structure J simply says, following the diagram above, that $d\Pi$ is surjective. If we now regard J within an arbitrary family J_B , this no longer needs to be the case. It will then suffice that the cokernel of $d\Pi^l$ is covered by the variation of J in the direction of B. In subsection 2.2 we will see that, when we count rational maps, an equivalent criterion of parametric regularity will be given by the usual regularity in some suitable ambient space.

There exist a large subset of parametric regular families of almost complex structures inside $(\mathcal{A}_I, \mathcal{A}_{I,D}^c)$. This is because one can employ Sard-Smale theorem [16] and show that any map $i : (B, \partial B) \to (\mathcal{A}_I, \mathcal{A}_{I,D}^c)$ in the prop (2.5) can be perturbed to an i' such that $i' \pitchfork \Pi$.

Definition 2.6 We will say that $(J_B, \partial J_B)$ satisfies hypothesis H_2 if it is a D-parametric regular family of almost complex structures.

Compactness

We have already mentioned that H_1 is not verified for all parametric regular families of almost complex structures. Moreover, we cannot guarantee that for an arbitrary choice of a regular family, the image of the evaluation map (5) is a pseudocycle. That is because we could potentially have nonsimple elements in the compactification $\overline{\mathcal{M}}_{0,k}(M, D, (J_B, \partial J_B))$ which would yield boundary strata of high dimension.

In the situation that B = pt there are various procedures [Li-Tian ([8]), Ruan ([14]), Fukaya-Ono ([5]) to build up a theory which would provide a virtual moduli cycle, that is, an object which carries a fundamental class required for the definition of the invariants.

Roughly speaking, locally one needs to consider here all the stable holomorphic maps as well as small perturbations of these. There are then various procedures to pass to a global object with the required properties. These go through without essential changes if one considers parameter spaces with no boundary [see Leung-Bryan([4], Ruan([14])].

To make this more precise, let's first give the following

Definition 2.7 [8] A stable smooth rational map is given by a tuple $(f, \Sigma, x_1, \ldots, x_k)$ satisfying:

1) $\Sigma = \bigcup_{i=1}^{m} \Sigma_i$ is a connected rational curve with normal crossing singularities and x_1, \ldots, x_k are distinct smooth points in Σ

2) f is continuous and each restriction $f_{|\Sigma_i|}$ lifts to a smooth map from the normalization $\overline{\Sigma_i}$ to M;

3) If $f_{|\Sigma_i|}$ is constant then Σ_i contains at least three special points. Here, a special point is either a singular point or a marked point.

We will quotient the space of stable maps by the group of automorphisms of the domain; define $\overline{\mathcal{F}}_D(M, 0, k)$ to be the space of equivalence classes $[f, \Sigma, x_1, \ldots, x_k]$. The bundle \mathcal{E} can be then viewed as a generalized bundle over $\overline{\mathcal{F}}_D(M, 0, k)$ and together with the section Φ defines a generalized Fredholm bundle in the sense of Li-Tian, of index $2n + 2c_1(D) + 2k - 6$.

In our situation we need to make sure that the boundary causes no problem. The following lemma basically states that if we consider an appropriately small open neighborhood of $\overline{\mathcal{M}}_{0,k}(M, D, (J_B, \partial J_B))$ consisting of almost holomorphic maps, then its projection onto J_B stays away from ∂J_B .

Lemma 2.8 For any compact set $J_B \in \mathcal{A}_I$ such that $\partial J_B \subset \mathcal{A}_{I,D}^c \exists a \ \delta > 0$ and $\epsilon(\delta) > 0$ for which there is no stable map $[f, \Sigma, x_1, \ldots, x_k]$ such that $\bar{\partial}_J f = \nu$, when $d(J, \partial J_B) < \delta$ and $\nu \in L^p(\Lambda^{0,1} \otimes_J f^*TM)$ with $|\nu| \leq \epsilon(\delta)$.

Proof: We will prove this by assuming the opposite. Assume that we have a sequence J_i , ν_i and f_i such that $d(J_i, \partial J_B) \to 0$, $|\nu_i| = \epsilon_i \to 0$ and each f_i is a stable map in class D with the property that $\bar{\partial}_{J_i} f_i = \nu_i$. Since J_B is compact we find a convergent subsequence J_i , whose limit J_{∞} is in ∂J_B . But this would lead to a contradiction because by the Gromov compactness theorem there is a subsequence of f_i which converges to a J_{∞} stable holomorphic map in class D. This will contradict the fact that $J_{\infty} \in \partial J_B \subset \mathcal{A}_{I,D}^c$.

With this lemma the following theorems hold exactly as in [8].

Theorem 2.9 The section $\Phi : B \times \overline{\mathcal{F}}_D(M, 0, k) \to \mathcal{E}$ defined by equation 6 gives rise to a generalized Fredholm orbifold bundle with the natural orientation and of index $2c_1(D) + 2k + 2n - 6 + \dim B$.

also

Theorem 2.10 Consider two homotopic maps $i : (B, \partial B) \to (\mathcal{A}, \mathcal{A}_D^c)$ and $i' : (B', \partial B') \to (\mathcal{A}, \mathcal{A}_D^c)$ that represent the same element in $\pi_*(\mathcal{A}, \mathcal{A}_D^c)$. Suppose also that there is a deformation between ω_B and $\omega_{B'}$. If Φ' and Φ are defined by equation 6 for the pairs $(J_{B'}, \omega_{B'})$ and (J_B, ω_B) respectively, then Φ is homotopic to Φ' as generalized Fredholm orbifold bundles.

Theorem 2.9 shows that there exist an Euler class

$$e([\Phi:B\times\overline{\mathcal{F}}_D(M,0,k)\to\mathcal{E}])\in H_r(B\times\overline{\mathcal{F}}_D(M,g,k)$$

which gives the virtual moduli cycle for the moduli space of equivalence classes of stable parametric holomorphic maps in the class D, $\overline{\mathcal{M}}_{0,k}^*(M, D, (J_B, \partial J_B))$. This is independent of the choice of (J_B, ω_B) by 2.10.

We will denote by $[\mathcal{M}]^{vir}$ the virtual cycle. In order to define the invariant we consider the evaluation maps $ev_i: B \times \overline{\mathcal{F}}_D(M, 0, k) \to M$ given by

$$ev_i(b, [f, \Sigma, x_1, \dots, x_k]) = f(x_i)$$

We then can define

$$PGW_{D,0,k}^{M,(J_B,J_{\partial B})} : \bigoplus_{i=1}^{k} H^{a_i}(M,\mathbb{Q})^k \to \mathbb{Q}$$

by

$$PGW_{D,0,k}^{M,(J_B,J_{\partial B})}(\alpha_1,\ldots,\alpha_k) = ev_1^*(\alpha_1) \wedge \ldots \wedge ev_k^*(\alpha_k)[\mathcal{M}]^{viv}$$

which are zero unless

$$\sum_{i=1}^{k} a_i = 2c_1(D) + 2k + 2n - 6 + \dim B$$
(10)

We should also point out that if one changes the orientation of B we obtain the same invariant but with a negative sign.

We have the following theorem:

Theorem 2.11 i) The invariants $PGW_{D,0,k}^{M,(J_B,J_{\partial B})}$ are symplectic deformation invariants and depend only on the relative homotopy class of $(J_B, J_{\partial B})$.

ii) For a fixed choice of k, D and α_i the map $\Theta_{0,k,\alpha_1,\ldots,\alpha_k} : \pi_*(\mathcal{A}_I, \mathcal{A}_{I,D}^c) \to \mathbb{Q}$, given by $\Theta_{k,\alpha_1,\ldots,\alpha_k}([(J_B, J_{\partial B})] = PGW_{D,0,k}^{M,(J_B,J_{\partial B})}(\alpha_1,\ldots,\alpha_k)$ is a homomorphism.

Proof: Point (i) follows from theorem 2.9 and 2.10. The fact that the morphism Θ defined in point (ii) is well defined follows from theorem 2.10. To show that it is a homeomorphism, we choose $(B_1, \partial B_1)$, and $(B_2, \partial B_2)$ representing 2 elements β_1 and β_2 inside $\pi_*(\mathcal{A}, \mathcal{A}_D^c)$. We can choose them such that we can concatenate them through a deformation process in which the *J*'s in the intersection do not admit any ϵ holomorphic maps in class D. We can therefore see that the new virtual cycle corresponding to the classes $\beta_1 + \beta_2$ will be a disjoint union of the virtual neighborhoods corresponding to β_1 and β_2 . But this implies that the parametric invariants corresponding to the new class $\beta_1 + \beta_2$ are the sum of the PGW corresponding to β_1 and β_2 . Therefore Θ is a homomorpism. \Box

We should point out that in the situation that the family $(J_B, J_{\partial B})$ satisfies H_1 and H_2 , then the integer valued invariants we obtain by intersecting the image of the pseudocycle $ev_*(\overline{\mathcal{M}}_{0,k}^*(M, D, (J_B, \partial J_B)))$ with the classes $(PD(\alpha_1), \ldots, PD(\alpha_k))$ in $H_*(M)^k$ are the same as $PGW_{D,0,k}^{M,(J_B,J_{\partial B})}(\alpha_1, \ldots, \alpha_k)$ defined above. Moreover, they can be obtained by counting the number of J_b holomorphic maps in class D with k marked points which intersect generic cycles representing $(PD(\alpha_1), \ldots, PD(\alpha_k))$ in $f(z_i)$.

2.2 A criterion of parametric regularity for rational maps

Consider a family of pairs $(J_{b'}, \omega_{b'})_{b' \in B'}$ where each $J_{b'}$ is an almost complex structure tamed by the symplectic forms $\omega_{b'}$.

Since the regularity of a holomorphic map is a local statement within B and it only concerns the almost complex structure data, we claim that for each $b \in \int B'$ we can restrict our attention to a neighborhood B such that the family J_B descends from a fibration in the following sense. We say that the family (J_B, ω_B) descends from a fibration $M \to \widetilde{M} \to B$ if \widetilde{M} comes with an almost complex structure \widetilde{J} such that the restriction to each fiber $M \times b$ is an almost complex structure J_b . Moreover \widetilde{M} admits a closed two form $\widetilde{\omega}$ which restricts on each fiber $M \times b$ to a symplectic form ω_b that tames J_b . Likewise, we can choose a trivialization of the fibration such that smoothly $\widetilde{M} = B \times M$ and π is just the projection on the first factor. In the following theorem we consider the family of parameters to be a subspace of \mathbb{C}^m and we denote by z the parameters in \mathbb{C}^m . **Theorem 2.12** Let $(J_z, \omega_z)_{z \in \mathbb{C}^m}$ be a family on M descending from the symplectic fibration $(\widetilde{M}, \widetilde{J}, \widetilde{\omega})$

Suppose that $f: \Sigma \longrightarrow M$ is a J_0 holomorphic map and consider the composite map

$$\widetilde{f}=i\circ f, \widetilde{f}:\Sigma\longrightarrow M\times 0\subset \widetilde{M}$$

which is \tilde{J} -holomorphic. If \tilde{f} is regular then f is (J_z) parametric regular. Moreover, if $\Sigma = S^2$ then the reverse statement holds.

The proof of the theorem occupies the rest of the section. Let $T_{|_{\pi^{-1}(0)}}\widetilde{M}$ be the tangent space along the preimage of $0 \in \mathbb{C}^m$. We will denote by H the subbundle of $T_{|_{\pi^{-1}(0)}}\widetilde{M}$ which is $\widetilde{\omega}$ orthogonal to the fiber $\{0\} \times M$. We would like H to coincide with the horizontal space of $T\widetilde{M}$ with respect to the trivialization π and to be \widetilde{J} invariant. This can be arranged by deforming the form $\widetilde{\omega}$ so that near the zero fiber $\{0\} \times M$ it is given by

$$\widetilde{\omega} = \omega_0 + \pi^*(\sigma_{base}),$$

where σ_{base} is a standard symplectic two form on the holomorphic base B. Throughout this deformation process \tilde{J} is still $\tilde{\omega}$ tamed.

Let g_0 be a metric on M_0 and ∇ be the Levi-Civita connection on M associated with it. ∇^{st} will be the standard Levi-Civita connection on \mathbb{C}^m . We will denote from now on $\widetilde{\nabla} = \nabla \times \nabla^{st}$, the product connection on $\widetilde{M} \simeq \mathbb{C}^m \times M$. The regularity of $\widetilde{f} : \Sigma \longrightarrow \widetilde{M}$ is by definition, equivalent to the fact that $D_{\widetilde{f}}$ is surjective, where $D_{\widetilde{f}}$ is the linearization of $\overline{\partial}$,

$$D_{\widetilde{f}}: C^{\infty}(\widetilde{f}^*T\widetilde{M}) \longrightarrow \Omega^{0,1}_{\widetilde{J}}(\Sigma, \widetilde{f}^*T\widetilde{M}).$$

Using the connection $\widetilde{\nabla}$ we will derive formulas for $D_{\widetilde{f}}$ and express them in terms of the linearization $D\Phi$.

Since $\widetilde{M} \simeq \mathbb{C}^m \times M$ and $\operatorname{im} \widetilde{f} \subset \{0\} \times M$, we have the following relations:

$$\widetilde{f}^*\left(T\widetilde{M}\right) = \widetilde{f}^*\left(T\widetilde{M}_{\pi^{-1}(0)}\right) = \widetilde{f}^*(H \oplus TM) = triv \oplus f^*(TM)$$

where by triv we denote the trivial *m*-dimensional complex bundle over Σ . This gives

$$C^{\infty}(f^*TM) \simeq C^{\infty}(triv) \oplus C^{\infty}(f^*TM)$$
(12)

Given that each fiber is \widetilde{J} invariant, and that H is \widetilde{J} invariant along $\pi^{-1}(0)$, we obtain

$$\Omega^{0,1}_{\widetilde{J}}(\Sigma, \widetilde{f}^*T\widetilde{M}) \simeq \Omega^{0,1}_J(\Sigma, f^*TM) \oplus \Omega^{0,1}_{\widetilde{J}}(\Sigma, H)$$
(13)

From (12) and (13) we obtain

$$D_{\widetilde{f}}: C^{\infty}(triv) \oplus C^{\infty}(f^{*}TM) \xrightarrow{} \Omega^{0,1}_{J}(\Sigma, f^{*}TM) \oplus \Omega^{0,1}_{\widetilde{J}}(\Sigma, H)$$

and by considering the appropriate restrictions we obtain the following operators

$$\begin{array}{rcl} D_{1,vert}: & C^{\infty}(triv) & \longrightarrow & \Omega_{J}^{0,1}\left(\Sigma,f^{*}TM\right) \\ D_{1,hor}: & C^{\infty}(triv) & \longrightarrow & \Omega_{\widetilde{J}}^{0,1}\left(\Sigma,H\right) \\ D_{2,vert}: & C^{\infty}\left(f^{*}TM\right) & \longrightarrow & \Omega_{J}^{0,1}(\Sigma,f^{*}TM) \\ D_{2,hor}: & C^{\infty}\left(f^{*}TM\right) & \longrightarrow & \Omega_{\widetilde{J}}^{0,1}\left(\Sigma,H\right) \end{array}$$

We will sometimes use $D_k = (D_{k,vert}, D_{k,hor}), \ k = 1, 2.$

To compute the formulas for these operators we will use the following general method (see [1]). Consider $\xi \in C^{\infty}(\Sigma, \tilde{f}^*T\widetilde{M})$ and $\widetilde{F}_{\xi} : [0,1] \times \Sigma \longrightarrow \widetilde{M}$ given by $\widetilde{F}_{\xi}(t,x) = \exp_{\tilde{f}(x)}^{\widetilde{\Sigma}}(t\xi(x))$, for ξ sufficiently small. Let $s : \Sigma \longrightarrow T\Sigma$ be a section and \widetilde{s} its lift to $T([0,1] \times \Sigma)$. We denote $\frac{\partial}{\partial t}$ the vector field in $T([0,1] \times \Sigma)$ corresponding to the parameter in [0,1]. Define $\tilde{f}_t(x) := \widetilde{F}_{\xi}(t,x)$. For any $x \in \Sigma$, define the path $\widetilde{\gamma}_x^{\xi} : [0,1] \longrightarrow \widetilde{M}$ given by $\widetilde{\gamma}_x^{\xi}(t) = \widetilde{F}_{\xi}(t,x)$, the image under \widetilde{F}_{ξ} of $[0,1] \times x$ in \widetilde{M} . By the definition of $\widetilde{F}_{\xi}, \widetilde{\gamma}_x^{\xi}$ is a geodesic path in \widetilde{M} relative to the connection $\widetilde{\nabla}$. Denote by $\tau_{t,x}^{\xi} : T_{\gamma_x(t)}\widetilde{M} \longrightarrow T_{\gamma_x(0)}\widetilde{M}$ the parallel transport in \widetilde{M} along the curve $\gamma_x := \widetilde{\gamma}_x^{\xi}$. To compute $D_{\tilde{f}}(\xi)(s)$ in general, one needs to consider the expression $\frac{1}{2}\tau_{t,x}^{\xi}(d\tilde{f}_t(s) + \widetilde{J}d\tilde{f}_t(js))$ and take its derivative with respect to t at t = 0 i.e.

$$D_{\widetilde{f}}(\xi)(s) = \frac{1}{2} \frac{\partial}{\partial t} \left(\tau_{t,x}^{\xi}(d\widetilde{f}_t(s) + \widetilde{J}d\widetilde{f}_t(js))) \right)_{|_{t=0}}$$
(14)

We define *Const* to be the subspace of $C^{\infty}(triv)$ made out of constant sections. For the proof of the theorem, we are particularly interested in computing $D_{1,hor}$ and the restriction of $D_{1,vert}$ to *Const*.

In order to simplify the notation, we denote by x the coordinate on Σ and write the points in $\mathbb{C}^m \times M$ as (z_1, \ldots, z_m, y) where $z_1 = w_1 + iv_1$ and so on. For simplicity we denote the vector field in *Const* by $\frac{\partial}{\partial w_k} = \partial_{w_k}$ and so on. Since we are going to work with an arbitrary choice of w_k and v_k we will refer to them simply as ∂_w , unless we need to be more specific.

Lemma 2.13 The following relations hold:

i) $D_{2,hor} = 0$ ii) $D_{2,vert} = D_f$ iii) $D_{1,hor}(\xi) = \bar{\partial}_{\mathbb{C}^m}(\xi), \forall \xi \in C^{\infty}(triv), where \bar{\partial}_{\mathbb{C}^m} is the delbar operator in <math>\mathbb{C}^m$. v) $(D_{1,vert})(\partial_z)(s) = \frac{1}{2} \frac{\partial}{\partial z} (J(z))_{|z=0} (df(js) \text{ for } \partial_z a \text{ typical vector field in Const } \subset C^{\infty}(triv).$

Proof: Since $\tilde{f} = f \circ i \subset \{0\} \times M$ we can naturally view any $\xi \in C^{\infty}(f^*TM)$ as an element in $C^{\infty}(\tilde{f}^*T\widetilde{M})$ with values in the vertical direction tangent to $\{0\} \times M$. We have that $\tilde{F}_{\xi}(t,x) = \exp_{\tilde{f}(x)}^{\tilde{\nabla}}(t\xi) = \exp_{f(x)}^{\nabla}(t\xi)$, with $\operatorname{im} \tilde{F} \subset \{0\} \times M$. This implies that the $d\tilde{f}_t(s)$ are also vertical vector fields supported in $\{0\} \times M$ and, since \tilde{J} keeps $T(\{0\} \times M)$ invariant, we have as well that the $\tilde{J}d\tilde{f}_t(js)$ are vertical vector fields in $\{0\} \times M$. Similarly, $\tilde{F}_{\xi}^* \frac{\partial}{\partial t}$ is a

vertical section in $T\widetilde{M}$ supported in $\{0\} \times M$ and parallel transport along $\widetilde{f}(x)$ with respect to $\widetilde{\nabla}$ is the same as parallel transport with respect to ∇ .

A direct application of (14) is that

$$\left(D_{\tilde{f}}\xi\right)(s) = \frac{1}{2}\frac{\partial}{\partial t} \left(\tau_{t,x}^{\xi}d\tilde{f}_{t}(s) + \tau_{t,x}^{\xi}\tilde{J}df_{t}(js)\right)_{|_{t=0}} = \left(D_{f}\xi\right)(s),$$

which proves (i). Relation (ii) follows immediately from the formula above, taking into account that $D_{\tilde{f}}\xi = D_{2,vert}(\xi)$, and that $\mathrm{im}D_{\tilde{f}}|_{C^{\infty}f*TM} \subset \Omega_{J}^{0,1}(\Sigma, f^{*}TM)$. For the proofs of (iii) and (v) we now consider $\xi \in C^{\infty}(triv)$. We can assume $\xi = \phi(x)\partial_w$

For the proofs of (iii) and (v) we now consider $\xi \in C^{\infty}(triv)$. We can assume $\xi = \phi(x)\partial_w$ where $\phi : \Sigma \leftarrow \mathbb{C}^m$. In this situation, $\widetilde{F}_{\xi}(t,x) = \exp_{\widetilde{f}(x)}^{\widetilde{\nabla}}(t\partial_w) = (\phi(x)t, 0, \dots, 0, f(x))$. It then follows that the paths γ_x are straight lines in $\mathbb{C}^n \times f(x) \subset \widetilde{M}$ and therefore the parallel transport along $\gamma_x, \tau_{t,x} : T_{(t,f(x))}\widetilde{M} \longrightarrow T_{0,f(x)})\widetilde{M}$ is the identity. We are also going to consider the coordinates $x \in \Sigma$ of the type $x = x_1 + ix_2$, and do our computations for $s = \partial_{x_1}$.

If $\widetilde{J}(t)$ is the almost complex structure at $\widetilde{\gamma}_x^{\xi}(t)$ then $\widetilde{J}(t)$ has the form $\begin{pmatrix} A_t & 0 \\ B_t & J_t \end{pmatrix}$ with respect to the product structure $\mathbb{C}^m \times M$. Moreover along $\pi^{-1}(0)$ we have $\widetilde{J}(0) = \begin{pmatrix} J_{\mathbb{C}^m} & 0 \\ 0 & J_t \end{pmatrix}$. Therefore $\frac{\partial}{\partial t}\widetilde{J}(t)$ preserves the fibers, the same as $\widetilde{J}(t)$ does. Moreover, along $\{0\} \times M$, $\widetilde{J}(0)$ preserves the splitting into TM and H. As we have seen parallel transport along $\widetilde{\gamma}_x^{\xi}(t)$ is just the identity.

Considering local coordinates on $\Sigma x = x_1 + ix_2$ and taking $s = \partial_{x_1}$, we have:

$$D_{1,hor}(\phi\partial_w)(\partial_{x_1}) = \frac{1}{2}proj_H \frac{\partial}{\partial t} \left(\tau_{t,x}^{\xi} d\widetilde{f}_t(\partial_{x_1}) + \frac{1}{2} \tau_{t,x}^{\xi} \widetilde{J} d\widetilde{f}_t(j\partial_{x_1}) \right)_{|_{t=0}}$$

$$= \frac{1}{2}proj_H \frac{\partial}{\partial t} \left(d\widetilde{f}_t(\partial_{x_1}) + \frac{1}{2} \widetilde{J} d\widetilde{f}_t(\partial_{x_2}) \right)_{|_{t=0}}$$

$$= \frac{1}{2} \frac{\partial}{\partial t} \left(\partial_{x_1}(\phi(x))t, 0, \dots, 0 \right)_{|_{t=0}} + \frac{1}{2}proj_H \frac{\partial}{\partial t} \left(\widetilde{J}_t \right)_{|_{t=0}} df(\partial_{x_2}) + \frac{1}{2}proj_H \widetilde{J}_0 \frac{\partial}{\partial t} \left(\partial_{x_2}(\phi(x))t, 0, \dots, 0, df(x) \right)_{|_{t=0}}$$

where, as mentioned before, $\phi: \Sigma \to \mathbb{C}^m$. But here the middle term vanishes because $df(\partial_{x_2})$ is a vertical vector and $\frac{\partial}{\partial t}\widetilde{J}$ preserves fibers so we get that $\frac{\partial}{\partial t}\left(\widetilde{J}_t\right)_{|t=0} df(\partial_{x_2})$ is also a vertical vector. Then

$$D_{1,hor}(\phi\partial_w)(\partial_{x_1}) = \frac{1}{2}\partial_{x_1}\phi(x) + \frac{1}{2}J_{\mathbb{C}^m}(\partial_{x_2})\phi(x)$$
(15)

For the last expression we have to use that along $\pi^{-1}(0)$, \widetilde{J}_0 preserves the horizontal space H, so $proj_H \circ \widetilde{J}_0 = \widetilde{J}_{\mathbb{C}^m} \circ proj_H$. Therefore, the conclusion follows that $D_{1,hor} = \overline{\partial}_{\mathbb{C}^m}$.

To prove point (v) of the theorem we need to consider now $\xi = \partial_w$, that is $\xi \in Const$. Under this assumption we have $\tau_{t,x}^{\partial_w} d\tilde{f}_t = df_0$. Thus

$$\frac{\partial}{\partial t}\tau_{t,x}^{\partial_w}d\widetilde{f}_t(s) = 0$$

As before, s is a just a section in $T\Sigma$. We then have

$$D_{1,vert}(\partial_{w})(s) = \frac{1}{2}proj_{V}\frac{\partial}{\partial t} \left(\tau_{t,x}^{\partial w}d\widetilde{f}_{t}(s) + \frac{1}{2}\tau_{t,x}^{\partial w}\widetilde{J}d\widetilde{f}_{t}(js)\right)_{|_{t=0}}$$

$$= \frac{1}{2}proj_{V}\frac{\partial}{\partial t} \left(\tau_{t,x}^{\partial w}d\widetilde{f}_{t}(s)\right)_{|_{t=0}} + \frac{1}{2}proj_{V}\frac{\partial}{\partial t} \left(\tau_{t,x}^{\partial w}\widetilde{J}(\tau_{t,x}^{\partial w})^{-1}\right)_{|_{t=0}} \cdot df(js)$$

$$+ \frac{1}{2}proj_{V}\widetilde{J}_{0} \left(\frac{\partial}{\partial t}\tau_{t,x}^{\partial w}d\widetilde{f}_{t}(js)\right)_{|_{t=0}} = \frac{1}{2}proj_{V} \left(\widetilde{\nabla}_{\partial_{w}}\widetilde{J}\right) df(js)$$

where we denote by $proj_V$ the projection onto the fibers. Recall that $\frac{\partial}{\partial t}\widetilde{J}$ takes vertical vector fields into vertical vector fields. Therefore

$$\frac{1}{2} proj_V \widetilde{\nabla}_{\partial_w} \widetilde{J} df(js) = \frac{1}{2} \frac{\partial}{\partial w} (J(z)) (df(js))$$

precisely because df(js) is a vertical vector field and because the covariant derivative along horizontal vector fields was chosen to be the standard connection in \mathbb{C}^m . Applying the same reasoning for $i\partial_v$ we see that

$$(D_{1,vert})(\partial_z)(s) = \frac{1}{2} \frac{\partial}{\partial z} (J(z))_{|z=0} (df(js))$$

It is worth to point out that $\frac{\partial}{\partial z}(J(z)|_{z=0} = d\psi_0^*(\frac{\partial}{\partial z}).$

Proof of the theorem: Implication " \Rightarrow " Using lemma 2.13, point (v) we get the commutativity of the following diagram

$$T_{0}\mathbb{C}^{m} \xrightarrow{d\psi} T_{J}\mathcal{A}_{I}$$

$$\downarrow^{i} \qquad \downarrow^{R}$$

$$Const \xrightarrow{D_{1,vert}} \Omega_{J}^{0,1}(\Sigma, f^{*}TM),$$

$$(16)$$

where $i: T_0 \mathbb{C}^n \to Const \subset C^{\infty}(triv)$ is the natural identification map and ψ is the morphism from the parameter space to the space of almost complex structures. R is, as mentioned before, given by $R(Y) = \frac{1}{2}Y \circ df \circ j$.

Since $D_{\tilde{f}}$ is surjective by hypothesis of, this means that $D_1 \oplus D_2$ is surjective. We therefore have, by lemma (2.13) (i),(ii),

$$D_1 = (D_{1,vert}, D_{1,hor}) : C^{\infty}(triv) \longrightarrow \operatorname{coker} D_f \oplus \Omega^{0,1}_{\widetilde{J}}(\Sigma, H)$$
(17)

is surjective. Since the kernel of the $\bar{\partial}_{\mathbb{C}^m}$ operator on \mathbb{C}^m consists precisely of constant sections, lemma 2.13 (iii) implies that $D_{1,hor}^{-1}(0) = Const$. Therefore we have that the operator

 $(D_{1,vert})_{|_{Const}}: Const \longrightarrow coker D_f$ is surjective. But this will imply that

 $D_{1,vert}|_{Const} \circ i: T_0\mathbb{C}^m \longrightarrow \operatorname{coker} D_f$. But as we saw in the proof of 2.5, R induces an isomorphism $\widetilde{R}: \widetilde{\operatorname{coker} d\Pi} \longrightarrow \operatorname{coker} D_2$ and moreover the diagram 16 will be still commutative if we restrict $d\psi$ and $D_{1,vert}$ to $\operatorname{coker} d\Pi$ and $\operatorname{coker} D_2$ respectively. Therefore

 $d\psi: T_0\mathbb{C}^n \longrightarrow \operatorname{coker} d\Pi$ is surjective. By proposition 2.5, this yields exactly the parametric regularity.

For the inverse implication, we notice that since $D_{1,hor}$ is $\partial_{\mathbb{C}^m}$, it will cover the space $\Omega^{0,1}_{\tilde{j}}(\Sigma, H)$ when $\Sigma = S^2$. By hypothesis we have that $d\psi: T_0\mathbb{C}^n \longrightarrow \operatorname{coker} d\Pi$. is surjective and the above observation implies that

$$D_1 = (D_{1,vert}, D_{1,hor}) : C^{\infty}(triv) \longrightarrow \operatorname{coker} D_f \oplus \Omega^{0,1}_{\widetilde{J}}(\Sigma, H)$$
(18)

is also surjective. Therefore $D_{\widetilde{f}}$ is a surjective operator.

3 Resolutions of singularities and relative PGW

3.1 Quotient singularities

In this subsection we will give an overview of work of Kronheimer [7] and Abreu-McDuff [3] on how to construct special families of almost complex structures arising from the study of the total spaces of deformations for some quotient singularities. In the end of the section we will explain how these families serve our purpose of counting nontrivial PGW. The local picture is as follows(see Kronheimer [7]):

We consider the particular type of Hirzebruch-Jung singularity $Y_0 = \mathbb{C}^2/C_{2\ell}$, given by the diagonal action by scalars of $C_{2\ell}$ on \mathbb{C}^2 , where $C_{2\ell}$ is the cyclic group of order 2ℓ . This admits a resolution $\sigma_0: \widetilde{Y}_0 \to Y_0$ where \widetilde{Y}_0 is the total space of the line bundle of degree -2ℓ over $\mathbb{C}P^1$. The exceptional curve of the resolution, we will call it E, is a curve of selfintersection -2ℓ and is the zero section of \widetilde{Y}_0 . This resolution admits a $2\ell - 1$ complex dimensional parameter family of deformations , $\widetilde{Y}_t, t \in \mathbb{C}^{2\ell-1}$. With the exception of the case $\ell = 2$ the total space $\widetilde{Y} = \bigcup \widetilde{Y}_t$ of the family of deformations is the total space of the vector bundle $\mathcal{O}(-1)^{2\ell}$. More precisely, we consider the exact sequence of bundles

$$\mathcal{O}(-2\ell) \rightarrow \mathcal{O}(-1)^{2\ell} \xrightarrow{r} \mathcal{O}^{2\ell-1}$$
 (19)

where r is given by evaluating at $2\ell - 1$ generic sections of the dual of \widetilde{Y} , $\widetilde{Y}^* = \mathcal{O}(1)^{2\ell}$. Since holomorphically $\mathcal{O}^{2\ell-1}$ is trivial, we can project it to its fiber $\mathbb{C}^{2\ell-1}$ and hence we obtain a submersion $\widetilde{q} : \mathcal{O}(-1)^{2\ell} \to \mathbb{C}^{2\ell-1}$ with $\widetilde{Y}_t = \widetilde{q}^{-1}(t)$. Also it can be seen that \widetilde{Y} is smoothly isomorphic with $\widetilde{Y}_0 \times \mathbb{C}^{2\ell-1}$ and a choice of trivialization provides an isomorphism

$$\theta: \widetilde{Y} \stackrel{C^{\infty}}{\cong} \widetilde{Y}_0 \times \mathbb{C}^{2\ell-1}$$
(20)

We consider now a 4ℓ -dimensional basis of sections in the dual \widetilde{Y}^* . Here the space of holomorphic sections is given by $\bigoplus_{i=1}^{2\ell} H^0(CP^1, \mathcal{O}(1)) \cong (\mathbb{C}^2)^{2\ell}$. Denote by Y the subspace of $(\mathbb{C}^2)^{2\ell}$ consisting of 2ℓ -tuples of vector in \mathbb{C}^2 which span either zero or a line. By evaluating all the 4ℓ section we obtain a map

$$\sigma: \widetilde{Y} \to Y \subset \mathbb{C}^{4\ell}$$

which contracts E to a point $\gamma_0 = \sigma(E)$. Moreover, γ_0 is the only singular point of Y. and the morphism is one to one outside E. We also define a map $q: Y \to \mathbb{C}^{2\ell-1}$ by evaluating at the original $2\ell - 1$ generic sections. The following diagram commutes



We can obtain a 2-form τ on Y by pulling back a Kähler form from $\mathbb{C}^{4\ell}$. Via σ^* this can be seen as a two form on \widetilde{Y} which restricts to a Kähler form τ_t on each fiber \widetilde{Y}_t if $t \neq 0$ but degenerates along E when t = 0. If we further push forward through θ , these forms can be seen as a family of forms on the manifold Y_0 .

As in [3], we can choose an appropriate compactification of the local picture as follows : Let $B^{4\ell-2}$ be the unit ball in $\mathbb{C}^{2\ell-1}$. We have a family $(\overline{Y}_t, J_t^{\ell}, \tau_t)_{t \in B^{4\ell-2}}$, where each $(\overline{Y}_t, J_t^{\ell}, \tau_t^{\ell}), t \neq 0$ is a Kähler manifold diffeomorphic with $S^2 \times S^2$, and, $(\overline{Y}_0, J_0^{\ell})$ is a complex manifold, also diffeomorphic with $S^2 \times S^2$ and τ_0 degenerates along $A - \ell F$. The total space of the family has the following properties:

a) The space $\overline{Y} = \bigcup_{t \in B^{4\ell-2}} \overline{Y}_t$ is smoothly diffeomorphic with $S^2 \times S^2 \times B^{4\ell-2}$. Moreover \overline{Y} is a complex manifold with a complex structure J^{ℓ} which restricts to each fiber \overline{Y}_t to the complex structure J_t^{ℓ} . Also, \overline{Y} has a closed (1,1) form τ which is satisfies all the properties of a Kähler form outside the zero fiber and restricts at each fiber to the forms, τ_t .

b) The form τ restricted to $\overline{Y_0}$ degenerates along the exceptional curve $A - \ell F$

Since the forms τ_t are obtained by restricting the closed form τ to fibers it is immediate that they are all in the same cohomology class. Moreover, since $(\tau_0)_{|A-\ell F} = 0$ we obtain that $\forall t \in B^{4\ell-2}, [\tau_t^\ell] = [\omega_\ell].$

From (a) we see that there is a holomorphic projection $\pi : \overline{Y} \to S^2 \times B^{4\ell-2}$. This is because every \overline{Y}_t is a ruled surface therefore it fibers over S^2 . If we denote be α the area form on S^2 we can construct a two form

$$\tau^{\lambda} = \tau + (\lambda - \ell)\pi^*(\alpha)$$

For $\lambda > \ell$ these forms are Kähler forms and moreover they restrict to each \overline{Y}_t to symplectic forms in the class $[\omega_{\lambda}]$. This proves that any J_t^{ℓ} is tamed by a form isotopic with ω_{λ} as long as $\lambda > \ell$. We now follow a similar procedure to construct a family of symplectic forms $\omega_t, t \in B^{4\ell-2}$ such that each ω_t tames J_t^{ℓ} . We will now change the forms τ_t by perturbing with a positive factor of $\pi^*(\alpha)$ only around t = 0 and smoothen with a cut-off function. By this procedure we obtain symplectic forms ω_t with variable cohomology classes.

In conclusion, we have pairs $(S^2 \times S^2, J_t^{\ell}, \omega_t)_{t \in B^{4\ell-2}}$ where ω_t is a symplectic structure on $S^2 \times S^2$ that tames J_t^{ℓ} . Moreover $[\omega_t]_{t \in S^{4\ell-3}} = [\omega_{\ell}]$. This gives a family of almost complex structures which we denote by abuse of notation B_{ℓ} such that $(B_{\ell}, \partial B_{\ell}) \in (\mathcal{A}_{[\ell,\ell+\epsilon]}, A_{\ell})$ for any $\epsilon > 0$. More importantly, only J_0^{ℓ} admits the exceptional curve in the class $A - \ell F$.

We will then obtain a family of almost complex structures on $(S^2 \times S^2 \times X)$ by taking $(J_t^{\ell} \times J_{st})$, and by abuse of notation, we will call this family also B_{ℓ} . Therefore we just produced on $(S^2 \times S^2 \times X)$ pairs $(B_\ell, \partial B_\ell) \subset (\mathcal{A}_{[\ell,\ell+\epsilon]}, A_\ell)$, with $\epsilon > 0$ that represents an element β_{ℓ} in $\pi_*(\mathcal{A}_{[\ell,\ell+\epsilon]}, A_{\ell})$. Moreover each $B_{\ell} \subset \mathcal{A}_{\ell+\epsilon}$ for any small $\epsilon > 0$.

From the choice of the J's we know that the only structure which admits $A - \ell F$ curves is $J_0 \times J_{st}$.

3.2 The computation of PGW

Here we prove that (H_1) and (H_2) are satisfied for the family $(B_\ell, \partial B_\ell)$, and therefore the invariant is integer valued and can be obtained by counting holomorphic maps intersecting generic cycles of appropriate dimension.

Claim 1. The family $(B_{\ell}, \partial B_{\ell})$ satisfies (H_1) .

Proof of claim 1: This is proved by inspection. Only $J_0^{\ell} \times J_{st}$ admits $A - \ell F$ stable maps, and the only maps in this class are copies of the imbedded map E in any fiber $S^2 \times S^2 \times pt$. Hence there are no decomposable J_b holomorphic maps. We should point out that for other almost complex structures J on $S^2 \times S^2 \times X$ one could have decomposable J-holomorphic maps in the class $A - \ell F$.

Claim 2. The family $(B_{\ell}, \partial B_{\ell})$ satisfies H_2 .

Proof of claim 2: From the sequence (19) we have that the map E, which is \tilde{J}^{ℓ} -holomorphic has the normal bundle $\mathcal{O}(-1)^{2\ell}$ and therefore we can apply lemma 3.5.1 pg 38 in [12] for the integrable almost complex structure \tilde{J} . If follows that E is \tilde{J}^{ℓ} regular inside \overline{Y} . If we consider now $\overline{Y} \times X$ and $\tilde{J}^{\ell} \times J_{st}$, the curve E lies entirely inside \overline{Y} and therefore the normal bundle inside $\overline{Y} \times X$ is $\mathcal{O}(-1)^{2\ell} \times trivial$, and therefore the curve is is $\tilde{J}^{\ell} \times J_{st}$ regular. This splitting and therefore regularity use the fact that the map E is of genus zero. Theorem 2.12 implies parametric regularity and therefore (H_2) holds.

We can therefore conclude that the invariants

$$PGW_{A-\ell F,0,k}^{S^2 \times S^2 \times X, (B_\ell, \partial B_\ell)} : \bigoplus_{i=1}^k H^{a_i} (S^2 \times S^2 \times X, \mathbb{Q})^k \to \mathbb{Z}$$

are integer valued. We have two situations. First, if X = pt then the moduli space of unparametrized curves has dimension 0 so we would count isolated curves. This follows immediately from the fact that $c_1(A - \ell F) = -4\ell + 2$ (adjunction formula) and therefore

$$\dim \mathcal{M}_{0,0}^*(S^2 \times S^2, A - \ell F, (B_\ell, \partial B_\ell)) = 2 \times 2 + 2c_1(A - \ell F) + \dim B^\ell - 6$$
$$= 4 - 4\ell + 4 + 4\ell - 2 - 6 = 0.$$

Moreover, the invariant $PGW_{A-\ell F,0,0}^{S^2 \times S^2 \times X, (B_{\ell},\partial B_{\ell})}([\text{pt}]) = 1$ because it counts E, the only $J_{b_{\ell}}$ map in the class $A - \ell F$.

In the situation that dim X = n > 0, we will count maps with one marked point. $c_1(A - \ell F)$ will be the same since the holomorphic maps in class $A - \ell F$ will have the image entirely in the fibers $S^2 \times S^2 \times pt$. we therefore have

$$\dim \mathcal{M}_{0,1}^*(S^2 \times S^2 \times X, A - \ell F, (B_\ell, \partial B_\ell)) = 2 \times (2+n) + 2c_1(A - \ell F) + \dim B^\ell - 6 + 2$$
$$= 2n + 2$$

We will consider a cycle in the homology class F which will lie in a fiber $S^2 \times S^2 \times pt$ inside $S^2 \times S^2 \times X$. It easily follows that the only J_{b_ℓ} holomorphic map with one marked point which intersect this cycle transversely is a copy of the map E inside the fiber $S^2 \times S^2 \times pt$. We obtain that

$$PGW_{A-\ell F,0,1}^{S^2 \times S^2 \times X, (B_\ell, \partial B_\ell)}(PD([F]) = 1.$$

Applying theorem 2.11 we obtain that the morphism Θ in both situations is nontrivial and therefore there is a nonzero element

$$\beta_{\ell} \in \pi_{4\ell-2}((\mathcal{A}_{[\ell,\ell+\epsilon]}, A_{\ell}) \text{ for all } \epsilon > 0$$
(22)

that is represented by the cycle $(B_{\ell}, \partial B_{\ell}) \subset (\mathcal{A}_{\ell+\epsilon}, \mathcal{A}_{\ell+\epsilon,D}^c)$.

4 Almost complex structures and symplectomorphism groups

4.1 Almost complex structures and symplectomorphisms; deformations along compact subsets

In this subsection we will give a quick overview of what can be said about the behavior of spaces of almost complex structures and about the symplectomorphisms groups as the symplectic form varies.

We will restrict our attention to variations of the symplectic form ω along a line L inside \mathcal{K} , parametrized by the real parameter λ . If L happens to be a ray $\lambda \omega, \lambda > 0$ then G_{λ} is independent of λ . It will therefore make sense to consider $L \neq ray$.

If $M = S^2 \times S^2$, a great deal is known about the structure of \mathcal{A}_{λ} see [10]. For example, one can establish that there is a direct inclusion $\mathcal{A}_{\lambda} \subset \mathcal{A}_{\lambda'}$, for $\lambda < \lambda'$. Moreover, the homotopy type of the spaces \mathcal{A}_{λ} changes only as λ strictly passes an integer ℓ .

None of this is known to hold for M an arbitrary symplectic manifold. Nevertheless, as a consequence of the fact that taming is an open condition, we are able to establish the following lemma, which we use in the proof of the theorem 2.12

Lemma 4.1 i) Let K' to be an arbitrary compact subset of \mathcal{A}_{λ} . Then there is an $\epsilon_{K'} > 0$ such that K' is contained in $\mathcal{A}_{\lambda+\epsilon}$, for $|\epsilon| < \epsilon_{K'}$.

ii) Consider K an arbitrary compact set in G_{λ} . Then there is an $\epsilon_K > 0$ and a map $h : [-\epsilon_K, \epsilon_K] \times K \to \mathcal{G}_{|L}$ such that the following diagram commutes

$$h: [-\epsilon_K, \epsilon_K] \times K \longrightarrow \mathcal{G}_{|L}$$

$$\downarrow^{pr_1} \qquad \qquad \downarrow^{pr_2}$$

$$[-\epsilon_K, \epsilon_K] \xrightarrow{incl} (-\infty, \infty).$$

$$(23)$$

Moreover, for any two such maps h and h' which coincide on $0 \times K$, there exist, for an ϵ' small enough, a homotopy $H: [0,1] \times [-\epsilon', \epsilon'] \times K \to \mathcal{G}_{|L}$ between them which satisfies

$$H: [0,1] \times [-\epsilon',\epsilon'] \times K \longrightarrow \mathcal{G}_{|L}$$

$$\downarrow^{pr_1} \qquad \qquad \downarrow^{pr_2}$$

$$[-\epsilon',\epsilon'] \xrightarrow{incl} (-\infty,\infty).$$

$$(24)$$

Proof: Subpoint (i) is an immediate consequence of the openness of the taming condition.

For the proof of (ii), let's first notice that, since the symplectic condition is an open condition, there is a convex neighborhood U of ω_{λ} inside the space of 2-forms such that any closed ω' in U is still symplectic. Moreover for any $g_{k_0} \in K \subset G_{\lambda}$ and any symplectomorphism $g_k \in K$ which is sufficiently close to g_{k_0} we can choose $\epsilon(k_0) > 0$ small enough, such that $g_k^* \omega_{\lambda+\epsilon}$ is still inside U, for all $0 \leq \epsilon < \epsilon(k_0)$. Since K is compact we can do this process finitely many times such that in the end we have an $\epsilon(K) > 0$ such that for any $g_k \in K$ $g_k^* \omega_{\lambda+\epsilon} \in U$, for all $0 \leq \epsilon < \epsilon(K)$. We will construct the elements $h(\epsilon, k)$ as follows. For $t \in [0, 1]$ the forms

$$\omega_{k,\lambda+\epsilon}^t := tg_k^*\omega_{\lambda+\epsilon} + (1-t)\omega_{\lambda+\epsilon}$$

are symplectic since both $g_k^* \omega_{\lambda+\epsilon}$ and $\omega_{\lambda+\epsilon}$ are inside the convex set U. We now apply Moser's argument and obtain a family of diffeomorphisms $\xi_{k,\lambda+\epsilon,t}$ with the property that $\xi_{k,\lambda+\epsilon,t}^* \omega_{k,\lambda+\epsilon}^t = \omega_{\lambda+\epsilon}$. We will now define $h(\epsilon,k) := g_k \circ \xi_{k,\lambda+\epsilon,1}$. Then h has the required properties.

For an arbitrary $h: K \times [-\epsilon, \epsilon]$ satisfying (23) we take a homotopy $F: [0,1] \times [-\epsilon, \epsilon] \times K \to \mathbb{R} \times \text{Diff} M$ given by $F(t, \epsilon, k) := (\epsilon, h(t\epsilon, k))$. This gives a homotopy between h and $h_0: [-\epsilon, \epsilon] \times K \to \mathbb{R} \times \text{Diff} M$, where $h_0(\epsilon', k) = h(0, k)$. We similarly obtain a homotopy F' between h' and h_0 , where h' also satisfies (23). By concatenating one homotopy with the opposite of the other we obtain a homotopy between h and h' which we call $G: [0, 1] \times [-\epsilon_1, \epsilon_1] \times K \to \mathbb{R} \times \text{Diff} M$. Denote by $g_{s,\epsilon,k} := G(s,\epsilon,k)$. We will now follow the same procedure as before. Namely, we restrict to a short interval $[-\epsilon', \epsilon']$ such that, if we call

$$\omega_{s,k,\lambda+\epsilon}^t := tg_{s,\epsilon,k}^*\omega_{\lambda+\epsilon} + (1-t)\omega_{\lambda+\epsilon}$$

then these are symplectic, $\forall 0 \leq |\epsilon| < \epsilon'$ and $\forall t, s \in [0, 1]$. This is possible because $\omega_{s,k,\lambda}^t = \omega_{\lambda}$. Applying Moser's argument again we obtain diffeomorphisms $\xi_{s,k,\lambda+\epsilon,t}$ with the property that $\xi_{s,k,\lambda+\epsilon,t}^* \omega_{s,k,\lambda+\epsilon}^t = \omega_{\lambda+\epsilon}$. We will now define $H(s,\epsilon,k) := g_{s,\epsilon,k} \circ \xi_{s,k,\lambda+\epsilon,1}$. Then H has the required properties.

Definition 4.2 Let $\rho: B \to G_{\lambda}$ be a cycle in G_{λ} . An extension ρ^{ϵ} of ρ is a smooth family of cycles $\rho^{\epsilon}: B \to G_{\lambda+\epsilon}$ defined for $|\epsilon| \leq \epsilon_0$ such that $\rho^0 = \rho$ and satisfying (24). Using 4.1 (i) every cycle ρ has an extension.

Observation : Consider two extensions $\rho_1^{\epsilon}, 0 \leq |\epsilon| < \epsilon_1$ and $\rho_2^{\epsilon}, 0 \leq |\epsilon| < \epsilon_2$. By (24) there is an $\epsilon' > 0$ and a homotopy between ρ_1^{ϵ} and ρ_2^{ϵ} defined for all $0 \leq \epsilon \leq \epsilon'$. Hence any extension provides well defined elements in $\pi_* G_{\lambda+\epsilon}$ for small values of ϵ . Therefore each $[\rho] \in \pi_*(G_{\lambda}^X)$ has an extension $[\rho^{\epsilon}] \in \pi_*(G_{\lambda+\epsilon}^X)$ whose **germ** at $\epsilon = 0$ is independent of the choices of ρ .

Definition 4.3 We say that a smooth family of elements $[\rho^{\epsilon}] \in \pi_* G_{\lambda+\epsilon}, 0 < \epsilon < \epsilon_{\rho}$ is new if it is not the extension for $\epsilon > 0$ of any element $[\rho] \in \pi_* G_{\lambda}$.

In the next section we will use the same letter ρ to refer both to cycles as well as to the homotopy class they represent.

4.2 The relation between almost complex structures and symplectomorphism groups; the role of PGW

Consider (M, ω_{λ}) symplectic structures on M such that as before, the symplectic forms ω_{λ} span a line in a positive cone \mathcal{K} inside $H^2(M, \mathbb{R})$. Denote by

$$\mathcal{A}_{\ell^+} = \{ J \mid \text{ there is an } \epsilon_J > 0 \text{ s.t. } J \in \mathcal{A}_{\ell+\epsilon} \text{ forall } 0 < \epsilon < \epsilon_J \}$$
(25)

Definition 4.4 Consider a nontrivial element $\beta_{\ell} \in \pi_*(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$. We say that β_{ℓ} is a **persistent** element if its image under the natural morphism

$$i_*\pi_*(\mathcal{A}_{\ell^+},\mathcal{A}_{\ell}) \to \pi_*(\mathcal{A}_{[\ell,\ell+\epsilon]},\mathcal{A}_{\ell})$$

is nonzero for any ϵ arbitrary small.

Proof of the theorem 1.3: We will consider the long exact sequence of relative homotopy groups of the pair $(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$

 $\cdots \longrightarrow \pi_k \mathcal{A}_{\ell^+} \longrightarrow \pi_k (\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell}) \longrightarrow \pi_{k-1} \mathcal{A}_{\ell} \longrightarrow \pi_{k-1} \mathcal{A}_{\ell^+} \longrightarrow \cdots$

Since by construction $\beta_{\ell} \in \pi_k(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$ is nontrivial, then one of the two following cases can happen:

Case 1 $\beta_{\ell} \mapsto \gamma_{\ell} \neq 0 \in \pi_{k-1} \mathcal{A}_{\ell}$

Case 2 $\beta_{\ell} \mapsto 0 \in \pi_{k-1} \mathcal{A}_{\ell}$. In this situation, there is an element $0 \neq \alpha_{\ell} \in \pi_k \mathcal{A}_{\ell^+}$ where $\alpha_{\ell} \mapsto \beta_{\ell}$.

We will do the analysis case by case for our situation:

Case 1 If we are in this case then we consider the fibration (2), that yields

 $G_{\ell} \longrightarrow \operatorname{Diff}_{0}(M) \longrightarrow \mathcal{A}_{\ell}$

We consider the long exact sequence in homotopy

 $\cdots \longrightarrow \pi_{k-1}(G_\ell) \longrightarrow \pi_{k-1}\mathrm{Diff}_0(M) \longrightarrow$

$$\longrightarrow \pi_{k-1}\mathcal{A}_{\ell} \longrightarrow \pi_{k-2}G_{\ell} \longrightarrow \pi_{k-2}\mathrm{Diff}_0(M) \longrightarrow \cdots$$

Again, there are two possibilities:

i) $\gamma_{\ell} \to \theta_{\ell} \neq 0 \in \pi_{k-2}G_{\ell}$. In this situation, we have a nontrivial element $\theta_{\ell} \in \pi_{k-2}G_{\ell}$, such that $\theta_{\ell} \mapsto 0 \in \pi_{k-2}\text{Diff}_0(M)$. Then we are in case **A**.

This element is *fragile*. This can be proved by contrapositive. Assume that θ_{ℓ} can be extended by $\theta_{\ell+\epsilon}$ which yields nontrivial classes in $\pi_{k-2}G_{\ell+\epsilon}$. Then $\theta_{\ell+\epsilon} \mapsto 0 \in \pi_{k-2}\text{Diff}_0(M)$ as well. Therefore it appears as a boundary of an element $\gamma_{\ell+\epsilon} \in \pi_{k-1}\mathcal{A}_{\ell+\epsilon}$ which is homotopic with γ_{ℓ} . But by construction and lemma (4.1), we know that γ_{ℓ} is a contractible cycle inside $\mathcal{A}_{\ell+\epsilon}$. This contradicts the existence of $\gamma_{\ell+\epsilon}$.

Observation : Direct construction of the elements $\eta_{\ell+\epsilon}$

If the fragile element $\theta_{\ell}\pi_{k-2}G_{\ell}$ is trivial, then the elements $\eta_{\ell+\epsilon} \in \pi_{k-1}G_{\ell+\epsilon}$ are constructed as follows.

Since θ_{ℓ} is trivial, it can be represented by a cycle θ_{ℓ} which is the boundary of a k-1dimensional disc $D_{\ell} \in G_{\ell}$. By lemma (4.1) point(ii) we can extend D_{ℓ} to discs $D_{\ell+\epsilon}$ inside $G_{\ell+\epsilon}$. Alternatively, we can push by extensions the cycles θ_{ℓ} into cycles $\theta_{\ell+\epsilon}$ inside $G_{\ell+\epsilon}$. These cycles are null homotopic therefore bound discs $C_{\ell+\epsilon}$ inside $G_{\ell+\epsilon}$. Due to (24), one can see that $\theta_{\ell+\epsilon}$ and $\partial D_{\ell+\epsilon}$ are homotopic for small ϵ , therefore we can glue $C_{\ell+\epsilon}$ and $D_{\ell+\epsilon}$ along their boundaries, and obtain a cycle which we denote by $\eta_{\ell+\epsilon}$. In what will follow we basically show that if θ_{ℓ} is trivial then $\eta_{\ell+\epsilon}$ gives nontrivial elements in homotopy. \Box

ii) $\gamma_{\ell} \mapsto 0 \in \pi_{k-2}G_{\ell}$. Then γ_{ℓ} is in the image of the morphism $\pi_{k-1}\text{Diff}_0(M) \to \pi_{k-1}\mathcal{A}_{\ell}$, and therefore there is an element, $\gamma'_{\ell} \in \pi_{k-1}\text{Diff}_0(M)$ such that $0 \neq \gamma'_{\ell} \mapsto \gamma_{\ell}$.

In this situation, we can choose a cycle $S \subset \mathcal{A}_{\ell}$ representing $\gamma_{\ell} \in \pi_{k-1}(\mathcal{A}_{\ell})$, and, using lemma (4.1), there is an $\epsilon_S > 0$ such that for any ϵ such that $0 < \epsilon < \epsilon_S, S \subset \mathcal{A}_{\ell+\epsilon}$. We make the following

Claim (1) $0 = [S] \in \pi_{k-1} \mathcal{A}_{\ell+\epsilon}$.

By hypothesis S is the boundary of a cycle B_{ℓ} such that $B_{\ell} \subset \mathcal{A}_{\ell+\epsilon}$ for all small $\epsilon > 0$. Therefore we have a k dimensional ball inside $\mathcal{A}_{\ell+\epsilon}$ whose boundary is S, which proves the claim. We therefore have:

Here, from the first row, since γ'_{ℓ} is in the kernel of the map $\pi_{4\ell-3}\text{Diff}_0(M) \to \pi_{k-1}\mathcal{A}_{\ell+\epsilon}$, it has to be in the image of the map $\pi_{k-1}(G_{\ell+\epsilon}) \to \pi_{k-1}\text{Diff}_0(M)$, and therefore we are able to produce an element $0 \neq \eta_{\ell+\epsilon} \in \pi_{k-1}(G_{\ell+\epsilon})$ such that $\eta_{\ell+\epsilon}$ persists in the topology of the group of diffeomorphisms. Thus we are in case **B**.

The elements we obtain here are *new*. This follows easily by assuming the opposite. That is, if we consider that there is an element $0 \neq \eta_{\ell} \in \pi_{k-1}G_{\ell}$ whose germ is given by $\eta_{\ell+\epsilon}$, then the image of η_{ℓ} in $\text{Diff}_0(M)$ has to be γ'_{ℓ} . But this contradicts the fact that $\gamma'_{\ell} \mapsto \gamma_{\ell} \neq 0$. **Case 2.** In this situation we have a nontrivial element $\alpha_{\ell} \in \pi_k \mathcal{A}_{\ell^+}$. We then have the following :

Claim (2) There is an ϵ such that for $0 < \delta < \epsilon \alpha_{\ell}$ has a representative C inside $\mathcal{A}_{\ell+\delta}$, $0 \neq [C] \in \pi_k \mathcal{A}_{\ell+\delta}$. The proof of this statement follows from the construction of α_{ℓ} . Namely, since $\beta_{\ell} \mapsto 0 \in \pi_{k-1} \mathcal{A}_{\ell}$ we conclude that there exist a k-dimensional disk D inside \mathcal{A}_{ℓ} whose boundary is ∂B_{ℓ} ; by lemma (4.1) (i) this can be viewed inside $\mathcal{A}_{\ell+\delta}$ for small δ . We can now glue B_{ℓ} and D along their boundary ∂B_{ℓ} . In this manner we get a cycle $C \subset \mathcal{A}_{\ell+\delta}$ which represents the class α_{ℓ} . We can therefore consider again the sequence

 $\cdots \longrightarrow \pi_k(G_{\ell+\delta}) \longrightarrow \pi_k \operatorname{Diff}_0(M) \longrightarrow$

$$\longrightarrow \pi_k \mathcal{A}_{\ell+\delta} \longrightarrow \pi_{k-1} G_{\ell+\delta} \longrightarrow \pi_{k-1} \text{Diff}_0(M) \longrightarrow \cdots$$

Claim (3) [C] doesn't lift to a nontrivial element in $\pi_k \text{Diff}_0(M)$.
Proof: of claim (3) We should first make the observation that there is a map

$$\pi_k \operatorname{Diff}_0(M) \to \pi_k \mathcal{A}_\lambda$$
 (26)

for any λ and moreover as λ varies this maps are homotopic in \mathcal{A}_I . If C did lift, the map $\pi_{4\ell-2}\mathrm{Diff}_0(M) \to \pi_k \mathcal{A}_\ell$ would produce a cycle $[B] \in \mathcal{A}_\ell$, which by means of lemma (4.1) can be viewed inside all $\mathcal{A}_{\ell+\epsilon}$ for small ϵ and which moreover is homotopic with C inside $\mathcal{A}_{[\ell,\ell+\epsilon]}$. Therefore [C] would map to $0 \in \pi_k(\mathcal{A}_{\ell^+}, \mathcal{A}_\ell)$, which would contradict its definition. Since [C] cannot be in the image of the map $\pi_k \text{Diff}_0(M) \to \pi_k \mathcal{A}_{\ell+\delta}$, we know that [C] must have nonzero image $[C] \mapsto \eta_{\ell+\delta} \neq 0$ in $\pi_{k-1}G_{\ell+\delta}$. Moreover form the obvious properties of exact sequences again, $\eta_{\ell+\delta} \to 0$ through the natural inclusion map $\pi_{k-1}G_{\ell+\delta} \to \pi_{k-1}\text{Diff}_0(M)$. The fact that this elements are *new* follows again by assuming the opposite. If they would form the **germ** of an element η_ℓ in $\pi_{k-1}G_\ell$, then η_ℓ would also be null homotopic inside $\text{Diff}_0(M)$ so it would therefore come from a class [C'] in $\pi_k \mathcal{A}_\ell$. Moreover, C' would be homotopic with C inside $\mathcal{A}_{[\ell,\ell+\delta]}$ therefore also in $(\mathcal{A}_{[\ell,\ell+\delta]}, \mathcal{A}_\ell)$ which is false given that C has to yield a nontrivial element in $\pi_k(\mathcal{A}_{[\ell,\ell+\delta]}, \mathcal{A}_\ell)$. Thus we are in the case **B** of the theorem.

With this, we have exhausted all the possible cases given by the nontrivial PGW. \Box

Assume that there is an ℓ such that there is no J in \mathcal{A}_{ℓ} which can be represented by a J-holomorphic curve in the class D. Then we have the following proposition

Proposition 4.5 Assume that no J in \mathcal{A}_{ℓ} admits J- holomorphic stable maps in class D. Consider an element $0 \neq \beta_{\ell} \in \pi_*(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$ obtained by counting nontrivial parametric Gromov-Witten invariants. Then β_{ℓ} is a persistent element.

The proof follows directly from the theorem (2.11).

Now consider the manifold $(S^2 \times S^2 \times X, \omega_\lambda \oplus \omega_{st})$. As explained in (22) the cycles $(B^\ell, \partial B^\ell)$ satisfy the definition (25), so they give by Prop (4.5) **persistent** elements in $\pi_{4\ell-2}(\mathcal{A}_{\ell^+}, \mathcal{A}_{\ell})$. Therefore theorem (1.2) applies and so the corollary (1.4) holds

Therefore theorem (1.3) applies and so the corollary (1.4) holds.

References

- B. Aebischer, M. Borer, M. Kalin, Ch. Leuenberger, H. M. Reimann Symplectic geometry, Progress in Mathematics vol 124, Birkhäuser, Basel, (1994)
- [2] M. Abreu, Topology of symplectomorphism groups of $S^2 \times S^2$, Inv. Math., **131** (1998), 1-23.
- [3] M. Abreu and D. McDuff, Topology of symplectomorphism groups of rational ruled surfaces, J. Amer. Math. Soc. 13, no.4 (2000), 971-1009.
- [4] J. Bryan and N. C. Leung, The enumerative geometry of K3 surfaces and modular forms, J. Amer. Math Soc. 13 (2000) no. 2, 371-410.
- [5] K. Fukaya and K. Ono, Arnold conjecture and Gromov-Witten invariant, *Topology*, 5 (1999), 933-1048.
- [6] M. Gromov, Pseudo holomorphic curves in symplectic manifolds, *Inv. Math.*, 82 (1985), 307-347.
- [7] P. Kronheimer, Some non-trivial families of symplectic structures, Harvard preprint, (1998).
- [8] T.J. Li and G.Tian, Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds, Topics in symplectic 4-manifolds (Irvine, CA, 1996), *First Int. Press Lect. Ser.*, I, (Cambridge, MA, 1998) 47-83.
- [9] H.V. Le and K. Ono Parameterized Gromov-Witten and topology of symplectomorphism groups, preprint, (2001)

- [10] D. McDuff, Almost complex structures on $S^2 \times S^2$, Duke. Math Journal **101** (2000), 135–177.
- [11] D. McDuff, Symplectomorphism groups and almost complex structures, preprint, (2000).
- [12] D. McDuff and D.A. Salamon, *J-holomorphic curves and quantum cohomology*, University Lecture Series 6, American Mathematical Society, Providence, RI, 1994.
- [13] D. McDuff and D.A. Salamon, Introduction to Symplectic Topology, 2nd edition, Oxford University Press, 1998.
- [14] Y. Ruan, Virtual neighborhoods and pseudo-holomorphic curves, Proceedings of 6th Gökova Geometry-Topology Conference Turkish J. Math., 23 no. 1 (1998), 161-231.
- [15] P. Seidel, On the group of symplectic automorphisms of CP^m × CPⁿ, Amer. Math. Soc. Trans., ser. 2 #196, Nothern California Symplectic Geometry Seminar, (1999), 237-250
- [16] S. Smale, An infinite dimensional version of Sard's Theorem, Amer. J. Math. 87 (1965), 861-866.