

EQUATING DECOMPOSITION NUMBERS FOR DIFFERENT PRIMES

GORDON JAMES AND ANDREW MATHAS

ABSTRACT. This paper shows that certain decomposition numbers for the Hecke algebras and q -Schur algebras at different roots of unity in characteristic zero are equal. To prove our results we first establish the corresponding theorem for the canonical basis of the level one Fock space and then apply deep results of Ariki and Varagnolo and Vasserot.

1. Introduction. Throughout this note we adopt the standard notation for the modular representation theory of the symmetric groups, as can be found in [6, 14].

Consider the following two submatrices of the p -modular decomposition matrices of the symmetric groups \mathfrak{S}_n .

$$\begin{array}{l}
 10, 1 \\
 9, 2 \\
 7, 4 \\
 6, 5 \\
 7, 2^2 \\
 4^2, 3 \\
 6, 2^2, 1 \\
 4^2, 2, 1
 \end{array}
 \left| \begin{array}{ccccccc}
 1 & & & & & & \\
 1 & 1 & & & & & \\
 . & 1 & 1 & & & & \\
 . & . & 1 & 1 & & & \\
 1 & 1 & 1 & . & 1 & & \\
 . & . & 1 & 1 & 1 & 1 & \\
 1 & 1 & 1 & 1 & 1 & . & 1 \\
 . & 1 & 1 & \mathbf{2} & 1 & 1 & 1 & 1
 \end{array} \right.
 \begin{array}{l}
 n = 11 \text{ and } p = 3
 \end{array}
 \qquad
 \begin{array}{l}
 18, 3 \\
 17, 4 \\
 13, 8 \\
 12, 9 \\
 13, 4^2 \\
 8^2, 5 \\
 12, 4^2, 1 \\
 8^2, 4, 1
 \end{array}
 \left| \begin{array}{cccccccc}
 1 & & & & & & & \\
 1 & 1 & & & & & & \\
 . & 1 & 1 & & & & & \\
 . & . & 1 & 1 & & & & \\
 1 & 1 & 1 & . & 1 & & & \\
 . & . & 1 & 1 & 1 & 1 & 1 & \\
 1 & 1 & 1 & 1 & 1 & . & 1 & \\
 . & 1 & 1 & \mathbf{1} & 1 & 1 & 1 & 1
 \end{array} \right.
 \begin{array}{l}
 n = 21 \text{ and } p = 5
 \end{array}$$

The two matrices are identical except for the labelling and the two bold faced entries (omitted entries are zero).

This paper was motivated by our attempts to compute the bold faced entry in the matrix above for $n = 21$ and $p = 5$. At the outset we knew that this number was either 1 or 2; however, we were unable to determine which of these possibilities was correct. Lübeck and Müller [12] have shown that this multiplicity is equal to 1 using computer calculations.

Note that the partitions $(4^2, 2, 1)$ and $(8^2, 4, 1)$ both have p -weight 3 and p -core $(p - 2, p - 2)$ for $p = 3$ and $p = 5$ respectively. Martin and Russell [13] have claimed that all of the p -modular decomposition numbers of the symmetric group of p -weight 3 are 0 or 1 when $p > 3$; unfortunately, their proof contains a gap when dealing with partitions with p -core $(p - 2, p - 2)$. This particular case is still open when $p > 5$; as a consequence, the claim in [13] that the decomposition numbers are always 0 or 1 for partitions of p -weight 3 when $p > 3$ is in doubt.

In this paper we prove a general theorem which indicates why the matrices above are very similar. This result is not about the decomposition matrices of the symmetric groups but rather about the closely related decomposition matrices of the Iwahori-Hecke algebras $\mathcal{H}_{C,q}(\mathfrak{S}_n)$ of the symmetric group at a complex root

1991 *Mathematics Subject Classification.* 20C08, 20C30, 33D80.

of unity. Let $q \in \mathbb{C}$ be a primitive e^{th} root of unity, where $e \geq 2$, and let $q' \in \mathbb{C}$ be a primitive e'^{th} root of unity where $e' > e$. Our result shows that certain decomposition numbers of $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ and $\mathcal{H}_{\mathbb{C},q'}(\mathfrak{S}_m)$ are equal for specified $m > n$. In particular, for the Hecke algebras all of the decomposition multiplicities above for $(n, e) = (11, 3)$ and $(m, e') = (21, 5)$ are equal (in the Hecke algebra case the multiplicities $d_{(4^2, 2, 1), (6, 5)}$, when $e = 3$, and $d_{(8^2, 4, 1), (12, 9)}$, when $e = 5$, are both equal to 1).

In section 4 we also show that the corresponding result holds for the q -Schur algebras in characteristic zero.

2. Abacuses and the Iwahori–Hecke algebra. In order to state our results we recall the abacus notation for partitions introduced in [7]. Fix an integer $e \geq 2$. An e -abacus is an abacus with e runners, which we label from left to right as $\rho_0, \dots, \rho_{e-1}$. Number the bead positions on the abacus by $0, 1, 2, \dots$ reading from left to right and then top to bottom; so the bead positions on ρ_r are numbered $r + me$ for $m \geq 0$. We order the beads on a given e -abacus according to their bead positions.

Let $\lambda = (\lambda_1, \lambda_2, \dots)$ be a partition. The length of λ is the smallest integer $\ell(\lambda)$ such that $\lambda_i = 0$ for all $i > \ell(\lambda)$. If $k \geq \ell(\lambda)$ then λ has a (unique) e -abacus representation with k beads; namely, the e -abacus with beads at positions

$$\lambda_k, \lambda_{k-1} + 1, \dots, \lambda_2 + k - 2, \lambda_1 + k - 1.$$

The bead positions on an abacus for λ encode the first column hook lengths, so this gives a natural bijection between the abacuses with k beads and the partitions of length at most k . For our purposes it is important that the m^{th} bead on an abacus for λ corresponds to row $l = k - m + 1$ of λ (row l of λ is empty if $l > \ell(\lambda)$).

In what follows we fix $k \geq 0$ and (with few exceptions) consider only abacuses with k beads; or, equivalently, partitions of length at most k .

We want to compare e -abacuses with $(e + 1)$ -abacuses. Fix an integer α with $0 \leq \alpha \leq e$. If λ is a partition with $\ell(\lambda) \leq k$ then λ can be represented on an e -abacus with k beads. Let λ^+ be the partition corresponding to the $(e + 1)$ -abacus obtained by inserting an *empty* runner before ρ_α in the e -abacus for λ (if $\alpha = e$ we insert an empty e^{th} runner). Let $\rho_0^+, \dots, \rho_e^+$ be the runners of the $(e + 1)$ -abacus of λ^+ ; then $\rho_r^+ = \rho_r$ if $r < \alpha$, ρ_α^+ is empty, and $\rho_r^+ = \rho_{r-1}$ if $r > \alpha$.

Although our notation does not reflect this the partition λ^+ does depend upon both the choice of α and the choice of k .

2.1. Example. Suppose that $e = 3$, $k = 4$ and $\alpha = 2$. Let $\lambda = (4^2, 3)$. Then the abacuses (with 4 beads) for λ , λ^+ and $\lambda^{++} = (\lambda^+)^+$ are as follows.

$e = 3$	$e' = 4$	$e'' = 5$
$\begin{array}{ccc} 2 & 0 & 1 \\ \hline \bullet & \cdot & \cdot \\ \cdot & \bullet & \cdot \\ \bullet & \bullet & \cdot \\ \cdot & \cdot & \cdot \end{array}$	$\begin{array}{cccc} 0 & 1 & 2 & 3 \\ \hline \bullet & \cdot & \cdot & \cdot \\ \cdot & \bullet & \cdot & \cdot \\ \bullet & \bullet & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{array}$	$\begin{array}{ccccc} 1 & 2 & 3 & 4 & 0 \\ \hline \bullet & \cdot & \cdot & \cdot & \cdot \\ \cdot & \bullet & \cdot & \cdot & \cdot \\ \bullet & \bullet & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \end{array}$
$\lambda = (4^2, 3)$	$\lambda^+ = (6^2, 4)$	$\lambda^{++} = (8^2, 5)$

We have labeled the runners by their residues which will be introduced below.

The reader is invited to check that the partitions which label the decomposition matrix for $n = 21$ and $p = 5$ in the introduction are precisely the partitions λ^{++}

as λ runs over the corresponding partitions of 11. We emphasize that the empty runner can be inserted anywhere in the abacus.

We are now almost ready to describe our main result. As in the introduction let q be a primitive e^{th} root of unity in \mathbb{C} and let $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ be the Iwahori–Hecke algebra of the symmetric group \mathfrak{S}_n defined over the complex numbers with parameter q ; see, for example, [14].

For each partition λ of n Dipper and the first author [2] defined a right $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ -module S_q^λ , called a **Specht module**. There is a natural bilinear form $\langle \cdot, \cdot \rangle$ on S_q^λ and $\text{Rad } S_q^\lambda = \{x \in S_q^\lambda \mid \langle x, y \rangle = 0 \text{ for all } y \in S_q^\lambda\}$ is an $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ -submodule of S_q^λ ; set $D_q^\lambda = S_q^\lambda / \text{Rad } S_q^\lambda$.

Recall that a partition λ is e -regular if no e non-zero parts of λ are equal. Dipper and James showed that D_q^λ is an absolutely irreducible $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ -module whenever λ is e -regular and, further, that every irreducible $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ -module arises uniquely in this way. (Conversely, if λ is not e -regular then $D_q^\lambda = 0$.)

If μ is e -regular let $[S_q^\lambda : D_q^\mu]$ be the multiplicity of the simple module D_q^μ as a composition factor of S_q^λ .

Let q' be a primitive $(e+1)^{\text{st}}$ root of unity in \mathbb{C} . Then we also have Hecke algebras $\mathcal{H}_{\mathbb{C},q'}(\mathfrak{S}_m)$ and modules $S_{q'}^\nu$ and $D_{q'}^\nu$. We shall prove the following.

2.2. Theorem. *Suppose that λ and μ be partitions of n of length at most k and suppose that μ is e -regular. Then $[S_q^\lambda : D_q^\mu] = [S_{q'}^{\lambda^+} : D_{q'}^{\mu^+}]$.*

It may not be clear to the reader that this result is really saying that the decomposition matrices of the blocks containing S_q^λ and $S_{q'}^{\lambda^+}$ are equal on those rows indexed by partitions of length at most k , when we order the rows of these matrices first by length and then by dominance. This follows from Lemma 3.3 below.

It is tempting to speculate that there is some form of category equivalence underpinning this result. However, in general, there are a different number of simple modules in the blocks for λ and λ^+ , so these blocks are certainly not Morita equivalent.

Rather than prove our comparison theorem for decomposition numbers directly we prove a stronger result relating the LLT-polynomials [9]. We now recall the notation needed to describe this.

3. The Fock space and $U_v(\widehat{\mathfrak{sl}}_e)$. Let v be an indeterminate over \mathbb{C} . The **Fock space** is the infinite rank free $\mathbb{C}[v, v^{-1}]$ -module $\mathcal{F} = \bigoplus_{n \geq 0} \bigoplus_{\lambda \vdash n} \mathbb{C}[v, v^{-1}]^\lambda$. The Fock space has a natural structure as a module for the affine quantum group $U_v(\widehat{\mathfrak{sl}}_e)$; we will describe how the positive part $U_v^+(\widehat{\mathfrak{sl}}_e)$ of $U_v(\widehat{\mathfrak{sl}}_e)$ acts on \mathcal{F} since this is all we shall need (full details can be found in [9, 14]).

The **diagram** of a partition λ is the set $[\lambda] = \{(c, d) \in \mathbb{N}^2 \mid d \leq \lambda_c\}$. A **node** is any ordered pair $(c, d) \in \mathbb{N}^2$; in particular, all of the elements of $[\lambda]$ are nodes. The e -**residue** of the node $x = (c, d)$ is $\text{res}_e(x) = d - c \pmod{e}$; x is an i -**node** if $\text{res}_e(x) = i$.

A node x is an **addable** node of λ if $[\lambda] \cup \{x\}$ is the diagram of a partition (and $x \notin [\lambda]$); similarly, x is **removable** if $[\lambda] \setminus \{x\}$ is the diagram of a partition (and $x \in [\lambda]$). For $i = 0, \dots, e-1$ let $A_i(\lambda)$ be the set of addable i -nodes for λ and $R_i(\lambda)$ be the set of removable i -nodes. Given two nodes $x = (c, d)$ and $y = (a, b)$ say that y is **above** x if $c > a$; if y is above x we write $y \succ x$.

In order to define the action of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ on \mathcal{F} for $i = 0, \dots, e-1$ write $\lambda \xrightarrow{i} \nu$ if ν is a partition of $n+1$ and $[\nu] = [\lambda] \cup \{x\}$ for some addable i -node x . Finally, we set $N_i(\lambda, \nu) = \#A_i(\lambda, \nu) - \#R_i(\lambda, \nu)$ where $A_i(\lambda, \nu) = \{y \in A_i(\lambda) \mid y \succ x\}$ and $R_i(\lambda, \nu) = \{y \in R_i(\lambda) \mid y \succ x\}$.

Let F_0, \dots, F_{e-1} be the Chevalley generators of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$. Then the action of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ on \mathcal{F} is determined by

$$(3.1) \quad F_i \lambda = \sum_{\lambda \xrightarrow{i} \nu} v^{N_i(\lambda, \nu)} \nu,$$

for $0, 1, \dots, e-1$.

Let $\Lambda_0, \dots, \Lambda_{e-1}$ be the fundamental weights of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ and let $L(\Lambda_0)$ be the irreducible integrable highest weight module of high weight Λ_0 . Then $L(\Lambda_0) \cong \mathbf{U}_v(\widehat{\mathfrak{sl}}_e)\varnothing = \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)\varnothing$ as a $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ -module [9], where $\varnothing \in \mathcal{F}$ is the empty partition.

Let $\bar{}$ be the bar involution on $\mathbf{U}_{\mathcal{A}}(\widehat{\mathfrak{sl}}_e)$, Lusztig's \mathcal{A} -form of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ (where $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$). Then Lascoux, Leclerc and Thibon [9] showed that $L(\Lambda_0)$ has a basis $\{B_\mu \mid \mu \text{ } e\text{-regular}\}$ which is uniquely determined by the requirements that $\overline{B_\mu} = B_\mu$ and

$$B_\mu = \sum_{\substack{\lambda \vdash n \\ \mu \geq \lambda}} b_{\lambda\mu}(v) \lambda$$

for some polynomials $b_{\lambda\mu}(v) \in \mathbb{Z}[v]$ such that $b_{\mu\mu}(v) = 1$ and $b_{\lambda\mu}(v) \in v\mathbb{Z}[v]$ whenever $\lambda \neq \mu$. This basis is the Kashiwara–Lusztig canonical basis of $L(\Lambda_0)$.

In particular, note that Lascoux, Leclerc and Thibon [9] showed that $b_{\lambda\mu}(v) = 0$ if either $|\lambda| \neq |\mu|$ or if λ and μ have different e -cores.

We want to compare the actions of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ and $\mathbf{U}_v(\widehat{\mathfrak{sl}}_{e+1})$ on the Fock space \mathcal{F} . In order to distinguish between these two algebras let F_0^+, \dots, F_e^+ be the Chevalley generators of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{e+1})$, let $\Lambda_0^+, \dots, \Lambda_e^+$ be its fundamental weights, and let $\mathcal{F}^+ \cong \mathcal{F}$ be the Fock space for the $\mathbf{U}_v(\widehat{\mathfrak{sl}}_{e+1})$ -action. Then $L(\Lambda_0^+) \cong \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{e+1})\varnothing$ as a $\mathbf{U}_v(\widehat{\mathfrak{sl}}_{e+1})$ -module. Given an $(e+1)$ -regular partition ν let

$$B_\nu^+ = \sum_{\nu \geq \sigma} b_{\sigma\nu}^+(v) \sigma$$

be the corresponding canonical basis element of $L(\Lambda_0^+) \subseteq \mathcal{F}^+$.

We can now state a stronger version of our main result.

3.2. Theorem. *Suppose that λ and μ are partitions of n of length at most k and suppose that μ is e -regular. Then $b_{\lambda\mu}(v) = b_{\lambda^+\mu^+}^+(v)$.*

Ariki [1, Prop. 4.3(2)] has shown that the polynomials $b_{\lambda\mu}(v)$ at $v = 1$ compute the decomposition multiplicities; explicitly, $[S_q^\lambda : D_q^\mu] = b_{\lambda\mu}(1)$ and $[S_q^{\lambda^+} : D_q^{\mu^+}] = b_{\lambda^+\mu^+}^+(1)$. Consequently, Theorem 3.2 implies Theorem 2.2. The result also hints at additional structure because, conjecturally, the polynomials $b_{\lambda\mu}(v)$ and $b_{\lambda^+\mu^+}^+(v)$ also describe the Jantzen filtrations of S_q^λ and $S_q^{\lambda^+}$; see [8, 9].

We begin the proof with the following Lemma which is largely book keeping. For example, the statement of Theorem 3.2 requires that μ^+ is $(e+1)$ -regular.

3.3. Lemma. *Let λ and μ be partitions with n of length at most k . Then*

- (i) μ^+ is $(e+1)$ -regular;
- (ii) λ and μ have the same e -core if and only if λ^+ and μ^+ have the same $(e+1)$ -core;
- (iii) if λ and μ have the same e -core then $|\lambda^+| = |\mu^+|$; and,
- (iv) $\lambda \supseteq \mu$ if and only if $\lambda^+ \supseteq \mu^+$

Proof. A partition is $(e+1)$ -regular if and only if its $(e+1)$ -abacus does not contain a string of $e+1$ consecutive beads. Hence, μ^+ is $(e+1)$ -regular since the runner ρ_α^+ is empty; this proves (i). (In fact, if μ is e -regular then so is μ^+ .)

Next, recall that the e -abacus for the e -core of λ is obtained by rearranging the beads on each runner of the e -abacus for λ in such a way that no bead has an empty bead position above it. Hence, if κ is the e -core of λ then κ^+ is the $(e+1)$ -core of λ^+ , so (ii) follows.

For (iii) define w by $|\lambda| = |\kappa| + we$; in other words, w is the e -weight of λ . Now, w can be read off the e -abacus for λ by adding up, for each bead β , the number of empty bead positions which are above β and also on the same runner. Consequently, w is also the $(e+1)$ -weight of λ^+ ; hence, $|\lambda^+| = |\kappa^+| + w(e+1)$. Lastly, since κ is also the e -core of μ it follows that μ is also a partition of e -weight w and that $|\mu^+| = |\kappa^+| + w(e+1) = |\lambda^+|$, as required.

Finally, let $\beta_1^\lambda < \dots < \beta_k^\lambda$ and $\beta_1^\mu < \dots < \beta_k^\mu$ be the positions of the beads on the e -abacuses for λ and μ respectively. Then it is easy to see that $\lambda \supseteq \mu$ if and only if $\sum_{s=t}^k \beta_s^\lambda \geq \sum_{s=t}^k \beta_s^\mu$ for $t = 1, \dots, k$. Rephrasing this condition in terms of the $(e+1)$ -abacuses of λ^+ and μ^+ proves (iv). \square

We remark that if λ and μ are partitions of n with different e -cores then, in general, it is *not* true that $|\lambda^+| = |\mu^+|$.

Let $\mathcal{F}_{>k}$ be the $\mathbb{C}[v, v^{-1}]$ -submodule of \mathcal{F} spanned by the partitions of length strictly greater than k . By (3.1) $\mathcal{F}_{>k}$ is a $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ -submodule of \mathcal{F} ; it is not, however, a $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ -submodule. Therefore, $\mathcal{F}_k = \mathcal{F}/\mathcal{F}_{>k}$ is a $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ -module. We abuse notation and identify the elements of \mathcal{F} with their images in \mathcal{F}_k ; with this understanding, $\{\lambda \mid \ell(\lambda) \leq k\}$ is a basis of \mathcal{F}_k .

Similarly, $\mathcal{F}_k^+ = \mathcal{F}^+/\mathcal{F}_{>k}^+$ is a $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{e+1})$ -module. We want to compare the action of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ on \mathcal{F}_k with the action of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{e+1})$ on \mathcal{F}_k^+ ; to do this we reinterpret (3.1) in terms of abacuses.

Suppose λ is a partition with $\ell(\lambda) \leq k$ and consider the e -abacus of λ . For $0 \leq r < e$ define the e -residue of the runner ρ_r to be the integer $\text{res}_e(\rho_r)$ determined by the following two conditions.

- (i) The e -residue of the runner which holds the last bead is $\lambda_1 - 1 \pmod{e}$.
- (ii) Modulo e , the e -residues of the runners increase by 1 from left to right.

In Example 2.1 the runners are labelled by their f -residues for $f = 3, 4$ and 5 respectively. Similarly, we define the $(e+1)$ -residues $\text{res}_{e+1}(\rho_r^+)$, for $0 \leq r \leq e$, of the runners of the $(e+1)$ -abacus for λ^+ .

The e -residue of a bead β is defined to be the e -residue of the corresponding runner. As we have seen, the k beads on the e -abacus correspond to the first k rows of λ (in reverse order); it is easy to see that the e -residue of a bead is equal to the e -residue of the node at the end of the corresponding row of λ . In particular, this implies that e -residues of the runners depend only on the e -core of λ .

The operator F_i acts on a partition λ by adding nodes of e -residue i . Because the e -residues of the runners correspond to the e -residues of nodes at the end of the rows of λ , this is the same as moving a bead on the e -abacus of λ from the runner with e -residue $i - 1$ to an adjacent empty position on the runner with e -residue i .

Recall that in the definition of λ^+ we have fixed an integer α with $0 \leq \alpha < e$. We now introduce a $\mathbb{C}[v, v^{-1}]$ -linear map ${}^\alpha F_i : \mathcal{F}_k \rightarrow \mathcal{F}_k^+$ for $i = 0, \dots, e - 1$. To define ${}^\alpha F_i$ it is enough to describe ${}^\alpha F_i \lambda$ for each partition λ with $\ell(\lambda) \leq k$. As above, let $\rho_0, \dots, \rho_{e-1}$ be the runners of the e -abacus of λ (with k beads) and let $\rho_0^+, \dots, \rho_e^+$ be the runners of the $(e + 1)$ -abacus for λ^+ . There is a unique r such that $i = \text{res}_e(\rho_r)$ (and $0 \leq r < e$); set $j = \text{res}_{e+1}(\rho_r^+)$. Define

$${}^\alpha F_i \lambda = \begin{cases} F_j^+ \lambda^+, & \text{if } 0 \leq r < \alpha, \\ F_{j+1}^+ F_j^+ \lambda^+, & \text{if } r = \alpha, \\ F_{j+1}^+ \lambda^+, & \text{if } \alpha < r < e, \end{cases}$$

where $j + 1$ is understood modulo e . Similarly, we define the divided powers ${}^\alpha F_i^{(a)}$ for $a \geq 1$; for example, when $r = \alpha$ we set ${}^\alpha F_i^{(a)} \lambda = F_{j+1}^{+(a)} F_j^{+(a)} \lambda^+$.

For our final piece of notation observe that $\ell(\lambda^+) \geq \ell(\lambda)$ for any partition λ . Therefore, we have a well-defined $\mathbb{C}[v, v^{-1}]$ -linear map $\Theta : \mathcal{F}_k \rightarrow \mathcal{F}_k^+$ determined by $\Theta(\lambda) = \lambda^+$ for $\ell(\lambda) \leq k$. As with λ^+ , we emphasize that Θ depends upon both α and k . The map Θ is injective but not surjective, having image the span of those partitions of length at most k which have an $(e + 1)$ -abacus with k beads and with an empty runner ρ_α^+ .

3.4. Lemma. *Suppose that $0 \leq i < e$. Then the following diagram commutes.*

$$\begin{array}{ccc} \mathcal{F}_k & \xrightarrow{F_i} & \mathcal{F}_k \\ & \searrow {}^\alpha F_i & \downarrow \Theta \\ & & \mathcal{F}_k^+ \end{array}$$

Proof. It suffices to verify the lemma for a partition $\lambda \in \mathcal{F}_k$. As above, let ρ_r be the runner in the e -abacus for λ for which $i = \text{res}_e(\rho_r)$ and set $j = \text{res}_{e+1}(\rho_r^+)$.

First consider $F_i \lambda = \sum_\nu v^{N_i(\lambda, \nu)} \nu^+$. Recall that the beads on the e -abacus for λ are naturally indexed by the rows of λ and that the e -residue of a bead is defined to be the e -residue of the node which is at the end of the corresponding row. Therefore, an addable i -node of λ corresponds to a node on runner $r - 1$ of the e -abacus which can be moved to the adjacent position on runner r (which must therefore be empty). Similarly, a removable i -node corresponds to a node on runner r which can be moved back to the adjacent position on runner $r - 1$; here, $r \pm 1$ is to be understood modulo e . The addable and removable nodes of λ^+ have analogous descriptions.

Fix a partition ν with $\lambda \xrightarrow{i} \nu$ and write $[\nu] = [\lambda] \cup \{x\}$. Then there exists a node at position β_x on the runner $r - 1$ of the e -abacus for λ which can be moved to the adjacent position on runner r so as to give the e -abacus for ν . Then $N_i(\lambda, \nu) = A - B$, where $A = \#\{y \in A_i(\lambda) \mid y \succ x\}$ and $B = \#\{y \in R_i(\lambda) \mid y \succ x\}$. If y is an addable or removable node of λ then $y \succ x$ if and only if it corresponds to a bead at position β_y with $\beta_y > \beta_x$. Hence, A is equal to the number of beads on runner ρ_{r-1} which come after β_x such that the adjacent position on ρ_r is vacant;

similarly, B is equal to the number of beads on ρ_r which are after β_x and for which the adjacent position on runner $r - 1$ is vacant.

Now consider the $(e + 1)$ -abacuses for λ^+ and ν^+ . Assume first that $r < \alpha$. Then the runners ρ_{r-1} and ρ_r for λ are the same as the runners ρ_{r-1}^+ and ρ_r^+ for λ^+ and so the last paragraph shows that the addable and removable i -nodes for λ correspond exactly to the addable and removable j -nodes for λ^+ . Hence, $N_j(\lambda^+, \nu^+) = A - B = N_i(\lambda, \nu)$. Similarly, when $r > \alpha$ the addable and removable i -nodes for λ correspond to the addable and removable $(j + 1)$ -nodes for λ^+ and $N_i(\lambda, \nu) = N_{j+1}(\lambda^+, \nu^+)$.

Finally, consider the case when $r = \alpha$. This time runner ρ_{r-1} is equal to ρ_{r-1}^+ and runner ρ_r is equal to ρ_{r+1}^+ ; whereas runner $\rho_r^+ = \rho_\alpha^+$ of λ^+ is empty. Therefore, the addable and removable i -nodes of λ again correspond to addable and removable j -nodes of λ^+ except that this time there are additional addable j -nodes of λ^+ corresponding to the adjacent pairs of beads on the runners ρ_{r-1} and ρ_r of the e -abacus of λ . Let σ be the partition such that $\lambda^+ \xrightarrow{j} \sigma \xrightarrow{j+1} \nu^+$. Since ρ_r^+ is empty, λ^+ has no removable j -nodes. Therefore, if we let l be the number of pairs of adjacent beads on runners ρ_{r-1} and ρ_r which are above β_x then $N_j(\lambda^+, \sigma) = A + l$. Next observe that σ has a single addable $(j + 1)$ -node (corresponding to the bead which we just moved), and that the removable $(j + 1)$ -nodes of σ correspond to the removable i -nodes of λ together with the l beads on runner ρ_r which we have already paired with an adjacent bead on ρ_{r-1} ; therefore, $N_{j+1}(\sigma, \nu^+) = -(B + l)$. Consequently, $N_j(\lambda^+, \sigma) + N_{j+1}(\sigma, \nu^+) = A - B = N_i(\lambda, \nu)$ and so we have

$$F_{j+1}^+ F_j^+ \lambda^+ = \sum_{\nu^+} v^{N_i(\lambda, \nu)} \nu^+ = \Theta(F_i \lambda),$$

where the sum is over those partitions ν^+ for which there exists a partition σ such that $\lambda^+ \xrightarrow{j} \sigma \xrightarrow{j+1} \nu^+$. Note that there are additional terms in the expansion of $F_j \lambda^+$ (corresponding to the pairs of adjacent beads on runners ρ_{r-1} and ρ_r of the e -abacus for λ); however, they all disappear when F_{j+1} is applied because these extra partitions do not have any addable $(j + 1)$ -nodes. This completes the proof. \square

Now consider $L(\Lambda_0)_k = L(\Lambda_0)/(L(\Lambda_0) \cap \mathcal{F}_k)$. If μ is an e -regular partition with $\ell(\mu) \leq k$ let $\tilde{B}_\mu = B_\mu + \mathcal{F}_{>k}$. As noted by Goodman and Wenzl [5, Lemma 4.1], the elements $\{B_\mu \mid \mu \text{ is } e\text{-regular and } \ell(\mu) \leq k\}$ give a basis of $L(\Lambda_0)_k$.

The bar involution induces a well-defined map on \mathcal{F}_k via $\overline{a + \mathcal{F}_k} = \bar{a} + \mathcal{F}_k$ for all $a \in \mathcal{F}$. It is easy to see that \tilde{B}_μ is the unique element of \mathcal{F}_k which is bar invariant and of the form $\mu + \sum_\lambda b_{\lambda\mu}(v)\lambda$ for some polynomials $b_{\lambda\mu}(v) \in v\mathbb{Z}[v]$, the sum being over the partitions of length at most k .

Similarly, $\{\tilde{B}_\nu^+ \mid \nu \text{ is } (e + 1)\text{-regular and } \ell(\nu) \leq k\}$, where $\tilde{B}_\nu^+ = B_\nu^+ + \mathcal{F}_{>k}^+$, is a basis of the $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{e+1})$ -module $L(\Lambda_0^+)_k = L(\Lambda_0^+)/(L(\Lambda_0^+) \cap \mathcal{F}_k^+)$.

3.5. Proposition. *Suppose that μ is an e -regular partition with at most k rows. Then $\tilde{B}_{\mu^+}^+ = \Theta(\tilde{B}_\mu)$.*

Proof. Looking at the definitions, $\Theta(\tilde{B}_\mu) = \mu^+$ plus a $q\mathbb{Z}[q]$ -linear combination of less dominant terms. Therefore, it is enough to show that $\Theta(\tilde{B}_\mu)$ is a bar invariant element of \mathcal{F}_k^+ .

Let $\emptyset \in \mathcal{F}_k$ be the image of the empty partition in \mathcal{F}_k . Following Lascoux, Leclerc and Thibon [9, Lemma 6.4] let $(r_1^{a_1}, \dots, r_s^{a_s})$ be the e -residue sequence of μ

corresponding to the e -ladders in the diagram of μ . Then $A_\mu = F_{r_s}^{(a_s)} \dots F_{r_1}^{(a_1)} \emptyset$ is a bar invariant element of \mathcal{F}_k of the form $A_\mu = \mu + \sum_\lambda a_{\lambda\mu}(v)\lambda$ where $a_{\lambda\mu}(v) \in \mathbb{Z}[v, v^{-1}]$ and the sum is over partitions λ of length at most k such that $\mu \triangleright \lambda$. Therefore, there exist uniquely determined polynomials $\alpha_{\sigma\mu}(v) \in \mathbb{Z}[v]$ such that $\tilde{B}_\mu = A_\mu - \sum_\sigma \alpha_{\sigma\mu}(v)\tilde{B}_\sigma$, where the sum is over e -regular partitions σ such that $\mu \triangleright \sigma$ and $\ell(\sigma) \leq k$.

Now consider the element $A_\mu^+ = \Theta(A_\mu) = \mu + \sum_\lambda a_{\lambda\mu}(v)\lambda^+$ in \mathcal{F}_k^+ . By the Lemma, $A_\mu^+ = {}^\alpha F_{r_s}^{(a_s)} \dots {}^\alpha F_{r_1}^{(a_1)} \emptyset$; hence, A_μ^+ is bar invariant. By induction on dominance $\tilde{B}_\sigma^+ = \Theta(\tilde{B}_\sigma)$ for $\mu \triangleright \sigma$. Therefore, the element $\Theta(\tilde{B}_\mu) = A_\mu^+ - \sum_\sigma \alpha_{\sigma\mu}(v)\tilde{B}_{\sigma^+}$ is also bar invariant. Consequently, $\tilde{B}_{\mu^+}^+ = \Theta(\tilde{B}_\mu)$ as we wanted to show. \square

It is easy to see [9] that the polynomials $b_{\sigma\tau}^+(v)$ are non-zero only if σ and τ have the same $(e+1)$ -core. Therefore, if μ is an e -regular partition with $\ell(\mu) \leq k$ then $\tilde{B}_{\mu^+}^+ = \sum_{\mu \triangleright \lambda} b_{\lambda^+\mu^+}^+(v)\lambda^+$; on the other hand, $\Theta(B_\mu) = \sum_{\mu \triangleright \lambda} b_{\lambda\mu}(v)\lambda^+$, where in both sums $\ell(\lambda) \leq k$. Hence, Theorem 3.2 follows from the Proposition.

Goodman and Wenzl [5, Prop. 5.14] have proved a result very similar to Proposition 3.5; however, it is not the same because we do not require that the weight $\mu + \rho$ belong to a proper face of an alcove. Our result is also more explicit. Nonetheless, the origin for these results is that the $b_{\lambda\mu}(v)$ are parabolic Kazhdan–Lusztig polynomials for the parabolic subgroup \mathfrak{S}_k of the extended affine Weyl group $\hat{\mathfrak{S}}_k$ [5, 11, 15]; in turn the parabolic Kazhdan–Lusztig polynomials are naturally indexed by the alcoves and, generically, the alcove geometry does not depend on k or e .

4. The q -Schur algebras. In this section we extend Theorem 2.2 to the decomposition matrices of the q -Schur algebras $S_{\mathbb{C},q}(n, n)$ of Dipper and the first author [3].

The Fock space also admits an action from a Heisenberg algebra \mathcal{H}_e [11]. The action of \mathcal{H}_e on \mathcal{F} commutes with the action of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ and it is useful because \mathcal{F} is irreducible when considered as a module for the algebra generated by the actions of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ and \mathcal{H}_e on \mathcal{F} . In addition, the action of \mathcal{H}_e allowed Leclerc and Thibon [11, §7.9] to extend the bar involution to the whole of \mathcal{F} ; in turn, this enabled them to extend the canonical basis of $L(\Lambda_0)$ to give a basis $\{B_\mu \mid \mu \text{ a partition}\}$ of \mathcal{F} where, the element B_μ is again uniquely determined by the two conditions that $\overline{B}_\mu = B_\mu$ and

$$B_\mu = \sum_{\substack{\lambda \vdash n \\ \mu \triangleright \lambda}} b_{\lambda\mu}(v)\lambda$$


for some polynomials $b_{\lambda\mu}(v) \in \mathbb{Z}[v]$ such that $b_{\mu\mu}(v) = 1$ and $b_{\lambda\mu}(v) \in v\mathbb{Z}[v]$ whenever $\lambda \neq \mu$. We will show that Theorem 3.2 generalizes to the non-regular case.

As in the previous sections we are only interested in the action of a subalgebra \mathcal{H}_e^+ of \mathcal{H}_e ; for the full story see [11, §7.5]. The algebra \mathcal{H}_e^+ is generated by elements \mathcal{V}_m for $m \geq 0$; before we can describe how \mathcal{V}_m acts on \mathcal{F} we need some more notation.

An e -ribbon is a connected strip of e -nodes which does not contain a 2×2 square; more precisely, an e -ribbon is a set of e nodes $R = \{(a_1, b_1), \dots, (a_e, b_e)\}$ such that (a_{i+1}, b_{i+1}) is either $(a_i + 1, b_i)$ or $(a_i, b_i - 1)$, for $i = 1, \dots, e - 1$. The head of R is the node $\text{head}(R) = (a_1, b_1)$ and $\text{spin}_e(R) = \#\{1 \leq i < e \mid a_{i+1} = a_i + 1\} - 1$ is the e -spin of R .

If λ and ν are partitions then we write $\lambda \xrightarrow{m:e} \nu$ if $[\lambda] \subseteq [\nu]$ and $[\nu] \setminus [\lambda]$ is a disjoint union of m e -ribbons such that the head of each ribbon is either in the first row of λ or is of the form (i, j) where $(i-1, j) \in [\lambda]$. Lascoux, Leclerc and Thibon (see [11, §4.1]), call ν/λ an e -ribbon tableau of weight (m) and they note that there is a unique way of writing $[\nu] \setminus [\lambda]$ as a disjoint union of ribbons; we will see this below when we reinterpret ribbons in terms of abacuses. Finally, if $\lambda \xrightarrow{m:e} \nu$ then $\text{spin}_e(\nu/\lambda)$, the e -spin of ν/λ , is the sum of the e -spins of the ribbons in $[\nu] \setminus [\lambda]$.

For example, if $\lambda = (3)$ and $e = 2$ then the partitions ν with $\lambda \xrightarrow{2:2} \nu$ are



with spins -2 , -2 and 2 respectively.

The algebra \mathcal{H}_e^+ is the subalgebra of \mathcal{H}_e generated by elements \mathcal{V}_m for $m \geq 1$. For each m , \mathcal{V}_m acts on the Fock space as the $\mathbb{C}[v, v^{-1}]$ -linear map determined by

$$\mathcal{V}_m \lambda = \sum_{\lambda \xrightarrow{m:e} \nu} (-q)^{\text{spin}_e(\nu/\lambda)} \nu$$

for all partitions λ . Observe that $\mathcal{F}_{>k}$ is a \mathcal{H}_e^+ -module; hence, there is a well-defined action of \mathcal{V}_m on the quotient space \mathcal{F}_k .

Similarly, there is an action of the positive Heisenberg algebra \mathcal{H}_{e+1}^+ on the Fock space \mathcal{F}^+ and this induces an action on \mathcal{F}_k^+ . We denote the generators of \mathcal{H}_{e+1}^+ by \mathcal{V}_m^+ for $m \geq 1$.

4.1. Lemma. *Suppose that λ and ν are partitions of n of length at most k .*

- (i) *If $\lambda \xrightarrow{m:e} \nu$ then λ and μ have the same e -core.*
- (ii) *We have $\lambda \xrightarrow{m:e} \nu$ if and only if $\lambda^+ \xrightarrow{m:e+1} \mu^+$; moreover, if $\lambda \xrightarrow{m:e} \nu$ then $\text{spin}_e(\nu/\lambda) = \text{spin}_{e+1}(\nu^+/\lambda^+)$.*

Proof. The Lemma will follow once we reinterpret the condition $\lambda \xrightarrow{m:e} \nu$ in terms of abacuses. Suppose that $\lambda \xrightarrow{m:e} \nu$. Then $[\nu] \setminus [\lambda]$ is a disjoint union of e -ribbons. Extend the partial order \succ on the set of nodes to a total order by defining $(a, b) \succ (c, d)$ if either $c > a$ or $c = a$ and $b > d$. Totally order the ribbons R_1, \dots, R_m in $[\nu] \setminus [\lambda]$ so that $i > j$ whenever $\text{head}(R_i) \succ \text{head}(R_j)$. Then the condition that the head of R_i is of the form (a, b) with either $a = 1$ or $(a-1, b) \in [\lambda]$ is equivalent to saying that $[\nu] \setminus (R_1 \cup \dots \cup R_i)$ is the diagram of a partition for $i = 1, \dots, m$. Hence, it is enough to treat the case $m = 1$. So let $R = R_1$ where $[\nu] = [\lambda] \cup R$.

Now the ribbon R is a rim hook and it is well-known that removing a rim hook of length e from ν is the same as moving a bead β on an e -abacus for ν to the (empty) bead position on the same runner which is in the preceding row. Further, by definition, $\text{spin}_e(\nu/\lambda)$ is the leg length of R minus one and, in terms of the e -abacus, the leg length of R is equal to the number of beads on the abacus which are between the old and new positions of β . Similarly, the condition $\lambda^+ \xrightarrow{m:e+1} \mu^+$ depends only on the $(e+1)$ -abacuses of ν^+ and λ^+ . As the e and $(e+1)$ abacuses differ only by the insertion of an empty runner, all of the assertions of the Lemma now follow. \square

4.2. Corollary. *Suppose that $m \geq 1$. Then the following diagram commutes.*

$$\begin{array}{ccc} \mathcal{F}_k & \xrightarrow{\mathcal{V}_m} & \mathcal{F}_k \\ \Theta \downarrow & & \downarrow \Theta \\ \mathcal{F}_k^+ & \xrightarrow{\mathcal{V}_m^+} & \mathcal{F}_k^+ \end{array}$$

Proof. As with Lemma 4.1 it suffices to check the result for a partition λ of length at most k . By the definitions and the previous Lemma,

$$\begin{aligned} \Theta(\mathcal{V}_m \lambda) &= \sum_{\lambda \xrightarrow{m:e} \nu} (-q)^{\text{spin}_e(\nu/\lambda)} \Theta(\nu) \\ &= \sum_{\lambda^+ \xrightarrow{m:e+1} \nu^+} (-q)^{\text{spin}_{e+1}(\nu^+/\lambda^+)} \nu^+ \\ &= \mathcal{V}_m^+ \lambda^+. \end{aligned}$$

Therefore, $\mathcal{V}_m^+(\Theta(\lambda)) = \Theta(\mathcal{V}_m \lambda)$ and we're done. \square

Let μ be a partition. As in the last section let \tilde{B}_μ and \tilde{B}_μ^+ denotes the images of $B_\mu \in \mathcal{F}$ and $B_\mu^+ \in \mathcal{F}^+$ in \mathcal{F}_k and \mathcal{F}_k^+ respectively. Then a basis of \mathcal{F}_k is given by the \tilde{B}^μ as μ runs over all partitions of length at most k and similarly for \mathcal{F}_k^+ .

By Proposition 3.5 we know that $\tilde{B}_{\mu^+}^+ = \Theta(B_\mu)$ whenever μ is an e -regular partition with $\ell(\mu) \leq k$. We can now drop the requirement that μ should be e -regular.

4.3. Proposition. *Suppose that μ is a partition with $\ell(\mu) \leq k$. Then $\tilde{B}_{\mu^+}^+ = \Theta(\tilde{B}_\mu)$.*

Proof. As before, the element $\Theta(\tilde{B}_\mu)$ is equal to μ^+ plus a $q\mathbb{Z}[q]$ -linear combination of less dominant terms, so it is enough to show that $\Theta(\tilde{B}_\mu)$ is invariant under the bar involution since $\tilde{B}_{\mu^+}^+$ is the unique bar invariant element in \mathcal{F}_k of this form.

By [11, Prop. 7.6] the bar involution on \mathcal{F} is completely determined by the conditions $\overline{\emptyset} = \emptyset$, $\overline{F_i^{(m)} x} = F_i^{(m)} \overline{x}$ and $\overline{\mathcal{V}_m x} = \mathcal{V}_m \overline{x}$, for all $x \in \mathcal{F}$, $0 \leq i < e$ and $m \geq 1$. For each partition $\tau = (\tau_1, \dots, \tau_s)$ let $\mathcal{V}_\tau = \mathcal{V}_{\tau_1} \dots \mathcal{V}_{\tau_s}$. Then

$$\mathcal{F} = \bigoplus_{\tau} \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e) \mathcal{V}_\tau \emptyset$$

is a decomposition of \mathcal{F} into a direct sum of irreducible $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules (where τ runs over all partitions of all integers); see [11, §7.5]. Moreover, the modules $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e) \mathcal{V}_\tau \emptyset$, for different τ , are all isomorphic as $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ -modules. Therefore, there exists a bar invariant basis of \mathcal{F} of the form $A_{\sigma\tau} = F_\sigma \mathcal{V}_\tau \emptyset$ where $F_\sigma \in \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$, σ is an e -regular partition and τ is an arbitrary partition (the elements F_σ are defined in terms of e -residue sequences as in the proof of Proposition 3.5). Consequently, we can write $\tilde{B}_\mu = \sum_{\sigma,\tau} a_{\sigma\tau}(v) A_{\sigma\tau}$ for some bar invariant Laurent polynomials $a_{\sigma\tau}(v) \in \mathbb{Z}[v, v^{-1}]$. Now, $A_{\sigma\tau}^+ = \Theta(A_{\sigma\tau}) = \Theta(F_\sigma \mathcal{V}_\tau \emptyset) = {}^\alpha F_\sigma \mathcal{V}_\tau^+ \emptyset$ by Lemma 3.4 and Corollary 4.2; therefore, $A_{\sigma\tau}^+$ is a bar invariant element of \mathcal{F}_k^+ . Hence, $\Theta(\tilde{B}_\mu) = \sum_{\sigma,\tau} a_{\sigma\tau}(v) A_{\sigma\tau}^+$ is a bar invariant element of \mathcal{F}_k^+ , as we needed to show. \square

4.4. *Remark.* For each composition $\tau = (\tau_1, \dots, \tau_s)$ Leclerc and Thibon [11] show that the action of the element $\mathcal{V}_\tau = \mathcal{V}_{\tau_1} \dots \mathcal{V}_{\tau_s}$ upon \mathcal{F} is described by certain polynomials associated with the ribbon tableaux of weight τ . This is completely analogous to the way in which the action of $F_{r_s}^{(a_s)} \dots F_{r_1}^{(a_1)}$ on \mathcal{F} can be described in terms of polynomials associated with standard tableaux.

Comparing the coefficient of λ^+ in $\tilde{B}_{\mu^+}^+$ and $\Theta(\tilde{B}_\mu)$ we obtain the following.

4.5. Corollary. *Suppose that λ and μ are partitions of length at most k . Then $b_{\lambda\mu}(v) = b_{\lambda^+\mu^+}(v)$.*

In order to compute the polynomials $b_{\lambda\mu}(v)$ when μ is not e -regular it is necessary to first invert the “ R -matrix” which describes the bar involution on the basis of \mathcal{F} given by the set of partitions. Computationally, this is quite time consuming; in comparison the regular case is much easier, being essentially Gaussian elimination. Corollary 4.5 therefore gives a slightly more efficient way of computing the polynomials $b_{\lambda\mu}(v) = b_{\lambda^+\mu^+}(v)$ since μ^+ is an $(e+1)$ -regular partition by Lemma 3.3(i).

Next, let W_q^λ and F_q^μ be the Weyl modules and simple modules, respectively, for the q -Schur algebra $S_{\mathbb{C},q}(n, n)$. Similarly, let $W_{q'}^{\lambda^+}$ and $F_{q'}^{\mu^+}$ be the Weyl modules and simple modules of the q' -Schur algebra $S_{\mathbb{C},q'}(n, n)$. Varagnolo and Vasserot [15] have shown that $[W_q^\lambda : F_q^\mu] = b_{\lambda\mu}(1)$; similarly, $[W_{q'}^{\lambda^+} : F_{q'}^{\mu^+}] = b_{\lambda^+\mu^+}(1)$.

4.6. Corollary. *Suppose that λ and μ are partitions of length at most k . Then*

$$[W_q^\lambda : F_q^\mu] = [W_{q'}^{\lambda^+} : F_{q'}^{\mu^+}] = [S_{q'}^{\lambda^+} : D_{q'}^{\mu^+}].$$

Proof. That $[W_q^\lambda : F_q^\mu] = [W_{q'}^{\lambda^+} : F_{q'}^{\mu^+}]$ follows directly from the previous Corollary and the remarks above. For the second claim, observe that the partition μ^+ is $(e+1)$ -regular by Lemma 3.3(i); therefore, $D_{q'}^{\mu^+} \neq 0$. Consequently, $[W_{q'}^{\lambda^+} : F_{q'}^{\mu^+}] = [S_{q'}^{\lambda^+} : D_{q'}^{\mu^+}]$ by Schur–Weyl reciprocity. \square

Standard Schur functor arguments yield the corresponding statements for the q -Schur algebras $S_{\mathbb{C},q}(n, r)$ and $S_{\mathbb{C},q'}(n, r)$; we leave the details to the reader.

The last result is interesting because it shows that every decomposition number for $S_{\mathbb{C},q}(n, n)$ is also a decomposition number for some Hecke algebra $\mathcal{H}_{\mathbb{C},q'}(\mathfrak{S}_m)$. In contrast Erdmann [4] has shown that in a given characteristic knowing all decomposition numbers for the classical Schur algebras (i.e. $q = 1$) is equivalent to knowing all decomposition numbers for the symmetric groups (for all n and for a fixed p). Leclerc [10] has proved the analogous result relating the decomposition numbers of the q -Schur algebras $S_{\mathbb{C},q}(n, n)$ and the Hecke algebras $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ (for all n and for a fixed q). No such result is known in the cross characteristic case (i.e. positive characteristic with $q \neq 1$).

Finally, we remark that the full action of $\mathbf{U}_v(\widehat{\mathfrak{sl}}_e)$ on \mathcal{F}_k and $\mathbf{U}_v(\widehat{\mathfrak{sl}}_{e+1})$ on \mathcal{F}_k^+ are compatible via the map Θ (in order to make this statement precise \mathcal{F}_k should be considered as a submodule of \mathcal{F} for the action of $\mathbf{U}_v^-(\widehat{\mathfrak{sl}}_e)$). This can be proved using similar arguments or, more simply, by invoking [11, Prop. 7.9] which says that the actions of $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_e)$ and $\mathbf{U}_v^-(\widehat{\mathfrak{sl}}_e)$ on \mathcal{F} are adjoint with respect to a natural scalar product on \mathcal{F} . The same argument also proves the corresponding statements for the Heisenberg algebras \mathcal{H}_e and \mathcal{H}_{e+1} .

5. Examples. Below we give part of the “crystallized” decomposition matrices $(b_{\lambda\mu}(v))$ of the Hecke algebras $\mathcal{H}_{\mathbb{C},q}(\mathfrak{S}_n)$ for $(e, n) = (2, 6), (3, 11), (4, 16)$ and $(5, 21)$. By our results, taking $k = 4$ and $\alpha = 2$, these submatrices are all the same.

\cdots	$e = 5$	$e = 4$	$e = 3$	$e = 2$							
	18, 3	14, 2	10, 1	6	1						
	17, 4	13, 3	9, 2	5, 1	v	1					
\cdots	13, 8	11, 5	7, 4	4, 2	.	v	1				
	12, 9	9, 7	6, 5	3^2	.	.	v	1			
	13, 4^2	10, 3^2	7, 2^2	4, 1^2	v	v^2	v	.	1		
\cdots	$8^2, 5$	$6^2, 4$	$4^2, 3$	2^3	.	.	v^2	v	v	1	
	12, $4^2, 1$	9, $3^2, 1$	6, $2^2, 1$	3, 1^3	v^2	v	v^2	v	v	.	1
	$8^2, 4, 1$	$6^2, 3, 1$	$4^2, 2, 1$	$2^2, 1^2$.	v^2	v^3	v^2	v^2	v	1

Note that when $e = 2$ some of the rows are indexed by partitions which are not 2-regular; in this case the matrix above is a submatrix of the decomposition matrix for the corresponding q -Schur algebra with $e = 2$ and $n = 6$. Setting $v = 2$ we recover the decomposition matrices of the Iwahori–Hecke algebras corresponding to those of the introduction.

To emphasize the dependence on k we again start with $(e, n) = (2, 6)$ but now take $k = 6$ (and $\alpha = 2$); this yields the following matrices.

$e = 5$	$e = 4$	$e = 3$	$e = 2$									
21, $6, 3^2$	16, $4, 2^2$	11, $2, 1^2$	6	1								
20, $7, 3^2$	15, $5, 2^2$	10, $3, 1^2$	5, 1	v	1							
16, $11, 3^2$	12, $8, 2^2$	8, $5, 1^2$	4, 2	.	v	1						
16, $7^2, 3$	12, $5^2, 2$	8, $3^2, 1$	4, 1^2	v	v^2	v	1					
15, $12, 3^2$	11, $9, 2^2$	7, $6, 1^2$	3, 3	.	.	v	.	1				
15, $7^2, 4$	11, $5^2, 3$	7, $3^2, 2$	3, 1^3	v^2	v	v^2	v	v	1			
11 $^2, 8, 3$	$8^2, 6, 2$	$5^2, 4, 1$	2^3	.	.	v^2	v	v	.	1		
11 $^2, 7, 4$	$8^2, 5, 3$	$5^2, 3, 2$	$2^2, 1^2$.	v^2	v^3	v^2	v^2	v	v	1	
11, $7^2, 4^2$	8, $5^2, 3^2$	5, $3^2, 2^2$	2, 1^4	v^2	v^3	.	v	.	v^2	.	v	1
10, $7^2, 4^2, 1$	7, $5^2, 3^2, 1$	4, $3^2, 2^2, 1$	1^6	v^3	.	.	v^2	.	.	.	v	1

Notice that in the first example the lengths of the partitions in the columns for $e = 2$ and $e = 5$ were always the same; this is no longer true in this example. For these partitions, the dominance order is a total order when $e \geq 3$; for this reason we have ordered the rows of this matrix according to dominance for the partitions with $e \geq 3$; this accounts for the different ordering of the rows for $e = 2$ in the two examples.

A consequence of Lemma 3.3 is that all of these matrices are the rows of decomposition matrices of the corresponding blocks which are indexed by partitions with at most k rows (where the partitions indexing the rows are ordered first by length and then by dominance).

References.

- [1] S. ARIKI, *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$* , J. Math. Kyoto Univ., **36** (1996), 789–808.
- [2] R. DIPPER AND G. JAMES, *Representations of Hecke algebras of general linear groups*, Proc. L.M.S. (3), **52** (1986), 20–52.
- [3] ———, *The q -Schur algebra*, Proc. L.M.S. (3), **59** (1989), 23–50.

- [4] K. ERDMANN, *Decomposition numbers for symmetric groups and composition factors of Weyl modules*, J. Algebra, **180** (1996), 316–320.
- [5] F. M. GOODMAN AND H. WENZL, *Crystal bases of quantum affine algebras and Kazhdan–Lusztig polynomials*, Int. Math. Res. Notes, **5** (1999), 251–275.
- [6] G. D. JAMES, *The representation theory of the symmetric groups*, SLN, **682**, Springer–Verlag, New York, 1978.
- [7] ———, *Some combinatorial results involving Young diagrams*, Proc. Camb. Phil. Soc., **83** (1978), 1–10.
- [8] G. D. JAMES AND A. MATHAS, *A q -analogue of the Jantzen–Schaper theorem*, Proc. L.M.S. (3), **74** (1997), 241–274.
- [9] A. LASCoux, B. LECLERC, AND J.-Y. THIBON, *Hecke algebras at roots of unity and crystal bases of quantum affine algebras*, Comm. Math. Phys., **181** (1996), 205–263.
- [10] B. LECLERC, *Decomposition numbers and canonical bases*, Algebr. Represent. Theory, **3** (2000), 277–287.
- [11] B. LECLERC AND J.-Y. THIBON, *Littlewood–Richardson coefficients and Kazhdan–Lusztig polynomials*, in Combinatorial methods in representation theory, M. Kashiwara et al., eds., Adv. Pure Math., **28**, 2000, 155–220.
- [12] F. LÜBECK AND J. MÜLLER, *personal communication*, 2000.
- [13] S. MARTIN AND L. RUSSELL, *Defect 3 blocks of symmetric group algebras*, J. Algebra, **213** (1999), 304–339.
- [14] A. MATHAS, *Hecke algebras and Schur algebras of the symmetric group*, Univ. Lecture Notes, **15**, A.M.S., 1999.
- [15] M. VARAGNOLO AND E. VASSEROT, *On the decomposition matrices of the quantized Schur algebra*, Duke J. Math., **100** (1999), 267–297.

G. J. DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE
 180 QUEEN’S GATE, LONDON SW7 2BZ. U.K.
g.james@ic.ac.uk www.ma.ic.ac.uk/~gdj/

A. M. SCHOOL OF MATHEMATICS F07, UNIVERSITY OF SYDNEY
 SYDNEY N.S.W. 2006. a.mathas@maths.usyd.edu.au
 AUSTRALIA. www.maths.usyd.edu.au/u/mathas/