Toric Fano varieties and birational morphisms

Cinzia Casagrande

Smooth toric Fano varieties are classified up to dimension 4. In dimension 2 there are five toric Del Pezzo surfaces: \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$ and S_i , the blow-up of \mathbb{P}^2 in *i* points, for i = 1, 2, 3. There are 18 toric Fano 3-folds [2, 20] and 124 toric Fano 4-folds [4, 17]. In higher dimensions, little is known about them: many properties that hold in low dimensions, are not known to hold in general.

Let X be a toric Fano variety of dimension n. We recall that in X linear and algebraic equivalence for divisors coincide, so the Picard number ρ_X is the rank of the Picard group of X. By means of some basic combinatorial properties of Σ_X , we show that for every irreducible invariant divisor $D \subset$ X, we have $0 \leq \rho_X - \rho_D \leq 3$. Moreover, if $\rho_X - \rho_D = 3$, then X is a toric S₃-bundle over a lower-dimensional toric Fano variety; if $1 \leq \rho_X - \rho_D \leq 2$ (with some additional hypotheses in the case $\rho_X - \rho_D = 1$), we give an explicit birational description of X. This gives a structure theorem for a large class of toric Fano varieties (theorem 2.4).

The best known bound for the Picard number of X is due to O. Debarre [8]: $\rho_X \leq 2\sqrt{2n^3}$. On the other hand, we know that the maximal Picard number in dimension n is at least 2n for n even, 2n - 1 for n odd. In fact, up to dimension 4, this is exactly the maximal Picard number. We show that the same is true in dimension 5: if X is a toric Fano 5-fold, $\rho_X \leq 9$ (theorem 3.2).

In the second part of the paper, we study equivariant birational morphisms $f: X \to Y$ whose source X is Fano. We give some combinatorial conditions on the possible positions of new generators in Σ_X , and we show that for every irreducible invariant divisor $D \subset X$, we have $\rho_Y - \rho_{f(D)} \leq 3$ (proposition 4.1).

Then we consider two different special cases of this situation: the case where f is a blow-up, and the case where the dimension is 4.

In the case of f a smooth equivariant blow-up, we are particularly interested in studying under which hypotheses Y can be non-projective. In fact, up to now it is not known whether a toric complete, smooth, non-projective variety Y can become Fano after a smooth equivariant blow-up (while this happens in the non-toric context: see [5], example 1.3). We show that if Y is non-projective, the center $A \subset Y$ of the blow-up must have dimension dim $A \ge 3$ and can not be a projective space (theorem 4.4). Hence the first possible case of such a behaviour should be in dimension 6, with dim A = 3.

Finally, we consider the case of f an equivariant birational morphism $f: X \to Y$ with X Fano and dim $X = \dim Y = 4$. We prove a factorization result which was announced in [7]: every equivariant birational morphism

between toric 4-folds, whose source is Fano, is a composite of smooth equivariant blow-ups between toric 4-folds (theorem 5.2). This result is obtained through the complete classification of the possible subdivisions in Σ_X of a maximal cone $\sigma \in \Sigma_Y$ (proposition 5.3).

The approach we use here to study fans of toric varieties is the language of primitive collections and primitive relations, which was introduced by V. Batyrev. This approach is very useful to deal with toric Mori theory and with fans of toric Fano varieties: it led to Batyrev's classification in dimension 4 [4], and was also used by H. Sato in [17] in the same context, giving some interesting factorization properties of equivariant birational maps and morphisms between toric Fano 3-folds and 4-folds.

Acknowledgments. This work is part of my Ph.D. thesis. I wish to thank my advisor, Lucia Caporaso, for constantly guiding me, and for all she taught me in these years.

I am also very grateful to Laurent Bonavero, who first pointed out to me Sato's paper on toric Fano varieties [17], for many helpful discussions and hints, and for the kind hospitality at the Institut Fourier of Grenoble.

Finally, I thank Olivier Debarre for some valuable conversations.

Contents:

1.	Preliminaries: primitive relations and toric Mori theory
2.	Primitive collections of order two and invariant divisors
3.	Picard number of toric Fano varieties15
4.	Equivariant birational morphisms whose source is Fano
5.	Equivariant birational morphisms in dimension 4: analysis of the possible
\mathbf{su}	bdivisions
Re	eferences

1. Preliminaries: primitive relations and toric Mori theory

In this section, we recall the basic definitions and properties of primitive collections and primitive relations, and some results of toric Mori theory. For a more detailed account, we refer to [11, 15] for properties of toric varieties, to [3, 4, 17] for primitive collections and primitive relations and to [16] for toric Mori theory.

An *n*-dimensional toric variety X is described by a finite fan Σ_X in the vector space $N_{\mathbb{Q}} = N \otimes_{\mathbb{Z}} \mathbb{Q}$, where N is a free abelian group of rank n. Throughout the paper we will deal with toric varieties that are smooth and complete; this is equivalent to ask that the support of Σ_X is the whole space $N_{\mathbb{Q}}$ and every cone in Σ_X is generated by a part of a basis of N.

We recall that for each r = 0, ..., n there is a bijection between the cones of dimension r in Σ_X and the orbits of codimension r in X; we'll denote by $V(\sigma)$ the closure of the orbit corresponding to $\sigma \in \Sigma_X$ and $V(x) = V(\langle x \rangle)$ in case of 1-dimensional cones.

For each 1-dimensional cone $\rho \in \Sigma_X$, let $v_{\rho} \in \rho \cap N$ be its primitive generator, and

$$G(\Sigma_X) = \{ v_\rho \, | \, \rho \in \Sigma_X \}$$

the set of all generators of Σ_X .

The following definitions were introduced by V. Batyrev in [3]:

Definition 1.1. A subset $P = \{x_1, \ldots, x_h\} \subseteq G(\Sigma_X)$ is a *primitive collection* for Σ_X if $\langle x_1, \ldots, x_h \rangle \notin \Sigma_X$, and $\langle x_1, \ldots, \check{x}_i, \ldots, x_h \rangle \in \Sigma_X$ for each $i = 1, \ldots, h$. We denote by $PC(\Sigma_X)$ the set of all primitive collections of Σ_X .

Definition 1.2. Let $P = \{x_1, \ldots, x_h\} \subseteq G(\Sigma_X)$ be a primitive collection and let $\sigma_P = \langle y_1, \ldots, y_k \rangle$ be the unique cone in Σ_X such that $x_1 + \cdots + x_h \in$ Rel Int σ_P . Then we get a linear relation

$$x_1 + \dots + x_h - (a_1y_1 + \dots + a_ky_k) = 0$$

with a_i a positive integer for each i = 1, ..., k. This is the *primitive relation* associated to P. The *degree* of P is the integer deg $P = h - \sum_{i=1}^{k} a_i$.

Let $\mathcal{A}_1(X)$ be the group of algebraic 1-cycles on X modulo numerical equivalence, $\mathcal{N}_1(X) = \mathcal{A}_1(X) \otimes_{\mathbb{Z}} \mathbb{Q}$, and $\operatorname{NE}(X) \subset \mathcal{N}_1(X)$ the cone of Mori, generated by classes of effective curves. We recall that there is an exact sequence:

$$0 \longrightarrow \mathcal{A}_1(X) \longrightarrow \mathbb{Z}^{G(\Sigma_X)} \longrightarrow N \longrightarrow 0, \qquad (\heartsuit)$$

where the map $\mathcal{A}_1(X) \to \mathbb{Z}^{G(\Sigma_X)}$ is given by $\gamma \mapsto \{\gamma \cdot V(x)\}_{x \in G(\Sigma_X)}$. Hence the group $\mathcal{A}_1(X)$ is canonically isomorphic to the lattice of integral relations among the elements of $G(\Sigma_X)$: a relation

$$\sum_{x \in G(\Sigma_X)} a_x x = 0, \qquad a_x \in \mathbb{Z},$$

corresponds to a 1-cycle that has intersection a_x with V(x) for all $x \in G(\Sigma_X)$. We will often identify classes in $\mathcal{N}_1(X)$ and the associated relations. We remark that since the canonical class on X is given by $K_X = -\sum_{x \in G(\Sigma_X)} V(x)$, if $\gamma \in \mathcal{N}_1(X)$ corresponds to $\sum_x a_x x = 0$, then $-K_X \cdot \gamma = \sum_x a_x$. In particular, for every primitive collection $P \in PC(\Sigma_X)$, the associated primitive relation defines a class $r(P) \in \mathcal{A}_1(X)$, and $-K_X \cdot r(P) = \deg P$.

For convenience, we will always write primitive relations as

$$x_1 + \dots + x_h = a_1 y_1 + \dots + a_k y_k$$

instead of $x_1 + \cdots + x_h - (a_1y_1 + \cdots + a_ky_k) = 0$, i. e. writing elements with negative coefficient on the right side. This must not be confused with the relation $-(x_1 + \cdots + x_h) + a_1y_1 + \cdots + a_ky_k = 0$, which is the opposite element in $\mathcal{A}_1(X)$.

The cone of effective curves in a smooth complete toric variety has been studied by M. Reid in [16], where he shows that in this case NE(X) is closed and polyhedral, generated by classes of invariant curves ([16], proposition 1.6). Moreover, he gives a precise description of the geometry of the fan around a cone corresponding to an extremal curve:

Theorem 1.3 (Reid [16]). Suppose that X is projective; let R be an extremal ray of NE(X) and γ a primitive element in $R \cap A_1(X)$. Then the relation associated to γ has the form

$$x_1 + \dots + x_h = a_1 y_1 + \dots + a_k y_k,$$

with $a_i \in \mathbb{Z}_{>0}$; $P = \{x_1, \ldots, x_h\}$ is a primitive collection and $\gamma = r(P)$. Moreover, for each cone $\langle z_1, \ldots, z_t \rangle$ such that

$$\{z_1,\ldots,z_t\} \cap \{y_1,\ldots,y_k,x_1,\ldots,x_h\} = \emptyset \text{ and } \langle y_1,\ldots,y_k,z_1,\ldots,z_t \rangle \in \Sigma_X,$$

we have

$$\langle x_1, \ldots, \check{x}_i, \ldots, x_h, y_1, \ldots, y_k, z_1, \ldots, z_t \rangle \in \Sigma_X$$
 for all $i = 1, \ldots, h$.

As a consequence of this result, we have an important description of the cone of effective curves for projective toric varieties:

Proposition 1.4 (Batyrev [3]). Suppose that X is projective: then the cone of effective curves NE(X) is generated by primitive relations.

Proof. It is an easy consequence of theorem 1.3 and lemma 2.2 below. \Box

This gives the following simple characterization of toric Fano varieties:

Proposition 1.5 (Batyrev [4]). Suppose that X is projective. Then X is Fano if and only if all primitive collections in Σ_X have strictly positive degree.

Finally, we recall the fundamental result:

Proposition 1.6. Let X be a toric Fano variety and $\gamma \in NE(X)$. If γ has anticanonical degree one, then it is extremal.

We notice that this result seems well-known by experts in toric geometry (see V. Batyrev [4], theorem 2.3.3 and corollary 2.3.4, and H. Sato [17], the proof of lemma 5.4), but we couldn't find an explicit proof of it written anywhere. Hence, we refer to [6], proposition 4.3.

2. Primitive collections of order two and invariant divisors

Let X be a toric Fano variety of dimension n. All toric varieties are birationally equivalent and we know, after the the weak factorization theorem [14, 1], that any two smooth, complete toric varieties can be obtained one from the other by a sequence of smooth equivariant blow-ups and blowdowns.

Here we give an *explicit* birational description of X when Σ_X has at least one pair of symmetric generators x, -x. This will be linked to some properties of irreducible invariant divisors on X.

Our starting point is the following lemma:

Lemma 2.1. Suppose that $\{x, -x\}$ is a primitive collection in Σ_X . Then for any other primitive collection P containing x, the associated primitive relation is

$$r(P): \quad x+y_1+\cdots+y_h=z_1+\cdots+z_h;$$

moreover, $P' = \{-x, z_1, \ldots, z_h\}$ is also a primitive collection, with relation

 $r(P'): (-x) + z_1 + \dots + z_h = y_1 + \dots + y_h.$

Both P and P' have degree 1, they are extremal, and $2h \leq n$. Furthermore, for any other primitive collection $Q = \{x, u_1, \ldots, u_m\}$ different from P and from $\{x, -x\}$, we have $V(\langle u_1, \ldots, u_m \rangle) \cap V(\langle z_1, \ldots, z_h \rangle) = V(\langle u_1, \ldots, u_m \rangle) \cap V(\langle y_1, \ldots, y_h \rangle) = \emptyset$.

For the proof of lemma 2.1, we need to recall the following condition for effectiveness for a numerical class (see [6], lemma 1.4):

Lemma 2.2. Let X be a smooth, complete toric variety and $\gamma \in \mathcal{A}_1(X)$ given by the relation

$$a_1x_1 + \dots + a_hx_h - (b_1y_1 + \dots + b_ky_k) = 0$$

with $a_i, b_j \in \mathbb{Z}_{>0}$ for each i, j. If $\langle y_1, \ldots, y_k \rangle \in \Sigma_X$, then $\gamma \in NE(X)$.

Proof of lemma 2.1. Suppose that the primitive relation associated to P is

$$x + y_1 + \dots + y_h = a_1 z_1 + \dots + a_k z_k.$$

Since X is Fano, $1 \leq \deg P = 1 + h - \sum_i a_i$, thus $\sum_i a_i \leq h$. On the other hand, consider the class γ in $\mathcal{N}_1(X)$ corresponding to the relation:

$$(-x) + a_1 z_1 + \dots + a_k z_k - (y_1 + \dots + y_h) = 0.$$

Since $\langle y_1, \ldots, y_h \rangle \in \Sigma_X$, lemma 2.2 implies that $\gamma \in \text{NE}(X)$, hence $1 \leq -K_X \cdot \gamma = 1 + \sum_i a_i - h$, namely $\sum_i a_i \geq h$: therefore we get $\sum_i a_i = h$ and

deg $P = \gamma \cdot (-K_X) = 1$. Then r(P) and γ are extremal: in particular, γ is primitive, which means $a_i = 1$ for all i and k = h.

Suppose now that $Q = \{x, u_1, \ldots, u_m\}$ is a primitive collection different from P and from $\{x, -x\}$. We want to show that $\langle u_1, \ldots, u_m, z_1, \ldots, z_h \rangle$ and $\langle u_1, \ldots, u_m, y_1, \ldots, y_h \rangle$ are not in Σ_X . Suppose that $\langle u_1, \ldots, u_m, z_1, \ldots, z_h \rangle \in$ Σ_X . Since r(P) is extremal, by theorem 1.3 we get that either $\langle x, u_1, \ldots, y_m \rangle \in$ Σ_X , or P = Q, in both cases a contradiction. Suppose now that the cone $\langle u_1, \ldots, u_m, y_1, \ldots, y_h \rangle$ is in Σ_X ; then since r(P') is extremal and $-x \notin Q$, theorem 1.3 implies that also $\langle u_1, \ldots, u_m, z_1, \ldots, z_h \rangle$ is in Σ_X , which gives a contradiction. \Box

Fibration in \mathbb{P}^1 : basic construction. We want to explain the basic geometric construction which is an immediate consequence of lemma 2.1. Even if the precise statement of our result will be given in theorem 2.4, we think it is useful for the reader to explain this construction here.

We fix two symmetric generators $x, -x \in G(\Sigma_X)$. If the class x + (-x) = 0 is extremal, then X is a toric \mathbb{P}^1 -bundle over a toric Fano variety W and V(x), V(-x) are the invariant sections of the bundle. In this case x and -x are not contained in any primitive collection different from $\{x, -x\}$. Otherwise, if there exists a primitive collection $P \neq \{x, -x\}$ containing x, the class x + (-x) = 0 is not extremal: a decomposition of this class in NE(X) is given by r(P) + r(P'), where P' is as in lemma 2.1.

For every such P, we define E_P to be the union of the exceptional loci of r(P) and r(P'). By lemma 2.1, for two different P, Q containing x, the loci E_P and E_Q are disjointed. The locus E_P has pure codimension #P-1, hence the divisorial ones are exactly the ones for P of order two.

Fact: outside the loci of the form E_P , X is fibered in \mathbb{P}^1 .

We think of the loci E_P as the "obstructions" to X being a \mathbb{P}^1 -bundle. We can "eliminate" these obstructions by a finite number of flips and blowdowns, getting a smooth projective toric variety which is a toric \mathbb{P}^1 -bundle over $W \simeq V(x)$. More precisely:

• for every primitive collection $P = \{x, y\}$ of order two, we blow-down the divisor V(y) with respect to the extremal relation r(P'): (-x) + z = y;

• for every primitive collection $P = \{x, y_1, \dots, y_h\}$ with $h \ge 2$, we flip the relation $r(P'): (-x) + z_1 + \dots + z_h = y_1 + \dots + y_h$: namely, we blow-up the subvariety $V(\langle y_1, \dots, y_h \rangle) \subset E_P$ and then blow-down the exceptional divisor V(v) with respect to the extremal class $(-x) + z_1 + \dots + z_h = v$; in this way we get a new smooth projective (non-Fano) toric variety with an extremal class given by the primitive relation $y_1 + \dots + y_h = (-x) + z_1 + \dots + z_h$. The flip exchanges an extremal class of anticanonical degree 1 with an extremal class of anticanonical degree -1.

We perform this for all primitive collections containing x (different from $\{x, -x\}$). In the end we get a smooth projective toric variety \overline{X} which is a toric \mathbb{P}^1 -bundle over a smooth toric variety W. The images of the divisors

V(x), V(-x) in \overline{X} are the invariant sections of the bundle; since the divisor V(x) in X is disjoint from the exceptional loci of the flips and of the blowdowns, we have $V(x) \simeq W$.

Geometric description in dimension 4: in dimension 4, when E_P is not divisorial, it is the union of two \mathbb{P}^2 which intersect in one point. Each \mathbb{P}^2 intersects one (and only one) of the divisors V(x), V(-x) along a \mathbb{P}^1 . Blowing-up the \mathbb{P}^2 that touches V(-x), we get an exceptional divisor isomorphic to $\mathbb{P}^2 \times \mathbb{P}^1$ with normal bundle $\mathcal{O}(-1, -1)$. The other \mathbb{P}^2 is blown-up in one point. When we blow-down the exceptional divisor on \mathbb{P}^1 , the image of E_P is S_1 , fibered in \mathbb{P}^1 on the intersections with the images of the two divisors V(x), V(-x). The divisor V(-x) is "flopped".



In general, the loci $V(\langle y_1, \ldots, y_h \rangle)$ and $V(\langle z_1, \ldots, z_h \rangle)$ are toric \mathbb{P}^h bundles over a smooth toric variety T of dimension n - 2h, and intersect along a common invariant section. Fiberwise on T, the same description as in the 4-dimensional case holds (with \mathbb{P}^h instead of \mathbb{P}^2).

Remark: we recall that a toric bundle is locally trivial on Zariski open subsets; see [10], theorem 6.7 on page 246.

Our next lemma concerns primitive collections of order two in Σ_X . Due to the fact that X is Fano, these primitive collections must have a particularly simple primitive relation. In fact, if $\{x, y\}$ is a primitive collection, its degree must be positive by proposition 1.5, hence there are only two possible primitive relations: x + y = 0 or x + y = z. In this last case, the relation has degree 1, so by proposition 1.6 the corresponding class in NE(X) is extremal. Therefore, for each pair of generators $x, y \in G(\Sigma_X)$, only three cases can occur:

- (i) $\langle x, y \rangle \in \Sigma_X$
- (ii) y = -x
- (iii) $z = x + y \in G(\Sigma_X)$, and the relation x + y = z is extremal, hence by theorem 1.3 the cones $\langle x, z \rangle$ and $\langle y, z \rangle$ are in Σ_X .

This has a nice geometric interpretation. We recall that two irreducible invariant divisors V(x), V(y) in X are disjoint if and only if the cone $\langle x, y \rangle$ is not in Σ_X . Hence, if $V(x) \cap V(y) = \emptyset$, there are only two possibilities:

- (a) x + y = 0, hence X is generically fibered in \mathbb{P}^1 as described in the basic construction;
- (b) there exists a third irreducible invariant divisor D that is a toric \mathbb{P}^1 bundle over a smooth toric variety Z; $D \cap V(x)$ and $D \cap V(y)$ are the sections of the \mathbb{P}^1 -bundle. The normal bundle of D restricted to a fiber is $\mathcal{O}_{\mathbb{P}^1}(-1)$, hence there exists a smooth projective variety Y and a smooth equivariant blow-up $X \to Y$ with exceptional divisor D. In Ythe images of V(x) and V(y) intersect along a codimension 2 subvariety which is the center of the blow-up; Y can fail to be Fano.

Lemma 2.3. If Σ_X has two different primitive relations x + y = z and x + w = v, then w = -x - y and v = -y. Therefore there are at most two primitive collections of order 2 and degree 1 containing x, and the associated primitive relations are x + y = (-w) and x + w = (-y).

Proof. Suppose $y \neq w$. The two relations x + y = z and x + w = v are extremal, thus the cones $\langle x, z \rangle$, $\langle y, z \rangle$, $\langle x, v \rangle$, $\langle w, v \rangle$ are in Σ_X .

If $\langle z, v \rangle \in \Sigma_X$, then $\langle z, v, y \rangle$ and $\langle z, v, w \rangle$ are in Σ_X , which is impossible, because z + w = v + y. Therefore $\{z, v\}$ is a primitive collection.

If the associated primitive relation is z + v = 0, we have 2x + y + w = 0. We claim that this is impossible. Indeed, if 2x + y + w = 0 the set $\{y, w\}$ can not be a part of a basis of the lattice, because x is not an integral linear combination of y and w. Since X is smooth, the cone $\langle y, w \rangle$ is not in Σ_X , so $\{y, w\}$ is a primitive collection. Its primitive relation can not be y + w = 0 (it would imply x = 0) nor y + w = t (it would be t = -2x), so its degree can not be positive, which is a contradiction because X is Fano.

Therefore the primitive relation associated to $\{z, v\}$ is z + v = u. We want to show that u = x, which gives the statement.

If u = y, we get 2x + w = 0, which is impossible because the generators are primitive elements in N. The same if u = w.

Suppose that $u \notin \{x, y, w\}$. Since the degree of the relation z + v = u is 1, this class is extremal: so $\langle u, v \rangle$ and $\langle u, z \rangle$ are in Σ_X , and since x + y = z and x + w = v are extremal, all the cones $\langle u, v, x \rangle$, $\langle u, v, w \rangle$, $\langle u, z, x \rangle$, $\langle u, z, y \rangle$ are in Σ_X .



Now consider the pair $\{y, v\}$: since y + 2v = u + w, the cone $\langle y, v \rangle$ can not be in Σ_X , hence $\{y, v\}$ is a primitive collection. It can not be y + v = 0, because it would imply u = x. Moreover, there can not be a primitive relation y + v = k, because this would imply $\langle k, v \rangle \in \Sigma_X$, which is impossible since k + v = u + w. Therefore we get a contradiction.

We are now ready to state the principal result of this section.

Theorem 2.4. For every irreducible invariant divisor $D \subset X$, we have $0 \le \rho_X - \rho_D \le 3$.

2.4.1. Case $\rho_X - \rho_D = 1$ and D = V(x) with $x, -x \in G(\Sigma_X)$.

There exists a sequence

$$X = X_1 \xrightarrow{\varphi_1} X_2 \xrightarrow{\varphi_2} \cdots \xrightarrow{\varphi_r} X_{r+1} \xrightarrow{\psi} \overline{X}$$
 (\diamondsuit)

where for all *i*, X_i is a smooth projective toric variety and φ_i is a toric flip that exchanges an extremal class having anticanonical degree 1 in NE(X_i) with an extremal class having anticanonical degree -1 in NE(X_{i+1}).

In $(\diamondsuit) \psi$ is an isomorphism and $\overline{X} = X_{r+1}$ is a toric \mathbb{P}^1 -bundle over a smooth toric variety $W \simeq D$. The exceptional loci of the flips are disjoint from D, and the image of D in \overline{X} is a section of the \mathbb{P}^1 -bundle. Moreover, D is Fano if and only if r = 0.

2.4.2. Case $\rho_X - \rho_D = 2$.

There exists a sequence as (\diamondsuit) such that one of the following occurs:

2.4.2a. the morphism ψ is an isomorphism and $\overline{X} = X_{r+1}$ is a toric S_2 bundle over a smooth toric Fano variety Z.

There is an irreducible invariant Fano divisor $D' \subset X$ with $\rho_{D'} = \rho_D$ such that the exceptional loci of the flips are disjoint from D'. The image \overline{D}' of D' in \overline{X} is a toric \mathbb{P}^1 -bundle over Z:



and for all $z \in Z \ \overline{D}' \cap \pi^{-1}(z)$ is the exceptional curve in $S_2 \simeq \pi^{-1}(z)$ given by the proper transform of the line in \mathbb{P}^2 through the two blown-up points. 2.4.2b. the variety \overline{X} is a toric \mathbb{P}^1 -bundle over a smooth toric variety $W \simeq D$ and ψ is a smooth equivariant blow-up of a codimension 2 subvariety of \overline{X} .

The exceptional loci of the flips and of the blow-down are disjoint from D, and the image of D in \overline{X} is a section of the \mathbb{P}^1 -bundle. Moreover, D is Fano if and only if r = 0.

2.4.3. Case $\rho_X - \rho_D = 3$.

X is a toric S_3 -bundle over a smooth toric Fano variety Z.

There are six invariant Fano divisors $D_1 = D, D_2, \ldots, D_6$ in X, all having Picard number $\rho_X - 3$, that are toric \mathbb{P}^1 -bundles over Z:



For all $z \in Z$, $D_i \cap \pi^{-1}(z)$ are the six exceptional curves in $S_3 \simeq \pi^{-1}(z)$.

Remark: an irreducible invariant divisor $D = V(x) \subset X$ has always $-K_D$ nef; D is Fano if and only if x is not contained in any primitive collection of degree 1 and order greater than 2, namely if and only if there are no curves $C \subset D$ such that $-K_X \cdot C = D \cdot C = 1$ (see [4], proposition 2.4.4 and corollary 2.4.5). We are interested in finding Fano divisors in X because this would allow to show some properties of X using induction. We will come back to this in the next section.

Tables 1 and 2 on page 11 describe the situation in dimensions 3 and 4. The number in every box is the number of varieties verifying the corresponding case, and the number inside the parentheses in the first row and in the left column is the total number of varieties verifying that case.

For the 3-dimensional case, we refer to [15], pages 90-92. We call F the toric Fano 3-fold with Picard number 5 which is not a product; it is obtained from $\mathbb{P}_{\mathbb{P}^1}(\mathcal{O} \oplus \mathcal{O} \oplus \mathcal{O}(1))$ blowing up the three invariant sections, and is a toric S_3 -bundle on \mathbb{P}^1 . We see in table 1 that the situation is very simple: the 18 toric Fano 3-folds are all either a toric bundle with fiber \mathbb{P}^1 , \mathbb{P}^2 , \mathbb{P}^3 , S_2 or S_3 , or they are obtained from a \mathbb{P}^1 -bundle with a single blow-up (case 2.4.2b, r=0).

In table 2, the letters refer to Batyrev's table of toric Fano 4-folds in [4]. The variety missing in Batyrev's table is Sato's 4-fold, see [17]. We remark first that with only two exceptions, in all cases described by theorem 2.4 there is at most one flip. The two exceptions are the Del Pezzo variety V^4 and the pseudo-Del Pezzo variety \tilde{V}^4 ; these two varieties are special from many points of view: see [19, 9, 17, 8].

We remark also that theorem 2.4 describes 107 of the 124 toric Fano 4-folds. Among the remaining 4-folds, 9 are projective bundles; the classes for which we really lack a description are the last two.

Proof of theorem 2.4. Let D = V(x). We recall that from the exact sequence (\heartsuit) on page 3 we have $\rho_X = \#G(\Sigma_X) - n$ for the Picard number of X, and since 1-dimensional cones in the fan Σ_D are given exactly by

	$\rho = 1$	$\rho = 2$	$\rho = 3$	$\rho = 4$	$\rho = 5$
	(1)	(4)	(7)	(4)	(2)
2.4.1, r = 0		3	5		
2.4.2a, r = 0				4	
2.4.2b, r = 0			2		
2.4.3					$S_3 \times \mathbb{P}^1, F$
$\mathbb{P}^s\text{-bundles}, s>1$	\mathbb{P}^3	1			

Table 1: toric Fano 3-folds.

	a — 1	a — 0	a — 2	$\rho = 4$	- F	a – C	a 7	o — 9
	$\rho = 1$	$\rho = 2$	$\rho = 3$	'	$\rho = 5$	$\rho = 6$	$\rho = 7$	$\rho = 8$
<u> </u>	(1)	(9)	(28)	(47)	(27)	(10)	(1)	(1)
2.4.1, r = 0		4	16	16				
(36)		(B)	(D)	(I,L)				
2.4.1, r = 1			2	4				
(6)			(G_1, G_3)	(M)				
2.4.1, r = 3					1			
(1)					(\widetilde{V}^4)			
2.4.1, r = 6						1		
(1)						(V^4)		
2.4.2a, r = 0				10	17	$S_2 \times S_2$		
(28)				(H)	(Q)			
2.4.2a, r = 1					3			
(3)					(R)			
2.4.2b, r = 0			3	12	1			
(16)			(E)	(I)	(3.4.1)			
2.4.2b, r = 1				2				
(2)				(J)				
2.4.3					4	8	$S_3 \times S_2$	$S_3 \times S_3$
(14)					(K)	(U)		
\mathbb{P}^{s} -bundles,	\mathbb{P}^4	5	3					
s > 1 (9)		(B, C)	(D)					
no symm. gen.			4	1	1			
(6)			(G)	(M_5)	(Sato's)			
no prim. coll.				2				
of order 2 (2)				(Z)				

Table 2: toric Fano 4-folds.

2-dimensional cones of Σ_X containing x, we get:

$$\begin{split} \rho_D &= \#\{y \in G(\Sigma_X) \mid y \neq x \text{ and } \langle y, x \rangle \in \Sigma_X\} - (n-1) \\ &= \#G(\Sigma_X) - \#\{y \in G(\Sigma_X) \mid \{y, x\} \text{ is a primitive collection}\} - n \\ &= \rho_X - \#\{y \in G(\Sigma_X) \mid \{y, x\} \text{ is a primitive collection}\}. \end{split}$$

Hence the difference $\rho_X - \rho_D$ is equal to the number of primitive collections of order two containing x. Let's show that x is contained in at most three primitive collections of order two. By lemma 2.3, there are at most two primitive collections of order two and degree 1 containing x, and the only primitive collection of order two and degree 2 containing x can be $\{x, -x\}$. So we have $\rho_X - \rho_D \leq 3$.

For the second part of the theorem, the idea of the proof is the following. Suppose that we have a pair $x, -x \in G(\Sigma_X)$. Then X is generically fibered in \mathbb{P}^1 as described in the basic construction. Moreover, we know by lemma 2.3 that there are at most two divisorial E_P . We are going to show that if there are exactly two, then there are no other obstructions; in this case there are three pairs of symmetric generators, which give a toric bundle in surfaces S_3 . If there is only one divisorial obstruction, we have two pairs of symmetric generators, and the non-divisorial obstructions to the two pairs are "compatible": hence with a finite number of flips we get a toric bundle in surfaces S_2 .

1) Since $\rho_X - \rho_D = 1$, x not contained in primitive collections of order two different from $\{x, -x\}$. Hence the statement follows from the basic construction: none of the E_P is divisorial.

2) D = V(x) with x contained in two primitive collections of order two. We consider first the case where both collections have degree 1: then by lemma 2.3, the primitive relations are x + y = (-w), x + w = (-y). Thus, by lemma 2.1, we know the following primitive relations of Σ_X :

$$(-y) + (-w) = x,$$
 $y + (-y) = 0,$ $w + (-w) = 0.$

Moreover, we know that $\langle y, w \rangle \in \Sigma_X$, because if $\{y, w\}$ were a primitive collection, the primitive relation should be y + w = (-x), and this would give a third primitive collection $\{x, -x\}$ of order two containing x. Hence in the plane $H \subset N_{\mathbb{Q}}$ spanned by x and y we get a fan of the Del Pezzo surface S_2 as in the following figure:



We remark that the generators x, y, -y, w, -w can not be contained in other primitive collections of order two beyond the ones already given.

Claim 1: if P is a primitive collection such that $\#P \ge 3$ and $y \in P$, then it is also $w \in P$.

Proof of claim 1. Let $P = \{y, z_1, \ldots, z_h\}$. By lemma 2.3, we know that the cone $\langle -y, z_1, \ldots, z_h \rangle$ is in Σ_X . If $w \notin P$, the extremality of the relation (-y) + (-w) = x implies by theorem 1.3 that $\langle -w, z_1, \ldots, z_h \rangle \in \Sigma_X$; again, using the relation x + y = (-w), we get $\langle P \rangle \in \Sigma_X$, a contradiction.

Suppose that $P = \{y, w, z_1, \ldots, z_h\}$ with $h \ge 1$. Clearly $\{z_1, \ldots, z_h\} \cap \{x, -y, -w\} = \emptyset$, otherwise P would contain another primitive collection. We know by lemma 2.1 that deg P = 1 and that Σ_X contains the following extremal relations:

$$y + w + z_1 + \dots + z_h = v_1 + \dots + v_{h+1},$$

$$(-y) + v_1 + \dots + v_{h+1} = w + z_1 + \dots + z_h,$$

$$(-w) + v_1 + \dots + v_{h+1} = y + z_1 + \dots + z_h.$$

Moreover, it is easy to see that also

$$x + v_1 + \dots + v_{h+1} = z_1 + \dots + z_h \tag{(\clubsuit)}$$

is a primitive relation; it has degree 2 and it is not extremal.

Claim 2: if P is a primitive collection such that $\#P \ge 3$ and $x \in P$, then P is obtained as (\clubsuit) , so in particular it has degree 2.

Proof of claim 2. Let $P = \{x, p_1, \ldots, p_r\}$. Then it is easy to see that also $\{-y, p_1, \ldots, p_r\}$ is primitive, so by lemma 2.1 we get extremal relations:

$$(-y) + p_1 + \dots + p_r = q_1 + \dots + q_r, \quad y + q_1 + \dots + q_r = p_1 + \dots + p_r.$$

By claim 1, we can assume $w = q_r$; then r(P) is $x + p_1 + \cdots + p_r = q_1 + \cdots + q_{r-1}$, which is like (\clubsuit) .

Clearly, for symmetry, claim 1 implies that if P is a primitive collection such that $\#P \ge 3$ and $w \in P$, then it is also $y \in P$. Now consider the two pairs y, -y and w, -w. Applying the basic construction, by claim 1 we see that we can eliminate the obstructions for y + (-y) = 0 and for w + (-w) = 0 simultaneously. Namely, we flip all the relations of type $y + w + z_1 + \cdots + z_h = v_1 + \cdots + v_{h+1}$, and get a smooth toric variety \overline{X} . Consider the blow-down $\overline{X} \to Y$ with respect to the relation (-y) + (-w) =x. In Y the classes y + (-y) = 0 and w + (-w) = 0 are both extremal, thus Y is a toric $(\mathbb{P}^1 \times \mathbb{P}^1)$ -bundle over a smooth toric variety Z. The center of the blow-up $\overline{X} \to Y$ is an invariant section, hence \overline{X} is a toric S_2 -bundle over Z. Moreover, D is the exceptional divisor of this blow-up, and the center of the blow-up is isomorphic to Z, so D is a \mathbb{P}^1 -bundle over Z. Claim 2 implies that D is Fano, because in Σ_X there are no primitive collections containing x and having degree 1 and order greater than 2. This implies that Z is Fano (see [18]; in the toric case it is easy to see this directly). Hence we have shown 2.4.2a.

We consider now the case where the two primitive collections containing x are $\{x, -x\}$ and $\{x, y\}$, with relations x + (-x) = 0 and x + y = v.

If $-y \in G(\Sigma_X)$, then we have primitive relations v + (-y) = x and v + (-x) = y. Hence v is contained in two primitive collection of order 2 and degree 1, and moreover $-v \notin G(\Sigma_X)$, because otherwise $\{x, -v\}$ would be a third primitive collection of order two containing x. Then we can consider v instead of x and get again 2.4.2a.

In the same way, if $-y \notin G(\Sigma_X)$ but $-v \in G(\Sigma_X)$, y is contained in two primitive collections of order 2 and degree 1, so we get again 2.4.2a.

Therefore we can assume $-y, -v \notin G(\Sigma_X)$:



In this case we can perform the basic contruction and get 2.4.2b; we remark that there is one single blow-down corresponding to the relation (-x) + v = y. Moreover, we remark that D is Fano if and only if there are no other obstructions, namely if and only if r = 0.

3) D = V(x) with x contained in three primitive collections of order two. By lemma 2.3, the relations are

$$x + (-x) = 0$$
, $x + y = (-w)$, $x + w = (-y)$.

Moreover, applying lemma 2.1 to the pairs $\{x, -x\}$, $\{y, -y\}$, $\{w, -w\}$ we get the following relations:

$$(-x) + (-y) = w$$
 $y + w = (-x)$ $y + (-y) = 0$
 $(-x) + (-w) = y$ $(-y) + (-w) = x$ $w + (-w) = 0$

The six generators x, y, w, -x, -y, -w lie in the same plane $H \subset N_{\mathbb{Q}}$ as in the following figure:



Since all relations having degree 1 are extremal, the cones $\langle x, -y \rangle$, $\langle x, -w \rangle$, $\langle w, -y \rangle$, $\langle -w, y \rangle$, $\langle -x, y \rangle$ and $\langle -x, w \rangle$ are all in Σ_X . These cones give a fan of the Del Pezzo surface S_3 .

We claim that for every primitive collection P in Σ_X different from the nine already given, we have $P \cap \{x, y, w, -x, -y, -w\} = \emptyset$. Indeed, suppose that $P \cap \{x, y, w, -x, -y, -w\} \neq \emptyset$. Since P can not contain another primitive collection, this intersection will have order 1 or 2. If the order is 2, we can suppose $P = \{x, -y, z_1, \ldots, z_h\}$ with $\{z_1, \ldots, z_h\} \cap \{x, y, w, -x, -y, -w\} = \emptyset$; but $\langle x, z_1, \ldots, z_h \rangle \in \Sigma_X$ implies $\langle x, -y, z_1, \ldots, z_h \rangle \in \Sigma_X$, a contradiction. If $P \cap \{x, y, w, -x, -y, -w\}$ has order 1, we can suppose $P = \{x, z_1, \ldots, z_h\}$ with $\{z_1, \ldots, z_h\} \cap \{x, y, w, -x, -y, -w\}$ are suppose $P = \{x, z_1, \ldots, z_h\}$ with $\{z_1, \ldots, z_h\} \cap \{x, y, w, -x, -y, -w\}$ has order 1, we can suppose $P = \{x, z_1, \ldots, z_h\}$ with $\{z_1, \ldots, z_h\} \cap \{x, y, w, -x, -y, -w\} = \emptyset$. Then by lemma 2.1 we have two primitive extremal relations:

$$x + z_1 + \dots + z_h = v_1 + \dots + v_h, \quad (-x) + v_1 + \dots + v_h = z_1 + \dots + z_h.$$

For what preceeds, we know that $\{v_1, \ldots, v_h\} \cap \{x, y, w, -x, -y, -w\} = \emptyset$. Then, using theorem 1.3 and the primitive extremal relations (-y)+(-w) = x, x + y = (-w), (-x) + (-w) = y, we get that $\langle -x, v_1, \ldots, v_h \rangle \in \Sigma_X$, a contradiction.

Applying the basic construction to the two pairs x, -x and y, -y, we see that the only obstructions are the divisorial ones. This means that blowingdown on X the two relations x + y = (-w) and (-x) + (-y) = w, we get a toric $(\mathbb{P}^1 \times \mathbb{P}^1)$ -bundle, hence X itself is a toric S_3 -bundle over a toric Fano variety Z.

We remark that the role of the six generators $\{x, y, w, -x, -y, -w\}$ is perfectly symmetric. The six invariant divisors $D_1 = D, D_2, \ldots, D_6$ corresponding to them are all Fano, hence also Z is Fano.

3. Picard number of toric Fano varieties

A first bound for the Picard number of *n*-dimensional toric Fano varieties was due to V. E. Voskresenskiĭ and A. A. Klyachko [19]:

$$o \le n^2 - n + 1.$$

Recently, O. Debarre [8] has found a better bound:

$$\rho \le 2\sqrt{2n^3}.$$

Moreover, there are some conjectures due to G. Ewald [9] for a linear bound in n.

Looking at tables 1 and 2 on page 11, we see that in dimension 3 and 4 the toric Fano varieties having maximal Picard number are actually S_3 -bundles over a toric Fano variety of lower dimension (and they are very few). Assuming that this holds in all dimensions, applying induction would give the following:

Conjecture 3.1. $\rho \leq 2n$ for n even, $\rho \leq 2n - 1$ for n odd.

This bound would be sharp: just consider $(S_3)^{\frac{n}{2}}$ for n even, $(S_3)^{\frac{n-1}{2}} \times \mathbb{P}^1$ or $(S_3)^{\frac{n-3}{2}} \times F$ for n odd, where F is the toric Fano 3-fold with $\rho_F = 5$ which is not a product (see table 1 on page 11).

In this section we show that conjecture 3.1 is true for n = 5:

Theorem 3.2. Let X be a toric Fano variety of dimension 5. Then $\rho_X \leq 9$. Moreover, $\rho_X = 9$ if and only if $X \simeq S_3 \times S_3 \times \mathbb{P}^1$ or $X \simeq S_3 \times F$.

Since $\rho_X = \#G(\Sigma_X) - n$, giving a bound for the Picard number is equivalent to giving a bound for the number of generators of Σ_X . Let f_i be the number of (i + 1)-dimensional cones in Σ_X : thus f_0 is the number of generators of the fan and f_1 the number of 2-dimensional cones. One has $f_1 \leq {f_0 \choose 2}$, and ${f_0 \choose 2} - f_1$ is exactly the number of primitive collections of order two.

As they count the faces of a simplicial convex polytope, the numbers f_i have to satisfy many relations; we refer to [13] for a survey on this. In particular, by the Dehn-Sommerville equations, $f_{[\frac{n}{2}]}, \ldots, f_{n-1}$ are completely determined by $f_0, \ldots, f_{[\frac{n}{2}]-1}$. In dimension 5 we have (see [13], page 112):

$$\begin{array}{rcl} f_2 &=& 4f_1 - 10f_0 + 20, \\ f_3 &=& 5f_1 - 15f_0 + 30, \\ f_4 &=& 2f_1 - 6f_0 + 12. \end{array}$$

For toric Fano varieties, we have a further relation, due to V. Batyrev:

Theorem 3.3 ([4], **2.3.7**). Let X be an n-dimensional toric Fano variety. Then

$$12f_{n-3} \ge (3n-4)f_{n-2}.$$

In dimension 5 this gives $12f_2 \ge 11f_3$, that together with the Dehn-Sommerville equations gives:

$$7f_1 \le 45(f_0 - 2).$$
 (**(**)

Using the results of the preceeding sections, we can get a new relation under the hypothesis that X does not have many primitive collections of order two:

Lemma 3.4. Let X be an n-dimensional toric Fano variety. Suppose that for any generator $x \in G(\Sigma_X)$, there are at most two primitive collections of order two containing x, and if there are exactly two, they are $\{x, -x\}$ and $\{x, y\}$ with $-y, -x - y \notin G(\Sigma_X)$. Then Σ_X has at most $\frac{3}{4} f_0$ primitive collections of order two, namely:

$$\binom{f_0}{2} - f_1 \le \frac{3}{4}f_0.$$

Proof. The maximal number of primitive collections of order two in X is given by the maximal number of configurations like in theorem 2.4.2b, where we have three primitive collections of order two involving four generators (see figure on page 14). Hence we can have $\left[\frac{f_0}{4}\right]$ of such configurations, giving $3\left[\frac{f_0}{4}\right]$ primitive collections of order two. It is easy to see that considering also the remaining $f_0 - 4\left[\frac{f_0}{4}\right]$ generators, the maximal number of primitive collections of order two is $\frac{3}{4}f_0$.

Proposition 3.5. Let X be as in lemma 3.4, and suppose that dim X = 5. Then $\rho_X \leq 8$.

Proof. By lemma 3.4, we have

$$f_1 \ge \frac{1}{2}f_0(f_0 - 1) - \frac{3}{4}f_0.$$

On the other hand, we have (\spadesuit) : $7f_1 \leq 45(f_0 - 2)$. These relations give $14f_0^2 - 215f_0 + 360 \leq 0$, hence $f_0 \leq 13$ and $\rho_X \leq 8$.

We remark that in this way we can not get a similar result in higher dimensions, because with theorem 3.3 and the Dehn-Sommerville equations we get a relation among $f_0, \ldots, f_{[\frac{n}{2}]-1}$.

Proof of theorem 3.2. Suppose that X has three primitive collections of order two having a common element. Then, by theorem 2.4, X is a toric S₃-bundle over a toric Fano 3-fold: in particular, $\rho_X \leq 9$.

Suppose now that X has two primitive collections of order two having a common element and that the configuration is as described in theorem 2.4.2a. Then X has the same Picard number as a toric S_2 -bundle over a toric Fano 3-fold; hence $\rho_X \leq 8$.

Finally, if X does not fit in the preceeding cases, then proposition 3.5 applies, so $\rho_X \leq 8$.

Therefore if $\rho_X = 9$, X is a toric S_3 -bundle over a toric Fano 3-fold Z with $\rho_Z = 4$. Looking at table 1 on page 11, we see that either $Z \simeq S_3 \times \mathbb{P}^1$ or $Z \simeq F$. In both cases, the only possibility for X being Fano is $X \simeq S_3 \times Z$.

4. Equivariant birational morphisms whose source is Fano

Throughout this section, X will be an n-dimensional toric Fano variety, Y an n-dimensional, smooth, complete toric variety, and $f: X \to Y$ an equivariant birational morphism. The fan of X is a subdivision of the fan of Y, so in particular $G(\Sigma_Y) \subseteq G(\Sigma_X)$. The new generators of X, namely the ones in $G(\Sigma_X) \setminus G(\Sigma_Y)$, correspond to the irreducible components of the exceptional divisor of f. We want to study which conditions are imposed on the possible positions of new generators by the fact that X is Fano. We fix an irreducible invariant divisor $E = V(x) \subset X$ and consider $A = f(E) = V(\eta) \subset Y$, where $\eta \in \Sigma_Y$. Consider $\operatorname{Star}(\eta) = \{\sigma \in \Sigma_Y | \sigma \supseteq \eta\}$ and

$$\Lambda_{\eta} = \operatorname{Supp} \operatorname{Star}(\eta) = \bigcup_{\sigma \supseteq \eta} \sigma \subset N_{\mathbb{Q}}.$$

 Λ_{η} is a closed, connected subset of $N_{\mathbb{Q}}$, and it has non-empty interior: we think of its boundary as a partition of the vector space $N_{\mathbb{Q}}$. We define the sets of generators:

$$\mathcal{G}_{\eta} = \{ z \in G(\Sigma_X) | z \in \operatorname{Int} \Lambda_{\eta} \}, \quad \mathcal{H}_{\eta} = \{ z \in G(\Sigma_X) | z \notin \Lambda_{\eta} \}.$$

 \mathcal{G}_{η} are the generators of Σ_X lying inside Λ_{η} , \mathcal{H}_{η} the generators lying outside: we have $G(\Sigma_X) = \mathcal{G}_{\eta} \cup \mathcal{H}_{\eta} \cup \{z \in G(\Sigma_X) | z \in \partial \Lambda_{\eta}\}.$

Proposition 4.1. In the above setting, we have:

- 1. either $\mathcal{H}_{\eta} = \emptyset$, or $|\mathcal{G}_{\eta} \cup \mathcal{H}_{\eta}| \leq 4$;
- 2. for all $z_1 \in \mathcal{G}_{\eta}$ and $z_2 \in \mathcal{H}_{\eta}$, either $z_1 + z_2 = 0$, or $z_1 + z_2 \in G(\Sigma_X) \cap \partial \Lambda_{\eta}$;
- 3. if $|\mathcal{G}_{\eta} \cup \mathcal{H}_{\eta}| = 4$, then X is a toric S₃-bundle over a toric Fano variety.

Remarks:

1) if $\mathcal{H}_{\eta} \neq \emptyset$, the majority of the new generators (the ones in $G(\Sigma_X) \setminus G(\Sigma_Y)$) must lie on the boundary of Λ_{η} .

2) All generators in \mathcal{G}_{η} are new generators: indeed, they forcely lie inside some cone of Σ_Y . So they all correspond to components of the exceptional divisor. Instead, the generators in \mathcal{H}_{η} can either be new generators, or they can be generators of Σ_Y which do not belong to $\operatorname{Star}(\eta)$; such generators are exactly $\rho_Y - \rho_A$, so we get $\rho_Y - \rho_A \leq |\mathcal{H}_{\eta}| \leq 3$.

3) The sets \mathcal{G}_{η} and \mathcal{H}_{η} can be seen in a more geometric way: let $z \in G(\Sigma_X)$ and $\tau \in \Sigma_Y$ such that $z \in \operatorname{RelInt} \tau$. Then $f(V(z)) = V(\tau)$ in Y, and we have:

$$z \in \operatorname{Int} \Lambda_{\eta} \quad \Leftrightarrow \quad \tau \supseteq \eta \quad \Leftrightarrow \quad f(V(z)) \subseteq A;$$
$$z \notin \Lambda_{\eta} \quad \Leftrightarrow \quad \tau + \eta \notin \Sigma_{Y} \quad \Leftrightarrow \quad f(V(z)) \cap A = \emptyset.$$

Therefore we have:

$$\mathcal{G}_{\eta} = \{ z \in G(\Sigma_X) | f(V(z)) \subseteq A \}, \quad \mathcal{H}_{\eta} = \{ z \in G(\Sigma_X) | f(V(z)) \cap A = \emptyset \}.$$

Corollary 4.2. Let X be Fano, Y a smooth complete toric variety and $f: X \to Y$ an equivariant birational morphism. Then for any irreducible invariant divisor $E \subset X$, we have $\rho_Y - \rho_{f(E)} \leq 3$.

If f(E) is a point $p \in Y$, then $\rho_Y \leq 3$, and E is the only component of the exceptional divisor contracted to p, unless $Y \simeq \mathbb{P}^n$. In the case $Y \simeq \mathbb{P}^n$, there can be at most another component contracted to p.

In particular, if one component of the exceptional divisor is contracted to a curve, then $\rho_Y \leq 4$.

The proof of proposition 4.1 will be an easy consequence of the following lemma:

Lemma 4.3. Let X be Fano and suppose that $\mathcal{G}, \mathcal{H} \subset G(\Sigma_X)$ are two nonempty, disjoint sets of generators such that for all $x \in \mathcal{G}$ and $y \in \mathcal{H}$, the set $\{x, y\}$ is a primitive collection in Σ_X . Then $|\mathcal{G} \cup \mathcal{H}| \leq 4$; moreover, if $|\mathcal{G} \cup \mathcal{H}| = 4$, X is a toric S₃-bundle over a toric Fano variety, and one of the following two cases occurs:

(i)
$$\mathcal{G} = \{x_1, x_2, x_3\}, \mathcal{H} = \{-x_1\}$$
 with primitive relations $(-x_1) + x_1 = 0, (-x_1) + x_2 = (-x_3)$ and $(-x_1) + x_3 = (-x_2);$

(*ii*) $\mathcal{G} = \{x_1, x_2\}, \mathcal{H} = \{-x_1, -x_2\}$ with primitive relations $(-x_1) + x_1 = 0, (-x_2) + x_2 = 0, (-x_1) + x_2 = v, x_1 + (-x_2) = (-v).$

Remark: in particular, this implies that the union of m irreducible invariant divisors on X is always a connected set for $m \ge 5$.

Proof of lemma 4.3. We already know from theorem 2.4 that $|\mathcal{G}| \leq 3$ and $|\mathcal{H}| \leq 3$: we have to show that if $|\mathcal{G}| = 3$, then $|\mathcal{H}| = 1$ (thus by symmetry we also have that $|\mathcal{H}| = 3$ implies $|\mathcal{G}| = 1$).

We set $\mathcal{G} = \{x_1, x_2, x_3\}$, and let $y \in \mathcal{H}$. Since $\{y, x_i\}$ is a primitive collection for i = 1, 2, 3, it must be $y = -x_i$ for some i. Therefore $\mathcal{H} \subseteq \{-x_1, -x_2, -x_3\}$. Suppose now that $-x_1 \in \mathcal{H}$: then by lemma 2.3, we have two primitive relations $(-x_1) + x_2 = (-x_3)$ and $(-x_1) + x_3 = (-x_2)$; having degree 1, these relations are extremal, hence the cones $\langle -x_3, x_2 \rangle$ and $\langle -x_2, x_3 \rangle$ are in Σ_X , so $\mathcal{H} = \{-x_1\}$. This gives case (i), and since $-x_1$ is contained in three primitive collection of order two, X is a toric S₃-bundle over a toric Fano variety by theorem 2.4.

It remains to show that when $|\mathcal{G}| = |\mathcal{H}| = 2$ the primitive relations are as described in case (*ii*). Let $\mathcal{G} = \{x_1, x_2\}$: if there exists $y \in \mathcal{H}$ such that $y \neq -x_1$ and $y \neq -x_2$, then we get primitive relations $x_1 + y = (-x_2)$, $x_2+y = (-x_1)$; in particular we see that $y = -x_1-x_2$ is uniquely determined by x_1 and x_2 , and the cones $\langle x_1, -x_2 \rangle$, $\langle -x_1, x_2 \rangle$ are in Σ_X : hence $\mathcal{H} = \{y\}$. Therefore, the only possibility if $|\mathcal{H}| = 2$ is that $\mathcal{H} = \{-x_1, -x_2\}$. Clearly in this case we have primitive relations as in (*ii*). Moreover, since $x_1 + v = x_2$, also $\{x_1, v\}$ is a primitive collection: then x_1 is contained in three primitive collections of order 2, and again by theorem 2.4 X is a toric S_3 -bundle over a toric Fano variety.

Proof of proposition 4.1. We remark that \mathcal{G}_{η} is non-empty, because $f(V(x)) = V(\eta)$ means that $x \in \text{Rel Int } \eta$: then x is in the interior of Λ_{η} , so $x \in \mathcal{G}_{\eta}$.

Let $y \in \mathcal{G}_{\eta}$ and $z \in \mathcal{H}_{\eta}$: the cone $\langle y, z \rangle$ crosses the boundary of Λ_{η} , which is composed of cones in Σ_Y . Since Σ_X is a subdivision of Σ_Y , it must be $\langle y, z \rangle \notin \Sigma_X$. Hence $\{y, z\}$ is a primitive collection for Σ_X .

Therefore, either $\mathcal{H}_{\eta} = \emptyset$, or lemma 4.3 applies to \mathcal{G}_{η} and \mathcal{H}_{η} , and we get (1) and (3). Also (2) is clear, because if we have a primitive relation $z_1 + z_2 = v$, then it has degree 1, so it is extremal, and the cones $\langle z_1, v \rangle$ and $\langle z_2, v \rangle$ are in Σ_X . Since they can not cross the boundary of Λ_{η} , it must be $v \in \partial \Lambda_{\eta}$.

Proof of corollary 4.2. Let $p = V(\eta) \in Y$. Since dim $\eta = n$, Σ_Y must have at least one generator outside η , hence $\mathcal{H}_{\eta} \neq \emptyset$; so $|\mathcal{G}_{\eta} \cup \mathcal{H}_{\eta}| \leq 4$ by theorem 4.1. Notice that $|\mathcal{G}_{\eta}|$ is exactly the number of the components of the exceptional divisor that are contracted to p.

We claim that the cases $|\mathcal{G}_{\eta}| = 3$, $|\mathcal{H}_{\eta}| = 1$ and $|\mathcal{G}_{\eta}| = |\mathcal{H}_{\eta}| = 2$ can not happen. Indeed, since dim $\eta = n$, we have that Star $\eta = \eta$ is strictly convex. Now, looking carefully at the primitive relations given by (i) and (ii) of lemma 4.3, one can easily see that if $|\mathcal{G}_{\eta}| = 3$, $|\mathcal{H}_{\eta}| = 1$ or $|\mathcal{G}_{\eta}| = |\mathcal{H}_{\eta}| = 2$, then there exists some generator v such that $v, -v \in \partial\eta$, which is impossible.

Therefore, we have in any case that $|\mathcal{G}_{\eta}| \leq 2$, and if $|\mathcal{G}_{\eta}| = 2$ then it must be $|\mathcal{H}_{\eta}| = 1$, which clearly implies $Y \simeq \mathbb{P}^n$.

From now on we will consider the case where the morphism $f: X \to Y$ is a smooth equivariant blow-up. We want to apply the preceeding results to study under which conditions it is possible that Y is non-projective. In fact, as far as we know, there are no known examples of a non-projective toric variety Y that becomes Fano after a smooth equivariant blow-up $X \to Y$.

Let $A \subset Y$ be the center of the blow-up. If A is a point, Y is always projective; if dim A = n - 2, a line in a non-trivial fiber of the blow-up has anticanonical degree 1, so it is extremal by proposition 1.6, and again Y is projective.

The result we obtain is the following:

Theorem 4.4. Let Y be a smooth complete non-projective toric variety of dimension $n = \dim Y \ge 4$, and $A \subset Y$ an irreducible invariant subvariety. If the blow-up of Y along A is Fano, then $3 \le \dim A \le n-3$, $n \ge 6$, $\rho_Y - \rho_A \le 2$ and $\rho_A > 1$.

Theorem 4.4 will be a consequence of proposition 4.5 and proposition 4.7.

Proposition 4.5. Let X be an n-dimensional toric Fano variety, $f: X \rightarrow Y$ a smooth equivariant blow-up and $E \subset X$ the exceptional divisor of f. If

 $\rho_X - \rho_E = 3$, or if $\rho_X - \rho_E = 2$ and X is a toric S₂-bundle $\pi: X \to Z$ with $\pi_{|E}: E \to Z$ a \mathbb{P}^1 -bundle, then Y is projective.

Proof. We will show that Y is projective when $\rho_X - \rho_E = 3$; in the same way one can prove the other case. By 2.4.1, there is a toric S_3 -bundle $\pi: X \to Z$ over a toric Fano variety Z. Let E = V(x): keeping the notation of theorem 2.4, we know that in Σ_X there are the nine primitive relations:

$$\begin{aligned} x + (-x) &= 0 \quad y + (-y) = 0 \quad w + (-w) = 0 \quad x + y = (-w) \quad x + w = (-y) \\ (-x) + (-y) &= w \quad y + w = (-x) \quad (-x) + (-w) = y \quad (-y) + (-w) = x. \end{aligned}$$

Let $\mathcal{K} \subset PC(\Sigma_X)$ be the set of these nine primitive collections; we also know that there is a bijection between $PC(\Sigma_X) \setminus \mathcal{K}$ and $PC(\Sigma_Z)$, induced by the surjective morphism $\pi_* \colon NE(X) \to NE(Z)$.

We claim that for every primitive collection $P \in PC(\Sigma_X) \setminus \mathcal{K}$, r(P)is extremal in NE(X) if and only if $\pi_*(r(P))$ is extremal in NE(Z). Indeed, clearly if r(P) decomposes as a sum of other primitive collections, the decomposition holds for $\pi_*(r(P))$, and conversely.

Now consider the class $\omega = r(Q)$ coming from the blow-up f; Y is projective if and only if ω is extremal in NE(X). If ω is in \mathcal{K} , then it has degree 1, hence it is extremal. So we can suppose that $\omega \in PC(\Sigma_X) \setminus \mathcal{K}$; we want to show that $\pi_*(\omega)$ is extremal in NE(Z). We remark that $\omega_{|E}$ is extremal in E, because it corresponds to a \mathbb{P}^r -bundle on the center of the blow-up f. But this implies that $\pi_*(\omega)$ is extremal in Z, because it is the image of $\omega_{|E}$ under the restriction $\pi_{|E} \colon E \to Z$, which is a \mathbb{P}^1 -bundle. \Box

In the case dim $A \leq 2$ it is sufficient to suppose that Y is not Fano to get strong conditions on A. This result is due to J. Wiśniewski [21], and it is particularly interesting because it holds in a general (non-toric) setting:

Theorem 4.6 ([21], 3.5 and 3.6). Let X be a Fano manifold of dimension $n \ge 4$ and $f: X \to Y$ the blow-up of a smooth variety Y along a smooth subvariety A, with dim $A \le 2$; let E be the exceptional divisor of f. Suppose that Y is not Fano. Then one of the following cases occurs:

- (i) $A \simeq \mathbb{P}^1$, $N_{A/Y} \simeq \mathcal{O}(-1)^{\oplus (n-1)}$ and consequently $E \simeq \mathbb{P}^1 \times \mathbb{P}^{n-2}$;
- (ii) $A \simeq \mathbb{P}^2$ and $N_{A/Y}$ is either $\mathcal{O}(-1)^{\oplus (n-2)}$ and $n \geq 5$, or $\mathcal{O}(-2) \oplus \mathcal{O}(-1)^{\oplus (n-3)}$, or $T\mathbb{P}^2(-3) \oplus \mathcal{O}(-1)^{\oplus (n-4)}$, where $T\mathbb{P}^2$ is the tangent bundle to \mathbb{P}^2 ;
- (iii) $A \simeq \mathbb{P}^1 \times \mathbb{P}^1$ and $N_{A/Y} \simeq \mathcal{O}(-1, -1)^{\oplus (n-2)}$ and consequently $E \simeq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^{n-3}$;
- (iv) A is a surface with a \mathbb{P}^1 -bundle structure $A \to C$ over a smooth curve C. In this case X admits another blow-down $X \to Y_1$ which contracts the divisor E to a smooth codimension 2 subvariety of Y_1 .

Proposition 4.7. Let $X \to Y$ be the blow-up of a smooth complete non-Fano toric variety Y along a smooth invariant surface $A \subset Y$; let $E \subset X$ be the exceptional divisor. Suppose that X is Fano, $\rho_X = 5$ and $\rho_A = 2$. Then Y is projective.

Moreover, $A \simeq \mathbb{P}^1 \times \mathbb{P}^1$ or $A \simeq S_1$, $E \simeq A \times \mathbb{P}^{n-3}$, and X is either a toric S_2 -bundle over $\mathbb{P}^1 \times \mathbb{P}^{n-3}$, or the blow-up of a toric \mathbb{P}^1 -bundle over $A \times \mathbb{P}^{n-3}$ along a codimension 2 smooth invariant subvariety contained in a section.

Proof. We know, by theorem 4.6, that $A \simeq \mathbb{F}_a$ with $a \in \mathbb{Z}_{\geq 0}$. We treat the case a > 0; the case a = 0 is very similar and easier, because the normal bundle $N_{E/X}$ is known.

Let E = V(x): we know by theorem 4.6 that E has a \mathbb{P}^1 -bundle structure over an (n-2)-dimensional toric variety. Hence in Σ_E we have primitive relations:

$$u_0 + u_1 = 0$$
, $v_0 + v_1 = au_0$, $z_0 + \dots + z_{n-3} = 0$,

while in Σ_X we know the relations $u_0 + u_1 = x$ and $z_0 + \cdots + z_{n-3} = x$. Moreover, either $\{v_0, v_1\}$ is a primitive collection in Σ_X , or $\{x, v_0, v_1\}$ is. Since $-K_E$ is nef, it must be a = 1 or a = 2.

We have $\rho_E = 3$ and $\rho_X = 5$, thus x is contained in exactly two primitive collections of order two $\{x, w_0\}$ and $\{x, w_1\}$, and

$$G(\Sigma_X) = \{x, u_0, u_1, v_0, v_1, w_0, w_1, z_0, \dots, z_{n-3}\}.$$

We examine the possible primitive relations of $\{x, w_0\}$ and $\{x, w_1\}$.

1) $x + w_0 = 0$, $x + w_1 = z_0$.



We remark that if $\{x, v_0, v_1\}$ is a primitive collection in Σ_X , then by proposition 2.1 the associated relation should be $x + v_0 + v_1 = p_0 + p_1$. Since the degree is 1, we would get $\langle x, p_0, p_1 \rangle \in \Sigma_X$, hence this would give in Σ_E the primitive relation $v_0 + v_1 = p_0 + p_1$, a contradiction. Hence $\{v_0, v_1\}$ is a primitive collection in Σ_X , with relation $v_0 + v_1 = au_0$, so a = 1. Then we get primitive relations:

$$z_0 + w_0 = w_1$$
, $w_1 + z_1 + \dots + z_{n-3} = 0$, $v_0 + v_1 = u_0$.

We easily see that X is the blow-up along $V(\langle z_0, w_0 \rangle)$ of the variety X' given by the primitive relations:

 $x + w_0 = 0$, $v_0 + v_1 = u_0$, $u_0 + u_1 = x$, $z_0 + \dots + z_{n-3} = x$.

X' is a \mathbb{P}^1 -bundle over $\mathbb{P}^{n-3} \times S_1$. Clearly the relation $z_0 + \cdots + z_{n-3} = x$ is extremal in NE(X), thus Y is projective.

2) $x + w_0 = 0$, $x + w_1 = v_1$. Exactly as in the preceding case we see that a = 1, $v_0 + v_1 = u_0$ is a primitive relation in Σ_X and X is the blow-up of X' along $V(\langle v_1, w_0 \rangle)$.

3) $x + w_0 = 0$, $x + w_1 = u_1$. As in case 1), we have that a = 1 and $v_0 + v_1 = u_0$ is a primitive relation in Σ_X . Moreover we know the primitive relations $u_0 + w_1 = 0$ and $u_1 + w_0 = w_1$. We are in the hypothesis of 2.4.2a, and x can not appear in other primitive collections, because this would give other primitive collections in E: hence X is toric S_2 -bundle over $\mathbb{P}^{n-3} \times \mathbb{P}^1$. In particular, Y is projective, by proposition 4.5.

4) $x + w_0 = 0$, $x + w_1 = u_0$. This is exactly as the preceeding case, except for the fact that the primitive relation associated to $\{v_0, v_1\}$ can be either $v_0 + v_1 = u_0$ or $v_0 + v_1 = w_1$. In both cases X is a toric S₂-bundle over $\mathbb{P}^{n-3} \times \mathbb{P}^1$ and Y is projective.

5)
$$x + w_0 = u_0$$
, $x + w_1 = u_1$.



This is the only case with $-x \notin G(\Sigma_X)$, and we are again in the hypothesis of 2.4.2a. This implies that E is Fano, hence a = 1. If $\{x, v_0, v_1\}$ is a primitive collection in Σ_X , the primitive relation could be $x + v_0 + v_1 = u_0$ or $x + v_0 + v_1 = w_0$, but both cases are impossible: indeed, we have seen in the proof of 2.4.2a that any primitive relation involving x with coefficient 1 can not contain any other generator of the fan of S_2 . Therefore we get the same as case 4), X is toric S_2 -bundle over $\mathbb{P}^{n-3} \times \mathbb{P}^1$ and Y is projective. \Box

Proof of theorem 4.4. Since Y is non-projective, it must be $\rho_Y \ge 4$ (see [12]). Moreover, by proposition 4.5 we have $\rho_Y - \rho_A \le 2$, hence $\rho_A \ge 2$. So it clearly can not be dim A = 1. If dim A = 2, the possible cases are described by theorem 4.6. Again, it can not be $A \simeq \mathbb{P}^2$; in the other cases we have $\rho_A = 2$ and $\rho_X = 5$: then proposition 4.7 applies and Y is projective. Hence, it must be dim $A \ge 3$.

5. Equivariant birational morphisms in dimension 4: analysis of the possible subdivisions

In [17], H. Sato shows that every equivariant birational morphism between toric Fano 3-folds is a composite of smooth equivariant blow-ups between toric Fano 3-folds. We recall that in general an equivariant birational morphism f between smooth complete toric varieties doesn't admit a factorization in a sequence of smooth blow-ups, if the dimension is greater than two (f will have a factorization as a sequence of smooth equivariant blow-ups and blow-downs; see [14] and [1]). In case of an equivariant birational morphism between toric Fano 3-folds, not only is there a factorization in smooth equivariant blow-ups, but also all the intermediate 3-folds are Fano. Such a result is proven in two steps:

1) if the source is Fano, then the morphism is a composite of smooth equivariant blow-ups;

2) if also the target is Fano, then all the intermediate 3-folds are Fano.

In [7] the author has shown, with an explicit counterexample, that the same result does not hold in dimension 4:

Proposition 5.1 ([7], 3.1). There exist two toric Fano 4-folds X and Y and a birational equivariant morphism $f: X \to Y$, such that f doesn't admit a decomposition in smooth equivariant blow-ups between toric Fano 4-folds.

In this section we show that at least 1) holds in dimension 4:

Theorem 5.2. Let X, Y be two toric 4-folds and $f: X \to Y$ an equivariant birational morphism. Suppose that X is Fano. Then f factorizes as a sequence of smooth equivariant blow-ups between toric 4-folds.

The proof of theorem 5.2 is analogous to Sato's proof in dimension 3: we fix a maximal cone $\sigma \in \Sigma_Y$ and study the possible subdivisions of σ in Σ_X , applying the results of the preceeding section about the possible positions of new generators. Geometrically, this means that we are considering the restriction of f to $f^{-1}(\mathcal{U}_{\sigma})$, where $\mathcal{U}_{\sigma} \subset Y$ is an invariant open subset isomorphic to \mathbb{A}^n . The result we obtain is the following:

Proposition 5.3. Let X and Y be two toric 4-folds and $f: X \to Y$ an equivariant birational morphism. Suppose that X is Fano and let $\sigma = \langle y_1, y_2, y_3, y_4 \rangle \in \Sigma_Y$. Then the possible subdivisions of σ in Σ_X are 17 and are all given by a sequence of at most 3 star-subdivisions. These 17 subdivisions are described in the following list, where we give for each subdivision: the sequence of cones whose star-subdivisions (in the given order) give the subdivision; the primitive relations inside the subdivision. The last four cases can occur only if $Y \simeq \mathbb{P}^4$.

- 1) $\sigma \in \Sigma_X$, i. e. σ is not subdivided.
- 2) $\langle y_1, y_2 \rangle$; the only primitive relation is $y_1 + y_2 = x_1$.

- 3) $\langle y_1, y_2, y_3 \rangle$; the only primitive relation is $y_1 + y_2 + y_3 = x_1$.
- 4) (y_1, y_2, y_3, y_4) ; the only primitive relation is $y_1 + y_2 + y_3 + y_4 = x_1$.
- 5) $\langle y_1, y_2 \rangle$, $\langle y_3, y_4 \rangle$; primitive relations: $y_1 + y_2 = x_1$, $y_3 + y_4 = x_2$.
- 6) $\langle y_1, y_2, y_3 \rangle$, $\langle y_1, x_1 \rangle$; primitive relations: $y_1 + y_2 + y_3 = x_1$, $y_1 + x_1 = x_2$, $y_2 + y_3 + x_2 = 2x_1$.

7) $\langle y_1, y_2, y_3 \rangle$, $\langle y_1, y_4 \rangle$; primitive relations: $y_1 + y_2 + y_3 = x_1$, $y_1 + y_4 = x_2$, $y_2 + y_3 + x_2 = x_1 + y_4$.

- 8) $\langle y_1, y_2, y_3 \rangle$, $\langle y_1, y_2 \rangle$; primitive relations: $y_1 + y_2 = x_2$, $x_2 + y_3 = x_1$.
- 9) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2, y_3 \rangle$; primitive relations: $y_1 + y_2 + y_3 = x_2$, $x_2 + y_4 = x_1$.

10) $\langle y_1, y_2, y_3 \rangle$, $\langle y_2, y_3, y_4 \rangle$, $\langle y_4, x_2 \rangle$; primitive relations: $y_1 + y_2 + y_3 = x_2$, $y_2 + y_3 + y_4 = x_3$, $y_1 + x_3 = x_1$, $y_4 + x_2 = x_1$, $y_2 + y_3 + x_1 = x_2 + x_3$.

11) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2 \rangle$; primitive relations: $y_1 + y_2 = x_2, y_3 + y_4 + x_2 = x_1$.

12) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2 \rangle$, $\langle y_3, y_4 \rangle$; primitive relations: $y_1+y_2 = x_2, y_3+y_4 = x_3, x_2 + x_3 = x_1$.

13) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2 \rangle$, $\langle x_2, y_3 \rangle$; primitive relations: $y_1 + y_2 = x_2$, $y_3 + x_2 = x_3$, $y_4 + x_3 = x_1$.

14) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, x_1 \rangle$; primitive relations: $y_1 + y_2 + y_3 + y_4 = x_1$, $y_1 + x_1 = x_2$, $x_2 + y_2 + y_3 + y_4 = 2x_1$.

15) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2, y_3 \rangle$, $\langle y_1, x_1 \rangle$; primitive relations: $y_1 + y_2 + y_3 = x_2$, $x_2 + y_4 = x_1$, $y_1 + x_1 = x_3$, $y_2 + y_3 + x_3 = x_1 + x_2$.

16) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2 \rangle$, $\langle y_3, x_1 \rangle$; primitive relations: $y_1 + y_2 = x_2$, $y_3 + y_4 + x_2 = x_1$, $y_3 + x_1 = x_3$, $y_4 + x_2 + x_3 = 2x_1$.

17) $\langle y_1, y_2, y_3, y_4 \rangle$, $\langle y_1, y_2 \rangle$, $\langle x_1, x_2 \rangle$; primitive relations: $y_1 + y_2 = x_2$, $y_3 + y_4 + x_2 = x_1$, $x_1 + x_2 = x_3$, $y_3 + y_4 + x_3 = 2x_1$.

































Remark: in the case $Y\simeq \mathbb{P}^4,$ it is easy to see that all these subdivisions can actually occur.

The proofs of proposition 5.3 and theorem 5.2 will take all the rest of

the section.

This first lemma is lemma 5.4 in [17]; it is a result of linear algebra, we give the proof for completeness:

Lemma 5.4. Let X, Y be two complete, smooth, n-dimensional toric varieties, and $f: X \to Y$ an equivariant birational morphism. Fix a cone $\tau = \langle y_1, \ldots, y_m \rangle$ in Σ_Y and let $\tilde{\tau} = \langle x_1, \ldots, x_k \rangle$ be the unique cone in Σ_X such that $y_1 + \cdots + y_m \in \text{Rel Int}(\tilde{\tau})$. Then there exist a partition J_1, \ldots, J_k of the set $\{1, \ldots, m\}$ such that

$$x_i = \sum_{j \in J_i} y_j \qquad \forall i = 1, \dots, k$$

Proof. Since X is complete, the point $y_1 + \cdots + y_m$ actually lies in some cone of Σ_X . Moreover, Σ_X is a subdivision of Σ_Y , thus it must be $\tilde{\tau} \subseteq \tau$. Then for all $i = 1, \ldots, k$ we have

$$x_i = a_{i1}y_1 + \dots + a_{im}y_m$$

with $a_{ij} \in \mathbb{Z}_{\geq 0}$. On the other hand, since $y_1 + \cdots + y_m \in \operatorname{RelInt} \tilde{\tau}$, there exist $b_1, \ldots, b_k \in \mathbb{Z}_{>0}$ such that

$$y_1 + \dots + y_m = \sum_{i=1}^k b_i x_i = \sum_{i=1}^k b_i \left(\sum_{h=1}^m a_{ih} y_h\right) = \sum_{h=1}^m \left(\sum_{i=1}^k b_i a_{ih}\right) y_h.$$

Since Y is smooth, the set $\{y_1, \ldots, y_m\}$ is a part of a basis of the lattice: hence we have

$$\sum_{i=1}^{k} b_i a_{ih} = 1 \quad \forall h = 1, \dots, m.$$

Since $b_i \in \mathbb{Z}_{>0}$ and $a_{ij} \in \mathbb{Z}_{\geq 0}$, for each $h = 1, \ldots, m$ there is an index $i_h = 1, \ldots, k$ such that $b_{i_h} = a_{i_h h} = 1$ and $a_{i_h} = 0$ for all $i \neq i_h$. This means that y_h appears only in x_{i_h} , with coefficient equal to 1.

In order to study the possible subdivisions, we need to know how nonextremal relations decompose in NE(X). Since in the general the coefficients of such a decomposition are not integral, but rational, we need to introduce a new type of classes in NE(X), which will allow us to work with integral decompositions. We refer to [6] for a more detailed account on this subject.

Definition 5.5. Let Z be an n-dimensional, complete, smooth toric variety, and $\gamma \in NE(Z)$. We say that γ is *contractible* if the two following conditions hold:

(i) γ is primitive, i. e. there exists a primitive collection P in Σ_Z such that $\gamma = r(P)$. Therefore we have

$$\gamma: \quad x_1 + \dots + x_h = a_1 y_1 + \dots + a_k y_k,$$

with $k \ge 0$, $a_i \in \mathbb{Z}_{>0}$ for all i and $\langle y_1, \ldots, y_k \rangle \in \Sigma_Z$.

(ii) if $\nu = \langle z_1, \dots, z_t \rangle$ is such that $\langle y_1, \dots, y_k \rangle + \nu \in \Sigma_Z$ and $\{z_1, \dots, z_t\} \cap \{x_1, \dots, x_h, y_1, \dots, y_k\} = \emptyset$, then

$$\langle x_1, \dots, \check{x}_i, \dots, x_h, y_1, \dots, y_k \rangle + \nu \in \Sigma_Z$$
 for all $i = 1, \dots, h$.

Comparing with theorem 1.3, we see that this is exactly Reid's description of the geometry of the fan around the walls corresponding to an extremal class. This description is actually equivalent to the existence of an equivariant morphism $\varphi_{\gamma} \colon Z \to Z_{\gamma}$ with connected fibers such that for all curves $C \subset Z$

$$\varphi_{\gamma}(C) = \{pt\} \quad \Longleftrightarrow \quad [C] \in \mathbb{Q}_{\geq 0}\gamma.$$

In projective varieties, contractible classes have the following interesting property:

Theorem 5.6 ([6]). Let Z be an n-dimensional, smooth, projective toric variety. Then for every $\eta \in \mathcal{A}_1(Z) \cap NE(Z)$ there is a decomposition

$$\eta = m_1 \gamma_1 + \dots + m_r \gamma_r$$

with γ_i contractible and $m_i \in \mathbb{Z}_{>0}$ for all $i = 1, \ldots, r$.

Lemma 5.7. Let X be a Fano 4-fold and $\{x_1, x_2, x_3\}$ a primitive collection in X with relation $x_1 + x_2 + x_3 = x$. Suppose that the relation is not contractible. Then there are three possible kinds of decomposition as a sum of primitive relations of degree 1:

- (A) $x_1 + x = y, y + x_2 + x_3 = 2x$. Then the cones $\langle x, x_2, x_3 \rangle$, $\langle x, x_2, y \rangle$, $\langle x, x_3, y \rangle$, $\langle y, x_1, x_2 \rangle$, $\langle y, x_1, x_3 \rangle$ are in Σ_X .
- (B) $x_1 + w = z$, $z + x_2 + x_3 = w + x$. Then the cones $\langle w, x, x_2, x_3 \rangle$, $\langle w, x, z, x_2 \rangle$, $\langle w, x, z, x_3 \rangle$, $\langle z, x, x_1, x_2 \rangle$, $\langle z, x, x_1, x_3 \rangle$ are in Σ_X .
- (C) $x_1 + z + w = 2x$, $x + x_2 + x_3 = z + w$. In this case, the cone $\langle z, w \rangle$ is in Σ_X and crosses $\langle x, x_2, x_3 \rangle$.



Proof. Since the relation $x_1 + x_2 + x_3 = x$ has degree two and it is not contractible, by theorem 5.6 it must be the sum of two relations of degree one. There are four types of primitive relations of degree one in a toric Fano 4-fold:

- a. $y_1 + y_2 = y_3;$
- b. $y_1 + y_2 + y_3 = 2y_4;$
- c. $y_1 + y_2 + y_3 = y_4 + y_5;$
- d. $y_1 + y_2 + y_3 + y_4 = 3y_5$.

We have to examine all possible combinations.

- (a+a): $y_1 + y_2 y_3 + w_1 + w_2 w_3 = x_1 + x_2 + x_3 x$. Then we have $y_3 = w_1, w_3 = x, y_1 = x_1, y_2 = x_2$ which implies $\langle x_1, x_2 \rangle \notin \Sigma_X$, a contradiction.
- (a+b): $y_1 + y_2 y_3 + w_1 + w_2 + w_3 2w_4 = x_1 + x_2 + x_3 x$. Then we have $y_1 = w_4 = x$, $y_3 = w_1$, $y_2 = x_1$, $w_2 = x_2$, $w_3 = x_3$; this gives (A).
- (a+c): $y_1 + y_2 y_3 + w_1 + w_2 + w_3 w_4 w_5 = x_1 + x_2 + x_3 x$. Then we have $w_5 = x$, $y_1 = w_4$, $y_3 = w_1$, $y_2 = x_1$, $w_2 = x_2$, $w_3 = x_3$; this gives (B).

(a+d): $y_1 + y_2 - y_3 + w_1 + w_2 + w_3 + w_4 - 3w_5 = x_1 + x_2 + x_3 - x$, impossible.

- (b+b): $y_1 + y_2 + y_3 2y_4 + w_1 + w_2 + w_3 2w_4 = x_1 + x_2 + x_3 x$, impossible.
- (b+c): $y_1 + y_2 + y_3 2y_4 + w_1 + w_2 + w_3 w_4 w_5 = x_1 + x_2 + x_3 x$. Then we have $y_4 = w_1 = x$, $y_1 = w_4$, $y_2 = w_5$, $y_3 = x_1$, $w_2 = x_2$, $w_3 = x_3$; this gives (C).
- (b+d): $y_1 + y_2 + y_3 2y_4 + w_1 + w_2 + w_3 + w_4 3w_5 = x_1 + x_2 + x_3 x$, impossible.

- (c+c): $y_1 + y_2 + y_3 y_4 y_5 + w_1 + w_2 + w_3 w_4 w_5 = x_1 + x_2 + x_3 x$. Then we have $y_5 = x$, $y_4 = w_1$, $w_4 = y_1$, $w_5 = y_2$, $y_3 = x_1$, $w_2 = x_2$, $w_3 = x_3$. We get the relations: $y_1 + y_2 + x_1 = y_4 + x$ and $y_4 + x_2 + x_3 = y_1 + y_2$. Since they both have degree 1, they are extremal, so the cones $\langle y_1, y_2, y_4, x \rangle$, $\langle y_1, y_2, y_4, x_2 \rangle$ and $\langle y_1, y_2, y_4, x_3 \rangle$ are in Σ_X , which is impossible.
- (c+d): $y_1 + y_2 + y_3 y_4 y_5 + w_1 + w_2 + w_3 + w_4 3w_5 = x_1 + x_2 + x_3 x$, impossible.
- (d+d): $y_1 + y_2 + y_3 + y_4 3y_5 + w_1 + w_2 + w_3 + w_4 3w_5 = x_1 + x_2 + x_3 x$, impossible.

Proof of proposition 5.3. Consider the unique cone $\tilde{\sigma} \in \Sigma_X$ such that $y_1 + y_2 + y_3 + y_4 \in \text{Rel Int } \tilde{\sigma}$, where $\sigma = \langle y_1, y_2, y_3, y_4 \rangle$. The generators of $\tilde{\sigma}$ are described in lemma 5.4.

(I) If dim $\tilde{\sigma} = 4$, then $\tilde{\sigma} = \sigma$; this is (1).

(II) If dim $\tilde{\sigma} = 3$, by lemma 5.4 the only possibility is $x_1 = y_1 + y_2 \in G(\Sigma_X)$ and $\tilde{\sigma} = \langle x_1, y_3, y_4 \rangle$. In $\Sigma_X y_1 + y_2 = x_1$ is a primitive relation of degree 1, so it is extremal; $\tilde{\sigma} \in \Sigma_X$ implies $\langle x_1, y_1, y_3, y_4 \rangle \in \Sigma_X$, $\langle x, y_2, y_3, y_4 \rangle \in \Sigma_X$; this gives (2).

(III) Suppose dim $\tilde{\sigma} = 2$. By lemma 5.4, there are two possibilities for $\tilde{\sigma}: \tilde{\sigma} = \langle y_1 + y_2, y_3 + y_4 \rangle$ or $\tilde{\sigma} = \langle y_1 + y_2 + y_3, y_4 \rangle$.

(III.1) Let $\tilde{\sigma} = \langle x_1, x_2 \rangle$ with $x_1 = y_1 + y_2$, $x_2 = y_3 + y_4$. Then $y_1 + y_2 = x_1$ and $y_3 + y_4 = x_2$ are primitive relations of degree 1, hence they are extremal. So $\tilde{\sigma} \in \Sigma_X$ implies that the cones $\langle x_1, x_2, y_1, y_3 \rangle$, $\langle x_1, x_2, y_1, y_4 \rangle$, $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$ are in Σ_X and this gives (5).

(III.2) Let $\tilde{\sigma} = \langle x_1, y_4 \rangle$ with $x_1 = y_1 + y_2 + y_3$. Then $\langle y_1, y_2, y_3 \rangle \notin \Sigma_X$, so $\{y_1, y_2, y_3\}$ contains a primitive collection.

(III.2.1) Suppose $\{y_1, y_2, y_3\}$ is primitive: then the primitive relation is $y_1 + y_2 + y_3 = x_1$.

(III.2.1.1) If the relation $y_1 + y_2 + y_3 = x_1$ is contractible, $\tilde{\sigma} \in \Sigma_X$ implies that $\langle x_1, y_1, y_2, y_4 \rangle$, $\langle x_1, y_1, y_3, y_4 \rangle$ and $\langle x_1, y_2, y_3, y_4 \rangle$ are in Σ_X and we get (3).

(III.2.1.2) If $\{y_1, y_2, y_3\}$ is primitive but $y_1 + y_2 + y_3 = x_1$ is not contractible, then it must be the sum of two relations of degree 1, and the possible cases are described in lemma 5.7. We remark that the case (C) never occurs, because Σ_X is a subdivision of Σ_Y .

(III.2.1.2.1) If the sum is like in (A), then we have two extremal relations: $y_1 + x_1 = x_2$ and $y_2 + y_3 + x_2 = 2x_1$. Since $\tilde{\sigma} = \langle x_1, y_4 \rangle$ is in Σ_X , we get that the cones $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$, $\langle x_1, x_2, y_3, y_4 \rangle$, $\langle x_2, y_1, y_2, y_4 \rangle$, $\langle x_2, y_1, y_3, y_4 \rangle$ are in Σ_X too. This is (6).

31

(III.2.1.2.2) If the sum is like in (B), we have the extremal relations $y_1 + x_3 = x_2$ and $x_2 + y_2 + y_3 = x_1 + x_3$. Examining the 4-dimensional cones in Σ_X given by these relations, we see that the 2-dimensional cones of Σ_X contained in σ and containing x_1 are $\langle x_1, y_1 \rangle$, $\langle x_1, y_2 \rangle$, $\langle x_1, y_3 \rangle$, $\langle x_1, x_2 \rangle$ and $\langle x_1, x_3 \rangle$. Since $\tilde{\sigma} = \langle x_1, y_4 \rangle \in \Sigma_X$, it must be $y_4 = x_2$ or $y_4 = x_3$. If $y_4 = x_2$, then $x_3 = y_4 - y_1 \notin \sigma$, which is impossible; hence $y_4 = x_3$ and $x_2 = y_1 + y_4$, and we get (7).

(III.2.2) If $\{y_1, y_2, y_3\}$ is not primitive, it must contain a primitive collection of order 2: we can suppose that it is $\{y_1, y_2\}$. The primitive relation can not be $y_1 + y_2 = 0$, because y_1 and y_2 both belong to σ . Therefore we get an extremal relation $y_1 + y_2 = x_2$. Moreover, since $y_3 + x_2 = x_1$, $\{y_3, x_2\}$ is a primitive collection of degree 1, which gives another extremal relation. Hence $\tilde{\sigma} = \langle x_1, y_4 \rangle \in \Sigma_X$ implies that $\langle x_1, x_2, y_1, y_4 \rangle$ and $\langle x_1, x_2, y_2, y_4 \rangle$ are in Σ_X . We claim that also $\langle x_1, y_1, y_3, y_4 \rangle$ and $\langle x_1, y_2, y_3, y_4 \rangle$ are in Σ_X , so the subdivision is (8). Indeed, suppose there exists $z \in G(\Sigma_X)$, $z \in \langle x_1, y_1, y_3, y_4 \rangle$, different from the vertexs. Then $\{z, x_2\}$ should be a primitive collection, which is impossible, because it could not have positive degree.

(IV) Finally, suppose dim $\tilde{\sigma} = 1$, i. e. $x_1 = y_1 + y_2 + y_3 + y_4 \in G(\Sigma_X)$.

(IV.1) We first consider the case where $\{y_1, y_2, y_3, y_4\}$ is primitive.

(IV.1.1) If $\{y_1, y_2, y_3, y_4\}$ is primitive and the relation $y_1 + y_2 + y_3 + y_4 = x_1$ is contractible, then $\langle x_1, y_1, y_2, y_3 \rangle$, $\langle x_1, y_1, y_2, y_4 \rangle$, $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$ are in Σ_X , so we get (4).

(IV.1.2) If $\{y_1, y_2, y_3, y_4\}$ is primitive and $y_1 + y_2 + y_3 + y_4 = x_1$ is not contractible, then by lemma 4.2 it must be $Y \simeq \mathbb{P}^4$: indeed, all the 3-dimensional faces of σ are in Σ_X , so there must be at least one generator of Σ_X different from x_1 in the interior of σ . Again by lemma 4.2, we know that Σ_X has exactly two generators in the interior of σ , x_1 and x_2 , with primitive relations $(-x_1)+x_2 = y_1$ and $x_1+y_1 = x_2$. We get two effective relations, $x_1+y_1 = x_2$ and $y_2 + y_3 + y_4 + x_2 = 2x_1$, whose sum is $y_1 + y_2 + y_3 + y_4 = x_1$. Since there are no other new generators in σ , these relations must be both contractible, therefore the cones $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$, $\langle x_1, x_2, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_2, y_1, y_2, y_3 \rangle$, $\langle x_2, y_1, y_2, y_4 \rangle$, $\langle x_2, y_1, y_3, y_4 \rangle$ are in Σ_X and we get (14).

(IV.2) Suppose now that $\{y_1, y_2, y_3\}$ is a primitive collection. The primitive relation can not be $y_1 + y_2 + y_3 = z_1 + z_2$, because the cone $\langle z_1, z_2 \rangle$ would cross $\langle y_1, y_2, y_3 \rangle$ and Σ_X would not be a subdivision of Σ_Y . Neither can the relation be $y_1 + y_2 + y_3 = 2x_2$, because $\{y_1, y_2, y_3\}$ is a part of a basis of the lattice. Thefore the primitive relation is $y_1 + y_2 + y_3 = x_2$. Then $\{x_2, y_4\}$ is a primitive collection with relation $x_2 + y_4 = x_1$, which is extremal.

(IV.2.1) If $y_1 + y_2 + y_3 = x_2$ is contractible, then the subdivision is given by the cones $\langle x_1, x_2, y_1, y_2 \rangle$, $\langle x_1, x_2, y_1, y_3 \rangle$, $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, y_1, y_2, y_4 \rangle$, $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$, and this gives (9).

(IV.2.2) If $y_1 + y_2 + y_3 = x_2$ is not contractible, then it is sum of two

primitive relations of degree 1, and we have to examine the cases given by lemma 5.7; we remark again that case (C) is not possible here, because Σ_X is a subdivision of Σ_Y .

(IV.2.2.1) If the sum is like in (A), then we have primitive relations $y_1 + x_2 = x_3$ and $x_3 + y_2 + y_3 = 2x_2$. We claim that this is impossible. Indeed, by the extremality of $x_2 + y_4 = x_1$ and $y_1 + x_2 = x_3$, we get that $\langle x_1, y_1 \rangle \in \Sigma_X$; on the other hand, we have $y_4 + x_3 = x_1 + y_1$, so $\{y_4, x_3\}$ would be a primitive collection of degree zero.

(IV.2.2.2) We have two possible decompositions as in case (B) of lemma 5.7:

- (B1) $y_1 + x_3 = x_1, y_2 + y_3 + x_1 = x_2 + x_3$. Then the cones $\langle x_1, y_1, y_2, y_4 \rangle$, $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, x_2, y_1, y_2 \rangle$, $\langle x_1, x_2, y_1, y_3 \rangle$, $\langle x_1, x_3, y_2, y_4 \rangle$, $\langle x_1, x_3, y_3, y_4 \rangle$, $\langle x_2, x_3, y_2, y_3 \rangle$, $\langle x_1, x_2, x_3, y_2 \rangle$, $\langle x_1, x_2, x_3, y_3 \rangle$ are in Σ_X , and we get (10).
- (B2) $y_1 + x_1 = x_3, y_2 + y_3 + x_3 = x_1 + x_2$. Then the cones $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_3, y_1, y_2, y_4 \rangle$, $\langle x_3, y_1, y_3, y_4 \rangle$, $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, x_3, y_2, y_4 \rangle$, $\langle x_1, x_3, y_3, y_4 \rangle$, $\langle x_2, x_3, y_1, y_2 \rangle$, $\langle x_2, x_3, y_1, y_3 \rangle$, $\langle x_1, x_2, x_3, y_2 \rangle$, $\langle x_1, x_2, x_3, y_3 \rangle$ are in Σ_X , and we get (15).

The last subdivision can occur only if $Y \simeq \mathbb{P}^4$, because there are two generators of Σ_X in the interior of σ .

(IV.3) Finally, suppose $\{y_1, y_2, y_3, y_4\}$ does not contain primitive relations of order 3. Then we can assume $\{y_1, y_2\}$ primitive, with relation $y_1 + y_2 = x_2$. Since $y_3 + y_4 + x_2 = x_1$, the cone $\langle y_3, y_4, x_2 \rangle$ can not be in Σ_X , so $\{y_3, y_4, x_2\}$ contains a primitive collection.

(IV.3.1) If $\{y_3, y_4\}$ is primitive, with relation $y_3 + y_4 = x_3$, then $x_2 + x_3 = x_1$ is also a primitive relation. Hence the subdivision is given by the cones $\langle x_1, x_2, y_1, y_3 \rangle$, $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, x_2, y_1, y_4 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$, $\langle x_1, x_3, y_1, y_3 \rangle$, $\langle x_1, x_3, y_2, y_3 \rangle$, $\langle x_1, x_3, y_2, y_4 \rangle$, and we get (12).

(IV.3.2) If $\{y_3, x_2\}$ is primitive, with relation $y_3 + x_2 = x_3$, then $y_4 + x_3 = x_1$ is also a primitive relation. Hence the subdivision is given by the cones $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_1, x_2, y_1, y_4 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$, $\langle x_1, x_3, y_1, y_3 \rangle$, $\langle x_1, x_3, y_2, y_3 \rangle$, $\langle x_1, x_2, x_3, y_1 \rangle$, $\langle x_1, x_2, x_3, y_2 \rangle$; this gives (13).

(IV.3.3) The last possibility is that $\{y_3, y_4, x_2\}$ is a primitive collection. (IV.3.3.1) If $y_3+y_4+x_2 = x_1$ is contractible, then the cones $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_1, x_2, y_1, y_3 \rangle$, $\langle x_1, x_2, y_2, y_3 \rangle$, $\langle x_1, x_2, y_1, y_4 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle$ are in Σ_X and this gives (11).

(IV.3.3.2) If $\{y_3, y_4, x_2\}$ is primitive but $y_3 + y_4 + x_2 = x_1$ is not contractible, then it must decompose as a sum of two primitive relations of degree 1 as described in lemma 5.7.

We have two possible decompositions as in case (A) of lemma 5.7:

(A1) $y_3 + x_1 = x_3, y_4 + x_2 + x_3 = 2x_1$. Then the cones $\langle x_1, x_2, y_1, y_4 \rangle$, $\langle x_1, x_2, y_2, y_4 \rangle, \quad \langle x_3, y_1, y_3, y_4 \rangle, \quad \langle x_3, y_2, y_3, y_4 \rangle, \quad \langle x_1, x_3, y_1, y_4 \rangle,$ $\langle x_1, x_3, y_2, y_4 \rangle$, $\langle x_2, x_3, y_1, y_3 \rangle$, $\langle x_2, x_3, y_2, y_3 \rangle$, $\langle x_1, x_2, x_3, y_1 \rangle$, $\langle x_1, x_2, x_3, y_2 \rangle$ are in Σ_X , and we get (16).

(A2) $x_1 + x_2 = x_3, y_3 + y_4 + x_3 = 2x_1$. Then the cones $\langle x_1, y_1, y_3, y_4 \rangle$, $\langle x_1, y_2, y_3, y_4 \rangle$, $\langle x_1, x_3, y_1, y_3 \rangle$, $\langle x_2, x_3, y_1, y_3 \rangle$, $\langle x_1, x_3, y_2, y_3 \rangle$, $\langle x_2, x_3, y_2, y_3 \rangle$, $\langle x_1, x_3, y_1, y_4 \rangle$, $\langle x_2, x_3, y_1, y_4 \rangle$, $\langle x_1, x_3, y_2, y_4 \rangle$, $\langle x_2, x_3, y_2, y_4 \rangle$ are in Σ_X , and we get (17).

We remark that both subdivisions can occur only when $Y \simeq \mathbb{P}^4$, because there are two generators of Σ_X in the interior of σ .

We claim that $y_3 + y_4 + x_2 = x_1$ can not decompose as in case (B) of lemma 5.7. Indeed, either the decomposition is given by $y_3 + w = z$ and $y_4 + x_2 + z = w + x_1$, or by $x_2 + w = z$ and $y_3 + y_4 + z = w + x_1$; in both cases we get $\langle z, y_3, x_1, x_2 \rangle \in \Sigma_X$. Now, $y_1 + y_2 = x_2$ is extremal, so it must be $z = y_1$ or $z = y_2$. This implies $w \notin \sigma$, a contradiction.

Proof of theorem 5.2. We know by proposition 5.3 that the subdivision of every 4-dimensional cone of Σ_Y is given by a sequence of star-subdivisions. We just have to check that we can order these star-subdivisions in a way which is compatible with the orders in all 4-dimensional cones of Σ_Y .

We order the cones corresponding to centers of star-subdivisions in such a way that:

- (i) the dimensions of the cones are non-increasing
- (ii) each cone appears in the fan obtained with the preceeding star-subdivisions.

It is easy to see directly, looking at the subdivisions given by proposition 5.3, that this order works. More precisely, in all the sequences of star-subdivisions given by proposition 5.3, except for (5) and (10), there is a unique cone of maximal dimension. In (5) and (10) both cones of maximal dimension can be chosen for the first star-subdivision. Then we can use induction.

References

- Dan Abramovich, Kenji Matsuki, and Suliman Rashid. A note on the factorization theorem of toric birational maps and its toroidal extension. *Tohoku Mathematical Journal*, 51:489–537, 1999.
- [2] Victor V. Batyrev. Toroidal Fano 3-folds. Mathematics of the USSR Izvestiya, 19:13–25, 1982.
- [3] Victor V. Batyrev. On the classification of smooth projective toric varieties. Tohoku Mathematical Journal, 43:569–585, 1991.

- [4] Victor V. Batyrev. On the classification of toric Fano 4-folds. Journal of Mathematical Sciences (New York), 94:1021–1050, 1999.
- [5] Laurent Bonavero and Shigeharu Takayama. Some boundedness results for Fano-like Moishezon manifolds. *Documenta Mathematica*, 5:141–150, 2000.
- [6] Cinzia Casagrande. Contractible classes in smooth complete toric varieties. *Preprint*, 2001.
- [7] Cinzia Casagrande. On the birational geometry of toric Fano 4-folds. *Comptes Rendus de l'Académie des Sciences*, 12:1093–1098, 2001.
- [8] Olivier Debarre. Toric Fano varieties. Preprint, 2001.
- [9] Günter Ewald. On the classification of toric Fano varieties. *Discrete Computational Geometry*, 3:49–54, 1988.
- [10] Günter Ewald. Combinatorial Convexity and Algebraic Geometry, volume 168 of Graduate Texts in Mathematics. Springer-Verlag, 1996.
- [11] William Fulton. Introduction to Toric Varieties. Number 131 in Annals of Mathematics Studies. Princeton University Press, 1993.
- [12] Peter Kleinschmidt and Bernd Sturmfels. Smooth toric varieties with small Picard number are projective. *Topology*, 30(2):289–299, 1991.
- [13] Peter McMullen and Geoffrey C. Shephard. Convex Polytopes and the Upper Bound Conjecture, volume 3 of London Mathematical Society Lecture Note Series. Cambridge University Press, 1971.
- [14] Robert Morelli. The birational geometry of toric varieties. Journal of Algebraic Geometry, 5:751–782, 1996.
- [15] Tadao Oda. Convex Bodies and Algebraic Geometry An Introduction to the Theory of Toric Varieties. Springer-Verlag, 1988.
- [16] Miles Reid. Decomposition of toric morphisms. In Arithmetic and Geometry, vol. II: Geometry, number 36 in Progress in Mathematics, pages 395–418, 1983.
- [17] Hiroshi Sato. Toward the classification of higher-dimensional toric Fano varieties. *Tohoku Mathematical Journal*, 52:383–413, 2000.
- [18] Michal Szurek and Jaroslaw A. Wiśniewski. Fano bundles over \mathbb{P}^3 and Q_3 . *Pacific Journal of Mathematics*, 141(1):197–208, 1990.
- [19] V. E. Voskresenskii and Alexander Klyachko. Toroidal Fano varieties and roots systems. *Mathematics of the USSR Izvestiya*, 24:221–244, 1985.
- [20] K. Watanabe and M. Watanabe. The classification of Fano 3-folds with torus embeddings. *Tokyo Journal of Mathematics*, 5:37–48, 1982.
- [21] Jaroslaw A. Wiśniewski. On contractions of extremal rays of Fano manifolds. Journal für die Reine und Angewandte Mathematik, 417:141–157, 1991.

Cinzia Casagrande Dipartimento di Matematica Università di Roma "La Sapienza" piazzale Aldo Moro, 5 00185 Roma ccasagra@mat.uniroma1.it