

**AN EXTENSION THEOREM FOR
SEPARATELY HOLOMORPHIC FUNCTIONS
WITH PLURIPOLAR SINGULARITIES**

MAREK JARNICKI (KRAKÓW) ^(†) PETER PFLUG (OLDENBURG) ^(‡)

ABSTRACT. Let $D_j \subset \mathbb{C}^{n_j}$ be a pseudoconvex domain and let $A_j \subset D_j$ be a locally pluriregular set, $j = 1, \dots, N$. Put

$$X := \bigcup_{j=1}^N A_1 \times \cdots \times A_{j-1} \times D_j \times A_{j+1} \times \cdots \times A_N \subset \mathbb{C}^{n_1} \times \cdots \times \mathbb{C}^{n_N} = \mathbb{C}^n.$$

Let $U \subset \mathbb{C}^n$ be an open neighborhood of X and let $M \subset U$ be a relatively closed subset of U . For $j \in \{1, \dots, N\}$ let Σ_j be the set of all $(z', z'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N)$ for which the fiber $M_{(z', z'')} := \{z_j \in \mathbb{C}^{n_j} : (z', z_j, z'') \in M\}$ is not pluripolar. Assume that $\Sigma_1, \dots, \Sigma_N$ are pluripolar. Put

$$X' := \bigcup_{j=1}^N \{(z', z_j, z'') \in (A_1 \times \cdots \times A_{j-1}) \times D_j \times (A_{j+1} \times \cdots \times A_N) : (z', z'') \notin \Sigma_j\}.$$

Then there exists a relatively closed pluripolar subset $\widehat{M} \subset \widehat{X}$ of the ‘envelope of holomorphy’ $\widehat{X} \subset \mathbb{C}^n$ of X such that:

- $\widehat{M} \cap X' \subset M$,
 - for every function f separately holomorphic on $X \setminus M$ there exists exactly one function \widehat{f} holomorphic on $\widehat{X} \setminus \widehat{M}$ with $\widehat{f} = f$ on $X' \setminus M$, and
 - \widehat{M} is singular with respect to the family of all functions \widehat{f} .
- Some special cases were previously studied in [Jar-Pfl 2001c].

1. Introduction. Main Theorem. Let $N \in \mathbb{N}$, $N \geq 2$, and let

$$\emptyset \neq A_j \subset D_j \subset \mathbb{C}^{n_j},$$

1991 *Mathematics Subject Classification.* 32D15, 32D10.

^(†) Research partially supported by the KBN grant No. 5 P03A 033 21.

^(‡) Research partially supported by the Niedersächsisches Ministerium für Wissenschaft und Kultur, Az. 15.3 – 50 113(55) PL.

where D_j is a domain, $j = 1, \dots, N$. We define an N -fold cross

$$\begin{aligned} X &= \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N) \\ &:= \bigcup_{j=1}^N A_1 \times \dots \times A_{j-1} \times D_j \times A_{j+1} \times \dots \times A_N \subset \mathbb{C}^{n_1 + \dots + n_N} = \mathbb{C}^n. \end{aligned}$$

Observe that X is connected.

Let $\Omega \subset \mathbb{C}^n$ be an open set and let $A \subset \Omega$. Put

$$h_{A,\Omega} := \sup\{u : u \in \mathcal{PSH}(\Omega), u \leq 1 \text{ on } \Omega, u \leq 0 \text{ on } A\},$$

where $\mathcal{PSH}(\Omega)$ denotes the set of all functions plurisubharmonic on Ω . Define

$$\omega_{A,\Omega} := \lim_{k \rightarrow +\infty} h_{A \cap \Omega_k, \Omega_k}^*,$$

where $(\Omega_k)_{k=1}^\infty$ is a sequence of relatively compact open sets $\Omega_k \subset \Omega_{k+1} \subset \subset \Omega$ with $\bigcup_{k=1}^\infty \Omega_k = \Omega$ (h^* denotes the upper semicontinuous regularization of h). Observe that the definition is independent of the exhausting sequence $(\Omega_k)_{k=1}^\infty$. Moreover, $\omega_{A,\Omega} \in \mathcal{PSH}(\Omega)$. Recall that if Ω is bounded, then $\omega_{A,\Omega} = h_{A,\Omega}^*$.

For an N -fold cross $X = \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N)$ put

$$\widehat{X} := \{(z_1, \dots, z_N) \in D_1 \times \dots \times D_N : \sum_{j=1}^N \omega_{A_j, D_j}(z_j) < 1\}.$$

Observe that if D_1, \dots, D_N are pseudoconvex, then \widehat{X} is a pseudoconvex open set in \mathbb{C}^n .

We say that a subset $\emptyset \neq A \subset \mathbb{C}^n$ is *locally pluriregular* if $h_{A \cap \Omega, \Omega}^*(a) = 0$ for any $a \in A$ and for any open neighborhood Ω of a (in particular, $A \cap \Omega$ is non-pluripolar).

Note that if A_1, \dots, A_N are locally pluriregular, then $X \subset \widehat{X}$ and \widehat{X} is connected ([Jar-Pfl 2001c], Lemma 4).

Let U be an open neighborhood of X and let $M \subset U$ be a relatively closed set. We say that a function $f: X \setminus M \rightarrow \mathbb{C}$ is *separately holomorphic* ($f \in \mathcal{O}_s(X \setminus M)$) if for any $(a_1, \dots, a_N) \in A_1 \times \dots \times A_N$ and $j \in \{1, \dots, N\}$ the function $f(a_1, \dots, a_{j-1}, \cdot, a_{j+1}, \dots, a_N)$ is holomorphic in the open set

$$D_j \setminus M_{(a_1, \dots, a_{j-1}, \cdot, a_{j+1}, \dots, a_N)},$$

where

$$M_{(a_1, \dots, a_{j-1}, \cdot, a_{j+1}, \dots, a_N)} := \{z_j \in \mathbb{C}^{n_j} : (a_1, \dots, a_{j-1}, z_j, a_{j+1}, \dots, a_N) \in M\}.$$

Suppose that $S_j \subset A_1 \times \cdots \times A_{j-1} \times A_{j+1} \times \cdots \times A_N$, $j = 1, \dots, N$, and define the *generalized N -fold cross*

$$T = \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; S_1, \dots, S_N) \\ := \bigcup_{j=1}^N \{(z', z_j, z'') \in (A_1 \times \cdots \times A_{j-1}) \times D_j \times (A_{j+1} \times \cdots \times A_N) : (z', z'') \notin S_j\}.$$

It is clear that $T \subset X$. Observe that

$$\mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N) = \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; \emptyset, \dots, \emptyset).$$

Moreover, if $N = 2$, then $\mathbb{T}(A_1, A_2; D_1, D_2; S_1, S_2) = \mathbb{X}(A_1 \setminus S_2, A_2 \setminus S_1; D_1, D_2)$. Consequently, any generalized 2-fold cross is a 2-fold cross.

Let $S \subset \Omega$ be a relatively closed pluripolar subset of an open set $\Omega \subset \mathbb{C}^n$. Let $\mathcal{F} \subset \mathcal{O}(\Omega \setminus S)$. We say that S is *singular with respect to \mathcal{F}* if for any point $a \in M$ there exists a function $f_a \in \mathcal{F}$ which is not holomorphically extendible to a neighborhood of a (cf. [Jar-Pfl 2000], § 3.4). Equivalently: the set S is minimal in the sense that there is no relatively closed set $S' \subsetneq S$ such that any function from \mathcal{F} extends holomorphically to $\Omega \setminus S'$. It is clear that for any relatively closed pluripolar set $S \subset \Omega$ and for any family $\mathcal{F} \subset \mathcal{O}(\Omega \setminus S)$ there exists a relatively closed set $S' \subset S$ such that any function $f \in \mathcal{F}$ extends to an $f' \in \mathcal{O}(\Omega \setminus S')$ and S' is singular with respect to the family $\{f' : f \in \mathcal{F}\}$.

The main result of our paper is the following extension theorem for separately holomorphic functions.

Main Theorem. *Let $D_j \subset \mathbb{C}^{n_j}$ be a pseudoconvex domain, let $A_j \subset D_j$ be a locally pluriregular set, $j = 1, \dots, N$, and let U be an open neighborhood of the N -fold cross*

$$X := \mathbb{X}(A_1, \dots, A_N; D_1, \dots, D_N).$$

Let $M \subset U$ be a relatively closed subset of U such that for each $j \in \{1, \dots, N\}$ the set

$$\Sigma_j = \Sigma_j(A_1, \dots, A_N; M) \\ := \{(z', z'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N) : M_{(z', \cdot, z'')} \text{ is not pluripolar}\}$$

is pluripolar. Put

$$X' := \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; \Sigma_1, \dots, \Sigma_N).$$

Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:

- $\widehat{M} \cap X' \subset M$,
- for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X' \setminus M$,

- \widehat{M} is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$, and
- $\widehat{X} \setminus \widehat{M}$ is pseudoconvex.

In particular, $\widehat{X} \setminus \widehat{M}$ is the envelope of holomorphy of $X \setminus M$ with respect to the space of separately holomorphic functions.

Notice that if $M \subset U$ is a pluripolar set, then $\Sigma_1, \dots, \Sigma_N$ are always pluripolar (cf. Lemma 8(a)).

The case where $N = 2$, $n_1 = n_2 = 1$, $D_1 = D_2 = \mathbb{C}$ was studied in [Jar-Pff 2001b], Th. 2.

Corollary 1. *Let $D_j, A_j, j = 1, \dots, N, X$, and U be as in the Main Theorem. Assume that $M \subset U$ is a relatively closed set such that for any $(a_1, \dots, a_N) \in A_1 \times \dots \times A_N$ and $j \in \{1, \dots, N\}$ the fiber $M_{(a_1, \dots, a_{j-1}, \cdot, a_{j+1}, \dots, a_N)}$ is pluripolar ⁽¹⁾.*

Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:

- $\widehat{M} \cap X \subset M$,
- for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X \setminus M$, and
- the domain $\widehat{X} \setminus \widehat{M}$ is pseudoconvex.

The case where $N = 2$, $D_2 = \mathbb{C}^{n_2}$, and A_2 is open was studied in [Chi-Sad 1988] (for $n_2 = 1$) and [Kaz 1988] (for arbitrary n_2).

The proof of the Main Theorem will be presented in Sections 3 (for $N = 2$) and 4 (for arbitrary N).

The following two examples illustrate the role played by the sets Σ_j and show that the assertion of the Main Theorem is in some sense optimal.

Example 2. Let $n_1 = n_2 = 1$, $D_1 = D_2 = \mathbb{C}$, $A_1 = A_2 = E :=$ the unit disc, $X := \mathbb{X}(E, E; \mathbb{C}, \mathbb{C}) = (E \times \mathbb{C}) \cup (\mathbb{C} \times E)$.

(a) Let $M := \{0\} \times E$. Then $\Sigma_1 = \emptyset$, $\Sigma_2 = \{0\}$, $X' = \mathbb{X}(E \setminus \{0\}, E; \mathbb{C}, \mathbb{C})$, $\widehat{M} = \{0\} \times \mathbb{C}$.

Put $f_0(z, w) := 1/z$, $z \neq 0$. Then $f_0 \in \mathcal{O}_s(X \setminus M)$ and \widehat{M} is singular with respect to f_0 .

(b) Let $M := \{0\} \times \{|w| = r\}$ for some $0 < r < 1$. Then $\Sigma_1 = \emptyset$, $\Sigma_2 = \{0\}$, $X' = \mathbb{X}(E \setminus \{0\}, E; \mathbb{C}, \mathbb{C})$, $\widehat{M} = \emptyset$.

Put

$$f_0(z, w) := \begin{cases} w & \text{if } z \neq 0 \text{ or } (z = 0 \text{ and } |w| > r) \\ 0 & \text{if } z = 0 \text{ and } |w| < r \end{cases}, \quad (z, w) \in X \setminus M.$$

Then $f_0 \in \mathcal{O}_s(X \setminus M)$, $\widehat{f}_0(z, w) \equiv w$, and $\widehat{f}_0(0, w) \neq f_0(0, w)$, $0 < |w| < r$.

2. Auxiliary results.

In the case $M = \emptyset$ the problem of extension of separately holomorphic functions was studied by many authors (under various assumptions on $(D_j, A_j)_{j=1}^N$), e.g. [Sic 1969], [Zah 1976], [Sic 1981], [Shi 1989], [Ngu-Zer 1991], [Ngu 1997], [Ale-Zer 2001] (for $N = 2$), and [Sic 1981], [Ngu-Zer 1995], [Jar-Pff 2001c] (for arbitrary N).

⁽¹⁾ That is $\Sigma_1 = \dots = \Sigma_N = \emptyset$.

Theorem 3 ([Ngu-Zer 1995], [Ale-Zer 2001]). *Let $(D_j, A_j)_{j=1}^N$ and X be as in the Main Theorem. Then any function from $\mathcal{O}_s(X)$ extends holomorphically to the pseudoconvex domain \widehat{X} .*

The case where M is analytic was studied in [Ökt 1998], [Ökt 1999], [Sic 2000], [Jar-Pfl 2001a]. The problem was completely solved in [Jar-Pfl 2001c].

Theorem 4 ([Jar-Pfl 2001b]). *Let $(D_j, A_j)_{j=1}^N$ and X be as in the Main Theorem. Let $M \subsetneq U$ be an analytic subset of an open connected neighborhood U of X . Then there exists an analytic set $\widehat{M} \subset \widehat{X}$ such that:*

- $\widehat{M} \cap U_0 \subset M$ for an open neighborhood U_0 of X , $U_0 \subset U$,
- for every $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X \setminus M$, and
- the domain $\widehat{X} \setminus \widehat{M}$ is pseudoconvex.

Remark 5. It is a natural idea to try to obtain Theorem 4 from the Main Theorem. More precisely, let $(D_j, A_j)_{j=1}^N$, X , U , and M be as in Theorem 4. Then, by the Main Theorem, there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ which has all the properties listed in the Main Theorem. We would like to know whether there is a direct argument showing that \widehat{M} must be analytic.

The following two results will play the fundamental role in the sequel.

Theorem 6 ([Chi 1993]). *Let $D \subset \mathbb{C}^n$ be a domain and let \widehat{D} be the envelope of holomorphy of D . Assume that S is relatively closed pluripolar subset of D . Then there exists a relatively closed pluripolar subset \widehat{S} of \widehat{D} such that $\widehat{S} \cap D \subset S$ and $\widehat{D} \setminus \widehat{S}$ is the envelope of holomorphy of $D \setminus S$.*

Theorem 7 ([Jar-Pfl 2001b]). *Let $A \subset E^{n-1}$ be locally pluriregular, let*

$$X := \mathbb{X}(A, E; E^{n-1}, \mathbb{C})$$

(notice that $\widehat{X} = E^{n-1} \times \mathbb{C}$), and let $U \subset E^{n-1} \times \mathbb{C}$ be an open neighborhood of X . Let $M \subset U$ be a relatively closed set such that $M \cap E^n = \emptyset$ and for any $a \in A$ the fiber $M_{(a, \cdot)}$ is polar. Then there exists a relatively closed pluripolar set $S \subset E^{n-1} \times \mathbb{C}$ such that

- $S \cap X \subset M$,
- any function from $\mathcal{O}_s(X \setminus M)$ extends holomorphically to $E^{n-1} \times \mathbb{C} \setminus S$, and
- $E^{n-1} \times \mathbb{C} \setminus S$ ⁽²⁾ is pseudoconvex.

Notice that the above result is a special case of our Main Theorem with $N = 2$, $n_1 = n - 1$, $D_1 = E^{n-1}$, $A_1 = A$, $n_2 = 1$, $D_2 = \mathbb{C}$, $A_2 = E$, $\Sigma_1 = \Sigma_2 = \emptyset$.

⁽²⁾ Here and in the sequel to simplify notation we write $P_1 \times \cdots \times P_k \setminus Q$ instead of $(P_1 \times \cdots \times P_k) \setminus Q$.

Proof. It is known (cf. [Chi-Sad 1988]) that each function $f \in \mathcal{O}_s(X \setminus M)$ has the univalent domain of existence $G_f \subset E^{n-1} \times \mathbb{C}$ ⁽³⁾. Let G denote the connected component of $\text{int} \bigcap_{f \in \mathcal{O}_s(X \setminus M)} G_f$ that contains E^n and let $S := E^{n-1} \times \mathbb{C} \setminus G$. It remains to show that S is pluripolar.

Take $(a, b) \in A \times \mathbb{C} \setminus M$. Since $M_{(a, \cdot)}$ is polar, there exists a curve $\gamma: [0, 1] \rightarrow \mathbb{C} \setminus M_{(a, \cdot)}$ such that $\gamma(0) = 0$, $\gamma(1) = b$. Take an $\varepsilon > 0$ so small that

$$\Delta_a(\varepsilon) \times (\gamma([0, 1]) + \Delta_0(\varepsilon)) \subset U \setminus M,$$

where $\Delta_{z_0}(r) = \Delta_{z_0}^k(r) \subset \mathbb{C}^k$ denotes the polydisc with center $z_0 \in \mathbb{C}^k$ and radius $r > 0$. Put $V_b := E \cup (\gamma([0, 1]) + \Delta_0(\varepsilon))$ and consider the cross

$$Y := \mathbb{X}(A \cap \Delta_a(\varepsilon), E; \Delta_a(\varepsilon), V_b).$$

Then $f \in \mathcal{O}_s(Y)$ for any $f \in \mathcal{O}_s(X \setminus M)$. Consequently, by Theorem 3, we get $\widehat{Y} \subset G_f$, $f \in \mathcal{O}_s(X \setminus M)$. Hence $\widehat{Y} \subset G$. In particular, we conclude that $\{a\} \times (\mathbb{C} \setminus M_{(a, \cdot)}) \subset G$.

Thus $S_{(a, \cdot)} \subset M_{(a, \cdot)}$ for all $a \in A$. Consequently, by Lemma 5 from [Chi-Sad 1988], S is pluripolar. \square

Lemma 8. (a) *Let $S \subset \mathbb{C}^p \times \mathbb{C}^q$ be pluripolar. Then the set*

$$A := \{z \in \mathbb{C}^p : S_{(z, \cdot)} \text{ is not pluripolar}\}$$

is pluripolar.

(b) *Let $M \subset \mathbb{C}^p \times \mathbb{C}^q$ be such that for each $a \in \mathbb{C}^p$ the fiber $M_{(a, \cdot)}$ is pluripolar. Let $C \subset \mathbb{C}^p \times \mathbb{C}^q$ be such that the set $\{z \in \mathbb{C}^p : C_{(z, \cdot)} \text{ is not pluripolar}\}$ is not pluripolar (e.g. $C = C' \times C''$, where $C' \subset \mathbb{C}^p$, $C'' \subset \mathbb{C}^q$ are non-pluripolar). Then $C \setminus M$ is non-pluripolar.*

(c) *Let $M \subset \mathbb{C}^p \times \mathbb{C}^q$ be such that for each $a \in \mathbb{C}^p$ the fiber $M_{(a, \cdot)}$ is pluripolar. Let $A \subset \mathbb{C}^p$ be locally pluriregular. Let $C := \{(a, b') \in A \times \mathbb{C}^{q-1} : M_{(a, b', \cdot)} \text{ is polar}\}$. Then C is locally pluriregular.*

Proof. (a) Let $v \in \mathcal{PSH}(\mathbb{C}^{p+q})$, $v \not\equiv -\infty$, be such that $S \subset v^{-1}(-\infty)$. Define

$$u(z) := \sup\{v(z, w) : w \in \overline{E}^q\}, \quad z \in \mathbb{C}^p.$$

Then $A \subset u^{-1}(-\infty)$. Moreover, $u \in \mathcal{PSH}(\mathbb{C}^p)$ and $u \not\equiv -\infty$.

(b) Suppose that $C \setminus M$ is pluripolar. Then, by (a), there exists a pluripolar set $A \subset \mathbb{C}^p$ such that the fiber $(C \setminus M)_{(a, \cdot)}$ is pluripolar, $a \in \mathbb{C}^p \setminus A$. Consequently, the fiber $C_{(a, \cdot)}$ is pluripolar, $a \in \mathbb{C}^p \setminus A$; contradiction.

(c) Fix a point $(a_0, b'_0) \in C$ and a neighborhood $U := \Delta_{(a_0, b'_0)}(r)$. We have to show that $h_{C \cap U, U}^*(a_0, b'_0) = 0$.

⁽³⁾ We like to thank Professor Evgeni Chirka for explaining us some details of the proof of Theorem 1 in [Chi-Sad 1988].

Let $u := h_{C \cap U, U}^* \in \mathcal{PSH}(U)$. Then for any $a \in A \cap \Delta_{a_0}(r)$ the function $u(a, \cdot)$ is plurisubharmonic on $\Delta_{b'_0}(r)$ and $u(a, \cdot) \leq 0$ on the set

$$(C \cap U)_{(a, \cdot)} = \{b' \in \Delta_{b'_0}(r) : (M_{(a, \cdot)})_{(b', \cdot)} \text{ is polar}\}.$$

By (a) (applied to the set $M_{(a, \cdot)}$), the set $\Delta_{b'_0}(r) \setminus (C \cap \Delta_{a_0}(r))_{(a, \cdot)}$ is pluripolar. Hence $u(a, \cdot) \leq 0$ on $\Delta_{b'_0}(r)$. In particular, $u(\cdot, b'_0) \leq 0$ on $A \cap \Delta_{a_0}(r)$, which implies that

$$h_{C \cap U, U}^*(a_0, b'_0) = u(a_0, b'_0) \leq h_{A \cap \Delta_{a_0}(r), \Delta_{a_0}(r)}^*(a_0) = 0.$$

□

Lemma 9. *Let $D_j, A_j, j = 1, \dots, N$, and X be as in the Main Theorem. Let*

$$S_j \subset A_1 \times \cdots \times A_{j-1} \times A_{j+1} \times \cdots \times A_N$$

be pluripolar, $j = 1, \dots, N$. Put

$$T := \mathbb{T}(A_1, \dots, A_N; D_1, \dots, D_N; S_1, \dots, S_N).$$

Then any function $f \in \mathcal{O}_s(T) \cap \mathcal{C}(T)$ ⁽⁴⁾ extends holomorphically to \widehat{X} .

If $N = 2$, then the result is true for any function $f \in \mathcal{O}_s(T)$ (see the proof). In the case where $N \geq 3$ we do not know whether the result is true for arbitrary $f \in \mathcal{O}_s(T)$.

Proof. We apply induction on N . The case $N = 2$ follows from Theorem 3 and the fact that $\widehat{X} = \widehat{T}$ (recall that if $N = 2$, then T is a 2-fold cross). Moreover, if $N = 2$, then the result is true for any $f \in \mathcal{O}_s(T)$.

Assume that the result is true for $N - 1 \geq 2$. Take an $f \in \mathcal{O}_s(T) \cap \mathcal{C}(T)$. Let Q denote the set of all $z_N \in A_N$ for which there exists a $j \in \{1, \dots, N - 1\}$ such that the fiber $(S_j)_{(\cdot, z_N)}$ is not pluripolar. Then, by Lemma 8(a), Q is pluripolar. Take a $z_N \in A_N \setminus Q$ and define

$$T_{z_N} := \mathbb{T}(A_1, \dots, A_{N-1}; D_1, \dots, D_{N-1}; (S_1)_{(\cdot, z_N)}, \dots, (S_{N-1})_{(\cdot, z_N)}).$$

Then $f(\cdot, z_N) \in \mathcal{O}_s(T_{z_N}) \cap \mathcal{C}(T_{z_N})$. By the inductive assumption, the function $f(\cdot, z_N)$ extends to an $\widehat{f}_{z_N} \in \mathcal{O}(\widehat{Y})$, where $Y = \mathbb{X}(A_1, \dots, A_{N-1}; D_1, \dots, D_{N-1})$.

Let $A' := A_1 \times \cdots \times A_{N-1}$. Consider the 2-fold cross

$$Z := \mathbb{T}(A', A_N; \widehat{Y}, D_N; S_N, Q) = ((A' \setminus S_N) \times D_N) \cup (\widehat{Y} \times (A_N \setminus Q)).$$

Let $g: Z \rightarrow \mathbb{C}$ be given by the formulae:

⁽⁴⁾ We say that a function $f: T \rightarrow \mathbb{C}$ is *separately holomorphic* if for any $j \in \{1, \dots, N\}$ and $(a', a'') \in (A_1 \times \cdots \times A_{j-1}) \times (A_{j+1} \times \cdots \times A_N) \setminus S_j$ the function $f(a', \cdot, a'')$ is holomorphic in D_j .

$$g(z', z_N) := f(z', z_N), (z', z_N) \in (A' \setminus S_N) \times D_N,$$

$$g(z', z_N) := \widehat{f}_{z_N}(z'), (z', z_N) \in \widehat{X}' \times (A_N \setminus Q).$$

Observe that g is well-defined.

Indeed, let $(z', z_N) \in ((A' \setminus S_N) \times D_N) \cap (\widehat{Y} \times (A_N \setminus Q))$. If $z' \in T_{z_N}$, then obviously $\widehat{f}_{z_N}(z') = f(z', z_N)$. Suppose that $z' \notin T_{z_N}$. Then $z' \in P_{z_N} := \bigcap_{j=1}^{N-1} (S_j)_{(\cdot, z_N)}$; P_{z_N} is pluripolar. Take a sequence $A' \setminus (S_N \cup P_{z_N}) \ni z'^\nu \rightarrow z'$. Then $z'^\nu \in T_{z_N}$. Thus $\widehat{f}_{z_N}(z'^\nu) = f(z'^\nu, z_N)$. Hence, by continuity, $\widehat{f}_{z_N}(z') = f(z', z_N)$ ⁽⁵⁾.

Moreover, $g \in \mathcal{O}_s(Z)$. Put $V := \mathbb{X}(A', A_N; \widehat{Y}, D_N) \supset Z$. Since the result is true for $N = 2$ (without the continuity), we get a holomorphic extension of g to \widehat{V} . It remains to observe that $\widehat{V} = \widehat{X}$; cf. [Jar-Pfl 2001c], the proof of Step 3. \square

Lemma 10. *Let $D \subset \mathbb{C}^p$, $G \subset \mathbb{C}^q$ be pseudoconvex domains, let $A \subset D$, $B \subset G$ be locally pluriregular, and let $M \subset U$ be a relatively closed subset of an open neighborhood U of the cross $X := \mathbb{X}(A, B; D, G)$. Let $A' \subset A$, $B' \subset B$ be such that $A \setminus A'$, $B \setminus B'$ are pluripolar and for any $(a, b) \in A' \times B'$ the fibers $M_{(a, \cdot)}$, $M_{(\cdot, b)}$ are pluripolar. Let $(D_j)_{j=1}^\infty, (G_j)_{j=1}^\infty$ be sequences of pseudoconvex domains, $D_j \Subset D$, $G_j \Subset G$, with $D_j \nearrow D$, $G_j \nearrow G$, such that $A'_j := A' \cap D_j \neq \emptyset$, $B'_j := B' \cap G_j \neq \emptyset$, $j \in \mathbb{N}$. We assume that for each $j \in \mathbb{N}$, $a \in A'_j$, and $b \in B'_j$, there exist:*

- polydiscs $\Delta_a(r_{a,j}) \subset D_j$, $\Delta_b(s_{b,j}) \subset G_j$ and
- relatively closed pluripolar sets $S_{a,j} \subset \Delta_a(r_{a,j}) \times G_j$, $S^{b,j} \subset D_j \times \Delta_b(s_{b,j})$

such that:

- $(\Delta_a(r_{a,j}) \times G_j) \cup (D_j \times \Delta_b(s_{b,j})) \subset U \cap \widehat{X}$,
- $((A' \cap \Delta_a(r_{a,j})) \times G_j) \cap S_{a,j} \subset M$, $(D_j \times (B' \cap \Delta_b(s_{b,j}))) \cap S^{b,j} \subset M$,
- for any $f \in \mathcal{O}_s(X \setminus M)$ there exist functions $f_{a,j} \in \mathcal{O}(\Delta_a(r_{a,j}) \times G_j \setminus S_{a,j})$, $f^{b,j} \in \mathcal{O}(D_j \times \Delta_b(s_{b,j}) \setminus S^{b,j})$ with
 - $f_{a,j} = f$ on $(A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M$,
 - $f^{b,j} = f$ on $D_j \times (B' \cap \Delta_b(s_{b,j})) \setminus M$,
- $S_{a,j}$ is singular with respect to the family $\{f_{a,j} : f \in \mathcal{O}_s(X \setminus M)\}$, $S^{b,j}$ is singular with respect to the family $\{f^{b,j} : f \in \mathcal{O}_s(X \setminus M)\}$.

Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:

- $\widehat{M} \cap X' \subset M$, where $X' := \mathbb{X}(A', B'; D, G)$,
- for any $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X' \setminus M$,
- the set \widehat{M} is singular with respect to the family $\{\widehat{f} : f \in \mathcal{O}_s(X \setminus M)\}$.

Proof. Fix a $j \in \mathbb{N}$. Put

$$\begin{aligned} \widetilde{U}_j &:= \bigcup_{a \in A'_j, b \in B'_j} (\Delta_a(r_{a,j}) \times G_j) \cup (D_j \times \Delta_b(s_{b,j})), \\ X_j &:= ((A \cap D_j) \times G_j) \cup (D_j \times (B \cap G_j)), \\ X'_j &:= (A'_j \times G_j) \cup (D_j \times B'_j). \end{aligned}$$

⁽⁵⁾ Here is the only place where the continuity of f is used.

Note that $X'_j \subset \widetilde{U}_j$. Take an $f \in \mathcal{O}_s(X \setminus M)$. We like to glue the sets $(S_{a,j})_{a \in A'_j}$, $(S^{b,j})_{b \in B'_j}$ and the functions $(f_{a,j})_{a \in A'_j}$, $(f^{b,j})_{b \in B'_j}$ to obtain a global holomorphic function $f_j := \bigcup_{a \in A'_j, b \in B'_j} f_{a,j} \cup f^{b,j}$ on $\widetilde{U}_j \setminus S_j$ where $S_j := \bigcup_{a \in A'_j, b \in B'_j} S_{a,j} \cup S^{b,j}$.

Let $a \in A'_j$, $b \in B'_j$. Observe that

$$\begin{aligned} f_{a,j} &= f \text{ on } (A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M, \\ f^{b,j} &= f \text{ on } D_j \times (B' \cap \Delta_b(s_{b,j})) \setminus M. \end{aligned}$$

Thus $f_{a,j} = f^{b,j}$ on the non-pluripolar set $(A' \cap \Delta_a(r_{a,j})) \times (B' \cap \Delta_b(s_{b,j})) \setminus M$ (cf. Lemma 8(b)). Hence

$$f_{a,j} = f^{b,j} \text{ on } \Delta_a(r_{a,j}) \times \Delta_b(s_{b,j}) \setminus (S_{a,j} \cup S^{b,j}).$$

Using the minimality of $S_{a,j}$ and $S^{b,j}$, we conclude that

$$S_{a,j} \cap (\Delta_a(r_{a,j}) \times \Delta_b(s_{b,j})) = S^{b,j} \cap (\Delta_a(r_{a,j}) \times \Delta_b(s_{b,j})).$$

Now let $a', a'' \in A'_j$ be such that $C := \Delta_{a'}(r_{a',j}) \cap \Delta_{a''}(r_{a'',j}) \neq \emptyset$. Fix a $b \in B'_j$. We know that $f_{a',j} = f^{b,j} = f_{a'',j}$ on $C \times \Delta_b(s_{b,j}) \setminus (S_{a',j} \cup S^{b,j} \cup S_{a'',j})$. Hence, by the identity principle, we conclude that $f_{a',j} = f_{a'',j}$ on $C \times G_j \setminus (S_{a',j} \cup S_{a'',j})$ and, moreover,

$$S_{a',j} \cap (C \times G_j) = S_{a'',j} \cap (C \times G_j).$$

The same argument works for $b', b'' \in B' \cap G_j$.

Let U_j be the connected component of $\widetilde{U}_j \cap \widehat{X}'_j$ with $X'_j \subset U_j$. We have constructed a relatively closed pluripolar set $S_j \subset U_j$ such that:

$S_j \cap X'_j \subset M$ and

for any $f \in \mathcal{O}_s(X \setminus M)$ there exists (exactly one) $f_j \in \mathcal{O}(U_j \setminus S_j)$ with $f_j = f$ on $X'_j \setminus M$.

Recall that $X'_j \subset U_j \subset \widehat{X}'_j$. Hence the envelope of holomorphy \widehat{U}_j coincides with \widehat{X}'_j (cf. [Jar-Pfl 2001b], the proof of Step 4).

Applying the Chirka theorem (Theorem 6), we find a relatively closed pluripolar set $\widehat{M}_j \subset \widehat{X}'_j$ such that:

$\widehat{M}_j \cap U_j \subset S_j$,

for any $f \in \mathcal{O}_s(X \setminus M)$ there exists (exactly one) function $\widehat{f}_j \in \mathcal{O}(\widehat{X}'_j \setminus \widehat{M}_j)$ with $\widehat{f}_j = f_j$ on $U_j \setminus S_j$ (in particular, $\widehat{f}_j = f$ on $X'_j \setminus M$),

the set \widehat{M}_j is singular with respect to the family $\{\widehat{f}_j : f \in \mathcal{O}_s(X \setminus M)\}$.

Since $A \setminus A'$, $B \setminus B'$ are pluripolar, we get

$$\begin{aligned} \widehat{X}'_j &= \{(z, w) \in D_j \times G_j : h_{A' \cap D_j, D_j}^*(z) + h_{B' \cap G_j, G_j}^*(w) < 1\} \\ &= \{(z, w) \in D_j \times G_j : h_{A \cap D_j, D_j}^*(z) + h_{B \cap G_j, G_j}^*(w) < 1\} = \widehat{X}_j. \end{aligned}$$

So, in fact, $\widehat{f}_j \in \mathcal{O}(\widehat{X}_j \setminus M)$. Observe that $\bigcup_{j=1}^{\infty} X_j = X$, $\widehat{X}_j \subset \widehat{X}_{j+1}$, and $\bigcup_{j=1}^{\infty} \widehat{X}_j = \widehat{X}$. Using again the minimality of the \widehat{M}_j 's (and gluing the \widehat{f}_j 's), we get a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ which satisfies all the required conditions. \square

Lemma 11. *Let $A \subset E^{n-1}$ be locally pluriregular, let $G \subset \mathbb{C}$ be a domain with $E \Subset G$, let $X := \mathbb{X}(A, E; E^{n-1}, G)$, and let $U \subset E^{n-1} \times G$ be an open neighborhood of X . Let $M \subset U$ be a relatively closed set such that $M \cap E^n = \emptyset$ and for any $a \in A$ the fiber $M_{(a, \cdot)}$ is polar. Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:*

- $\widehat{M} \cap X \subset M$,
- for any $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X \setminus M$,
- the set \widehat{M} is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$.

Notice that the above result is a special case of our Main Theorem with $N = 2$, $n_1 = n - 1$, $D_1 = E^{n-1}$, $A_1 = A$, $n_2 = 1$, $D_2 = G$, $A_2 = E$, $\Sigma_1 = \Sigma_2 = \emptyset$.

Proof. By Lemma 10, it suffices to show that for any $a_0 \in A$ and for any domain $G' \Subset G$ with $E \Subset G'$ there exist $r > 0$ and a relatively closed pluripolar set $S \subset \Delta_{a_0}(r) \times G' \subset U$ such that:

- $S \cap X \subset M$ and
- any function from $\mathcal{O}_s(X \setminus M)$ extends holomorphically to $\Delta_{a_0}(r) \times G' \setminus S$.

Fix a_0 and G' . For $b \in G$, let $\rho = \rho_b > 0$ be such that $\Delta_b(\rho) \Subset G$ and $M_{(a_0, \cdot)} \cap \partial \Delta_b(\rho) = \emptyset$ (cf. [Arm-Gar 2001], Th. 7.3.9). Take $\rho^- = \rho_b^- > 0$, $\rho^+ = \rho_b^+ > 0$ such that $\rho^- < \rho < \rho^+$, $\Delta_b(\rho^+) \Subset G$, and $M_{(a_0, \cdot)} \cap \overline{P} = \emptyset$, where

$$P = P_b := \{w \in \mathbb{C}: \rho^- < |w| < \rho^+\}.$$

Let $\gamma: [0, 1] \rightarrow G \setminus M_{(a_0, \cdot)}$ be a curve such that $\gamma(0) = 0$, $\gamma(1) \in \partial \Delta_b(\rho)$. There exists an $\varepsilon = \varepsilon_b > 0$ such that

$$\Delta_{a_0}(\varepsilon) \times ((\gamma([0, 1]) + \Delta_0(\varepsilon)) \cup P) \subset U \setminus M.$$

Put $V = V_b := E \cup (\gamma([0, 1]) + \Delta_0(\varepsilon)) \cup P$ and consider the cross

$$Y = Y_b := \mathbb{X}(A \cap \Delta_{a_0}(\varepsilon), E; \Delta_{a_0}(\varepsilon), V).$$

Then $f \in \mathcal{O}_s(Y)$ for any $f \in \mathcal{O}_s(X \setminus M)$. Consequently, by Theorem 3, any function from $\mathcal{O}_s(X \setminus M)$ extends holomorphically to $\widehat{Y} \supset \{a_0\} \times V$. Shrinking ε and V , we may assume that any function $f \in \mathcal{O}_s(X \setminus M)$ extends to a function $\widetilde{f} = \widetilde{f}_b \in \mathcal{O}(\Delta_{a_0}(\varepsilon) \times W)$, where

$$W = W_b := \Delta_0(1 - \varepsilon) \cup (\gamma([0, 1]) + \Delta_0(\varepsilon)) \cup P.$$

In particular, \tilde{f} is holomorphic in $\Delta_{a_0}(\varepsilon) \times P$, and therefore may be represented by the Hartogs–Laurent series

$$\tilde{f}(z, w) = \sum_{k=0}^{\infty} \tilde{f}_k(z)(w-b)^k + \sum_{k=1}^{\infty} \tilde{f}_{-k}(z)(w-b)^{-k} =: \tilde{f}^+(z, w) + \tilde{f}^-(z, w),$$

$$(z, w) \in \Delta_{a_0}(\varepsilon) \times P,$$

where $\tilde{f}^+ \in \mathcal{O}(\Delta_{a_0}(\varepsilon) \times \Delta_b(\rho^+))$ and $\tilde{f}^- \in \mathcal{O}(\Delta_{a_0}(\varepsilon) \times (\mathbb{C} \setminus \overline{\Delta}_b(\rho^-)))$. Recall that for any $a \in A \cap \Delta_{a_0}(\varepsilon)$ the function $\tilde{f}(a, \cdot)$ extends holomorphically to $G \setminus M_{(a, \cdot)}$. Consequently, for any $a \in A \cap \Delta_{a_0}(\varepsilon)$ the function $\tilde{f}^-(a, \cdot)$ extends holomorphically to $\mathbb{C} \setminus (M_{(a, \cdot)} \cap \overline{\Delta}_b(\rho^-))$. Now, by Theorem 7, there exists a relatively closed pluripolar set $S = S_b \subset \Delta_{a_0}(\varepsilon) \times \overline{\Delta}_b(\rho^-)$ such that:

$$S \cap ((A \cap \Delta_{a_0}(\varepsilon)) \times \overline{\Delta}_b(\rho^-)) \subset M \text{ and}$$

any function \tilde{f}^- extends holomorphically to a function $\tilde{\tilde{f}}^- \in \mathcal{O}(\Delta_{a_0}(\varepsilon) \times \mathbb{C} \setminus S)$.

Since $\tilde{f} = \tilde{f}^+ + \tilde{f}^-$, the function \tilde{f} extends holomorphically to a function $\widehat{f} = \widehat{f}_b \in \mathcal{O}(\Delta_{a_0}(\varepsilon) \times \Delta_b(\rho^+) \setminus S)$. We may assume that the set S is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$.

Using the identity principle and the minimality of the S_b 's, one can easily show that for $b', b'' \in G$, if $B := \Delta_{b'}(\rho_{b'}^+) \cap \Delta_{b''}(\rho_{b''}^+) \neq \emptyset$, then

$$S_{b'} \cap (\Delta_{a_0}(\eta) \times B) = S_{b''} \cap (\Delta_{a_0}(\eta) \times B), \quad \widehat{f}_{b'} = \widehat{f}_{b''} \text{ on } \Delta_{a_0}(\eta) \times B,$$

where $\eta := \min\{\varepsilon_{b'}, \varepsilon_{b''}\}$. Thus the functions $\widehat{f}_{b'}$, $\widehat{f}_{b''}$ and sets $S_{b'}$, $S_{b''}$ may be glued together.

Now, select $b_1, \dots, b_k \in G$ so that $G' \subset \bigcup_{j=1}^k \Delta_{b_j}(\rho_{b_j}^+)$. Put

$$r := \min\{\varepsilon_{b_j}: j = 1, \dots, k\}.$$

Then $S := (\Delta_{a_0}(r) \times G') \cap \bigcup_{j=1}^k S_{b_j}$ gives the required relatively closed pluripolar subset of $\Delta_{a_0}(r) \times G'$ such that $S \cap X \subset M$ and for any $f \in \mathcal{O}_s(X \setminus M)$, the function $\widehat{f} := \bigcup_{j=1}^k \widehat{f}_{b_j}$ extends holomorphically f to $\Delta_{a_0}(r) \times G' \setminus S$. \square

Lemma 12. *Let $A \subset E^p$ be locally pluriregular, let $R > 1$, let*

$$X := \mathbb{X}(A, E^q; E^p, \Delta_0^q(R)),$$

and let $U \subset E^p \times \Delta_0^q(R)$ be an open neighborhood of X . Let $M \subset U$ be a relatively closed set such that $M \cap E^{p+q} = \emptyset$ and for any $a \in A$ the fiber $M_{(a, \cdot)}$ is pluripolar.

Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:

- $\widehat{M} \cap X \subset M$,
- for any $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X \setminus M$,

- the set \widehat{M} is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$.

Notice that the above result is a special case of our Main Theorem with $N = 2$, $n_1 = p$, $D_1 = E^p$, $A_1 = A$, $n_2 = q$, $D_2 = \Delta_0^q(R)$, $A_2 = E^q$, $\Sigma_1 = \Sigma_2 = \emptyset$.

Proof. The case $q = 1$ follows from Lemma 11. Thus assume that $q \geq 2$. By Lemma 10, it suffices to show that for any $a_0 \in A$ and for any $R' \in (1, R)$ there exist $r = r_{R'} > 0$ and a relatively closed pluripolar set $S = S_{R'} \subset \Delta_{a_0}(r) \times \Delta_0^q(R') \subset U$ such that

- $S \cap X \subset M$,
- any function from $\mathcal{O}_s(X \setminus M)$ extends holomorphically to $\Delta_{a_0}(r) \times \Delta_0^q(R') \setminus S$.

Fix an $a_0 \in A$ and let R'_0 be the supremum of all $R' \in (0, R)$ such that $r_{R'}$ and $S_{R'}$ exist. Note that $1 \leq R'_0 \leq R$. It suffices to show that $R'_0 = R$.

Suppose that $R'_0 < R$. Fix $R'_0 < R'' < R$ and choose $R' \in (0, R'_0)$ such that $\sqrt[q]{R'^{q-1}R''} > R'_0$. Let $r := r_{R'}$, $S := S_{R'}$.

Write $w = (w', w_q) \in \mathbb{C}^q = \mathbb{C}^{q-1} \times \mathbb{C}$. Let C denote the set of all $(a, b') \in (A \cap \Delta_{a_0}(r)) \times \Delta_0^{q-1}(R')$ such that the fiber $(M \cup S)_{(a, b', \cdot)}$ is polar. By Lemma 8(a,c), C is pluriregular. Now, by Lemma 11 applied to the cross

$$Y_q := \mathbb{X}(C, \Delta_0(R'); \Delta_{a_0}(r) \times \Delta_0^{q-1}(R'), \Delta_0(R))$$

and the set $M_q := M \cup S$, we conclude that there exists a closed pluripolar set $S_q \subset \widehat{Y}_q$ such that $S_q \cap Y_q \subset M_q$ and any function $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to $\widehat{Y}_q \setminus S_q$. Using the product property of the relative extremal function (cf. [Ngu-Sic 1991]), we get

$$\begin{aligned} \widehat{Y}_q &= \{(z, w', w_q) \in \Delta_{a_0}(r) \times \Delta_0^{q-1}(R') \times \Delta_0(R) : \\ &\quad h_{C, \Delta_{a_0}(r) \times \Delta_0^{q-1}(R')}^*(z, w') + h_{\Delta_0(R'), \Delta_0(R)}^*(w_q) < 1\} \\ &= \{(z, w', w_q) \in \Delta_{a_0}(r) \times \Delta_0^{q-1}(R') \times \Delta_0(R) : \\ &\quad h_{(A \cap \Delta_{a_0}(r)) \times \Delta_0^{q-1}(R'), \Delta_{a_0}(r) \times \Delta_0^{q-1}(R')}^*(z, w') + h_{\Delta_0(R'), \Delta_0(R)}^*(w_q) < 1\} \\ &= \{(z, w', w_q) \in \Delta_{a_0}(r) \times \Delta_0^{q-1}(R') \times \Delta_0(R) : \\ &\quad \max\{h_{A \cap \Delta_{a_0}(r), \Delta_{a_0}(r)}^*(z), h_{\Delta_0^{q-1}(R'), \Delta_0^{q-1}(R')}^*(w')\} + h_{\Delta_0(R'), \Delta_0(R)}^*(w_q) < 1\} \\ &= \{(z, w', w_q) \in \Delta_{a_0}(r) \times \Delta_0^{q-1}(R') \times \Delta_0(R) : \\ &\quad h_{A \cap \Delta_{a_0}(r), \Delta_{a_0}(r)}^*(z) + h_{\Delta_0(R'), \Delta_0(R)}^*(w_q) < 1\}. \end{aligned}$$

Since $R'' < R$, we find a $r_q \in (0, r]$ such that any function $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to a function \widetilde{f}_q on $\Delta_{a_0}(r_q) \times \Delta_0^{q-1}(R') \times \Delta_0(R'') \setminus S_q$. We may assume that S_q is singular with respect to the family $\{\widetilde{f}_q : f \in \mathcal{O}_s(X \setminus M)\}$.

Repeating the above argument for the coordinates w_ν , $\nu = 1, \dots, q-1$, and gluing the obtained sets, we find an $r_0 \in (0, r]$ and a relatively closed pluripolar

set $S_0 := \bigcup_{j=1}^q S_j$ such that any function $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to a function $\tilde{f}_0 := \bigcup_{j=1}^q \tilde{f}_j$ holomorphic in $\Delta_{a_0}(r_0) \times \Omega \setminus S_0$, where

$$\Omega := \bigcup_{\nu=1}^q \Delta_0^{j-1}(R') \times \Delta_0(R'') \times \Delta_0^{q-j}(R').$$

Let $\widehat{\Omega}$ denote the envelope of holomorphy of Ω . Applying the Chirka theorem (Theorem 6), we find a relatively closed pluripolar subset \widehat{S}_0 of $\Delta_{a_0}(r_0) \times \widehat{\Omega}$ such that any function $f \in \mathcal{O}_s(X \setminus M)$ extends to a function \widehat{f} holomorphic on $\Delta_{a_0}(r_0) \times \widehat{\Omega} \setminus \widehat{S}_0$. Let $R''' := \sqrt[q]{R'^{q-1}R''}$. Observe that $\Delta_0(R''') \subset \widehat{\Omega}$. Recall that $R''' > R'_0$. We may assume that \widehat{M} is singular with respect to the family $\{\widehat{f} : f \in \mathcal{O}_s(X \setminus M)\}$. To get a contradiction it suffices to show that $\widehat{M} \cap X \subset M$. We argue as in the proof of Lemma 11:

Take $(a, b) \in (A \cap \Delta_{a_0}(r_0)) \times \Delta_0^q(R''') \setminus M$. Since $M_{(a, \cdot)}$ is pluripolar, there exists a curve $\gamma: [0, 1] \rightarrow \Delta_0(R''') \setminus M_{(a, \cdot)}$ such that $\gamma(0) = 0$, $\gamma(1) = b$. Take an $\varepsilon > 0$ so small that

$$\Delta_a(\varepsilon) \times (\gamma([0, 1]) + \Delta_0^q(\varepsilon)) \subset \Delta_{a_0}(r) \times \Delta_0^q(R''') \setminus M.$$

Put $V_b := E^q \cup (\gamma([0, 1]) + \Delta_0^q(\varepsilon))$ and consider the cross

$$Y := \mathbb{X}(A \cap \Delta_a(\varepsilon), E^q; \Delta_a(\varepsilon), V_b).$$

Then $f \in \mathcal{O}_s(Y)$ for any $f \in \mathcal{O}_s(X \setminus M)$. Consequently, by Theorem 3, $\widehat{Y} \subset \Delta_{a_0}(r) \times \Delta_0^q(R''') \setminus \widehat{M}$, which implies that $\widehat{M}_{(a, \cdot)} \cap \Delta_0^q(R''') \subset M_{(a, \cdot)}$. \square

3. Proof of the Main Theorem for $\mathbf{N} = 2$. To simplify notation put: $p := n_1$, $D := D_1$, $A := A_1$, $A' := A \setminus \Sigma_2$, $q := n_2$, $G := D_2$, $B := A_2$, $B' := B \setminus \Sigma_1$.

It suffices to verify the assumptions of Lemma 10. Let $(D_j)_{j=1}^\infty, (G_j)_{j=1}^\infty$ be approximation sequences: $D_j \Subset D_{j+1} \Subset D$, $G_j \Subset G_{j+1} \Subset G$, $D_j \nearrow D$, $G_j \nearrow G$, $A' \cap D_j \neq \emptyset$, and $B' \cap G_j \neq \emptyset$, $j \in \mathbb{N}$.

Fix a $j \in \mathbb{N}$, $a \in A' \cap D_j$ and let Ω_j be the set of all $b \in G_{j+1}$ such that there exist a polydisc $\Delta_{(a,b)}(r_b) \subset D_j \times G_{j+1}$ and a relatively closed pluripolar set $S_b \subset \Delta_{(a,b)}(r_b)$ such that:

$$S_b \cap ((A' \cap \Delta_a(r_b)) \times \Delta_b(r_b)) \subset M,$$

any function $f \in \mathcal{O}_s(X \setminus M)$ extends to a function $\tilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r_b) \setminus S_b)$ with $\tilde{f}_b = f$ on $(A' \cap \Delta_a(r_b)) \times \Delta_b(r_b) \setminus M$,

and S_b is singular with respect to the family $\{\tilde{f}_b : f \in \mathcal{O}_s(X \setminus M)\}$.

It is clear that Ω_j is open. Observe that $\Omega_j \neq \emptyset$. Indeed, since $B \cap G_j \setminus M_{(a, \cdot)} \neq \emptyset$, we find a point $b \in B \cap G_j \setminus M_{(a, \cdot)}$. Therefore there is a polydisc $\Delta_{(a,b)}(r) \subset D_j \times G_j \setminus M$. Put

$$Y := \mathbb{X}(A \cap \Delta_a(r), B \cap \Delta_b(r); \Delta_a(r), \Delta_b(r)).$$

By Theorem 3, we find an $r_b \in (0, r)$ such that any function $f \in \mathcal{O}_s(X \setminus M)$ extends to $\tilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r_b))$ with $\tilde{f}_b = f$ on $\Delta_{(a,b)}(r_b) \cap Y \supset (A \cap \Delta_a(r_b)) \times \Delta_b(r_b)$. Consequently, $b \in \Omega_j$.

Moreover, Ω_j is relatively closed in G_{j+1} . Indeed, let c be an accumulation point of Ω_j in G_{j+1} and let $\Delta_c(3R) \subset G_{j+1}$. Take a point $b \in \Omega_j \cap \Delta_c(R) \setminus M_{(a,\cdot)}$ and let $r \in (0, r_b]$, $r < 2R$, be such that $\Delta_{(a,b)}(r) \cap M = \emptyset$. Observe that $\tilde{f}_b \in \mathcal{O}(\Delta_{(a,b)}(r))$ and $\tilde{f}_b(z, \cdot) = f(z, \cdot) \in \mathcal{O}(\Delta_b(2R) \setminus M_{(z,\cdot)})$ for any $z \in A' \cap \Delta_a(r)$. Hence, by Lemma 12 (with $R' := R$), there exists a relatively closed pluripolar set $S \subset \Delta_a(\rho') \times \Delta_b(R)$ with $\rho' \in (0, r)$ such that any f has an extension $\widehat{f}_b \in \mathcal{O}(\Delta_a(\rho') \times \Delta_b(R) \setminus S)$. Take an $r_c > 0$ so small that $\Delta_{(a,c)}(r_c) \subset \Delta_a(\rho') \times \Delta_b(R)$ and put $S_c := S \cap \Delta_{(a,c)}(r_c)$, $\tilde{f}_c := \widehat{f}_b$ on $\Delta_{(a,c)}(r_c) \setminus S_c$. Obviously $\tilde{f}_c = \widehat{f}_b = f$ on $(A' \cap \Delta_a(r_c)) \times \Delta_c(r_c) \setminus M$. Hence $c \in \Omega_j$.

Thus $\Omega_j = G_{j+1}$. There exists a finite set $T \subset \overline{G}_j$ such that

$$\overline{G}_j \subset \bigcup_{b \in T} \Delta_b(r_b).$$

Define $r_{a,j} := \min\{r_b : b \in T\}$. Take $b', b'' \in T$ with $C := \Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''}) \neq \emptyset$. Then $\tilde{f}_{b'} = f = \tilde{f}_{b''}$ on $(A' \cap \Delta_a(r_{a,j})) \times (\Delta_{b'}(r_{b'}) \cap \Delta_{b''}(r_{b''})) \setminus M$. Consequently, $\tilde{f}_{b'} = \tilde{f}_{b''}$ on $\Delta_a(r_{a,j}) \times C \setminus (S_{b'} \cup S_{b''})$. In particular, using the minimality of the sets $S_{b'}$ and $S_{b''}$, we conclude that they coincide on $\Delta_a(r_{a,j}) \times C$ and that the functions $f_{b'}$ and $f_{b''}$ glue together. Thus we get a relatively closed pluripolar set $S_{a,j} \subset \Delta_a(r_{a,j}) \times G_j$ such that $S_{a,j} \cap ((A' \cap \Delta_a(r_{a,j})) \times G_j) \subset M$ and any function $f \in \mathcal{O}_s(X \setminus M)$ extends holomorphically to an $f_{a,j} \in \mathcal{O}(\Delta_a(r_{a,j}) \times G_j \setminus S_{a,j})$ with $f_{a,j} = f$ on $(A' \cap \Delta_a(r_{a,j})) \times G_j \setminus M$.

Changing the role of z and w , we get $S^{b,j}$ and $f^{b,j}$, $b \in B' \cap G_j$. \square

The above proof of the Main Theorem for $N = 2$ shows that the following generalization of Lemma 12 is true.

Theorem 13. *Let $D \subset \mathbb{C}^p$, $G \subset \mathbb{C}^q$ be pseudoconvex domains, let $A \subset D$ be locally pluriregular, let $B \subset G$ be open and non-empty, and let $M \subset U$ be a relatively closed subset of an open neighborhood U of the cross $X := \mathbb{X}(A, B; D, G)$ such that $M \cap (D \times B) = \emptyset$ and for any $a \in A$ the fiber $M_{(a,\cdot)}$ is pluripolar. Then there exists a relatively closed pluripolar set $\widehat{M} \subset \widehat{X}$ such that:*

- $\widehat{M} \cap X \subset M$,
- for any $f \in \mathcal{O}_s(X \setminus M)$ there exists exactly one $\widehat{f} \in \mathcal{O}(\widehat{X} \setminus \widehat{M})$ with $\widehat{f} = f$ on $X \setminus M$,
- the set \widehat{M} is singular with respect to the family $\{\widehat{f} : f \in \mathcal{O}_s(X \setminus M)\}$.

Observe that if $G = \mathbb{C}^q$, then $\widehat{X} = D \times \mathbb{C}^q$. Consequently, Theorem 6 generalizes also Theorem 7.

Proof. We apply Lemma 10 (as in the proof of the Main Theorem for $N = 2$). The functions $f_{a,j}$ are constructed exactly as in that proof (with $A' = A$). The

functions $f^{b,j}$ are simply given as $f^{b,j} := f|_{D_j \times \Delta_b(s_{b,j})}$ with $\Delta_b(s_{b,j}) \subset B \cap D_j$ ($S^{b,j} := \emptyset$). \square

4. Proof of the Main Theorem. First observe that, by Lemma 8(b), the set $X' \setminus M$ is not pluripolar. Consequently, the function \widehat{f} is uniquely determined.

We proceed by induction on N . The case $N = 2$ is proved.

Let $D_{j,k} \nearrow D_j$, $D_{j,k} \Subset D_{j,k+1} \Subset D_j$, where $D_{j,k}$ are pseudoconvex domains with $A_{j,k} := A_j \cap D_{j,k} \neq \emptyset$, $j = 1, \dots, N$. Put

$$\begin{aligned} X_k &:= \mathbb{X}(A_{1,k}, \dots, A_{N,k}; D_{1,k}, \dots, D_{N,k}) \subset X, \\ \Sigma_{j,k} &:= (A_{1,k} \times \dots \times A_{j-1,k} \times A_{j+1,k} \times \dots \times A_{N,k}) \cap \Sigma_j, \quad j = 1, \dots, N, \\ X'_k &:= \mathbb{T}(A_{1,k}, \dots, A_{N,k}; D_{1,k}, \dots, D_{N,k}; \Sigma_{1,k}, \dots, \Sigma_{N,k}) \subset X_k. \end{aligned}$$

It suffices to show that for each $k \in \mathbb{N}$ the following condition (*) holds.

(*) There exists a domain U_k , $X'_k \subset U_k \subset \widehat{X}_k$ and a relatively closed pluripolar set $M_k \subset U_k$, such that:

$$M_k \cap X'_k \subset M \text{ and}$$

for any $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\widetilde{f}_k \in \mathcal{O}(U_k \setminus M_k)$ with $\widetilde{f}_k = f$ on $X'_k \setminus M$.

Indeed, fix a $k \in \mathbb{N}$ and observe that, by Lemma 9, \widehat{X}_k is the envelope of holomorphy of U_k . Hence, in virtue of the Chirka theorem (Theorem 6), there exists a relatively closed pluripolar set \widehat{M}_k of \widehat{X}_k , $\widehat{M}_k \cap U_k \subset M_k$, such that $\widehat{X}_k \setminus \widehat{M}_k$ is the envelope of holomorphy of $U_k \setminus M_k$. In particular, for each $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\widehat{f}_k \in \mathcal{O}(\widehat{X}_k \setminus \widehat{M}_k)$ with $\widehat{f}_k|_{U_k \setminus M_k} = \widetilde{f}_k$. We may assume that \widehat{M}_k is singular with respect to the family $\{\widehat{f}_k: f \in \mathcal{O}_s(X \setminus M)\}$.

In particular, $\widehat{M}_{k+1} \cap \widehat{X}_k = \widehat{M}_k$. Consequently:

$\widehat{M} := \bigcup_{k=1}^{\infty} \widehat{M}_k$ is a relatively closed pluripolar subset of \widehat{X} with $\widehat{M} \cap X' \subset M$, for each $f \in \mathcal{O}_s(X \setminus M)$, the function $\widehat{f} := \bigcup_{k=1}^{\infty} \widehat{f}_k$ is holomorphic on $\widehat{X} \setminus \widehat{M}$ with $\widehat{f} = f$ on $X' \setminus M$, and

\widehat{M} is singular with respect to the family $\{\widehat{f}: f \in \mathcal{O}_s(X \setminus M)\}$.

It remains to prove (*). Fix a $k \in \mathbb{N}$. For any

$$a = (a_1, \dots, a_N) \in A_{1,k} \times \dots \times A_{N,k} \setminus M$$

let $\tau = \tau_k(a)$ be such that $\Delta_a(\tau) \subset D_{1,k} \times \dots \times D_{N,k} \setminus M$. Consider the N -fold cross

$$Y_a := \mathbb{X}(A_1 \cap \Delta_{a_1}(\tau), \dots, A_N \cap \Delta_{a_N}(\tau); \Delta_{a_1}(\tau), \dots, \Delta_{a_N}(\tau)).$$

Observe that any function from $\mathcal{O}_s(X \setminus M)$ belongs to $\mathcal{O}_s(Y_a)$. Consequently, by Theorem 3, any function from $\mathcal{O}_s(X \setminus M)$ extends holomorphically to \widehat{Y}_a . Let $\rho = \rho_k(a) \in (0, \tau]$ be such that $\Delta_a(\rho) \subset \widehat{Y}_a$.

If $N \geq 4$, then we additionally define $(N - 2)$ -fold crosses

$$Y_{k,\mu,\nu} := \mathbb{X}(A_{1,k}, \dots, A_{\mu-1,k}, A_{\mu+1,k}, \dots, A_{\nu-1,k}, A_{\nu+1,k}, \dots, A_{N,k}; \\ D_{1,k}, \dots, D_{\mu-1,k}, D_{\mu+1,k}, \dots, D_{\nu-1,k}, D_{\nu+1,k}, \dots, D_{N,k}), \\ 1 \leq \mu < \nu \leq N,$$

and we assume that ρ is so small that

$$\Delta_{(a_1, \dots, a_{\mu-1}, a_{\mu+1}, \dots, a_{\nu-1}, a_{\nu+1}, \dots, a_N)}(\rho) \subset \widehat{Y}_{k,\mu,\nu}, \quad 1 \leq \mu < \nu \leq N.$$

For $j \in \{1, \dots, N\}$, define the 2-fold crosses:

$$Z'_{k,a,j} := \left\{ (z', z_j, z'') : ((A_1 \cap \Delta_{a_1}(\rho)) \times \dots \times (A_{j-1} \cap \Delta_{a_{j-1}}(\rho))) \times D_{j,k+1} \right. \\ \left. \times ((A_{j+1} \cap \Delta_{a_{j+1}}(\rho)) \times \dots \times (A_N \cap \Delta_{a_N}(\rho))) : (z', z'') \notin \Sigma_j \right\} \cup \Delta_a(\rho),$$

$$Z_{k,a,j} := \left((A_1 \cap \Delta_{a_1}(\rho)) \times \dots \times (A_{j-1} \cap \Delta_{a_{j-1}}(\rho)) \times D_{j,k+1} \right. \\ \left. \times (A_{j+1} \cap \Delta_{a_{j+1}}(\rho)) \times \dots \times (A_N \cap \Delta_{a_N}(\rho)) \right) \cup \Delta_a(\rho).$$

Now, we apply Theorem 13 to the 2-fold cross $Z'_{k,a,j}$ and the set M . We find a relatively closed pluripolar set $S_{k,a,j} \subset \widehat{Z}'_{k,a,j} = \widehat{Z}_{k,a,j}$ such that:

$$S_{k,a,j} \cap Z'_{k,a,j} \subset M,$$

for any function $f \in \mathcal{O}_s(X \setminus M)$ there exists an $\widetilde{f}_{k,a,j} \in \mathcal{O}(\widehat{Z}_{k,a,j} \setminus S_{k,a,j})$ such that $\widetilde{f}_{k,a,j} = f$ on $Z'_{k,a,j} \setminus M$,

$$S_{k,a,j} \text{ is singular with respect to the space } \{\widetilde{f}_{k,a,j} : f \in \mathcal{O}_s(X \setminus M)\}.$$

Observe that $\{(a_1, \dots, a_{j-1})\} \times \overline{D}_{j,k} \times \{(a_{j+1}, \dots, a_N)\} \Subset \widehat{Z}_{k,a,j}$. Consequently, we find $r = r_k(a) \in (0, \rho]$ such that

$$V_{k,a,j} := \Delta_{(a_1, \dots, a_{j-1})}(r) \times D_{j,k} \times \Delta_{(a_{j+1}, \dots, a_N)}(r) \subset \widehat{Z}_{k,a,j}, \quad j = 1, \dots, N.$$

Let

$$V_k := \bigcup_{\substack{a \in A_{1,k} \times \dots \times A_{N,k} \setminus M \\ j \in \{1, \dots, N\}}} V_{k,a,j}.$$

Note that $X'_k \subset V_k$. Let U_k be the connected component of $V_k \cap \widehat{X}_k$ that contains X_k .

It remains to glue the sets $S_{k,a,j}$ and functions $\widetilde{f}_{k,a,j}$. Then

$$S_k := \bigcup_{\substack{a \in A_{1,k} \times \dots \times A_{N,k} \setminus M \\ j \in \{1, \dots, N\}}} S_{k,a,j} \cap U_k, \quad \widetilde{f}_k := \bigcup_{\substack{a \in A_{1,k} \times \dots \times A_{N,k} \setminus M \\ j \in \{1, \dots, N\}}} \widetilde{f}_{k,a,j}|_{V_{k,a,j} \cap U_k \setminus S_k}$$

will satisfy (*).

To check that the gluing process is possible, let $a, b \in A_{1,k} \times \cdots \times A_{N,k} \setminus M$, $i, j \in \{1, \dots, N\}$ be such that $V_{k,a,i} \cap V_{k,b,j} \neq \emptyset$. We have the following two cases:

(a) $i \neq j$: We may assume that $i = N - 1$, $j = N$. Write $w = (w', w'') \in \mathbb{C}^{n_1 + \cdots + n_{N-2}} \times \mathbb{C}^{n_{N-1} + n_N}$. Observe that

$$V_{k,a,N-1} \cap V_{k,b,N} = \left(\Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b)) \right) \times \Delta_{b_{N-1}}(r_k(b)) \times \Delta_{a_N}(r_k(a)).$$

We consider the following three subcases:

$N = 2$ (cf. the proof of Lemma 10): Then $V_{k,a,1} \cap V_{k,b,2} = \Delta_{b_1}(r_k(b)) \times \Delta_{a_2}(r_k(a))$. We know that $\tilde{f}_{k,a,1} = \tilde{f}_{k,b,2}$ on the non-pluripolar set

$$(A_1 \cap \Delta_{b_1}(r_k(b)) \setminus \Sigma_2) \times (A_2 \cap \Delta_{a_2}(r_k(a)) \setminus \Sigma_1) \setminus M;$$

cf. Lemma 8(b). Hence, by the identity principle, $\tilde{f}_{k,a,1} = \tilde{f}_{k,b,2}$ on $V_{k,a,1} \cap V_{k,b,2} \setminus (S_{k,a,1} \cup S_{k,b,2})$. Consequently, the sets $S_{k,a,1}$, $S_{k,b,2}$ and the functions $\tilde{f}_{k,a,1}$, $\tilde{f}_{k,b,2}$ glue together.

$N = 3$: Then $V_{k,a,2} \cap V_{k,b,3} = (\Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b)) \times \Delta_{b_2}(r_k(b)) \times \Delta_{a_3}(r_k(a)))$. Let

$$C'' := (A_2 \cap \Delta_{b_2}(r_k(b))) \times (A_3 \cap \Delta_{a_3}(r_k(a))) \setminus \Sigma_1.$$

Recall that for any $c'' \in C''$ the fiber $M_{(\cdot, c'')}$ is pluripolar. We have $\tilde{f}_{k,a,2}(\cdot, c'') = f(\cdot, c'') = \tilde{f}_{k,b,3}(\cdot, c'')$ on $\Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b)) \setminus M_{(\cdot, c'')}$.

Now, let C' denote the set of all $c' \in \Delta_{a_1}(r_k(a)) \cap \Delta_{b_1}(r_k(b))$ such that the fiber $(S_{k,a,2} \cup S_{k,b,3})_{(c', \cdot)}$ is pluripolar. Recall that the complement of C' is pluripolar (Lemma 8(a)). If $c' \in C'$, then $\tilde{f}_{k,a,2}(c', \cdot) = \tilde{f}_{k,b,3}(c', \cdot)$ on $C'' \setminus (S_{k,a,2} \cup S_{k,b,3})_{(c', \cdot)}$. Consequently, by the identity principle, $\tilde{f}_{k,a,2}(c', \cdot) = \tilde{f}_{k,b,3}(c', \cdot)$ on $\Delta_{b_2}(r_k(b)) \times \Delta_{a_3}(r_k(a)) \setminus (S_{k,a,2} \cup S_{k,b,3})_{(c', \cdot)}$, $c' \in C'$. Finally, $\tilde{f}_{k,a,2} = \tilde{f}_{k,b,3}$ on $V_{k,a,2} \cap V_{k,b,3} \setminus (S_{k,a,2} \cup S_{k,b,3})$. Consequently, the sets $S_{k,a,2}$, $S_{k,b,3}$ and the functions $\tilde{f}_{k,a,2}$, $\tilde{f}_{k,b,3}$ glue together.

If $N \in \{2, 3\}$, then we jump directly to (b) and we conclude that the Main Theorem is true for $N \in \{2, 3\}$.

$N \geq 4$: Here is the only place where the induction over N is used. We assume that the Main Theorem is true for $N - 1 \geq 3$.

Let

$$C'' := \{c'' \in (A_{N-1} \cap \Delta_{b_{N-1}}(r_k(b))) \times (A_N \cap \Delta_{a_N}(r_k(a))) : (\Sigma_s)_{(\cdot, c'')} \text{ is pluripolar, } s = 1, \dots, N - 2\};$$

note that, by Lemma 8(a), C'' is non-pluripolar. For any $c'' \in C''$ the function $f_{c''} := f(\cdot, c'')$ is separately holomorphic on $Y_{k,N-1,N} \setminus M_{(\cdot, c'')}$. Moreover, the set

$M_{(\cdot, c'')}$ satisfies all the assumptions of the Main Theorem. Indeed,

$$\Sigma_s(A_{1,k}, \dots, A_{N-2,k}; M_{(\cdot, c'')}) = (\Sigma_s(A_{1,k}, \dots, A_{N,k}; M))_{(\cdot, c'')} \subset (\Sigma_s)_{(\cdot, c'')},$$

$s = 1, \dots, N-2.$

By the inductive assumption, the function $f_{c''}$ extends to a function

$$\widehat{f}_{c''} \in \mathcal{O}(\widehat{Y}_{k,N-1,N} \setminus \widehat{M}(c'')),$$

where $\widehat{M}(c'')$ is relatively closed pluripolar subset of $\widehat{Y}_{k,N-1,N}$ such that $\widehat{M}(c'') \cap Y'_{k,N-1,N} \subset M_{(\cdot, c'')}$. Recall that

$$\Delta_{a'}(r_k(a)) \cup \Delta_{b'}(r_k(b)) \subset \widehat{Y}_{k,N-1,N}.$$

Since $\widetilde{f}_{k,a,N-1}(\cdot, c'') = f_{c''}$ on $\Delta_{a'}(r_k(a)) \cap Y'_{k,N-1,N} \setminus M_{(\cdot, c'')}$ and $\widetilde{f}_{k,b,N}(\cdot, c'') = f_{c''}$ on $\Delta_{b'}(r_k(b)) \cap Y'_{k,N-1,N} \setminus M_{(\cdot, c'')}$, we conclude that $\widetilde{f}_{k,a,N-1}(\cdot, c'') = \widehat{f}_{c''} = \widetilde{f}_{k,b,N}(\cdot, c'')$ on $\Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b)) \setminus M_{(\cdot, c'')}$.

Let $c' \in \Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b))$ be such that the fiber $(S_{k,a,N-1} \cup S_{k,b,N})_{(c', \cdot)}$ is pluripolar. Then $\widetilde{f}_{k,a,N-1}(c', \cdot) = \widetilde{f}_{k,b,N}(c', \cdot)$ on $C'' \setminus (S_{k,a,N-1} \cup S_{k,b,N})_{(c', \cdot)}$. Consequently, by the identity principle, $\widetilde{f}_{k,a,N-1}(c', \cdot) = \widetilde{f}_{k,b,N}(c', \cdot)$ on $(\Delta_{b_{N-1}}(r_k(b)) \times \Delta_{a_N}(r_k(a))) \setminus (S_{k,a,N-1} \cup S_{k,b,N})_{c'}$ and, finally, $\widetilde{f}_{k,a,N-1} = \widetilde{f}_{k,b,N}$ on $(V_{k,a,N-1} \cap V_{k,b,N}) \setminus (S_{k,a,N-1} \cup S_{k,b,N})$. Consequently, the sets $S_{k,a,N-1}$, $S_{k,b,N}$ and the functions $\widetilde{f}_{k,a,N-1}$, $\widetilde{f}_{k,b,N}$ glue together.

(b) $i = j$: We may assume that $i = j = N$. Observe that

$$V_{k,a,N} \cap V_{k,b,N} = \left(\Delta_{(a_1, \dots, a_{N-1})}(r_k(a)) \cap \Delta_{(b_1, \dots, b_{N-1})}(r_k(b)) \right) \times D_{N,k}.$$

By (a) we know that:

$$\begin{aligned} \widetilde{f}_{k,a,N} &= \widetilde{f}_{k,a,N-1} \text{ on } V_{k,a,N} \cap V_{k,a,N-1} \setminus (S_{k,a,N} \cup S_{k,a,N-1}), \\ \widetilde{f}_{k,a,N-1} &= \widetilde{f}_{k,b,N} \text{ on } V_{k,a,N-1} \cap V_{k,b,N} \setminus (S_{k,a,N-1} \cup S_{k,b,N}). \end{aligned}$$

Hence (we write $w = (w', w_N) \in \mathbb{C}^{n_1 + \dots + n_{N-1}} \times \mathbb{C}^{n_N}$):

$$\begin{aligned} \widetilde{f}_{k,a,N} &= \widetilde{f}_{k,b,N} \text{ on } V_{k,a,N} \cap V_{k,a,N-1} \cap V_{k,b,N} \setminus (S_{k,a,N-1} \cup S_{k,a,N} \cup S_{k,b,N}) \\ &= \left(\Delta_{a'}(r_k(a)) \cap \Delta_{b'}(r_k(b)) \right) \times \Delta_{a_N}(r_k(a)) \setminus (S_{k,a,N-1} \cup S_{k,a,N} \cup S_{k,b,N}), \end{aligned}$$

and finally, by the identity principle,

$$\widetilde{f}_{k,a,N} = \widetilde{f}_{k,b,N} \text{ on } V_{k,a,N} \cap V_{k,b,N} \setminus (S_{k,a,N} \cup S_{k,b,N}).$$

Consequently, the sets $S_{k,a,N}$, $S_{k,b,N}$ and the functions $\widetilde{f}_{k,a,N}$, $\widetilde{f}_{k,b,N}$ glue together.

The proof of the Main Theorem is completed. \square

REFERENCES

- [Ale-Zer 2001] O. Alehyane, A. Zeriahi, *Une nouvelle version du théorème d'extension de Hartogs pour les applications séparément holomorphes entre espaces analytiques*, Ann. Polon. Math. 76 (2001), 245–278.
- [Arm-Gar 2001] D. H. Armitage, S. J. Gardiner, *Classical Potential Theory*, Springer Verlag, 2001.
- [Chi 1993] E. M. Chirka, *The extension of pluripolar singularity sets*, Proc. Steklov Inst. Math. 200 (1993), 369–373.
- [Chi-Sad 1988] E. M. Chirka, A. Sadullaev, *On continuation of functions with polar singularities*, Math. USSR-Sb. 60 (1988), 377–384.
- [Jar-Pfl 2000] M. Jarnicki, P. Pflug, *Extension of Holomorphic Functions*, de Gruyter Expositions in Mathematics 34, Walter de Gruyter, 2000.
- [Jar-Pfl 2001a] M. Jarnicki, P. Pflug, *Cross theorem*, IMUJ Preprint 2001/08 (to appear in Ann. Polon. Math. 77 (2001)).
- [Jar-Pfl 2001b] M. Jarnicki, P. Pflug, *An extension theorem for separately holomorphic functions with singularities*, IMUJ Preprint 2001/27 (2001).
- [Jar-Pfl 2001c] M. Jarnicki, P. Pflug, *An extension theorem for separately holomorphic functions with analytic singularities* (to appear).
- [Kaz 1988] M. V. Kazaryan, *On holomorphic continuation of functions with pluripolar singularities*, Dokl. Akad. Nauk Arm. SSR 87 (1988), 106–110.
- [Ngu 1997] Nguyen Thanh Van, *Separate analyticity and related subjects*, Vietnam J. Math. 25 (1997), 81–90.
- [Ngu-Sic 1991] Nguyen Thanh Van, J. Siciak *Fonctions plurisousharmoniques extrémales et systèmes doublement orthogonaux de fonctions analytiques*, Bull. Sc. Math. 115 (1991), 235–244.
- [Ngu-Zer 1991] Nguyen Thanh Van, A. Zeriahi, *Une extension du théorème de Hartogs sur les fonctions séparément analytiques*, Analyse Complexe Multivariables, Récents Développement, A. Meril (ed.), EditEl, Rende, 1991, pp. 183–194.
- [Ngu-Zer 1995] Nguyen Thanh Van, A. Zeriahi, *Systèmes doublement orthogonaux de fonctions holomorphes et applications*, Banach Center Publ. 31 (1995), 281–297.
- [Ökt 1998] O. Öktem, *Extension of separately analytic functions and applications to range characterization of exponential Radon transform*, Ann. Polon. Math. 70 (1998), 195–213.
- [Ökt 1999] O. Öktem, *Extending separately analytic functions in \mathbb{C}^{n+m} with singularities*, Extension of separately analytic functions and applications to mathematical tomography (Thesis), Dep. Math. Stockholm Univ., 1999.
- [Shi 1989] B. Shiffman, *On separate analyticity and Hartogs theorem*, Indiana Univ. Math. J. 38 (1989), 943–957.
- [Sic 1969] J. Siciak, *Separately analytic functions and envelopes of holomorphy of some lower dimensional subsets of \mathbb{C}^n* , Ann. Polon. Math. 22 (1969–1970), 147–171.
- [Sic 1981] J. Siciak, *Extremal plurisubharmonic functions in \mathbb{C}^N* , Ann. Polon. Math. 39 (1981), 175–211.
- [Sic 2000] J. Siciak, *Holomorphic functions with singularities on algebraic sets*, IMUJ Preprint 2000/21 (to appear in Univ. Iag. Acta Math. 75 (2001)).
- [Zah 1976] V. P. Zahariuta, *Separately analytic functions, generalizations of Hartogs theorem, and envelopes of holomorphy*, Math. USSR-Sb. 30 (1976), 51–67.

UNIwersytet Jagielloński
 Instytut Matematyki
 30-059 Kraków, Reymonta 4, Poland
E-mail: jarnicki@im.uj.edu.pl

CARL VON OSSIEZKY UNIVERSITÄT OLDENBURG
 FACHBEREICH MATHEMATIK
 POSTFACH 2503
 D-26111 OLDENBURG, GERMANY
E-mail: pflug@mathematik.uni-oldenburg.de