

# WELL-POSEDNESS FOR THE LINEARIZED MOTION OF THE FREE SURFACE OF A COMPRESSIBLE FLUID

HANS LINDBLAD

## 1. INTRODUCTION

We consider Euler's equations

$$(1.1) \quad \rho(\partial_t + V^k \partial_k) v_j = -\partial_j p, \quad j = 1, \dots, n \quad \text{in } \mathcal{D}, \quad \text{where } \partial_i = \partial/\partial x^i$$

describing the motion of a perfect compressible fluid in vacuum:

$$(1.2) \quad (\partial_t + V^k \partial_k) \rho + \rho \operatorname{div} V = 0, \quad \operatorname{div} V = \partial_k V^k \quad \text{in } \mathcal{D}$$

where the velocity  $v = (v_1, \dots, v_n)$ , the density  $\rho$  and the domain  $\mathcal{D} \subset [0, T] \times \mathbb{R}^n$  are to be determined. Here the pressure  $p = p(\rho)$  is assumed to be a given function of the density which is strictly increasing at the boundary where the density is assumed to be constant. Furthermore  $V^k = \delta^{ki} v_i = v_k$  and we have used the summation convention that repeated upper and lower indices are summed over. We also require the boundary conditions on the free boundary  $\partial\mathcal{D}$ ;

$$(1.3) \quad \begin{cases} p = 0, & \text{on } \partial\mathcal{D} \\ (\partial_t + V^k \partial_k)|_{\partial\mathcal{D}} \in T(\partial\mathcal{D}), \end{cases}$$

The first condition says that the pressure vanishes outside the domain. The second condition says that the boundary should move with the velocity of the fluid particles at the boundary.

Given a simply connected bounded domain  $\mathcal{D}_0 \subset \mathbb{R}^n$  and initial data  $v_0$  and  $\rho_0$ , we want to find a set  $\mathcal{D} \subset [0, T] \times \mathbb{R}^n$  and a vector field  $v$  solving (1.1)-(1.3) and satisfying the initial conditions

$$(1.4) \quad \begin{cases} \{x; (0, x) \in \mathcal{D}\} = \mathcal{D}_0 \\ v = v_0, \quad \rho = \rho_0 \quad \text{on } \{0\} \times \mathcal{D}_0 \end{cases}$$

Let  $\mathcal{D}_t = \{x; (t, x) \in \mathcal{D}\}$  and  $\mathcal{N}$  be the exterior unit normal to the free surface  $\partial\mathcal{D}_t$ . Christodoulou[C2] conjectured the initial value problem (1.1)-(1.4), is well posed in Sobolev spaces under the assumption

$$(1.5) \quad \nabla_{\mathcal{N}} p \leq -c_0 < 0, \quad \text{on } \partial\mathcal{D}, \quad \text{where } \nabla_{\mathcal{N}} = \mathcal{N}^i \partial_{x^i}.$$

(1.5) is a natural *physical condition* since the pressure  $p$  has to be positive in the interior of the fluid. It is essential for the well posedness. In the incompressible irrotational case (1.5) always hold, see [W1,2,CL], and [W1,W2] proved existence in this case. In general Ebin[E1] showed that the incompressible problem is ill posed when (1.5) is not satisfied. In [CL], together with Christodoulou, we proved *a priori* bounds in Sobolev spaces in the general incompressible case of non vanishing curl and in [L1] we showed existence

---

The author was supported in part by the National Science Foundation.

for the linearized equations, assuming that (1.5) hold. In [L1] we also announced an existence proof for the nonlinear problem using the Nash-Moser technique.

For the compressible case there are however no previous existence or well-posedness results. This is due to that the methods used for the irrotational incompressible case reduced the equations to equations on the boundary and this does not work in the compressible case since the divergence is nonvanishing and the pressure satisfies a wave equation in the interior. To be able to deal with the compressible case one therefore needs the techniques developed in [CL,L1] since those use interior estimates.

Here we show existence for the linearized equations and estimates for these in Sobolev spaces in the general compressible case, assuming that (1.5) hold. In a forthcoming paper we use improvements of the existence and estimates here to prove existence for the nonlinear problem using the Nash-Moser technique also in the compressible case. The existence proof here uses the orthogonal decomposition of a vector field into a divergence free part and a gradient. For the divergence free part we get an equation of the type studied in [L1] and for the divergence we get a wave equation on a bounded domain with Dirichlet boundary conditions. The interaction terms between these equations are lower order so if we put up an iteration, the equations decouple for the the new iterate and the previous iterates only enter in the lower order terms.

Existence of solutions for the wave equation on a bounded domain is well known. However dealing with the divergence free part of the equation requires the techniques developed in [L1]. Here we use a generalization of the existence theorem in [L1] to the case when the divergence of the solution we linearize around is nonvanishing. In [L1] we showed that the linearized incompressible Euler's equations becomes an evolution equation for what we called the normal operator. The normal operator is unbounded and not elliptic in the case of nonvanishing curl. It is however positive assuming the physical condition (1.5) and this leads to existence. Up to lower order terms, the projection of the linearized compressible Euler's equations onto divergence free vector fields becomes the linearized incompressible Euler's equations.

As pointed out above, the positive of the pressure (1.5) leads to the positive of the normal operator, introduced in [L1]. It appears that this condition is needed for the well-posedness also in the compressible case since the divergence part essentially decouples from the divergence. In fact, the compressible case was the main motivation for formulating the conjecture (1.5) since in that case it is clear that the pressure has to be positive and in nature one expects fluids to be slightly compressible.

In section 2 we express Euler's equations in the Lagrangian coordinates and derive the linearized equations. In section 3 we define the orthogonal projection and the normal operator. In section 4 we use the projection to decompose the equation into an evolution equation for the normal operator for the divergence free part and a wave equation for the divergence. In section 4 we also give the statement of the existence theorem. In section 5 we give the existence and estimates used for the the divergence free part. In section 6 we give the existence and estimates used for the divergence part. In section 7 we give the *a priori* estimates for the linearized system. In section 8 we discuss the initial and compatibility conditions. Then finally in section 9 we construct an iteration to obtain a solution for the coupled system.

## 2. LAGRANGIAN COORDINATES AND THE LINEARIZED EQUATION.

Let us introduce Lagrangian coordinates in which the boundary becomes fixed. Let  $\Omega$  be a domain in  $\mathbf{R}^n$  and let  $f_0 : \Omega \rightarrow \mathcal{D}_0$  be a diffeomorphism. Assume that  $v(t, x)$ ,  $p(t, x)$ ,  $(t, x) \in \mathcal{D}$  are given satisfying the boundary conditions (1.3)-(1.4). The Lagrangian coordinates  $x = x(t, y) = f_t(y)$  are given by solving

$$(2.1) \quad dx/dt = V(t, x(t, y)), \quad x(0, y) = f_0(y), \quad y \in \Omega$$

Then  $f_t : \Omega \rightarrow \mathcal{D}_t$  is a diffeomorphism, and the boundary becomes fixed in the new  $y$  coordinates. Let us introduce the notation

$$(2.2) \quad D_t = \frac{\partial}{\partial t} \Big|_{y=\text{constant}} = \frac{\partial}{\partial t} \Big|_{x=\text{constant}} + V^k \frac{\partial}{\partial x^k},$$

for the material derivative and

$$(2.3) \quad \partial_i = \frac{\partial}{\partial x^i} = \frac{\partial y^a}{\partial x^i} \frac{\partial}{\partial y^a}.$$

for the partial derivatives. In these coordinates Euler's equation (1.1) become

$$(2.4) \quad \rho D_t^2 x_i + \partial_i p = 0, \quad (t, y) \in [0, T] \times \Omega$$

and the continuity equation (1.2) become

$$(2.5) \quad D_t \rho + \rho \operatorname{div} V = 0, \quad (t, y) \in [0, T] \times \Omega$$

Here the pressure  $p = p(\rho)$  is assumed to be a function of the density  $\rho$ . With  $h$ , the enthalpy, i.e.  $h'(\rho) = p'(\rho)/\rho$  and  $h = 0$  when  $p = 0$ , (2.4) becomes

$$(2.6) \quad D_t^2 x_i + \partial_i h = 0, \quad (t, y) \in [0, T] \times \Omega$$

and the boundary condition (1.3) become

$$(2.7) \quad h|_{\partial\Omega} = 0$$

Since  $\partial_i h = \rho^{-1} \partial_i p$ , Christodoulou's physical condition (1.5) become

$$(2.8) \quad \nabla_{\mathcal{N}} h \leq -c_0 < 0, \quad \text{on } \partial\Omega, \quad \text{where } \nabla_{\mathcal{N}} = \mathcal{N}^i \partial_{x^i}.$$

The enthalpy  $h$  is a function of the density  $\rho$  that satisfies (2.5) but since  $1/\kappa$ , where

$$(2.9) \quad \kappa = \det(\partial x / \partial y),$$

also satisfies (2.5) it follows that  $\rho = \rho_0 \kappa_0 / \kappa$ , where  $\rho_0$  and  $\kappa_0$  are the initial values. By a theorem of [DM] one can arbitrarily prescribe the volume form  $\kappa_0$  up to a constant so we may take  $\kappa_0 = 1/\rho_0$  and  $\Omega$  to be the unit ball, by composing with a diffeomorphism. Hence  $\rho = 1/\kappa$  so  $h$  is a function of  $1/\kappa$ , so we can think of it as a function of  $\sigma = \ln \kappa$ , changing notation:  $h = h(\sigma)$ . Since  $h$  is a function of  $\rho$  we can think of  $\rho = \rho(h)$  as function of  $h$ . Hence with  $e(h) = \ln \rho(h)$  (2.5) instead become

$$(2.10) \quad D_t e(h) + \operatorname{div} V = 0.$$

We have now reduced the system (1.1)-(1.2) to (2.6) and (2.10) and we have reduced the boundary condition (1.3) to (2.7). We have reduced the physical condition (1.5) to (2.8). However, as we will see in below, in order for it to be solvable there are also a couple of other conditions that enter (2.13)-(2.14).

Taking the divergence of (2.6) using (2.10), and that  $[D_t, \partial_i] = -(\partial_i V^k) \partial_k$ , gives a wave equation for  $h$  with Dirichlet boundary conditions:

$$(2.11) \quad D_t^2 e(h) - \Delta h - (\partial_i V^k) \partial_k V^i = 0, \quad h|_{\partial\Omega} = 0.$$

Here

$$(2.12) \quad \Delta h = \sum_i \partial_i^2 h = \kappa^{-1} \partial_a (\kappa g^{ab} \partial_b h), \quad \text{where } g_{ab} = \delta_{ij} \frac{\partial x^i}{\partial y^a} \frac{\partial x^j}{\partial y^b},$$

is the metric in the Lagrangian coordinates and  $g^{ab}$  is its inverse. Here  $\partial_a = \partial/\partial y^a$  and we use the convention that differentiation with respect to the Eulerean coordinates is denoted by letters  $i, j, k, l, m, n$  and with respect to the Lagrangian coordinates is denoted by  $a, b, c, d, e, f$ .

In order for (2.11) to be solvable we must have that

$$(2.13) \quad 0 < c_1 \leq e' \leq 1/c_1 < \infty, \quad c_1 |\xi|^2 \leq g^{ab} \xi_a \xi_b \leq |\xi|^2 / c_1$$

for some constant  $c_1 > 0$ . The first condition is related to that the pressure is assumed to be a strictly increasing function of the pressure. The second condition is related to that the geometry of the boundary  $\partial \mathcal{D}_t$  is bounded. Without getting further into the physical motivation of the condition (2.13) we remark that from a mathematical point of view it is necessary for the well-posedness of the problem since one can not solve the wave equation (2.13) without this condition.

In order for (2.11) to have a formal solution in  $C^{r-1}$  we must also assume that the following  $r$  compatibility conditions on initial data are satisfied

$$(2.14) \quad D_t^j h|_{\partial \Omega} = 0, \quad \text{when } t = 0, \quad \text{for } j = 0, \dots, m-1.$$

For  $j = 0, 1$  this simply means that the initial conditions should vanish on the boundary, for  $j = 2$  this means that the sum of the last two terms (2.11) should vanish on the boundary and for larger  $j$  one gets more conditions from taking time derivatives of (2.11).

Let us now derive the linearized equations. The calculations that follows below are partly identical to those in [L1] since the equation (2.6) mathematically is the same as the equation for the incompressible case, its only a difference of notation since the enthalpy  $h$  plays the same that the pressure  $p$  played in the incompressible case. We therefore refer the reader to [L1] for more details of the calculations.

If  $\delta$  is a variation in the Lagrangian coordinates, i.e. derivative w.r.t. a parameter when  $t$  and  $y$  are fixed, then  $[\delta, D_t] = 0$ ,  $[\delta, \partial_i] = -(\partial_i \delta x^k) \partial_k$  so  $[\delta - \delta x^k \partial_k, \partial_i] = 0$ . Applying  $\delta - \delta x^k \partial_k$  to (2.6) gives:

$$(2.15) \quad D_t^2 \delta x_i - (\partial_k D_t^2 x_i) \delta x^k - \partial_i (\delta x^k \partial_k h - \delta h) = 0$$

Using that also  $[\delta - \delta x^k \partial_k, \Delta] = 0$  we get from applying  $\delta - \delta x^k \partial_k$  to (2.11)

$$(2.16) \quad D_t^2 (e'(h) \delta h) - \delta x^k \partial_k D_t^2 e(h) - \Delta (\delta h - \delta x^k \partial_k h) - 2(\partial_i V^k) \partial_k (\delta V^i - \delta x^l \partial_l V^i) = 0, \quad \delta h|_{\partial \Omega} = 0.$$

where  $\delta V^i = D_t \delta x^i$ . Since we are considering a variation within families of solutions satisfying the  $m$  compatibility conditions (2.14) this then translates into the same  $m$  compatibility conditions for the variation in order for the last part of (2.16) to hold. These are

$$(2.17) \quad D_t^j \delta h|_{\partial \Omega} = 0, \quad \text{when } t = 0, \quad \text{for } j = 0, \dots, m-1.$$

Since

$$(2.18) \quad \begin{aligned} [\partial_i, D_t^2] \delta x^i &= D_t ([\partial_i, D_t] \delta x^i) + [\partial_i, D_t] D_t \delta x^i = 2(\partial_i V^k) (\partial_k \delta V^i - (\partial_k V^l) \partial_l \delta x^i) + (\partial_i D_t V^k) \partial_k \delta x^i \\ &= 2(\partial_i V^k) \partial_k (\delta V^i - \delta x^l \partial_l V^i) + \delta x^l \partial_l ((\partial_i V^k) \partial_k V^i) + \partial_k ((\partial_i D_t V^k) \delta x^i) - (\partial_i \partial_k D_t V^k) \delta x^i \\ &= 2(\partial_i V^k) \partial_k (\delta V^i - \delta x^l \partial_l V^i) + \partial_k ((\partial_i D_t V^k) \delta x^i) - (\partial_i D_t \operatorname{div} V) \delta x^i \end{aligned}$$

we get by taking the divergence of (2.15):

$$(2.19) \quad D_t^2 \operatorname{div} \delta x - \delta x^i \partial_i D_t \operatorname{div} V - \Delta(\delta x^k \partial_k h - \delta h) + 2(\partial_i V^k) \partial_k (\delta V^i - \delta x^l \partial_l V^i) = 0$$

Hence if we add (2.19) and (2.16) and use (2.10) we get

$$(2.20) \quad D_t^2 (\operatorname{div} \delta x + e'(h) \delta h) = 0,$$

It follows that

$$(2.21) \quad \operatorname{div} \delta x = -e' \delta h,$$

Actually, this also follows from our original assumption that  $h$  was a function of the Jacobian  $\kappa$  since if  $\sigma = \ln \kappa$  then  $\delta \sigma = \operatorname{div} \delta x$ , see [L1], so  $\delta h(\sigma) = h' \delta \sigma = h' \operatorname{div} \delta x = \dots = -\operatorname{div} \delta x / e'$ .

The basic assumption in solving the system (2.15) and (2.16) is that one should assume that  $\operatorname{div} \delta x$  has the same space regularity as  $\delta x$ .

Let us now express also the vector field in the Lagrangian frame. Let

$$(2.22) \quad W^a = \frac{\partial y^a}{\partial x^i} \delta x^i, \quad \dot{g}_{ab} = D_t g_{ab}, \quad \omega_{ab} = \partial_a v_b - \partial_b v_a,$$

where  $v_a = (\partial x^i / \partial y^a) D_t x_i$ , then since the divergence is invariant

$$(2.23) \quad \operatorname{div} \delta x = \operatorname{div} W = \kappa^{-1} \partial_a (\kappa W^a)$$

The linearized equation (2.15), see (2.9)-(2.20) in [L1], become

$$(2.24) \quad g_{ab} D_t^2 W^b - \partial_a ((\partial_c h) W^c - \delta h) = -(\dot{g}_{ac} - \omega_{ac}) D_t W^c + F^a,$$

where we also have added an inhomogeneous term  $F$ . The reason for adding the inhomogeneous term is because its simpler to move the initial conditions into an inhomogeneous term. If we have initial conditions satisfying constraint conditions at the boundary up to a certain order one can form a power series satisfying the equation up to that order as  $t \rightarrow 0$ . Subtracting of the power series yields an equation with vanishing initial conditions and instead an inhomogeneous term vanishing to that order minus two as  $t \rightarrow 0$ .

The idea is now to decompose the equation into a divergence free part, to be dealt with using the methods from the incompressible case in [L1] and a divergence part, to be dealt with using estimates and existence for wave equations on a bounded domain. The interaction terms turn out to be lower order.

### 3. THE PROJECTION AND THE NORMAL OPERATOR.

We will now make an orthogonal decomposition:  $\mathcal{H} = L^2 = \mathcal{H}_d \oplus \mathcal{H}_g$  into divergence free vector fields and gradients of functions in  $H_0^1(\Omega)$ . Let us therefore define the orthogonal projection  $P$  onto divergence free vector fields by

$$(3.1) \quad PU^a = U^a - g^{ab} \partial_b p_U, \quad \Delta p_U = \operatorname{div} U, \quad p_U|_{\partial\Omega} = 0$$

(Here  $\Delta q = \kappa^{-1} \partial_a (\kappa g^{ab} \partial_b q)$ . )  $P$  is the orthogonal projection in the inner product, see [L1],

$$(3.2) \quad \langle U, W \rangle = \int_{\Omega} g_{ab} U^a W^b \kappa dy$$

Note also that, with

$$(3.3) \quad \|W(t)\|_r = \|W(t, \cdot)\|_{H^r(\Omega)} = \sum_{k=0}^r \|\partial^k W(t, \cdot)\|_{L^2(\Omega)}, \quad \|W(t)\|_{r,s} = \sum_{k=0}^s \|\hat{D}_t^k W(t, \cdot)\|_{H^r}$$

denoting the Sobolev norms for fixed time with space and time differentiation of order  $r$  and  $s$ , where  $\hat{D}_t$  is given by (4.2), we have

$$(3.4) \quad \|PW\|_{r,s} \leq C\|W\|_{r,s}, \quad \|(I - P)W\|_{r,s} \leq C\|\operatorname{div} W\|_{r-1,s}$$

since its just a matter of solving the Dirichlet problem and commuting through time derivatives.

We will now define the normal operator, which is a symmetric operator on divergence free vector fields introduced in [L1]. For a function  $f$  that vanishes on the boundary define  $A_f W^a = g^{ab} \underline{A}_f W_b$ , where

$$(3.5) \quad \underline{A}_f W_a = -\partial_a ((\partial_c f) W^c - q), \quad \Delta((\partial_c f) W^c - q) = 0, \quad q|_{\partial\Omega} = 0$$

and we restrict our attention to divergence free vector fields. If  $U$  and  $W$  are divergence free then

$$(3.6) \quad \langle U, A_f W \rangle = \int_{\partial\Omega} n_a U^a (-\partial_c f) W^c dS$$

where  $n$  is the unit connormal. If  $f|_{\partial\Omega} = 0$  then  $-\partial_c f|_{\partial\Omega} = (-\nabla_N f) n_c$ . It follows that  $A_f$  is a symmetric operator on divergence free vector fields, and in particular the normal operator

$$(3.7) \quad A = A_h$$

where  $h$  is the enthalpy is positive since we assumed the physical condition that  $-\nabla_N h \geq c_0 > 0$  on the boundary. The normal operator is order one, by (3.4)

$$(3.8) \quad \|AW\|_{r,s} \leq C\|W\|_{r+1,s}.$$

The normal operator has certain delicate commutator properties with vector fields and positivity properties which were essential for the existence proof in [L1]. The main difficulty being that it is not elliptic acting on vector fields with nonvanishing curl. Furthermore, in order to prove existence one had to replace it by a sequence of bounded operator which uniformly had the same commutator and positivity properties. Here we will only use a generalization of the existence proof in [L1] and we will explain what needs to be changed.

#### 4. DECOMPOSITION OF THE EQUATION INTO A DIVERGENCE FREE PART AND A DIVERGENCE PART AND STATEMENT OF THE THEOREM.

We now want to decompose the linearized equations (2.24)

$$(4.1) \quad D_t^2 W^a - g^{ab} \partial_b ((\partial_c h) W^c - \delta h) = -g^{ab} (\dot{g}_{bc} - \omega_{bc}) D_t W^c + F^a, \quad \delta h = -\operatorname{div} W / e'$$

into an equation for the divergence free part and an equation for the divergence. Now,  $D_t$  does not commute with taking the divergence, if  $\operatorname{div} V \neq 0$ , so we will replace it by a modified time derivative that does:

$$(4.2) \quad \hat{D}_t = D_t + \operatorname{div} V, \quad \text{ie} \quad \hat{D}_t W^a = D_t W^a + (\operatorname{div} V) W^a = \kappa^{-1} D_t (\kappa W^a)$$

Then it follows from (2.23) that  $\hat{D}_t \operatorname{div} W = \operatorname{div} \hat{D}_t W$ , since if  $\sigma = \ln \kappa$  then  $\dot{\sigma} = D_t \sigma = \operatorname{div} V$ , see [L1]. With this notation we have  $\hat{D}_t^2 = (D_t + \operatorname{div} V)(D_t + \operatorname{div} V) = D_t^2 + 2\dot{\sigma} D_t + \dot{\sigma}^2 + \ddot{\sigma} = D_t^2 + 2\dot{\sigma} \hat{D}_t + \ddot{\sigma} - \dot{\sigma}^2$  so

$$(4.3) \quad D_t^2 = \hat{D}_t^2 - 2\dot{\sigma} \hat{D}_t + \dot{\sigma}^2 - \ddot{\sigma}, \quad D_t = \hat{D}_t - \dot{\sigma}$$

Hence, with  $\dot{W} = \hat{D}_t W$  and  $\ddot{W} = \hat{D}_t^2 W$ , we can write the equation

$$(4.4) \quad \ddot{W}^a - g^{ab} \partial_b ((\partial_c h) W^c - \delta h) = B^a(W, \dot{W}) + F^a, \quad \delta h = -\operatorname{div} W / e'$$

where

$$(4.5) \quad B^a(W, \dot{W}) = -g^{ab} (\dot{g}_{bc} - \omega_{bc}) (\dot{W}^c - \dot{\sigma} W^c) + 2\dot{\sigma} \dot{W}^a + (\ddot{\sigma} - \dot{\sigma}^2) W^a$$

Taking the divergence of (4.4) gives

$$(4.6) \quad \hat{D}_t^2 (e' \delta h) - \Delta (\delta h - (\partial_c h) W^c) = -\operatorname{div} B(W, \dot{W}) - \operatorname{div} F$$

where we assumed that  $0 < c_1 < e' < 1/c_1 < \infty$  for some constant  $c_1$ . We get

$$(4.7) \quad e' \hat{D}_t^2 \delta h - \Delta (\delta h - (\partial_c h) W^c) = G(W, \dot{W}, \partial W, \partial \dot{W}, \delta h, \hat{D}_t \delta h) - \operatorname{div} F, \quad \operatorname{div} W_1 = -e' \delta h$$

where

$$(4.8) \quad G(W, \dot{W}, \partial W, \partial \dot{W}, \delta h, \hat{D}_t \delta h) = -\operatorname{div} B(W, \dot{W}) + 2(D_t e') \hat{D}_t \delta h + (D_t^2 e') \delta h, \quad W = W_0 + W_1$$

We now, also want to project (4.4) onto the divergence free vector fields. Let us therefore decompose into a divergence free part and a divergence part.

$$(4.9) \quad W_0 = W - W_1, \quad W_1^a = g^{ab} \partial_b p_W, \quad \Delta p_W = \operatorname{div} W, \quad p_W|_{\partial\Omega} = 0$$

Note that by (3.4)

$$(4.10) \quad \|W_1\|_{r+1} \leq C \|\operatorname{div} W_1\|_r, \quad \|\dot{W}_1\|_r \leq C \|\hat{D}_t \operatorname{div} W\|_{r-1}$$

If  $W_0$  is divergence free then obviously  $\ddot{W}_0$  is divergence free so if  $P$  denotes the orthogonal projection onto the divergence free vector fields then we need to calculate  $P(\hat{D}_t^2 W_1)$ , where  $W_1^a = (I - P)W^a = g^{ab} \partial_b q$  and  $\Delta q = \operatorname{div} W$ ,  $q|_{\partial\Omega} = 0$ . With  $k^{ab} = \hat{D}_t g^{ab}$  and  $\dot{k}^{ab} = \hat{D}_t k^{ab}$  and  $\dot{q} = D_t q$ ,  $\ddot{q} = D_t^2 q$  we get

$$(4.11) \quad \begin{aligned} \hat{D}_t^2 W_1^a &= \hat{D}_t^2 (g^{ab} \partial_b q) = \dot{k}^{ab} \partial_b p_W + 2k^{ab} \partial_b D_t \dot{q} + g^{ab} \partial_b \ddot{q} \\ &= \dot{k}^{ab} \partial_b q + 2k^{ab} g_{bc} (\hat{D}_t (g^{cd} \partial_b q) - k^{cd} \partial_d q) + g^{ab} \partial_b \ddot{q} = \dot{k}^a_c W^c + 2k^a_b (\dot{W}_1^b - k_{bd} W^d) + g^{ab} \partial_b \ddot{q} \end{aligned}$$

If we project to the divergence free vector fields then the last term vanishes since  $\tilde{q}|_{\partial\Omega} = 0$  so we get

$$(4.12) \quad P(\hat{D}_t^2 W^a) = \hat{D}_t^2 W_0^a + P(\dot{k}^a_c W_1^c + 2k^a_b(\dot{W}_1^b - k_{bd}W_1^d)), \quad k^{ab} = \hat{D}_t g^{ab}, \quad \dot{k}^{ab} = \hat{D}_t^2 g^{ab}$$

We now want to project (4.4) onto the divergence free vector fields. The most important thing in doing to so is that the projection of the gradient of  $\delta h$  vanishes since  $\delta h|_{\partial\Omega} = 0$ . We therefore get

$$(4.13) \quad \ddot{W}_0^a + AW_0^a = PH^a(W_0, \dot{W}_0, W_1, \dot{W}_1, \partial W_1) + PF^a$$

where  $AW^a = -P(g^{ab}\partial_b((\partial_c h)W_0^c))$  is the normal operator and

$$(4.14) \quad H^a(W_0, \dot{W}_0, W_1, \dot{W}_1, \partial W_1) = g^{ab}\partial_b((\partial_c h)W^c) - g^{ab}(\dot{g}_{bc} - \omega_{bc})(\dot{W}^c - \dot{\sigma}W^c) + 2\dot{\sigma}\dot{W}^a + (\ddot{\sigma} - \dot{\sigma}^2)W^a + \dot{k}^a_c W_1^c + 2k^a_b(\dot{W}_1^b - k_{bd}W_1^d)$$

The equations (4.1) for  $W$  are hence replaced by (4.7) and (4.13) for  $W_0$  and  $W_1$ . We will solve this system for  $(W_0, W_1)$  nad we hence get a solution  $W = W_0 + W_1$  of (4.1).

We will now prove existence for the linearized equations

$$(4.15) \quad \ddot{W}^a - g^{ab}\partial_b((\partial_c h)W^c - \delta h) = B^a(W, \dot{W}) + F^a, \quad \text{where} \quad \delta h = -\text{div } W / e'$$

and  $B$  is given by (4.5), with initial conditions

$$(4.16) \quad W|_{t=0} = \tilde{W}^0, \quad \dot{W}|_{t=0} = \tilde{W}^1$$

assuming that  $F$  vanishes to infinite order as  $t \rightarrow 0$  and the compatibility conditions for initial data described below hold to any order.

Suppose that we have a solution of (4.15)-(4.16) where  $F$  vanishes to any order as  $t \rightarrow 0$ . Then, if we move the second term in the left of (4.15) into the right hand side and differentiate with respect to time we get

$$(4.17) \quad \hat{D}_t^{k+2}W = B_k(W, \cdot, \hat{D}_t^{k+1}W, \partial W, \dots, \partial \hat{D}_t^k W, \partial^2 W, \dots, \partial^2 \hat{D}_t^k W) + \hat{D}_t^k F,$$

for some function  $B_k$ , where  $\hat{D}_t^k F|_{t=0} = 0$ . Let us therefore define functions of space only by

$$(4.18) \quad \tilde{W}^{k+2} = B_k(\tilde{W}^0, \dots, \tilde{W}^{k+1}, \partial \tilde{W}^0, \dots, \partial \tilde{W}^k, \partial^2 \tilde{W}^0, \dots, \partial^2 \tilde{W}^k)|_{t=0}, \quad k \geq 0$$

Since  $\hat{D}_t^k \text{div } W = \text{div } \hat{D}_t^k W$ , the compatibility conditions for solving the wave equation (2.17) become:

$$(4.19) \quad \text{div } \tilde{W}^k|_{\partial\Omega} = 0, \quad \text{for} \quad k \leq m-1$$

(4.19) is called the  $m^{th}$  order compatibility condition. Since  $\tilde{W}^k$  are determined from initial data (4.16) this gives some compatibility conditions on initial data. We now assume that initial data in (4.16) are smooth and that the compatibility conditions hold for all orders  $m = 0, 1, \dots$

We will prove the following theorem:



**Theorem 4.1.** *Suppose that  $(x, h, e')$  are smooth and that (2.7), (2.8) and (2.13) hold for  $0 \leq t \leq T$ , Suppose that the inhomogeneous term  $F$  in (4.15) is smooth for  $0 \leq t \leq T$  and vanishes to any order as  $t \rightarrow 0$ . Suppose also that the initial conditions (4.16) are smooth and satisfy all the  $m^{\text{th}}$  order compatibility conditions (4.19), for  $m = 0, 1, \dots$ . Then the linearized equations (4.15)-(4.16) have a smooth solution for  $0 \leq t \leq T$ .*

*Furthermore the solution satisfy the following energy estimates. Let  $W = W_0 + W_1$  be the decomposition (4.9) and set*

$$(4.20) \quad E_r(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r} + \|\operatorname{div} W_1(t)\|_r + \|\operatorname{div} \dot{W}_1(t)\|_{r-1}$$

*where  $\dot{W} = \hat{D}_t W = D_t W + (\operatorname{div} V)W$  and  $\langle W_0 \rangle_{A,r}$  is defined in the next section. Then if  $r \geq 1$  there is a constant  $C$  depending only on  $(x, h, e')$ ,  $r$  and  $T$  such that for  $0 \leq t \leq T$  we have*

$$(4.21) \quad E_r(t) \leq C \left( E_r(0) + \int_0^t \|F\|_r d\tau \right)$$

*Here  $\|F\|_r$  is defined in the next section, (5.5). Furthermore*

$$(4.22) \quad E_1(t) \leq C \left( E_1(0) + \int_0^t \|F\|_1 d\tau \right)$$

*and if*

$$(4.23) \quad \tilde{E}_r(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r} + \|\operatorname{div} W_1(t)\|_r$$

*then we also have:*

$$(4.24) \quad \tilde{E}_r(t) \leq C \left( E_r(0) + \int_0^t \|F\|_r d\tau \right)$$

*Remark.* We have for simplicity stated the result for infinitely differentiable initial data that satisfy the compatibility conditions (4.18) for all orders  $m = 0, 1, \dots$ . The proof actually gives that if the  $r^{\text{th}}$  order compatibility condition is satisfied then we get a solution in the spaces associated with the  $r^{\text{th}}$  energy provided that initial data has some additional regularity  $(\tilde{W}^0, \tilde{W}^1) \in H^{2r+1} \times H^{2r}$ . However one should be able to just assume that the  $r^{\text{th}}$  order energy (4.20) of data is bounded initially, and recover the loss of regularity of the solution afterwards, see [Ev]. In our application of this result to the nonlinear problem using Nash-Moser, we have vanishing initial data and the compatibility conditions are fulfilled to any order.

In the proof that follows we assume that  $(x, h, e')$  are fixed smooth functions satisfying the conditions (2.7), (2.8) and (2.13). The proof of existence will use energy norms to any order  $r$ . All constants  $C$  in the proof that follows will depend on some  $C^{r'}$  norms of  $(x, h, e')$ , for some  $r'$  depending on the order  $r$  of the energy norm of  $W$  that we are estimating, but we will not calculate the explicit dependence here.

## 5. EXISTENCE AND ESTIMATES IN THE DIVERGENCE FREE CLASS.

In [L1] we proved existence for solutions of

$$(5.1) \quad \hat{D}_t^2 W_0 + A W_0 = H, \quad W_0|_{t=0} = \tilde{W}_0^0, \quad \dot{W}_0|_{t=0} = \tilde{W}_0^1$$

We have

**Theorem 5.1.** Suppose that  $x, h \in C^{r+2}([0, T] \times \Omega)$ ,  $h|_{\partial\Omega} = 0$ ,  $\nabla_N h|_{\partial\Omega} \leq -c_0 < 0$ . Then if initial data in (5.1) and the inhomogeneous term in (5.1) are divergence free and satisfy

$$(5.2) \quad (\tilde{W}_0^0, \tilde{W}_0^1) \in N^r(\Omega) \times H^r(\Omega), \quad H \in L^1([0, T], H^r(\Omega))$$

the linearized equations (5.1) have a solution

$$(5.3) \quad (W_0, \dot{W}_0) \in C([0, T], N^r(\Omega) \times H^r(\Omega)).$$

Furthermore, with a constant  $C$  depending only on the norm of  $x$  and  $h$ ,  $T$  and the constant  $c_0$  we have

$$(5.4) \quad E_{0r}(t) \leq C \left( E_{0r}(0) + \int_0^t \|H(\tau)\|_r d\tau \right), \quad E_{0r}(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r}$$

Here

$$(5.5) \quad \langle W \rangle_{A,r} = \sum_{|I| \leq r, I \in \mathcal{S}} \langle \mathcal{L}_S^I W, A \mathcal{L}_S^I W \rangle^{1/2}$$

where  $\mathcal{S}$  is a spanning set of tangential space derivatives,  $\mathcal{L}_S^I$  is a product of  $|I|$  Lie derivatives with respect to these vector fields.  $A$  is the normal operator which is positive so  $\langle W \rangle_{A,r}$  is a seminorm. Here  $H^r(\Omega)$  is a space which is equivalent to the usual Sobolev spaces.  $N^r(\Omega)$  is the completion of the  $C^\infty$  divergence free vector fields in the norm  $\|W\|_{N^r} = \|W\|_{H^r} + \langle W \rangle_{A,r}$ .  $W \in N^r(\Omega)$  is equivalent to that  $W$  is divergence free,  $W \in H^r(\Omega)$  and the normal component  $W_N = N_a W^a$  is in  $H^r(\partial\Omega)$ .

*Proof.* In case  $\text{div } V = 0$  this was proven in [L1] and the proof there can be easily modified by multiplying or dividing by  $\kappa = \det(\partial x / \partial y)$ . Let us now indicate what needs to be changed in [L1]. We can use the same set of tangential vector fields as in [L1], but now they are no longer divergence free so it is no longer true that a Lie derivative with respect to these preserves the divergence free condition. But one can easily modify the Lie derivative so it preserves the divergence free condition. The modified Lie derivative applied to a vector field is

$$(5.6) \quad \hat{\mathcal{L}}_T W = \mathcal{L}_T W + (\text{div } T)W,$$

It satisfies  $\text{div } \hat{\mathcal{L}}_T W = \hat{T} \text{div } W$ , where  $\hat{T} = T + (\text{div } T)$ . One then has to make it so one always apply this modified Lie derivative to vector fields. However we use the usual Lie derivative, when applied to one forms since we use that it commutes with covariant differentiation. In deriving the estimates for all components of a vector field in terms of the divergence, the curl and the tangential components we then use that  $\mathcal{L}_T(g_{ab}W^b) = g_{ab}\hat{\mathcal{L}}_T W^b + (\check{\mathcal{L}}_T g_{ab})W^b$ , where  $\check{\mathcal{L}}_T g_{ab} = \mathcal{L}_T g_{ab} - (\text{div } T)g_{ab}$ . Let us now examine how the critical commutator with the normal operator, (3.5), is changed from what it was in [L1]. We have

$$(5.7) \quad \mathcal{L}_T \underline{A}_f W_a = \mathcal{L}_T \partial_a ((\partial_c f)W^c - q) = \partial_a ((\partial_c f)\hat{\mathcal{L}}_T W + \partial_c(\check{T}f)W^c - Tq + f(\partial_c \text{div } T)W^c)$$

where  $\check{T}f = Tf - (\text{div } T)f$ . When we project again the last two terms vanish since they vanish on the boundary so the commutator relation in [L1] will be replaced by

$$(5.8) \quad P \mathcal{L}_T \underline{A} W = \underline{A}_f \hat{\mathcal{L}}_T W + \underline{A}_{\check{T}f} W$$

The issue of how to deal with the initial conditions in case  $\text{div } V \neq 0$  is discussed in section 8.  $\square$

Now, the norms used in Theorem 5.1 are natural for the initial value problem. However, when solving the wave equation with Dirichlet boundary conditions it is more natural to first look on norms with many time derivatives. Because of the coupling between the two equations we must therefore also estimate more time derivatives of the divergence free part. From differentiating the equation (5.1) we get:

**Lemma 5.2.** *Let  $E_{0,r}$  be as in Theorem 5.1 and let*

$$(5.9) \quad |||W(t)|||_r = \sum_{s+k \leq r} \|W(t)\|_{k,s} \quad \text{where} \quad \|W(t)\|_{r,s} = \sum_{k=0}^s \|\hat{D}_t^k W(t)\|_r$$

Then

$$(5.10) \quad |||\dot{W}_0|||_r + |||W_0|||_r \leq C(E_{0,r} + |||H|||_{r-1}), \quad r \geq 1$$

*Proof.* The proof is just differentiation of (5.1) using that  $A$  is order one, (3.8);

$$(5.11) \quad |||\ddot{W}_0|||_{r-1} \leq C(|||W_0|||_r + |||H|||_{r-1}), \quad r \geq 1$$

which proves (5.10) for  $r = 1$  so we may assume that  $r \geq 2$  in (5.10). We have

$$(5.12) \quad |||W_0|||_r \leq \|W_0\|_r + \|\dot{W}_0\|_{r-1} + |||\ddot{W}_0|||_{r-2}, \quad r \geq 2$$

so together with (5.11) we get

$$(5.13) \quad |||W_0|||_r \leq C(E_{0,r} + |||W_0|||_{r-1} + |||H|||_{r-2}), \quad r \geq 2$$

Since also  $|||W_0|||_1 \leq E_{0,1}$  we can use induction in  $r$  to prove that

$$(5.14) \quad |||W_0|||_r \leq C(E_{0,r} + |||H|||_{r-2}), \quad r \geq 2$$

Similarly

$$(5.15) \quad |||\dot{W}_0|||_r \leq \|\dot{W}_0\|_r + |||\ddot{W}_0|||_{r-1}, \quad r \geq 1$$

so by (5.11) again

$$(5.16) \quad |||\dot{W}_0|||_r \leq C(E_{0,r} + |||W_0|||_r + |||H|||_{r-1}), \quad r \geq 1$$

(5.10) for  $r \geq 2$  now follows from (5.14) and (5.16).  $\square$

## 6. EXISTENCE AND ESTIMATES FOR THE WAVE EQUATION.

Existence of solutions for a wave equation with Dirichlet boundary conditions and initial conditions satisfying some compatibility conditions is well known, see e.g. [H]. The result in [H] is stated with vanishing initial condition but we will reduce to that case by subtracting of the  $m^{th}$  order Taylor series. We consider the Cauchy problem for the wave equation on a bounded domain with Dirichlet boundary conditions:

$$(6.1) \quad f\hat{D}_t^2\psi - \triangle\psi = \triangle\phi + G = \tilde{G}, \quad \text{in } \Omega, \quad \psi|_{\partial\Omega} = 0,$$

$$(6.2) \quad \psi|_{t=0} = \psi_0, \quad \hat{D}_t\psi|_{t=0} = \psi_1$$

Here

$$(6.3) \quad \triangle\psi = \frac{1}{\sqrt{\det g}} \partial_a \left( \sqrt{\det g} g^{ab} \partial_b \psi \right),$$

where  $g^{ab}$  is the inverse of the metric  $g_{ab}$  and  $\det g = \det\{g_{ab}\}$ . We assume that  $g^{ab}$  and  $f$  are smooth satisfying:

$$(6.4) \quad c_1 < f < 1/c_1, \quad c_1|\xi|^2 \leq g^{ab}\xi_a\xi_b \leq |\xi|^2/c_1$$

for some constant  $0 < c_1 < \infty$ .

In order for this to be solvable initial data must be compatible with the boundary condition. If we move the Laplacian in (6.1) over to the right hand side and differentiate (6.1) with respect to time we get

$$(6.5) \quad D_t^{k+2}\psi = G_k(\psi, \dots, D_t^{k+1}\psi, \partial\psi, \dots, \partial D_t^k\psi, \partial^2\psi, \dots, \partial^2 D_t^k\psi)$$

for some functions  $G_k$ . We therefore define functions of the space variables only  $\psi_k$

$$(6.6) \quad \psi_{k+2} = G_k(\psi_0, \dots, \psi_{k+1}, \partial\psi_0, \dots, \partial\psi_k, \partial^2\psi_0, \dots, \partial^2\psi_k)$$

where  $\psi_0$  and  $\psi_1$  are as in (6.2). For this to be compatible with the boundary conditions we must have

$$(6.7) \quad \psi_k|_{\partial\Omega} = 0, \quad \text{for } k \leq m-1 \quad \text{or} \quad \psi_k \in H_0^1(\Omega) \quad \text{for } k \leq m-1$$

(6.5) is called the  $m^{th}$  order compatibility condition. Since  $\psi_k$  are determined from the initial conditions  $\psi_0$  and  $\psi_1$  this gives some compatibility conditions on the initial conditions. The last alternative in (6.7) is the weak form, see [Ev] of the compatibility conditions if initial data  $(\psi_0, \psi_1) \in H^r \times H^{r-1}$ . For the first form in (6.7) we of course have to assume that  $\psi_k$  are continuous as well close to the boundary, so we have to assume some additional regularity.

The result in [H] is stated with vanishing initial conditions but, if the compatibility conditions are satisfied, one can reduce to that case by subtracting of the Taylor series:

$$(6.8) \quad \tilde{\psi}_m = \sum_{k=0}^{m-1} t^k \psi_k / k!,$$

and instead get vanishing initial conditions but an inhomogeneous term. If  $\bar{\psi} = \psi - \tilde{\psi}_k$  then

$$(6.9) \quad f \hat{D}_t^2 \bar{\psi} - \Delta \bar{\psi} = -f \hat{D}_t^2 \tilde{\psi}_k + \Delta \tilde{\psi}_k + \tilde{G} = \bar{G}$$

The new  $\bar{G}$  must then vanish to order  $m-3$  as  $t \rightarrow 0$ . In [H] there is an existence proof in Sobolev spaces of order  $m-3$  for this case. For simplicity, we will however state the result for infinitely differentiable initial data, satisfying (6.5) for all orders  $m = 0, 1, \dots$ . It then follows from [H] that we have a solution in  $C^k$  as well for any  $k$  by subtracting off the Taylor series (6.6) to sufficiently high order.

We will use an energy estimate, where  $\phi$  in the right hand side of (6.1) is treated in a special way so that we require one less space derivative but one more time derivative of  $\phi$ . This because the term  $\phi$  will be a coupling from the equation for the divergence free part and for that one we control more space than time regularity, see Theorem 5.1. This is needed in order to get the lowest order energy estimates for the coupled system.

**Theorem 6.1.** Suppose that  $g^{ab}$  and  $f$  are smooth and satisfy (6.4) and that  $\tilde{G}$  is smooth. Suppose also that the initial condition  $(\psi_0, \psi_1)$  in (6.2) are smooth and satisfy the compatibility conditions (6.5) for all  $m = 0, 1, \dots$ . Then (6.1)-(6.2) has a smooth solution  $\psi$ . Let

$$(6.10) \quad E_{1,r}(t) = \|\psi(t)\|_{0,r} + \|\psi(t)\|_{1,r-1}, \quad \text{where} \quad \|\psi(t)\|_{r,s} = \sum_{k=0}^s \|\hat{D}_t^k \psi(t)\|_r$$

Then for each  $r \geq 1$  there is a constant  $C$  such that for  $0 \leq t \leq T$  we have

$$(6.11) \quad E_{1,r}(t) \leq C \left( E_{1,r}(0) + \|\phi(0)\|_{1,r-1} + \int_0^t (\|\phi(\tau)\|_{1,r} + \|G(\tau)\|_{0,r-1}) d\tau \right)$$

*Proof.* The existence was just discussed above so we will only need to prove the estimate. The proof is going to be an application of Lemma 6.2 below. However, in the energy  $E$  in (6.12) we are missing an estimate of  $\|\psi\|_{L^2}$  and we only have estimates for  $\|\psi + \phi\|_{1,r-1}$  instead of  $\|\psi\|_{1,r-1}$ . The first problem is solved by replacing  $E$  by  $\tilde{E} = E + \|\psi\|_{L^2}^2$ . Since  $d(\|\psi(t)\|_{L^2}^2)/dt \leq \|D_t \psi(t)\|_{L^2}^2 \leq E$  (6.14) still holds for  $\tilde{E}$ . The second problem that in the energy  $\tilde{E}$  we only have estimates for  $\|\psi + \phi\|_{1,r-1}$  instead of  $\|\psi\|_{1,r-1}$ , is solved by adding the term:

$$(6.12) \quad \|\phi(t)\|_{1,r-1} \leq \|\phi(0)\|_{1,r-1} + \int_0^t \|\phi(\tau)\|_{1,r} d\tau \quad \square$$

**Lemma 6.2.** Suppose that  $\psi$  is a solution of (5.1). Let  $\nabla^a = g^{ab} \partial_b$ .

$$(6.13) \quad E(t) = \left( \sum_{s \leq r-1} \frac{1}{2} \int_{\Omega} f |\hat{D}_t^{s+1} \psi|^2 + |\hat{D}_t^s \nabla(\psi + \phi)|^2 \kappa dy \right)^{1/2},$$

where  $|\hat{D}_t^s \nabla \chi|^2 = g_{ab} (\hat{D}_t^s \nabla^a \chi) (\hat{D}_t^s \nabla^b \chi)$ . Then

$$(6.14) \quad \frac{dE}{dt} \leq C \left( E + \sum_{s \leq r-1} (\|\nabla \hat{D}_t^{s+1} \phi\|_{L^2} + \|\hat{D}_t^s G\|_{L^2}) \right)$$

*Proof.* We will prove that  $dE^2/dt$  is bounded by  $E$  times the right hand side of (6.13) and (6.13) follows from this since  $dE/dt = (dE^2/dt)/(2E)$ . With the notation  $\hat{D}_t = D_t + \text{div } V$  and  $\check{D}_t = D_t - \text{div } V$  we have  $\hat{D}_t(fg^2) = (\check{D}_t f)g^2 + 2fg\hat{D}_t g$ , and since also  $D_t \kappa / \kappa = \text{div } V$  it follows that

$$(6.15) \quad \begin{aligned} \frac{dE^2}{dt} = & \sum_{s \leq r-1} \int_{\Omega} \left( f (\hat{D}_t^{s+1} \psi) (\hat{D}_t^{s+2} \psi) + g_{ab} (\hat{D}_t^s \nabla^a (\psi + \phi)) (\hat{D}_t^{s+1} \nabla^b (\psi + \phi)) \right) \kappa dy \\ & + \frac{1}{2} \int_{\Omega} \left( (\check{D}_t f) (\hat{D}_t^{s+1} \psi)^2 + (\check{D}_t g_{ab}) (\hat{D}_t^s \nabla^a (\psi + \phi)) (\hat{D}_t^s \nabla^b (\psi + \phi)) \right) \kappa dy \end{aligned}$$

Here the terms on the second row are obviously bounded by a constant times  $E^2$  so it remains to look on the terms on the first row. The commutator  $[\hat{D}_t^{s+1}, \nabla^b] \chi$  is bounded by a constant times  $|\hat{D}_t^j \nabla \chi|$

for  $j \leq s$  so if we commute through the time derivative in the last term on the first row we get a commutator which is controlled by the  $E^2$ . After commuting we get if we integrate by parts

$$(6.16) \quad \frac{dE^2}{dt} = \sum_{s \leq r-1} \int_{\Omega} \left( f(\hat{D}_t^{s+1}\psi)(\hat{D}_t^{s+2}\psi) + (\hat{D}_t^s \nabla^a(\psi + \phi))(\partial_a(\hat{D}_t^{s+1}\psi) + \partial_a(\hat{D}_t^{s+1}\phi)) \right) \kappa dy + O(E^2) \\ = \sum_{s \leq r} \int_{\Omega} \left( (\hat{D}_t^{s+1}\psi)(f\hat{D}_t^{s+2}\psi - \kappa^{-1}\partial_a(\kappa\hat{D}_t^s \nabla^a(\phi + \psi)) + (\hat{D}_t^s \nabla^a(\psi + \phi))(\partial_a(\hat{D}_t^{s+1}\phi)) \right) \kappa dy + O(E^2)$$

Here the last term in the integrand on the second row is controlled by  $CE \sum_{s \leq r} \|\nabla \hat{D}_t^{s+1}\phi\|$ , so it only remains to look on the first term. Now  $\hat{D}_t g = \kappa^{-1} D_t(\kappa g)$  so differentiating  $f^2 \hat{D}_t^2 \psi - \Delta(\psi + \phi) = G$ :

$$(6.17) \quad \hat{D}_t^s \left( f\hat{D}_t^2 \psi - \kappa^{-1} \partial_a(\kappa \nabla^a(\psi + \phi)) \right) = f\hat{D}_t^{s+2} \psi - \kappa^{-1} \partial_a(\kappa \hat{D}_t^s \nabla^a(\psi + \phi)) + \sum_{k=0}^{s-1} c_k (D_t^{s-k} f)(\hat{D}_t^{k+2} \psi)$$

for some constants  $c_k$ . Since the  $L^2$  norm of the last term is bounded by  $CE$  the lemma follows.  $\square$

One can get additional space regularity from taking time derivatives of the equation (6.1) and solving the Dirichlet problem for the Laplacian.

**Lemma 6.3.** *Let  $E_{1,r}$  be as in Theorem 6.1 and let  $\|\psi\|_r = \sum_{s+k \leq r} \|D_t^s \psi\|_k$ . Then*

$$(6.18) \quad \|\psi\|_r \leq C(E_{1,r} + \|G\|_{r-2} + \|\phi\|_r), \quad r \geq 2, \quad \|\psi\|_1 \leq CE_{1,1}$$

*Proof.* Solving the Dirichlet problem:

$$(6.19) \quad \Delta \psi = f\hat{D}_t^2 \psi - \Delta \phi - G, \quad \psi|_{\partial\Omega} = 0$$

gives

$$(6.20) \quad \|\psi\|_{k+2} \leq C(\|\hat{D}_t^2 \psi\|_k + \|G\|_k + \|\phi\|_{k+2})$$

and commuting through time derivatives in (6.19) gives

$$(6.21) \quad \Delta \hat{D}_t^s \psi = \sum_{j=0}^s c_k D_t^{s-j} f \hat{D}_t^{2+j} \psi - \Delta \hat{D}_t^s \phi - D_t^s G - [\hat{D}_t^s, \Delta] \psi + [\hat{D}_t^s, \Delta] \phi \quad \hat{D}_t^s \psi|_{\partial\Omega} = 0$$

so solving the Dirichlet problem gives

$$(6.22) \quad \|\hat{D}_t^s \psi\|_{k+2} \leq C \left( \sum_{j=0}^s \|\hat{D}_t^{2+j} \psi\|_k + \|\hat{D}_t^s G\|_k + \sum_{j=0}^s \|\hat{D}_t^j \phi\|_{k+2} + \sum_{j=0}^{s-1} \|\hat{D}_t^j \psi\|_{k+2} \right)$$

Here the last term is lower order and is absent if  $s = 0$  so using induction in  $s$  we get

$$(6.23) \quad \sum_{j=0}^s \|\hat{D}_t^j \psi\|_{k+2} \leq C \sum_{j=0}^s (\|\hat{D}_t^{2+j} \psi\|_k + \|\hat{D}_t^j G\|_k + \|\hat{D}_t^j \phi\|_{k+2})$$

or with  $\|\psi\|_{r,s} = \sum_{k=0}^s \|D_t^k \psi\|_r$

$$(6.24) \quad \|\psi\|_{s+2, r-s-2} \leq C(\|\psi\|_{s, r-s} + \|G\|_{s, r-s-2} + \|\phi\|_{s+2, r-s-2}), \quad 0 \leq s \leq r-2$$

Since  $\|\psi\|_r = \sum_{s=0}^r \|\psi\|_{r-s, s}$  it therefore inductively follows that

$$(6.25) \quad \|\psi\|_r \leq C(\|\psi\|_{0, r} + \|\psi\|_{1, r-1} + \|G\|_{r-2} + \|\phi\|_r) \quad \square$$

We can similarly turn this around and get estimates for  $E_r$  from more space derivatives instead which can be used for the initial conditions.

**Lemma 6.4.** *Let  $\psi$  be the solution of (6.1) and let  $E_{2,r}(t) = \|\psi(t)\|_{r,0} + \|\psi(t)\|_{r-1,1}$ . Then*

$$(6.26) \quad \|\psi\|_r \leq C(E_{2,r} + \|G\|_{r-2} + \|\phi\|_r), \quad r \geq 2, \quad \|\psi\|_1 \leq CE_{2,1}$$

*Proof.* The proof is exactly the same as that of Lemma 6.3 apart from that the Laplacian and the second derivative with respect to time changed role. We obtain the result simply by differentiating

$$(6.27) \quad \hat{D}_t^2 \psi = f^{-1}(\Delta \psi + \Delta \phi + G)$$

with respect to time.

$$(6.28) \quad \|\psi\|_{r-k,k} \leq C\left(\|\psi\|_{r,0} + \|\psi\|_{r-1,1} + \sum_{j=2}^k \|G + \Delta \phi\|_{r-j,j-2}\right)$$

This follows from (6.5) using induction in  $k$ .  $\square$

## 7. A PRIORI BOUNDS FOR THE LINEARIZED SYSTEM.

Let us first consider the equation (4.13) for the divergence free part:

$$(7.1) \quad \ddot{W}_0 + AW_0 = PH(W_0, \dot{W}_0, W_1, \dot{W}_1, \partial W_1) + PF^a$$

Associated with (7.1) are energies

$$(7.2) \quad E_{0,r}(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r}$$

Let us now consider the wave equation for the divergence  $\delta h$

$$(7.3) \quad e' \hat{D}_t^2 \delta h - \Delta(\delta h - (\partial_c h)W^c) = G(W, \dot{W}, \partial W, \partial \dot{W}, \delta h, \hat{D}_t \delta h) - \operatorname{div} F,$$

where  $W = W_0 + W_1$  and  $W_1^a = g^{ab} \partial_b q$  is given by solving the Dirichlet problem

$$(7.4) \quad \Delta q = -e' \delta h, \quad q|_{\partial\Omega} = 0$$

so that  $\operatorname{div} W_1 = -e' \delta h$ . Associated with (7.3) there are energies

$$(7.5) \quad E_{1,r}(t) = \|\operatorname{div} W_1\|_{1,r-1} + \|\operatorname{div} W_1\|_{0,r},$$

where  $\|\chi\|_{r,s} = \sum_{k=0}^s \|\hat{D}_t^k \chi\|_r$ .

Through differentiating the equations (7.1) and (7.3) we can then control all components of the appropriate order through these energies. The energy (7.2) contains all space derivatives so we get additional time derivatives from differentiating (7.1) with respect to time, see Lemma 5.2. The energy (7.5) on the other hand contain all time derivatives so we get additional space derivatives through differentiating (7.3) and solving the Dirichlet problem for the Laplacian, which was done in Lemma 6.3. This leads to:

**Lemma 7.1.** *Let  $|||\chi|||_r = \sum_{s+k \leq r} \|\hat{D}_t^s \chi\|_k$ . Then*

$$(7.6) \quad |||\dot{W}_0|||_r + |||W_0|||_r + |||\operatorname{div} W_1|||_r \leq C(E_{0,r} + E_{1,r} + |||F|||_{r-1}), \quad r \geq 1$$

*Proof.* By Lemma 5.2 applied to (7.1), using (3.4) and (3.8):

$$(7.7) \quad |||\dot{W}_0|||_r + |||W_0|||_r \leq C(E_{0,r} + |||\dot{W}_0|||_{r-1} + |||W_0|||_{r-1} + |||\operatorname{div} W_1|||_{r-1} + |||F|||_{r-1})$$

Since also  $|||\dot{W}_0|||_0 + |||W_0|||_0 \leq E_{0,0}$  it follows by induction in  $r$  that

$$(7.8) \quad |||\dot{W}_0|||_r + |||W_0|||_r \leq C(E_{0,r} + |||\operatorname{div} W_1|||_{r-1} + |||F|||_{r-1})$$

We get from Lemma 6.3 applied to this (7.3), with  $\psi = -\operatorname{div} W/e'$ ,  $\phi = -(\partial_c h)W_0^c$  and  $f = e'$ :

$$(7.9) \quad |||\operatorname{div} W_1|||_r \leq C(E_{1,r} + |||\operatorname{div} W_1|||_{r-1} + |||W_0|||_r + |||F|||_{r-1}), \quad r \geq 2$$

Since also  $|||\operatorname{div} W_1|||_1 \leq E_{1,1}$  it follows from induction in  $r$ :

$$(7.10) \quad |||\operatorname{div} W_1|||_r \leq C(E_{1,r} + |||W_0|||_r + |||F|||_{r-1}),$$

Since also  $|||\operatorname{div} W_1|||_0 \leq E_{1,1}$  (7.6) now follows from (7.8) and (7.10)  $\square$

**Lemma 7.2.** *If  $r \geq 1$  then*

$$(7.11) \quad E_{0,r}(t) + E_{1,r}(t) \leq C\left(E_{0,r}(0) + E_{1,r}(0) + \int_0^t (E_{0,r}(\tau) + E_{1,r}(\tau) + |||F(\tau)|||_r) d\tau\right),$$

*Proof.* Using the estimate in Theorem 5.1 applied to (7.1) and the estimate for the projection (3.4) and for the normal operator (3.8) we get for  $0 \leq t \leq 1$ :

$$(7.12) \quad E_{0,r}(t) \leq CE_{0,r}(0) + C \int_0^t (\|\operatorname{div} W_1\|_{r,0} + \|\operatorname{div} W_1\|_{r-1,1} + \|W_0\|_{r,0} + \|W_0\|_{r,1} + \|F\|_r) d\tau$$

It follows from Theorem 6.1 applied to (7.3) with  $\phi = -(\partial_c h)W_0^c$  respectively  $\phi = 0$

$$(7.13) \quad E_{1,r}(t) \leq C\|W_0(t)\|_{1,r-1} + CE_{1,r}(0) + C \int_0^t (\|W_0\|_{1,r} + \|\operatorname{div} W_1\|_{0,r} + \|\operatorname{div} W_1\|_{1,r-1} + \|F\|_{1,r-1}) d\tau$$

$$(7.14) \quad E_{1,r}(t) \leq CE_{1,r}(0) + C \int_0^t (\|W_0\|_{2,r-1} + \|\operatorname{div} W_1\|_{0,r} + \|\operatorname{div} W_1\|_{1,r-1} + \|F\|_{1,r-1}) d\tau$$

Here we will use (7.14) for  $r \geq 2$  and (7.13) for  $r = 1$ . If  $r \geq 2$  then by Lemma 7.1 the integrands in (7.12) and (7.14) are bounded by  $C(E_{0,r} + E_{1,r})$  since then  $\|W\|_{2,r-1} \leq C|||\dot{W}_0|||_r$ , so (7.11) for  $r \geq 2$  follows. If  $r = 1$  then  $\|W_0(t)\|_{1,0} \leq E_{0,1}(t)$  so if  $r = 1$  we can replace  $\|W_0(t)\|_{1,r-1}$  in (7.13) by the right hand side of (7.12). Since by Lemma 7.1 the integrands in (7.13) and (7.12) are bounded by  $C(E_{0,r} + E_{1,r})$ , (7.11) follows also for  $r = 1$ .  $\square$



It follows from Lemma 7.2 that for  $0 \leq t \leq 1$

$$(7.15) \quad E_{0,r}(t) + E_{1,r}(t) \leq C(E_{0,r}(0) + E_{1,r}(0) + \int_0^t \|F(\tau)\|_r d\tau)$$

The proof of this is a standard argument. Let  $f(t)$  be the right hand side of (7.16). Then by (7.16)  $f'(t) \leq Cf(t) + C\|F(t)\|_r$  so it follows that  $(fe^{-Ct})' \leq C\|F(t)\|_r e^{-Ct}$ . Integrating this inequality gives  $f(t) \leq e^{Ct}(f(0) + \int_0^t \|F(\tau)\|_r d\tau)$ .

Now we want to have bounds in terms of a more pleasant energy not involving so many time derivatives. Let

$$(7.16) \quad E_{2,r}(t) = \|\operatorname{div} W_1\|_{r,0} + \|\operatorname{div} W_1\|_{r-1,1}$$

and let

$$(7.17) \quad E_r(t) = E_{0,r}(t) + E_{2,r}(t)$$

Then by Lemma 7.1  $E_r(t) \leq C(E_{0,r}(t) + E_{1,r}(t))$ . It remains to get an inequality in the other direction for the initial conditions. In order to do this we now have to turn it back and get estimates for high time derivatives from the equation:

**Lemma 7.3.**

$$(7.18) \quad \|\dot{W}_0\|_r + \|W_0\|_r + \|\operatorname{div} W_1\|_r \leq C(E_{0,r} + E_{2,r}), \quad r \geq 1$$

*Proof.* The proof is the same as that of Lemma 7.1 apart from that we replace  $E_{1,r}$  everywhere by  $E_{2,r}$ . The critical inequality (7.9) with  $E_{1,r}$  replaced by  $E_{2,r}$  comes from using Lemma 6.4 instead of Lemma 6.3.  $\square$

Finally we arrive at:

**Theorem 7.4.** *Let*

$$(7.19) \quad E_r(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r} + \|\operatorname{div} W_1(t)\|_r + \|\operatorname{div} \dot{W}_1(t)\|_{r-1}$$

where  $\dot{W} = \hat{D}_t W = D_t W + (\operatorname{div} V)W$ . Then there is a constant  $C$  such that for  $0 \leq t \leq T$

$$(7.20) \quad E_r(t) \leq C\left(E_r(0) + \int_0^t \|F\|_r d\tau\right)$$

Furthermore

$$(7.21) \quad E_1(t) \leq C\left(E_1(0) + \int_0^t \|F\|_1 d\tau\right)$$

If

$$(7.22) \quad \tilde{E}_r(t) = \|\dot{W}_0(t)\|_r + \|W_0(t)\|_r + \langle W_0(t) \rangle_{A,r} + \|\operatorname{div} W_1(t)\|_r$$

then

$$(7.23) \quad \tilde{E}_r(t) \leq C\left(E_r(0) + \int_0^t \|F\|_r d\tau\right)$$

## 8. INITIAL AND COMPATIBILITY CONDITIONS.

Assuming that the compatibility conditions are satisfied we can get all higher time derivatives by differentiating the equation with an inhomogeneous term:  $F$  that vanishes to any order as  $t \rightarrow 0$ :

$$(8.1) \quad \ddot{W}^a = g^{ab} \partial_b ((\partial_c h) W^c + \operatorname{div} W / e') + B^a(W, \dot{W}) + F^a, \quad W|_{t=0} = \tilde{W}^0, \quad \dot{W}|_{t=0} = \tilde{W}^1,$$

where  $\dot{W} = \hat{D}_t W = \kappa^{-1} D_t(\kappa W)$ . so

$$(8.2) \quad \hat{D}_t^{k+2} W = B_k(W, \dots, \hat{D}_t^{k+1} W, \partial W, \dots, \partial \hat{D}_t^k W, \partial^2 W, \dots, \partial^2 \hat{D}_t^k W) + \hat{D}_t^k F,$$

for some function  $B_k$ , where  $\hat{D}_t^k F|_{t=0} = 0$ . Let us therefore define functions of space only by

$$(8.3) \quad \tilde{W}^{k+2} = B_k(\tilde{W}^0, \dots, \tilde{W}^{k+1}, \partial \tilde{W}^0, \dots, \partial \tilde{W}^k, \partial^2 \tilde{W}^0, \dots, \partial^2 \tilde{W}^k)|_{t=0}, \quad k \geq 0$$

We now define the approximate power series solution by

$$(8.4) \quad \tilde{W}(t, y) = \frac{\kappa(0, y)}{\kappa(t, y)} \sum_{k=0}^{m-1} \tilde{W}^k(y) t^k / k!$$

defines a formal power series solutions. What we are doing is just expanding  $\kappa W$  in a formal power series in  $t$ , since  $D_t(\kappa W) = \kappa \hat{D}_t W$ . Now the compatibility conditions are that

$$(8.5) \quad \hat{D}_t^k \operatorname{div} W|_{\partial\Omega} = 0, \quad \text{when } t = 0 \quad k \leq m-1$$

Since  $\hat{D}_t \operatorname{div} W = \operatorname{div} \hat{D}_t W$  this it follows that this is equivalent to

$$(8.6) \quad \operatorname{div} \tilde{W}^k|_{\partial\Omega} = 0, \quad k \leq m-1,$$

and it then follows that

$$(8.7) \quad \operatorname{div} \tilde{W}|_{\partial\Omega} = 0$$

for all  $t$ .

Suppose that  $\overline{W}$  is a solution of (8.1) and let  $W = \overline{W} - \tilde{W}$ . Then  $\overline{W}$  satisfies the linearized equations with an inhomogeneous term:

$$(8.8) \quad \ddot{W}^a = g^{ab} \partial_b ((\partial_c h) W^c + \operatorname{div} W / e') + B^a(W, \dot{W}) + F^a,$$

$$(8.9) \quad W|_{t=0} = 0, \quad \dot{W}|_{t=0} = 0, \quad \operatorname{div} W|_{\partial\Omega} = 0$$

where

$$(8.10) \quad F^a = \ddot{W}^a - g^{ab} \partial_b ((\partial_c h) \tilde{W}^c + \operatorname{div} \tilde{W} / e') - B^a(\tilde{W}, \dot{\tilde{W}})$$

We are now going to find a solution to the equation with the inhomogeneous term (8.9) and adding  $\tilde{W}$  we then get a solution to (8.1). Here  $F$  vanishes to order  $m-2$  as  $t \rightarrow 0$ . We remark that the condition that (8.7) holds for all  $t$  is essential since in the next section we will find a solution to (8.8) with  $\operatorname{div} W = 0|_{\partial\Omega} = 0$  and because of (8.7) this will also be true for the sum.

## 9. EXISTENCE FOR THE LINEARIZED SYSTEM.

We now want to find a solution to the linearized equations (8.8)-(8.9) with the inhomogeneous term (8.10). This is equivalent to solving the system to solving the coupled system for  $(W_0, W_1)$ :

$$(9.1) \quad \ddot{W}_0 + AW_0 = PH(W_0, \dot{W}_0, W_1, \dot{W}_1, \partial W_1) + PF,$$

$$(9.2) \quad W_0|_{t=0} = 0, \quad \dot{W}_0|_{t=0} = 0$$

and

$$(9.3) \quad f\hat{D}_t^2\delta h - \Delta(\delta h - (\partial_c h)W^c) = G(W, \dot{W}, \partial W, \partial \dot{W}, \delta h, \hat{D}_t\delta h) - \operatorname{div} F,$$

$$(9.4) \quad \delta h|_{\partial\Omega} = 0, \quad \delta h|_{t=0} = 0, \quad \hat{D}_t\delta h|_{t=0} = 0$$

where  $W = W_0 + W_1$  and  $W_1^a = g^{ab}\partial_b q$  is given by solving the Dirichlet problem

$$(9.5) \quad \Delta q = -\delta h/e' \quad q|_{\partial\Omega} = 0$$

so that  $\operatorname{div} W_1 = -\delta h/e'$ .

If we have a solution  $(W_0, W_1)$  of the system (9.1)-(9.5) then  $W = W_0 + W_1$  is a solution of the equation (8.8)-(8.9). For  $W_1$  given we have existence for (9.1)-(9.2) by [L1] and for  $W_0$  given we have existence for (9.3)-(9.5) by [H]. In order to get a solution for the coupled system we put up an iteration so the (9.1)-(9.1) and (9.3)-(9.5) are only coupled through previous iterates. Hence existence follows for the iterates. The iteration is constructed as follows. Let  $(W_0^0, W_1^0) = (0, 0)$  for all times. We now define  $(W_0^{k+1}, W_1^{k+1})$  for  $k \geq 0$  to be the solution of the system

$$(9.6) \quad \ddot{W}_0^{k+1} + AW_0^{k+1} = PH(W_0^k, \dot{W}_0^k, W_1, \dot{W}_1^k, \partial W_1^k) + PF,$$

$$(9.7) \quad W_0^{k+1}|_{t=0} = 0, \quad \dot{W}_0^{k+1}|_{t=0} = 0$$

and

$$(9.8) \quad f\hat{D}_t^2\delta h^{k+1} - \Delta(\delta h^{k+1} - (\partial_c h)W^{kc}) = G(W^k, \dot{W}^k, \partial W^k, \partial \dot{W}^k, \delta h^k, \hat{D}_t\delta h^k) - \operatorname{div} F,$$

$$(9.9) \quad \delta h^{k+1}|_{\partial\Omega} = 0, \quad \delta h^{k+1}|_{t=0} = 0, \quad \hat{D}_t\delta h^{k+1}|_{t=0} = 0$$

where  $W^k = W_0^k + W_1^k$  and  $W_1^{ka} = g^{ab}\partial_b q^k$  is given by solving the Dirichlet problem

$$(9.10) \quad \Delta q^k = -\delta h^k/e' \quad q^k|_{\partial\Omega} = 0$$

so that  $\operatorname{div} W_1^k = -\delta h^k/e'$ .

It now remains to get uniform bounds for the iterates and convergence. To obtain convergence we consider the differences

$$(9.11) \quad \overline{W}_i^k = W_i^k - W_i^{k-1}$$

for  $k \geq 1$ . Since (9.6)-(9.10) hold for  $k \geq 0$  it follows that (9.6)-(9.10) with  $W_i^j$  replaced by  $\overline{W}_i^j$  hold for  $k \geq 1$  but without the inhomogeneous term  $F$ . For  $k = 0$ , then instead  $\overline{W}_i^1 = W_i^1$  so then (9.6)-(9.10) hold as well for the differences, but with the inhomogeneous term. Altogether, this can be phrased as that (9.6)-(9.10) hold for the differences for  $k \geq 0$  if we multiply the inhomogeneous term by  $\delta_{0k}$ . In the rest of this section we will prove:

**Theorem 9.1.** *We have*

$$(9.12) \quad \sum_{j=1}^{\infty} \|\dot{\bar{W}}_0^j\|_r + \|\bar{W}_0^j\|_r + \|\operatorname{div} \bar{W}_1^j\|_r + \langle \bar{W}_0^j J \rangle_{A,r} \leq C(\|F\|_{r-1} + \int_0^t \|F\|_r d\tau)$$

*Proof of Theorem 9.1.* The theorem follows from Lemma 9.4 and Lemma 9.3..  $\square$

*Proof of Theorem 4.1.* . If the initial conditions in (8.1) satisfy the  $m^{th}$  order compatibility condition (8.6) then the Taylor polynomial (8.4) is in  $(\tilde{W}, D_t \tilde{W}) \in C([0, T], H^{m+2} \times H^{m+1})$  if in addition the initial data are in  $(\tilde{W}^0, \tilde{W}^1) \in H^{2m+2} \times H^{2m+1}$ . The initial data therefore produces an inhomogeneous term (8.1) which is in  $C([0, T], H^m)$ . By the existence result in [L1] for (9.6)-(9.7) and in [H] for (9.8)-(9.10) we have a solution in the energy space (4.20) for  $r = m$ . By theorem (9.1) we in fact have convergence in this space as well. This means that the boundary condition will be satisfied in a weak sense if we look on weak solutions. And if  $r > n/2 + 2$  then we in fact have convergence in  $C^{r-[n/2]}$  to a classical solution. Since  $r$  was arbitrary, we in fact get a solution in  $C^k$ , for any  $k$ .  $\square$

Associated with (9.6) and (9.8)-(9.10) there are the energies:

$$(9.13) \quad E_{0,r}^k(t) = \|\dot{W}_0^k(t)\|_r + \|W_0^k(t)\|_r + \langle W_0^k(t) \rangle_{A,r}$$

$$(9.14) \quad E_{1,r}^k(t) = \|W_1^k(t)\|_{0,r} + \|W_1^k(t)\|_{1,r-1}$$

Similarly, associated with the differences there are energies:

$$(9.15) \quad \bar{E}_{0,r}^k = \|\dot{\bar{W}}_0\|_r + \|\bar{W}_0\|_r + \langle \bar{W}_0 \rangle_{A,r}$$

$$(9.16) \quad \bar{E}_{1,r}^k = \|\operatorname{div} \bar{W}_1\|_{0,r} + \|\operatorname{div} \bar{W}_1\|_{1,r-1}$$

We want to prove that the energies of the differences tend to zero, and in fact we will prove that their sum is uniformly bounded. We therefore define:

$$(9.17) \quad \bar{E}_{i,r}^{k*} = \sum_{j=1}^k \bar{E}_{i,r}^j, \quad \text{for } i = 0, 1$$

**Lemma 9.2.** *There is a constant  $C$  independent of  $k$  such that for  $0 \leq t \leq 1$  we have:*

$$(9.18) \quad \bar{E}_{0,1}^{k*} + \bar{E}_{1,1}^{k*} \leq C \int_0^t \|F\|_1 d\tau$$

*Proof.* Since the differences satisfies (9.6)-(9.10) but without the inhomogeneous term  $F$  for  $k \geq 1$ , we get as in the proof of Lemma 7.2 for  $k \geq 1$ :

$$(9.19) \quad \bar{E}_{0,r}^{k+1}(t) \leq C \int_0^t (\|\operatorname{div} \bar{W}_1^k\|_{r,0} + \|\operatorname{div} \bar{W}_1^k\|_{r-1,1} + \|\bar{W}_0^k\|_{r,0} + \|\bar{W}_0^k\|_{r,1}) d\tau,$$

$$(9.20) \quad \bar{E}_{1,r}^{k+1}(t) \leq C \|\bar{W}_0^k(t)\|_{1,r-1} + C \int_0^t (\|\bar{W}_0^k\|_{1,r} + \|\operatorname{div} \bar{W}_1^k\|_{0,r} + \|\operatorname{div} \bar{W}_1^k\|_{1,r-1}) d\tau$$

$$(9.21) \quad E_{1,r}^{k+1}(t) \leq C \int_0^t (\|\bar{W}_0^k\|_{2,r-1} + \|\operatorname{div} \bar{W}_1^k\|_{0,r} + \|\operatorname{div} \bar{W}_1^k\|_{1,r-1}) d\tau,$$

For  $k = 0$ ,  $(\overline{W}_0^1, \overline{W}_1^1) = (W_0^1, W_1^1)$  since  $(W_0^0, W_1^0) = (0, 0)$  and it follows that the difference  $(\overline{W}_0^1, \overline{W}_1^1)$  satisfies (9.6)-(9.10), now with the inhomogeneous term present but instead the other terms involving the previous iterate  $(W_0^0, W_1^0)$  vanishing. As in the proof of Lemma 7.2 we therefore get

$$(9.22) \quad \overline{E}_{i,r}^1 \leq C \int_0^t (\|F\|_{1,r-1} + \|F\|_{r,0}) d\tau, \quad \text{for } i = 0, 1.$$

since the previous iterates are zero.

Let us now assume that  $r = 1$ . By (9.22):

$$(9.23) \quad \overline{E}_{0,1}^1 + \overline{E}_{1,1}^1 \leq C \int_0^t \|F\|_1 d\tau$$

and for  $k \geq 1$  we have by (9.19) and (9.20):

$$(9.24) \quad \overline{E}_{0,1}^{k+1}(t) \leq C \int_0^t (\overline{E}_{0,1}^k + \overline{E}_{1,1}^k) d\tau$$

$$(9.25) \quad \overline{E}_{1,1}^{k+1}(t) \leq C \overline{E}_{0,1}^k(t) + C \int_0^t (\overline{E}_{0,1}^k + \overline{E}_{1,1}^k) d\tau$$

For  $k \geq 1$  we can replace the first term in the right of (9.25) by right hand side of (9.24) for  $k$  replaced by  $k - 1$  so it follows that

$$(9.26) \quad \overline{E}_{0,1}^{k+1}(t) + \overline{E}_{1,1}^{k+1}(t) \leq C \int_0^t (\overline{E}_{0,1}^k + \overline{E}_{0,1}^{k-1} + \overline{E}_{1,1}^k + \overline{E}_{1,1}^{k-1}) d\tau$$

for  $k \geq 1$ . Adding up (9.23) and (9.26) for  $k = 1, \dots$  we get

$$(9.27) \quad \overline{E}_{0,1}^{k+1*}(t) + \overline{E}_{1,1}^{k+1*}(t) \leq C \int_0^t (\overline{E}_{0,1}^{k*} + \overline{E}_{1,1}^{k*} + \|F\|_1) d\tau$$

and (9.18) follows from this. In fact if  $G$  denotes the right hand side then  $0 \leq G' \leq C(G + \|F\|_r)$  so  $(GE^{-Ct})' \leq Ce^{-Ct}\|F\|_r$  and since  $G(0) = 0$  we get  $G(t) \leq Ce^{Ct} \int_0^t \|F\|_r d\tau$ .  $\square$

**Lemma 9.3.** *For  $r \geq 2$  there is a constant  $C$  independent of  $k$  such that*

$$(9.28) \quad \sum_{j \leq k} \|\dot{\overline{W}}_0^j\|_r + \|\overline{W}_0^j\|_r + \|\operatorname{div} \overline{W}_1^j\|_r \leq C(\overline{E}_{0,r}^{k*} + \overline{E}_{1,r}^{k*} + \|F\|_{r-1})$$

*Proof.* As in the proof of Lemma 7.1 we see that for  $k \geq 2$  we have

$$(9.29) \quad \|\dot{\overline{W}}_0^k\|_r + \|\overline{W}_0^k\|_r \leq C(\overline{E}_{0,r}^k + \|\dot{\overline{W}}_0^{k-1}\|_{r-1} + \|\overline{W}_0^{k-1}\|_{r-1} + \|\operatorname{div} \overline{W}_1^{k-1}\|_{r-1})$$

$$(9.30) \quad \|\operatorname{div} \overline{W}_1^k\|_r \leq C(\overline{E}_{1,r}^k + \|\operatorname{div} \overline{W}_1^{k-1}\|_{r-1} + \|\overline{W}_0^{k-1}\|_r),$$

Now, since for  $k = 1$ ,  $(\overline{W}_0^1, \overline{W}_1^1) = (W_0^1, W_1^1)$  and since  $(W_0^0, W_1^0) = (0, 0)$  we get as in the proof of Lemma 7.1 that

$$(9.31) \quad \|\dot{\overline{W}}_0^1\|_r + \|\overline{W}_0^1\|_r \leq C(\overline{E}_{0,r}^1 + \|F\|_{r-1})$$

$$(9.32) \quad \|\operatorname{div} \overline{W}_1^1\|_r \leq C(\overline{E}_{1,r}^1 + \|F\|_{r-1}),$$

since the previous iterates are zero but instead we have the inhomogeneous term. Let

$$(9.33) \quad M_r^{k*} = \sum_{j \leq k} (\|\dot{\overline{W}}_0^j\|_r + \|\overline{W}_0^j\|_r), \quad N_r^{k*} = \sum_{j \leq k} \|\operatorname{div} \overline{W}_1^j\|_r$$

Then summing over all  $k \geq 2$  in (9.29)-(9.30) and adding  $k = 1$  from (9.31)-(9.32) gives:

$$(9.34) \quad M_r^{k*} \leq C(\overline{E}_{0,r}^{k*} + M_{r-1}^{k*} + N_{r-1}^{k*}) + M_r^1$$

$$(9.35) \quad N_r^{k*} \leq C(\overline{E}_{0,r}^{k*} + M_r^{k*} + N_{r-1}^{k*}) + N_r^1$$

Using induction in  $r$  it follows that

$$(9.36) \quad M_r^{k*} + N_r^{k*} \leq C(\overline{E}_{0,r}^{k*} + \overline{E}_{0,r}^{k*} + M_r^1 + N_r^1)$$

so the lemma follows.  $\square$

**Lemma 9.4.** *There is a constant  $C$  independent of  $k$  such that*

$$(9.37) \quad \overline{E}_{0,r}^{k*} + \overline{E}_{1,r}^{k*} \leq C \int_0^t \|\|F\|\|_r d\tau$$

*Proof.* If we sum up (9.19) and (9.21) for all  $k \geq 1$  and (9.22) we get if  $r \geq 2$ :

$$(9.38) \quad \overline{E}_{0,r}^{k+1*} + \overline{E}_{1,r}^{k+1*} \leq C \int_0^t \sum_{j \leq k} (\|\dot{\overline{W}}_0^j\|_r + \|\overline{W}_0^j\|_r + \|\operatorname{div} \overline{W}_1^j\|_r) d\tau + C \int_0^t \|\|F\|\|_r d\tau$$

Using Lemma 9.3 we get

$$(9.39) \quad \overline{E}_{0,r}^{k+1*} + \overline{E}_{0,r}^{k+1*} \leq C \int_0^t (\overline{E}_{0,r}^{k*} + \overline{E}_{1,r}^{k*} + \|\|F\|\|_r) d\tau$$

and the lemma follows from this, see the end of the proof of Lemma 9.2.  $\square$

**Acknowledgments.** I would like to thank Demetrios Christodoulou and Kate Okikiolu for enlightening discussions.

## REFERENCES

- [BG]. M.S. Baouendi and C. Gouaouic, *Remarks on the abstract form of nonlinear Cauchy-Kovalevsky theorems*, Comm. Part. Diff. Eq. **2** (1977), 1151-1162.
- [C1]. D. Christodoulou, *Self-Gravitating Relativistic Fluids: A Two-Phase Model*, Arch. Rational Mech. Anal. **130** (1995), 343-400.
- [C2]. D. Christodoulou, *Oral Communication* (August 95).
- [CK]. D. Christodoulou and S. Klainerman, *The Nonlinear Stability of the Minkowski space-time*, Princeton Univ. Press, 1993.
- [CL]. D. Christodoulou and H. Lindblad, *On the motion of the free surface of a liquid.*, Comm. Pure Appl. Math. **53** (2000), 1536-1602.

- [DM]. B. Dacorogna and J. Moser, *On a partial differential equation involving the Jacobian determinant.*, Ann. Inst. H. Poincaré Anal. Non. Linéaire **7** (1990), 1-26.
- [E1]. D. Ebin, *The equations of motion of a perfect fluid with free boundary are not well posed.*, Comm. Part. Diff. Eq. **10** (1987), 1175–1201.
- [E2]. D. Ebin, *Oral communication* (November 1997).
- [Ev]. C. Evans, *Partial Differential Equations*, AMS.
- [H]. Hörmander, *The analysis of Linear Partial Differential Operators III*, Springer Verlag.
- [L1]. H. Lindblad, *Well posedness for the linearized motion of the free surface of a liquid.*, preprint (January 2001), URL: <http://arXiv.org/abs/math/0112030>.
- [L2]. ———, *Well posedness for the linearized motion of the free surface of a compressible fluid.*, preliminary preprint (March 2001).
- [L3]. ———, *Well posedness for the motion of the free surface of a liquid.*, In preparation.
- [Na]. V.I. Nalimov,, *The Cauchy-Poisson Problem (in Russian)*,, Dynamika Splosh. Sredy **18** (1974,), 104-210.
- [Ni]. T. Nishida, *A note on a theorem of Nirenberg*, J. Diff. Geometry **12** (1977), 629-633.
- [W1]. S. Wu, *Well-posedness in Sobolev spaces of the full water wave problem in 2-D*, Invent. Math. **130** (1997), 39-72.
- [W2]. S. Wu, *Well-posedness in Sobolev spaces of the full water wave problem in 3-D*, J. Amer. Math. Soc. **12** (1999), 445-495.
- [Y]. H. Yosihara, *Gravity Waves on the Free Surface of an Incompressible Perfect Fluid* **18** (1982), Publ. RIMS Kyoto Univ., 49-96.

UNIVERSITY OF CALIFORNIA AT SAN DIEGO

*E-mail address:* `lindblad@math.ucsd.edu`