

Endomorphisms of Stable Continuous-Trace C^* -algebras

Ilan Hirshberg

Abstract

We classify spectrum-preserving endomorphisms of stable continuous-trace C^* -algebras up to inner automorphism by a surjective multiplicative invariant taking values in finite dimensional vector bundles over the spectrum. Specializing to automorphisms, this gives a different approach to results of Phillips and Raeburn. We also give a decomposition result for non spectrum-preserving endomorphisms.

2000 Mathematics Subject Classification: Primary: 46L05; Secondary: 46M20

1 Introduction

The structure of automorphisms of continuous-trace C^* -algebras was studied by Lance ([L]) and Smith ([S]) (the case of spectrum preserving automorphisms of $C(X) \otimes \mathcal{K}$, or more precisely $C(X) \otimes \mathcal{B}(\mathcal{H})$), and subsequently by Phillips and Raeburn ([PhR], see also [Ro],[RW]) for general continuous trace C^* -algebras.

This paper is concerned with the question of generalizing some of those results to the situation of ‘unital’ endomorphisms of stable continuous-trace C^* -algebras. We refer the reader to [RW] and [Di] for general references on the theory of continuous-trace C^* -algebras.

Throughout the paper, X will denote a locally compact Hausdorff space; \mathcal{H} will denote a separable Hilbert space and $\mathcal{B}(\mathcal{H})$ will denote the bounded operators on \mathcal{H} ; $\mathcal{M}(\mathcal{A})$ will denote the multiplier algebra of \mathcal{A} ; \mathcal{K} will denote the algebra of compact operators on a separable Hilbert space; automorphisms, endomorphisms and homomorphisms between C^* -algebras and C^* -bundles will always be $*$ -homomorphisms; $\delta(\mathcal{A})$ will denote the Dixmier-Douady invariant of a continuous-trace C^* -algebra. By a unital homomorphism between two non-unital C^* -algebras we mean a homomorphism which maps approximate units to approximate units.

We recall that every stable continuous-trace C^* -algebra for which all the irreducible representations act on separable space is isomorphic to the algebra of sections a locally trivial \mathcal{K} -bundle over a locally compact Hausdorff space. Let E be a locally trivial \mathcal{K} -bundle over X , and let $\mathcal{A} = \Gamma_0(E)$. Denote by $\text{Aut}_X(\mathcal{A})$ the automorphisms of \mathcal{A} which commute with the multiplication action

of $C_0(X)$ on \mathcal{A} , and by $\text{Inn}(\mathcal{A})$ the group of inner automorphisms of \mathcal{A} , i.e. those automorphisms of \mathcal{A} which are given by $\alpha(A) = UAU^*$ for some $U \in \mathcal{M}(\mathcal{A})$. Denote by $\text{Homeo}_\delta(X)$ the group of homeomorphisms of X which preserve $\delta(A) \in H^3(X; \mathbb{Z})$, where cohomology here is taken to mean Čech cohomology. The results of Phillips and Raeburn, specialized to this case, can be summarized in the following two exact sequences.

$$1 \rightarrow \text{Inn}(\mathcal{A}) \rightarrow \text{Aut}_X(\mathcal{A}) \rightarrow H^2(X; \mathbb{Z}) \rightarrow 1 \quad (1)$$

$$1 \rightarrow \text{Aut}_X(\mathcal{A}) \rightarrow \text{Aut}(\mathcal{A}) \rightarrow \text{Homeo}_\delta(X) \rightarrow 1 \quad (2)$$

Of course, some of this structure must be lost when generalizing to endomorphisms. The fact that the endomorphisms do not form a group means that speaking of exact sequences won't make much sense.

The bulk of the paper will be devoted providing an adequate generalization to sequence (1). Instead of looking for sheaf-cohomological invariant, our invariant takes values in vector bundles over X . We will show that the associated vector bundle of a spectrum-preserving endomorphism α , denoted $\text{vect}(\alpha)$ classifies α up to inner automorphism (i.e. $\text{vect}(\alpha) \cong \text{vect}(\beta) \iff \alpha = \text{Ad}U \circ \beta$); that all locally trivial vector bundles over X arise in this way; and that we have a product formula $\text{vect}(\alpha \circ \beta) \cong \text{vect}(\alpha) \otimes \text{vect}(\beta)$ (in other words, vect is a semigroup homomorphism from the semigroup of endomorphisms under composition to vector bundles under tensor products).

The invariant for spectrum-preserving automorphisms in this approach will be a complex line bundle over X . So if we specialize the above results to automorphisms, then we get a similar sequence to (1) with the group of complex line bundles over X instead of $H^2(X, \mathbb{Z})$. But those two groups are isomorphic, so all is well.

The structure of this paper is as follows. In section 2, we study the space of endomorphisms of \mathcal{K} . In section 3, we discuss the case of spectrum preserving endomorphisms of $C_0(X) \otimes \mathcal{K}$. In section 4, we generalize the main results of section 3 to algebras with possibly non-trivial Dixmier-Douady invariant. In section 5 we obtain a decomposition result which gives some form of generalization for sequence (2).

2 Endomorphisms of \mathcal{K}

Let α be an endomorphism of \mathcal{K} which is unital, in the sense that it maps approximate units to approximate units.

Let n be the trace of $\alpha(P)$ for a minimal projection $P \in \mathcal{K}$, where by trace we mean the unique (unbounded) trace on \mathcal{K}_+ whose value on minimal projections is 1. This n clearly is independent of the choice of P , and will be called $\text{index}(\alpha)$.

The following simple lemma and its short proof was conveyed to the author by W. B. Arveson.

Lemma 2.1. *There are isometries $S_1, \dots, S_n \in \mathcal{M}(\mathcal{K}) \cong \mathcal{B}(\mathcal{H})$ with*

$$\sum_{i=1}^n S_i S_i^* = 1$$

such that for all $A \in \mathcal{K}$,

$$\alpha(A) = \sum_{i=1}^n S_i A S_i^*$$

Proof: Letting \mathcal{K} act irreducibly on some space \mathcal{H} , we observe that α gives a representation of \mathcal{K} on \mathcal{H} . This representation decomposes into a direct sum of n irreducible representations on pairwise orthogonal subspaces $\mathcal{H}_1 \oplus \dots \oplus \mathcal{H}_n = \mathcal{H}$, and there are isometries S_i from \mathcal{H} onto \mathcal{H}_i , $i = 1, \dots, n$ which implement $\alpha|_{\mathcal{H}_i}$, i.e. $S_i A S_i^* \xi_i = \alpha(A) \xi_i$ for $\xi_i \in \mathcal{H}_i$. The lemma follows from this immediately. \square

Notice that we have $S_i A = \alpha(A) S_i$, $i = 1, \dots, n$. Let

$$V_\alpha = \{T \in \mathcal{B}(\mathcal{H}) \mid TA = \alpha(A)T \ \forall A \in \mathcal{K}\}$$

V_α is clearly a vector space. It also has a natural inner product on it. If $T, S \in V_\alpha$ then $T^* S A = A T^* S$ for all $A \in \mathcal{K}$, and therefore $T^* S$ is a scalar multiple of the identity, and indeed $\langle S, T \rangle = T^* S$ is an inner product.

Lemma 2.2. *The S_i 's are an orthonormal basis for V_α .*

Proof: Let $T \in V_\alpha$ then $T = (\sum_{i=1}^n S_i S_i^*) T = \sum_{i=1}^n \langle S_i, T \rangle S_i$, as required. Orthonormality is immediate. \square

If T_1, \dots, T_m are isometries such that $\alpha(A) = \sum_{i=1}^m T_i A T_i^*$ for all $A \in \mathcal{K}$ then clearly $m = \text{index}(\alpha) = n$, and the T_i 's form another orthonormal basis for V_α . Conversely, if T_1, \dots, T_n are an orthonormal basis for V_α then it is straightforward to verify that $\alpha(A) = \sum_{i=1}^n T_i A T_i^*$ for all $A \in \mathcal{K}$.

Denote by $\text{End}(\mathcal{K})$ the space of unital endomorphisms of \mathcal{K} , endowed with the point-norm topology, and by $\text{End}^n(\mathcal{K})$ the subset of endomorphisms of index n . Notice that $\text{End}(\mathcal{K})$ (and hence all the $\text{End}^n(\mathcal{K})$'s) is metrizable, hence paracompact (if A_1, A_2, \dots is a dense sequence in the unit sphere of \mathcal{K} , then $d(\alpha, \beta) = \sum_k \|\alpha(A_k) - \beta(A_k)\|/2^k$ is a metric inducing the point-norm topology).

Proposition 2.3. *$\text{index} : \text{End}(\mathcal{K}) \rightarrow \mathbb{N}$ is continuous (so $\text{End}^n(\mathcal{K})$ is closed and open in $\text{End}(\mathcal{K})$ for all n).*

Proof: Let $\alpha_k \rightarrow \alpha$. Let P be a minimal projection, then for sufficiently large k , we have $\|\alpha_k(P) - \alpha(P)\| < 1$. Therefore, $\alpha_k(P)$ and $\alpha(P)$ are Murray-von-Neumann equivalent, and in particular they have the same trace, i.e. $\text{index}(\alpha_k) \rightarrow \text{index}(\alpha)$. (cf [Pr] Proposition 2.3) \square

We denote by $\mathcal{C}_n \subseteq \mathcal{B}(\mathcal{H})^n$ the space of all n -tuples of isometries satisfying the Cuntz relations, equipped with the strong operator topology. We remark that \mathcal{C}_n is metrizable (since the unit sphere of $\mathcal{B}(\mathcal{H})$ is metrizable in the SOT), hence paracompact. We described above a surjective map $\pi : \mathcal{C}_n \rightarrow \text{End}^n(\mathcal{K})$, given by $\pi(S_1, \dots, S_n)(A) = \sum_{i=1}^n S_i x S_i^*$. Suppose $(S_1^{(k)}, \dots, S_n^{(k)}) \rightarrow (S_1, \dots, S_n)$ in \mathcal{C}_n , and let $A = \xi \otimes \bar{\eta} \in \mathcal{K}$ a rank 1 operator, then

$$\begin{aligned} \pi(S_1^{(k)}, \dots, S_n^{(k)})(A) &= \sum_{i=1}^n S_i^{(k)} A S_i^{(k)*} = \\ &= \sum_{i=1}^n S_i^{(k)} \xi \otimes \overline{S_i^{(k)*} \eta} \rightarrow \sum_{i=1}^n S_i^* \xi \otimes \overline{S_i^* \eta} = \pi(S_1, \dots, S_n)(A) \end{aligned}$$

so π is continuous.

We define the *universal intertwining bundle*

$$\begin{array}{c} \mathbb{C}^n \\ \downarrow \\ \mathcal{I}_n \\ \downarrow \\ \text{End}^n(\mathcal{K}) \end{array}$$

to be the sub-bundle of $\text{End}^n(\mathcal{K}) \times \mathcal{B}(\mathcal{H})$ whose fiber over α is V_α (where the topology on $\mathcal{B}(\mathcal{H})$ is the strong operator topology).

$$\begin{array}{c} \mathcal{U}_n \\ \downarrow \\ \mathcal{C}_n \\ \pi \downarrow \\ \text{End}^n(\mathcal{K}) \end{array}$$

is the corresponding principal \mathcal{U}_n -bundle (the bundle of orthonormal frames).

Lemma 2.4. $\pi : \mathcal{C}_n \rightarrow \text{End}^n(\mathcal{K})$ has local cross sections. In other words, the bundles above are locally trivial.

Proof: (cf [RW], Proposition 1.6) We think of \mathcal{K} as acting irreducibly on \mathcal{H} . Fix $\alpha \in \text{End}^n(\mathcal{K})$, and let $P = \nu \otimes \bar{\nu} \in \mathcal{K}$ be a fixed minimal projection. Let $\omega_1, \dots, \omega_n$ be an orthonormal basis for $\alpha(P)\mathcal{H}$, and define $S_i \in \mathcal{B}(\mathcal{H})$, $i = 1, \dots, n$ by $S_i \xi = \alpha(\xi \otimes \bar{\nu})\omega_i$.

For any $i, j \in 1, \dots, n$, and $\xi, \eta \in \mathcal{H}$, we have

$$\langle S_i \xi, S_j \eta \rangle = \langle \alpha(\xi \otimes \bar{\nu})\omega_i, \alpha(\eta \otimes \bar{\nu})\omega_j \rangle = \langle \alpha(\eta \otimes \bar{\nu})^* \alpha(\xi \otimes \bar{\nu})\omega_i, \omega_j \rangle =$$

$$= \langle \alpha((\eta \otimes \bar{\nu})^* \xi \otimes \bar{\nu}) \omega_i, \omega_j \rangle = \langle \xi, \eta \rangle \langle \alpha(\nu \otimes \bar{\nu}) \omega_i, \omega_j \rangle = \langle \xi, \eta \rangle \langle \omega_i, \omega_j \rangle = \langle \xi, \eta \rangle \delta_{ij}$$

so S_1, \dots, S_n are isometries with orthogonal ranges.

Let $A = \xi \otimes \bar{\eta}$, and let $\zeta \in \mathcal{H}$, then

$$\begin{aligned} S_i A \zeta &= (S_i \xi \otimes \bar{\eta}) \zeta = \langle \zeta, \eta \rangle S_i \xi = \langle \zeta, \eta \rangle \alpha(\xi \otimes \bar{\nu}) \omega_i = \\ &= \alpha((\xi \otimes \bar{\eta})(\zeta \otimes \bar{\nu})) \omega_i = \alpha(A) \alpha(\zeta \otimes \bar{\nu}) \omega_i = \alpha(A) S_i \zeta \end{aligned}$$

so S_i is an intertwiner.

Therefore, $\pi(S_1, \dots, S_n) = \alpha$. Now, define $f_1, \dots, f_n : \text{End}^n(\mathcal{K}) \rightarrow \mathcal{H}$ by

$$f_i(\beta) = \beta(P) \omega_i$$

The f_i are all clearly continuous, and $f_i(\alpha) = \omega_i$. Therefore, there is an open neighborhood N of α in which f_1, \dots, f_n are linearly independent. Let $g_1, \dots, g_n : N \rightarrow \mathcal{H}$ be obtained from f_1, \dots, f_n via the Gram-Schmidt orthonormalization process, then the g_i 's are continuous as well. Now, let $s_1, \dots, s_n : N \rightarrow \mathcal{B}(\mathcal{H})$ be given by

$$s_i(\beta) \xi = \beta(\xi \otimes \bar{\nu}) g_i(\beta), \quad \xi \in \mathcal{H}$$

then the s_i 's are clearly continuous (when $\mathcal{B}(\mathcal{H})$ is given the strong operator topology), and from the above, we have $(s_1(\beta), \dots, s_n(\beta)) \in \mathcal{C}_n$ for all β , and $\pi(s_1(\beta), \dots, s_n(\beta)) = \beta$, so we have a local cross section, as required. \square

Theorem 2.5. \mathcal{C}_n is contractible.

Proof: Let $v_t : L^2(\mathbb{R}_+) \rightarrow L^2(\mathbb{R}_+)$ be the unilateral shifts, $p_t = 1 - v_t v_t^*$, (T'_1, \dots, T'_n) some fixed n -tuple of isometries satisfying the Cuntz relations on a separable Hilbert space \mathcal{H}' . Identify $\mathcal{H} = \mathcal{H}' \otimes L^2(\mathbb{R}_+)$, and let $V_t = 1 \otimes v_t$, $P_t = 1 \otimes p_t$, $T_i = T'_i \otimes 1$, $i = 1, \dots, n$. Define a family of maps $\Phi_t : \mathcal{C}_n \rightarrow \mathcal{C}_n$ by

$$\Phi_t(S_1, \dots, S_n) = (\dots, V_t S_i V_t^* + P_t T_i P_t, \dots)$$

then it is easy to see Φ_t is SOT-continuous in t , that $\Phi_t(S_1, \dots, S_n) \in \mathcal{C}_n$ for all $(S_1, \dots, S_n) \in \mathcal{C}_n$, and that $\text{SOT} - \lim \Phi_t(S_1, \dots, S_n) = (T_1, \dots, T_n)$ for any $(S_1, \dots, S_n) \in \mathcal{C}_n$, so Φ_t gives us a contraction as required. \square

Recall that an open cover $\{U_\lambda\}$ of a topological space Y is said to be *numerable* if there is a locally finite partition of unity $\{f_\mu\}$ such that the open cover $\{f_\mu^{-1}((0, 1])\}$ refines $\{U_\lambda\}$. Notice that if Y is paracompact then any open cover is numerable. A principal \mathcal{U}_n -bundle ζ over Y is said to be *numerable* if there is a numerable open cover $\{U_\lambda\}$ of Y such that $\zeta|_{U_\lambda}$ is trivial for all λ . If Y is paracompact then any locally trivial bundle is numerable. The pull back of a numerable bundle is also numerable. A numerable principal \mathcal{U}_n -bundle over a space B is said to be *universal* if

1. Any numerable principal \mathcal{U}_n -bundle over some space Y is isomorphic to a pull-back bundle of this bundle via a map $Y \rightarrow B$.

2. Two maps $Y \rightarrow B$ give rise to isomorphic pull-back bundles if and only if they are homotopic.

A universal Hermitian vector bundle is defined in the analogous way, and a Hermitian vector bundle is universal if and only if its associated principal \mathcal{U}_n -bundle is universal.

By a theorem of Dold ([Do] Theorem 7.5), a numerable principal \mathcal{U}_n -bundle is universal if and only if the total space is contractible.

Corollary 2.6. *The above bundles are universal.*

Remark 2.7. For any fixed $(S_1, \dots, S_n) \in \mathcal{C}_n$ we can define a map $\mathcal{U}(\mathcal{H}) \rightarrow \mathcal{C}_n$ by $U \mapsto (US_1, \dots, US_n)$. This map can readily be seen to be a homeomorphism, so in fact all the \mathcal{C}_n 's are homeomorphic.

Remark 2.8. The focus of this paper is ordinary (=complex) C^* -algebras, however it is worth noticing that the analogous results of this section along with their proofs hold for endomorphisms of the real C^* -algebra of compact operators on a real separable Hilbert space as well (replacing complex bundles by real bundles, unitaries by orthogonals and so on).

3 Spectrum Preserving Endomorphisms of $C_0(X) \otimes \mathcal{K}$

We now turn to study endomorphisms of $C_0(X) \otimes \mathcal{K}$ which commute with the multiplier action of $C_0(X)$. We note that any endomorphism α of $C_0(X) \otimes \mathcal{K}$ induces a map $\alpha^* : X \rightarrow X$, so in this section we are restricting our attention precisely to those endomorphisms α such that $\alpha^* = id$. We refer to those as *spectrum preserving endomorphisms*.

Lemma 3.1. *Let $\sigma : X \rightarrow End(\mathcal{K})$ be a continuous map, then there is a spectrum preserving endomorphism α of $C_0(X) \otimes \mathcal{K}$ given by $\alpha(f)(x) = \sigma(x)(f(x))$. Furthermore, any spectrum preserving endomorphism α of $C_0(X) \otimes \mathcal{K}$ is of this form.*

Proof: The proof is straightforward (and virtually identical to Lemma 4.28 in [RW], using endomorphisms instead of automorphisms), and the details are left to the reader.

We henceforth denote this map by $x \mapsto \alpha_x$.

Definition 3.2.

$$vect(\alpha) = \{(x, T) \mid TA = \alpha_x(A)T \ \forall A \in \mathcal{K}\} \subseteq X \times \mathcal{B}(\mathcal{H})$$

Notice that $vect(\alpha)$ is the pull-back of the universal intertwining bundle(s) via the map $x \rightarrow \alpha_x$ (where possibly the dimension may be different on different components of X ; by Proposition 2.3, the dimension will be locally constant). In particular, $vect(\alpha)$ is numerable.

Corollary 3.3. *For any numerable Hermitian vector bundle v there exists a spectrum preserving endomorphism α such that $\text{vect}(\alpha) \cong v$. Furthermore, $\text{vect}(\alpha) \cong \text{vect}(\beta)$ if and only if α is connected to β via a path of spectrum preserving endomorphisms.*

Theorem 3.4. *Let α, β be spectrum preserving endomorphisms, then*

$$\text{vect}(\alpha \circ \beta) \cong \text{vect}(\alpha) \otimes \text{vect}(\beta)$$

Proof: Define a map

$$\text{vect}(\alpha) \otimes \text{vect}(\beta) \rightarrow \text{vect}(\alpha \circ \beta)$$

by

$$(x, T \otimes S) \mapsto (x, TS)$$

Verifying that this map is a well defined bundle isomorphism is immediate. \square

The associated vector bundle to a spectrum preserving endomorphism classifies the spectrum preserving endomorphism up to homotopy. A more natural equivalence relation on spectrum preserving endomorphisms is classification up to inner automorphism, i.e., $\alpha \sim \beta$ if there is a unitary U in $\mathcal{M}(C_0(X) \otimes \mathcal{K})$ such that $\text{Ad } U \circ \alpha = \beta$. The following proposition shows that the two equivalence relations coincide.

Observation 3.5. *$\text{Hom}(\text{vect}(\alpha), \text{vect}(\beta))$ is isomorphic to the sub-bundle of the $X \times \mathcal{B}(\mathcal{H})$ whose fiber over x is $\text{span}\{TS^* \mid T \in \text{vect}(\beta), S \in \text{vect}(\alpha)\}$.*

Theorem 3.6. *$\text{vect}(\alpha) \cong \text{vect}(\beta)$ if and only if $\alpha = \text{Ad } U \circ \beta$ for some unitary U .*

We give two proofs of this theorem. The first proves the theorem directly, using the preceding observation. The second proof uses a modified argument from [Po] (Theorem 2.4) to show that equivalence up to inner automorphism coincides with equivalence up to homotopy.

Proof 1: If $\alpha = \gamma \circ \beta$ where γ is an inner automorphism, then by theorem 3.4 we have $\text{vect}(\alpha) \cong \text{vect}(\gamma) \otimes \text{vect}(\beta)$. But $\text{vect}(\gamma)$ is the trivial line bundle, so $\text{vect}(\gamma) \otimes \text{vect}(\beta) \cong \text{vect}(\beta)$.

For the converse, $\text{vect}(\alpha) \cong \text{vect}(\beta)$ means that there is an isomorphism between them, i.e. a unitary element in $\text{Hom}(\text{vect}(\alpha), \text{vect}(\beta))$, which by observation 3.5 can be viewed as a sub-bundle of $X \times \mathcal{B}(\mathcal{H})$. So we know that there is a unitary section U of this sub-bundle (and in particular, $U \in \mathcal{M}(C_0(X) \otimes \mathcal{K})$), and any such section can immediately be seen to satisfy $U\alpha(A)U^* = \beta(A)$ for all $A \in C_0(X) \otimes \mathcal{K}$, i.e. $\alpha = \text{Ad } U \circ \beta$, as required. \square

Proof 2: Suppose U implements an equivalence between α and β . U can be thought of as a function $u : X \rightarrow \mathcal{U}(\mathcal{H})$. We know that $\mathcal{U}(\mathcal{H})$ is contractible (by theorem 2.5 for example, observing that $\mathcal{U}(\mathcal{H}) \cong \mathcal{C}_1$). Therefore, there

is a path of functions $u_t : X \rightarrow \mathcal{U}(\mathcal{H})$, continuous when viewed as functions $X \times [0, 1] \rightarrow \mathcal{U}(\mathcal{H})$, such that $u_0 = u$, $u_1 \equiv 1$. Let U_t be the corresponding path of unitaries in $\mathcal{M}(C_0(X) \otimes \mathcal{K})$, then $\alpha_t = \text{Ad}U_t \circ \alpha$ implements a homotopy between α and β .

Conversely, suppose α and β are homotopic. Let e_{ij} ($i, j \in \mathbb{N}$) be matrix units for \mathcal{K} , and let $E_{ij} = 1 \otimes e_{ij} \in C_b(X) \otimes \mathcal{K} \subseteq \mathcal{M}(C_0(X) \otimes \mathcal{K})$. So the projections $\alpha(E_{11})$, $\beta(E_{11})$ are homotopic (where by that we mean the canonical extensions of α , β applied to E_{11}). Since homotopic projections are unitarily equivalent (see, for example, [B] chapter 4), we can find a unitary V such that $V\alpha(E_{11})V^* = \beta(E_{11})$, and therefore we can find a partial isometry $W \in \mathcal{M}(C_0(X) \otimes \mathcal{K})$ with $W^*W = \alpha(E_{11})$, $WW^* = \beta(E_{11})$. Let $U = \sum_{i=1}^{\infty} \beta(E_{i1})W\alpha(E_{1i})$, the sum taken in the strict topology. It is straightforward to verify that U is unitary. Now, the span of the elements of the form $f \otimes e_{ij}$, where $f \in C_0(X)$ is dense in $C_0(X) \otimes \mathcal{K}$, and

$$\begin{aligned} U\alpha(f \otimes e_{kl})U^* &= \sum_{i,j} \beta(E_{i1})W\alpha(E_{1i})\alpha(f \otimes e_{kl})\alpha(E_{j1})W^*\beta(E_{1j}) = \\ &= \beta(E_{k1})W\alpha(f \otimes e_{11})W^*\beta(E_{1l}) = \beta(E_{k1})Wf \cdot \alpha(E_{11})W^*\beta(E_{1l}) = \\ &= f \cdot \beta(E_{k1})WW^*WW^*\beta(E_{1l}) = f \cdot \beta(E_{k1})\beta(E_{11})\beta(E_{1l}) = \\ &= f \cdot \beta(E_{kl}) = \beta(f \otimes e_{kl}) \end{aligned}$$

□

Remark 3.7. Notice that α is an automorphism if and only if $\text{vect}(\alpha)$ is a line bundle. So, if we restrict our attention to automorphisms, we get a map from the group of spectrum preserving automorphisms onto the group of Hermitian line bundles, whose kernel is the inner automorphisms. The group of line bundles is isomorphic to $H^2(X; \mathbb{Z})$ (at least in the case in which X is paracompact; see for example [K], I.3). So, we recover the exact sequence of Phillips and Raeburn mentioned in the introduction.

Composition of spectrum preserving endomorphisms maps under vect to tensor products. One might ask if there is a natural operation on spectrum preserving endomorphisms which is taken to the direct (Whitney) sum. We can indeed construct such an operation, as follows. Let $s_1, s_2 \in \mathcal{M}(C_0(X) \otimes \mathcal{K})$ be two isometries such that $s_1s_1^* + s_2s_2^* = 1$, and let α, β be spectrum preserving endomorphisms of $C_0(X) \otimes \mathcal{K}$, then we can define $\alpha \oplus_{(s_1, s_2)} \beta$ by

$$(\alpha \oplus_{(s_1, s_2)} \beta)(A) = s_1\alpha(A)s_1^* + s_2\beta(A)s_2^*$$

While the definition depends, of course, on the choice of s_1, s_2 , this direct sum construction is unique up to homotopy, and hence by theorem 3.6, up to equivalence via an inner automorphism. The uniqueness up to homotopy follows easily from the fact that each pair (s_1, s_2) corresponds in the obvious way to a map $X \rightarrow \mathcal{C}_2$, and any two such maps are homotopic, since \mathcal{C}_2 is contractible (theorem 2.5).

Define a map

$$\Psi : vect(\alpha) \oplus vect(\beta) \rightarrow vect(\alpha \oplus_{(s_1, s_2)} \beta)$$

by

$$\Psi(x, T_1 \oplus T_2) = (x, s_1(x)T_1 + s_2(x)T_2)$$

where $s_1(x), s_2(x)$ are the two isometries obtain by evaluating the map $X \rightarrow \mathcal{C}_2$ at x . To verify that this is a bundle map, we first have to see that $\Psi(x, T_1 \oplus T_2) \in vect(\alpha \oplus_{(s_1, s_2)} \beta)$. Let us denote the maps $X \rightarrow End(\mathcal{K})$ corresponding to α, β by $x \mapsto \alpha_x, \beta_x$, respectively, and fix $x \in X$, then for any $A \in \mathcal{K}$,

$$\begin{aligned} \Psi(T_1 \oplus T_2)A &= s_1(x)T_1A + s_2(x)T_2A = s_1(x)\alpha_x(A)T_1 + s_2(x)\beta_x(A)T_2 = \\ &= (\alpha \oplus_{(s_1, s_2)} \beta)_x(A)(s_1(x)T_1 + s_2(x)T_2) = (\alpha \oplus_{(s_1, s_2)} \beta)_x(A)\Psi(T_1 \oplus T_2) \end{aligned}$$

so indeed $\Psi(x, T_1 \oplus T_2) \in vect(\alpha \oplus_{(s_1, s_2)} \beta)$. Ψ is clearly a bundle map, so now it remains to show that it is an isomorphism. Since $s_1(x), s_2(x)$ are isometries, Ψ is injective. For surjectivity, let $(x, T) \in vect(\alpha \oplus_{(s_1, s_2)} \beta)$. Let $T_1 = s_1(x)^*T, T_2 = s_2(x)^*T$ then for any $A \in \mathcal{K}$, we have

$$\begin{aligned} T_1A &= s_1(x)^*TA = s_1(x)^*(\alpha \oplus_{(s_1, s_2)} \beta)_x(A)T = \\ &= s_1(x)^*(s_1(x)\alpha_x(A)s_1^* + s_2(x)\beta_x(A)s_2^*)T = \alpha_x(A)s_1^*T = \alpha_x(A)T_1 \end{aligned}$$

and similarly we have $T_2A = \beta_x(A)T_2$, so $(x, T_1 \oplus T_2) \in vect(\alpha) \oplus vect(\beta)$ and $\Psi(x, T_1 \oplus T_2) = (x, T)$, so Ψ is surjective.

We conclude the section with a summary of the main results.

Conclusion: To each spectrum preserving endomorphism α of $C_0(X) \otimes \mathcal{K}$ we associate the bundle $vect(\alpha)$ of intertwining operators.

1. $vect(\alpha)$ is numerable, and any numerable vector bundle arises as $vect(\alpha)$ for some spectrum preserving endomorphism α .
2. $vect(\alpha) \cong vect(\beta)$ if and only if $\alpha = AdU \circ \beta$ for some $U \in \mathcal{M}(C_0(X) \otimes \mathcal{K})$.
3. $vect(\alpha \circ \beta) = vect(\alpha) \otimes vect(\beta)$
4. $vect(\alpha)$ is a trivial bundle if and only if the endomorphism is inner, in the sense that there exist isometries $S_1, \dots, S_n \in \mathcal{M}(C_0(X) \otimes \mathcal{K})$ such that $\alpha(A) = \sum S_i A S_i^*$.

Remark 3.8. As in the previous section, we remark that the analogous results for real C^* -algebras can be obtained in a straightforward way, by going through the proofs, replacing the complex terminology by the real terminology.

4 Spectrum Preserving Endomorphisms of Stable Continuous-Trace C^* -algebras

Let E be a numerable fiber bundle over X with fiber \mathcal{K} and structure group $\text{Aut}(\mathcal{K})$, and let $\Gamma_0(E)$ be the C^* -algebra of sections. The purpose of this section is to generalize the main results of the previous section to this case.

Note that $\mathcal{M}(\Gamma_0(E))$ can be identified with the sections of a bundle with fiber $\mathcal{B}(\mathcal{H})$ and the same Dixmier-Douady class (see [PhR], Proposition 2.15 and the following remark). We refer to this bundle henceforth as the *multiplier bundle of E* . Denote by \mathcal{B}_x the fiber of the multiplier bundle over a point $x \in X$. For a spectrum preserving endomorphism α , we denote by α_x the action of α on the fiber over x (coming from the induced action of α on the quotient algebra corresponding to the primitive ideal represented by x), and then as before, we can define

$$\text{vect}(\alpha) = \{(x, T) \mid T \in \mathcal{B}_x, TA = \alpha_x(A)T \ \forall A \in E_x\}$$

Since E is locally trivial, we see that the restriction of $\text{vect}(\alpha)$ to each trivialization of E is locally trivial, and hence it is also locally trivial.

$\text{vect}(\alpha)$ determines α as above: if $N \subseteq X$ is such that $\text{vect}(\alpha)|_N, E|_N$ are trivial, then choose orthonormal sections S_1, \dots, S_n of the restricted bundle, and then $\alpha|_N(A) = \sum S_i A S_i^*$ for $A \in \Gamma(E|_N)$; those restrictions determine α .

If there is a unitary $U \in \mathcal{M}(\Gamma_0(E))$ such that $\alpha(A) = U^* \beta(A) U$ for all $A \in \Gamma_0(E)$ then $TA_x = \alpha_x(A_x)T$ if and only if $(U_x T)A_x = \beta_x(A_x)(U_x T)$, so left multiplication by U gives a vector bundle isomorphism $\text{vect}(\alpha) \cong \text{vect}(\beta)$.

Conversely, suppose $\text{vect}(\alpha) \cong \text{vect}(\beta)$. Note that as before, $\text{Hom}(\text{vect}(\alpha), \text{vect}(\beta))$ is isomorphic to the sub-bundle of the multiplier bundle whose fiber over x is $\text{span}\{TS^* \mid T \in \text{vect}(\beta), S \in \text{vect}(\alpha)\}$. Since there exists a unitary element in $\text{Hom}(\text{vect}(\alpha), \text{vect}(\beta))$, we know that there is a unitary section U of the above sub-bundle (and in particular, $U \in \mathcal{M}(\Gamma_0(E))$), and any such U satisfies $U\alpha(A)U^* = \beta(A)$ for all $A \in \Gamma_0(E)$.

So, as in the case of spectrum preserving endomorphisms of $C_0(X) \otimes \mathcal{K}$, we constructed a map vect from spectrum preserving endomorphisms to vector bundles such that $\text{vect}(\alpha) \cong \text{vect}(\beta)$ if and only if α and β are equivalent via an inner automorphism. The product formula from theorem 3.4 extends as well to this situation, with essentially the same proof.

If E_1, E_2 are two locally trivial \mathcal{K} -bundles over X , then $E_1 \otimes E_2$ is a locally trivial \mathcal{K} -bundle over X , and if α_1, α_2 are spectrum preserving endomorphisms of $\Gamma(E_1), \Gamma(E_2)$ respectively then we can have a spectrum preserving endomorphism $\alpha_1 \otimes \alpha_2$ of $\Gamma(E_1 \otimes E_2)$.

Lemma 4.1. *Let E_1, E_2 be two locally trivial \mathcal{K} -bundles over X , and let α_1, α_2 be spectrum preserving endomorphisms of $\Gamma(E_1), \Gamma(E_2)$, respectively.*

1. $\text{vect}(\alpha \otimes_X \beta) \cong \text{vect}(\alpha) \otimes \text{vect}(\beta)$
2. If $\varphi : \Gamma(E_1) \rightarrow \Gamma(E_2)$ is an isomorphism commuting with the multiplier action of $C_0(X)$ then $\text{vect}(\varphi \alpha \varphi^{-1}) \cong \text{vect}(\alpha)$

(cf [RW] lemma 6.6)

Proof: Let M_1, M_2 denote the multiplier bundles of E_1, E_2 respectively. We denote by $M_1 \otimes M_2$ the fiber-wise tensor product bundle, where the tensor product is taken to be the spatial tensor product, and observe that $M_1 \otimes M_2$ is canonically the multiplier bundle of $E_1 \otimes E_2$. $\text{vect}(\alpha \otimes \beta)$ is a sub-bundle of $M_1 \otimes M_2$, as is $\text{vect}(\alpha) \otimes \text{vect}(\beta)$, and it is immediate to verify that they are in fact the same. That proves (1). (2) follows immediately from the product formula (the extension of theorem 3.4). \square

We can now obtain surjectivity of vect as follows (cf [RW], p. 160). Suppose v is a given numerable vector bundle. Let E be our given \mathcal{K} -bundle, and let E_0 be the trivial bundle. By the results of the previous section, there exists a spectrum preserving endomorphism α_0 of $\Gamma(E_0) \cong C_0(X) \otimes \mathcal{K}$ with $\text{vect}(\alpha_0) \cong v$. Note that $\text{vect}(id_{\Gamma_0(E)})$ is the trivial line bundle ϵ_0 over X . Define a spectrum preserving endomorphism α of $\Gamma_0(E)$ by $\alpha = \alpha_0 \otimes id_{\Gamma_0(E)}$, then

$$\text{vect}(\alpha) \cong \text{vect}(\alpha_0) \otimes \text{vect}(id_{\Gamma_0(E)}) \cong v \otimes \epsilon_0 \cong v$$

If α is a given endomorphism $\Gamma_0(E)$, let E^{op} be the opposite bundle, and let $\alpha_0 = \alpha \otimes id_{E^{op}}$, then α_0 is an endomorphism of $E_0 \cong E \otimes E^{op}$, and as above, $\text{vect}(\alpha) \cong \text{vect}(\alpha_0)$, and we know that $\text{vect}(\alpha_0)$ is numerable, hence $\text{vect}(\alpha)$ is numerable as well.

5 Decomposition of General Endomorphisms

We now consider the case of a general endomorphism of $\Gamma_0(E)$. We first consider the case of the trivial bundle $\Gamma_0(E) \cong C_0(X) \otimes \mathcal{K}$, and then generalize the result to non-trivial bundles.

Any endomorphism α of $C_0(X) \otimes \mathcal{K}$ induces a (contravariant) continuous proper map $\alpha^* : X \rightarrow X$. Conversely, given a continuous proper map $f : X \rightarrow X$, we can define an endomorphism of $\gamma_f = f^* \otimes id$ of $C_0(X) \otimes \mathcal{K}$ (where f^* is the induced map on $C_0(X)$). Clearly, though, in general $\gamma_f \neq \alpha$ for $f = \alpha^*$.

Proposition 5.1. *Any endomorphism α of $C_0(X) \otimes \mathcal{K}$ decomposes into $\beta \circ \gamma_f$, where γ_f is as above and β is a spectrum preserving endomorphism.*

Proof: Let $f = \alpha^*$, and let $F \in C_0(X) \otimes \mathcal{K}$, thought of as a function $X \rightarrow \mathcal{K}$. So, for each $x \in X$, we have

$$\alpha(F)(x) = \sigma_x F(f(x))$$

for some (unital) endomorphism σ_x of \mathcal{K} . If $x_\nu \rightarrow x_0$ is a convergent net, then $\alpha(F)(x_\nu) \rightarrow \alpha(F)(x_0)$, and therefore $\sigma_{x_\nu} F(f(x_\nu)) \rightarrow \sigma_{x_0} F(f(x_0))$. Therefore (since F is continuous), we conclude that $\sigma_{x_\nu} \rightarrow \sigma_{x_0}$, so $x \mapsto \sigma_x$ is a continuous map from X to $\text{End}(\mathcal{K})$. Thus, this map $x \mapsto \sigma_x$ gives rise to a spectrum preserving endomorphism β , and clearly we have $\beta \circ \gamma_f = \alpha$, as required. \square

We now turn to the case of a non-trivial bundle. As in the case of $C_0(X) \otimes \mathcal{K}$, each endomorphism α of $\Gamma_0(E)$ induces a (contravariant) continuous proper map on the $\alpha^* : X \rightarrow X$. We henceforth denote this map by f_α .

We know that f_α lifts to a bundle map $E \rightarrow E$ (by its definition), and therefore $f_\alpha^* E \cong E$. By naturality of the Dixmier-Douady invariant (see [Ro] for example) we must have $f_\alpha^*(\delta(\Gamma_0(E))) = \delta(\Gamma_0(E))$.

In the other direction, let $f : X \rightarrow X$ be a continuous proper map, then it induces a pull-back homomorphism $\gamma_f : \Gamma_0(E) \rightarrow \Gamma_0(f^*E)$. If we know furthermore that $f^*(\delta(\Gamma_0(E))) = \delta(\Gamma_0(E))$, then we know that $f^*E \cong E$, and therefore f^*E is bundle-isomorphic to E (i.e. via an isomorphism which fixes X). Let $\varphi : \Gamma_0(f^*E) \rightarrow \Gamma_0(E)$ be a isomorphism which preserves X , so $\gamma = \varphi \circ \gamma_f$ is an endomorphism of $\Gamma_0(E)$ such that $f = f_\gamma$.

Let α now be a general endomorphism of $\Gamma_0(E)$, and let $f = f_\alpha$, $\gamma_f : \Gamma_0(f^*E) \rightarrow \Gamma_0(E)$ be as above. If $F \in \Gamma_0(E)$, $x \in X$ then $\alpha(F)(x) = \sigma_x((f_\alpha^*F)(x))$ for some homomorphism $\sigma_x : (f^*E)_x \rightarrow E_x$. Verifying that σ gives a continuous section of the bundle $Hom(f^*E, E)$ (the bundle of fiber-wise homomorphisms) is straightforward, and therefore they give rise to a homomorphism $\beta : \Gamma_0(f^*E) \rightarrow \Gamma_0(E)$ which commutes with the multiplier action of $C_0(X)$, and clearly $\alpha = \beta \circ \gamma_f$, giving us a decomposition analogous to that of proposition 5.1.

References

- [B] Blackadar, B., *K-Theory for Operator Algebras*, 2nd edition, MSRI publications **5**, Cambridge University Press, 1998.
- [Di] Dixmier, J., *C*-algebras*, North Holland, 1977.
- [Do] Dold, A., Partitions of unity in the theory of fibrations, *Ann. Math.*, **78** no. 2 (1963), 223–255.
- [K] Karoubi, M., *K-theory*, Grundlehren der mathematischen Wissenschaften **226**, Springer-Verlag, 1978.
- [L] Lance, E.C., Automorphisms of certain operator algebras, *Amer. J. Math.*, **91** (1969), 160–174.
- [PhR] Phillips, J. and Raeburn, I., Automorphisms of C^* -algebras and second Čech cohomology, *Indiana Univ. Math. J.* **29**, no. 6 (1980), 799–822.
- [Po] Powers, R.T., An index theory for semigroups of $*$ -endomorphisms of $\mathcal{B}(\mathcal{H})$ and type II_1 factors, *Can. J. Math.*, **40** no. 1 (1988), 86–114.
- [Pr] Price, G.L., Endomorphisms of certain operator algebras, *Publ. Res. Inst. Math. Sci.*, **25** (1989), 45–57.
- [RW] Raeburn, I. and Williams, D.P., *Morita Equivalence and Continuous-Trace C*-algebras*, Mathematical Surveys and Monographs **60**, The American Mathematical Society, 1998.

- [Ro] Rosenberg, J., Continuous-trace algebras from the bundle theoretic point of view, *J. Austral. Math. Soc. Ser. A* **47**, no. 3 (1989), 368–381.
- [S] Smith, M.S.B., On automorphism groups of C^* -algebras, *Trans. Amer. Math. Soc.*, **152** (1970), 623–648.

Ilan Hirshberg
Department of Mathematics
University of California at Berkeley
Berkeley, CA 94720
USA

email: *ilan@math.berkeley.edu*