

GENUS 2 CURVES WITH (3,3)-SPLIT JACOBIAN AND LARGE AUTOMORPHISM GROUP

T. SHASKA

ABSTRACT. Let \mathcal{C} be a genus 2 curve defined over k , $\text{char}(k) = 0$. If \mathcal{C} has a (3,3)-split Jacobian then we show that the automorphism group $\text{Aut}(\mathcal{C})$ is isomorphic to one of the following: \mathbb{Z}_2, V_4, D_8 , or D_{12} . There are exactly six \mathbb{C} -isomorphism classes of genus two curves \mathcal{C} with $\text{Aut}(\mathcal{C})$ isomorphic to D_8 (resp., D_{12}). We show that exactly four (resp., three) of these classes with group D_8 (resp., D_{12}) have representatives defined over \mathbb{Q} . We discuss some of these curves in detail and find their rational points.

1. INTRODUCTION

Let \mathcal{C} be a genus 2 curve defined over an algebraically closed field k , of characteristic zero. We denote by $K := k(\mathcal{C})$ its function field and by $\text{Aut}(\mathcal{C}) := \text{Aut}(K/k)$ the automorphism group of \mathcal{C} . Let $\psi : \mathcal{C} \rightarrow \mathcal{E}$ be a degree n maximal covering (i.e. does not factor through an isogeny) to an elliptic curve \mathcal{E} defined over k . We say that \mathcal{C} has a *degree n elliptic subcover*. Degree n elliptic subcovers occur in pairs. Let $(\mathcal{E}, \mathcal{E}')$ be such a pair. It is well known that there is an isogeny of degree n^2 between the Jacobian $J_{\mathcal{C}}$ of \mathcal{C} and the product $\mathcal{E} \times \mathcal{E}'$. We say that \mathcal{C} has **(n,n)-split Jacobian**. The locus of such \mathcal{C} (denoted by \mathcal{L}_n) is an algebraic subvariety of the moduli space \mathcal{M}_2 of genus two curves. For the equation of \mathcal{L}_2 in terms of Igusa invariants, see [19]. Computation of the equation of \mathcal{L}_3 was the main focus of [18]. For $n > 3$, equations of \mathcal{L}_n have not yet been computed.

Equivalence classes of degree 2 coverings $\psi : \mathcal{C} \rightarrow \mathcal{E}$ are in 1-1 correspondence with conjugacy classes of non-hyperelliptic involutions in $\text{Aut}(\mathcal{C})$. In any characteristic different from 2, the automorphism group $\text{Aut}(\mathcal{C})$ is isomorphic to one of the following: $\mathbb{Z}_2, \mathbb{Z}_{10}, V_4, D_8, D_{12}, \mathbb{Z}_3 \rtimes D_8, GL_2(3)$, or 2^+S_5 ; see [19]. Here V_4 is the Klein 4-group, D_8 (resp., D_{12}) denotes the dihedral group of order 8 (resp., 12), and $\mathbb{Z}_2, \mathbb{Z}_{10}$ are cyclic groups of order 2 and 10. For a description of other groups, see [19]. If $\text{Aut}(\mathcal{C}) \cong \mathbb{Z}_{10}$ then \mathcal{C} is isomorphic to $Y^2 = X^6 - X$. Thus, if \mathcal{C} has extra automorphisms and it is not isomorphic to $Y^2 = X^6 - X$ then $\mathcal{C} \in \mathcal{L}_2$. We say that a genus 2 curve \mathcal{C} has **large automorphism group** if the order of $\text{Aut}(\mathcal{C})$ is bigger than 4.

In section 2, we describe the loci for genus 2 curves with $\text{Aut}(\mathcal{C})$ isomorphic to D_8 or D_{12} in terms of Igusa invariants. From these invariants we are able to determine the field of definition of a curve \mathcal{C} with $\text{Aut}(\mathcal{C}) \cong D_8$ or D_{12} . Further, we find the equation for this \mathcal{C} and j -invariants of degree 2 elliptic subcovers in terms of i_1, i_2, i_3 (cf. section 2). This determines the fields of definition for these elliptic subcovers.

Let \mathcal{C} be a genus 2 curve with (3,3)-split Jacobian. In section 3 we give a short description of the space \mathcal{L}_3 . Results described in section 3 follow from [18], even though sometimes nontrivially. We find equations of degree 3 elliptic subcovers

in terms of the coefficients of \mathcal{C} . In section 4, we show that $\text{Aut}(\mathcal{C})$ is one of the following: \mathbb{Z}_2 , V_4 , D_8 , or D_{12} . Moreover, we show that there are exactly six \mathbb{C} -isomorphism classes of genus two curves $\mathcal{C} \in \mathcal{L}_3$ with automorphism group D_8 (resp., D_{12}). We explicitly find the absolute invariants i_1, i_2, i_3 which determine these classes. For each such class we give the equation of a representative genus 2 curve \mathcal{C} . We notice that there are four cases (resp., three) such that the triple of invariants $(i_1, i_2, i_3) \in \mathbb{Q}^3$ when $\text{Aut}(\mathcal{C}) \cong D_8$ (resp., $\text{Aut}(\mathcal{C}) \cong D_{12}$). Using results from section 2, we determine that there are exactly four (resp., three) genus 2 curves $\mathcal{C} \in \mathcal{L}_3$ (up to \mathbb{Q} -isomorphism) with group D_8 (resp., D_{12}) defined over \mathbb{Q} and list their equations in Table 1. We discuss these curves and their degree 2 and 3 elliptic subcovers in more detail in section 5. Our focus is on the cases which have elliptic subcovers defined over \mathbb{Q} . In some of these cases we are able to use these subcovers to find the rational points of the genus 2 curve. This technique has been used before by Flynn and Wetherell [5] for degree 2 elliptic subcovers.

Curves of genus 2 with degree 2 elliptic subcovers (or with elliptic involutions) were first studied by Legendre and Jacobi. The genus 2 curve with the largest known number of rational points has automorphism group isomorphic to D_{12} ; thus it has degree 2 elliptic subcovers. It was found by Keller and Kulesz and it is known to have at least 588 rational points; see [10]. Using degree 2 elliptic subcovers Howe, Leprevost, and Poonen [8] were able to construct a family of genus 2 curves whose Jacobians each have large rational torsion subgroups. Similar techniques probably could be applied using degree 3 elliptic subcovers. Curves of genus 2 with degree 3 elliptic subcovers have already occurred in the work of Clebsch, Hermite, Goursat, Burkhardt, Brioschi, and Bolza in the context of elliptic integrals. For a history of this topic see Krazer [11] (p. 479). For more recent work see Kuhn [12] and [18]. More generally, degree n elliptic subfields of genus 2 fields have been studied by Frey [6], Frey and Kani [7], Kuhn [12], and this author [17].

Acknowledgements: The author wants to thank G. Cardona for pointing out the parametrizations in equations (5) and (6) which made some of the computations easier.

2. GENUS TWO CURVES AND THE MODULI SPACE \mathcal{M}_2 .

Let k be an algebraically closed field of characteristic zero and \mathcal{C} a genus 2 curve defined over k . Then \mathcal{C} can be described as a double cover of $\mathbb{P}^1(k)$ ramified in 6 places w_1, \dots, w_6 . This sets up a bijection between isomorphism classes of genus 2 curves and unordered distinct 6-tuples $w_1, \dots, w_6 \in \mathbb{P}^1(k)$ modulo automorphisms of $\mathbb{P}^1(k)$. An unordered 6-tuple $\{w_i\}_{i=1}^6$ can be described by a binary sextic, (i.e. a homogenous equation $f(X, Z)$ of degree 6). Let \mathcal{M}_2 denote the moduli space of genus 2 curves; see [16]. To describe \mathcal{M}_2 we need to find polynomial functions of the coefficients of a binary sextic $f(X, Z)$ invariant under linear substitutions in X, Z of determinant one. These invariants were worked out by Clebsch and Bolza in the case of zero characteristic and generalized by Igusa for any characteristic different from 2; see [1], [9].

Consider a binary sextic i.e. homogeneous polynomial $f(X, Z)$ in $k[X, Z]$ of degree 6:

$$f(X, Z) = a_6 X^6 + a_5 X^5 Z + \dots + a_0 Z^6$$

Classical invariants $\{J_{2i}\}$ of $f(X, Z)$ are homogeneous polynomials of degree $2i$ in $k[a_0, \dots, a_6]$, for $i = 1, 2, 3, 5$; see [9], [19] for their definitions. Here J_{10} is simply the

discriminant of $f(X, Z)$. It vanishes if and only if the binary sextic has a multiple linear factor. These J_{2i} are invariant under the natural action of $SL_2(k)$ on sextics. Dividing such an invariant by another one of the same degree gives an invariant under $GL_2(k)$ action.

Two genus 2 fields K (resp., curves) in the standard form $Y^2 = f(X, 1)$ are isomorphic if and only if the corresponding sextics are $GL_2(k)$ conjugate. Thus if I is a $GL_2(k)$ invariant (resp., homogeneous $SL_2(k)$ invariant), then the expression $I(K)$ (resp., the condition $I(K) = 0$) is well defined. Thus the $GL_2(k)$ invariants are functions on the moduli space \mathcal{M}_2 of genus 2 curves. This \mathcal{M}_2 is an affine variety with coordinate ring

$$k[\mathcal{M}_2] = k[a_0, \dots, a_6, J_{10}^{-1}]^{GL_2(k)} = \text{subring of degree 0 elements in}$$

$k[J_2, \dots, J_{10}, J_{10}^{-1}]$; see Igusa [9]. The *absolute invariants*

$$(1) \quad i_1 := 144 \frac{J_4}{J_2^2}, \quad i_2 := -1728 \frac{J_2 J_4 - 3J_6}{J_2^3}, \quad i_3 := 486 \frac{J_{10}}{J_2^5}$$

are even $GL_2(k)$ -invariants. Two genus 2 curves with $J_2 \neq 0$ are isomorphic if and only if they have the same absolute invariants. If $J_2 = 0$ then we can define new invariants as in [18]. For the rest of this paper if we say ‘‘there is a genus 2 curve \mathcal{C} defined over k ’’ we will mean the k -isomorphism class of \mathcal{C} .

One can define $GL_2(k)$ invariants with J_{10} in the denominator which will be defined everywhere. However, this is not efficient in doing computations since the degrees of these rational functions in terms of the coefficients of \mathcal{C} will be multiples of 10 and therefore higher than degrees of i_1, i_2, i_3 . For the purposes of this paper defining i_1, i_2, i_3 as above is not a restriction as it will be seen in the proof of the theorem 1. For the proofs of the following two lemmas, see [19].

Lemma 1. *The automorphism group G of a genus 2 curve \mathcal{C} in characteristic $\neq 2$ is isomorphic to $\mathbb{Z}_2, \mathbb{Z}_{10}, V_4, D_8, D_{12}, \mathbb{Z}_3 \rtimes D_8, GL_2(3)$, or $2^+ S_5$. The case when $G \cong 2^+ S_5$ occurs only in characteristic 5. If $G \cong \mathbb{Z}_3 \rtimes D_8$ (resp., $GL_2(3)$) then \mathcal{C} has equation $Y^2 = X^6 - 1$ (resp., $Y^2 = X(X^4 - 1)$). If $G \cong \mathbb{Z}_{10}$ then \mathcal{C} has equation $Y^2 = X^6 - X$.*

Remark 2. *It is worth mentioning that the analogue of the above lemma has been settled for all $2 \leq g \leq 48$ in characteristic zero; see [14]. This was a major computational effort using the computer algebra system GAP.*

For the rest of this paper we assume that $\text{char}(k) = 0$.

Lemma 3. *i) The locus \mathcal{L}_2 of genus 2 curves \mathcal{C} which have a degree 2 elliptic subcover is a closed subvariety of \mathcal{M}_2 . The equation of \mathcal{L}_2 is given by equation (17) in [19].*

ii) The locus of genus 2 curves \mathcal{C} with $\text{Aut}(\mathcal{C}) \cong D_8$ is given by the equation of \mathcal{L}_2 and

$$(2) \quad 1706J_4^2 J_2^2 + 2560J_4^3 + 27J_4 J_2^4 - 81J_2^3 J_6 - 14880J_2 J_4 J_6 + 28800J_6^2 = 0$$

ii) The locus of genus 2 curves \mathcal{C} with $\text{Aut}(\mathcal{C}) \cong D_{12}$ is

$$(3) \quad \begin{aligned} & -J_4 J_2^4 + 12J_2^3 J_6 - 52J_4^2 J_2^2 + 80J_4^3 + 960J_2 J_4 J_6 - 3600J_6^2 = 0 \\ & 864J_{10} J_2^5 + 3456000J_{10} J_4^2 J_2 - 43200J_{10} J_4 J_2^3 - 2332800000J_{10}^2 - J_4^2 J_2^6 \\ & \quad - 768J_4^4 J_2^2 + 48J_4^3 J_2^4 + 4096J_4^5 = 0 \end{aligned}$$

We will refer to the locus of genus 2 curves \mathcal{C} with $\text{Aut}(\mathcal{C}) \cong D_8$ (resp., $\text{Aut}(\mathcal{C}) \cong D_8$) as D_8 -locus (resp., D_{12} -locus).

Each genus 2 curve $\mathcal{C} \in \mathcal{L}_2$ has a non-hyperelliptic involution $v_0 \in \text{Aut}(\mathcal{C})$. There is another non-hyperelliptic involution $v'_0 := v_0 w$, where w is the hyperelliptic involution. Thus, degree 2 elliptic subcovers come in pairs. We denote the pair of degree 2 elliptic subcovers by (E_0, E'_0) . If $\text{Aut}(\mathcal{C}) \cong D_8$ then $E_0 \cong E'_0$ or E_0 and E'_0 are 2-isogenous. If $\text{Aut}(\mathcal{C}) \cong D_{12}$, then E_0 and E'_0 are isogenous of degree 3. See [19] for details.

Lemma 4. *Let \mathcal{C} be a genus 2 curve defined over k . Then,*

i) *$\text{Aut}(\mathcal{C}) \cong D_8$ if and only if \mathcal{C} is isomorphic to*

$$(4) \quad Y^2 = X^5 + X^3 + tX$$

for some $t \in k \setminus \{0, \frac{1}{4}, \frac{9}{100}\}$.

ii) *$\text{Aut}(\mathcal{C}) \cong D_{12}$ if and only if \mathcal{C} is isomorphic to*

$$(5) \quad Y^2 = X^6 + X^3 + t$$

for some $t \in k \setminus \{0, \frac{1}{4}, -\frac{1}{50}\}$.

Proof. i) $\text{Aut}(\mathcal{C}) \cong D_8$: Then \mathcal{C} is isomorphic to

$$Y^2 = (X^2 - 1)(X^4 - \lambda X^2 + 1)$$

for $\lambda \neq \pm 2$; see [19]. Denote $\tau := \sqrt{-2\frac{\lambda+6}{\lambda-2}}$. The transformation

$$\phi : (X, Y) \rightarrow \left(\frac{\tau x - 1}{\tau x + 1}, \frac{4\tau}{(\tau x + 1)^3} \cdot \frac{(\lambda + 6)^2}{\lambda - 2} \right)$$

gives

$$Y^2 = X^5 + X^3 + tX$$

where $t = (\frac{\lambda-2}{2(\lambda-6)})^2$ and $t \neq 0, \frac{1}{4}$. If $t = \frac{9}{100}$ then $\text{Aut}(\mathcal{C})$ has order 24.

Conversely, the absolute invariants i_1, i_2, i_3 of a genus 2 curve \mathcal{C} isomorphic to

$$Y^2 = X^5 + X^3 + tX$$

satisfy the locus as described in lemma 2, part ii). Thus, $\text{Aut}(\mathcal{C}) \cong D_8$.

ii) $\text{Aut}(\mathcal{C}) \cong D_{12}$: In [19] it is shown that \mathcal{C} is isomorphic to

$$Y^2 = (X^3 - 1)(X^3 - \lambda)$$

for $\lambda \neq 0, 1$ and $\lambda^2 - 38\lambda + 1 \neq 0$. Then,

$$\phi : (X, Y) \rightarrow ((\lambda + 1)^{\frac{1}{3}} X, (\lambda + 1) Y)$$

transforms \mathcal{C} to the curve with equation

$$Y^2 = X^6 + X^3 + t$$

where $t = \frac{\lambda}{(\lambda+1)^2}$ and $t \neq 0, \frac{1}{4}$. If $t = -\frac{1}{50}$ then $\text{Aut}(\mathcal{C})$ has order 48.

The absolute invariants i_1, i_2, i_3 of a genus 2 curve \mathcal{C} isomorphic to

$$Y^2 = X^6 + X^3 + t$$

satisfy the locus as described in lemma 2, part iii). Thus, $\text{Aut}(\mathcal{C}) \cong D_{12}$. This completes the proof. \square

The following lemma determines a genus 2 curve for each point in the D_8 or D_{12} locus.

Lemma 5. Let $\mathbf{p} := (J_2, J_4, J_6, J_{10})$ be a point in \mathcal{L}_2 such that $J_2 \neq 0$ and (i_1, i_2, i_3) the corresponding absolute invariants.

i) If \mathbf{p} is in the D_8 -locus, then the genus two curve \mathcal{C} corresponding to \mathbf{p} is given by:

$$Y^2 = X^5 + X^3 + -\frac{3}{4} \frac{345i_1^2 + 50i_1i_2 - 90i_2 - 1296i_1}{2925i_1^2 + 250i_1i_2 - 9450i_2 - 54000i_1 + 139968} X$$

ii) If \mathbf{p} is in the D_8 -locus, then the genus two curve \mathcal{C} corresponding to \mathbf{p} is given by:

$$Y^2 = X^6 + X^4 + \frac{1}{4} \frac{540i_1^2 + 100i_1i_2 - 1728i_1 + 45i_2}{2700i_1^2 + 1000i_1i_2 + 204525i_1 + 40950i_2 - 708588}$$

Proof. i) By previous lemma every genus 2 curve \mathcal{C} with automorphism group D_8 is isomorphic to $Y^2 = X^5 + X^3 + tX$. Since $J_2 \neq 0$ then $t \neq -\frac{3}{20}$ and the absolute invariants are:

$$(6) \quad i_1 = -144t \frac{(20t-9)}{(20t+3)^2}, \quad i_2 = 3456t^2 \frac{(140t-27)}{(20t+3)^3}, \quad i_3 = 243t^3 \frac{(4t-1)^2}{(20t+3)^5}$$

From the above system we have

$$t = -\frac{3}{4} \frac{345i_1^2 + 50i_1i_2 - 90i_2 - 1296i_1}{2925i_1^2 + 250i_1i_2 - 9450i_2 - 54000i_1 + 139968}$$

ii) By previous lemma every genus 2 curve \mathcal{C} with automorphism group D_{12} is isomorphic to $Y^2 = X^6 + X^3 + t$. The absolute invariants are:

$$(7) \quad i_1 = 1296 \frac{t(5t+1)}{(40t-1)^2}, \quad i_2 = -11664 \frac{t(20t^2+26t-1)}{(40t-1)^3}, \quad i_3 = \frac{729}{16} \frac{t^2(4t-1)^3}{(40t-1)^5}$$

From the above system we have

$$t = \frac{1}{4} \frac{540i_1^2 + 100i_1i_2 - 1728i_1 + 45i_2}{2700i_1^2 + 1000i_1i_2 + 204525i_1 + 40950i_2 - 708588}$$

This completes the proof. \square

Note: If $J_2 = 0$ then there is exactly one isomorphism class of genus 2 curves with automorphism group D_8 (resp., D_{12}) given by $Y^2 = X^5 + X^3 - \frac{3}{20}X$ (resp., $Y^2 = X^6 + X^3 - \frac{1}{40}$).

Remark 6. If the invariants $i_1, i_2, i_3 \in \mathbb{Q}$ then from above lemma there is a \mathcal{C} corresponding to these invariants defined over \mathbb{Q} . If a genus 2 curve does not have extra automorphisms (i.e. $\text{Aut}(\mathcal{C}) \cong \mathbb{Z}_2$), then an algorithm of Mestre determines if the curve is defined over \mathbb{Q} .

If the order of the automorphism group $\text{Aut}(\mathcal{C})$ is divisible by 4, then \mathcal{C} has degree 2 elliptic subcovers. These elliptic subcovers are determined explicitly in [19]. Do these elliptic subcovers of \mathcal{C} have the same field of definition as \mathcal{C} ? In general the answer is negative. The following lemma determines the field of definition of these elliptic subcovers when $\text{Aut}(\mathcal{C})$ is isomorphic to D_8 or D_{12} .

Lemma 7. Let \mathcal{C} be a genus 2 curve defined over k , $\text{char}(k) = 0$.

i) If \mathcal{C} has equation

$$Y^2 = X^5 + X^3 + tX$$

where $t \in k \setminus \{\neq 0, \frac{1}{4}, \frac{9}{100}\}$, then its degree 2 elliptic subfields have j -invariants given by

$$j^2 - 128 \frac{2000t^2 + 1440t + 27}{(4t-1)^2} + 4096 \frac{(100t-9)^3}{(4t-1)^3} = 0$$

ii) If \mathcal{C} has equation

$$Y^2 = X^6 + X^3 + t$$

where $t \in k \setminus \{\neq 0, \frac{1}{4}, -\frac{1}{50}\}$, then its degree 2 elliptic subfields have j -invariants given by

$$j^2 - 13824t \frac{500t^2 + 965t + 27}{(4t-1)^3} + 47775744t \frac{(25t-4)^3}{(4t-1)^4} = 0$$

Proof. The proof is elementary and follows from [19]. \square

3. CURVES OF GENUS 2 WITH DEGREE 3 ELLIPTIC SUBCOVERS

In this section we will give a brief description of spaces \mathcal{L}_2 and \mathcal{L}_3 . In case $J_2 \neq 0$ we take these spaces as equations in terms of i_1, i_2, i_3 , otherwise as homogeneous equations in terms of J_2, J_4, J_6, J_{10} . By a point $\mathfrak{p} \in \mathcal{L}_3$ we will mean a tuple (J_2, J_4, J_6, J_{10}) which satisfies equation of \mathcal{L}_3 . When it is clear that $J_2 \neq 0$ then $\mathfrak{p} \in \mathcal{L}_3$ would mean a triple $(i_1, i_2, i_3) \in \mathcal{L}_3$. As before k is an algebraically closed field of characteristic zero.

Definition 8. A **non-degenerate pair** (resp., **degenerate pair**) is a pair $(\mathcal{C}, \mathcal{E})$ such that \mathcal{C} is a genus 2 field with a degree 3 elliptic subcover \mathcal{E} where $\psi : \mathcal{C} \rightarrow \mathcal{E}$ is ramified in two (resp., one) places. Two such pairs $(\mathcal{C}, \mathcal{E})$ and $(\mathcal{C}', \mathcal{E}')$ are called isomorphic if there is a k -isomorphism $\mathcal{C} \rightarrow \mathcal{C}'$ mapping $\mathcal{E} \rightarrow \mathcal{E}'$.

If $(\mathcal{C}, \mathcal{E})$ is a non-degenerate pair, then \mathcal{C} can be parameterized as follows

$$(8) \quad Y^2 = (\mathfrak{v}^2 X^3 + \mathfrak{u}\mathfrak{v}X^2 + \mathfrak{v}X + 1)(4\mathfrak{v}^2 X^3 + \mathfrak{v}^2 X^2 + 2\mathfrak{v}X + 1)$$

where $\mathfrak{u}, \mathfrak{v} \in k$ and the discriminant

$$\Delta = -16 \mathfrak{v}^{17} (\mathfrak{v} - 27) (27\mathfrak{v} + 4\mathfrak{v}^2 - \mathfrak{u}^2\mathfrak{v} + 4\mathfrak{u}^3 - 18\mathfrak{u}\mathfrak{v})^3$$

of the sextic is nonzero. We let $R := (27\mathfrak{v} + 4\mathfrak{v}^2 - \mathfrak{u}^2\mathfrak{v} + 4\mathfrak{u}^3 - 18\mathfrak{u}\mathfrak{v}) \neq 0$. For $4\mathfrak{u} - \mathfrak{v} - 9 \neq 0$ degree 3 coverings are given by

$$U_1 = \frac{\mathfrak{v}X^2}{\mathfrak{v}^2 X^3 + \mathfrak{u}\mathfrak{v}X^2 + \mathfrak{v}X + 1}, \quad U_2 = \frac{(\mathfrak{v}X + 3)^2 ((4\mathfrak{u} - \mathfrak{v} - 9)X + 3\mathfrak{u} - \mathfrak{v})}{\mathfrak{v}(4\mathfrak{u} - \mathfrak{v} - 9)(4\mathfrak{v}^2 X^3 + \mathfrak{v}^2 X^2 + 2\mathfrak{v}X + 1)}$$

and elliptic curves have equations:

$$(9) \quad \begin{aligned} \mathcal{E} : \quad V_1^2 &= R U_1^3 - (12\mathfrak{u}^2 - 2\mathfrak{u}\mathfrak{v} - 18\mathfrak{v})U_1^2 + (12\mathfrak{u} - \mathfrak{v})U_1 - 4 \\ \mathcal{E}' : \quad V_2^2 &= c_3 U_2^3 + c_2 U_2^2 + c_1 U_2 + c_0 \end{aligned}$$

where

$$(10) \quad \begin{aligned} c_0 &= -(9\mathfrak{u} - 2\mathfrak{v} - 27)^3 \\ c_1 &= (4\mathfrak{u} - \mathfrak{v} - 9)(729\mathfrak{u}^2 + 54\mathfrak{u}^2\mathfrak{v} - 972\mathfrak{u}\mathfrak{v} - 18\mathfrak{u}\mathfrak{v}^2 + 189\mathfrak{v}^2 + 729\mathfrak{v} + \mathfrak{v}^3) \\ c_2 &= -\mathfrak{v}(4\mathfrak{u} - \mathfrak{v} - 9)^2(54\mathfrak{u} + \mathfrak{u}\mathfrak{v} - 27\mathfrak{v}) \\ c_3 &= \mathfrak{v}^2(4\mathfrak{u} - \mathfrak{v} - 9)^3 \end{aligned}$$

The above facts can be deduced from lemma 1 of [18]. The case $4u - v - 9 = 0$ is treated separately in [18]. There is an automorphism $\beta \in \text{Gal}_{k(u,v)/k(i_1, i_2, i_3)}$

$$(11) \quad \begin{aligned} \beta(u) &= \frac{(v - 3u)(324u^2 + 15u^2v - 378uv - 4uv^2 + 243v + 72v^2)}{(v - 27)(4u^3 + 27v - 18uv - u^2v + 4v^2)} \\ \beta(v) &= -\frac{4(v - 3u)^3}{4u^3 + 27v - 18uv - u^2v + 4v^2} \end{aligned}$$

which permutes the j -invariants of \mathcal{E} and \mathcal{E}' . The map

$$\theta : (u, v) \rightarrow (i_1, i_2, i_3)$$

defined when $J_2 \neq 0$ and $\Delta \neq 0$ has degree 2. Denote by J_θ the Jacobian matrix of θ . Then $\det(J_\theta) = 0$ consist of the (non-singular) curve \mathfrak{X} given by

$$(12) \quad \mathfrak{X} : 8v^3 + 27v^2 - 54uv^2 - u^2v^2 + 108u^2v + 4u^3v - 108u^3 = 0$$

and 6 isolated (u, v) solutions. These solutions correspond to the following values for (i_1, i_2, i_3) :

$$(13) \quad \left(-\frac{8019}{20}, -\frac{1240029}{200}, -\frac{531441}{100000}\right), \left(\frac{729}{2116}, \frac{1240029}{97336}, \frac{531441}{13181630464}\right), \left(81, -\frac{5103}{25}, -\frac{729}{12500}\right)$$

We denote the image of \mathfrak{X} in the \mathcal{L}_3 locus by \mathfrak{Y} . The map θ restricted to \mathfrak{X} is unirational. The curve \mathfrak{Y} can be computed as an affine curve in terms of i_1, i_2 . For each point $\mathfrak{p} \in \mathfrak{Y}$ degree 3 elliptic subcovers are isomorphic. If \mathfrak{p} is an ordinary point in \mathfrak{Y} and $\mathfrak{p} \neq \mathfrak{p}_6$ (cf. Table 1) then the corresponding curve $\mathcal{C}_{\mathfrak{p}}$ has automorphism group V_4 .

If $(\mathcal{C}, \mathcal{E})$ is a degenerate pair then \mathcal{C} can be parameterized as follows

$$Y^2 = (3X^2 + 4)(X^3 + X + c)$$

for some c such that $c^2 \neq -\frac{4}{27}$; see [18]. We define $\mathfrak{w} := c^2$. The map

$$\mathfrak{w} \rightarrow (i_1, i_2, i_3)$$

is injective as it was shown in [18].

Definition 9. Let \mathfrak{p} be a point in \mathcal{L}_3 . We say \mathfrak{p} is a **generic point** in \mathcal{L}_3 if the corresponding $(\mathcal{C}_{\mathfrak{p}}, \mathcal{E})$ is a non-degenerate pair. We define

$$e_3(\mathfrak{p}) := \begin{cases} |\theta^{-1}(\mathfrak{p})|, & \text{if } \mathfrak{p} \text{ is a generic point} \\ 1 & \text{otherwise} \end{cases}$$

In [18] it is shown that the pairs (u, v) with $\Delta(u, v) \neq 0$ bijectively parameterize the isomorphism classes of non-degenerate pairs $(\mathcal{C}, \mathcal{E})$. Those \mathfrak{w} with $\mathfrak{w} \neq -\frac{4}{27}$ bijectively parameterize the isomorphism classes of degenerate pairs $(\mathcal{C}, \mathcal{E})$. Thus, the number $e_3(\mathfrak{p})$ is the number of isomorphism classes of such pairs $(\mathcal{C}, \mathcal{E})$. In [18] it is shown that $e_3(\mathfrak{p}) = 0, 1, 2$, or 4. The following lemma describes the locus \mathcal{L}_3 . For details see [18].

Lemma 10. *The locus \mathcal{L}_3 of genus 2 curves with degree 3 elliptic subcovers is the closed subvariety of \mathcal{M}_2 defined by the equation*

$$(14) \quad C_8 J_{10}^8 + \cdots + C_1 J_{10} + C_0 = 0$$

where coefficients $C_0, \dots, C_8 \in k[J_2, J_6, J_{10}]$ are displayed in [18].

As noted above, with the assumption $J_2 \neq 0$ equation (14) can be written in terms of i_1, i_2, i_3 .

4. AUTOMORPHISM GROUPS OF GENUS 2 CURVES WITH DEGREE 3 ELLIPTIC SUBCOVERS

Let $\mathcal{C} \in \mathcal{L}_3$ be a genus 2 curve defined over an algebraically closed field k , $\text{char}(k) = 0$. The following theorem determines the automorphism group of \mathcal{C} .

Theorem 11. *Let \mathcal{C} be a genus two curve which has a degree 3 elliptic subcover. Then the automorphism group of \mathcal{C} is one of the following: \mathbb{Z}_2, V_4, D_8 , or D_{12} . Moreover, there are exactly six curves $\mathcal{C} \in \mathcal{L}_3$ with automorphism group D_8 and six curves $\mathcal{C} \in \mathcal{L}_3$ with automorphism group D_{12} .*

Proof. We denote by $G := \text{Aut}(\mathcal{C})$. None of the curves $Y^2 = X^6 - X$, $Y^2 = X^6 - 1$, $Y^2 = X^5 - X$ have degree 3 elliptic subcovers since their J_2, J_4, J_6, J_{10} invariants don't satisfy equation (14). From lemma 1 we have the following cases:

i) If $G \cong D_8$, then \mathcal{C} is isomorphic to

$$Y^2 = X^5 + X^3 + tX$$

as in Lemma 3. Classical invariants are:

$$J_2 = 40t + 6, \quad J_4 = 4t(9 - 20t), \quad J_6 = 8t(22t + 9 - 40t^2), \quad J_{10} = 16t^3(4t - 1)^2.$$

Substituting in the equation (14) we have the following equation:

$$(15) \quad (196t - 81)^4(49t - 12)(5t - 1)^4(700t + 81)^4(490000t^2 - 136200t + 2401)^2 = 0$$

For

$$t = \frac{81}{196}, \frac{12}{49}, \frac{1}{5}, -\frac{81}{700}$$

the triple (i_1, i_2, i_3) has the following values respectively:

$$\left(\frac{729}{2116}, \frac{1240029}{97336}, \frac{531441}{13181630464}\right), \quad \left(\frac{4288}{1849}, \frac{243712}{79507}, \frac{64}{1323075987}\right), \\ \left(\frac{144}{49}, \frac{3456}{8575}, \frac{243}{52521875}\right), \quad \left(-\frac{8019}{20}, -\frac{1240029}{200}, -\frac{531441}{10000}\right)$$

If

$$490000t^2 - 136200t + 2401 = 0$$

then we have two distinct triples (i_1, i_2, i_3) which are in $\mathbb{Q}(\sqrt{2})$. Thus, there are exactly 6 genus 2 curves $\mathcal{C} \in \mathcal{L}_3$ with automorphism group D_8 and only four of them have rational invariants.

ii) If $G \cong D_{12}$ then \mathcal{C} is isomorphic to a genus 2 curve in the form

$$Y^2 = X^6 + X^3 + t$$

as in Lemma 3. Then, $J_2 = -6(40t - 1)$ and

$$J_4 = 324t(5t + 1), \quad J_6 = -162t(740t^2 + 62t - 1), \quad J_{10} = -729t^2(4t - 1)$$

Then the equation of \mathcal{L}_3 becomes:

$$(16) \quad (25t - 4)(11t + 4)^3(20t - 1)^6(111320000t^3 - 60075600t^2 + 13037748t + 15625)^3 = 0$$

For

$$t = \frac{4}{25}, -\frac{4}{11}, \frac{1}{20}$$

the corresponding values for (i_1, i_2, i_3) are respectively:

$$\left(\frac{64}{5}, \frac{1088}{25}, \frac{1}{84375}\right), \quad \left(\frac{576}{361}, \frac{60480}{6859}, \frac{243}{2476099}\right), \quad \left(81, -\frac{5103}{25}, -\frac{729}{12500}\right)$$

If

$$111320000t^3 - 60075600t^2 = 13037748t + 15625 = 0$$

then there are three distinct triples (i_1, i_2, i_3) none of which is rational. Hence, there are exactly 6 classes of genus 2 curves $\mathcal{C} \in \mathcal{L}_3$ with $\text{Aut}(\mathcal{C}) \cong D_{12}$ of which three have rational invariants.

iii) $G \cong V_4$. There is a 1-dimensional family of genus 2 curves with a degree 3 elliptic subcover and automorphism group V_4 given by \mathfrak{Y} .

iv) Generically genus 2 curves \mathcal{C} have $\text{Aut}(\mathcal{C}) \cong \mathbb{Z}_2$. For example, every point $\mathfrak{p} \in \mathcal{L}_3 \setminus \mathcal{L}_2$ correspond to a class of genus 2 curves with degree 3 elliptic subcovers and automorphism group isomorphic to \mathbb{Z}_2 . This completes the proof. \square

The theorem determines that there are exactly 12 genus 2 curves $\mathcal{C} \in \mathcal{L}_3$ with automorphism group D_8 or D_{12} . Only seven of them have rational invariants. From Lemma 4, we have the following:

Corollary 12. *There are exactly four (resp., three) genus 2 curves \mathcal{C} defined over \mathbb{Q} (up to $\bar{\mathbb{Q}}$ -isomorphism) with a degree 3 elliptic subcover which have automorphism group D_8 (resp., D_{12}). They are listed in Table 1.*

	\mathcal{C}	$\mathfrak{p} = (i_1, i_2, i_3)$	$e_3(\mathfrak{p})$	$\text{Aut}(\mathcal{C})$
\mathfrak{p}_1	$196X^5 + 196X^3 + 81X$	$i_1 = \frac{729}{2116}, i_2 = \frac{1240029}{97336}, i_3 = \frac{531441}{13181630464}$	2	D_8
\mathfrak{p}_2	$49X^5 + 49X^3 + 12X$	$i_1 = \frac{4288}{1849}, i_2 = \frac{243712}{79507}, i_3 = \frac{64}{1323075987}$	1	D_8
\mathfrak{p}_3	$5X^5 + 5X^3 + X$	$i_1 = \frac{144}{49}, i_2 = \frac{3456}{8575}, i_3 = \frac{243}{52521875}$	2	D_8
\mathfrak{p}_4	$700X^5 + 700X^3 - 81X$	$i_1 = -\frac{8019}{20}, i_2 = -\frac{1240029}{200}, i_3 = -\frac{531441}{10000}$	2	D_8
\mathfrak{p}_5	$25X^6 + 25X^3 + 4$	$i_1 = \frac{64}{5}, i_2 = -\frac{1088}{25}, i_3 = -\frac{1}{84375}$	1	D_{12}
\mathfrak{p}_6	$11X^6 + 11X^3 - 4$	$i_1 = \frac{576}{361}, i_2 = \frac{60480}{6859}, i_3 = \frac{243}{2476099}$	1	D_{12}
\mathfrak{p}_7	$20X^6 + 20X^3 + 1$	$i_1 = 81, i_2 = -\frac{5103}{25}, i_3 = -\frac{729}{12500}$	2	D_{12}

TABLE 1. Rational points $\mathfrak{p} \in \mathcal{L}_3$ with $|\text{Aut}(\mathfrak{p})| > 4$

Remark 13. *All points \mathfrak{p} in Table 1 are in the locus $\det(J_\theta) = 0$. We have already seen cases $\mathfrak{p}_1, \mathfrak{p}_4$, and \mathfrak{p}_7 as the exceptional points of $\det(J_\theta) = 0$; see equation (13). The class \mathfrak{p}_3 is a singular point of order 2 of \mathfrak{Y} , \mathfrak{p}_2 is the only point which belong to the degenerate case, and \mathfrak{p}_6 is the only ordinary point in \mathfrak{Y} such that the order of $\text{Aut}(\mathfrak{p})$ is greater than 4.*

5. COMPUTING ELLIPTIC SUBCOVERS

Next we will consider all points \mathfrak{p} in Table 1 and compute j-invariants of their degree 2 and 3 elliptic subcovers. To compute j-invariants of degree 2 elliptic subcovers we use lemma 5 and the values of t from the proof of theorem 1. We recall that for $\mathfrak{p}_1, \dots, \mathfrak{p}_4$ there are four degree 2 elliptic subcovers which are two and two isomorphic. We list the j-invariant of each isomorphic class. They are 2-isogenous as mentioned before. For $\mathfrak{p}_5, \mathfrak{p}_6, \mathfrak{p}_7$ there are two degree 2 elliptic subcovers which are 3-isogenous to each other. To compute degree 3 elliptic subcovers for each \mathfrak{p} we find the pairs (u, v) in the fiber $\theta^{-1}(\mathfrak{p})$ and then use equations (9). We focus on cases which have elliptic subcovers defined over \mathbb{Q} . There are techniques for computing rational points of a genus two curves which have degree 2 subcovers

defined over \mathbb{Q} as in Flynn and Wetherell [5]. Sometimes degree 3 elliptic subcovers are defined over \mathbb{Q} even though degree 2 elliptic subcovers are not; see examples 2 and 6. These degree 3 subcovers help determine rational points of genus 2 curves as illustrated in examples 2, 4, 5, and 6.

Example 1. $\mathfrak{p} = \mathfrak{p}_1$: The j -invariants of degree 3 elliptic subcovers are $j = j' = 66^3$. A genus 2 curve \mathcal{C} corresponding to \mathfrak{p} is

$$\mathcal{C} : Y^2 = X^6 + 3X^4 - 6X^2 - 8.$$

Claim: The equation above has no rational solutions.

Indeed, two of degree 2 elliptic subcovers (isomorphic to each other) have equations

$$\begin{aligned} \mathcal{E}_1 : Y^2 &= x^3 + 3x^2 - 6x - 8 \\ \mathcal{E}_2 : Y^2 &= -8x^3 - 6x^2 + 3x + 1 \end{aligned}$$

where $x = X^2$ (i.e. $\phi : \mathcal{C} \rightarrow \mathcal{E}_1$ of degree 2 such that $\phi(X, Y) = (X^2, Y)$). The elliptic curve \mathcal{E}_1 has rank 0. Thus, the rational points of \mathcal{C} are the preimages of the torsion points of \mathcal{E}_1 . The torsion group of \mathcal{E}_1 has order 4 and is given by

$$\text{Tor}(\mathcal{E}_1) = \{\mathcal{O}, (-1, 0), (2, 0), (-4, 0)\}$$

None of the preimages is rational. Thus, \mathcal{C} has no rational points except the point at infinity.

Example 2. $\mathfrak{p} = \mathfrak{p}_2$: The j -invariants of degree 2 elliptic subcovers are

$$76771008 \pm 44330496\sqrt{3}.$$

The point \mathfrak{p}_2 belongs to the degenerate locus with $\mathfrak{w} = 0$. Thus, the equation of the genus 2 curve \mathcal{C} corresponding to \mathfrak{p} is

$$\mathcal{C} : Y^2 = (3X^2 + 4)(X^3 + X).$$

Indeed, this curve has both pairs $(\mathcal{C}, \mathcal{E})$ and $(\mathcal{C}, \mathcal{E}')$ degenerate pairs. It is the only such genus 2 curve defined over \mathbb{Q} . This fact was noted in [12] and [17]. Both authors failed to identify the automorphism group. The degree 3 coverings are

$$U_1 = \phi_1(X) = X^3 + X, \quad U_2 = \phi_2(X) = \frac{X^3}{3X^2 + 4}$$

and equations of elliptic curves:

$$\mathcal{E} : V_1^2 = 27U_1^3 + 4U_1, \quad \text{and} \quad \mathcal{E}' : V_2^2 = U_2^3 + U_2.$$

\mathcal{E} and \mathcal{E}' are isomorphic with j -invariant 1728. They have rank 0 and rational torsion group of order 2, $\text{Tor}(\mathcal{E}) = \{\mathcal{O}, (0, 0)\}$. Thus, the only rational points of \mathcal{C} are in fibers $\phi_1^{-1}(0)$ and $\phi_2^{-1}(\infty)$. Hence, $\mathcal{C}(\mathbb{Q}) = \{(0, 0), \infty\}$.

Example 3. $\mathfrak{p} = \mathfrak{p}_3$: All degree 2 and 3 elliptic subcovers are defined over $\mathbb{Q}(\sqrt{5})$.

Example 4. $\mathfrak{p} = \mathfrak{p}_4$: Degree 2 elliptic subcovers have j -invariants

$$\frac{1728000}{2809} \pm \frac{17496000}{2809}\sqrt{I}$$

where $I^2 = -1$. Thus, we can't recover any information from degree 2 subcovers. One corresponding value for $(\mathfrak{u}, \mathfrak{v})$ is $(\frac{25}{2}, \frac{250}{9})$. Then \mathcal{C} is

$$\mathcal{C} : 3^8 \cdot Y^2 = (100X + 9)(2500X^2 + 400X + 9)(25X + 9)(2500X^2 + 225X + 9)$$

Degree 3 elliptic subcovers are have equations

$$(17) \quad \begin{aligned} \mathcal{E} : V_1^2 &= -\frac{1}{81}(10U_1 - 3)(8575U_1^2 - 2940U_1 + 108) \\ \mathcal{E}' : V_2^2 &= -\frac{686}{59049}(1700U_2 - 441)(1445000U_2^2 - 696150U_2 + 83853) \end{aligned}$$

where

$$(18) \quad \begin{aligned} U_1 = \phi_1(X) &= 2250 \frac{X^2}{(25X + 9)(2500X^2 + 400X + 9)} \\ U_2 = \phi_2(X) &= \frac{1}{340} \frac{(250X + 27)^2(340X + 9)}{(100X + 9)(2500X^2 + 400X + 9)} \end{aligned}$$

The curve \mathcal{E}' has rank 0 and torsion group

$$\text{Tor}(\mathcal{E}') = \{\mathcal{O}, (\frac{441}{1700}, 0)\}.$$

Thus, the rational points of \mathcal{C} are in $\phi_2^{-1}(\frac{441}{1700})$ and $\phi_2^{-1}(\infty)$. Thus,

$$\mathcal{C}(\mathbb{Q}) = \{(-\frac{9}{100}, 0), (-\frac{9}{25}, 0)\}.$$

Example 5. $\mathfrak{p} = \mathfrak{p}_5$: Degree 2 j -invariants are $j_1 = 0$, $j_2 = -1228800$ and degree 3 j -invariants as shown below are $j = j' = 0$. Let \mathcal{C} be genus 2 curve with equation

$$\mathcal{C} : Y^2 = (X^3 + 1)(4X^3 + 1)$$

corresponding to \mathfrak{p} . The case is treated separately in [18]. Degree 3 elliptic subcovers have equations

$$\mathcal{E} : V_1^2 = -27U_1^3 + 4, \quad \mathcal{E}' : V_2^2 = -16(27U_2^3 - 1)$$

where

$$U_1 = \phi_1(X) = \frac{X^2}{X^3 + 1}, \quad U_2 = \phi_2(X) = \frac{X}{4X^3 + 1}.$$

The rank of both \mathcal{E} and \mathcal{E}' is zero. Thus, the rational points of \mathcal{C} are the preimages of the rational torsion points of \mathcal{E} and \mathcal{E}' . The torsion points of \mathcal{E} are $\text{Tor}(\mathcal{E}) = \{\mathcal{O}, (0, 2), (0, -2)\}$. Then $\phi_1^{-1}(0) = \{0, \infty\}$ and $\phi_1^{-1}(\infty) = \{-1, \frac{1}{2} \pm \frac{\sqrt{-3}}{2}\}$. Thus,

$$\mathcal{C}(\mathbb{Q}) = \{(0, 1), (0, -1), (-1, 0)\}$$

Example 6. $\mathfrak{p} = \mathfrak{p}_6$: This point is in \mathfrak{M} and it is not a singular point of \mathfrak{M} . It has isomorphic degree 3 elliptic subcovers; see [18]. The corresponding (\mathbf{u}, \mathbf{v}) pair is $(\mathbf{u}, \mathbf{v}) = (20, 16)$ and $e_3(\mathfrak{p}) = 1$. Then, the genus 2 curve has equation:

$$\mathcal{C} : Y^2 = (256X^3 + 320X^2 + 16X + 1)(1024X^3 + 256X^2 + 32X + 1)$$

Degree 3 elliptic subcovers have j -invariants $j = j' = -32768$ and equations

$$(19) \quad \begin{aligned} \mathcal{E} : V_1^2 &= 4(-5324U_1^3 + 968U_1^2 - 56U_1^2 + 1) \\ \mathcal{E}' : V_2^2 &= 11^3(-32000U_2^3 + 35200U_2^2 - 12320U_2 + 11^3) \end{aligned}$$

where

$$U_1 = 16 \frac{X^2}{256X^3 + 320X^2 + 16X + 1}, \quad U_2 = \frac{1}{20} \frac{(16X + 3)^2(20X + 1)}{1024X^3 + 256X^2 + 32X + 1}$$

Both elliptic curves have trivial torsion but rank > 0 . One can try to adapt more sophisticated techniques in this case as Flynn and Wetherell have done for degree 2 subcovers. This is the only genus 2 curve (up to \mathbb{C} -isomorphism) with automorphism

group D_{12} and isomorphic degree 2 elliptic subcovers. Indeed all degree 2 and 3 elliptic subcovers are \mathbb{C} -isomorphic with j -invariants -32768 . Degree 2 elliptic subcovers also have rank 1 which does not provide any quick information about rational points of \mathcal{C} .

Example 7. $\mathfrak{p} = \mathfrak{p}_7$: All degree 2 and 3 elliptic subcovers are defined over $\mathbb{Q}(\sqrt{5})$.

Throughout this paper we have made use of several computer algebra packages as APECS, MAPLE, and GAP. The interested reader can check [19] and [18] for more details on loci \mathcal{L}_2 and \mathcal{L}_3 . The equations for these spaces, j -invariants of elliptic subcovers of degree 2 and 3, and other computational aspects of genus 2 curves can be down loaded from author's web site.

REFERENCES

- [1] O. BOLZA, On binary sextics with linear transformations into themselves. *Amer. J. Math.* **10**, 47-70.
- [2] J. CASSELS AND V. FLYNN, Prolegomena to a Middlebrow Arithmetic of Curves of Genus Two, LMS, 230, 1996.
- [3] A. CLEBSCH, Theorie der Binären Algebraischen Formen, Verlag von B.G. Teubner, Leipzig, (1872).
- [4] T. EKEDAHL, J. P. SERRE, Exemples de courbes algébriques jacobienne complètement décomposable. *C. R. Acad. Sci. Paris Sr. I Math.*, 317 (1993), no. 5, 509–513.
- [5] V. FLYNN AND J. WETHERELL, Finding rational points on bielliptic genus 2 curves, *Manuscripta Math.* **100**, 519-533 (1999).
- [6] G. FREY, On elliptic curves with isomorphic torsion structures and corresponding curves of genus 2. *Elliptic curves, modular forms, and Fermat's last theorem (Hong Kong, 1993)*, 79-98, Ser. Number Theory, I, *Internat. Press, Cambridge, MA*, (1995).
- [7] G. FREY AND E. KANI, Curves of genus 2 covering elliptic curves and an arithmetic application. *Arithmetic algebraic geometry (Texel, 1989)*, 153-176, *Progr. Math.*, **89**, Birkhäuser Boston, Boston, MA, (1991).
- [8] E. HOWE, F. LEPRÉVOST, AND B. POONEN, Large torsion subgroups of split Jacobians of curves of genus two or three. *Forum. Math.* **12** (2000), no. 3, 315-364.
- [9] J. IGUSA, Arithmetic Variety Moduli for genus 2. *Ann. of Math. (2)*, **72**, 612-649, (1960).
- [10] W. KELLER, L. KULESZ, Courbes algébriques de genre 2 et 3 possédant de nombreux points rationnels. *C. R. Acad. Sci. Paris Sr. I Math.* **321** (1995), no. 11, 1469–1472.
- [11] A. KRAZER, Lehrbuch der Thetafunctionen, Chelsea, New York, (1970).
- [12] M. R. KUHN, Curves of genus 2 with split Jacobian. *Trans. Amer. Math. Soc* **307** (1988), 41-49
- [13] H. LANGE, Über die Modulvarietät der Kurven vom Geschlecht 2. *J. Reine Angew. Math.*, **281**, 80-96, 1976.
- [14] K. MAGAARD, T. SHASKA, S. SHPECTOROV, AND H. VÖLKLEIN, The automorphism group of a Riemann surface, *Proceedings of the Conference on Communications in Arithmetic Fundamental Groups and Galois Theory*, Kyoto University, Springer, 2001 (to appear).
- [15] P. MESTRE, Construction de courbes de genre 2 à partir de leurs modules. In T. Mora and C. Traverso, editors, *Effective methods in algebraic geometry*, volume 94. *Prog. Math.*, 313-334. Birkhäuser, 1991. Proc. Congress in Livorno, Italy, April 17-21, (1990).
- [16] D. MUMFORD, The Red Book of Varieties and Schemes, Springer, 1999.
- [17] T. SHASKA, Genus 2 curves with (n,n)-decomposable Jacobians, *Jour. Symb. Comp.*, Vol 31, **no. 5**, pg. 603-617, 2001.
- [18] T. SHASKA, Genus 2 fields with degree 3 elliptic subfields, (submitted for publication).
- [19] T. SHASKA AND H. VÖLKLEIN, Elliptic Subfields and automorphisms of genus 2 function fields. *Proceeding of the Conference on Algebra and Algebraic Geometry with Applications: The celebration of the seventieth birthday of Professor S.S. Abhyankar*, Springer-Verlag, 2001.

UNIVERSITY OF CALIFORNIA AT IRVINE, IRVINE, CA 92697.
E-mail address: tshaska@math.uci.edu