

# HYPERELLIPTIC JACOBIANS WITHOUT COMPLEX MULTIPLICATION, DOUBLY TRANSITIVE PERMUTATION GROUPS AND PROJECTIVE REPRESENTATIONS

YURI G. ZARHIN

## 1. INTRODUCTION

Let  $K$  be a field of characteristic zero,  $K_a$  its algebraic closure,  $n \geq 5$  an integer,  $f(x)$  an irreducible polynomial over  $K$  of degree  $n$ , whose Galois group  $\text{Gal}(f)$  acts doubly transitively on the set  $\mathfrak{R}$  of roots of  $f$ . Let  $C : y^2 = f(x)$  be the corresponding hyperelliptic curve and  $X=J(C)$  its jacobian defined over  $K$ . Earlier, the author [20], [21], [25] has proven that the ring  $\text{End}(X)$  of all  $K_a$ -endomorphisms coincides with  $\mathbf{Z}$  if  $\text{Gal}(f)$  is either the full symmetric group  $\mathbf{S}_n$  or the corresponding alternating group  $\mathbf{A}_n$  or a *small* Mathieu group  $M_n$  (with  $n = 11$  or  $12$ ) or  $\mathfrak{R}$  could be identified with the projective space  $\mathbf{P}^{m-1}(\mathbf{F}_q)$  over a finite field  $\mathbf{F}_q$  of *odd* characteristic in such a way that  $\text{Gal}(f)$  contains the projective special linear group  $\text{PSL}_m(\mathbf{F}_q)$  while  $m \geq 3$  and  $(m, q) \neq (4, 3)$ . (Similar results were obtained when  $\text{Gal}(f) = \mathbf{L}_2(2^r)$ ,  $\text{Sz}(2^{2r+1})$  or  $\text{U}_3(2^r)$  [21], [24].) The proof was based on an observation that in all these cases the natural (faithful) representation of  $\text{Gal}(f)$  in  $X_2$  is *very simple*; in particular, it is *absolutely irreducible*. (See [21], [22], [23] for the definition and basic properties of very simple representations.)

We refer the reader to [9], [10], [4], [5], [8], [20], [21], [22], [24], [25] for a discussion of known results about, and examples of, hyperelliptic jacobians without complex multiplication.

In the present paper we suggest a new approach to already known examples when  $\text{End}(X) = \mathbf{Z}$  from [20], [21] and [25]. Namely, instead of very simplicity, we use a theorem of Feit-Tits [2] complemented by results of Kleidman-Liebeck [6]. Besides obtaining new proofs of already known results, we get new examples when  $\text{End}(X) = \mathbf{Z}$  with *reducible* (but still absolutely indecomposable)  $\text{Gal}(f)$ -module  $X_2$ . Namely, we prove that  $\text{End}(X) = \mathbf{Z}$  when  $\text{Gal}(f)$  is a *big* Mathieu group (with  $n = 22, 23$  or  $24$ ) or  $\mathfrak{R}$  could be identified with the projective space  $\mathbf{P}^{m-1}(\mathbf{F}_q)$  over a finite field  $\mathbf{F}_q$  of characteristic 2 in such a way that  $\text{Gal}(f)$  becomes either the projective special linear group  $\mathbf{L}_m(q) := \text{PSL}_m(\mathbf{F}_q)$  or the projective linear group  $\text{PGL}_m(\mathbf{F}_q)$  with  $m > 2$  (except  $(m, q) = (4, 2), (3, 4)$ ). We refer to Theorem 5.1 (and Definitions 3.1 and 3.6) for a justification for the long title of the present article.

The paper is organized as follows. In §2 we state the main results. Section 3 contains auxiliary results from representation theory. In §4 we discuss fields of definitions for endomorphisms of abelian varieties. Section 5 contains proofs of the main results.

---

Partially supported by the NSF.

## 2. MAIN RESULTS

Throughout this paper we assume that  $K$  is a field of characteristic 0. We fix its algebraic closure  $K_a$  and write  $\text{Gal}(K)$  for the absolute Galois group  $\text{Aut}(K_a/K)$ .

**Theorem 2.1.** *Let  $K$  be a field of characteristic zero,  $K_a$  its algebraic closure,  $f(x) \in K[x]$  an irreducible polynomial of degree  $n$ . Suppose  $n = 11, 12, 22, 23$  or  $24$  and the Galois group of  $f$  is the corresponding Mathieu group  $M_n$ . Let  $C_f$  be the hyperelliptic curve  $y^2 = f(x)$ . Let  $J(C_f)$  be its jacobian,  $\text{End}(J(C_f))$  the ring of  $K_a$ -endomorphisms of  $J(C_f)$ . Then  $\text{End}(J(C_f)) = \mathbf{Z}$ .*

**Remark 2.2.** The case of small Mathieu groups was done in Th. 7.13 on p. 489 of [21] (see also [25]). However, in this paper we give a unified proof for all Mathieu groups.

**Theorem 2.3.** *Let  $K$  be a field of characteristic zero,  $K_a$  its algebraic closure,  $f(x) \in K[x]$  an irreducible polynomial of degree  $n$  and  $\mathfrak{R} \subset K_a$  the set of its roots. We write  $\text{Gal}(f)$  for the Galois group of  $f$ . Suppose there exist integers  $m > 2$  and  $r \geq 1$  such that  $n = \frac{q^m - 1}{q - 1}$  where  $q = 2^r$  and  $(m, q) \neq (2, 2), (4, 2), (3, 4)$ . Assume, in addition, that  $\mathfrak{R}$  could be identified with the projective space  $\mathbf{P}^{m-1}(\mathbf{F}_q)$  over the finite field  $\mathbf{F}_q$  in such a way that  $\text{Gal}(f)$  contains  $\mathbf{L}_m(q) := \text{PSL}_m(\mathbf{F}_q)$  as a subgroup. (E.g.,  $\text{Gal}(f) = \mathbf{L}_m(q)$  or  $\text{PGL}_m(\mathbf{F}_q)$ .) Let  $C_f$  be the hyperelliptic curve  $y^2 = f(x)$ . Let  $J(C_f)$  be its jacobian,  $\text{End}(J(C_f))$  the ring of  $K_a$ -endomorphisms of  $J(C_f)$ . Then  $\text{End}(J(C_f)) = \mathbf{Z}$ .*

**Remark 2.4.** In the case of  $m = 2$  this assertion is proven in [21].

**Theorem 2.5.** *Let  $K$  be a field of characteristic zero,  $K_a$  its algebraic closure,  $f(x) \in K[x]$  an irreducible polynomial of degree  $n$  and  $\mathfrak{R} \subset K_a$  the set of its roots. We write  $\text{Gal}(f)$  for the Galois group of  $f$ . Assume that there exist an odd power prime  $q \geq 3$  and an integer  $m \geq 3$  such that  $n = \frac{q^m - 1}{q - 1}$  and the set  $\mathfrak{R}$  could be identified with the projective space  $\mathbf{P}^{m-1}(\mathbf{F}_q)$  over the prime field  $\mathbf{F}_q$  in such a way that  $\text{Gal}(f)$  contains  $\mathbf{L}_m(q) := \text{PSL}_m(\mathbf{F}_q)$  as a subgroup. (E.g.,  $\text{Gal}(f) = \mathbf{L}_m(q)$  or  $\text{PGL}_m(\mathbf{F}_q)$ .) Let  $C_f$  be the hyperelliptic curve  $y^2 = f(x)$ . Let  $J(C_f)$  be its jacobian,  $\text{End}(J(C_f))$  the ring of  $K_a$ -endomorphisms of  $J(C_f)$ . Then  $\text{End}(J(C_f)) = \mathbf{Z}$ .*

**Remark 2.6.** When  $(m, q) \neq (4, 3)$  this assertion is already proven in [25]. However, in this paper we give a unified proof for all  $(m, q)$ .

## 3. GROUP THEORY

**Definition 3.1.** Suppose  $\mathcal{G} \neq \{1\}$  is a perfect finite group and  $g > 1$  is an integer. We say that  $\mathcal{G}$  is  $g$ -unbounded if it enjoys the following properties:

- (i) Each homomorphism  $\mathcal{G} \rightarrow \text{PSL}(g - 1, \mathbf{C})$  is trivial;
- (ii) Either  $g$  is odd and each homomorphism  $\mathcal{G} \rightarrow \text{PSL}(g, \mathbf{Q}) = \text{SL}(g, \mathbf{Q})$  is trivial or  $g$  is even and each homomorphism  $\mathcal{G} \rightarrow \text{PSL}(g, \mathbf{R}) = \text{SL}(g, \mathbf{R})/\{\pm 1\}$  is trivial.

**Remark 3.2.** (i) Clearly, if every nontrivial irreducible projective representation of  $\mathcal{G}$  in characteristic zero has dimension  $> g$  then all nontrivial projective representations of  $\mathcal{G}$  in characteristic zero have dimension  $> g$  and therefore  $\mathcal{G}$  is  $g$ -unbounded.

- (ii) Clearly, if  $\mathcal{H}$  is a simple non-abelian group isomorphic to a subgroup of  $\mathcal{G}$  then the  $g$ -unboundness of  $\mathcal{H}$  implies the  $g$ -unboundness of  $\mathcal{G}$ .
- (iii) Clearly, if every nontrivial projective representation of  $\mathcal{G}$  in characteristic 2 has dimension  $> g$  then all nontrivial projective representations of  $\mathcal{G}$  in characteristic zero have dimension  $> g$  and therefore  $\mathcal{G}$  is  $g$ -unbounded.
- (iv) Every  $\mathcal{G}$  is 2-unbounded. Indeed, it suffices to check that each homomorphism from  $\mathcal{G}$  to  $\mathrm{PSL}(2, \mathbf{R})$  is trivial. But each finite subgroup in  $\mathrm{PSL}(2, \mathbf{R})$  is the image of a finite subgroup in  $\mathrm{SL}(2, \mathbf{R})$ . Since each finite subgroup of  $\mathrm{SL}(2, \mathbf{R})$  is commutative, the image  $\pi(\mathcal{G})$  of each homomorphism  $\pi : \mathcal{G} \rightarrow \mathrm{PSL}(2, \mathbf{R})$  is commutative. The perfectness of  $\mathcal{G}$  implies that  $\pi(\mathcal{G})$  is also perfect. This implies that  $\pi(\mathcal{G}) = \{1\}$ , i.e.,  $\pi$  is trivial.

**Example 3.3.** Suppose  $g \geq 4$  is an integer. Then the alternating groups  $\mathbf{A}_{2g+1}$  and  $\mathbf{A}_{2g+2}$  are  $g$ -bounded. Indeed,  $2g + 1 > 8$  and, by a theorem of Wagner [19], every nontrivial linear representation of  $\mathbf{A}_{2g+1}$  in characteristic 2 has dimension  $\geq 2g$ . Since the Schur multiplier of  $\mathbf{A}_{2g+1}$  is 2, one may easily deduce that every nontrivial projective representation of  $\mathbf{A}_{2g+1}$  in characteristic 2 also has dimension  $\geq 2g > g$ . By Remark 3.2((iii),  $\mathbf{A}_{2g+1}$  is  $g$ -unbounded. Since  $\mathbf{A}_{2g+1}$  is isomorphic to a subgroup of  $\mathbf{A}_{2g+2}$ , the group  $\mathbf{A}_{2g+2}$  is also  $g$ -unbounded.

**Examples 3.4.** (a) The Mathieu groups  $M_{23}$  and  $M_{24}$  are 11-unbounded. Indeed, it follows from the Tables in [1] that all nontrivial irreducible projective representations of  $M_{23}$  in characteristic zero have dimension  $\geq 22$ . This implies that  $M_{23}$  is  $d$ -unbounded for all  $d < 22$ . Since  $M_{23}$  is isomorphic to a subgroup of  $M_{24}$ , the group  $M_{24}$  is also  $d$ -unbounded for all  $d < 22$ .

(b) The Mathieu group  $M_{22}$  is 10-unbounded. Indeed, it follows from the Tables in [1] that in characteristic zero all nontrivial irreducible projective representations of  $M_{22}$  have dimension  $\geq 10$  and there are no nontrivial irreducible *linear* 10-dimensional representations. This implies that all homomorphisms  $M_{22} \rightarrow \mathrm{PGL}(9, \mathbf{C})$  are trivial. Now in order to establish the 10-unboundness of  $M_{22}$  we need only to check that each homomorphism from  $M_{22}$  to  $\mathrm{PSL}(10, \mathbf{R})$  is trivial. Let us assume that  $\rho : M_{22} \rightarrow \mathrm{PSL}(10, \mathbf{R}) = \mathrm{SL}(10, \mathbf{R})/\{\pm 1\}$  is a nontrivial group homomorphism. Clearly,  $\rho$  lifts to a nontrivial homomorphism

$$\rho' : M'_{22} \rightarrow \mathrm{SL}(10, \mathbf{R})$$

where  $M'_{22}$  sits in a central extension

$$\{1\} \rightarrow \{\pm 1\} \hookrightarrow M'_{22} \twoheadrightarrow M_{22} \rightarrow \{1\}.$$

This extension is non-splittable, since there are no nontrivial homomorphisms from  $M_{22}$  to  $\mathrm{SL}(10, \mathbf{R})$  [1]. On the other hand, it also follows from the Tables in [1] that there are no 10-dimensional linear irreducible representations of  $M'_{22}$  defined over  $\mathbf{R}$ . This implies easily that there are no nontrivial 10-dimensional linear representations of  $M'_{22}$  defined over  $\mathbf{R}$ . This gives us the desired contradiction and proves the 10-unboundness of  $M_{22}$ .

- (c) Suppose  $r \geq 1$  and  $m \geq 2$  are positive integers and assume that  $(m, r) \neq (2, 1), (2, 2), (3, 2), (4, 2)$ . Let us put  $q = 2^r$ . We define a positive integer  $g$  as follows.

$$g = q/2 \text{ if } m = 2 \text{ and } g = \frac{1}{2} \frac{q^m - q}{q - 1} \text{ if } m > 2.$$

Then the group  $\mathbf{L}_m(q) = \mathrm{PSL}_m(\mathbf{F}_q)$  is  $g$ -unbounded. Indeed, it is known [18] that under our assumptions on  $(m, q)$  all nontrivial irreducible projective representations of  $\mathbf{L}_m(q)$  in characteristic zero have dimension  $\geq \frac{q^m - q}{q - 1} > g$  if  $m > 2$ . It is also known [18] that all nontrivial irreducible projective representations of  $\mathbf{L}_2(q)$  in characteristic zero have dimension  $\geq q - 1 > g$  if  $q > 4$ .

(d) Suppose  $q \geq 3$  is an odd power prime,  $m \geq 3$  is an integer. Let us put

$$g := \left\lfloor \frac{1}{2} \left( \frac{q^m - 1}{q - 1} - 1 \right) \right\rfloor = \left\lfloor \frac{1}{2} \left( \frac{q^m - q}{q - 1} \right) \right\rfloor.$$

Then the group  $\mathbf{L}_m(q) = \mathrm{PSL}_m(\mathbf{F}_q)$  is  $g$ -bounded. Indeed, it is known [18] that if  $(q, m) \neq (3, 4)$  then all nontrivial irreducible projective representations of  $\mathbf{L}_m(q)$  in characteristic zero have dimension  $\geq \frac{q^m - 1}{q - 1} - 1 > g$ .

If  $(q, m) = (3, 4)$  then  $g = 19$ . It follows from the Tables in [1] that all nontrivial irreducible projective representations of  $\mathbf{L}_4(3)$  in characteristic zero have dimension  $\geq 26 > 19$ .

**Example 3.5.** The Mathieu groups  $M_{11}$  and  $M_{12}$  are 5-unbounded. Indeed, it follows from the Tables in [1] that all nontrivial irreducible projective representations of  $M_{11}$  in characteristic zero have dimension  $\geq 10$ . This implies that  $M_{11}$  is  $d$ -unbounded for all  $d < 10$ . Since  $M_{11}$  is isomorphic to a subgroup of  $M_{12}$ , the group  $M_{12}$  is also  $d$ -unbounded for all  $d < 10$ .

**Definition 3.6.** Suppose  $\mathcal{G}$  is a simple non-abelian group,  $\ell$  is a prime. Suppose

$$\{1\} \rightarrow N \hookrightarrow \mathcal{G}' \xrightarrow{\gamma} \mathcal{G} \rightarrow \{1\}$$

is a short exact sequence of finite groups where  $N$  is a group of exponent 1 or  $\ell$  and no proper subgroup of  $\mathcal{G}'$  maps onto  $\mathcal{G}$ . (In particular,  $\mathcal{G}'$  is perfect.) Then  $\mathcal{G}'$  is called a *minimal cover* of  $\mathcal{G}$ .

In addition, if either  $N = \{1\}$  (i.e.,  $\mathcal{G}' = \mathcal{G}$ ) or the exponent of  $N$  is  $\ell$  then we say that  $\mathcal{G}'$  is a *minimal  $\ell$ -cover* of  $\mathcal{G}$ .

**Remark 3.7.** Clearly, the minimal cover  $\mathcal{G}'$  is always perfect. It is also clear that each normal subgroup in  $\mathcal{G}'$  except  $\mathcal{G}'$  itself lies in  $N$ . This implies easily that if  $\rho : \mathcal{G}' \rightarrow M$  is a nontrivial group homomorphism then  $\ker(\rho)$  lies in  $N$  and therefore the image  $\rho(\mathcal{G}')$  is also a minimal cover of  $\mathcal{G}$ . In addition, if  $\mathcal{G}'$  is a minimal  $\ell$ -cover then  $\rho(\mathcal{G}')$  is also one.

**Remark 3.8.** Suppose  $\mathcal{G}$  is a simple non-abelian group and  $\gamma : H \twoheadrightarrow \mathcal{G}$  is a surjective homomorphism of finite groups. Let  $\mathcal{G}'$  be a subgroup of smallest order among the subgroups  $H'$  of  $H$  such that  $\gamma(H') = \mathcal{G}$ . (Clearly, such  $\mathcal{G}'$  always exists.) Then  $\gamma : \mathcal{G}' \rightarrow \mathcal{G}$  is a minimal cover. In addition, if the kernel of  $\gamma$  is either trivial or has exponent  $\ell$  then  $\gamma : \mathcal{G}' \rightarrow \mathcal{G}$  is a minimal  $\ell$ -cover.

**Examples 3.9.** (i) Suppose  $\mathcal{G}$  is a simple non-abelian group isomorphic either to  $\mathbf{A}_5 \cong \mathbf{L}_2(4)$  or to  $\mathbf{A}_6$ . Suppose  $\mathcal{G}'$  is a minimal cover of  $\mathcal{G}$ . Then, by Remark 3.2(iv), the perfect group  $\mathcal{G}'$  is 2-bounded.

- (ii) Suppose  $\mathcal{G}$  is a simple non-abelian group isomorphic either to  $\mathbf{L}_3(2) \cong \mathbf{L}_2(7)$  or to  $\mathbf{A}_7$  or to  $\mathbf{A}_8 \cong \mathbf{L}_4(2)$ . Notice that in all these cases the order of  $\mathcal{G}$  is divisible by 7.

Suppose  $\mathcal{G}'$  is a minimal cover of  $\mathcal{G}$ . Then  $\mathcal{G}'$  is 3-bounded. Indeed, if  $\pi_2 : \mathcal{G}' \rightarrow \mathrm{PSL}(2, \mathbf{C})$  is a *nontrivial* homomorphism then, by Remark 3.7, its image  $H' := \pi(\mathcal{G}') \subset \mathrm{PSL}(2, \mathbf{C})$  is also a minimal cover of  $\mathcal{G}$ . In particular,  $H_2$  is a perfect finite group having a quotient isomorphic to  $\mathcal{G}$ . This implies that nonsolvable  $H_2 \subset \mathrm{PSL}(2, \mathbf{C})$  is not isomorphic to  $\mathbf{A}_5$  which could not be true ([17], Th. 6.17 on p. 404). The obtained contradiction implies that there are no nontrivial homomorphisms from  $\mathcal{G}'$  to  $\mathrm{PSL}(2, \mathbf{C})$ .

Now assume that there exists a *nontrivial* homomorphism  $\pi_3 : \mathcal{G}' \rightarrow \mathrm{SL}(3, \mathbf{Q})$ . As above, the image  $H_3 := \pi(\mathcal{G}') \subset \mathrm{SL}(3, \mathbf{Q})$  is also a minimal cover of  $\mathcal{G}$ . In particular,  $H_3$  is a finite group having a quotient isomorphic to  $\mathcal{G}$ ; in particular, 7 divides the order of  $H_3$  and therefore  $H_3 \subset \mathrm{SL}(3, \mathbf{Q})$  contains an element of order 7. But this is not true, since the degree of the 7th cyclotomic field over  $\mathbf{Q}$  is  $6 > 3$ . The obtained contradiction ends the proof of the 3-unboundness of  $\mathcal{G}'$ .

We will use the following result of Feit-Tits ([2], Theorem on pp. 1092–1093 and Prop. 4.1 on p. 1098) concerning complex projective representations of minimum degree. (See also [6].)

**Theorem 3.10** (Feit-Tits Theorem). *Suppose  $\mathcal{G}$  is a known simple non-abelian group that is not a group of Lie type in characteristic 2. Suppose  $\mathcal{G}' \xrightarrow{\gamma} \mathcal{G}$  is a minimal cover of  $\mathcal{G}$  and  $d$  is the smallest positive integer such that there exists a nontrivial homomorphism  $\mathcal{G}' \rightarrow \mathrm{PGL}(d, \mathbf{C})$ . Then the kernel of each homomorphism  $\mathcal{G}' \rightarrow \mathrm{PGL}(d, \mathbf{C})$  contains  $\ker(\gamma)$ . In particular,  $\mathcal{G}$  is isomorphic to a subgroup of  $\mathrm{PGL}(d, \mathbf{C})$ .*

**Remark 3.11.** Kleidman and Liebeck [6] studied the case of simple groups of Lie type in characteristic 2. In particular, they proved the following assertion (ibid, Th. 3 on p. 316). Suppose  $q$  is a power of 2 and  $m \geq 2$  is an integer such that  $(m, q) \neq (2, 2)$  (i.e.,  $\mathbf{L}_m(q)$  is a simple non-abelian group). If  $g$  is a positive integer and  $\mathcal{G}'$  is a minimal cover of  $\mathbf{L}_m(q)$  such that  $\mathcal{G}'$  is isomorphic to a subgroup of  $\mathrm{PGL}(g, \mathbf{C})$  then either  $m = 4$  and  $g \geq q^3$  or  $\mathbf{L}_m(q)$  is isomorphic to a subgroup of  $\mathrm{PGL}(g, \mathbf{C})$ .

**Lemma 3.12.** *Suppose  $q \geq 2$  is an integral power of 2 and  $m \geq 2$  is an integer such that  $(m, q) \neq (2, 2), (2, 4), (3, 2), (4, 2)$ . Suppose  $\mathcal{G}'$  is a minimal cover of  $\mathbf{L}_m(q)$ .*

- (i) *If  $m > 2$  then  $\mathcal{G}'$  is  $\frac{1}{2} \frac{q^m - q}{q - 1}$ -unbounded.*
- (ii) *If  $m > 2$  then  $\mathcal{G}'$  is  $\frac{q}{2}$ -unbounded.*

*Proof.* Let us start with the case of  $m = 2$ . Let us put  $g = \frac{q}{2}$ . Let  $\rho : \mathcal{G}' \rightarrow \mathrm{PGL}(g, \mathbf{C})$  is a *nontrivial* group homomorphism. By Remark 3.7, the image  $\rho(\mathcal{G}') \subset \mathrm{PGL}(g, \mathbf{C})$  is also a minimal cover of  $\mathbf{L}_2(q)$ . Applying Remark 3.11 to  $\rho(\mathcal{G}')$ , we conclude that  $\mathbf{L}_2(q)$  is isomorphic to a subgroup of  $\mathrm{PGL}(g, \mathbf{C})$ . By Example 3.4(c), this is not true. The obtained contradiction implies that there are no nontrivial homomorphisms from  $\mathcal{G}'$  to  $\mathrm{PGL}(g, \mathbf{C})$ .

Now assume that  $m > 2$ . Let us put  $g = \frac{1}{2} \frac{q^m - q}{q - 1}$ . Notice that if  $m = 4$  then

$$g = \frac{1}{2}(q^3 + q^2 + q) < q^3.$$

Let  $\rho : \mathcal{G}' \rightarrow \mathrm{PGL}(g, \mathbf{C})$  is a *nontrivial* group homomorphism. Again, the image  $\rho(\mathcal{G}') \subset \mathrm{PGL}(g, \mathbf{C})$  is also a minimal cover of  $\mathbf{L}_m(q)$ . Applying Remark 3.11 to  $\rho(\mathcal{G}')$ , we conclude that either  $m = 4$  and  $g \geq q^3$  or  $\mathbf{L}_m(q)$  is isomorphic to a subgroup of  $\mathrm{PGL}(g, \mathbf{C})$ . But we have already seen that if  $m = 4$  then  $g < q^3$ . This implies that  $\mathbf{L}_m(q)$  is isomorphic to a subgroup of  $\mathrm{PGL}(g, \mathbf{C})$ . By Example 3.4(c), this is not true. The obtained contradiction implies that there are no nontrivial homomorphisms from  $\mathcal{G}'$  to  $\mathrm{PGL}(g, \mathbf{C})$ .  $\square$

**Lemma 3.13.** *Suppose a simple non-abelian finite group  $\mathcal{G}$  and an integer  $g > 1$  enjoy one of the following properties:*

- (a)  $g = 11$  and  $\mathcal{G} = \mathrm{M}_{23}$  or  $\mathrm{M}_{24}$ ;
- (b)  $g = 10$  and  $\mathcal{G} = \mathrm{M}_{22}$ ;
- (c)  $g = 5$  and  $\mathcal{G} = \mathrm{M}_{11}$  or  $\mathrm{M}_{12}$ ;
- (d)  $g = \left\lfloor \frac{1}{2} \left( \frac{q^m - 1}{q - 1} - 1 \right) \right\rfloor$  where  $q \geq 3$  is an odd power prime,  $m \geq 3$  is an integer and  $\mathcal{G} = \mathbf{L}_m(q)$ ;
- (e)  $\mathcal{G} = \mathbf{A}_{2g+1}$  or  $\mathbf{A}_{2g+2}$ ;

*If  $\mathcal{G}'$  is a minimal cover of  $\mathcal{G}$  then  $\mathcal{G}'$  is  $g$ -unbounded.*

*Proof.* Case (a). We have seen that there are no nontrivial homomorphisms to  $\mathrm{PSL}(11, \mathbf{C})$  either from  $\mathrm{M}_{23}$  or  $\mathrm{M}_{24}$ . It follows from the Feit–Tits theorem that the same is true for the minimal cover  $\mathcal{G}'$ .

Case (c). We have seen that there are no nontrivial homomorphisms to  $\mathrm{PSL}(5, \mathbf{C})$  either from  $\mathrm{M}_{11}$  or  $\mathrm{M}_{12}$ . It follows from the Feit–Tits theorem that the same is true for the minimal cover  $\mathcal{G}'$ .

Case (d). It follows from the Feit–Tits theorem combined with Example 3.4(d) that there are no nontrivial homomorphisms from  $\mathcal{G}'$  to  $\mathrm{PSL}(g, \mathbf{C})$ .

Case (b). Since every homomorphism from  $\mathrm{M}_{22}$  to  $\mathrm{PSL}(9, \mathbf{C})$  is trivial, we conclude that, thanks to the Feit–Tits theorem, that every homomorphism from  $\mathcal{G}'$  to  $\mathrm{PSL}(9, \mathbf{C})$  is also trivial. In order to finish the proof we have to check that every homomorphism from  $\mathcal{G}'$  to  $\mathrm{PSL}(10, \mathbf{R})$  is also trivial. Let us assume that  $\rho' : \mathcal{G}' \rightarrow \mathrm{PSL}(10, \mathbf{R}) = \mathrm{SL}(10, \mathbf{R})/\{\pm 1\}$  is a nontrivial group homomorphism. Let us consider the composition

$$\pi : \mathcal{G}' \xrightarrow{\rho'} \mathrm{PSL}(10, \mathbf{R}) \subset \mathrm{PSL}(10, \mathbf{C})$$

of  $\rho'$  and the natural embedding  $\mathrm{PSL}(10, \mathbf{R}) \subset \mathrm{PSL}(10, \mathbf{C})$ . Clearly, the composition  $\pi : \mathcal{G}' \rightarrow \mathrm{PSL}(10, \mathbf{C})$  is a *nontrivial* group homomorphism. Since  $\mathcal{G}'$  is a minimal cover of  $\mathrm{M}_{22}$  and 10 is the smallest dimension of a nontrivial projective representation of  $\mathrm{M}_{22}$  over  $\mathbf{C}$  [1], the Feit–Tits theorem implies that  $\ker(\pi)$  contains  $\ker(\mathcal{G}' \twoheadrightarrow \mathrm{M}_{22})$ . Since the image of  $\pi$  lies in  $\mathrm{PSL}(10, \mathbf{R}) \subset \mathrm{PSL}(10, \mathbf{C})$ , we conclude that  $\pi$  gives rise to a nontrivial homomorphism  $\mathrm{M}_{22} \rightarrow \mathrm{PSL}(10, \mathbf{R})$ . Contradiction.

Case (e) follows easily from the Feit–Tits theorem combined with Examples 3.9 and Lemma 3.12.  $\square$

**Theorem 3.14.** *Suppose  $g > 1$  is an integer,  $D$  a finite-dimensional semisimple  $\mathbf{Q}$ -algebra enjoying the following properties:*

- (i) *Let us split  $D$  into a direct sum*

$$D = \oplus_{i=1}^r D_i$$

*of simple  $\mathbf{Q}$ -algebras  $D_i$ . Then the number  $r$  of summands does not exceed  $g$ ;*

- (ii) Let us present a summand  $D_i$  as the algebra of square matrices of size  $d_i$  over a division  $\mathbf{Q}$ -algebra  $T_i$ . Then every  $T_i$  admits a positive involution. In addition,

$$\bigoplus_{i=1}^r d_i \leq g.$$

- (iii) If  $r = 1$  (i.e.,  $D = D_1$  is simple) then  $n_1 \cdot \dim_{\mathbf{Q}}(T_1)$  divides  $2g$ . In addition, the center of  $D_1$  is either a totally real number field of degree dividing  $g$  or a CM-field of degree dividing  $2g$ . Also, if  $n_1 \cdot \dim_{\mathbf{Q}}(T_1) = 2g$  and  $T_1$  is a quaternion  $\mathbf{Q}$ -algebra then it is indefinite.

Suppose  $H$  is a  $g$ -unbounded group and  $\rho : H \rightarrow \text{Aut}(D)$  is a group homomorphism such that the subalgebra

$$D^H = \{u \in D \mid \rho(h)u = u \quad \forall h \in H\}$$

of  $H$ -invariants coincides with  $\mathbf{Q}$ . Then  $D = \mathbf{Q}$ .

*Proof.* Let  $C_i$  be the center of  $D_i$ . Then  $C_i$  is either a totally real number field or a CM-field. Clearly,

$$C = \bigoplus_{i=1}^r C_i$$

is the center of  $D$ . This implies that  $C$  is  $H$ -stable and the action of  $H$  permutes  $C_i$ 's. This gives rise to a homomorphism from  $H$  to the group  $\mathbf{S}_r$  of permutations in  $r$  letters which must be trivial. Indeed, the perfectness of  $H$  implies that its image in  $\mathbf{S}_r$  lies in the alternating subgroup  $\mathbf{A}_r$  which embeds into  $\text{PSL}(r-1, \mathbf{C})$  if  $r > 2$  and the inequality  $r \leq g$  and the triviality of homomorphisms implies in this case that  $H \rightarrow \mathbf{S}_r$  is trivial. If  $r \leq 2$  then  $\mathbf{A}_r$  is itself trivial. So,  $H$  leaves stable each  $C_i$ . This implies easily that  $\bigoplus_{i=1}^r \mathbf{Q}$  consists of  $H$ -invariants. Since  $D^H = \mathbf{Q}$ , we conclude that  $r = 1$  and therefore  $D = D_1$  and  $C = C_1$  is also the center of  $M_1$ . So,  $C$  is either a totally real number field of degree dividing  $g$  or a purely imaginary quadratic extension of a totally real number field  $C^+$  where  $[C^+ : \mathbf{Q}]$  divides  $g$ . In the case of totally real  $C$  let us put  $C^+ := C$ . Clearly, in both cases  $C^+$  is the largest totally real subfield of  $C$  and therefore the action of  $H$  leaves  $C^+$  stable. Let us put  $d := [C^+ : \mathbf{Q}]$ . I claim that  $d = 1$ , i.e.,  $C^+ = \mathbf{Q}$ . Indeed, suppose  $d > 1$ . Clearly, one may identify  $\text{Aut}(H)$  with a subgroup of  $\text{GL}(d-1, \mathbf{Q})$  and therefore the action of  $H$  on  $C^+$  gives us a homomorphism

$$H \rightarrow \text{Aut}(C^+) \subset \text{GL}(d-1, \mathbf{Q}),$$

whose triviality we need to check. Assume the contrary. The perfectness of  $H$  implies that the *nontrivial* image of the composition  $H \rightarrow \text{GL}(d-1, \mathbf{Q})$  is, in fact, a perfect subgroup of  $\text{SL}(d-1, \mathbf{Q})$ . This perfectness implies, in turn, that the image of  $H$  in  $\text{PSL}(d-1, \mathbf{Q})$  is also nontrivial. Taking into account the inequality  $d \leq g$  and the inclusion  $\text{PSL}(d-1, \mathbf{Q}) \subset \text{PSL}(d-1, \mathbf{C})$ , we obtain a nontrivial homomorphism  $H \rightarrow \text{PSL}(d-1, \mathbf{C})$ . This contradicts to the  $g$ -unboundness of  $H$ . Hence  $d = 1$  and  $C^+ = \mathbf{Q}$ .

Now I claim that  $C = \mathbf{Q}$ . Indeed, if  $C \neq \mathbf{Q}$  then  $C$  is an imaginary quadratic field and  $\text{Aut}(C)$  is a cyclic group of order 2. The perfectness of  $H$  implies that  $H \rightarrow \text{Aut}(C)$  is trivial and therefore  $C$  consists of  $H$ -invariants. Since  $D^H = \mathbf{Q}$ , we get a contradiction. Hence  $C = \mathbf{Q}$ .

So,  $T_1$  is a central simple  $\mathbf{Q}$ -algebra with a positive involution. Hence either  $T_1 = \mathbf{Q}$  or a quaternion  $\mathbf{Q}$ -algebra.

Assume that  $T_1 = \mathbf{Q}$ . Then  $D = D_1$  is the matrix algebra of size  $d_1$  over  $\mathbf{Q}$ . Clearly,  $d_1 \leq g$ . By Skolem-Noether theorem,  $\text{Aut}(D)$  is  $\text{PGL}(d_1, \mathbf{Q})$ . So, perfect  $H$

acts on  $D$  via  $\rho : H \rightarrow \text{Aut}(D) = \text{PGL}(d_1, \mathbf{Q})$ , whose image must lie in  $\text{PSL}(d_1, \mathbf{Q})$ . Since  $d_1 \leq g$ ,  $\text{PSL}(d_1, \mathbf{Q})$  is a subgroup of  $\text{PSL}(g, \mathbf{Q})$ , we obtain the triviality of  $\rho : H \rightarrow \text{Aut}(D) = \text{PGL}(d_1, \mathbf{Q}) \subset \text{PSL}(g, \mathbf{Q})$ . This implies that the whole  $D$  consists of  $H$ -invariants. Since  $D^H = \mathbf{Q}$ , we conclude that  $D = \mathbf{Q}$ .

Now assume that  $T_1$  is a quaternion  $\mathbf{Q}$ -algebra. Then  $D \neq \mathbf{Q}$  and therefore  $\rho : H \rightarrow \text{Aut}(D)$  is nontrivial. We need to arrive to a contradiction.

We have  $\dim_{\mathbf{Q}}(D_1) = 4$  and  $4n_1 \leq 2g$ .

Assume that  $4n_1 = 2g$ . Then  $g = 2n_1$  is even and  $T_1 \otimes_{\mathbf{Q}} \mathbf{R}$  is isomorphic to the matrix algebra of size 2 over  $\mathbf{R}$ . This implies that  $D_{\mathbf{R}} := D \otimes_{\mathbf{Q}} \mathbf{R}$  is the matrix algebra of size  $g$  and

$$\text{Aut}(D) \subset \text{Aut}_{\mathbf{R}}(D_{\mathbf{R}}) = \text{PGL}(g, \mathbf{R}).$$

Therefore the nontrivial  $\rho$  gives rise to a nontrivial homomorphism  $H \rightarrow \text{PGL}(g, \mathbf{R})$ . Again, the perfectness of  $H$  implies that the image lies in  $\text{PSL}(g, \mathbf{R})$  and we get a nontrivial homomorphism  $H \rightarrow \text{PSL}(g, \mathbf{R})$ . Contradiction.

Assume that  $4n_1 < 2g$ . Then  $D_{\mathbf{C}} := D \otimes_{\mathbf{Q}} \mathbf{C}$  is the matrix algebra over  $\mathbf{C}$  of size  $2n_1 < g$  and

$$\text{Aut}(D) \subset \text{Aut}_{\mathbf{C}}(D_{\mathbf{C}}) = \text{PGL}(2n_1, \mathbf{C}).$$

Therefore the nontrivial  $\rho$  gives rise to a nontrivial homomorphism  $H \rightarrow \text{PGL}(g-1, \mathbf{C})$ . Again, the perfectness of  $H$  implies that the image of  $H$  lies in  $\text{PSL}(g-1, \mathbf{C})$  and we get a nontrivial homomorphism  $H \rightarrow \text{PSL}(g-1, \mathbf{C})$ . Contradiction.  $\square$

Let  $B$  be a finite set consisting of  $n \geq 5$  elements. We write  $\text{Perm}(B)$  for the group of permutations of  $B$ . A choice of ordering on  $B$  gives rise to an isomorphism

$$\text{Perm}(B) \cong \mathbf{S}_n.$$

Let us consider the permutation module  $\mathbf{F}_2^B$ : the  $\mathbf{F}_2$ -vector space of all functions  $\varphi : B \rightarrow \mathbf{F}_2$ . The space  $\mathbf{F}_2^B$  carries a natural structure of  $\text{Perm}(B)$ -module and contains the stable line  $\mathbf{F}_2 \cdot 1_B$  of constant functions and the stable hyperplane  $(\mathbf{F}_2^B)^0$  of functions  $\varphi$  with  $\sum_{\alpha \in B} \varphi(\alpha) = 0$ . Clearly,  $(\mathbf{F}_2^B)^0$  contains  $\mathbf{F}_2 \cdot 1_B$  if and only if  $n$  is even. Let us put  $Q_B := (\mathbf{F}_2^B)^0$  if  $n$  is odd and  $Q_B := (\mathbf{F}_2^B)^0 / (\mathbf{F}_2 \cdot 1_B)$  if  $n$  is even. Clearly,  $Q_B$  carries a natural structure of faithful  $\text{Perm}(B)$ -module. For each permutation group  $H \subset \text{Perm}(B)$  the corresponding  $H$ -module is called the *heart* of the permutation representation of  $H$  on  $B$  over  $\mathbf{F}_2$  [7], [11], [3].

**Lemma 3.15.**  *$\text{End}_H(Q_B) = \mathbf{F}_2$  if either  $n$  is odd and  $H$  acts 2-transitively on  $B$  or  $n$  is even and  $H$  acts 3-transitively on  $B$ .*

*Proof.* See Satz 4 in [7].  $\square$

**Lemma 3.16.** *Suppose  $q > 2$  is an integral odd power prime,  $m \geq 3$  is an integer,  $B = \mathbf{P}^{m-1}(\mathbf{F}_q)$  is the corresponding  $(m-1)$ -dimensional projective space over the finite field  $\mathbf{F}_q$  and  $H = \mathbf{L}_m(q) = \text{PSL}_m(\mathbf{F}_q) \subset \text{Perm}(B)$  is the corresponding projective special linear group over  $\mathbf{F}_q$  acting naturally and faithfully on the projective space. Then the  $H$ -module  $Q_B$  is absolutely simple. In particular,  $\text{End}_H(Q_B) = \mathbf{F}_2$ .*

*Proof.* See [11], Table 1 on page 2.  $\square$



**Remark 3.17.** Suppose  $q = 2^r$  is an integral power of 2 and  $m \geq 2$  is an integer such that either  $m \geq 3$  or  $q > 4$ . Suppose  $B = \mathbf{P}^{m-1}(\mathbf{F}_q)$  is the corresponding  $(m-1)$ -dimensional projective space over the finite field  $\mathbf{F}_q$  and  $H = \mathbf{L}_m(q) = \mathrm{PSL}_m(\mathbf{F}_q) \subset \mathrm{Perm}(B)$  is the corresponding projective special linear group over  $\mathbf{F}_q$  acting naturally and faithfully on the projective space. Then  $\mathrm{End}_H(Q_B) = \mathbf{F}_2$ . Indeed, it is well-known that  $H = \mathrm{PSL}_m(\mathbf{F}_q)$  acts doubly transitively on  $B = \mathbf{P}^{m-1}(\mathbf{F}_q)$ . Clearly,  $\#(B) = \frac{q^m-1}{q-1}$  is odd and therefore the assertion follows from Lemma 3.15. Notice that if  $m \geq 3$  then the  $H$ -module  $Q_B$  is reducible [11]; see §5 of [3] for details. If  $m = 2$  then the  $H$ -module  $Q_B$  is absolutely simple [11] (and even very simple [21]).

#### 4. ENDOMORPHISMS OF ABELIAN VARIETIES

Let  $K$  be a field of characteristic zero. We fix its algebraic closure  $K_a$  and write  $\mathrm{Gal}(K)$  for the absolute Galois group  $\mathrm{Aut}(K_a/K)$ . Let  $X$  is an abelian variety of positive dimension defined over  $K$ . Then the group  $X(K_a)$  of its algebraic points has a natural structure of  $\mathrm{Gal}(K)$ -module. If  $m$  is a positive integer then we write  $X_m$  for the kernel of multiplication by  $m$  in  $X(K_a)$ . It is well-known [12] that  $X_m$  is a free  $\mathbf{Z}/m\mathbf{Z}$ -module of rank  $2\dim(X)$  provided with the structure of  $\mathrm{Gal}(K)$ -module inherited from  $X(K_a)$ . We denote by  $\tilde{\rho}_{m,X}$  the corresponding homomorphism

$$\tilde{\rho}_{m,X} : \mathrm{Gal}(K) \rightarrow \mathrm{Aut}(X_m) \cong \mathrm{GL}(2\dim(X), \mathbf{Z}/m\mathbf{Z})$$

which defines the structure of Galois module on  $X_m$ . We have

$$\tilde{\rho}_{m,X}(\sigma)(x) = \sigma(x) \quad \forall \sigma \in \mathrm{Gal}(K), x \in X_m \subset X(K_a).$$

We write

$$\tilde{G}_{m,X,K} = \tilde{\rho}_{m,X}(\mathrm{Gal}(K))$$

for the image of  $\mathrm{Gal}(K)$  in  $\mathrm{Aut}(X_m)$ . If  $K(X_m)$  is the field of definition of all points of order  $m$  on  $X$  then it is a finite Galois extension of  $K$ , whose Galois group  $\mathrm{Gal}(K(X_m)/K) = \tilde{G}_{m,X,K}$ .

Suppose  $m > 1$ . Clearly,  $X_m = mX_{m^2}$  coincides with the kernel of multiplication by  $m$  in  $X_{m^2}$ . In particular, every endomorphism of the commutative group  $X_{m^2}$  leaves  $X_m$  stable. Therefore the restriction to  $X_m$  gives rise to a natural (obviously surjective) ring homomorphism (the reduction modulo  $m$ )

$$\mathrm{red}_m : \mathrm{End}(X_{m^2}) \rightarrow \mathrm{End}(X_m); \quad \mathrm{red}_m(u)(x) = ux \quad \forall u \in \mathrm{End}(X_{m^2}), x \in X_m.$$

Clearly,  $\ker(\mathrm{red}_m) = m\mathrm{End}(X_{m^2})$ . This implies that each  $v \in \ker(\mathrm{red}_m)$  satisfies  $v^2 = 0 = mv$ .

Restricting  $\mathrm{red}_m$  to the automorphism group  $\mathrm{Aut}(X_{m^2})$  of  $X_{m^2}$ , we obtain the (obviously surjective) group homomorphism

$$\mathrm{red}_m^* : \mathrm{Aut}(X_{m^2}) \rightarrow \mathrm{Aut}(X_m); \quad \mathrm{red}_m^*(u)(x) = ux \quad \forall u \in \mathrm{Aut}(X_{m^2}), x \in X_m.$$

Clearly,  $\ker(\mathrm{red}_m^*) = \mathrm{Id} + m\mathrm{End}(X_{m^2})$ . (Here  $\mathrm{Id}$  is the identity automorphism of  $X_{m^2}$ .) This implies that each  $u \in \ker(\mathrm{red}_m^*)$  is of the form  $\mathrm{Id} + v$  with  $v^2 = 0 = mv$ . This implies that  $u^m = \mathrm{Id}$ .

**Remark 4.1.** Notice that the homomorphisms  $\mathrm{red}_m$  and  $\mathrm{red}_m^*$  do not depend on the choice of the field of definition  $K$  for  $X$ . In particular, they both are  $\mathrm{Gal}(K)$ -equivariant.

Clearly,

$$\tilde{\rho}_{m^2, X} = \text{red}_m^* \tilde{\rho}_{m, X};$$

in particular,

$$\tilde{G}_{m, X, K} = \text{red}_m^*(\tilde{G}_{m^2, X, K}).$$

There is an important special case when  $m = \ell$  is a prime. Then  $X_m = X_\ell$  is a  $2\dim(X)$ -dimensional  $\mathbf{F}_\ell$ -vector space provides with the structure of  $\text{Gal}(K)$ -module inherited from  $X(K_a)$ . We have  $u^\ell = \text{Id}$  for each  $u \in \ker(\text{red}_\ell^*)$  and therefore the exponent of the nontrivial finite group  $\ker(\text{red}_\ell^*)$  is  $\ell$ . We also have

$$\tilde{G}_{\ell, X, K} = \text{red}_\ell^*(\tilde{G}_{\ell^2, X, K})$$

and therefore the kernel of the surjective group homomorphism

$$\text{red}_\ell^* : \tilde{G}_{\ell^2, X, K} \rightarrow \tilde{G}_{\ell, X, K}$$

is either trivial or a finite group of exponent  $\ell$ .

**Remark 4.2.** Assume that  $\tilde{G}_{\ell, X, K}$  contains a simple non-abelian subgroup  $\mathcal{G}$ . Let  $H \subset \tilde{G}_{\ell^2, X, K}$  be the preimage of  $\mathcal{G}$  with respect to  $\text{red}_\ell^* : \tilde{G}_{\ell^2, X, K} \rightarrow \tilde{G}_{\ell, X, K}$ . Clearly,  $\text{red}_\ell^* : H \rightarrow \mathcal{G}$  is a surjective homomorphism, whose kernel is either trivial or has exponent  $\ell$ . According to Remark 3.8 there exists a subgroup

$$\mathcal{G}' \subset H \subset \tilde{G}_{\ell^2, X, K}$$

such that  $\text{red}_\ell^* : \mathcal{G}' \rightarrow \mathcal{G}$  is a minimal  $\ell$ -cover.

Clearly,  $\{\sigma \in \text{Gal}(K) \mid \rho_{\ell^2, X}(\sigma) \in \mathcal{G}'\}$  is an open subgroup of finite index in  $\text{Gal}(K)$  and therefore coincides with the Galois group of certain finite separable algebraic extension  $L = L_{\mathcal{G}}$  of  $K$ . Clearly, we have

$$\rho_{\ell^2, X}(\text{Gal}(L)) = \mathcal{G}', \quad \rho_{\ell, X}(\text{Gal}(L)) = \text{red}_\ell^*(\mathcal{G}') = \mathcal{G}.$$

In other words,

$$\tilde{G}_{\ell^2, X, L} = \mathcal{G}', \quad \tilde{G}_{\ell, X, L} = \mathcal{G}$$

and  $\text{red}_\ell^* : \tilde{G}_{\ell^2, X, L} \rightarrow \tilde{G}_{\ell, X, L}$  is a minimal  $\ell$ -cover.

We write  $\text{End}(X)$  for the ring of all  $K_a$ -endomorphisms of  $X$  and  $\text{End}_K(X)$  for the ring of all  $K$ -endomorphisms of  $X$ . We have

$$\mathbf{Z} = \mathbf{Z} \cdot \text{Id}_X \subset \text{End}_K(X) \subset \text{End}(X)$$

where  $\text{Id}_X$  is the identity automorphism of  $X$ .

Since  $X$  is defined over  $K$ , one may associate with every  $u \in \text{End}(X)$  and  $\sigma \in \text{Gal}(K)$  an endomorphism  ${}^\sigma u \in \text{End}(X)$  such that

$${}^\sigma u(x) = \sigma u(\sigma^{-1}x) \quad \forall x \in X(K_a).$$

In fact, there is a group homomorphism

$$\kappa_X : \text{Gal}(K) \rightarrow \text{Aut}(\text{End}(X)); \quad \kappa_X(\sigma)(u) = {}^\sigma u \quad \forall \sigma \in \text{Gal}(K), u \in \text{End}(X).$$

It is well-known that  $\text{End}_K(X)$  coincides with the subring of  $\text{Gal}(K)$ -invariants in  $\text{End}(X)$ , i.e.,

$$\text{End}_K(X) = \{u \in \text{End}(X) \mid {}^\sigma u = u \quad \forall \sigma \in \text{Gal}(K)\}.$$

It is also well-known, that  $\text{End}(X)$ , viewed as a group (with respect to addition) is a free commutative group of finite rank and  $\text{End}_K(X)$  is its *pure* subgroup, i.e., the

quotient-group  $\text{End}(X)/\text{End}_K(X)$  is also a free commutative group of finite rank. It is also well-known that there exists a finite Galois extension  $K'$  of  $K$  such that all the endomorphisms of  $X$  are defined over  $K'$ , i.e.

$$\text{Gal}(K') \subset \ker(\kappa_X) \subset \text{Gal}(K).$$

**Remark 4.3.** It is proven in [15] that all the endomorphisms of  $X$  are defined over  $K(X_4) \cap K(X_\ell)$  where  $\ell$  is an arbitrary odd prime. This means that  $\ker(\tilde{\rho}_{4,X}) \subset \ker(\kappa_X)$  and  $\ker(\tilde{\rho}_{\ell,X}) \subset \ker(\kappa_X)$  for all odd primes  $\ell$ . This implies that if  $\Gamma_K = \kappa_X(\text{Gal}(K)) \subset \text{Aut}(\text{End}(X))$  then there exists a surjective homomorphism

$$\tilde{G}_{4,X} \twoheadrightarrow \Gamma_K \subset \text{Aut}(\text{End}(X))$$

and surjective homomorphisms

$$\tilde{G}_{\ell,X} \twoheadrightarrow \Gamma_K \subset \text{Aut}(\text{End}(X))$$

for all odd primes  $\ell$ . Notice that

$$\text{End}_K(X) = \text{End}(X)^{\Gamma_K}.$$

**Remark 4.4.** (i) Let us put  $\text{End}^0(X) := \text{End}(X) \otimes \mathbf{Q}$ . It is well-known ([12], §21) that  $\text{End}^0(X)$  is a semisimple finite-dimensional  $\mathbf{Q}$ -algebra. Clearly, the natural map

$$\text{Aut}(\text{End}(X)) \rightarrow \text{Aut}(\text{End}^0(X))$$

is an embedding.

- (ii) Recall that  $X$  is isogenous over  $K_a$  to a product  $\prod_{i=1}^r Y_i^{d_i}$  where  $Y_i$ 's are mutually non-isogenous absolutely simple abelian varieties (of positive dimension) over  $K_a$ . Then

$$g = \dim(X) = \sum_{i=1}^r d_i \cdot \dim(Y_i) \geq \sum_{i=1}^r d_i.$$

Let us put  $T_i := \text{End}^0(Y_i)$  and denote by  $D_i$  the algebra of square matrices of size  $d_i$  over  $T_i$ . Then each  $T_i$  is a division  $\mathbf{Q}$ -algebra admitting a positive involution ([12], §21). Let us denote by  $C_i$  the center of  $T_i$ . Then either  $C_i$  is a totally real number field and  $[C_i : \mathbf{Q}]$  divides  $\dim(Y_i)$  or  $C_i$  is a CM-field and  $[C_i : \mathbf{Q}]$  divides  $2\dim(Y_i)$ . It is also clear that  $C_i$  is the center of  $D_i$ .

Since  $\text{char}(K_a) = 0$ , the number  $\dim_{\mathbf{Q}}(T_i)$  divides  $2\dim(Y_i)$  ([12], §21, p. 202). Clearly,  $D_i = \text{End}^0(Y_i^{d_i})$  and

$$\text{End}^0(X) = \oplus_{i=1}^r D_i.$$

- (iii) Assume now that  $r = 1$ , i.e.,  $X$  is isogenous to  $Y_1^{d_1}$  and  $\text{End}^0(X) = D_1$ . Then  $g = \dim(X) = d_1 \cdot \dim(Y_1)$ . Hence either  $C_1$  is totally real number field and  $[C_1 : \mathbf{Q}]$  divides  $g$  or  $C_1$  is a CM-field and  $[C_1 : \mathbf{Q}]$  divides  $2g$ . It is also clear that  $d_1 \dim_{\mathbf{Q}}(T_1)$  divides  $d_1 \cdot 2\dim(Y_1) = 2\dim(X) = 2g$ . If  $C_1 = \mathbf{Q}$  then  $T_1$  is either  $\mathbf{Q}$  or a quaternion  $\mathbf{Q}$ -algebra.
- (iv) We continue to assume that  $r = 1$ . If  $T_1$  a quaternion  $\mathbf{Q}$ -algebra and  $d_1 \dim_{\mathbf{Q}}(T_1) = 2g$  then, taking into account that  $\dim_{\mathbf{Q}}(T_1) = 4$ , we conclude that  $g$  is even,  $d_1 = g/2$  and  $Y_1$  is an absolutely simple abelian surface. Since in characteristic zero the endomorphism algebra of an absolutely simple abelian surface is either a field or an indefinite quaternion  $\mathbf{Q}$ -algebra [13] (see also [14]), we conclude that  $T_1 = \text{End}^0(Y_1)$  is an *indefinite* quaternion  $\mathbf{Q}$ -algebra.

**Theorem 4.5.** *Suppose  $K$  is a field of characteristic 0, suppose  $X$  is an abelian variety over a  $K$  of dimension  $g > 1$ . Suppose  $\ell$  is a prime,*

$$\tilde{G}_{\ell,X,K} = \tilde{\rho}_{\ell,X}(\text{Gal}(K)) \subset \text{Aut}(X_\ell)$$

*is the image of  $\text{Gal}(K)$  in  $\text{Aut}(X_\ell)$ . Let us put  $g = \dim(X)$  and assume that  $g > 1$  (i.e.,  $X$  is not an elliptic curve). Assume that  $\tilde{G}_{\ell,X,K}$  contains a simple non-abelian subgroup  $\mathcal{G}$  such that*

$$\text{End}_{\mathcal{G}}(X_\ell) = \mathbf{F}_\ell$$

*and one of the following conditions holds:*

- (a)  $\ell$  is odd and  $\mathcal{G}$  is  $g$ -unbounded.
- (b)  $\ell = 2$  and every 2-minimal cover of  $\mathcal{G}$  is  $g$ -unbounded.

*Then the ring  $\text{End}(X)$  of all  $K_a$ -endomorphisms of  $X$  coincides with  $\mathbf{Z}$ .*

*Proof.* First, using Remark 4.2 we may replace  $K$  by its finite separable algebraic extension  $L$  in such a way that  $\tilde{G}_{\ell,X,L} = \mathcal{G}$  and

$$\text{red}_\ell^* : \tilde{G}_{\ell^2,X,L} \hookrightarrow \tilde{G}_{\ell,X,L} = \mathcal{G}$$

is a minimal  $\ell$ -cover. Clearly,  $\tilde{G}_{\ell,X,L}$  is  $g$ -unbounded if  $\ell$  is odd. If  $\ell = 2$  then it follows from Remark 4.2 that  $\tilde{G}_{\ell,X,L}$  is a minimal 2-cover of  $\tilde{G}_{2,X,L} = \mathcal{G}$  and therefore is  $g$ -unbounded.

Second, I claim that  $\text{End}_L(X) = \mathbf{Z}$ . Indeed, it is well-known that there is an embedding.

$$\text{End}_L(X) \otimes \mathbf{Z}/\ell\mathbf{Z} \hookrightarrow \text{End}_{\text{Gal}(K)}(X_\ell).$$

On the other hand, since  $\rho_{\ell,X}(\text{Gal}(L)) = \tilde{G}_{\ell,X,L}$ ,

$$\text{End}_{\text{Gal}(K)}(X_\ell) = \text{End}_{\tilde{G}_{\ell,X,L}}(X_\ell) = \text{End}_{\mathcal{G}}(X_\ell) = \mathbf{F}_\ell,$$

the rank of free commutative group  $\text{End}_L(X)$  is either 0 or 1. Clearly, it must be 1 and this implies that  $\text{End}_L(X) = \mathbf{Z}$ .

Now let us put  $D := \text{End}^0(X)$ . Clearly,  $\text{End}(X)$  is a  $\mathbf{Z}$ -lattice in the  $\mathbf{Q}$ -vector space  $D$ . Let  $\text{Aut}(D)$  be the group of automorphisms of the  $\mathbf{Q}$ -algebra  $D$ . We have  $\text{Aut}(\text{End}(X)) \subset \text{Aut}(D)$ . We have

$$\kappa_X(\text{Gal}(L)) = \Gamma_L \subset \text{Aut}(\text{End}(X)) \subset \text{Aut}(D).$$

Clearly, we have

$$D^{\Gamma_L} = \text{End}(X)^{\Gamma_L} \otimes \mathbf{Q} = \text{End}_L(X) \otimes \mathbf{Q} = \mathbf{Z} \otimes \mathbf{Q} = \mathbf{Q}.$$

We are going to finish the proof, using Theorem 3.14. Let us put  $H := \tilde{G}_{\ell,X,L}$  if  $\ell$  is odd and  $H := \tilde{G}_{4,X,L}$  if  $\ell = 2$ . Clearly, in both cases  $H$  is  $g$ -unbounded. Thanks to Remark 4.3, there exists a surjective homomorphism

$$\rho : H \twoheadrightarrow \Gamma_L \subset \text{Aut}(D).$$

Clearly,

$$D^H = D^{\Gamma_L} = \mathbf{Q}.$$

In light of Remark 4.4 the semisimple  $\mathbf{Q}$ -algebra  $D = \text{End}^0(X)$  satisfies all the conditions of Theorem 3.14 with  $g = \dim(X)$ . Applying Theorem 3.14, we conclude that  $D = \mathbf{Q}$ , i.e.,  $\text{End}^0(X) = \mathbf{Q}$  and therefore  $\text{End}(X) = \mathbf{Z}$ .  $\square$

## 5. HYPERELLIPTIC JACOBIANS

**Theorem 5.1.** *Let  $K$  be a field of characteristic zero,  $K_a$  its algebraic closure,  $f(x) \in K[x]$  a polynomial of degree  $n \geq 5$  and  $\mathfrak{R} \subset K_a$  the set of its roots. let  $K(\mathfrak{R}) \subset K_a$  be the splitting field of  $f$  and  $\text{Gal}(f) := \text{Gal}(K(\mathfrak{R})/K)$  the Galois group of  $f$ , viewed as a subgroup of the group  $\text{Perm}(\mathfrak{R})$  of all permutations of  $\mathfrak{R}$ . Suppose  $\text{Gal}(f)$  contains a simple non-abelian group  $\mathcal{G}$  enjoying one of the following two properties:*

- (i)  *$n$  is odd and  $\mathcal{G}$  acts 2-transitively on  $\mathfrak{R}$ . In addition, every 2-minimal cover of  $\mathcal{G}$  is  $\frac{n-1}{2}$ -bounded.*
- (ii)  *$n$  is even and  $\mathcal{G}$  acts 3-transitively on  $\mathfrak{R}$ . In addition, every 2-minimal cover of  $\mathcal{G}$  is  $\frac{n-2}{2}$ -bounded.*
- (iii)  *$n$  is even and  $\text{End}_{\mathcal{G}}(Q_{\mathfrak{R}}) = \mathbf{F}_2$ . In addition, every 2-minimal cover of  $\mathcal{G}$  is  $\frac{n-2}{2}$ -bounded.*

*Let  $J(C_f)$  be the jacobian of the hyperelliptic curve  $C = C_f : y^2 = f(x)$ . Then the ring  $\text{End}(J(C_f))$  of all  $K_a$ -endomorphisms of  $J(C_f)$  coincides with  $\mathbf{Z}$ .*

*Proof.* Suppose  $f(x) \in K[x]$  is a polynomial of degree  $n \geq 5$  without multiple roots and  $X := J(C_f)$  is the jacobian of  $C = C_f : y^2 = f(x)$ . It is well-known that  $g := \dim(X) = \frac{n-1}{2}$  if  $n$  is odd and  $g := \dim(X) = \frac{n-2}{2}$  if  $n$  is even. It is also well-known (see for instance Sect. 5 of [21]) that  $\tilde{G}_{2,X,K} \cong \text{Gal}(f)$ . More precisely, let  $K(\mathfrak{R})$  be the splitting field of  $f$  and  $\text{Gal}(f) := \text{Gal}(K(\mathfrak{R})/K)$  the Galois group of  $f$ , viewed as a subgroup of the group  $\text{Perm}(\mathfrak{R})$  of all permutations of  $\mathfrak{R}$ . We have

$$\text{Gal}(f) \subset \text{Perm}(\mathfrak{R})$$

and the action of  $\text{Gal}(f)$  on  $\mathfrak{R}$  is transitive if and only if  $f$  is irreducible.

Now let us consider the heart (end of §3) of the permutation representation of  $\text{Gal}(f)$  on  $\mathfrak{R}$ : the faithful  $\text{Gal}(f)$ -module  $Q_{\mathfrak{R}}$ . It is well-known (see for instance, Th. 5.1 on p. 478 of [21]) that the homomorphism  $\rho_{2,X} : \text{Gal}(K) \rightarrow \text{Aut}(X_2)$  factors through the canonical surjection  $\text{Gal}(K) \twoheadrightarrow \text{Gal}(K(\mathfrak{R})/K) = \text{Gal}(f)$  and the  $\text{Gal}(f)$ -modules  $X_2$  and  $Q_{\mathfrak{R}}$  are isomorphic. In particular,

$$\text{Gal}(f) = \rho_{2,X}(\text{Gal}(K)) = \tilde{G}_{2,X,K}.$$

Now assume that  $f$  satisfies the conditions of Theorem 5.1 and let us put

$$H := \mathcal{G} \subset \text{Gal}(f) \subset \text{Perm}(\mathfrak{R}) \subset \text{Aut}(Q_{\mathfrak{R}}) = \text{Aut}(X_2).$$

It follows easily from Lemma 3.15 that we always have

$$\text{End}_{\mathcal{G}}(X_2) = \text{End}_H((Q_{\mathfrak{R}})) = \mathbf{F}_2.$$

Now the assertion of Theorem 5.1 follows readily from Theorem 4.5.  $\square$

*Proof of Theorem 2.1.* It is well-known that all Mathieu groups  $M_n \subset \mathbf{S}_n$  are, at least, 3-transitive permutation groups. Now Theorem 2.1 becomes an immediate corollary of Theorem 5.1 combined with Lemma 3.13(a-c).  $\square$

*Proof of Theorem 2.3.* Recall that  $q$  is a power of 2. It is well-known that  $\mathbf{L}_m(q) = \text{PSL}_m(\mathbf{F}_q)$  acts doubly transitively on  $\mathbf{P}^{m-1}(\mathbf{F}_q) = \mathfrak{R}$ . It is also clear that  $n = \#(\mathbf{P}^{m-1}(\mathbf{F}_q)) = \frac{q^m-1}{q-1}$  is odd. Now Theorem 2.3 becomes an immediate corollary of Theorem 5.1 combined with Examples 3.9 and Lemma 3.12.  $\square$

*Proof of Theorem 2.5.* It is an immediate corollary of Theorem 5.1 combined with Lemma 3.13(d) and Lemma 3.16. (Notice that in this case one may check that the  $\text{Gal}(f)$ -module  $J(C_f)_2$  is very simple.)  $\square$

**Remark 5.2.** Combining Remark 3.13(e) and Lemma 3.15 with Theorem 5.1, we obtain immediately that  $\text{End}(J(C_f)) = \mathbf{Z}$  if  $n \geq 5$  and  $\text{Gal}(f)$  contains  $\mathbf{A}_n$ . This assertion was proven by a different method (based on the very simplicity of  $J(C_f)_2$ ) in [20].

#### REFERENCES

- [1] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, *Atlas of finite groups*. Clarendon Press, Oxford, 1985.
- [2] W. Feit, J. Tits, *Projective representations of minimum degree of group extensions*. *Canad. J. Math.* **30** (1978), 1092–1102.
- [3] A. A. Ivanov, Ch. E. Praeger, *On finite affine 2-Arc transitive graphs*. *Europ. J. Combinatorics* **14** (1993), 421–444.
- [4] N. Katz, *Monodromy of families of curves: applications of some results of Davenport-Lewis*. In: *Séminaire de Théorie des Nombres, Paris 1979-80* (ed. M.-J. Bertin); *Progress in Math.* **12**, pp. 171–195, Birkhäuser, Boston-Basel-Stuttgart, 1981.
- [5] N. Katz, *Affine cohomological transforms, perversity, and monodromy*. *J. Amer. Math. Soc.* **6** (1993), 149–222.
- [6] P. B. Kleidman, M. W. Liebeck, *On a theorem of Feit and Tits*. *Proc. AMS* **107** (1989), 315–322.
- [7] M. Klemm, *Über die Reduktion von Permutationsmoduln*. *Math. Z.* **143** (1975), 113–117.
- [8] D. Masser, *Specialization of some hyperelliptic jacobians*. In: *Number Theory in Progress* (eds. K. Györy, H. Iwaniec, J. Urbanowicz), vol. I, pp. 293–307; de Gruyter, Berlin-New York, 1999.
- [9] Sh. Mori, *The endomorphism rings of some abelian varieties*. *Japanese J. Math.* **2**(1976), 109–130.
- [10] Sh. Mori, *The endomorphism rings of some abelian varieties*. II. *Japanese J. Math.* **3**(1977), 105–109.
- [11] B. Mortimer, *The modular permutation representations of the known doubly transitive groups*. *Proc. London Math. Soc.* (3) **41** (1980), 1–20.
- [12] D. Mumford, *Abelian varieties*, Second edition, Oxford University Press, London, 1974.
- [13] F. Oort, *Endomorphism algebras of abelian varieties*. *Algebraic Geometry and Commutative Algebra in Honor of M. Nagata* (1987, Ed. H. Hijikata et al), Kinokuniya Co., Tokyo 1988; Vol. II, pp. 469 - 502.
- [14] F. Oort, Yu. G. Zarhin, *Endomorphism algebras of complex tori*. *Math. Ann.* **303** (1995), 11–29.
- [15] A. Silverberg, *Fields of definition for homomorphisms of abelian varieties*. *J. Pure Appl. Algebra* **77** (1992), 253–262.
- [16] A. Silverberg, Yu. G. Zarhin, *Variations on a theme of Minkowski and Serre*. *J. Pure and Applied Algebra* **111** (1996), 285–302.
- [17] M. Suzuki, *Group Theory I*. Springer-Verlag, 1982.
- [18] Pham Huu Tiep, A. E. Zalesskii, *Minimal characters of the finite classical groups*. *Comm. Algebra* **24**(1996), 2093–2167.
- [19] A. Wagner, *The faithful linear representations of  $\mathbf{S}_n$  and  $\mathbf{A}_n$  over a field of characteristic 2*. *Math. Z.* **151** (1976), 127–137.
- [20] Yu. G. Zarhin, *Hyperelliptic jacobians without complex multiplication*. *Math. Res. Letters* **7** (2000), 123–132.
- [21] Yu. G. Zarhin *Hyperelliptic jacobians and modular representations*. In: *Moduli of abelian varieties* (C. Faber, G. van der Geer, F. Oort, eds.), pp. 473–490, *Progress in Math.*, Vol. 195, Birkhäuser, Basel–Boston–Berlin, 2001.
- [22] Yu. G. Zarhin, *Hyperelliptic jacobians without complex multiplication in positive characteristic*. *Math. Res. Letters* **8** (2001), 429–435.

- [23] Yu. G. Zarhin, *Cyclic covers of the projective line, their jacobians and endomorphisms*, <http://xxx.lanl.gov/abs/math.AG/0003002>, to appear in J. reine angew. Math.
- [24] Yu. G. Zarhin, *Hyperelliptic jacobians and simple groups  $U_3(2^m)$* . Proc. AMS, to appear.
- [25] Yu. G. Zarhin, *Very simple 2-adic representations and hyperelliptic jacobians*, <http://arXiv.org/abs/math.AG/0109014>.

Department of Mathematics, Pennsylvania State University,  
University Park, PA 16802, USA

Institute for Mathematical Problems in Biology,  
Russian Academy of Sciences, Pushchino, Moscow Region, 142292, RUSSIA

*E-mail address:* zarhin@math.psu.edu