

EXPLICIT DESCENT OVER $X(3)$ AND $X(5)$

CATHERINE O'NEIL

ABSTRACT. Let E be an elliptic curve over a field K , and let p denote either 3 or 5. In the case that all of the p -torsion points of E are defined over K , we give an explicit description of p -descent on E in two parts. The first part consists of an explicit map from the finite abelian group $E(K)/pE(K)$ to $Sel(E)[p]$, and the second part is an explicit identification of $Sel(E)[p]$ with models of principal homogeneous spaces of E inside \mathbb{P}^{p-1} .

1. INTRODUCTION

This paper represents a complete solution, in a special case, to the problem of performing explicit p -descent on elliptic curves for the primes $p = 3$ and $p = 5$. Namely, we work with the assumption that all of the p -torsion points of the elliptic curve E are defined over the field of definition of E . From this vantage point, an obstruction (called the ‘period-index obstruction’) to constructing smooth models of genus one curves in low-dimensional projective spaces is extremely concrete and in some sense is the only obstruction to performing descent. The general case, when there are no rationality assumptions on the p -torsion, also contains this obstruction and much more; we believe it is instructive to isolate this obstruction, i.e. to consider this a kind of ‘base case’ for a general method.

A few remarks are warranted. First, 2-descent over \mathbb{Q} is completely known and implemented for the computer with no rationality assumptions on the 2-torsion points. References for 2-descent include John Cremona’s ‘mwrnk’ program and [5] (see pages 84 and 85 for the manifestation of the period-index obstruction in the case of 2-descent), based on the methods explained in [2]. For complete 2-descent over a general number field see [12]. An algorithmic method for p -descent with no rationality assumption and which generalises to some higher genus cases can be found in [14].

Next, why do 3-descent if 2-descent is known? Loosely speaking, when there is nontrivial 2-torsion in III , 2-descent is not effective, but 3-descent may be, if the 3-torsion of III is trivial. In other words the obstruction to the effectiveness of p -descent is $\text{III}[p]$. Since III is conjecturally finite, descent will be ‘eventually effective.’ Certainly it is useful to have two or three primes completely understood, and maybe more if possible.

Date: October 25, 2018.

Next, we know (by the Galois equivariance of the Weil pairing) that no elliptic curve defined over \mathbb{Q} satisfies the assumption that all of its 3 or 5 torsion is rational. Therefore we must be taking elliptic curves over larger number fields. However, given an elliptic curve E over \mathbb{Q} , it is still may be worthwhile to ‘base-change’ to a number field K where the assumption does hold; performing descent over K may determine, for example, the Mordell-Weil rank of E over K , thus bounding the Mordell-Weil rank of E over \mathbb{Q} .

Work is currently in progress, involving the author and others, to explicitly perform 3-descent on E (in Weierstrass form) without any assumptions of rationality of the 3-torsion points of E .

The paper is organised as follows: first we will give the general set-up, reviewing the mechanics of p -descent on elliptic curves and our approach, which will consist of two basic parts. Next we will work through the cases $p = 3$ and $p = 5$ separately.

2. THE GENERAL SET-UP

Fix a number field K and an elliptic curve $(E, \mathcal{O}_E \in E(K))$ defined over K , where $\mathcal{O}_E \in E(K)$ is the origin of the group law for $E(K)$. Then the Mordell-Weil group $E(K)$ is known to be a finitely generated abelian group, so we can write it as a product of a free part and a torsion part: $E(K) = \mathbb{Z}^r \times E(K)_{tors}$. For a prime $p \in \mathbb{N}$, we then know that $E(K)/pE(K) = (\mathbb{Z}/p\mathbb{Z})^r \times E[p](K)$ is finite, and its dimension as an \mathbb{F}_p -vector space is $r + \epsilon$, where $\epsilon = 0, 1$ or 2 . A major and historical goal of number theorists is to compute r . The theory of p -descent is one approach.

Indeed, ‘(The first) p -descent on E ’ can be described as an attempt to compute the finite group $Sel(E)[p]$; this group sits in a commutative diagram whose rows are exact sequences and where each group on the top row is a subgroup of the corresponding group on the bottom row:

$$\begin{array}{ccccccccc} 1 & \rightarrow & E(K)/pE(K) & \rightarrow & Sel(E)[p] & \rightarrow & \text{III}(E)[p] & \rightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \rightarrow & E(K)/pE(K) & \rightarrow & H^1(K, E[p]) & \rightarrow & H^1(K, E) & \rightarrow & 1 \end{array},$$

The elements of the torsion cohomological group $H^1(K, E)$ have long been (abstractly) identified with (isomorphism classes of) principal homogeneous spaces for E . A nontrivial class of $H^1(K, E)$ is represented by a genus one curve with no points defined over K and whose Jacobian elliptic curve is isomorphic to E . The right top group $\text{III}(E)$ is a subgroup of $H^1(K, E)$ and as such, an element of $\text{III}(E)$ is characterized as a class represented by a curve C for which, for every place v of K , there is a K_v -rational point of C . A famous conjecture states that $\text{III}(E)$ is finite.

If in fact $\text{III}(E)[p]$ is trivial, then ‘knowing $Sel(E)[p]$ ’ is equivalent to ‘knowing $E(K)/pE(K)$,’ which by the above discussion is very close to computing r ; indeed if one has a handle on the p -torsion of E , then it is the same. If on the other hand

$\text{III}(E)[p]$ is non-trivial, then one proceeds to the second level of descent involving the following commutative diagram:

$$\begin{array}{ccccccccc} 1 & \rightarrow & E(K)/p^2E(K) & \rightarrow & \text{Sel}(E)[p^2] & \rightarrow & \text{III}(E)[p^2] & \rightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \rightarrow & E(K)/pE(K) & \rightarrow & \text{Sel}(E)[p] & \rightarrow & \text{III}(E)[p] & \rightarrow & 1 \end{array}.$$

The key point is that the right-most vertical arrow is ‘multiplication by p ,’ so if $\text{III}[p^2] = \text{III}[p]$, then the image of the middle vertical map all comes from the left. In other words, we can think of the initial contribution from $\text{III}[p]$ as noise which we’ve filtered out by going to the second level of descent. Notice if $\text{III}(E)$ is finite, descent will eventually be effective.

Since we’ve already understood 2-descent, it might seem natural to next try to understand 4-descent. However, another approach to ‘filtering out’ the noise of $\text{III}(E)[2]$ is to switch *primes*; that is, we hope that if $\text{III}(E)[2]$ is non-trivial, than $\text{III}(E)[3]$ will be trivial. Moreover, the actual computations grow very quickly; we therefore save ourselves work by restricting to $p = 3$.

A more refined description of the goal of p -descent is then to not only compute the middle group $\text{Sel}(E)[p]$ but to piece together what part ‘comes from the left’ and what part ‘contributes to the right;’ thus descent breaks naturally into two basic parts. The first part is simply an explicit description of the first map. The second part is an explicit identification of elements of $\text{Sel}(E)[p]$ with smooth models of the appropriate genus one curves inside projective space. The reason we want explicit models is that we can then actually look for ‘local points,’ that is, points over the field K_v , and thus determine whether that curve contributes to the right.

Now assume K is large enough so that $E[p](K) = E[p](\overline{K})$, and choose an \mathbb{F}_p basis S, T of $E[p](K)$ with fixed Weil pairing. Then the Galois cohomology group $H^1(K, E[p])$ can be identified with the group $K^*/K^{*p} \times K^*/K^{*p}$ (see Lemma 3.1 on page 4 of [9]), and its subgroup $\text{Sel}(E)[p]$ can be identified with a finite subgroup of $K^*/K^{*p} \times K^*/K^{*p}$.

For the first part of descent (see Chapter X of ([11])), we find a pair of rational functions (f_S, f_T) on E which, when evaluated at a point of $E(K)$, give its image in $H^1(G, E[p]) \cong K^*/K^{*p} \times K^*/K^{*p}$. Note this will depend on the chosen basis above. By Theorem 1.1d on page 278 of [11], it is sufficient for the functions f_S and f_T to satisfy $\text{div}(f_S) = p \cdot (S) - p \cdot (O_E)$ and $\text{div}(f_T) = p \cdot (T) - p \cdot (O_E)$ respectively; moreover, they can be chosen to satisfy $f_S \circ [p] = g_S^p$ and $f_T \circ [p] = g_T^p$ for some rational functions g_S and g_T .

Lemma 2.1. *The expansions of f_S and f_T with respect to a local parameter at O_E can be chosen to have leading coefficients which are perfect p th powers, i.e. so $f_S = \frac{a}{t^p} + \dots$ and $f_T = \frac{b}{t^p} + \dots$ where both a and b are perfect p th powers and where $t \in \mathcal{O}_{E, O_E}$ is a parameter in the local ring at O_E . Here the ‘ \dots ’ refer to ‘higher order terms.’*

Proof. First we prove that locally the expansions look like:

$$f_S = \frac{a}{t^p} + \dots, \quad g_S = \frac{c}{t} + \dots, \quad \text{and} \quad t \circ [p] = d \cdot t + \dots$$

The first is because we know f_S has a pole of order p at O_E , the second because g_S has a simple pole at O_E ; in fact $\text{div}(g_S) = \sum_{pP_i=S}(P_i) - \sum_{pQ_i=O}(Q_i)$, and we get only one copy of O_E on the right. Finally, $t \circ [p]$ has a simple zero at O_E since $[p]O_E = O_E$ and $[p]$ is étale when $\text{char}(K) \neq p$. We know that $f_T \circ [p] = g_T^p$, and a quick calculation shows this is equivalent to a being a perfect p th power. \square

The background for the second part is to be found in [9], and is summarized as follows.

We want to find an efficient model of C . However, there isn't always a degree three line bundle on C , so we can't always model C as a smooth cubic in \mathbb{P}^2 . The order of C as an element of $H^1(K, E)$ (called the 'period' of C) is closely related to the smallest degree of a line bundle \mathcal{L} on C (called the 'index' of C). To explain this better, we will view the Selmer group $\text{Sel}(E)[p]$ as a subgroup of the cohomology group $H^1(K, E[n])$. By Proposition 2.2 on page 3 of [9], elements of $H^1(K, E[n])$ correspond to diagrams $C \rightarrow S$ where C is a period n principal homogeneous spaces of the elliptic curve E as above, and where S is a Brauer-Severi variety of dimension $n - 1$. These diagrams are twists of a fixed diagram $E \rightarrow \mathbb{P}^{n-1}$, given by the divisor $n \cdot O_E$. Under the natural map from $H^1(K, E[p])$ to $H^1(K, E)$, the diagram $C \rightarrow S$ goes to C . The other forgetful map, sending $C \rightarrow S$ to S , is a quadratic map from $H^1(K, E[p])$ to $H^1(K, \text{PGL}_p) \subset H^2(K, \mathcal{G}_m)[p] = \text{Br}(K)[p]$ (for this inclusion see p. 158 of [10]) and is called the *period-index obstruction map*. The obstruction is trivial exactly when S is isomorphic over K to \mathbb{P}^{p-1} , and in this case we say that the diagram $C \rightarrow S$ has its period equal to its index. In our situation we will make use of our identification $H^1(K, E[p]) \cong K^*/K^{*p} \times K^*/K^{*p}$; then the period-index obstruction for an element $(a, b) \in K^*/K^{*p} \times K^*/K^{*p}$ is (Proposition 3.4 on page 6 of [9]) its associated 'Hilbert symbol' $(a, b)_{\text{Hilb}, p}$. In particular, $(a, b)_{\text{Hilb}, p}$ is trivial exactly when b is in the image of the norm map from the field $K(\alpha)$ to K , where α is chosen such that $\alpha^p = a$.

A crucial fact that we will take advantage of is that the elements of $\text{Sel}(E)[p]$, which a priori correspond to diagrams $C \rightarrow S$, always have $S \cong_K \mathbb{P}^{p-1}$; in other words, diagrams $C \rightarrow S$ where $C \in \text{III}(E)$ *always* have trivial period-index obstruction (for a proof see the Remark on page 3 of [9]).

In this article we will perform the second part of descent not only for elements of $\text{Sel}(E)[p]$ but for any element of $H^1(K, E[p])$ which has trivial period-index obstruction. Indeed we will start with a "trivialisation" of a Hilbert symbol (the data of an element $\beta \in K[x]/(x^p - a)$ whose norm is a), and produce a model for the corresponding genus one curve under the isomorphism $H^1(K, E[p]) \cong K^*/K^{*p} \times K^*/K^{*p}$.

3. EXPLICIT DESCENT OVER $X(3)$

3.1. The image of $E(K)/3E(K)$. Let K be a field whose characteristic is different from 3 and which contains a primitive 3rd root of unity ζ . Let E be an elliptic curve over K with full 3-torsion given by the following equation:

$$E : X^3 + Y^3 + Z^3 + \lambda XYZ = 0.$$

Define the origin O_E to be the point $(1; -1; 0)$ and a basis of 3-torsion points $\langle S, T \rangle$ to be $S = (1; \zeta; 0), T = (1; 0; -1)$, where $\zeta \in K$ is a fixed cube root of unity.

By results in [8], there exist matrices in PGL_3 which act as “translation by S and T ” on E and which we will denote by M_S and M_T . We leave it to the reader to prove these matrices are D_3 and $M_{1,3}$ respectively, where D_n is the diagonal matrix $\mathrm{diag}(1, \zeta_n, \zeta_n^2, \dots, \zeta_n^{n-1})$ for some primitive n root of unity ζ_n , (here take $\zeta_3 = \zeta$) and where

$$M_{a,n} = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ a & 0 & 0 & \dots & 0 \end{pmatrix}.$$

We will find a pair of rational functions (f_S, f_T) on E which when evaluated at a point of $E(K)$ gives its image in $H^1(G, E[3]) \cong K^*/K^{*3} \times K^*/K^{*3}$. By Lemma 2.1, the function f_S needs to satisfy the condition that $\mathrm{div}(f_S) = 3 \cdot (S) - 3 \cdot (O_E)$; this is satisfied trivially by the quotient of the hyperplane at S by the hyperplane at O_E , since they are both flex points. The hyperplane at O_E is easily seen to be given by $3X + 3Y - \lambda Z$, and the other hyperplanes are in fact translates of this one by the above matrices; this observation doesn't save much time in \mathbb{P}^2 , since hyperplanes are easy to compute, but will in the $n = 5$ case. Similarly f_T is, up to a constant, a ratio of hyperplanes, and we get $f_S = \kappa_S \frac{3\zeta^2 X + 3\zeta Y - \lambda Z}{3X + 3Y - \lambda Z}$ and $f_T = \kappa_T \frac{3X - \lambda Y + 3Z}{3X + 3Y - \lambda Z}$ for some constants κ_S and κ_T .

Proposition 3.1. $f_S = (\lambda^3 + 27) \frac{3\zeta^2 X + 3\zeta Y - \lambda Z}{3X + 3Y - \lambda Z}$ and $f_T = (\lambda^2 - 3\lambda + 9) \frac{3X - \lambda Y + 3Z}{3X + 3Y - \lambda Z}$.

Proof. By Lemma 2.1, we want then to compute the expansion of $\frac{3\zeta^2 X + 3\zeta Y - \lambda Z}{3X + 3Y - \lambda Z}$ at the origin $O_E = (1; -1; 0)$. Here we can work in affine coordinates by setting $X = 1$, since this is true locally. Then $F : 1 + Y^3 + Z^3 + \lambda YZ = 0$. Moreover, since O_E takes on a non-zero value at the hyperplane at S , we can just evaluate there to get $(3\zeta^2 X + 3\zeta Y - \lambda Z)|_{O_E} = 3\zeta^2 - 3\zeta$. Note that $\zeta^2 - \zeta = \sqrt{-3}$, so $3\zeta^2 - 3\zeta = (-\sqrt{-3})^3$ is a cube in K . So in fact we can completely ignore this term. We're left with $\frac{1}{3X + 3Y - \lambda Z}$ which has a triple zero at O_E . In other words

$$3 + 3Y - \lambda Z \cong aZ^3 + \dots,$$

where \cong signifies that we are working in \mathcal{O}_{E, O_E} . Multiply the above by the function Y , a nonzero function at O_E , to get $3Y + 3Y^2 - \lambda YZ \cong aYZ^3 + \dots$; since $-\lambda YZ \cong 1 + Y^3 + Z^3$ (we're working modulo F) we substitute to get

$$1 + 3Y + 3Y^2 + Y^3 + Z^3 = (1 + Y)^3 + Z^3 \cong aYZ^3 + \dots$$

The function $1 + Y$ has a zero at O_E , so it's an alternative parameter for the local ring $\mathcal{O}_{E, O_E} : 1 + Y \cong bZ + \dots$ but we already have the tangent line equation which tells us that $3(1 + Y) \cong \lambda Z + \dots$, i.e. $b = \lambda/3$. Replace $(1 + Y)^3$ above now by $(\lambda/3)Z^3$:

$$Z^3(1 + (\lambda/3)^3) \cong aYZ^3.$$

Evaluate at $Y = -1$ to get $a = -(1 + (\lambda/3)^3)$. Our original constant then is $1/a$, which modulo cubes is seen to be $\frac{1}{27 + \lambda^3}$. Finally, to normalize we want the function f_S to have a leading coefficient which is 1. In other words, we need to multiply the ratio of the two hyperplanes by the constant $27 + \lambda^3$. For f_T , we are already almost done- the only difference is the value of the numerator at the origin: $(3X - \lambda Y + 3Z)|_{O_E} = 3 + \lambda$. The leading coefficient then is $\frac{3 + \lambda}{27 + \lambda^3} = \frac{1}{\lambda^2 - 3\lambda + 9}$. \square

3.2. Models of Genus One curves in \mathbb{P}^2 . Let E be an elliptic curve over the field K . Assume $\text{char}(K) \neq 3$ and that $E[3](K) = E[3](\overline{K})$. Choose a basis $\langle S, T \rangle$ of $E[3]$ and identify $H^1(K, E[3])$ with $K^*/K^{*3} \times K^*/K^{*3}$. Take $(a, b) \in H^1(K, E[3])$ whose corresponding Hilbert symbol is trivial. If both a and b are cubes, the element (a, b) represents E . We will first assume that a is not a cube in K ; the other case will be dealt with at the end of this section. Define $\alpha, \beta \in \overline{K}$ such that $\alpha^3 = a$ and $\beta = \sum_{i=0}^2 \beta_i \alpha^i$ is such that $\mathbb{N}_{K(\alpha)/K}(\beta) = b$. Moreover define $\sigma \in G(K(\alpha)/K)$ to be the generator of that cyclic Galois group such that $\sigma(\alpha) = \alpha\zeta$. Let Tr be the trace function of $K(\alpha)$ to K , and let $u = \beta/\sigma^2(\beta)$.

Theorem 3.2. *The curve (a, b) is given by*

$$\begin{aligned} &(Tr(u) + \lambda)(a^2X^3 + aY^3 + Z^3) + 3Tr(\alpha u)(aX^2Z + aY^2X + Z^2Y) + \\ &3Tr(\alpha^2u)(aX^2Y + Y^2Z + Z^2X) + 3(2aTr(u) - \lambda a)XYZ = 0. \end{aligned}$$

Proof. By results in [8], the action of the 3-torsion points of E on a genus one curve C embedded in \mathbb{P}_K^2 can be represented as automorphisms of \mathbb{P}_K^2 , in other words as elements of $\text{PGL}_3(K)$. Moreover, if C is represented in $H^1(G, E[3])$ by the pair (a, b) which depends on a chosen basis $\langle S, T \rangle$ then the determinants of the matrices representing “translation by S ” and “translation by T ” are a and b respectively. Moreover the Weil pairing is given by the commutator of lifts to $\text{GL}_3(K)$. With that in mind, we will search for $C \leftrightarrow (a, b)$ by finding cubics which are invariant under the action of standard matrices representing this 3-torsion action. Define M_S to be the matrix $M_{a,3}$ and M_T to be the matrix $D_3[\beta_0 I + \beta_1 M_S + \beta_2 M_S^2]$, if $\beta = \beta_0 + \beta_1 \alpha + \beta_2 \alpha^2$. The determinant of M_S is a , the determinant of M_T is b , and

the commutator $[M_S, M_T]$ is ζI . A cubic which is invariant under the action of M_S but with no fixed points must be of the form

$$F = A(a^2X^3 + aY^3 + Z^3) + B(aX^2Z + aY^2X + Z^2Y) + \\ C(aX^2Y + Y^2Z + Z^2X) + 3DXYZ = 0.$$

This is because M_S acts linearly on the 10-dimensional space of cubics. There are 3 eigenspaces of dimensions 3, 3, and 4, and the first two have zeroes at the fixed points of M_S , whereas the last eigenspace does not and is generated by the above four cubics.

On the other hand we also insist that F be invariant under the action of M_T . To ease computations we introduce the following notation: fix eigenvectors $v_i = (1, \alpha\zeta^i, \alpha^2\zeta^{2i})$ of M_S . Then $M_S v_i = \alpha\zeta^i v_i$. The four coefficients A, B, C , and D are linear combinations of $F(v_0), F(v_1), F(v_2)$, and $\mathcal{T}(v_0, v_1, v_2)$, where \mathcal{T} is the trilinear form associated to F , as follows:

$$\begin{pmatrix} F(v_0) \\ F(v_1) \\ F(v_2) \\ \mathcal{T}(v_0, v_1, v_2) \end{pmatrix} = \begin{pmatrix} 3a^2 & 3a^{5/3} & 3a^{4/3} & 3a \\ 3a^2 & 3a^{5/3}\zeta^2 & 3a^{4/3}\zeta & 3a \\ 3a^2 & 3a^{5/3}\zeta & 3a^{4/3}\zeta^2 & 3a \\ 18a^2 & 0 & 0 & -9a \end{pmatrix} \begin{pmatrix} A \\ B \\ C \\ D \end{pmatrix}.$$

Now it is easy to see how M_T acts on the $F(v_i)$:

$$F^{M_T}(v_i) = F(M_T v_i) = F(D[\beta_0 I + \beta_1 M_S + \beta_2 M_S^2]v_i) = \\ F(D(\beta_0 + \beta_1 \alpha \zeta^i + \beta_2 \alpha^2 \zeta^{2i})v_i) = F(\sigma^i(\beta) D v_i) = \sigma^i(\beta)^3 F(v_{i+1}).$$

Similarly,

$$\mathcal{T}^{M_T}(v_0, v_1, v_2) = \mathcal{T}(\beta v_1, \sigma(\beta)v_2, \sigma^2(\beta)v_0) = b \mathcal{T}(v_0, v_1, v_2).$$

The fact that F is invariant under the action of M_T is equivalent to the projective point $P = (F(v_0); F(v_1); F(v_2); \mathcal{T}(v_0, v_1, v_2))$ being fixed by M_T . We have seen that $M_T(P) = (\beta^3 F(v_1); \sigma(\beta)^3 F(v_2); \sigma^2(\beta)^3 F(v_0); b \mathcal{T}(v_0, v_1, v_2))$. For some $\mu \neq 0$ we have $F(v_1) = \frac{\mu}{\beta^3} F(v_0)$ and $F(v_2) = \frac{\sigma^2(\beta)^3}{\mu} F(v_0)$. Moreover, the Jacobian of the above curve is (see [8]):

$$X^3 + Y^3 + \prod_{i=0}^2 F(v_i) Z^3 + \mathcal{T}(v_0, v_1, v_2) XYZ = 0,$$

in other words

$$X^3 + Y^3 + \frac{F(v_0)^3 \sigma^2(\beta)^3}{\beta^3} Z^3 + \mathcal{T}(v_0, v_1, v_2) XYZ = 0;$$

setting $F(v_0) = \frac{\beta}{\sigma^2(\beta)}$ (note that $F(v_0) \neq 0$ because F has no fixed points under the action of M_S) and renaming $\mathcal{T}(v_0, v_1, v_2) = \lambda$, the Jacobian is exactly E . Note that $\sigma(F(v_i)) = F(\sigma(v_i)) = F(v_{i+1})$, so $F(v_1) = \frac{\sigma(\beta)}{\beta}$ and $F(v_2) = \frac{\sigma^2(\beta)}{\sigma(\beta)}$. To finish the

proof, we need to invert the above matrix to find the coefficients A, B, C , and D in terms of the $F(v_i)$'s and \mathcal{T} :

$$\begin{pmatrix} A \\ B \\ C \\ D \end{pmatrix} = \frac{1}{27a^2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 3\alpha & 3\alpha\zeta & 3\alpha\zeta^2 & 0 \\ 3\alpha^2 & 3\alpha^2\zeta^2 & 3\alpha^2\zeta & 0 \\ 2a & 2a & 2a & -a \end{pmatrix} \begin{pmatrix} F(v_0) \\ F(v_1) \\ F(v_2) \\ \mathcal{T}(v_0, v_1, v_2) \end{pmatrix}.$$

We can ignore the factor of $\frac{1}{27a^2}$, since we're working projectively. Then $A = F(v_0) + F(v_1) + F(v_2) + \mathcal{T}(v_0, v_1, v_2) = \frac{\beta}{\sigma^2(\beta)} + \frac{\sigma(\beta)}{\beta} + \frac{\sigma^2(\beta)}{\beta} + \lambda = \text{Tr}(u) + \lambda$, and similarly for the other coefficients. \square

When a is a cube The formula in Theorem 3.2 was computed assuming a is not a cube in the base field K , but it holds in the case that a is a cube as well. We can view the formula as holding over the algebra $K[\alpha]/(\alpha^3 - a)$, which is a field if a is not a cube; in other words we view α as strictly a symbol. The norm function on this algebra takes an element $s = s_0 + s_1\alpha + s_2\alpha^2$ to the expression $\mathbb{N}(s) = \prod_{i=0}^2 (s_0 + s_1\alpha\zeta^i + s_2\alpha^2\zeta^{2i}) = s_0^3 + s_1^3a + s_2^3a^2 - 3s_0s_1s_2a$. Similarly, the trace function has $\text{Tr}(s) = \sum_{i=0}^2 (s_0 + s_1\alpha\zeta^i + s_2\alpha^2\zeta^{2i}) = 3 \cdot s_0$, since $1 + \zeta + \zeta^2 = 0$. In other words, we can define the linear automorphism ' σ ' to take α to $\zeta\alpha$, and $\mathbb{N}(s) = s\sigma(s)\sigma^2(s)$. In the case of representing a curve corresponding to the pair (a, b) for a a cube, we may take $a = 1$ and we want to find $\beta = \beta_0 + \beta_1\alpha + \beta_2\alpha^2$ such that $\mathbb{N}(\beta) = \beta\sigma(\beta)\sigma^2(\beta) = b$. There are many solutions: for example, we can take $\beta_0 = \frac{b+2}{3}$ and $\beta_1 = \beta_2 = \frac{b-1}{3}$ if those are nonzero quantities. Then as above we can formally compute $\text{Tr}(u) = (b-1)^2/b + 3$, $\text{Tr}(\alpha u) = (b-1)(b-\zeta^2)/b$, and $\text{Tr}(\alpha^2 u) = (b-1)(b-\zeta)/b$.

4. EXPLICIT DESCENT OVER $X(5)$

4.1. The image of $E(K)/5E(K)$. Let K be a field whose characteristic is different from 5 and which contains a primitive 5th root of unity ζ . Let $E = E_\lambda$ be the elliptic curve over K given by the equations $\lambda x_i^2 + \lambda^2 x_{i-2}x_{i+2} - x_{i-1}x_{i+1}$ for i between 0 and 4 and whose origin is given by $O_E = (0; \lambda; 1; -1; -\lambda)$.

Remark. The curves E_λ for $n = 3, 5$ are also considered in [6], where the Cassels-Tate pairing is used to compute $S^{(n)}(E_\lambda/\mathbb{Q})$ for all non-cuspidal $\lambda \in \mathbb{Q}$. Equations for E_λ for $n = 5$ in Weierstrass form may be found in [6] or [1].

The curve E as above has full 5-torsion over K and the matrices which act as translation by a S and T , for a basis $\langle S, T \rangle$, are given by D_5 and $M_{1,5}$ respectively (see notation on page 5).

We will find a pair of rational functions (f_S, f_T) on E which when evaluated at a point of $E(K)$ gives its image in $H^1(G, E[5]) \cong K^*/K^{*5} \times K^*/K^{*5}$. By Lemma 2.1, the function f_S can be chosen to satisfy $\text{div}(f_S) = 5 \cdot (S) - 5 \cdot (O_E)$; then we can

take f_S to be a scalar multiple of the quotient of the hypertangent plane at S by the hypertangent plane at O_E , since both S and O_E are hyperflex points (to see this, note that E is a degree 5 curve in \mathbb{P}_K^4 and that the hyperplane $x_0 = 0$ goes through the points $i \cdot S$ for i between 0 and 4, which means that the divisor giving the embedding $E \rightarrow \mathbb{P}^4$ is linearly equivalent to $(O_E) + (S) + (2S) + (3S) + (4S) \equiv 5 \cdot (O_E) \equiv 5 \cdot (S)$).

Proposition 4.1. *The hypertangent plane at the origin of E is given by*

$$H_{O_E} : \alpha x_0 + \beta(x_1 + x_4) + \gamma(x_2 + x_3),$$

where $\alpha = \lambda^{10} - 14\lambda^5 - 1$, $\beta = -5\lambda^2(1 + 2\lambda^5)$, and $\gamma = 5\lambda^3(\lambda^5 - 2)$.

Proof. An easy calculation (in Maple for example) verifies that the above hyperplane intersects E only at O_E . In order to find the above, we computed a local parameterization of the curve E in the local ring \mathcal{O}_{E,O_E} using the equations for E and Maple. \square

To finish finding the equations f_S and f_T , we simply find the hypertangent planes at S and T (these are translates of H_{O_E} by the 5-torsion matrices D_5 and $M_{1,5}$) and evaluate them at O_E . We then scale H_S/H_{O_E} by the appropriate function of λ so that its leading coefficient in the expansion at O_E is a perfect fifth power:

Proposition 4.2. *The hypertangent planes at S and at T are given by*

$$H_S : \alpha x_0 + \beta(x_1\zeta^4 + x_4\zeta) + \gamma(x_2\zeta^3 + x_3\zeta^2)$$

and

$$H_T : \alpha x_4 + \beta(x_0 + x_3) + \gamma(x_1 + x_2).$$

The rational functions f_S and f_T are given by

$$f_S = \frac{[(\lambda^2 + \lambda - 1)(\lambda^4 - 3\lambda^3 + 4\lambda^2 - 2\lambda + 1)(\lambda^4 + 2\lambda^3 + 4\lambda^2 + 3\lambda + 1)]^2}{5 \lambda (\zeta - \zeta^4) (\lambda^5 - 2)} \frac{H_S}{H_{O_E}},$$

$$f_T = \frac{\lambda^2(\lambda^4 - 3\lambda^3 + 4\lambda^2 - 2\lambda + 1)^2(\lambda^4 + 2\lambda^3 + 4\lambda^2 + 3\lambda + 1)}{(\lambda^2 + \lambda - 1)} \frac{H_T}{H_{O_E}}.$$

4.2. Models of Genus One curves in \mathbb{P}^4 . Let E be an elliptic curve over the field K . Assume $\text{char}(K) \neq 5$ and that $E[5](K) = E[5](\overline{K})$. Choose a basis $\langle S, T \rangle$ of $E[5]$ and identify $H^1(K, E[5])$ with $K^*/K^{*5} \times K^*/K^{*5}$. For $(a, b) \in H^1(K, E[5])$ whose corresponding Hilbert symbol is trivial, assume as in the previous section that a is not a perfect fifth power in K , and define $\alpha, \beta \in \overline{K}$ such that $\alpha^5 = a$ and $\beta = \sum_{i=0}^4 \beta_i \alpha^i$ such that $\mathbb{N}_{K(\alpha)/K}(\beta) = b$. Moreover define $\sigma \in G(K(\alpha)/K)$ to be the generator of that cyclic Galois group such that $\sigma(\alpha) = \alpha\zeta$.

Theorem 4.3. *There exists a unique 5-dimensional K -vector space of quadrics in the variables $x_i, i = 0, \dots, 4$, denoted $V_{\alpha, \beta}$, satisfying the following:*

- $V_{\alpha, \beta}$ defines a smooth degree 5 genus one curve $C_{\alpha, \beta}$ in \mathbb{P}^4 .

- The Jacobian of $C_{\alpha,\beta}$ is isomorphic over K to E , and in fact the diagram $C_{\alpha,\beta} \rightarrow \mathbb{P}^4$ corresponds to the pair $(a, b) \in H^1(G, E[n](\overline{K}))$.
- The action of $E[n]$ on $C_{\alpha,\beta}$, a principal homogeneous space of E , is determined by the action of S and T , and these are given as restrictions of the following automorphisms of \mathbb{P}^4 : ‘action by S ’ = $M_S = M_{\alpha,5}$, and ‘action by T ’ = $M_T = D_5 \cdot \sum_{i=0}^4 \beta_i M_S^i$.

Proof. By Proposition 2.2 on page 3 of [9], the cohomology group $H^1(G, E[n](\overline{K}))$ parameterizes diagrams $C \rightarrow \mathbb{P}^{n-1}$ up to isomorphism, where an isomorphism is a map defined over K from C to C which extends to a K -map on \mathbb{P}^{n-1} , i.e. an element of $\text{PGL}_n(K)$. Then $V_{\alpha,\beta}$ determines such a diagram, and the prescribed action of $E[n]$ fixes the model: modifying a diagram $C \rightarrow \mathbb{P}^{n-1}$ by an element of $\text{PGL}_n(K)$ has the effect of conjugating M_S and M_T . Since the group generated by M_S and M_T (a twist of the Heisenberg group) is its own centraliser, there is no non-trivial modification. \square

Our goal of this section is to determine $V_{\alpha,\beta}$.

Definition 4.1. Define, for $i = 0, \dots, 4$, v_i to be points in \mathbb{A}_K^5 such that $M_S v_i = \alpha \zeta^i v_i$. In particular we see that $\sigma(v_i) = v_{i+1}$.

Definition 4.2. Define the action of a matrix M on a function f to be such that $f^M(x) = f(Mx)$. Then $f^{M_1 M_2} = (f^{M_1})^{M_2}$.

Lemma 4.4. With notation as above, we can choose a quadric $Q \in V_{\alpha,\beta}$ such that $Q(v_0) = Q(v_1)$ and such that $\langle Q^{M_S^i} \rangle_{i=0\dots 4}$ forms a basis of $V_{\alpha,\beta}$. Moreover, such a Q is unique up to K -scaling. Therefore to determine $V_{\alpha,\beta}$ it suffices to determine Q .

Remark. We are actually taking a fixed quadric Q (not defined up to a scalar), since we need to make sense of the nonzero quantities $Q(v_0)$ and $Q(v_1)$.

Proof. We can span $V_{\alpha,\beta}$ by translates of any K -rational quadric Q by the action of powers of M_S since the eigenvalues of M_S acting on the space of quadrics are the fifth roots of a^2 , not defined over K . Note that $Q(v_i) \neq 0$ for all i since if so we would have $Q^{M_S^i}(v_i) = Q(M_S^i v_i) = 0$ as well, in other words we would have a point on C fixed by M_S , namely the projectivization of v_i . Next, note that $Q^{M_S^i}(v_0) = Q(M_S^i v_0) = Q(\alpha^i v_0) = \alpha^{2i} Q(v_0)$. Therefore if we replace Q by the quadric $Q' = \sum_{i=0}^4 a_i Q^{M_S^i}$, for $a_i \in K$, then $Q'(v_0) = \sum_{i=0}^4 a_i Q^{M_S^i}(v_0) = (\sum_{i=0}^4 a_i \alpha^{2i}) Q(v_0)$, and likewise $Q'(v_1) = \sum_{i=0}^4 a_i Q^{M_S^i}(v_1) = (\sum_{i=0}^4 a_i \alpha^{2i} \zeta^{2i}) Q(v_1)$. So in order to have $Q'(v_0) = Q'(v_1)$, we need to find $a = \sum_{i=0}^4 a_i \alpha^{2i}$ so that $Q(v_0)/Q(v_1) = \sigma(a)/a$. This is possible by Hilbert’s Theorem 90, since $Q(v_0)/Q(v_1) = Q(v_0)/\sigma(Q(v_0))$ is in the kernel of the norm map. Next, Q' and its translates under M_S also generate $V_{\alpha,\beta}$. The K -rational quadric Q' is nonzero since its value at v_0 is nonzero; by the above comment its

translates by powers of M_S span $V_{\alpha,\beta}$. Finally, such a Q' is unique up to an element of K , since if we had both Q and Q' such that $Q(v_0)/Q(v_1) = Q'(v_0)/Q'(v_1) = 1$, we could write $Q' = \sum_{i=0}^4 a_i Q^{M_S^i}$ for $a_i \in K$ to get $Q'(v_0)/Q'(v_1) = a/\sigma(a) \cdot Q(v_0)/Q(v_1)$, i.e. we would have $a = \sigma(a)$, or in other words $a \in K$. \square

Our newly defined goal is to find Q as in Lemma 4.4. We will choose a matrix M by which to modify Q as in Lemma 4.4 so that the coefficients of Q^M are easy to manipulate. Define $M = (v_0 v_1 v_2 v_3 v_4)$, a 5×5 matrix defined over $K(\alpha)$. Then

$$Q^M(x) = Q(Mx) = Q\left(\sum_{i=0}^4 v_i x_i\right) = \sum_{i=0}^4 Q(v_i) x_i^2 + \sum_{0 \leq i < j \leq 4} B(v_i, v_j) x_i x_j,$$

where $B(w, v) = Q(w + v) - Q(w) - Q(v)$.

Define $M'_S = M^{-1} M_S M$ and $M'_T = M^{-1} M_T M$. Then we have $(Q^M)^{M'_S} = (Q^{M_S})^M$ and $(Q^M)^{M'_T} = (Q^{M_T})^M$. A calculation shows us $M'_S = \alpha D_5$ and $M'_T = M_{1,5}^{-1} \cdot \text{diag}(\beta, \sigma(\beta), \sigma^2(\beta), \sigma^3(\beta), \sigma^4(\beta))$.

Next, note that by assumption M_T fixes $V_{\alpha,\beta}$. Moreover, since M_T is defined over K , the image of Q under the action by M_T is again defined over K . Thus there exist numbers $\gamma_i \in K$ such that $Q^{M_T} = \sum_{i=0}^4 \gamma_i Q^{M_S^i}$. Acting on that equation by M we get (here let $Q' = Q^M$):

$$Q'^{M'_T} = \sum_{i=0}^4 \gamma_i Q'^{(M'_S)^i}.$$

Since we know the M'_T and M'_S explicitly, we can compute the left and right sides of this equation and compare them.

$$Q'^{M'_T}(x) = \sum_{i=0}^4 Q(v_i) \sigma^{i-1}(\beta)^2 x_{i-1}^2 + \sum_{0 \leq i < j \leq 4} B(v_i, v_j) \sigma^{i-1}(\beta) \sigma^{j-1}(\beta) x_{i-1} x_{j-1}$$

and

$$Q'^{(M'_S)^i}(x) = \alpha^{2i} \left(\sum_{j=0}^4 Q(v_j) \zeta^{2ji} x_j^2 + \sum_{0 \leq j < k \leq 4} B(v_j, v_k) \zeta^{ji+ki} x_j x_k \right).$$

Proposition 4.5. *For every $i = 0, \dots, 4$, $Q(v_i) = Q(v_0)$, $B(v_{i+1}, v_{i-1}) = B(v_1, v_4) \cdot \prod_{j=1}^i \sigma^{j-1} \left(\frac{\beta^2}{\sigma(\beta)\sigma^4(\beta)} \right)$, and $B(v_{i+2}, v_{i-2}) = B(v_2, v_3) \cdot \prod_{j=1}^i \sigma^{j-1} \left(\frac{\beta^2}{\sigma^2(\beta)\sigma^3(\beta)} \right)$.*

Proof. We compare the coefficient of x_0^2 on both sides of the above equation. On the left, we get $Q(v_1)\beta^2$. On the right we get $\sum_{i=0}^4 \gamma_i \alpha^{2i} Q(v_0)$. Define $\gamma = \sum_{i=0}^4 \gamma_i \alpha^{2i}$, then we have $Q(v_1)\beta^2 = \gamma Q(v_0)$. On the other hand we have chosen Q as in Lemma 4.4 so that $Q(v_0) = Q(v_1) \neq 0$. Therefore $\gamma = \beta^2$. We can determine the rest of the $Q(v_i)$'s now since $1 = \sigma\left(\frac{Q(v_1)}{Q(v_0)}\right) = \frac{Q(v_2)}{Q(v_1)}$ etc. so all of the $Q(v_i)$'s are equal to $Q(v_0)$.

Next, compare the coefficients of x_1x_4 : on the left we get $B(v_2, v_0)\sigma(\beta)\sigma^4(\beta)$ and on the right we get $\sum_{i=0}^4 \gamma_i \alpha^{2i} B(v_1, v_4)$. In other words we have $\frac{B(v_2, v_0)}{B(v_1, v_4)} = \frac{\beta^2}{\sigma(\beta)\sigma^4(\beta)}$. Acting by σ on both sides gives $\frac{B(v_3, v_1)}{B(v_2, v_0)} = \sigma\left(\frac{\beta^2}{\sigma(\beta)\sigma^4(\beta)}\right)$, so $\frac{B(v_3, v_1)}{B(v_1, v_4)} = \frac{\beta^2}{\sigma(\beta)\sigma^4(\beta)} \cdot \sigma\left(\frac{\beta^2}{\sigma(\beta)\sigma^4(\beta)}\right)$. We continue in this way. Similarly, comparing coefficients of x_2x_3 on both sides we get $\frac{B(v_3, v_4)}{B(v_2, v_3)} = \frac{\beta^2}{\sigma^2(\beta)\sigma^3(\beta)}$ and we finish by acting on both sides by σ and solving for $\frac{B(v_{i+2}, v_{i-2})}{B(v_2, v_3)}$. \square

Now we have only to determine the three unknowns $Q(v_0)$, $B(v_1, v_4)$, and $B(v_2, v_3)$; moreover, since we are actually working projectively, we only need to know two of them, or more precisely it is adequate to know the ratios $\frac{Q(v_0)}{B(v_1, v_4)}$ and $\frac{Q(v_0)}{B(v_2, v_3)}$. Define E_λ to be the elliptic curve given by the equations $\lambda x_i^2 + \lambda^2 x_{i-2} x_{i+2} - x_{i-1} x_{i+1}$ for i between 0 and 4 and whose origin is given by $\mathcal{O}_E = (\lambda, -1; 1, -\lambda; 0)$.

Theorem 4.6. *The Jacobian of $C_{\alpha, \beta}$ is E_λ , where*

$$\lambda = -\frac{Q(v_0)}{B(v_2, v_3)} \frac{\sigma^3(\beta)\sigma^4(\beta)}{\beta\sigma(\beta)}.$$

Proof. By [8], Theorem 4.2 on page 37, the Jacobian of $C_{\alpha, \beta}$ is given as E_A given by the quadrics $S_0 = x_0^2 - x_2x_3 + x_1x_4$, $S_1 = x_1^2 - x_0x_2 + Ax_3x_4$, $S_2 = x_2^2 - x_1x_3 - Ax_0x_4$, $S_3 = x_3^2 - x_0x_1 - x_2x_4$, and $S_4 = Ax_4^2 + x_1x_2 - x_0x_3$ ¹ for a parameter

$$A = \prod_{i=0}^4 \frac{Q(v_i)}{B(v_i, v_{i+1})}.$$

A calculation using Proposition 4.5 shows that the above A in this case is the fifth power of $\frac{Q(v_0)}{B(v_2, v_3)} \frac{\sigma^3(\beta)\sigma^4(\beta)}{\beta\sigma(\beta)}$. Finally, When A is a perfect fifth power, say of $-\lambda$, the K -rational map $\text{diag}(1, -\lambda, \lambda, -1, \lambda^{-2})$ maps E_A to E_λ . \square

Lemma 4.7.

$$\frac{Q(v_0)^2}{B(v_1, v_4)B(v_2, v_3)} = -\frac{\beta^2\sigma(\beta)}{\sigma^3(\beta)\sigma^4(\beta^2)}.$$

Proof. This follows from Lemma 4.3 on page 139 of [8] and using Proposition 4.5. Loosely speaking, this is a condition on the intersection of five quadrics to form a smooth genus one curve; note that five quadrics in general position do not intersect. \square

¹This corrects a minus sign error in that paper, namely the coefficient of x_4^2 in S_4 .

Corollary 4.8.

$$\frac{B(v_1, v_4)}{Q(v_0)} = \lambda \cdot \frac{\sigma^4(\beta)}{\beta}.$$

We now have all of the coefficients of Q' in terms of $Q(v_0)$. We can divide out by this non-zero term to get

Proposition 4.9. *Let $u_1 = \frac{\sigma^4(\beta)}{\beta}$ and $u_2 = \frac{\sigma^3(\beta)\sigma^4(\beta)}{\beta\sigma(\beta)}$. Then Q' is given by:*

$$\sum_{i=0}^4 x_i^2 + \lambda \sum_{i=0}^4 x_i x_{i+2} \cdot \sigma^{i+1} u_1 - \lambda^{-1} \sum_{i=0}^4 x_i x_{i+1} \cdot \sigma^{i+3} u_2.$$

Finally, since $Q' = Q^M$, we can recover Q as $Q'^{M^{-1}}$. A calculation shows:

Theorem 4.10. *Write*

$$Q = \sum_{0 \leq i \leq j \leq 4} a_{i,j} x_i x_j.$$

Then the $a_{i,j}$ are given as follows. For $i = j$ we have

$$a_{i,i} = \text{Tr} \left(\frac{1}{\alpha^{2i}} \right) + \lambda \text{Tr} \left(\frac{u_1}{\alpha^{2i}} \right) - \lambda^{-1} \text{Tr} \left(\frac{u_2}{\alpha^{2i}} \right)$$

and for $i \neq j$ we have

$$a_{i,j} = \text{Tr} \left(\frac{2}{\alpha^{i+j}} \right) + \lambda \text{Tr} \left((\zeta^{i-j} + \zeta^{j-i}) \frac{u_1}{\alpha^{i+j}} \right) - \lambda^{-1} \text{Tr} \left((\zeta^{2i-2j} + \zeta^{2j-2i}) \frac{u_2}{\alpha^{i+j}} \right).$$

Remark. The question of solving “norm equations,” that is, developing an algorithm to find β as in the above theorem, has been studied extensively by Denis Simon [13], written up in the new book by Henri Cohen [4], and implemented in Pari.

REFERENCES

- [1] Beaver, C., *5-torsion in the Shafarevich-Tate group of a family of elliptic curves*, J. Number Theory, 82 (2000), pp. 25-46.
- [2] Birch, B. and Swinnerton-Dyer, H., *Notes on Elliptic Curves (I)*, JREIA, 212 (1963), pp. 7-25.
- [3] Cassels, J.W.S., *Lectures on Elliptic Curves*, Cambridge University Press, Cambridge (1991).
- [4] Cohen, H., *Advanced Topics in Computational Number Theory*, Graduate Texts in Mathematics, 193, Springer-Verlag (2000).
- [5] Cremona, J., *Classical Invariants and 2-descent on elliptic curves*, J. of Symb. Comp., 31 (2001), pp.71-87.
- [6] Fisher, T., *The Cassels-Tate pairing and the Platonic solids*, submitted for publication.
- [7] Lang, S. and Tate, J., *Principal homogeneous spaces over abelian varieties*, Amer. J. Math. 80 (1958), pp. 659-684.
- [8] O’Neil, C., *Jacobians of Genus One Curves*, Math. Res. Lett. 8 (2001), no. 1-2, pp. 125-140.
- [9] O’Neil, C., *The Period-Index Obstruction for Elliptic Curves*, to appear in J. of Number Theory.

- [10] Serre, J.-P., *Local Fields*, Graduate Texts in Mathematics, 67, Springer-Verlag, New York (1979).
- [11] Silverman, Joseph H., *The Arithmetic of Elliptic Curves*, Graduate Texts in Mathematics 106, Springer-Verlag, New York (1986).
- [12] Simon, D., *Computing the Rank of Elliptic Curves over Number Fields*, submitted for publication.
- [13] Simon, D., *Solving Norm Equations Using S -Units*, submitted to Math. Comp. (1999).
- [14] Schaefer, E., and Stoll, M., *How to Do a p -descent on an elliptic curve*, submitted for publication.