

# Geometric structures on $G_2$ and $Spin(7)$ -manifolds

Jae-Hyoun Lee and Naichung Conan Leung

October 24, 2018

## Abstract

This article studies the geometry of moduli spaces of  $G_2$ -manifolds, associative cycles, coassociative cycles and deformed Donaldson-Thomas bundles. We introduce natural symmetric cubic tensors and differential forms on these moduli spaces. They correspond to Yukawa couplings and correlation functions in M-theory.

We expect that the Yukawa coupling characterizes (co-)associative fibrations on these manifolds. We discuss the Fourier transformation along such fibrations and the analog of the Strominger-Yau-Zaslow mirror conjecture for  $G_2$ -manifolds.

We also discuss similar structures and transformations for  $Spin(7)$ -manifolds.

The *mirror symmetry conjecture* for Calabi-Yau threefolds, which is originated from the study of the string theory on  $X \times \mathbb{R}^{3,1}$ , has attracted much attention in both the physics and mathematics communities. Certain parts of the conjecture should hold true for Calabi-Yau manifolds of any dimensions. But others are designed only for threefolds, for example, the special Kähler geometry on the moduli space, the holomorphic Chern-Simons theory and relationships to knot invariants. These features should be interpreted in the realm of the geometry of the  $G_2$ -manifold,  $M = X \times S^1$ . From a physical point of view, they come from the studies of M-theory on  $M \times \mathbb{R}^{3,1}$ .

In mathematics, manifolds with  $G_2$  holonomy group have been studied for a while. In the Berger's list of holonomy groups of Riemannian manifolds, there are a couple of exceptional holonomy groups, namely  $G_2$  and  $Spin(7)$ . Manifolds with these holonomy groups are Einstein manifolds of dimension seven and eight respectively. As cousins of Kähler manifolds, the analogs of complex structures, complex submanifolds and Hermitian-Yang-Mills bundles are vector cross products (Gray [7]), calibrated submanifolds (Harvey and Lawson [8]) and Donaldson-Thomas bundles ([6]). These geometries are studied by many people in the Oxford school including Donaldson, Hitchin, Joyce, Thomas and others.

A manifold  $M$  with  $G_2$ -holonomy, or simply a  $G_2$ -manifold, can be characterized by the existence of a parallel positive three form  $\Omega$ . Because of the natural inclusion of Lie groups  $SU(3) \subset G_2$ , the product of a Calabi-Yau threefold  $X$  with a circle  $S^1$  has a canonical  $G_2$ -structure with  $\Omega = \text{Re } \Omega_X - \omega_X \wedge dt$ , where  $\Omega_X$  and  $\omega_X$  are the holomorphic volume form and the Calabi-Yau Kähler form on  $X$ .

The mirror symmetry conjecture for Calabi-Yau manifolds roughly says that there is a duality transformation from the symplectic geometry (or the A-model) to the complex geometry (or the B-model) between a pair of Calabi-Yau threefolds. For instance natural cubic structures, called the A- and B-Yukawa couplings, on their moduli spaces can be identified. This gives highly non-trivial predictions for the enumerative geometry of  $X$ .

From physical considerations, it is more natural to understand the *M-theory* on  $M \times \mathbb{R}^{3,1}$  with  $M$  being a  $G_2$ -manifold. This is studied recently by Acharya, Atiyah, Vafa, Witten, Yau, Zaslow and others (e.g. [1], [2], [3], [4]). To better understand the M-theory and its duality transformations, or (equivalently) the geometry of  $G_2$ -manifolds, we need to study natural geometric structures on various moduli spaces attached to  $M$ . For example we introduce the analog of the Yukawa coupling on the moduli space of  $G_2$ -metrics as follows,

$$\mathcal{Y}_M(\phi) = \int_M \Omega(\hat{\phi}, \hat{\phi}, \hat{\phi}) \wedge * \Omega,$$

where  $\phi \in H^3(M, \mathbb{R})$ . This cubic structure comes from the exceptional Jordan algebra structure for  $E_6$  and it seems to be even more natural than those A- and B-Yukawa couplings for Calabi-Yau threefolds. It will be used later to characterize different kinds of fibration structures on  $M$ .

The geometry of a  $G_2$ -manifold is reflected by its calibrated submanifolds ([8]) and Yang-Mills bundles ([6]). Calibrated submanifolds are always volume minimizing and there are two different types in a  $G_2$ -manifold  $M$ : (i) coassociative submanifolds  $C$  of dimension four, calibrated by  $*\Omega$  and (ii) associative submanifolds  $A$  of dimension three, calibrated by  $\Omega$ . Sometimes it is important to include both points and the whole manifold in the above list, together they are calibrated by different components of  $e^{\Omega+*\Omega}$ .

In physics, one needs to study *supersymmetric cycles* [14] which are calibrated submanifolds together with *deformed Yang-Mills bundles* over them. In the  $G_2$  case, there are three different types: *Coassociative cycle*  $(C, D_E)$ , with  $D_E$  an ASD connection on a coassociative submanifold  $C$ ; (ii) *Associative cycle*  $(A, D_E)$ , with  $D_E$  a unitary flat connection on an associative submanifold  $A$ , it is a critical point of the following functional,

$$\int_{A \times [0,1]} Tr \left[ e^{*\Omega + \tilde{F}} \right].$$

(iii) *Deformed Donaldson-Thomas connection*  $D_E$  which satisfies the equation

$$F \wedge *\Omega + F^3/6 = 0,$$

and it is a critical point of the following functional,

$$\int_{M \times [0,1]} Tr \left[ e^{*\Omega + \tilde{F}} \right].$$

Moduli spaces of these cycles are denoted as  $\mathcal{M}^{coa}(M)$ ,  $\mathcal{M}^{ass}(M)$  and  $\mathcal{M}^{bdl}(M)$  respectively, or simply  $\mathcal{M}^{coa}$ ,  $\mathcal{M}^{ass}$  and  $\mathcal{M}^{bdl}$ . We will define canonical three forms and four forms on them using the Clifford multiplication on spinor bundles and the Lie algebra structure on the space of self-dual two forms. In physics languages, they correspond to *correlation functions* in quantum field theory. We will also introduce natural symmetric cubic tensors on these moduli spaces.

In the flat situation such canonical three forms determine  $G_2$ -structures on both  $\mathcal{M}^{coa}$  and  $\mathcal{M}^{ass}$ . In general the moduli space of associative (resp. coassociative) submanifolds can be regarded as a coassociative (resp. associative) subspace of  $\mathcal{M}^{ass}$  (resp.  $\mathcal{M}^{coa}$ ).

We summarize these structures on various moduli spaces in following tables:

	moduli	Yukawa coupling	Metric tensor	Prepotential
$\mathcal{M}^{G_2}$	$G_2$ -structures	$\mathcal{Y}$	$\mathcal{G}$	$\mathcal{F}$

Table: Structures on the moduli space of  $G_2$ -metric with unit volume.

	moduli of	3 form	4 form	cubic tensor
$M$	points	$\Omega$	$\Theta$	
$\mathcal{M}^{coa}(M)$	coassociative cycles	$\Omega_{\mathcal{M}^{coa}(M)}$	$\Theta_{\mathcal{M}^{coa}(M)}$	$\mathcal{Y}_{\mathcal{M}^{coa}(M)}$
$\mathcal{M}^{ass}(M)$	associative cycles	$\Omega_{\mathcal{M}^{ass}(M)}$	$\Theta_{\mathcal{M}^{ass}(M)}$	$\mathcal{Y}_{\mathcal{M}^{ass}(M)}$
$\mathcal{M}^{bdl}(M)$	deformed DT bundles	$\Omega_{\mathcal{M}^{bdl}(M)}$	$\Theta_{\mathcal{M}^{bdl}(M)}$	$\mathcal{Y}_{\mathcal{M}^{bdl}(M)}$

Table: Structures on the moduli space of various cycles on  $M$ .

In section 3 we study structures of calibrated fibrations on  $M$ , they are associative  $T^3$ -fibration, coassociative  $T^4$ -fibration and coassociative K3-fibration. Then we propose a duality transformation along such fibrations on  $G_2$ -manifolds analogous to the Fourier-Mukai transformation in the geometric mirror symmetry conjecture by Strominger, Yau and Zaslow [18]. Roughly speaking it should be given by a fiberwise Fourier transformation on a coassociative  $T^4$ -fibration on  $M$ . We partially verify our proposed conjecture in the flat case, in the spirit of [11] and [13].<sup>1</sup>

For instance the transformation of an ASD connection over a coassociative torus fiber is another ASD connection over the dual torus fiber, as studied earlier by Schenk [17], Braam and van Baal [5].

We also study the Fourier transformation on an associative  $T^3$ -fibration on  $M$ . The Fourier transformation on a coassociative  $T^4$ -fibration takes cycles calibrated by  $e^\Theta$  (resp.  $*e^\Theta$ ) back to themselves. On an associative  $T^3$ -fibration, the Fourier transformation takes cycles calibrated by  $e^\Theta$  to those calibrated by  $*e^\Theta$ .

In section 4, we study the geometry of  $Spin(7)$ -manifolds in a same manner, and therefore our discussions will be brief.

## 1 $G_2$ -manifolds and their moduli

### 1.1 Definitions

We first review some basic structures on  $G_2$ -manifolds (see [10] or [16] for more details). Any  $G_2$ -manifold  $M$  admits a parallel positive three form  $\Omega_M$  (or simply  $\Omega$ ) and its Ricci curvature is zero. When

$$M = \mathbb{R}^7 = \text{Im } \mathbb{H} \oplus \mathbb{H} = \text{Im } \mathbb{O}$$

with the standard metric and orientation  $dx^{123}dy^{0123}$  we have,

$$\Omega = dx^{123} - dx^1(dy^{23} + dy^{10}) - dx^2(dy^{31} + dy^{20}) - dx^3(dy^{12} + dy^{30}).$$

---

<sup>1</sup>Those authors are also familiar with these transformations in the  $G_2$  case.

Remark: Since the Lie group  $G_2$  preserves a vector cross product structure on  $\mathbb{R}^7$ , any  $G_2$ -manifold inherits such a product structure  $\times$  on its tangent spaces given by,

$$\langle u \times v, w \rangle = \Omega(u, v, w).$$

The parallel four form  $*\Omega$  will be denoted as  $\Theta$ ,

$$\Theta = dy^{0123} + dx^{23}(dy^{23} + dy^{10}) + dx^{31}(dy^{31} + dy^{20}) + dx^{12}(dy^{12} + dy^{30}).$$

When  $M$  is the product of a Calabi-Yau threefold  $X$  and  $S^1$  with its reversed orientation, then the holonomy group equals  $Hol(M) = SU(3) \subset G_2$ . Thus  $M = X \times S^1$  is a  $G_2$ -manifold. In this case we have

$$\begin{aligned}\Omega &= \text{Re } \Omega_X - \omega_X \wedge d\theta \\ \Theta &= \text{Im } \Omega_X \wedge d\theta - \omega_X^2/2.\end{aligned}$$

where  $\omega_X$  and  $\Omega_X$  are the Ricci flat Kähler form and a holomorphic volume form on  $X$ .

Using the  $G_2$  action, we can decompose differential forms (or cohomology classes) on  $M$  into irreducible components. For example,

$$\begin{aligned}\Lambda^1 &= \Lambda_7^1 \\ \Lambda^2 &= \Lambda_7^2 \oplus \Lambda_{14}^2 \\ \Lambda^3 &= \Lambda_1^3 \oplus \Lambda_7^3 \oplus \Lambda_{27}^3.\end{aligned}$$

When  $Hol(M) = G_2$ , we have  $H_7^k(M) = 0$  for all  $k$  and  $H_1^3(M)$  (resp.  $H_1^4(M)$ ) is generated by  $\Omega$  (resp.  $\Theta$ ). Furthermore the moduli space of  $G_2$ -metrics is smooth with tangent space equals  $H^3(M) = H_1^3(M) \oplus H_{27}^3(M)$ . If we normalize the volume to be one, the tangent space to this moduli space  $\mathcal{M}^{G_2}$  becomes  $H_{27}^3(M)$ .

## 1.2 $G_2$ -Analog of Yukawa couplings

In this subsection we introduce several symmetric tensors on the moduli space of  $G_2$ -manifolds. They are the Yukawa coupling, a metric tensor and the prepotential. First we recall a tensor  $\chi \in \Omega^3(M, T_M)$  which is defined by  $\iota_v \Theta = \langle \chi, v \rangle \in \Omega^3(M)$  for every tangent vector  $v$ .

**Definition 1** Suppose  $M$  is a compact  $G_2$ -manifold, we define

$$\mathcal{C}_M : \bigotimes^3 H^3(M, \mathbb{R}) \rightarrow \mathbb{R}$$

$$\mathcal{C}_M(\phi_1, \phi_2, \phi_3) = \int_M \Omega(\hat{\phi}_1, \hat{\phi}_2, \hat{\phi}_3) \wedge \Theta,$$

where  $\hat{\phi} = *(\phi \wedge \chi) \in \Omega^1(M, T_M)$  and  $\Omega(\hat{\phi}_1, \hat{\phi}_2, \hat{\phi}_3) \in \Omega^3(M)$  is the evaluation of the three form  $\Omega$  on the vector components of  $\hat{\phi}_j$ 's.

It is clear that  $\mathcal{C}_M$  is symmetric. By using  $\mathcal{C}_M$  we define  $\mathcal{Y}_M$ ,  $\mathcal{G}_M$  and  $\mathcal{F}_M$  on  $H_{27}^3(M)$  as follows: <sup>2</sup>

Yukawa coupling	$\mathcal{Y}_M(\phi) = \mathcal{C}_M(\phi, \phi, \phi) = C \int_M \Omega \wedge * \Omega$
Metric tensor	$\mathcal{G}_M(\phi) = \mathcal{C}_M(\phi, \phi, \Omega) = C' \int_M \phi \wedge * \phi$
Prepotential	$\mathcal{F}_M = \mathcal{C}_M(\Omega, \Omega, \Omega)$ .

Table: Symmetric tensors on the moduli space of  $G_2$ -manifolds. Here  $\phi \in H_{27}^3(M)$ .

We will use this Yukawa coupling and  $p_1(M)$  in section 3.1 to describe structures of calibrated fibrations on  $M$ .

### Reduction to CY threefolds

When  $M = X \times S^1$  with  $Hol(X) = SU(3)$ , i.e.  $X$  is a Calabi-Yau threefold, any deformation of the Einstein structure on  $M$  remains a product. There is a natural isomorphism of complex vector spaces,

$$H_{27}^3(M, \mathbb{C}) \cong H^{1,1}(X) + H^{2,1}(X) + H^{1,2}(X).$$

For example if  $\phi$  is a primitive form in  $H^{1,1}(X)$  (resp.  $H^{2,1}(X)$ ) then  $\phi \wedge dt$  (resp.  $\phi$ ) is in  $H_{27}^3(M, \mathbb{C})$ . Moreover  $\mathcal{Y}_M$  restricts to the Yukawa couplings on  $H^{1,1}(X)$ ,  $H^{2,1}(X)$  and  $H^{1,2}(X)$  (e.g. [13]).

Remark: The cubic structure on the  $G_2$ -module  $\Lambda_{27}^3$  is the exceptional Jordan algebra structure on  $\mathbb{R}^{27}$ , whose automorphism group determines an exceptional Lie group of type  $E_6$ . When we reduce from  $G_2$  to  $SU(3)$ , i.e.  $M = X \times S^1$ , the Kähler form  $\omega_X$  is naturally an element in  $\Lambda_{27}^3$  which splits  $\Lambda_{27}^3 = \mathbf{27}$  into  $\mathbf{1} + \mathbf{26}$ . In this case the automorphism group of the algebraic structure of this 26 dimensional vector space determines an exceptional Lie group of type  $F_4$ . Roughly speaking the Yukawa coupling on a  $G_2$ -manifold arises from a  $E_6$  structure. When the holonomy group reduces to  $SU(3)$ , then the  $E_6$  structure also reduces to a  $F_4$  structure.

### Including $B$ -fields

From physical considerations, Acharya [1] argues that when  $M$  is a smooth  $G_2$  manifold, then the low energy theory for M-theory on  $M \times \mathbb{R}^{3,1}$  is a  $N = 1$  supergravity theory with  $b_2(M)$  Abelian vector multiplets plus  $b_3(M)$  neutral chiral multiplets. We also include one scalar field and consider the enlarged moduli space  $\mathcal{M}^{G_2+B} = \mathcal{M}^{G_2} \times H^2(M, U(1)) \times H^0(M, U(1))$ . Note  $b_0(M) = 1$ .

---

<sup>2</sup>Recall  $H_{27}^3(M) = H^3(M) \cap Ker(\wedge \Omega) \cap Ker(\wedge \Theta)$ .

We are going to construct the corresponding Yukawa coupling for this moduli space. We define (i) a symmetric cubic tensor  $\mathcal{C}'$  on  $H^2(M, \mathbb{R})$ ,

$$\begin{aligned}\mathcal{C}'_M : \bigotimes^3 H^2(M) &\rightarrow \mathbb{R} \\ \mathcal{C}'_M(\beta_1, \beta_2, \beta_3) &= \int_M \Omega(\hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3) \wedge \Theta,\end{aligned}$$

where  $\hat{\beta} = \iota_{\beta} \chi \in \Omega^1(M, T_M)$  and (ii) a symmetric bilinear tensor  $\mathcal{Q}_M$  on  $H^2(M, \mathbb{R})$ ,

$$\begin{aligned}\mathcal{Q}_M : \bigotimes^2 H^2(M) &\rightarrow \mathbb{R} \\ \mathcal{Q}_M(\beta_1, \beta_2) &= \int_M \beta_1 \wedge \beta_2 \wedge \Omega.\end{aligned}$$

Now the Yukawa coupling on the enlarged moduli space  $\mathcal{M}^{G_2} \times H^2(M, U(1)) \times H^0(M, U(1))$  of  $G_2$ -manifolds (with  $Hol = G_2$ ) is defined as a combination of  $\mathcal{Y}_M$ ,  $\mathcal{C}'_M$  and  $\mathcal{Q}_M$  as follow,

$$\tilde{\mathcal{Y}}_M : \bigotimes^3 (H_{27}^3(M) + H_{14}^2(M) + H^0(M)) \rightarrow \mathbb{R}.$$

When  $M = X \times S^1$  for a Calabi-Yau threefold  $X$  we have  $H_{14}^2(M) = H_{prim}^{1,1}(X, \mathbb{R})$  which corresponds to the decomposition  $\mathfrak{g}_2 = \mathfrak{su}(3) + \mathbb{C}^3 + \mathbb{C}^{3*}$  because of  $H^{2,0}(X) = H^{0,2}(X) = 0$  for a manifold with  $SU(3)$  holonomy. Therefore

$$H_{14}^2(M, U(1)) + H^0(M, U(1)) \cong H^2(X, U(1)),$$

and an element in it is usually called a *B-field* on the Calabi-Yau manifold  $X$ . Our Yukawa coupling  $\mathcal{Y}'$  thus reduces to the usual one on Calabi-Yau manifolds coupled with B-fields.

Remark: If we do not fix the volume of  $M$ , then the moduli space of  $G_2$ -metrics with  $B$ -fields is locally isomorphic to  $H^{\leq 3}(M, \mathbb{R})$ . In M-theory (see e.g. [4]) it is desirable to complexify this moduli space and this implies that the resulting space will be locally isomorphic to the total cohomology group  $H^*(M, \mathbb{R})$  of  $M$ .

## 2 Supersymmetric cycles

### 2.1 Deformed Donaldson-Thomas bundles

In [6], Donaldson and Thomas study the following Yang-Mills equation for Hermitian connections  $D_E$  on a complex vector bundle  $E$  over  $M$ ,

$$F_E \wedge \Theta = 0 \in \Omega^6(M, ad(E)).$$

This is the Euler-Lagrange equation for the following Chern-Simons type functional,

$$\mathcal{A}(E) \rightarrow \mathbb{R}$$

$$D_E \rightarrow \int_M CS_3(D_0, D_E) \wedge \Theta,$$

where  $D_0$  is any fixed background connection. This functional is equivalent to the following functional

$$\int_{M \times [0,1]} Tre^{\tilde{F}} \wedge \Theta,$$

where  $\tilde{F}$  is the curvature of a connection on  $M \times [0, 1]$  formed by the affine path of connections on  $E$  joining  $D_0$  and  $D_E$ .

On a  $G_2$ -manifold we introduce the following *deformed Donaldson-Thomas equation* which should have the corresponding effect of preserving supersymmetry in M-theory [14],

$$F_E \wedge \Theta + F_E^3/6 = 0 \in \Omega^6(M, ad(E)).$$

Equivalently this equals,

$$[e^{\Theta+F_E}]^{[6]} = 0 \in \Omega^6(M, ad(E)).$$

It is the Euler-Lagrangian equation for the following Chern-Simons type functional,

$$D_E \rightarrow \int_{M \times [0,1]} Tr[e^{\Theta+\tilde{F}}] = \int_M CS(D_0, D_E) e^{\Theta}.$$

The mirror transformation along a coassociative  $T^4$ -fibration on a flat  $G_2$ -manifold, which we will discuss in section 3.3, takes an associative section to a deformed Donaldson-Thomas bundles on the mirror manifold.

### Geometric structures on $\mathcal{M}^{bdl}(M)$

We define natural differential forms and a symmetric cubic tensor on the moduli space of deformed DT bundles  $\mathcal{M}^{bdl}(M)$  as follows,

3-form:	$\Omega_{\mathcal{M}^{bdl}(M)} = \int_M Tr[\alpha \wedge \beta \wedge \gamma]_{skew} e^{\Theta+F_E}$
4-form:	$\Theta_{\mathcal{M}^{bdl}(M)} = \int_M Tr[\alpha \wedge \beta \wedge \gamma \wedge \delta]_{skew} * e^{\Theta+F_E}$
Cubic:	$\mathcal{Y}_{\mathcal{M}^{bdl}(M)} = \int_M \langle \alpha, [\beta, \gamma] \rangle_{ad} e^{\Theta+F_E}.$

Table: Natural geometric structures on the moduli space of deformed DT bundles on M. Here  $\alpha, \beta, \gamma, \delta \in \Omega^1(M, ad(E))$ .



Here  $\langle \cdot, \cdot \rangle_{ad}$  is the Killing form on  $\mathfrak{g}$ , the Lie algebra of the gauge group. In particular the symmetric cubic tensor  $\mathcal{Y}_{\mathcal{M}^{bdl}(M)}$  on  $\mathcal{M}^{bdl}(M)$  is trivial when  $G$  is Abelian. Note  $e^{\Theta+F_E}$  in the cubic form works as  $[e^{\Theta+F_E}]^{[6]}$ .

Remark: When  $M$  is a flat torus  $T = \mathbb{R}^7/\Lambda$ , the moduli space  $\mathcal{M}^{bdl}(M)$  of flat  $U(1)$ -bundles is canonically isomorphic to the dual flat torus  $T^* = \mathbb{R}^{7*}/\Lambda^*$ . Under this natural identification, we have

$$\begin{aligned}\Omega_{\mathcal{M}^{bdl}(T)} &= \Omega_{T^*}, \\ \Theta_{\mathcal{M}^{bdl}(T)} &= \Theta_{T^*}.\end{aligned}$$

Moreover this transformation from  $T$  to  $\mathcal{M}^{bdl} = T^*$  is involutive, i.e.  $\mathcal{M}^{bdl}(T^*) = T$ .

Remark: The tangent bundle of  $M$  is a  $G_2$ -bundle, moreover the curvature tensor of the Levi-Civita connection on  $M$  satisfies the DT-equation,

$$F \wedge \Theta = 0.$$

This equation is equivalent to  $F \in \Omega_{14}^2(M, ad(T_M))$  and it follows from the fact that  $F \in \Omega^2(M, \mathfrak{g}_2(T_M))$  and the torsion freeness of the connection.

Infinitesimal deformations of  $T_M$  as a DT-bundle are parametrized by the first cohomology group  $H^1(M, \mathfrak{g}_2(T_M))$  of the following elliptic complex [6]:

$$0 \rightarrow \Omega^0(M, \mathfrak{g}_2(T_M)) \xrightarrow{D_E} \Omega^1(M, \mathfrak{g}_2(T_M)) \xrightarrow{\Theta D_E} \Omega^6(M, \mathfrak{g}_2(T_M)) \xrightarrow{D_E} \Omega^7(M, \mathfrak{g}_2(T_M)) \rightarrow 0.$$

When  $M = X \times S^1$  with  $X$  a Calabi-Yau threefold with  $SU(3)$  holonomy, then there is a natural isomorphism,

$$H^1(M, \mathfrak{g}_2(T_M)) \cong H^{1,1}(X) + H^{2,1}(X) + H^1(X, End_0(T_X)).$$

Furthermore the above cubic tensor  $\mathcal{Y}_{\mathcal{M}^{bdl}(M)}$  restricts to  $\mathcal{Y}_A$  on  $H^{1,1}(X)$ ,  $\mathcal{Y}_B$  on  $H^{2,1}(X)$  and a similar Yukawa coupling  $\mathcal{Y}_C$  on  $H^1(X, End_0(T_X))$ . All three seemingly unrelated Yukawa couplings on a Calabi-Yau threefold combine in a natural way when we study the  $G_2$ -manifold  $X \times S^1$ !

## 2.2 Associative cycles

### Associative submanifolds

There are two types of nontrivial calibrations on a  $G_2$ -manifold  $M$  as studied by Harvey and Lawson ([8]), they are (i) associative submanifolds and (ii) coassociative submanifolds. They are always absolute minimal submanifolds in  $M$ .

A three dimensional submanifold  $A$  in  $M$  is called an *associative* submanifold if the restriction of  $\Omega$  to  $A$  equals the volume form on  $A$  of the induced metric.

It has the following two equivalent characterizations: (1) the restriction of  $\chi \in \Omega^3(M, T_M)$  to  $A$  is zero;

$$\chi|_A = 0 \in \Omega^3(A, T_M|_A).$$

(2) the vector cross product on  $M$  preserves  $T_A$ , i.e.  $u, v \in T_A$  implies that  $u \times v \in T_A$ . As a corollary, two associative submanifolds can not intersect along a two dimensional subspace.

McLean studies the deformation theory of associative submanifolds in [15] and identifies their infinitesimal deformations as certain twisted harmonic spinors. Unlike coassociative submanifolds, deformations of an associative submanifold can be *obstructed*. For example, if we take any smooth isolated rational curve  $C$  with normal bundle  $O \oplus O(-2)$  in a Calabi-Yau threefold  $X$ , then  $C \times S^1$  is an associative submanifold in  $M = X \times S^1$  with obstructed deformations. That is the moduli space of associative submanifolds, denoted  $\mathcal{B}^{ass}(M)$ , can be singular.

### Associative cycles

From a physical perspective, one would couple associative submanifolds with gauge fields. This is analogous to coupling of gauge fields with special Lagrangian submanifolds of a Calabi-Yau manifold. From a mathematical perspective, as we will see, the moduli space of associative submanifolds coupled with gauge fields has very rich geometric structures.

An *associative cycle* is defined to be any pair  $(A, D_E)$  with  $A$  an associative submanifold in  $M$  and  $D_E$  a unitary flat connection on a bundle  $E$  over  $A$ . The flatness equation,  $F_E = 0$  for connections over a three manifold is the Euler-Lagrange equation of the standard Chern-Simons functional. Analogously the associativity condition for a pair  $(A, D_E)$  is the Euler-Lagrange equation of the following functional:

$$\begin{aligned} \text{Map}(A, M) \times \mathcal{A}(E) &\rightarrow \mathbb{R}/\mathbb{Z} \\ CS(A, D_E) &= \int_{A \times [0,1]} \text{Tr} \left[ e^{\Theta + \tilde{F}} \right]. \end{aligned}$$

The first term in the above functional is a direct analog of a functional used by Thomas [19] for special Lagrangian submanifolds in a Calabi-Yau threefold.

Next we are going to study the moduli space of associative cycles, namely the critical set of  $CS$ , and we denote it as  $\mathcal{M}^{ass}(M)$ . This space has richer structure than the moduli space of associative submanifolds  $\mathcal{B}^{ass}(M)$ .

The tangent space of  $\mathcal{M}^{ass}(M)$  at an associative cycle  $(A, D_E)$  can be identified as (see [15]),

$$T_{(A, D_E)} \mathcal{M}^{ass} = \text{Ker} \mathbf{D} \oplus H^1(A, \text{ad}(E)).$$

We define a three form  $\Omega_{\mathcal{M}^{ass}(M)}$ , four form  $\Theta_{\mathcal{M}^{ass}(M)}$  and a symmetric cubic tensor  $\mathcal{Y}_{\mathcal{M}^{ass}(M)}$  on  $\mathcal{M}^{ass}(M)$  as follows:

3-form:	$\Omega_{\mathcal{M}^{ass}(M)} = \begin{cases} \int_A Tr \alpha \wedge \beta \wedge \gamma \\ - \int_A \langle \alpha \cdot \bar{\phi}, \bar{\eta} \rangle \Omega \end{cases}$
4-form:	$\Theta_{\mathcal{M}^{ass}(M)} = \begin{cases} \int_A \langle \bar{\phi}, *(\alpha \wedge \beta) \cdot \bar{\eta} \rangle \Omega \\ \int_A \det(\phi, \eta, \xi, \zeta) \Omega \end{cases}$
Cubic:	$\mathcal{Y}_{\mathcal{M}^{ass}(M)} = \int_A \langle [\alpha, \beta], \gamma \rangle \Omega.$

Table: Natural geometric structures on the moduli space of associative cycles on  $M$ . Here  $\alpha, \beta, \gamma, \delta \in \Omega^1(A, ad(E))$  and  $\phi, \eta, \xi, \zeta \in Ker D$ .

In the definition of the four form, we use the fact that the twisted spinor bundle  $S$  is a complex rank two bundle and therefore, as a real vector bundle  $S_{\mathbb{R}}$ , there is a natural trivialization of  $\Lambda^4 S_{\mathbb{R}} \cong \mathbb{R}$ . The action of  $\text{Im } \mathbb{H}$  on  $\mathbb{H}$  has used to define 3 and 4-form.

### An example

In this example, the above three form  $\Omega_{\mathcal{M}^{ass}(M)}$  defines a  $G_2$  structure on the moduli space  $\mathcal{M}^{ass}(M)$ . We consider a flat example  $M = T^3 \times T^4$  with  $T^3 = \mathbb{R}^3/\Lambda_3$  and  $T^4 = \mathbb{R}^4/\Lambda_4$ . The projection to the second factor  $M \rightarrow B = T^4$  is an associative fibration. The moduli space  $\mathcal{M}^{ass}(M)$  of  $(A, D_E)$  with  $A$  a fiber and  $D_E$  a flat  $U(1)$ -connection over  $A$ , can be identified with  $T^{3*} \times T^4$  where  $T^{3*} = \mathbb{R}^{3*}/\Lambda_3^*$  is the dual torus to  $T^3$ . Furthermore  $\Omega_{\mathcal{M}^{ass}(M)}$  and  $\Theta_{\mathcal{M}^{ass}(M)}$  are precisely the natural calibration three and four forms on the flat  $G_2$ -manifold  $T^{3*} \times T^4$ . This is a simple but fun exercise.

In particular,  $\chi_{\mathcal{M}^{ass}(T^3 \times T^4)} = \chi_{\mathcal{M}^{ass}(T^{3*} \times T^4)}$  and therefore,  $T^{3*} \times T^4 \rightarrow T^4$  defines an associative fibration structure on  $\mathcal{M}^{ass}(T^3 \times T^4) \rightarrow \mathcal{B}^{ass}(T^3 \times T^4)$  with a coassociative section, namely,  $\Omega_{\mathcal{M}^{ass}(T^3 \times T^4)}$  (resp.  $\chi_{\mathcal{M}^{ass}(T^3 \times T^4)}$ ) vanishes on the section (resp. fibers) of this fibration.

In the next paragraph, we will explain that such structures exist on every  $\mathcal{M}^{ass}(M)$ . However, when  $M$  is not flat,  $\Omega_{\mathcal{M}^{ass}(M)}$  and  $\Theta_{\mathcal{M}^{ass}(M)}$  are not parallel forms, thus they do not define  $G_2$ -structure on  $\mathcal{M}^{ass}(M)$  and we can only discuss their (co-)associativity in terms of vanishing of appropriate tensors.

### Moduli space of associative submanifolds is coassociative

Every associative submanifold  $A$  defines an associative pair  $(A, D_E)$  where  $D_E$  is simply the trivial connection  $d$ . Thus we have an embedding of  $\mathcal{B}^{ass}(M)$  inside  $\mathcal{M}^{ass}(M)$ . It is not difficult to see that the restriction of  $\Omega_{\mathcal{M}^{ass}(M)}$  to it is zero. As we will see in the next subsection, this property characterizes a coassociative submanifold is  $\Omega_{\mathcal{M}^{ass}(M)}$  defines a  $G_2$  structure on  $\mathcal{M}^{ass}(M)$ . Furthermore we have a natural fibration structure,  $\mathcal{M}^{ass}(M) \rightarrow \mathcal{B}^{ass}(M)$  by forgetting the connections. Fibers are associative submanifolds in the sense that the restriction  $\chi_{\mathcal{M}^{ass}(M)}$  to them are zero. Here  $\chi_{\mathcal{M}^{ass}(M)}$  is defined the same way as  $\chi$  for  $M$ . The reason is fiber directions for  $\mathcal{M}^{ass}(M) \rightarrow \mathcal{B}^{ass}(M)$  corresponds to  $H^1(A, ad(E))$  and each component in  $\Theta_{\mathcal{M}^{ass}(M)}$  involves at least 2  $Ker D$  components in its variables. Therefore  $\chi_{\mathcal{M}^{ass}}$  must vanish on fibers.

## 2.3 Coassociative cycles

### Coassociative submanifolds

A four dimensional submanifold  $C$  in  $M$  is called a *coassociative submanifold* if it is calibrated by  $\Theta$ , i.e.  $\Theta|_C = \text{vol}_C$ . It can be characterized by

$$\Omega|_C = 0 \in \Omega^3(C, \mathbb{R}).$$

Another characterization is the vector cross product on  $M$  preserves  $N_{C/M} \subset TM$ . As a corollary of this, any two coassociative submanifolds in  $M$  can not intersect along a three dimension subspace.

The normal bundle of  $C$  in  $M$  can be identified with the bundle of anti-self-dual two forms on  $C$ ,

$$N_{C/M} = \Lambda_+^2(C).$$

Furthermore the space of infinitesimal deformations of  $C$  inside  $M$  as a coassociative submanifold is isomorphic to the space of self-dual harmonic two forms on  $C$ ,  $H_+^2(C, \mathbb{R})$ . In fact the deformation theory of coassociative submanifold is unobstructed [15] and therefore their moduli space, denoted  $\mathcal{B}^{coa}(M)$ , is always smooth.

### Coassociative cycles

Similar to the associative cases, we will also couple coassociative submanifolds with certain gauge fields. A *coassociative cycle* is a pair  $(C, D_E)$  with  $C$  a coassociative submanifold in  $M$  and  $E$  an anti-self-dual (ASD) connection on  $C$  with respect to the induced metric. Recall that a connection is called ASD if its curvature 2-form  $F_E$  lies in  $\Omega_-^2(C, \text{ad}(E))$ , i.e.  $F_E + *F_E = 0$ . The tangent space of their moduli space  $\mathcal{M}^{coa}(M)$  can be identified as  $H_+^2(C) + H^1(C, \text{ad}(E))$  [15]. We define a three form  $\Omega_{\mathcal{M}^{coa}(M)}$ , a four form  $\Theta_{\mathcal{M}^{coa}(M)}$  and a symmetric cubic tensor  $\mathcal{Y}_{\mathcal{M}^{coa}(M)}$  on  $\mathcal{M}^{coa}(M)$  as follows:

3-form:	$\Omega_{\mathcal{M}^{coa}(M)} = \begin{cases} \int_C [\phi, \eta] \wedge \xi \\ - \int_C \text{Tr}(\phi \wedge \alpha \wedge \beta) \end{cases}$
4-form:	$\Theta_{\mathcal{M}^{coa}(M)} = \begin{cases} - \int_C \text{Tr}(\alpha \wedge \beta \wedge \gamma \wedge \delta)_{skew} \\ \int_C \text{Tr}([\phi, \eta] \wedge \alpha \wedge \beta) \end{cases}$
Cubic:	$\mathcal{Y}_{\mathcal{M}^{coa}(M)} = \int_A \text{Tr} \phi [\alpha, \beta]_{\text{ad}(E)}.$

Table: Natural geometric structures on the moduli space of coassociative cycles on  $M$ . Here  $\alpha, \beta, \gamma, \delta \in \Omega^1(C, \text{ad}(E))$  and  $\phi, \eta, \xi \in H_+^2(C)$ .

Here, Lie algebra structure on space of 2-forms is given by  $\wedge^2 \mathbb{R}^4 \simeq \mathfrak{so}(\mathbb{R}^4)$ .

### An example

In this example, the above three form  $\Omega_{\mathcal{M}^{coa}(M)}$  defines a  $G_2$  structure on the moduli space  $\mathcal{M}^{coa}(M)$ . We consider a flat example  $M = T^3 \times T^4$  with

$T^3 = \mathbb{R}^3/\Lambda_3$  and  $T^4 = \mathbb{R}^4/\Lambda_4$ . The projection to the first factor  $M \rightarrow B = T^3$  is a coassociative fibration. The moduli space  $\mathcal{M}^{coa}$  of  $(C, D_E)$  with  $C$  a fiber and  $D_E$  a flat  $U(1)$ -connection over  $C$ , can be identified with  $T^3 \times T^{4*}$  where  $T^{4*} = \mathbb{R}^{4*}/\Lambda_4^*$  is the dual torus to  $T^4$ . Moreover  $\Omega_{\mathcal{M}^{coa}}$  and  $\Theta_{\mathcal{M}^{coa}}$  are precisely the natural calibration three and four form on the flat  $G_2$ -manifold  $T^3 \times T^{4*}$ . This is another simple but fun exercise.

### Moduli space of coassociative submanifolds is associative

The fibration  $\mathcal{M}^{coa}(M) \rightarrow \mathcal{B}^{coa}(M)$  is a 'coassociative fibration' and the section embeds  $\mathcal{B}^{coa}(M)$  inside  $\mathcal{M}^{coa}(M)$  behaves as an 'associative' submanifold in an appropriate sense. The arguments are parallel to the case for  $\mathcal{B}^{ass}(M)$  in  $\mathcal{M}^{ass}(M)$  as we discussed in section 2.2 and hence omitted.

## 2.4 Reductions to Calabi-Yau threefolds

In this section we assume  $M = X \times S^1$  with  $X$  a Calabi-Yau threefold and

$$\Omega = \text{Re } \Omega_X - \omega \wedge dt,$$

where  $\Omega_X$  and  $\omega_X$  are the holomorphic three form and the Kähler form on  $X$  respectively. We will see that every moduli space of cycles (with  $U(1)$ -connections) on  $M$  and its natural differential forms inherit similar decompositions. That is,

$\mathcal{M}(M) = \mathcal{M}(X) \times S^1,$
$\Omega_{\mathcal{M}(M)} = \text{Re } \Omega_{\mathcal{M}(X)} - \omega_{\mathcal{M}(X)} \wedge dt.$

The manifold  $M$  itself can be regarded as the moduli space of zero dimensional cycles in  $M$ .

Remark : The reduction from a  $G_2$ -manifold  $M$  to a Calabi-Yau threefold  $X$  has a physical significance. Namely we reduce the studies of M-theory on an eleven dimensional space time  $\mathbb{R}^{3,1} \times M$  to the string theory on ten dimensional space time  $\mathbb{R}^{3,1} \times X$ .

### On the moduli of deformed DT bundles

The *deformed Hermitian Yang-Mills* connection on a bundle  $E$  on  $X$  is introduced in [14] as a supersymmetric cycle. Its curvature tensor satisfies the following equation,

$$\begin{aligned} F \wedge \omega^2 &= F^3/3, \\ F^{0,2} &= 0. \end{aligned}$$

Their moduli space  $\mathcal{M}^{bdl}(X)$  has a natural holomorphic three form  $\Omega_{\mathcal{M}^{bdl}(X)}$  and an almost symplectic form  $\omega_{\mathcal{M}^{bdl}(X)}$ ,

$$\begin{aligned}\Omega_{\mathcal{M}^{bdl}(X)}(\alpha, \beta, \gamma) &= \int_X \alpha \wedge \beta \wedge \gamma \wedge \Omega_X \\ \omega_{\mathcal{M}^{bdl}(X)}(\alpha, \beta) &= \int_X \alpha \wedge \bar{\beta} \wedge e^{\omega + F_E}.\end{aligned}$$

Here  $\alpha, \beta, \gamma \in H^1(X, O_X) = H^{0,1}(X)$ , the tangent space of  $\mathcal{M}^{bdl}(M)$  at  $D_E$ .

The pullback of any deformed Hermitian Yang-Mills connection  $D_E$  on  $X$  is a deformed DT connection on  $M$ . This is because the condition  $F^{0,2} = 0$  for the holomorphicity of  $E$  on a threefold is equivalent to  $F \wedge \text{Re} \Omega_X = 0$  in  $\Omega^5(X)$ . We can obtain other elements in  $\mathcal{M}^{bdl}(M)$  by tensoring with flat  $U(1)$ -connection on  $S^1$ . The moduli space of flat connections on  $S^1$  is the dual circle, also denoted as  $S^1$ . It is not difficult to verify the above mentioned decompositions for  $\mathcal{M}^{bdl}(M)$  and  $\Omega_{\mathcal{M}^{bdl}(M)}$  in this case.

### On the moduli of associative cycles

Suppose that an associative submanifold  $A$  in  $M = X \times S^1$  is of product types, then

Case (1): $A = \Sigma \times S^1$	$\Sigma$ holomorphic curve in $X$ .
Case (2): $A = L \times \{t\}$	$L$ special Lagrangian with phase = 0.

Case (1):  $A = \Sigma \times S^1$ . We denote the moduli space of holomorphic curves  $\Sigma$  in  $X$  as  $\mathcal{B}^{cx}(X)$ . Similarly we consider the moduli space of pairs  $(\Sigma, D_E)$  where  $D_E$  is a flat  $U(1)$ -connection on  $\Sigma$ , and denote it as  $\mathcal{M}^{cx}(X)$ . Gopakumar and Vafa conjecture that the cohomology group of  $\mathcal{M}^{cx}(X)$  admits an  $\mathfrak{sl}(2) \times \mathfrak{sl}(2)$  action whose multiplicities determine Gromov-Witten invariants of  $X$  of every genus for the class  $[\Sigma]$ . The tangent space of  $\mathcal{M}^{cx}(X)$  at  $(\Sigma, D_E)$  equals  $H^0(\Sigma, N_{\Sigma/X}) \oplus H^1(\Sigma, O_\Sigma)$ . It carries natural holomorphic three form  $\Omega_{\mathcal{M}^{cx}(X)}$  and symplectic form  $\omega_{\mathcal{M}^{cx}(X)}$  as follow,

holomorphic 3 form:	$\Omega_{\mathcal{M}^{cx}(X)} = \int_\Sigma \widetilde{(\phi \wedge \eta)} \wedge \alpha,$
symplectic form:	$\omega_{\mathcal{M}^{cx}(X)} = \int_\Sigma \alpha \wedge \bar{\beta} \pm \langle \phi, \eta \rangle \omega_X.$

Here  $\alpha, \beta \in H^1(\Sigma, O_\Sigma)$  and  $\phi, \eta \in H^0(\Sigma, N_{\Sigma/X})$  and  $\widetilde{(\phi \wedge \eta)} \in T_\Sigma^*$  is the image  $\phi \wedge \eta \in \Lambda^2 N_{\Sigma/X}$  under the natural identification  $\Lambda^2 N_{\Sigma/X} \cong T_\Sigma^*$  because of  $\Lambda^3 T_X \cong O_X$ .

Note that any deformation of  $A = \Sigma \times S^1$  as an associative submanifold in  $M$  must still be of the same form, i.e.  $\mathcal{B}^{ass}(M) = \mathcal{B}^{cx}(X)$ . However there are more flat  $U(1)$  bundles on  $A$  than on  $\Sigma$ , i.e.  $H^1(A, \mathbb{R}/\mathbb{Z}) \cong H^1(\Sigma, \mathbb{R}/\mathbb{Z}) \times S^1$ . We have the decompositions for  $\mathcal{M}^{coa}(M)$  and  $\Omega_{\mathcal{M}^{coa}(M)}$ .

Case (2):  $A = L \times \{t\}$  with  $L$  a special Lagrangian submanifold of zero phase in  $X$ , i.e. calibrated by  $\text{Re } \Omega$ .

We denote the moduli space of  $L$  (resp.  $(L, D_E)$  with  $D_E$  a flat  $U(1)$ -connection on  $L$ ) in  $X$  as  $\mathcal{B}^{cx}(X)$  (resp.  $\mathcal{M}^{SL}(X)$ ). McLean [15] introduces a three form on  $\mathcal{B}^{cx}(X)$ . On  $\mathcal{M}^{SL}(X)$ , there are a holomorphic three form  $\Omega_{\mathcal{M}^{SL}(X)}$  and a symplectic form  $\omega_{\mathcal{M}^{SL}(X)}$  as follow

holomorphic 3 form:	$\Omega_{\mathcal{M}^{SL}(X)} = \int_L \alpha \wedge \beta \wedge \gamma,$
symplectic form:	$\omega_{\mathcal{M}^{SL}(X)} = \int_L \langle \alpha, J\beta \rangle \text{Re } \Omega_X.$

Here  $\alpha, \beta, \gamma \in H^1(L, \mathbb{C})$ , the tangent space of  $\mathcal{M}^{SL}(X)$  at  $(L, D_E)$  ([15]).

It is not difficult to verify that  $\mathcal{B}^{ass}(M) = \mathcal{B}^{SL}(X) \times S^1$  and the decomposition for  $\mathcal{M}^{ass}(M)$  and  $\Omega_{\mathcal{M}^{ass}(M)}$ .

### On the moduli of coassociative cycles

Suppose  $C$  is a coassociative submanifold of product type in  $M = X \times S^1$  then

Case (1): $C = L \times S^1$	$L$ special Lagrangian with phase $= \pi/2$
Case (2): $C = S \times \{t\}$	$S$ complex surface in $X$ .

Case (1):  $C = L \times S^1$ . We can argue as in the previous situation to obtain decompositions for  $\mathcal{M}^{coa}(M)$  and  $\Omega_{\mathcal{M}^{coa}(M)}$ .

Case (2):  $C = S \times \{t\}$ . We denote the moduli space of complex surfaces  $S$  (resp.  $(S, D_E)$  with  $D_E$  a flat  $U(1)$ -connection on  $S$ ) in  $X$  as  $\mathcal{B}^{cx}(X)$  (resp.  $\mathcal{M}^{cx}(X)$ ). The tangent space of  $\mathcal{M}^{cx}(X)$  can be identified as  $H^{2,0}(S) \oplus H^{0,1}(S)$  because of  $K_X = O_X$ . We have natural differential forms on  $\mathcal{M}^{cx}(X)$ :

holomorphic 3 form:	$\Omega_{\mathcal{M}^{cx}(X)} = \int_S \phi \wedge \alpha \wedge \beta,$
symplectic form:	$\omega_{\mathcal{M}^{cx}(X)} = \int_S (\langle \phi, J\eta \rangle + \langle \alpha, J\beta \rangle) \omega^2/2.$

Here  $\phi, \eta \in H^{2,0}(S)$  and  $\alpha, \beta \in H^{0,1}(S)$ .

Again we can verify the decompositions for  $\mathcal{M}^{coa}(M)$  and  $\Omega_{\mathcal{M}^{coa}(M)}$ .

### 3 Fibrations and Fourier transforms

We review briefly the Strominger-Yau-Zaslow's construction [18] of the conjectural mirror manifold of a Calabi-Yau threefold: If  $X$  is close to a large complex and Kähler structure limit point, then it should admit a special Lagrangian fibration,

$$\pi_X : X \rightarrow B,$$

calibrated by  $\text{Im } \Omega_X$ . Moreover there is a special Lagrangian section, calibrated by  $\text{Re } \Omega_X$ . The mirror manifold is conjectured to be the dual torus fibration over  $B$ . Moreover the fiberwise Fourier transformation is expected to play an important role in the mirror transformation of the complex (resp. symplectic) geometry of  $X$  to the symplectic (resp. complex) geometry of its mirror manifold.

In this section we study calibrated fibrations on  $G_2$ -manifolds and Fourier transformations along them.

#### 3.1 Calibrated fibrations

##### Coassociative fibrations

Starting with a special Lagrangian fibration with a section on  $X$ , the induced fibration on the  $G_2$ -manifold  $M = X \times S^1$ ,

$$\pi : M \rightarrow B,$$

becomes a coassociative  $T^4$ -fibration with an associative section.

Recall that a general fiber in a Lagrangian fibration on any symplectic manifold is necessarily a torus. This is not true for a general fiber in a coassociative fibration on a  $G_2$ -manifold. For example, given any holomorphic fibration on a Calabi-Yau threefold,

$$p_X : X \rightarrow \mathbb{CP}^1 \cong S^2,$$

its general fiber is necessarily a K3 surface. The induced fibration on the  $G_2$ -manifold  $M = X \times S^1$  over  $S^2 \times S^1$  is again a coassociative fibration, which is now a K3 fibration.

We expect that a general fiber in a coassociative fibration is always a torus or a K3 surface<sup>3</sup>. This is indeed the case provided that a general fiber admits an Einstein metric: The normal bundle of a coassociative submanifold  $F$  is always the bundle of self-dual two forms on  $F$ . When  $F$  is a fiber of  $\pi$  then the normal bundle is necessarily trivial, i.e.

$$\Lambda_+^2(F) = F \times \mathbb{R}^3.$$

This implies that

$$3\tau(F) + 2\chi(F) = 0,$$

where  $\tau(F)$  and  $\chi(F)$  are the signature and the Euler characteristic of  $F$  respectively.

---

<sup>3</sup>We learn this from S.-T. Yau.



From physical considerations we expect that there should be a family of  $G_2$ -metrics together with coassociative fibrations parametrized by  $(t_0, \infty)$  such that the limit as  $t \rightarrow \infty$  would imply the second fundamental form of each smooth fiber would go to zero. Combining with the fact the Ricci tensor of a  $G_2$ -metric is trivial, this suggest that every smooth fiber should admit a Einstein metric. However characteristic numbers for  $F$  saturate the Hitchin inequality for Einstein four manifolds, this implies that  $F$  is either flat or covered by a K3 surface.

### Associative fibrations

Suppose  $M$  has an associative fibration,

$$\pi : M \rightarrow B.$$

As in the previous case, we expect a general fiber to admit a metric with zero Ricci curvature in the adiabatic limit, thus it must be a three torus  $T^3$ .

### Conjectural characterizations

Calibrated fibrations on a Calabi-Yau threefold  $X$  can be either special Lagrangian  $T^3$ -fibrations or holomorphic fibrations. In the latter case a generic fiber can be an elliptic curve  $T^2$ , an Abelian surface  $T^4$  or a K3 surface. Wilson [20] studies such fibrations in terms of the ring structure on  $H^2(X)$  and the linear form,

$$\int_X p_1(X) \cup (\cdot) : H^2(X, \mathbb{R}) \rightarrow \mathbb{R}.$$

Notice that such linear form is trivial on  $H^3(X, \mathbb{R})$ .

We conjecture that similar characterizations should hold true for  $G_2$ -manifolds. To be precise, let  $\Omega_t$  be a family of  $G_2$ -structures on  $M$  for  $t \in (0, 1)$  such that as  $t \rightarrow 0$  the volume of  $M$  goes to zero while the diameter remains constant. For small  $t$ ,  $M$  should admit a calibrated fibration, the structure of this fibration should be determined by the Yukawa coupling  $\mathcal{Y}$  and  $p_1(M)$  as follows,

On $M$ ,	Characterizations:
Associative $T^3$ -fibration	$\mathcal{Y}(\Omega_t, \Omega_t, H^3) \rightarrow 0$
Coassociative $T^4$ -fibration	$\mathcal{Y}(\Omega_t, \Omega_t, H^3) \rightarrow 0$ & $\int p_1 \cup \Omega_t \rightarrow 0$
Coassociative K3-fibration	$\mathcal{Y}(\Omega_t, \Omega_t, H^3) \rightarrow 0$ & $\int p_1 \cup \Omega_t \rightarrow 0$ .

It is also possible that  $\mathcal{Y}(\Omega_t, \Omega_t, H^3)$  becomes unbounded, but this should only happen when  $\tilde{M} = X \times \mathbb{R}$ .

Remark: We also expect that in above situations,  $\lim_{t \rightarrow 0} \Omega_t$  is of infinite distance from any point in the moduli space of  $G_2$ -metrics on  $M$ . On the other hand finite distance boundary points in this moduli space (i.e. incompleteness) might be related to contractions of (i) an ADE configuration of associative  $S^3$ 's or (ii) an associative family of ADE configuration of  $S^2$ 's and so on. C.L. Wang further conjectures that such incompleteness of the moduli space should be equivalent to degenerating families of  $G_2$ -structures on  $M$  with both the volume and the diameter bounded away from zero and infinity uniformly.

### 3.2 Analog of the SYZ conjecture

We are going to propose an analog of the SYZ conjecture for  $G_2$ -manifolds, namely a transformation of  $G_2$ -geometry along a calibrated fibration. Acharya [1] is the first to use these  $T^4$ - or  $T^3$ -fibrations to study the mirror symmetry for  $G_2$ -manifolds from a physics point of view.

#### The $G_2$ mirror conjecture

There is a *reasonably* large class of  $G_2$ -manifolds such that every such  $M$  satisfies the followings:

(1) *Mirror pair*: There is a family of  $G_2$ -metric  $g_t$  together with a coassociative  $T^4$ -fibration and an associative section on  $M$ . As  $t$  goes to infinity the diameter and the curvature tensor of each fiber go to zero. The moduli space  $\mathcal{M}^{coa}(M)$  of coassociative cycles  $(C, D_E)$  in  $M$  when  $C$  is a fiber admits a suitable compactification  $W_t$  as a  $G_2$ -manifold with  $\Omega_{\mathcal{M}^{coa}(M)}$  the calibrating three form (after suitable instanton corrections from associative cycles). The relation between  $M$  and  $W$  is involutive.

(2) *Yukawa couplings*: The Yukawa couplings  $\mathcal{Y}_M$  and  $\mathcal{Y}_W$  for  $M$  and  $W$  on  $H_{27}^3(M)$  and  $H_{27}^3(W)$  should be naturally identified after suitable corrections coming from associative cycles in  $M$  and  $W$ .

(3) *Coassociative geometry*: There is an identification between moduli spaces of coassociative pairs or points in  $M$  and moduli spaces of corresponding objects in  $W$ .

(4) *Associative geometry*: There is an identification between moduli spaces of associative pairs or deformed Donaldson-Thomas bundles in  $M$  and moduli spaces of corresponding objects in  $W$ .

The mirror symmetry conjecture for  $G_2$ -manifolds can be summarized in the following table:

Coassociative geometry on $M$ $\Leftrightarrow$ Associative geometry on $W$ ,
Associative geometry on $M$ $\Leftrightarrow$ Coassociative geometry on $W$ .

### 3.3 Fourier transform on coassociative $T^4$ -fibrations

In this subsection we verify various parts of the above conjecture for *flat*  $G_2$ -manifolds, i.e.  $M = B \times T$  with  $B = \mathbb{R}^3$  and  $T = \mathbb{R}^4/\Lambda$  a flat torus. The projection map

$$\pi : M \rightarrow B,$$

is a coassociative  $T^4$ -fibration on  $M$ . The large structure limit on  $M$  can be obtained by rescaling  $\Lambda$ . If we denote  $\mathcal{M}^{coa}(M)$  the moduli space of coassociative cycles  $(C, D_E)$  on  $M$  with  $[C]$  representing a fiber class and  $D_E$  is a flat  $U(1)$ -connection on  $C$ , then it can be naturally identified with the total space  $W = B \times T^*$  of the dual torus fibration

$$\pi' : W \rightarrow B.$$

Moreover, under this identification, the canonical three form and four form on the moduli space correspond to the calibrating three form and four form for the  $G_2$ -structure on  $W$ :

$$\begin{aligned}\Omega_W &= \Omega_{\mathcal{M}^{coa}(M)}, \\ \Theta_W &= \Theta_{\mathcal{M}^{coa}(M)}.\end{aligned}$$

We are going to perform fiberwise Fourier transformation on  $M$ . As usual we denote the coordinates on  $B$  (resp.  $T$  and  $T^*$ ) as  $x^1, x^2, x^3$  (resp.  $y^0, \dots, y^3$  and  $y_0, \dots, y_3$ ). The dual torus  $T^*$  is treated as the moduli space of flat  $U(1)$  connections on  $T$ . The universal line bundle over  $T \times T^*$  is called the Poincaré bundle  $P$ . We normalize it so that its restriction to both  $T \times 0$  and  $0 \times T^*$  are trivial. It has a universal connection whose curvature two form equals

$$\mathbb{F} = \sum_{j=0}^3 dy^j \wedge dy_j \in \Omega^2(T \times T^*).$$

The Fourier transformation, or the Fourier-Mukai transformation, from  $M$  to  $W$  can be described roughly as follows:

$$\mathbf{F}(\bullet) = p'_*(p^*(\bullet) \otimes P),$$

where  $p$  and  $p'$  is defined by the following commutative diagram,

$$\begin{array}{ccc} M \times_B W & \xrightarrow{p'} & W \\ p \downarrow & & \downarrow \pi' \\ M & \xrightarrow{\pi} & B. \end{array}$$

and  $p^*$  is the pull back and  $p'_*$  is the push forward operator or equivalently integration along fibers of  $p' : M \times_B W \rightarrow W$ . We can discuss this transformation on different levels, including differential forms, cohomology, K-theory and so on. For differential forms, we have

$$\begin{aligned}\mathbf{F} : \Omega^*(M) &\longrightarrow \Omega^*(W) \\ \mathbf{F}(\varphi) &= \int_{M/B} p^*(\varphi) \wedge e^{\frac{\sqrt{-1}}{2\pi} \mathbb{F}}\end{aligned}$$

where  $\mathbb{F}$  is the universal curvature 2-form on the Poincaré bundle  $P$ . Moreover  $\varphi$  could be a differential form supported on a submanifold in  $M$ , and the same for  $\mathbf{F}(\varphi)$ , too. For example the above Fourier transformation preserves the calibrating three form and four form, more precisely we have,

$\mathbf{F}(e^{\Theta_M}) = e^{\Theta_W},$
$\mathbf{F}(*e^{\Theta_M}) = *e^{\Theta_W}.$

These follow from the following integrals with orientation  $y^{0123}$ ,

$$\int_{M/B} e^{\Theta_M} \wedge e^{\mathbb{F}} = e^{\Theta_W} \text{ and } \int_{M/B} (*e^{\Theta_M}) \wedge e^{\mathbb{F}} = (*e^{\Theta_W}).$$

Similarly, on the level of cohomology classes, we have,

$$\begin{aligned} \mathcal{F} : H^*(M) &\longrightarrow H^*(W) \\ \mathcal{F}(\varphi) &= \int_{M/B} p^*([\varphi]) \cup ch(P). \end{aligned}$$

We can also transform a connection  $D_E$  on a bundle  $E$  over a submanifold  $C$  in  $M$  to one in  $W$ . To do this, we consider tensor product of the pullback connection on the pullback bundle  $p^*E \longrightarrow C \times_B W$  with the universal connection on the Poincaré bundle restricted to  $C \times_B W$ . Under good circumstances, the pushforward sheaf under  $p' : C \times_B W \rightarrow W$  gives a vector bundle over a submanifold in  $W$ . For instance, a flat  $U(1)$  connection over  $T^4 \times \{p\}$  in  $M = T^4 \times T^3$  is transformed to a point in  $W = T^{4*} \times T^3$ , as discussed earlier.

## Transforming coassociative cycles

### Transforming ASD connections on $T^4$ fibers

Suppose that we relax the flatness assumption on  $D_E$  and we assume that it is an ASD connection of a torus fiber  $C = T$ . The moduli space of such coassociative cycles  $(C, D_E)$  has the form,

$$\mathcal{M}^{coa}(M) = B \times \mathcal{M}^{ASD}(T),$$

where  $\mathcal{M}^{ASD}(T)$  denotes the moduli space of ASD connections on the flat torus  $T$ .

When  $D_E$  has no flat factor, the Fourier transformation of an ASD connection over a flat torus  $T$  is studied by Schenk [17], Braam and van Baal [5] and

it produces another ASD connection over the dual torus  $T^*$ . Moreover it is an isometry from  $\mathcal{M}^{ASD}(T)$  to  $\mathcal{M}^{ASD}(T^*)$ . Using their results we have,

$$\mathbf{F} : \mathcal{M}^{coa}(M) \xrightarrow{\sim} \mathcal{M}^{coa}(W).$$

Moreover it is not difficult to identify the natural three forms  $\Omega_{\mathcal{M}^{coa}}$  on these moduli spaces under  $\mathbf{F}$ .

### Describing other coassociative cycles in $M$

Suppose  $C$  is any coassociative submanifold in  $M$  which is not a fiber. The intersection of  $C$  with a general fiber can not have dimension 3 (see section 2.3). The intersection can not have dimension 1 neither because otherwise the normal bundle of  $C$  in  $M$ , which is associative, lies inside the relative tangent bundle of the coassociative fibration, a contradiction. Therefore the intersection of  $C$  with a general fiber must have dimension 2.

We assume that the coassociative submanifold  $C$  in  $M$  is *semi-flat* in the sense that it intersects each fiber along a flat subtorus. As subtori in  $T^4$ , they are all translations of each other. Using the fact that the vector cross product on  $M$  preserves the normal directions of  $C$ , one can show that the image of  $C$  in  $B \cong \mathbb{R}^3$  must be an affine plane. Therefore, up to coordinate changes, we can assume that  $C$  is of the form

$$C : \begin{cases} x^3 = 0, \\ y^0 = B^0(x^1, x^2), \\ y^3 = B^3(x^1, x^2), \end{cases}$$

for some smooth functions  $B^0$  and  $B^3$  in variables  $x^1$  and  $x^2$  satisfying

$$-\frac{\partial}{\partial x^2} B^3 + \frac{\partial}{\partial x^1} B^0 = \frac{\partial}{\partial x^1} B^3 + \frac{\partial}{\partial x^2} B^0 = 0.$$

Suppose  $D_E$  is an ASD connection on  $C$  which is *semi-flat* in the sense that  $D_E$  is invariant under translations along fiber directions, we write

$$D_E = d + a_1(x^1, x^2) dx^1 + a_2(x^1, x^2) dx^2 + D_1(x^1, x^2) dy^1 + D_2(x^1, x^2) dy^2,$$

and the ASD condition is equivalent to

$$\begin{aligned} \frac{\partial}{\partial x^2} a_1 - \frac{\partial}{\partial x^1} a_2 &= 0, \\ -\frac{\partial}{\partial x^2} D_1 + \frac{\partial}{\partial x^1} D_2 &= \frac{\partial}{\partial x^1} D_1 + \frac{\partial}{\partial x^2} D_2 = 0. \end{aligned}$$

On each  $T^4$  fiber of  $\pi : M \rightarrow B$  over a point  $(x^1, x^2, 0) \in B$ , we obtain a flat 2-torus defined by

$$\{(y^0, y^1, y^2, y^3) \in T^4 \mid y^0 = B^0(x^1, x^2), y^3 = B^3(x^1, x^2)\}$$

together with a flat  $U(1)$  connection  $D_E = d + D_1(x^1, x^2) dy^1 + D_2(x^1, x^2) dy^2$  on it, because  $D_1(x^1, x^2)$  and  $D_2(x^1, x^2)$  are constant on each fiber.

### Transforming coassociative families of subtori

Applying the Fourier transformation from this  $T^4$  fiber to  $T^{4*}$ , it is not difficult to check that the above flat connection ( resp. flat 2-torus ) in  $T^4$  will be transformed to a flat 2-torus ( resp. flat connection ) in  $T^{4*}$  ( see [13] for more details ). Putting these together over various fibers of  $\pi : M \rightarrow B$ , we obtain the Fourier transformation of  $(C, D_E)$  in  $M$ , which is the pair  $(C', D'_E)$  on  $W$  with

$$C' : \begin{cases} x^3 = 0, \\ y_1 = D_1(x^1, x^2), \\ y_2 = D_2(x^1, x^2). \end{cases}$$

and

$$D'_E = d + a_1(x^1, x^2) dx^1 + a_2(x^1, x^2) dx^2 + B^0(x^1, x^2) dy_0 + B^3(x^1, x^2) dy_3.$$

$(C', D'_E)$  is again a semi-flat coassociative cycle in  $W$  and notice that the ASD connection for  $D_E$  (resp. coassociative condition for  $C$ ) implies the coassociative condition for  $C$  (resp. ASD condition for  $D'_E$ ). Furthermore the Fourier transformation induces a bijection on the semi-flat part of the moduli spaces,

$$\mathbf{F} : \mathcal{M}^{coa}(M)_{semi\flat} \rightarrow \mathcal{M}^{coa}(W)_{semi\flat}.$$

It is easy to check that when  $C$  is horizontal, i.e.  $B^0 = B^3 = 0$ , then  $\mathbf{F}$  preserves the natural three forms  $\Omega_{\mathcal{M}^{coa}}$  on these moduli spaces and we expect it continues to hold in general.

## Transforming associative cycles and DT bundles

### Describing associative cycles in $M$

Suppose  $A$  is any associative submanifold in  $M$  then it can only intersect a general fiber in dimension 2 or 0. In the former case, we can argue as before, the image of  $A$  in  $B \cong \mathbb{R}^3$  must be an affine line provided that  $A$  is *semi-flat*. We can write, up to coordinate changes,

$$A : \begin{cases} x^2 = x^3 = 0, \\ y^2 = B^2(x^1), \\ y^3 = B^3(x^1). \end{cases}$$

The associative condition on  $A$  implies that

$$\frac{\partial}{\partial x^1} B^2 = \frac{\partial}{\partial x^1} B^3 = 0.$$

That is, both  $B^2(x^1)$  and  $B^3(x^1)$  are constant functions.

Suppose  $D_E$  is a unitary connection on  $A$  which is invariant under translations along fiber directions, we can write,

$$D_E = a(x^1) dx^1 + D^0(x^1) dy_0 + D^1(x^1) dy_1.$$

If  $(A, D_E)$  is an associative cycle, then the flatness of  $D_E$  is equivalent to

$$\frac{\partial}{\partial x^1} a = \frac{\partial}{\partial x^1} D^0 = \frac{\partial}{\partial x^1} D^1 = 0.$$

That is  $a, D^0$  and  $D^1$  are all constant functions.

### Transforming associative families of tori

As in the earlier situation, the Fourier transformation of a semi-flat cycle  $(A, D_E)$  on  $M$ , which is not necessarily associative, is a pair  $(A', D'_E)$  on  $W$  with

$$A' : \begin{cases} x^2 = 0, \\ x^3 = 0, \\ y_0 = D^0(x^1), \\ y_1 = D^1(x^1). \end{cases}$$

and

$$D'_E = a(x^1) dx^1 + B^2(x^1) dy_2 + B^3(x^1) dy_3$$

It is obvious that  $(C, D_E)$  is associative if and only if  $(C', D'_E)$  is associative.

Remark: Recall that associative cycles in  $M$  are critical points of the Chern-Simons functional. Using arguments in [13], we can show that the Fourier transformation preserves the Chern-Simons functional of semi-flat three cycles on  $M$  and  $W$  which are *not* necessarily associative.

### Transforming associative sections to deformed DT bundles

Any section of the (trivial) coassociative fibration  $\pi$  on  $M$  can be expressed as a function  $f$ ,

$$f : B \rightarrow T = \frac{\mathbb{H}}{\Lambda}.$$

Harvey and Lawson [8] show that the associative condition is equivalent to the following equation,

$$-\frac{\partial f}{\partial x^1} i - \frac{\partial f}{\partial x^2} j - \frac{\partial f}{\partial x^3} k = \frac{\partial f}{\partial x^1} \times \frac{\partial f}{\partial x^2} \times \frac{\partial f}{\partial x^3}.$$

If we write  $f = {}^t(f^0, f^1, f^2, f^3) \in \mathbb{R}^4/\Lambda$ ,  $I = {}^t(1, i, j, k)$  and  $f_{x^j}^i = \frac{\partial f^i}{\partial x^j}$ , then the above equation can be rewritten as follows,

$$\det(I, f_{x^1}, f_{x^2}, f_{x^3})_{4 \times 4} = (f_{x^1}^1 + f_{x^2}^2 + f_{x^3}^3)1 + (-f_{x^1}^0 + f_{x^2}^3 - f_{x^3}^2)i + (-f_{x^1}^3 - f_{x^2}^0 + f_{x^3}^1)j + (f_{x^1}^2 - f_{x^2}^1 - f_{x^3}^0)k.$$

If  $D_E$  is a unitary flat connection on  $A$ , we can write

$$D_E = d + a_1 dx^1 + a_2 dx^2 + a_3 dx^3$$

where  $a_k$ 's are matrix valued functions of  $(x^1, x^2, x^3)$ .

As before, or from [13], the Fourier transformation of  $(A, D_E)$  is a unitary connection  $D'_E$  on the whole manifold  $W$  given by

$$D'_E = d + a_1 dx^1 + a_2 dx^2 + a_3 dx^3 + f^0 dy_0 + f^1 dy_1 + f^2 dy_2 + f^3 dy_3.$$

We are going to show that  $D'_E$  satisfies the deformed DT equation, i.e.

$$F_{E'} \wedge \Theta_W + F_{E'}^3 / 6 = 0.$$

First we notice that the curvature of  $D'_E$  is given by

$$F_{E'} = \sum_{j=1}^3 \sum_{k=0}^3 f_{x^i}^k dx^i \wedge dy_k.$$

and  $a_j$ 's do not occur here because of the flatness of  $D_E$ . By direct but lengthy computations, we can show that the deformed DT equation for  $D'_E$  is reduced to the flatness of  $D_E$  together with the associativity of  $A$ . Therefore the Fourier transformation in this case gives,

$$\mathbf{F} : \mathcal{M}^{ass}(M) \rightarrow \mathcal{M}^{bdl}(W).$$

In fact this transformation  $\mathbf{F}$  preserves the Chern-Simons functionals (see section 2.1, 2.2) on  $M$  and  $W$ . Namely suppose  $(A, D_E)$  is any pair on  $M$  which is not necessarily an associative cycle, then we have

$$\int_{A \times [0,1]} Tr \left[ e^{\Theta_M + \tilde{F}} \right] = (const) \int_{W \times I} Tr \left[ e^{\Theta_W + \tilde{F}'} \right]$$

We expect that  $\mathbf{F}$  preserves the natural three forms on  $\mathcal{M}^{ass}(M)$  and  $\mathcal{M}^{bdl}(W)$ .

### 3.4 Fourier transform on associative $T^3$ -fibrations

Suppose  $M$  has an associative  $T^3$ -fibration,

$$\pi : M \rightarrow B.$$

The fiberwise Fourier transformation  $\mathbf{F}$  in this case should interchange the two types of geometry on  $G_2$ -manifolds,

$$\text{Associative Geometry on } M \xleftarrow{\mathbf{F}} \text{Coassociative Geometry on } W.$$

Here  $W$  is the dual three torus fibration to  $\pi$  on  $M$ , it can be identified as the moduli space  $\mathcal{M}^{ass}(M)$  of associative cycles  $(A, D_E)$  on  $M$  with  $D_E$  being a flat  $U(1)$ -connection over a fiber  $A$ .



In the *flat* case, we have  $M = T \times \mathbb{R}^4$  and  $\mathcal{M}^{ass}(M) \cong W = T^* \times B$ . Under these identifications we have,

$$\begin{aligned}\Omega_W &= \Omega_{\mathcal{M}^{ass}(M)}, \\ \Theta_W &= \Theta_{\mathcal{M}^{ass}(M)}.\end{aligned}$$

We can analyze semi-flat associative and coassociative cycles in  $M$  and their Fourier transformations as before. For example, semi-flat coassociative submanifolds in  $M$  are either sections or a family of affine two tori over an affine plane in  $B$ . Semi-flat associative submanifolds in  $M$  are either fibers or a family of affine two tori over an affine line in  $B$ .

Suppose  $(C, D_E)$  is a coassociative cycle in  $M$  with  $C$  being a section of the (trivial) associative fibration on  $M$ . The coassociative condition of  $C$  can be expressed in term of a differential equation for a function  $g = (g^1, g^2, g^3) : \mathbb{R}^4 \rightarrow T = \mathbb{R}^3/\Lambda$  (see [8]),

$$\nabla g^1 i + \nabla g^2 j + \nabla g^3 k = \nabla g^1 \times \nabla g^2 \times \nabla g^3.$$

If we write the ASD connection as

$$D_E = d + a_0 dy^0 + a_1 dy^1 + a_2 dy^2 + a_3 dy^3,$$

then the Fourier transformation of  $(C, D_E)$  will be the following connection  $D'_E$  on  $W$ ,

$$D'_E = d + a_0 dy^0 + a_1 dy^1 + a_2 dy^2 + a_3 dy^3 + g^1 dx_1 + g^2 dx_2 + g^3 dx_3$$

Again direct but lengthy computations show that the deformed DT equation on  $D'_E$  is reduced to the above coassociativity condition of  $C$  and the ASD equation for  $D_E$ .

## 4 $Spin(7)$ -geometry

There are only two kinds of exceptional holonomy groups for Riemannian manifolds, namely  $G_2$  for dimension 7 and  $Spin(7)$  in dimension 8. Both of them play important roles in M-theory. Almost all of our earlier discussions for  $G_2$ -geometry has a  $Spin(7)$  analog. As a result, we will only indicate what geometric structures we have in  $Spin(7)$ -geometry and leave out those derivations which are the same as in their  $G_2$ -counterparts.

An eight dimensional Riemannian manifold  $Z$  with  $Spin(7)$ -holonomy has a closed self-dual four form  $\Theta_Z$  which can be expressed as follows,

$$\begin{aligned}\Theta_Z = & -dy^{0123} - dx^{0123} - (dx^{10} + dx^{23})(dy^{10} + dy^{23}) \\ & - (dx^{20} + dx^{31})(dy^{20} + dy^{31}) - (dx^{30} + dx^{12})(dy^{30} + dy^{12}),\end{aligned}$$

when  $Z = \mathbb{R}^8$ . As before, we can decompose differential forms on  $Z$  into  $Spin(7)$  irreducible components (see e.g. [10]), for example  $\Lambda^4 = \Lambda_1^4 + \Lambda_7^4 + \Lambda_{27}^4 + \Lambda_{35}^4$ .

### Holonomy reduction

For example if  $M$  is a  $G_2$ -manifold, then  $Z = M \times S^1$  has a  $Spin(7)$ -metric with

$$\Theta_Z = \Omega_M \wedge dt - \Theta_M,$$

This corresponds to the natural inclusion  $G_2 \subset Spin(7)$ . We can also consider the reduction of holonomy to the subgroup  $Spin(6) \subset Spin(7)$ . Notice that  $Spin(6) = SU(4)$ , namely such a  $Z$  would be a Calabi-Yau fourfold and

$$\Theta_Z = -\frac{\omega_Z^2}{2} + \text{Re} \Omega_Z,$$

where  $\omega_Z$  and  $\Omega_Z$  are the Ricci flat Kähler form and the holomorphic volume form on  $Z$ . We can compare their decompositions of differential forms as follows,

$$\begin{aligned}\Lambda_{35}^4 &= \Lambda_{prim}^{3,1} + \Lambda_{prim}^{1,1} + \Lambda_{prim}^{1,3}, \\ \Lambda_{27}^4 &= \Lambda_{prim}^{2,2} + \Lambda^{0,2} + \Lambda^{0,4}.\end{aligned}$$

We can reduce the holonomy group to smaller subgroups of  $Spin(7)$  and give the following tables,

Holonomy	Manifold
$Spin(7)$	$Spin(7)$ -manifold
$Spin(6) = SU(4)$	Calabi-Yau 4-fold
$Spin(5) = Sp(2)$	Hyperkähler 4-fold
$Spin(4) = SU(2)^2$	$K3 \times K3$
$Spin(3) = SU(2)$	$K3 \times T^4$

## 4.1 *Spin* (7)-Analog of Yukawa coupling

### An analog of the Yukawa coupling

We consider the  $(3, 1)$ -tensor  $\chi_Z \in \Omega^3(Z, T_Z)$  constructed by raising an index of  $\Theta_Z$  using the metric tensor.

We define a symmetric quartic tensor  $\mathcal{Q}$  on  $H^4(Z, \mathbb{R})$  as follows,

$$\begin{aligned} \mathcal{Q} : \bigotimes^4 \Omega^4(Z, \mathbb{R}) &\rightarrow \mathbb{R} \\ \mathcal{Q}(\phi_1, \phi_2, \phi_3, \phi_4) &= \int_Z \Theta_Z \left( \hat{\phi}_1, \hat{\phi}_2, \hat{\phi}_3, \hat{\phi}_4 \right) \wedge \Theta_Z. \end{aligned}$$

Here  $\hat{\phi} = *(\phi \wedge \chi) \in \Omega^1(Z, T_Z)$ . Note that  $\hat{\phi}$  is zero if  $\phi \in \Omega_{27}^4(Z, \mathbb{R})$ . We define the *Yukawa coupling*  $\mathcal{Y}$  to be the restriction of  $\mathcal{Q}$  to  $H_{35}^4(Z, \mathbb{R})$ . We can also define a cubic form, a quadratic form and a linear form on  $H_{35}^4(Z, \mathbb{R})$  by evaluating  $\mathcal{Q}$  on one, two and three  $\Theta_Z$  before restricting it to  $H_{35}^4(Z, \mathbb{R})$ .

## 4.2 Deformed DT bundles and Cayley cycles

### Deformed DT bundles

As in the  $G_2$  case, we define a *deformed DT connection* to be a connection  $D_E$  over  $Z$  whose curvature tensor  $F_E$  satisfies

$$*F_E + \Theta_Z \wedge F_E + F_E^3/6 = 0,$$

or equivalently,

$$[e^{\Theta_Z + F_E + *F_E}]^{[6]} = 0.$$

Its moduli space  $\mathcal{M}^{ddl}(Z)$  carries a natural four form  $\Theta_{\mathcal{M}^{ddl}(Z)}$  defined by,

$$\Theta_{\mathcal{M}^{ddl}(Z)}(\alpha, \beta, \gamma, \delta) = \int_Z \text{Tr}_E [\alpha \wedge \beta \wedge \gamma \wedge \delta]_{skew} \wedge e^{\Theta_Z + F_E}.$$

with  $\alpha, \beta, \gamma, \delta \in H^1(Z, \text{ad}(E))$ , the first cohomology group of the elliptic complex,

$$0 \rightarrow \Omega^0(Z, \text{ad}(E)) \xrightarrow{D_E} \Omega^1(Z, \text{ad}(E)) \xrightarrow{\pi_7 \circ D_E} \Omega_7^2(Z, \text{ad}(E)) \rightarrow 0.$$

### Cayley cycles

A submanifold  $C$  in  $Z$  of dimension four is called a *Cayley submanifold* if it is calibrated by  $\Theta_Z$ , i.e. The restriction on  $\Theta_Z$  to  $C$  equals the volume form on  $C$  with respect to the induced metric ([8]). A *Cayley cycle* is defined to be any pair  $(C, D_E)$  with  $C$  a Cayley submanifold in  $Z$  and  $D_E$  is an ASD connection on a bundle  $E$  over  $C$ . The moduli space  $\mathcal{M}^{Cay}(Z)$  of Cayley cycles in  $Z$  also have a natural four form  $\Theta_{\mathcal{M}^{Cay}(Z)}$  defined by

$$\Theta_{\mathcal{M}^{Cay}(Z)} = \begin{cases} \int_C \text{Tr} \alpha \wedge \beta \wedge \gamma \wedge \delta \\ - \int_C \left\langle \phi, \text{Tr}(\widehat{\alpha \wedge \beta}) \cdot \eta \right\rangle \Theta_Z \\ - \int_C \det(\phi, \eta, \xi, \zeta) \Theta_Z \end{cases}$$

where  $\alpha, \beta, \gamma, \delta \in \Omega^1(C, ad(E))$  and  $\phi, \eta, \xi, \zeta \in Ker \mathbf{D}$  are harmonic spinors and

$$\widehat{\alpha \wedge \beta} = \alpha \wedge \beta + *(\alpha \wedge \beta) \in \Omega_+^2(C, ad(E)) = \text{Im}(\mathbb{H}) \otimes ad(E)$$

We leave it to our readers to verify that the Fourier transformation along Cayley  $T^4$ -fibration on a flat  $Spin(7)$ -manifold  $Z$  will transform a Cayley section cycle  $(C, D_E)$  on  $Z$  to a deformed DT bundle over the dual torus fibration. Transformations of certain non-section Cayley cycles on  $Z$  can also be analyzed as in the  $G_2$  case.

Acknowledgments: The second author is partially supported by NSF/DMS-0103355 and he thanks R. Bryant, R. Thomas, C.L. Wang, S.-T. Yau and E. Zaslow for insightful and valuable discussions. He is grateful to S.-T. Yau who brought us to the attention of this subject.

## References

- [1] B.S. Acharya, *On Mirror Symmetry for Manifolds of Exceptional Holonomy*, Nucl.Phys. **B524** 269 (1998).
- [2] M. Aganagic, C. Vafa, *Mirror symmetry and a  $G_2$  flop*, J. High Energy Phys. 2003, no. 5, 061, 16 pp.
- [3] M. Atiyah, J. Maldacena, C. Vafa, *An M-theory flop as a large  $N$  duality*, J. Math. Phys. 42 (2001), no. 7, 3209–3220.
- [4] M. Atiyah, E. Witten, *M-Theory Dynamics On A Manifold Of  $G_2$  Holonomy*, Adv. Theor. Math. Phys. 6 (2002), no. 1, 1–106.
- [5] P.J. Braam, P. van Baal, *Nahm's transformation for instantons*, Comm. Math. Phys., 122 (1989) pp. 267-280.
- [6] S. K. Donaldson and R. P. Thomas, *Gauge Theory in Higher Dimensions, The Geometric Universe* (Oxford, 1996), Oxford Univ. Press, Oxford (1998) pp. 31-47.
- [7] A. Gray, *Vector cross products on manifolds*, Trans. Amer. Math. Soc. 141 (1969) 465–504.
- [8] R. Harvey, B. Lawson, *Calibrated geometries*, Acta Math. 148 (1982), 47-157.
- [9] N. Hitchin, *The geometry of three-forms in six and seven dimensions*, J. Differential Geom. 55 (2000), no. 3, 547–576.
- [10] D. Joyce, *Compact manifolds with special holonomy*, Oxford Univ. Press, 2000.

- [11] N.C. Leung, *Mirror symmetry without corrections*, Comm. Anal. Geom. 13 (2005), no. 2, 287–331.
- [12] N.C. Leung, *Symplectic structures on gauge theory*, Comm. Math. Phys., 193 (1998) 47-67.
- [13] N.C. Leung, S.Y. Yau, E. Zaslow, *From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform*, Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), 209–225, AMS/IP Stud. Adv. Math., 23, Amer. Math. Soc., Providence, RI, 2001.
- [14] M. Marino, R. Minasian, G. Moore, and A. Strominger, *Nonlinear Instantons from supersymmetric p-Branes*, J. High Energy Phys. 2000, no. 1, Paper 5, 32 pp.
- [15] R.C. McLean, *Deformation of Calibrated Submanifolds*, Commun. Anal. Geom. **6** (1998) 705-747.
- [16] S. Salamon, *Riemannian geometry and holonomy groups*, John Wiley & Sons, 1989.
- [17] H. Schenk, *On a generalised Fourier transform for instantons over flat tori*, Comm. Math. Phys., 116 (1988) pp. 177-183.
- [18] A. Strominger, S.-T. Yau, and E. Zaslow, *Mirror Symmetry is T-Duality*, Nuclear Physics **B479** (1996) 243-259.
- [19] R. Thomas, *Moment maps, monodromy and mirror manifolds*; Symplectic geometry and mirror symmetry (Seoul, 2000), 467–498, World Sci. Publ., River Edge, NJ, 2001.
- [20] P. Wilson, *The role of in Calabi-Yau classification - a preliminary survey*, Mirror Symmetry II, edited by B. Greene and S.-T. Yau, International Press, (1997) 381-392.

Addresses:

Naichung Conan Leung (leung@ims.cuhk.edu.hk)

Institute of Mathematical Sciences and Department of Mathematics, The Chinese University of Hong Kong, Hong Kong

Jae-Hyuk Lee (jhlee@math.wustl.edu)

Department of Mathematics, Washington University in St. Louis, U.S.A.