A note on $H^1(\text{Emb}(M, N))$

Olaf Müller

November 21, 2018

Max-Planck-Institute for Mathematics in the Sciences Inselstrasse 22-26 D-04103 Leipzig, Europe

Abstract

Aim of this note is to gain cohomological information about the manifold $\operatorname{Emb}(M, N)$ from the topology of the target manifold N. For special conditions, a monomorphism $H^1(N) \to H^1(Emb(M, N))$ is constructed.

Let $\operatorname{Emb}_{g}(M, N)$ be the manifold of all embeddings of a manifold M into a manifold N that differ from a fixed reference embedding g only within a compact subset of M. Note that if M is compact then $\operatorname{Emb}_{g}(M, N) = \operatorname{Emb}(M, N)$, the space of all embeddings. Now and in the following M should be nonempty and N at least one-dimensional to give sense to the notion of 1-forms. $\operatorname{Emb}_{g}(M, N)$ is infinite-dimensional iff $\dim(M) \geq 1$.

Note. The following theorem 2 is a straightforward consequence within the framework of fiber integration (cf. [2], [1]), nevertheless it may be instructive to see another proof for the particular case.

For our construction, a crucial tool is the identification of a tangent vector V resp. the value of a vector field V on $\text{Emb}_{g}(M, N)$ at a fixed map f with a vector field along f, i.e. a section of $f^{*}TN$:

$$\Gamma|_f : \operatorname{Vect}(\operatorname{Emb}_{\mathfrak{g}}(\mathcal{M}, \mathcal{N})) \to \Gamma(f^* \mathcal{T} \mathcal{N}),$$

$$V|_f(p) := \mathcal{L}_V ev_p$$

or equivalently,

$$\hat{V}|_f : p \mapsto \partial_t(f(p))$$

where $p \in M$, f_t a curve representing V(f). This means, we just fix a point $p \in M$ and note the direction in which it is moved infinitesimally by the family of maps f_t . $\hat{V}|_f$ has compact support because $\bigcup_{t \in [-1;1]} \{f_t \neq g\}$ is compact.

Conversely, a vector field V on N maps to a vector field \tilde{V} on $\text{Emb}_{g}(M, N)$ by

 \sim : Vect(N) \rightarrow Vect(Emb_g(M, N))

 $\tilde{V}: f \mapsto \partial_t|_{t=0} (Fl_t^V \circ f),$

where Fl_t^V is the flow of V.

Lemma 1 (i) Both maps $\tilde{}$ and $\hat{}|_f$ are smooth for all f; moreover, $\hat{}$ depends smoothly on f.

- (ii) $|_f \circ = f^* \quad \forall f \in \operatorname{Emb}_g(M, N),$ i.e. the composition is the pull-back along f.
- (iii) The Lie bracket is preserved by $\tilde{}$:

$$[\tilde{V}, \tilde{W}] = [\widetilde{V, W}].$$

Proof. The infinite-dimensional manifold of vector fields on a manifold is in particular a Frêchet space, and the flow of a vector field depends smoothly on it, so both maps are smooth. Looking deeply at the definitions, one sees that they are inverse to each other in the sense of (ii). For a proof of the third statement we define

$$A : \operatorname{Diff}(N) \to \operatorname{Diff}(\operatorname{Emb}_{g}(M, N)),$$

$$h \mapsto (f \mapsto h \circ f).$$

As this map is a differentiable Lie group homomorphism, its differential $dA|_1$: $T_1 \text{Diff}(N) \to T_1 \text{Diff}(\text{Emb}_{g}(M, N))$ preserves the Lie bracket in the sense that [dA(V), dA(W)] = dA[V, W]. Now the source resp. target of dA is as Lie algebra isomorphic to Vect(N) resp. $\text{Vect}(\text{Emb}_{g}(M, N))$ via Φ resp. Ψ , and $\Psi \circ dA \circ \Phi^{-1} = \tilde{}$ by the chain rule \Box

In order to get a p-form $\hat{\omega}$ on $\text{Emb}_{g}(M, N)$ from one on N, for each p vector fields $X_1, ..., X_p \in \text{Vect}(\text{Emb}_{g}(M, N))$ we define

$$\hat{\omega}|_f(X_1,...X_p) := \int_M (\omega \circ f)(\hat{X}_1|_\gamma,...\hat{X}_p|_\gamma).$$

Note that for this definition we need a volume form on M.

Theorem 2 $: \Lambda^*(N) \to \Lambda^*(\text{Emb}(M, N))$ is an injective chain map.

Proof. Obviously, $\hat{\omega}$ is a p-form. We have to show that $d\hat{\omega} = \widehat{d\omega}$. We calculate $d\hat{\omega}(X_1, \dots, X_{p+1})$ at a section f using

$$d\Omega(X_1, \dots, X_{n+1}) = \sum_{i=1}^{n+1} ((-1)^{i=1} X_i(\Omega(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_p))) - \sum_{i < j} \Omega(-1)^{i+j}([X_i, X_j], X_1 \dots, X_{i-1}, X_{i+1}, \dots, X_{j-1}, X_{j+1}, \dots, X_n).$$

Now the crucial point is that because of tensoriality of $d\hat{\omega}$ we can choose $X_i = \tilde{x_i}$ for some vector fields x_i on N. Lemma 1 then implies $\widehat{[X_i, X_j]} = f^*([x_i, x_j])$, and because of $\hat{X_i}|_f = f^*x_i$ we have

$$X(\hat{\omega}(Y,Z))|_f = X \int_M (\omega \circ f)(|\hat{Y}|_f, |\hat{Z}|_f) = X \int_M (\omega \circ f)(y,z) = \int_M x(\omega(y,z)).$$

Using these equations we get the result transferring all terms to Lie brackets and derivatives in N.

Up to this point, we could have used any linear function on $C^{\infty}(M)$ instead of the integral. Then, if ω is nonzero, there is a point $m \in M$ and vector fields \hat{Y}_i such that $\omega|_m(\hat{Y}_1...\hat{Y}_n) \neq 0$. If we scale e.g. \hat{Y}_1 with a function F on M, then $\Omega(F\hat{Y}_1...\hat{Y}_n) = \int_M F \cdot (\omega \circ f)(\hat{Y}_1...\hat{Y}_n)$, and the statement follows from the nondegeneracy of the scalar product of $L_2(M)$

Therefore $\hat{}$ induces a homomorphism $\hat{}^*$ of the corresponding cohomology groups. If we knew that for each exact $\hat{\omega} = d\alpha$ also the form ω is exact or, equivalently, that $\alpha = \hat{\beta}$ for some β we would know that $\hat{}^*$ is injective on the cohomology level. This is asserted for the *first* cohomology group by the following theorem:

Theorem 3 If M is compact and if there is an embedding $M \to \mathbb{R}^{\dim N}$, then the map^{**} is a monomorphism $H^1(N) \to H^1(\operatorname{Emb}(M, N))$.

Note. Cohomology means here and in the following always de Rham cohomology.

Proof. Given a 1-form θ on N we have to show that if $\hat{\theta}$ is the differential of a function F on Emb(M, N) then θ is the differential of some function f on N (this implies $F = \hat{f}$ by Theorem 2). It is enough to show that $\int_c \theta = 0$ for all closed curves c in N. Given such a curve, we can construct a tubular neighborhood T of c and a smooth vector field on N such that $Fl_V^t(T) \subset T \forall t$ and c is the integral curve for V with Fl_V^1 being the identity on T (just take the standard circular flow vector field on the solid torus diffeomorphic to a small tubular neighborhood and scale it down radially to zero in its complement in a larger tubular neighborhood). Then the flow of V on c together with the distinction of a starting point c(0) fixes a parametrisation of the curve to which we will refer in the following. Consider curves K_{δ} in Emb(M, N) defined by $K_{\delta}(t) = Fl_V^t \circ \epsilon_{\delta}$ with $Im(\epsilon_{\delta}) \subset B_{\delta}(c(0)) \subset T$. Then we observe

$$\partial_t|_{t_0} \int_0^t dt' \,\hat{\theta}(\dot{K}_{\delta}(t')) = \hat{\theta}|_{K_{\delta}(t_0)}(\tilde{V}) = \int_M (\theta \circ K_{\delta}(t_0))(V \circ K_{\delta}(t_0)) = \int_M (\theta(V)) \circ K_{\delta}(t_0)$$

Therefore

$$\partial_t|_{t_0} \int_0^t dt' \ \hat{\theta}(\dot{K}_{\delta}(t')) \xrightarrow[\delta \to 0]{} vol(M) \cdot \theta(\dot{c}(t_0)) = vol(M) \cdot \partial_t|_{t_0} \int_0^t dt' \ \theta(\dot{c}(t'))$$

This convergence is uniform in t_0 : Define $D_{\delta} := max_{t \in [0,1]} diam(Im(K_{\delta}(t))),$ $H := max_{p \in T} \parallel grad(\theta(V))(p) \parallel$. Then we have

$$\left|\int_{M} (\theta(V)) \circ K_{\delta}(t_{0}) - vol(M) \cdot \theta(\dot{c}(t_{0}))\right| \leq D_{\delta} \cdot H$$

with $D_{\delta} \to 0$ for $\delta \to 0$ which implies uniform convergence. Therefore $\int_{K_{\delta}} \hat{\theta} \to vol(M) \cdot \int_{c} \theta$ and $\int_{c} \theta = 0$ \Box

One easily sees that one cannot just drop the condition of the existence of an embedding $M \to \mathbb{R}^{\dim N}$ by recalling the following fact about surfaces:

Let M be an orientable compact two-dimensional manifold without boundary distinct from $\mathbb{S}^1 \times \mathbb{S}^1$ and \mathbb{S}^2 . Then the identity component (and therefore all connected components) of the diffeomorphism group Diff(M) (= Emb(M, M)) is homotopy equivalent to a point (cf. [3], [4]).

From Lemma 2.1. in [2] it is clear that $\hat{\omega}$ is invariant under the natural action of volume-preserving diffeomorphisms of M on Emb(M, N). In the case of M being the distant union of finitely many points, all diffeomorphisms (permutations) are volume-preserving if we give the same weight to each point. Therefore ω is well-defined under the usual Σ_n -action in the definition of the configuration space and we get:

Corollary 4 Let $C_m(N)$ be the n-point configuration space of a manifold N, dim $N \ge 1$. Then there is a monomorphism $H^1(N) \to H^1(C_m(N))$

This result can also be compared to computations of the homology of configuration spaces as in [5].

A result analogous to Theorem 3 for the space of differentiable maps from M to N (without any injectivity condition) is found in [6].

References

- R.A. Abraham, J.E. Marsden, T.Ratiu: Manifolds, TensorAnalysis, and Applications, Springer-Verlag 1988
- [2] Tim Swift: A note on the space of lagrangian submanifolds of a symplectic manifold, Journal of Geometry and Physics 35 (2000) 183-192
- [3] C.J. Earl, J. Eells: The diffeomorphism group of a compact Riemann surface, Bulletin of the American Mathematical Society 73 (1967) 557-559
- [4] A. Grammain: Groupe des difféomorphismes et espace de Teichmüller d'une surface, Springer Lecture Notes in Mathematics 383 (1974) 157-169
- [5] C.-F. Bödigheimer, F.Cohen, L.Taylor: On the homology of configuration spaces, Topology 28 (1989) 111-123
- [6] Hông Vân Lê: Geometry of a class of soliton solutions of nonlinear Schrödinger equations, preprint, Leipzig 2001