# An estimate for the volume entropy of nonpositively curved graph manifolds

S. Buyalo \*

#### Abstract

Let M be a closed 3-dimensional graph manifold. We prove that h(g) > 1 for each geometrization g of M, where h(g) is the topological entropy of geodesic flow of g.

# 1 Introduction

Asymptotic geometry of a nonpositively curved (NPC for brevity) graph manifold is a complicated mixture of flat and hyperbolic parts which both nontrivially contribute in general picture. Recall that a NPC metric on a closed 3-dimensional graph manifold M recovers the JSJ-decomposition of M in the following sense. There is the unique (up to isotopy) minimal finite collection E of flat geodesically embedded tori and Klein bottles which are pairwise disjoint, and the metric completion of each connected component of the complement to E is a Seifert space called a block of M. Each block  $M_v$ is fibered over a 2-orbifold  $S_v$  with negative Euler characteristic,  $\chi(S_v) < 0$ . Furthermore, the metric locally splitts as  $U \times (-\varepsilon, \varepsilon)$  along the interior of each block, where U is a NPC surface, the splitting is compatible with the fibration, the fibers are closed geodesics, and the regular fibers have one and the same length  $l_v > 0$  depending only on the block.

Since we are interested in asymptotic properties, which are certainly the same for any finite covering of M, we may assume for simplicity that M is orientable, the collection E consists of tori, and each block is a trivial  $S^1$ -bundle over a compact surface  $S_v$  with boundary,  $M_v = S_v \times S^1$ . We also assume that the graph manifold structure of M is nontrivial, i.e. M itself is not a Seifert fibered space (though it may consist of one block).

The flat part of the asymptotic geometry of M studied in [BS], [CK], see also [HS]. Roughly speaking, it can be described by very special geodesic rays  $[0, \infty) \to M$ , which terminate in no block, skip through separating tori  $e \in E$  almost tangently spending most of the time near tori and moreover this time rapidly increases with each step. Though the set of such rays is

<sup>\*</sup>Partially supported by RFFI Grants 99-01-00104 and 00-15-96024.

a negligible part all of the rays, it contains an important information about geometry of M: it was shown in [CK] how this information allows to recover (up to scaling) the marked length spectrum of closed geodesics on  $S_v$  and the fiber length for each block  $M_v$ .

Here we study the hyperbolic part of the asymptotic geometry of M, assuming that the surface U from the local splitting  $U \times (-\varepsilon, \varepsilon)$  above has the constant curvature K = -1. In other words, each block fibers over a *hyperbolic* orbifold (surface)  $S_v$ . A NPC metric on M satisfying this condition is called a *geometrization* of M (because the metric of each block is modelled on  $\mathrm{H}^2 \times \mathbb{R}$ ). Note that any geometrization of M is only  $C^{1,1}$ -smooth being analytic along the interior of each block. It is known [L] that M admits a NPC metric if and only if it admits a geometrization. Necessary and sufficient topological conditions for M to carry a NPC metric were found in [BK].

The relevant metric invariant which measures hyperbolicity of a space is the volume entropy h. Let  $\pi : X \to M$  be the universal covering,  $x_0 \in X$ . Recall that h = h(X) is defined by

$$h = \lim_{R \to \infty} \frac{1}{R} \ln \operatorname{vol} B_R(x_0),$$

where  $B_R(x)$  is the ball in X of radius R centered at x. It is well known [M] that the limit exists, h is independent of the choice of  $x_0$  and if M is NPC then h coincides with the topological entropy of geodesic flow of M. The volume entropy scales as  $l^{-1}$ , l is the length. Thus the choice of a geometrization g of M also serves as a normalization. The sectional curvatures of g satisfy  $-1 \leq K \leq 0$ . Hence  $h(g) \leq 2$  by comparison with H<sup>3</sup>. Our main result is this.

**Theorem 1.1.** For any geometrization g of a graph manifold M we have h(g) > 1.

Remark 1.2. Though the universal covering X of M looks much more complicated than the model space  $\mathrm{H}^2 \times \mathbb{R}$ , even the estimate  $h(g) \geq 1 = h(\mathrm{H}^2 \times \mathbb{R})$  is not obvious and nontrivial: X contains no isometrically and geodesically embedded  $\mathrm{H}^2$ , which would lead to  $h(g) \geq 1$ ; on the other hand, the attempt to compare  $X \to \mathrm{H}^2 \times \mathbb{R}$  via exponential maps identifying some tangent spaces fails because the Jacobian of this map is > 1 at some points. Finally, the estimate from [BW] for the measure theoretic entropy of geodesic flow, which is always  $\leq h(g)$ , gives only  $\pi/4 < 1$  as a lower bound for any geometrization g of M (if one ignores the fact that  $C^{1,1}$ -smoothness of g is not sufficient to apply this estimate).

To prove Theorem 1.1 we use the well known fact that h(g) coincides with the critical exponent of a Poincaré series

$$\mathcal{P}(t) = \sum_{\gamma \in \Gamma} e^{-t|x_0 - \gamma x_0|},$$

where the fundamental group  $\Gamma = \pi_1(M)$  isometrically acts on X as the deck transformation group. Actually, instead of  $\mathcal{P}$  we use a modified Poincaré series  $\mathcal{P}_W$ , where summation is taken over a set W of walls in X. Our proof involves three ingredients: (i) a local estimate, which is technical and used in (ii); this estimate is obtained in section 2. (ii) An accumulating procedure, which is an inductive construction of appropriately choosen broken geodesics in X between the base point  $x_0$  and the walls from W; the choice of these paths is the key point of the proof. The outcome of the accumulating procedure is a generating set for  $\mathcal{P}_W$  used in (iii); the procedure is described in sect. 3. (iii) A self-similarity type argument. This part of the proof uses a standard idea from self-similarity theory to show that  $\mathcal{P}_W(h)$ diverges for some h > 1 using the generating set obtained in (ii). This is done in sect. 4.

Acknowledgment. The author is grateful to W. Ballmann for useful discussions of the topic of this note.

# 2 Local estimate

Let F be the universal covering of a compact hyperbolic surface S with geodesic boundary. We identify F with a convex subset  $F \subset H^2$  bounded by countably many disjoint geodesic line and fix  $o \in H^2 \setminus F$ . Let  $w_0$  be the boundary line of F closest to  $o, o_0 \in w_0$  be the point on  $w_0$  closest to o, so that  $|o - o_0| = \operatorname{dist}(o, F) =: l > 0$ .

Let A be the set of the boundary lines of F different from  $w_0$ . For each  $w \in A$  we denote by  $o_w \in w$  the point closest to o. Then the geodesic segment  $oo_w$  intersects  $w_0$  at some point  $t_w$ , and for  $\tilde{l}_w = |o - o_w|$  we have

$$\widetilde{l}_w = \widetilde{l}'_w + l''_w,$$

where  $\tilde{l}'_w = |o - t_w|, \, l''_w = |t_w - o_w|$  (all distances are taken in H<sup>2</sup>).

Next, we identify  $\mathrm{H}^2$  with  $\mathrm{H}^2 \times 0 \subset \mathrm{H}^2 \times \mathbb{R}$ , so that F becomes a subset of  $\mathrm{H}^2 \times \mathbb{R}$ , and we use for it the notations introduced above. Note that the point  $o_0$  is closest to o from the wall  $w_0 \times \mathbb{R}$ . We take a geodesic line  $\sigma \subset w_0 \times \mathbb{R}$  through  $o_0$  which is not horizontal, i.e.,  $\sigma \neq w_0 \times 0$ , and take  $s_w \in \sigma$  with  $|s_w - o_0| = |t_w - o_0|$ . Now, we put  $l'_w := |o - s_w|$  (the distance is taken in  $\mathrm{H}^2 \times \mathbb{R}$ ),  $\Delta_w := \tilde{l'_w} - l'_w$ .

In other words, we replace the distance  $\tilde{l}'_w$  between o and  $t_w$  is the hyperbolic plane  $\mathrm{H}^2$  by the distance  $l'_w$  in  $\mathrm{H}^2 \times \mathbb{R}$ , which is shorter by comparison: the triangles  $oo_0 t_w \subset \mathrm{H}^2 \times 0$ ,  $oo_0 s_w \subset \mathrm{H}^2 \times \mathbb{R}$  both have the right angles at  $o_0$ ,  $\angle (oo_0 t_w) = \frac{\pi}{2} = \angle (oo_0 s_w)$ , the common side  $oo_0$  and equal sides  $|o_0 - t_w| = |o_0 - s_w|$ . Since  $oo_0 t_w$  lies in the hyperbolic plane  $\mathrm{H}^2 \times 0$  but  $oo_0 s_w$  not, we have  $\Delta_w > 0$  except the case  $t_w = o_0 = s_w$ . Now, we want to estimate accumulation of the differences  $\Delta_w$  from below. The precise statement is this. **Lemma 2.1.** Given  $l_0 > 0$ ,  $\alpha_0 \in (0, \pi/2]$ , there exists  $\lambda_0 > 1$ , which depends only on  $l_0$ ,  $\alpha_0$  and the compact surface S, so that

$$\lambda(F, l, \alpha) := e^l \sum_{w \in A} e^{\Delta_w} e^{-\tilde{l}_w} \ge \lambda_0,$$

whenever  $l = \text{dist}(o, w_0) \ge l_0$  and the angle  $\alpha$  between the lines  $w_0 \times 0$  and  $\sigma$  is at least  $\alpha_0, \alpha_0 \le \alpha \le \pi/2$ .

*Proof.* By well known formula of hyperbolic geometry we have  $e^l = (\tan \frac{\psi}{4})^{-1}$ ,  $e^{-\tilde{l}_w} = \tan \frac{\psi_w}{4}$ , where  $\psi$ ,  $\psi_w$  are the angles under which  $w_0$  respectively  $w \in A$  are observed in H<sup>2</sup> from o. The boundary at infinity  $\partial_{\infty} F \subset \partial_{\infty} H^2 = S^1$  coincides with the limit set of  $\pi_1(S)$  represented in Iso(H<sup>2</sup>) as a Fuchsian group of second kind. It is well known that the Hausdorff dimension of  $\partial_{\infty} F$  (with respect to the angle metric) is < 1, in particular, the Lebesgue measure of  $\partial_{\infty} F$  is zero. Thus  $\psi = \sum_{w \in A} \psi_w$ . Therefore,  $\tan \frac{\psi}{4} \leq \sum_{w \in A} \tan \frac{\psi_w}{4}$  and

$$e^l \sum_{w \in A} e^{-\tilde{l}_w} = \sum_{w \in A} \tau_w \ge 1,$$

where  $\tau_w = e^{l-l_w}$ . However, this sum,  $\sum_{w \in A} \tau_w$ , can be close to 1 as much as we like (taking for instance  $l \to \infty$ ). As we noticed above,  $\Delta_w > 0$  unless  $s_w = t_w$ . So we always have  $\lambda(F, l, \alpha) > 1$ . The point is that  $\lambda(F, l, \alpha)$  is separated from 1 uniformly over all  $l \ge l_0$ ,  $\alpha \ge \alpha_0$ .

Consider  $A_0 \subset A$  consisting of all  $w \in A$  with  $|t_w - o_0| \ge 1$ . Then  $\Delta_w \ge \delta_0 > 0$  for all  $w \in A_0$ , where  $\delta_0$  depends only on  $l_0$ ,  $\alpha_0$ . We claim that

$$\sum_{w \in A_0} \tau_w \ge m_0 > 0,\tag{1}$$

where  $m_0 = m_0(\tilde{F}, l_0)$  is independent of l. Assuming (1), we have

$$\begin{aligned} \lambda(F,l,\alpha) &= \sum_{w \in A} e^{\Delta_w} \tau_w \ge \sum_{w \in A_0} e^{\Delta_w} \tau_w + \sum_{w \in A \setminus A_0} \tau_w \\ &\ge e^{\delta_0} \sum_{w \in A_0} \tau_w + \sum_{w \in A \setminus A_0} \tau_w \\ &= (e^{\delta_0} - 1) \sum_{w \in A_0} \tau_w + \sum_{w \in A} \tau_w \ge (e^{\delta_0} - 1)m_0 + 1 =: \lambda_0 > 1 \end{aligned}$$

It remains to prove (1). Let  $o'_w \in w$  be the point closest to  $o_0 \in w_0$ . Then  $\tilde{l}_w \leq |o - o'_w| \leq l + |o_0 - o'_w|$ , thus  $\tilde{l}_w - l \leq \operatorname{dist}(o_0, w)$ . Hence  $\tau_w \geq e^{-\operatorname{dist}(o_0,w)} \geq \frac{\tilde{\psi}_w}{4}$ , where w is observed from  $o_0$  under the angle  $\tilde{\psi}_w$ . The Lebesgue measure class on  $\partial_{\infty} \mathrm{H}^2$  is independent of the choice of origin, thus  $\sum_{w \in A} \widetilde{\psi}_w = \pi$ .

Since  $l \geq l_0$ , for a sufficiently small  $m_0 = m_0(S, l_0) > 0$  the sectors  $S^+(m_0)$ ,  $S^-(m_0)$ , defined below, intersect no  $w \in A \setminus A_0$ . Here is the definition of  $S^{\pm}(m_0)$ . The common vertex  $o_0$  of  $S^{\pm}(m_0)$  divides the line  $w_0$  into two opposite rays  $w_0^{\pm}$ . The sectors  $S^{\pm}(m_0) \subset \mathrm{H}^2$  are bounded by the rays  $w_0^{\pm}$ ,  $s^{\pm}(m_0)$ , where  $\angle_{o_0}(s^{\pm}(m_0), w_0^{\pm}) = 2m_0$ , and  $s^{\pm}(m_0) \cap F \neq \emptyset$ .

Therefore, it follows from  $\sum_{w \in A} \widetilde{\psi}_w = \pi$  that  $\sum_{w \in A_0} \tau_w \ge m_0$ , which completes the proof.

# **3** Accumulating procedure

To describe the accumulating procedure we need some information about metric structure of the universal covering X of (M, g), where g is a geometrization.

#### 3.1 Metric structure of the universal covering

Recall (see, for example, [BS], [CK]) that X can be represented as the countable union  $X = \bigcup_v X_v$  of blocks, where each  $X_v$  is a closed convex subset in X isometric to the metric product  $F_v \times \mathbb{R}$ , and  $F_v$  is the universal covering of a compact hyperbolic surface  $S_v$  with geodesic boundary. Every two blocks are either disjoint or intersect over a boundary component which is a 2-flat in X separating them and consequently no three blocks have a point in common. The 2-flats in X, which separate blocks, are called the walls. A wall w common for blocks  $X_v, X_{v'}$  covers a 2-torus  $e \subset M$ , which separates (may be locally) the blocks  $M_v = \pi(X_v), M_{v'} = \pi(X_{v'})$  of M. The metric decompositions  $X_v = F_v \times \mathbb{R}, X_{v'} = F_{v'} \times \mathbb{R}$  do not agree on w, and their  $\mathbb{R}$ -factors induce two fibrations of w by parallel geodesics. We denote by  $\alpha_w$  the angle between these fibrations,  $0 < \alpha_w \leq \pi/2$ . Since M is compact and the set E of separating tori in M is finite, we have  $\alpha_0 := \inf_w \alpha_w > 0$ , where the infimum is taken over all the walls in X.

## 3.2 Modified Poincaré series

We fix a wall  $w^* \subset X$ , take a block  $X_{v^*} \subset X$ , for which  $w^*$  is a boundary wall, and take a base point  $x_0 \in w^*$ . We denote by  $W_0$  the set of the boundary walls of  $X_{v^*}$  different from  $w^*$ ; by  $W_n$ ,  $n \ge 1$  the set of walls in X at the combinatorial distance n + 1 from  $w^*$ , that is,  $w \in W_n$  if and only if any geodesic segment in X between  $x_0$  and w intersets n walls over its interior, including a wall from  $W_0$ . Note that  $W = \bigcup_{n\ge 0} W_n$  consists of all walls in X, which lie on one and the same side of  $w^*$  as  $X_{v^*}$ . Now, we define a modified Poincaré series as follows

$$\mathcal{P}_W(t) = \sum_{w \in W} e^{-t \operatorname{dist}(x_0, w)}$$

Comparing  $P_W$  with  $\mathcal{P}(t) = \sum_{\gamma \in \Gamma} e^{-t|x_0 - \gamma x_0|}$ , one easily obtains from triangle inequality that  $\mathcal{P}(t) \ge e^{-D} \mathcal{P}_W(t)$ , where D > 0 is the maximal diameter of the tori  $e \in E$ . Recall that the critical exponent of  $\mathcal{P}$  is defined as the infimum of  $t \in \mathbb{R}$  for which  $\mathcal{P}(t) < \infty$ . Therefore, the critical exponent  $\overline{h}$  of  $\mathcal{P}_W$  satisfies  $\overline{h} \le h(g)$ , and to prove Theorem 1.1 it suffices to show that  $\overline{h} > 1$ .

#### 3.3 Special broken-geodesic paths

**Proposition 3.1.** For each  $n \ge 0$  we have

$$\mathcal{P}_n(1) := \sum_{w \in W_n} e^{-\operatorname{dist}(x_0, w)} \ge \frac{\pi}{4} \lambda_0^n,$$

where  $\lambda_0 > 1$  is the constant from Lemma 2.1.

*Proof.* Induction over n. For each  $n \ge 0$  and each wall  $w \in W_n$  we produce a broken geodesic  $\xi_w$  in X between  $x_0$  and w as follows. For  $w \in W_0$  we put  $\xi_w = x_0 x_w$ , where  $x_w \in w$  is the point closest to  $x_0$ . Note that  $x_0 x_w$  lies in a horizontal slice  $F_{v^*} \times \{r_0\}$  of the block  $X_{v^*} = F_{v^*} \times \mathbb{R}$ .

Assume that for all  $k, 0 \leq k \leq n-1$  and all  $w \in W_k$  the broken geodesic  $\xi_w$  is already defined,  $\xi_w$  is a juxtaposition  $\eta_{w_0}\eta_{w_1}\ldots\eta'_{w_k}$  of geodesic segments with the sequence  $w_i \in W_i$  of walls leading to  $w = w_k$ . Moreover, we assume that each segment of  $\xi_w$  lies in a block and connects its different boundary components, the last segment  $\eta'_{w_k}$  lies in a horizontal slice of the block, which contains it, and  $\eta'_{w_k}$  is orthogonal to the wall w.

Take  $w \in W_n$ . Then there is a unique  $\overline{w} \in W_{n-1}$  which precedes w. By the assumption, the last edge  $\eta'_{\overline{w}} = s_{\overline{w}} x_{\overline{w}}$  of  $\xi_{\overline{w}}$  lies in a horizontal slice  $F_{v'} \times \{r_{v'}\}$  of its block  $X_{v'} = F_{v'} \times \mathbb{R}$  and it is orthogonal to the wall  $\overline{w}$  (at the end point  $x_{\overline{w}}$ ). The important feature of our construction is that  $\xi_w$  contains all segments of  $\xi_{\overline{w}}$  but the last one  $\eta'_{\overline{w}}$ .

Let  $X_v \subset X$  be the other block adjacent to  $\overline{w}$ , in particular, the walls w,  $\overline{w}$  are its boundary components. Recall that the metric splitting  $X_v = F_v \times \mathbb{R}$ does not agree with that of  $X_{v'}$  along of  $\overline{w}$ . Let  $F_v \times \{r_v\}$  be the horizontal slice of  $X_v$  which contains  $x_{\overline{w}}$  on the corresponding boundary component. The boundary lines of the slices  $F_{v'} \times \{r_{v'}\}$ ,  $F_v \times \{r_v\}$ , which lie in  $\overline{w}$ , contain  $x_{\overline{w}}$  and form an angle  $\alpha_{\overline{w}} \in (\alpha_0, \pi/2]$ .

To construct  $\xi_w$  we do the following. We take an isometric copy  $F_v \subset \mathrm{H}^2 \times \{r_{v'}\}$  of  $F_v \times \{r_v\}$  (by rotating the last by the angle  $\alpha_{\overline{w}}$ ), where  $F_{v'} \times \{r_{v'}\} \subset \mathrm{H}^2 \times \{r_{v'}\}$ , so that  $F_{v'} \times \{r_{v'}\}$  and  $F_v$  are sitting in the hyperbolic plane  $\mathrm{H}^2 \times \{r_{v'}\}$  and they are adjacent along the common boundary

component, for which we use the same notation  $\overline{w}$ . Now, we connect the initial point  $s_{\overline{w}}$  of the last edge  $\eta'_{\overline{w}} \subset \xi_{\overline{w}}$  with the boundary component of  $F_v$  corresponding to w by the shortest geodesic segment  $s_{\overline{w}}x'_w \subset \mathrm{H}^2 \times \{r_{v'}\}$  and take  $t_w = s_{\overline{w}}x'_w \cap \overline{w}$ . The segment  $t_wx'_w$  turned back to  $F_v \times \{r_v\}$  gives the last segment  $\eta'_w = s_wx_w$  of  $\xi_w$ . So  $s_w \in \overline{w} \cap F_v \times \{r_v\}$ ,  $|s_w - x_{\overline{w}}| = |t_w - x_{\overline{w}}|$  and  $\eta'_w \subset F_v \times \{r_v\}$  is orthogonal to w at  $x_w$ . To complete the construction of  $\xi_w$ , we delete the last segment  $\eta'_w \subset \xi_{\overline{w}}$  replacing it by  $\eta_{\overline{w}}\eta'_w$ , where  $\eta_{\overline{w}} = s_{\overline{w}}s_w \subset X_{v'}$ . Clearly, so constructed  $\xi_w$  has all properties advertised above.

Let  $l = |s_{\overline{w}} - x_{\overline{w}}| = L(\eta'_{\overline{w}})$  be the length of the last segment of  $\xi_{\overline{w}}$ ,  $l'_w = L(\eta_{\overline{w}}), l''_w = L(\eta'_w)$ . Then we have

$$L(\xi_w) = L(\xi_{\overline{w}}) - l + l'_w + l''_w = L(\xi_{\overline{w}}) - l + \widetilde{l}_w - \Delta_w,$$

where  $\tilde{l}_w = L(s_{\overline{w}}x'_w)$ ,  $\tilde{l}'_w = L(s_{\overline{w}}t_w)$  so that  $\tilde{l}_w = \tilde{l}'_w + l''_w$ , and  $\Delta_w = \tilde{l}'_w - l'_w$ . We obtained the same configuration which was studied in sect. 2, and we are going to apply Lemma 2.1 to estimate  $\mathcal{P}_n(1)$  from below. Note that l being the length of a segment in X connecting different boundary components of a block is separated from 0 by some positive constant  $l_0$ , which depends only on M,  $l \geq l_0 > 0$ .

For n = 0 we have

$$\mathcal{P}_0(1) = \sum_{w \in W_0} e^{-|x_0 - x_w|} = \sum_{w \in W_0} \tan \frac{\psi_w}{4} \ge \frac{\pi}{4},$$

where  $\psi_w$  is the angle under which the boundary component w of  $X_{v^*}$  is observed from  $x_0$  (in the horizontal direction).

By inductive assumption we have

$$\mathcal{P}_{n-1}(1) \ge \sum_{w \in W_{n-1}} e^{-L(\xi_w)} \ge \frac{\pi}{4} \lambda_0^{n-1}.$$

Represent  $W_n = \bigcup_{\overline{w}} W_{n,\overline{w}}$ , where the union is taken over all  $\overline{w} \in W_{n-1}$  and each  $w \in W_{n,\overline{w}}$  follows  $\overline{w}$ . Then

$$\mathcal{P}_n(1) \ge \sum_{w \in W_n} e^{-L(\xi_w)} = \sum_{\overline{w} \in W_{n-1}} e^{-L(\overline{\xi_w})} e^l \sum_{w \in W_{n,\overline{w}}} e^{\Delta_w} e^{-\widetilde{l}_w}.$$

Applying Lemma 2.1 with  $A = W_{n,\overline{w}}$ , we obtain

$$\mathcal{P}_n(1) \ge \lambda_0 \sum_{\overline{w} \in W_{n-1}} e^{-L(\xi_{\overline{w}})} \ge \frac{\pi}{4} \lambda_0^n,$$

which completes the proof.

## 4 Self-similarity argument

The constant  $\lambda_0 > 1$  from Proposition 3.1 depends only on some metric data of M. Thus there is  $n \in \mathbb{N}$ , n = n(M), so that  $\frac{\pi}{4}\lambda_0^n > 1$ . It follows from Proposition 3.1 that

$$\mathcal{P}_n(\overline{h}) = \sum_{w \in W_n} e^{-\overline{h}\operatorname{dist}(x_0,w)} \ge 1$$

for some  $\overline{h} > 1$ . Furthermore, taking *n* sufficiently large, we can find  $\overline{h} > 1$  with  $\mathcal{P}_n(\overline{h}) \geq 1$  for any choice of the initial block  $X_{v^*}$ , its wall  $w^*$  and the base point  $x_0 \in w^*$ , since the set of choices up to isometries of X is compact.

We fix  $n \in \mathbb{N}$  with this property and select a subset  $W^* \subset W$ ,  $W^* = \bigcup_{k \geq 1} W_k^*$ , where  $W_k^* = W_{kn}$ . The set  $W_1^*$  serves as the generating set for  $W^*$ . Connecting  $x_0$  with each wall  $w \in W_1^*$  by the shortest geodesic segment  $x_0 x_w$ , we obtain new base points  $x_w \in w$  (these  $x_w$  may be different from  $x_w$  constructed in the proof of Proposition 3.1). By induction, we find a base point  $x_w \in w$  for each  $w \in W_k^*$ ,  $k \geq 1$  with the property dist $(x_{\overline{w}}, w) = |x_{\overline{w}} - x_w|$ , where  $\overline{w} \in W_{k-1}^*$  precedes w. Furthermore, by the choice of n and  $\overline{h}$  we have

$$\sum_{w} e^{-\overline{h}|x_{\overline{w}} - x_w|} \ge 1$$

for each  $\overline{w} \in W_{k-1}^*$ , where the summation is taken over all  $w \in W_k^*$  which follow  $\overline{w}$ . Since  $\operatorname{dist}(x_0, w) \leq \operatorname{dist}(x_0, x_w) \leq \operatorname{dist}(x_0, x_{\overline{w}}) + |x_{\overline{w}} - x_w|$ , we obtain

$$\mathcal{P}_{W_k^*}(\overline{h}) = \sum_{w \in W_k^*} e^{-\overline{h}\operatorname{dist}(x_0, w)}$$
  
$$\geq \sum_{w \in W_{k-1}^*} e^{-\overline{h}\operatorname{dist}(x_0, x_w)} \geq \sum_{w \in W_n} e^{-\overline{h}\operatorname{dist}(x_0, x_w)} \geq 1$$

for each  $k \geq 1$ . Therefore, the modified Poincaré series

$$\mathcal{P}_W(\overline{h}) \ge \mathcal{P}_{W^*}(\overline{h}) = \sum_{k \ge 1} \mathcal{P}_{W^*_k}(\overline{h})$$

diverges at  $\overline{h}$  and hence  $h(g) \ge \overline{h} > 1$ . This completes the proof of Theorem 1.1.

## References

[BW] W. Ballmann and M. Wojtkowski, An estimate for the measure theoretic entropy of geodesic flow, Ergod. Th. & Dynam. Sys. 9 (1989), 271–279.

- [BK] S. Buyalo and V. Kobel'skii, Geometrization of graph-manifolds. II. Isometric geometrization, St. Petersburg Math. J. 7 (1996), 387–404.
- [BS] S. Buyalo and V. Schroeder, On the asymptotic geometry of nonpositively curved graph manifolds, Trans. Amer. Math. Soc. 353 (2001), 853–875.
- [CK] C. Croke and B. Kleiner, The geodesic flow and a nonpositively curved graph manifold, arXiv:math.DG/9911170, 1999.
- [HS] C. Hummel and V. Schroeder, Tits geometry of cocompact realanalytic Hadamard manifolds of dimension 4. Differential Geom. Appl. 11 (1999), 129–143.
- B. Leeb, 3-manifolds with(out) metrics of nonpositive curvature. Invent. Math. 122 (1995), 277–289.
- [M] A. Manning, Topological entropy for geodesic flows. Ann. of Math. (2) 110 (1979), 567–573.