

AN ALGORITHM FOR COMPUTING THE WEIERSTRASS NORMAL FORM OF HYPERELLIPTIC CURVES

Mark van Hoeij

Abstract

An algorithm is given to compute a normal form for hyperelliptic curves. The elliptic case has been treated in [2]. In this paper the hyperelliptic case is treated.

1 Introduction

The problem we consider is the same as in [2], but then for the hyperelliptic case instead of the elliptic case. Let C be an algebraic curve given by an irreducible polynomial $f \in \mathbb{C}[x, y]$. The function field of C is the field of fractions of $\mathbb{C}[x, y]/(f)$, and is denoted as $\mathbb{C}(x, y)$. We want to decide if C is hyperelliptic or not. If C is hyperelliptic, we want to find a Weierstrass normal form for C (i.e. an equation $y_0^2 = F(x_0)$ for some squarefree polynomial $F(x_0) \in \mathbb{C}[x_0]$). We want to find a birational map from C to this normal form, and its inverse. This means expressing x, y in terms of x_0, y_0 and vice versa. Let g be the genus of C .

The problems to be solved are the following

- Decide if C is hyperelliptic or not. By definition, C is hyperelliptic if $g > 1$ and there exists $x_0 \in \mathbb{C}(x, y)$ such that $\mathbb{C}(x_0)$ is a subfield of $\mathbb{C}(x, y)$ with index 2. If C is hyperelliptic then the field $K := \mathbb{C}(x_0)$ is unique.
- Find a generator x_0 of K .
- Find an element y_0 of $\mathbb{C}(x, y)$, not in K . Take for example $y_0 = x$ (interchange x, y if it turns out that $x \in K$). Compute the minimal polynomial of $y_0 = x$ over K . This is done by writing x_0 as a quotient v_1/v_2 with $v_1, v_2 \in \mathbb{C}[x, y]$ and computing the resultant $\text{Res}_y(v_2\tilde{x}_0 - v_1, f)$ where \tilde{x}_0 is the name of a variable that plays the role of x_0 . Then add an element of K to y_0 to change its minimal polynomial into the form $y_0^2 - F(x_0)$ for some rational function $F(x_0) \in K$. Then multiply y_0 by an element of K to turn F_0 into a squarefree polynomial.

- At this point the functions x_0, y_0 are expressed as elements of $\mathbb{C}(x, y)$ and their algebraic relation $y_0^2 - F(x_0)$ is in Weierstrass form. Now view x_0, y_0 as variables \tilde{x}_0, \tilde{y}_0 , and express x, y as rational functions in these variables. This can be done in the same way as in [2].

2 Decide if C is hyperelliptic

If $f \in \mathbb{C}[x, y]$ is irreducible, then we can decide if C is elliptic by computing the genus g . The curve is elliptic if and only if $g = 1$. For the hyperelliptic case more work needs to be done. An algorithm for this is implemented in Maple 6. To view the it, issue the following commands in Maple 6 or higher

```
interface(verboseproc=2): op('algcures/is_hyperelliptic');
```

and the following code appears

```
proc(f, x, y)
local a, b, i, j, g, z;
option remember, 'Copyright (c) 1999 Waterloo Maple Inc. All rights\
reserved. Author: M. van Hoeij';
i := degree(f, {x, y});
j := min(degree(f, x), degree(f, y), i - ldegree(f, {x, y}));
if i < 4 or j < 2 then false
elif j = 2 then evalb(1 < 'algcures/genus'(f, x, y))
else
b := 'algcures/differentials'(f, x, y, skip_dx);
g := nops(b);
j := 1/2*(i - 1)*(i - 2) - g;
if g = 2 then true
elif g < 2 or j < 2 or j < 4 and
{seq(z[3], z = 'algcures/singularities'(f, x, y))} = {1}
then false
else
z := {seq(a[i], i = 1 .. g)};
j := {coeffs(expand(numer(subs(RootOf(f, y) = y, evala(
Expand(
subs(y = RootOf(f, y), add(a[i]*b[i], i = 1 .. g)^2))))),
z)};
z := {seq(a[i], i = 1 .. nops(j))};
z := nops(j) - nops(indets(subs('solve/linear'({coeffs(
expand(add(a[i]*j[i], i = 1 .. nops(j))), {x, y})), z)
, z)) intersect z);
if z < 2*g - 1 then error "should not happen"
else evalb(z = 2*g - 1)
end if
end if
end if
end proc
```

We will not explain all steps, but the main steps are the following.

- Let V be the vector space of holomorphic differentials. An algorithm described in [1] is used to compute a basis b_1, \dots, b_g for V .
- If $g < 2$ then the curve is not hyperelliptic and the algorithm stops.
- Let W be the span of all $b_i b_j$, $1 \leq i \leq j \leq g$.
- If C is hyperelliptic, and if t is any generator of K , i.e. $K = \mathbb{C}(t)$, then it is known that

$$t^i dt, \quad i = 0, \dots, g-1 \quad (1)$$

is a basis of V . Then $t^i d^2 t$, $i = 0, \dots, 2g-2$ must be a basis of W . So if C is hyperelliptic then the dimension of W must be $2g-1$. It is not hard to show that if C is not hyperelliptic and $g > 1$ then $\dim(W) > 2g-1$. In the algorithm, $z = \dim(W)$. Testing if $z = 2g-1$ decides if C is hyperelliptic or not.

3 Compute a function of degree 2

Once we have a generator x_0 for K , it is not difficult to find y_0 , see section 1. This is implemented in Maple in the procedure ‘`algcurves/genus2`’ (to view the code type similar commands as in the previous section). After that, the remaining steps are the same as in [2].

So the main remaining problem to be solved is to compute some generator x_0 of K . We have a basis b_1, \dots, b_g of V . We want to find two particular elements of V , whose quotient is a generator of K . For example, if t is any generator of K , then dt , $t dt$ will do, and so will $t^{g-2} dt$, $t^{g-1} dt$. We will describe how to find a subspace $V_2 \subseteq V$ of the form $\mathbb{C} t^{g-2} dt + \mathbb{C} t^{g-1} dt$ for some generator t of K . If v_1, v_2 is any basis of V_2 , then we can take $x_0 = v_1/v_2$ as our generator of K .

The idea for finding V_2 is the following. Take an arbitrary nonsingular point P on the curve. Since t may be any generator of K , we may assume without loss of generality that $t(P) = 0$. Then $t^{g-1} dt$ has the highest valuation at P of any nonzero element of V , and $t^{g-2} dt$ has the second highest valuation. So we can find V_2 from V by solving linear equations: As long as $\dim(V) > 2$, let m be the smallest valuation at P of any element of V . Then compute the subspace of V of those elements that have valuation $> m$, and replace V by that subspace. This way the dimension decreases, and after a number of steps V will equal V_2 . Then take $x_0 = v_1/v_2$ where v_1, v_2 is any basis of V_2 . This is implemented in Maple in the procedure ‘`algcurves/index2subfield`’.

4 An example

$$f = y^9 + 3x^2 y^6 + 3x^4 y^3 + x^6 + y^2$$

This curve has genus 3 and is hyperelliptic. The result of the Maple commands:
`with(algcurves): Weierstrassform(f,x,y,x0,y0);`
 is

$$[y_0^2 + 2 - 7x_0 + 35x_0^4 - 21x_0^5 + 7x_0^6 - x_0^7 + 21x_0^2 - 35x_0^3, \\ \frac{y(y^3 + x^2 + 1)}{y^6 + 2x^2y^3 + x^4 + y}, -\frac{x(y^3 + x^2)}{y}, -(-1 + x_0)y_0, 1 - 3x_0 + 3x_0^2 - x_0^3]$$

The first expression is the Weierstrassform $y_0^2 - F(x_0)$. The second expression is a generator for the field K , i.e. a function of degree 2 in $\mathbb{C}(x, y)$. Under the birational map this corresponds to x_0 . The third expression is the image of y_0 under this birational map. The 4'th and 5'th expression are x, y expressed in terms of x_0, y_0 .

References

- [1] B. Deconinck and M. van Hoeij. Computing Riemann matrices of algebraic curves, *PhysicaD*, **152**, p. 28–46, (2001)
- [2] M. van Hoeij. An algorithm for computing the Weierstrass normal form. *ISSAC '95 Proceedings*, p. 90–95, (1995). Available from <http://www.math.fsu.edu/~hoeij/papers.html>