

**Period mappings and differential
equations. From \mathbb{C} to \mathbb{C}_p**

TÔHOKU-HOKKAIDÔ LECTURES IN ARITHMETIC GEOMETRY

Yves André

(with appendices by F. Kato and N. Tsuzuki)

Contents

Chapter I. Analytic aspects of p -adic periods.	9
(Analysis)	
1. Analytic continuation; topological point of view.	10
2. Analytic continuation; algebraic approach.	21
3. The tale of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$.	35
4. Abelian periods as algebraic integrals.	43
5. Periods as solutions of the Gauss-Manin connection.	52
Chapter II. Introduction to the theory of p -adic period mappings.	65
(Geometry)	
1. A survey of moduli and period mappings over \mathbb{C} .	66
2. Preliminaries on p -divisible groups.	72
3. A stroll in the crystalline world.	75
4. Moduli problems for p -divisible groups.	83
5. p -adic period domains.	87
6. The p -adic period mapping and the Gauss-Manin connection.	90
7. p -adic uniformization of Shimura varieties.	96
Chapter III. p -adic orbifolds and monodromy.	111
(Group theory)	
1. Étale coverings and fundamental groups in p -adic geometry.	112
2. Temperate fundamental groups.	126
3. Local and global monodromy of p -adic differential equations.	141
4. Non-archimedean orbifolds, uniformizing differential equations and period mappings.	158
5. p -adic triangle groups.	175
6. The tale of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ (Escher's triangle group and its diadic and triadic twins).	190
Appendix A. Rapid Course in p -adic Analysis.	207

by F. Kato

1. Introduction.	207
2. Rigid analytic spaces.	209
3. Relation with Formal Geometry.	212
4. Topology of Rigid Analytic Space.	216
5. Berkovich' approach to non-archimedean analysis	218
Appendix B. Overview of theory of p -adic uniformization.	221
	by F. Kato
1. Bruhat-Tits building.	221
2. Drinfeld symmetric space.	222
3. Uniformization.	226
4. Examples.	227
5. Mumford's fake projective plane.	228
Appendix C. p -adic symmetric domains and Totaro's theorem	231
	by N. Tsuzuki
1. Weakly admissible filtered isocrystals.	231
2. Filtered isocrystals with G -structure.	232
3. Totaro's theorem.	233
4. p -adic symmetric domains.	236
Bibliography	239

Introduction.

Purpose of this book.

This book stems from lectures given at the Universities of Tōhoku and Hokkaidō.

Its main purpose is to introduce the reader to p -adic analytic geometry and to the theory of p -adic analytic functions and differential equations, by focusing on the theme of *period mappings*. Of course, this approach is not meant to replace more systematic expositions of p -adic analysis or geometry found in a number of good treatises, but to complement them.

It is our general rule, in this book, to follow as closely as possible the complex theory, and to go back and forth between the complex and the p -adic worlds. We hope that this will make the text of interest both to some complex geometers and to some arithmetic geometers.

In the course of Chapter I, this approach will eventually become a kaleidoscope of half-correspondences and broken echoes. We hope that the reader will then have gained enough hindsight and wariness about these analogies, and will enjoy seeing unity being restored at a deeper level in Chapters II and III.

We have chosen the theme of period mappings because of its central role in the nineteenth-century mathematics as a fertile place of interaction between differential equations, group theory, algebraic and differential geometry, topology (and even number theory). In fact, it was a guiding thread in the early harmonious development of these branches of mathematics, from Gauss and Riemann to Klein and Poincaré (*cf.* [Gray86]).

The origins lie in Gauss' largely unpublished work on elliptic functions on one hand (rediscovered and extended by Abel and by Jacobi), and on the hypergeometric differential equation in the complex domain on the other hand. Gauss knew the connection between the two topics: the inverse of the indefinite integral $\int \frac{dx}{\sqrt{(1-x^2)(1-k^2x^2)}}$ is a single-valued function with two independent periods ω_1 and ω_2 which are solutions of the hypergeometric differential equation with parameters $(\frac{1}{2}, \frac{1}{2}; 1)$ in the variable k^2 .

Riemann's point of view of the 'Riemann surface' of a multivalued complex function has given a geometric framework for all of complex analysis. He applied this idea with equal success to Jacobi's inversion problem for more general indefinite algebraic integrals on one hand; and to the elucidation of the paradoxical polymorphism of hypergeometric functions which had puzzled Gauss and Kummer by introducing the concept of monodromy, on the other hand. He studied in detail the monodromy of the multivalued map $k^2 \mapsto \tau = \omega_2/\omega_1$, the first and basic example of a 'period mapping', whose inverse is single-valued. He also showed (and Schwarz rediscovered)

that under certain conditions, the quotient of two solutions of more general hypergeometric equations maps the upper half-plane onto a curvilinear triangle.

Group-theoretic aspects of monodromy were studied by Jordan, and the geometrization of complex analysis around the concept of automorphic function was carried on by Klein, who once described his work as ‘blending Galois with Riemann’⁽¹⁾. The achievement fell to Poincaré. Starting from Fuchs’ problem of finding all second-order differential equations for which a quotient of solutions τ admits a single-valued inverse, Poincaré founded the theory of ‘uniformizing differential equations’ (in modern terminology), and eventually recognized that every Riemann surface of genus > 1 should arise from the action of a discrete group of non-euclidean moves on the upper half-plane.

What about non-archimedean analogues of this saga?

It is clear that the development of p -adic analysis and geometry took a strikingly different route.

The very beginnings looked similar, indeed: J. Tate introduced rigid geometry as a proper framework for the uniformization $\mathbb{C}_p^\times/q^{\mathbb{Z}}$ of elliptic curves with multiplicative reduction; B. Dwork developed p -adic analysis starting from the hypergeometric differential equation with parameters $(\frac{1}{2}, \frac{1}{2}; 1)$, where he discovered the essential notion of ‘Frobenius structure’. But it was immediately clear that since solutions of p -adic differential equations do not converge up to the next singularity, no faithful counterpart of complex monodromy could take place.

From there on, different p -adic theories grew apart, with their own languages, most of them relying on sophisticated parts of contemporary algebraic geometry, and all claiming some analogy with “the complex case”:

- the theory of differential equations matured slowly (with a strong orientation toward applications to exponential sums), struggling with the problems of singularities, without being able to tackle global problems,
- crystalline theory offered a global viewpoint on differential equations (oriented toward the cohomology of varieties in positive characteristic), but did not help to understand singularities,
- the theory of p -adic representations and p -adic Hodge theory developed independently of differential equations, as did the several avatars of rigid geometry, motivated by idiosyncratic problems.

However, it is the author’s opinion that the situation has somewhat changed over the last years, that isolated branches are merging by fits and

⁽¹⁾quoted from [Gray86, p. 179].

starts⁽²⁾, and that a synthesis is gradually emerging. This encourages to hope that, after many twists and turns, period mappings can indeed become a unifying theme in the p -adic context. This book is intended to be a contribution in this direction, by bringing closer the p -adic theory to its complex precursor — from periods and monodromy up to triangle groups. Much remains to be done in order to achieve comparable harmony and clarity.

Contents of this book.

Chapter I is preparatory. It deals with the problems of analytic continuation — with emphasis on the case of solutions of differential equations — and periods of abelian integrals, in the p -adic context.

Multivalued complex-analytic functions can be handled in two different, but essentially equivalent, ways:

- 1) in a geometric way using Riemann surfaces, coverings and paths;
- 2) as limits of algebraic functions; this less orthodox way leads to a more Galois-theoretic viewpoint on analytic continuation.

Both ways are practicable in the p -adic context, but eventually lead to completely different theories of analytic continuation.

Following the first way demands to have at disposal p -adic spaces which are locally arcwise connected. Surprisingly enough in view of the fractal nature of p -adic numbers, such a nice p -adic geometry does exist: it has been built by V. Berkovich (from the categorical viewpoint, it is essentially equivalent to rigid geometry, in the sense of Tate and Raynaud). In this framework, the monodromy of differential equations can be analyzed in the usual way; but a new phenomenon occurs: it is no longer true that there exists a basis of solutions around every ordinary point. This approach is therefore limited to a rather special class of connections. We shall see in the sequel how the theory of p -adic period mappings provides interesting examples in this class.

Following way 2), one encounters Dwork's notion of Frobenius structure, which has often been considered as a plausible substitute for monodromy in the p -adic context. We discuss so-called unit-root crystals and overconvergence, and illustrate these notions by Dwork's treatment of the p -adic Gamma function Γ_p , the Gross-Koblitz formula, and by a detailed study of the p -adic hypergeometric function $F(\frac{1}{2}, \frac{1}{2}, 1; z)$.

Abelian periods show themselves in two different ways:

- 1) as integrals of algebraic differentials over loops,

⁽²⁾let us mention notably the incursion of rigid geometry into the crystalline viewpoint and into the geometric theory of finite coverings, the maturity of index theory and its applications to algebraic geometry in positive characteristic, the new connections between p -adic representations and differential equations on annuli...

2) when the integrand depends on a parameter z , as solutions of certain linear differential equations in z (Picard-Fuchs, or Gauss-Manin).

Both ways are passable in the p -adic context, and again lead to completely different theories.

Following up the first way — integrals being interpreted as Riemann sums — leads to P. Colmez' construction of abelian p -adic periods (in the sense of Fontaine-Messing). These periods relate the Tate module (p -primary torsion points) to the first De Rham cohomology module of a given abelian variety over a p -adic field. They do not live in \mathbb{C}_p , but in a certain p -adic ring B_{dR}^+ (of which \mathbb{C}_p is a quotient), and it is not possible in general to express them function-theoretically, when the abelian variety moves in a family. Nevertheless, we stress some particular cases where this can actually be done, throwing a bridge between ways 1) and 2).

The second way — Dwork's viewpoint of periods as solutions of Gauss-Manin connections — will prevail at the modular level, when p -divisible groups and filtered De Rham modules attached to families of abelian varieties will be compared not directly, but via their moduli spaces by the period mapping.

This way of looking at periods lacks arithmetic structure: namely, a rational, or at least a \mathbb{Q} -structure, on the \mathbb{C}_p -space of solutions, to convey some arithmetical meaning to the periods. We discuss some cases where such a canonical structure exists (' p -adic Betti lattices'), notably the case of abelian varieties with supersingular reduction. We get here well-defined p -adic periods (completely different from the Fontaine-Messing periods), and compute them in terms of Γ_p -values in the case of elliptic curves with complex multiplication (p -adic analogue of the Lerch-Chowla-Selberg formula).

Chapter II is an introduction to the geometric theory of p -adic period mappings, in the sense of Drinfeld-Rapoport-Zink.

We have tried to keep prerequisites at a minimum, and to emphasize as much as possible the analogies between the complex and p -adic contexts. Basic definitions about p -divisible groups and crystals are recalled, and the proof of some basic results is sketched.

The theory of period mappings attaches to a family of algebraic varieties its periods, viewed abstractly as a moving point in a suitable grassmannian. Due to constraints of Riemann-type, the period mapping actually takes its values in an open subset of the grassmannian, the period domain, which is a symmetric domain. It is multivalued, the ambiguity being described by the projective monodromy of the Gauss-Manin connection.

The younger p -adic theory is far less advanced: at present, there is a theory of p -adic period mappings only for p -divisible groups or closely related geometric objects. For want of wide range, it has nevertheless gained richness and depth.

In the presentation of Drinfeld-Rapoport-Zink, one starts by constructing a moduli space \mathcal{M} for p -divisible groups which are (quasi-)isogenous to a given one in characteristic p . The p -adic period mapping \mathcal{P} then relates this moduli space to a suitable grassmannian which parametrizes the Hodge filtration in the Dieudonné module. There are again constraints of Riemann-type, which force the period mapping to take its values in an open subset of the grassmannian, the period domain, which is a ‘symmetric domain’. The best known example of such a period domain is the Drinfeld space $\mathbb{C}_p \setminus \mathbb{Q}_p$, a p -adic analogue of the double-half-plane $\mathbb{C} \setminus \mathbb{R}$.

In the situation where the p -divisible groups are algebraizable, *i.e.* come from p -primary torsion of abelian varieties parametrized by a certain Shimura variety $\mathcal{S}h$, the moduli space \mathcal{M} provides a uniformization of some tubular region in the Shimura variety $\mathcal{S}h$.

We show that, just as in the complex case, \mathcal{P} can be described in terms of quotients of solutions of the associated Gauss-Manin connection (this feature does not seem to appear in the literature, except in the old special case of Dwork’s period mapping for elliptic curves with ordinary reduction). This allows to give explicit formulas ‘à la Dwork’ for \mathcal{P} in many cases.

The most interesting cases, investigated by Drinfeld-Rapoport-Zink, arise when $\mathcal{S}h$ itself (more accurately: a whole connected component) is uniformized by \mathcal{M} : $\mathcal{S}h$ is then a quotient of \mathcal{M} by an arithmetic discrete group Γ . We show that, up to replacing $\mathcal{S}h$ by a finite covering (to kill torsion in Γ), the solutions of the Gauss-Manin connection extend to global multivalued functions: in other words, the Gauss-Manin connection has global monodromy in the sense of chapter I, and the projective monodromy group coincides with Γ .

We review the case of Shimura curves attached to quaternion algebras over totally real fields (Čerednik-Drinfeld-Boutot-Zink): there is a global p -adic uniformization when p divides the discriminant of the quaternion algebra. We then construct, using the p -adic Betti lattices of chapter I, a canonical $\overline{\mathbb{Q}}$ -space of solutions of the Gauss-Manin connection which is stable under global monodromy.

Chapter III explores the group-theoretic aspects of the theory of period mappings, in the p -adic context.

The modern presentation of ramified coverings, uniformization, and of the Gauss-Riemann-Fuchs-Schwarz theory uses the notions of orbifold and uniformizing differential equation. We develop p -adic counterparts of these notions.

The right notion of ramified covering to adopt is not obvious, because étale coverings are only exceptionally topological coverings, in the p -adic context. Berkovich’s geometry provides a very convenient framework for this kind of problems, since his spaces are locally arcwise connected. In the first section, we discuss the formalism of fundamental groups attached to

categories of (possibly infinite) étale coverings satisfying simple axioms. Although the definition of these topological groups is simple and natural, their topology itself may be very complicated (not locally compact in general).

It turns out that there are too few topological coverings and too many étale coverings, in general, to provide a workable theory of monodromy. To remedy this, we introduce in the second section the notion of *temperate étale covering*. Such coverings are essentially built from (possibly infinite) topological coverings of finite étale coverings. They are classified by temperate fundamental groups, which seem to be the right p -adic equivalents, in many ways, of fundamental groups of complex manifolds. These groups are not discrete in general, but nevertheless often possess many infinite discrete quotients. We present a large selection of examples.

Temperate étale coverings are well-suited to the definition of p -adic orbifold charts and of the category of *p -adic orbifolds* (they replace the unramified coverings, over \mathbb{C}). As in the complex case, orbifolds and orbifold fundamental groups are the tools for a theory of ramified coverings, developed in section 4.

Before turning to the p -adic analogue of the *uniformizing differential equation* attached to an orbifold of dimension one (via Schwarzian derivatives), we discuss local and global monodromy of p -adic differential equations in section 3.

We outline the Christol-Mebkhout theory of p -adic slopes of differential equations over annuli, and the relation with Galois representations of local fields of characteristic p .

We then define and study the *p -adic étale Riemann-Hilbert functor*, which attaches a vector bundle with integrable connection to any discrete representation of the étale fundamental group of a p -adic manifold; connections in the image are characterized by the fact that the étale sheaf of germs of solutions is locally constant. This is a vast generalization of the phenomena of global monodromy studied in Chapter I; for instance, the differential equation $y' = y$ over the affine line belong to this class.

Uniformizing differential equations of orbifolds of dimension one also belong to this class, and the representation actually factors through the temperate fundamental group. The case of a Shimura orbifold is of special interest: the period mapping (complex or p -adic) is given by a quotient of two solutions of a fuchsian differential equation defined over a number field, which can be interpreted as uniformizing differential equation either over \mathbb{C} or over \mathbb{C}_p .

In Section 5, we examine the case of Schwarz orbifolds: the projective line with $0, 1, \infty$ as branched points (endowed with suitable finite multiplicities). Over \mathbb{C} , the uniformizing differential equations are of hypergeometric type, with projective monodromy group identified with the orbifold fundamental group, namely with a (cocompact) triangle group.

In the p -adic case, we define *p -adic triangle groups* to be projective monodromy groups of those hypergeometric differential equations which are in

the image of the p -adic étale Riemann-Hilbert functor: thus by definition, there exists a finite étale covering of $\mathbb{C}_p \setminus \{0, 1\}$ over which the hypergeometric function extends to a global multivalued analytic function.

We give a purely geometric description of these discrete subgroups of $PSL_2(\mathbb{C}_p)$ (without reference to differential equations; it is perhaps here that we are closest to Fuchs and Schwarz). From this, and recent combinatorial work by F. Kato, it follows that infinite p -adic triangle groups exist only for $p \leq 5$.

We then construct the p -adic analogues of Takeuchi's list of arithmetic triangle groups, *i.e.* the list of all "arithmetic" p -adic triangle groups for every p , using the Čerednik-Drinfeld-Boutot-Zink uniformization of Shimura curves and p -adic period mapping. Special values of the corresponding hypergeometric functions at CM points are expressed in terms of Γ_p -values.

On the style.

The first chapter is rather down-to-earth. It has kept something of the informal style of lecture notes, and this also holds to a lesser extent for the second chapter; the level is inhomogeneous, the pace may sometimes be brisk, proofs are often omitted and replaced by references. In contrast, the last and longest chapter is devoted to a systematic exposition of new material.

We hope that our constant function-theoretic viewpoint brings some unity to the exposition, throughout the chapters. We have tried to keep them (and even the sections) as logically independent as possible. Thus, for example:

- only subsections 1.5 and 5.3 of chapter I are needed in the sequel;
- until section 7, chapter II is almost self-contained;
- until subsection 4.7, chapter III is almost self-contained;
- reading fragments of III should be enough to grasp the ins and outs of p -adic triangle groups.

Sections I.3 and III.6 are small pieces of 'computational mathematics' (without computer) intended as testing ground for notions developed in the preceding sections.

Acknowledgments. It is a pleasure to acknowledge here our main sources of inspiration:

- first of all, V. Berkovich's views on p -adic geometry: without this extremely convenient way of seeing p -adic spaces, our systematic to-ings and fro-ings between complex and p -adic worlds would have been impossible;

- Rapoport-Zink's book on period spaces for p -divisible groups has been a constant reference for Chapter II;

- in Chapter III, our treatment has been much influenced by the work of M. van der Put on p -adic discontinuous groups, by J. De Jong's study

of étale coverings, and no less by M. Yoshida's orbifold viewpoint on the (complex) Gauss-Schwarz theory.

Concerning this book, I am much indebted to F. Kato in many ways: not only did he organize the Tōhoku and the Hokkaidō lectures series, asking me to speak on these matters, but he also suggested the very project of writing these notes, and influenced a lot both the content and the style by his numerous questions and comments; I have been much encouraged in writing them by his constant interest and by his own recent work on p -adic triangle groups.

The Tōhoku lectures, as well as the birth of some ideas presented in III (especially p -adic triangle groups), took place during my stay in Japan thanks to a fellowship from the Japan Society for the promotion of Science. I am very grateful to this institution, and to H. Shiga who arranged the stay.

I thank M. Matignon, F. Baldassarri and S. Bosch for invitations to lecture on these topics at the Universities of Bordeaux, Padova, and Münster respectively.

I am grateful to F. Kato and N. Tsuzuki for writing the appendices of this book. I am also very grateful to S. Matsuda who kindly took care, with great patience, of the nightmarish electronic harmonization of the main text and the appendices, and drawing the figures.

Thanks are also due to P. Colmez and O. Gabber for several useful comments, and to J.F. Boutot for sending me overseas his joint preprint with T. Zink about uniformization of Shimura curves.

I also thank very much P. Bradley for his careful reading of a version of chapter III and for kindly providing a long list of errata (of course, I am responsible for any remaining error).

CHAPTER I

Analytic aspects of p -adic periods.

Contents

1. Analytic continuation; topological point of view.	10
2. Analytic continuation; algebraic approach.	21
3. The tale of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$.	35
4. Abelian periods as algebraic integrals.	43
5. Periods as solutions of the Gauss-Manin connection.	52

1. Analytic continuation; topological point of view.

ABSTRACT: Thanks to Berkovich's presentation of p -adic analytic geometry, it is possible to make sense of the familiar monodromy principle in the exotic world of p -adic manifolds. Its application is however more limited than in the classical case, because (1) sheaves of solutions of linear differential equations are usually not locally constant, and (2) many spaces (for instance, annuli) are simply connected.

1.1. The Monodromy principle.

Let S be a topological space, and \mathcal{F} an abelian sheaf on S . For an open set $U \subseteq S$ and a section $f \in \Gamma(U, \mathcal{F})$, the support of f is the subset $\text{Supp}(f) = \{u \in U \mid f_u \neq 0\}$, which is easily seen to be closed in U .

Definition 1.1.1. We say that the sheaf \mathcal{F} satisfies the *principle of unique continuation* if for any open set $U \subseteq S$ and any section $f \in \Gamma(U, \mathcal{F})$ the support $\text{Supp}(f)$ is open in U .

Note that the principle of unique continuation is a local property.

Lemma 1.1.2. *If \mathcal{F} satisfies the principle of unique continuation, then any two sections $f, g \in \Gamma(U, \mathcal{F})$ on a connected open subset $U \subseteq S$ coincide if (and only if) their germs f_s, g_s at some point $s \in U$ coincide. The converse also holds if S is locally connected.*

PROOF. The first assertion is clear. For the converse, let u be a point adherent to the complement of $\text{Supp}(f)$. There exists a sufficiently small connected open neighborhood V of u contained in U . Since $f_s = 0$ at some point $s \in V$, $f \equiv 0$ on V . Hence $u \notin \text{Supp}(f)$. This shows that $\text{Supp}(f)$ is open. \square

Lemma 1.1.3. *Let $p : F \rightarrow S$ be the local homeomorphism canonically attached to \mathcal{F} : $\mathcal{F}(U)$ is the set of (continuous) sections of p over U . If F is separated (i.e. Hausdorff), then \mathcal{F} satisfies the principle of unique continuation. The converse also holds if S is separated.*

PROOF. If F is separated, then for any two sections f, g of p over an open subset $U \subset S$, the set of $s \in U$ such that $f(s) = g(s)$ is closed. Applying this to the zero-section $g = 0$, we see that the support of f is open.

Conversely, let x, y be two points of F . If $p(x) \neq p(y)$, there are disjoint open neighborhoods V_x, V_y of $p(x)$ and $p(y)$ respectively in the separated space S ; then $p^{-1}V_x$ and $p^{-1}V_y$ are disjoint open neighborhoods of x and y respectively. If $p(x) = p(y)$, there are open neighborhoods U_x, U_y of x and y respectively, an open subset $V \subset S$, and sections f, g of p over V , such that $f(V) = U_x$, $g(V) = U_y$. By the principle of unique continuation, $f \neq g$ defines an open subset $W \subset V$ containing $p(x) = p(y)$; then $f(W)$ and $g(W)$ are disjoint open neighborhoods of x and y respectively. \square

Definition 1.1.4. We say that the sheaf \mathcal{F} satisfies the *monodromy principle* if it has the following property:

let $\Gamma : [a, b] \times [0, 1] \rightarrow S$ be any continuous map with $\Gamma(\{a\} \times [0, 1]) = \{x_a\}$, $\Gamma(\{b\} \times [0, 1]) = \{x_b\}$. Let f_{x_a} be an element of the stalk \mathcal{F}_{x_a} . Assume that for any $t \in [0, 1]$, f_{x_a} extends to a global section f^{Γ_t} of $\Gamma_t^* \mathcal{F}$ on $[a, b]$. Then this extension is unique and $f^{\Gamma_t}(b) \in \mathcal{F}_{x_b}$ is independent of t .

In this situation, the section f^{Γ_t} of $\Gamma_t^* \mathcal{F}$ is called the *continuation of f_{x_a} along the path Γ_t* , and $f^{\Gamma_t}(b) \in \mathcal{F}_{x_b}$ its *value at x_b* .

Proposition 1.1.5. *If \mathcal{F} satisfies the principle of unique continuation, then it satisfies the principle of monodromy. The converse also holds if S is locally arcwise connected.*

PROOF. Assume that \mathcal{F} satisfies the principle of unique continuation. This guarantees the uniqueness of the continuation f^{Γ_t} of f_{x_a} along the path Γ_t for any t . Moreover, by a special case of the proper base change theorem (applied to the first projection $[a, b] \times [0, 1] \rightarrow [a, b]$, cf. [Iv86, IV, 1.4]) f^{Γ_t} extends to a section of $\Gamma^* \mathcal{F}$ on a suitable subset of the form $[a, b] \times (t - \epsilon, t + \epsilon)$. By unicity, these sections glue together to a global section of $\Gamma^* \mathcal{F}$. The restriction of this sheaf to $\{b\} \times [0, 1]$ is the constant sheaf with stalk \mathcal{F}_{x_b} . Therefore the value $f^{\Gamma_t}(b) \in \mathcal{F}_{x_b}$ is independent of t .

Conversely, let U be an arcwise connected open subset of S , let x_a, x_b be two points of U , and let f be a section of \mathcal{F} over U . Let $\gamma : [a, b] \rightarrow S$ be any path from x_a to x_b . It follows from the unicity of continuation of f_{x_a} along γ (requested in the monodromy principle) that if f_{x_a} is 0, so is f_{x_b} . \square

Example 1.1.6. (1) Any locally constant abelian sheaf \mathcal{F} on a topological space S satisfies the principle of unique continuation, hence the principle of monodromy.

In fact, for any $\Gamma : [a, b] \times [0, 1] \rightarrow S$ as in Lemma 1.1.3, the inverse image $\Gamma^* \mathcal{F}$ is locally constant, hence constant and canonically isomorphic to the constant sheaf attached to \mathcal{F}_{x_a} . Therefore the extension f^γ of any $f_{x_a} \in \mathcal{F}_{x_a}$ along any path γ exists, and the value at the other extremity $x_b = \gamma(b)$ depends only on the *homotopy class* of γ .

(2) When S is a complex manifold, the structure sheaf \mathcal{O}_S satisfies the principle of unique continuation, hence the principle of monodromy.

1.1.7. Let S be a topological space, connected and locally arcwise (or simply) connected, and \mathcal{F} a locally constant abelian sheaf on S . We fix a point $s \in S$, and denote by $\pi_1(S, s)$ the fundamental group based at s .⁽¹⁾ To any loop $\gamma : [0, 1] \rightarrow S$ based at s , let us associate the so-called *monodromy* along γ , defined by the composite

$$\mathcal{F}_s \xrightarrow{\sim} (\gamma^* \mathcal{F})_0 \xrightarrow{\sim} \Gamma([0, 1], \gamma^* \mathcal{F}) \xrightarrow{\sim} (\gamma^* \mathcal{F})_1 \xrightarrow{\sim} \mathcal{F}_s.$$

⁽¹⁾to remove any ambiguity, let us say that we adopt Deligne's convention: the composition in $\pi_1(S, s)$ is induced by the juxtaposition of loops in the reverse order. As such, $\pi_1(S, s)$ acts *on the right* on the pointed universal covering.

By 1.1.6 (1), the monodromy $\mathcal{F}_s \xrightarrow{\sim} \mathcal{F}_s$ depends only on the class of γ in $\pi_1(S, s)$, hence gives rise to a left $\mathbb{Z}[\pi_1(S, s)]$ -module structure on \mathcal{F}_s . This construction yields an equivalence of the categories

$$\{\text{locally constant sheaves on } S\} \xrightarrow{\sim} \{\text{left } \mathbb{Z}[\pi_1(S, s)]\text{-modules}\} :$$

giving a locally constant sheaf amounts to giving its value at s together with the monodromy action.

1.2. Rigid geometry and the problem of unique continuation.

1.2.1. For p a prime number, let \mathbb{Q}_p denote as usual the completion of \mathbb{Q} for the p -adic absolute value $|\cdot|_p : |p^n \frac{a}{b}|_p = p^{-n}$ if the rational integers a, b are prime to p . This ultrametric absolute value extends in a unique way to each finite extension of \mathbb{Q}_p . These finite extensions are locally compact and totally disconnected. They are all complete, but “the” algebraic closure $\overline{\mathbb{Q}_p}$ of \mathbb{Q}_p itself is not complete. Its completion, denoted by \mathbb{C}_p , turns out to be algebraically closed, and plays the role of \mathbb{C} in p -adic analysis.

In the sequel, $D(a, r^+)$ (*resp.* $D(a, r^-)$) stands for the disk — archimedean or not — of radius r centered at a with (*resp.* without) circumference.

It might be surprising at first that geometries can be built upon p -adic numbers, whose “fractal” nature makes them hardly amenable to intuition as a continuum. Nevertheless, Bourbaki’s presentation of analytic geometry [Bou83] treats the real, complex and p -adic cases on equal footing. This approach is based on a local definition of analytic functions as sums of convergent power series. Its major drawback is that these analytic functions fail to satisfy the principle of unique continuation, essentially because two ultrametric disks are either concentric or disjoint (like drops of mercury)




FIGURE 1

1.2.2. M. Krasner had the idea to remedy this by using a definition of analytic functions *à la Runge*. He properly founded ultrametric analysis by introducing his *analytic elements* defined as uniform limits of rational functions, a global notion which overcomes, to some extent, problems stemming from the disconnectedness of the p -adics.

The next step is due to J. Tate. In order to deal with more general spaces than just subsets of the line, he introduced and developed the so-called *rigid analytic geometry* (as opposed to Bourbaki’s “wobbly” analytic geometry), based on affinoid algebras (topological K -algebras isomorphic to quotients of rings of restricted formal power series, *i.e.* whose coefficients tends to 0) and a suitable Grothendieck topology [Ta71].

In a connected rigid analytic variety S , one has the following avatar of analytic continuation [Be96, 0.1.13]: assume that there is no admissible covering formed by two disjoint non-empty open subsets, and let f be an analytic function on S . If there is a connected open subset U over which f vanishes, then $f = 0$.

1.2.3. The first achievement of rigid geometry was Tate’s representation of an elliptic curve with “bad reduction at p ” as a rigid analytic quotient $\mathbb{C}_p^\times/q^{\mathbb{Z}}$ of the multiplicative group by the discrete group generated by q . Here q is given by the usual series $\frac{1}{j} + \dots$ in the j -invariant of the elliptic curve ($|j| > 1$), interpreted p -adically. The quotient is obtained by gluing two affinoid annuli of width $|q|^{-1/2}$ along their boundary; the inverse image in \mathbb{C}_p^\times of each of these annuli consists of countably many disjoint copies of it. This is analogous to Jacobi’s partial uniformization $\mathbb{C}^\times/q^{\mathbb{Z}}$ of a complex elliptic curve.

1.2.4. This result was extended by D. Mumford to curves of higher genus [Mu72b]. Here, the partial uniformization which serves as a complex model is the Schottky uniformization of complex curves of genus $g > 1$, which we briefly recall (cf. [Heh75]). Let D be an open subset of $\mathbb{P}^1(\mathbb{C})$, limited by $2g$ disjoint circles $C_1, C'_1, \dots, C_g, C'_g$. One assumes the existence, for any $i = 1 \dots, g$, of an element $\gamma_i \in PGL_2(\mathbb{C})$, with two distinct fixed points, which sends C_i to C'_i and D outside itself. Then the subgroup Γ generated by the γ_i ’s is discrete and freely generated by them, the complement $\Omega \subset \mathbb{P}^1(\mathbb{C})$ of the topological closure of the set of fixed points of Γ is an open dense subset ($= \bigcup_{\gamma \in \Gamma} \gamma(D)$), and Ω/Γ is a projective smooth complex curve of genus g .

When \mathbb{C} is replaced by \mathbb{C}_p , the same construction applies. The rigid analytic quotient Ω/Γ exists and is called a Mumford curve. Among projective smooth curves of genus g over \mathbb{C}_p , Mumford curves are characterized by the existence of a reduction over the residue field $\overline{\mathbb{F}}_p$ such that every irreducible component is isomorphic to \mathbb{P}^1 and intersect the others at double points, cf. [GvdP80].

Let us mention that T. Ichikawa has proposed a unified theory of the archimedean and non-archimedean Schottky-Mumford uniformizations, cf. [Ic97].

1.3. Berkovich geometry and the principle of monodromy

1.3.1. Rigid analytic spaces are endowed with a Grothendieck generalized topology, and their structure sheaf is a sheaf with respect to this topology. Hence it cannot be said to satisfy the principle of unique continuation in the strict sense of Definition 1.1.1. Moreover, there is no non-trivial path in such spaces. Therefore, rigid geometry is not a suitable setting for discussing p -adic analytic continuation in an intuitive way.

In contrast, Berkovich’s viewpoint on p -adic geometry does not suffer from these drawbacks: Berkovich’s analytic spaces are genuine locally ringed

topological spaces, which are even locally arcwise connected. We refer to [Ber99] for a compact technical presentation.

1.3.2. The buildings blocks are the same: affinoid algebras A (called strictly affinoid algebras by Berkovich), *i.e.* topological K -algebras isomorphic to quotients of rings of restricted power series. However, instead of attaching to A its maximal spectrum $\mathrm{Spm}(A)$ as in rigid geometry, Berkovich analytic geometry deals with the *affinoid space* $M(A)$ of all bounded multiplicative seminorms on A .

This “spectrum” contains “more points” than the maximal spectrum $\mathrm{Spm}(A)$, namely something like “generic points”, which “complete” $\mathrm{Spm}(A)$. There is a natural inclusion $\mathrm{Spm}(A) \subseteq M(A)$ which identifies $\mathrm{Spm}(A)$ with the subspace of all seminorms $|\cdot|$ with $A/\mathrm{Ker}|\cdot| = \mathbb{C}_p$. This “completion”, in fact, simplifies the topology; *e.g.* Berkovich analytic spaces are locally arcwise connected.

1.3.3. Berkovich’s affinoids. Let us be a little more precise about the definition of *affinoid spaces* in Berkovich geometry. Let A be an affinoid algebra over a complete subfield K of \mathbb{C}_p .

- (i) A point of $M(A)$ is a bounded multiplicative seminorm on A .
- (ii) The topology of $M(A)$ is the weakest one so that the mapping $M(A) \ni \chi \mapsto \chi(f) \in \mathbb{R}_{\geq 0}$ is continuous for any $f \in A$.
- (iii) The sheaf of rings $\mathcal{O}_{M(A)}$ is defined by $\Gamma(U, \mathcal{O}_{M(A)}) = \varprojlim A_V$, where V runs over finite unions $\bigcup V_i$ of affinoid domains contained in U , and $A_V = \mathrm{Ker}(\prod_i A_{V_i} \rightrightarrows \prod_{ij} A_{V_i \cap V_j})$, A_{V_i} being the affinoid algebra attached to the affinoid domain V_i .

The value of a “function” $f \in A$ at a point $\chi \in M(A)$ is its image in the field $A/\mathrm{Ker}\chi$. This field inherits the absolute value induced by the seminorm χ , and its completion is denoted by $\mathcal{H}(\chi)$.

Example 1.3.4. Let $K\{t\}$ denote the ring of restricted power series in one variable. Let us assume, for simplicity, that K is algebraically closed. In rigid geometry, the maximal spectrum $\mathrm{Spm}(K\{t\})$ is just the closed disk $D(0, 1^+)$ in K of radius 1 in the usual sense. In $M(K\{t\})$, the following four kinds of points occur:

- (1) classical points (*i.e.* those coming from $\mathrm{Spm}(K\{t\})$): $x \in D(0, 1^+)$, $\chi_x(f) = |f(x)|_K$ for $f \in K\{t\}$,
- (2) generic points of disks: $\chi = \eta_{x,r}$ for $0 < r \leq 1$ with $r \in |K^\times|$, $\chi(f) = |f|_{D(x,r^+)}$ (the sup-norm),
- (3) the same, for $r \notin |K^\times|$,
- (4) generic points of infinite decreasing families $\{D_\alpha\}$ of closed disks with radius ≤ 1 : $\chi(f) = \inf |f|_{D_\alpha}$.

Therefore, two arbitrary distinct points in $M(K\{t\})$ can be connected by a unique path. For example, two points of type (1), x and y , are connected by a path consisting of points of type (2) and (3) associated to the disks




FIGURE 2. Generic points and paths on a Berkovich space.

$D(x, r^+)$ and $D(y, r^+)$ for $0 < r \leq |x - y|$. The complement of a point of type (1) or (4) is connected. The affine line \mathbb{A}^1 is the union of the affinoid spaces associated to the algebras of power series convergent in $D(0, r^+)$, and the projective line \mathbb{P}^1 is the Alexandroff compactification of \mathbb{A}^1 .

Remark 1.3.5. Adding generic points to rigid spaces to come up with Berkovich spaces simplifies the topology in contrast to what happens in algebraic geometry, when generic points are added to varieties to produce schemes. Actually Berkovich’s generic points are closed, unlike Grothendieck’s ones.

1.3.6. The construction of (strictly) analytic K -spaces by gluing affinoid spaces together is a little delicate (one is not gluing open subspaces); we refer to [Ber93]. These spaces are *locally compact, locally countable at infinity, and locally arcwise connected*.

There is a fully faithful functor between Berkovich’s Hausdorff (strictly) analytic K -spaces and rigid analytic varieties: at the level of underlying sets, this functor sends a space S to the subset of “classical points”, *i.e.* points s for which $[\mathcal{H}(s) : K] < \infty$.

This functor establishes *an equivalence of categories between the category of paracompact (strictly) analytic K -spaces and quasi-separated rigid spaces over K having an admissible affinoid covering of finite type*.

This equivalence respects the notion of dimension (topological dimension in the case of Berkovich analytic spaces), as well as the standard properties of local rings such as: reduced, normal, smooth..., and the properties: finite, étale... of morphisms. Also, a paracompact analytic space is connected (in the usual sense) if and only if the corresponding rigid space does not admit an admissible covering by two disjoint non-empty open subsets.

Furthermore, there is a canonical functor “analytification”

$$\left\{ \begin{array}{l} \text{separated schemes} \\ \text{locally of finite type} \\ \text{over } K \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{paracompact (strictly)} \\ \text{analytic } K\text{-spaces} \end{array} \right\}$$

and a canonical functor “generic fiber”

$$\left\{ \begin{array}{l} \text{separated formal schemes locally} \\ \text{finitely presented over } \mathcal{O}_K \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{paracompact (strictly)} \\ \text{analytic } K\text{-spaces} \end{array} \right\}.$$

1.3.7. p -adic manifolds. We shall be mainly concerned with paracompact (strictly) K -analytic spaces S which satisfy the following assumption:

any $s \in S$ has a neighborhood $U(s)$ which is isomorphic to an affinoid subdomain of some space V_s which admits locally an étale morphism to the affine space $\mathbb{A}^{\dim S}$.

For convenience, we shall call such a space S a K -manifold, and simply a p -adic manifold if $K = \mathbb{C}_p$.

It is an important result of V. Berkovich [Ber99] that p -adic manifolds are locally contractible. Therefore they have universal coverings.

Proposition 1.3.8. *The structure sheaf of any p -adic manifold S satisfies the principle of unique continuation, and (equivalently) the principle of monodromy.*

PROOF. Since the principle of unique continuation is local, we may replace S by the neighborhood $U(s)$ of any point s given in advance. Thus we may assume that S itself is affinoid: $S = M(A)$, and connected, and we have to show that the homomorphism $\iota_s : A \rightarrow \mathbb{A}_s \simeq \varinjlim_{V \ni s} A_V$ is injective (V runs over the affinoid neighborhoods of s). This follows from the fact that the homomorphism of completion at $s : A \rightarrow \widehat{A}_s = \varprojlim_n A/\mathcal{I}_s^n$ is injective and factors through ι_s . □

In the case of curves (*i.e.* p -adic manifolds of dimension 1), analytic continuation is particularly intuitive, because there is a basis of open subsets \mathcal{U} with finite boundary such that two arbitrary points in \mathcal{U} are connected by a unique geometric path lying in \mathcal{U} .

1.4. Topological coverings and étale coverings.

1.4.1. Complex manifolds are locally contractible, and have universal coverings. There is no need to distinguish between topological coverings and étale coverings (finite or infinite).

In p -adic rigid geometry, the situation is more complicated. It is natural to call topological covering any morphism $f : Y \rightarrow X$ such that there is an admissible cover (X_i) of X and an admissible cover (Y_{ij}) of $f^{-1}(X_i)$ with disjoint Y_{ij} isomorphic to (X_i) via f . Indeed, such topological coverings correspond to locally constant sheaves of sets on X . It is still true that *topological coverings are étale, but the converse is wrong*, even if one restricts to finite surjective morphisms. Indeed, the Kummer covering $z \mapsto z^n$ of the punctured unit disk is an étale covering, but not a topological covering, if $n > 1$.

1.4.2. It is again more convenient to tackle these questions in the framework of Berkovich's geometry. For instance, one sees immediately that a Kummer covering $z \mapsto z^n$ as above is not a topological covering because a classical point has n preimages, while the "generic point" $\eta_{0,1}$ (corresponding to the sup-norm on the disk) is its own single preimage. Topological coverings of a p -adic manifold X are defined in the usual way; they correspond to locally

constant sheaves of sets on X . They coincide [Ber90, 3.3.4] with topological coverings of the rigid analytic variety associated to X .

1.4.3. There are only three one-dimensional simply connected complex manifolds up to isomorphism: the projective line $\mathbb{P}^1(\mathbb{C})$, the affine line \mathbb{C} , and the disk $\mathbb{D} (\simeq \mathfrak{h}$, the complex upper half plane). In contrast, there are plenty of *simply-connected* one-dimensional p -adic manifolds, for instance: any annulus, the line deprived from a finite number of points, more generally any connected p -adic manifold homeomorphic to a *subset of* $\mathbb{P}^1(\mathbb{C}_p)$, any smooth projective curve with “good reduction”... (all these Berkovich spaces look like “bushy trees”).

1.4.4. One defines the (discrete) topological fundamental group $\pi_1^{\text{top}}(S, s)$ of a pointed p -adic manifold (S, s) in the usual way. The general topological theory of coverings applies: (S, s) is naturally isomorphic to the quotient of the pointed universal covering (\tilde{S}, \tilde{s}) by $\pi_1^{\text{top}}(S, s)$, and $\pi_1^{\text{top}}(S, s)$ classify the topological coverings of (S, s) . In the one-dimensional case, the topological fundamental group is a discrete *free* group (more precisely, it is isomorphic to the fundamental group of the dual graph of the so-called semistable reduction of S ([dJ95b, 5.3])).

For instance, if $S = \Omega/\Gamma$ is the uniformization of a Mumford curve (cf. 1.2), Ω is the universal covering of S and the Schottky group Γ is isomorphic to the topological fundamental group.

On the other hand, one can define the (profinite) algebraic fundamental group $\pi_1^{\text{alg}}(S, s)$ à la Grothendieck, classifying the finite étale coverings of (S, s) . In contrast to the complex situation, the natural map $\pi_1^{\text{alg}}(S, s) \rightarrow \pi_1^{\text{top}}(S, s)^\wedge$ to the profinite completion of $\pi_1^{\text{top}}(S, s)$ is *generally not injective* (e.g. for annuli).

1.5. Connections with locally constant sheaves of solutions.

1.5.1. Let us briefly recall the complex situation. Let S be a complex connected manifold, (\mathcal{M}, ∇) a vector bundle of rank r with integrable connection on S . The classical Cauchy theorem shows that for any $s \in S$, the solution space $(\mathcal{M} \otimes \mathcal{O}_{S,s})^\nabla$ at s has dimension r . Analytic continuation along paths gives rise to a homomorphism $\pi_1^{\text{top}}(S, s) \rightarrow \text{Aut}_{\mathbb{C}}((\mathcal{M} \otimes \mathcal{O}_{S,s})^\nabla)$ (the *monodromy*). The sheaf of germs of solutions \mathcal{M}^∇ is locally constant on S : its pull-back over the universal covering \tilde{S} of S is constant. Conversely, any complex representation V of $\pi_1^{\text{top}}(S, s)$ of dimension r gives rise naturally to a vector bundle \mathcal{M} of rank r with integrable connection ∇ : $\mathcal{M} = (V \times \tilde{S})/\pi_1^{\text{top}}(S, s)$, $\nabla(V) = 0$. This sets up an equivalence of categories:

$$\left\{ \begin{array}{l} \text{finite dimensional} \\ \text{representations of } \pi_1^{\text{top}}(S, s) \end{array} \right\} \simeq \left\{ \begin{array}{l} S\text{-vector bundles with} \\ \text{integrable connection} \end{array} \right\}.$$

1.5.2. Let (S, s) be now a pointed connected p -adic manifold. It is still true that any \mathbb{C}_p -linear representation V of $\pi_1^{\text{top}}(S, s)$ of dimension r gives

rise to a vector bundle $\mathcal{M} = \mathcal{M}_V$ a vector bundle of rank r with integrable connection $\nabla = \nabla_V$ (same formula). The functor

$$\left\{ \begin{array}{c} \text{finite dimensional} \\ \text{representations of } \pi_1^{\text{top}}(S, s) \end{array} \right\} \rightarrow \left\{ \begin{array}{c} S\text{-vector bundles with} \\ \text{integrable connection} \end{array} \right\}$$

is still fully faithful, but *no longer surjective*; its essential image consists of those connections whose sheaf of solutions is locally constant, *i.e.* becomes constant over \tilde{S} . In fact, the classical “*Cauchy theorem*” according to which the solution space $(\mathcal{M} \otimes \mathcal{O}_{S,s})^\nabla$ at s has dimension r holds for every classical point s of S — which corresponds to a point of the associated rigid variety —, but *does not hold* for non-classical points s of the Berkovich space S in general.

1.5.3. Any non-trivial connection over the projective line minus a few points gives an example when the p -adic analogue of the “*Cauchy theorem*” does not hold: indeed, in this case, the topological fundamental group is trivial.

However, a p -adic variant of Cauchy’s theorem in the neighborhood of a non-classical point s may be restored as follows (Dwork’s technique of generic points): it suffices to extend the scalars from \mathbb{C}_p to a complete algebraically closed extension of $\mathcal{H}(s)$. This transforms s into a classical point, and neighborhoods of s after scalar extensions are “smaller” than before.

1.5.4. When “*Cauchy’s theorem*” holds at every point of S , one can continue the local solutions along paths and get the monodromy representation just as in the complex situation.

This nice category of p -adic connections has not yet attracted much attention.

Example 1.5.5. Let us consider the case when S is a Tate elliptic curve: $S = \mathbb{C}_p^\times / q^\mathbb{Z}$, with $s =$ its origin. Then $\pi_1^{\text{top}}(S, s) = q^\mathbb{Z}$, and the connections on S which arise from representations of $q^\mathbb{Z}$ are those which become trivial over $\tilde{S} = \mathbb{C}_p^\times$. In this correspondence, the multivalued solutions \vec{y} (*i.e.* horizontal sections) of such a connection are the solutions of the associated linear q -difference equation $\vec{y}(qt) = M(q)\vec{y}(t)$, where t is the standard coordinate on \mathbb{C}_p^\times and the matrix $M(q)$ is the image of q in the monodromy representation.

The simplest example is given by the obvious one-dimensional representation $M(q) = q$. It corresponds to $\mathcal{M} = \mathcal{O}_S$, $\nabla(1) = \omega_{\text{can}}$ (the canonical differential on S induced by dt/t); here the q -difference equation is $t.dy = y.dt$ with obvious solution $y = t$.

Let us now look at the representation $M(q) = \sqrt{q}$. The associated q -difference equation is $t.dy = \frac{1}{2}y.dt$. There is the obvious solution \sqrt{t} , which leads to $\mathcal{M} = \mathcal{O}_S$, $\nabla(1) = \frac{1}{2}\omega_{\text{can}}$. Here we encounter an interesting paradox: \sqrt{t} is not a multivalued analytic function on S (*i.e.* it is not an analytic function on the universal covering \mathbb{C}_p^\times). In the complex situation, a similar paradox arose in the work of G. Birkhoff in his theory of q -difference

equations, which was pointed out and analyzed by M. van der Put and M. Singer in the last chapter of their book [SvdP97]. The solution of the paradox is that the vector bundle \mathcal{M} associated to the representation $q \rightarrow \sqrt{q}$ of $q^{\mathbb{Z}}$ (or to the q -difference equation $y(qt) = \sqrt{q}.t$) is in fact a non-trivial vector bundle of rank one, and the basic solution is not \sqrt{t} , but $\frac{\theta(t/\sqrt{q})}{\theta(t)}$, where $\theta(t) = \prod_{n>0}(1 - q^n t) \prod_{n\leq 0}(1 - q^n/t)$.

1.5.6. In the previous example, it is easily seen that rank-one vector bundles with connection on S which arise from a representation of $\pi_1^{\text{top}}(S, s)$ form a space of dimension one, while the space of all rank-one vector bundles with connection on S has dimension two.

We next consider the case of a p -adic manifold S which “is” a geometrically irreducible algebraic \mathbb{C}_p -curve. Let \overline{S} be its projective completion. It follows from the Van Kampen theorem, together with the fact that punctured disks are simply-connected, that $\pi_1^{\text{top}}(S, s) \rightarrow \pi_1^{\text{top}}(\overline{S}, s)$ is an isomorphism. It follows that the vector bundles with connection attached to representations of $\pi_1^{\text{top}}(S, s)$ automatically extend to \overline{S} . Hence we may assume without loss of generality that S is compact.

Vector bundles with connection on S are algebrizable, and one can use C. Simpson’s construction ([Si94]) to define the *moduli space of connections* of rank r over S , denoted by $M_{\text{dR}}(S, r)$. On the other hand, we have seen that the topological fundamental group $\pi_1^{\text{top}}(S, s)$ is free on $b_1(\Delta)$ generators, being isomorphic to the fundamental group of the dual graph Δ of the semistable reduction of S . Simpson has also studied the *moduli space of representations of dimension r* of such a group. We denote it by $M_{\text{B}}(S, r)$; in fact, it depends only on the couple of integers $(b_1(\Delta), r)$.

Let us assume that S is of genus $g \geq 2$. Simpson shows that $M_{\text{dR}}(S, r)$ is algebraic irreducible of dimension $2(r^2(g - 1) + 1)$. On the other hand, $M_{\text{B}}(S, r)$ is an algebraic irreducible affine variety of dimension $(r^2(b_1(\Delta) - 1) + 1)$. We note that this dimension is maximal when the Betti number $b_1(\Delta)$ takes its maximal value, namely g . This corresponds to the case where S is a Mumford curve (a curve with totally degenerate reduction).

It turns out that the functor which associates a vector bundle with connection to any finite-dimensional representation of the topological fundamental group induces an injective analytic map of moduli spaces $\iota : M_{\text{B}}(S, r) \rightarrow M_{\text{dR}}(S, r)$.

[The map is induced by the functor $V \rightarrow (\mathcal{M}_V, \nabla_V)$, and its injectivity follows from the faithfulness of this functor. The difficulty in showing that ι is analytic is that $M_{\text{dR}}(S, r)$ is a priori a moduli space for algebraic connections, not for all analytic connections; we shall not pursue here in this direction].

In the complex situation, the corresponding map ι would always be an analytic isomorphism (Riemann-Hilbert-Simpson). In the p -adic cases, we see that the connections which satisfy Cauchy’s theorem at every point

(classical or not) form a stratum of maximal dimension half of the dimension of the total moduli space.

We shall leave the closer analysis of this kind of p -adic differential equations with global monodromy until later chapters, where we show how they arise in the context of period mappings. In the next section, we shall deal with a very different kind of differential equations, which play a distinguished role in p -adic analysis under the name of *unit-root F -crystals*.

2. Analytic continuation; algebraic approach.

ABSTRACT: An equivalent, but more algebraic, approach to analytic continuation consists in interpreting complex analytic multivalued functions as limits of algebraic functions. Since the concept of topological coverings and that of étale coverings do not coincide in the p -adic setting, the algebraic approach leads in that case to a theory quite different from that of the previous section. It turns out to be well-suited to the function-theoretic study of so-called unit-root F -isocrystals, and induces us to revisit two fundamental notions due to Dwork: Frobenius structure and overconvergence.

2.1. Limits of rational functions.

2.1.1. Over \mathbb{C} . Let us briefly survey the complex analytic situation. Let S be a connected complex analytic curve, and U an open set of S (not necessarily connected) which is holomorphically convex; *i.e.* $S \setminus U$ has no compact connected component. Then $\mathcal{O}(S)$ is dense in $\mathcal{O}(U)$ for the topology of uniform convergence on every compact set. We denote this situation by $\mathcal{O}(U) = \widehat{\mathcal{O}(S)}^U$ (*cf.* *e.g.* [Re89, 13]).

Moreover, if S is the Riemann surface coming from an affine algebraic curve S^{alg} , then we also have $\mathcal{O}(U) = \widehat{\mathcal{O}(S^{\text{alg}})}^U$, which generalizes the theorem of Runge on approximation by rational functions. Indeed, considering an embedding $S^{\text{alg}} \hookrightarrow (\mathbb{A}^N)^{\text{alg}}$, we may extend analytic functions on S to analytic functions on $(\mathbb{A}^N)^{\text{alg}}$ (due to the vanishing of the first cohomology group of the coherent ideal sheaf defining S). An analytic functions on $(\mathbb{A}^N)^{\text{alg}}$ can be approximated by polynomials, which we restrict to S .

2.1.2. Over the p -adics. We have an analogous situation. Let K be a complete subfield of \mathbb{C}_p , and let S be an analytic curve coming from a smooth affine algebraic curve over K . We consider a closed immersion $S \hookrightarrow \mathbb{A}^N$. For any $r > 0$, $S_r := S \cap D_{\mathbb{A}^N}(0, r^+)$ is an affinoid domain in S . The same argument of polynomial approximation used in 2.1.1 shows that $\mathcal{O}(S_r)$ is the completion of $\mathcal{O}(S^{\text{alg}})$.

For any compact $Z \subset S$ (*e.g.* an affinoid domain), we define the topological ring $\mathcal{H}(Z)$ of *analytic elements* on Z to be the completion of $\Gamma(Z, \mathcal{O}_S) = \varinjlim_{Z \subset U: \text{open}} \Gamma(U, \mathcal{O}_S)$ under the sup-norm. Whenever Z is contained in a S_r , $\mathcal{H}(Z)$ is also the completion of $\mathcal{O}(S_r)_Z$, where the subscript Z denotes the localization with respect to the set of elements which do not vanish on Z (*cf.* [Ber90] and [Ray94] for a more precise version of the Runge theorem).

The holomorphic convexity condition on Z is that for some r , Z is the intersection of affinoid neighborhoods defined by inequalities of the form $|f_i| \leq 1$ in S_r . If this condition is satisfied, $\mathcal{H}(Z)$ is in fact the completion of $\mathcal{O}(S_r)$ itself.

2.1.3. The Krasner-Dwork viewpoint. For $S = \mathbb{A}^1$, and $K = \mathbb{C}_p$, we deduce $\mathcal{H}(Z) = \widehat{(K[z]_Z)}$, which is nothing but the M. Krasner's original definition of analytic elements on Z .

Let $\widehat{K}(z)$ be the completion of $K(z)$ by means of Gauss norm. One can interpret its elements as the analytic elements on a generic disk, and then specialize in the complement $Z \subseteq \mathbb{P}^1$ of a finite union of disks $D(a_i, 1^-)$, ($i = 1, 2, \dots$). This viewpoint leads us to understand analytic continuation as a *specialization*.

2.2. Limits of algebraic functions; complex case.

Let S be a Riemann surface coming from a complex affine algebraic curve S^{alg} , smooth and connected. Let \tilde{S} be the universal covering of S (here we tacitly fix a base point $s \in S$). We endow $\mathcal{O}(\tilde{S})$ with the topology of uniform convergence on every compact; note that the group $\pi_1(S)$ acts (continuously) on $\mathcal{O}(\tilde{S})$.

On the other hand, let $\mathcal{O}(S^{\text{alg}})^{\text{et}}$ be the maximal unramified integral extension of $\mathcal{O}(S^{\text{alg}})$. Elements in $\mathcal{O}(S^{\text{alg}})^{\text{et}}$ can be regarded as unramified algebraic functions on S^{alg} ; thus we have $\mathcal{O}(S^{\text{alg}})^{\text{et}} \subset \mathcal{O}(\tilde{S})$.

Proposition 2.2.1. $\mathcal{O}(S^{\text{alg}})^{\text{et}}$ is dense in $\mathcal{O}(\tilde{S})$.

For example, when $S^{\text{alg}} = \mathbb{P}^1 \setminus \{0, \infty\}$, the function $\log z \in \mathcal{O}(\tilde{S})$ can be written as $\lim_{n \rightarrow \infty} n(z^{1/n} - 1)$, uniformly on every compact subset of $\tilde{S} \simeq \mathbb{A}^1$. We can use this formula to compute $\log_\gamma 1 = \lim n(\zeta_n - 1) = 2i\pi$, where γ is the counter-clockwise loop around 0 with the base point 1.



FIGURE 3

PROOF OF 2.2.1. (after O. Gabber). Consider a countable covering of \tilde{S} ($\simeq \mathbb{P}^1(\mathbb{C}), \mathbb{C}$ or $D(0, 1^-)$) by relatively compact contractible open subsets which are oriented manifolds U_n (e.g. disks), such that $\overline{U}_n \subset U_{n+1}$. For any n , let us consider the set

$$\mathcal{F}_n = \{\gamma \in \pi_1(S, s) \mid \gamma \neq 1 \text{ and } \gamma \overline{U}_n \cap \overline{U}_n \neq \emptyset\}.$$

If this set were infinite, we could find a sequence of pairwise distinct elements $\gamma_m \in \pi_1(S, s)$ and a sequence of points $s_m \in \overline{U}_n$ such that $\gamma_m s_m \in \overline{U}_n$. By compactness of \overline{U}_n , we might assume that the sequences s_m and $\gamma_m s_m$ converge to points s' and s'' respectively. Then the sequence $\gamma_m s'$ converges to s'' , which contradicts the discreteness of the $\pi_1(S, s)$ -orbits in \tilde{S} . So \mathcal{F}_n is finite.

On the other hand, S is topologically a surface of genus g with $t \geq 1$ punctures, so $\pi_1(S)$ is generated by $2g + t$ generators γ_i subject to the relation

$$[\gamma_1, \gamma_{g+1}] \cdots [\gamma_g, \gamma_{2g}] \gamma_{2g+1} \cdots \gamma_{2g+t} = 1.$$

Thus $\pi_1(S)$ is free with $2g + t - 1$ generators. Therefore it is residually finite (*cf.* [Bou64, p. 150, ex. 34]); so we can find a subgroup $\Gamma_n \subseteq \pi_1(S)$ of finite index which avoids \mathcal{F}_n .

We immediately see that $((\Gamma_n \setminus \{1\}) \cdot \overline{U}_n) \cap \overline{U}_n = \emptyset$. The restriction to \overline{U}_n of the canonical projection $\tilde{S} \rightarrow S_n := \tilde{S}/\Gamma_n$ is thus an embedding. We identify \overline{U}_n with its image.

The part of the exact sequence of cohomologies with compact support

$$\cdots \longrightarrow H^1(\overline{U}_n, \mathbb{Z}) \longrightarrow H_c^2(S_n \setminus \overline{U}_n, \mathbb{Z}) \longrightarrow H_c^2(S_n, \mathbb{Z}) \longrightarrow \cdots$$

which can also be written as

$$\cdots \rightarrow H_1(\overline{U}_n, \mathbb{Z}) = 0 \rightarrow H_0(S_n \setminus \overline{U}_n, \mathbb{Z}) \rightarrow H_0(S_n, \mathbb{Z}) = \mathbb{Z} \rightarrow \cdots$$

and from which we deduce that $S_n \setminus \overline{U}_n$ is connected. Because $\partial U_n \subset \overline{U}_{n+1} \setminus \overline{U}_n$, we have $S_n \setminus \overline{U}_n = S_n \setminus U_n$, hence $S_n \setminus U_n$ is connected. Since $S_n \setminus U_n$ is not compact, U_n is holomorphically convex in S_n . Riemann's existence theorem assures that S_n is the analytification of an affine algebraic curve S_n^{alg} . Here we can apply 2.1.1 to deduce $\mathcal{O}(U_n) = \widehat{\mathcal{O}(S_n^{\text{alg}})}_{U_n}$. Note that $\mathcal{O}(S_n^{\text{alg}}) \subset \mathcal{O}(S^{\text{alg}})^{\text{et}}$. Since every compact set of \tilde{S} is contained in some U_n , we conclude the desired equality $\mathcal{O}(\tilde{S}) = \widehat{\mathcal{O}(S^{\text{alg}})}^{\text{et}}$. \square

2.2.2. It follows from Proposition 2.2.1 that if Ω/Γ is a Schottky partial uniformization of S , then the intersection $\mathcal{O}(S^{\text{alg}})^{\text{et}} \cap \mathcal{O}(\Omega)$ is dense in $\mathcal{O}(\Omega)$. A variant of this statement (with a similar proof) holds, in the p -adic case, for a Mumford curve. We leave it to the reader.

2.3. Limits of algebraic functions; p -adic case.

2.3.1. In the p -adic situation, we have seen that there are “much more” étale coverings than topological coverings. For instance, the unit disk $\mathbb{D} = \mathbb{D}(0, 1^+)$ is simply-connected, but admits many non-trivial finite étale coverings, *e.g.* the Artin-Schreier covering $\mathbb{D} \rightarrow \mathbb{D}$, $y \mapsto z = y^p - y$.

Proposition 2.2.1 suggests to replace, in the p -adic case, the ring $\mathcal{O}(\tilde{S})$, which is often too small, by some kind of completion of $\mathcal{O}(S)^{\text{et}} = \varinjlim \mathcal{O}(S')$, where S' runs over the finite étale connected coverings of S .

Let us now assume that S is an affinoid curve with good reduction (hence simply connected). Then there is a Gauss p -adic norm on $\mathcal{O}(S)$, which extends uniquely to $\mathcal{O}(S)^{\text{et}}$. The completion $\widehat{\mathcal{O}(S)^{\text{et}}}$ is however pathological in several senses: for instance, it is difficult to give a function-theoretic meaning to its elements, and the continuous derivations of $\mathcal{O}(S)$ extends to $\mathcal{O}(S)^{\text{et}}$ but not to $\widehat{\mathcal{O}(S)^{\text{et}}}$ in a natural way.

2.3.2. To bypass such problems, we are going to look for some convenient *subring* of $\widehat{\mathcal{O}(S)}^{\text{et}}$. For simplicity, let us limit ourselves to the following situation:

- $K = \widehat{\mathbb{Q}}_p^{\text{ur}} \subset \mathbb{C}_p$ is the completion of the maximal unramified algebraic extension of \mathbb{Q}_p ,
- $\mathfrak{v} = \widehat{\mathbb{Z}}_p^{\text{ur}}$ is its ring of integers (the Witt ring of $\overline{\mathbb{F}}_p$),
- S_0 is the affine line over $\overline{\mathbb{F}}_p$, deprived from finitely many points $\overline{\zeta}_1 \dots, \overline{\zeta}_\nu$; we choose a point s_0 on S_0 ,
- \mathcal{R} is the p -adic completion of $\mathfrak{v}[z, \frac{1}{(z-\zeta_1)\dots(z-\zeta_\nu)}]$, where $\zeta_1 \dots \zeta_\nu$ are liftings of $\overline{\zeta}_1 \dots, \overline{\zeta}_\nu$ in \mathfrak{v} ; its residue ring is $\mathcal{O}(S_0)$,
- $S = \mathcal{M}(\widehat{\mathcal{R}}[\frac{1}{p}]) = \text{D}(0, 1^+) \setminus \bigcup \text{D}(\zeta_i, 1^-)$, the associated affinoid domain over K . There is a natural specialization map $\text{sp} : S \rightarrow S_0$ from characteristic 0 to characteristic p .

Let us consider the integral closure \mathcal{R}^{et} of \mathcal{R} in $\mathcal{O}(S)^{\text{et}}$ and its p -adic completion $\widehat{\mathcal{R}^{\text{et}}} \subset \widehat{\mathcal{O}(S)^{\text{et}}}$. It is thus endowed with (a natural extension of) the p -adic valuation, and its residue ring is $\mathcal{O}(S_0)^{\text{et}}$. Moreover, there is a natural structure of “differential ring” (better: a connection) on $\widehat{\mathcal{R}^{\text{et}}}$ with \mathfrak{v} as ring of constants. The “remarkable equivalence of categories” of A. Grothendieck [EGA IV, 18.1.2] allows to identify $\text{Aut}_{\text{cont}}(\widehat{\mathcal{R}^{\text{et}}}/\mathcal{R})$ with the algebraic fundamental group $\pi_1^{\text{alg}}(S_0, s_0)$.

2.3.3. “Analytic continuation” as specialization. Let $\text{D}(s_0, 1^-) = \text{sp}^{-1}\{s_0\}$ be the residue class of s_0 in S . The morphism $\mathcal{R} \rightarrow \mathcal{O}(\text{D}(s_0, 1^-))$ extends non-canonically to a continuous morphism $\widehat{\mathcal{R}^{\text{et}}}[\frac{1}{p}] \rightarrow \mathcal{O}(\text{D}(s_0, 1^-))$, determined *only up to the action of* $\pi_1^{\text{alg}}(S_0, s_0)$. This allows to interpret elements of $\widehat{\mathcal{R}^{\text{et}}}[\frac{1}{p}]$ as certain multivalued locally analytic functions (in the “wobbly” sense).

The main difference here with Krasner’s analytic continuation (*cf.* §§2.1) is the ambiguity coming from $\pi_1^{\text{alg}}(S_0, s_0)$. This provides a kind of multivalued analytic continuation, which may be interpreted as analytic continuation in characteristic 0 along an “étale path” in characteristic p (figure 4).




FIGURE 4. analytic continuation along an underground path

This is very close to the Robba-Christol theory of “algebraic elements” [C86], in which the main player is the *complete* subalgebra of $H^\infty(\mathbb{D}_{\mathbb{C}_p}(0, 1^-))$ (the algebra of bounded analytic functions) generated the algebraic functions analytic on $\mathbb{D}_{\mathbb{C}_p}(0, 1^-)$. Underlying this “algebraic” analytic continuation, there is a combinatorics of *automata* (whereas a combinatorics of graphs underlies the “topological” analytic continuation, as we have seen).⁽²⁾

Remark 2.3.4. The complex formula $\log z = \lim_n n(z^{\frac{1}{n}} - 1)$ has the following p -adic analogue: $\log z = \lim_n p^{-n}(z^{p^n} - 1)$ for $z \in \mathbb{D}(1, 1^-)$. However, the convergence is not uniform on $\mathbb{D}(1, 1^-)$, and the analytic function $\log z$ is not bounded on that disk. Nevertheless $\log : \mathbb{D}(1, 1^-) \rightarrow \mathbb{A}^1$ defines an infinite Galois étale covering of the p -adic affine line (not at all a topological covering), with Galois group μ_{p^∞} , the p -primary torsion in \mathbb{C}_p .

2.4. $\widehat{\mathcal{R}}^{\text{et}}$ and unit-root F -crystals.

2.4.1. Let us assume for technical simplicity that $p \neq 2$. Let \mathcal{M} be a free \mathcal{R} -module of finite rank μ , endowed with a connection $\nabla : \mathcal{M} \rightarrow \Omega_{\mathcal{R}} \otimes_{\mathcal{R}} \mathcal{M}$, where $\Omega_{\mathcal{R}}$ is the rank-one module of continuous differentials (relative to \mathfrak{v}).

In general, solutions make sense only very locally: typically, analytic solutions exist in disks of radius $p^{\frac{-1}{p-1}}$ (the radius of p -adic convergence of the exponential function, which is the basic example). A very important criterion, due to B. Dwork, for the convergence of analytic solutions in any open disk of radius 1, is the existence of a so-called Frobenius structure.

2.4.2. F -crystals. Let σ be the Frobenius automorphism of \mathfrak{v} lifting the p th-power map in $\overline{\mathbb{F}}_p$. There are many σ -linear endomorphisms ϕ of \mathcal{R} which reduce to the p th-power map of $\mathcal{O}(S_0)$ in characteristic p . For instance, we can take $\phi(z) = z^p$, so that $\phi((z - \zeta_1) \dots (z - \zeta_\nu)) = (z^p - \sigma(\zeta_1)) \dots (z^p - \sigma(\zeta_\nu)) \equiv ((z - \zeta_1) \dots (z - \zeta_\nu))^p \pmod{p}$. One can show that any lifting ϕ extends uniquely to a $\pi_1^{\text{alg}}(S_0, s_0)$ -equivariant endomorphism of $\widehat{\mathcal{R}}^{\text{et}}$.

A Frobenius structure on (\mathcal{M}, ∇) is a rule F which associates to every lifting ϕ a homomorphism $F(\phi) : \phi^* \mathcal{M} \rightarrow \mathcal{M}$ (that is to say, a ϕ -linear endomorphism of \mathcal{M}) such that:

- (i) $F(\phi) \otimes \mathbb{Q}$ is an *isomorphism*,
- (ii) $F(\phi)$ is *horizontal*, i.e. compatible with the connections $\phi^* \nabla$ and ∇ respectively,
- (iii) for any two liftings ϕ and ϕ' , the homomorphisms $F(\phi)$ and $F(\phi')$ are related by $F(\phi') = F(\phi) \circ \chi(\phi', \phi)$, where $\chi(\phi', \phi) : \phi'^* \mathcal{M} \rightarrow \phi^* \mathcal{M}$

⁽²⁾for these and further aspects of the Krasner, resp. Christol-Robba, analytic continuation, we refer to [MoRo69] and to the mimeographed notes of the numerous conferences of the Groupe d'étude d'analyse ultramétrique, Paris, devoted to this subject (1973-1980).

is the ‘‘Taylor isomorphism’’ given by the formulas

$$\begin{aligned}\chi(\phi', \phi)(\phi'^* m) &= \sum_{n \geq 0} \phi^* \left(\nabla \left(\frac{d}{dz} \right)^n (m) \right) (\phi'(z) - \phi(z))^n / n! \\ &= \sum_{n \geq 0} \phi^* \left(\nabla \left(z \frac{d}{dz} \right)^n (m) \right) \left(\log \frac{\phi'(z)}{\phi(z)} \right)^n / n!.\end{aligned}$$

In virtue of (iii), a Frobenius structure amounts to the data of the semi-linear horizontal endomorphism $F(\phi)$, for the standard $\phi(z) = z^p$.

The triple (\mathcal{M}, ∇, F) is called an F -crystal, cf. [Ka73]⁽³⁾. The very existence of a horizontal isomorphism $\phi^* \mathcal{M} \otimes \mathbb{Q} \simeq \mathcal{M} \otimes \mathbb{Q}$ implies that the radius of convergence ρ of any local solution of (\mathcal{M}, ∇) satisfies $\min(1, \rho) = \min(1, \rho^p)$, that is to say: $\rho \geq 1$ (since $\rho \neq 0$).

2.4.3. Unit-root F -crystals. This is the case where $F(\phi)$ (not only $F(\phi) \otimes \mathbb{Q}$) is an isomorphism (for one, or equivalently, for all ϕ). The name comes from their first appearance in Dwork’s computation of the p -adic units among the reciprocal zeroes of the zeta-function of hypersurfaces in characteristic p .

N. Katz has constructed a functor:

$$\left(\begin{array}{c} \text{continuous } \mathbb{Z}_p\text{-representations} \\ \text{of } \pi_1^{\text{alg}}(S_0, s_0) \end{array} \right) \longrightarrow (\text{unit-root } F\text{-crystals})$$

which associates to any free \mathbb{Z}_p -module V of rank r with a continuous action of $\pi_1^{\text{alg}}(S_0, s_0)$ a unit-root F -crystal $(\mathcal{U}_V, \nabla_V, F_V)$ over \mathcal{R} of rank r . Let us recall its definition at the level of objects. For any $m \in \mathbb{N}$, we set $S_m = \text{Spec } \mathcal{R}/p^m \mathcal{R}$. Starting from a representation ρ of $\pi_1^{\text{alg}}(S_0, s_0)$, let G_n denote the image of ρ in $GL(V/p^n V)$. The homomorphism $\pi_1^{\text{alg}}(S_0, s_0) \rightarrow G_n$ corresponds to an étale covering $S_{n,0} \rightarrow S_0$, which has a unique étale lifting $\pi_{n,m} : S_{n,m} \rightarrow S_m$. The action of G_n on $S_{n,0}$ extends uniquely to $S_{n,m}$; on the other hand, the action of ϕ on S_m extends uniquely to $S_{n,m}$, and the ϕ - and G_n -actions commute. The opposite action makes $\mathcal{O}_{S_{n,m}}$ into a right G_n -module. If we set $\mathcal{U}_n = \pi_{n,n*} \mathcal{O}_{S_{n,n}} \otimes_{(\mathfrak{v}/p^n \mathfrak{v})[G_n]} V$, we then have a compatible system of isomorphisms

$$\Phi_n = \phi \otimes \text{id} : \phi^* \mathcal{U}_n \rightarrow \mathcal{U}_n$$

and a compatible system of connections

$$\nabla_n = d \otimes \text{id} : \mathcal{U}_n \rightarrow \pi_{n,n*} \Omega_{S_{n,n}} \otimes_{\mathcal{R}/p^n \mathcal{R}} \mathcal{U}_n \simeq \Omega_{\mathcal{R}} \otimes_{\mathcal{R}} \mathcal{U}_n$$

The unit-root crystal attached to ρ is given by $\mathcal{U} = \varprojlim_n \mathcal{U}_n$, $\nabla = \varprojlim_n \nabla_n$, $\Phi = \varprojlim_n \Phi_n$.

The next statement summarizes results of Katz and R. Crew [Cr85].

Proposition 2.4.4. (i) *The Katz functor is an equivalence of categories;*

⁽³⁾the endomorphism F_{S_0} of S_0 given by the p th-power map, as well as σ , ϕ and F are all called ‘‘Frobenius’’ in the usual jargon, without causing too much confusion, it seems...

- (ii) any unit-root F -crystal (\mathcal{U}, ∇, F) is solvable in $\widehat{\mathcal{R}}^{\text{et}}$, i.e. : $\mathcal{U} \otimes_{\mathcal{R}} \widehat{\mathcal{R}}^{\text{et}} \simeq (\mathcal{U} \otimes_{\mathcal{R}} \widehat{\mathcal{R}}^{\text{et}})^{\nabla} \otimes_{\mathfrak{v}} \widehat{\mathcal{R}}^{\text{et}}$;
- (iii) the rule $(\mathcal{U}, \nabla, F) \mapsto (\mathcal{U} \otimes_{\mathcal{R}} \widehat{\mathcal{R}}^{\text{et}})^{\nabla=0, F(\phi)=\text{id}}$ (with Galois action coming from that on $\widehat{\mathcal{R}}^{\text{et}}$) provides an inverse of the Katz functor.

More precisely, we have the action of ϕ and d/dz on $\widehat{\mathcal{R}}^{\text{et}}$ commute with the action of $\pi_1^{\text{alg}}(S_0, s_0)$, and we have a canonical isomorphism

$$\mathcal{U} \otimes_{\mathcal{R}} \widehat{\mathcal{R}}^{\text{et}} \simeq V \otimes_{\mathbb{Z}_p} \widehat{\mathcal{R}}^{\text{et}}$$

compatible with $\phi, d/dz$ and $\pi_1^{\text{alg}}(S_0, s_0)$ (diagonal action of $\phi, d/dz$ on the left hand side, diagonal action of $\pi_1^{\text{alg}}(S_0, s_0)$ on the right hand side), which allows to reconstruct the representation V from the unit-root F -crystal and conversely. In fact, the connection as well as V can be reconstructed from (\mathcal{U}, Φ) alone.

This proposition may be compared with 1.5.4, though it applies to a quite different type of p -adic connections. In 1.5.4 (for dimension 1), the analytic curve S had typically bad reduction and the main player was the *discrete fundamental group* $\pi_1(\Delta)$ of the dual graph of the semistable reduction, together with the $\pi_1(\Delta)$ -module $\mathcal{O}(\widetilde{S})$. Here the curve S has good reduction S_0 and the main player is the *compact fundamental group* $\pi_1^{\text{alg}}(S_0)$, together with the $\pi_1^{\text{alg}}(S_0)$ -module $\widehat{\mathcal{R}}^{\text{et}}$.

Before presenting one of Dwork's classical unit-root F -crystals, let us just mention that the above theory extends with little change to the case when the base ring \mathfrak{v} is a finite ramified extension of $\widehat{\mathbb{Z}}_p^{\text{ur}}$: one has to fix an extension of σ to \mathfrak{v} , to replace \mathbb{Z}_p by \mathfrak{v}^{σ} , and to impose some mild nilpotence constraint on ∇ if the ramification index is $\geq p-1$ (also, it is customary to extend the definition of Frobenius structure on replacing ϕ by some power).

Example 2.4.5. Dwork's exponential. We denote by π a fixed $(p-1)$ th root of $-p$, and set $\mathfrak{v} = \widehat{\mathbb{Z}}_p^{\text{ur}}[\pi]$, with $\sigma(\pi) = \pi$. Let us consider the differential equation

$$(*) \quad f'(z) = -\pi f(z)$$

over $S = \text{D}(0, 1^+)$, which has the analytic solution $f_a = e^{-\pi(z-a)}$ in any residue class $\text{D}(a, 1^-) \subset \text{D}(0, 1^+)$. The change of variable $z \mapsto z^p$ leads to the differential equation

$$(*_p) \quad g'(z) = -p\pi z^{p-1}g(z) = \pi^p z^{p-1}g(z)$$

Dwork's exponential function is

$$E_{\pi}(z) = e^{\pi(z-z^p)} \in \mathbb{Z}_p[\pi][[z]].$$

This is an invertible element of \mathcal{R} , the π -adic completion of $\mathfrak{v}[z]$. This function provides the unit-root Frobenius structure which relates $(*)$ and $(*_p)$.

Let us reformulate this in the setting of 2.4.3: the relevant F -crystal is (\mathcal{R}, ∇, F) , with $\nabla(1) = \pi dz$, $F(\phi)(1) = E_\pi(z)$ for the standard ϕ . This is the unit-root F -crystal attached to the finite character of $\pi_1^{\text{alg}}(\mathbb{A}_{\mathbb{F}_p}^1, 0)$ corresponding to the Artin-Schreier covering $z = y^p - y$. The solution f_0 of $(*)$ belongs to $\widehat{\mathcal{R}^{\text{et}}}$: indeed, $e^{-\pi z} = E_\pi(y) \in \widehat{\mathfrak{v}[y]} \subset \widehat{\mathcal{R}^{\text{et}}}$; explicitly, $e^{-\pi z} = \lim_n (1 - \pi p^n z)^{p^{-n}}$.

Dwork has computed the value of his exponential for any $(p-1)$ th root of unity ζ_{p-1} (e.g. $\zeta_{p-1} = 1$): this is

$$E_\pi(\zeta_{p-1}) = \zeta_p^{\zeta_{p-1}}$$

where ζ_p is the unique p th root of unity $\equiv 1 + \pi \pmod{\pi}$ (cf. [La90, chap. 14]). In the spirit of 2.3.3, this may be understood as follows:

$\zeta_p^{\zeta_{p-1}}$ = the value at $(z = 0)$ of the analytic continuation of f_0 along the “wild loop” corresponding to the path from $(y = 0)$ to $(y = \zeta_{p-1})$ on the Artin-Schreier covering $z = y^p - y$ in characteristic p (figure 5). If one changes ζ_{p-1} , this value $\zeta_p^{\zeta_{p-1}}$ is multiplied by some p th root of unity.




FIGURE 5. a wild underground loop

In our special case, the discussion of 2.3.3 tells us that $e^{-\pi z}$ admits an extension to any disk $D(a, 1^-) \subset D(0, 1^+)$, analytic in that disk, and well-defined up to multiplication by some p th root of unity. It is easy to find a formula for this extension: it must be proportional to $e^{\pi(a-z)}$, and by evaluation at a , we find that it is $e^{\pi(a-z)} E_\pi(b)$, where b is any solution of the equation $b^p - b = a$. Its p th power is, as expected, $e^{-p\pi z}$ itself.

Therefore, this multivalued exponential $e^{-\pi z}$ on $D(0, 1^+)$ appears as a multivalued section of the logarithmic étale covering $\frac{-1}{\pi} \log : D(1, p^{-\frac{1}{p-1}+}) \rightarrow D(0, 1^+)$. This is just the opposite way from the complex situation, where the logarithm is a multivalued section of the étale covering of \mathbb{C}^\times given by the exponential.

2.5. p -adic chiaroscuro: overconvergence.

2.5.1. In fact, Dwork's exponential E_π is more than just an element of the π -adic completion of $\mathfrak{v}[z]$: it is *overconvergent*, i.e. extends to an analytic function on a disk of radius $\rho > 1$ ($\rho = p^{(p-1)/p^2}$).

Overconvergence is a fundamental notion in p -adic analysis: dimming the contours of an affinoid turns out to be the key to the finiteness properties of p -adic cohomology, as was recognized by Dwork, and subsequently developed by P. Monsky, G. Washnitzer, P. Berthelot (rigid cohomology, \mathcal{D}^\dagger -modules)...




FIGURE 6

2.5.2. Let again S be $D(0, 1^+) \setminus \bigcup D(\zeta_i, 1^-)$. For any $\rho > 1$, let us consider the bigger affinoids $S_\rho = D(0, \rho^+) \setminus \bigcup D(\zeta_i, \frac{1}{\rho}^-)$. The ring of *overconvergent analytic functions* on S is

$$\mathcal{H}^\dagger(S) = \bigcup_{\rho > 1} \mathcal{O}(S_\rho).$$

Its relevance to the algebraic viewpoint on analytic continuation comes from the following result [**BDR80**]:

Proposition 2.5.3. *Let $f \in \mathcal{O}(S)$ satisfy a monic polynomial equation with coefficients in $\mathcal{H}^\dagger(S)$. Then $f \in \mathcal{H}^\dagger(S)$.*

2.5.4. A F -crystal (\mathcal{M}, ∇, F) is *overconvergent* if (\mathcal{M}, ∇) as well as $F(\phi)$ extends over some S_ρ . This is the case in example 2.4.5. For unit-root F -crystals, Crew has given the following characterization:

Proposition 2.5.5. *The unit-root F -crystal $(\mathcal{U}_V, \nabla_V, F_V)$ attached to a p -adic representation V is overconvergent if and only if the images in $GL(V)$ of the inertia groups at the missing points $\bar{\zeta}_1, \dots, \bar{\zeta}_\nu, \infty$ are finite.*

2.6. (Overconvergence and Frobenius) Dwork's derivation of the p -adic Gamma function and exponential sums.

2.6.1. We come back to the situation of 2.4.5. For any $\alpha \in \mathbb{Z}_p$, let us consider $M_\alpha^\dagger := z^\alpha e^{\pi z} \mathcal{H}^\dagger(S)$ endowed with the derivation $z \frac{d}{dz}$. A simple computation shows that

$$z^\alpha e^{\pi z} z^{k+1} \equiv -\frac{\alpha + k}{\pi} z^\alpha e^{\pi z} z^k$$

in $M_\alpha^\dagger/z\frac{d}{dz}M_\alpha^\dagger$, from which one deduces that this cokernel has dimension 1 over K and is generated by $[z^\alpha e^{\pi z}]$.

Following Dwork, let us introduce the operator ψ defined by

$$\psi\left(\sum a_n z^n\right) = \sum a_{pn} z^n.$$

This is a left inverse of the Frobenius operator induced by $\phi : z \mapsto z^p$. It acts on $\mathcal{H}^\dagger(S)$ and commutes with $z\frac{d}{dz}$ up to multiplication by p : $\psi \circ z\frac{d}{dz} = pz\frac{d}{dz} \circ \psi$. The operator ψ can also be applied to any element of M_α^\dagger for $\alpha \in \mathbb{N}$: one finds that

$$\psi(z^\alpha e^{\pi z} f) = z^\beta e^{\pi z} \psi(z^{\alpha-p\beta} E_\pi(z) f)$$

where β is the *successor* of $\alpha \in \mathbb{Z}_p$, *i.e.* the unique p -adic integer β such that $p\beta - \alpha \in \mathbb{Z} \cap [0, p[$ (note that since $\alpha - p\beta > -p$, the terms containing a negative power of z disappear when applying ψ). This formula makes sense for any $\alpha \in \mathbb{Z}_p$, and we can see that ψ applies M_α^\dagger into M_β^\dagger , and commutes with $z\frac{d}{dz}$. Let us write the induced map of one-dimensional cokernels $[\psi] : M_\alpha^\dagger/z\frac{d}{dz}M_\alpha^\dagger \rightarrow M_\beta^\dagger/z\frac{d}{dz}M_\beta^\dagger$ in the form

$$[\psi]([z^\alpha e^{\pi z}]) = \pi^{p\beta-\alpha} \Gamma_p(\alpha) [z^\beta e^{\pi z}].$$

2.6.2. This function Γ_p is then nothing but Morita's p -adic Gamma function, characterized by its continuity and the functional equation

$$\Gamma_p(0) = 1, \quad \Gamma_p(\alpha + 1)/\Gamma_p(\alpha) = \begin{cases} -\alpha & \text{if } \alpha \text{ is a unit,} \\ -1 & \text{if } |\alpha|_p < 1. \end{cases}$$

Let us check this functional equation:

- for $\alpha = 0$, one has $[z^k e^{\pi z}] = 0$ if $k > 0$; hence $\Gamma_p(0)[e^{\pi z}] = [e^{\pi z}][\psi(E_\pi(z))]$ $= [e^{\pi z}]$;
- if $|\alpha|_p = 1$, β is also the successor of $\alpha + 1$, and $\Gamma_p(\alpha + 1)[z^\beta e^{\pi z}] = \pi^{\alpha-p\beta+1}[\psi]([z^{\alpha+1} e^{\pi z}]) = \pi^{\alpha-p\beta+1}[\psi](-\frac{\alpha}{\pi}[z^\alpha e^{\pi z}]) = -\alpha\Gamma_p(\alpha)[z^\beta e^{\pi z}]$;
- finally, if $|\alpha|_p < 1$, one has $\alpha = p\beta$, and $\beta + 1$ is the successor of $\alpha + 1$; one gets $\Gamma_p(\alpha+1)[z^{\beta+1} e^{\pi z}] = \pi^{-p+1}[\psi]([z^{\alpha+1} e^{\pi z}]) = -\frac{1}{p}[\psi](-\frac{\alpha}{\pi}[z^\alpha e^{\pi z}]) = \frac{\beta}{\pi}\Gamma_p(\alpha)[z^\beta e^{\pi z}] = -\Gamma_p(\alpha)[z^{\beta+1} e^{\pi z}]$.

2.6.3. Let us now check the continuity of Γ_p on \mathbb{Z}_p , or better, its analyticity on any disk $D(-k, |p|^+)$, $k = 0, 1, \dots, p-1$. Let us write $E_\pi(z) = e^{\pi(z-z^p)} = \sum_0^\infty e_n z^n$.

For any $\alpha \in \mathbb{Z}_p \cap D(-k, 1^-)$, one has

$$\begin{aligned} \psi(z^\alpha e^{\pi z}) &\equiv \pi^k \Gamma_p(\alpha) z^\beta e^{\pi z} = z^\beta e^{\pi z} \psi(z^{-k} E_\pi(z)) \\ &= z^\beta e^{\pi z} \sum_{n=0}^\infty e_{pn+k} z^n \equiv z^\beta e^{\pi z} \sum_{n=0}^\infty e_{pn+k} (-\pi)^{-n} (\beta)_n, \end{aligned}$$

where $(\beta)_n = \left(\frac{\alpha+k}{p}\right)_n$ is the Pochhammer symbol. Easy estimates now show that $\Gamma_p(\alpha) = \sum_0^\infty e_{pn+k} (-\pi)^{-n-k} \left(\frac{\alpha+k}{p}\right)_n$ is analytic on $D(-k, |p|^+)$.

2.6.4. Gross-Koblitz formula: for $k = 0, 1, \dots, p-2$, one has

$$\Gamma_p\left(\frac{k}{p-1}\right) = -\pi^{-k} \sum_{\zeta_{p-1}} \zeta_{p-1}^{-k} \zeta_p^{\zeta_{p-1}} \in \mathbb{Q}_p[\pi] \cap \overline{\mathbb{Q}}$$

where ζ_{p-1} runs over the $(p-1)$ th roots of unity, and $\zeta_p^{\zeta_{p-1}}$ is as before the unique p th root of unity $\equiv 1 + \zeta_{p-1}\pi \pmod{\pi}$.

Let us sketch the proof. We choose $\alpha = \frac{k}{p-1}$, so that $\alpha = \beta$ and ψ acts on the ind-Banach-space M_α^\dagger via the formula: $\psi(z^{-k}E_\pi(z)f) = (z^\alpha e^{\pi z})^{-1}\psi(z^\alpha e^{\pi z}f)$. Coming back to the definition of ψ , one observes that for any $g \in \mathcal{H}^\dagger(S)$,

$$\psi(z^{-k}g)(z) = \frac{1}{p} \sum_{t \in \phi^{-1}(z)} t^{-k}g(t).$$

Using the fact that the domain of analyticity of this function is bigger than the domain of analyticity of g itself, one shows that ψ is a nuclear operator of M_α^\dagger . In particular, it has a trace, which is the trace of the “composition operator” $\Psi : g \in \mathcal{H}^\dagger(S) \mapsto \frac{1}{p} \sum_{t \in \phi^{-1}(z)} t^{-k}E_\pi(t)g(t)$. The computation of this trace is done by approximating E_π by polynomials and studying the resulting action on the subspace of polynomials. One finds $\text{Tr } \Psi = \frac{1}{p-1} \sum_{\zeta_{p-1}} \zeta_{p-1}^{-k} E_\pi(\zeta_{p-1})$.

At last, because $\psi \circ z \frac{d}{dz} = pz \frac{d}{dz} \circ \psi$, one has

$$\begin{aligned} \pi^k \Gamma_p\left(\frac{k}{p-1}\right) &= \text{Tr}\left([\psi] \middle| M_\alpha^\dagger / z \frac{d}{dz} M_\alpha^\dagger\right) = (1-p) \text{Tr}(\psi \middle| M_\alpha^\dagger) \\ &= - \sum_{\zeta_{p-1}} \zeta_{p-1}^{-k} E_\pi(\zeta_{p-1}) = - \sum_{\zeta_{p-1}} \zeta_{p-1}^{-k} \zeta_p^{\zeta_{p-1}}. \end{aligned}$$

We refer to [CR94] for a detailed account. A more general form of the Gross-Koblitz, proved along the same lines, shows that for any $1 \leq k < p^r$, the product

$$\prod_0^{r-1} \Gamma_p\left(\frac{p^i k}{p^r - 1}\right)$$

belongs to the cyclotomic field $\mathbb{Q}(\zeta_{p^r})$.

2.6.5. A more straightforward and elementary proof has been discovered by A. Robert. It goes as follows.

As we have just seen, the right hand side of the Gross-Koblitz formula can be written

$$\begin{aligned} & -\pi^{-k} \sum_{\zeta_{p-1}} \zeta_{p-1}^{-k} E_\pi(\zeta_{p-1}) = -\pi^{-k} \sum_{n=0}^{\infty} \left(\sum_{\zeta_{p-1}} \zeta_{p-1}^{n-k} \right) e_n \\ & = (1-p)\pi^{-k} \sum_{m=0}^{\infty} e_{(p-1)m+k}. \end{aligned}$$

Using the expansion of $\Gamma(\alpha)$ given in 2.6.3, in the case of $\alpha = \frac{k}{p-1}$, we thus have to show that

$$\sum_{n=0}^{\infty} e_{pn+k} (-\pi)^{-n} \left(\frac{k}{p-1} \right)_n = (1-p) \sum_{m=0}^{\infty} e_{(p-1)m+k}.$$

Denote the left hand side by G_k (for any $k \in \mathbb{N}$). Due to the overconvergence of E_π , it is not difficult to see that $\lim_{k \rightarrow \infty} G_k = 0$. One has Robert's formula:

$$G_k - G_{p-1+k} = (1-p)e_k.$$

Summing up consecutive expressions, one gets a telescoping sum which yields

$$\text{the desired equality } G_k = G_k - \lim_{m \rightarrow \infty} G_{(p-1)m+k} = (1-p) \sum_{m=0}^{\infty} e_{(p-1)m+k}.$$

It remains to prove Robert's formula. One first observes that $z \frac{d}{dz} E_\pi = (\pi z - p\pi z^p) E_\pi(z)$, which yields the relation $ne_n = \pi(e_{n-1} - pe_{n-p})$ for $n \geq p$, hence $\pi e_{p-1+m} = p\pi e_m + (m+p)e_{m+p}$ for $m \geq 0$. Then

$$\begin{aligned} & G_k - G_{p-1+k} \\ & = e_k + \sum_0^{\infty} e_{p(n+1)+k} (-\pi)^{-n-1} \left(\frac{k}{p-1} \right)_{n+1} \\ & \quad - \sum_0^{\infty} e_{p-1+pn+k} (-\pi)^{-n} \left(\frac{k}{p-1} + 1 \right)_n \\ & = e_k + \sum_0^{\infty} \left[\frac{k}{p-1} e_{p(n+1)+k} + \pi e_{p-1+pn+k} \right] (-\pi)^{-n-1} \left(\frac{k}{p-1} + 1 \right)_n \\ & = e_k + \sum_0^{\infty} \left[p \frac{(k+(n+1)(p-1))}{p-1} e_{p(n+1)+k} + p\pi e_{pn+k} \right] (-\pi)^{-n-1} \left(\frac{k}{p-1} + 1 \right)_n \\ & = (1-p)e_k + p \sum_0^{\infty} \frac{(k+(n+1)(p-1))}{p-1} e_{p(n+1)+k} (-\pi)^{-n-1} \left(\frac{k}{p-1} + 1 \right)_n \\ & \quad - p \sum_1^{\infty} \frac{k+n(p-1)}{p-1} e_{pn+k} (-\pi)^{-n} \left(\frac{k}{p-1} + 1 \right)_{n-1} \\ & = (1-p)e_k. \end{aligned}$$

2.6.6. The right hand side of the Gross-Koblitz formula is a special case of an *exponential sum*, *i.e.* an expression of the form

$$S_r(\bar{f}, \bar{g}, \bar{h}) = \sum_{x_1, \dots, x_d \in \mathbb{F}_{p^r}, \bar{h}(\underline{x}) \neq 0} \chi(N_{\mathbb{F}_{p^r}/\mathbb{F}_p} \bar{g}(\underline{x})) \exp\left(\frac{2\pi i}{p} \text{Tr}_{\mathbb{F}_{p^r}/\mathbb{F}_p} \bar{f}(\underline{x})\right)$$

where \bar{h} is a polynomial with coefficients in \mathbb{F}_{p^r} , \bar{f} and \bar{g} are rational functions with coefficients in \mathbb{F}_{p^r} with no pole where \bar{h} vanishes, and where χ is a character of \mathbb{F}_p^\times . This includes Gauss, Jacobi and Kloosterman sums as special cases. In fact, it is known, after the classical works of Gauss, Artin, Weil, that counting solutions of systems of polynomial equations in finite fields amounts to the computation of exponential sums $S_r(\bar{f}, \bar{g}, \bar{h})$. They are studied via their generating series, the so-called *L-series*:

$$L(\bar{f}, \bar{g}, \bar{h}; t) = \exp\left(\sum_{r \geq 1} S_r(\bar{f}, \bar{g}, \bar{h}) \frac{t^r}{r}\right).$$

Dwork's methods, refined by P. Robba and others, allow to tackle the questions of the rationality of L , its degree, and of the functional equation relating $L(\bar{f}, \bar{g}, \bar{h}; t)$ to $L(\bar{f}, \bar{g}, \bar{h}; \frac{1}{p^r t})$. In the one-dimensional case, the solution is elementary and follows the pattern sketched in 2.6.4. Namely [CR94]:

- (i) One considers liftings f, g, h of $\bar{f}, \bar{g}, \bar{h}$ in characteristic zero, and one introduces the affinoid set S defined by $|h| = 1$. One sets $F = g(z)^{1/p-1} \exp(\pi f(z))$. Then F'/F is a rational function, and one shows that the differential operator $\frac{d}{dz} + \frac{F'}{F}$ has an index in $\mathcal{H}^\dagger(S)$ (in a slightly generalized sense, and which can be computed thanks to the work of Robba). This is the crucial point. Hence the cohomology spaces of the de Rham complex: $\Omega^0 = F\mathcal{H}^\dagger(S) \rightarrow \Omega^1 = F\mathcal{H}^\dagger(S)dz$ are finite-dimensional over \mathbb{C}_p .
- (ii) One observes that $E := F^{\phi - \text{id}}$ (*i.e.* $E(z) = F(z^p)/F(z)$) is an element of $\mathcal{H}^\dagger(S)$. This allows to define two endomorphisms of the de Rham complex by setting:

$$\phi^0(F.g) = FEg^\phi, \quad \phi^1(F.gdz) = FE\phi'g^\phi dz \quad (\text{with } \phi'(z) = pz^{p-1});$$

$$\psi^0(F.g)(z) = F(z) \sum_{\phi(t)=z} \frac{g(t)}{E(t)}, \quad \psi^1(F.gdz) = \sum_{\phi(t)=z} \frac{g(t)}{E(t)\phi'(t)}.$$

One has $\psi^* \circ \phi^* = p \cdot \text{id}$. In particular, the operators $H^i(\psi^*)$, $i = 0, 1$, are invertible on the cohomology spaces. The same argument as in 2.6.4 shows that ψ^i is nuclear on Ω^i . In particular, it has a trace.

- (iii) Polynomial approximation allows to prove the trace formula:

$$\text{tr}(\psi^1) - \text{tr}(\psi^0) = \text{tr}(H^1(\psi^*)) - \text{tr}(H^0(\psi^*)) = S_r(\bar{f}, \bar{g}, \bar{h}).$$

It follows that

$$L(\bar{f}, \bar{g}, \bar{h}; t) = \frac{\det(\text{id} - tH^1(\psi^*))}{\det(\text{id} - tH^0(\psi^*))}.$$

This is a rational function since the cohomology is finite-dimensional. Moreover, since the $H^i(\psi^*)$ are invertible, the computation of its degree amounts to the computation of the index of $\frac{d}{dz} + \frac{F'}{F}$ (the computation of the dimension of H^0 , 0 or 1, being essentially trivial).

- (iv) The functional equation follows from a topological “dual theory” in which the transpose of ϕ and ψ play the roles of ψ and ϕ respectively.

Let us remark at last that this method has an *archimedean analogue* in the so-called thermodynamic formalism —in dimension one. The analogy is especially striking in the presentation of D. Mayer [May91]: exponential sums correspond to “partition functions”, L to the Ruelle zeta-function, Frobenius to the “shift”, Dwork’s operator ϕ^* to the “transfer operator” (or Ising-Perron-Frobenius-Ruelle operator) and is given by a “composition operator” (loc. cit. 7.2.2), its nuclearity is established by the same argument, and the trace formula has the same form (loc. cit. 7.17).

In view of this close analogy, one could dream of an archimedean proof of the rationality of Weil zeta functions, parallel to Dwork’s p -adic proof...

3. The tale of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$.

ABSTRACT: This is a detailed concrete illustration of the somewhat abstract non-archimedean notions discussed in the previous section. We refer to [Yo97] (and [Hu87, 9]) for an excellent account of the archimedean tale of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$.

3.1. Dwork's hypergeometric function.

Let us consider the Legendre pencil of elliptic curves with parameter $z \neq 0, 1, \infty$, given in inhomogeneous coordinates by

$$y^2 = x(x-1)(x-z).$$

As a scheme over $\mathbb{Z}[z, \frac{1}{2z(1-z)}]$, we denote it by X . The first de Rham cohomology module $H_{\text{dR}}^1(X)$ is free of rank 2 over $\mathbb{Z}[z, \frac{1}{2z(1-z)}]$, and endowed with the Gauss-Manin connection ∇ (derivation with respect to the parameter z); it is generated by the class ω of $\frac{dx}{y}$, and $\nabla(\frac{d}{dz})(\omega)$. The canonical symplectic form (cup-product) satisfies $\langle \omega, \nabla(\frac{d}{dz})(\omega) \rangle = \frac{2}{z(z-1)}$. The Gauss-Manin connection is given by the hypergeometric differential equation with parameters $(\frac{1}{2}, \frac{1}{2}, 1)$

$$\nabla(L_{\frac{1}{2}, \frac{1}{2}, 1})(\omega) = 0, \text{ with } L_{\frac{1}{2}, \frac{1}{2}, 1} = z(1-z)\frac{d^2}{dz^2} + (1-2z)\frac{d}{dz} - \frac{1}{4}.$$

Let p be an odd prime. The *Hasse invariant* is the polynomial $h_p(z) \in \mathbb{Z}[\frac{1}{2}][z]$ obtained by truncating the hypergeometric series $(-1)^{\frac{p-1}{2}} F(\frac{1}{2}, \frac{1}{2}, 1; z)$ at order $\frac{p-1}{2}$. It enjoys the following well-known properties:

- functional equations: $h_p(z) \equiv (-1)^{\frac{p-1}{2}} h_p(1-z) \equiv z^{\frac{p-1}{2}} h_p(\frac{1}{z}) \pmod{p}$,
- for any $\bar{\zeta} \in \overline{\mathbb{F}}_p \setminus \{0, 1\}$, one has $h_p(\bar{\zeta}) = 0$ if and only if the elliptic curve $X_{\bar{\zeta}}$ over $\overline{\mathbb{F}}_p$ is *supersingular*, i.e. has no geometric point of order p , cf. [Hu87, 13.3].

The roots of $h_p \pmod{p}$ are distinct and lie in \mathbb{F}_{p^2} : we denote them by $\bar{\zeta}_1, \dots, \bar{\zeta}_{(p-1)/2}$, and choose liftings $\zeta_1, \dots, \zeta_{(p-1)/2}$ in $\widehat{\mathbb{Z}}_p^{\text{ur}}$.

Non-supersingular elliptic curves are called *ordinary*; they have exactly p points of order p .

Dwork's hypergeometric function is

$$f_p(z) = (-1)^{\frac{p-1}{2}} \frac{F(\frac{1}{2}, \frac{1}{2}, 1; z)}{F(\frac{1}{2}, \frac{1}{2}, 1; z^p)} \in \mathbb{Z}\left[\frac{1}{2}\right][[z]].$$

One has $f_p(z) \equiv h_p(z) \pmod{p}$. Dwork discovered that although the p -adic radius of convergence of this series is exactly 1, f_p does extend to a p -adic analytic function on \mathbb{A}^1 deprived from the supersingular disks $D(\zeta_j, 1^-)$; we denote this extension by the same symbol f_p .

In terms of this function, he obtained in 1958 his famous p -adic formula for the number of rational points of an ordinary elliptic curve defined over \mathbb{F}_{p^n} :

for any $s_0 \in \mathbb{F}_{p^n}$, $s_0 \neq 0, 1, \bar{\zeta}_j$, the number of \mathbb{F}_{p^n} -points of X_{s_0} is

$$1 - \prod_{k=0}^{n-1} f_p(\omega^{p^k}) + p^n \left(1 - \prod_{k=0}^{n-1} f_p(\omega^{p^k})^{-1} \right),$$

where ω denotes the unique $(p^n - 1)$ th root of unity $\equiv s_0 \pmod{p}$.

3.2. The ordinary unit-root F -crystal.

On completing $\widehat{\mathbb{Z}_p^{\text{ur}}}[z, \frac{1}{2z(1-z)}]$ p -adically, $(H_{\text{dR}}^1(X), \nabla)$ gives rise to an overconvergent F -crystal (\mathcal{H}, ∇, F) . We do not discuss here the construction of the Frobenius structure, which is a general p -adic feature of Gauss-Manin connections. In fact, various analytic or geometric constructions are available, but in our present case, the Frobenius structure can be made quite explicit, cf. [Dw69].

Let us introduce a few notations:

- \mathcal{R} = the p -adic completion of $\widehat{\mathbb{Z}_p^{\text{ur}}}[z, \frac{1}{h_p(z)}]$
- \mathcal{R}_{ord} = the p -adic completion of $\widehat{\mathbb{Z}_p^{\text{ur}}}[z, \frac{1}{z(1-z)h_p(z)}]$
- $S = M(\mathcal{R}[\frac{1}{p}]) = D(0, 1^+) \setminus (\bigcup D(\zeta_j, 1^-))$
- $S_{\text{ord}} = M(\mathcal{R}_{\text{ord}}[\frac{1}{p}]) = S \setminus (D(0, 1^-) \cup D(1, 1^-))$ is the *ordinary locus*
- $S_{\text{nss}} = \mathbb{A}^1 \setminus (\bigcup D(\zeta_j, 1^-))$ is the *non-supersingular locus*.

The restriction of (\mathcal{H}, ∇, F) to the ordinary locus possesses a unique non-zero unit-root sub- F -crystal (\mathcal{U}, ∇, F) . This unit-root F -crystal extends over S (and even over S_{nss} as an F -isocrystal with logarithmic singularity at ∞ in the sense of [Scho85]) cf. also [O00]; we use the same symbol for the extension. It can be described along the following lines:

- \mathcal{U} is the unique rank-one horizontal submodule of \mathcal{H} .
- $\mathcal{U} \otimes_{\mathcal{R}_{\text{ord}}} \widehat{\mathcal{R}_{\text{ord}}^{\text{et}}} = (\mathcal{H} \otimes_{\mathcal{R}_{\text{ord}}} \widehat{\mathcal{R}_{\text{ord}}^{\text{et}}})^{\nabla} \otimes_{\widehat{\mathbb{Z}_p^{\text{ur}}}} \widehat{\mathcal{R}_{\text{ord}}^{\text{et}}}$. (Here $\widehat{\mathcal{R}_{\text{ord}}^{\text{et}}}$ is defined as in 2.3.2.)
- for any ordinary $s_0 \in \overline{\mathbb{F}_p}$, the associated p -adic representation of $\pi_1^{\text{alg}}(S_0, s_0)$ is $(\mathcal{H} \otimes_{\mathcal{R}_{\text{ord}}} \widehat{\mathcal{R}_{\text{ord}}^{\text{et}}})^{\nabla=0, F(\phi)=\text{id}} \simeq H_{\text{et}}^1(X_{s_0}, \mathbb{Z}_p)$.
- For any $u \in (\mathcal{U} \otimes_{\mathcal{R}} \widehat{\mathcal{R}^{\text{et}}})^{\nabla}$, “the” image of u in $\mathcal{O}(D(s_0, 1^-))$ is a bounded solution of the differential operator $L_{\frac{1}{2}, \frac{1}{2}, 1}$ on $D_{\mathbb{C}_p}(s_0, 1^-)$; this also characterizes \mathcal{U} .
- $\mathcal{U}|_{D(0, 1^-)}^{\nabla}$ has a canonical \mathbb{Z} -submodule, which can be identified with $H^1((X_z)^{\text{an}}, \mathbb{Z})$ for any $z \in D(0, 1^-) \setminus \{0\}$; for a generator u , one has

$$\langle \omega, u \rangle = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \text{ in } \mathcal{O}(D(0, 1^-))$$

($\sqrt{-1}$ appears as residue of $\frac{dx}{y}|_{z=0}$ at $x=0$), cf. [And90]. Similarly, $\mathcal{U}|_{\mathbb{D}(1,1^-)}^\nabla$ has a canonical \mathbb{Z} -submodule; for a generator u , one has

$$\langle \omega, u \rangle = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; 1-z\right) \text{ in } \mathcal{O}(\mathbb{D}(1,1^-)).$$

The existence of the unit-root F -crystal (\mathcal{U}, ∇, F) over \mathcal{R} then amounts to the two function-theoretic facts:

$$\text{dlog } F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \text{ and } f_p(z) = \left(\sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)\right)^{1-\sigma}$$

both extend to units in \mathcal{R} .

The inertia at any supersingular point maps *onto* $\text{Aut } H_{\text{et}}^1(X_{s_0}, \mathbb{Z}_p) \simeq \mathbb{Z}_p^\times$ (J.-I. Igusa, cf. e.g. [vdP87]). The inertia at ∞ acts as ± 1 . According to Crew's criterion, the F -crystal (\mathcal{U}, ∇, F) is *not* overconvergent.

We refer to [SPM91] for a study in the same spirit of more general hypergeometric equations and other differential equations “coming from geometry”.

3.3. Analytic continuation: a précis.

3.3.1. We have seen that Dwork's hypergeometric function extends to an analytic function on the whole of S_{nss} . Some of its special values have been computed. Let us mention [Ko79], [You92]

- (i) $f_p(1) = 1$ (Koblitz),
- (ii) if $p \equiv 1 \pmod{4}$,

$$f_p(-1) = (-1)^{\frac{p-1}{4}} \frac{\Gamma_p(1/4)^2}{\Gamma_p(1/2)} = \frac{\Gamma_p(1/4)}{\Gamma_p(1/2)\Gamma_p(3/4)} \quad (\text{Young}).$$

(the condition $p \equiv 1 \pmod{4}$ ensures that -1 is an ordinary modulus); $f_p(-1)$ is a Gauss integer (Van Hamme).

3.3.2. We have seen that the logarithmic derivative $\text{dlog } F(\frac{1}{2}, \frac{1}{2}, 1; z)$ also extends to an analytic function on S_{nss} . It satisfies the functional equations

$$\begin{aligned} \text{dlog } F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) &= -\text{dlog } F\left(\frac{1}{2}, \frac{1}{2}, 1; 1-z\right) \\ &= -\frac{1}{2z} - \frac{1}{z^2} \text{dlog } F\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{1}{z}\right). \end{aligned}$$

3.3.3. Let us turn to the more subtle case of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ itself. The general discussion of 2.3.3/2.4.3 applies to the ordinary unit-root F -crystal and tells us that $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ admits an extension to any unit disk $\mathbb{D}(s, 1^-) \subset S$, analytic in that disk, and well-defined up to multiplication by a unit in \mathbb{Z}_p . In other words, $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ is the specialization of an element (in fact, a unit) of $\widehat{\mathcal{R}}_{\text{nss}}^{\text{et}}$. It may be suggestive to denote such an element by $F(\frac{1}{2}, \frac{1}{2}, 1; \eta)$, where η stands for the Berkovich generic point of S corresponding to the sup-norm).

In $D(1, 1^-)$, this extension is $F(\frac{1}{2}, \frac{1}{2}, 1; 1-z)$ (up to \mathbb{Z}_p^\times). The refined structure of F -isocrystal with logarithmic singularity at ∞ allows to construct an extension of $F(\frac{1}{2}, \frac{1}{2}, 1; z)^2$ to $D(\infty, 1^-)$, analytic in that disk: it is $\frac{1}{z}F(\frac{1}{2}, \frac{1}{2}, 1; \frac{1}{z})^2$ (up to \mathbb{Z}_p^\times); notice the square, which comes from the exponent $\pm 1/2$ at ∞ .

3.3.4. The Krasner representation of $f_p(z)$ and $\mathrm{dlog} F(\frac{1}{2}, \frac{1}{2}, 1; z)$ as uniform limits of rational functions without pole (nor zero) on S can be made explicit: for $n > 0$, let g_n be the polynomial obtained by truncating $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ at order $p^n - 1$; then $f_p(z) = \lim(-1)^{\frac{p-1}{2}} \frac{g_{n+1}(z)}{g_n(z^p)}$ and $\mathrm{dlog} F(\frac{1}{2}, \frac{1}{2}, 1; z) = \lim \mathrm{dlog} g_n(z)$ [Dw69, 3.4]. On the other hand, $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ cannot be approximated by rational functions on S , but $L_{\frac{1}{2}, \frac{1}{2}, 1}(g_n)$ tends uniformly to 0 on $D(0, 1^+)$ [Ro75]. One has

$$\sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \equiv (h_p(z))^{-\frac{1}{p-1}} \pmod{p}$$

but we do not know any explicit representation of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ as a p -adic limit of algebraic functions on S .

3.3.5. Any solution of $L_{\frac{1}{2}, \frac{1}{2}, 1}$ on an ordinary disk $D(s, 1^-)$, which is not proportional to (the extended) $F(\frac{1}{2}, \frac{1}{2}, 1; z)$, is not bounded in $D_{\mathbb{C}_p}(s, 1^-)$, hence does not extend to neighboring disks in any reasonable *analytic* sense. In spite of this obstruction to continuation, one can nevertheless “jump from disk to disk” (see figure 7) and extend them as locally analytic functions, in




FIGURE 7

the following way.

For each $\bar{\zeta} \in S_0 \setminus \{0, 1\}$, there is a *canonical lifting* $\zeta_{\mathrm{can}} \in \mathbb{Z}_p^{\mathrm{ur}}$: namely, the modulus ζ_{can} for which $X_{\zeta_{\mathrm{can}}}$ has complex multiplication by the quadratic order $\mathrm{End}(X_{\bar{\zeta}})$. Let us fix a branch of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ in $D(\zeta_{\mathrm{can}}, 1^-)$, *i.e.* an analytic solution $F_{\bar{\zeta}} \in \widehat{\mathbb{Z}_p^{\mathrm{ur}}}[[z - \zeta_{\mathrm{can}}]]$ of $L_{\frac{1}{2}, \frac{1}{2}, 1}$ in $D(\zeta_{\mathrm{can}}, 1^-)$ which is a specialization of $F(\frac{1}{2}, \frac{1}{2}, 1; \eta) \in (\widehat{\mathcal{R}^{\mathrm{et}}})^\times$ (any other branch is of the form $c \cdot F_{\bar{\zeta}}$ with $c \in \mathbb{Z}_p^\times$). Let $u_{\bar{\zeta}} \in \mathcal{U}_{D(\zeta_{\mathrm{can}}, 1^-)}^\nabla$ be defined by

$$\langle \omega, u_{\bar{\zeta}} \rangle = \sqrt{-1}F_{\bar{\zeta}} \text{ in } \mathcal{O}(\mathbb{D}(\zeta_{\text{can}}, 1^-)).$$

On the other hand, let $v_{\bar{\zeta}} \in \mathcal{M}|_{\mathbb{D}(\zeta_{\text{can}}, 1^-)}^{\nabla}$ be defined by

$$\sqrt{-1}F_{\bar{\zeta}}(\zeta_{\text{can}}).v_{\bar{\zeta}} = \omega(\zeta_{\text{can}}), \text{ so that } \langle v_{\bar{\zeta}}, u_{\bar{\zeta}} \rangle = 1.$$

Then $\langle \omega, v_{\bar{\zeta}} \rangle$ defines an unbounded solution of $L_{\frac{1}{2}, \frac{1}{2}, 1}$ in $\mathbb{D}_{\mathbb{C}_p}(\zeta_{\text{can}}, 1^-)$.

3.3.6. Finally, in any supersingular disk $\mathbb{D}_{\mathbb{C}_p}(\zeta_i, 1^-)$, there is no non-zero bounded solution of $L_{\frac{1}{2}, \frac{1}{2}, 1}$.

3.4. The Tate and Dwork-Serre-Tate parameters.

3.4.1. It turns out that the formal group $\widehat{X}_{(\widehat{\mathcal{R}}_{\text{ord}}^{\text{et}})}$ of X over $\widehat{\mathcal{R}}_{\text{ord}}^{\text{et}}$ is isomorphic to $\widehat{\mathbb{G}}_m$ (cf. [vdM89, I]). Any such isomorphism transforms the canonical differential ω_{can} on $\widehat{\mathbb{G}}_m$ into $\Theta \frac{dx}{y}$, for a suitable element $\Theta \in \widehat{\mathcal{R}}_{\text{ord}}^{\text{et}}$ well-defined up to multiplication by an element of \mathbb{Z}_p^\times . It is possible to arrange normalizations so that the following relation hold (loc. cit.):

$$\Theta = \left(\sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; \eta\right) \right)^{-1}.$$

Let $\zeta \in \widehat{\mathbb{Z}}_p^{\text{ur}}, \zeta \neq 0, 1$, be a point of S . By specialization, the element $\Theta(\zeta) \in \widehat{\mathbb{Z}}_p^{\text{ur}\times}$ (well-defined up to \mathbb{Z}_p^\times) links up ω_{can} and $\frac{dx}{y}$ on $\widehat{X}_\zeta \simeq \widehat{\mathbb{G}}_m$; it is called the *Tate parameter* or *Tate constant* of X_ζ . It is well-defined up to multiplication by an element of \mathbb{Z}_p^\times .

This is the p -adic analogue of the following familiar situation: let us consider the two-step uniformization of a complex elliptic curve

$$\mathbb{C} \xrightarrow{\exp\left(\frac{2i\pi}{\omega_1}\cdot\right)} \mathbb{C}^\times \longrightarrow \mathbb{C}^\times/q^{\mathbb{Z}} \simeq \mathbb{C}/(\omega_1\mathbb{Z} + \omega_2\mathbb{Z})$$

with $q = \exp(2i\pi\frac{\omega_2}{\omega_1})$. If the elliptic curve is X_ζ with $\zeta \in \mathbb{D}(0, 1^-) \setminus \{0\}$, and if ω_1 and ω_2 are fundamental periods of $\frac{dx}{y}$ (ω_1 being the period attached to the vanishing cycle), then $\frac{2i\pi}{\omega_1}$ appears as the analogue of $\Theta(\zeta)$, and it is well-known that $\omega_1/2i\pi = iF(\frac{1}{2}, \frac{1}{2}, 1; \zeta)$ (up to sign).

3.4.2. We come back to the symplectic basis $v_{\bar{\zeta}}, u_{\bar{\zeta}}$ and write $\frac{1}{\sqrt{-1}F_{\bar{\zeta}}}\omega$ in the form $v_{\bar{\zeta}} + \tau u_{\bar{\zeta}}$, where

$$\tau = -\frac{\langle \omega, v_{\bar{\zeta}} \rangle}{\langle \omega, u_{\bar{\zeta}} \rangle} \in (z - \zeta_{\text{can}})\widehat{\mathbb{Q}}_p^{\text{ur}}[[z - \zeta_{\text{can}}]].$$

This defines an unbounded element of $\mathcal{O}(\mathbb{D}_{\mathbb{C}_p}(\zeta_{\text{can}}, 1^-))$ (notice that another choice of $u_{\bar{\zeta}}$ multiplies τ by an element of \mathbb{Z}_p^\times).

Applying $\nabla(d/dz)$ to $\frac{1}{\sqrt{-1}F_\zeta}\omega = v_\zeta + \tau u_\zeta$ and using $\langle \omega, \nabla(\frac{d}{dz})(\omega) \rangle = \frac{2}{z(z-1)}$, one derives:

$$\frac{d\tau}{dz} = \frac{2}{z(1-z)F_\zeta^2}.$$

The exponential of τ is called the *Dwork-Serre-Tate parameter*. We shall recall later its meaning as a parameter for p -divisible groups. It satisfies the following remarkable integrality property [Dw69, Th.4], [Ka73]

$$q = e^\tau \in 1 + (z - \zeta_{\text{can}})\widehat{\mathbb{Z}}_p^{\text{ur}}[[z - \zeta_{\text{can}}]].$$

This suggests the following question: does q arise from an element of $\widehat{\mathcal{R}}_{\text{ord}}^{\text{et}}$? The answer is *no*. This may be seen by considering $q \pmod{p}$: the formula for $d\tau/dz = \text{dlog}(q)$ shows that $q \pmod{p}$ is non-constant ($\text{dlog}(q) \equiv \frac{2(h_p)^{2/p-1}}{z(z-1)} \pmod{p}$). If q comes from an element of $\widehat{\mathcal{R}}_{\text{ord}}^{\text{et}}$ which specializes to 1 at every canonical modulus ζ_{can} , one has $q \equiv 1 \pmod{p}$, a contradiction.

3.5. Complex counterpart: the supersingular locus.

Let us now revert things and try to understand the complex situation from the p -adic viewpoint!

Let D^+ and D^- be the (complex-conjugate) connected components of $D(\frac{1}{2}, \frac{3}{2}) \setminus (D(\frac{-1}{4}, \frac{3}{4}) \cup D(\frac{5}{4}, \frac{3}{4}))$. Let S be the complement of $D^+ \cup D^-$ in the complex plane. Note that S is closed and arcwise connected, and that $\pi_1^{\text{top}}(S)$ is a free group with two generators γ^+, γ^- (see figure 8). Its interior S° has three connected (simply-connected) components $D_0 = D(\frac{-1}{4}, \frac{3}{4})$, $D_1 = D(\frac{5}{4}, \frac{3}{4})$ and D_∞ .




FIGURE 8

Let $\mathcal{O}(S)$ denote the ring of continuous functions on S analytic in S° . According to Mergelyan's theorem [Mer54], they are uniform limits of rational functions on every compact $K \subset S$ such that $\pi_0(\mathbb{C} \setminus K, 0)$ is finite. This ring is however not stable under differentiation, and we consider its differential closure \mathcal{R} in $\mathcal{O}(S^\circ)$. We also consider the integral closure \mathcal{R}^{et} of \mathcal{R} in $\mathcal{O}(S^\circ)$. Every element of \mathcal{R}^{et} defines a multivalued locally analytic function on S , *i.e.* an analytic germ which may be analytically continued along any path of S not ending at $-1, \frac{1}{2}, 2$, in such a way that the germ at the other extremity is analytic.

For instance, $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ may be viewed as an element of \mathcal{R}^{et} : in fact, $F(\frac{1}{2}, \frac{1}{2}, 1; z)^4 \in \mathcal{O}(S)$. An explicit continuous extension of $F(\frac{1}{2}, \frac{1}{2}, 1; z)^4$ from D_0 to S is given by $F(\frac{1}{2}, \frac{1}{2}, 1; 1-z)^4$ in D_1 , $\frac{1}{z^2} F(\frac{1}{2}, \frac{1}{2}, 1; \frac{1}{z})^4$ in D_∞ , and the values $\frac{\Gamma(1/4)^8}{64\pi^6}$, $\frac{\Gamma(1/4)^8}{16\pi^6}$, $\frac{\Gamma(1/4)^8}{64\pi^6}$ at $-1, \frac{1}{2}, 2$ respectively [Car61, p.189]. On the other hand, note that $\text{dlog} F(\frac{1}{2}, \frac{1}{2}, 1; z)$ is in \mathcal{R} but not in $\mathcal{O}(S)$.

The de Rham cohomology of the Legendre elliptic pencil gives rise to an $\mathcal{R}[1/z(1-z)]$ -module with connection (\mathcal{H}, ∇) . It admits a *unique* non-zero horizontal submodule \mathcal{U} which extends to S ; we use the same symbol for the extension. It can be described along the following lines:

- For any $u \in (\mathcal{U} \otimes_{\mathcal{R}} \mathcal{R}^{\text{et}})^\nabla$, “the” image of u in $\mathcal{O}(D_j)$ ($j = 0, 1$, or ∞), is a bounded solution of the differential operator $L_{\frac{1}{2}, \frac{1}{2}, 1}$.
- $\mathcal{U} \otimes_{\mathcal{R}} \mathcal{R}^{\text{et}}[\frac{1}{z(1-z)}] = (\mathcal{H} \otimes_{\mathcal{R}[\frac{1}{z(1-z)}]} \mathcal{R}^{\text{et}}[\frac{1}{z(1-z)}])^\nabla \otimes_{\mathbb{C}} \mathcal{R}^{\text{et}}[\frac{1}{z(1-z)}]$.
- $(\mathcal{H} \otimes_{\mathcal{R}[\frac{1}{z(1-z)}]} \mathcal{R}^{\text{et}}[\frac{1}{z(1-z)}])^\nabla$ has a canonical $\mathbb{Z}[i]$ -submodule ($i = \sqrt{-1}$) which can be locally identified with the part $H^1(X_z^{\text{an}}, \mathbb{Z}[i])_{\text{isotriv}}$ of $H^1(X_z^{\text{an}}, \mathbb{Z}[i])$ where $\pi_1^{\text{top}}(S, z)$ acts through a finite group.
- $\mathcal{U}|_{D_0}^\nabla$ has a canonical \mathbb{Z} -submodule, which can be identified with the part of $H^1(X_z^{\text{an}}, \mathbb{Z}[i])_{\text{isotriv}}$ invariant under complex conjugation. One of the two generators u satisfies

$$\left\langle \omega, \frac{u}{2i\pi} \right\rangle = iF\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \text{ in } \mathcal{O}(D_0).$$

Similarly, $\mathcal{U}|_{D_1}^\nabla$ has a canonical \mathbb{Z} -submodule, and for one of the two generators u , one has

$$\left\langle \omega, \frac{u}{2i\pi} \right\rangle = iF\left(\frac{1}{2}, \frac{1}{2}, 1; 1-z\right) \text{ in } \mathcal{O}(D_1).$$

Note the occurrence of $2i\pi$ in these formulas.

- The local monodromy γ^+ around D^+ maps *onto*

$$\text{Aut } H^1(X_s, \mathbb{Z}[i])_{\text{isotriv}} \simeq \mathbb{Z}/4\mathbb{Z}.$$

Same for γ^- . The local monodromy at ∞ acts as ± 1 .

- In D^+ and in D^- , there is no non-zero bounded analytic solution of $L_{\frac{1}{2}, \frac{1}{2}, 1}$.

This is quite similar to §§3.2, §§3.3, $D^+ \cup D^-$ playing the role of the *supersingular locus*. This picture is assuredly very different from the traditional view of monodromy for $F(\frac{1}{2}, \frac{1}{2}, 1; z)$, and may shed some light upon the divergences between the topological and algebraic approaches to analytic continuation in the p -adic case (here of course, the coexistence of the two pictures is explained by the fact that the principle of unique continuation (Definition 1.1.1) fails for the sheaf of germs of continuous functions on S , analytic on S°).

The tale of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ is not finished: we have not yet explored the islands of supersingularity. We shall reach them in §§5.2.

4. Abelian periods as algebraic integrals.

ABSTRACT: We discuss periods of abelian varieties. Their p -adic counterparts live naturally in Fontaine's ring \mathbb{B}_{dR} . We present Colmez' construction of abelian p -adic periods, which relies on p -adic integration and reflects as closely as possible the complex picture. We also deal with the concrete evaluation of elliptic p -adic periods.

4.1. Over \mathbb{C} .

Let A be an abelian variety over \mathbb{C} of dimension g , and $\omega_1, \dots, \omega_g$ a basis of invariant differential forms. Let Λ be the image of the map

$$\iota : H_1(A(\mathbb{C}), \mathbb{Z}) \rightarrow \mathbb{C}^g ; \quad \gamma \mapsto \left(\int_{\gamma} \omega_i \right)_i .$$

One has a canonical isomorphism

$$\mathbb{C}^g / \Lambda \xrightarrow{\sim} A(\mathbb{C}).$$

Let us denote the projection $\mathbb{C}^g \rightarrow A(\mathbb{C})$ by pr . For any differential one-form ω of the second kind, one can consider a primitive function f_{ω} of $\text{pr}^*\omega$; it is meromorphic (univalued) on \mathbb{C}^g , because all residues of ω are 0 by definition, and it is unique up to addition of an arbitrary constant. Then $f_{\omega}(z_1 + z_2 + z_3) - f_{\omega}(z_1 + z_2) - f_{\omega}(z_1 + z_3) + f_{\omega}(z_1)$ defines a periodic function on $(\mathbb{C}^g)^3$, hence induces a meromorphic function on $A(\mathbb{C})^3$, which we denote by F_{ω}^3 . Note that the function F_{ω}^3 can also be defined purely algebraically by the following conditions:

- $F_{\omega}^3(z_1, 0, z_3) = F_{\omega}^3(z_1, z_2, 0) = 0$.
- $dF_{\omega}^3 = m_{123}^*\omega - m_{12}^*\omega - m_{13}^*\omega + m_1^*\omega$,

where m_{123} is the addition $A^3 \rightarrow A$ sending (z_1, z_2, z_3) to $z_1 + z_2 + z_3$, etc. For $\gamma \in H_1(A(\mathbb{C}), \mathbb{Z})$ the value $f_{\omega}(i(\gamma) + a) - f_{\omega}(a)$ does not depend on $a \in \mathbb{C}^g$ (a being chosen so that $i(\gamma) + a$ and a are not poles of f_{ω}), and one has the equality

$$f_{\omega}(i(\gamma) + a) - f_{\omega}(a) = \int_{\gamma} \omega,$$

which describes the period pairing

$$H_{\text{dR}}^1(A) \times H_1(A(\mathbb{C}), \mathbb{Z}) \longrightarrow \mathbb{C}$$

(or equivalently, the isomorphism $H_{\text{dR}}^1(A) \otimes \mathbb{C} \xrightarrow{\sim} H_{\text{B}}^1(A(\mathbb{C}), \mathbb{Z}) \otimes \mathbb{C}$.) Here $H_{\text{dR}}^1(A)$ denotes the first *algebraic* de Rham cohomology group of A , which coincides with the group of differential forms of the second kind modulo exact forms.

4.2. Over the p -adics; prolegomena

If we try to translate this into the p -adic setting, one has to face at once the problem: what is *integration over a loop*?

4.2.1. One tentative way is via Berkovich's theory, where loops do exist. Let us for instance consider the Legendre elliptic curve X_z . Viewed as a p -adic space, for $z \in D(0, 1^-) \setminus \{0\}$ ($p \neq 2$), this is a Tate curve: $X_z^{\text{an}} \simeq \mathbb{C}_p^\times / q^{\mathbb{Z}}$. The canonical differential ω_{can} inherited from \mathbb{C}_p^\times and $\frac{dx}{y}$ are proportional:

$$\omega_{\text{can}} = \Theta(z) \cdot \frac{dx}{y} \quad \text{with} \quad 1/\Theta(z) = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right).$$

The basic element γ of $H_1(X_z^{\text{an}}, \mathbb{Z})$ can be identified with the generator q of $q^{\mathbb{Z}}$, and it is natural to set $\int_\gamma \omega_{\text{can}} = \int_0^q \frac{dt}{t} = \log(q)$ (choosing a branch of the p -adic logarithm). One then has

$$\omega_2^{(p)} := \int_\gamma^{(p)} \frac{dx}{y} = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \log(q).$$

This formula, as well as

$$\begin{aligned} \sqrt{q} &= \frac{z}{16} e^{\frac{F^*\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)}{F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)}} \\ &\left(\text{with } F^*\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) = 4 \sum_{n>0} \binom{2n}{n}^2 \left(\sum_1^{2n} \frac{(-1)^{m-1}}{m}\right) \left(\frac{z}{16}\right)^n\right) \end{aligned}$$

are the same as those encountered in the complex situation. This leaves the open problem: how can one construct “the other period” $\omega_1^{(p)}$, *i.e.* the p -adic analogue of $\omega_1 = 2\pi\sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)$, since the corresponding loop is missing?

4.2.2. One simple tentative answer would be to replace the topological covering $\mathbb{C}_p^\times \rightarrow X_z^{\text{an}}$ by other étale coverings. Natural candidates for this purpose are étale coverings of order p^n , especially those corresponding to torsion points of order p^n which are close to the origin. Let ζ_{p^n} be a p^n th root of unity in \mathbb{C}_p^\times and let x_n be its image in X_z^{an} . In the corresponding complex case, taking $\zeta_{p^n} = e^{2i\pi/p^n}$ would give the right answer $p^n \int_0^{x_n} \frac{dx}{y} = \omega_1$. In the p -adic case, we get instead $p^n \int_0^{x_n} \frac{dx}{y} = 0$. Indeed, already in the case of the multiplicative group, we have $p^n \int_1^{\zeta_{p^n}} \frac{dt}{t} = p^n \log(1 + (\zeta_{p^n} - 1)) = 0$ p -adically: in other words, $2i\pi$ “*is missing*”.

4.2.3 (*Riemann-Shnirelman sums.*). It is appropriate to evoke here Shnirelman's approach to integration over loops, indeed one of the earliest works in p -adic analysis. This is an adaptation to the p -adic case of the computation of integrals by Riemann sums

$$\frac{1}{2i\pi} \int_{C(a,r)} f(z) dz = \int_0^1 f(a + re^{2i\pi\theta}) r e^{2i\pi\theta} d\theta = \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{\zeta^m=1} f(a + r\zeta) r\zeta.$$

Now, let $a, r \in \mathbb{C}_p$ and let f be a \mathbb{C}_p -valued function on the circumference $C(a, |r|)$. Let us denote the limit

$$\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{\zeta^m=1} f(a + r\zeta) \cdot r\zeta$$

symbolically by $\int_{C(a,r)} f(z) dz$. For better convergence, Shnirelman actually restricted the limit to those integers m prime to p . In the case of an analytic function f on $C(a, |r|)$, this restriction is unnecessary (one can even take $m = p^k$) and one has the following analogue of Cauchy's theorem of residues (*cf.* [Ko80, app.])

Lemma 4.2.4. *Assume that f is a meromorphic function on $D(a, |r|^+)$, and that its poles z_1, \dots, z_ν all lie in $D(a, |r|^-)$. Then $\int_{C(a,r)} f(z) dz$ exists and equals $\sum \text{Res}_{z_i} f$.*

In the (trivial) special case $f = 1/z$, we get $\int_{C(0,1)} dz/z = 1$ (not 0 as in 4.2.2!), but $2i\pi$ is still missing.

4.2.5. Actually, there is no way to remedy this if one remains in \mathbb{C}_p . A deeper reason for that, due to Tate, is that while $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ acts on the inverse system of p^n th roots of unity ($n \geq 0$) through the cyclotomic character χ , there is no element $(2i\pi)_p$ in \mathbb{C}_p such that $g((2i\pi)_p) = \chi(g)(2i\pi)_p$ for every $g \in \text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$.

It turns out that $(2i\pi)_p$ is in a sense the only “missing piece”: there is a good theory of p -adic periods (due to J.M. Fontaine, W. Messing [FM87], G. Faltings) which lives in some \mathbb{Q}_p -algebra isomorphic to $\mathbb{C}_p[[(2i\pi)_p]]$, and which we shall now describe in the case of abelian varieties.

4.3. The Fontaine ring B_{dR} .

4.3.1. *The ring R .* Let $\mathcal{O}_{\mathbb{C}_p}$ be the ring of integers of \mathbb{C}_p , *i.e.* $\{x \in \mathbb{C}_p, |x|_p \leq 1\}$. Let us set

$$R := \varprojlim_{x \mapsto x^p} \mathcal{O}_{\mathbb{C}_p},$$

i.e. R is the set of all series $(x^{(n)})_{n \in \mathbb{N}}$ such that $(x^{(n+1)})^p = x^{(n)}$. This is in fact a ring of characteristic p with

$$\left((x^{(n)}) + (y^{(n)}) \right)^{(n)} = \lim_{m \rightarrow \infty} \left(x^{(n+m)} + y^{(n+m)} \right)^{p^m}$$

and

$$\left((x^{(n)}) \cdot (y^{(n)}) \right)^{(n)} = x^{(n)} \cdot y^{(n)}.$$

Let $W(R)$ be the Witt ring with coefficients in R . for $x \in R$, let $[x]$ denote the Teichmüller representative in $W(R)$.

4.3.2. *The ring B_{dR}^+ .* To any element $(x_0, x_1, \dots, x_n, \dots) = \sum p^n [x_n^{p^{-n}}]$ in $W(\mathbb{R})$, we associate

$$\theta((x_n)) = \sum_{n=0}^{\infty} p^n x_n^{(n)}.$$

This defines a surjective homomorphism $\theta: W(\mathbb{R}) \rightarrow \mathcal{O}_{\mathbb{C}_p}$, whose kernel is principal. It extends to a homomorphism $\theta: W(\mathbb{R}) \left[\frac{1}{p} \right] \rightarrow \mathbb{C}_p$, and B_{dR}^+ is defined as the $(\text{Ker } \theta)$ -adic completion

$$B_{\text{dR}}^+ := \varprojlim W(\mathbb{R}) \left[\frac{1}{p} \right] / (\text{Ker } \theta)^n.$$

By continuity, θ further extends to a homomorphism $\theta: B_{\text{dR}}^+ \rightarrow \widehat{\mathbb{C}_p}$. Then B_{dR}^+ is a complete discrete valuation ring with maximal ideal $\text{Ker } \theta$ and residue field \mathbb{C}_p . Moreover, the Galois group $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ acts on B_{dR}^+ in such a way that θ is equivariant with respect to this Galois action,

$$\text{Gr}^* B_{\text{dR}}^+ \simeq \bigoplus_{r \in \mathbb{N}} \mathbb{C}_p(r),$$

where Gr^* refers to the filtration by the powers of $\text{Ker } \theta$, and where the ‘‘twist’’ (r) indicates that the $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ -action is twisted by the r th power of the cyclotomic character.

It turns out that B_{dR}^+ contains naturally a copy of $\overline{\mathbb{Q}_p}$ (which θ maps isomorphically to $\overline{\mathbb{Q}_p} \subset \mathbb{C}_p$). More precisely, P. Colmez [Co94] has shown that B_{dR}^+ is the separated *completion* of $\overline{\mathbb{Q}_p}$ with respect to the topology defined by taking $(p^n \mathcal{O}_{\overline{\mathbb{Q}_p}}^{(k)})_{n,k}$ as a basis of neighborhoods of 0, where $\mathcal{O}_{\overline{\mathbb{Q}_p}}^{(k)}$ denotes the subring of $\overline{\mathbb{Z}_p}$ defined inductively as

$$\mathcal{O}_{\overline{\mathbb{Q}_p}}^{(0)} = \overline{\mathbb{Z}_p}, \quad \mathcal{O}_{\overline{\mathbb{Q}_p}}^{(k)} = \text{Ker} \left(d: \mathcal{O}_{\overline{\mathbb{Q}_p}}^{(k-1)} \rightarrow \Omega_{\mathcal{O}_{\overline{\mathbb{Q}_p}}^{(k-1)}/\mathbb{Z}_p}^1 \otimes \overline{\mathbb{Z}_p} \right).$$

It is easy to deduce from this description that B_{dR}^+ contains $\widehat{\overline{\mathbb{Z}_p^{\text{ur}}}}$.

4.3.3. *Some remarkable elements of B_{dR}^+ .* For any $x \in B_{\text{dR}}^+$ such that $|\theta(x) - 1|_p < 1$, the series $\log(x) = -\sum_{n>0} \frac{(1-x)^n}{n}$ converges in B_{dR}^+ . In particular, let $\underline{z} = (\dots, z^{(1)}, z^{(0)})$ be an element of \mathbb{R} such that $z^{(0)} \in \overline{\mathbb{Z}_p}$ or $\widehat{\overline{\mathbb{Z}_p^{\text{ur}}}}$. Then one can define the element $\text{Log } \underline{z} = \log\left(\frac{z^{(0)}}{|\underline{z}|}\right)^{(4)}$. Note that $\theta(\text{Log } \underline{z}) = 0$.

In the special case where each $z^{(n)} = \zeta_{p^n}$ is a primitive p^n th root of unity, $\text{Log}(\dots, \zeta_{p^n}, \dots, 1)$ is the element $(2i\pi)_p$ we were looking for. This is a generator of $\text{Ker } \theta$, and $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ acts on it through the cyclotomic character; in other words, we have a $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ -equivariant isomorphism

$$\text{Gr}^* B_{\text{dR}}^+ \simeq \mathbb{C}_p[(2i\pi)_p].$$

⁽⁴⁾we follow Colmez’ sign convention; Fontaine’s LOG is $\log - \text{Log}$.

Note that another choice of $(\dots, \zeta_{p^n}, \dots, 1)$ changes $(2i\pi)_p$ by multiplication by a unit in \mathbb{Z}_p . Similarly, up to addition of an element of $(2i\pi)_p \mathbb{Z}_p$, $\text{Log } \underline{z}$ depends only on $z^{(0)}$; it is sometimes simply denoted by $\text{Log } z^{(0)}$.

By choosing a double embedding of $\overline{\mathbb{Q}}$ into \mathbb{C} and $\overline{\mathbb{Q}}_p$, one obtains a canonical element $(2i\pi)_p$, attached to the sequence $(\dots, e^{2i\pi/p^n}, \dots, 1)$. Similarly, if $z^{(0)} \in \overline{\mathbb{Q}}$, then $\text{Log } z^{(0)}$ is well-defined up to addition of an element of $(2i\pi)_p \mathbb{Z}$.

4.4. Colmez' construction of abelian p -adic periods.

Let A be an abelian variety defined over a p -adic local field K . The Fontaine-Messing p -adic period pairing is a pairing

$$\int^{(p)} : H_{\text{dR}}^1(A) \otimes T_p(A_{\overline{K}}) \longrightarrow B_{\text{dR}}^+$$

where $T_p(A_{\overline{K}}) = \varprojlim \text{Ker}([p^n] : A_{\overline{K}} \rightarrow A_{\overline{K}})$ is the Tate module (a \mathbb{Z}_p -module of rank $2 \dim A$ with $\text{Gal}(\overline{K}/K)$ -action).

We present Colmez' construction, which is parallel to 4.1.

Let ω be a differential form on A of second kind, and F_ω^3 the function on A^3 determined as in 4.1. Then:

Proposition 4.4.1 ([Co92, 4.1]). *There exists a locally meromorphic function F_ω on $A(B_{\text{dR}})$, unique up to constant, such that:*

- (1) $dF_\omega = \omega$.
- (2) $F_\omega(z_1 + z_2 + z_3) - F_\omega(z_1 + z_2) - F_\omega(z_1 + z_3) + F_\omega(z_1) = F_\omega^3(z_1, z_2, z_3)$.
- (3) If $\omega = dF$, then $F_\omega = F$.

Moreover, if $\alpha : A_1 \rightarrow A_2$ is a morphism of abelian varieties and ω is a differential form of second kind on A_2 , then $F_{\alpha^*\omega} = \alpha^*F_\omega$.

Note that the function F_ω is *not* multivalued; this fact comes from the following lemma specific to the p -adic case:

Lemma 4.4.2 ([Co91, 4.3]). *For any neighborhood V of 0 in $A(B_{\text{dR}}^+)$, there exists an open subgroup U of $A(B_{\text{dR}}^+)$ contained in V such that $A(B_{\text{dR}}^+)/U$ is a torsion group.*

Take a proper model \mathcal{A} of A over \mathcal{O}_K . Let $\gamma = (\dots, u_n, \dots, u_2, u_1 = 0) \in T_p(A_{\overline{K}})$ with each $u_n \in \mathcal{A}(\mathcal{O}_{\mathbb{C}_p})$, and choose $a_n \in \mathcal{A}(B_{\text{dR}}^+)$ so that neither a_n nor $a_n +_A \widehat{u}_n$ is close to a pole of ω . For suitable liftings $\widehat{u}_n \in \mathcal{A}(B_{\text{dR}}^+)$ of u_n (i.e. $\theta(\widehat{u}_n) = u_n$), the following holds.

Theorem 4.4.3 ([Co91, 5.2]). *The limit*

$$\oint_\gamma^{(p)} \omega := \lim_{n \rightarrow \infty} p^n (F_\omega(a_n) - F_\omega(a_n +_A \widehat{u}_n))$$

converges to an element in \mathbf{B}_{dR}^+ , and it defines a non degenerate bilinear pairing

$$\oint^{(p)} : \mathbf{H}_{\text{dR}}^1(A) \otimes T_p(A_{\overline{K}}) \longrightarrow \mathbf{B}_{\text{dR}}^+,$$

compatible with the Galois action and the filtrations.

Remark 4.4.4. On composing the period pairing $\oint^{(p)}$ with θ , one gets a bilinear map $\mathbf{H}_{\text{dR}}^1(A) \otimes T_p(A_{\overline{K}}) \longrightarrow \mathbb{C}_p$ which sends $\Omega^1(A) \subset \mathbf{H}_{\text{dR}}^1(A)$ to 0. This degenerate pairing describes “half” of the (Hodge)-Tate-Raynaud decomposition

$$\mathbf{H}_{\text{et}}^1(A_{\overline{K}}, \mathbb{Q}_p) \otimes \mathbb{C}_p \simeq \Omega^1(A) \otimes \mathbb{C}_p(-1) \oplus \mathbf{H}^1(\mathcal{O}(A)) \otimes \mathbb{C}_p$$

of $\text{Gal}(\overline{K}/K)$ -modules.

4.5. Some computations of elliptic p -adic periods.

4.5.1. Tate elliptic curves. In this case, $T_p(A_{\overline{K}})$ sits in an exact sequence

$$0 \rightarrow \mathbb{Z}_p(1) \rightarrow T_p(A_{\overline{K}}) \rightarrow \mathbb{Z}_p \rightarrow 0.$$

Let us take for γ_1 the image of $(2i\pi)_p \in \mathbb{Z}_p(1)$ in $T_p(A_{\overline{K}})$, and for γ_2 any lifting of $1 \in \mathbb{Z}_p$. For concreteness, let $A = X_z$ be in Legendre form as in 4.2.1. Let $\eta \in \mathbf{H}_{\text{dR}}^1$ satisfy $\langle \omega, \eta \rangle = 1$ (e.g. $\frac{x dx}{4y}$). We set $\omega_i^{(p)} = \int_{\gamma_i}^{(p)} \omega$, $\eta_i^{(p)} = \int_{\gamma_i}^{(p)} \eta$. Then one has the “Legendre relation”

$$\omega_1^{(p)} \eta_2^{(p)} - \eta_1^{(p)} \omega_2^{(p)} = (2i\pi)_p$$

and the formulas (cf. [And90], [And96])

$$\frac{\omega_1^{(p)}}{(2i\pi)_p} = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) = 1/\Theta(z), \quad \omega_2^{(p)} = \sqrt{-1}F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right) \text{Log } q$$

(compare with the complex case in 3.4.1, and also [Car61, p.406]). Note that the “period” we found in 4.2.1 by integration along the Berkovich loop is *not* a Fontaine-Messing period (there is a log instead of Log).

4.5.2. Elliptic curves with ordinary reduction. We now assume that $z \in \mathbb{Z}_p^{\text{ur}}$ and that $A = X_z$ has good ordinary reduction (with the notation of §§3.1, this means that $z(1-z)h_p(z) \neq 0$). Again, $T_p(A_{\overline{\mathbb{Q}_p}})$ sits in an exact sequence ([Ser68a, A.2.4])

$$0 \rightarrow \mathbb{Z}_p \gamma_1 \rightarrow T_p(A_{\overline{\mathbb{Q}_p}}) \rightarrow T_p(A \bmod p) \rightarrow 0.$$

Then one still has the formula $\Theta(z) = \frac{(2i\pi)_p}{\omega_1^{(p)}}$, where $\Theta(z) \in \widehat{\mathbb{Z}_p^{\text{ur}}}$ is the Tate constant discussed in 3.4.1 (compare [And90], [dSh87, 4.3]). In particular, $\frac{\omega_1^{(p)}}{(2i\pi)_p}$ is given by the evaluation at z of “the” extension of $F(\frac{1}{2}, \frac{1}{2}, 1; ?)$ discussed in §§3.4.

On the other hand, let $\gamma_2 \in T_p(A_{\overline{\mathbb{Q}}_p})$ be such that the Weil pairing $\langle \gamma_1, \gamma_2 \rangle \in \mathbb{Z}_p(1) = \mathbb{Z}_p(2i\pi)_p$ is the chosen generator $(2i\pi)_p$. Then one has the “Legendre relation”, and $\omega_2^{(p)} = \omega_1^{(p)} \cdot \frac{\text{Log } q}{(2i\pi)_p}$, where $q \in \widehat{\mathbb{Z}}_p^{\text{ur}}$ denotes here the so-called *Dwork(-Serre-Tate) parameter* of A introduced in 3.4.2. If $q = 1$, *i.e.* when A is the *canonical lifting* of $A \pmod{p}$, then one can choose γ_2 such the corresponding value of $\text{Log } 1$ is 0, and then $\omega_2^{(p)} = 0$.

4.6. Periods of CM elliptic curves and values of the Gamma function.

4.6.1. The case of an elliptic curve A with *supersingular reduction* is more delicate. If A does not have complex multiplication (CM), one can show, using [Ser68a, A.2.2], that the four basic periods $\omega_1^{(p)}, \omega_2^{(p)}, \eta_1^{(p)}, \eta_2^{(p)}$ are algebraically independent over $\overline{\mathbb{Q}}_p$, so that one cannot expect “formulas” for the periods.

On the other hand, it is well-known that the complex periods of an elliptic curve with complex multiplication can be expressed in terms of special values of the Γ function. So one may ask for a p -adic analogue involving Γ_p (*cf.* 2.6). Such a formula has been found by A. Ogus, actually not for the periods, but for the action of Frobenius.

4.6.2. *The Lerch-Chowla-Selberg-Ogus formulas.* Let A be an elliptic curve with complex multiplication by $\mathbb{Q}(\sqrt{-d})$ over $\overline{\mathbb{Q}}$ ($-d$ denotes a fundamental discriminant). Let $\epsilon = \left(\frac{-d}{\cdot}\right)$ be the quadratic character $(\mathbb{Z}/d)^\times \rightarrow \mathbb{Z}/2$ induced by the embedding $\mathbb{Q}(\sqrt{-d}) \subset \mathbb{Q}(\zeta_d)$, h = the class number of $\mathbb{Q}(\sqrt{-d})$, and w = the number of roots of unity in $\mathbb{Q}(\sqrt{-d})$. For any $u \in (\mathbb{Z}/d)^\times$, we denote by $\langle \frac{u}{d} \rangle$ the unique rational number in $]0, 1]$ such that $d \langle \frac{u}{d} \rangle \equiv u$.

Let v be a place of $\overline{\mathbb{Q}}$ of residue characteristic p , with associated embedding $\overline{\mathbb{Q}} \subset \overline{\mathbb{Q}}_p$. To any lifting ψ_v to $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ of the Frobenius element in $\text{Gal}(\mathbb{Q}_p^{\text{ur}}/\mathbb{Q}_p)$, one attaches a ψ_v -linear endomorphism Ψ_v of $H_{\text{dR}}^1(A/\overline{\mathbb{Q}}) \otimes \overline{\mathbb{Q}}_p$ ⁽⁵⁾. On the other hand, it is well-known that A has supersingular reduction at v if and only if $\epsilon(p) = -1$ or 0.

Theorem 4.6.3. *There exists a basis (ω, η) of $H_{\text{dR}}^1(A/\overline{\mathbb{Q}})$ of eigenvectors under the action of $\sqrt{-d}$ (ω being the class of a regular differential), and an element $\gamma \in H_1(A(\mathbb{C}), \mathbb{Q})$, such that*

$$\int_{\gamma} \omega = \sqrt{2i\pi} \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma \left\langle \frac{u}{d} \right\rangle \right)^{\epsilon(u)w/4h}$$

$$\int_{\gamma} \eta = \sqrt{2i\pi} \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma \left\langle -\frac{u}{d} \right\rangle \right)^{\epsilon(u)w/4h}$$

⁽⁵⁾using its crystalline interpretation, *cf.* [Be083, 4]; we refer to [Ch198] for a nice recent survey of crystalline cohomology.

and such that for every place of $\overline{\mathbb{Q}}$ of residue characteristic p satisfying $\epsilon(p) = -1$,

$$\begin{aligned}\Psi_v^*(\omega) &= p \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/4h} \eta \\ \Psi_v^*(\eta) &= \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle -p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/4h} \omega\end{aligned}$$

up to multiplication by some root of unity.

This is a concatenation of [O90, 3.15, 3.9], taking into account the formula $2hd/w = -\sum_1^d \epsilon(u)u$, cf. [H81, VII].

Remark. The last two formulas extend to the case when $\epsilon(p) = +1$ if one interchanges ω and η in the right hand sides (for $d = 4$ and $p \equiv 1 \pmod{4}$), this is compatible with Young's formula in §§3.3, since $\Gamma_p(1/2)^4 = 1$, and with the formula $F(\frac{1}{2}, \frac{1}{2}, 1; -1) = \frac{1}{2} \frac{\Gamma(1/4)}{\Gamma(1/2)\Gamma(3/4)}$.

In that case, the expressions

$$\prod \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/4h}, \quad \prod \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/4h}$$

are algebraic numbers: indeed, let r be the order of the subgroup $\langle p \rangle$ of $(\mathbb{Z}/d)^\times$ generated by p , and $k = p^r - 1/d$; then

$$\prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)} = \prod_{w \in (\mathbb{Z}/d)^\times / \langle p \rangle} \prod_1^r \Gamma_p \left(\frac{p^i w k}{p^r - 1} \right)^{-\epsilon(w)}$$

and one concludes by the Gross-Koblitz formula.

4.6.4. The ramified case. In the ramified case, *i.e.* when $\epsilon(p) = 0$, there is again an analogue of the last two formulas for $p \neq 2$. This relies on Coleman's computation of the Frobenius matrix of Fermat curves of degree divisible by p — which have arboreal reduction modulo p . The result takes the same form as in the case $\epsilon(p) = -1$: Ψ is now attached to an element ψ of degree one in the Weil group (*i.e.* a lifting of the Frobenius element in $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$), the expressions $p \frac{u}{d}$ have to be replaced by $\psi \frac{u}{d}$, and Γ_p must be extended to \mathbb{Q}_p [Col90, 6.5].

Coleman's complicated expressions have been simplified by F. Urfels in his thesis (Strasbourg, 1998; unpublished). If one passes to the Iwasawa logarithm \log_p (at the cost of rational powers of p), the result is that one should replace $\log_p \Gamma_p \left\langle \psi \frac{u}{d} \right\rangle$ in Ogus' formula by $G_p(\left\langle \psi \frac{u}{d} \right\rangle) - G_p(\left\langle \frac{u}{d} \right\rangle)$, where G_p denotes Diamond's LogGamma function

$$G_p(x) = \lim_{m \rightarrow \infty} \frac{1}{p^m} \sum_{n=0, \dots, p^m-1} (x+n) \log_p(x+n) - (x+n).$$

Example. We take $p = 3$. Let $\mathbb{Q}[\sqrt{-3n}]$ be an imaginary quadratic field with fundamental discriminant $-3n$, n prime to 3. Let A be an elliptic curve with complex multiplication by some order in $\mathbb{Q}[\sqrt{-3n}]$.

Let ψ be an element of the Weil group which acts by -1 on $\mathbb{Q}_3/\mathbb{Z}_3 \cong \mu_{3^\infty}$. One has a formula $\Psi_v^*(\omega) = \kappa\eta$ with

$$\log_3 \kappa = -(\epsilon(u)w/4h) \sum_{u \in (\mathbb{Z}/3n)^\times} \left(G_3 \left(\left\langle \psi \frac{u}{3n} \right\rangle \right) - G_3 \left(\left\langle \frac{u}{3n} \right\rangle \right) \right)$$

We notice that for $u \in (\mathbb{Z}/3n)^\times$, $\epsilon \left(3n \left\langle \psi \frac{u}{3n} \right\rangle \right) = -\epsilon(u)$, whence

$$\log_3 \kappa = (w/2h) \sum_{u \in (\mathbb{Z}/3n)^\times} \epsilon(u) G_3 \left(\left\langle \frac{u}{3n} \right\rangle \right).$$

On the other hand, for $p = 3$, the Teichmüller character is nothing but the Legendre symbol $\left(\frac{\cdot}{3} \right)$. Using this, the latter expression can be rewritten, by the Ferrero-Greenberg formula (*cf.* [La90, chap. 17]), as

$$\begin{aligned} \log_3 \kappa &= (w/2h) \cdot (L'_3(0, \epsilon) + L_3(0, \epsilon) \log_3 n) \\ &= (w/2h) \cdot \sum_{v \in (\mathbb{Z}/n)^\times} \left(\frac{n}{v} \right) \log_3 \Gamma_3 \left\langle \frac{v}{n} \right\rangle. \end{aligned}$$

When $\left(\frac{n}{3} \right) = 1$, then $L_3(0, \epsilon) = 0$ and the Gross-Koblitz formula shows that $L'_3(0, \epsilon) = \sum \left(\frac{n}{v} \right) \log_3 \Gamma_3 \left\langle \frac{v}{n} \right\rangle$ is the Iwasawa logarithm of an algebraic number (a Gauss sum, *cf. loc. cit.*), hence $\kappa \in \overline{\mathbb{Q}}$.

A particularly simple case is $n = 1$: one finds $\log_3 \kappa = 0$, thus $\kappa \in 3^{\mathbb{Q}} \cdot \mu_\infty \subset \overline{\mathbb{Q}}$.

For $n = 8$, a contrario, one has $\left(\frac{n}{3} \right) = -11$: one finds

$$\log_3 \kappa = (1/2) \cdot \left[\log_3 \Gamma_3 \left\langle \frac{1}{8} \right\rangle - \log_3 \Gamma_3 \left\langle \frac{3}{8} \right\rangle - \log_3 \Gamma_3 \left\langle \frac{5}{8} \right\rangle + \log_3 \Gamma_3 \left\langle \frac{7}{8} \right\rangle \right] = 0$$

by the functional equation of Γ_3 , so that again $\kappa \in 3^{\mathbb{Q}} \cdot \mu_\infty \subset \overline{\mathbb{Q}}$.

4.6.5. Colmez' product formula. There is a natural extension of $|\cdot|_p$ on $\overline{\mathbb{Q}}_p$ to B_{dR}^+ (however not as an absolute value), such that $|(2i\pi)_p|_p = p^{-\frac{1}{p-1}}$ *cf.* [Co91].

Colmez has remarked that the logarithm of the product $|2i\pi| \prod_p |(2i\pi)_p|_p = 2\pi \prod_p p^{-\frac{1}{p-1}}$ is formally equal to $\log(2\pi) + \zeta'(1)/\zeta(1)$, a divergent sum which can be renormalized using the functional equation of ζ : setting $\zeta'(1)/\zeta(1) = -\zeta'(0)/\zeta(0) = -\log(2\pi)$, the renormalized product is

$$|2i\pi| \prod_p' |(2i\pi)_p|_p = 1.$$

He has given an amazing generalization of this product formula to periods of CM elliptic curves and many other CM abelian varieties (*loc. cit.*).

5. Periods as solutions of the Gauss-Manin connection.

ABSTRACT: The variation of the periods in a family of complex algebraic varieties is controlled by the Picard-Fuchs differential equation. We discuss the p -adic situation, present Dwork's general viewpoint on p -adic periods. We then discuss the question of the existence of an arithmetic structure on the space of solutions, which would give an intrinsic meaning to the period modulo $\overline{\mathbb{Q}}^\times$. We present such an arithmetic structure (analogous to the Betti lattices in the complex situation) in the case of abelian varieties with either multiplicative reduction or supersingular reduction. In the latter case, we relate the periods, in the presence of complex multiplication, to special values of the p -adic Gamma function.

5.1. Stokes.

We come back to the basic question of the meaning of *integration over a loop*, already discussed in §§4.2. A different approach is based on the Stokes lemma: integrating *exact* differentials on a loop gives zero. This is the approach favored by K. Aomoto in the complex case and by B. Dwork in the p -adic case.

5.1.1. In order to see how this idea can be implemented, let us consider the Hankel expression for the gamma function

$$\frac{1}{\Gamma(1-\alpha)} = \frac{1}{2i\pi} \int_{\gamma} x^{\alpha} e^x \frac{dx}{x}$$

where γ is the following loop based at $-\infty$.



FIGURE 9

Stokes' lemma suggests to attach to this integral the following complex

$$x^{\alpha} e^x \mathcal{O} \xrightarrow{xd/dx} x^{\alpha} e^x \mathcal{O}$$

where \mathcal{O} is a suitable ring of analytic functions. From the point of view of index theory (Malgrange-Ramis), a natural choice is

$$\mathcal{O} = \mathbb{C}[[x]]_{-1,1^-} = \left\{ \sum_{n \geq 0} a_n x^n \mid \exists \kappa > 0, \exists r \in]0, 1[, |a_n| \leq \kappa r^n / n! \right\}$$

identified with the ring of entire functions of *exponential order* $O(e^{r|x|})$ for some $r < 1$ ⁽⁶⁾. Using the formula $x^{\alpha} e^x x^k = \frac{-1}{\alpha+k} x^{\alpha} e^x x^{k+1} + xd/dx(\frac{1}{\alpha+k} x^{\alpha} e^x x^k)$, a simple computation then shows that $H^0 = 0$ and that H^1 is of dimension 1; it is generated by $[\frac{1}{2i\pi} x^{\alpha} e^x]$, whose integration along γ gives $\frac{1}{\Gamma(1-\alpha)}$.

⁽⁶⁾we use the traditional notation for rings of Gevrey series; the value $(-1, 1^-)$ of the index is "characteristic", *i.e.* an extremal value for which H^1 is of dimension one.

The p -adic analogue

$$\mathbb{C}_p[[x]]_{-1,1^-} = \left\{ \sum_{n \geq 0} a_n x^n \mid \exists \kappa > 0, \exists r \in]0, 1[, |a_n| \leq \kappa r^n / |n!|_p \right\}$$

is nothing but the ring of *overconvergent* analytic functions on $\mathbb{D}(0, |\pi|_p)$ (here π is Dwork's constant). By change of variable $z = \pi x$, we thus recover the complex $M_\alpha^\dagger \xrightarrow{zd/dz} M_\alpha^\dagger$ of 2.6, where $\Gamma_p(\alpha)$ ($= \pm \frac{1}{\Gamma_p(1-\alpha)}$) appeared.

5.1.2. This approach inspired by Stokes' lemma is well-suited for studying many kinds of hypergeometric functions (confluent or not). Basically, the integral will satisfy difference equations with respect to the exponents (α in the previous example), and differential algebraically on parameters. In the complex situation, one of the main problems is the construction of nice loops (such as γ); this is the object of a so-called "topological intersection theory" generalizing the usual Betti homology and period pairing. In the p -adic case, the focus has been more on the construction and properties of the Frobenius structure, in particular its analyticity with respect to the exponents ("Boyarsky principle" [Dw83]).

5.1.3. When no exponential is involved and when the exponents are rational, the integrand is algebraic; if it depends algebraically on parameters, the period integral is a solution of a *Gauss-Manin connection*.

Let us say a few words about the classical case of $F(a, b, c; z)$. We assume that $a, b, c \in \mathbb{Z}_p$, and that $c - a, c - b, b, a$ all lie outside \mathbb{Z} . We set

$$f_{a,b,c;z} = x^b(1-x)^{c-b}(1-zx)^{-a}$$

where, for simplicity, z is limited to $|z(1-z)| = 1$. Let Z be the affinoid $\mathbb{D}(0, 1^+) \setminus (\mathbb{D}(0, 1^-) \cup \mathbb{D}(1, 1^-) \cup \mathbb{D}(1/z, 1^-))$. The relevant complex is

$$f_{a,b,c;z} \mathcal{H}^\dagger(Z) \xrightarrow{xd/dx} f_{a,b,c;z} \mathcal{H}^\dagger(Z).$$

Its $H^1 = H_{a,b,c}^1$ is of dimension 2; it is generated by $[f_{a,b,c;z}]$ and $[\frac{f_{a,b,c;z}}{1-x}]$. The Gauss-Manin connection is

$$\nabla \left(\frac{d}{dz} \right) \begin{pmatrix} [f_{a,b,c;z}] \\ [\frac{f_{a,b,c;z}}{1-x}] \end{pmatrix} = \begin{pmatrix} -\frac{c}{z} & \frac{c-b}{z} \\ \frac{c-a}{1-z} & \frac{a+b-c}{1-z} \end{pmatrix} \begin{pmatrix} [f_{a,b,c;z}] \\ [\frac{f_{a,b,c;z}}{1-x}] \end{pmatrix}$$

cf. [Dw82, 3.1, 1.2]. The Frobenius structure relates $H_{a,b,c}^1$ to $H_{a',b',c'}^1$, where a', b', c' are the successors of a, b, c respectively (*cf.* 2.6). For all these hypergeometric series, there is a story similar to that of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$. In this context, it is fruitful to combine the Boyarsky principle and the contiguity relations.

5.1.4. In this sketch of Dwork's approach, "periods" are just analytic solutions of the Gauss-Manin connection. One can ask more: namely, one can ask for a dual theory of p -adic cycles and a period pairing as in the complex

case⁽⁷⁾. At best, we can expect these “ p -adic Betti lattices” to be locally horizontal, functorial with respect to the endomorphisms of the geometric fibres, and defined over \mathbb{Z} or at least over some number field.

It turns out that this can be done in some cases, *e.g.* in the cases of abelian varieties with multiplicative reduction, and of abelian varieties with supersingular reduction ([And90], [And95]). Let us outline the results.

5.2. p -adic Betti lattices for abelian varieties with multiplicative reduction.

5.2.1. Over \mathbb{C} . We first recall the “multiplicative uniformization” of complex abelian varieties. Let A be an abelian variety of dimension g over \mathbb{C} . One has the analytic representation

$$A(\mathbb{C}) = T(\mathbb{C})/M,$$

where T is a torus of dimension g , and M is a lattice of rank g . Set $M' := \text{Hom}(T, \mathbb{G}_m) (= M_{A^\vee})$. Then we have the exact sequence

$$0 \longrightarrow 2i\pi M'^\vee \longrightarrow \Lambda \longrightarrow M \longrightarrow 0,$$

where Λ is the period lattice (of rank $2g$), and this sequence splits by choosing a branch of \log . The inclusion $M \hookrightarrow T$ defines a pairing $M \times M' \rightarrow \mathbb{C}^\times$. A polarization, on the other hand, induces $M \rightarrow M'$. These data give rise to the pairing (“multiplicative period”)

$$q = (q_{ij}): M \otimes M \longrightarrow \mathbb{C}^\times$$

and $-\log |q|$ is a scalar product on $M_{\mathbb{R}}$.

5.2.2. Over the p -adics. Let K be a p -adic local field and A an abelian variety over K having (split) multiplicative reduction. One has a similar analytic representation [Mu72a]

$$A(K) = T(K)/M$$

Set again $M' := \text{Hom}(T, \mathbb{G}_m)$. We have the following identifications

$$M \simeq H_1(A^{\text{rig}}, \mathbb{Z}) \quad \text{and} \quad M' \simeq H_1(A^{\vee \text{rig}}, \mathbb{Z}),$$

where A^{rig} denotes the associated rigid analytic variety over K to A . Then, just as in the complex case, one can construct a natural exact sequence:

$$0 \longrightarrow (2i\pi)_p M'^\vee \longrightarrow \Lambda \longrightarrow M \longrightarrow 0,$$

split by the choice of a branch of the p -adic logarithm \log_p , and a non-degenerate pairing

$$\int^p : H_{\text{dR}}^1(A) \times \Lambda \longrightarrow K[(2i\pi)_p].$$

⁽⁷⁾Dwork has developed a “dual theory” and a pairing given by residues, which play somehow the role of “ p -adic cycles” (*cf.* [Ro86, 5]); however, this dual theory is an avatar of de Rham cohomology with proper supports and does not enjoy the properties of discreteness and horizontality that we are looking for.

(The construction involves one-motives and Frobenius [And90]). Restricted to $(2i\pi)_p M'^V$ which is in a canonical way a subgroup of $T_p(A_{\overline{K}})$, this pairing coincides with the Fontaine-Messing pairing.

5.2.3. Degenerating abelian pencils. Let $\mathcal{A} \rightarrow \mathbb{A}^1 \setminus \{0, \zeta_1, \dots, \zeta_r\}$ be a pencil of polarized abelian varieties defined over a number field k . We consider an element ω of the relative H_{dR}^1 and its associated Picard-Fuchs differential equation $\mathcal{L}\omega = 0$. We assume that the connected component of identity of the special fibre at 0 of the Neron model of \mathcal{A} is a split torus T over k . The dual abelian pencil then has the same property, with a torus T' . We set $M = \text{Hom}(T', \mathbb{G}_m)$, $M' = \text{Hom}(T, \mathbb{G}_m)$.

Let us fix $k \hookrightarrow \mathbb{C}$. The constant subsheaf of $R_1(f_{\mathbb{C}}^{\text{an}})_* \mathbb{Z}$ in the neighborhood of 0 identifies with $2i\pi M'^V$: its fibre at any $z (\neq 0, \zeta_i)$ is the lattice $2i\pi M'_z{}^V$ attached to \mathcal{A}_z (5.2.1). Similarly for M . On the other hand, the choice of a branch of log identifies $H_{1B}(\mathcal{A}_z, \mathbb{Z}) \simeq (R_1(f_{\mathbb{C}}^{\text{an}})_* \mathbb{Z})_z$ with $\Lambda = 2i\pi M'_z{}^V \oplus M_z$. Then for any $\gamma \in 2i\pi M'^V$, $y(z) = \frac{1}{2i\pi} \int_{\gamma_z} \omega_z$ is a solution of \mathcal{L} in $k[[z]]$.

More generally, for any $\gamma^* \in \Lambda$, $\frac{1}{2i\pi} \int_{\gamma_z^*} \omega_z$ is a solution of the Picard-Fuchs differential equation of the form $y(z) \frac{\log az^n}{2i\pi} + y^*(z)$, with $y^*(z) \in k[[z]]$, $a \in k$, $n \in \mathbb{Z}$. (In the case of the Legendre pencil, we have already met this situation with $y(z) = iF(\frac{1}{2}, \frac{1}{2}, 1; z)$, $y^*(z) = 2F^*(\frac{1}{2}, \frac{1}{2}, 1; z)$, $a = 2^{-8}$, $n = 2$).

Let now fix an embedding $k \hookrightarrow K \subset \mathbb{C}_p$. For any $z \neq 0$ close enough to 0, M and M' identify respectively with the lattices M_z , M'_z attached to \mathcal{A}_z (5.2.2), and the choice of a determination of the p -adic logarithm identifies Λ with Λ_z . It turns out that the p -adic pairing \int^p of 5.2.2 extends to a horizontal pairing over a punctured disk around 0, which is given by the “same” formula as in the complex case

$$\frac{1}{(2i\pi)_p} \int_{\gamma_z^*}^p \omega_z = y(z) \frac{\log_p az^n}{(2i\pi)_p} + y^*(z),$$

where $y(z), y^*(z)$ are the *same formal series as above, evaluated p -adically* (loc. cit.). The computation of 4.2.1 can be considered as a special case. This is also closely related to the work of T. Ichikawa on “universal periods” for Mumford curves [Ic97].

5.2.4. Relation to the Fontaine-Messing periods. Since the restriction of \int^p to $(2i\pi)_p M'_z{}^V \subset T_p(\mathcal{A}_{z, \overline{K}})$ is the Fontaine-Messing pairing, we obtain $(2i\pi)_p y(z)$ as a Fontaine-Messing period of \mathcal{A}_z . More generally, $y(z) \text{Log}_p az^n + (2i\pi)_p y^*(z)$ appears as a period (Log instead of log). We conclude that in this degenerating case, the Fontaine-Messing periods behave relatively well with respect to the Gauss-Manin connection. A similar observation applies to the pairing composed with θ (Hodge-Tate periods).

5.3. p -adic Betti lattices for abelian varieties with supersingular reduction.

5.3.1. Any supersingular abelian variety over $\overline{\mathbb{F}}_p$ is isogenous to a power of a supersingular elliptic curve. We refer to [Wa69, 4] for a detailed study of supersingular elliptic curves over finite fields (including the classification of isogeny classes and isomorphism classes, refining Deuring's classical work), and to [LiO98] for the higher dimensional case. Let us simply recall a few basic facts:

- there is exactly one $\overline{\mathbb{F}}_p$ -isogeny class of supersingular elliptic curves.
- If A_0 is any supersingular elliptic curve over \mathbb{F}_{p^n} , then $\mathbf{D} := \text{End}(A_0)_{\overline{\mathbb{F}}_p}$ is a maximal order in a quaternion algebra over \mathbb{Q} , ramified exactly at p, ∞ ; we set $\mathbf{D} = \mathbf{D}_{\mathbb{Q}}$.
- A_0 is $\overline{\mathbb{F}}_p$ -isomorphic (but not necessarily \mathbb{F}_{p^n} -isomorphic) to an elliptic curve defined over \mathbb{F}_{p^2} .
- If $n = 1$, $\text{End}(A_0) \otimes \mathbb{Q} = \mathbb{Q}[Fr_p]$, an imaginary quadratic field; if $p \neq 2$, this is $\mathbb{Q}[\sqrt{-p}]$. More precisely, the order $\text{End}(A_0)$ can be either $\mathbb{Z}[Fr_p]$ or the maximal order in $\text{End}(A_0) \otimes \mathbb{Q}$ (which coincide if $p \equiv 3 \pmod{4}$). If $p = 2$, there is another possibility, namely $\text{End}(A_0) = \mathbb{Z}[Fr_2] \simeq \mathbb{Z}[\sqrt{-1}]$.
- There exists a supersingular elliptic curve over \mathbb{F}_p whose Frobenius endomorphism satisfies $(Fr_p)^2 = -p \cdot \text{id}$ and whose endomorphism algebra is the maximal order in $\mathbb{Q}[Fr_p]$. Such elliptic curves belong to a single \mathbb{F}_p -isogeny class.
- If $p \neq 2, 3$, there is exactly one \mathbb{F}_p -isogeny class of supersingular elliptic curves over \mathbb{F}_p (but several $\overline{\mathbb{F}}_p$ -isomorphism classes in general).

5.3.2. Let A_0 be a supersingular abelian variety of dimension g over \mathbb{F}_{p^n} . By functoriality, the elements of $\text{End}((A_0)_{\overline{\mathbb{F}}_p}) \otimes \mathbb{Q} \simeq M_g(\mathbf{D})$ act linearly on the crystalline cohomology $H_{\text{crys}}^*((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p) = H_{\text{crys}}^*((A_0)_{\overline{\mathbb{F}}_p}/\widehat{\mathbb{Z}}_p^{\text{ur}}) \otimes_{\widehat{\mathbb{Z}}_p^{\text{ur}}} \overline{\mathbb{Q}}_p$. In degree one, this provides an embedding of $M_g(\mathbf{D})$ into the endomorphism ring of $H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$.

On the other hand, one has a canonical isomorphism

$$\bigwedge^2 H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p) \simeq H_{\text{crys}}^2((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$$

and a canonical \mathbb{Q} -structure in $H_{\text{crys}}^2((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p) \simeq \overline{\mathbb{Q}}_p^{g(2g-1)}$ coming from the fact that the whole cohomology in degree 2 is generated by algebraic cycles (since $(A_0)_{\overline{\mathbb{F}}_p}$ is isogenous to the g th power of a supersingular elliptic curve, one reduces easily to the case $g = 2$, in which case this fact is well-known).

Let F be either $\overline{\mathbb{Q}}$ or a splitting number field for \mathbf{D} : $\mathbf{D} \otimes_{\mathbb{Q}} F \simeq M_2(F)$. In the latter case, this amounts to saying that F is totally imaginary, and for any place v above p , $[F_v : \mathbb{Q}_p]$ is even. We fix an embedding of F in $\overline{\mathbb{Q}}_p$.

Via this embedding, the $M_g(\mathbf{D})$ -action extends to a F -linear $M_{2g}(F)$ -action on $H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$.

We consider an irreducible cyclic sub- $M_{2g}(F)$ -module $M_{2g}(F) \cdot u \subset H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$. Obviously, such submodules exist, and have F -dimension $2g$. We may and shall choose u in such a way that its exterior square of this F -space coincides with the canonical F -structure on $H_{\text{crys}}^2((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$ (this can be achieved by replacing u by a suitable multiple).

Proposition 5.3.3. *Up to a homothety by a factor in $\sqrt{F^\times}$, the normalized $M_{2g}(F)$ -submodule $M_{2g}(F) \cdot u \subset H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$ depends only on $F \subset \overline{\mathbb{Q}}_p$, not on the choice of u . In particular, for $F = \overline{\mathbb{Q}}$, this defines a canonical $\overline{\mathbb{Q}}$ -structure in $H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$, stable under $\text{End}((A_0)_{\overline{\mathbb{F}}_p})$.*

PROOF. Indeed, two such $M_{2g}(F)$ -submodules are related by some $h \in GL_{2g}(\overline{\mathbb{Q}}_p)$, such that $\wedge^2(h) \in GL_{g(2g-1)}(F)$. Now $M_{2g}(F) \cdot hu = h(M_{2g}(F) \cdot u)$ implies that h normalizes $GL_{2g}(F)$. It follows that the image of h in $PGL_{2g}(\overline{\mathbb{Q}}_p)$ lies in $PGL_{2g}(F)$.

(When $g > 1$, the proposition is not surprising, since $\wedge^2 V$ is a faithful representation of $PSL(V)$ for any space V of dimension > 1). \square

Remarks 5.3.4. (a) A minimal choice for F is a splitting quadratic field $\mathbb{Q}(\sqrt{-d})$ (the splitting property amounts to saying that $d > 0$ and p ramifies or remains prime in $\mathbb{Q}(\sqrt{-d})$. A natural choice is $d = p$).
 (b) Let \mathcal{O}_F be the ring of integers of F . Then $\text{End}((A_0)_{\overline{\mathbb{F}}_p}) \otimes \mathcal{O}_F$ is an order in $M_{2g}(F)$, and there is a full $(\text{End}((A_0)_{\overline{\mathbb{F}}_p}) \otimes \mathcal{O}_F)$ -lattice in $(M_{2g}(F) \cdot u)$. As an \mathcal{O}_F -module, it is projective of rank $2g$. If $F = \overline{\mathbb{Q}}$, i.e. $\mathcal{O}_F = \overline{\mathbb{Z}}$, then this module is *canonically determined* by the normalization of its top exterior power, taking into account the canonical isomorphism

$$\bigwedge^{2g} H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Z}}_p) \simeq H_{\text{crys}}^{2g}((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Z}}_p) \simeq \overline{\mathbb{Z}}_p,$$

the latter isomorphism coming from the trace map.

Hence there is a *canonical $\overline{\mathbb{Z}}$ -structure* in the crystalline cohomology of supersingular abelian varieties (not used in the sequel).

5.3.5. Let A be an abelian variety defined over a subfield $k \subset \mathbb{C}_p$, which has good *supersingular reduction* A_0 over the residue field of k . There is a canonical embedding $\text{End}(A) \hookrightarrow \text{End}((A_0)_{\overline{\mathbb{F}}_p})$.

The canonical $\overline{\mathbb{Q}}$ -subspace of $H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$ defined in 5.3.3 will be denoted by

$$H_{\text{B}}^1(A_{\mathbb{C}_p}, \overline{\mathbb{Q}}) \subset H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}}_p}/\overline{\mathbb{Q}}_p)$$

because, as we shall see, it shares many properties with the Betti space $H_{\text{B}}^1(A_{\mathbb{C}}, \overline{\mathbb{Q}}) = H_{\text{B}}^1(A_{\mathbb{C}}, \mathbb{Z}) \otimes \overline{\mathbb{Q}}$ of a complex abelian variety $A_{\mathbb{C}}$. Its dual will be denoted by $H_{1,\text{B}}(A_{\mathbb{C}_p}, \overline{\mathbb{Q}})$.

Similarly, we shall write $H_B^1(A_{\mathbb{C}_p}, F)$ for the F -subspace described 5.3.3 (well-defined up to a homothety in $\sqrt{F^\times}$), and $H_{1,B}(A_{\mathbb{C}_p}, F)$ for its F -dual.

Assume that $k \subset \overline{\mathbb{Q}_p}$. There is a functorial isomorphism [BeO83]

$$H_{\text{dR}}^1(A) \otimes_k \overline{\mathbb{Q}_p} \simeq H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}_p}}/\overline{\mathbb{Q}_p}),$$

from which one derives a canonical isomorphism

$$H_{\text{dR}}^1(A) \otimes_k \overline{\mathbb{Q}_p} \simeq H_B^1(A_{\mathbb{C}_p}, \overline{\mathbb{Q}}) \otimes_{\overline{\mathbb{Q}}} \overline{\mathbb{Q}_p},$$

which is functorial with respect to homomorphisms of abelian varieties with supersingular reduction

This isomorphism, which is far from being tautological, as we shall see, can be translated into a $\overline{\mathbb{Q}_p}$ -valued pairing between $H_{1,B}(A_{\mathbb{C}_p}, F)$ and $H_{\text{dR}}^1(A)$. This pairing is conveniently expressed in the form of a “*period matrix*” $\Omega = (\omega_{ij})$ with entries in $\overline{\mathbb{Q}_p}$, depending on the choice of a basis $(\gamma_i)_{i=1, \dots, 2g}$ of $H_{1,B}(A_{\mathbb{C}_p}, F)$ and a basis $(\omega_j)_{j=1, \dots, 2g}$ of $H_{\text{dR}}^1(A)$ (if A is principally polarized, we choose symplectic bases).

Warning: these p -adic periods attached to abelian varieties with supersingular reduction have little to do with Fontaine-Messing periods (a motivic interpretation of these periods is proposed in [And95]).

5.3.6. Horizontality. If we let A move in a family A_z , $H_{\text{crys}}^1((A_0)_{\overline{\mathbb{F}_p}}/\overline{\mathbb{Q}_p})$ may be identified with a space of horizontal sections of the de Rham cohomology localized in the disk parameterizing the liftings A_z of $(A_0)_{\overline{\mathbb{F}_p}}$ [BeO83]. Thus the period matrix $\Omega(z)$ is a fundamental matrix of solutions of the Gauss-Manin connection.

In the sequel of this subsection, we assume for simplicity that $g = 1$, *i.e.* A is an elliptic curve.

5.3.7. CM periods, Γ_p -values, and transcendence.

- Let A_0 be as before a supersingular elliptic curve over \mathbb{F}_p^n . Let A be a CM-lifting of A_0 , *i.e.* an elliptic curve defined over some number field $k \subset \overline{\mathbb{Q}} \subset \overline{\mathbb{Q}_p}$ with residue field \mathbb{F}_p^n , such that $E = \text{End}(A) \otimes \mathbb{Q}$ is quadratic. Note that E is ipso facto a subfield of \mathcal{D} . We denote by c the conductor of the order $\text{End}(A)$ (in particular, $ce \in \text{End}(A)$).

- Let (γ_1, γ_2) be any symplectic basis of $H_{1,B}(A, F)$. We can write $ce \cdot \gamma_1 = a\gamma_1 + b\gamma_2$ with $a, b \in F$, so that

$$\tau := \frac{\omega_{12}}{\omega_{11}} = \frac{-a + c\sqrt{-d}}{b} \in \mathbb{P}^1(FE).$$

In particular, $\tau \in \mathbb{P}^1(\overline{\mathbb{Q}})$. We conjecture the following converse, which is a p -adic analogue of T. Schneider’s theorem [Schn37] (in the complex case):

Conjecture 5.3.8. *Assume that A is a lifting of A_0 defined over $\overline{\mathbb{Q}}$, and that $\tau = \frac{\omega_{12}}{\omega_{11}} \in \mathbb{P}^1(\overline{\mathbb{Q}})$. Then A has complex multiplication.*

• Let $E = \mathbb{Q}[\sqrt{-d}]$ and $E' = \mathbb{Q}[\sqrt{-d'}]$ be two distinct subfields of \mathbf{D} (d, d' squarefree; we do not exclude the case $d = d'$). According to [Vig80, II 1.4], $E \cap \mathbf{D} = \mathcal{O}_E$, $E' \cap \mathbf{D} = \mathcal{O}_{E'}$. The elements $e = \sqrt{-d} \in E$ and $e' = \sqrt{-d'} \in E'$ generate the algebra \mathbf{D} . We remark that $ee' + e'e$ commutes to e and e' , hence is an integer $m \in \mathbb{Z}$ (an even integer if d or $d' \equiv 3 \pmod{4}$, which is divisible by 4 if both $d, d' \equiv 3 \pmod{4}$). We have $(ee' - e'e)^2 = m^2 - 4dd'$. Since $ee' - e'e$ anticommutes with e , it is not an integer, hence $|m| < 2\sqrt{dd'}$. On the other hand, the images of e and e' in $\mathbf{D} \otimes \mathbb{F}_p$ commute ($ee' - e'e$ acts trivially on regular differentials in characteristic p), therefore p divides $4dd' - m^2$.

• On the other hand, there exists a unique symplectic basis of eigenvectors for e in $H_{\text{dR}}^1(A)$ of the form $(\omega_1, \omega_2 + \sigma\omega_1)$, with $\sigma \in k$. We get the relation

$$\omega_{22} + \sigma\omega_{12} = \tilde{\tau}(\omega_{21} + \sigma\omega_{11}), \text{ with } \tilde{\tau} = \frac{-a - c\sqrt{-d}}{b}.$$

Let now $E' = \mathbb{Q}[\sqrt{-d'}] \neq E$ be another subfield of \mathbf{D} , and assume that (γ_1, γ_2) is a basis of eigenvectors for E' : $e'\gamma_1 = \sqrt{-d'}\gamma_1$. We get $(ee' + e'e)\gamma_1 = \frac{2a\sqrt{-d'}}{c}\gamma_1 = m\gamma_1$, so that

$$\frac{\tilde{\tau}}{\tau} = \frac{m + 2\sqrt{dd'}}{m - 2\sqrt{dd'}} \in \mathbb{Q}[\sqrt{dd'}]$$

the sign of the square root being chosen in such a way that $p \mid 2\sqrt{dd'} - m$ (in the unramified case).

• The previous relations, together with $\det \Omega = 1$, show that the transcendence degree over $\overline{\mathbb{Q}}$ of the entries of Ω is at most one in the CM case.

• We now describe the p -adic number ω_{11} modulo $\overline{\mathbb{Q}}^\times$, keeping the notation of 4.6.2 and Theorem 4.6.3 (ϵ is the Dirichlet character, w the number of roots of unity, h the class number).

Theorem 5.3.9. *In case p does not ramify in the field of complex multiplications $\mathbb{Q}[\sqrt{-d}] = \text{End}(A) \otimes \mathbb{Q}$, one has*

$$\omega_{11} \sim \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/8h} \text{ in } \overline{\mathbb{Q}}_p^\times / \overline{\mathbb{Q}}^\times.$$

Examples. If A is the Legendre curve $X_{1/2}$ ($y^2 = x(x-1)(x-1/2)$) and $p \equiv 3 \pmod{4}$, we find $\omega_{11} \sim \Gamma_p(1/4)$. Similarly, if $\text{End}(A) \otimes \mathbb{Q} = \mathbb{Q}(\sqrt{-3})$ and $p \equiv 2 \pmod{3}$, then $\omega_{11} \sim \Gamma_p(1/3)^{3/2}$.

PROOF. We may assume that the number field k ($\subset \overline{\mathbb{Q}}_p$) is Galois over \mathbb{Q} . We denote by $k' = k \cap \mathbb{Q}_p$ the subfield of k fixed by the local Galois group $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$. Let us consider the Weil restriction $B := \mathcal{R}_{k/k'}(A)$ (cf. [BLR90, 7.6]). This is an abelian variety over $k' \subset \mathbb{Q}_p$, with good, supersingular reduction at p (like A), and we have $B_k \simeq \prod_{\tau \in \text{Gal}(k/k')} A^\tau$.

On the other hand, let ψ be any element of degree one in the Weil group (*i.e.* any lifting of the Frobenius element in $\text{Gal}(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$), and let Ψ_{A^τ} (*resp.* Ψ_B) be the corresponding semilinear endomorphism of $H_{\text{dR}}^1(A^\tau/k) \otimes \overline{\mathbb{Q}}_p$ (*resp.* $H_{\text{dR}}^1(B/k') \otimes \overline{\mathbb{Q}}_p$). If Φ_B denotes the canonical linear action induced by $\text{Fr}_{\mathbb{F}_p}$ on $H_{\text{crys}}^1(B_0/\mathbb{Q}_p) \otimes \overline{\mathbb{Q}}_p \simeq H_{\text{dR}}^1(B)_{\overline{\mathbb{Q}}_p}$, one has the formula $\Psi_B = \Phi_B \circ (\text{id} \otimes \psi)$ (*cf.* [BeO83, 4]). Moreover, $\Psi_B = \bigoplus \Psi_{A^\tau}$ with respect to the decomposition $H_{\text{dR}}^1(B, \overline{\mathbb{Q}}_p) \simeq \bigoplus_{\tau} H_{\text{dR}}^1(A^\tau, \overline{\mathbb{Q}}_p)$.

If (ω_1, ω_2) is any symplectic basis of eigenvectors for $\mathbb{Q}[\sqrt{-d}]$ in $H_{\text{dR}}^1(A)_{\overline{\mathbb{Q}}}$, with $\omega_1 \in \Omega^1(A)_{\overline{\mathbb{Q}}}$, we can write $\Psi_A(\omega_1) = \kappa \cdot \omega_2$, where κ modulo $\overline{\mathbb{Q}}^\times$ is given by Ogus' formula in Theorem 4.6.3

$$\kappa \sim \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle p \frac{u}{d} \right\rangle \right)^{-\epsilon(u)w/4h} \text{ in } \overline{\mathbb{Q}}_p^\times / \overline{\mathbb{Q}}^\times.$$

Because the conjugates A^τ are isogenous to each other over k (and since isogenies preserve the regular differential forms), we derive that the constant κ , *resp.* the ‘‘period’’ ω_{11} , is the same modulo $\overline{\mathbb{Q}}^\times$ for each of them. Therefore, for any $\omega \in \Omega^1(B) \subset H_{\text{crys}}^1(B_0/\mathbb{Q}_p)$, we can write $\Phi_B(\omega) = \Psi_B(\omega) = \kappa \cdot \eta$ for some $\eta \in H_{\text{dR}}^1(B)_{\overline{\mathbb{Q}}}$ whose pairing with any element of $H_{1B}(B, \overline{\mathbb{Q}})$ belongs to $\omega_{21} \cdot \overline{\mathbb{Q}}$. Since $\omega_{11}^{-1} \omega \in H_{1B}^1(B, \overline{\mathbb{Q}})$, $\omega_{21}^{-1} \eta \in H_{1B}^1(B, \overline{\mathbb{Q}})$, and since Φ_B respects $H_{1B}(B, \overline{\mathbb{Q}})$, we get $\kappa \sim \frac{\omega_{11}}{\omega_{21}} \sim \omega_{11}^2$, whence the result. \square

Conjecture 5.3.10. *Assume that A is a lifting of A_0 defined over $\overline{\mathbb{Q}}$, and that p does not ramify $\text{End}(A) \otimes \mathbb{Q}$ (which is obviously the case if A does not have complex multiplication). Then $H_{\text{dR}}^1(B)_{\overline{\mathbb{Q}}} \neq H_{1B}^1(B, \overline{\mathbb{Q}})$.*

In the absence of complex multiplication, this follows from the previous conjecture. In the presence of complex multiplication, it amounts to the transcendence of ω_{11} .

Taking into account 5.3.9, the conjecture would imply, for instance, the transcendence of the adic numbers $\Gamma_3(1/4)$, $\Gamma_7(1/4)$, $\Gamma_2(1/3)$, $\Gamma_5(1/3)$ (note, in contrast, that $\Gamma_5(1/4)$ and $\Gamma_7(1/3)$ are algebraic numbers, according to the Gross-Koblitz formula (2.6.4)).

- Actually, it may happen, in the ramified CM case, that $H_{\text{dR}}^1(A)_{\overline{\mathbb{Q}}} = H_{1B}^1(A, \overline{\mathbb{Q}})$.

Let us first notice that the argument given in the proof of theorem 5.3.9 still works in the ramified case, and allows to conclude that $\kappa \sim \omega_{11}^2$. The computation of $\kappa \pmod{\overline{\mathbb{Q}}^\times}$ (for p odd) according to Ogus-Coleman-Urfels was explained in 4.6.4. In the example of an elliptic curve with complex multiplication by $\mathbb{Q}[\sqrt{-3}]$, we have seen that $\kappa \in \overline{\mathbb{Q}}$, hence $\omega_{11} \in \overline{\mathbb{Q}}$ and $H_{\text{dR}}^1(A)_{\overline{\mathbb{Q}}} = H_{1B}^1(A, \overline{\mathbb{Q}})$.

This is also the case for an elliptic curve with complex multiplication by $\mathbb{Q}[\sqrt{-3n}]$ if the Legendre symbol $(\frac{n}{3})$ is 1, but probably not in general if $(\frac{n}{3}) = -1$ (although this happens for $n = 8$, after the last example in 4.6.4).

5.3.11. $L_{\frac{1}{2}, \frac{1}{2}, 1}$ in a supersingular disk. For concreteness, we consider the special case of the supersingular Legendre elliptic curve with parameter $z = 1/2$ in characteristic $p \equiv 3 \pmod{4}$. A basis of solutions of the hypergeometric differential operator $L_{\frac{1}{2}, \frac{1}{2}, 1} = HGDO(\frac{1}{2}, \frac{1}{2}, 1)$ in the supersingular disk $D(1/2, 1^-)$ is given by

$$F\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{2}; (1-2z)^2\right), \quad (1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right).$$

- Let us consider the symplectic basis $\omega_1 = [\frac{dx}{2y}], \omega_2 = [\frac{(2x-1)dx}{4y}]$ of \mathcal{M} (de Rham cohomology). At $z = 1/2$, this is a basis of eigenvectors for the action of $E' := \text{End}(X_{1/2}) \otimes \mathbb{Q} = \mathbb{Q}(\sqrt{-1})$. The Gauss-Manin connection satisfies

$$\omega_2 = 2z(z-1)\nabla\left(\frac{d}{dz}\right)\omega_1 + \frac{4z-5}{6}\omega_1.$$

The fundamental solution matrix Y of the Gauss-Manin connection (expressed in the basis (ω_1, ω_2)), normalized by $Y(1/2) = \text{id}$, is then given by

$$\begin{aligned} y_{11} &= F\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{2}; (1-2z)^2\right) + \frac{1}{2}(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right), \\ y_{12} &= (1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right), \\ y_{2i} &= 2z(z-1)\frac{dy_{1i}}{dz} + \frac{4z-5}{6}y_{1i}, \quad i = 1, 2. \end{aligned}$$

- We choose a symplectic basis (γ_1, γ_2) of eigenvectors for E' in $H_{1B}^1(X_{1/2}, F)$ (note that $\mathbb{Q}(\sqrt{-1})$ is always contained in F). We have $H_{1B}^1(X_{1/2}, F) = H_{1B}^1(X_z, F)$ for any point z in the supersingular disk $D(1/2, 1^-)$. Because the period matrix $\Omega(z)$ is another solution matrix of the Gauss-Manin connection, we have the relation $Y(z) = \Omega(z).\Omega(1/2)^{-1}$.

Let us consider a CM-point ζ in the supersingular disk $D(1/2, 1^-)$: $\text{End}(X_\zeta) = E = \mathbb{Q}(\sqrt{-d})$ (and $E \neq E'$ in \mathbf{D}).




FIGURE 10

Let σ and m be as before (in the present case, the integer m is even: $m = 2n$). Our period relations can be easily expressed in terms of $Y(\zeta)$. Apart from the obvious relation $\det Y(\zeta) = 1$, we find a relation

$$\left. \frac{y_{11}(y_{22} + \sigma y_{12})}{y_{12}(y_{21} + \sigma y_{11})} \right|_{z=\zeta} = \frac{\tilde{\tau}}{\tau} = \frac{n + \sqrt{d}}{n - \sqrt{d}}.$$

Changing the viewpoint, one can consider ζ as fixed, and vary the prime $p \neq 2$ (or more precisely the place of $\mathbb{Q}(\zeta, \sqrt{-1}, \sqrt{-d})$). This relation between values of p -adic hypergeometric functions at ζ holds whenever $|\zeta - 1/2|_p < 1$; it depends on p via the integer n ($|n| < \sqrt{d}$, $p \mid d - n^2$). A relation of the same kind holds at the places at infinity (derived along similar lines, using the usual Betti lattices). As a specific example, we can take $p = 3$, $\zeta =$ a primitive 6th root of unity. Then X_ζ has complex multiplication by $\mathbb{Z}[\zeta]$, $\sigma = \frac{-\zeta^2}{2(1+\zeta)}$, $n = 0$ and $\frac{\tilde{\tau}}{\tau} = -1$.

- On the other hand, one can combine Theorem 5.3.9 and the formula $Y(\zeta) = \Omega(\zeta) \cdot \Omega(1/2)^{-1}$ in order to express the value mod $\overline{\mathbb{Q}}^\times$ of the p -adic hypergeometric functions y_{ij} in terms of Γ_p (*supersingular avatar of Young's formulas*).

For instance, if $p = 7$, $\zeta = 2(\sqrt{2} - 1)$ (complex multiplication by $\mathbb{Z}[\sqrt{-2}]$), one has $\omega_{11}(\zeta) \sim (\Gamma_7(1/8)\Gamma_7(3/8))^{1/2}$, $\omega_{11}(1/2) \sim \Gamma_7(1/4)$, from which one derives the 7-adic evaluation

$$F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; 5 - 4\sqrt{2}\right) \sim (\Gamma_7(1/8)\Gamma_7(3/8))^{1/2} \Gamma_7(1/4)^{-1}.$$

5.4. Conclusion and vista.

We have seen many instances of the following scenario: some familiar notion or phenomenon from the complex realm shows itself in two different aspects; each of these aspects has a natural p -adic counterpart; these counterparts are not complementary aspects of the same p -adic entity, but belong to totally different theories.

In II, we shall see that the theory of p -adic period mappings is a privileged field where this semantic splitting process integrates into an harmonious picture, where the complementarity between the p -adic counterparts is restored at a deeper level.

The p -adic period mappings which we shall deal with relate some deformation spaces of p -divisible groups to certain grassmannians. It will turn out that they can be described by quotients of analytic solutions of the Gauss-Manin connection, just as in the complex case. In particular, at the *modular level*, the Fontaine-Messing periods (which links up directly the étale and crystalline representations of p -divisible groups) go offstage or remain at the background.

⁽⁸⁾F. Beukers has also found these relations independently, by a different method.

On the other hand, we shall see that the differential equations studied in section 1 (*resp.* 2) arise in the context of deformations of supersingular (*resp.* ordinary) p -divisible groups.

CHAPTER II

Introduction to the theory of p -adic period mappings.

Contents

1. A survey of moduli and period mappings over \mathbb{C} .	66
2. Preliminaries on p -divisible groups.	72
3. A stroll in the crystalline world.	75
4. Moduli problems for p -divisible groups.	83
5. p -adic period domains.	87
6. The p -adic period mapping and the Gauss-Manin connection.	90
7. p -adic uniformization of Shimura varieties.	96

1. A survey of moduli and period mappings over \mathbb{C} .

ABSTRACT: Review of moduli, Shimura varieties, Hodge structures, period mappings, period domains, Gauss-Manin connections.

1.1. Moduli spaces.

1.1.1. The term “moduli” was coined by Riemann to describe the continuous essential parameters of smooth complex compact curves of a given genus g . The use of this term has been extended to many classification problems in analytic or algebraic geometry: typically, a classification problem consists of a discrete part (the dimension, and other numerical invariants...) and a continuous part (the moduli).

One can distinguish between local and global moduli spaces:

- (i) local moduli are the parameters of the most general small (or infinitesimal) deformation of a given analytic manifold or algebraic variety. Let for instance X be a compact analytic manifold. Then a famous theorem of Kuranishi asserts that there is a local deformation $\pi : \underline{X} \rightarrow S$ with base point s ($X = \underline{X}_s$) such that any other deformation comes, locally around s , from π by base change; moreover the tangent space of S at s is isomorphic to $H^1(X, (\Omega_X^1)^\vee)$. For curves, the dimension of this space is $3g-3$ if $g \geq 2$ (the number of Riemann’s complex moduli).
- (ii) Global moduli spaces, when they exist, provide the solution of the continuous part of moduli problems. For instance, Douady has constructed the global moduli space for closed analytic subspaces of a given compact complex manifold. Although global moduli spaces were already studied in the nineteenth century (in the case of curves), the concept was first put on firm foundations by Grothendieck in terms of representable functors.

The existence of a global moduli space is often obstructed by the presence of automorphisms of the objects under classification. One possibility is to rule out these automorphisms by imposing some extra structure (rigidification). When all automorphism groups are finite, another possibility is to look for moduli orbifolds rather than moduli spaces

1.1.2. The most classical moduli problems concern complex elliptic curves. These are classified by their j -invariant. However the affine j -line \mathbb{A}^1 is not a fine moduli space: indeed, in any family of elliptic curves parametrized by the whole affine line, all the fibers are isomorphic.

One can rigidify the problem by introducing a “level N structure” on elliptic curves E , *i.e.* fixing an identification $\text{Ker}[N]_A \cong (\mathbb{Z}/N\mathbb{Z})^2$ such that the Weil pairing is given by $(\xi, \eta) \mapsto \exp(2\pi i(\xi \wedge \eta))$ (here $[N]_A$ denotes the multiplication by N on the elliptic curve A).

For $N \geq 3$, this rules out automorphisms of A , and the corresponding moduli functor is representable. For $N = 3$, the moduli space is $\mathbb{A}^1 \setminus \{\mu^3 =$

1}); the universal elliptic curve with level 3 structure is the Dixon elliptic curve

$$x^3 + y^3 + 3\mu xy = 1 \text{ (in affine coordinates } x, y)$$

with $\text{Ker}[3]_A =$ the set of flexes.

For $N = 2$, the automorphism $[-1]_A$ remains; there is an elliptic curve with level 2 structure over $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, namely the Legendre elliptic curve

$$y^2 = x(x-1)(x-\lambda) \text{ (in affine coordinates } x, y)$$

with $\text{Ker}[2]_A = \{(0, 0), (1, 0), (\lambda, 0), (\infty, \infty)\}$, but it is not universal: another one is given by $\lambda y^2 = x(x-1)(x-\lambda)$.

1.1.3. These moduli problems admit two natural generalizations: moduli problems for curves of higher genus, and moduli problems for polarized abelian varieties (polarizations ensure that the automorphism groups are finite).

There is a moduli orbifold for smooth compact complex curves of genus $g > 0$, denoted by \mathcal{M}_g , which can be described as follows. The set of complex structures on compact connected oriented surface S_g of genus g is in a natural way a contractible complex manifold of dimension $3g - 3$ (*resp.* the Poincaré upper half plane \mathfrak{h} if $g = 1$), the Teichmüller space \mathcal{T}_g . The mapping class group Γ_g , *i.e.* the component of connected components of the group of orientation-preserving self-homeomorphisms of S_g , acts on right on \mathcal{T}_g , and \mathcal{M}_g is the quotient \mathcal{T}_g/Γ_g . For $g = 1$, one recovers $\mathcal{M}_1 = \mathfrak{h}/SL(2, \mathbb{Z})$.

There is a moduli orbifold for principally polarized abelian varieties of dimension $g > 0$, denoted by \mathcal{A}_g , which can be described as follows. For any abelian variety A , one has an exact sequence

$$0 \rightarrow L \rightarrow \text{Lie } A \rightarrow A \rightarrow 0$$

and a principal polarization corresponds to a positive definite hermitian form H on $\text{Lie } A$ whose imaginary part induces a perfect alternate form on L . Choosing a \mathbb{C} -basis $\omega_1, \dots, \omega_g$ of $\Omega^1(A) \cong (\text{Lie } A)^\vee$ and a symplectic \mathbb{Z} -basis $\gamma_1^+, \gamma_1^-, \dots, \gamma_g^+, \gamma_g^-$ of L gives rise to a period matrices $\Omega^+ = (\int_{\gamma_j^+} \omega_i)$, $\Omega^- = (\int_{\gamma_j^-} \omega_i)$, and $\tau = (\Omega^-)^{-1}\Omega^+$ does not depend on the ω_i 's. By the properties of H , τ belongs to the Siegel upper half space \mathfrak{h}_g of symmetric matrices with positive definite imaginary part (Riemann relations); it is well-defined up to right multiplication by a matrix in $Sp(2g, \mathbb{Z})$ (which is the same as the standard left action on \mathfrak{h}_g of the transposed matrix). This construction identifies \mathcal{A}_g with $\mathfrak{h}_g/Sp(2g, \mathbb{Z})$.

The construction which associates to a curve its jacobian variety with the principal polarization given by the theta divisor gives rise to an immersion $\mathcal{M}_g \hookrightarrow \mathcal{A}_g$, the Torelli map.

1.1.4. Following G. Shimura, one also studies refined moduli problems for “decorated” (principally polarized) abelian varieties, by prescribing in addition that the endomorphism algebra contains a given simple \mathbb{Q} -algebra,

and imposing a level N structure. This gives rise to moduli orbifolds (*resp.* moduli spaces if $N \geq 3$) of “PEL type”.

More precisely, let B be a simple finite-dimensional \mathbb{Q} -algebra, with a positive involution $*$. Let V be a B -module of finite type, endowed with an alternate \mathbb{Q} -bilinear form such that $\langle bv, w \rangle = \langle v, b^*w \rangle$. One denotes by G the algebraic \mathbb{Q} -group of B -linear symplectic similitudes of V .

Let \mathcal{B} be a maximal order in B stable under $*$, and let L be a lattice in V , stable under \mathcal{B} and autodual for $\langle \cdot, \cdot \rangle$ (this will represent the structure of $H^1(A, \mathbb{Z})$ for the abelian varieties A under consideration). We choose in addition a lagrangian subspace $F_0^1 \subset V_{\mathbb{C}}$ stable under B (which will be $\Omega^1(A_0) \cong (\text{Lie } A_0)^\vee$ for one particular abelian variety A_0 under consideration).

There is a moduli orbifold (*resp.* moduli space if $N \geq 3$) for principally $*$ -polarized⁽¹⁾ abelian varieties A , together with a given action $\iota : \mathcal{B} \rightarrow \text{End } A$ such that $\det(\iota(b)|\Omega^1(A)) = \det(b|F_0^1)$ for all $b \in \mathcal{B}$ (Shimura type condition), and a given $\mathcal{B}/N\mathcal{B}$ -linear symplectic similitude $A[N] = \text{Ker}[N]_A \rightarrow \text{Hom}(L, \mathbb{Z}/N\mathbb{Z})$.

It amounts to the same to consider abelian varieties A up to isogeny, with B -action (with Shimura type condition), together with a \mathbb{Q} -homogeneous principal $*$ -polarization and a class of B -linear symplectic similitudes $H^1(A, \hat{\mathbb{Z}} \otimes \mathbb{Q}) \rightarrow \hat{\mathbb{Z}} \otimes V$ modulo the group $\{g \in G(\hat{\mathbb{Z}} \otimes \mathbb{Q}) \mid (g-1)(\hat{\mathbb{Z}} \otimes L) \subset \hat{\mathbb{Z}} \otimes L\}$.

These moduli orbifolds are called Shimura orbifolds of “PEL type”. They are algebraic, and defined over the number field $E = \mathbb{Q}[\text{tr}(b|F_0^1)]$ (the so-called reflex field). Their connected components are defined over finite abelian extensions of E , *cf.* [Del71].

1.2. Period mappings.

1.2.1. In the case of curves, the construction of period mappings belongs to the nineteenth century. The idea is to study the deformations of a curve by looking at the variation of its periods. Let $f : \underline{X} \rightarrow S$ be an algebraic family of projective smooth (connected) curves of genus g over \mathbb{C} . Let \tilde{S} be the universal covering of S . Then the homology groups $H_1(-, \mathbb{Z})$ of the fibers form a constant local system on \tilde{S} . Let us choose a symplectic basis $\gamma_1^+, \gamma_1^-, \dots, \gamma_g^+, \gamma_g^-$. Choosing, locally, an auxiliary \mathbb{C} -basis $\omega_1, \dots, \omega_g$ of $f_*\Omega_{\underline{X}/S}^1$ gives rise as above to a period matrices $\Omega^+ = (\int_{\gamma_j^+} \omega_i)$, $\Omega^- = (\int_{\gamma_j^-} \omega_i)$, and $\mathcal{P} = (\Omega^-)^{-1}\Omega^+$ does not depend on the ω_i 's and is well-defined on \tilde{S} .

We get a commutative square

⁽¹⁾a principal $*$ -polarization is a symmetric \mathcal{B} -linear isomorphism between the abelian variety and its dual, endowed with the transposed action of \mathcal{B} twisted by $*$

$$\begin{array}{ccc}
\tilde{S} & \xrightarrow{\mathcal{P}} & \mathfrak{h}_g \\
\downarrow & & \downarrow \\
S & \xrightarrow{\mathcal{P}} & \mathfrak{h}_g/Sp(2g, \mathbb{Z})
\end{array}$$

Applying this construction to the universal case $S = \mathcal{M}_g$ ($\mathcal{T}_g = \tilde{\mathcal{M}}_g$ in the orbifold sense), the bottom line of this commutative square is nothing but an analytic interpretation of the Torelli map.

1.2.2. An important feature of the period mapping \mathcal{P} is that it is a *quotient of solutions of a fuchsian differential system*, with rational exponents at infinity. This comes out as follows. The vector bundle $f_*\Omega^1_{\underline{X}/S}$ is a subbundle (of rank g) of a vector bundle \mathcal{H} (of rank $2g$) whose fibers are the De Rham cohomology spaces $H^1_{\text{DR}}(\underline{X}_s)$. This bundle \mathcal{H} carries an integrable connection, the Gauss-Manin connection

$$\nabla_{\text{GM}} : \mathcal{H} \rightarrow \Omega^1_S \otimes \mathcal{H}.$$

The local system of germs of horizontal analytic sections is $R^1 f_* \mathbb{C}$ (with fibers $H^1(\underline{X}_s, \mathbb{C})$). Completing, locally, the basis $\omega_1, \dots, \omega_g$ of sections of $f_*\Omega^1_{\underline{X}/S}$ into a basis $\omega_1, \dots, \omega_g, \eta_1, \dots, \eta_g$ of sections \mathcal{H} allows to write the connection in form of a differential system, and a full *solution matrix* is given by the “full $2g \times 2g$ -period matrix” $\begin{pmatrix} \Omega^+ & \Omega^- \\ N^+ & N^- \end{pmatrix}$, while $\mathcal{P} = (\Omega^-)^{-1}\Omega^+$.

In the case of the Legendre elliptic curve over $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ (cf. I.3), one finds that \mathcal{P} is given by the quotient τ of two solutions of the hypergeometric differential equation with parameters $(\frac{1}{2}, \frac{1}{2}, 1)$.

1.2.3. The theory of period mappings has been generalized to any projective smooth morphism $f : \underline{X} \rightarrow S$ by P. Griffiths [Gri71]. We assume that S and the fibers of f are connected for simplicity. The idea is the following: by abstracting the structure of the periods of the fibers of f , one arrives at the notion of polarized Hodge structure; one then constructs a map from the universal covering \tilde{S} to a classifying space \mathcal{D} of polarized Hodge structures.

For a fixed n , the real cohomology spaces $V_s^{\mathbb{R}} := H^n(\underline{X}_s, \mathbb{R})$ form a local system $V^{\mathbb{R}}$ on S . We denote by $V^{\mathbb{C}}$ its complexification. The vector bundle $\mathcal{O}_S \otimes V^{\mathbb{C}}$ is the analytification of an algebraic vector bundle \mathcal{H} on S , and $V^{\mathbb{C}}$ is the local system of germs of analytic solutions of an algebraic fuchsian integrable connection ∇_{GM} on \mathcal{H} , the Gauss-Manin connection.

On the other hand, the spaces $V_s^{\mathbb{R}}$ carry a natural algebraic representation of \mathbb{C}^\times (Hodge structure), which amounts to a decomposition $V_s^{\mathbb{C}} = \bigoplus_{p+q=n, p \geq 0} V^{p,q}$, $\overline{V^{p,q}} = V^{q,p}$. When s varies, the filtrations

$$F_s^p := \bigoplus_{p'+q=n, p' \geq p} V^{p',q}$$

are the fibers of a decreasing filtration F_s^p of \mathcal{H} by vector subbundles (the Hodge filtration). The Gauss-Manin connection fails to preserve the Hodge filtration by just one notch: $\nabla_{\text{GM}}(F^p) \subset \Omega_S^1 \otimes F^{p-1}$.

The last piece of data is the polarization: a bilinear form Q on $V_s^{\mathbb{R}}$, $Q(v, v') = (-1)^n Q(v', v)$, such that $(-i)^n Q(v, \bar{v})$ is hermitian on $V_s^{\mathbb{C}}$, definite (of sign $(-1)^p$) on each $V^{p,q} = F_s^p \cap \overline{F}_s^q$, and such that F_s^p is the orthogonal of F_s^{n-p-1} in $V_s^{\mathbb{C}}$.

Let $G_{\mathbb{R}}$ denote the real group of similitudes of Q . The space of isotropic flags of type $0 \subset \dots \subset F_s^p \subset F_s^{p-1} \subset \dots \subset V_s^{\mathbb{C}}$ is a homogeneous space

$$\mathcal{D}^{\vee} = P \backslash G_{\mathbb{C}}$$

for a suitable parabolic subgroup P .

The flags satisfying the above positivity condition (on each $F_s^p \cap \overline{F}_s^q$) are classified by an analytic open submanifold, the *period domain*,

$$\mathcal{D} \subset \mathcal{D}^{\vee}; \quad \mathcal{D} \cong K \backslash G_{\mathbb{R}}^{\text{ad}}(\mathbb{R})^0$$

for a suitable compact subgroup K (\mathcal{D}^{\vee} is called the compact dual of \mathcal{D}).

Over \tilde{S} , the local system $V^{\mathbb{C}}$ becomes constant, and the construction which attaches to any $s \in \tilde{S}$ the classifying point of the corresponding flag $0 \subset \dots \subset F_s^p \subset F_s^{p-1} \subset \dots \subset V_s^{\mathbb{C}}$ gives rise to a holomorphic mapping, the *period mapping*

$$\tilde{S} \xrightarrow{\mathcal{P}} \mathcal{D}.$$

Via Plücker coordinates, \mathcal{P} is again given by quotients of solutions of the Gauss-Manin connection ∇_{GM} . The projective monodromy group of ∇_{GM} is a subgroup $\Gamma \subset G_{\mathbb{R}}^{\text{ad}}$ (well-defined up to conjugation), and we get a commutative square

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\mathcal{P}} & \mathcal{D} \subset \mathcal{D}^{\vee} \\ \downarrow & & \downarrow \\ S & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma. \end{array}$$

1.2.4. In the simple case of a family of elliptic curves and $n = 1$, $G_{\mathbb{R}} = GL_{2,\mathbb{R}}$, the flag space is just $\mathcal{D}^{\vee} = \mathbb{P}_{\mathbb{C}}^1$, $K = PSO(2)$ (the isotropy group of $i \in \mathbb{P}_{\mathbb{C}}^1$ in $PSL_2(\mathbb{R}) = G_{\mathbb{R}}^{\text{ad}}(\mathbb{R})^0$), and $\mathcal{D} \cong K \backslash G_{\mathbb{R}}^{\text{ad}}(\mathbb{R})^0$ is the Poincaré upper half plane \mathfrak{h} .

More generally, for a polarized abelian scheme, $G_{\mathbb{R}}$ is the group of symplectic similitudes, and \mathcal{D}^{\vee} is the grassmannian of lagrangian subspaces of \mathbb{C}^{2g} , and \mathcal{D} is the Siegel upper half space.

When one deals with refined moduli problems for abelian schemes with PEL decoration as in 1.1.4, one should work with the group $G_{\mathbb{R}}$ of B -linear

symplectic similitudes of $V_{\mathbb{R}}$ (notation as in 1.1.4). In that case, the corresponding period domain \mathcal{D} is a symmetric domain.

More precisely, the datum of our B -stable lagrangian subspace $F_0^1 \subset V_{\mathbb{C}}$ (attached to our particular A_0 with $H^1(A_0, \mathbb{C}) = V_{\mathbb{C}}$) amounts to the datum of an algebraic \mathbb{R} -homomorphism $h_0 : \mathbb{C}^\times \rightarrow G_{\mathbb{R}} \subset GL(V_{\mathbb{R}})$ such that h_0 acts by the characters z and \bar{z} on $V_{\mathbb{C}}$ (F^1 is the subspace where h_0 acts through z), and such that the symmetric form $\langle v, h(i)w \rangle$ is positive definite on $V_{\mathbb{R}}$. For any abelian variety A with PEL structure of the correct type, together with a fixed isomorphism $H^1(A, \mathbb{C}) \cong V_{\mathbb{C}}$, the corresponding subspace $F^1 H^1(A, \mathbb{C}) = \Omega^1(A) \subset V_{\mathbb{C}}$ is an eigenspace for some conjugate of h_0 . The set of conjugates of h_0 under $G(\mathbb{R})$ is a finite union of copies of a symmetric domain \mathcal{D} .

For any component S of the *Shimura orbifold*, we thus get a commutative square

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\mathcal{P}} & \mathcal{D} \subset \mathcal{D}^\vee \\ \mathcal{Q} \downarrow & & \downarrow \mathcal{Q} \\ S & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma \end{array}$$

where the horizontal maps are *isomorphisms*, \mathcal{Q} denotes the quotient maps, and Γ is a congruence subgroup of level N in the semi-simple group G^{ad} (a conjugate of the standard one).

The complex manifold \tilde{S} admits the following modular description. Let U be the oriented real Lie group $\text{Hom}(L, (\mathbb{R}/\mathbb{Z})^\times)$ (with \mathcal{B} -action and polarization), and let \hat{U} be the associated infinitesimal Lie group. The automorphism group J of the decorated \hat{U} coincides with $G(\mathbb{R})^0$. Then \tilde{S} is a moduli space for *marked* decorated abelian varieties, where the marking is an isomorphism $\rho : \hat{A} \rightarrow \hat{U}$ of associated infinitesimal decorated oriented real Lie groups (note that, by taking duals of Lie algebras, ρ amounts to an B -linear symplectic isomorphism $H^1(A, \mathbb{R}) \cong V^{\mathbb{R}}$). Note that the standard right action of J on \tilde{S} is obtained by functoriality.

2. Preliminaries on p -divisible groups.

ABSTRACT: Definitions and basic theorems about p -divisible groups, quasi-isogenies, liftings and deformations.

2.1. p -divisible groups and quasi-isogenies.

2.1.1. We begin with four definitions. Let S be a scheme. Let Λ and Λ' be commutative group schemes over S .

A homomorphism $f : \Lambda \rightarrow \Lambda'$ is called an *isogeny* if it is an (f.p.p.f.) epimorphism with finite locally free kernel.

Let p be a prime number. Λ is a *p -divisible group* (or Barsotti-Tate group) if $\Lambda = \varinjlim_n \text{Ker}[p^n]$ and for all n , the multiplication $[p^n]$ by p^n on Λ is an isogeny.

If S is connected, the rank of $\text{Ker}[p^n]$ is then of the form p^{hn} , where h is an integer called the height of Λ .

From the fact that $[p]$ is an isogeny, it follows that for p -divisible groups, $\text{Hom}_S(\Lambda, \Lambda')$ is a torsion-free \mathbb{Z}_p -module.

A *quasi-isogeny* of p -divisible groups Λ, Λ' is a global section ρ of the Zariski sheaf $\underline{\text{Hom}}_S(\Lambda, \Lambda') \otimes_{\mathbb{Z}} \mathbb{Q}$ such that there exists locally an integer n for which $p^n \rho$ is an isogeny. By abuse, one writes $\rho : \Lambda \rightarrow \Lambda'$ as for homomorphisms. We denote by $\text{qisog}_S(\Lambda, \Lambda')$ the \mathbb{Q}_p -space of quasi-isogenies.

The objects of the category of *p -divisible groups on S up to isogeny* are the p -divisible groups over S ; morphisms between Λ and Λ' are global sections of $\underline{\text{Hom}}_S(\Lambda, \Lambda') \otimes_{\mathbb{Z}} \mathbb{Q}$. This is a \mathbb{Q}_p -linear category.

2.1.2. The most important example of a p -divisible group is $\Lambda = A[p^\infty]$, the (inductive system of) p -primary torsion of an abelian scheme A over S ([Ta66]); in this case, the height h is twice the relative dimension of A .

2.1.3. When p is locally nilpotent on S , any p -divisible group Λ over S is formally smooth; the completion $\hat{\Lambda}$ of a p -divisible group Λ along the zero section is a formal Lie group [Gro74, VI,3.1]. However $\hat{\Lambda}$ is not necessarily itself a p -divisible group; it is if and only if the separable rank of the fibers of $\text{Ker}[p]$ is a locally constant function on S (in which case Λ is an extension

$$1 \rightarrow \hat{\Lambda} \rightarrow \Lambda \rightarrow \Lambda^{\text{et}} \rightarrow 1$$

of an ind-etale p -divisible group Λ^{et} by the infinitesimal p -divisible group $\hat{\Lambda}$, cf. [Gro74, III,7.4]). An example of an ind-etale (*resp.* infinitesimal) p -divisible group of height $h = 1$ is $\mathbb{Q}_p/\mathbb{Z}_p = \varinjlim_n \mathbb{Z}/p^n\mathbb{Z}$ (*resp.* $\hat{\mathbb{G}}_m$).

2.2. Two theorems on p -divisible groups.

2.2.1. A considerable amount of work has been devoted to p -divisible groups and their applications to the p -adic study of abelian varieties and their local moduli. Among the pioneers, let us mention: Barsotti, Tate, Serre, Grothendieck, Lubin, Messing...

In the sequel, we shall outline the main concepts and results of the Rapoport-Zink theory of period mappings for p -divisible groups, emphasizing the analogy with the complex case. We refer to their book [RZ96] for details and proofs. We shall also emphasize the relation to differential equations, which does not appear in [RZ96].

The whole theory relies upon three basic theorems on p -divisible groups and their Dieudonné modules, which we shall recall below:

- (i) the rigidity theorem for p -divisible groups up to isogeny,
- (ii) the Serre-Tate theorem, and
- (iii) the Grothendieck-Messing theorem,

which deal with infinitesimal deformations of p -divisible groups.

2.2.2. Let \mathfrak{v} be a complete discrete valued ring \mathfrak{v} of mixed characteristic $(0, p)$. Let $\text{Nil}_{\mathfrak{v}}$ denote the category of locally noetherian \mathfrak{v} -schemes S on which p is locally nilpotent. For any S in $\text{Nil}_{\mathfrak{v}}$, the ideal of definition $\mathcal{J} = \mathcal{J}_{S_{\text{red}}}$ of the closed subscheme S_{red} (of characteristic p) is locally nilpotent.

Theorem 2.2.3 (Rigidity theorem for p -divisible groups up to isogeny). *Every homomorphism $\bar{\rho} : \Lambda \times_S S_{\text{red}} \rightarrow \Lambda' \times_S S_{\text{red}}$ of p -divisible groups up to isogeny admits a unique lifting $\rho : \Lambda \rightarrow \Lambda'$. Moreover, ρ is a quasi-isogeny if $\bar{\rho}$ is.*

PROOF. (following an argument of V. Drinfeld). We may assume that S is affine, and that \mathcal{J} is nilpotent. Then the connected part $\hat{\Lambda}'$ is a formal Lie group, hence there is a power p^n of p such that for any affine S -scheme S' , $\hat{\Lambda}'(S')$ is killed by $[p^n]$ (in fact if $\mathcal{J}^{r+1} = 0$ on S , one can take $n = r^2$).

We observe that $\text{Ker}(\Lambda'(S') \rightarrow \Lambda'(S'_{\text{red}})) = \hat{\Lambda}'(S')$. We have

$$\begin{aligned} & \text{Ker}(\text{Hom}(\Lambda(S'), \Lambda'(S')) \rightarrow \text{Hom}(\Lambda(S'_{\text{red}}), \Lambda'(S'_{\text{red}}))) \\ &= \text{Hom}(\Lambda(S'), \text{Ker}(\Lambda'(S') \rightarrow \Lambda'(S'_{\text{red}}))) \end{aligned}$$

which is zero because $\text{Ker}(\Lambda'(S') \rightarrow \Lambda'(S'_{\text{red}}))$ is killed by $[p^n]$ while Λ is p -divisible.

This implies the injectivity of $\text{qisog}_S(\Lambda, \Lambda') \rightarrow \text{qisog}_{S_{\text{red}}}(\Lambda \times_S S_{\text{red}}, \Lambda' \times_S S_{\text{red}})$ (taking into account the fact that $\text{Hom}(\Lambda(S'_{\text{red}}), \Lambda'(S'_{\text{red}}))$ is torsion-free).

For the surjectivity, let us first show that for any $\bar{f} \in \text{Hom}(\Lambda(S'_{\text{red}}), \Lambda'(S'_{\text{red}}))$, there is a lifting $g \in \text{Hom}(\Lambda(S'), \Lambda'(S'))$ of $p^n \bar{f}$. For any $x \in \Lambda(S')$, let \bar{x} denote its image in $\Lambda(S'_{\text{red}})$. Since Λ' is formally smooth, $\bar{f}(\bar{x})$ admits a lifting $y \in \Lambda'(S')$. Since $\text{Ker}(\Lambda'(S') \rightarrow \Lambda'(S'_{\text{red}}))$ is killed by p^n , $g(x) := [p^n]y$ is well-defined. This construction provides the desired lifting g .

It remains to show that g is an isogeny if \bar{f} is. Let \bar{f}' be a quasi-inverse of \bar{f} ($\bar{f}'\bar{f} = [p^m]$), and g' be a lifting of $p^n \bar{f}'$. Then $g'g = [p^{2n+m}]$ by unicity of liftings. Thus g is an epimorphism, and the subscheme $\text{Ker } g$ of $\text{Ker}[p^{2n+m}]$ is finite over S . On the other hand, Λ is flat over S and the fibers of g' are flat (being isogenies). The criterium of flatness fiber by fiber [EGA IV, 11.3.10] implies that g' is flat, and we conclude that it is an isogeny. \square

Let us observe that there is variant of both the statement and the proof of 2.2.3 for abelian schemes, instead of p -divisible groups.

The next theorem asserts that deforming an abelian scheme of characteristic p is equivalent to deforming its p -divisible group. For any S in Nil_\flat , let $\text{Def}_{p\text{-div}}(S)$ be the category of triples $(\overline{A}, \Lambda, \epsilon)$ consisting of an abelian scheme over S_{red} , a p -divisible group Λ over S and an isomorphism $\epsilon : \overline{A}[p^\infty] \cong \Lambda \times_S S_{\text{red}}$.

Theorem 2.2.4 (The Serre-Tate theorem). *The functor*

$$A \mapsto (A_{\text{red}}, \Lambda = A[p^\infty], \epsilon : A_{\text{red}}[p^\infty] \cong \Lambda \times_S S_{\text{red}})$$

induces an equivalence of categories between the category of abelian schemes over S and the category $\text{Def}_{p\text{-div}}(S)$.

We again follow Drinfeld's argument [Dri76], cf. also [Ka81]. We may assume that S is affine and that $\mathcal{J}^{n+1} = 0$. We begin with the full faithfulness: given a homomorphism $f^\infty : A[p^\infty] \rightarrow B[p^\infty]$ such that $f^\infty \pmod{\mathcal{J}}$ comes from a homomorphism $\overline{f} : A_{\text{red}} \rightarrow B_{\text{red}}$ of abelian schemes over S_{red} , we have to show that there exists a unique homomorphism $f : A \rightarrow B$ compatible with f^∞ and \overline{f} . The unicity follows from the injectivity of $\text{Hom}(A, B) \rightarrow \text{Hom}(A_{\text{red}}, B_{\text{red}})$ (2.2.3). Furthermore, 2.2.3 shows the existence of a homomorphism g compatible with $p^n f^\infty$ and $p^n \overline{f}$. To show the existence of an f such that $g = p^n f = f \circ [p^n]$, the point is to show that g kills $A[p^n]$, which can be seen on g^∞ .

For the essential surjectivity, we have to construct an abelian scheme A over S compatible with a given datum $(\overline{A}, \Lambda, \epsilon)$ in $\text{Def}_{p\text{-div}}(S)$. By unicity of liftings, it suffices to do so locally on S . hence we may assume that S is affine, and choose an abelian scheme A' over S and an isogeny $A'_{\text{red}} \rightarrow \overline{A}$. By the above argument, p^n times the given isogeny $A'_{\text{red}} \rightarrow \overline{A}$ lifts to a (unique) isogeny $e : A'[p^\infty] \rightarrow \Lambda$ of p -divisible groups over S . Its kernel is a finite locally free subgroup scheme of A' , and we can form the abelian scheme $A'' := A' / \text{Ker } e$. The isogeny $A''[p^\infty] \rightarrow \Lambda$ induced by e is an isomorphism $\pmod{\mathcal{J}}$, hence is an isomorphism.

2.2.5. In the sequel, we shall have to deal occasionally with p -divisible groups over formal schemes rather than over schemes. Our formal schemes \mathfrak{X} will be adic, locally noetherian; hence there is a largest ideal of definition $\mathcal{J} \subset \mathcal{O}_{\mathfrak{X}}$, and $\mathfrak{X} = \varprojlim \mathfrak{X}_n$, $\mathfrak{X}_n = \text{Spec}(\mathcal{O}_{\mathfrak{X}}/\mathcal{J}^{n+1})$ ($\mathfrak{X}_{\text{red}} = \mathfrak{X}_0$).

A p -divisible group Λ over \mathfrak{X} will be an inductive system of p -divisible groups Λ_n over \mathfrak{X}_n such that $\Lambda_{n+1} \times_{\mathfrak{X}_{n+1}} \mathfrak{X}_n \cong \Lambda_n$.

3. A stroll in the crystalline world.

ABSTRACT: Review of Dieudonné modules, crystals, Grothendieck-Messing theory, convergent isocrystals.

3.1. From Dieudonné modules to crystals.

3.1.1. At first, Dieudonné theory presents itself as an analogue of Lie theory for formal Lie groups over a perfect field k of characteristic $p > 0$. In practice, its scope is limited to the case of *commutative* formal Lie groups (in which case the classical counterpart is not very substantial). The theory extends to the case of p -divisible groups over k .

Let W be the Witt ring of k , with its Frobenius automorphism σ . The theory associates to any p -divisible group Λ over k a *Dieudonné module*, *i.e.* a finitely generated free module over W endowed with a σ -linear Frobenius F and a σ^{-1} -linear Verschiebung V satisfying $FV = VF = p \cdot \text{id}$ (we work with the contravariant theory: $\mathbf{D}(\Lambda) = \text{Hom}(\Lambda, \mathcal{W}_k)$, where \mathcal{W}_k is the Witt scheme of k). This provides an *anti-equivalence of categories between p -divisible groups over k and Dieudonné modules*. The rank of $\mathbf{D}(\Lambda)$ is the height h of Λ .

Examples 3.1.2. • $\mathbf{D}(\mathbb{Q}_p/\mathbb{Z}_p) = W$, with $F = \sigma$, $V = p\sigma^{-1}$.

• $\mathbf{D}(\hat{\mathbb{G}}_m) = W(-1)$, *i.e.* W as an underlying module, with $F = p\sigma$, $V = \sigma^{-1}$.

• When $\Lambda = A[p^\infty]$ for an abelian variety A over k , $\mathbf{D}(\Lambda)$ may be identified with $H_{\text{cris}}^1(A/W)$.

However, the structure of Dieudonné modules is rather complicated in general unless one inverts p .

3.1.3. When k is algebraically closed, Dieudonné modules $\otimes \mathbb{Q}$ were completely classified by Dieudonné (this provides a classification of p -divisible groups over k up to isogeny). They are classified by their *slopes* $\lambda \in [0, 1] \cap \mathbb{Q}$ and multiplicities $m_\lambda \in \mathbb{Z}_{>0}$ (these data are better recorded in the form of a Newton polygon). This comes as follows: each Dieudonné module \mathbf{D} has a canonical increasing finite filtration by Dieudonné submodules \mathbf{D}_λ , such that the associated graded W -module is free, and with the following property: if $\lambda = a/b$ in irreducible form, $\text{Gr}_\lambda \otimes \mathbb{Q}$ admits a basis of m_λ vectors x satisfying $F^b x = p^a x$.

For instance, the slope of $W(-1)$ is 1; the slope of the Dieudonné module attached to (the p -divisible group of) a supersingular elliptic curve over k is $1/2$, with multiplicity 2.

3.1.4. The problem of generalizing Dieudonné theory to p -divisible groups over more general bases S (over which p is nilpotent) has been tackled and advertised by Grothendieck [**Gro74**], and further studied by many geometers.

What should be the right substitutes for the W -modules $\mathbf{D}(\Lambda)$ (*resp.* for the $W[\frac{1}{p}]$ -spaces $\mathbf{D}(\Lambda)_{\mathbb{Q}}$)?

Grothendieck's proposal was to define $\mathbf{D}(\Lambda)$ as a F -crystal on the crystalline site of S .

The right substitute for $W[\frac{1}{p}]$ -spaces was proposed later by P. Berthelot and A. Ogus under the name of *convergent F -isocrystal*. Let us describe these notions.

3.2. The Dieudonné crystal of a p -divisible group.

3.2.1. Let \mathfrak{v} be a finite extension of W of degree e . We assume that $e < p$, and that ϖ is a uniformizing parameter of \mathfrak{v} such that σ extends to \mathfrak{v} by setting $\sigma(\varpi) = \varpi$.

Let S be in $\text{Nil}_{\mathfrak{v}}$. An S -divided power thickening $T_0 \hookrightarrow T$ is given by

- (i) an S -scheme T_0 ,
- (ii) a \mathfrak{v} -scheme T on which p is locally nilpotent,
- (iii) a closed immersion $T_0 \hookrightarrow T$ over \mathfrak{v} ,
- (iv) a collection of maps “ $\frac{x^n}{n!}$ ” from the ideal of definition of T_0 in T to \mathcal{O}_T , which satisfy the formal properties of the divided powers.

S -divided power thickenings form a category $(S/\mathfrak{v})_{\text{cris}}$ (a fppf site). A sheaf on $(S/\mathfrak{v})_{\text{cris}}$ is the data, for every $T_0 \hookrightarrow T$, of a fppf sheaf $\mathcal{E}_{T_0, T}$ on T , and for every morphism $f : (T_0 \hookrightarrow T) \rightarrow (T'_0 \hookrightarrow T')$ in $(S/\mathfrak{v})_{\text{cris}}$, of a homomorphism $f^* \mathcal{E}_{T_0, T} \rightarrow \mathcal{E}_{T'_0, T'}$ satisfying the obvious transitivity condition. Example: $\mathcal{O}_{S_0/\mathfrak{v}}$, defined by $(\mathcal{O}_{S/\mathfrak{v}})_{T_0, T} = \mathcal{O}_T$. Another example: any fppf sheaf \mathcal{F} on S_0 gives rise to a sheaf $\mathcal{F}_{\text{cris}}$ on $(S/\mathfrak{v})_{\text{cris}}$ by $(\mathcal{F}_{\text{cris}})_{T_0, T} = \mathcal{F}_{T_0}$.

A *crystal* on $(S/\mathfrak{v})_{\text{cris}}$ (or, abusively, on S) is a sheaf \mathcal{E} of $\mathcal{O}_{S/\mathfrak{v}}$ -modules on $(S/\mathfrak{v})_{\text{cris}}$ such that the homomorphisms $f^* \mathcal{F}_{T_0, T} \rightarrow \mathcal{F}_{T'_0, T'}$ are isomorphisms (in Grothendieck's terms: “crystals grow and are rigid”). One denotes by \mathcal{E}_T the \mathcal{O}_T -module obtained by evaluating the crystal on $(\text{id} : T \hookrightarrow T) \in (S/\mathfrak{v})_{\text{cris}}$.

Because of this rigidity, and using the canonical divided powers on the ideal $\varpi\mathcal{O}_S$ (since $e < p$, $\frac{\varpi^n}{n!} \in \mathfrak{v}$ for any n), one gets an equivalence of categories between crystals on S and crystals on the scheme $S_0 = S/\varpi$ of characteristic p .

3.2.2. The Dieudonné crystal attached to a p -divisible group Λ over S is the crystalline $\mathcal{E}xt$ -sheaf $\mathcal{E}xt^1(\Lambda_{\text{cris}}, \mathcal{O}_{S/\mathfrak{v}})$. It can be shown that $\mathbf{D}(\Lambda)$ is a finite locally free crystal on S , of rank the height of Λ . It depends functorially in Λ (in a contravariant way).

An important, though formal, consequence of the crystalline local character of this definition is that *the formation of $\mathbf{D}(\Lambda)$ commutes with base change in $(S/\mathfrak{v})_{\text{cris}}$* , and for any S -divided power thickening $(T_0 \hookrightarrow T)$, there is a canonical isomorphism of \mathcal{O}_T -modules $\mathbf{D}(\Lambda)_{T_0, T} \cong \mathbf{D}(\Lambda_{T_0})_{T_0, T}$.

It follows that the datum of $\mathbf{D}(\Lambda)$ amounts to the datum of $\mathbf{D}(\Lambda \times_S S_0)$: the Dieudonné crystal depends only on the p -divisible group modulo ϖ .

3.2.3. The pull-back of a crystal \mathcal{E} on S_0 via the absolute Frobenius F_{S_0} is denoted by $\mathcal{E}^{(p)}$. An F -crystal structure on \mathcal{E} is the datum of a morphism of crystals $F : \mathcal{E}^{(p)} \rightarrow \mathcal{E}$ which admits an inverse up to a power of p .

If Λ is a p -divisible group over S_0 , $\mathbf{D}(\Lambda)$ is endowed by functoriality with a structure of F -crystal. The classical Dieudonné module is obtained for $S_0 = T_0 = \text{Spec}(k)$, $T = \text{Spf}(W) = \varinjlim \text{Spec}(W/p^n)$.

Although we shall not use this fact, let us mention that the “Dieudonné functor” \mathbf{D} from p -divisible groups on S_0 to F -crystals is fully faithful under further mild assumptions on S_0 (and even an equivalence with values in the category of “Dieudonné crystals”, *i.e.* finite locally free F -crystals for which F admits an inverse up to p — the *Verschiebung*); without any further assumption on S_0 , it induces an equivalence of categories between ind-étale p -divisible groups on S_0 and unit-root F -crystals (*i.e.* finite locally free F -crystals for which F is an isomorphism), *cf.* [BM79], [dJ95a].

3.3. The Hodge filtration.

3.3.1. Let Λ^\vee be the (Serre) dual of Λ (the \varinjlim of the Cartier duals of the $\Lambda[p^n]$'s). There is a canonical, locally split, exact sequence of \mathcal{O}_S -modules

$$0 \rightarrow F^1 \rightarrow \mathbf{D}(\Lambda)_S \rightarrow \text{Lie } \Lambda^\vee \rightarrow 0$$

where $F^1 = \text{CoLie } \Lambda = \omega_\Lambda$ may be identified with the module of invariant differentials on the formal Lie group $\hat{\Lambda}$. In case Λ is the p -divisible group attached to an abelian S -scheme \underline{A} , this exact sequence reduces to the Hodge exact sequence

$$0 \rightarrow F^1 \rightarrow H_{\text{DR}}^1(\underline{A}/S) \rightarrow \text{Lie } \underline{A}^\vee \rightarrow 0.$$

3.3.2. Now let $S_0 \hookrightarrow S$ be a *nilpotent* divided power thickening (this means that the products $\frac{x_1^{n_1}}{n_1!} \dots \frac{x_m^{n_m}}{n_m!}$ vanish for $n_1 + \dots + n_m \gg 0$); this is for instance the case if $S_0 = S/\varpi$ as before and $e < p - 1$.

The next theorem asserts that deformations of p -divisible groups are controlled by the variation of the Hodge filtration in the Dieudonné module [Mes72]. For any p -divisible group $\bar{\Lambda}$ on S_0 , and any lifting Λ over S , we have seen that $\mathbf{D}(\Lambda)_S$ identifies with $\mathbf{D}(\bar{\Lambda})_S$, which is a lifting of $\mathbf{D}(\bar{\Lambda})_{S_0}$ to S .

Theorem 3.3.3 (The Grothendieck-Messing theorem). *The functor*

$$\Lambda \mapsto (\Lambda \times_S S_0, \omega_\Lambda)$$

induces an equivalence of categories between the category of p -divisible groups on S , and the category of pairs consisting of a p -divisible group $\bar{\Lambda}$ over S_0 (Zariski-locally liftable to S) and a locally direct factor vector subbundle of $\mathbf{D}(\bar{\Lambda})_S$ which lifts $\omega_{\bar{\Lambda}} \subset \mathbf{D}(\bar{\Lambda})_{S_0}$.

We have already seen the faithfulness in 2.2.3. We just indicate here the principle of proof of essential surjectivity and fullness. Working with nilpotent divided power thickenings allows one to construct, by the exponential

method, a formally smooth group scheme $\mathbb{E}(\overline{\Lambda})$ over S together with a canonical isomorphism $\mathbf{D}(\overline{\Lambda})_{S_0} = \text{Lie}(\mathbb{E}(\overline{\Lambda}) \times_S S_0)$, in such a way that the canonical exact sequence

$$0 \rightarrow \omega_{\overline{\Lambda}} \rightarrow \mathbf{D}(\overline{\Lambda})_{S_0} \rightarrow \text{Lie } \overline{\Lambda}^\vee \rightarrow 0$$

lifts to an exact sequence of formally smooth group schemes

$$0 \rightarrow \omega_{\overline{\Lambda}} \rightarrow \mathbb{E}(\overline{\Lambda}) \times_S S_0 \rightarrow \overline{\Lambda}^\vee \rightarrow 0$$

Via the exponential, it is equivalent to give a local summand F^1 of $\mathbf{D}(\overline{\Lambda})_S$ which lifts $\omega_{\overline{\Lambda}}$, or to give a vector sub-group scheme V of $\mathbb{E}(\overline{\Lambda})$ which lifts $\omega_{\overline{\Lambda}}$. It then turns out that $\mathbb{E}(\overline{\Lambda})/V$ is a p -divisible group Λ^\vee , whose Serre dual is the looked for p -divisible group Λ over S : $\Lambda \times_S S_0 = \overline{\Lambda}$ by duality, and $\omega_\Lambda = F^1$.

The formation of $\mathbb{E}(\overline{\Lambda})$ is functorial (contravariant in $\overline{\Lambda}$). If we have two p -divisible groups $\overline{\Lambda}, \overline{\Lambda}'$ and a morphism $\overline{f}: \overline{\Lambda} = \Lambda \times_S S_0 \rightarrow \overline{\Lambda}' = \Lambda' \times_S S_0$ such that $\mathbf{D}(f)$ respects the F^1 's, then the homomorphism $\mathbb{E}(\overline{f}): \mathbb{E}(\overline{\Lambda}) \rightarrow \mathbb{E}(\overline{\Lambda}')$ preserves the vector subgroup schemes, hence induces a homomorphism $f: \mathbb{E}(\overline{\Lambda})/V \cong \Lambda \rightarrow \mathbb{E}(\overline{\Lambda}')/V' \cong \Lambda'$ which lifts \overline{f} .

The exponential construction becomes simpler when the ideal of S_0 in S is of square zero (and endowed with the trivial divided powers). This is the case which will be used in the sequel.

3.4. Crystals and connections.

3.4.1. We have already encountered the terms “ F -crystal”, “unit-root F -crystal” in I.2.4.2, 2.4.3 in the more down-to-earth context of connections. This comes as follows. Assume for simplicity that S_0 is affine of characteristic p and lifts to a formally smooth p -adic affine formal scheme \mathcal{S} over $\mathfrak{v} = W$. Then there is an equivalence of categories between finite locally free crystals on $(S_0/W)_{\text{cris}}$ and finite locally free $\mathcal{O}(\mathcal{S})$ -modules with integrable topologically nilpotent connection (this interpretation of crystals extends to much more general situations, *cf.* [dJ95a, 2.2.2]).

Using this equivalence of categories, the Katz (covariant) functor

$$\left(\begin{array}{c} \text{continuous } \mathbb{Z}_p\text{-representations} \\ \text{of } \pi_1^{\text{alg}}(S_0, s_0) \end{array} \right) \longrightarrow (\text{unit-root } F\text{-crystals})$$

is the composite of the following (anti)equivalences of categories

$$\left(\begin{array}{c} \text{continuous } \mathbb{Z}_p\text{-representations} \\ \text{of } \pi_1^{\text{alg}}(S_0, s_0) \end{array} \right) \longrightarrow (\text{ind-etale } p\text{-divisible groups over } S_0) \\ \xrightarrow{\mathbf{D}} (\text{unit-root } F\text{-crystals over } S_0).$$

3.4.2. Let us briefly recall the construction of the connection attached to a finite locally free crystal \mathcal{E} on $(S_0/W)_{\text{cris}}$. It depends on the technique of formally adding divided powers to an ideal (the pd-hull). Let Δ_0 be the

diagonal in $S_0 \times S_0$, and let $\hat{\Delta}$ be the p -completion of the pd-hull of the diagonal in $\mathcal{S} \hat{\times} \mathcal{S}$, endowed with the two projections p_1, p_2 to \mathcal{S} . Then by rigidity of crystals, we have for any n an isomorphism

$$p_2^*(\mathcal{E}_{S_0, \mathcal{S}/p^n}) \cong \mathcal{E}_{\Delta_0, \hat{\Delta}/p^n} \cong p_1^*(\mathcal{E}_{S_0, \mathcal{S}/p^n})$$

and at the limit $n \rightarrow \infty$, an isomorphism of finite locally free $\mathcal{O}(\hat{\Delta})$ -modules

$$\epsilon : p_2^*(\mathcal{E}_{S_0, \mathcal{S}}) \cong p_1^*(\mathcal{E}_{S_0, \mathcal{S}})$$

satisfying a cocycle condition. Such an isomorphism is the ‘‘Taylor series’’

$$\epsilon(1 \otimes e) = \sum_{n_1, \dots, n_d \geq 0} \left(\nabla \left(\frac{\partial}{\partial t_1} \right)^{n_1} \dots \nabla \left(\frac{\partial}{\partial t_d} \right)^{n_d} \right) (e) \otimes \prod_i \frac{(1 \otimes t_i - t_i \otimes 1)^{n_i}}{n_i!}$$

of an integrable connection ∇ on the finite locally free $\mathcal{O}(\mathcal{S})$ -module $\mathcal{E}_{S_0, \mathcal{S}}$ (here t_1, \dots, t_d denote local coordinates on \mathcal{S}).

3.4.3. In general, the Taylor series converges only in polydiscs of radius $|p|^{1/p-1}$, due to the presence of the factorials. One says that ∇ is *convergent* if its Taylor series converges in (open) unit polydiscs. A well-known argument due to Dwork shows that F -crystals give rise to convergent connections ∇ ([Ka73]).

In order to put the discussion of convergence on proper foundations, we need the notion of tube in analytic geometry.

3.5. Interlude : analytic spaces associated with formal schemes and tubes.

3.5.1. Let us first review the Raynaud-Berthelot construction of the analytic space attached to a \mathfrak{v} -formal scheme \mathfrak{X} . It will be convenient for later purpose to formulate the construction in the frame of Berkovich spaces rather than rigid geometry (we refer to [Ber93, 1.6] for the translation).

As in 2.2.5, we assume that \mathfrak{X} is adic, locally noetherian; moreover, we assume that $\mathfrak{X}_{\text{red}} = \text{Spec}(\mathcal{O}_{\mathfrak{X}}/\mathcal{J})$ is a separated k -scheme locally of finite type (in particular, $p \in \mathcal{J}$). However, we *do not* assume that \mathfrak{X} is p -adic, *i.e.* that the topology of $\mathcal{O}_{\mathfrak{X}}$ is p -adic.

We describe the construction in the affine case $\mathfrak{X} = \text{Spf}(\mathcal{A})$; the general case follows by gluing. One chooses generators f_1, \dots, f_r of $J = \Gamma(\mathfrak{X}, \mathcal{J})$, and defines, for any $n > 0$,

$$\mathcal{B}_n = \mathcal{A}\{T_1, \dots, T_r\} / (f_1^n - \varpi T_1, \dots, f_r^n - \varpi T_r)$$

where $\mathcal{A}\{T_1, \dots, T_r\}$ stands for the ϖ -adic completion of $\mathcal{A}[T_1, \dots, T_r]$. One has $\mathcal{B}_n / \varpi \mathcal{B}_n \cong (\mathcal{A} / (\varpi, f_1^n, \dots, f_r^n)) [T_1, \dots, T_r]$, a k -algebra of finite type; hence \mathcal{B}_n is a \mathfrak{v} -algebra topologically of finite type (*i.e.* quotient of an algebra of the form $\mathcal{A}\{T_1, \dots, T_r\}$), so that $\mathcal{B}_n[\frac{1}{p}]$ is an affinoid algebra. Let $\mathcal{M}(\mathcal{B}_n)$ be its Berkovich spectrum. The homomorphism $\mathcal{B}_{n+1} \rightarrow \mathcal{B}_n$ sending T_i to $f_i T_i$ identifies $\mathcal{M}(\mathcal{B}_n)$ with an affinoid subdomain of $\mathcal{M}(\mathcal{B}_{n+1})$ [Be96, 0.2.6].

The analytic space \mathfrak{X}^{an} is defined as the union $\bigcup_n \mathcal{M}(\mathcal{B}_n)$. It does not depend on the choice of f_1, \dots, f_r . This is a *paracompact (strictly) analytic space* over the fraction field K of \mathfrak{v} (one can replace the nested “ball-like” $\mathcal{M}(\mathcal{B}_n)$ by nested “annulus-like” spaces in order to obtain locally finite coverings).

Examples 3.5.2. (i) if $\mathfrak{X} = \text{Spf}(\mathfrak{v}\{t\})$, \mathfrak{X}^{an} is the closed unit disc.

(ii) if $\mathfrak{X} = \text{Spf}(\mathfrak{v}[[t]])$, $\mathcal{J} = (\varpi, t)$, then $\mathcal{M}(\mathcal{B}_n)$ is the closed disc $\mathbb{D}_K(0, |\varpi|^{1/n})$, and \mathfrak{X}^{an} is the open unit disc.

3.5.3. Let S_0 be a closed subscheme of $\mathfrak{X}_{\text{red}}$, and let \mathfrak{X}_{S_0} be the formal completion of \mathfrak{X} along S_0 . The *tube of S_0 in \mathfrak{X}* is $\mathfrak{X}_{S_0}^{\text{an}}$. It is denoted by $]S_0[_{\mathfrak{X}}$ (and considered as a subspace of \mathfrak{X}^{an}). It depends only on $S_{0,\text{red}}$.

Example: if \mathfrak{X} is as in example (i) above (so that $\mathfrak{X}_{\text{red}} = \mathbb{A}_k^1$), and S_0 is the point 0, then \mathfrak{X}_{S_0} is as in example (ii) above: $]0[_{\mathfrak{X}} = \mathbb{D}(0, 1^-) \subset \mathbb{D}(0, 1^+)$. More generally, the polydiscs mentioned in 3.4.3 are the tubes of closed points.

3.6. Convergent isocrystals.

3.6.1. Let S_0 be a separated k -scheme of finite type, and $S_0 \hookrightarrow \mathfrak{Y}$ a closed immersion into a flat p -adic formal \mathfrak{v} -scheme that is formally smooth in a neighborhood of S_0 . Let E be a vector bundle on the tube $]S_0[_{\mathfrak{Y}}$. An integrable connection ∇ on E is *convergent* if its Taylor series

$$\epsilon(1 \otimes e) = \sum_{n_1, \dots, n_d \geq 0} \left(\nabla \left(\frac{\partial}{\partial t_1} \right)^{n_1} \dots \nabla \left(\frac{\partial}{\partial t_d} \right)^{n_d} \right) (e) \otimes \prod_i \frac{“(1 \otimes t_i - t_i \otimes 1)^{n_i}”}{n_i!}$$

is induced by an isomorphism on the tube of the diagonal $]S_0[_{\mathfrak{Y} \times \mathfrak{Y}}$

$$\epsilon : p_2^*(E) \cong p_1^*(E)$$

It amounts to the same to say that for any affine U in \mathfrak{Y} with local coordinates t_1, \dots, t_d , any section $e \in \Gamma(U^{\text{an}}, E)$ and any $\eta < 1$, one has

$$\left\| \nabla \left(\frac{\partial}{\partial t_1} \right)^{n_1} \dots \nabla \left(\frac{\partial}{\partial t_d} \right)^{n_d} (e) \right\|_{\eta^{n_1 + \dots + n_d}} \rightarrow 0$$

where $\| \cdot \|$ denotes a Banach norm on $\Gamma(U^{\text{an}}, E)$.

According to Berthelot, *the category of vector bundles with convergent connection depends only on S_0 — even only on $S_{0,\text{red}}$ — (up to canonical equivalence), and is functorial in S_0/K ; this is the category of *convergent isocrystals on S_0/K* (or, abusively, on S_0).*

Since any separated k -scheme locally of finite type can be locally embedded into a \mathfrak{X} as above, one can define the category of convergent isocrystals on such a scheme by gluing [Be96, 2.3.2].

3.6.2. There is a natural functor from the category of F -isocrystals on S_0/\mathfrak{v} to the category of convergent isocrystals on S_0/K [Be96, 2.4]. The composite of this functor with the Dieudonné functor is denoted by $\mathbf{D}(-)_{\mathbb{Q}}$. It

factors through a \mathbb{Q}_p -linear functor, still denoted by $\mathbf{D}(-)_{\mathbb{Q}}$:

$$\left\{ \begin{array}{l} p\text{-divisible groups over } S_0 \\ \text{up to isogeny} \end{array} \right\} \longrightarrow \{\text{convergent isocrystals on } S_0\}$$

3.6.3. The advantage of convergent isocrystals over “crystals up to isogeny” lies in their strong “continuity property”, which allows to remove the restriction $e < p$ on ramification (as we shall do henceforth), and work conveniently with singular schemes S_0 (while divided powers present certain “pathologies” in the singular case). In our applications, S_0 will be a countable (non-disjoint) union of irreducible projective varieties, and \mathfrak{v} may be highly ramified.

3.6.4. If S_0 is the reduced underlying scheme $\mathfrak{X}_{\text{red}}$ of a formal scheme \mathfrak{X} as in 3.5.1, there is a notion of *evaluation of E on \mathfrak{X}^{an}* [Be96, 2.3.2]: this is a vector bundle on \mathfrak{X}^{an} with an integrable connection.

By functoriality of the construction, it will be enough to deal with the affine case $\mathfrak{X} = \text{Spf } \mathcal{A}$ and to assume that there is a closed immersion $S_0 = (\text{Spf } \mathcal{A})_{\text{red}} \hookrightarrow \mathfrak{Y}$ into a formally smooth p -adic formal \mathfrak{v} -scheme. One can construct inductively a compatible family of \mathfrak{v} -morphisms $\text{Spec}(\mathcal{O}_{\mathfrak{X}}/\mathcal{I}^n) \rightarrow \mathfrak{Y}$, whence a \mathfrak{v} -morphism $\mathfrak{X} \rightarrow \mathfrak{Y}$. Let $v :]S_0[_{\mathfrak{X}} = \mathfrak{X}^{\text{an}} \rightarrow]S_0[_{\mathfrak{Y}}$ be the associated analytic map. The looked for evaluation of E on \mathfrak{X}^{an} is the pull-back by v of the vector bundle with integrable connection on $]S_0[_{\mathfrak{Y}}$ defined by E .

3.6.5. Let now S be a separated scheme in $\text{Nil}_{\mathfrak{v}}$ (it may also be viewed in Nil_W). Let \mathcal{E} be a finite locally free F -crystal on S/W . This provides a vector bundle \mathcal{E}_S on S , which only depends on the inverse image $\mathcal{E} \times_W k$ of \mathcal{E} on the k -scheme S/p (cf. 3.2.1). On the other hand, the convergent isocrystal E on S/p attached to F -crystal $\mathcal{E} \times_W k$ (cf. 3.6.2) depends only on S_{red} .

3.6.6. Let us apply this to Dieudonné crystals. Let \mathfrak{X} be a formal scheme as in 3.5.1, and let $\Lambda = \varinjlim \Lambda_n$ be a p -divisible on $\mathfrak{X} = \varinjlim \mathfrak{X}_n$. This gives rise to a vector bundle $\mathbf{D}(\Lambda)_{\mathfrak{X}} = \varinjlim \mathbf{D}(\Lambda_n)_{\mathfrak{X}_n}$ on \mathfrak{X} . On the other hand, we have the vector bundle $\mathbf{D}(\Lambda_0)_{\mathfrak{X}}$ on \mathfrak{X} and a natural homomorphism $\mathbf{D}(\Lambda_0)_{\mathfrak{X}} \rightarrow \mathbf{D}(\Lambda)_{\mathfrak{X}}$ obtained by pull-back $\mathfrak{X}_{\text{red}} = \mathfrak{X}_0 \rightarrow \mathfrak{X}$.

When $e \geq p$, this is not an isomorphism in general. However, *the associated morphism of analytic vector bundles $(\mathbf{D}(\Lambda_0)_{\mathfrak{X}})^{\text{an}} \rightarrow (\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$ is an isomorphism. Moreover, $(\mathbf{D}(\Lambda_0)_{\mathfrak{X}})^{\text{an}}$ is the analytic vector bundle underlying the evaluation of $\mathbf{D}(\Lambda_0)_{\mathbb{Q}}$ on \mathfrak{X}^{an} .*

Indeed, working locally, and taking in account the very construction of \mathfrak{X}^{an} , we reduce to the case when \mathfrak{X} is p -adic. Then \mathfrak{X}/p is a k -scheme. Using the canonical divided powers on (p) and the rigidity of crystals, $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathfrak{X}} \cong \mathbf{D}(\Lambda)_{\mathfrak{X}}$. It is thus enough to see that $\mathbf{D}(\Lambda_0)^{\text{an}} \cong \mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathfrak{X}}^{\text{an}}$. This follows from the fact that the convergent isocrystals $\mathbf{D}(\Lambda_0)_{\mathbb{Q}}$ and $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathbb{Q}}$ are “the same”, since $\mathfrak{X}_0 = (\mathfrak{X}/p)_{\text{red}}$: indeed, it is clear that $(\mathbf{D}(\Lambda_0)_{\mathfrak{X}})^{\text{an}}$ (resp. $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathfrak{X}}^{\text{an}}$) coincides with the evaluation of

the convergent isocrystal $\mathbf{D}(\Lambda_0)_{\mathbb{Q}}$ (*resp.* $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathbb{Q}}$) on \mathfrak{X}^{an} (*cf.* also [dJ95b, 6.4], in a slightly less general context).

3.6.7. Let us draw from this two consequences:

- (i) taking into account the fact that the functor $\mathbf{D}(-)_{\mathbb{Q}}$ factors through the category of p -divisible groups up to isogeny, we see that if we are given a quasi-isogeny $\Lambda_0 \rightarrow \Lambda'_0$, then there is a canonical isomorphism of vector bundles $(\mathbf{D}(\Lambda'_0)_{\mathfrak{X}})^{\text{an}} \rightarrow (\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$.
- (ii) Through the isomorphism $(\mathbf{D}(\Lambda_0)_{\mathfrak{X}})^{\text{an}} \rightarrow (\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$ (or equivalently, through the isomorphism $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathfrak{X}}^{\text{an}} \rightarrow (\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$), the analytic vector bundle $(\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$ is naturally endowed with an integrable connection, which deserves to be called the *Gauss-Manin* connection attached to Λ .

Indeed, when $\Lambda = \underline{A}[p^\infty]$ for an abelian scheme \underline{A} over \mathfrak{X} , $(\mathbf{D}(\Lambda)_{\mathfrak{X}})^{\text{an}}$ may then be identified with the De Rham cohomology bundle $H_{\text{DR}}^1(\underline{A}^{\text{an}}/\mathfrak{X}^{\text{an}})$ together with its Gauss-Manin connection. Here, one can use the comparison theorem between crystalline and De Rham cohomology, or the theory of universal vectorial extensions and Grothendieck's \natural -structures, *cf.* [MM74], [BBM82].

In the first alternative, one uses the divided powers on (p) and construct a canonical isomorphism of vector bundles with connection $\mathbf{D}(\Lambda \times_{\mathfrak{X}} \mathfrak{X}/p)_{\mathfrak{X}} \cong H_{\text{DR}}^1(\underline{A}/\mathfrak{X})$.

In the second alternative, one uses Grothendieck's construction of a canonical connection on the universal vectorial extension $\mathbb{E}(\Lambda)$ of Λ^\vee — *cf.* 3.3.2; the induced connection on the Lie algebra $\mathbf{D}(\Lambda)_{\mathfrak{X}}$ turns out to be the Gauss-Manin connection on $H_{\text{DR}}^1(\underline{A}/\mathfrak{X})$.

The Gauss-Manin connection is especially significant when \mathfrak{X}^{an} is smooth (even though \mathfrak{X} itself may not be formally smooth).

4. Moduli problems for p -divisible groups.

ABSTRACT: Review of moduli spaces of p -divisible groups quasi-isogenous to a fixed one modulo p , with examples.

4.1. Moduli problem for p -divisible groups quasi-isogenous to a fixed one modulo p .

4.1.1. Here, we work over the discrete valuation ring $\mathfrak{v} = W = \widehat{\mathbb{Z}_p^{\text{ur}}}$ (the completion of the maximal unramified extension of \mathbb{Z}_p), with uniformizing parameter $\varpi = p$. Recall that $\text{Nil}_{\mathfrak{v}}$ denote the category of locally noetherian \mathfrak{v} -schemes S on which p is locally nilpotent.

Let $\overline{\Lambda}$ be a fixed p -divisible group over $k = \overline{\mathbb{F}_p}$. The following theorem is due to M. Rapoport and T. Zink [RZ96, 2.16, 2.32].

Theorem 4.1.2. *The functor*

$$\text{Nil}_{\mathfrak{v}} \rightarrow \text{Sets}$$

$$S \mapsto \left\{ (\Lambda, \rho) \mid \begin{array}{l} \Lambda: p\text{-divisible group on } S, \\ \rho \in \text{qisog}(\Lambda \times_S S_{\text{red}}, \overline{\Lambda} \times_{\overline{\mathbb{F}_p}} S_{\text{red}}) \end{array} \right\} / \cong$$

is representable by a formal scheme \mathfrak{M} over \mathfrak{v} ; $\mathfrak{M}_{\text{red}}$ is locally of finite type over $\overline{\mathbb{F}_p}$ and its irreducible components are projective $\overline{\mathbb{F}_p}$ -varieties.

Of course, the rigidity of p -divisible groups up to isogeny (2.2.3) is crucial here. The main difficulty in proving 4.1.2 is to control the powers of p which are needed in the process of lifting isogenies.

The separated formal scheme \mathfrak{M} is not formally smooth in general, but has been conjectured to be flat over \mathfrak{v} . The group J of self-quasi-isogenies of $\overline{\Lambda}$ acts (on the right) on \mathfrak{M} by $(\Lambda, \rho).j = (\Lambda, \rho \circ j)$. This group J is actually the group of \mathbb{Q}_p -points of an algebraic group over \mathbb{Q}_p .

We denote by \mathcal{M} the Berkovich analytic space over \mathbb{C}_p attached to \mathfrak{M} : $\mathcal{M} = \mathfrak{M}^{\text{an}} \hat{\otimes}_K \mathbb{C}_p$ (cf. 3.5). This is a paracompact (strictly) analytic space. We shall see later that it is smooth, and a p -adic manifold in the sense of I.1.3.7.

4.2. Examples.

We take $\overline{\Lambda} = \overline{A}[p^\infty]$, the p -primary torsion of an elliptic curve \overline{A} over $\overline{\mathbb{F}_p}$ ($h = 2$).

4.2.1. *The ordinary case, i.e. $\overline{A}[p](\overline{\mathbb{F}_p}) \neq 0$.*

Then $\overline{\Lambda} \cong \hat{\mathbb{G}}_m \oplus \mathbb{Q}_p/\mathbb{Z}_p$, and $J = (\mathbb{G}_{m\mathbb{Q}_p})^2$. For any (Λ, ρ) in $\mathfrak{M}(S)$, the separable rank of $\text{Ker}[p]$ is constant (due to the quasi-isogeny ρ). Therefore Λ is an extension

$$\hat{\Lambda} \rightarrow \Lambda \rightarrow \Lambda^{\text{et}}.$$

The quasi-isogeny ρ respects this extension, hence splits into two parts $(\hat{\rho}, \rho^{\text{et}})$. We have $\hat{\Lambda} \cong \hat{\mathbb{G}}_m$, $\Lambda^{\text{et}} \cong \mathbb{Q}_p/\mathbb{Z}_p$. Up to isomorphism, the pair

$(\hat{\rho}, \rho^{\text{et}})$ thus amounts to an element of $(\mathbb{Q}_p^\times/\mathbb{Z}_p^\times)^2 \cong \mathbb{Z}^2$. Therefore

$$\mathfrak{M} = \coprod_{\mathbb{Z}^2} \mathfrak{M}^0$$

is a disjoint sum of copies of a formal (group) scheme \mathfrak{M}^0 which parametrizes extensions of $\mathbb{Q}_p/\mathbb{Z}_p$ by $\hat{\mathbb{G}}_m$. It is known that

$$\mathfrak{M}^0 = \hat{\mathbb{G}}_m$$

This identification is given by the following recipe. Recall that if $S = \text{Spec}(R)$, R local artinian with residue field $\overline{\mathbb{F}}_p$ and radical \mathfrak{m} ($\mathfrak{m}^{n+1} = 0$), then $\hat{\mathbb{G}}_m(S) = 1 + \mathfrak{m}$; in particular $\hat{\mathbb{G}}_m(S)$ is killed by $[p^n]$. Let us choose any lift q_n of $1/p^n \in \mathbb{Q}_p/\mathbb{Z}_p$ in Λ . Then $[p^n]q_n$ is a well-defined element q in $\hat{\Lambda}(S) \cong 1 + \mathfrak{m}$, which is unchanged if n is replaced by a bigger integer (but another choice of the isomorphism $\hat{\Lambda} \cong \hat{\mathbb{G}}_m$ would change q into q^a , $a \in \mathbb{Z}_p^\times$). As a formal W -scheme, $\mathfrak{M}^0 = \text{Spf}(\mathfrak{v}[[q-1]])$, with the $(p, q-1)$ -adic topology. The associated analytic space \mathcal{M}^0 over \mathbb{C}_p is the open unit disc $|q-1| < 1$.

The parameter q is thus the local modulus for deformations of $\overline{A}[p^\infty]$. On the other hand, if $p \neq 2$, we have an algebraic local modulus for deformations of \overline{A} (say, with level two structure): the Legendre parameter $z = \lambda$ around the Legendre parameter ζ_{can} of the canonical lifting A_{can} of \overline{A} (cf. I.3.4.2). The p -divisible group $A_{\text{can}}[p^\infty]$ splits: its parameter is $q = 1$. Therefore, in this special case, the Serre-Tate theorem 2.2.4 asserts that $q \in 1 + (z - \zeta_{\text{can}})\hat{\mathbb{Z}}^{\text{ur}}[[z - \zeta_{\text{can}}]]$. It has been proved by W. Messing and N. Katz that q is the Dwork-Serre-Tate parameter discussed in I.3.4, ([Mes76], [Ka81]) — which explains the terminology.

4.2.2. *The supersingular case, i.e. $\overline{A}[p](\overline{\mathbb{F}}_p) = 0$.*

In this case $\overline{\Lambda} = \hat{\Lambda}$ is a so-called Lubin-Tate formal group, $\text{End } \overline{\Lambda} = \mathcal{B}_p$, “the” maximal order in the (non-split) quaternion algebra B_p over \mathbb{Q}_p , and $J = B_p^\times$.

Here again, there is a discrete invariant: the “height” of the quasi-isogeny ρ , which amounts to the p -adic valuation of the norm of its image in $J = B_p^\times$. Therefore

$$\mathfrak{M} = \coprod_{\mathbb{Z}} \mathfrak{M}^0$$

is a disjoint sum of copies of a formal scheme \mathfrak{M}^0 which parametrizes deformations of the Lubin-Tate group $\hat{\Lambda}$. According to Lubin-Tate, $\mathfrak{M}^0 \cong \text{Spf}(\mathfrak{v}[[t]])$, with the (p, t) -adic topology. The associated analytic space \mathcal{M}^0 over \mathbb{C}_p is the open unit disc $|t| < 1$.

Assume for simplicity that \overline{A} is defined over \mathbb{F}_p . Then one can describe the universal formal deformation Λ over $\text{Spf } \mathbb{Z}_p[[t]]$ through its logarithm

$\sum b_n(t)X^{p^{2n}}$, which is given recursively by

$$b_0(t) = 1, \quad b_n(t) = \frac{1}{p} \sum_{0 \leq m < n} b_m(t)t_{n-m}^{p^{2m}}$$

The so-called canonical lifting, corresponding to $t = 0$, has formal complex multiplication by \mathbb{Z}_{p^2} , cf. [GH94a], [GH94b].

4.3. Decorated variants.

4.3.1. Just as in the complex case 1.1.4, it is useful to consider variants of the moduli problem 4.1 for “decorated” p -divisible groups, with prescribed endomorphisms and/or polarization.

Instead of the \mathbb{Q} -algebra B of 1.1.4, we shall consider a finite-dimensional semi-simple \mathbb{Q}_p -algebra B_p , since the category of p -divisible groups up to isogeny is \mathbb{Q}_p -linear. Let V_p be a B_p -module of finite type, and let $G_{\mathbb{Q}_p}$ be the group of B_p -linear endomorphisms of V_p .

Let \mathcal{B}_p be a maximal order in B_p , and let L_p be a lattice in V_p , stable under \mathcal{B}_p .

4.3.2. We assume that \mathcal{B}_p acts on our p -divisible group $\bar{\Lambda}$. One then redefines J to be the group of self-quasi-isogenies of $\bar{\Lambda}$ which respect the \mathcal{B}_p -action.

For technical reasons, one picks up a lifting $\tilde{\Lambda}$ of $\bar{\Lambda}$ with \mathcal{B}_p -action to some finite extension of W , and looks at the \mathcal{B}_p -module $F_0^1 := \omega_{\tilde{\Lambda}}$. We denote by \mathfrak{v} the finite extension $W[\mathrm{tr}(b|F_0^1)]$ of W .

One strengthens the moduli problem 4.1 by imposing to our pairs (Λ, ρ) that Λ is endowed with a \mathcal{B}_p -action $\iota : \mathcal{B}_p \rightarrow \mathrm{End} \Lambda$, ρ respects the \mathcal{B}_p -action, and moreover a “Shimura type condition” $\det(\iota(b)|\omega(\Lambda)) = \det(b|F_0^1)$.

It follows easily from 4.1.2 [RZ96, 3.25] that this moduli problem is representable by a formal scheme, still denoted by \mathfrak{M} , over \mathfrak{v} . It is acted on by J . $\mathfrak{M}_{\mathrm{red}}$ is locally of finite type over $\overline{\mathbb{F}}_p$ and its irreducible components are projective $\overline{\mathbb{F}}_p$ -varieties.

4.3.3. Let us say a word about the polarized variant (here, it is safer to assume $p \neq 2$). One assumes that \mathcal{B}_p is endowed with an involution $*$, and that L_p is autodual for an alternate \mathbb{Z}_p -bilinear form such that $\langle bv, w \rangle = \langle v, b^*w \rangle$. Of course, here, $G_{\mathbb{Q}_p}$ denotes the algebraic \mathbb{Q}_p -group of B_p -linear symplectic similitudes of V_p . One assumes that $\bar{\Lambda}$ is endowed with a $*$ -polarization ⁽²⁾, and modify J to respect the polarization up to a multiple in \mathbb{Q}_p .

The corresponding moduli problem is again representable by a formal scheme with the same properties. This formal scheme has been conjectured to be flat by Rapoport-Zink; although this has been verified in many cases (cf. e.g. [Gor01]), there are some counter-examples (cf. [P00]).

⁽²⁾a $*$ -polarization is a symmetric \mathcal{B}_p -linear quasi-isogeny between the p -divisible and its Serre dual, endowed with the transposed action of \mathcal{B}_p twisted by $*$

Example 4.3.4 (fake elliptic curves at a critical prime p). Let us fix a indefinite quaternion algebra B over \mathbb{Q} and a maximal order \mathcal{B} in B (after Eichler, they are all conjugate). A prime p is called *critical* if it divides the discriminant of B , *i.e.* if $B_p = B \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is the (non-split) quaternion algebra over \mathbb{Q}_p . Note that B_p then contains copies of the unramified quadratic extension on \mathbb{Q}_p ; we denote by \mathbb{Q}_{p^2} one of them. The maximal order (unique up to conjugation) can be written $\mathcal{B}_p = \mathcal{B} \otimes \mathbb{Z}_p = \mathbb{Z}_{p^2}[\Pi]$, $\Pi^2 = p$, $\Pi.a = a^\sigma \Pi$ for $a \in \mathbb{Z}_{p^2}$. We fix a critical prime p .

We consider abelian surfaces with quaternionic multiplication by \mathcal{B} . They are sometimes called “*fake elliptic curves*” because they share many features with elliptic curves, *cf.* [BC91, III, 1].

We fix a fake elliptic curve \bar{A} over $\bar{\mathbb{F}}_p$, and set $\Lambda = \bar{A}[p^\infty]$. This is a special formal \mathcal{B}_p -modules of height 4 in the following sense.

Let S be in Nil_p (or else an inductive limit of such schemes — as a formal scheme), and let Λ be a p -divisible group over S . After Drinfeld, one says that Λ is a *special formal \mathcal{B}_p -modules of height 4* if

- (i) Λ is infinitesimal ($\Lambda = \hat{\Lambda}$) of height $h = 4$,
- (ii) \mathcal{B}_p acts on Λ ,
- (iii) via this action, ω_Λ is a locally free $(\mathbb{Z}_{p^2} \otimes_{\mathbb{Z}_p} \mathcal{O}_S)$ -module of rank 1.

We take $V_p = B_p$ where B_p acts by left multiplication. In this situation, $G_{\mathbb{Q}_p} \cong B_p^\times$. When $S = \text{Spec}(\bar{\mathbb{F}}_p)$, there is a unique class of \mathcal{B}_p -isogeny of special formal \mathcal{B}_p -modules of height 4. The group J of self-quasi-isogenies of a special formal \mathcal{B}_p -module of height 4 is $J = GL_2(\mathbb{Q}_p)$. This can be easily read off the Dieudonné module $\otimes \mathbb{Q}$.

In this example, there is again a discrete invariant: the “height” of the quasi-isogeny ρ , which amounts to the p -adic valuation of the determinant of its image in J . Therefore

$$\mathfrak{M} = \coprod_{\mathbb{Z}} \mathfrak{M}^0$$

is a disjoint sum of copies of a formal scheme \mathfrak{M}^0 which parametrizes deformations of the special formal module $\hat{\Lambda}$. This formal scheme has been described by Drinfeld, *cf.* 6.3.4.

4.3.5. This example and examples 4.2 were the sources of the general theory. Let us at once suggest a vague analogy with the archimedean situation — especially 1.1.4, 1.2.4 — which will become more and more precise in the next sections: one can see the p -divisible group $\bar{\Lambda}$ in characteristic p (*resp.* J) as an analogue of the infinitesimal oriented real Lie group \hat{U} (*resp.* $G(\mathbb{R})^0$).

5. p -adic period domains.

ABSTRACT: Review of p -adic symmetric domains and period domains via stability theory.

5.1. p -adic flag spaces.

5.1.1. We begin with the p -adic analogue of the flag space \mathcal{D}^\vee of 1.2.3, 1.2.4. In 1.2.4, this flag space parametrized the lagrangian subspaces F^1 in the H_{DR}^1 of abelian varieties of a certain type, marked by a fixed isomorphism $H_{\text{DR}}^1 \cong V^{\mathbb{C}}$.

In the p -adic case, \mathcal{D}^\vee will parametrize the Hodge filtration F^1 of Dieudonné modules of p -divisible groups of a certain type (3.3.1).

5.1.2. More precisely, let us consider a moduli problem for p -divisible groups as in 4.1, possibly decorated as in 4.3 (from which we follow the notation). The decoration reflects on the Dieudonné module $\mathbf{D}(\overline{\Lambda})_{\mathbb{Q}} = \mathbf{D}(\overline{\Lambda}) \otimes \mathbb{Q}$: one has an isomorphism of B_p -module (with symplectic structure, in the polarized case):

$$\mathbf{D}(\overline{\Lambda})_{\mathbb{Q}} \cong V_p \otimes_{\mathbb{Q}_p} \widehat{\mathbb{Q}_p^{\text{ur}}}.$$

We fix such an isomorphism, and consider that $G_{\mathbb{Q}_p} \otimes_{\mathbb{Q}_p} \widehat{\mathbb{Q}_p^{\text{ur}}}$ acts on $\mathbf{D}(\overline{\Lambda})_{\mathbb{Q}}$. But $\mathbf{D}(\overline{\Lambda})_{\mathbb{Q}}$ has an extra structure: the σ -linear Frobenius automorphism. The group J is nothing but the *subgroup* of $G_{\mathbb{Q}_p}(\widehat{\mathbb{Q}_p^{\text{ur}}})$ of elements commuting with Frobenius.

5.1.3. Let \mathfrak{v} be the finite extension of $W = \mathbb{Z}_p^{\text{ur}}$ considered in 4.3.2 and let K be its fraction field. We denote here by $F_0^1 \subset V_p^{\mathbb{C}_p} = V_p \otimes \mathbb{C}_p$ the \mathbb{C}_p -span of the space $\omega_{\overline{\Lambda}} \subset V_p \otimes_{\mathbb{Q}_p} K$ introduced in 4.3.2. We set $G_{\mathbb{C}_p} = G_{\mathbb{Q}_p} \otimes_{\mathbb{Q}_p} \mathbb{C}_p$. The flag variety \mathcal{D}^\vee of all $G_{\mathbb{C}_p}$ -conjugates of F_0^1 in $V_p^{\mathbb{C}_p} = \mathbf{D}(\overline{\Lambda}) \otimes_W \mathbb{C}_p$ is a homogeneous space

$$\mathcal{D}^\vee = P \backslash G_{\mathbb{C}_p}$$

for a suitable parabolic subgroup P . There is an ample line bundle \mathcal{L} on \mathcal{D}^\vee which is homogeneous under the derived group $G_{\mathbb{C}_p}^{\text{der}}$; we denote by $\mathcal{D}^\vee \subset \mathbb{P}(W)$ an associated $G_{\mathbb{C}_p}^{\text{der}}$ -equivariant projective embedding (if $G_{\mathbb{C}_p} = GL_{2g}$, $\mathcal{D}^\vee = \text{Grass}(g, 2g)$, one can just take the SL_{2g} -equivariant Plücker embedding).

Note that, there is an obvious right action of J on \mathcal{D}^\vee , which factors through the adjoint group $J^{\text{ad}} := \text{Im}(J \rightarrow G_{\mathbb{C}_p}^{\text{ad}})$.

5.1.4. In our three basic examples (4.2.1, 4.2.2, 4.3.4), we have $G_{\mathbb{C}_p} = GL_2$, hence $\mathcal{D}^\vee = \mathbb{P}_{\mathbb{C}_p}^1$.

5.2. Symmetric domains via stability theory.

5.2.1. In the p -adic case, one cannot define symmetric spaces by a positivity condition as in 1.2.3, 1.2.4. One uses instead the Hilbert-Mumford notion of semi-stability, following an idea of M. van der Put and H. Voskuil [vdPV92].

5.2.2. We set $J' = J \cap G_{\mathbb{C}_p}^{\text{der}}$. This is again the group of \mathbb{Q}_p -points of an algebraic group over \mathbb{Q}_p . Let $T \cong (\mathbb{G}_m)_{\mathbb{Q}_p}^j$ be a maximal \mathbb{Q}_p -split torus in J' . The *semi-stable locus* of \mathcal{D}^\vee with respect to T -action is the open subset of points $x \in \mathcal{D}^\vee$ such that there is a T -invariant function of $\mathbb{P}(W)$ which does not vanish on x .

One can check whether x is semi-stable with respect to T using the Hilbert criterion: let λ be any cocharacter (= one-parameter subgroup) of T . By properness of \mathcal{D}^\vee , the map $z \in \mathbb{G}_m \mapsto x \cdot \lambda(z) \in \mathcal{D}^\vee$ extends to \mathbb{A}^1 , *i.e.* the limit $x_0 = \lim_{z \rightarrow 0} x \cdot \lambda(z)$ exists. It is clearly fixed by \mathbb{G}_m , thus the fiber \mathcal{L}_{x_0} corresponds to a character $z \mapsto z^{-\mu(x, \lambda)}$ of \mathbb{G}_m . Then x is semi-stable if and only if the Mumford invariant $\mu(x, \lambda)$ is ≥ 0 for every λ .

5.2.3. One defines the *symmetric space* \mathcal{D} to be the intersection of the semi-stable loci of \mathcal{D}^\vee with respect to all maximal \mathbb{Q}_p -split tori T in J' . This is a p -adic manifold in the sense of I.1.3.6. It is also called the *p -adic period domain* (associated with the moduli problem).

Since J acts on the set of tori T by conjugation, it is clear that \mathcal{D} is stable under the J -action on \mathcal{D}^\vee .

5.3. Examples.

5.3.1. In our three basic examples, we have $G_{\mathbb{C}_p} = GL_2$, $\mathcal{D}^\vee = \mathbb{P}^1$.

In example 4.2.1, $J' \subset J = (\mathbb{G}_m)^2$ is the one-dimensional torus T with elements (z^{-1}, z) . Its action on a point x is by z^2 -scaling; the limit point x_0 is always 0, except if $x = \infty$. The only non-semi-stable point is ∞ , hence $\mathcal{D} = \mathbb{A}^1$.

In example 4.2.2, $J' = SL_1(B_p) \subset J = B_p^\times$. There is no non-trivial \mathbb{Q}_p -split torus in J' , hence $\mathcal{D} = \mathcal{D}^\vee = \mathbb{P}^1$.

5.3.2. We now turn to example 4.3.4. In this example, $J = GL_2(\mathbb{Q}_p)$ and $J' = SL_2(\mathbb{Q}_p)$. All maximal \mathbb{Q}_p -split tori are conjugate $g^{-1}Tg$ of the above T . Therefore, $\mathcal{D} = \mathbb{P}^1 \setminus \bigcup \infty.g$ is the *Drinfeld space* over \mathbb{C}_p

$$\mathcal{D} = \Omega_{\mathbb{C}_p} := \mathbb{P}_{\mathbb{C}_p}^1 \setminus \mathbb{P}^1(\mathbb{Q}_p)$$

To be consistent, we consider the right action of PGL_2 -action on \mathbb{P}^1 , which is deduced from the customary left action by the rule $x.g = g^{-1}.x$.

The Drinfeld space is actually defined over \mathbb{Q}_p , *i.e.* is a \mathbb{Q}_p -analytic manifold Ω . It is closely related to the *Bruhat-Tits tree* \mathcal{T} of $J^{\text{ad}} = PGL_2(\mathbb{Q}_p)$. The vertices of this tree correspond to closed discs in \mathbb{Q}_p ; two such vertices are connected by an edge if for the corresponding discs D', D'' , one has

$D' \subset D''$ and the radius of D'' is p times the radius of D' (each vertex has $p + 1$ neighbors). The group $PGL_2(\mathbb{Q}_p)$ acts on \mathcal{T} (on the right, say).




FIGURE 1. \mathcal{T} for $p = 2$

For any disc D centered in \mathbb{Q}_p , let η_D denote its Berkovich “generic point” (which is a point in Berkovich’s affine line \mathbb{A}^1 over \mathbb{Q}_p). Then the geometric realization of \mathcal{T} may be identified with the closed subset

$$|\mathcal{T}| = \{\eta_{D(a,r^+)} \mid a \in \mathbb{Q}_p, 0 < r < \infty\} \subset \Omega$$

(a $PGL_2(\mathbb{Q}_p)$ -equivariant embedding), and the set of vertices with the subset

$$\text{ver } \mathcal{T} = \{\eta_{D(a,r^+)} \mid a \in \mathbb{Q}_p, r \in p^{\mathbb{Z}}\} \subset |\mathcal{T}|$$

According to Berkovich, $|\mathcal{T}|$ is a retract of Ω [Ber90]; in particular, Ω is simply connected.

On the other hand, Ω is the generic fiber of a p -adic formal scheme $\hat{\Omega}$ whose reduction modulo p is an infinite tree of projective lines, with dual graph \mathcal{T} (cf. [BC91, I]).

The Drinfeld space Ω is a p -adic analogue of the Poincaré double half-plane $\mathbb{P}_{\mathbb{C}}^1 \setminus \mathbb{P}^1(\mathbb{R})$. Note that the latter space could also be defined using stability, in the same way (with $J' = SL_2(\mathbb{R})$). Another similarity: according to [Ber90], there is a $PGL_2(\mathbb{Q}_p)$ -invariant metric on Ω (which extends the standard metric on $|\mathcal{T}|$).

6. The p -adic period mapping and the Gauss-Manin connection.

ABSTRACT: Rapoport-Zink's period mapping as classifying map for the Hodge filtration in the Dieudonné module. Basic properties: equivariance, étaleness. Its relation to the Gauss-Manin connection. Examples and formulas.

6.1. Construction.

6.1.1. The construction of the p -adic period mapping is parallel to the complex case: one attaches to a point $s \in \mathcal{M}$ representing a pair (Λ, ρ) the point $\mathcal{P}(s)$ of \mathcal{D}^\vee which parametrizes the notch F^1 of the Hodge filtration of the Dieudonné module of Λ . The idea of the construction is already present in Grothendieck's talk [Gro70].

6.1.2. More precisely, notation being as in 4.1.1–4.3.1, let us consider the universal object $(\underline{\Lambda}, \rho)$ on the moduli formal scheme \mathfrak{M} , and the Dieudonné crystal $\mathbf{D}(\underline{\Lambda})$. Its evaluation $\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}}$ on \mathfrak{M} is a vector bundle on \mathfrak{M} .

Lemma 6.1.3. *The quasi-isogeny $\underline{\rho}$ induces a trivialization of the associated analytic vector bundle on $\mathcal{M} = \mathfrak{M}_{\mathbb{C}_p}^{\text{an}}$:*

$$(\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}} \cong V_p^{\mathbb{C}_p} \otimes_{\mathbb{C}_p} \mathcal{O}_{\mathcal{M}}$$

PROOF. ρ is a quasi-isogeny between $\underline{\Lambda} \times_{\mathfrak{M}} \mathfrak{M}_{\text{red}}$ and $\overline{\Lambda} \times_{\overline{\mathbb{F}}_p} \mathfrak{M}_{\text{red}}$. According to 3.6.7 (i), by the rigidity of convergent isocrystals, it induces a canonical isomorphism $(\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}} \cong (\mathbf{D}(\overline{\Lambda} \times_{\overline{\mathbb{F}}_p} \mathfrak{M}_{\text{red}})_{\mathfrak{M}})^{\text{an}}$. On the other hand, since the formation of $\mathbf{D}(\Lambda)$ commutes with base change (3.2.2), we have

$$(\mathbf{D}(\overline{\Lambda} \times_{\overline{\mathbb{F}}_p} \mathfrak{M}_{\text{red}})_{\mathfrak{M}})^{\text{an}} = \mathbf{D}(\overline{\Lambda}) \otimes_W \mathcal{O}_{\mathcal{M}} = V_p \otimes_{\mathbb{Q}_p} \mathcal{O}_{\mathcal{M}}.$$

□

6.1.4. On the other hand, we have a locally direct summand $F^1 = (\omega_{\underline{\Lambda}})^{\text{an}}$ in $(\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}} \cong V_p \otimes_{\mathbb{Q}_p} \mathcal{O}_{\mathcal{M}}$ (3.3.1). This determines an \mathcal{M} -point of the flag space \mathcal{D}^\vee , *i.e.* an analytic mapping

$$\mathcal{P} : \mathcal{M} \longrightarrow \mathcal{D}^\vee$$

This is called the *period mapping* for the moduli problem (decorated or not). It is clear from the construction that \mathcal{P} is J -equivariant.

If \mathcal{M}^0 is any connected component of \mathcal{M} , it is stable under $J' \subset J$, hence the induced period mapping $\mathcal{P} : \mathcal{M}^0 \longrightarrow \mathcal{D}^\vee$ is J' -equivariant (the J' -action factors through J'^{ad}).

6.2. Properties.

Theorem 6.2.1. *The period mapping \mathcal{P} is étale. Its image lies in the period domain \mathcal{D} .*

The first assertion reduces, via an infinitesimal criterion of étaleness, to the “essential surjectivity part” of the Grothendieck-Messing theorem

(3.3.3), in the case of an ideal of square zero, *cf.* [RZ96, 5.17]. It corresponds to what is called the “local Torelli property” in the complex case.

The second assertion is a consequence of a theorem of B. Totaro on the equivalence of semi-stability and “weak admissibility” [Tot96]. The second assertion may be viewed as a p -adic analogue of the “Riemann relations” (1.1.3).

Corollary 6.2.2. *\mathcal{M} is smooth, and any component \mathcal{M}^0 is a connected p -adic manifold in the sense of I.1.3.7.*

6.2.3. It is very likely that \mathcal{P} , viewed as a mapping $\mathcal{M} \rightarrow \mathcal{D}$, is surjective. This is closely related to Fontaine’s conjecture on the realizability of filtered Dieudonné modules by p -divisible groups. Although recent work by C. Breuil [Bre99], [Bre00] settles the Fontaine conjecture at least for finite residue fields of characteristic $\neq 2$ (and in other cases), it is not clear to the author to which extent the surjectivity conjecture is now established.

On the other hand, the fibers of the period mapping are relatively well-understood, *cf.* [RZ96, 5.37].

6.2.4. The argument in the proof of 6.1.3 shows a little more. Recall from 3.6.7 (ii) that the analytic vector bundle $(\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}}$ on \mathcal{M} comes together with an integrable connection ∇_{GM} (the Gauss-Manin connection).

Lemma 6.2.5. *The trivialization*

$$(\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}} \cong V_p^{\mathbb{C}_p} \otimes_{\mathbb{C}_p} \mathcal{O}_{\mathcal{M}}$$

of 6.1.3 induces a trivialization of the Gauss-Manin connection:

$$((\mathbf{D}(\underline{\Lambda})_{\mathfrak{M}})^{\text{an}})^{\nabla_{\text{GM}}} \cong V_p^{\mathbb{C}_p}.$$

The point is that the convergent isocrystal $\mathbf{D}(\Lambda \times_{\mathfrak{M}} \mathfrak{M}_{\text{red}})$ is constant, since it comes by base change, from $\mathbf{D}(\overline{\Lambda})_{\mathbb{Q}}$.

This useful lemma will allow us to give explicit expression of the period mapping in terms of quotients of solutions of differential equations, as in the complex case.

6.3. Examples and formulas.

6.3.1. We start with example 4.1.1 (p -divisible groups of elliptic curves with ordinary reduction). The extension

$$1 \rightarrow \hat{\mathbb{G}}_m \rightarrow \Lambda \rightarrow \mathbb{Q}_p/\mathbb{Z}_p \rightarrow 1$$

parametrized by $q \in 1 + pW$ gives rise, by contravariance, to an extension of Dieudonné modules (*cf.* 3.1.1)

$$0 \rightarrow W \rightarrow \mathbf{D}(\Lambda) \rightarrow W(-1) \rightarrow 0$$

which splits canonically due to the rigidity of Dieudonné crystals: $\mathbf{D}(\Lambda) = \mathbf{D}(\overline{\Lambda}) = W \oplus W(-1)$. We denote by e_0, e_{-1} the corresponding canonical basis of $\mathbf{D}(\Lambda)$. The image $\tau = \mathcal{P}(q) \in W[\frac{1}{p}]$ of q under the period mapping

is, by definition, the unique element such that $\tau e_0 + e_{-1}$ generates F^1 . It turns out that

$$\tau = \log q,$$

thus

$$\mathcal{M}^0 = \mathbf{D}(1, 1^-) \xrightarrow{\mathcal{P}=\log} \mathbb{A}^1 = \mathcal{D}.$$

Let us indicate a sketch of proof of this remarkable formula (the formula goes back to Messing-Katz [Mes76], [Ka81]). The argument is more transparent if we think in terms of one-motives instead of elliptic curves: that is, rather than viewing Λ as the p -divisible attached to some elliptic curve over W , one remarks that it is also the p -divisible group attached to the one-motive $M = [\mathbb{Z} \xrightarrow{1 \mapsto q} \mathbb{G}_m]$. Then $\mathbf{D}(\Lambda)$ can be identified with $H_{\text{DR}}(M)$, the dual of the Lie algebra of G^{\natural} , where $[\mathbb{Z} \rightarrow G^{\natural}]$ denotes the universal vectorial extension of M (cf. [Del74]); this extension splits canonically ($G^{\natural} = \mathbb{G}_a \times \mathbb{G}_m$), and this splitting agrees with the above splitting of $\mathbf{D}(\Lambda)$. We are thus reduced to prove the following purely algebraic fact: let us view $M = [\mathbb{Z} \xrightarrow{1 \mapsto q} \mathbb{G}_m]$ as a one-motive over $K((q-1))$; then in terms of the canonical basis e_0, e_{-1} of $H_{\text{DR}}(M)$, $F^1 \subset H_{\text{DR}}(M)$ is generated by the vector $(\log q)e_0 + e_{-1}$. This well-known fact is easily proved by transcendental means, replacing K by \mathbb{C} .

6.3.2. Since the basis e_0, e_{-1} is horizontal under the Gauss-Manin connection ∇ on $\mathbf{D}(1, 1^-)$ (6.2.5)⁽³⁾, τ is a quotient of solutions of the Gauss-Manin connection. In terms of the Legendre parameter $z = \lambda$ (around the parameter ζ_{can} of the canonical lifting A_{can} of the fixed ordinary elliptic curve \overline{A}), the equality $\tau = \log q$ means that $q \in 1 + (z - \zeta_{\text{can}})W[[z - \zeta_{\text{can}}]]$ is the Dwork-Serre-Tate parameter discussed in I.3.4 (for $p \neq 2$).

This allows to give explicit formulas for \mathcal{P} as a function of the algebraic parameter z rather than q . For instance, let us assume that $p \equiv 1 \pmod{4}$. The disc $|z - \frac{1}{2}| < 1$ is an ordinary disc. The canonical lifting is $\zeta_{\text{can}} = \frac{1}{2}$ (complex multiplication by $\mathbb{Z}[\sqrt{-1}]$). The Gauss-Manin connection is represented by the hypergeometric differential operator with parameter $(\frac{1}{2}, \frac{1}{2}, 1)$. Let us recall the setting, which we have already met in I.5.3.11.

We consider the symplectic basis $\omega_1 = [\frac{dx}{2y}]$, $\omega_2 = [\frac{(2x-1)dx}{4y}]$ of relative De Rham cohomology. It has the virtue to be, at $z = \frac{1}{2}$, a basis of eigenvectors for the complex multiplication. The Gauss-Manin connection satisfies

$$\omega_2 = 2z(z-1)\nabla\left(\frac{d}{dz}\right)\omega_1 + \frac{4z-5}{6}\omega_1.$$

⁽³⁾which is induced by the usual Gauss-Manin connection attached to the Legendre elliptic pencil, cf. 3.6.6 (ii).

The first row of the solution matrix Y normalized by $Y(\frac{1}{2}) = \text{id}$ is given by

$$\begin{aligned} y_{11} &= F\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{2}; (1-2z)^2\right) + \frac{1}{2}(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right), \\ y_{12} &= (1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right). \end{aligned}$$

We have to relate $\omega_{1|_{1/2}}$, $\omega_{2|_{1/2}}$ to the symplectic basis e_0, e_{-1} . Because $z = 1/2$ gives the canonical lifting, the canonical splitting of the Dieudonné module is given by the complex multiplication: e_0, e_{-1} is a basis of eigenvectors. In fact, taking into account the fact that we deal with symplectic bases, we find

$$\omega_{1|_{1/2}} = \frac{\Theta}{2}e_{-1}, \quad \omega_{2|_{1/2}} = \frac{2}{\Theta}e_0,$$

where $\Theta \in W^\times$ denotes the Tate constant (for $z = \frac{1}{2}$) discussed in I.3.4.1, 4.5. (the ambiguity — by a factor in \mathbb{Z}_p^\times — is removed since the basis e_0, e_{-1} has been fixed). The period mapping $\tau = \tau(z)$ is given by the quotient of the entries in the first row of the matrix Y . $\begin{pmatrix} \Theta/2 & 0 \\ 0 & 2/\Theta \end{pmatrix}$, that is to say:

$$\tau(z) = \left(\frac{\Theta^2}{4}\right) \frac{(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right)}{F\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{2}; (1-2z)^2\right) + \frac{1}{2}(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right)}$$

Note that since the target of the period mapping is \mathbb{A}^1 , the denominator does not vanish on $D(\frac{1}{2}, 1^-)$. Note also that, in analytic terms, the étaleness of \mathcal{P} (6.2.1) reflects the fact that $\frac{d\tau}{dz}$ does not vanish, being the quotient of linearly independent solutions of a linear differential equation of rank two.

6.3.3. We now turn to example 4.1.2 (p -divisible groups of elliptic curves with supersingular reduction). In this case, the period mapping

$$\mathcal{M}^0 \cong D(0, 1^-) \xrightarrow{\mathcal{P}} \mathbb{P}^1 = \mathcal{D}$$

was thoroughly investigated by Gross-Hopkins [GH94a], [GH94b]. In the special case considered in 4.1.2. ii) (the fixed supersingular elliptic curve \overline{A} being defined over $\overline{\mathbb{F}}_p$), they obtained the following formula

$$\mathcal{P}(t) = \lim_{n \rightarrow \infty} p^n b_{2n}(t) / p^{n+1} b_{2n+1}(t) \in \mathbb{Q}_p[[t]]$$

and a closed formula was proposed by J. K. Yu in [Yu95, 11].

In terms of the algebraic Legendre parameter $z = \lambda$ in a supersingular disc, it is still true, via 6.2.5, that $\tau = \mathcal{P}(z)$ is a quotient of solutions of the Gauss-Manin connection. This allows to give explicit formulas for \mathcal{P} as a function of the algebraic parameter z . For instance, let us assume that $p \equiv 3 \pmod{4}$. The disc $|z - \frac{1}{2}| < 1$ is a supersingular disc. In terms of a symplectic basis of eigenvectors for the action of $[\sqrt{-1}]$ on $V_p^{\text{C}p}$, we get the following formula

for the Gross-Hopkins period mapping

$$\tau(z) = \kappa \frac{(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right)}{F\left(\frac{1}{4}, \frac{1}{4}, \frac{1}{2}; (1-2z)^2\right) + \frac{1}{2}(1-2z)F\left(\frac{3}{4}, \frac{3}{4}, \frac{3}{2}; (1-2z)^2\right)}$$

where κ is a constant depending on the choice of the basis (a natural choice would be to take this basis inside the p -adic Betti lattice discussed in I.5.3; the corresponding constant κ is then the one occurring in the proof of I.Theorem 5.3.9.

Note that since the target of the period mapping is \mathbb{P}^1 , the denominator vanishes somewhere on $D(\frac{1}{2}, 1^-)$.

Note also that the period mapping in the complex situation⁽⁴⁾ is given, up to a constant factor, by *the same hypergeometric formula, viewed as a complex analytic function*.

6.3.4. At last, let us turn to example 4.3.4 (p -divisible groups of fake elliptic curves at a critical prime p). In this case, the period mapping

$$\mathcal{M}^0 \xrightarrow{\mathcal{P}} \Omega_{\mathbb{C}_p} = \mathcal{D}$$

is an *isomorphism*. This is a reinterpretation of Drinfeld's work, cf. [RZ96] (and [BC91, II8]). Actually, Drinfeld establishes an isomorphism at the level of formal schemes

$$\mathfrak{M}^0 \cong \hat{\Omega} \hat{\otimes}_{\mathbb{Z}_p} \widehat{\mathbb{Z}_p^{\text{ur}}}$$

which is J^1 -equivariant (where J^1 denotes the image in $J^{\text{ad}} = PGL_2(\mathbb{Q}_p)$ of the elements of J whose determinant is a p -adic unit).

6.3.5. Recall that the p -adic manifold $\Omega_{\mathbb{C}_p}$ is simply connected (in the usual topological sense). However, the above isomorphism and the modular property of \mathfrak{M} , allowed Drinfeld to construct a tower of finite etale connected Galois coverings of $\Omega_{\mathbb{C}_p}$ (or even of $\Omega_{\widehat{\mathbb{Q}_p^{\text{ur}}}}$), as follows.

One considers the universal special formal module $\underline{\Lambda}$ of height 4 over $\mathfrak{M}^0 \cong \hat{\Omega} \hat{\otimes}_{\mathbb{Z}_p} \widehat{\mathbb{Z}_p^{\text{ur}}}$. For any $n \geq 1$, $\text{Ker}[p^n]$ is a finite locally free formal group scheme of rank p^{4n} over $\hat{\Omega} \hat{\otimes}_{\mathbb{Z}_p} \widehat{\mathbb{Z}_p^{\text{ur}}}$. This gives rise to a finite etale covering of the analytic space $\Omega_{\widehat{\mathbb{Q}_p^{\text{ur}}}}$ (fibered in rank-one $\mathcal{B}_p/p^n \mathcal{B}_p$ -modules).

To get a connected Galois covering, one looks at the Π -action (recall from 4.3.4 that $\mathcal{B}_p = \mathbb{Z}_{p^2}[\Pi]$, $\Pi^2 = p$): one has $\text{Ker}[p^n] = \text{Ker}[\Pi^{2n}]$, and the complement of $\text{Ker}[\Pi^{2n-1}]$ in $\text{Ker}[\Pi^{2n}]$ provides a finite etale Galois covering Σ^n of $\Omega_{\widehat{\mathbb{Q}_p^{\text{ur}}}}$ with group $(\mathcal{B}_p/p^n \mathcal{B}_p)^\times$. When n grows, these form a projective system of Galois coverings (with Galois group the profinite completion of \mathcal{B}_p^\times). This tower is equivariant with respect to the $GL_2(\mathbb{Q}_p)$ -action.

Over \mathbb{C}_p , Σ^n decomposes into finitely many copies of a connected p -adic manifold Υ^n which is a finite etale Galois covering of $\Omega_{\mathbb{C}_p}$ with group

⁽⁴⁾for the Legendre elliptic pencil, and computed with help of a basis of cycles — with coefficients in the Gauss integers — which are eigenvectors for complex multiplication at $z=1/2$.

$SL_1(\mathcal{B}_p/p^n\mathcal{B}_p)$. This can be seen either by using various maximal commutative \mathbb{Q}_p -subalgebras of B_p and reasoning as in [Var98a, 1.4.6], or by global means, using the Čerednik-Drinfeld theorem (7.4.7), or else by the properties of the so-called determinant map [BZ95, II].

Up to now, there is no other method for constructing non-abelian finite étale coverings of $\Omega_{\mathbb{C}_p}$. The coverings Υ^n remain rather mysterious: are they simply-connected? Is any connected finite étale covering of $\Omega_{\mathbb{C}_p}$ a quotient of some Υ^n ? Does there exist an infinite étale Galois covering of analytic manifolds $\Upsilon \rightarrow \Omega_{\mathbb{C}_p}$ with Galois group $SL_1(\mathcal{B}_p)$?

Drinfeld’s construction works for any GL_n over any local field L , replacing Ω by the complement ${}_L\Omega^{(n-1)}$ of L -rational hyperplanes in \mathbb{P}^{n-1} .

6.4. De Jong’s viewpoint on p -adic period mappings.

J. de Jong [dJ95b] has proposed an interpretation of the period mapping which clarifies a lot the nature of its fibers. We briefly present this interpretation under the following mild simplifying assumption:

- * that there is a family of affinoid domains (X_i) of \mathcal{M}^0 such that the $\mathcal{P}(X_i)$ (which are finite unions of affinoid domains in \mathcal{D} since \mathcal{P} is étale) form an admissible covering of \mathcal{D} .

This assumption is fulfilled in our three basic examples (but not always, cf. [RZ96, 5.53]).

Under (*), De Jong shows (*loc. cit.* ⁽⁵⁾, intr. and 7.2) that there is an étale local system of \mathbb{Q}_p -spaces \mathcal{V} on \mathcal{D} such that \mathcal{M}^0 is a component of the space of \mathbb{Z}_p -lattices in \mathcal{V} . Here, an “étale local system of \mathbb{Q}_p -spaces” is given by the data

$$\mathcal{V} = (\{U_i \hookrightarrow \mathcal{D}\}, \mathcal{V}_i, \phi_{ij})$$

where $U_i \hookrightarrow \mathcal{D}$ are open immersions, $\mathcal{V}_i = \varprojlim_n \mathcal{V}_i/p^n\mathcal{V}_i$ are étale local systems of \mathbb{Z}_p -lattices (each $\mathcal{V}_i/p^n\mathcal{V}_i$ being a finite locally free sheaf of $\mathbb{Z}/p^n\mathbb{Z}$ -modules on the étale site of U_i), and $\phi_{ij} : (\mathcal{V}_i \otimes \mathbb{Q}_p)|_{U_i \times_{\mathcal{D}} U_j} \rightarrow (\mathcal{V}_j \otimes \mathbb{Q}_p)|_{U_i \times_{\mathcal{D}} U_j}$ are isomorphisms satisfying the usual cocycle condition (cf. [dJ95b, 4]).

Of course, in the “decorated case”, \mathcal{V} inherits the relevant decoration.

It follows from this interpretation that in example 4.1.2, the fibers of the period mapping are in natural bijection with the set of vertices of the Bruhat-Tits tree \mathcal{T} .

Similarly, in example 4.3.4, one can see that the fibers of \mathcal{P} are in bijection with the one-point set $SL_1(B_p)/SL_1(\mathcal{B}_p)$, cf. 7.4.4.

⁽⁵⁾of course $p = \ell$ throughout §§6 and 7 of *loc. cit.*

7. p -adic uniformization of Shimura varieties.

ABSTRACT: Application of the theory of p -adic period mappings: Rapoport-Zink's p -adic uniformization theorem. The case of global uniformization. Čerednik-Drinfeld uniformization of Shimura curves and other examples.

7.1. p -integral models of Shimura varieties.

7.1.1. Let $\mathcal{S}h$ be a Shimura variety of PEL type as in 1.1.4. We recall the setting of 1.1.4: B is a simple finite-dimensional \mathbb{Q} -algebra, with a positive involution $*$, V is a B -module of finite type, endowed with an alternate \mathbb{Q} -bilinear form such that $\langle bv, w \rangle = \langle v, b^*w \rangle$; G denotes the \mathbb{Q} -group of B -linear symplectic similitudes of V ; \mathcal{B} is a maximal order in B stable under $*$, and L is a lattice in V , stable under \mathcal{B} and autodual for $\langle \cdot, \cdot \rangle$.

The (non-connected) Shimura variety $\mathcal{S}h$ is defined over the reflex field E (which is a number field in \mathbb{C}). There are variants of $\mathcal{S}h$ in which the principal congruence subgroup of level N of $G(\hat{\mathbb{Z}})$ is replaced by any open compact subgroup C of $G(\mathbb{A}_f)$, where $\mathbb{A}_f = \hat{\mathbb{Z}} \otimes \mathbb{Q}$ is the ring of finite adeles of \mathbb{Q} . The complex points of $\mathcal{S}h$ admit the following adelic description

$$\mathcal{S}h(\mathbb{C}) = C \backslash ((G(\mathbb{R})^{\text{ad}} \cdot h_0) \times G(\mathbb{A}_f)) / G(\mathbb{Q})$$

(cf. 1.2.4 about the one-parameter group h_0).

7.1.2. We set $B_p = B \otimes_{\mathbb{Q}} \mathbb{Q}_p$, $\mathcal{B}_p = \mathcal{B} \otimes_{\mathbb{Z}} \mathbb{Z}_p$, and so on. We denote by C_p the stabilizer of L_p in $G(\mathbb{Q}_p)$

We assume that C is of the form $C^p C_p$ for some open compact subgroup C^p of $G(\mathbb{A}_f^p)$, where $\mathbb{A}_f^p = (\prod_{\ell \neq p} \mathbb{Z}_{\ell}) \otimes \mathbb{Q}$.

In the case of a congruence subgroup of level N , this condition means that *the level is prime to p* .

We fix an embedding $v : E \hookrightarrow \mathbb{C}_p$, denote by E_v the v -completion of E , by \mathfrak{v} the discrete valuation ring $\widehat{\mathbb{Z}}^{\text{ur}} \cdot \mathcal{O}_{E_v} = \widehat{\mathbb{Z}}^{\text{ur}} \cdot \mathcal{O}_E$ (compositum in \mathbb{C}_p), and by K the fraction field of \mathfrak{v} .

7.1.3. Let S be a scheme. The objects of the the category of S -abelian schemes up to prime to p isogeny are the abelian schemes over S ; morphisms between A and A' are global sections of $\underline{\text{Hom}}_S(A, A') \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$, where $\mathbb{Z}_{(p)}$ is the ring of rational numbers with denominator prime to p . This is a $\mathbb{Z}_{(p)}$ -linear category. Its relation to p -divisible groups is given by the following elementary lemma:

Lemma 7.1.4. *The faithful functor $A \mapsto A[p^{\infty}]$*

*$\{S\text{-abelian schemes up to prime to } p \text{ isogeny}\} \rightarrow \{p\text{-divisible groups over } S\}$
is conservative, i.e. reflects isomorphisms.*

The following result (which we state in a rather vague manner) is due to R. Kottwitz [Kot92] (Compare with 1.2.4.).

Proposition 7.1.5. *For sufficiently small C^p , there is a moduli scheme Sh over \mathcal{O}_{E_v} for isomorphism classes of abelian varieties A up to prime to p isogeny, with \mathcal{B} -action (with Shimura type condition), together with a \mathbb{Q} -homogeneous principal $*$ -polarization and a class of B -linear symplectic similitudes $H_{\text{et}}^1(A, \mathbb{A}_f^p) \rightarrow \mathbb{A}_f^p \otimes V$ modulo the group C^p . The generic fiber $Sh \otimes_{\mathcal{O}_{E_v}} E_v$ is a finite sum of copies of $Sh \otimes_E E_v$.*

In the case of a congruence subgroup of level N , this condition that C^p is sufficiently small means that N is sufficiently large (and kept prime to p).

7.2. The uniformization theorem.

7.2.1. One fixes a point of $Sh(\overline{\mathbb{F}}_p)$, hence a $\overline{\mathbb{F}}_p$ -abelian variety \overline{A} up to prime to p isogeny with \mathcal{B} -action, principal $*$ -polarization, and level structure. We consider the p -divisible group $\overline{\Lambda} = \overline{A}[p^\infty]$ (with \mathcal{B}_p -action, principal $*$ -polarization, level structure). We denote by \overline{G} the \mathbb{Q} -group of self-quasi-isogenies of \overline{A} respecting the additional structure. It is clear that $\overline{G}(\mathbb{Q}_p) \subset J$; on the other hand $\overline{G}(\mathbb{A}_f^p) \subset G(\mathbb{A}_f^p)$.

We also consider an auxiliary lifting $\tilde{\overline{A}}$ of \overline{A} (with \mathcal{B} -action) over some finite extension $\mathcal{O}_{E'}$ of \mathcal{O}_E in \mathfrak{v} , and set $\tilde{\overline{\Lambda}} = \tilde{\overline{A}}[p^\infty]$. We can then identify $\mathbf{D}(\overline{\Lambda}) \otimes_W K$ with $H_{\text{DR}}^1(\tilde{\overline{A}}) \otimes_{\mathcal{O}_{E'}} K$, and the submodule $\omega_{\tilde{\overline{\Lambda}}}$ with $F^1 H_{\text{DR}}^1(\tilde{\overline{A}}) \otimes_{\mathcal{O}_{E'}} K$. Thus if the complex lagrangian space F_0^1 of 1.1.4 is taken to be $F^1 H_{\text{DR}}^1(\tilde{\overline{A}}) \otimes_{\mathcal{O}_{E'}} \mathbb{C}$ (for some complex embedding of E' extending the natural complex embedding of E), the complex and p -adic Shimura type conditions “agree” (1.1.4, 4.3.2). We can consider the corresponding decorated moduli problem for p -divisible groups, and the formal moduli scheme \mathfrak{M} over \mathfrak{v} as in 4.3. Moreover, the flags spaces \mathcal{D}^\vee considered in the complex and p -adic situations agree, *i.e.* come from the same flag space defined over the reflex field E .

7.2.2. One defines a morphism of functors on $\text{Nil}_{\mathfrak{v}}$:

$$\mathfrak{M} \rightarrow Sh$$

as follows [RZ96, 6.14]. Let S be in $\text{Nil}_{\mathfrak{v}}$ and let us consider a pair $(\Lambda, \rho) \in \mathfrak{M}(S)$. By rigidity of p -divisible groups up to isogeny (2.2.3), the \mathcal{B}_p -quasi-isogeny $\rho \in \text{qisog}(\Lambda \times_S S_{\text{red}}, \overline{\Lambda} \times_{\overline{\mathbb{F}}_p} S_{\text{red}})$ lifts to a unique \mathcal{B}_p -quasi-isogeny $\tilde{\rho} \in \text{qisog}(\Lambda, \tilde{\overline{\Lambda}} \times_{\mathfrak{v}} S)$. This can be algebraized in a weak sense: there is an S -abelian scheme $A_{(\Lambda, \rho)}$ up to prime to p isogeny, with \mathcal{B} -action and with p -divisible group Λ , and a quasi-isogeny $A_{(\Lambda, \rho)} \rightarrow \tilde{\overline{A}} \times_{\mathcal{O}_{E'}} S$ which induces $\tilde{\rho}$ at the level of p -divisible groups. Moreover $A_{(\Lambda, \rho)}$ is unique in the category of S -abelian schemes up to prime to p isogeny with \mathcal{B} -action (7.1.4), and its formation is functorial in (Λ, ρ) . One then defines the required morphism of functors on $\mathfrak{M}(S)$ by setting $(\Lambda, \rho) \mapsto A_{(\Lambda, \rho)}$ (endowed with the polarization and level structure inherited from \overline{A}).

This defines a morphism of formal schemes, hence a morphism of associated analytic spaces over \mathbb{C}_p :

$$\mathcal{Q} : \mathcal{M} \rightarrow \mathcal{S}h_{\mathbb{C}_p}^{\text{an}}.$$

7.2.3. It turns out that the decorated abelian varieties (up to prime to p isogeny) which are *isogenous* to \overline{A} form in a natural way a *countable union* $Z = \bigcup Z_n$ of *projective irreducible subvarieties* Z_n of $\mathcal{S}h_{\overline{\mathbb{F}}_p}$, each of them meeting only finitely many members of the family ([RZ96, 6.23, 6.34]). One can slightly generalize the notion of tubes (3.5) and define the tube $]Z[_\subset \mathcal{S}h_{\mathbb{C}_p}^{\text{an}}$. We denote by S the connected component of $]Z[_$ which contains the modular point s of our chosen lifting \tilde{A} .

Theorem 7.2.4. *For sufficiently small C^p , there is a connected component \mathcal{M}^0 of \mathcal{M} such that the restriction $\mathcal{Q} : \mathcal{M}^0 \rightarrow S$ is a topological covering of S . More precisely, S is the (right) quotient of \mathcal{M}^0 by some torsion-free discrete subgroup of J .*

cf. [RZ96, 6.23, 6.31]. Roughly speaking, the proof of 7.2.4 involves two steps: using the Serre-Tate theorem 2.2.4, one shows that the morphism of \mathfrak{v} -formal schemes $\mathfrak{M} \rightarrow \widehat{\mathcal{S}h}$ defined above is formally etale. One concludes by a careful study of this morphism on geometric points.

In our examples 4.2, S is an open unit disc, and the map \mathcal{Q} is an isomorphism (in 4.2.1, it is induced by $q \mapsto z = \lambda$).

7.2.5. Let us consider the universal decorated abelian scheme \underline{A} over $\mathcal{S}h$, and the vector bundle $\mathcal{H} = H_{\text{DR}}^1(\underline{A}/\mathcal{S}h)$ with its Gauss-Manin connection $\nabla_{\text{GM}} : \mathcal{H} \rightarrow \Omega_{\mathcal{S}h}^1 \otimes \mathcal{H}$.

Corollary 7.2.6. *Viewed as a p -adic connection and restricted to S , the Gauss-Manin connection $\nabla_{\text{GM}|_S}$ comes from a representation of $\pi_1^{\text{top}}(S, s)$ on $V^{\mathbb{C}_p}$ (hence it satisfies Cauchy's theorem, cf. I.1.5).*

PROOF. Let \tilde{S} be the universal covering of S . According to I.1.5), we have to show that the pull-back of ∇_{GM} on \tilde{S} is trivial ($\cong V^{\mathbb{C}_p} \otimes \mathcal{O}_{\tilde{S}}$). By 7.2.4 and 6.2.5, this already holds on the quotient \mathcal{M}^0 of \tilde{S} . \square

Remark. Using the Weil descent data of [RZ96, 6.21], one can show that the representation is defined over a finite unramified extension E'_v of E_v , i.e. comes from a representation of $\pi_1^{\text{top}}(S, s)$ on $V \otimes E'_v$.

7.2.7. The assumption $C = C^p C_p$ is necessary to define the morphism $\mathcal{Q} : \mathcal{M}^0 \rightarrow S$. The assumption “ C^p small enough” is much less important. Theorem 7.2.4. holds without it, except that the discrete subgroup of J is then no longer torsion-free [the point is that there is always a normal subgroup C'^p of finite index which is small enough, one applies the theorem to C'^p , and one passes to the quotient by C'^p/C^p]. However, to give a modular interpretation of the (ramified) quotient of \mathcal{M}^0 in this case, one needs

the notion of p -adic orbifold (to be discussed in III); moreover the Gauss-Manin connection may acquire logarithmic singularities along the branched locus.

7.3. Global uniformization

We say that there is *global uniformization* in case S is (the analytification of) a whole connected component of the Shimura variety Sh . This occurs when the reduction mod p of the decorated abelian varieties A parametrized by Sh form a *single isogeny class*.

In the situation of global uniformization, the group \overline{G} of self-quasi-isogenies of \overline{A} (with decoration) is an *inner form of G* .

In view of our analogy 4.3.5, this situation can be expected to be closest to the complex case (in the notation of 1.2.4, the \hat{A} 's are all isomorphic to \hat{U}). Indeed, we then get a commutative diagram of p -adic manifolds (similar to 1.2.4):

$$\begin{array}{ccc} \mathcal{M}^0 & \xrightarrow{\mathcal{P}} & \mathcal{D} \subset \mathcal{D}^\vee \\ \mathcal{Q} \downarrow & & \downarrow \mathcal{Q} \\ S & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma. \end{array}$$

where the horizontal maps are étale, \mathcal{Q} denotes the quotient maps, and Γ is an arithmetic subgroup in the semi-simple group \overline{G}^{ad} . In all known examples, \mathcal{P} is actually an isomorphism and $\mathcal{M}^0 = \hat{S}$ [RZ96, 6], [BZ95], [Var98a], [Var98b].

Here, the archimedean—non archimedean correspondence goes beyond a mere analogy: indeed, here and in 1.2.4, what is denoted by S (*resp.* \mathcal{D}^\vee) is the — complex or p -adic — analytification of the same algebraic variety.

Remark. To avoid any ambiguity, let us point out that we consider here only phenomena of global uniformization which can be interpreted in terms of moduli spaces of p -divisible groups. This does not account for all cases of global p -adic uniformization of Shimura varieties. For instance, it is known that the elliptic modular curve $X_0(p)$ is a Mumford curve over \mathbb{Q}_p , whereas the p -adic uniformization of 7.2.4 (or variants with level p structure) is local.

7.4. Example: the Čerednik-Drinfeld uniformization of Shimura curves at a critical prime.

7.4.1. The first historical example of global uniformization in the above sense is the Čerednik uniformization of “Shimura curves” (modular curves for fake elliptic curves) at a critical prime p , *cf.* [Čer76]. Čerednik’s method was of group-theoretic nature and did not involve formal groups. A modular proof was subsequently proposed by Drinfeld [Dri76] (*cf.* also [BC91, III]). For examples, and related topics, we refer to [vdP89], [vdP92a], [vdP92b].

7.4.2. We fix an indefinite quaternion algebra B over \mathbb{Q} , a maximal order \mathcal{B} in B , and a positive involution $*$ of \mathcal{B} . We consider the standard representation $V = B$ of B , so that $G = GL_B(V) \cong B^\times$.

We refer to the foundational paper [Shi67] for the properties of the (connected) Shimura curve $\mathcal{X}^+(N)$ of level N attached to B . It parametrizes principally $*$ -polarized abelian surfaces with multiplication by \mathcal{B} (fake elliptic curves, *cf.* 4.3.4), with level N structure. We recall the following properties of $\mathcal{X}^+(N)$, which actually characterize it (*loc. cit.* 3.2):

- (a) $\mathcal{X}^+(N)$ is a projective geometrically connected smooth curve defined over the cyclotomic field $\mathbb{Q}(\zeta_N)$,
- (b) $\mathcal{X}^+(N)(\mathbb{C}) = \mathfrak{h}/\Gamma^+(N)$, that is to say: $\mathcal{X}^+(N)$ is the quotient of the Poincaré upper half plane by the image $\Gamma^+(N) = PSL_1(\mathcal{B})(N)$ of the group $\{g \in (1 + N\mathcal{B})^\times, \text{Nr}(g) > 0\}$ in $PSL_2(\mathbb{R})$, where Nr stands for the reduced norm,
- (c) for any imaginary quadratic field K in B such that $\mathcal{O}_K \subset \mathcal{B}$, let z be its fixed point in \mathfrak{h} . Then the extension of $\mathbb{Q}(\zeta_N)$ generated by the image of z in $\mathcal{X}^+(N)$ is the class field over K with conductor $N\mathcal{B} \cap \mathcal{O}_K$.

The commutative diagram of 1.2.4, involving the complex period mapping, specializes to the following

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\mathcal{P}} & \mathcal{D} = \mathfrak{h} \subset \mathcal{D}^\vee = \mathbb{P}^1 \\ \mathfrak{q} \downarrow & & \downarrow \mathfrak{q} \\ S = \mathcal{X}^+(N)_{\mathbb{C}}^{\text{an}} & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma^+(N) \end{array}$$

where the horizontal maps are isomorphisms (and the vertical maps are topological coverings if $\Gamma_p^+(N)$ is torsion-free⁽⁶⁾).

There is a useful variant \mathcal{X}^* of $\mathcal{X}^+ = \mathcal{X}^+(1)$, already considered by Shimura ([Shi67, 3.13], in which Γ^+ is replaced by the image Γ^* in $PSL_2(\mathbb{R})$ of the group $\{g \in B^\times \mid g\mathcal{B} = \mathcal{B}g, \text{Nr}(g) > 0\}$:

$$\mathcal{X}^*(\mathbb{C}) = \mathfrak{h}/\Gamma^*.$$

It is defined over \mathbb{Q} . More generally, there is a Shimura curve \mathcal{X}_Γ attached to any congruence subgroup Γ in $B^{\times+}/\mathbb{Q}^\times$, where $B^{\times+}$ denotes the group of elements of positive reduced norm.

7.4.3. It is known that $\mathcal{X}^+(N)$ has good reduction at any prime p which does not divide the discriminant $d(B)$ of B nor the level N (Y. Morita). If instead p is critical (but still *does not divide* N), as we shall assume henceforth, Čerednik's theorem shows that $\mathcal{X}^+(N)_{\mathbb{C}_p}$ is a *Mumford curve* uniformized by the Drinfeld space $\Omega_{\mathbb{C}_p} = \mathbb{P}_{\mathbb{C}_p}^1 \setminus \mathbb{P}^1(\mathbb{Q}_p)$ (at least if N is large

⁽⁶⁾if this group is not torsion-free \tilde{S} is the universal covering in the sense of orbifolds.

enough); in fact, $\mathcal{S}h$ becomes a Mumford curve over some finite unramified extension of \mathbb{Q}_p (note that $\mathbb{Q}(\zeta_N) \subset \mathbb{Q}_p^{\text{ur}}$).

To be more explicit, let \overline{B} be the definite quaternion \mathbb{Q} -algebra which is ramified at the same primes as B except p : $\overline{B}_p \cong M_2(\mathbb{Q}_p)$. Let $\overline{\mathcal{B}}$ be a maximal order in \overline{B} . Let $\Gamma_p^+(N)$ be the image in $PGL_2(\mathbb{Q}_p)$ of the group $\{g \in (1 + N\overline{\mathcal{B}}[\frac{1}{p}])^\times \mid \text{ord}_p(\text{Nr}(g)) \text{ even}\}$. Let Γ_p^* be the image in $PGL_2(\mathbb{Q}_p)$ of the normalizer $\{g \in \overline{B}^\times \mid g\overline{\mathcal{B}}[\frac{1}{p}] = \overline{\mathcal{B}}[\frac{1}{p}]g\}$. These are discrete subgroups, and one has the Čerednik uniformization:

$$\begin{aligned}\mathcal{X}^+(N)_{\mathbb{C}_p}^{\text{an}} &\cong \Omega_{\mathbb{C}_p}/\Gamma_p^+(N), \\ \mathcal{X}^*_{\mathbb{C}_p} &\cong \Omega_{\mathbb{C}_p}/\Gamma_p^*.\end{aligned}$$

For recent applications to p -adic L -functions, Heegner points, etc... cf. [BD98].

7.4.4. The case of \mathcal{X}^* is the one originally considered in [Čer76], except that Čerednik uses the adelic language. In order to make the translation, it is useful to keep in mind the following few facts (cf. [Vig80, pp. 40, 99 and passim]).

For critical p , the unique maximal order \mathcal{B}_p of B_p is $\{b \in B_p \mid \text{Nr}(b) \in \mathbb{Z}_p\}$. Hence, we have an equality $SL_1(\mathcal{B}_p) = SL_1(B_p)$ and a chain of compact normal subgroups

$$PSL_1(\mathcal{B}_p) = PSL_1(B_p) \subset PGL_1(\mathcal{B}_p) \subset PGL_1(B_p)$$

with $PGL_1(B_p)/PSL_1(\mathcal{B}_p) \cong \mathbb{Q}_p^\times/(\mathbb{Q}_p^\times)^2$ (which is of type $(2, 2)$ if $p \neq 2$, of type $(2, 2, 2)$ if $p = 2$).

On the other hand, the normalizer of \mathcal{B}_ℓ in B_ℓ^\times is B_ℓ^\times if ℓ is critical, and $\mathcal{B}_\ell^\times \cdot \mathbb{Q}_\ell^\times$ otherwise. It follows that

$$\begin{aligned}\Gamma^* &= \{g \in B^{\times+} \mid \forall \ell \nmid d(B), g \in \mathcal{B}_\ell^\times \cdot \mathbb{Q}_\ell^\times\}/\mathbb{Q}^\times \\ \Gamma_p^* &= \{g \in \overline{B}^\times \mid \forall \ell \nmid d(B), g \in \overline{\mathcal{B}}_\ell^\times \cdot \mathbb{Q}_\ell^\times\}/\mathbb{Q}^\times \\ \Gamma^+(N) &= \{g \in B^{\times+} \mid \forall \ell, g \in (1 + N\mathcal{B}_\ell)^\times \cdot \mathbb{Q}_\ell^\times\}/\mathbb{Q}^\times \\ \Gamma_p^+(N) &= \left\{g \in \overline{B}^\times \mid \begin{array}{l} \text{ord}_p(\text{Nr}(g)) \text{ even;} \\ \forall \ell \neq p, g \in (1 + N\overline{\mathcal{B}}_\ell)^\times \cdot \mathbb{Q}_\ell^\times \end{array} \right\}/\mathbb{Q}^\times\end{aligned}$$

If N is prime to $d(B)$, this can also be rewritten

$$\begin{aligned}\Gamma^+(N) &= \left\{g \in B^{\times+} \mid \begin{array}{l} \forall \ell \mid d(B), \text{ord}_\ell(\text{Nr}(g)) \text{ even;} \\ \forall \ell \nmid d(B), g \in (1 + N\mathcal{B}_\ell)^\times \cdot \mathbb{Q}_\ell^\times \end{array} \right\}/\mathbb{Q}^\times \\ \Gamma_p^+(N) &= \left\{g \in \overline{B}^\times \mid \begin{array}{l} \forall \ell \mid d(B), \text{ord}_\ell(\text{Nr}(g)) \text{ even;} \\ \forall \ell \nmid d(B), g \in (1 + N\overline{\mathcal{B}}_\ell)^\times \cdot \mathbb{Q}_\ell^\times \end{array} \right\}/\mathbb{Q}^\times\end{aligned}$$

In particular, for $N = 1$, we see that there are canonical bijections

$$\Gamma^*/\Gamma^+ = \Gamma_p^*/\Gamma_p^+ = (\mathbb{Z}/2\mathbb{Z})^{\#\{\ell \mid d(B)\}}$$

7.4.5. We now give some indications on Drinfeld’s modular approach to Čerednik’s uniformization.

Lemma 7.4.6. (i) *Fake elliptic curves have potentially good reduction at every prime; the reduction is supersingular reduction at the critical prime p . In particular, over $\overline{\mathbb{F}}_p$, they form a unique isogeny class.*
(ii) *Fake elliptic curves have a unique principal $*$ -polarization.*

PROOF. (i): due to the presence of many endomorphisms, the potential good reduction follows easily from the semi-stable reduction theorem for abelian schemes; also, the non-split quaternion algebra B_p acts on the p -divisible group up to isogeny, and this forces the slopes to be $1/2$; for details and (ii), cf. [BC91, III]. \square

It follows from (i) that any fake elliptic curve over $\overline{\mathbb{F}}_p$ is isogenous to the square of a supersingular elliptic curve \overline{E} . However, while there are only finitely many supersingular elliptic curves up to isomorphism, fake elliptic curves over $\overline{\mathbb{F}}_p$ have continuous moduli! The moduli spaces are chains of \mathbb{P}^1 's.⁽⁷⁾

Let us introduce the quaternion \mathbb{Q} -algebra D ramified only at p and ∞ : $D \cong \text{End}^{\mathbb{Q}}(\overline{E})$. Let \overline{A} be a fake elliptic curve over $\overline{\mathbb{F}}_p$. It follows from 7.4.6 (i) that $\text{End}^{\mathbb{Q}}(\overline{A}) \cong M_2(D)$, and $\text{End}_{\mathcal{B}}^{\mathbb{Q}}(\overline{A}) \cong \overline{B}$.

In particular, the inner form \overline{G} of G considered in 7.3 is \overline{B}^{\times} , and $\overline{G}(\mathbb{Q}_p) = GL_2(\mathbb{Q}_p) = J$.

It follows from 7.4.6 (ii) that we may disregard polarizations. The p -divisible groups attached to fake elliptic curves with \mathcal{B} -multiplication are special formal \mathcal{B}_p -modules of height 4; recall from 4.3.4 the corresponding formal moduli space $\mathfrak{M} = \coprod_{\mathbb{Z}} \mathfrak{M}^0$. The period mapping is an isomorphism

(6.3.4, 6.4), and the commutative diagram of 7.3 specializes to

$$\begin{array}{ccc} \tilde{S} = \mathcal{M}^0 & \xrightarrow{\mathcal{P}} & \mathcal{D} = \Omega_{\mathbb{C}_p} \subset \mathcal{D}^{\vee} = \mathbb{P}^1 \\ \varrho \downarrow & & \downarrow \varrho \\ S = \mathcal{X}^+(N)_{\mathbb{C}_p}^{\text{an}} & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma_p^+(N) \end{array}$$

where the horizontal maps are isomorphisms (and the vertical maps are topological coverings if $\Gamma_p^+(N)$ is torsion-free), in complete analogy with 7.2.

7.4.7. Let us end this section by considering the case where p divides the level N : $N = p^n \cdot M$, $p \nmid M$.

⁽⁷⁾ Another curious feature of the higher dimensional case is that the product of $n > 1$ supersingular elliptic curves over $\overline{\mathbb{F}}_p$ form a single isomorphism class (Deligne, Ogus)

In this case, $\Gamma_p^+(N) = \Gamma_p^+(M)$. The finite étale covering $\mathcal{X}^+(N)_{\mathbb{C}_p}^{\text{an}} \rightarrow \mathcal{X}^+(M)_{\mathbb{C}_p}^{\text{an}}$ is not a topological covering anymore (even if $\Gamma_p^+(M)$ is torsion-free). As was shown by Drinfeld, it is hidden in the finite étale covering Υ^n of $\Omega_{\mathbb{C}_p}$ (6.3.5).

Let us assume for simplicity that $M > 2$, so that $-1 \notin \Gamma^+(M)$. This implies that $\Gamma^+(N)/\Gamma^+(M) \cong SL_1(\mathcal{B}/p^n\mathcal{B})$. Then Υ^n is a topological covering of $\mathcal{X}^+(N)_{\mathbb{C}_p}^{\text{an}}$, and more precisely, one has the commutative diagram

$$\begin{array}{ccc} \mathcal{X}^+(N)_{\mathbb{C}_p}^{\text{an}} & \xrightarrow{\sim} & \Upsilon^n/\Gamma_p^+(M) \\ \downarrow & & \downarrow \\ \mathcal{X}^+(M)_{\mathbb{C}_p}^{\text{an}} & \xrightarrow{\mathcal{P}} & \Omega_{\mathbb{C}_p}/\Gamma_p^+(M) \end{array}$$

where the vertical maps are finite étale Galois covering maps with group $SL_1(\mathcal{B}/p^n\mathcal{B})$, cf. [BC91, III.5.5], [Var98a]. Similarly, there is such a p -adic global uniformization for any Shimura curve \mathcal{X}_Γ (cf. 7.4.1).

7.5. Other examples of global p -adic uniformization.

7.5.1. Shimura's paper [Shi67] deals not only with quaternion algebra over \mathbb{Q} , but also with quaternion algebras B/E over a totally real number field E . When B splits at a single real place, he constructs curves $\mathcal{X}^+(N)$ and \mathcal{X}^* satisfying properties similar to (a),(b),(c) of 7.4.2.

In this context, N may be an arbitrary ideal of \mathcal{O}_E , $\mathbb{Q}(\zeta_N)$ has to be replaced by the ray class field with conductor $N\cdot\infty$, and the condition $N(g) > 0$ means that the reduced norm of g is totally positive, or, what amounts to the same, is positive at the split real place ∞_0 .

The novelty, however, is that these Shimura curves have *no direct interpretation* as moduli spaces for decorated abelian varieties, since B is not in the Albert list of endomorphism algebras of abelian varieties when $[E : \mathbb{Q}] > 1$.

To overcome this problem, Shimura introduces the following remarkable trick. Let K be a totally imaginary quadratic extension of E . Then $B^\bullet = B \otimes_E K$ is in Albert's list (we endow it with the involution which is the tensor product of $*$ on B and complex conjugation on K). It turns out that the Shimura varieties of PEL type for abelian varieties of dimension $g = 4[E : \mathbb{Q}]$ ($*$ -polarized, with level structure) with complex multiplication by B^\bullet are twisted forms of the same curve. The Shimura curve attached to B is obtained by Weil descent from the Shimura curves attached to $B \otimes_E K$, for sufficiently many quadratic extensions K .

7.5.2. Čerednik's original p -adic uniformization theorem already applies to Shimura curves attached to quaternion algebras over a totally real number field E . His method has been further developed in [Var98a], [Var98b]. On

the other hand, Drinfeld’s modular approach has been extended by Boutot-Zink [BZ95], using Shimura’s trick. Let us sketch the principle of the construction.

Let G be the multiplicative group of B^\times , viewed as an algebraic group over \mathbb{Q} . Let Γ be a congruence subgroup of $G^{\text{ad}}(\mathbb{Q})$ (alternatively, one can consider an open compact subgroup C of $G(\mathbb{A}_f)$). Let us first assume that Γ is *maximal at p* (alternatively C is of the form $C^p C_p$ for some open compact subgroup C^p of $G(\mathbb{A}_f^p)$, cf. 7.1.2). The (non-connected) Shimura curve Sh given by

$$Sh_{\mathbb{C}}^{\text{an}} = C \backslash ((\mathbb{P}_{\mathbb{C}}^1 \setminus \mathbb{P}^1(\mathbb{R})) \times G(\mathbb{A}_f)) / G(\mathbb{Q})$$

is defined over the reflex field, which is E (embedded into \mathbb{R} via ∞_0). Over some finite abelian extension E'/E , it decomposes as a disjoint union of geometrically connected Shimura curves \mathcal{X}_{Γ} (\mathfrak{h}/Γ over \mathbb{C} ; one possible Γ is the image of $G(\mathbb{Q}) \cap C$ in $G^{\text{ad}}(\mathbb{Q})$).

Let $v = v_0$ be a *critical* finite place of E , *i.e.* such that B_v is a non-split quaternion algebra. Let p be the residue characteristic of v_0 and let v_1, \dots, v_m be the other places of E above p .

Let K be a totally imaginary quadratic extension of E , such that every place v_i splits in K . We denote by w_i, \bar{w}_i the two places of K above v_i . The quaternion algebra $B^\bullet = B \otimes_E K$ is ramified at $w = w_0$: $B_w^\bullet \cong B_v$.

We fix an extension $\infty_0 : K \hookrightarrow \mathbb{C}$ of the real embedding ∞_0 of E , and a double embedding $\mathbb{C} \hookrightarrow \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ whose restriction to K is compatible with (∞_0, w) . The embeddings $K \hookrightarrow \mathbb{C}_p$ which factor through some w_i but not through some \bar{w}_i form a CM type Ψ : $\Psi \coprod \bar{\Psi} = \text{Hom}(K, \mathbb{C}_p)$ (identified with $\text{Hom}(K, \mathbb{C})$ through our chosen double embedding).

There is a natural \mathbb{Q} -group G^\bullet such that $G^\bullet(\mathbb{Q}) = \{b \in B^\bullet \mid bb^* \in E\}$, acting on $V^\bullet = V \otimes_E K$. One can attach to C an open compact subgroup C^\bullet of $G^\bullet(\mathbb{A}_f)$, maximal at p , and a Shimura curve Sh^\bullet such that Sh is open and closed in Sh^\bullet ([BZ95, 3.11]). Alternatively, one can attach to Γ a congruence subgroup Γ^\bullet in $G^{\bullet \text{ad}}(\mathbb{Q})$ and a corresponding connected Shimura curve $\mathcal{X}_{\Gamma^\bullet}$ which is a twisted form of \mathcal{X}_{Γ} (defined over some abelian extension of K). The point is that this new Shimura curve Sh^\bullet is of PEL type. More precisely, for C small enough, it is a moduli space for $*$ -polarized abelian varieties of dimension $g = 4[E : \mathbb{Q}]$ with multiplication by $\mathcal{B}^\bullet = \mathcal{B} \otimes_{\mathcal{O}_E} \mathcal{O}_K$, Shimura type Ψ , and level structure. As in 7.1.5, it admits a model Sh^\bullet over \mathcal{O}_{E_v} .

Let $\underline{A} \rightarrow Sh^\bullet$ be the universal abelian scheme up to prime to p isogeny. Due to the \mathcal{O}_K -action, the p -divisible group of \underline{A} splits as $\underline{A}[p^\infty] \cong \prod_i \Lambda_{w_i} \times \Lambda_{\bar{w}_i}$, where Λ_{w_i} and $\Lambda_{\bar{w}_i}$ are p -divisible groups of height $4[E_{v_i} : \mathbb{Q}_p]$ with formal multiplication by E_{v_i} . Moreover, $\Lambda_w = \Lambda_{w_0}$ is a special formal \mathcal{B}_v -module of height $4.[E_v : \mathbb{Q}_p]$ (defined as in 4.3.4, replacing \mathbb{Q}_p by E_v), and $\Lambda_{\bar{w}}$ is the Serre dual of Λ_w ; the other Λ_{w_i} , $i > 0$, are etale, and the $\Lambda_{\bar{w}_i}$ are

their respective Serre duals. Hence the Newton polygon of the Dieudonné module of $\underline{A}[p^\infty]$ has slopes $0, \frac{1}{2}, 1$ (only $\frac{1}{2}$ if $m = 0$).




FIGURE 2

Let \mathfrak{v} be the discrete valuation ring $\widehat{\mathbb{Z}^{\text{ur}}}\mathcal{O}_{E_v}$ as in 7.1.2. The moduli problem for \mathcal{B}_v -module of height $4.[E_v : \mathbb{Q}_p]$ is representable by the p -adic formal scheme $\prod_{\mathbb{Z}} (E_v \hat{\Omega})_{\mathfrak{v}}$, where $E_v \hat{\Omega}$ is the analogue for (and over) \mathcal{O}_{E_v} of the formal scheme $\hat{\Omega}$, with associated analytic space $(E_v \Omega)_{C_p} = \mathbb{P}_{C_p}^1 \setminus \mathbb{P}^1(E_v)$, cf. 6.3.5.

7.5.3. This is the modular way taken by Boutot and Zink to reprove (and strengthen) Čerednik’s theorem. As was remarked by Varshavsky [Var98b, 3.13], one obtains a simpler adelic formulation if one works with a twisted version ${}^{\text{ad}}Sh$ of Sh (corresponding to a different normalization of h_0 in 1.2.4); this does not change the adelic description of the associated complex space

$${}^{\text{ad}}Sh_{\mathbb{C}}^{\text{an}} = C \setminus ((\mathbb{P}_{\mathbb{C}}^1 \setminus \mathbb{P}^1(\mathbb{R})) \times G(\mathbb{A}_f)) / G(\mathbb{Q}),$$

but makes the analogy with the following description of the associated p -adic space more striking.

Let \overline{B} be the quaternion algebra over E obtained from B by interchanging the invariants at ∞_0, v , and let \overline{G} be the \mathbb{Q} -algebraic group attached to \overline{B}^{\times} . Let us identify $\overline{G}(\mathbb{A}_f^p)$ with $G(\mathbb{A}_f^p)$ via an anti-isomorphism $\overline{B} \otimes_E \mathbb{A}_f^p \rightarrow B \otimes_E \mathbb{A}_f^p$, and set $\overline{C} = GL_2(E_v)C^p$ viewed as a subgroup of $\overline{G}(\mathbb{A}_f)$. Then

$$\begin{aligned} {}^{\text{ad}}Sh_{C_p}^{\text{an}} &= C^p \setminus ((E_v \Omega)_{C_p} \times G(\mathbb{A}_f^p)) / \overline{G}(\mathbb{Q}) \\ &= \overline{C} \setminus ((\mathbb{P}_{C_p}^1 \setminus \mathbb{P}^1(E_v)) \times \overline{G}(\mathbb{A}_f)) / \overline{G}(\mathbb{Q}) \end{aligned}$$

where $\overline{G}(\mathbb{Q})$ acts on $(E_v \Omega)_{C_p}$ through $\overline{G}^{\text{ad}}(\mathbb{Q}_p) \cong PGL_2(E_v)$.

Moreover, one can drop the assumption that C is maximal at p , on replacing $(E_v \Omega)_{C_p}$ by some finite étale covering as in 7.4.7. [BZ95, 3], [Var98b, 5].

7.5.4. Rapoport, Zink and Boutot on one hand (by modular methods à la Drinfeld), and Varshavsky on the other hand (by group-theoretic methods à la Čerednik), have pointed out similar global p -adic uniformizations in much more general cases.

Let us only mention a remarkable example: Mumford's *fake projective plane* [Mu79]. This is a smooth projective complex surface S with the same Betti numbers as \mathbb{P}^2 , but not isomorphic to \mathbb{P}^2 . Actually, it is a surface of general type. Mumford's construction is of diadic nature. It is a quotient of the two-dimensional generalization $\Omega^{(2)}$ of Ω by an explicit arithmetic group. This $\Omega^{(2)}$ is the complement of the lines defined over \mathbb{Q}_2 in the projective plane.

Using the last theorem of [RZ96] or [Var98b], one can see that S is actually a Shimura surface of PEL type, parametrizing polarized abelian varieties of dimension 9, with multiplication by an order in a simple \mathbb{Q} -algebra B of dimension 18 with center $\mathbb{Q}(\sqrt{-7})$. The invariants of B at the two primes of $\mathbb{Q}(\sqrt{-7})$ above 2 are $1/3$ and $2/3$ (cf. a recent preprint by F. Kato for more detail). The associated group G is a group of unitary similitudes of signature $(1, 2)$.

The “diadic ball” $\Omega^{(2)}$ (or rather its formal avatar) has an modular interpretation for certain p -divisible groups (with multiplication by B_2). These p -divisible groups split as a product of three p -divisible groups of slope $1/3$ and height 3, and three p -divisible groups of slope $2/3$ and height 3.

The associated Gauss-Manin connection splits into six factors of rank 3 (in two variables). It resembles the Appell-Lauricella connection studied in [Ter73], [DM93], [Yo87], but is different.

7.6. Application of the theory of p -adic Betti lattices.

7.6.1. We consider more closely the Gauss-Manin connection $(\mathcal{H}, \nabla_{\text{GM}})$ in the case of Shimura curves.

We begin with the case $E = \mathbb{Q}$ as in 7.4. Then \mathcal{H} is a vector bundle of rank 4, and the B -action on \mathcal{H} obtained by functoriality of De Rham cohomology commutes with ∇_{GM} .

Let us choose an imaginary quadratic field $F = \mathbb{Q}(\sqrt{-d})$ inside B . Then $(\mathcal{H}, \nabla_{\text{GM}}) \otimes F$ splits into two parts: $(\mathcal{H}, \nabla_{\text{GM}})_+ \oplus (\mathcal{H}, \nabla_{\text{GM}})_-$ (on which $F \subset B$ acts through identity and complex conjugation respectively).

On the other hand, let us fix embeddings $F \subset \overline{\mathbb{Q}} \subset \mathbb{C}$, and let us fix a base point $s \in \mathcal{X}_\Gamma(\overline{\mathbb{Q}})$ of our Shimura curve. This is the moduli point of a decorated fake elliptic curve $A/\overline{\mathbb{Q}}$. We have $V^{\mathbb{C}} = H_B^1(A_{\mathbb{C}}, \mathbb{C}) \cong (\mathcal{H}_{\mathbb{C}}^{\text{an}})^{\nabla_{\text{GM}}}$, and this again splits into \pm parts. The Betti cohomology F -space $H_B^1(A_{\mathbb{C}}, F)$ already splits into two parts: $H_B^1(A_{\mathbb{C}}, F)_+ \oplus H_B^1(A_{\mathbb{C}}, F)_-$, both of rank two and stable under the action of $B \otimes F \cong M_2(F)$.

The flag space \mathcal{D}^{\vee} is actually defined over the reflex field \mathbb{Q} . Over F , it becomes isomorphic to \mathbb{P}^1 , or more precisely, to $\mathbb{P}(H_B^1(A_{\mathbb{C}}, F)_+)$. The period mapping describes the “slope” of the Hodge line $F^1 \cap V_+^{\mathbb{C}}$ with respect to a basis of $H_B^1(A_{\mathbb{C}}, F)_+ \subset ((\mathcal{H}_+)^{\text{an}})^{\nabla_{\text{GM}}}$, and is thus given by a quotient of solutions of the (partial) Gauss-Manin connection $\nabla_{\text{GM}+}$.

Assume for simplicity that $\Gamma \subset B^{\times}/\mathbb{Q}^{\times}$ is torsion-free. Then Γ may be identified with the projective monodromy group, *i.e.* with the quotient by

the homotheties of the image of the monodromy representation

$$\pi_1((\mathcal{X}_\Gamma)_\mathbb{C}, s) \rightarrow GL(H_B^1(A_\mathbb{C}, F)_+).$$

7.6.2. Let us now turn to the p -adic side. We fix an embedding $\overline{\mathbb{Q}} \subset \mathbb{C}_p$. We shall limit ourselves to the case of a prime p of *supersingular reduction* for A . Note that this includes the case of critical primes (7.4.6).

Let D be the definite quaternion algebra over \mathbb{Q} which is ramified only at p , and let \overline{B} be the definite quaternion algebra over \mathbb{Q} which is ramified at the same primes as B except p . Then, whether p is critical or not, the group J (of self-quasi-isogenies of $\overline{A}[p^\infty]$ commuting with B) is \overline{B}_p^\times .

By the local criterion for splitting fields [Vig80, III.3.5], F is also a splitting field for the quaternion algebra D (and \overline{B} as well). We can therefore apply the theory of I.5.3, and get an F -structure $H_B^1(A_{\mathbb{C}_p}, F)$ inside $H_{\text{cris}}^1(\overline{A}, / \mathbb{C}_p) \cong V^{\mathbb{C}_p} = \mathbf{D}(\overline{\Lambda}) \otimes \mathbb{C}_p$. By construction, it is stable under the action of $M_2(D) \otimes F \cong M_4(F) \cong B \otimes \overline{B} \otimes F$.

Using the embedding $F \otimes 1 \otimes F \subset B \otimes \overline{B} \otimes F$, one splits this F -structure $H_B^1(A_{\mathbb{C}_p}, F)$ into two parts $H_B^1(A_{\mathbb{C}_p}, F)_+ \oplus H_B^1(A_{\mathbb{C}_p}, F)_-$ (on which $F \otimes 1 \otimes 1 \subset B \otimes \overline{B} \otimes F$ acts through identity and complex conjugation respectively), both of rank two and stable under the action of $\overline{B} \otimes F \subset M_4(F)$, hence under J .

The p -adic flag space \mathcal{D}^\vee is again defined over the reflex field \mathbb{Q} . Over F , it becomes isomorphic to $\mathbb{P}(H_B^1(A_{\mathbb{C}_p}, F)_+)$ (note that the scaling ambiguity of $H_B^1(A_{\mathbb{C}_p}, F)_+$ by a factor in $\sqrt{F^\times}$ disappears here). The J -equivariant period mapping \mathcal{P} describes the ‘‘slope’’ of the Hodge line $F^1 \cap V_+^{\mathbb{C}_p}$ with respect to a fixed basis (e_0, e_1) of $H_B^1(A_{\mathbb{C}_p}, F)_+ \subset ((\mathcal{H}_+)_\mathbb{C}_p)^{\text{an}} \nabla^{\text{GM}}$, and is thus given by a quotient of solutions of $\nabla_{\text{GM}+}$.

Note that if B is split at p , then \mathcal{P} is of Gross-Hopkins type, and one has only a local p -adic uniformization of \mathcal{X}_Γ around s (in a supersingular disk).

If instead p is critical, then \mathcal{P} is of Drinfeld type, and the group $\Gamma_p \subset J^1$ appearing in the global p -adic uniformization of \mathcal{X}_Γ may be identified with the quotient by the homotheties of the image of the monodromy representation (in the sense of 7.2.6)

$$\pi_1^{\text{top}}((\mathcal{X}_\Gamma)_{\mathbb{C}_p}, s) \rightarrow GL(H_B^1(A_{\mathbb{C}_p}, F)_+).$$

A slight refinement provides a rank two $\mathcal{O}_F[\frac{1}{p}]$ -lattice $H_B^1(A_{\mathbb{C}_p}, \mathcal{O}_F[\frac{1}{p}])_+$ in $((\mathcal{H}_+)_\mathbb{C}_p)^{\text{an}} \nabla^{\text{GM}}$ stable under p -adic monodromy.

7.6.3. One can go one step further and see that the phenomenon encountered in 6.3.3 shows up here again: ‘‘the’’ quotient τ of solutions of solutions of ∇_{GM} which expresses locally the period mapping \mathcal{P} is essentially given by the same formula in the complex and the p -adic cases.

More precisely, assume that A has complex multiplication by an order in $M_2(F)$ (so that A is isogenous to the square of an elliptic curve with

complex multiplication by $\mathcal{O}_{\mathbb{Q}(\sqrt{-d})}$ with [fundamental] discriminant $-d$), and identify $\text{End}_B(A)$ with F . We denote by ϵ the Dirichlet character, w the number of roots of unity, h the class number attached to $\mathcal{O}_{\mathbb{Q}(\sqrt{-d})}$.

We may choose our symplectic basis e_0, e_1 of $H_B^1(A_{\mathbb{C}}, F)_+$ (*resp.* $H_B^1(A_{\mathbb{C}_p}, F)_+$) in such a way that $[\sqrt{-d}]^*e_0 = -\sqrt{-d}e_0$, $[\sqrt{-d}]^*e_1 = \sqrt{-d}e_1$ (here $[\sqrt{-d}]$ is viewed as an element of $\text{End}_B(A) = F$).

Let ω be a section of \mathcal{H}_+ in a Zariski neighborhood U of s , which corresponds to a relative differential of the first kind (no pole) on the universal abelian scheme. Then ω is a cyclic vector for \mathcal{H}_+ over U with respect to ∇_{GM} , so that ∇_{GM} amounts to a concrete linear differential equation of order two over U .

We calculate τ with respect the basis e_0, e_1 of $H_B^1(A_{\mathbb{C}}, F)_+$ (*resp.* $H_B^1(A_{\mathbb{C}_p}, F)_+$) in a small neighborhood of the base point s . Denoting by z an algebraic local parameter at s in U , and extending e_0, e_1 by horizontality, we then have the proportionality

$$\omega \sim \tau(z)e_0 + e_1$$

Theorem 7.6.4. *There is a quotient $y = y_1/y_2$ of solutions of ∇_{GM} in $\overline{\mathbb{Q}}[[z]]$ such that τ takes the form $\tau(z) = \kappa^{-1}y(z)$ for a suitable constant κ .*

In the complex case, $\kappa \sim \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma \left\langle \frac{u}{d} \right\rangle \right)^{\epsilon(u)w/2h} \in \mathbb{C}^\times / \overline{\mathbb{Q}}^\times$.

In the p -adic case, assuming moreover that $p \nmid d$,

$$\kappa \sim \prod_{u \in (\mathbb{Z}/d)^\times} \left(\Gamma_p \left\langle \frac{pu}{d} \right\rangle \right)^{-\epsilon(u)w/4h} \in \mathbb{C}_p^\times / \overline{\mathbb{Q}}^\times.$$

PROOF. Let us write $\omega|_{z=0} = \omega_{11}e_1$. Then $\tau(z) = \kappa^{-1}y(z)$, with $\kappa = \omega_{11}^2/2\pi i$ in the complex case, $\kappa = \omega_{11}^2$ in the p -adic case (the factor $2\pi i$ which arises in the complex case is the factor of proportionality of the symplectic forms in De Rham and in Betti cohomology respectively). The evaluation of ω_{11} in the complex case is given by the Lerch-Chowla-Selberg formula (*cf.* I.4.6.3). In the p -adic case, it is given by I.5.3.9. \square

7.6.5. This discussion generalizes without much complication to the case of Shimura curves attached to a quaternion algebra D over a totally real number field E of degree > 1 . We concentrate on the case of a critical place v and assume, for simplicity, that E is Galois over \mathbb{Q} and that v is the unique place of E with the same residue characteristic p (*i.e.* $m = 0$ in the notation of 7.5.2).

Let F be a totally imaginary quadratic extension of E contained in B . Then F is a splitting field for B, \overline{B} , and D . Let us fix a double embedding $\mathbb{C} \hookrightarrow \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ as in 7.5.2, and an extension $FK \hookrightarrow \overline{\mathbb{Q}}$ of ∞_0 . For C small enough, $\mathcal{S}h^\bullet$ carries a universal abelian scheme of relative dimension $g = [FK : \mathbb{Q}]$. After tensoring $\otimes_K \overline{\mathbb{Q}}$, its Gauss-Manin connection splits into pieces of rank two, indexed by the embeddings of FK into $\overline{\mathbb{Q}}$. We select the piece $(\mathcal{H}, \nabla_{\text{GM}})_+$ corresponding to ∞_0 .

Let us fix a base point $s \in \mathcal{S}h(\overline{\mathbb{Q}}) \subset \mathcal{S}h^\bullet(\overline{\mathbb{Q}})$ of our Shimura curve. This is the moduli point of a decorated abelian variety $A^\bullet/\overline{\mathbb{Q}}$ (of dimension $g = [FK : \mathbb{Q}]$ and multiplication by \mathcal{B}^\bullet). We have $V^\bullet = V \otimes_E K \cong H_B^1(A_{\mathbb{C}}^\bullet, \mathbb{Q})$. Note that $H_B^1(A_{\mathbb{C}}, F) \cong V \otimes_{\mathbb{Q}} FK$ splits into pieces of FK -rank two, indexed by the embeddings of F into $\overline{\mathbb{Q}}$. We select the piece $H_B^1(A_{\mathbb{C}}^\bullet, F)_+$ corresponding to ∞_0 . We then have a natural embedding $H_B^1(A_{\mathbb{C}}^\bullet, F)_+ \subset ((\mathcal{H}_+^{\text{an}})_{\mathbb{C}}^{\text{an}})^{\nabla_{\text{GM}}}$, and $H_B^1(A_{\mathbb{C}}^\bullet, F)_+$ is stable under monodromy.

On the other hand, it follows from 7.4.6, that A^\bullet has supersingular reduction \overline{A}^\bullet at v . We can construct the F -structure $H_B^1(A_{\mathbb{C}_p}^\bullet, F)$ inside $H_{\text{cris}}^1(\overline{A}^\bullet, / \mathbb{C}_p) \cong V^{\mathbb{C}_p} = \mathbf{D}(\overline{\Lambda}^\bullet) \otimes \mathbb{C}_p$. By construction, it is stable under the action of $M_g(D) \otimes F \cong M_{2g}(F)$, which contains naturally $B \otimes_{\mathbb{Q}} \overline{B} \otimes_{\mathbb{Q}} F$.

Using the embedding $F \otimes 1 \otimes F \subset B \otimes \overline{B} \otimes F$, one splits this F -structure $H_B^1(A_{\mathbb{C}_p}, F)$ into parts of FK -rank two, indexed by the embeddings of F into $\overline{\mathbb{Q}}$. We select the piece $H_B^1(A_{\mathbb{C}_p}^\bullet, F)_+$ corresponding to ∞_0 . We then have a natural embedding $H_B^1(A_{\mathbb{C}_p}^\bullet, F)_+ \subset ((\mathcal{H}_+^{\text{an}})_{\mathbb{C}_p}^{\text{an}})^{\nabla_{\text{GM}}}$, and $H_B^1(A_{\mathbb{C}_p}^\bullet, F)_+$ is stable under p -adic monodromy.

7.7. Conclusion.

Roughly speaking, the period mapping \mathcal{P} attaches to each member of an algebraic family of algebraic complex varieties (with additional structure) a point in some flag space, which encodes the Hodge filtration in the cohomology of this variety. This mapping can be expressed in terms of quotients of solutions of the fuchsian differential equation (Picard-Fuchs/Gauss-Manin) which controls the variation of the cohomology.

We have described the theory of period mappings for p -divisible groups in somewhat similar terms.

Sometimes, it is possible to go through the looking-glass of sheer analogies. This occurs when we restrict our attention to p -divisible groups attached to polarized abelian varieties with prescribed endomorphisms and fixed ‘‘Shimura type’’. In this situation, we have three algebraic objects defined over a number field (the reflex field):

- (a) the Shimura variety $\mathcal{S}h$ parametrizing certain ‘‘decorated’’ abelian varieties,
- (b) the Gauss-Manin connection ∇_{GM} describing the variation of these abelian varieties over $\mathcal{S}h$, and
- (c) the flag variety \mathcal{D}^\vee .

These objects are transcendently related, both in the complex and p -adic sense.

The period mapping may be viewed as multivalued and defined on p -adic open domains in $\mathcal{S}h$; it is given as in the complex case by quotients of solutions of the Gauss-Manin connection (as a p -adic connection).

In most situations, these open domains are small and \mathcal{P} is single-valued.

However, it happens in some remarkable cases that \mathcal{P} is a global multi-valued function. This occurs when the abelian varieties under consideration form a *single isogeny class modulo p* . In this situation, the complex and p -adic transcendental relations between Sh, ∇_{GM} are completely similar: Sh is uniformized by an analytic space which is étale over some open subset \mathcal{D} of \mathcal{D}^\vee , and the group of deck transformations Γ is isomorphic to the projective global monodromy group of ∇_{GM} . In III, we shall analyze in detail the group-theoretic aspects of this situation.

CHAPTER III

p-adic orbifolds and monodromy.

Contents

1. Étale coverings and fundamental groups in <i>p</i> -adic geometry.	112
2. Temperate fundamental groups.	126
3. Local and global monodromy of <i>p</i> -adic differential equations.	141
4. Non-archimedean orbifolds, uniformizing differential equations and period mappings.	158
5. <i>p</i> -adic triangle groups.	175
6. The tale of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ (Escher's triangle group and its diadic and triadic twins).	190

1. Étale coverings and fundamental groups in p -adic geometry.

ABSTRACT: In this section, inspired by De Jong’s work [dJ95b], we discuss various kinds of finite or infinite étale coverings in non-archimedean analytic geometry, and the associated fundamental groups. The temperate fundamental group of a p -adic manifold encapsulates information on the bad reduction properties of all its finite étale coverings.

1.1. More on the topological fundamental group.

1.1.1. We shall work throughout in the context of Berkovich analytic geometry, over a fixed complete ultrametric field $(K, |\cdot|)$.

What has been said about Berkovich spaces in I.1 remains true in this broader context (*i.e.* without any further restriction on K).

More precisely, we shall work with *paracompact strictly K -analytic spaces* in the sense of [Ber93]. We recall that these form a category equivalent to the category of quasi-separated rigid spaces over K having an admissible covering of finite type.

The main advantage of these spaces over rigid spaces lies in their topology: this is a topology in the usual sense (not a Grothendieck topology); moreover, each point has a fundamental system of open neighborhoods which are locally compact, countable at infinity and arcwise connected.

We shall deal mostly with what we have called, for convenience, (analytic) *K -manifolds*, *i.e.* paracompact strictly K -analytic spaces which satisfy the following condition (I.1.3.7):

*any $s \in S$ has a neighborhood $U(s)$ which is isomorphic to an affinoid subdomain of some smooth space (in the sense of [Ber93] *i.e.* a space which admits locally an étale morphism to the affine space $\mathbb{A}^{\dim S}$).*

Remarks 1.1.2. a) Any affinoid domain of a K -manifold is rigid-analytically smooth, *i.e.* satisfies the jacobian criterion (*cf.* [Ber93, 2.2]; note also that “smooth” in the sense of Berkovich amounts to “rigid-analytically smooth and having no boundary”); a fortiori, an affinoid K -manifold is rigid-analytically smooth. Conversely, a rigid-analytically smooth (strictly) K -affinoid space S is a K -manifold, *cf.* [Ber99, rem. 9.7].

b) This notion is slightly stronger than the notion of spaces locally embeddable in a smooth space considered in [Ber99]. The main difference is that we impose that S is a *good space* in the sense of [Ber93], *i.e.* that any point has an affinoid neighborhood, which is not automatic for non-classical points (this will be useful in § 3).

c) It follows immediately from remark a) that *a K -manifold is nothing but a good and rigid-analytically smooth (paracompact strictly K -analytic) space.*

d) Let $\phi : S' \rightarrow S$ be an étale morphism of paracompact strictly K -analytic spaces. If S is a K -manifold, so is S' .

e) For an algebraic K -variety X , one has X smooth $\iff X^{\text{an}}$ smooth $\iff X^{\text{an}}$ is a K -manifold. A morphism f is étale $\iff f^{\text{an}}$ is étale.

It follows from [Ber99, cor. 9.5] that K -manifolds are locally arcwise-connected and *locally simply-connected*, hence subject to the familiar theory of topological coverings (the difficulty is to show that non-classical points admit contractible neighborhoods). In particular, any pointed K -manifold (S, s) admits a *universal (pointed) topological covering* (\tilde{S}, \tilde{s}) , such that $S \cong \tilde{S}/\pi_1^{\text{top}}(S, s)$; \tilde{S} is a (pointed) K -manifold, and $\pi_1^{\text{top}}(S, s)$ is a discrete group of automorphisms of \tilde{S} .

Let us also recall that in dimension one, $\pi_1^{\text{top}}(S, s)$ is a free group isomorphic to the fundamental group of the dual graph of the semistable reduction of S (cf. [dJ95b, 5.3]). In higher dimension, there is no such restriction on the discrete groups $\pi_1^{\text{top}}(S, s)$: actually, any differentiable manifold is homotopy equivalent to some K -manifold [Ber90, 6.1.8].

1.1.3. Berkovich's analysis of the homotopy of analytic spaces of higher dimension makes use of De Jong's alteration theorem. This allows him to reduce the problem to the case of the generic fiber of so-called polystable fibrations of formal schemes over K^0 (the ring of integers in K). He associates to such fibrations a simplicial set which encodes the combinatorics of incidence of the intersection strata in the special fiber, and shows that the geometric realization of this simplicial set is homotopy equivalent to the generic fiber (viewed as an analytic K -manifold) [Ber99, 5.4].

In particular, in the case of (the generic fiber of) an algebraic polystable fibration, the simplicial set is finite, hence the topological fundamental group is finitely generated. Using De Jong's theorem and the following result, one can deduce that *the topological fundamental group of an algebraic K -manifold is finitely generated*.

Proposition 1.1.4. *Let \bar{S} be a K -manifold, and let Z be a Zariski-closed, nowhere dense, reduced analytic subset. Then any topological covering of $S := \bar{S} \setminus Z$ extends uniquely to a topological covering of \bar{S} . Equivalently (since S and \bar{S} are locally simply-connected), the natural homomorphism $\pi_1^{\text{top}}(S, s) \rightarrow \pi_1^{\text{top}}(\bar{S}, s)$ is an isomorphism.*

PROOF. Extensions of topological coverings from S to \bar{S} are unique if they exist, since \bar{S} is normal (note that S is connected, like \bar{S} , by normality of \bar{S} [Ber90, 3.3.16]). For the existence, we argue by induction on the dimension of Z (starting from the trivial case where Z is empty). This reduces the question to extending topological coverings from S to $\bar{S} \setminus Z^{\text{sing}}$ (where Z^{sing} denotes the non-smooth locus of Z , which is closed and of dimension less than the dimension of Z , cf. [Ber93, 2.2]); equivalently, this reduces the question to the case where Z is (rigid-analytically) smooth.

By unicity, the question of existence of extensions is local, *i.e.* is a question at the neighborhood in S of each point z of Z ; we may assume

that \overline{S} and Z are (strictly) affinoid, rigid-analytically smooth and equidimensional (Z of codimension $r > 0$ in \overline{S}). The statement is clear if z has a neighborhood $U(z)$ which is a finite union of affinoid domains U_i 's such that $U_i \setminus (U_i \cap Z)$ is simply-connected. This is indeed the case, by Kiehl's theorem on the existence of tubular neighborhoods [Kie67a]: translated into Berkovich geometry, it asserts that (for S affinoid and Z a Zariski-closed subset, both rigid-analytically smooth and equidimensional) Z has a neighborhood which is a finite union of affinoid domains U_i 's isomorphic to $(U_i \cap Z) \times \mathbb{D}^r$, where \mathbb{D}^r is an r -dimensional polydisk, and $(U_i \cap Z)$ corresponds to $(U_i \cap Z) \times \{0\}$. Since $U_i \cap Z$ is a rigid-analytically smooth affinoid domain, it is locally simply connected according to Berkovich; on the other hand, in the non-archimedean situation, $\mathbb{D}^r \setminus \{0\}$ is simply-connected. Whence the result. \square

1.2. Étale versus topological coverings (again).

1.2.1. Let S be a connected K -manifold. Let $S' \xrightarrow{f} S$ be a finite étale morphism (S' is then a K -manifold). Its degree n may be defined as the rank of the locally free \mathcal{O}_S -module $f_*\mathcal{O}_{S'}$. Assume that K is algebraically closed. Then one has the formula

$$n = \sum_{y \in f^{-1}(x)} [\mathcal{H}(y) : \mathcal{H}(x)]$$

where $\mathcal{H}(x)$ stands for the residue field of the point x (if $S = \mathbf{M}(\mathcal{A})$ is affinoid and x corresponds to the bounded multiplicative seminorm $|\cdot|_x$, this is just $\mathcal{H}(x) = Q(\widehat{\mathcal{A}/\text{Ker}|\cdot|_x})$).

Then f is a *topological covering* if and only if all fibers have the same cardinality, which amounts to the algebraic notion of being “completely decomposed”: $\mathcal{H}(y) = \mathcal{H}(f(y))$ for any $y \in S'$, cf. [Ber90, 3.2.7], [Ber93, 6.3.1].

This criterion shows at once that the Kummer étale covering $z \mapsto z^n$ (with $\text{char}(K) \nmid n$) of the punctured disk is *not* a topological covering.

1.2.2. A *geometric point* \overline{s} of S is a point with value in some complete algebraically closed extension $(\Omega, |\cdot|)$ of $(K, |\cdot|)$, i.e. a morphism $\overline{s} : \mathbf{M}(\Omega) \rightarrow S$ in the category of analytic spaces over K [Ber90, p.48]. For any finite étale covering there are exactly n liftings $\overline{s}' : \mathbf{M}(\Omega) \rightarrow S'$ of \overline{s} , i.e. n geometric points of S' above S . We usually denote by s the point of S which is the image of \overline{s} .

Definition 1.2.3 (Berkovich, De Jong). A morphism $f : S' \rightarrow S$ is a *covering* (resp. *étale covering*, resp. *topological covering*) if S is covered by open subsets U such that $\coprod V_j = f^{-1}U \rightarrow U$ and the restriction of f to every V_j is finite (resp. étale finite, resp. an isomorphism).

There is an obvious notion of morphism between coverings of S . The category of étale coverings, *resp.* finite étale coverings, *resp.* topological coverings, will be denoted by $\text{Cov}_S^{\text{ét}}$, *resp.* $\text{Cov}_S^{\text{alg}}$, *resp.* $\text{Cov}_S^{\text{top}}$.

There are inclusions (fully faithful embeddings)

$$\text{Cov}_S^{\text{alg}} \hookrightarrow \text{Cov}_S^{\text{ét}}, \quad \text{Cov}_S^{\text{top}} \hookrightarrow \text{Cov}_S^{\text{ét}}.$$

Remarks 1.2.4. (i) If $S' \rightarrow S$ is an étale covering, then S' is a K -manifold.

(ii) It is worth noticing that the notions of étale and of topological coverings are local on the base S , but not the notion of algebraic coverings (except if S is compact).

(iii) Even if S is covered by finitely many U 's as in the definition, it does not follow that S' is a disjoint union of finite algebraic coverings of S . For instance, a Tate elliptic curve (with universal covering $\tilde{S} \cong \mathbb{G}_m$) has a covering by two open subsets U_1, U_2 such that $\tilde{S}|_{U_i}$ is a disjoint union of copies of S .

(iv) It is an immediate but important fact that these categories are stable under taking fiber product (over S). It is also stable under taking finite disjoint unions.

(v) One should pay attention to the fact that unlike $\text{Cov}_S^{\text{top}}$, $\text{Cov}_S^{\text{ét}}$ is not stable under arbitrary disjoint unions, nor under composition (in the sense that the composition of an étale covering S'/S and an étale covering S''/S' may not be an étale covering S''/S). However, we have:

Lemma 1.2.5. *Any morphism which is the composition of a covering (resp. étale covering) followed or preceded by a finite (resp. finite étale) morphism is a covering (resp. étale covering).*

PROOF. Let us consider the case of a covering $g : S'' \rightarrow S'$ followed by a finite covering $f : S' \rightarrow S$ (the reverse situation is immediate from the definition). Let s be a point of S . Then for any s' in the finite set $f^{-1}(s)$, there exists an open neighborhood $U_{s'} \subset S'$ of s' such that $g^{-1}(U_{s'})$ is a disjoint union of spaces $U_{s',i}$, each being finite over $U_{s'}$ via g . We may assume that the $U_{s'}$ are pairwise disjoint. Since f is finite, it is in particular a closed continuous map, hence S' admits a basis of open neighborhoods of $f^{-1}(\bar{s})$ of the form $f^{-1}(V)$. In particular, there is an open neighborhood V_s of s such that $\prod_{s' \in f^{-1}(s)} U_{s'}$ contains $f^{-1}(V_s)$. We may replace each $U_{s'}$ by

its intersection with $f^{-1}(V_s)$ (which is a union of connected components of $f^{-1}(V_s)$). It is then clear that $g \circ f$ induces a finite morphism from $U_{s',i}$ to V_s . Hence $g \circ f$ is a covering. The étale variant is similar. \square

Examples 1.2.6. (i) The prototype of an étale covering is given by the logarithm map

$$\log : D_{\mathbb{C}_p}(1, 1^-) \longrightarrow (\mathbb{A}_{\mathbb{C}_p}^1)^{\text{an}}.$$

For any $m \in \mathbb{N}$, the inverse image of the open disk

$$D_m = D\left(0, (|p|^{-m + \frac{1}{p-1}})^-\right) \subset (\mathbb{A}_{\mathbb{C}_p}^1)^{\text{an}}$$

is a disjoint union

$$\coprod_{\zeta \in \mu_{p^\infty}} \left\{ q \in D(1, 1^-) \mid \zeta q^{p^m} \in D\left(1, (|p|^{\frac{1}{p-1}})^-\right) \right\}$$

and each of these components is isomorphic to D_m if $m = 0$, and finite étale onto D_m in general (we are using here the factorization $q \mapsto \zeta q^{p^m} = t \mapsto \log t \mapsto p^{-m} \log t = \log q$ of the logarithm).

(ii) This example may be interpreted as a Dwork period mapping (II.6.3.1). In fact, all examples of period mappings considered in II.6.3 are étale coverings of the period domain \mathcal{D} . For instance, the Gross-Hopkins period mapping (II.6.3.2) is an étale covering of $(\mathbb{P}_{\mathbb{C}_p}^1)^{\text{an}}$, cf. [dJ95b, 7.2].

Any period mapping is étale, but it may happen in some cases that it is surjective without being an étale covering, cf. [RZ96, 5.41] (this somewhat mysterious phenomenon is due to the existence of bijective non-quasi-compact étale morphisms).

(iii) Any (possibly infinite) topological covering of a finite étale covering is an étale covering after 1.2.5.

The following two lemmas are due to V. Berkovich and J. De Jong (and not stated in the most general form).

Lemma 1.2.7. (i) *An étale presheaf \mathcal{F} on S which is representable by an étale covering S' is a sheaf. For any subdomain $U \subset S$, $\mathcal{F}|_U$ is representable by $S' \times_S U$.*

(ii) *Let \mathcal{F} be an étale sheaf. Let U_i an open covering of S . If for every i , $\mathcal{F}|_{U_i}$ is representable by an étale covering, so is \mathcal{F} .*

For (i), cf. [Ber93, 4.1.3, 4.1.4, 4.1.5]. Here we take advantage of working with *strict* analytic spaces. Gluing — using [Ber93, 1.3.3] — gives (ii) (cf. also [dJ95b, 2.3]).

Lemma 1.2.8. *Let $S' \rightarrow S$ be an étale covering, and let $R \subset S' \times_S S'$ be a union of connected components which is an equivalence relation on S' over S . Then S'/R (viewed as an étale sheaf on S) is representable by an étale covering $S'' \rightarrow S$.*

cf. [dJ95b, 2.4]: by the previous lemma, the question is local on S ; the proof is by reduction to the case of a finite étale covering of affinoid spaces.

In the situation of 1.2.8, we say that the $S'' \rightarrow S$ is a *quotient étale covering* of $S' \rightarrow S$.

1.3. Existence of étale paths.

1.3.1. Let us fix a geometric point \bar{s} of S . We consider the fiber functor

$$F_{S,\bar{s}}^{\text{et}} = F_{\bar{s}}^{\text{et}} : \text{Cov}_S^{\text{et}} \longrightarrow \text{Sets}, \quad S' \mapsto \{\text{geometric points } \bar{s}' \text{ of } S' \text{ above } \bar{s}\}$$

and its restrictions

$$F_{S,\bar{s}}^{\text{alg}} = F_{\bar{s}}^{\text{alg}} : \text{Cov}_S^{\text{alg}} \longrightarrow \text{Sets},$$

(considered in [SGA 1]) and

$$F_{S,\bar{s}}^{\text{top}} = F_{\bar{s}}^{\text{top}} : \text{Cov}_S^{\text{top}} \longrightarrow \text{Sets}.$$

An *étale path* (resp. *algebraic path*) from \bar{s} to another geometric point \bar{t} of S is an isomorphism of functors $F_{\bar{s}}^{\text{et}} \xrightarrow{\sim} F_{\bar{t}}^{\text{et}}$ (resp. $F_{\bar{s}}^{\text{alg}} \xrightarrow{\sim} F_{\bar{t}}^{\text{alg}}$).

The set of étale paths is topologized by taking as fundamental open neighborhoods of an étale path α the set $\text{Stab}_{S',\bar{s}'} \circ \alpha$, where $\text{Stab}_{S',\bar{s}'}$ runs among the stabilizers in $\text{Aut}(F_{\bar{s}}^{\text{et}})$ of arbitrary geometric points \bar{s}' above \bar{s} in arbitrary étale coverings S'/S .

If we do the same for algebraic paths, we get the profinite topology considered in [SGA 1].

On the other hand, an isomorphism of functors $F_{\bar{s}}^{\text{top}} \xrightarrow{\sim} F_{\bar{t}}^{\text{top}}$ amounts to a path up to homotopy in the usual sense between s and t in the arcwise connected space S (this is compatible with composition of paths, which is juxtaposition in the reverse order according to our convention); they form a discrete set — in fact, a principal homogeneous space under $\pi^{\text{top}}(S, s)$ acting on the left.

The inclusions $\text{Cov}_S^{\text{alg}} \hookrightarrow \text{Cov}_S^{\text{et}}$, $\text{Cov}_S^{\text{top}} \hookrightarrow \text{Cov}_S^{\text{et}}$ induce continuous maps from the space of étale paths between \bar{s} and \bar{t} to the space of algebraic paths between \bar{s} and \bar{t} (resp. to the discrete set of paths up to homotopy between s and t).

If $f : S \rightarrow X$ is any morphism of K -manifolds, and if α is an étale path from \bar{s} to \bar{t} , there is an obvious notion of push-forward $f_*\alpha$, which is an étale path from $f \circ \bar{s}$ to $f \circ \bar{t}$; it is compatible with composition of étale paths (as composition of isomorphisms of functors).

1.3.2. It follows from [SGA 1, exp. V] that algebraic paths exist. Using the nice topology of Berkovich spaces, A.J. De Jong was able to prove [dJ95b, 2.9]:

Key lemma 1.3.3. *Étale paths exist. More precisely, any Berkovich path γ (up to homotopy) between s and t lifts to an étale path $\bar{\gamma}$ between \bar{s} and \bar{t} .*

PROOF. Up to introducing intermediate geometric points, we may assume that S is affinoid (hence compact). For any *finite* open cover $\mathcal{U} = (U_i)$ of S , let us denote by $\text{Cov}_{S,\mathcal{U}}^{\text{et}}$ the full subcategory of Cov_S^{et} of étale coverings

$f : S' \rightarrow S$ such that $f^{-1}(U_i) = \coprod V_{ij}$ and V_{ij} is étale finite over U_i via f . We thus have

$$\mathrm{Cov}_S^{\mathrm{et}} = \varinjlim_{\mathcal{U}} \mathrm{Cov}_{S,\mathcal{U}}^{\mathrm{et}}, \text{ and } \mathrm{Isom}(F_{\bar{s}}^{\mathrm{et}}, F_{\bar{t}}^{\mathrm{et}}) = \varprojlim_{\mathcal{U}} \mathrm{Isom}(F_{\bar{s},\mathcal{U}}^{\mathrm{et}}, F_{\bar{t},\mathcal{U}}^{\mathrm{et}})$$

where $F_{\bar{s},\mathcal{U}}^{\mathrm{et}}, F_{\bar{t},\mathcal{U}}^{\mathrm{et}}$ denote the restrictions of $F_{\bar{s}}^{\mathrm{et}}, F_{\bar{t}}^{\mathrm{et}}$ to $\mathrm{Cov}_{S,\mathcal{U}}^{\mathrm{et}}$.

We have to show that $\mathrm{Isom}(F_{\bar{s}}^{\mathrm{et}}, F_{\bar{t}}^{\mathrm{et}}) \neq \emptyset$. The idea is to construct non-empty compact subsets $K_{\mathcal{U}}$ of $\mathrm{Isom}(F_{\bar{s},\mathcal{U}}^{\mathrm{et}}, F_{\bar{t},\mathcal{U}}^{\mathrm{et}})$ (topologized as above) such that if \mathcal{U}' refines \mathcal{U} , $K_{\mathcal{U}'}$ maps to $K_{\mathcal{U}}$ under the natural map $\mathrm{Isom}(F_{\bar{s},\mathcal{U}'}^{\mathrm{et}}, F_{\bar{t},\mathcal{U}'}^{\mathrm{et}}) \rightarrow \mathrm{Isom}(F_{\bar{s},\mathcal{U}}^{\mathrm{et}}, F_{\bar{t},\mathcal{U}}^{\mathrm{et}})$; it will follow that $\mathrm{Isom}(F_{\bar{s}}^{\mathrm{et}}, F_{\bar{t}}^{\mathrm{et}}) \supset \varinjlim_{\mathcal{U}} K_{\mathcal{U}} \neq \emptyset$.

In fact, it suffices to do so for a cofinal system of \mathcal{U} 's, which are chosen as follows. Since S is *arcwise connected*, there is a continuous embedding $I = [0, 1] \hookrightarrow S$ with $0 \mapsto s$, $1 \mapsto t$. One considers finite open coverings $\mathcal{U} = (U_1, \dots, U_n)$ such that

$$U_1 \cap I = [0, r_1[, \dots, U_i \cap I =]r_i, t_i[, \dots, U_m \cap I =]r_m, 1], U_{m+1} \cap I = \emptyset, \dots$$

for some points $r_i < t_{i-1} < r_{i+1} < t_i$ in I ($i = 2, \dots, m-1$) and some $m \leq n$.

Given \mathcal{U} , one chooses s_i in $U_i \cap U_{i+1} \cap I$ (which amounts to a real number between r_{i+1} and t_i), and a geometric point \bar{s}_i above s_i ($i = 1, \dots, m-1$); one completes this collection of geometric points by setting $\bar{s}_0 = \bar{s}$, $\bar{s}_m = \bar{t}$. Let us look at the diagram

$$\mathrm{Cov}_{S,\mathcal{U}}^{\mathrm{et}} \xrightarrow{\sigma_i} \mathrm{Ind} \mathrm{Cov}_{\mathcal{U}_{i+1}}^{\mathrm{alg}} \begin{array}{c} \xrightarrow{F_{\bar{s}_i, U_{i+1}}^{\mathrm{alg}}} \\ \xrightarrow{F_{\bar{s}_{i+1}, U_{i+1}}^{\mathrm{alg}}} \end{array} \mathrm{Sets}.$$

One has $F_{\bar{s}_j, U_{i+1}}^{\mathrm{alg}} \circ \sigma_i = F_{\bar{s}_i, \mathcal{U}}^{\mathrm{et}}$, $j = i, i+1$, whence continuous maps

$$\mathrm{Isom}(F_{\bar{s}_i, U_{i+1}}^{\mathrm{alg}}, F_{\bar{s}_{i+1}, U_{i+1}}^{\mathrm{alg}}) \rightarrow \mathrm{Isom}(F_{\bar{s}_i, \mathcal{U}}^{\mathrm{et}}, F_{\bar{s}_{i+1}, \mathcal{U}}^{\mathrm{et}}).$$

Since algebraic paths exist, $\mathrm{Isom}(F_{\bar{s}_i, U_{i+1}}^{\mathrm{alg}}, F_{\bar{s}_{i+1}, U_{i+1}}^{\mathrm{alg}})$ is a non-empty compact set for any $i = 0, \dots, m$. On composing such isomorphisms for $i = 0, \dots, m$, one gets a continuous map

$$\sigma_{\mathcal{U}, (\bar{s}_i)} : \prod_{i=0}^m \mathrm{Isom}(F_{\bar{s}_i, U_{i+1}}^{\mathrm{alg}}, F_{\bar{s}_{i+1}, U_{i+1}}^{\mathrm{alg}}) \rightarrow \mathrm{Isom}(F_{\bar{s}, \mathcal{U}}^{\mathrm{et}}, F_{\bar{t}, \mathcal{U}}^{\mathrm{et}})$$

whose image is a non-empty compact set $K_{\mathcal{U}}$.

In fact, while the map $\sigma_{\mathcal{U}, (\bar{s}_i)}$ depends on the choice of the geometric points \bar{s}_i , its image $K_{\mathcal{U}}$ does not. Indeed, for any other choice \bar{s}'_i , there is an algebraic path

$$\gamma_i : F_{\bar{s}_i, U_{i+1} \cap U_i}^{\mathrm{alg}} \xrightarrow{\sim} F_{\bar{s}'_i, U_{i+1} \cap U_i}^{\mathrm{alg}}, \quad \gamma_0 = \mathrm{id}, \quad \gamma_m = \mathrm{id}.$$

We denote by the same letter the induced algebraic paths on U_i and on U_{i+1} . Then $\sigma_{\mathcal{U},(\bar{s}_i)}(\alpha_0, \dots, \alpha_m) = \sigma_{\mathcal{U},(\bar{s}_i)}(\gamma_1 \alpha_0 \gamma_0^{-1}, \dots, \gamma_m \alpha_m \gamma_{m-1}^{-1})$, whence the independence.

From this observation, it is straightforward to conclude that if \mathcal{U}' refines \mathcal{U} , $K_{\mathcal{U}'}$ maps to $K_{\mathcal{U}}$ under the natural map $\text{Isom}(F_{\bar{s}, \mathcal{U}'}^{\text{et}}, F_{\bar{t}, \mathcal{U}'}^{\text{et}}) \rightarrow \text{Isom}(F_{\bar{s}, \mathcal{U}}^{\text{et}}, F_{\bar{t}, \mathcal{U}}^{\text{et}})$. This establishes the first assertion.

For the second one (the surjectivity of $\text{Isom}(F_{\bar{s}}^{\text{et}}, F_{\bar{t}}^{\text{et}}) \rightarrow \text{Isom}(F_{\bar{s}}^{\text{top}}, F_{\bar{t}}^{\text{top}})$), one may argue as follows: let $\bar{\bar{s}}$ be a lifting of \bar{s} to the universal topological covering $\pi : \bar{S} \rightarrow S$. Then $\gamma \cdot \bar{\bar{s}}$ is a lifting of \bar{t} to the universal topological covering. Let α be an étale path between $\bar{\bar{s}}$ and $\gamma \cdot \bar{\bar{s}}$. Then $\bar{\gamma} := \pi_*(\alpha)$ is an étale path between \bar{s} and \bar{t} which maps to the ordinary path γ up to homotopy, viewed as an isomorphism $F_{\bar{s}}^{\text{top}} \rightarrow F_{\bar{t}}^{\text{top}}$. \square

1.4. The formalism of fundamental groups.

1.4.1. Let Cov_S^\bullet be a full subcategory of Cov_S^{et} which is *stable under taking connected components, fiber products (over S) and quotients* (in the sense of 1.2.8.). Examples: Cov_S^{et} , $\text{Cov}_S^{\text{top}}$, $\text{Cov}_S^{\text{alg}}$.

We denote by $F_{S, \bar{s}}^\bullet = F_{\bar{s}}^\bullet$ the restriction of $F_{\bar{s}}^{\text{et}}$ to Cov_S^\bullet , and set

$$\pi_1^\bullet(S, \bar{s}) = \text{Aut } F_{\bar{s}}^\bullet$$

topologized by considering as fundamental open neighborhoods of 1 the stabilizers $\text{Stab}_{S', \bar{s}'}^\bullet$ in $\pi_1^\bullet(S, \bar{s})$ of arbitrary geometric points \bar{s}' above \bar{s} in arbitrary étale coverings $S' \rightarrow S$ in Cov_S^\bullet (here we see $\pi_1^\bullet(S, \bar{s})$ as acting on the left on $F_{\bar{s}}^\bullet(S')$).

That the topology is indeed compatible with the group law is due to the fact that the system of subgroups $\text{Stab}_{S', \bar{s}'}^\bullet$ is stable under intersection (since Cov_S^\bullet is stable under taking fiber products) and under conjugation (which amounts to a change of \bar{s}').

In the case $\text{Cov}_S^\bullet = \text{Cov}_S^{\text{et}}$, $\pi_1^{\text{et}}(S, \bar{s})$ is the étale fundamental group introduced and studied by De Jong [dJ95b].

In the case $\text{Cov}_S^\bullet = \text{Cov}_S^{\text{top}}$, $\pi_1^{\text{top}}(S, \bar{s}) = \pi_1^{\text{top}}(S, s)$ is (an incarnation of) the topological fundamental group.

In the case $\text{Cov}_S^\bullet = \text{Cov}_S^{\text{alg}}$, $\pi_1^{\text{alg}}(S, \bar{s})$ is a profinite group, which will be called the *algebraic fundamental group*. The theory is embodied in [SGA 1, V]. If S is the analytification of an algebraic smooth K -variety S^{alg} and if $\text{char}(K) = 0$, then this group coincides with Grothendieck's algebraic fundamental group. This follows from the *Gabber-Lütkebohmert theorem*, according to which the GAGA functor is an equivalence between the categories of finite étale coverings of S^{alg} and S respectively [Lüt93].

Lemma 1.4.2. *The natural continuous map*

$$\phi : \pi_1^\bullet(S, \bar{s}) \rightarrow \varprojlim_{\text{Stab}_{S', \bar{s}'}^\bullet} \pi_1^\bullet(S, \bar{s}) / \text{Stab}_{S', \bar{s}'}^\bullet$$

is a homeomorphism. In particular, $\pi_1^\bullet(S, \bar{s})$ is a separated pro-discrete (hence totally discontinuous) topological space.

PROOF. ϕ sends any neighborhood of $\text{id} \in \pi_1^\bullet(S, \bar{s})$ to a neighborhood of $\phi(\text{id}) \in \text{im}(\phi)$. Since ϕ is equivariant with respect to the continuous action of $\pi_1^\bullet(S, \bar{s})$ by left translation, it is open onto its image. Hence it suffices to show that it is bijective. Let $\underline{\gamma} := (\gamma_{S', \bar{s}'})$ be a compatible family of elements of $\pi_1^\bullet(S, \bar{s}) / \text{Stab}_{S', \bar{s}'}$, i.e. represent an element of $\varprojlim \pi_1^\bullet(S, \bar{s}) / \text{Stab}_{S', \bar{s}'}$.

For any S' in Cov_S^\bullet and any $\bar{s}' \in F_{\bar{s}}^\bullet(S')$, $\gamma_{S', \bar{s}'}$ is another well-defined element of $F^\bullet(S')$, which we denote by $\gamma \cdot \bar{s}'$. From the compatibility of the $\gamma_{S', \bar{s}'}$'s, it follows that γ defines an endomorphism of the functor $F_{\bar{s}}^\bullet$.

On the other hand, one has a homeomorphism

$$\text{inv} : \varprojlim \pi_1^\bullet(S, \bar{s}) / \text{Stab}_{S', \bar{s}'} \rightarrow \varprojlim \text{Stab}_{S', \bar{s}'} \setminus \pi_1^\bullet(S, \bar{s})$$

induced by $g \mapsto g^{-1}$. If we copy the construction $\underline{\gamma} \mapsto \gamma$ starting from $\text{inv}(\underline{\gamma})$ and using right actions instead of left actions, the resulting endomorphism of $F_{\bar{s}}^\bullet$ is nothing but the inverse of γ . Hence γ is an element of $\pi_1^\bullet(S, \bar{s})$, and it is clear that $\underline{\gamma} \mapsto \gamma$ is inverse to the natural map $\pi_1^\bullet(S, \bar{s}) \rightarrow \varprojlim \pi_1^\bullet(S, \bar{s}) / \text{Stab}_{S', \bar{s}'}$. \square

1.4.3. It is clear from the definitions that $F_{\bar{s}}^\bullet$ may be enriched to a functor

$$\text{Cov}_S^\bullet \longrightarrow \pi_1^\bullet(S, \bar{s})\text{-Sets}, \quad S' \mapsto F_{\bar{s}}^\bullet(S')$$

where $\pi_1^\bullet(S, \bar{s})\text{-Sets}$ is the category of discrete sets endowed with a continuous left action of $\pi_1^\bullet(S, \bar{s})$.

Proposition 1.4.4. *Up to isomorphism (unique up to composition by inner automorphisms), $\pi_1^\bullet(S, \bar{s})$ does not depend on \bar{s} .*

PROOF. The space $\text{Isom}(F_{\bar{s}}^\bullet, F_{\bar{t}}^\bullet)$, non-empty due to 1.3.3, is formally a $\pi_1^{\text{ét}}(S, \bar{t}) - \pi_1^{\text{ét}}(S, \bar{s})$ -bitorsor, and the result follows. \square

Theorem 1.4.5. *The enriched functor $F_{\bar{s}}^\bullet$*

$$\text{Cov}_S^\bullet \longrightarrow \pi_1^\bullet(S, \bar{s})\text{-Sets}$$

is fully faithful, and extends to an equivalence of categories

$$\{\text{disjoint unions of objects of } \text{Cov}_S^\bullet\} \longrightarrow \pi_1^\bullet(S, \bar{s})\text{-Sets}.$$

Connected coverings correspond to $\pi_1^\bullet(S, \bar{s})$ -orbits.

PROOF. (cf. [dJ95b, 2.10] in the étale case). If $f : S' \rightarrow S$ is an étale covering in Cov_S^\bullet and \bar{s}', \bar{s}'' are elements of $F_{\bar{s}}^\bullet(S')$ in the same connected component of S' , proposition 1.3.3 ensures the existence of an étale path α from \bar{s}' to \bar{s}'' . Then $f_*(\alpha) \in \pi_1^{\text{ét}}(S, \bar{s})$, and $f_*(\alpha) \cdot \bar{s}' = \bar{s}''$; of course, we may replace here $f_*(\alpha)$ by its image in $\pi_1^\bullet(S, \bar{s})$. This shows that $\pi_1^\bullet(S, \bar{s})$ -orbits in $F_{\bar{s}}^\bullet(S')$ correspond bijectively to connected components of S' . In particular

$$F_{\bar{s}}^{\bullet}(S') = \pi_1^{\bullet}(S, \bar{s}) / \text{Stab}_{S', \bar{s}'}^{\bullet}$$

if S' is connected.

This implies that the enriched functor $F_{\bar{s}}^{\bullet}$ is fully faithful, because if S' is connected, a morphism between étale coverings $S' \rightarrow S''$ of S may be identified via its graph with a connected component T of the étale covering $S' \times_S S''$ (which belongs to Cov_S^{\bullet} by assumption) such that $F_{\bar{s}}^{\bullet}(T) \xrightarrow{\sim} F_{\bar{s}}^{\bullet}(S')$ (this can be checked locally on S , and this reduces to a well-known fact about finite algebraic coverings).

It remains to show that any $\pi_1^{\bullet}(S, \bar{s})$ -orbit comes from a (connected) étale covering. By definition, such an orbit may be written $\pi_1^{\bullet}(S, \bar{s})/H$, where H is a subgroup containing the stabilizer $\text{Stab}_{S', \bar{s}'}^{\bullet}$ of some geometric point \bar{s}' above \bar{s} in some connected étale covering S' of S . As we have seen, the connected components of $S' \times_S S'$ correspond bijectively to $\pi_1^{\bullet}(S, \bar{s})$ -orbits in $F_{\bar{s}}^{\bullet}(S' \times_S S') = F_{\bar{s}}^{\bullet}(S')^2$. The union $R \subset S' \times_S S'$ of connected components which corresponds to the union of points $(\bar{s}', \bar{s}' \cdot h)$, $h \in H$, is an equivalence relation on S' over S . By lemma 1.2.8, the quotient $S'' = S'/R$ is an étale covering of S which belongs to Cov_S^{\bullet} by assumption. If \bar{s}'' denotes the image of \bar{s}' in S'' , it is clear that $H = \text{Stab}_{S'', \bar{s}''}^{\bullet}$. \square

1.4.6. A connected étale covering $S' \rightarrow S$ is *Galois* if S is the quotient $S' / (\text{Aut}_S S')$; $\text{Aut}_S S'$ is called the *Galois group* of the covering. By [Ber93, 4.1.9], an étale Galois covering of S with group G is the same as a connected principal homogeneous space under the constant S -group $G_S^{(1)}$. Through the dictionary of 1.4.5, a Galois étale covering in Cov_S^{\bullet} corresponds to a surjective continuous homomorphism

$$\pi_1^{\bullet}(S, \bar{s}) \rightarrow G,$$

with G discrete ($\cong \text{Aut}_S S'$). In other words, a (pointed) covering (S', \bar{s}') in Cov_S^{\bullet} is Galois if and only if $\text{Stab}_{S', \bar{s}'}^{\bullet} \subset \pi_1^{\bullet}(S, \bar{s})$ is normal.

Example: the logarithm (1.2.6. (i)) is a Galois étale covering of $(\mathbb{A}_{\mathbb{C}_p}^1)^{\text{an}}$ with group $\mu_{p^\infty} \cong \mathbb{Q}_p / \mathbb{Z}_p$.

The following is also clear:

Corollary 1.4.7. $\pi_1^{\bullet}(S, \bar{s})$ is a pro-discrete group if and only if in Cov_S^{\bullet} any connected covering is dominated by a Galois covering.

Remark. This property holds in $\text{Cov}_S^{\text{top}}$ and $\text{Cov}_S^{\text{alg}}$, but not in Cov_S^{et} in general. Let for instance $S = \mathbb{P}_{\mathbb{C}_p}^1$, and let us consider the Galois covering given by the Gross-Hopkins period mapping (II.6.3.3). According to [dJ95b, 7.4] (using the viewpoint sketched in II.6.4), the corresponding discrete quotient of $\pi_1^{\text{et}}(\mathbb{P}^1, \bar{s})$ is the composition

⁽¹⁾ *loc. cit.* 4.1.4, which is used in the proof, is available since we are dealing with strictly analytic spaces, *cf. loc. cit.* 4.1.5.

$$\pi_1^{\text{et}}(\mathbb{P}^1, \bar{s}) \rightarrow SL_2(\mathbb{Q}_p) \rightarrow SL_2(\mathbb{Q}_p)/SL_2(\mathbb{Z}_p)$$

where the first map is a continuous surjective homomorphism, and the quotient topology on $SL_2(\mathbb{Q}_p)$ is non-discrete (*loc. cit.* p. 116). Since the biggest normal subgroup of $SL_2(\mathbb{Q}_p)$ contained in $SL_2(\mathbb{Z}_p)$ is $\{\pm \text{id}\}$, we see that its preimage in $\pi_1^{\text{et}}(\mathbb{P}^1, \bar{s})$ cannot be open. Thus $\pi_1^{\text{et}}(\mathbb{P}^1, \bar{s})$ is *not* a pro-discrete group.

Corollary 1.4.8. *Let $\text{Cov}_S^{\bullet\bullet}$ be a full subcategory of Cov_S^\bullet stable under taking connected components, fiber products (over S) and quotients. Then the corresponding continuous homomorphism $\pi_1^\bullet(S, \bar{s}) \rightarrow \pi_1^{\bullet\bullet}(S, \bar{s})$ has dense image (it is neither strict⁽²⁾ nor surjective in general).*

In particular, if $\text{Cov}_S^{\text{alg}} \subset \text{Cov}_S^\bullet$, then $\pi_1^{\text{alg}}(S, \bar{s})$ is the profinite completion⁽³⁾ of $\pi_1^\bullet(S, \bar{s})$.

PROOF. By lemma 1.4.2, it suffices to prove that for any S' in $\text{Cov}_S^{\bullet\bullet}$ and any $\bar{s}' \in F_{\bar{s}}^\bullet(S') = F_{\bar{s}}^{\bullet\bullet}(S')$, the natural map $\pi_1^\bullet(S, \bar{s}) \rightarrow \pi_1^{\bullet\bullet}(S, \bar{s})/\text{Stab}_{S', \bar{s}'}^{\bullet\bullet}$ is surjective. We may replace S' by the connected component of s' . According to the previous theorem, $\pi_1^{\bullet\bullet}(S, \bar{s})/\text{Stab}_{S', \bar{s}'}^{\bullet\bullet}$ and $\pi_1^\bullet(S, \bar{s})/\text{Stab}_{S', \bar{s}'}^\bullet$ are then identified to $F_{\bar{s}}^\bullet(S') = F_{\bar{s}}^{\bullet\bullet}(S')$, whence the first assertion. The second assertion is clear, since $\pi_1^{\text{alg}}(S, \bar{s})$ is profinite.

To see that the image of $\pi_1^\bullet(S, \bar{s}) \rightarrow \pi_1^{\bullet\bullet}(S, \bar{s})$ is neither strict nor surjective in general, it suffices to take S such that $\pi_1^{\text{top}}(S, s)$ is infinite (*e.g.* a Tate curve). Then $\pi_1^{\text{et}}(S, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s})$ is neither surjective nor strict. This follows from the commutative square of topological groups

$$\begin{array}{ccc} \pi_1^{\text{et}}(S, \bar{s}) & \longrightarrow & \pi_1^{\text{alg}}(S, \bar{t}) \\ \downarrow & & \downarrow \\ \pi_1^{\text{top}}(S, s) & \longrightarrow & \widehat{\pi_1^{\text{top}}(S, t)} \end{array}$$

where the roof denotes the profinite completion; the coimage (*resp.* image) of the “south-east map” $\pi_1^{\text{top}}(S, \bar{s}) \rightarrow \widehat{\pi_1^{\text{top}}(S, t)}$ is infinite discrete (*resp.* compact). \square

In this context, the following general lemma is useful:

Lemma 1.4.9. *Let G, H be separated topological groups such that any neighborhood of 1 contains an open subgroup, and let $f : G \rightarrow H$ be a continuous homomorphism.*

⁽²⁾recall that a continuous homomorphism between topological groups is strict if the induced continuous bijective homomorphism between the coimage and the image is an isomorphism, i.e. bicontinuous; the coimage is the quotient of the source by the kernel, with the quotient topology.

⁽³⁾the profinite completion of a topological group is the inverse limit of its quotients by *open* normal subgroups of finite index.

- (i) If the induced functor $f^* : H\text{-Sets} \rightarrow G\text{-Sets}$ is essentially surjective, then f is injective.
- (ii) If $f^* : H\text{-Sets} \rightarrow G\text{-Sets}$ is an equivalence, then f is injective, strict and has dense image.
- (iii) If $f^* : H\text{-Sets} \rightarrow G\text{-Sets}$ is an equivalence and $G = \varprojlim G/U$ (the limit running over open subgroups of G), then f is an isomorphism.
- (iv) If G is complete and if the normal open subgroups are cofinal among the open subgroups U , then $G = \varprojlim G/U$.

PROOF. (i) otherwise, let $g \neq 1$ be in the kernel of f . Because G is separated, there exists a neighborhood U of 1 not containing g , and by hypothesis, one may assume that U is an open subgroup of G . By assumption the (connected) G -set G/U corresponds via f^* to a (connected) H -set X . Since $f(g) = 1$, g acts trivially on $f^*(X)$, in contradiction with the fact that $g \notin U$.

(ii) To show that f has dense image, it suffices to show that for $h \in H$ and any neighborhood V of 1, $f(G).h \cap V \neq \emptyset$. By hypothesis, one may assume that V is an open subgroup of H . Thus one has to show that $H = G.V$ for any open subgroup V of H , i.e. G acts transitively on H/V . If the action is not transitive, $f^*(H/V)$ can be written as a disjoint union of G -sets, which contradicts the the assumption that f^* is an equivalence. It remains to show that f is strict. But we have seen that f^* induces an equivalence on transitive sets. Therefore f^{-1} induces a bijection between open subgroups of H and open subgroups of G , whence the result.

(iii) Indeed one has a commutative diagram

$$\begin{array}{ccc} G & \xrightarrow{\subset} & H \\ \wr \downarrow & & \downarrow \\ \varprojlim G/U & \xrightarrow{\sim} & \varprojlim H/V \end{array}$$

and the right vertical map is injective since H is separated.

(iv) follows from [Bou74, III.7.2, prop. 2]. \square

1.4.10. Let $f : T \rightarrow S$ be any morphism of connected K -manifolds. We have the natural functor

$$\text{Cov}_S^{\text{et}} \rightarrow \text{Cov}_T^{\text{et}} : S' \mapsto S' \times_S T.$$

Let us *assume* that it induces a functor

$$f^{\bullet\bullet} : \text{Cov}_S^{\bullet} \rightarrow \text{Cov}_T^{\bullet}.$$

Since $F_{S,\bar{s}}^* = F_{T,\bar{t}}^* \circ f^{\bullet\bullet}$ if $\bar{s} = f(\bar{t})$, this gives rise to a continuous homomorphism

$$f_{\bullet} : \pi_1^{\bullet}(T, \bar{t}) \rightarrow \pi_1^{\bullet}(S, \bar{s} = f(\bar{t})).$$

Corollary 1.4.11. f_*^\bullet is an isomorphism of topological groups if and only if $f^{\bullet*}$ is an equivalence of categories.

PROOF. If f_*^\bullet is an isomorphism, it induces an equivalence $\pi_1^\bullet(S, \bar{s})\text{-Sets} \cong \pi_1^\bullet(T, \bar{t})\text{-Sets}$, and it follows from 1.4.5 that $f^{\bullet*}$ is an equivalence of categories. The other implication is immediate. \square

Corollary 1.4.12. Assume that f is an étale covering in Cov_S^\bullet , thus corresponding to the $\pi_1^\bullet(S, \bar{s})$ -orbit $\pi_1^\bullet(S, \bar{s})/\text{Stab}_{T, \bar{t}}^\bullet$ through the dictionary of 1.4.5.

- (a) Then $\text{Stab}_{T, \bar{t}}^\bullet$ is the closure of the image of f_*^\bullet . In particular, if f is a Galois covering, its Galois group is $\pi_1^\bullet(S, \bar{s})/(\text{Im } f_*^\bullet)^\ominus$.
- (b) Assume moreover that for any covering $T' \rightarrow T$ in Cov_T^\bullet , the composition with f gives rise to a covering $T' \rightarrow S$ in Cov_S^\bullet . Then f_*^\bullet is open and injective (hence $\pi_1^\bullet(T, \bar{t})$ may be identified with the open and closed subgroup $\text{Stab}_{T, \bar{t}}^\bullet$ of $\pi_1^\bullet(S, \bar{s})$).

PROOF. (a) Looking at the natural identifications

$$\pi_1^\bullet(S, \bar{s})/\text{Stab}_{T, \bar{t}}^\bullet \cong F_{\bar{s}}^\bullet(T) \cong F_{\bar{t}, \bar{t}}^\bullet(T \times_S T)$$

of $\pi_1^\bullet(S, \bar{s})$ -orbits. The connected component of $T \times_S T$ containing (\bar{t}, \bar{t}) is the diagonal $T \subset T \times_S T$, therefore the $\pi_1^\bullet(T, \bar{t})$ -orbit is reduced to the point (\bar{t}, \bar{t}) . This shows that $\text{Im } f_*^\bullet$ lies in $\text{Stab}_{T, \bar{t}}^\bullet$.

To show that it is dense, we have to show that for any connected S' in Cov_S^\bullet and any geometric point \bar{s}' of S' above \bar{s} , the map $\pi_1^\bullet(T, \bar{t}) \rightarrow \pi_1^\bullet(S, \bar{s})/\text{Stab}_{T, \bar{t}}^\bullet$ induced by f_*^\bullet is surjective. This follows from 1.4.5 as in 1.4.8.

(b) Under the extra assumption, one has functors $\text{Cov}_T^\bullet \rightarrow \text{Cov}_S^\bullet \rightarrow \text{Cov}_T^\bullet : T'/T \mapsto T'/S; S'/S \mapsto (S' \times_S T)/T$. For any open subgroup $H \subset \pi_1^\bullet(T, \bar{t})$ (corresponding as in 1.4.5 to a pointed connected covering (T', \bar{t}')), $\pi_1^\bullet(T, \bar{t})/H \cong F_{\bar{t}'}^\bullet(T')$ is a subset of $F_{\bar{t}', \bar{t}}^\bullet(T' \times_S T) \cong F_{\bar{s}}^\bullet(T') \cong \pi_1^\bullet(S, \bar{s})/\text{Stab}_{T', \bar{t}'}^\bullet$. In other words, f_*^\bullet induces an open injective homomorphism

$$\pi_1^\bullet(T, \bar{t})/H \hookrightarrow \pi_1^\bullet(S, \bar{s})/f_*^\bullet(H).$$

Indeed, $f_*^\bullet(H) = \text{Stab}_{T', \bar{t}'}^\bullet$ is open in $\pi_1^\bullet(S, \bar{s})$: such groups correspond to connected coverings of S which dominate T (*i.e.* which factorize over f). Since such coverings are cofinal in Cov_S^\bullet , we can pass to the limit, using 1.4.2:

$$\pi_1^\bullet(T, \bar{t}) = \varprojlim \pi_1^\bullet(T, \bar{t})/H \hookrightarrow \varprojlim \pi_1^\bullet(S, \bar{s})/f_*^\bullet(H) \cong \pi_1^\bullet(S, \bar{s})$$

and conclude that f_*^\bullet is an open injective homomorphism. \square

1.5. Example: Ramero's locally algebraic étale fundamental group.

1.5.1. When S is countable at infinity, L. Ramero has introduced an interesting category $\text{Cov}_S^{\text{loc.alg}}$ of étale coverings, called ‘locally algebraic’, which generalize the logarithmic covering of \mathbb{A}^1 , and which are intimately connected with the local study of meromorphic differential equations around singularities [Ram98, 4].

An étale covering is locally algebraic if the preimage of every *compact subdomain* $W \subset S$ is a disjoint union $\coprod W_i$ of subdomains W_i which are finite étale over W . Since S is (by assumption) a union of an increasing countable family of compact subdomains S_j , $j \geq 1$, it suffices to check the condition for $W = S_j, \forall j$.

This category is stable under taking connected components, fiber products (over S) and quotients. Moreover, it is stable by disjoint union and ‘composition’. Assuming that $s \in S_1$, one has a natural homomorphism

$$\varinjlim \pi_1^{\text{loc.alg}}(S_j, \bar{s}) = \varinjlim \pi_1^{\text{alg}}(S_j, \bar{s}) \rightarrow \pi_1^{\text{loc.alg}}(S, \bar{s}).$$

Proposition 1.5.2. *This is an isomorphism: $\pi_1^{\text{loc.alg}}(S, \bar{s}) \cong \varinjlim \pi_1^{\text{alg}}(S_j, \bar{s})$.*

PROOF. It is clear that an automorphism of $F_{S, \bar{s}}^{\text{loc.alg}}$ amounts to a compatible family of automorphisms of $F_{S_j, \bar{s}}^{\text{alg}}$, hence the homomorphism is bijective. Moreover, the induced map $\varinjlim \pi_1^{\text{alg}}(S_j, \bar{s})\text{-sets} \rightarrow \pi_1^{\text{loc.alg}}(S, \bar{s})\text{-sets}$ is an equivalence: both sides are equivalent to $\text{Cov}_S^{\text{loc.alg}}$, cf. 1.4.5 and [Ram98, 4.1.6]. Therefore open subgroups correspond to stabilizers as usual, and the bijection $\pi_1^{\text{loc.alg}}(S, \bar{s}) \cong \varinjlim \pi_1^{\text{alg}}(S_j, \bar{s})$ is bicontinuous. \square

It is not known whether $\pi_1^{\text{loc.alg}}(S, \bar{s})$ is a pro-discrete group (cf. in this direction *loc. cit.* 4.3).

Example. The logarithm of any Lubin-Tate formal group over a p -adic field gives rise to a locally algebraic étale covering of the analytic affine line, cf. *loc. cit.* 6.1.1.

On the other hand, the Gross-Hopkins covering of \mathbb{P}^1 is not locally algebraic, although there is a finite affinoid cover (V_i) of \mathbb{P}^1 such that the restriction of the covering to any V_i is a disjoint union of finite étale coverings.

2. Temperate fundamental groups.

ABSTRACT: We introduce the notion of temperate étale covering. Such coverings are essentially built from (possibly infinite) topological coverings of finite étale coverings; they are classified by temperate fundamental groups, which seem to be the right non-archimedean equivalents of fundamental groups of complex manifolds. These groups are not discrete, but nevertheless often possess many infinite discrete quotients.

2.1. Temperate étale coverings and temperate fundamental groups.

In the archimedean context, the categories $\text{Cov}_S^{\text{ét}}$ and $\text{Cov}_S^{\text{top}}$ coincide; in the non-archimedean situation, in contrast, $\text{Cov}_S^{\text{top}}$ is “too small” (for instance, it consists of disjoint unions of copies of S if S has dimension one and tree-like reduction, *e.g.* if S is an annulus), while $\text{Cov}_S^{\text{ét}}$ is “too big” (for $S = \mathbb{P}^1$, it contains many non-trivial objects).

We introduce an intermediate category, which seems to be a closer analogue of the category of unramified coverings in the archimedean context.

Definition 2.1.1. An étale covering $S' \rightarrow S$ is temperate if it is a quotient⁽⁴⁾ of a composite étale covering $T' \rightarrow T \rightarrow S$, where $T' \rightarrow T$ is a (possibly infinite) topological covering, and $T \rightarrow S$ is a finite étale covering.

Hence we have the commutative square

$$\begin{array}{ccc} & T' & \\ & \swarrow & \searrow \\ T & & S' \\ & \searrow & \swarrow \\ & S & \end{array}$$

with $T' \in \text{Cov}_T^{\text{top}}$, $T \in \text{Cov}_S^{\text{alg}}$, $S' \in \text{Cov}_S^{\text{ét}}$.

Remark. The notion of temperate covering is stable under base change as in 1.4.10.

Lemma 2.1.2. *Let $T \rightarrow S$ be a finite Galois étale covering, and let $\tilde{T} \rightarrow T$ be the universal topological covering of T . Then the composite $\tilde{T} \rightarrow S$ is a Galois temperate covering.*

PROOF. From 1.2.5, we know that it is an étale covering, automatically temperate. It remains to show that it is Galois. By the universal property of \tilde{T} , any automorphism of T lifts to \tilde{T} , hence there is an exact sequence of discrete groups:

$$1 \rightarrow (\pi_1^{\text{top}}(T, t) \cong) \text{Aut}_T \tilde{T} \rightarrow \text{Aut}_S \tilde{T} \rightarrow \text{Aut}_S T \rightarrow 1.$$

Since $\tilde{T}/(\text{Aut}_T \tilde{T}) = T$ and $T/(\text{Aut}_S T) = S$, it follows that $\tilde{T}/(\text{Aut}_S \tilde{T}) = S$. \square

⁽⁴⁾in the sense of 1.2.8.

We see from the lemma that in the definition of a temperate covering, we may assume that $T' \rightarrow S$ and $T' \rightarrow T$ are Galois.

Remarks 2.1.3. (i) It is *not true* that any quotient of a tower $T' \rightarrow T \rightarrow S$ as in definition 2.1.1 is again of this form. As a counter-example, let us consider a Tate elliptic curve E over $K = \mathbb{C}_p$, and set $T = E \setminus E[2]$ (2-torsion points removed). Then \tilde{T} is \mathbb{G}_m^{an} deprived from the countable subgroup $\pm(\sqrt{q})^{\mathbb{Z}}$, and $T \cong \tilde{T}/q^{\mathbb{Z}}$. The inversion $x \mapsto 1/x$ on \tilde{T} induces the inversion on E , and $\mathbb{G}_m/\{\text{id}, \text{inv}\} = \mathbb{A}^1$ (via $x \mapsto x + 1/x$), $E/\{\text{id}, \text{inv}\} = \mathbb{P}^1$. Let us consider the infinite temperate covering

$$S' = \tilde{T}/\{\text{id}, \text{inv}\} \rightarrow S = T/\{\text{id}, \text{inv}\} = \mathbb{P}^1 \setminus \{0, 1, \lambda, \infty\}.$$

Then S' is not a topological covering of any finite étale covering S'' of S . Indeed, otherwise \tilde{T} would be the universal topological covering of some connected component T' of $T \times_S S''$. The tower $\tilde{T} \rightarrow T' \rightarrow T$ would then extend to a tower of topological coverings $\mathbb{G}_m^{\text{an}} \rightarrow E' \rightarrow E$, where E' is another Tate curve isogenous to E (we use the fact that the topological fundamental group of T' and E is the same). Then $S'' \subset E'/\{\text{id}, \text{inv}\} \cong \mathbb{P}^1$. But any open subset in $\mathbb{P}_{\mathbb{C}_p}^1$ is simply-connected, whence $\tilde{T} = S''$, a contradiction.

This example also shows that *a temperate covering may become a topological covering after finite étale base change, without being itself a topological covering.*

(ii) The notion of temperate covering is not local on S . In fact, the Gross-Hopkins étale covering of $(\mathbb{P}_{\mathbb{C}_p}^1)^{\text{an}}$ is not temperate, although there is a finite open cover (U_i) of $(\mathbb{P}_{\mathbb{C}_p}^1)^{\text{an}}$ such that it becomes a disjoint union of finite étale coverings over each U_i (by compactness).

2.1.4. We denote the category of temperate (étale) coverings of S by $\text{Cov}_S^{\text{temp}}$. From the fact that topological and finite étale coverings respectively are stable under taking connected components and fiber products, it is not difficult to see that $\text{Cov}_S^{\text{temp}}$ is stable under taking connected components and fiber products (over S). By definition, it is stable under taking quotients.

According to the results of 1.4, there is thus defined a separated totally discontinuous group, the *temperate fundamental group* $\pi_1^{\text{temp}}(S, \bar{s})$, which does not depend on the geometric point \bar{s} up to isomorphism. Moreover:

Lemma 2.1.5. $\pi_1^{\text{temp}}(S, \bar{s})$ is a pro-discrete group (hence complete, cf. [Bou74, III. 57.7]). A basis of open neighborhoods of 1 is given by normal closed subgroups H such that $\pi_1^{\text{temp}}(S, \bar{s})/H$ is the Galois group of the universal topological covering of some finite étale Galois covering of S .

PROOF. This follows from 1.4.7. and 2.1.2 (connected temperate coverings admit a Galois closure). \square

The inclusion $\text{Cov}_S^{\text{top}} \hookrightarrow \text{Cov}_S^{\text{temp}}$ gives rise to a *surjective* homomorphism

$$\pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{top}}(S, s)$$

(*cf.* 1.4.8, taking into account the discreteness of $\pi_1^{\text{top}}(S, s)$), which shows that $\pi_1^{\text{temp}}(S, \bar{s})$, although not a discrete group, has many *infinite discrete quotients* in general; we say that $\pi_1^{\text{temp}}(S, \bar{s})$ is “lacunary”.

On the other hand, the inclusion $\text{Cov}_S^{\text{alg}} \hookrightarrow \text{Cov}_S^{\text{temp}}$ gives rise to a continuous homomorphism with dense image

$$\pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s}),$$

and $\pi_1^{\text{alg}}(S, \bar{s})$ may be identified with the profinite completion of $\pi_1^{\text{temp}}(S, \bar{s})$. This shows that $\pi_1^{\text{temp}}(S, \bar{s})$ is not “too small” (as opposed to $\pi_1^{\text{top}}(S, s)$), and the following two propositions show that it is not “too big” as well (as opposed to $\pi_1^{\text{et}}(S, \bar{s})$):

Proposition 2.1.6. *If $\dim S = 1$, the homomorphism $\pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s})$ is injective.*

PROOF. By 2.1.2 and 2.1.5, it suffices to see that if $T \rightarrow S$ is a finite Galois étale covering, the Galois group of the temperate covering $\tilde{T} \rightarrow S$ is residually finite. But this group contains $\pi_1^{\text{top}}(T, t)$ as a normal subgroup of finite index. The point is that in dimension one, topological fundamental groups are free, hence residually finite. \square

The lacunary subgroup $\pi_1^{\text{temp}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\})$ of the profinite free group on two generators $\pi_1^{\text{alg}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\})$ is especially interesting and mysterious, and will be the object of our investigations in later sections⁽⁵⁾.

Proposition 2.1.7. *If S is algebraic and K of characteristic zero, the Galois group of any Galois temperate covering of S is finitely generated. Moreover, if K has only countably many finite extensions in a fixed algebraic closure \bar{K} (e.g. if K is a p -adic field), then $\pi_1^{\text{temp}}(S, \bar{s})$ and all its quotients by closed normal subgroups are ‘polish’, i.e. metrizable of countable type and complete.*

PROOF. Let us consider the first assertion. By 2.1.2, it suffices to treat the case of the universal topological covering of a finite Galois étale covering. By the already mentioned Gabber-Lütkebohmert theorem, that finite étale covering is algebraic. The Galois group admits as a normal subgroup of finite index the topological fundamental group of that finite étale covering, which is finitely generated (*cf.* 1.1.3); whence the result.

Since $\pi_1^{\text{temp}}(S, \bar{s})$ is separated, its metrizability is equivalent to the existence of a countable fundamental system of neighborhood of 1 [Bou74, IX.3.1

⁽⁵⁾recall, on the other hand, that $\pi_1^{\text{top}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\})$ is trivial.

prop. 1]. By 2.1.5, it suffices to see that there are countably many finite étale coverings of S , which follows, by dévissage, from the assumption that $\text{Gal}(\overline{K}/K)$ has countably many open subgroups, and from the topological finite generation of the algebraic fundamental group over \overline{K} . This also shows that $\pi_1^{\text{temp}}(S, \overline{s})$ is not only a pro-discrete group (which already implies that it is complete), but even a countable inverse limit of discrete countable groups. It follows that the metrizable group $\pi_1^{\text{temp}}(S, \overline{s})$ admits a countable dense subset, hence its topology has a countable basis [Bou74, IX.2.8 prop. 12].

It follows that any closed subgroup of $\pi_1^{\text{temp}}(S, \overline{s})$ and any quotient of $\pi_1^{\text{temp}}(S, \overline{s})$ by a closed normal subgroup is metrizable and complete [Bou74, IX.3.1 prop. 4], and of countable type (for a quotient: because it has a countable dense subset). \square

Notice however that the first assertion does not imply that $\pi_1^{\text{temp}}(S, \overline{s})$ is topologically finitely generated. On the other hand, we shall show below that it is not locally compact in general (2.3.12).

Proposition 2.1.8. *Let $f : T \rightarrow S$ be an étale finite Galois covering of a connected K -manifold with group G , and \bar{t} be a geometric point of T . Then there is an exact sequence*

$$1 \rightarrow \pi_1^{\text{temp}}(T, \bar{t}) \rightarrow \pi_1^{\text{temp}}(S, f(\bar{t})) \rightarrow G \rightarrow 1.$$

PROOF. It is clear from the definition 2.1.1 that any temperate covering of T gives rise, by composition with f , to a temperate covering of S . Thus the proposition follows from 1.4.12.b. \square

Remark. We shall see below (2.3.3) that on the other hand, an étale covering of S which is a finite étale covering of a topological covering is not necessarily temperate.

The following result is an important criterion to recognize topological, *resp.* temperate, *resp.* locally algebraic coverings:

Theorem 2.1.9. *Let S' be a Galois étale covering of S with group G . Then*

- (a) *if G is torsion-free, S' is a topological covering of S ;*
- (b) *if G is virtually torsion-free, S' is a temperate covering of S ; more precisely, S' is a topological covering of some finite étale Galois covering of S ;*
- (c) *assume that S is countable at infinity; then S' is a locally algebraic covering of S if and only if G is the union of an increasing sequence of finite subgroups.*

PROOF. (a) (cf. also [dJ95b]) Let $\rho : \pi_1^{\text{et}}(S, \overline{s}) \rightarrow G$ be the continuous homomorphism corresponding $S' \rightarrow S$. Let \bar{t} be another geometric point of S , and let $\overline{\gamma}$ be an étale path from \bar{t} to \overline{s} . Under the composed continuous

homomorphism

$$\pi_1^{\text{et}}(t, \bar{t}) \rightarrow \pi_1^{\text{et}}(S, \bar{t}) \xrightarrow{\text{ad}(\bar{\gamma})} \pi_1^{\text{et}}(S, \bar{s}) \rightarrow G$$

the compact (profinite) group $\pi_1^{\text{et}}(t, \bar{t}) \cong \text{Gal}(\mathcal{H}(t)^{\text{sep}}/\mathcal{H}(t))$ is sent to 1 if G is torsion-free. This means that for any $\bar{t}' \in F_{\bar{t}}(S')$, $\mathcal{H}(t') = \mathcal{H}(t)$. Let U be an open neighborhood of t (which we may assume to be contractible thanks to Berkovich's theorem), such that $S' \times_S U \cong \coprod V_j$, with V_j finite étale over U . Since $\mathcal{H}(t') = \mathcal{H}(t)$ for any point t' in V_j above t , it follows from 1.2.1 that V_j is a topological covering of the contractible open set U , hence a disjoint union of finitely many copies of U . This holds for any point $t \in S$, therefore $S' \rightarrow S$ is a topological covering.

(b) If G is virtually torsion-free, let N be a torsion-free normal subgroup of finite index, and let $S' \rightarrow T \rightarrow S$ be the corresponding intermediate Galois covering. By *a*), $S' \rightarrow T$ is a topological covering, hence $S' \rightarrow S$ is a temperate étale covering.

(c) Assume that S' corresponds to a principal homogeneous space under G_S , classified by an element of $H^1(S, G)$. Let us write $S = \bigcup S_j$, with S_j compact. Assume that $G = \bigcup G_k$, with G_k finite. Then $H^1(S_j, G) = \varinjlim H^1(S_j, G_k)$, therefore the restriction of S' to any S_j is a disjoint union of finite étale coverings of S_j , hence $S' \rightarrow S$ is locally algebraic. Conversely, if $S' \rightarrow S$ is Galois locally algebraic, its group G is a discrete image of $\varinjlim \pi_1^{\text{alg}}(S_j, \bar{s})$, each $G_j := \text{Im}(\pi_1^{\text{alg}}(S_j, \bar{s}))$ is finite, and $G = \bigcup G_j$. \square

Corollary 2.1.10. *In dimension one, $\pi_1^{\text{temp}}(S, \bar{s})$ is the pro-VTFD-completion of $\pi_1^{\text{et}}(S, \bar{s})$.*

By pro-VTFD-completion of a topological group Γ , we mean the inverse limit of its virtually torsion-free discrete quotients. This is a filtered limit (if Γ/H' and Γ/H'' are VTFD, so is $\Gamma/(H' \cap H'')$). Pro-VTFD-completions are functorial (due to the fact that any subgroup of a VTF group is VTF).

PROOF. By 2.1.2, the coverings of the form $\tilde{T} \rightarrow T \rightarrow S$, with T finite étale Galois over S , are cofinal among temperate coverings. In dimension one, we have seen that the Galois groups of such Galois coverings are virtually torsion-free, whence the result (using 1.3.2). \square

Examples. (i) It follows from 2.1.10 that the logarithmic (locally algebraic) covering

$$\log : D_{\mathbb{C}_p}(1, 1^-) \longrightarrow \mathbb{A}_{\mathbb{C}_p}^1$$

of the affine line is *not* temperate, since its Galois group $\mathbb{Q}_p/\mathbb{Z}_p$ has no non-trivial virtually torsion-free quotient. This also follows from 2.1.6, since $\pi_1^{\text{alg}}(\mathbb{A}_{\mathbb{C}_p}^1) = 0$.

Remark. It is easy to draw from 2.1.9 that a connected étale covering which is both temperate and locally algebraic is (finite) algebraic.

(ii) It also follows from 2.1.9 that there is no étale Galois covering of the punctured disk with Galois group \mathbb{Z} . This fact has noteworthy consequences on the local monodromy of p -adic differential equations (*cf.* next sections).

Remark. In [And98], we have limited ourselves to temperate coverings for which the tower $T' \rightarrow T \rightarrow S$ occurring in definition 2.1.1 is Galois with VTF group. The fundamental attached to the category of such coverings is then the pro-VTFD-completion of $\pi_1^{\text{ét}}(S, \bar{s})$. The definition 2.1.1 seems however more natural (and is much more general in higher dimension). In [And98], another topological group is introduced, namely the coimage of $\pi_1^{\text{ét}}(S, \bar{s}) \rightarrow \pi_1^{\text{top}}(S, \bar{s}) \times \pi_1^{\text{alg}}(S, \bar{s})$, denoted by $\pi_1^{\text{red}}(S, \bar{s})$. In dimension one, there is a continuous injective homomorphism with dense image $\pi_1^{\text{red}}(S, \bar{s}) \rightarrow \pi_1^{\text{temp}}(S, \bar{s})$, but it not clear whether this is an isomorphism (the question is the openness of $\pi_1^{\text{ét}}(S, \bar{s}) \rightarrow \pi_1^{\text{temp}}(S, \bar{s})$).

Theorem 2.1.11. *Assume that K is algebraically closed of characteristic zero. Let \bar{S} be a (paracompact strictly) normal connected K -analytic space \bar{S} , and let Z be a closed analytic subset such that $S := \bar{S} \setminus Z$ is smooth. Assume that for any point s of \bar{S} , there is a covering (in the sense of 1.2.3) from a K -manifold onto a neighborhood $U(s)$ of s which is étale above $U(s) \cap S$. Then any temperate étale covering $S' \rightarrow S$ extends to a (possibly ramified) covering $\bar{S}' \rightarrow \bar{S}$, in the sense of 1.2.3: any point x has an open neighborhood $U(x)$ such that $S' \times_S U(x)$ is a disjoint union of finite (ramified) coverings V_j of $U(x)$. Moreover, one may assume that the V_j are normal; such an extension $\bar{S}' \rightarrow \bar{S}$ is then unique.*

Furthermore, if Z has codimension ≥ 2 and \bar{S} is smooth, any temperate étale covering $S' \rightarrow S$ extends to a unique temperate covering $\bar{S}' \rightarrow \bar{S}$.

PROOF. We start with the statement about unicity. Since it is local on \bar{S} , it is enough to establish it in the case of finite coverings (by definition 1.2.3). Let \bar{S}' and \bar{S}'' be two connected finite coverings of \bar{S} which coincide over S . The diagonal Δ is a connected component of $S' \times_S S''$. Its closure $\bar{\Delta}$ in the normalization of $\bar{S}' \times_{\bar{S}} \bar{S}''$ is again a connected component (using [Ber99, 3.3.16]). The projections $\bar{\Delta} \rightarrow \bar{S}'$, $\bar{\Delta} \rightarrow \bar{S}''$ are finite, and their restrictions to Δ are isomorphisms; hence they are themselves isomorphisms if \bar{S}' and \bar{S}'' are normal.

Let us turn to the existence of the extension (with the normality property). By unicity, this is a local question; we may and shall assume that Z is a hypersurface, and that \bar{S} is (strictly) affinoid, and that there exists a finite covering $\bar{T} \rightarrow \bar{S}$ by a rigid-analytically smooth affinoid \bar{T} , which restricts to a finite étale covering $T \rightarrow S$.

Let us first show that any quotient of a temperate covering $S' \rightarrow S$ which has the extension property inherits this extension property: by unicity, the question is again local on \bar{S} ; by definition of coverings, this reduces the question to the case when $S' \rightarrow S$ is finite. By glueing (*cf.* [Ber93, 1.3.3]), it is again enough to treat the affinoid (normal) case. The argument is similar to the above one for unicity and left to the reader.

This remark allows us to assume that $S' \rightarrow S$ is a topological covering of a finite étale covering of S . Let us first treat the case of a finite étale covering. In this case, the extension property is proved in Gabber-Lütkebohmert [Lüt93]. \square

The crucial case is the case of a punctured disk $S = \mathbb{D}^*$ which is worth stating separately:

Lemma 2.1.12 (Gabber, Lütkebohmert). *Any finite étale covering of \mathbb{D}^* becomes a finite disjoint union of Kummer coverings on a smaller concentric punctured disk.*

(Here the assumption that K is of characteristic zero is essential). From there, one draws the extension property, in the product case, when \overline{S} is a polydisk and Z a union of axes (*cf. loc. cit.*, 3.3). By Kiehl's theorem on the existence of tubular neighborhoods [Kie67b], the statement also holds if Z is a strict normal crossing divisor in \overline{S} . In the general case, one can invoke embedded resolution for Z , *cf.* [Sch99] (note that [Lüt93, §4] proposes an alternative strategy which uses embedded resolution only in dimension 2).

It remains to deal at last with the case of a topological covering $S' \rightarrow S$. In order to avoid problems of singularities, we consider a finite étale covering $T \rightarrow S$ as above, T being a rigid-analytically smooth affinoid. It will be more convenient to replace S' by $S' \times_S T$ (of which S' is a quotient) so that the new S' is a topological covering of T itself a finite étale covering of S . By 1.1.4, S' extends to a topological covering $\overline{S}' \rightarrow \overline{T}$, and composition with $\overline{T} \rightarrow \overline{S}$ provides the desired extension of S'/S to a covering $\overline{S}'/\overline{S}$.

The last statement of the theorem follows from the Nagata-Zariski purity theorem (which becomes easy in the case of a finite morphism of one-dimensional affinoids).

Remark. This extension property of temperate coverings does not hold for arbitrary étale coverings.

Indeed, one can show that the logarithmic covering of the affine line does not extend to a (ramified) covering of the projective line.

The previous theorem is completed by the following result, which is proven as in the complex case [SGA 1, exp. XII, 5.3]:

Proposition 2.1.13. *Let \overline{S} be a normal analytic space, and let Z be a closed analytic subset such that $S := \overline{S} \setminus Z$ is dense in \overline{S} . Then the functor which attaches to any finite normal covering of \overline{S} its restriction to S is fully faithful.*

It follows that in 2.1.11, the extension of coverings from temperate coverings of S to 'normal' coverings of \overline{S} 'is' a fully faithful functor.

2.2. Interlude: tangential base-points.

2.2.1. Instead of basing fundamental groups at geometric points, it is sometimes more convenient⁽⁶⁾ to base them at tangent vectors at points at infinity.

P. Deligne has explained the construction for affine algebraic curves in characteristic zero, for topological, algebraic (= profinite) and other kinds of fundamental groups [Del89]. In the profinite case, these tangential base-points amount to genuine geometric generic points, corresponding to embeddings of the function field of the curve into the field of Puiseux series at the chosen point at infinity (with respect to a fixed local parameter z at that point; however, the embedding depends only on the tangent vector $\partial/\partial z$).

If we try to imitate the construction in our non-archimedean setting, we face at once the following difficulty: let us assume for simplicity that the curve is the affine line; since there is no way to extend the Gauss norm on $K(z)$ to the field of Puiseux series with arbitrary coefficients in K , one does not get geometric (generic) points in the sense of 1.2.2. We do not know how to overcome this difficulty for (analytic) étale coverings in general (and we doubt that this is possible). For temperate étale coverings, however, we can circumvent the difficulty as follows.

2.2.2. We assume that K is algebraically closed of characteristic 0. Let \overline{S} be a K -manifold of dimension one, and let S be the complement of a *classical* point 0 in \overline{S} (so that 0 has a basis of open neighborhoods consisting in disks centered at 0). As above, it follows from 2.1.12 that *the restriction of any temperate covering of S to a punctured disk around 0 of sufficiently small radius splits into a disjoint union of Kummer coverings of bounded degree*. If we replace \overline{S} by its tangent space at 0 and 0 by the zero tangent vector, the same construction yields an equivalence of categories, with an obvious inverse given by ‘extension’. We thus get a *specialization functor*:

$$\mathrm{Cov}_S^{\mathrm{temp}} \rightarrow \mathrm{Cov}_{T_{\overline{S},0}^0}^{\mathrm{temp}}.$$

Any geometric point \vec{t} of $T_{\overline{S},0}^0$, e.g. any non-zero tangent vector at 0, then gives rise by composition to a fiber functor

$$F_{\vec{t}}^{\mathrm{temp}} : \mathrm{Cov}_S^{\mathrm{temp}} \rightarrow \mathrm{Sets}.$$

Proposition 2.2.3. *This fiber functor is isomorphic to $F_{\vec{t}}^{\mathrm{temp}}$ for any geometric point \vec{t} .*

PROOF. We follow the line of proof of 1.3.3. We may assume that \overline{S} is affinoid (hence compact). Let us consider finite open covers $\overline{\mathcal{U}} = (\overline{U}_i)_{i \geq 1}$ of S such that \overline{U}_1 is a disk centered at 0 and $0 \notin \overline{U}_i$ for $i > 1$.

Let us denote by $\mathcal{U} = (U_i)$ its trace on S , and by $\mathrm{Cov}_{S,\mathcal{U}}^{\mathrm{temp}}$ the full subcategory of $\mathrm{Cov}_S^{\mathrm{temp}}$ of temperate coverings $f : S' \rightarrow S$ such that $f^{-1}U_i = \coprod V_{ij}$ and V_{ij} is étale finite over U_i via f .

⁽⁶⁾for instance in the study of Galois actions on fundamental groups.

The point is that by 2.1.11 (or 2.2.2), we have $\text{Cov}_S^{\text{temp}} = \varinjlim_{\mathcal{U}} \text{Cov}_{S,\mathcal{U}}^{\text{temp}}$, where \mathcal{U} runs among the open covers of S satisfying the above properties. Also

$$\text{Isom}(F_{\vec{t}}^{\text{temp}}, F_{\vec{t}}^{\text{temp}}) = \varinjlim_{\mathcal{U}} \text{Isom}(F_{\vec{t},\mathcal{U}}^{\text{temp}}, F_{\vec{t},\mathcal{U}}^{\text{temp}})$$

where $F_{\vec{t},\mathcal{U}}^{\text{temp}}, F_{\vec{t},\mathcal{U}}^{\text{temp}}$ denote the restrictions of $F_{\vec{t}}^{\text{temp}}, F_{\vec{t}}^{\text{temp}}$ to $\text{Cov}_{S,\mathcal{U}}^{\text{temp}}$.

To show that $\text{Isom}(F_{\vec{t}}^{\text{temp}}, F_{\vec{t}}^{\text{temp}}) \neq \emptyset$, we construct non-empty compact subsets $K_{\mathcal{U}}$ of $\text{Isom}(F_{\vec{t},\mathcal{U}}^{\text{temp}}, F_{\vec{t},\mathcal{U}}^{\text{temp}})$ (topologized as before) such that if \mathcal{U}' refines \mathcal{U} , $K_{\mathcal{U}'}$ maps to $K_{\mathcal{U}}$ under the natural map $\text{Isom}(F_{\vec{t},\mathcal{U}'}^{\text{temp}}, F_{\vec{t},\mathcal{U}'}^{\text{temp}}) \rightarrow \text{Isom}(F_{\vec{t},\mathcal{U}}^{\text{temp}}, F_{\vec{t},\mathcal{U}}^{\text{temp}})$; it will follow that $\text{Isom}(F_{\vec{t}}^{\text{temp}}, F_{\vec{t}}^{\text{temp}}) \supset \varinjlim_{\mathcal{U}} K_{\mathcal{U}} \neq \emptyset$.

In fact, it suffices to do so for a cofinal system of \mathcal{U} 's. We now use a continuous embedding $I = [0, 1] \hookrightarrow \overline{S}$ with $0 \mapsto 0 \in \overline{S}$, $1 \mapsto t$. We consider open covers $\mathcal{U} = (U_1, \dots, U_n)$ as above, and such that

$$U_1 \cap I = 0, r_1[, \dots, U_i \cap I =]r_i, t_i[, \dots, U_m \cap I =]r_m, 1], U_{m+1} \cap I = \emptyset, \dots$$

for some points $r_i < t_{i-1} < r_{i+1} < t_i$ in I ($i = 2, \dots, m-1$) and some $m \leq n$.

Given \mathcal{U} , one chooses s_i in $U_i \cap U_{i+1} \cap I$ (which amounts to a real number between r_{i+1} and t_i), and a geometric point \overline{s}_i above s_i ($i = 1, \dots, m-1$); one completes this collection by setting $\overline{s}_0 = \vec{t}$, $\overline{s}_m = \vec{t}$.

Let us look at the diagram

$$\text{Cov}_{S,\mathcal{U}}^{\text{temp}} \xrightarrow{\sigma_i} \text{Ind Cov}_{\mathcal{U}_{i+1}}^{\text{alg}} \begin{array}{c} \xrightarrow{F_{\overline{s}_i, U_{i+1}}^{\text{alg}}} \\ \xrightarrow{F_{\overline{s}_{i+1}, U_{i+1}}^{\text{alg}}} \end{array} \longrightarrow \text{Sets}.$$

One has $F_{\overline{s}_j, U_{i+1}}^{\text{alg}} \circ \sigma_i = F_{\overline{s}_i, \mathcal{U}}^{\text{temp}}$, $j = i, i+1$, whence continuous maps

$$\text{Isom}(F_{\overline{s}_i, U_{i+1}}^{\text{alg}}, F_{\overline{s}_{i+1}, U_{i+1}}^{\text{alg}}) \rightarrow \text{Isom}(F_{\overline{s}_i, \mathcal{U}}^{\text{temp}}, F_{\overline{s}_{i+1}, \mathcal{U}}^{\text{temp}}).$$

But $\text{Isom}(F_{\overline{s}_i, U_{i+1}}^{\text{alg}}, F_{\overline{s}_{i+1}, U_{i+1}}^{\text{alg}})$ is a non-empty compact set for any $i = 0, \dots, m$: for $i = 0$ ($s_0 = \vec{t}$), one uses the analogue of 2.2.3 for the category of finite étale coverings, which follows from Deligne's discussion, *loc. cit* (or even directly from [SGA 1, V], interpreting $F_{\vec{t}, U_{i+1}}^{\text{alg}}$ as a fiber functor at a geometric generic point). The rest of the proof goes exactly as in 1.3.3. \square

2.2.4. This proposition allows to construct fundamental groups $\pi_1^\bullet(S, \vec{t})$ based at a tangential base-point \vec{t} for any full subcategory $\text{Cov}_S^\bullet \rightarrow \text{Cov}_S^{\text{temp}}$ stable under taking connected components, fiber products and quotients. The theory of 1.4 extends without change to this variant, as far as one restricts to curves over an algebraically closed field of characteristic zero.

For $\text{Cov}_S^{\text{top}} \rightarrow \text{Cov}_S^{\text{temp}}$, the interpretation of tangential base-points is not as transparent as in the complex-analytic context.

2.3. Description of some temperate fundamental groups.

In this subsection, we assume that K is *algebraically closed of characteristic 0* (typically $K = \mathbb{C}_p$).

2.3.1. It follows from 2.1.6 (together with 1.4.1) that \mathbb{P}^1 and \mathbb{A}^1 have trivial temperate fundamental groups⁽⁷⁾.

Let us next examine $\mathbb{G}_m = \mathbb{P}^1 \setminus \{0, \infty\}$. We have $\pi_1^{\text{top}}(\mathbb{G}_m, 1) = 0$; on the other hand, any connected finite étale covering is Kummer $\mathbb{G}_m \rightarrow \mathbb{G}_m$, $x \mapsto x^n$. It follows that

$$\pi_1^{\text{temp}}(\mathbb{G}_m, 1) = \pi_1^{\text{alg}}(\mathbb{G}_m, 1) = \widehat{\mathbb{Z}}(1) := \varprojlim_n \mu_n \cong \widehat{\mathbb{Z}} = \prod_{\ell} \mathbb{Z}_{\ell}.$$

Similarly, $\pi_1^{\text{temp}}(\mathbb{G}_m^n, 1) \cong \widehat{\mathbb{Z}}^n$.

2.3.2. Elliptic curves. Let us first consider the case of an *elliptic curve with good reduction* S (so that $\pi_1^{\text{top}}(S, 1) = \{1\}$). Then any finite étale covering is given by an isogeny $S' \rightarrow S$ and S' has good reduction. Hence any temperate covering is an isogeny, and

$$\pi_1^{\text{temp}}(S, 1) = \pi_1^{\text{alg}}(S, 1) \cong \widehat{\mathbb{Z}} \times \widehat{\mathbb{Z}}.$$

Let now $S = \mathbb{G}_m/q^{\mathbb{Z}}$ be instead a *Tate elliptic curve*. Then the homomorphisms

$$\pi_1^{\text{temp}}(\mathbb{G}_m, 1) \rightarrow \pi_1^{\text{temp}}(S, 1) \rightarrow q^{\mathbb{Z}} = \pi_1^{\text{top}}(S, 1)$$

actually give rise to a canonical exact sequence (1.4.12. (b) may be applied here)

$$1 \rightarrow \widehat{\mathbb{Z}}(1) \rightarrow \pi_1^{\text{temp}}(S, 1) \rightarrow \mathbb{Z} \rightarrow 1,$$

and since $\pi_1^{\text{temp}}(S, 1) \subset \pi_1^{\text{alg}}(S, 1) \cong \widehat{\mathbb{Z}} \times \widehat{\mathbb{Z}}$ is abelian, to a non-canonical isomorphism

$$\pi_1^{\text{temp}}(S, 1) \cong \widehat{\mathbb{Z}} \times \mathbb{Z}.$$

This is the simplest manifestation of the lacunary character of temperate fundamental groups.

(The case of a Tate curve deprived from its origin is more difficult).

2.3.3. Mumford curves. Let now S be a Mumford curve of genus $g > 1$: $S = \Omega/\Gamma$, $\Omega \cong \tilde{S}$, and $\pi_1^{\text{top}}(S, s) \cong \Gamma$, a Schottky group, free on g generators.

⁽⁷⁾in contrast, recall that \mathbb{A}^1 has non-trivial locally-algebraic fundamental group, and that \mathbb{P}^1 has trivial locally-algebraic fundamental group but non-trivial étale fundamental group.

We have homomorphisms

$$\pi_1^{\text{temp}}(\Omega, \bar{s}) \rightarrow \pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{top}}(S, s)$$

which give rise to an exact sequence

$$1 \rightarrow H \rightarrow \pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{top}}(S, s) \rightarrow 1$$

with $H := \text{Im}(\pi_1^{\text{temp}}(\Omega, \bar{s}) \rightarrow \pi_1^{\text{temp}}(S, \bar{s}))^- \subset \pi_1^{\text{temp}}(S, \bar{s})$. As an abstract group, $\pi_1^{\text{temp}}(S, \bar{s})$ is thus the semidirect product $H.\Gamma$.

Passing to the profinite completion, we get an exact sequence

$$\widehat{H} \rightarrow \pi_1^{\text{alg}}(S, \bar{s}) \rightarrow \widehat{\Gamma} \rightarrow 1,$$

and the homomorphism $\pi_1^{\text{alg}}(\Omega, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s})$ factors through \widehat{H} .

This homomorphism is very far from being injective. The point is that the étale covering of S obtained from an arbitrary finite Galois étale covering $\Omega' \rightarrow \Omega$ is not Galois (or, what amounts to the same by 2.1.9, not temperate). The finite Galois étale coverings $\Omega' \rightarrow \Omega$ which give rise to a Galois covering of S are called “equivariant” in [vdP83, 1.3]; by 2.1.9, they correspond to a topological Galois covering of some Galois étale covering $S' \rightarrow S$.

Moreover, there are examples [vdP83, 2.7] (one may choose for Ω the Drinfeld half-plane) where the degrees of Ω'/Ω and S'/S coincide, but Ω' is not the universal topological covering of S' ; it follows that Ω' is not simply-connected (in fact, its topological fundamental group is free of infinite type), and that the injection $\pi_1^{\text{temp}}(\Omega, \bar{s}) \rightarrow \pi_1^{\text{alg}}(\Omega, \bar{s})$ is not a bijection in general.

Let us return to the non-injectivity of $\pi_1^{\text{alg}}(\Omega, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s})$. Actually, the following stronger result holds: $\pi_1^{\text{alg}}(S, \bar{s})$ is metrizable, while $\pi_1^{\text{alg}}(\Omega, \bar{s})$ is not.

Recall that a profinite group is metrizable if and only if the set of its open subgroups is countable, cf. [Ser94, 1.3]. This is the case for $\pi_1^{\text{alg}}(S, \bar{s})$, which is even topologically finitely generated, but not for $\pi_1^{\text{alg}}(\Omega, \bar{s})$. In fact, one can show that for any $n \geq 1$, the set of open subgroups $H \subset \pi_1^{\text{alg}}(\Omega, \bar{s})$ endowed with an isomorphism $\pi_1^{\text{alg}}(\Omega, \bar{s})/H \rightarrow \mathbb{Z}/n\mathbb{Z}$ is already uncountable. Indeed, this set may be identified with $\mathcal{O}(\Omega)^\times/(\mathcal{O}(\Omega)^\times)^n$, or else with the set of $\mathbb{Z}/n\mathbb{Z}$ -valued currents⁽⁸⁾ on the tree associated to Ω (cf. [FvdP81, I.8.9, V.2.3]). Since $g \geq 2$, this tree has infinitely many vertices connected to more than two edges, and it follows that the $\mathbb{Z}/n\mathbb{Z}$ -valued currents do not form a countable set.

A similar picture holds for Mumford curves deprived from a finite set of points.

2.3.4. Affinoid curves with good reduction. In this case, one has

$$\pi_1^{\text{temp}}(S, \bar{s}) = \pi_1^{\text{alg}}(S, \bar{s}).$$

⁽⁸⁾ *i.e.* $\mathbb{Z}/n\mathbb{Z}$ -valued functions on the edges of the tree, such that the sum of its values at the edges starting from an arbitrary vertex is zero.

We may assume that s is the maximal point of S corresponding to the sup-norm. The assertion then follows from De Jong's observation [dJ95b, 7.5] that the map $\pi_1^{\text{ét}}(s, \bar{s}) = \pi_1^{\text{alg}}(s, \bar{s}) \rightarrow \pi_1^{\text{alg}}(S, \bar{s})$ is already surjective. We shall only need the case of the closed unit disk. The proof goes as follows (assuming that $S \subset \mathbb{P}^1$ as in *loc. cit.*).

One has to show that any finite étale covering $f : S' \rightarrow S$ of degree n such that $S'_s = \{t_1, \dots, t_n\}$ splits. The assumption implies that f is a local isomorphism around each t_i (*cf.* 1.2.1). Hence one can find an affinoid neighborhood V of s such that $S'_V = W_1 \amalg \dots \amalg W_n$, and $W_i \cong V$. Because V contains the maximal point s which is the boundary of S in \mathbb{P}^1 (in the sense of [Ber90, 3.1]), we get an admissible covering (for the Grothendieck topology) of $\mathbb{P}^1 = S \cup (\mathbb{P}^1 \setminus (S \setminus V))$ by closed analytic subdomains. One then glues S' and n copies of $\mathbb{P}^1 \setminus (S \setminus V)$ together via the isomorphisms $W_i \rightarrow V$ in order to get a finite étale covering of \mathbb{P}^1 which restricts to S' over S . Such a covering is trivial.

Concerning the algebraic fundamental group of the closed unit disk over \mathbb{C}_p , one has the following result, essentially due to M. Raynaud (not used in the sequel):

Theorem 2.3.5. *A finite group is a quotient of the profinite group $\pi_1^{\text{alg}}(\mathbb{D}_{\mathbb{C}_p}(0, 1^+), 0)$ if and only if it is generated by its p -Sylow subgroups.*

The necessity of the condition means that any finite Galois étale covering of $\mathbb{D}_{\mathbb{C}_p}(0, 1^+)$ of order prime to p is trivial, which is proven in [Ber93, 6.3.3] and [Lüt93, 2.11]. The converse is much harder and follows from Raynaud's solution of the Abhyankar conjecture on finite étale coverings of $\mathbb{A}_{\mathbb{F}_p}^1$, *cf.* [Ray94]. Of course, we may replace 0 by any geometric point or tangential base-point (*cf.* 2.2.3).

The standard example of a non-trivial finite Galois étale covering of $\mathbb{D}(0, 1^+)$ is the Artin-Schreier covering $\mathbb{D}(0, 1^+) \xrightarrow{z \mapsto z^p - z} \mathbb{D}(0, 1^+)$ (with group \mathbb{Z}/p); it splits on any smaller concentric disk. A less standard example, with non-solvable Galois group A_5 , is given by $\mathbb{D}(1, 1^+) \setminus \mathbb{D}(0, 1^-) \xrightarrow{z \mapsto z^3 - z^{-2}} \mathbb{D}(0, 1^+)$ for $p = 3$.

2.3.6. Punctured disks. Let us fix $r \in |K^\times|$, and set $\mathbb{D}_r = \mathbb{D}_K(0, r^+)$, $\mathbb{D}_r^* = \mathbb{D}_r \setminus \{0\}$. We let \bar{s} be any geometric point or tangential base-point of \mathbb{D}^* .

Proposition 2.3.7. *There are canonical isomorphisms*

$$\pi_1^{\text{temp}}(\mathbb{D}_r^*, \bar{s}) = \pi_1^{\text{alg}}(\mathbb{D}_r^*, \bar{s}) = \pi_1^{\text{alg}}(\mathbb{D}_r, \bar{s}) \times \widehat{\mathbb{Z}}(1).$$

PROOF. By homothety, we may assume that $r = 1$ (and drop the index r). These canonical isomorphisms are induced by the obvious fully faithful functors (*cf.* 1.4.8):

$$\text{Cov}_{\mathbb{D}^*}^{\text{kum}} \hookrightarrow \text{Cov}_{\mathbb{D}^*}^{\text{alg}}, \quad \text{Cov}_{\mathbb{D}}^{\text{alg}} \hookrightarrow \text{Cov}_{\mathbb{D}^*}^{\text{alg}} \hookrightarrow \text{Cov}_{\mathbb{D}^*}^{\text{temp}}$$

where $\text{Cov}_{\mathbb{D}^*}^{\text{kum}}$ denotes the category of finite disjoint unions of Kummer coverings of \mathbb{D}^* ; it is stable under taking connected components, fiber products (the fiber product of a Kummer covering of degree m and a Kummer covering of degree n is a finite disjoint union of Kummer coverings of degree $\text{lcm}(m, n)$) and quotients. The fundamental group $\pi_1^{\text{kum}}(\mathbb{D}^*, \bar{s})$ is clearly $\mathbb{Z}(1)$.

Assume for a while that s is the maximal point of \mathbb{D} (the statement does not depend on the choice of s). Then as we have seen in 2.3.4, the homomorphism $\pi_1^{\text{temp}}(s, \bar{s}) \rightarrow \pi_1^{\text{alg}}(\mathbb{D}, \bar{s})$ is surjective. Since it factors through $\pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s})$, it follows that $\pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s}) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}, \bar{s}) = \pi_1^{\text{alg}}(\mathbb{D}, \bar{s})$ is surjective.

Let us now take for s a tangential base-point at 0. Let us consider the maps $\mathbb{D}_r \rightarrow \mathbb{D}$ (for $r \in |K^\times|$, $r < 1$) and the corresponding homomorphism (cf. 1.4.10):

$$\pi_1^{\text{temp}}(\mathbb{D}_r^*, \bar{s}) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s}).$$

It follows from 2.1.12 that

$$\varprojlim_{r \rightarrow 0} \text{Im}(\pi_1^{\text{temp}}(\mathbb{D}_r^*, \bar{s}) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s})) = \mathbb{Z}(1),$$

the composition

$$\mathbb{Z}(1) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s}) \rightarrow \mathbb{Z}(1)$$

being identity. It remains to show that the sequence

$$1 \rightarrow \mathbb{Z}(1) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s}) \rightarrow \pi_1^{\text{temp}}(\mathbb{D}, \bar{s}) = \pi_1^{\text{alg}}(\mathbb{D}, \bar{s}) \rightarrow 1$$

is exact. The problem is only in the middle; since $\pi_1^{\text{temp}}(\mathbb{D}^*, \bar{s})$ is a pro-discrete group, it amounts to the following fact, which is a very special case of 2.1.12: a temperate Galois covering of \mathbb{D}^* extends to \mathbb{D} if and only if its restriction to a sufficiently small punctured disk centered at 0 splits. \square

On the other hand, the consideration of the logarithmic étale covering restricted to a punctured disk centered at infinity shows that $\pi_1^{\text{loc.alg}}(\mathbb{D}_r^*, \bar{s}) \neq \pi_1^{\text{alg}}(\mathbb{D}_r^*, \bar{s})$, and *a fortiori*,

$$\pi_1^{\text{et}}(\mathbb{D}_r^*, \bar{s}) \neq \pi_1^{\text{temp}}(\mathbb{D}_r^*, \bar{s}).$$

2.3.8. \mathbb{A}^1 minus a few points. Let $S = \mathbb{A}^1 \setminus \{\zeta_1, \dots, \zeta_m\}$. For any $i = 1, \dots, m$, let \vec{t}_i be a tangential base-point at ζ_i , and let α_i be a temperate path from \vec{t}_i to a fixed geometric point \bar{s} (or tangential base-point) of S . Then we have a canonical homomorphism corresponding to Kummer coverings induced by $z - \zeta_i \mapsto (z - \zeta_i)^n$ (cf. 2.3.6)

$$\widehat{\mathbb{Z}}(1) \hookrightarrow \pi_1^{\text{temp}}(S, \vec{t}_i) \rightarrow \widehat{\mathbb{Z}}(1)$$

whose composition is identity. By composition with $\text{ad}(\alpha_i)$, we get a monomorphism

$$\widehat{\mathbb{Z}}(1) \hookrightarrow \pi_1^{\text{temp}}(S, \bar{s})$$

which depends of course on α_i . Let us denote by γ_i the image in $\pi_1^{\text{temp}}(S, \bar{s})$ of a topological generator of $\widehat{\mathbb{Z}}(1) \cong \widehat{\mathbb{Z}}$. The ‘local monodromies’ γ_i are not at all canonical, but the *smallest closed normal subgroup* $\langle \gamma_i \rangle^-$ of $\pi_1^{\text{temp}}(S, \bar{s})$ which contains them is independent of all choices. In fact we have⁽⁹⁾:

Proposition 2.3.9. $\langle \gamma_i \rangle^- = \pi_1^{\text{temp}}(\mathbb{A}^1 \setminus \{\zeta_1, \dots, \zeta_m\}, \bar{s})$.

PROOF. This amounts to saying that a temperate (Galois, if one wishes) covering splits if its restriction to sufficiently small punctured disks around any of the ζ_i splits. But the latter condition implies that the covering extends to a temperate covering of \mathbb{A}^1 (cf. 2.1.11), which automatically splits (cf. 2.3.1). \square

By analogy with the complex situation, 2.3.9 suggests two questions:

Question 2.3.10. *Is it possible to choose the γ_i ’s in such a way that they are topological generators of $\pi_1^{\text{temp}}(S, \bar{s})$?*

Following the above construction, one can also introduce a local monodromy γ_∞ at ∞ .

Question 2.3.11. *Is it possible to choose the γ_i ’s and γ_∞ in such a way that their product (in suitable order) is identity?*

We do not know the answer to 2.3.10, but shall answer 2.3.11 in 6, when we have more tools at disposal.

For $m \geq 2$, $\pi_1^{\text{temp}}(S, \bar{s})$ is indeed a rather complicated topological group, far from being discrete and from being compact:

Proposition 2.3.12. *For $m \geq 2$, $\pi_1^{\text{temp}}(\mathbb{A}^1 \setminus \{\zeta_1, \dots, \zeta_m\}, \bar{s})$ is not locally compact.*

PROOF. Assume that $\pi_1^{\text{temp}}(S, \bar{s})$ is locally compact. Then because $\pi_1^{\text{temp}}(S, \bar{s})$ is a prodiscrete group, there would exist a *compact* (necessarily totally discontinuous, hence profinite) open normal subgroup H . Let $S' \rightarrow S$ be the corresponding Galois temperate covering with group $G = \pi_1^{\text{temp}}(S, \bar{s})/H$. Up to replacing H by a small subgroup, we may and shall assume that $S' = \tilde{T}$ is the topological universal covering of a finite Galois étale covering T of S . Via 2.1.8, we then have an exact sequence

$$1 \rightarrow H \rightarrow \pi_1^{\text{temp}}(T, \bar{t}) \rightarrow \pi_1^{\text{top}}(T, \bar{t}) \rightarrow 1.$$

Let T' be any finite étale covering of T . Then the universal topological covering \tilde{T}' is a temperate covering of T which dominates $S' = \tilde{T}$. Due to the compactness of H , the map $\tilde{T}' \rightarrow \tilde{T}$ must be finite.

⁽⁹⁾this is the correct version of cor. 5.3 in [And98], which was wrongly stated.

We now choose T' in order to contradict this property. Let \overline{X} be a projective smooth analytic curve over K such that $\text{rk } \pi_1^{\text{top}}(\overline{X}) > \text{rk } \pi_1^{\text{top}}(T)$. Assume that \overline{X} is endowed with a finite morphism to \mathbb{P}^1 which is unramified above S (since $m \geq 2$, this just amounts to saying that \overline{X} is defined over $\overline{\mathbb{Q}}$, according to a well-known theorem of Belyi). We set $X = \overline{X} \times_{\mathbb{P}^1} S \subset \overline{X}$, and choose for T' a connected component of $X \times_S T$. Note that since T is Galois over S , T' is Galois over X . Let Y be a connected component of $X \times_S \tilde{T}$ lying above X . Then \tilde{T}' is a topological covering of Y , and is finite over Y (if and) only if it coincides with Y . We then have a commutative square of Galois temperate coverings

$$\begin{array}{ccc}
 & Y = \tilde{T}' & \\
 \swarrow & & \searrow \\
 \tilde{X} & & T' \\
 \searrow & & \swarrow \\
 & X &
 \end{array}$$

Now $\text{Aut}_X Y$ admits the free group $\text{Aut}_{T'} Y$ as a normal subgroup of finite index, and admits the free group $\text{Aut}_X \tilde{X}$ as a quotient. Since $\text{rank } \text{Aut}_{T'} Y \leq \text{rank } \pi_1^{\text{top}}(T) < \text{rank } \pi_1^{\text{top}}(\overline{X}) = \text{rank } \pi_1^{\text{top}}(X)$, this is impossible. Therefore $\pi_1^{\text{temp}}(S)$ is not locally compact. \square

Remark. In [And], temperate fundamental groups of algebraic p -adic manifolds will be used to give a geometric description of the local Galois group $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$.

3. Local and global monodromy of p -adic differential equations.

ABSTRACT: We briefly review and compare the theory of singularities of ordinary linear differential equations in the formal, complex and p -adic contexts. We then outline the Christol-Mebkhout theory of p -adic slopes of differential modules over annuli. There is a close relation with Galois representations of local fields of characteristic p , expressed by Crew's p -adic local monodromy conjecture, which has been recently proved [And02b]. Complications occurring in the p -adic context reflect the fact that the étale fundamental groups of annuli are non-abelian.

We then define and study the non-archimedean Riemann-Hilbert functor, which attaches a vector bundle with integrable connection to any continuous representation with discrete coimage of the étale fundamental group of a non-archimedean manifold. Connections in the image are characterized by the fact that the étale sheaf of germs of solutions is locally constant.

3.1. Introduction.

3.1.1. Let us briefly review some well-known facts about singularities of ordinary linear differential equations. For concreteness, we consider a differential equation

$$(*) \quad \alpha_\mu(z)\partial^\mu y + \cdots + \alpha_0(z)y = 0, \quad \partial = z\frac{d}{dz},$$

with polynomial coefficients $\alpha_i(z)$, and concentrate on phenomena at infinity.

3.1.2. We begin the discussion with the formal setting. One defines the formal Newton polygon at ∞ , \widehat{NP}_∞ to be the convex hull of $\{(X \leq i, Y \geq \deg_z \alpha_0 - \deg_z \alpha_i)\}$ in the real plane. Its 'height'⁽¹⁰⁾ is called the (formal) irregularity of $(*)$ at ∞ . The equation is called regular at ∞ if the irregularity is 0, *i.e.* if the only finite slope is 0.

It is known that $(*)$ has a basis of 'formal' solutions of the form

$$\hat{y} = \sum \hat{u}_i \left(\frac{1}{z}\right) z^{-\hat{e}_i} (\log^{k_i} z) e^{P_i(z^{1/\mu^!})},$$

where the P_i 's are polynomials, and the \hat{u}_i are just formal power series in $\frac{1}{z}$. The degree of the 'ramified polynomials' $P_i(z^{1/\mu^!})$ actually corresponds to the slopes of \widehat{NP}_∞ . The \hat{e}_i 's are called the formal or Turrittin exponents; they can be computed algebraically.

Example. Let us illustrate these notions with the *Bessel equation*

$$(B)_\nu : \quad (\partial^2 + (z^2 - \nu^2))y = 0$$

⁽¹⁰⁾the height is the difference between the Y -coordinate of the highest vertex and the Y -coordinate of the lowest vertex.

A basis of ‘formal solutions’ at ∞ is given by

$$\hat{y}_{\pm} = z^{-1/2} e^{\pm\sqrt{-1}z} \cdot {}_2F_0\left(\frac{\nu+1}{2}, \frac{-\nu+1}{2}; \frac{\mp\sqrt{-1}}{2z}\right), \quad \hat{e}_i = 1/2,$$

and \widehat{NP}_{∞} is




FIGURE 1

3.1.3. Complex-analytic setting. Solutions at any non-singular point s (i.e. a point in \mathbb{C}^{\times} which is not a root of α_{μ}) converge up to the next singularity.

Let $\mathcal{C} = \mathcal{C}(]r', r[) = \{z, r' < |z| < r\}$ be an annulus which contains no singularity. Using the fact the $\pi_1(\mathcal{C}, s) \cong \mathbb{Z}$, it is classical to deduce that (*) has a basis of ‘analytic’ solutions of the form

$$y = \sum u_i \left(\frac{1}{z}\right) z^{-e_i} (\log^{k_i} z),$$

where the u_i are Laurent series in $\frac{1}{z}$ which converge in \mathcal{C} . The e_i ’s are called the analytic exponents.⁽¹¹⁾

The relation between the y ’s (analytic) and the \hat{y} ’s (formal) is not straightforward. If $r = \infty$, the \hat{y} ’s occur as asymptotic expansions of the y ’s on suitable sectors, and the theory of multisummability/acceleration provides a kind of canonical inverse process (again on suitable sectors). If ∞ is a regular singularity, the \hat{u}_i converge so that with $u_i = \hat{u}_i, e_i = \hat{e}_i$, the solutions y and \hat{y} coincide.

Example. For the Bessel equation (say with non-integral parameter ν to simplify), one has a basis of solutions on $\mathcal{C} = \mathbb{C}^{\times}$ given by

$$y_{\pm} = z^{\pm\nu} \sum_0^{\infty} \frac{(-1)^n z^{2n}}{4^n n! (\pm\nu + 1)_n}.$$

Moreover $e_i = \pm\nu$ (analytic exponents at $\infty = -$ analytic exponents at 0); here we are ‘lucky’, there are just two singularities, one of them is regular.

⁽¹¹⁾in terms of differential systems, this amounts to the fact that a system $\partial Y = AY$ on \mathcal{C} has a fundamental solution matrix Y of the form $U \cdot z^{-E}$, where U is analytic in \mathcal{C} and E is constant (Fuchs normal form).

In general, the computation of analytic exponents involves infinite determinants. Even in the so-called fuchsian case where there are only regular singularities, the computation of analytic exponents on an annulus which encloses several regular singularities is transcendental; e.g. the exponents of the differential equation of order two with solution ${}_2F_1(\frac{1}{2}, \frac{1}{2}, 1; z^2)$, on an annulus enclosing 0 and 1 but not -1 and ∞ , are transcendental numbers: $\frac{\log(3 \pm 2\sqrt{2})}{2\pi i}$ (12).

3.1.4. p -adic-analytic setting. The first problem one encounters when dealing with p -adic differential equations (cf. I.2.4) is that the solutions at non-singular points have small radii of convergence in general, and do not converge up to the next singularity.

Example. $\frac{d}{dz}y = y$, $e^z = \sum \frac{z^n}{n!}$ converges only in $D(0, |\pi|^-)$, $\pi^{p-1} = -p$.

This problem leads one to normalize somehow the situation by postulating

Dwork's condition: the radius of convergence of solutions of () at a generic point*⁽¹³⁾ t_1 of modulus 1 is ≥ 1 .

Example. One 'normalizes' Bessel's equation by the change of variable $z \mapsto 2\pi\sqrt{-1}z$. The normalized Bessel differential operator is then

$$B_{\nu, \pi} = \partial^2 - (4\pi^2 z^2 + \nu^2)$$

which is solvable in the p -adic generic disk whenever $\nu \in \mathbb{Z}_p$.

3.1.5. Let us now discuss problems at p -adic singularities, concentrating again on phenomena at infinity.

- The first one is the problem of 'small divisors': if some formal exponent $\hat{e}_i \in \mathbb{Z}_p$ (or rather some difference $\hat{e}_i - \hat{e}_j$) is a Liouville number, *i.e.* very closely approximated by infinitely many rational integers, the formal power series \hat{u}_i occurring in the formal solutions \hat{y} may diverge, even in the regular case. Otherwise, the \hat{u}_i converge, even in the irregular case [Bal82].

Example. if $\nu \in \mathbb{Z}_p$ is well-approximable by negative integers, $\sum_0^\infty \frac{(-1)^n z^{2n}}{4^n n! (\nu+1)_n}$ may diverge, although 0 is a regular singularity of the Bessel equation.

- Even if Liouville numbers do not show up (*e.g.* if everything is defined over $\overline{\mathbb{Q}}$) so that the \hat{u}_i 's converge, these series may not converge as far as one might expect (*e.g.* in a unit disk under the Dwork condition). This subtle phenomenon occurs in the normalized Bessel case $B_{\nu, \pi}$ for $p = 2$ (${}_2F_0(\frac{\nu+1}{2}, \frac{-\nu+1}{2}; \frac{\pm 1}{4\pi z})$ do not converge up to the unit circle): there is an

⁽¹²⁾ An independent solution is given by $i \cdot {}_2F_1(\frac{1}{2}, \frac{1}{2}, 1; 1 - z^2)$, and in this basis, the local monodromy matrix around 1 (*resp.* 0) is $\begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}$ (*resp.* $\begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix}$); the eigenvalues of the monodromy around \mathcal{C} are the eigenvalues of $\begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 4 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -7 & 4 \\ -2 & 1 \end{pmatrix}$, *i.e.* $3 \pm 2\sqrt{2}$.

⁽¹³⁾ a generic point of modulus ρ in the sense of Dwork is a point t_ρ in a complete extension Ω of K with $|t_\rho| = \rho$ and such that the disk $D_\Omega(t_\rho, \rho^-)$ does not contain any element of K .

annulus $\mathcal{C}(]1, r[)$ such that neither y_{\pm} nor \hat{y}_{\pm} provide any information about the solutions of $B_{\nu, \pi}$ in $\mathcal{C}(]1, r[)$.

In particular, the question arises whether there is a basis of solutions

$$\sum u_i \left(\frac{1}{z}\right) z^{-e_i} (\log^{k_i} z),$$

where the u_i are Laurent series in $\frac{1}{z}$ which converge in such an annulus $\mathcal{C}(]1, r[)$. More generally, concerning the existence of a ‘ p -adic Fuchs normal form’:

- Does there exist a basis of solutions of (*) of the form

$$y = \sum u_i \left(\frac{1}{z}\right) z^{-e_i} (\log^{k_i} z),$$

where the u_i are Laurent series which converge in a given annulus $\mathcal{C}(]r', r[)$?

A necessary condition has been put forward by P. Robba: *for any $\rho \in]r', r[$, the radius of convergence of the solutions at the generic point t_{ρ} of modulus ρ is $\geq \rho$* (in the variable z , not $1/z$).

The problems are: to which extent is the converse true? What can be said if Robba’s condition is not fulfilled?

These questions have recently become almost complete answers thanks to the Christol-Mebkhout theory of p -adic slopes of differential modules over annuli.

3.2. Local monodromy of p -adic differential equations. Outline of Christol-Mebkhout theory.

3.2.1. Let K be a complete subfield of \mathbb{C}_p . We assume for simplicity that the restriction of the p -adic valuation to K is discrete.

The Robba K -algebra at ‘infinity’ is

$$\mathcal{R} = \mathcal{R}_{z, K} = \{K\text{-analytic functions on some } \mathcal{C}(]1, r[)\}.$$

Endowed with $\partial = z d/dz$, this is a differential (integral) K -algebra. Although it is not simple as a differential algebra (there are many non-trivial differential ideals), it has the following useful property, which often allows one to replace \mathcal{R} by its fraction field $Q(\mathcal{R})$ (cf. [And02a]):

any $\mathcal{R}[\partial]$ -module M which is of finite presentation over \mathcal{R} is free over \mathcal{R} .

In the sequel, we simply refer to such an M as a *differential module* over \mathcal{R} .

3.2.2. For any $\rho > 1$ sufficiently close to 1, let us denote by $R(M, \rho)$ the infimum of ρ and the maximal radius R such that M is solvable (= has a basis of analytic solutions) in $D(t_{\rho}, R^-)$.

One says that M is *solvable* (at the inner boundary), if $\lim_{\rho \rightarrow 1^+} R(M, \rho) = 1$ (this is essentially Dwork’s condition, except that $R(M, 1)$ itself is not defined.)

3.2.3. A useful criterion (which goes back the Dwork's early studies) for M being solvable is the existence of a Frobenius structure.

Let σ be an automorphism of K inducing $t \mapsto t^{p^f}$ in characteristic p , and let φ_f be the σ -linear endomorphism of \mathcal{R} defined by $z \mapsto z^{p^f}$.

One says that M has a Frobenius structure if $\varphi_f^* M \cong M$. In what follows, the particular choice of such an isomorphism is irrelevant.

Example. Let $M_{\nu,\pi} = \mathcal{R}/\mathcal{R}B_{\nu,\pi}$ be the differential module over \mathcal{R} attached to the normalized Bessel operator. It turns out that, up to isomorphism, $M_{\nu,\pi}$ does not depend on $\nu \in \mathbb{Z}_p$ [And02a]. Choosing $\nu = 0$ for instance, the existence of a Frobenius structure is known by the interpretation of $M_{\nu,\pi}$ in terms of Kloosterman sums [Dw74] (in fact, for any $\nu \in \mathbb{Q} \cap \mathbb{Z}_p$, there is a relation to twisted Kloosterman sums, and a Frobenius structure has a geometric interpretation, cf. [Be84]).

3.2.4. We now turn to p -adic slopes⁽¹⁴⁾, as defined by G. Christol and Z. Mebkhout.

- A solvable M is of *greatest slope* $\lambda \in [0, \infty[$ if $R(M, \rho) = \rho^{1-\lambda}$ for ρ close to 1.
- M is *purely of slope* λ if any non-zero solution at t_ρ has radius of convergence $\rho^{1-\lambda}$ for ρ close to 1.

Theorem 3.2.5. *Any solvable M has a decreasing filtration $M_{>\lambda}$ such that $Gr_\lambda M$ is free and purely of slope λ .*

cf. [CM00]. Following Christol-Mebkhout, one defines the *p -adic Newton polygon* (at infinity) NP_∞ : the horizontal length of the segment of slope λ is $rk Gr_\lambda M$. In particular, the greatest slope of M in the above sense is the greatest slope of NP_∞ in the usual sense. The height of NP_∞ is called the *p -adic irregularity* of M .

A solvable module M is called *tame* (or *p -adically regular*) if the only finite slope of NP_∞ is zero. This amounts to say that M satisfies the Robba condition of 3.1.5: $R(M, \rho) = \rho$ when $\rho \rightarrow 1^+$.

One of the main results in the Christol-Mebkhout theory is ([CM00]):

Theorem 3.2.6. *NP_∞ has integral vertices.*

A useful complement in the computation of NP_∞ is given by ([And02a]):

Lemma 3.2.7. *Assume M comes from a differential module over the subring of $(K \cap \overline{\mathbb{Q}})((\frac{1}{x}))$ consisting of convergent series in $\mathcal{C}([1, \infty[)$. Then the p -adic greatest slope λ is \leq than the greatest formal slope λ_∞ .*

Example. It can be shown (using 3.2.8 below) that the Bessel module $M_{\nu,\pi}$ is not tame; 3.2.6 and 3.2.7 then leave only two possibilities for NP_∞ , and it turns out that both occur (cf. [And02a]).

⁽¹⁴⁾ which should not be confused with the so-called slopes of Frobenius which measure the p -adic size of a fixed Frobenius structure (when it exists); a unit-root F -crystal corresponding to the case of Frobenius slope zero.



FIGURE 2

As for the converse of Robba's criterion (*cf.* 3.1.5), one has (*cf.* [CM97]):

Theorem 3.2.8 (p -adic local monodromy theorem, tame case). *Assume M has a Frobenius structure⁽¹⁵⁾, and is purely of slope zero. Then M is an iterated extension of rank one differential modules of type $\partial - \alpha$, $\alpha \in \mathbb{Q} \cap \mathbb{Z}_p$ (called the p -adic exponents of M).*

With respect to the conclusion about the rationality of the p -adic exponents, this result may be seen as a p -adic analogue of Grothendieck's ℓ -adic local monodromy theorem for tame ℓ -adic coefficients.

The result, if not its proof, is closely related to the fact that a tame (= of degree prime to p) Galois étale finite covering of any annulus \mathcal{C} is Kummer (the tame algebraic fundamental group of \mathcal{C} is $\widehat{\mathbb{Z}}'(1)$, where the dash means as usual that the p -component is omitted, *cf.* [Ber93]). The relationship will appear more clearly below.

3.2.9. Let us return to our original differential equation (*). Let us assume that the only singularities at finite distance are in $\mathbb{D}(0, 1^+)$. Let us also assume that (*) satisfies Dwork's condition (so that the associated differential module over \mathcal{R} is solvable), and that the formal exponents at ∞ are rational. Then the p -adic behavior of (*) in a small punctured disk $\mathbb{D}(\infty, \epsilon)^*$ centered at ∞ is dictated by the formal behavior (*cf.* 3.1.5), while the p -adic behavior in a thin annulus $\mathcal{C}(]1, r[)$ is predicted by Christol-Mebkhout theory. However, in the intermediate region $\mathcal{C}([r, 1/\epsilon])$, nothing seems to be known.

We shall return to this problem in 3.5.

3.3. Local monodromy of p -adic differential equations and Galois representations.

3.3.1. The differential sub- K -algebra

$$\mathcal{E}^\dagger = \{\text{bounded } K\text{-analytic functions on some } \mathcal{C}(]1, r[)\}$$

⁽¹⁵⁾there is a more general version without Frobenius structure, in which one encounters some trouble with the problem of small divisors. The existence of a Frobenius structure simplifies the statement, if not the proof, since it implies a priori that if p -adic exponents exist, they are rational, hence non-Liouville.

of bounded functions in \mathcal{R} is actually a field. With the p -adic sup-norm, it is even a henselian field, with residue field $k((\frac{1}{z}))$ (where k stands for the residue field of K). Hence any Galois extension $k((\frac{1}{z'}))/k((\frac{1}{z}))$ gives rise to an unramified Galois extension $\mathcal{E}_{z',K'}^\dagger/\mathcal{E}_{z,K}^\dagger$ (and, further, to a finite integral extension $\mathcal{R}_{z',K'}/\mathcal{R}_{z,K}$).

3.3.2. In I.2.4.3, we have reviewed Katz' functor from representations of fundamental groups of algebraic curves in characteristic p to unit-root F -(iso)crystals. This construction has a local analogue, and one has the following theorem of N. Tsuzuki [Tsu98]:

Theorem 3.3.3. *There is a natural equivalence of categories between the category of continuous $Q(W(k))$ -linear representations of $\text{Gal}(k((\frac{1}{z}))^{\text{sep}}/k((\frac{1}{z})))$ with finite inertia and the category of overconvergent unit-root F -isocrystals on $\text{Spec } k((\frac{1}{z}))$.*

In particular, disregarding the precise Frobenius structure, one gets a functor

$$\left\{ \begin{array}{l} p\text{-adic representations of } k((\frac{1}{z})) \\ \text{with finite inertia} \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{differential modules over } \mathcal{E}^\dagger \\ \text{with Frobenius structure} \end{array} \right\}$$

hence, losing even more structure, a functor

$$\left\{ \begin{array}{l} \text{finite } p\text{-adic representations of } k((\frac{1}{z})) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{differential modules over } \mathcal{R}_{z,K} \text{ with Frobenius} \\ \text{structure, which become trivial over some} \\ \text{finite extension } \mathcal{R}_{z',K'} \text{ of } \mathcal{R}_{z,K} \text{ as above} \end{array} \right\}$$

Definition 3.3.4. (R. Crew) A differential module M over \mathcal{R} is quasi-unipotent if there is a Galois extension $k((\frac{1}{z'}))/k((\frac{1}{z}))$ such that after tensoring with the corresponding finite extension $\mathcal{R}_{z',K'}$ of $\mathcal{R}_{z,K}$, M becomes an iterated extension of trivial differential modules over $\mathcal{R}_{z',K'}$.

There is an extensive work by Crew, S. Matsuda and Tsuzuki relating Galois representation theory and ramification theory of $k((\frac{1}{z}))$ and Christol-Mebkhout theory in the case of quasi-unipotent differential modules, cf. [Cr00]. In particular, they establish a dictionary between Swan conductor and p -adic irregularity (in the above sense), break decomposition (in the sense of Katz) and p -adic slope decomposition...

Examples. (i) A tame module M with Frobenius structure is quasi-unipotent by 3.2.8; indeed, it suffices to take a Kummer extension $k((\frac{1}{z^{1/n}}))/k((\frac{1}{z}))$.

(ii) $\mathcal{R}/\mathcal{R}(\partial - \pi z)$ is quasi-unipotent: take an Artin-Schreier extension defined by $z' - z'^p = z$.

(iii) p -adic Bessel $M_{\nu,\pi}$ for odd p : a cyclic extension of $k((\frac{1}{z}))$ of degree $2p$ suffices. Indeed, it suffices to deal with the case $\nu = 1/2$ (cf. 3.2.3). Then $(\frac{1}{z})^{\pm 1/2} \cdot e^{\pm 2\pi z}$ is a basis of solutions, and one reduces to the previous two examples.

Conjecture 3.3.5 (Crew’s p -adic local monodromy conjecture). *Any differential module M over \mathcal{R} , with Frobenius structure, is quasi-unipotent.*

This conjecture, which can be seen as a p -adic analogue of Grothendieck’s ℓ -adic local monodromy theorem for wild ℓ -adic coefficients, and as a *non-abelian* extension of 3.2.8 to the wild case, has now been solved [And02b]. The proof relies heavily on theorems 3.2.5, 3.2.6, 3.2.8.

A crucial test [And02a], which is at the origin of the solution, is the case of the *diadic* normalized Bessel module $M_{\nu,2}$ (this test was suggested by Mebkhout as a potential counterexample). This example provides a typical non-abelian instance of 3.3.5:

Theorem 3.3.6. *For $K = \mathbb{Q}_4$, the normalized Bessel differential $\mathcal{E}_{x,K}^\dagger$ -module M is quasi-unipotent.*

More precisely, let E be the supersingular elliptic curve over \mathbb{F}_4 with affine equation $Y^2 + Y = X^3 + \omega$, ($\omega^3 = 1, \omega \neq 1$); then the covering $E \rightarrow E/\text{Aut}(E) \cong \mathbb{P}^1$ is wildly totally ramified at the origin of E , and defines an unramified Galois extension of $\mathcal{E}_{x,K}^\dagger$ of degree 24, with group $\text{Aut}(E) \cong SL_2(\mathbb{Z}/3\mathbb{Z})$, which contains all solutions of M . The underlying Galois representation of $G_{\mathbb{F}_4((1/z))}$ corresponds to the action of $\text{Aut}(E)$ on $H_{\text{cris}}^1(E)$.

3.4. A non-archimedean Riemann-Hilbert equivalence.

3.4.1. Let us return to a general ordinary linear differential equation with polynomial or analytic coefficients

$$(*) \quad \alpha_\mu(z)\partial^\mu y + \cdots + \alpha_0(z)y = 0 \quad \text{on } S \subset \mathbb{A}^1, \text{ with } \partial = z \frac{d}{dz}.$$

In the p -adic case, contrary to the complex situation, there does *not* exist, in general, any étale covering $S' \rightarrow S$ such that $(*)$ has a full set of solutions in $\mathcal{O}_{S'}$.

Example: $\partial^2 y = 0$, $S \supset \mathcal{C}(]1 - \epsilon, 1[)$; there is no “ p -adic Riemann surface of the logarithm” on S .

However, there are interesting global situations where this does happen.

Example: $\partial y - zy = 0$ on $S = \mathbb{A}^1 : S' = D(1, 1^-) \xrightarrow{\log} S$, the infinite Galois étale covering with group $\mu_{p^\infty} \cong \mathbb{Q}_p/\mathbb{Z}_p$, trivializes the equation.

The object of this subsection is to investigate this phenomenon in full generality.

3.4.2. Let K be a complete non-archimedean field. We note that by definition, K -manifolds are ‘good’ in the sense of [Ber93, 1.2.6], *i.e.* any point has an affinoid neighborhood. This property simplifies the theory of coherent

sheaves (in the dictionary between rigid-analytic spaces and Berkovich analytic spaces, the theories of coherent sheaves and vector bundles correspond, *cf.* [Ber93, 1.3.4, 1.6])⁽¹⁶⁾.

On the other hand, each coherent sheaf M on S gives rise to an étale sheaf $M_{\text{ét}}$ on S [Ber93, 4.1.2]. Moreover, one has étale descent for coherent sheaves: descent data are effective (*cf.* proof of [Ber93, 4.1.10], and also [dJvdP96, 3.2.3] in the rigid setting). We shall use this only for locally free coherent sheaves (vector bundles) of rank n . These are in bijection with $H_{\text{ét}}^1(S, GL_n)$.

In the sequel, S will be a connected K -manifold, and \bar{s} a geometric point of S . We assume that the point s is K -rational, *i.e.* is a classical point with residue field K .

3.4.3. A continuous K -linear representation ρ of a topological group Γ will be called *discrete* if its coimage $\Gamma/\ker \rho$ is discrete, or equivalently, if ρ factors through a discrete group.

An *étale local system* on S is a locally constant étale sheaf of K -vector spaces on S .

Proposition 3.4.4. *There is a natural equivalence of categories*

$$\{ \text{discrete } \pi_1^{\text{ét}}(S, \bar{s})\text{-representations} \} \xrightarrow{\sim} \{ \text{étale local systems on } S \}.$$

PROOF. Let ρ be a discrete representation in a K -vector space V . This amounts to a continuous homomorphism $\pi_1^{\text{ét}}(S, \bar{s}) \rightarrow \text{Aut}_K V$, where $\text{Aut}_K V$ denotes the discrete group of K -linear automorphisms of V . On the other hand, any étale local system \mathcal{V} on S gives rise in the usual way to an étale Čech cocycle with values in $\text{Aut}_K V$, with $V = \mathcal{V}_s$, and its isomorphism class gives a well-defined element of $H_{\text{ét}}^1(S, \text{Aut}_K V)$. Such a cohomology class may also be interpreted as usual as an $\text{Aut}_K V$ -principal homogeneous space \mathcal{F} (as an étale sheaf) over S up to isomorphism. What has to be shown is that such a principal homogeneous space is representable by an étale covering space. By lemma 1.2.7. (ii), this is a local question on S . We thus may assume that S is affinoid, and that for some finite étale covering $S' \rightarrow S$, $\mathcal{F}_{S'}$ is representable by an étale covering of S' . In that case, the representability of \mathcal{F} follows from the descent theory of schemes.

Starting from a representation V of $\pi_1^{\text{ét}}(S, \bar{s})$, we have thus attached the $\text{Aut}_K V$ -principal homogeneous space and étale sheaf \mathcal{F} ; the looked for étale local system is the associated étale sheaf \mathcal{V} of K -vector spaces, with respect

⁽¹⁶⁾Actually there are a number of technicalities involved in the analytic étale topology. First of all, since we restrict our attention to strictly analytic spaces in order to have the dictionary with rigid analytic varieties at disposal, Berkovich's definition of the étale site has to be slightly modified: this point is discussed in detail in R. Huber's book [Hub96, 8.3]. Translated into rigid geometry, one gets the 'partially proper étale' site of *loc. cit.*. The associated sheaves correspond to the 'overconvergent/conservative/constructible étale sheaves' of Van der Put, Schneider, and De Jong [dJvdP96], *cf.* [Hub96, 8.2.12]. We don't insist further on these subtleties.

to the canonical representation V of $\text{Aut}_K V$ (the pull-back of this étale local system on \mathcal{F} is the constant sheaf with value V).

Conversely, from an étale local system \mathcal{V} , one attaches the $\text{Aut}_K V$ -principal homogeneous space \mathcal{F} viewed as an étale covering space (with $V = \mathcal{V}_{\bar{s}}$); the structure of $\pi_1^{\text{ét}}(S, \bar{s})$ -set on its fiber at \bar{s} comes from a continuous homomorphism $\pi_1^{\text{ét}}(S, \bar{s}) \rightarrow \text{Aut}_K V$ (the cohomology class of \mathcal{F} in $H^1(\pi_1^{\text{ét}}(S, \bar{s}), \text{Aut}_K V)$ is nothing but the latter homomorphism up to conjugation). \square

In the sequel of this section, we assume that K is of *characteristic zero*.

3.4.5. Let M be a vector bundle (= locally free \mathcal{O}_S -module of finite rank) with a connection: $\nabla : M \rightarrow \Omega_S^1 \otimes M$ (*i.e.* a K -linear map satisfying Leibniz rule). This can also be interpreted \mathcal{O}_S -linearly, in terms of first-order jets, as a section $M \rightarrow \mathcal{P}^1(M)$ of the natural projection $\mathcal{P}^1(M) \rightarrow M$.

The vector bundles with connection (*resp.* with integrable connection) form a tannakian category over K . The unit for \otimes is the trivial connection $d : \mathcal{O}_S \rightarrow \Omega_S^1$.

Because s is a classical point with residue field K (of characteristic zero), the stalk at s of the sheaf $M^\nabla = \ker \nabla$ of germs of horizontal sections is a K -vector space of rank the rank of M , which can be identified with the fiber $M_{(s)}$ (note that since S has a K -rational point, $(\mathcal{O}_{S_{\text{ét}}})^{d_{S_{\text{ét}}}} = K$).

To M (*resp.* $\mathcal{P}^1(M)$), one attaches an étale sheaf $M_{\text{ét}}$ (*resp.* $\mathcal{P}^1(M)_{\text{ét}}$) — locally free $\mathcal{O}_{S_{\text{ét}}}$ -modules. Thus ∇ also correspond to a section of the natural projection $\mathcal{P}^1(M)_{\text{ét}} \rightarrow M_{\text{ét}}$, or equivalently, to a K -linear map $\nabla_{\text{ét}} : M_{\text{ét}} \rightarrow \Omega_{S_{\text{ét}}}^1 \otimes M_{\text{ét}}$. The kernel of the latter map is an étale sheaf of K -vector spaces denoted by $M_{\text{ét}}^\nabla$ (the étale sheaf of germs of horizontal sections). Let t be point of S . The fiber of $M_{\text{ét}}^\nabla$ at t (*cf.* [Ber93, 4.2]) is a \bar{K} -vector space endowed with a discrete $\text{Gal}(\overline{\mathcal{H}(t)}/\mathcal{H}(t))$ -action. If $M_{\text{ét}}^\nabla$ is locally constant, the dimension of this vector space is independent of t ; taking $t = s$, we find that it is the rank of M .

Remark. Any classical point admits a neighborhood over which M^∇ is constant of rank equal to the rank of M (by the non-archimedean analogue of Cauchy's theorem on solutions of analytic differential equations). The constancy of $M_{\text{ét}}^\nabla$ imposes a constraint only at non-classical points of the Berkovich space S .

Theorem 3.4.6. *There is a natural equivalence of tannakian categories (the non-archimedean étale Riemann-Hilbert functor):*

$$\left\{ \begin{array}{l} \text{discrete finite dimensional} \\ \pi_1^{\text{ét}}(S, \bar{s})\text{-representations} \end{array} \right\} \xrightarrow{\text{RH}^{\text{ét}}} \left\{ \begin{array}{l} \text{vector bundles with integrable} \\ \text{connection } (M, \nabla) \text{ on } S \text{ such that} \\ M_{\text{ét}}^\nabla \text{ is an étale local system} \end{array} \right\}.$$

In this correspondence, the subspace of $\pi_1^{\text{ét}}(S, \bar{s})$ -invariants corresponds to the space of global sections of M^∇ .

PROOF. Using the previous proposition and étale descent of coherent modules, it suffices to show that the natural maps

$$\mathcal{V} \mapsto (\mathcal{V} \otimes_K \mathcal{O}_{S_{\text{ét}}}, \text{id} \otimes d_{S_{\text{ét}}}), \quad (M, \nabla) \mapsto M_{\text{ét}}^\nabla,$$

induce quasi-inverse functors

$$\left\{ \begin{array}{l} \text{locally constant étale sheaves} \\ \text{of fin. dim. } K\text{-vector spaces} \end{array} \right\} \rightleftharpoons \left\{ \begin{array}{l} \text{vector bundles with integrable} \\ \text{connection } (M, \nabla) \text{ such that} \\ M_{\text{ét}}^\nabla \text{ is an étale local system} \end{array} \right\}$$

It is clear that these maps are functorial (and the first one is clearly compatible with the tensor structures). The point is to show that they are quasi-inverse.

On the one hand, it is clear that $\text{id} \otimes d_{S_{\text{ét}}}$ is an integrable connection, and that $(\mathcal{V} \otimes_K \mathcal{O}_{S_{\text{ét}}})^{\text{id} \otimes d_{S_{\text{ét}}}} = \mathcal{V}$ (taking into account that $(\mathcal{O}_{S_{\text{ét}}})^{d_{S_{\text{ét}}}} = K$). We have to show that if (M, ∇) is a vector bundle with integrable connection such that $M_{\text{ét}}^\nabla$ is locally constant, then the natural horizontal map

$$\phi : M_{\text{ét}}^\nabla \otimes_K \mathcal{O}_{S_{\text{ét}}} \rightarrow M_{\text{ét}}$$

is an isomorphism. By the remark preceding the theorem, the stalks of $M_{\text{ét}}^\nabla$ are \bar{K} -vector spaces of constant dimension, which is the rank of M . It follows that $M_{\text{ét}}^\nabla \otimes_K \mathcal{O}_{S_{\text{ét}}}$ is the étale sheaf attached to a vector bundle of the same rank as M . It is a general well-known fact about integrable connections⁽¹⁷⁾ that ϕ is a monomorphism of vector bundles, and that $\text{Im } \phi$ is locally a direct summand of $M_{\text{ét}}$ (because the cokernel inherits an integrable connection, hence is locally free). To show that this it is an epimorphism, it thus suffices to check that $M_{\text{ét}}^\nabla \otimes_K \mathcal{O}_{S_{\text{ét}}}$ and $M_{\text{ét}}$ have the same rank, which is the case by assumption. \square

3.4.7. The $\pi_1^{\text{ét}}(S, \bar{s})$ -representation attached to (M, ∇) is called the *monodromy representation*. Note that the underlying vector space is canonically identified with the fiber M_s . The image of the monodromy representation is called the *monodromy group*. It may be non-discrete (while the coimage is discrete by assumption); this phenomenon is not specific to the non-archimedean situation, and is indeed familiar in the complex case.

3.4.8. The étale Riemann-Hilbert functor unifies several earlier constructions. Indeed, if the monodromy representation factors through $\pi_1^{\text{top}}(S, s)$, we recover the ‘topological Riemann-Hilbert functor’ considered in I.1.5, whose image consists of connections with locally constant sheaves of solutions.

⁽¹⁷⁾proven by passing to the completion around classical points.

It also includes some cases of overconvergent unit-root isocrystals with finite monodromy already considered in I.2.5 (in that case, the monodromy factors through a discrete, hence finite, representation of $\pi_1^{\text{alg}}(S, \bar{s})$).

Of course, connections in the image of the étale Riemann-Hilbert functor are still rather special⁽¹⁸⁾, but it seems difficult to extend its scope without invoking “relative Fontaine rings”.

Remarks 3.4.9. (i) Whether $M_{\text{ét}}^{\nabla}$ is an étale local system or not can be checked locally on S . In particular, the property for a connection to be in the essential image of $RH^{\text{ét}}$ is local.

(ii) The étale fundamental group $\pi_1^{\text{ét}}(\mathbb{P}_{\mathbb{C}_p}^1, \bar{s})$, although highly non-trivial, does not have any non-trivial discrete finite-dimensional representation. Indeed, the corresponding connection on $\mathbb{P}_{\mathbb{C}_p}^1$ would be algebrizable, and necessarily trivial.

3.5. Examples and applications.

3.5.1. For any full subcategory Cov_S^{\bullet} of $\text{Cov}_S^{\text{ét}}$ which is stable under taking connected components, fiber products and quotients, we have seen that there is a natural continuous homomorphism

$$\pi_1^{\text{ét}}(S, \bar{s}) \rightarrow \pi_1^{\bullet}(S, \bar{s})$$

with dense image (1.4.8). Any discrete representation of $\pi_1^{\bullet}(S, \bar{s})$ gives rise to a discrete representation of $\pi_1^{\text{ét}}(S, \bar{s})$ with the same coimage. It follows that the étale Riemann-Hilbert functor induces a fully faithful functor

$$\left\{ \begin{array}{l} \text{discrete fin. dim.} \\ \pi_1^{\bullet}(S, \bar{s})\text{-representations} \end{array} \right\} \xrightarrow{RH^{\bullet}} \left\{ \begin{array}{l} \text{vector bundles with integrable} \\ \text{connection } (M, \nabla) \text{ on } S \end{array} \right\}.$$

Note that the monodromy group defined in 3.4.7 coincides with the image of $\pi_1^{\bullet}(S, \bar{s})$. We shall discuss the avatars RH^{\bullet} in the cases of $\text{Cov}_S^{\text{top}}$, $\text{Cov}_S^{\text{alg}}$, $\text{Cov}_S^{\text{loc.alg}}$, $\text{Cov}_S^{\text{temp}}$.

3.5.2. The functor RH^{top} was studied in I.1.5 (in the case $K = \mathbb{C}_p$). Its essential image consists of connections which satisfy ‘Cauchy’s theorem’ at every (possibly non-classical) point of S , or equivalently, which have a full set of multivalued analytic solutions, or else, such that M^{∇} is a locally constant sheaf on the topological space S .

3.5.3. The essential image of the functor RH^{alg} consists of objects (M, ∇) such that $M_{\text{ét}}^{\nabla}$ becomes constant on some finite étale covering of S .

Let us assume for instance that S is affinoid with good reduction S^0 , and that K is a finite extension of $\widehat{\mathbb{Q}_p^{\text{ur}}}$ as in I.2.4. Then among the finite

⁽¹⁸⁾for instance, it does not account for the differential equation of the logarithm, as we have said in 3.4.1; to see this, one can notice that otherwise the monodromy group would be contained in the additive group of K , hence torsion-free, but this implies that the étale covering which trivializes the connection is a topological covering (2.1.9), hence is trivial.

representations of $\pi_1^{\text{alg}}(S, \bar{s})$, one can distinguish those which factor through $\pi_1^{\text{alg}}(S_0, \bar{s}_0)$. The corresponding connections underlie overconvergent unit-root isocrystals with finite global monodromy. We have seen a standard example in I.2.4.5, given by the differential equation $\frac{d}{dz}y = -\pi y$ on $S = \mathbb{D}(0, 1^+)$, which is trivialized over an Artin-Schreier covering.

Let us now consider a case of bad reduction, namely the case of an annulus $S_r = \mathcal{C}(]1, r[)$, $r \in |K|, r > 1$ over any discretely valued p -adic field K . Let \mathcal{R} be the Robba ring at infinity. Then RH^{alg} induces a functor

$$\{\text{finite representations of } \pi_1^{\text{alg}}(S_r, \bar{s})\} \rightarrow \{\mathcal{R}\text{-differential modules}\}.$$

In relation to the local monodromy theorem 3.3.5, it would be very interesting to understand the essential image of this functor as $r \rightarrow 1^+$. Note however that there is a difficulty with base-points in passing to the limit $r \rightarrow 1^+$ (the problem disappears if one considers only abelian representations).

Nevertheless, any \mathcal{R} -differential module M extends to a vector bundle (M_r, ∇_r) with connection on some S_r . Let us assume that M is semisimple and has a Frobenius structure.

If M is tame, then 3.2.8 ensures that M is in the image of the above functor

$$\{\text{finite } \pi_1^{\text{alg}}(S_r, \bar{s})\text{-representations}\} \rightarrow \{\mathcal{R}\text{-differential modules}\}$$

for r close enough to 1. The local monodromy theorem 3.3.5 implies the same if M is wild, although the monodromy representation might be non-abelian in this case, as 3.3.6 shows.

3.5.4. The essential image of the functor $RH^{\text{loc.alg}}$ consists of objects (M, ∇) such that for every compact subdomain $W \subset S$, $(M_{\text{et}}^{\nabla})|_W$ becomes constant on some finite étale covering of W . Although not explicitly introduced in [Ram98], it is clear that these connections played an important heuristic role in that paper, especially those coming from abelian representations of the locally algebraic fundamental group (1.5).

The basic example is $(\mathcal{O}_{\mathbb{A}^1}, d-1)$ over \mathbb{C}_p (with solution e^z). It is trivialized by the logarithmic étale covering of \mathbb{A}^1 .

Similarly, for any ramified polynomial $P(z^{1/m})$, the differential equation with basic solution $e^{P(z^{1/m})}$ is trivialized over a locally algebraic étale covering of a sufficiently small punctured disk $\mathbb{D}_{\mathbb{C}_p}(\infty, \epsilon)^*$ around ∞ (the logarithmic covering of a Kummer covering). By 3.1.5, we see that any differential equation with polynomial coefficients — or more generally meromorphic singularity at ∞ — with *rational* formal exponents at ∞ , and with no logarithm in the formal solutions \hat{y} (cf. 3.1.2), corresponds to an object in the image of the locally algebraic Riemann-Hilbert functor

$$\left\{ \begin{array}{c} \text{finite dimensional} \\ \pi_1^{\text{loc.alg}}(\mathbb{D}(\infty, \epsilon)^*, \bar{s})\text{-representations} \end{array} \right\} \rightarrow \{\mathcal{O}(\mathbb{D}(\infty, \epsilon)^*)\text{-differential modules}\}$$

for ϵ small enough.

There is a certain analogy between the logarithmic étale covering of $D_{\mathbb{C}_p}(\infty, \epsilon)^*$ and the *real blow-up* of $D_{\mathbb{C}}(\infty, \epsilon)^*$ which plays such an important role in the study of irregular singularities of complex differential equations. In both cases, the point at infinity becomes a limiting circle⁽¹⁹⁾.

Let us return to a general vector bundle (M, ∇) with connection on $D(\infty, 1^-)^*$ which extends to a vector bundle with meromorphic connection at ∞ . We assume that it has a Frobenius structure near the unit circle, and that the formal exponents at infinity are rational. Regarding problem 3.2.9, a natural question is:

Question 3.5.5. *Under which further conditions, if any, does there exist a locally algebraic covering of $D(\infty, 1^-)^*$ over which the connection becomes an iterated extension of trivial connections?*

At any rate, in the cases where this holds, this would let Ramero's locally algebraic fundamental group play the same role, over \mathbb{C}_p , as Ramis' wild fundamental group (over \mathbb{C}) in the local study of meromorphic irregular singularities.

3.5.6. The essential image of the functor RH^{temp} consists of objects (M, ∇) such that $M_{\text{ét}}^{\nabla}$ becomes constant on some temperate étale covering of S . This implies that there is a finite étale Galois covering $f: T \rightarrow S$ such that the pull-back $(M', \nabla') = f^*(M, \nabla)$ has a full set of multivalued analytic solutions, *i.e.* $M'^{\nabla'}$ is locally constant. Let G be the group $\text{Aut}_S T$, and set $n = |G|$.

Assume that there are n distinct K -rational points t_i in T lying above s . Then the monodromy group admits the following *concrete interpretation*: let α_{ij} be a Berkovich path (up to homotopy) connecting t_i to t_j . Then $f_*(\alpha_{ij}) \in \pi_1^{\text{temp}}(S, \bar{s})$. When i, j and α_{ij} vary, they generate the image $\text{Aut}_S \tilde{T}$ of $\pi_1^{\text{temp}}(S, \bar{s})$ (note that, varying j , they generate the finite quotient G). It follows that the monodromy group consists of the images of the $f_*(\alpha_{ij})$'s in $GL(M_{(s)})$ by the monodromy representation.

Therefore, except for the occurrence of the finite étale covering, the picture is quite similar to the complex situation.

Proposition 3.5.7. *Let $S = \mathbb{A}^1 \setminus \{\zeta_1, \dots, \zeta_m\}$. Assume⁽²⁰⁾ that the residue characteristic p of K does not divide n . Then:*

- (a) *for any open or closed disk $D \subset S$, the restriction of ∇ to D is trivial (i.e. has a full set of solutions); in particular, ∇ satisfies the Robba condition;*
- (b) *for any open or closed annulus $\mathcal{C} \subset S$, the pull-back of ∇ to the Kummer covering of \mathcal{C} of degree n is trivial.*

⁽¹⁹⁾L. Ramero points out that the change from analytic to étale topology is essential in the p -adic case, while in the complex case, the real blow-up corresponds to completion for a different uniform structure of the punctured disk with no change of the topology.

⁽²⁰⁾this very strong assumption is rarely met in practice.

In particular, p -adic exponents in the sense of Christol-Mebkhout exist (in any annulus which does not contain any ζ_i), and they are rational numbers with denominator n (or a divisor of n).

PROOF. One is immediately reduced to considering closed disks and annuli. We know the pull-back of ∇ to \tilde{T} is trivial. Let D' (resp. C') be a connected component of $D \times_S \tilde{T}$ (resp. $C \times_S \tilde{T}$). Because n is assumed to be prime to p , the covering $D' \rightarrow D$ is trivial (resp. is a Kummer covering of degree dividing n), cf. [Ber93, 6.3.3, 6.3.5] (taking into account the fact that disks and annuli are simply-connected). The assertions follow. \square

Example. (see also the example at the end of 2.1.2). Let K be a p -adic field for any odd p . Let $T = E \setminus E[2]$ be a Tate curve deprived from its four 2-torsion points (assumed to be K -rational), written in Legendre form:

$$y^2 = x(x - 1)(x - \lambda), \quad |\lambda(\lambda - 1)| \neq 1.$$

Let f be the projection from T to $S = \mathbb{A}^1 \setminus \{0, 1, \lambda\}$. Here \tilde{T} is of the form $\mathbb{G}_m \setminus \pm(\sqrt{q})^{\mathbb{Z}}$, and $T \cong \tilde{T}/q^{\mathbb{Z}}$.

The sum of the canonical one-dimensional K -linear representation of $q^{\mathbb{Z}}$ (q acting by multiplication by a factor q) and its dual extends to a representation of the semi-direct product of $q^{\mathbb{Z}}$ and $\mathbb{Z}/2\mathbb{Z}$ (which is the Galois group of the covering $\tilde{T} \rightarrow S$). One can choose the local monodromies at $0, \lambda, 1, \infty$ respectively in such a way that they correspond to the matrices

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & q \\ q^{-1} & 0 \end{pmatrix}, \begin{pmatrix} 0 & q^{-1} \\ q & 0 \end{pmatrix}$$

respectively. It is then easy to check that the monodromy around an annulus surrounding 0 and λ alone ($|\lambda|$ being assumed < 1 for convenience) is trivial: the p -adic exponents are zero.

3.5.8. Much more interesting examples arise in the context of p -adic period mappings. We place ourselves in the setting of II.7.2. Assuming the congruence subgroup sufficiently small, the picture was the following:

$$\begin{array}{ccccc} \tilde{S} & \longrightarrow & \mathcal{M} & \xrightarrow{p} & \mathcal{D} \\ & \searrow & \downarrow & & \downarrow \\ & & S & \longrightarrow & \mathcal{D}/\Gamma \end{array}$$

where S was a suitable connected ‘tube-like’ domain in the Shimura variety $Sh_{\mathbb{C}_p}$ (\mathcal{M}^0 was a deformation space for p -divisible groups quasi-isogenous to a fixed one, and \mathcal{D} was the period domain).

The period mapping turned out to be expressable in terms of quotients of solutions of the Gauss-Manin connection (on S), and the Gauss-Manin connection was trivializable on \mathcal{M}^0 , hence on \tilde{S} (II.7.2.6).

Let us now abandon the assumption that the congruence subgroup is sufficiently small. Then a similar picture holds true: Γ is now only virtually torsion-free, and the covering $\mathcal{M}^0 \rightarrow S$ is now ramified. By removing the singularities, one obtains a *temperate étale covering*, which trivializes the Gauss-Manin connection (using II.7.2.6 after passing to a suitable finite étale covering). It follows that the *Gauss-Manin connection (restricted to S deprived from the singularities) is in the image of the temperate Riemann-Hilbert functor*.

The situation is therefore similar to the complex one, except that S is only a small domain in the Shimura variety in general. The most favorable situation occurs in the case of global uniformization (II.7.3), *i.e.* when S is a component of $\mathcal{S}h$.

3.5.9. Let us now give a criterion for a connection in the image of RH^{et} to be in the image of RH^{temp} .

Proposition 3.5.10. *Let $\rho : \pi_1^{\text{et}}(S, \bar{s}) \rightarrow GL(V)$ be a finite-dimensional discrete representation (corresponding to an étale local system \mathcal{V}). Let us consider the following conditions:*

- (a) ρ factors through a discrete representation of $\pi_1^{\text{temp}}(S, \bar{s})$,
- (b) $\text{Im } \rho$ is virtually torsion-free,
- (c) $\text{Im } \rho$ is finitely generated.

Then (c) \Rightarrow (b) \Rightarrow (a). If moreover S is the analytification of a smooth algebraic K -variety, then (a) \Rightarrow (c).

PROOF. (c) \Rightarrow (b): this is Selberg's lemma: any finitely generated subgroup of $GL(V)$ is virtually torsion-free.

(b) \Rightarrow (a) follows from 2.1.9. (b).

(a) \Rightarrow (c): under (a), $\text{Coim } \rho$ is the Galois group of a Galois temperate covering of S . If S is algebraic, $\text{Coim } \rho$ must be finitely generated by 2.1.7. \square

3.5.11. We finish the section by showing that the connections which lie in the image of the temperate Riemann-Hilbert functor are algebraizable (this is not automatic in the non-archimedean situation), and even fuchsian:

Theorem 3.5.12. *Assume that S is the analytification of a smooth algebraic K -variety S^{alg} . Let (M, ∇) be a vector bundle with integrable connection on S which comes from a discrete representation of $\pi_1^{\text{temp}}(S, \bar{s})$. Then (M, ∇) is the analytification of a (unique) algebraic vector bundle with integrable connection on S^{alg} , which has regular singularities and exponents in \mathbb{Q}/\mathbb{Z} at any divisorial valuation of $K(S)$.*

PROOF. By assumption, there is a finite Galois étale covering $S' \rightarrow S$ (with group G) such that (M, ∇) becomes trivial on the topological universal covering of S' . The pull-back $(M, \nabla)_{S'}$ corresponds to a (unique) representation of $\pi_1^{\text{top}}(S', s')$ (for any point s' above s). Let \bar{S}' be a smooth algebraic compactification of S' (Hironaka). We may replace K by the completion of its

algebraic closure. By 1.1.4, the homomorphism $\pi_1^{\text{top}}(S', s') \rightarrow \pi_1^{\text{top}}(\overline{S}', s')$ is an isomorphism, and it follows that $(M, \nabla)_{S'}$ extends (uniquely) to a vector bundle with integrable connection on \overline{S}' . By GAGA (Köpf), the latter is the analytification of a (unique) algebraic vector bundle with integrable connection on $\overline{S}'^{\text{alg}}$. It is now easy to conclude by Galois descent from S'^{alg} to S^{alg} (we refer to [AB01] for the details on regular singularities, exponents, and GAGA functor for connections). \square

3.5.13. Besides discrete representations of $\pi_1^{\text{temp}}(S, \overline{s})$, it is also useful to consider *projective discrete representations*, *i.e.* discrete representations with value in $PGL(V)$. They give rise to *projective connections* over S^{alg} , at least when S^{alg} is an *affine curve*.

Indeed, replacing S^{alg} by a finite Galois étale covering S'^{alg} with Galois group G , we may replace $\pi_1^{\text{temp}}(S')$ by $\pi_1^{\text{top}}(S')$, which is a free group, for which the lifting property is immediate. This provides a connection ∇' over S'^{alg} , and the associated projective connection is endowed with a G -action compatible with the G -action on S'^{alg} , hence descends to a projective connection on S^{alg} .

One has more:

Proposition 3.5.14. *Then any projective discrete representation $\overline{\rho} : \pi_1^{\text{temp}}(S, \overline{s}) \rightarrow PSL(V)$ lifts to a genuine discrete representation $\rho : \pi_1^{\text{temp}}(S, \overline{s}) \rightarrow SL(V)$, after replacing K by a finite extension.*

PROOF. We have seen that $\overline{\rho}$ corresponds to a projective connection on S^{alg} . Now it is well known that any projective connection lifts to a genuine connection after replacing K by a finite extension (because $H^2(S_{K,et}^{\text{alg}}, \mathbb{Z}/2\mathbb{Z}) = 0$, cf. [Ka90, 2.2.2.1]). It suffices to check that the pull-back ∇'' of this connection on S'^{alg} comes from a discrete representation $\rho : \pi_1^{\text{temp}}(S', \overline{s}') \rightarrow SL(V)$. But we have on the other hand our connection ∇' which is projectively equivalent to ∇'' , and which we may assume to come from a discrete representation $\pi_1^{\text{temp}}(S', \overline{s}') \rightarrow SL(V)$. Let S'' be a finite étale covering of S' such that the pull-back of any class in $H^1(S_{K,et}^{\text{alg}}, \mathbb{Z}/2\mathbb{Z})$ vanishes in $H^1(S''_{K,et}{}^{\text{alg}}, \mathbb{Z}/2\mathbb{Z})$. Since ∇' and ∇'' are projectively equivalent, their pull-backs on S'' are isomorphic (*loc. cit.*). It follows that ∇'' comes from a discrete representation $\pi_1^{\text{temp}}(S', \overline{s}') \rightarrow SL(V)$. \square

We leave it as an exercise to translate 3.5.10 in the context of projective connections.

4. Non-archimedean orbifolds, uniformizing differential equations and period mappings.

ABSTRACT: We review the complex case, and then define and study non-archimedean orbifolds, their associated orbifold fundamental groups, and uniformizing differential equations.

4.1. Review of complex orbifolds.

4.1.1. There are several variants of the notion of complex orbifold. We follow [DM93] and see orbifolds as being locally quotients of a complex manifold by a finite group acting faithfully. More precisely, a *complex orbifold* $\mathcal{S} = (\overline{S}, (Z_i, e_i))$ consists in

- a normal analytic space \overline{S} ,
- a locally finite collection of irreducible (Weil) divisors (Z_i) ,
- a positive integer⁽²¹⁾ e_i attached to each Z_i .

It is assumed that \overline{S} is covered by so-called *orbifold charts*, *i.e.* morphisms $W \xrightarrow{\phi} V \subset \overline{S}$ where

- W is a complex manifold,
- ϕ is a (ramified) covering in the sense of 1.2.3 (*i.e.* V is covered by open subsets U such that $\coprod V_j = \phi^{-1}U \rightarrow U$ and the restriction of ϕ to every V_j is finite),
- except for a closed subset Σ of codimension 2 in V , ϕ is étale above $V \setminus (V \cap \bigcup Z_i)$ (hence induces a topological covering of $V \setminus (V \cap \bigcup Z_i)$), and is *ramified with index e_i above Z_i* .⁽²²⁾

Note that if s is a non-singular point of \overline{S} not contained in $\bigcup Z_i$, then any smooth open neighborhood of s in \overline{S} disjoint from $\bigcup Z_i$ provides an orbifold chart around s . Hence we may and shall assume that Σ is contained in $\bigcup Z_i \cup \overline{S}^{\text{sing}}$ (where $\overline{S}^{\text{sing}}$ denotes the set of singular points of \overline{S}).

We shall often write $Z = \bigcup Z_i \cup \overline{S}^{\text{sing}}$, and $S = \overline{S} \setminus Z$ (which is a manifold).

Remark. By shrinking the W 's, one can assume that they are open subsets of \mathbb{C}^n ; this matches with the conventional use of the word ‘chart’. However, we do not impose here this condition which would have no non-archimedean counterpart.

4.1.2. A *morphism of orbifolds* $f : \mathcal{S}' \rightarrow \mathcal{S}$ is a morphism $\overline{S}' \rightarrow \overline{S}$, such that \overline{S}' is covered by orbifold charts $\phi' : W' \rightarrow V'$ with the property that $f \circ \phi'$ factors through some orbifold chart $\phi : W \rightarrow V$ of \mathcal{S} : $f \circ \phi' = \phi \circ f'$ (for some morphism $f' : W' \rightarrow W$).

⁽²¹⁾in other variants, one also allows $e_i = \infty$. This leads to ‘parabolic structures’ which have indeed non-archimedean counterparts in positive characteristic (cusps of Drinfeld modular curves...). But we shall be mostly interested in non-archimedean orbifolds in characteristic zero, where such ‘parabolic structures’ cannot occur.

⁽²²⁾it is easily seen that this definition is equivalent to that of [DM93], using *loc. cit.* 14.2.

Example. Let $(\overline{S}, (Z_i, e_i)_{i \in I})$ be an orbifold, and let $J \subset I$ be the subset of indices for which $e_i > 1$. Then the identity of \overline{S} induces an isomorphism $(\overline{S}, (Z_i, e_i)_{i \in I}) \cong (\overline{S}, (Z_i, e_i)_{i \in J})$. In other words, one can ‘forget’ the divisors with index $e_i = 1$.

Considering complex manifolds as orbifolds with an empty collection of divisors ($S = \overline{S}$), one can embed the category of complex manifolds as a full subcategory of the category of complex orbifolds.

It will be convenient to attach to any irreducible divisor $D \subset \overline{S}$ an index $e_D \in \mathbb{Z}_{>0}$ which is e_i if $D = Z_i$ and 1 if D is none of the Z_i ’s.

4.1.3. A morphism f is called an *orbifold-covering* if it is a covering in the sense of 1.2.3 and for any orbifold chart $\phi' : W' \rightarrow V'$ of \mathcal{S}' , $f \circ \phi' : W' \rightarrow V = f(V')$ is an orbifold chart of \mathcal{S} (in loose words: any lifting f' as in 4.1.2 is étale).

The restriction of f above S is then an étale covering: indeed, this is clear except above a closed subset of codimension ≥ 2 in S , thus above the whole of S by purity since S is smooth.

Example. $(\mathbb{A}^1, (0, e')) \xrightarrow{z \mapsto z^n} (\mathbb{A}^1, (0, e))$ is a morphism if and only if $e | ne'$, and is an orbifold-covering if and only if $e = ne'$.

Remark. Let $f : \mathcal{S}' \rightarrow \mathcal{S}$ be an orbifold-covering. As we just saw, we may assume that all the indices e'_i, e_j are > 1 . Then $f(Z') \subset Z$ and $f^{-1}(S) \subset S'$.

4.1.4. Following Thurston, an orbifold is said to be good or *uniformizable* if it admits a *global* orbifold chart (*i.e.* with $V = \overline{S}$). In that case, there exists a *universal* global orbifold chart, called the *universal orbifold-covering* (it is indeed an orbifold-covering in the above sense); its underlying space is a simply-connected complex manifold if \overline{S} is connected.

Example. By a classical result due to Fox, $(\mathbb{P}^1, (\zeta_i, e_i)_{i=0, \dots, n})$ is uniformizable if and only if $n \geq 2$ or ($n = 1$ and $e_0 = e_1$). Under these conditions, there exist algebraic global charts which are finite coverings of \mathbb{P}^1 .

4.1.5. It is proved in *loc. cit.* that orbifold charts are locally unique. Apart from a subset of codimension 2, they are given in local coordinates around $\phi^{-1}(Z_i)$ by: $(z_1, z_2, \dots, z_n) \mapsto (z_1^{e_i}, z_2, \dots, z_n)$.

It follows that there exist orbifold charts of the form $W \rightarrow V = W/G$ (*loc. cit.* 14.6), where W is an open subset of \mathbb{C}^n stable under a finite group G of linear transformations.

4.1.6. Let $\mathcal{S} = (\overline{S}, (Z_i, e_i))$ be a connected orbifold, and let Γ be a discrete group acting faithfully and properly by automorphisms on \mathcal{S} . The *orbifold quotient* \mathcal{S}/Γ is defined as follows: the underlying analytic space is \overline{S}/Γ . The ramifications divisors are the images of the Z_i ’s, and the irreducible divisors D with non-trivial fixer Γ_D ; the latter are endowed with the multiplicity $e_D \cdot |\Gamma_D|$.

Note that the projection $\mathcal{S} \rightarrow \mathcal{S}/\Gamma$ is an orbifold-covering.

4.1.7. One of the main applications of complex orbifolds is to represent moduli problems for isomorphism classes of objects with finite automorphism group. We have already alluded to this in II.1.1. Here we shall content ourselves with examining the case of complex elliptic curves.

Let H be the subgroup of $SL_3(\mathbb{R})$ consisting of matrices of the form $\begin{pmatrix} 1 & x_1 & x_2 \\ 0 & a & b \\ 0 & c & d \end{pmatrix}$

(this is a semi-direct product $SL_2(\mathbb{R}) \ltimes \mathbb{R}^2$). It acts on the product $\mathfrak{h} \times \mathbb{C}$ of the Poincaré upper half plane \mathfrak{h} and \mathbb{C} by the rule

$$\begin{pmatrix} 1 & x_1 & x_2 \\ 0 & a & b \\ 0 & c & d \end{pmatrix} \cdot (\tau, v) = \left(\frac{a\tau + b}{c\tau + d}, \frac{v + x_1\tau + x_2}{c\tau + d} \right).$$

As is well-known, elliptic curves E together with a symplectic isomorphism $H_1(E, \mathbb{Z}) \cong \mathbb{Z}^2$ are classified by points of the Poincaré upper half plane \mathfrak{h} , the universal object being the quotient of $\mathfrak{h} \times \mathbb{C}$ by the action of the subgroup $1 \cdot \mathbb{Z}^2 \subset H$ (mapping to \mathfrak{h} through the first projection).

Let $\text{Ker}(SL_2(\mathbb{Z}) \rightarrow SL_2(\mathbb{Z}/N\mathbb{Z}))$ be the full congruence group of level N , $\Gamma(N)$ its image in $PGL(2)$, and $\tilde{\Gamma}(N)$ be the subgroup

$$\text{Ker}(SL_2(\mathbb{Z}) \rightarrow SL_2(\mathbb{Z}/N\mathbb{Z})).\mathbb{Z}^2 \subset H.$$

For $N \geq 3$, the group $\tilde{\Gamma}(N)$ is torsion-free and acts freely on $\mathfrak{h} \times \mathbb{C}$; elliptic curves together with a level N structure (*cf.* II.1.1.2) are classified by $\mathfrak{h}/\Gamma(N)$, the universal object being $(\mathfrak{h} \times \mathbb{C})/\tilde{\Gamma}(N) \rightarrow \mathfrak{h}/\Gamma(N)$.

This still holds for $N = 1$ or 2 , provided one interprets these quotients as orbifold quotients. For $N = 2$, the element $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in \tilde{\Gamma}(N)$ of order 2 has fixed divisors in $\mathfrak{h} \times \mathbb{C}$: the quotient of $\{(\tau, x_1\tau + x_2) \mid x_1, x_2 \in (\frac{1}{2}\mathbb{Z})^2\}$ by $\tilde{\Gamma}(N)$, which is a disjoint union $\coprod_1^4 Z_i$ of four ‘horizontal’ curves (horizontal with respect to the projection to $\mathfrak{h}/\Gamma(2) = \mathbb{P}^1 \setminus \{0, 1, \infty\}$); they have the index $e_i = 2$ (the order of the fixer).

This orbifold is uniformizable: the Legendre elliptic pencil provides a global orbifold chart. Note however that, as an analytic variety, $\mathfrak{h} \times \mathbb{C}/\tilde{\Gamma}(2)$ is not an elliptic surface but a \mathbb{P}^1 -bundle.

The case $N = 1$ is similar: $\mathfrak{h}/\Gamma(1) = \mathbb{A}^1$ is the j -line. It should be considered as an orbifold, with divisor $\{j = 0 \text{ (index 3)}, j = 1728 \text{ (index 2)}\}$. In addition to the image of the horizontal divisors Z_i on $\mathfrak{h} \times \mathbb{C}/\tilde{\Gamma}(1)$, there are three vertical divisors with index 2, and two vertical divisors with index 3.

4.1.8. More generally, let us consider a Shimura variety of PEL type as in II.1.2.4. Recall from *loc. cit.* the commutative diagram

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\mathcal{P}} & \mathcal{D} \\ \mathcal{Q} \downarrow & & \downarrow \mathcal{Q} \\ S & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma \end{array}$$

where S is a component of the Shimura variety, the horizontal maps are *isomorphisms*, and Γ is a congruence subgroup. If Γ has torsion, it is appropriate to consider S as an orbifold quotient (defined by \mathcal{Q} as in 4.1.6).

4.2. Orbifold fundamental groups.

4.2.1. Another important application of orbifolds is the Galois theory of *ramified* coverings.

Let $\mathcal{S} = (\overline{S}, (Z_i, e_i))$ be a uniformizable connected orbifold. We denote by $\tilde{\mathcal{S}}$ the universal orbifold-covering (which is a simply-connected complex manifold, together with a (ramified) covering morphism $\tilde{\mathcal{S}} \rightarrow \overline{S}$). We fix a base-point s of $S = \overline{S} \setminus (\bigcup Z_i \cup \overline{S}^{\text{sing}})$, and a point \tilde{s} of $\tilde{\mathcal{S}}$ above s . Then the pair $(\tilde{\mathcal{S}}, \tilde{s})$ is unique up to unique isomorphism.

4.2.2. Each divisor Z_i defines a conjugacy class of elements γ_i in $\pi_1^{\text{top}}(S, s)$: the local monodromy at Z_i based at s .

The *orbifold fundamental group* $\pi_1^{\text{orb}}(\mathcal{S}, s)$ is the quotient of $\pi_1^{\text{top}}(S, s)$ by the *normal subgroup* $\langle \gamma_i^{e_i} \rangle$ generated by the $\gamma_i^{e_i}$'s.

This definition is justified by the following fact (*cf.* [DM93, 14.10]): one has a commutative square of functors

$$\begin{array}{ccc} \{\text{orbifold-coverings of } \mathcal{S}\} & \xrightarrow{\subset} & \{\text{étale (= topological) coverings of } S\} \\ \wr \downarrow & & \downarrow \wr \\ \{\pi_1^{\text{orb}}(\mathcal{S}, s)\text{-sets}\} & \xrightarrow{\subset} & \{\pi_1^{\text{top}}(S, s)\text{-sets}\} \end{array}$$

where the top horizontal functor is the (fully faithful) restriction functor, and the vertical functors are equivalences of categories.

The orbifold fundamental group may also be identified with the group of automorphisms of the (ramified) covering $\tilde{\mathcal{S}} \rightarrow \overline{S}$. Galois orbifold-coverings correspond to normal subgroups of π_1^{orb} .

Example. Let $\mathcal{S} = (\mathbb{P}^1, (\zeta_i, e_i)_{i=0, \dots, n})$ with $n \geq 2$ or ($n = 1$ and $e_0 = e_1$). Then $\pi_1^{\text{orb}}(\mathcal{S}, s)$ has generators $\gamma_0, \dots, \gamma_n$ and relations $\gamma_0^{e_0} = \dots = \gamma_n^{e_n} = \prod \gamma_i = 1$. In particular, it depends only, up to isomorphism, on the collection of integers e_i , and not on the ζ_i 's.

The universal orbifold-covering is $\mathbb{P}^1, \mathbb{A}^1$ or the open unit disk \mathbb{D} according to whether $\sum(1 - \frac{1}{e_i})$ is < 2 , $= 2$, or > 2 respectively.

Remark. The *finite* orbifolds coverings are classified by a profinite group $\pi_1^{\text{alg,orb}}(\mathcal{S}, s)$, which is actually the profinite completion of $\pi_1^{\text{orb}}(\mathcal{S}, s)$. When \overline{S} is algebraic, this profinite group has a purely algebraic definition.

4.2.3. Let $\phi : \overline{S}' \rightarrow \overline{S}$ be a global orbifold chart, such that the induced map $S' \rightarrow S$ is a Galois (topological) covering with group G . Let s' be a point of S' above s . Then one has an exact sequence

$$1 \rightarrow \pi_1^{\text{top}}(\overline{S}', s') \rightarrow \pi_1^{\text{orb}}(\mathcal{S}, s) \rightarrow G \rightarrow 1.$$

4.3. Uniformizing differential equations.

4.3.1. We follow the exposition in [Yo87, 4, 5.2] and [Kra96, 4]. We now assume that \overline{S} is a complex smooth projective *curve*. Then $Z = \{\zeta_0, \dots, \zeta_n\}$ is just a finite set of points and $S = \overline{S} \setminus Z$.

We assume that \mathcal{S} is uniformizable. By Riemann's uniformization theorem, the universal orbifold-covering $\tilde{\mathcal{S}}$ is then isomorphic to either $\mathbb{P}^1, \mathbb{A}^1$ or the open unit disk D . We fix such an isomorphism⁽²³⁾; the group $\text{Aut } \tilde{\mathcal{S}}$ is then a Lie subgroup of $PSL_2(\mathbb{C})$. The orbifold fundamental group $\pi_1^{\text{orb}}(\mathcal{S}, s)$ is a discrete subgroup of $\text{Aut } \tilde{\mathcal{S}}$ acting properly on $\tilde{\mathcal{S}}$, and the quotient space is $\tilde{\mathcal{S}}/\pi_1^{\text{orb}}(\mathcal{S}, s) = \overline{S}$.

4.3.2. We fix a non-constant rational function $z \in \mathbb{C}(\overline{S})$. We shall assume for simplicity that z has no pole on S . Having identified $\tilde{\mathcal{S}}$ with a domain of \mathbb{P}^1 allows us to consider the canonical map

$$\tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{S}} \subset \mathbb{P}^1$$

as an analytic⁽²⁴⁾ function τ on $\tilde{\mathcal{S}}$ (in loose terms, this is the multivalued inverse of the projection $\tilde{\mathcal{S}} \rightarrow \overline{S} = \tilde{\mathcal{S}}/\pi_1^{\text{orb}}(\mathcal{S}, s)$, restricted to S). Notice that the derivation $\frac{d}{dz}$ extends canonically to $\mathcal{O}(\tilde{\mathcal{S}})$.

Proposition 4.3.3. *There is a unique differential equation of the form*

$$y'' - qy = 0, \quad q \in \mathbb{C}(\overline{S}),$$

the uniformizing differential equation, such that τ is the quotient of two solutions in $\mathcal{O}(\tilde{\mathcal{S}})$ of this differential equation. It is fuchsian and has singularities only at $Z = \{\zeta_0, \dots, \zeta_n\}$. The monodromy group pointed at s is a discrete finitely generated subgroup of $SL_2(\mathbb{C})$ whose image in $PSL_2(\mathbb{C})$ is $\pi_1^{\text{orb}}(\mathcal{S}, s)$.

If $\overline{S} = \mathbb{P}^1$ and $\zeta_n = \infty$, the exponents are rational numbers given by the

⁽²³⁾in some cases, it will be more convenient to work with \mathfrak{h} instead of D .

⁽²⁴⁾in the case $\tilde{\mathcal{S}} = \mathbb{P}^1$, it is only meromorphic a priori. But in that case $\overline{S} = \mathbb{P}^1$ and the covering $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ is finite. It will be convenient to normalize things so that ∞ lies above ∞ and z is the standard coordinate on \mathbb{P}^1 . With this choice, τ has no pole.

Riemann scheme:

$$\begin{pmatrix} \zeta_0 & \cdots & \zeta_{n-1} & \zeta_n = \infty \\ \frac{1}{2}(1 + \frac{1}{e_0}) & \cdots & \frac{1}{2}(1 + \frac{1}{e_{n-1}}) & \frac{-1}{2}(1 + \frac{1}{e_n}) \\ \frac{1}{2}(1 - \frac{1}{e_0}) & \cdots & \frac{1}{2}(1 - \frac{1}{e_{n-1}}) & \frac{-1}{2}(1 - \frac{1}{e_n}) \end{pmatrix}.$$

cf. loc. cit.

PROOF. Existence: q is nothing but the Schwarz derivative

$$\{\tau, z\} = \left(\frac{d\tau}{dz}\right)^{1/2} \frac{d^2}{dz^2} \left[\left(\frac{d\tau}{dz}\right)^{-1/2} \right],$$

which is single-valued on \overline{S} . A basis of solutions of the uniformizing differential equation is given by $(\frac{d\tau}{dz})^{-1/2}$ and $\tau \cdot (\frac{d\tau}{dz})^{-1/2}$. The fact that $q \in \mathbb{C}(\overline{S})$ follows immediately from the invariance of the Schwarz derivative under homographic transformations.

Unicity: this is a purely algebraic fact: q is determined by the projective connection attached to the differential equation $y'' - qy = 0$.

The determination of the Riemann scheme comes from this expression and the fact that around $\zeta_{i < n}$ (*resp.* around ∞) τ is given by $z - \zeta_i \mapsto c_i \cdot (z - \zeta_i)^{1/e_i} + h.o.t.$ (*resp.* $1/z \mapsto c_n/z^{e_n} + h.o.t.$). \square

Remarks 4.3.4. (i) The case of three singularities will be detailed in the next section. In the case of more than three singularities, it is extremely difficult in general to write down explicitly the uniformizing differential equations.

(ii) In [Yo87], an extension of the theory of uniformizing differential equations to higher dimensions is developed. Its practical scope seems limited however to the case when \mathcal{S} is a symmetric domain.

(iii) The projective equivalence class of the uniformizing differential equation is a variant of what is called the Schwarz structure or canonical indigenous \mathbb{P}^1 -bundle over \overline{S} , *cf.* [Gun67], [Moc96].

4.4. Non-archimedean orbifolds.

4.4.1. We now work over a complete non-archimedean field K . The definition of orbifolds is parallel to the one which we have selected in the complex situation. Being mainly interested in orbifolds which are quotients of manifolds by finite groups, it is natural to replace everywhere étale (= topological) coverings in the complex case by *temperate* étale coverings in the non-archimedean context.

A K -orbifold $\mathcal{S} = (\overline{S}, (Z_i, e_i))$ consists in

- a normal K -analytic space \overline{S} (in the sense of Berkovich),
- a locally finite collection of irreducible divisors (Z_i) ,
- a positive integer e_i attached to each Z_i .

It is assumed that \overline{S} is covered by so-called *orbifold charts*, *i.e.* morphisms $W \xrightarrow{\phi} V \subset \overline{S}$ where

- W is a K -manifold,
- ϕ is a (ramified) covering in the sense of 1.2.3 (*i.e.* V is covered by open subsets U such that $\coprod V_j = \phi^{-1}U \rightarrow U$ and the restriction of ϕ to every V_j is finite),
- except for a closed subset Σ of codimension 2 in V , ϕ restricts to a temperate étale covering above $V \setminus (V \cap \bigcup Z_i)$, and is ramified with index e_i above Z_i .

Here again, we may and shall assume that Σ is contained in $\bigcup Z_i \cup \overline{S}^{\text{sing}}$ and set $Z = \bigcup Z_i \cup \overline{S}^{\text{sing}}$, $S = \overline{S} \setminus Z$ (which is a K -manifold).

Remarks 4.4.2. (i) At this point, because the notion of orbifold is local, it would not change if we had modified the definition of orbifold charts by replacing ‘temperate étale’ by ‘finite étale’ or by ‘étale’. In contrast, we insist that orbifold charts are defined in terms of temperate coverings.

(ii) Since we only prescribe the ramification indices and not the ramification groups, this notion of orbifold is only reasonable in the tame case, *i.e.* when $\text{char}(K) \nmid e_i$. We shall be mostly interested in the characteristic 0 case.

(iii) If $\dim \overline{S} = 1$, then $S = \overline{S} \setminus \bigcup Z_i$.

4.4.3. A morphism of orbifolds $f : \mathcal{S}' \rightarrow \mathcal{S}$ is a morphism $\overline{S}' \rightarrow \overline{S}$, such that \overline{S}' is covered by orbifold charts $\phi' : W' \rightarrow V'$ with the property that $f \circ \phi'$ factors through some orbifold chart $\phi : W \rightarrow V$ of \mathcal{S} : $f \circ \phi' = \phi \circ f'$.

Example. Let $(\overline{S}, (Z_i, e_i)_{i \in I})$ be an orbifold, and let $J \subset I$ be the subset of indices for which $e_i > 1$. Then the identity of \overline{S} induces an isomorphism $(\overline{S}, (Z_i, e_i)_{i \in I}) \cong (\overline{S}, (Z_i, e_i)_{i \in J})$. In other words, one can ‘forget’ the divisors with index $e_i = 1$.

Considering K -manifolds as orbifolds with an empty collection of divisors ($S = \overline{S}$), one can embed the category of K -manifolds as a full subcategory of the category of K -orbifolds.

It will be convenient to attach to any irreducible divisor $D \subset \overline{S}$ an index $e_D \in \mathbb{Z}_{>0}$ which is e_i if $D = Z_i$ and 1 if D is none of the Z_i 's.

4.4.4. A morphism f is called an *orbifold-covering* if it is a covering in the sense of 1.2.3 and for any orbifold chart $\phi' : W' \rightarrow V'$ of \mathcal{S}' , $f \circ \phi' : W' \rightarrow V = f(V')$ is an orbifold chart of \mathcal{S} (note that if \mathcal{S}' and \mathcal{S} are just K -manifolds, an orbifold-covering amounts to a temperate étale covering, by our definition of orbifold charts).

Orbifold-coverings of \mathcal{S} form a category in an obvious way, denoted by $\text{Cov}_{\mathcal{S}}$.

The restriction of f above S is then an étale covering: indeed, this is clear except above a closed subset of codimension ≥ 2 in S , thus above the whole of S by purity since S is smooth. However, we do not know in general whether it is temperate (due to the non-local nature of the definition of temperate coverings).

Example. Assume that $\text{char}(K) \nmid n$. Then $(\mathbb{A}^1, (0, e')) \xrightarrow{z \mapsto z^n} (\mathbb{A}^1, (0, e))$ is a morphism of orbifolds if and only if $e \mid ne'$, and is an orbifold-covering if and only if $e = ne'$.

Remark. Let $f : \mathcal{S}' \rightarrow \mathcal{S}$ be an orbifold-covering. As we just saw, we may assume that all the indices e'_i, e_j are > 1 . Under this assumption, $f(Z') \subset Z$ and $f^{-1}(S) \subset S'$.

4.4.5. An orbifold is said to be *uniformizable*⁽²⁵⁾ if it admits a *global* orbifold chart.

However, *universal global orbifold charts do not exist*; this is no surprise: there is no universal temperate covering of a K -manifold, except in trivial cases. For the same reason, *orbifold charts are not locally unique*.

Let $f : \mathcal{S}' \rightarrow \mathcal{S}$ be an orbifold-covering of a uniformizable orbifold \mathcal{S} . Then the *restriction of f to $f^{-1}(S)$ is a temperate étale covering* of S .

Indeed, let $W \rightarrow \overline{S}$ be a global orbifold chart. Then $W|_S \times_S f^{-1}(S) \rightarrow f'(W|_S \times_S f^{-1}(S)) = f^{-1}(S)$ is an orbifold chart for \mathcal{S}' , hence $W|_S \times_S f^{-1}(S) \rightarrow S$ is an orbifold chart for \mathcal{S} . Since its image is S itself, it is a temperate covering of S , of which $f^{-1}(S) \rightarrow S$ is a quotient.

Example. Assume that K is algebraically closed of characteristic zero. Then $(\mathbb{P}^1, (\zeta_i, e_i)_{i=0, \dots, n})$ is uniformizable if and only if $n \geq 2$ or $(n = 1$ and $e_0 = e_1)$, just as in the complex case. This actually follows from the complex case: one may assume that K has the same cardinality as \mathbb{C} , choose an abstract isomorphism $K \cong \mathbb{C}$, and use the fact that in the complex case, one can find a global algebraic orbifold chart which is a finite ramified covering of \overline{S} .

4.4.6. Let us again assume that K is of characteristic zero. Let $\mathcal{S} = (\overline{S}, (Z_i, e_i))$ be a connected K -orbifold, and let Γ be a virtually free discrete group acting faithfully and properly by automorphisms on \mathcal{S} . The *orbifold quotient* \mathcal{S}/Γ is defined as in the complex situation: the underlying analytic space is \overline{S}/Γ . The ramification divisors are the images of the Z_i 's, and the irreducible divisors D with non-trivial fixer Γ_D ; the latter are endowed with the multiplicity $e_D \cdot |\Gamma_D|$. Note that the projection $\mathcal{S} \rightarrow \mathcal{S}/\Gamma$ is an orbifold-covering.

Interesting examples with \overline{S} open in \mathbb{P}^1 are discussed in [vdP92a].

4.4.7. One of the applications of K -orbifolds would be to represent moduli problems for isomorphism classes of objects with finite automorphism groups.

Typical instances occur in the context of p -adic uniformization of Shimura varieties. Let us briefly mention how. Recall from II.7.2, 7.3 the following basic commutative diagram

⁽²⁵⁾we avoid the terminology ‘good’ in order to prevent confusion with good spaces in the sense of Berkovich, *cf.* last section.

$$\begin{array}{ccc}
\mathcal{M}^0 & \xrightarrow{\mathcal{P}} & \mathcal{D} \\
\mathcal{Q} \downarrow & & \downarrow \mathcal{Q} \\
S & \xrightarrow{\mathcal{P}} & \mathcal{D}/\Gamma
\end{array}$$

S is a tube-like domain in the p -adic analytification of a Shimura variety, the horizontal maps are étale, and Γ is an arithmetic subgroup in the semi-simple group \overline{G}^{ad} . Under the assumption that Γ is small enough, the maps \mathcal{Q} are topological covering maps.

If one drops that assumption, one has to consider S as an orbifold quotient (defined by \mathcal{Q} as in 4.4.6). This is especially relevant in the case of global uniformization, *i.e.* when S is a connected component of the Shimura variety.

The fake projective planes of Mumford, Ishida, Kato, provide interesting examples of this situation, *cf.* II.7.5 and [Kat98]. We shall see several other instances in the sequel.

4.5. Non-archimedean orbifold fundamental groups.

From now on, we assume that K is *algebraically closed of characteristic zero*.

4.5.1. Let $\mathcal{S} = (\overline{\mathcal{S}}, (Z_i, e_i))$ be a uniformizable connected orbifold.

As we have seen in 4.4.5, the restriction above S of an orbifold-covering is a temperate étale covering. This gives rise to a restriction functor

$$\text{Cov}_S \longrightarrow \text{Cov}_S^{\text{temp}}.$$

Proposition 4.5.2. *This functor is fully faithful. Its image consists of those temperate étale coverings $S' \rightarrow S$ which extend to ramified coverings $\overline{\mathcal{S}}' \rightarrow \overline{\mathcal{S}}$, with $\overline{\mathcal{S}}'$ normal, such that the ramification index at each irreducible divisor above Z_i divides e_i (for every i).*

This image is stable under taking connected components, fiber products (over S), quotients, finite disjoint unions, and Galois closure.

PROOF. From Kiehl's theorem on the existence of tubular neighborhoods and the example at the end of 4.4.4, it follows that \mathcal{S}' is determined by $\overline{\mathcal{S}}'$ and by the ramification index n_{ij} at each divisor Z_{ij} above Z_i (for every i): namely, $e'_{ij} = e_i/n_{ij}$. The first assertion then follows immediately from theorem 2.1.11. The second assertion is a straightforward consequence of this characterization of the image. \square

Let us fix a geometric base-point \overline{s} of $S = \overline{\mathcal{S}} \setminus (\bigcup Z_i \cup \overline{\mathcal{S}}^{\text{sing}})$ (in dimension one, we could also choose a tangential base-point in the sense of 2.2).

Definition 4.5.3. The orbifold fundamental group pointed at \overline{s}

$$\pi_1^{\text{orb}}(\mathcal{S}, \overline{s})$$

is the topological group attached to Cov_S viewed as a subcategory of Cov_S^{et} , *cf.* 1.4.

This is a prodiscrete group (1.4.7). By 1.4.5, one has a commutative square of functors

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{disjoint unions of} \\ \text{orbifold-coverings of } S \end{array} \right\} & \xrightarrow{\subset} & \left\{ \begin{array}{l} \text{disjoint unions of} \\ \text{temperate coverings of } S \end{array} \right\} \\ \downarrow \wr & & \downarrow \wr \\ \{\pi_1^{\text{orb}}(\mathcal{S}, s)\text{-sets}\} & \xrightarrow{\subset} & \{\pi_1^{\text{temp}}(S, s)\text{-sets}\}. \end{array}$$

Galois orbifold-coverings correspond to open normal subgroups of π_1^{orb} .

Example. For $r \in |K^\times|$, we set $D_r = D(0, r^+)$, $D_r^* = D_r \setminus \{0\}$, and consider the orbifold $\mathcal{S}_e = (D, (0, e))$ (multiplicity e at the origin). Then by the same calculation as in 2.3.7, one finds

$$\pi_1^{\text{orb}}(\mathcal{S}_e) = \pi_1^{\text{alg}}(D) \times \mathbb{Z}/e\mathbb{Z}$$

(we drop the base point in the notation since the group is abelian, *cf.* 1.4.4).

This can also be expressed in the following way: let γ be a topological generator of the $\widehat{\mathbb{Z}}(1)$ -factor in $\pi_1^{\text{temp}}(D^*)$ (the factor corresponding to Kummer coverings). Then

$$\pi_1^{\text{orb}}(\mathcal{S}_e) = \pi_1^{\text{temp}}(D^*, \bar{s}) / \langle \gamma^e \rangle^-,$$

where $\langle \gamma^e \rangle^-$ denotes the closure of the (normal) subgroup generated by γ .

4.5.4. By Kiehl's theorem, \bar{S} is locally isomorphic to a product $D^* \times Z_i^{\text{smooth}}$ in the neighborhood of the non-singular part of Z_i ; whence a homomorphism $\pi_1^{\text{temp}}(D^*) \rightarrow \pi_1^{\text{temp}}(S, \bar{s})$, well-defined up to conjugation. Let us denote by γ_i the image in $\pi_1^{\text{temp}}(S, \bar{s})$ of a topological generator of the $\widehat{\mathbb{Z}}(1)$ -factor of $\pi_1^{\text{temp}}(D^*)$ ("local monodromy" at Z_i). The closure $\langle \gamma_i^{e_i} \rangle^-$ of the normal subgroup of $\pi_1^{\text{temp}}(S, \bar{s})$ generated by the $\gamma_i^{e_i}$'s is a well-defined closed subgroup, independent of the choice of the γ_i 's.

Theorem 4.5.5. (a) *One has $\pi_1^{\text{temp}}(S, \bar{s}) \cong \varprojlim_{(e_i)} \pi_1^{\text{orb}}((\bar{S}, (Z_i, e_i)), \bar{s})$, where the limit is taken according to the ordering by divisibility.*

(b) *The homomorphism $\pi_1^{\text{temp}}(S, \bar{s}) \rightarrow \pi_1^{\text{orb}}(S, \bar{s})$ induces an injective strict homomorphism*

$$\pi_1^{\text{temp}}(S, \bar{s}) / \langle \gamma_i^{e_i} \rangle^- \rightarrow \pi_1^{\text{orb}}(S, \bar{s})$$

with dense image. This is an isomorphism if S is the analytification of a smooth algebraic K -variety S^{alg} ; in particular $\pi_1^{\text{temp}}(S, \bar{s}) / \langle \gamma_i \rangle^- \cong \pi_1^{\text{temp}}(\bar{S}, \bar{s})$.

PROOF. (a) reflects the fact that $\text{Cov}_S^{\text{temp}} = \varprojlim_{(\bar{S}, (Z_i, e_i), \bar{s})} \text{Cov}_{(\bar{S}, (Z_i, e_i), \bar{s})}$ which follows from the previous proposition.

(b) The previous proposition shows more precisely that a $\pi_1^{\text{temp}}(S, \bar{s})$ -set comes from a $\pi_1^{\text{orb}}(S, \bar{s})$ -set if and only if the $\gamma_i^{e_i}$'s act trivially.

We shall use lemma 1.4.9. By construction of the topology, open subgroups of $\pi_1^{\text{temp}}(S, \bar{s})$ and $\pi_1^{\text{orb}}(S, \bar{s})$ respectively form a fundamental system of neighborhoods of 1. In fact, this is even true for normal open subgroups (existence of Galois closures). It follows that $\pi_1^{\text{temp}}(S, \bar{s})/\langle \gamma_i^{e_i} \rangle^-$ has the same property. The first part of b) then follows from 1.4.9. (ii). The second part follows from 1.4.9. (iii) and (iv), since we have shown that the quotient of $\pi_1^{\text{temp}}(S, \bar{s})$ by any closed normal subgroup (*e.g.* $\langle \gamma_i^{e_i} \rangle^-$) is complete when S is algebraic (2.1.7). \square

Remarks 4.5.6. (a) 2.3.9 is a special case of the last assertion of 4.5.5.

(b) When S is algebraic, the profinite completion of $\pi_1^{\text{orb}}(S, \bar{s})$ admits an algebraic description.

(c) The situation is analogous to the classical one (over \mathbb{C} , *cf.* 4.2), with at least two noteworthy differences:

- since there is no universal orbifold-covering, the geometrically meaningful object is not π_1^{orb} itself, but its discrete quotients,
- the non-archimedean π_1^{orb} is a finer invariant which takes into account the relative “position” of the Z_i . For instance, if $\bar{S} = \mathbb{P}_{\mathbb{C}_p}^1$, the classical π_1^{orb} depends only on the numbers e_i , not on the branch points Z_i themselves. In the p -adic case, this is not the case. For instance, if $Z = \{0, 1, \lambda, \infty\}$ and every $e_i = 2$, then the Legendre elliptic curve with parameter λ provides a global orbifold chart, and it follows from 2.3.2 that the orbifold fundamental group is either
 - a (compact) semi-product of $\mathbb{Z}/2\mathbb{Z}$ by $\widehat{\mathbb{Z}} \times \widehat{\mathbb{Z}}$ if $|\lambda(1-\lambda)| = 1$,
 - a (non-compact) semi-product of $\mathbb{Z}/2\mathbb{Z}$ by $\mathbb{Z} \times \widehat{\mathbb{Z}}$ if $|\lambda(1-\lambda)| \neq 1$.

4.5.7. Let $\phi : \bar{S}' \rightarrow \bar{S}$ be a global orbifold chart, such that the induced map $S' \rightarrow S$ is a finite Galois étale covering with group G . Let \bar{s}' be a geometric point of S' above \bar{s} .

Proposition 4.5.8. *The universal topological covering $\widetilde{\bar{S}'}$ of \bar{S}' is a Galois orbifold-covering of S . Let $\Gamma = \text{Aut}(\widetilde{\bar{S}'}/S)$ denote the corresponding (discrete) Galois group. Then one has a commutative diagram of strict homomorphisms of topological groups, in which the rows are exact sequences*

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \pi_1^{\text{temp}}(\bar{S}', \bar{s}') & \longrightarrow & \pi_1^{\text{orb}}(S, \bar{s}) & \longrightarrow & G \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & \pi_1^{\text{top}}(\widetilde{\bar{S}'}, \bar{s}') & \longrightarrow & \Gamma & \longrightarrow & G \longrightarrow 1.
 \end{array}$$

PROOF. Let us denote by \mathcal{S}' the orbifold \bar{S}' with all $\phi^{-1}(Z_i)$ endowed with multiplicity $e_i = 1$. The pull-back by ϕ provides the vertical arrows of a

commutative diagram of functors

$$\begin{array}{ccc} \mathrm{Cov}_{\mathcal{S}} & \xrightarrow{\subset} & \mathrm{Cov}_{\mathcal{S}}^{\mathrm{temp}} \\ \downarrow & & \downarrow \\ \mathrm{Cov}_{\mathcal{S}'} = \mathrm{Cov}_{\tilde{\mathcal{S}}'}^{\mathrm{temp}} & \xrightarrow{\subset} & \mathrm{Cov}_{\mathcal{S}'}^{\mathrm{temp}}. \end{array}$$

One concludes by 1.4.12 that there is an exact sequence

$$1 \rightarrow \pi_1^{\mathrm{temp}}(\tilde{\mathcal{S}}', \tilde{s}') \rightarrow \pi_1^{\mathrm{orb}}(\mathcal{S}, \bar{s}) \rightarrow G \rightarrow 1.$$

On the other hand, it follows from 1.1.4 that the restriction of $\tilde{\mathcal{S}}'$ above \mathcal{S}' is nothing but the universal topological covering $\tilde{\mathcal{S}}'$ of \mathcal{S}' . By 2.1.2, $\tilde{\mathcal{S}}' \rightarrow \mathcal{S}'$ is a Galois temperate covering, and it follows from that and 4.5.2 that $\tilde{\mathcal{S}}' \rightarrow \mathcal{S}$ is a Galois orbifold-covering. It is then clear that the corresponding epimorphism $\pi_1^{\mathrm{orb}}(\mathcal{S}, \bar{s}) \rightarrow \Gamma$ gives rise to the above commutative diagram. \square

4.5.9. Assume that \mathcal{S} is the analytification of a smooth algebraic K -variety S^{alg} . According to 3.5.12, discrete representations of $\pi_1^{\mathrm{temp}}(\mathcal{S}, \bar{s})$ give rise to algebraic vector bundles with regular integrable connection (M, ∇) on S^{alg} .

Proposition 4.5.10. *An algebraic vector bundle with regular integrable connection (M, ∇) on S^{alg} comes from a discrete representation ρ of $\pi_1^{\mathrm{orb}}(\mathcal{S}, \bar{s})$ if and only if there is a finite global orbifold chart $\phi : \tilde{\mathcal{S}}' \rightarrow \tilde{\mathcal{S}}$ such that $(\phi^*(M))^{\nabla}$ extends to a locally constant sheaf on $\tilde{\mathcal{S}}'$.*

This follows from 3.5.6 and 4.5.5.

Remark. If \mathcal{S}'/\mathcal{S} is Galois, one can restrict ρ to the open normal subgroup $\pi_1^{\mathrm{temp}}(\tilde{\mathcal{S}}', s')$ of $\pi_1^{\mathrm{orb}}(\mathcal{S}, \bar{s})$ (3.5.7), and this restriction factors through $\pi_1^{\mathrm{top}}(\tilde{\mathcal{S}}', s')$.

4.6. p -adic uniformizing differential equations.

4.6.1. We now develop the non-archimedean counterpart of 4.3. We assume that $\tilde{\mathcal{S}}$ is a smooth projective curve over K . Then $Z = \{\zeta_0, \dots, \zeta_n\}$ is just a finite set of points and $\mathcal{S} = \tilde{\mathcal{S}} \setminus Z$.

We assume moreover that \mathcal{S} is uniformizable. To imitate the complex theory, we shall assume that *there exists a finite Galois global orbifold chart $\phi : \tilde{\mathcal{S}}' \rightarrow \tilde{\mathcal{S}}$ such that $\tilde{\mathcal{S}}'$ is a Mumford curve* (of any genus ≥ 0).

The universal topological covering $\tilde{\tilde{\mathcal{S}}}'$ of $\tilde{\mathcal{S}}'$ will replace the universal orbifold-covering, which does not exist in the non-archimedean situation. By assumption, there is an embedding

$$\tilde{\tilde{\mathcal{S}}}' \subset \mathbb{P}^1,$$

which we fix, so that the group $\mathrm{Aut} \tilde{\tilde{\mathcal{S}}}'$ becomes a Lie subgroup of $PGL_2(K)$.

The orbifold-covering $\tilde{\mathcal{S}}' \rightarrow \bar{\mathcal{S}}$ gives rise to a projective discrete representation

$$\bar{\rho}: \pi_1^{\text{orb}}(\mathcal{S}, s) \rightarrow PGL_2(K).$$

Its image Γ is contained in the normalizer in $PGL_2(K)$ of the Schottky group $\pi_1^{\text{top}}(\bar{\mathcal{S}}', s')$, hence discrete; it acts properly on $\bar{\mathcal{S}}'$, with quotient space $\bar{\mathcal{S}}'/\Gamma = \bar{\mathcal{S}}$.

4.6.2. We fix a non-constant rational function $z \in K(\bar{\mathcal{S}})$, and assume for simplicity that z has no pole on S .

Having identified $\tilde{\mathcal{S}}'$ with a domain of \mathbb{P}^1 allows us to consider $z: \tau \mapsto z(\tau)$ as a meromorphic function defined on that domain (and analytic on $\tilde{S}' \subset \tilde{\mathcal{S}}'$). Notice that the derivation $\frac{d}{dz}$ extends canonically to $\mathcal{O}(\tilde{\mathcal{S}}')$.

Proposition 4.6.3. *There is a unique differential equation of the form*

$$y'' - qy = 0, \quad q \in K(\bar{\mathcal{S}}),$$

the uniformizing differential equation, such that a local inverse $\tau(z)$ of the map $z(\tau)$ is given by the quotient of two solutions in $\mathcal{O}(\tilde{S}')$ of this differential equation. It is fuchsian and has singularities only at $Z = \{\zeta_0, \dots, \zeta_n\}$.

If $\bar{\mathcal{S}} = \mathbb{P}^1$ and $\zeta_n = \infty$, the exponents are rational numbers given by the Riemann scheme:

$$\left(\begin{array}{cccc} \zeta_0 & \dots & \zeta_{n-1} & \zeta_n = \infty \\ \frac{1}{2}(1 + \frac{1}{e_0}) & \dots & \frac{1}{2}(1 + \frac{1}{e_{n-1}}) & \frac{-1}{2}(1 + \frac{1}{e_n}) \\ \frac{1}{2}(1 - \frac{1}{e_0}) & \dots & \frac{1}{2}(1 - \frac{1}{e_{n-1}}) & \frac{-1}{2}(1 - \frac{1}{e_n}) \end{array} \right).$$

The proof is the same as in the complex case: q is nothing but the Schwarz derivative $\{\tau, z\} = (\frac{d\tau}{dz})^{1/2} \frac{d^2}{dz^2} [(\frac{d\tau}{dz})^{-1/2}]$, which is single-valued on $\bar{\mathcal{S}}$. A basis of solutions of the uniformizing differential equation is given by $(\frac{d\tau}{dz})^{-1/2}$ and $\tau \cdot (\frac{d\tau}{dz})^{-1/2}$.

Proposition 4.6.4. *The projective connection attached to the uniformizing differential corresponds to the projective discrete representation $\bar{\rho}$ in the sense of 3.5.13: it is in the image of the temperate étale Riemann-Hilbert functor and its projective monodromy group is the discrete group Γ .*

PROOF. A basis of solutions of the symmetric square of the uniformizing differential equation is given by $\frac{1}{d\tau/dz}, \frac{\tau}{d\tau/dz}, \frac{\tau^2}{d\tau/dz}$. These are elements of $\mathcal{O}(\tilde{\mathcal{S}}')$ (in fact multivalued meromorphic functions on $\bar{\mathcal{S}}'$). Criterion 4.5.10 then shows that this symmetric square comes from a discrete representation of $\pi_1^{\text{orb}}(\mathcal{S}, s)$. It is clear that this representation is nothing but the composition of $\bar{\rho}$ with the standard homomorphism $PGL_2 \rightarrow GL_3$. \square

[One could also argue by using an analogue of 4.5.10 for projective connections.]

Remark. The connection itself attached to the uniformizing differential equation corresponds to a discrete representation of $\pi^{\text{temp}}(S, \bar{s})$ (cf. 3.5.14), which does not necessarily factor through $\pi_1^{\text{orb}}(\mathcal{S}, s)$ [it factors through an orbifold fundamental group, up to replacing the e_i 's by appropriate multiples].

4.7. p -adic Shimura orbifolds and uniformizing differential equations.

4.7.1. We are now ready for a synthesis, relating the theory of p -adic period mappings surveyed in II with the theory of p -adic orbifolds.

Let us consider a maximal order \mathcal{B} in an indefinite quaternion algebra B over \mathbb{Q} , endowed with a positive involution $*$. “Fake elliptic curves” are $*$ -polarized abelian surfaces with multiplication by \mathcal{B} . They are classified by a smooth projective Shimura curve \mathcal{X}^+ defined over \mathbb{Q} , cf. II.7.4.2, which should actually be considered as an orbifold $(\mathcal{X}^+, (\zeta_i, e_i))$, because the action of Γ^+ on the Poincaré upper half-plane \mathfrak{h} has fixed points (the ζ_i 's are their images in \mathcal{X}^+).

The Gauss-Manin connection $(\mathcal{H}, \nabla_{\text{GM}})$ attached to the “universal” fake elliptic curve (in the sense of orbifolds) is a connection with logarithmic poles at the ζ_i 's. It has been analyzed in II.7.6. The choice of a quadratic splitting field $\mathbb{Q}(\sqrt{-d})/\mathbb{Q}$ for B decomposes $(\mathcal{H}, \nabla_{\text{GM}})$ into two pieces of rank two (the “+ and – eigenspaces” for the action of $\sqrt{-d}$). It follows from the theory of the complex period mapping that a multivalued section $\tau : \mathcal{X}^+(\mathbb{C}) \setminus \{\zeta_i\} \rightarrow \mathfrak{h}$ of the projection $\mathfrak{h} \rightarrow \mathcal{X}^+(\mathbb{C})$ is given by a quotient of two non-proportional solutions of $\nabla_{\text{GM},+}$. One derives that *the uniformizing differential equation for $(\mathcal{X}^+, (\zeta_i, e_i))$ is projectively equivalent to $\nabla_{\text{GM},+}$.*

On the other hand, there is a canonical $\overline{\mathbb{Q}}$ -structure in the \mathbb{C} -space of solutions at a given point s (the Betti $H^1(-/\overline{\mathbb{Q}})$ of the fake elliptic curve with parameter s), which is stable under (global) monodromy.

4.7.2. According to the Čerednik-Drinfeld theory surveyed in II.7.4, 7.6 (and the description of the p -adic period mapping given there as a quotient of solutions of the Gauss-Manin connection), there is an entirely parallel p -adic description of this geometric situation for any critical prime p , i.e. any p which divides the discriminant of B : $\mathcal{X}^+(\mathbb{C}_p)$ is a p -adic orbifold of the kind considered in 4.6, and *the p -adic uniformizing differential equation for $(\mathcal{X}^+, (\zeta_i, e_i))$ is projectively equivalent to $\nabla_{\text{GM},+}$, viewed as a p -adic connection (cf. II.7.2.6, II.7.2.7).*

By the theory of p -adic Betti lattices (cf. II.7.6.2), there is a canonical $\overline{\mathbb{Q}}$ -structure in the \mathbb{C}_p -space of solutions at a given point s , which is stable under (global) monodromy.

4.7.3. Following II.7.5, all this generalizes to Shimura curves defined by quaternion algebras B over totally real number fields E , and a “congruence

subgroup" Γ (here, we drop the general assumption of II.7 that Γ is small enough). We recall that these Shimura curves are not modular if $E \neq \mathbb{Q}$, but are twisted forms of modular Shimura curves defined by quaternion algebras $B^\bullet = B \otimes_E K$ over suitable totally imaginary quadratic extensions K/E . One can thus still attach a Gauss-Manin connection to such a situation, and this connection is related to the uniformizing differential equations both in the complex and in the v -adic sense⁽²⁶⁾, for *critical places* v of E , cf. II.7.5, II.7.6 (we fix embeddings $\mathbb{C} \leftarrow \overline{\mathbb{Q}} \rightarrow \mathbb{C}_p$ for every critical $v \mid p$, compatible with the privileged places ∞_0 and v of E , and denote by \overline{B} the quaternion algebra over E obtained from B by interchanging the local invariants at ∞_0 and v). We summarize these results as follows, using the terminology of orbifolds:

Theorem 4.7.4. *The Shimura curve \mathcal{X}_Γ is defined over $\overline{\mathbb{Q}}$, and may be considered, in a canonical way, both as a complex orbifold and as a p -adic orbifold (for any critical $v \mid p$).*

The complex uniformizing differential equation is defined over $\overline{\mathbb{Q}}$ and coincides with the p -adic uniformizing differential equation. It is projectively equivalent to a direct summand of the Gauss-Manin connection: the period mapping (resp. the p -adic period mapping) is given by a suitable quotient τ (resp. τ_p) of two non-proportional solutions. The projective monodromy group of the uniformizing differential equation over \mathbb{C} (resp. over \mathbb{C}_p) is Γ (resp. is a discrete arithmetic group Γ_p in $PGL_2(\mathbb{C}_p)$ attached to the quaternion algebra \overline{B}).

Moreover, there is a canonical $\overline{\mathbb{Q}}$ -structure in the space of solutions of the complex (resp. p -adic) uniformizing differential equation at any $\overline{\mathbb{Q}}$ -point, which is stable under global monodromy.

Let us recall that the p -adic projective monodromy group, based at a geometric point \bar{s} , can be understood in a sense close to the usual intuitive one, cf. 3.5.6. In loose words: there is a smooth projective curve S and a finite orbifold-covering $S \rightarrow \mathcal{X}_\Gamma$ such that the projective monodromy is given by the values taken by τ_p when one passes from a point s' lying above s to another s'' by all possible paths on the Berkovich space $S_{\mathbb{C}_p}^{\text{an}}$.

4.7.5. Let us illustrate this section with a concrete example (cf. [And98] for more details). We first define a quadrangle $ABCD$ with circular edges in \mathfrak{h} , as follows

L is the half line of slope $-1/2$ through the origin,

Σ (resp. Σ') is the unit half-circle centered at the origin (resp. at $2 - \sqrt{3}$),

D is the point $L \cap \Sigma'$,

⁽²⁶⁾if p divides the level of the congruence group, the orbifold is not quite of the kind considered in 4.6: the top space is not the set of ordinary points for a Schottky group, but a finite étale covering of such a space, cf. II.7.4.7; however, the construction of the uniformizing differential equation can still be done in the same way, due to the fact that the finite étale coverings of the Drinfeld space which occur are $PGL_2(E_v)$ -equivariant.




FIGURE 3

Σ'' is the half-circle centered on the horizontal axis, orthogonal to Σ and passing through D ,

B (resp. C) is the intersection point of the vertical axis with Σ (resp. Σ'),

A is the intersection point of Σ with Σ' (specifically, $A = -\frac{1+2i}{\sqrt{5}}$).

The quadrangle $ABCD$ is a hyperbolic quadrangle with angles $(\frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{2}, \frac{\pi}{6})$. The symmetries around the edges of this polygon generate a group of Möbius transformations which contains a subgroup Γ^* of index 2 of conformal transformations. This fuchsian group has already attracted the attention of many a mathematician (J.F. Michon, M.F. Vigneras [Vig80, p. 123], F. Beukers, D. Krammer [Kra96], N. Elkies [Elk98], D. Kohel...). It is $PSL_2(\mathbb{R})$ -conjugated to the discrete group generated by the following unimodular matrices (up to sign)

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -\frac{\sqrt{5}+1}{2} \\ \frac{\sqrt{5}-1}{2} & 0 \end{pmatrix}, \begin{pmatrix} -\sqrt{3} & 1+\sqrt{5} \\ 1-\sqrt{5} & \sqrt{3} \end{pmatrix}, \begin{pmatrix} \frac{\sqrt{3}+\sqrt{15}}{2} & -2 \\ 2 & \frac{\sqrt{3}-\sqrt{15}}{2} \end{pmatrix}$$

of respective orders 2, 2, 2, 6, whose product in this order is 1. The orbifold quotient of \mathfrak{h} by Γ^* has been determined by Krammer — and, later but independently, by Elkies: it is

$$(\mathbb{P}^1; (0, e_0 = 2), (1, e_1 = 2), (81, e_{81} = 2), (\infty, e_\infty = 6)),$$

the points A, B, C, D being mapped to $0, 1, 81, \infty$ respectively.

Moreover, these authors have computed the uniformizing equation⁽²⁷⁾. It is actually easier to write down the differential equation obtained from the uniformizing equation by multiplying all solutions by $P^{1/4}$, where $P = z(z-1)(z-81)$. It is

$$(*) \quad 18P.y'' + 9P'y' + (z-9).y = 0.$$

4.7.6. In general, it is quite hard to compute uniformizing differential equations (due to the presence of accessory parameters, when there are more than three singularities). The method of Krammer and Elkies takes advantage of the fact that the orbifold under consideration is actually a Shimura orbifold (the method, which seems to go back to H.P.F. Swinnerton-Dyer, exploits the Hecke correspondences).

In fact, Γ^* coincides with the group denoted by the same symbol in II.7.4.2, namely $\{g \in B^\times \mid g\mathcal{B} = \mathcal{B}g, \text{Nr}(g) > 0\}$, for the quaternion algebra B of discriminant 3.5 over \mathbb{Q} [Vig80, IV.3.B,C] (cf. [And98, 8] for more details on the coincidence of Γ with the group denoted by W^+ in [Kra96], up to conjugation), and is the orbifold fundamental group of the Shimura orbifold \mathcal{X}^* :

$$\mathcal{X}^*(\mathbb{C}) = \mathfrak{h}/\Gamma^*.$$

4.7.7. Let us now consider the critical prime 5 (similar phenomena occur for $p = 3$). Note that there is a ‘‘confluence’’ between the singularities 1 and 81 in characteristic 5. Let \overline{B} be the (definite) quaternion algebra with discriminant 3, and let $\overline{\mathcal{B}}$ be a maximal order ($\cong \mathbb{Z}[1, i, \frac{i+j}{2}, \frac{1+ij}{2}]$ with $i^2 = -1, j^2 = -3, ij = -ji$). Let Γ_5^* be the image in $PGL_2(\mathbb{Q}_5)$ of the normalizer $\{g \in \overline{B}^\times \mid g\overline{\mathcal{B}}[\frac{1}{5}] = \overline{\mathcal{B}}[\frac{1}{5}]g\}$. This is a discrete subgroup, and one has the Čerednik uniformization

$$\mathcal{X}^*(\mathbb{C}_p) = \Omega_{\mathbb{C}_5}/\Gamma_5^*$$

by the Drinfeld space (II.7.4.3), which represents \mathcal{X}^* as a p -adic orbifold.

It follows from what precedes that viewed as a 5-adic differential equation, (*) is obtained from the p -adic differential uniformizing equation attached to \mathcal{X}^* by multiplying all solutions by $(z(z-1)(z-81))^{1/4}$. In particular, it has projective global monodromy group Γ_5^* . This group is generated by the *conjugacy classes* of local monodromies at $0, 1, 81$, as follows from 2.3.9⁽²⁸⁾.

⁽²⁷⁾it was also found, earlier, by D. and G. Chudnovsky [CC89, p.193], who did not give the proof.

⁽²⁸⁾however, it is not clear that it is generated by three such elements, as is stated erroneously in loc. cit.

5. p -adic triangle groups.

ABSTRACT: In loose words, p -adic triangle groups are projective monodromy groups of those hypergeometric differential equations which are in the image of the étale Riemann-Hilbert functor. We give a purely geometric description of them, and then construct all “arithmetic” p -adic triangle groups using the Čerednik-Drinfeld uniformization of Shimura curves.

5.1. Review of Schwarz triangle groups.

5.1.1. A *Schwarz orbifold* is a complex orbifold of the form

$$\mathcal{S}(e_0, e_1, e_\infty) = (\mathbb{P}^1; (0, e_0), (1, e_1), (\infty, e_\infty)).$$

It is uniformizable. The universal orbifold-covering is $\mathbb{P}^1, \mathbb{A}^1$ or the open unit disk \mathbb{D} according to whether $\sum \frac{1}{e_i}$ is > 1 , $= 1$, or < 1 respectively.

We fix a base-point \bar{s} (which can be a tangential base-point).

The *Schwarz triangle group*⁽²⁹⁾ $\Delta(e_0, e_1, e_\infty)$ may be defined as $\pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty), \bar{s})$. As an abstract group, it has generators $\gamma_0, \gamma_1, \gamma_\infty$ and relations $\gamma_0^{e_0} = \gamma_1^{e_1} = \gamma_\infty^{e_\infty} = \Pi\gamma_i = 1$. A permutation of e_0, e_1, e_∞ does not change the group, only the orbifold by a homography interchanging 0, 1 and ∞ .

5.1.2. The uniformizing differential equation is fuchsian with singularities at 0, 1, ∞ , and Riemann scheme

$$\left(\begin{array}{ccc} 0 & 1 & \infty \\ \frac{1}{2}(1 + \frac{1}{e_0}) & \frac{1}{2}(1 + \frac{1}{e_1}) & \frac{-1}{2}(1 + \frac{1}{e_\infty}) \\ \frac{1}{2}(1 - \frac{1}{e_0}) & \frac{1}{2}(1 - \frac{1}{e_1}) & \frac{-1}{2}(1 - \frac{1}{e_\infty}) \end{array} \right).$$

Multiplying solutions of this equation by

$$z^{\frac{1}{2}(-1 + \frac{1}{e_0})} (z - 1)^{\frac{1}{2}(-1 + \frac{1}{e_1})}$$

transforms this equation into a (projectively equivalent) differential equation of hypergeometric type

$$HGDE(a, b, c) \quad z(z - 1)y'' + (c - (a + b + 1)z)y' - aby = 0,$$

with $1 - c = \frac{1}{e_0}$, $c - a - b = \frac{1}{e_1}$, $\pm(a - b) = \frac{1}{e_\infty}$, whose Riemann scheme is

$$\left(\begin{array}{ccc} 0 & 1 & \infty \\ 0 & 0 & \frac{1}{2}(1 - \frac{1}{e_0} - \frac{1}{e_1} + \frac{1}{e_\infty}) \\ \frac{1}{e_0} & \frac{1}{e_1} & \frac{1}{2}(1 - \frac{1}{e_0} - \frac{1}{e_1} - \frac{1}{e_\infty}) \end{array} \right).$$

The triangle group $\Delta(e_0, e_1, e_\infty)$ may thus be interpreted as the *projective monodromy group pointed at \bar{s} of this hypergeometric differential equation*.

⁽²⁹⁾in conformity with the previous chapter, we do not allow the e_i 's to be infinite. Thus we restrict our attention to the so-called cocompact triangle groups.

5.1.3. The quotient τ of a suitable basis of solutions of $HGDE(a, b, c)$ sends the upper half-plane bijectively onto a curvilinear triangle \mathcal{T} with vertices $\tau(0), \tau(1), \tau(\infty)$ and respective angles $\pi/e_0, \pi/e_1, \pi/e_\infty$. By the Schwarz reflection principle, the map τ can be extended to the lower half-plane through each of the “segments” $]0, 1[,]1, \infty[,]\infty, 0[$ by conformal reflections s_∞, s_0, s_1 respectively. The products $\gamma_0 = s_1 s_\infty, \gamma_1 = s_\infty s_0, \gamma_\infty = s_0 s_1$ give the standard generators of $\Delta(e_0, e_1, e_\infty)$. By repeating these reflections, the occurring curvilinear triangles do not overlap (except on their edges) and provide a tiling of the universal orbifold-covering ($\mathbb{P}^1, \mathbb{A}^1$ or \mathbb{D}).

5.1.4. A similar situation arises for any hypergeometric differential equation with rational parameters (a, b, c) . However, if some of the differences of the exponents $1 - c, c - a - b, a - b$ is not the inverse of an integer, the triangles which occur by iterating Schwarz reflections may overlap. Moreover, if the local projective monodromies are of respective order e_0, e_1, e_∞ at $0, 1, \infty$, the global projective monodromy group need not be $\Delta(e_0, e_1, e_\infty)$. This is related to the fact that a homomorphic image of $\Delta(e_0, e_1, e_\infty)$ which preserves the order of $\gamma_0, \gamma_1, \gamma_\infty$ need not be isomorphic to $\Delta(e_0, e_1, e_\infty)$ (cf. [Gre81] for a detailed discussion of homomorphic images of triangle groups⁽³⁰⁾).

5.1.5. The case $\sum \frac{1}{e_i} > 1$ (universal covering \mathbb{P}^1) corresponds to the Schwarz list of $HGDE$ with finite (projective) monodromy groups: the list of triples (e_0, e_1, e_∞) is, up to permutation:

- (2, 2, n) corresponding to a dihedral group,
- (2, 3, 3) corresponding to the tetrahedral group,
- (2, 3, 4) corresponding to the octahedral group,
- (2, 3, 5) corresponding to the icosahedral group.

In the case $\sum \frac{1}{e_i} = 1$ (universal covering \mathbb{A}^1), there is a global orbifold chart given by an elliptic curve with complex multiplication. There are three possibilities for (e_0, e_1, e_∞) , up to permutation: (2, 3, 6), (2, 4, 4), (3, 3, 3).

All the other triples belong to the “hyperbolic case”: $\Delta(e_0, e_1, e_\infty)$ is a fuchsian group.

For more detail on these topics, we refer to the good treatises [Car61], [Mag74], [Yo97].

5.1.6. As was remarked by J. Wolfart, Belyi’s criterion for algebraic curves to be defined over $\overline{\mathbb{Q}}$ can be expressed in terms of Schwarz orbifolds as follows:

a complex projective curve \overline{S} is defined over $\overline{\mathbb{Q}}$ if and only if it is the base of an orbifold-covering $(\overline{S}, (Z_i, e_i))$ of some Schwarz orbifold.

⁽³⁰⁾we are indebted to M. Kapovich for this reference.

5.2. p -adic Schwarz orbifolds and triangle groups.

5.2.1. A p -adic Schwarz orbifold is an orbifold over some complete subfield K of \mathbb{C}_p of the form

$$\mathcal{S}(e_0, e_1, e_\infty) = (\mathbb{P}^1, (0, e_0), (1, e_1), (\infty, e_\infty)).$$

Up to replacing K by a finite extension, it is uniformizable (cf. example in 4.4.5). However, there is no universal orbifold-covering (4.4.5).

We want our p -adic triangle groups to be *discrete* groups of fractional linear p -adic transformations. Thus it is not possible in general to define them as orbifold fundamental groups, but only as appropriate discrete quotients. This leads us to imitate the alternative definition 5.1.2 of triangle groups as projective hypergeometric monodromy groups.

5.2.2. Let e_0, e_1, e_∞ be integers > 1 , and define

$$a = \frac{1}{2} \left(1 - \frac{1}{e_0} - \frac{1}{e_1} + \frac{1}{e_\infty} \right), \quad b = \frac{1}{2} \left(1 - \frac{1}{e_0} - \frac{1}{e_1} - \frac{1}{e_\infty} \right), \quad c = 1 - \frac{1}{e_0}.$$

We now consider the hypergeometric differential equation $HGDE(a, b, c)$ as a p -adic differential equation.

Definition 5.2.3. Assume that $HGDE(a, b, c)$ is in the image of the étale Riemann-Hilbert functor, *i.e.* the étale sheaf of germs of solutions is locally constant on $\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\}$ (cf. 3.4.6). If the corresponding projective monodromy group

$$\text{Im}[\pi_1^{\text{ét}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \bar{s}) \rightarrow PGL_2(\mathbb{C}_p)]$$

is discrete and finitely generated, it is then called a p -adic triangle group, and denoted by $\Delta_p(e_0, e_1, e_\infty)$.

In the sequel, we say that “ $\Delta_p(e_0, e_1, e_\infty)$ exists” to express that the assumptions of 5.2.3 are fulfilled (this depends on the four numbers p, e_0, e_1, e_∞).

Proposition 5.2.4. (a) *If $\Delta_p(e_0, e_1, e_\infty)$ exists, there exists a finite Galois covering $\phi : \bar{S}' \rightarrow \mathbb{P}^1$ ramified above $\{0, 1, \infty\}$, such that the pull-back of $HGDE(a, b, c)$ by ϕ has a full set of multivalued analytic solutions over $S' = \phi^{-1}(\mathbb{P}^1 \setminus \{0, 1, \infty\})$.*

(b) *If $\Delta_p(e_0, e_1, e_\infty)$ exists, the homomorphism*

$$\text{Im}[\pi_1^{\text{ét}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \bar{s}) \rightarrow PGL_2(\mathbb{C}_p)]$$

factors through a surjective homomorphism

$$\pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty)) \rightarrow \Delta_p(e_0, e_1, e_\infty).$$

(c) *Conversely, if the projective connection attached to $HGDE(a, b, c)$ comes from a projective representation*

$$\pi_1^{\text{orb}}(\mathcal{S}) \rightarrow PGL_2(\mathbb{C}_p)$$

with discrete image, this image is a p -adic triangle group $\Delta_p(e_0, e_1, e_\infty)$.

- (d) Every p -adic triangle group $\Delta_p(e_0, e_1, e_\infty)$ is a discontinuous subgroup of $PGL_2(K)$ for some finite extension K of \mathbb{Q}_p .

PROOF. (a) Because the Wronskian of $HGDE(a, b, c)$ is algebraic, the determinant of the representation $\rho : \pi_1^{\text{et}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\}, \bar{s}) \rightarrow GL_2(\mathbb{C}_p)$ factors through a finite representation of $\pi_1^{\text{temp}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\}, \bar{s})$. Under the assumptions of 5.2.3, $\text{Im } \bar{\rho}$ is finitely generated, and it follows that $\text{Im } \rho$ is finitely generated. By 3.5.10, one derives that ρ factors through $\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \bar{s})$. This implies a), cf. 3.5.6 (where a concrete description of the monodromy is also given).

(b) We have just seen that ρ factors through $\pi_1^{\text{temp}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\}, \bar{s})$. To check that the associated projective representation factors through $\pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty))$ is a matter of projective local monodromy — or more plainly, of difference of exponents — at $0, 1, \infty$ (cf. 4.5); this is clear.

(c) Again because the Wronskian of $HGDE(a, b, c)$ is algebraic, the fact that the projective connection attached to $HGDE(a, b, c)$ comes from a projective representation of $\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \bar{s})$ implies that $HGDE(a, b, c)$ itself comes from a representation of $\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}, \bar{s})$ (cf. the discussion in 3.5.13). By 3.5.10, the image is finitely generated, and the assertion follows.

(d) If $\Delta_p(e_0, e_1, e_\infty)$ exists, there exists a finite Galois global orbifold chart $\bar{S}' \rightarrow \mathbb{P}^1$ such that pull-back of $HGDE(a, b, c)$ on the universal topological covering \tilde{S}' becomes trivial (as a connection). Notice that \bar{S}' is a projective curve defined over some number field, and a fortiori over some finite extension K/\mathbb{Q}_p . Up to replacing K by a finite extension, we may assume that \bar{S}' is also defined over K (and that the image in \bar{S}' of some geometric point above \bar{s} is K -rational). Thus ρ factors through a representation of $\pi_1^{\text{temp}}(\mathbb{P}_K^1 \setminus \{0, 1, \infty\}, \bar{s}) \rightarrow GL_2(K)$, which makes clear that $\Delta_p(e_0, e_1, e_\infty) \subset PGL_2(K)$. Since K is locally compact, any discrete subgroup of $PGL_2(K)$ is discontinuous [GvdP80, 1.6.4]. \square

Remark. By (b), if $\Delta_p(e_0, e_1, e_\infty)$ exists, there *exists* a finite Galois global orbifold chart $\phi : \bar{S}' \rightarrow \mathbb{P}^1$ such that the quotient of two non-proportional solutions of $HGDE(a, b, c)$ defines a multivalued meromorphic function on \bar{S}' . However, one should not believe that *any* finite Galois global orbifold chart $\phi : \bar{S}' \rightarrow \mathbb{P}^1$ has this property.

Indeed, we shall see below that the diadic triangle group $\Delta_2(4, 4, 4)$ exists. The Fermat curve F_4 over \mathbb{C}_2 provides a finite Galois global orbifold chart, with $\pi_1^{\text{top}}(F_4) = 1$ (because F_4 has tree-like reduction). But the corresponding projective hypergeometric connection does not become trivial over F_4 , since $\Delta(4, 4, 4)$ is infinite.

5.2.5. The case $\sum \frac{1}{e_i} > 1$ corresponds to the list of p -adic $HGDE$ with finite (projective) monodromy groups. This is the same as the Schwarz list:

$(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 4)$, $(2, 3, 5)$, up to permutation, and yields finite p -adic triangle groups for every p which are isomorphic to the corresponding finite Schwarz triangle groups.

On the other hand, there is *no p -adic triangle group with $\sum \frac{1}{e_i} = 1$* . Indeed, there is a global algebraic orbifold chart given by an elliptic curve with complex multiplication. Viewed over \mathbb{C}_p , these have good reduction, and it follows from 4.5.7 and 2.3.2 that the orbifold fundamental group is compact. Any discrete representation would then be finite, and it would follow that the corresponding *HGDE* has a basis of algebraic solutions. This is an algebraic property, which can be translated back over \mathbb{C} . One finds that such a *HGDE* would belong to the Schwarz list, a contradiction.

In the sequel, we restrict our attention to the “hyperbolic case” $\sum \frac{1}{e_i} < 1$. It is not clear a priori that there exists any hyperbolic p -adic triangle group.

5.3. Mumford-Schwarz orbifolds.

5.3.1. Our aim is to give a more geometric description of p -adic triangle groups, independent of the theory of monodromy of p -adic differential equations. This involves the notion of Mumford-Schwarz orbifold.

Definition 5.3.2. A Mumford-Schwarz orbifold is a p -adic Schwarz orbifold such that there exists a finite Galois global orbifold chart $\overline{\mathcal{S}}' \rightarrow \mathbb{P}^1$, $\overline{\mathcal{S}}'$ being a Mumford curve.

In other words, it is the orbifold quotient of a Mumford curve $\overline{\mathcal{S}}'$ by a finite group of automorphisms G such that $\overline{\mathcal{S}}'/G = \mathbb{P}^1$ and the Galois covering $\overline{\mathcal{S}}' \rightarrow \mathbb{P}^1$ is ramified exactly above $0, 1, \infty$ with ramification index e_0, e_1, e_∞ respectively.

According to 4.5.8, one has an exact sequence

$$1 \rightarrow \pi_1^{\text{temp}}(\overline{\mathcal{S}}', \overline{\mathcal{S}}') \rightarrow \pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty), \overline{\mathcal{S}}) \rightarrow G \rightarrow 1.$$

On the other hand, because $\overline{\mathcal{S}}'$ is a Mumford curve, $\pi_1(\overline{\mathcal{S}}', \overline{\mathcal{S}}')$ is a Schottky group embedded in $PGL_2(K)$, and the universal topological covering $\widetilde{\overline{\mathcal{S}}}'$ can be identified with the set $\Omega \subset \mathbb{P}_{\mathbb{C}_p}^1$ of ordinary points for $\pi_1(\overline{\mathcal{S}}', \overline{\mathcal{S}}')$. The subgroup Γ of all liftings of the elements of $G \subset \text{Aut}(\overline{\mathcal{S}}')$ in $PGL_2(K)$ is a subgroup of finite index inside the normalizer of $\pi_1(\overline{\mathcal{S}}', \overline{\mathcal{S}}')$ in $PGL_2(K)$; it sits in an extension

$$1 \rightarrow \pi_1^{\text{top}}(\overline{\mathcal{S}}', \overline{\mathcal{S}}') \rightarrow \Gamma \rightarrow G \rightarrow 1.$$

Remark. A Mumford-Schwarz orbifold is hyperbolic if and only if some (resp. every) curve $\overline{\mathcal{S}}'$ as above has genus ≥ 2 .

Proposition 5.3.3. *Let $\mathcal{S} = (\mathbb{P}^1; (0, e_0), (1, e_1), (\infty, e_\infty))$ be a Mumford-Schwarz orbifold. Then the p -adic triangle group $\Delta_p(e_0, e_1, e_\infty)$ exists, and*

coincides with the discrete group Γ (in particular, Γ does not depend on the choice of $\overline{S}'/\mathbb{P}^1$). Moreover, one has a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \pi_1^{\text{temp}}(\overline{S}', \overline{s}') & \longrightarrow & \pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty), \overline{s}) & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \pi_1^{\text{top}}(\overline{S}', \overline{s}') & \longrightarrow & \Delta_p(e_0, e_1, e_\infty) & \longrightarrow & G \longrightarrow 1 \end{array}$$

with surjective vertical maps and exact arrows.

PROOF. Let us consider the p -adic uniformizing differential equation attached to the orbifold-covering $\widetilde{S}' \rightarrow \mathcal{S}(e_0, e_1, e_\infty)$ (3.6). It is fuchsian with singularities at $0, 1, \infty$, and Riemann scheme

$$\left(\begin{array}{ccc} 0 & 1 & \infty \\ \frac{1}{2}(1 + \frac{1}{e_0}) & \frac{1}{2}(1 + \frac{1}{e_1}) & \frac{-1}{2}(1 + \frac{1}{e_\infty}) \\ \frac{1}{2}(1 - \frac{1}{e_0}) & \frac{1}{2}(1 - \frac{1}{e_1}) & \frac{-1}{2}(1 - \frac{1}{e_\infty}) \end{array} \right).$$

Multiplying solutions of this equation by

$$z^{\frac{1}{2}(-1+\frac{1}{e_0})}(z-1)^{\frac{1}{2}(-1+\frac{1}{e_1})}$$

transforms the equation into a (projectively equivalent) p -adic differential equation of hypergeometric type $HGDE(a, b, c)$ with $1 - c = \frac{1}{e_0}$, $c - a - b = \frac{1}{e_1}$, $\pm(a - b) = \frac{1}{e_\infty}$. By 4.6.4, the projective connection attached to $HGDE(a, b, c)$ is the projective connection attached to the projective representation given by the composed homomorphism $\pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty), \overline{s}) \rightarrow \Gamma \rightarrow PGL_2(K)$, which has discrete image. By 5.2.4. (b), we conclude that $\Delta_p(e_0, e_1, e_\infty)$ exists and coincides with Γ . The commutative diagram in the statement comes from 4.5.8. \square

5.3.4. F. Kato calls the p -adic triangle groups which arise in this way “ p -adic triangle groups of Mumford type” [Kat01]. Inspired by F. Herrlich’s study of automorphisms of Mumford curves (using graphs of groups), he has developed geometrico-combinatorial techniques to construct Mumford-Schwarz orbifolds, and even a complete classification scheme, including a description of p -adic triangle groups “of Mumford type” as explicit amalgamated sums. As a by-product, he has proved the following

Theorem 5.3.5 (F. Kato). *For $p = 2, 3$ and 5 , there are infinitely many hyperbolic Mumford-Schwarz orbifolds. For $p > 5$, there is none.*

5.3.6. The following theorem explains how to obtain all p -adic triangle groups from p -adic triangle groups “of Mumford type”.

Theorem 5.3.7. $\Delta_p(e_0, e_1, e_\infty)$ exists if and only if $\mathcal{S}(e_0, e_1, e_\infty)$ is a finite orbifold-covering of some Mumford-Schwarz orbifold $\mathcal{S}(e'_0, e'_1, e'_\infty)$. Moreover, the latter can be chosen in such a way that $\Delta_p(e_0, e_1, e_\infty) = \Delta_p(e'_0, e'_1, e'_\infty)$ as a subgroup of $PGL_2(\mathbb{C}_p)$.

PROOF. (a) Assume that we have a finite orbifold-covering $h : \mathcal{S} \rightarrow \mathcal{S}'$ of a Mumford-Schwarz orbifold by a Schwarz orbifold. The pull-back by h of the hypergeometric projective connection $HGPC(a', b', c')$ related to e'_0, e'_1, e'_∞ as above is the hypergeometric projective connection $HGPC(a, b, c)$ related to e_0, e_1, e_∞ . Because \mathcal{S}' is a Mumford-Schwarz orbifold, its uniformizing differential equation, hence $HGPC(a', b', c')$, comes from a representation of the temperate fundamental group, and it follows that so does $HGPC(a, b, c)$. Moreover, the monodromy group of $HGPC(a, b, c)$ is a subgroup of finite index of the monodromy group of $HGPC(a', b', c')$, which is the discrete finitely generated group $\Delta_p(e'_0, e'_1, e'_\infty)$. Therefore the monodromy group of $HGPC(a, b, c)$ is itself discrete finitely generated, and is $\Delta_p(e_0, e_1, e_\infty)$.

(b) Conversely, assume that $\Delta_p(e_0, e_1, e_\infty)$ exists. Let $\Sigma \rightarrow \mathbb{P}^1$ be the ramified covering corresponding to the quotient $\Delta_p(e_0, e_1, e_\infty)$ of $\pi_1^{\text{orb}}(\mathcal{S}, \bar{s})$. By definition of $\Delta_p(e_0, e_1, e_\infty)$ as projective monodromy group, any quotient of two non-proportional solutions of $HGDE(a, b, c)$ defines an étale map

$$\tau : \Sigma \rightarrow \mathbb{P}_{\mathbb{C}_p}^1$$

which is equivariant under $\Delta_p(e_0, e_1, e_\infty)$.

On the other hand, $\Delta_p(e_0, e_1, e_\infty)$ being a finitely generated discontinuous subgroup of $PGL_2(K)$ for some finite extension K/\mathbb{Q}_p , it admits a normal subgroup of finite index which is a Schottky group Γ . Let $\Omega \subset \mathbb{P}_{\mathbb{C}_p}^1$ denote the set of ordinary points for Γ . Since $\Delta_p(e_0, e_1, e_\infty)$ normalizes Γ , it preserves Ω and the quotient $H = \Delta_p(e_0, e_1, e_\infty)/\Gamma$ is a group of automorphisms of the Mumford curve Ω/Γ . We denote by T the quotient of this Mumford curve by H (a smooth projective curve).

We claim that

$$\tau(\Sigma) \subset \Omega.$$

Indeed, otherwise τ would have a value $\tau(s)$ which is a limit point for Γ , *i.e.* an accumulation point of an infinite sequence of points $\gamma_n(x)$, $x \in \mathbb{P}^1$. Since τ is étale, hence open, we may assume that $x \in \text{Im } \tau$, say $x = \tau(t)$. By Γ -equivariance of τ , we see that $\tau(s)$ is an accumulation point of the infinite sequence $\tau(\gamma_n(t))$. This contradicts the fact that the set $\{\gamma_n(t)\}$ is discrete in Σ and that τ is étale (hence locally invertible around s).

Next, being $\Delta_p(e_0, e_1, e_\infty)$ -equivariant, τ induces an analytic map

$$\bar{\tau} : \Sigma/\Delta_p(e_0, e_1, e_\infty) = \mathbb{P}^1 \rightarrow \Omega/\Delta_p(e_0, e_1, e_\infty) = T.$$

This map is necessarily algebraic and finite, and $T \cong \mathbb{P}^1$ (we normalize the latter isomorphism by imposing that $\bar{\tau}(\{0, 1, \infty\}) \subset \{0, 1, \infty\}$). Notice that

because τ is $\Delta_p(e_0, e_1, e_\infty)$ -equivariant, it follows that

$$\tau(\Sigma) = \Omega$$

and that τ is a finite étale covering (not a topological covering, unless $\Sigma = \Omega$). We have a cartesian square

$$\begin{array}{ccc} \Sigma & \longrightarrow & \Omega \\ \downarrow & & \downarrow \\ \mathbb{P}^1 & \longrightarrow & \mathbb{P}^1 \end{array}$$

in which the vertical maps are Galois orbifold-coverings with group $\Delta_p(e_0, e_1, e_\infty)$, and define orbifold quotients \mathcal{S} and \mathcal{S}' of Σ and Ω respectively. Since τ is étale, we see that \mathcal{S} is actually a finite orbifold-covering of \mathcal{S}' . This forces \mathcal{S}' to be a Schwarz orbifold (like \mathcal{S}), hence a Mumford-Schwarz orbifold by construction. \square

Corollary 5.3.8. (a) *Assume that $\Delta_p(e'_0, e'_1, e'_\infty)$ is a p -adic triangle group “of Mumford type”. Assume that $\Delta(e_0, e_1, e_\infty)$ is a subgroup of finite index of the corresponding “complex” triangle group $\Delta(e'_0, e'_1, e'_\infty)$. Then $\Delta_p(e_0, e_1, e_\infty)$ exists, and can be realized as a subgroup of finite index of $\Delta_p(e'_0, e'_1, e'_\infty)$.*

(b) *If $\Delta_p(e_0, e_1, e_\infty)$ exists, and if the corresponding “complex” triangle group $\Delta(e_0, e_1, e_\infty)$ is maximal in its commensurability class of triangle groups, then $\Delta_p(e_0, e_1, e_\infty)$ is “of Mumford type”.*

PROOF. (a) If $\Delta(e_0, e_1, e_\infty)$ is a subgroup of finite index of $\Delta(e'_0, e'_1, e'_\infty)$, there is a finite morphism $\mathbb{P}^1 \rightarrow \mathbb{P}^1$, defined over a number field, which underlies a finite orbifold-covering $\mathcal{S}(e_0, e_1, e_\infty) \rightarrow \mathcal{S}(e'_0, e'_1, e'_\infty)$ in the complex sense, but also in the p -adic sense (for every embedding of the number field in \mathbb{C}_p). The assertion comes from the “if” part of the theorem.

(b) If $\Delta_p(e_0, e_1, e_\infty)$ exists, then, similarly, by the “only if” part of 5.3.6, there is a finite morphism $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ defined over a number field, which underlies a finite orbifold-covering $\mathcal{S}(e_0, e_1, e_\infty) \rightarrow \mathcal{S}(e'_0, e'_1, e'_\infty)$ in the p -adic sense, and also in the complex sense. Hence $\Delta(e_0, e_1, e_\infty) \subset \Delta(e'_0, e'_1, e'_\infty)$ with equality if and only if $\mathcal{S}(e_0, e_1, e_\infty) = \mathcal{S}(e'_0, e'_1, e'_\infty)$. \square

Corollary 5.3.9. *There is no infinite p -adic triangle group for $p > 5$.*

PROOF. This follows from 5.3.5 and 5.3.6. \square

5.4. Arithmetic triangle groups. Review of Takeuchi’s list.

5.4.1. (Cocompact) arithmetic triangle groups are hyperbolic triangle groups which are *commensurable with the unit groups of quaternion algebras B over number fields*.

In order to make this definition more precise, let us first place ourselves in greater generality. Let E be a number field, and let B be a quaternion algebra over E . Let V be a finite set of places of E containing the places at

infinity and at least one place (possibly infinite) where B is unramified. Let us define

$$G_{(V)} = \prod B_v^\times / E_v^\times = \prod PGL_2(E_v)$$

where the product runs over all places $v \in V$ which are unramified for B . Let $\mathcal{O}_E(V)$ be the ring of so-called V -integers of E (elements which are integers at the places not in V), let $\mathcal{B}(V)$ be a maximal $\mathcal{O}_E(V)$ -order in B , and let $PSL_1(\mathcal{B}(V))$ be the image in $G_{(V)}$ of the group of elements of $\mathcal{B}(V)$ of reduced norm 1. This is a cocompact discrete subgroup of $G_{(V)}$, but the projection of $PSL_1(\mathcal{B}(V))$ in any proper factor of $G_{(V)}$ is dense [Vig80, IV.1.1].

Any subgroup of $G_{(V)}$ which is commensurable to $PSL_1(\mathcal{B}(V))$ is called an *arithmetic subgroup* of $G_{(V)}$.

In particular, arithmetic subgroups of $PGL_2(\mathbb{R})$ are subgroups which are commensurable with the unit groups $PSL_1(\mathcal{B})$ of quaternion algebras over *totally real* number fields E , which are *split at exactly one place at infinity* ∞_0 (and one can take $V = \{\infty_0\}$).

5.4.2. The complete list of cocompact arithmetic triangle groups (there are 76 such groups, up to permutation of e_0, e_1, e_∞), and of the corresponding quaternion algebras, has been established by K. Takeuchi; moreover, he has shown that two arithmetic triangle groups are commensurable with each other if and only if they come from the same quaternion algebra. Here is the list [Tak77, Table 1]:

E	$\text{disc}(B)$	(e_0, e_1, e_∞) up to permutation
\mathbb{Q}	2.3	(2, 4, 6) (2, 6, 6) (3, 4, 4) (3, 6, 6)
$\mathbb{Q}(\sqrt{2})$	v_2	(2, 3, 8) (2, 4, 8) (2, 6, 8) (2, 8, 8) (3, 3, 4) (3, 8, 8) (4, 4, 4) (4, 6, 6) (4, 8, 8)
$\mathbb{Q}(\sqrt{3})$	v_2	(2, 3, 12) (2, 6, 12) (3, 3, 6) (3, 4, 12) (3, 12, 12) (6, 6, 6)
$\mathbb{Q}(\sqrt{3})$	v_3	(2, 4, 12) (2, 12, 12) (4, 4, 6) (6, 12, 12)
$\mathbb{Q}(\sqrt{5})$	v_2	(2, 4, 5) (2, 4, 10) (2, 5, 5) (2, 10, 10) (4, 4, 5) (5, 10, 10)
$\mathbb{Q}(\sqrt{5})$	v_3	(2, 5, 6) (3, 5, 5)
$\mathbb{Q}(\sqrt{5})$	v_5	(2, 3, 10) (2, 5, 10) (3, 3, 5) (5, 5, 5)
$\mathbb{Q}(\sqrt{6})$	v_2	(3, 4, 6)
$\mathbb{Q}(\cos \pi/7)$	1	(2, 3, 7) (2, 3, 14) (2, 4, 7) (2, 7, 7) (2, 7, 14) (3, 3, 7) (7, 7, 7)
$\mathbb{Q}(\cos \pi/9)$	1	(2, 3, 9) (2, 3, 18) (2, 9, 18) (3, 3, 9) (3, 6, 18) (9, 9, 9)
$\mathbb{Q}(\cos \pi/9)$	$v_2.v_3$	(2, 4, 18) (2, 18, 18) (4, 4, 9) (9, 18, 18)

$\mathbb{Q}(\cos \pi/8)$	v_2	$(2, 3, 16) (2, 8, 16) (3, 3, 8) (4, 16, 16) (8, 8, 8)$
$\mathbb{Q}(\cos \pi/10)$	v_2	$(2, 5, 20) (5, 5, 10)$
$\mathbb{Q}(\cos \pi/12)$	v_2	$(2, 3, 24) (2, 12, 24) (3, 3, 12) (3, 8, 24) (6, 24, 24)$ $(12, 12, 12)$
$\mathbb{Q}(\cos \pi/15)$	v_3	$(2, 5, 30) (5, 5, 15)$
$\mathbb{Q}(\cos \pi/15)$	v_5	$(2, 3, 30) (2, 15, 30) (3, 3, 15) (3, 10, 30) (15, 15, 15)$
$\mathbb{Q}(\sqrt{2}, \sqrt{5})$	v_2	$(2, 5, 8) (4, 5, 5)$
$\mathbb{Q}(\cos \pi/11)$	1	$(2, 3, 11)$.

(v_p denotes the unique place of E above p . Notice that all fields E appearing in this table are Galois — even abelian — over \mathbb{Q}).

5.4.3. We do not know whether all arithmetic triangle groups are congruent subgroups (actually, unit groups of quaternion algebras seem to be unsolved hard cases of the congruence subgroup problem, *cf.* [Rap92]). Nevertheless, in each commensurability class of arithmetic triangle groups, there is a (unique) triangle group of the form

$$\Gamma^* = \{g \in B^\times \mid g\mathcal{B} = \mathcal{B}g, \text{Nr}(g) > 0\} :$$

this is the one which appears first in the each of the above rows, *cf.* [Tak77, Table 3]. The corresponding Schwarz orbifolds are Shimura orbifolds \mathcal{X}^* . More generally, any Schwarz orbifold whose orbifold fundamental group is a congruence group, is a Shimura orbifold.

5.5. Arithmetic p -adic triangle groups. A p -adic analogue of Takeuchi's list.

5.5.1. Arithmetic p -adic triangle groups are hyperbolic p -adic triangle groups which are commensurable with the p -unit groups of quaternion algebras \overline{B} over number fields.⁽³¹⁾

Let now V be the set of p -adic and infinite places of E . It follows from the discussion of 5.4.1 that arithmetic subgroups of $G_{(V)} = PGL_2(E_v)$ are subgroups which are commensurable with the p -unit groups $PSL_1(\overline{B}[\frac{1}{p}])$ of quaternion algebras over *totally real* number fields E , which are *totally definite at infinity and split at exactly one place v above p* .

5.5.2. Let $\Delta_p(e_0, e_1, e_\infty) \subset PGL_2(\mathbb{C}_p)$ be an arithmetic p -adic triangle group. Thus, there is a totally real number field $E \subset \mathbb{C}_p$, and a totally definite quaternion algebra \overline{B} over E , split at one place v above p , and a maximal order $\overline{\mathcal{B}}$, such that $\Delta_p(e_0, e_1, e_\infty) \subset PGL_2(E_v)$ is commensurable,

⁽³¹⁾In fact, these were the first p -adic triangle groups to be constructed [And98], and the existence of non-arithmetic hyperbolic p -adic triangle groups remained unclear for a while, before Kato's construction.

up to conjugation, to $PSL_1(\overline{B}[\frac{1}{p}])$. Instead of the latter, it is equivalent and more convenient to work with the image Γ_p^* in $PGL_2(E_v)$ of the normalizer $\{g \in \overline{B}^\times \mid g\overline{B}[\frac{1}{p}] = \overline{B}[\frac{1}{p}]g\}$.

Let us show that the corresponding complex triangle group $\Delta_p(e_0, e_1, e_\infty) \subset PGL_2(\mathbb{C})$ is arithmetic, attached to the same totally real number field E (embedded into \mathbb{C} via some real place ∞_0), and to the quaternion algebra B over E obtained from \overline{B} by interchanging the local invariants at ∞_0 and v (in particular B is definite at every place at infinity except ∞_0 , and *ramified at v* — which turns out to be the unique place v_p of E above p according to Takeuchi's classification).

In the case $\Delta_p(e_0, e_1, e_\infty) = \Gamma_p^*$, this follows from Čerednik's uniformization explained in II.7.4, II.7.5: in this case, the Schwarz orbifold is a Shimura orbifold \mathcal{X}^* and $\Delta(e_0, e_1, e_\infty) = \Gamma^*$.

In general, the Schwarz orbifold $\mathcal{S}(e_0, e_1, e_\infty)$ is close to be a Shimura orbifold. In fact, according to 5.3.7, the p -adic Schwarz orbifold $\mathcal{S}(e_0, e_1, e_\infty)$ is a finite orbifold-covering of a Mumford-Schwarz orbifold $\mathcal{S}(e'_0, e'_1, e'_\infty)$ with $\Delta_p(e_0, e_1, e_\infty) = \Delta_p(e'_0, e'_1, e'_\infty)$. Let Γ be a torsion-free subgroup of finite index of $\Delta_p(e'_0, e'_1, e'_\infty) \cap \Gamma_p^*$. Let $\Omega \subset \mathbb{P}_{\mathbb{C}_p}^1$ be the set of ordinary points for Γ . Then the Mumford curve $S = \Omega/\Gamma$ is simultaneously a finite orbifold covering of $\mathcal{S}(e'_0, e'_1, e'_\infty)$ and of \mathcal{X}^* (via the embeddings $\Gamma \subset \Delta_p(e'_0, e'_1, e'_\infty)$ and $\Gamma \subset \Gamma_p^*$ respectively).

All these finite orbifold-coverings are defined over number fields $\subset \mathbb{C}$. Looking at the corresponding complex orbifolds, it is now clear that $\pi_1^{\text{orb}}(\mathcal{S}(e_0, e_1, e_\infty)) = \Delta(e_0, e_1, e_\infty)$ is commensurable with $\pi_1^{\text{orb}}(\mathcal{X}^*) = \Gamma^*$, whence the claim.

5.5.3. Let us show, conversely, that if $\Delta(e_0, e_1, e_\infty)$ appears in Takeuchi's list, and if B/E is ramified at the place $v \mid p$, then $\Delta_p(e_0, e_1, e_\infty)$ exists and is an arithmetic p -adic triangle group; more precisely, it is commensurable with the p -unit group of the quaternion algebras \overline{B}/E obtained from B by interchanging the local invariants at ∞_0 (the indefinite place at infinity) and v_p (in particular \overline{B} is totally definite, and split at v_p).

In the case $\Delta(e_0, e_1, e_\infty) = \Gamma^*$, *i.e.* if (e_0, e_1, e_∞) is the first triple occurring on a row of Takeuchi's table, this follows from 4.7.4.

In general, $\Delta(e_0, e_1, e_\infty)$ is commensurable to an arithmetic triangle group $\Delta(e'_0, e'_1, e'_\infty)$ of the form Γ^* , and it is not difficult to deduce that $\Delta_p(e_0, e_1, e_\infty)$ exists and is commensurable to $\Delta_p(e'_0, e'_1, e'_\infty) = \Gamma_p^*$. The point is that the corresponding respective projective connections $HGPC(a, b, c)$ and $HGPC(a', b', c')$ become isomorphic after suitable finite pull-backs.

5.5.4. Putting 5.5.2 and 5.5.3 together, we conclude that the arithmetic p -adic triangle groups $\Delta_p(e_0, e_1, e_\infty)$ are exactly those corresponding to the triples (e_0, e_1, e_∞) in Takeuchi's list when v_p divides $\text{disc}(B)$. From that list, one gets:

Theorem 5.5.5. [And98]. *Up to permutation of the indices, there are 45 arithmetic 2-adic triangle groups, 16 arithmetic 3-adic triangle groups, 9*

arithmetic 5-adic triangle groups (and no arithmetic p -adic triangle group for $p > 5$).

More precisely, they are given by the following tables:

Arithmetic 2-adic triangle groups:

E	$\text{disc}(\overline{B})$	(e_0, e_1, e_∞) up to permutation
\mathbb{Q}	3	(2, 4, 6) (2, 6, 6) (3, 4, 4) (3, 6, 6)
$\mathbb{Q}(\sqrt{2})$	1	(2, 3, 8) (2, 4, 8) (2, 6, 8) (2, 8, 8) (3, 3, 4) (3, 8, 8) (4, 4, 4) (4, 6, 6) (4, 8, 8)
$\mathbb{Q}(\sqrt{3})$	1	(2, 3, 12) (2, 6, 12) (3, 3, 6) (3, 4, 12) (3, 12, 12) (6, 6, 6)
$\mathbb{Q}(\sqrt{5})$	1	(2, 4, 5) (2, 4, 10) (2, 5, 5) (2, 10, 10) (4, 4, 5) (5, 10, 10)
$\mathbb{Q}(\sqrt{6})$	1	(3, 4, 6)
$\mathbb{Q}(\cos \pi/9)$	v_3	(2, 4, 18) (2, 18, 18) (4, 4, 9) (9, 18, 18)
$\mathbb{Q}(\cos \pi/8)$	1	(2, 3, 16) (2, 8, 16) (3, 3, 8) (4, 16, 16) (8, 8, 8)
$\mathbb{Q}(\cos \pi/10)$	1	(2, 5, 20) (5, 5, 10)
$\mathbb{Q}(\cos \pi/12)$	1	(2, 3, 24) (2, 12, 24) (3, 3, 12) (3, 8, 24) (6, 24, 24) (12, 12, 12)
$\mathbb{Q}(\sqrt{2}, \sqrt{5})$	1	(2, 5, 8) (4, 5, 5)

Arithmetic 3-adic triangle groups:

\mathbb{Q}	2	(2, 4, 6) (2, 6, 6) (3, 4, 4) (3, 6, 6)
$\mathbb{Q}(\sqrt{3})$	1	(2, 4, 12) (2, 12, 12) (4, 4, 6) (6, 12, 12)
$\mathbb{Q}(\sqrt{5})$	1	(2, 5, 6) (3, 5, 5)
$\mathbb{Q}(\cos \pi/9)$	v_2	(2, 4, 18) (2, 18, 18) (4, 4, 9) (9, 18, 18)
$\mathbb{Q}(\cos \pi/15)$	1	(2, 5, 30) (5, 5, 15)

Arithmetic 5-adic triangle groups:

$\mathbb{Q}(\sqrt{5})$	1	(2, 3, 10) (2, 5, 10) (3, 3, 5) (5, 5, 5)
$\mathbb{Q}(\cos \pi/15)$	1	(2, 3, 30) (2, 15, 30) (3, 3, 15) (3, 10, 30) (15, 15, 15)

Remarks 5.5.6. (a) In these lists, one notices that p divides at least one of the numbers e_0, e_1, e_∞ . This means that the parameters a, b, c of the corresponding hypergeometric differential equation are not p -adic integers. In the literature on p -adic hypergeometric functions, on the contrary, a, b, c are always assumed to be p -adic integers, for obvious convergence reasons.

It is well-known that a fuchsian p -adic differential equation with Frobenius structure has exponents in $\mathbb{Q} \cap \mathbb{Z}_p$. Since this is not the case here, there is no Frobenius structure in the standard sense. However, there is some hidden Frobenius structure due to the “geometric origin” of the differential equation. We leave to the reader the intriguing question of finding out the right formulation of this hidden Frobenius structure (hint: the Frobenius structure is explicit at the level of the period mapping, *cf.* [BZ95]).

(b) Takeuchi has also given a table with the various inclusions of triangle groups in each commensurability class. When $\Delta(e_0, e_1, e_\infty)$ is maximal (and p ramified in the corresponding quaternion algebra), the corresponding p -adic triangle group $\Delta_p(e_0, e_1, e_\infty)$ is “of Mumford type”, according to 5.3.8. (b).

Arithmetic p -adic triangle groups not “of Mumford type” do exist, and are related to Shimura curves with *level divisible by p* . One can construct such p -adic triangle groups using Drinfeld’s finite étale coverings Υ^n of the Drinfeld “half space” Ω , which uniformize p -adically Shimura curves with level $p^n \cdot M$ (p prime to M), *cf.* II.7.4.7. One has $\Gamma_p^+(p^n M) = \Gamma_p^+(M)$, and the orbifold-quotient of a suitable intermediate covering Σ between Ω and Υ^n by $\Gamma_p^+(M)$ may be a Schwarz orbifold but not a Mumford-Schwarz orbifold. It seems that $\Delta_2(4, 4, 4)$ is of this type, for $p = 2, n = M = 1$ (we have not checked the details); the corresponding $\Gamma_2^+(1)$ is $\Delta_2(3, 3, 4)$.

(c) Let \bar{S} be a smooth projective curve defined over $\bar{\mathbb{Q}}$, and ζ_i be $\bar{\mathbb{Q}}$ -points on \bar{S} endowed with indices e_i . We then say that the orbifold $\mathcal{S} = (\bar{S}, (\zeta_i, e_i))$ is defined over $\bar{\mathbb{Q}}$. In general, there is no reason for the uniformizing differential equation to be defined over $\bar{\mathbb{Q}}$.

However, this occurs if either \mathcal{S} is a Schwarz orbifold (the uniformizing differential equation being essentially of hypergeometric type with rational parameters), or a Shimura orbifold (the uniformizing differential equation being essentially of Gauss-Manin type, *cf.* 4.7.4). On the other hand, it is not difficult to see that the property for the uniformizing differential equation to be defined over $\bar{\mathbb{Q}}$ depends only on the commensurability class of the orbifold fundamental group $\pi_1^{\text{orb}}(\mathcal{S})$.

A transcendence conjecture of Chudnovsky, modified by Krammer [Kra96, 12], predicts that the Schwarz and the Shimura cases are the only exceptions, up to commensurability. Namely:

Conjecture 5.5.7. (*Chudnovsky-Krammer*). *Let \mathcal{S} be a one-dimensional hyperbolic orbifold defined over $\bar{\mathbb{Q}}$. Assume that the uniformizing differential equation is defined over $\bar{\mathbb{Q}}$. Then $\pi_1^{\text{orb}}(\mathcal{S})$ is either commensurable to a triangle group, or is arithmetic.*

We *conjecture* the (straightforward) p -adic analogue of 5.5.7.

(d) By the theory of p -adic Betti lattices, there is a canonical $\overline{\mathbb{Q}}$ -structure in the \mathbb{C}_p -space of solutions at any point of $\mathbb{P}^1(\overline{\mathbb{Q}})$ of the hypergeometric differential equation $HDGE(a, b, c)$ attached to an arithmetic p -adic triangle group (*cf.* 4.7.4. If the base point is chosen to be one of the singularities $0, 1, \infty$, one applies 4.7.4 to a finite orbifold-covering of the Schwarz orbifold which is a Shimura curve attached to a torsion-free congruence subgroup).

This raises the following (open) questions:

Is there a $\overline{\mathbb{Q}}$ -structure in the \mathbb{C}_p -space of solutions of the hypergeometric differential equation $HDGE(a, b, c)$ attached to a non-arithmetic p -adic triangle group? Are non-arithmetic p -adic triangle groups related to some p -adic period mappings and suitable higher-dimensional Shimura orbifolds?

5.6. On special values of hypergeometric functions attached to arithmetic p -adic triangle groups.

5.6.1. Let a, b, c be rational numbers such that $c > a + b$. Then the hypergeometric series ${}_2F_1(a, b; c; z)$ converges at 1 and takes the value (Gauss' formula)

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}.$$

If one considers ${}_2F_1(a, b; c; z)$ as a multivalued analytic function on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, the values at 1 form a finitely generated subgroup of \mathbb{C} ; they generate a $\overline{\mathbb{Q}}$ -subspace of \mathbb{C} of dimension ≤ 2 , namely:

$$\frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}\overline{\mathbb{Q}} + \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}\overline{\mathbb{Q}}.$$

Similarly, its values at ∞ generate the $\overline{\mathbb{Q}}$ -subspace of \mathbb{C}

$$\frac{\Gamma(c)\Gamma(b-a)}{\Gamma(c-a)\Gamma(b)}\overline{\mathbb{Q}} + \frac{\Gamma(c)\Gamma(a-b)}{\Gamma(a)\Gamma(c-b)}\overline{\mathbb{Q}}.$$

5.6.2. If $HGDE(a, b, c)$ is the differential equation attached as above to an *arithmetic* triangle group $\Delta(e_0, e_1, e_\infty)$, these formulas are special cases of similar formulas expressing the special values of ${}_2F_1(a, b; c; z)$ at CM points, in terms of Gamma values, up to factors in $\overline{\mathbb{Q}}$.

The point is that there is a $\overline{\mathbb{Q}}$ -structure in the complex space of solutions which is stable under monodromy: the $H_{1,B}(\cdot, \overline{\mathbb{Q}})$ of the fake elliptic curves A_z parametrized by $z \in \mathbb{P}^1$ (\mathbb{P}^1 being viewed as a Shimura curve) if the quaternion algebra attached to $\Delta(e_0, e_1, e_\infty)$ is over \mathbb{Q} , *cf.* II.7.6.3 for the general case. For any point $z \in \mathbb{P}^1(\overline{\mathbb{Q}})$, let V_z be the $\overline{\mathbb{Q}}$ -structure in the complex space of solutions at z consisting of solutions whose Taylor coefficients are in $\overline{\mathbb{Q}}$. Then V_z is related to $H_{1,B}(A_{z,\mathbb{C}}, \overline{\mathbb{Q}})$ by the period matrix of A_z . At any CM point z (according to an old observation of Shimura, the branch points of a Shimura orbifold — here, $0, 1, \infty$ — are actually CM points), the Chowla-Selberg formula (I.4.6.3) thus relates the

two $\overline{\mathbb{Q}}$ -structures in terms of Γ -values. In particular, V_0 is generated by ${}_2F_1(a, b, c; z)$ and $z^{1-c}{}_2F_1(a+1-c, b+1-c, 2-c; z)$, and the branch point 0 corresponds to a CM point (for elliptic curves rather than fake elliptic curves, this was worked out in I.5.3.10).

5.6.3. There is an entirely analogous phenomenon for *arithmetic p -adic* triangle group $\Delta_p(e_0, e_1, e_\infty)$: thanks to II.7.6.4, there is a $\overline{\mathbb{Q}}$ -structure in the \mathbb{C}_p -space of solutions which is stable under p -adic monodromy: the “ p -adic Betti lattice” $H_{1,B}(\cdot, \overline{\mathbb{Q}})$ of the fake elliptic curves A_z if the quaternion algebra attached to $\Delta_p(e_0, e_1, e_\infty)$ is over \mathbb{Q} , *cf.* II.7.6.3 for the general case. “Periods” should be understood here in the sense of I.5.3; p -adic CM periods were computed in terms of Γ_p -values in I.4.6.3, I.4.6.4 (for $p \neq 2$), I.5.3.8.

Therefore, if $HGDE(a, b, c)$ is the differential equation attached as above to an *arithmetic p -adic* triangle group $\Delta_p(e_0, e_1, e_\infty)$, there is, at least for odd p , an expression in terms of Γ_p -values of the $\overline{\mathbb{Q}}$ -space of dimension ≤ 2 generated by the values taken at any given CM point, *e.g.* the point 1, by ${}_2F_1(a, b, c; z)$ viewed as a multivalued p -adic analytic function on a suitable finite étale covering of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$.

We shall see examples in the next section.

6. The tale of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ (Escher’s triangle group and its diadic and triadic twins).

ABSTRACT: We study the rich geometry related to the simplest cocompact arithmetic triangle group: associated Shimura curves, period mapping and monodromy — in the archimedean, diadic and triadic cases. At other primes, the story would be similar to that of $F(\frac{1}{2}, \frac{1}{2}, 1; z)$; we do not repeat it.

6.1. Escher’s triangle group $\Delta(2, 4, 6)$.

6.1.1. This is the symmetry group of the black-and-white tessellation of the unit disk by hyperbolic triangles with angles $(\frac{\pi}{2}, \frac{\pi}{4}, \frac{\pi}{6})$




FIGURE 4

M.C. Escher has engraved several celebrated variants of this Schwarz tessellation (“cirkel-limiet”, angels and demons), cf [Esc71]⁽³²⁾.

By gluing together two basic triangles, we get hyperbolic triangles with angles $(\frac{\pi}{3}, \frac{\pi}{4}, \frac{\pi}{4})$ or $(\frac{\pi}{2}, \frac{\pi}{6}, \frac{\pi}{6})$




FIGURE 5

⁽³²⁾he has also used another Schwartz tessellation in a picture — “cirkel-limiet III” — brilliantly analyzed by H.S.M. Coxeter [Cox79]

This exhibits $\Delta(3, 4, 4)$ and $\Delta(2, 6, 6)$ as subgroups of index 2 of $\Delta(2, 4, 6)$. Furthermore, by gluing two triangles with angles $(\frac{\pi}{2}, \frac{\pi}{6}, \frac{\pi}{6})$, one can get a triangle with angles $(\frac{\pi}{3}, \frac{\pi}{6}, \frac{\pi}{6})$, which exhibits $\Delta(3, 6, 6)$ as a subgroup of index 2 of $\Delta(2, 6, 6)$.

Similarly, the intersection of $\Delta(2, 6, 6)$ and $\Delta(3, 4, 4)$ in $\Delta(2, 4, 6)$ is a quadrangle group $\diamond(2, 2, 3, 3)$.

6.1.2. The vertices of one of the basic triangles are

$$O = (0, 0), \quad Q = (\sqrt{2} - 1, 0), \quad P = \left(\sqrt{3} \cos \frac{5\pi}{12}, \cos \frac{5\pi}{12} \right).$$

The matrix of the composition of the reflection along side QP , *resp.* PO , followed by the reflection along side OQ is $A = \sqrt{-1} \begin{pmatrix} \sqrt{2} & -1 \\ 1 & -\sqrt{2} \end{pmatrix}$, *resp.*

$$B = \frac{1}{2} \begin{pmatrix} \sqrt{3} + \sqrt{-1} & 0 \\ 0 & \sqrt{3} - \sqrt{-1} \end{pmatrix}.$$

These matrices satisfy $A^2 = B^6 = (AB)^4 = -1$ and, taken up to sign, generate the triangle group $\Delta(2, 4, 6)$, (*cf.* [Mag74, II5].)

6.1.3. The group $\Delta(2, 4, 6)$ enjoys the following “universal property”, discovered by Mennike in 1968:

for all but a finite number of indefinite ternary quadratic forms q with integral coefficients, the (integral) associated unitary group of q is contained in $\Delta(2, 4, 6)$.

It coincides with $\Delta(2, 4, 6)$ in the case of $q = 3x^2 - y^2 - z^2$, (*cf.* [Mag74, III3]). These unitary groups have been thoroughly investigated by Fricke and Klein.

6.2. Some Shimura curves attached to $D_{2,3}$.

6.2.1. The \mathbb{Q} -subalgebra of $M_2(\mathbb{Q}(\sqrt{-1}, \sqrt{2}, \sqrt{3}))$ generated by the matrices A and B is the quaternion algebra $B = B_{2,3}$ over \mathbb{Q} with discriminant 6. This quaternion algebra is unique up to isomorphism and admits standard generators i, j and relations $i^2 = -1$, $j^2 = 3$, $ij = -ji$.

It follows that $\Delta(2, 4, 6)$ is a subgroup of $B_{2,3}^\times / \mathbb{Q}^\times$. According to [Tak77], $\Delta(2, 4, 6)$, $\Delta(3, 4, 4)$, $\Delta(2, 6, 6)$ and $\Delta(3, 6, 6)$ are the only cocompact triangle groups which give rise to a quaternion algebra over \mathbb{Q} .

As any indefinite quaternion algebra over \mathbb{Q} , $B_{2,3}$ has a unique maximal order $\mathcal{B} = \mathcal{B}_{2,3}$ modulo conjugation, *cf.* [Vig80, p.99]. It can be described as $\mathbb{Z} + \mathbb{Z}i + \mathbb{Z}j + \mathbb{Z}\rho$, where $\rho = \frac{1+i+j+ij}{2}$. According to [Tak77], it can also be described as *the subring of $B_{2,3}$ generated by the elements of $\diamond(2, 2, 3, 3)$* (the subgroup of $\Delta(2, 4, 6)$ of index 4 considered above).

Following the notation of II.7.4, we introduce the following arithmetic subgroups of $B^\times / \mathbb{Q}^\times$:

- the image $\Gamma^+(N)$ of the group $\{g \in (1 + N\mathcal{B})^\times \mid \text{Nr}(g) = 1\}$ in $PSU(1, 1)(\mathbb{R}) = \text{Aut}(D(0, 1^-))$, where Nr stands for the reduced norm ($\Gamma^+(1) = PSL_1(\mathcal{B})$ is abbreviated into Γ^+),
- the image Γ^* in $PSU(1, 1)(\mathbb{R})$ of the group $N(\mathcal{B})^+ = \{g \in B^\times \mid g\mathcal{B} = \mathcal{B}g \mid \text{Nr}(g) > 0\}$, i.e. $N(\mathcal{B})^+/\mathbb{Q}^\times$.

According to [Tak77], one has:

- $\Delta(2, 4, 6) = \Gamma^*$,
- $\diamond(2, 2, 3, 3) = \Gamma^+$.

These groups also appear in [Vig80, p.122], together with the orbifolds $\mathcal{X}_{\mathbb{C}}^{+\text{an}} = D(0, 1^-)/\diamond(2, 2, 3, 3)$ and $\mathcal{X}_{\mathbb{C}}^{*\text{an}} = D(0, 1^-)/\Delta(2, 4, 6)$. The orbifold Euler characteristics are found to be $-\frac{1}{3}$ and $-\frac{1}{12}$ respectively.

6.2.2. Any element of $N(\mathcal{B})^+$ can be written uniquely as a product

$$q.g.(1+i)^{\epsilon_2}(3i+j+ij)^{\epsilon_3},$$

where $q \in \mathbb{Q}$, $g \in SL_1(\mathcal{B})$, $\epsilon_m = 0$ or 1 , cf. [Mic84]. The quotient group

$$\begin{aligned} & \Delta(2, 4, 6)/\diamond(2, 2, 3, 3) \\ &= \{1, w_2 = [1+i], w_3 = [3i+j+ij], w_6 = w_2w_3 = [-3+3i+2ij]\} \\ & \cong (\mathbb{Z}/2\mathbb{Z})^2 \end{aligned}$$

is called the Atkin-Lehner group. The fixpoints of the involutions w_m have been the object of thorough investigations, cf. *loc. cit.* and [Shi67] (cf. also II.7.4.4). In our case, the results are:

- the two fixpoints of w_2 on $D(0, 1^-)/\diamond(2, 2, 3, 3)$ have the same image in $D(0, 1^-)/\Delta(2, 4, 6)$, which is the image of the fixpoint in $D(0, 1^-)$ of any maximal embedding of $\mathbb{Z}[\sqrt{-1}]$ in \mathcal{B} ,
- the two fixpoints of w_3 on $D(0, 1^-)/\diamond(2, 2, 3, 3)$ have the same image in $D(0, 1^-)/\Delta(2, 4, 6)$, which is the image of the fixpoint in $D(0, 1^-)$ of any maximal embedding of $\mathbb{Z}[\sqrt[3]{1}]$ in \mathcal{B} ,
- the two fixpoints of w_2 on $D(0, 1^-)/\diamond(2, 2, 3, 3)$ have the same image in $D(0, 1^-)/\Delta(2, 4, 6)$, which is the image of the fixpoint in $D(0, 1^-)$ of any maximal embedding of $\mathbb{Z}[\sqrt{-6}]$ in \mathcal{B} ,

From easy considerations of ramification, one draws that

- $\langle \Gamma^+, (1+i) \rangle = \Delta(3, 4, 4)$
- $\langle \Gamma^+, (3i+j+ij) \rangle = \Delta(2, 6, 6)$
- $\langle \Gamma^+, (1+i) \rangle$ is the third intermediate group, which is a quadrangle group $\diamond(2, 2, 2, 3)$ (not to be confused with $\diamond(2, 2, 3, 3)$).

Lemma 6.2.3. *The congruence group $\Gamma^+(2)$ is torsion-free. The quotient $\Gamma^+/\Gamma^+(2)$ is the tetrahedral group $\Delta(2, 3, 3) \cong \mathfrak{A}_4$ (of order 12).*

PROOF. Recall that $\mathcal{B} = \mathbb{Z} + \mathbb{Z}i + \mathbb{Z}j + \mathbb{Z}\rho$. Any element $a + bi + cj + dij$ of finite order in the congruence subgroup $SL_1(\mathcal{B})(2)$ of level two must have

reduced trace $2a = \pm 2$ and reduced norm $a^2 + b^2 - 3c^2 - 3d^2 = 1$; it is easy to see that the only such elements are ± 1 , which implies the first assertion.

The quotient group $\Gamma^+/\Gamma^+(2)$ is represented by $\{1, i, j, ij, \frac{1 \pm i \pm j \pm ij}{2}\}$ modulo 2. It admits $\{1, \bar{i}, \bar{j}, \bar{ij}\}$ as a normal subgroup, and $\{1, \bar{\rho}, \bar{\rho}^2\}$ as a non-normal subgroup. its center is trivial. It is thus isomorphic to the tetrahedral group, the group of oriented symmetries of the tetrahedron. \square



FIGURE 6

Corollary 6.2.4. *The associated Shimura curve $\mathcal{X}^+(2)$ has genus $g = 3$.*

PROOF. By definition, $\mathcal{X}^+(2)_{\mathbb{C}}^{\text{an}} = \mathbb{D}(0, 1^-)/\Gamma^+(2)$ (here we prefer to identify the universal covering with the unit disk rather than the Poincaré upper half plane as in II.7.4). The assertion follows from the computation $2 - 2g = \chi(\mathbb{D}(0, 1^-)/\Gamma^+(2)) = 12 \cdot \chi_{\text{orb}}(\mathbb{D}(0, 1^-)/\Gamma^+) = -4$. \square

Lemma 6.2.5. *The congruence group $\Gamma^+(2)$ is a normal subgroup of $\Delta(2, 4, 6) = \Gamma^*$. The quotient $\Gamma^*/\Gamma^+(2)$ is the extended octahedral group $\Delta^*(2, 3, 4) \cong \mathfrak{S}_4 \times \mathbb{Z}/2\mathbb{Z}$ (of order 48). The generator of the factor $\mathbb{Z}/2\mathbb{Z}$ is given by the image \tilde{w}_3 of $(3i + j + ij)$.*

We refer to [For29, N. 58] and [Cox74, p.27] for a discussion of $\Delta^*(2, 3, 4)$, the group of all symmetries of the octahedron.




FIGURE 7

PROOF. Let us first show that $\Gamma^+(2)$ is a normal subgroup of $\Delta(2, 6, 6) = \langle \Gamma^+, (3i + j + ij) \rangle$. It makes sense to reduce the latter modulo 2, $(3i + j + ij)$ being sent to $1 \in \Delta(2, 3, 3)$; on the other hand, Nr induces a surjective

homomorphism $\Delta(2, 6, 6) \rightarrow 3^{\mathbb{Z}}/9^{\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z}$, and $(3i + j + ij)$ is sent to a generator of $\mathbb{Z}/2\mathbb{Z}$. It follows that $\Gamma^+(2)$ is a normal subgroup of $\Delta(2, 6, 6)$, with quotient group $\mathfrak{A}_4 \times \mathbb{Z}/2\mathbb{Z}$.

We next remark that every element of $\Gamma^+(2)$ can be written

$$\begin{aligned} a + bi + cj + dij, \text{ with } a, b, c, d \in \mathbb{Z}, b \equiv c \equiv d \not\equiv a \pmod{2}, \\ a^2 + b^2 - 3c^2 - 3d^2 = 1. \end{aligned}$$

Since $(1+i)(a+bi+cj+dij)(1+i)^{-1} = a+bi-dj+cij$, we see that $(1+i)$ normalizes $\Gamma^+(2)$, and conclude that $\Gamma^+(2)$ is normal in Γ^* . The quotient group sits in an extension

$$1 \rightarrow \Delta(2, 3, 3) \times \mathbb{Z}/2\mathbb{Z} \rightarrow G \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 1,$$

the quotient $\mathbb{Z}/2\mathbb{Z}$ being generated by $(1+i)$. Since the image of $(1+i)$ in G is not central, we conclude that $G \cong \mathfrak{S}_4 \times \mathbb{Z}/2\mathbb{Z}$. \square

Corollary 6.2.6. *The automorphism group of the curve $\mathcal{X}^+(2)_{\mathbb{C}}$ is $\Delta^*(2, 3, 4)$.*

PROOF. It follows from 6.2.4 and 6.2.5 that $\text{Aut } \mathcal{X}^+(2)_{\mathbb{C}}$ contains $\Delta^*(2, 3, 4)$. On the other hand, any automorphism of $\mathcal{X}^+(2)_{\mathbb{C}}$ lifts to an element of $SU(1, 1)(\mathbb{R})$ which normalizes \mathcal{B} . The image in $PSU(1, 1)(\mathbb{R})$ of such an element lies in Γ^* , whence $\text{Aut } \mathcal{X}^+(2)_{\mathbb{C}} \subset \Gamma^*/\Gamma^+(2) = \Delta^*(2, 3, 4)$. \square

By similar arguments, one shows that the groups $\Delta(3, 4, 4)/\Gamma^+(2)$ and $\diamond(2, 2, 2, 3)/\Gamma^+(2)$ are the octahedral group $\Delta(2, 3, 4) \cong \mathfrak{S}_4$.

One reads from the following table of ramification that \tilde{w}_3 is a hyperelliptic involution of $\mathcal{X}^+(2)_{\mathbb{C}}$.




FIGURE 8

6.2.7. Similar considerations apply to the congruence subgroup of level three $\Gamma^+(3)$. The Shimura curve $\mathcal{X}^+(3)_{\mathbb{C}}$ has genus 4 and its automorphism group is of order 72. We omit the details.

6.2.8. By the general theory of Shimura curves, $\mathcal{X}^+(2)$ is a moduli space for polarized abelian surfaces with quaternionic multiplication by $\mathcal{B}_{2,3}$ and level-two structure, *cf.* II.7.4. It is defined over \mathbb{Q} , as well as \mathcal{X}^* and the intermediate Shimura curves attached to $\Delta(3, 4, 4)$, $\Delta(2, 6, 6)$, $\diamond(2, 2, 2, 3)$ and $\diamond(2, 2, 3, 3)$. According to Y. Ihara, A. Kurihara [Kur79], A. Ogg and J.F. Michon, in each of the cases $\Delta(2, 4, 6)$, $\Delta(3, 4, 4)$, $\Delta(2, 6, 6)$, $\diamond(2, 2, 2, 3)$, the \mathbb{Q} -curve itself is $\mathbb{P}_{\mathbb{Q}}^1$, whereas for $\diamond(2, 2, 3, 3)$, it is the conic $x^2 + y^2 + 3z^2 = 0$.⁽³³⁾ The Atkin-Lehner involutions are defined over \mathbb{Q} .

6.2.9. Any CM point on $\mathcal{X}^+(2)$ parametrizes an abelian surface isogenous to the square of an elliptic curve with complex multiplication. It follows from 6.2.2 that the points of ramification of the “projection” $\mathcal{X}^+(2) \rightarrow \mathcal{X}^*$ are CM points. More precisely, the corresponding imaginary quadratic fields are

- $\mathbb{Q}[\sqrt{-1}]$ for the points of ramification index 4 (*e.g.* the image of the point $P \in D(0, 1^-)$, *cf.* 6.1.2)
- $\mathbb{Q}[\sqrt{-3}]$ for the points of ramification index 6 (*e.g.* the image of the point $O \in D(0, 1^-)$)
- $\mathbb{Q}[\sqrt{-6}]$ for the points of ramification index 2 (*e.g.* the image of the point $Q \in D(0, 1^-)$).

6.3. Triadic uniformization.

6.3.1. The Shimura curve $\mathcal{X}^+(2)$ has good reduction at any prime $p \neq 2, 3$. As for the critical primes 2 and 3, the situation for $p = 3$ is easier to describe than for $p = 2$, since 3 does not divide the level (II.7.4).

For the triadic Čerednik-Drinfeld uniformization, the relevant quaternion algebra is the Hamilton definite quaternion algebra $\overline{\mathcal{B}} = B_{2,\infty}$ over \mathbb{Q} with basis $(1, i, j, ij)$ and relations $i^2 = j^2 = -1, ij = -ji$. A maximal order (unique up to conjugation, *cf.* [Vig80, 5.11]) consists of the Hurwitz quaternions

$$\overline{\mathcal{B}} = \overline{\mathcal{B}}_{2,\infty} = \mathbb{Z} + \mathbb{Z}i + \mathbb{Z}j + \mathbb{Z}\rho, \quad \rho = \frac{1 + i + j + ij}{2}.$$

The group $PGL_1(\overline{\mathcal{B}}) = \overline{\mathcal{B}}^{\times} / \pm 1$ is finite, isomorphic to the tetrahedral group $\Delta(2, 3, 3) \cong \mathfrak{A}_4$.

Following the notation of II.7.4.3, we introduce the following $\{3, \infty\}$ -arithmetic subgroups of $\overline{\mathcal{B}}^{\times} / \mathbb{Q}^{\times}$:

⁽³³⁾ on the other hand, we do not know any hyperelliptic equation for $\mathcal{X}^+(2)$, even over \mathbb{C} .

- the image $\Gamma_3^+(N)$ of the group $\{g \in (1 + N\overline{\mathcal{B}}[\frac{1}{3}])^\times \mid \text{ord}_3 \text{Nr}(g) \text{ even}\}$ in $PGL_2(\mathbb{Q}_3) = \text{Aut}(\Omega_{\mathbb{C}_3})$ ($\Gamma_3^+(1)$ is abbreviated into Γ_3^+),

- the image Γ_3^* in $PGL_2(\mathbb{Q}_3)$ of the group $N(\mathcal{B}[\frac{1}{3}]) = \{g \in \overline{\mathcal{B}}^\times \mid g\overline{\mathcal{B}}[\frac{1}{3}] = \overline{\mathcal{B}}[\frac{1}{3}]g\}$, *i.e.* $N(\mathcal{B}[\frac{1}{3}])/\mathbb{Q}^\times$.

According to Čerednik-Drinfeld, one has:

- $\mathcal{X}^+(2)_{\mathbb{C}_3}^{\text{an}} = \Omega_{\mathbb{C}_3}/\Gamma_3^+(2)$

- $\mathcal{X}^+_{\mathbb{C}_3} = \Omega_{\mathbb{C}_3}/\Gamma_3^+$.

- $\mathcal{X}^*_{\mathbb{C}_3} = \Omega_{\mathbb{C}_3}/\Gamma_3^*$.

In terms of triadic triangle (*resp.* quadrangle) groups, one can thus write

- $\Delta_3(2, 4, 6) = \Gamma_3^*$,

- $\diamond_3(2, 2, 3, 3) = \Gamma_3^+$.

6.3.2. These uniformizations actually hold over \mathbb{Q}_9 , the quadratic unramified extension of \mathbb{Q}_3 . The corresponding Mumford curves over \mathbb{Q}_9 appear in [GvdP80, IX], without referring to the Čerednik theorem. The results of *loc. cit.* allow to give a concrete description of the arithmetic groups in terms of the generators $1, i, j, ij$ of $\overline{\mathcal{B}}$.

Let us fix a square root of -1 in \mathbb{Q}_9 . A convenient matrix representation for the elements of $\overline{\mathcal{B}} \otimes \mathbb{Q}_9$ is

$$a + bi + cj + dij \mapsto \begin{pmatrix} a + b\sqrt{-1} & c + d\sqrt{-1} \\ -c + d\sqrt{-1} & a - b\sqrt{-1} \end{pmatrix}$$

Of course, $\overline{\mathcal{B}} \otimes \mathbb{Q}_3 \cong M_2(\mathbb{Q}_3)$, and any such matrix representation is conjugate to the previous after $\otimes \mathbb{Q}_9$ one by some $u \in PGL_2(\mathbb{Q}_9)$; u is a kind of triadic Cayley transform which sends the Drinfeld space Ω to the complement $u\Omega$ of the “circle” $\{x + y\sqrt{-1} \mid x, y \in \mathbb{Q}_3, x^2 + y^2 = -1\}$ in the projective line (here, we follow the usual convention that PGL_2 acts on the left on \mathbb{P}^1).

After *loc. cit.*, the congruence level subgroup of level two $PGL_1(\overline{\mathcal{B}}[\frac{1}{3}])(2)$ is generated by

$$z_{\epsilon\epsilon'} = i + \epsilon j + \epsilon' ij, \quad \epsilon = \pm 1, \epsilon' = \pm 1$$

with the only relations $z_{\epsilon\epsilon'}^2 = 1$. The subgroup of index two $\Gamma_3^+(2)$ consisting of words of even length (which is also the kernel of the homomorphism $PGL_1(\overline{\mathcal{B}}[\frac{1}{3}])(2) \rightarrow 3^\mathbb{Z}/9^\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$ induced by Nr) is a *Schottky group of rank three* in $PGL_2(\mathbb{Q}_9)$, freely generated by

$$\gamma_1 = z_{++}z_{+-} = \begin{pmatrix} -1 - 2\sqrt{-1} & 2 \\ -2 & -1 + 2\sqrt{-1} \end{pmatrix},$$

$$\gamma_2 = z_{++}z_{-+} = \begin{pmatrix} -1 - 2\sqrt{-1} & -2\sqrt{-1} \\ -2\sqrt{-1} & -1 + 2\sqrt{-1} \end{pmatrix},$$

$$\gamma_3 = z_{++}z_{+-} = \begin{pmatrix} 1 & 2 - 2\sqrt{-1} \\ -2 - 2\sqrt{-1} & 1 \end{pmatrix}.$$

Moreover $PGL_1(\overline{\mathcal{B}}[\frac{1}{3}])$ is the semi-direct product of $PGL_1(\overline{\mathcal{B}}[\frac{1}{3}])(2)$ and the finite group $PGL_1(\overline{\mathcal{B}})$, and Γ_3^* is generated by $PGL_1(\overline{\mathcal{B}}[\frac{1}{3}])$ and the image of $(1 + i)$.

At last, the dual graph of the stable reduction modulo 3 of $\mathcal{X}^+(2)_{\mathbb{Q}_9}^{\text{an}}$ is




FIGURE 9

6.3.3. The case of the prime 2 is similar if one replaces $\mathcal{X}^+(2)$ by $\mathcal{X}^+(3)^{(34)}$. This is a Mumford curve over \mathbb{Q}_4 , the unramified quadratic extension of \mathbb{Q}_2 . The relevant quaternion algebra here is $B_{3,\infty}$. A maximal order (again unique up to conjugation) is given by $\mathbb{Z} + \mathbb{Z}i + \mathbb{Z}\frac{i+j}{2} + \mathbb{Z}\frac{1+ij}{2}$, $i^2 = -1$, $j^2 = -3$, $ij = -ji$.

- $\Delta_2(2, 4, 6) = \Gamma_2^*$,
- $\diamond_2(2, 2, 3, 3) = \Gamma_2^+$.

6.4. The non-archimedean triangle groups $\Delta_3(2, 4, 6)$, $\Delta_2(2, 4, 6)$ and Herrlich’s tree.

6.4.1. Our aim here is to explore the combinatorics of the finitely generated groups $\Delta_3(2, 4, 6)$ and $\Delta_2(2, 4, 6)$.

It follows from the above considerations that $\Delta_3(2, 4, 6)$ sits in an exact sequence

$$1 \rightarrow \langle \gamma_1, \gamma_2, \gamma_3 \rangle = \Gamma_3^+(2) \rightarrow \Delta_3(2, 4, 6) \rightarrow \mathfrak{S}_4 \times \mathbb{Z}/2\mathbb{Z} = \text{Aut}(\mathcal{X}^+(2)) \rightarrow 1.$$

This extension is described by the outer action of $\mathfrak{S}_4 \times \mathbb{Z}/2\mathbb{Z}$ on $\langle \gamma_1, \gamma_2, \gamma_3 \rangle$. Similarly, $\Delta_2(2, 4, 6)$ is an extension of a group of order 72, $\text{Aut}(\mathcal{X}^+(3))$, by a free group of rank four. In order to describe these extensions, we shall use F. Herrlich’s work on automorphisms of Mumford curves.

6.4.2. Let S be a Mumford curve over some finite extension K of \mathbb{Q}_p , and let $\Gamma \subset PGL_2(K)$ be the corresponding Schottky group. It is known that any automorphism of S lifts to $PGL_2(K)$: in fact, the normalizer $N(\Gamma)$ of Γ in sits $PGL_2(K)$ in an exact sequence [GvdP80, IV]:


$$1 \rightarrow \Gamma \rightarrow N(\Gamma) \rightarrow \text{Aut } S \rightarrow 1.$$

Herrlich constructs a tree \mathcal{T} endowed with an action of $N(\Gamma)$ as follows. One introduces a partition of the set of affinoid disks in $\mathbb{P}^1(K)$ into classes

⁽³⁴⁾to deal with the diadic uniformization of $\mathcal{X}^+(2)$ itself would involve a certain finite étale covering of degree 12 of Ω .

of the following form: a class contains a unique disk in $\mathbb{A}^1(K)$, of radius denoted by r and all affinoid disks of the form $|z - a| \geq r$. The vertices of \mathcal{T} are the classes of those disks which contain two fixpoints of some non-trivial element of $N(\Gamma)$, whose distance is the radius of the disk. Two vertices are related by an edge if they can be represented by disks which are either both maximal, or nested in such a way that one is a maximal proper subdisk of the other. Herrlich shows that the natural action of $N(\Gamma)$ on affinoid disks induce an action without inversion on \mathcal{T} . This allows to apply the Bass-Serre theory of graphs of groups [Ser77], exploiting the fact that the stabilizer of vertices or edges belong to the short list of finite subgroups of $PGL_2(K)$ [Her80a] [Her80b]. When $S/\text{Aut}(S)$ has genus 0, the quotient $\mathcal{T}/N(\Gamma)$ is a finite tree of groups. The group $N(\Gamma)$ can then be identified with its fundamental group, which is an amalgamated sum of the stabilizers of the vertices along the stabilizers of the edges.

6.4.3. Let us apply this to $S = \mathcal{X}^+(2)_{\mathbb{Q}_9}$ with $N(\Gamma) = \Delta_3(2, 4, 6)$. One has $(g-1)/\sharp(\text{Aut}(S)) = \frac{1}{24}$, and an inspection of the list of [Her80a] leaves only one possibility for the tree of groups $\mathcal{T}_3/\Delta_3(2, 4, 6)$, namely:



where \mathfrak{D}_n are the dihedral groups. In this case, the Herrlich tree \mathcal{T}_3 may be considered as a triadic analogue of the Schwarz-Escher tessellation:




FIGURE 10

6.4.4. In the case of $S = \mathcal{X}^+(3)_{\mathbb{Q}_4}$ with $N(\Gamma) = \Delta_2(2, 4, 6)$, one still has $(g-1)/\sharp(\text{Aut}(S)) = \frac{1}{24}$, and an inspection of the list of [Her80a] leaves only two possibilities for the tree of groups $\mathcal{T}_2/\Delta_2(2, 4, 6)$, namely:

$$\mathfrak{D}_6 \bullet \text{---} \circ \mathfrak{D}_4$$

\mathfrak{D}_2

and

$$\mathfrak{S}_4 \bullet \text{---} \circ \mathfrak{S}_4$$

\mathfrak{D}_4

In order to select the right one, we remark that $\Delta_2(2, 4, 6)$ contains the finite group generated by ij and the quotient of $(\mathcal{B}_{3,\infty})^\times = \{\pm 1, \pm i, \frac{\pm i \pm j}{2}, \frac{\pm 1 \pm ij}{2}\}$ by ± 1 . This group is isomorphic to $\mathfrak{D}_6 \cong \mathbb{Z}/2\mathbb{Z} \times \mathfrak{D}_3$, which is not contained in \mathfrak{S}_4 . Since any finite subgroup of the amalgamated sum of two groups is conjugated to a subgroup of one of these groups [Ser77, 4.3], we conclude that $\Delta_2(2, 4, 6) \cong \mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$, not $\mathfrak{S}_4 *_{\mathfrak{D}_4} \mathfrak{S}_4$

The Herrlich tree \mathcal{T}_2 may be considered as a diadic analogue of the Schwarz-Escher tessellation:




FIGURE 11

(this is the set of points u such that $j(\frac{e^{\pi i/3} - u}{1 - e^{\pi i/3} u}) \in [0, 1728]$, where j denotes the modular invariant).

It is important to pay attention to the fact that the symbol $\mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$ is not free from ambiguity: namely, there are essentially two non-equivalent ways of embedding \mathfrak{D}_2 into $\mathfrak{D}_6 \times \mathfrak{D}_4$:

1st way) the centers of \mathfrak{D}_6 and \mathfrak{D}_4 restrict to the same subgroup $\mathbb{Z}/2\mathbb{Z}$ of \mathfrak{D}_2 . On identifying \mathfrak{D}_6 and \mathfrak{D}_4 with the matrix groups

$$\left\langle \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix} \right\rangle \text{ and } \left\langle \begin{pmatrix} 0 & 1 \\ \pm 1 & 0 \end{pmatrix} \right\rangle$$

respectively, the corresponding amalgam $\mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$ is isomorphic to $GL_2(\mathbb{Z})$;

2nd way) the trace of the center of $\mathfrak{D}_6 \times \mathfrak{D}_4$ on \mathfrak{D}_2 is \mathfrak{D}_2 itself. The corresponding amalgam $\mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$ maps onto a semidirect product of $(\mathbb{Z}/3\mathbb{Z})^2$ and \mathfrak{D}_4 (the latter acting on $(\mathbb{Z}/3\mathbb{Z})^2$ as a matrix group $\langle \left(\begin{smallmatrix} 0 & 1 \\ \pm 1 & 0 \end{smallmatrix} \right) \rangle$).

The following theorem summaries our results and settles the remaining ambiguity:

Theorem 6.4.5. (i) $\Delta_3(2, 4, 6) \cong \mathfrak{D}_6 *_{\mathfrak{D}_3} \mathfrak{S}_4$.
(ii) $\Delta_2(2, 4, 6) \cong \mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$, where the embedding of \mathfrak{D}_2 into $\mathfrak{D}_6 \times \mathfrak{D}_4$ is such that the trace of the center of $\mathfrak{D}_6 \times \mathfrak{D}_4$ on \mathfrak{D}_2 is \mathfrak{D}_2 itself.

PROOF. Only the last assertion remains to be established. Let \mathcal{S} be the diadic Schwarz orbifold $(\mathbb{P}^1; (0, 2), (1, 4), (\infty, 6))$. There are surjective homomorphisms (3.5.5, 4.3.3)

$$\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \rightarrow \pi_1^{\text{orb}}(\mathcal{S}) \rightarrow \Delta_2(2, 4, 6).$$

The images $\bar{\gamma}_0, \bar{\gamma}_1, \bar{\gamma}_\infty \in \Delta_2(2, 4, 6)$ of local monodromy elements $\gamma_0, \gamma_1, \gamma_\infty \in \pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\})$ are elements of order 2, 4, 6 respectively. By 2.3.9, the smallest closed normal subgroup of $\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\})$ containing γ_1, γ_∞ is $\pi_1^{\text{temp}}(\mathbb{P}^1 \setminus \{0, 1, \infty\})$ itself. It follows that the smallest normal subgroup of the discrete group $\Delta_2(2, 4, 6)$ containing $\bar{\gamma}_1, \bar{\gamma}_\infty$ is $\Delta_2(2, 4, 6)$ itself. It thus suffices to show that if we were in the case $\mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4 \cong GL_2(\mathbb{Z})$, then $\bar{\gamma}_1, \bar{\gamma}_\infty \in SL_2(\mathbb{Z})$. This is clear, since any element of order 6 (resp. 4) in $GL_2(\mathbb{Z})$ is conjugate to $\pm \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$ (resp. to $\pm \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$). \square

Part (ii) of this theorem allows us to answer at last question 2.3.11 concerning local monodromies $\gamma_0, \gamma_1, \gamma_\infty$ at 0, 1, ∞ in the temperate fundamental group $\pi_1^{\text{temp}}(\mathbb{P}_{\mathbb{C}_p}^1 \setminus \{0, 1, \infty\})$:

Corollary 6.4.6. *Question 2.3.11 has a negative answer for $p = 2$: one cannot find local monodromies $\gamma_0, \gamma_1, \gamma_\infty$ such that $\gamma_0 \cdot \gamma_1 \cdot \gamma_\infty = 1$.*

PROOF. Assume that such local monodromies exist. By means of their images $\bar{\gamma}_0, \bar{\gamma}_1, \bar{\gamma}_\infty \in \Delta_2(2, 4, 6) \cong \mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$ (notation of the previous proof), it would be possible to construct a homomorphism

$$\phi : \Delta(2, 4, 6) \rightarrow \Delta_2(2, 4, 6),$$

where $\Delta(2, 4, 6)$ denotes the standard (archimedean) triangle group. Notice that the image of ϕ cannot be contained in any conjugate of \mathfrak{D}_6 or \mathfrak{D}_4 , since none of these finite groups contains elements of order 6 and of order 4 simultaneously.

According to [Ser77, 6.3.5], any (cocompact) triangle group Δ has the fixpoint property (FA): any action (without inversion) of Δ on any tree has a fixpoint. It follows that the quotient $\text{Im } \phi$ of $\Delta(2, 4, 6)$ would have property (FA) as well. Since $\text{Im } \phi$ is a subgroup of the amalgam $\mathfrak{D}_6 *_{\mathfrak{D}_2} \mathfrak{D}_4$,

this would imply, according to [Ser77, 6.2, prop. 21], that $Im \phi$ is contained in a conjugate of \mathfrak{D}_6 or \mathfrak{D}_4 , a contradiction. \square

Remark 6.4.7. The arithmetic p -adic triangle groups $\Delta(e_0, e_1, e_\infty)$ do *not* have property (FA). Indeed, their description as p -unit groups in quaternion algebras shows that they admit a two-dimensional representation for which the eigenvalues are non-integral in general. This property contradicts (FA), cf. [Ser77, 6.2, prop. 22]. Actually, Kato's description of p -adic triangle groups of Mumford type as amalgams, together with 5.3.7, implies that *no hyperbolic p -adic triangle group has property (FA)* (in contrast to archimedean triangle groups). Hence no hyperbolic p -adic triangle group can be generated by three elements g_0, g_1, g_∞ of finite order satisfying $g_0 g_1 g_\infty = 1$.

6.5. $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ and the period mapping. Global analytic triadic continuation.

6.5.1. Rather than $\Delta(2, 4, 8)$, we now prefer to work with $\Delta(6, 2, 4)$ (this amounts to permute $0, 1, \infty$ by $t \mapsto z = \frac{1}{1-t}$, and is more faithful to the first picture of this section which places the vertex O at the center of the disk).

The triangle group $\Delta(6, 2, 4)$ is the projective monodromy group of the hypergeometric differential operator $L_{\frac{1}{24}, \frac{7}{24}, \frac{5}{6}}$. More precisely, $L_{\frac{1}{24}, \frac{7}{24}, \frac{5}{6}}(y) = 0$ is a *uniformizing differential equation* for the orbifold \mathcal{X}^* viewed as the Schwarz orbifold

$$\left(\mathbb{P}^1, \begin{array}{ccc} 0 & 1 & \infty \\ e_0 = 6 & e_1 = 2 & e_\infty = 4 \end{array} \right)$$

More explicitly, $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ and $z^{1/6} F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)$ are independent solutions of $L_{\frac{1}{24}, \frac{7}{24}, \frac{5}{6}}$ and for a suitable constant κ , the mapping

$$\tau(z) = \kappa^{-1} \cdot \frac{z^{1/6} F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)}{F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)}$$

sends the upper half plane bijectively to the basic triangle OQP of the Schwarz tessellation, and $\tau(0) = O$, $\tau(1) = Q$, $\tau(\infty) = P$.




FIGURE 12

It extends to a multivalued analytic function on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ with value in $\mathcal{D}(0, 1^-)$. The inverse mapping $z(\tau) : \mathcal{D}(0, 1^-) \rightarrow \mathbb{P}^1$ is the (single-valued) $\Delta(2, 4, 6)$ -automorphic uniformizing mapping of the Shimura orbifold \mathcal{X}^* .

6.5.2. Actually, τ expresses the period mapping \mathcal{P} for “fake elliptic curves” with multiplication by $\mathcal{B}_{2,3}$, *cf.* II.1.2 and II.7.4: more precisely, the multivalued function $\tau(z)$ is related to the commutative diagram

$$\begin{array}{ccc} \widetilde{\mathcal{X}_{\mathbb{C}}^{*\text{an}}} & \xrightarrow{\overset{P}{\sim}} & \mathcal{D} = \mathcal{D}(0, 1^-) \subset \mathcal{D}^{\vee} = \mathbb{P}^1 \\ \mathcal{Q} \downarrow & & \downarrow \\ \left(\mathbb{P}^1, \begin{array}{ccc} 0 & 1 & \infty \\ 6 & 2 & 4 \end{array} \right) = \mathcal{X}_{\mathbb{C}}^{*\text{an}} & \longrightarrow & \mathcal{D}/\Delta(2, 4, 6) \end{array}$$

by

$$\tau(z) = \mathcal{P} \circ \mathcal{Q}^{-1}, \quad z(\tau) = \mathcal{Q} \circ \mathcal{P}^{-1}.$$

6.5.3. Using Gauss’ formula for the value of hypergeometric functions at 1, one can compute the constant κ , *cf.* [Car61, 392]:

$$\kappa = \frac{\Gamma(\frac{7}{6})}{\Gamma(\frac{5}{6})} \sqrt{\frac{\Gamma(\frac{1}{24})\Gamma(\frac{7}{24})\Gamma(\frac{13}{24})\Gamma(\frac{19}{24})}{\Gamma(\frac{5}{24})\Gamma(\frac{11}{24})\Gamma(\frac{17}{24})\Gamma(\frac{23}{24})}}.$$

Using the functional equations of Γ , one checks that $\kappa \sim \left(\frac{\Gamma(\frac{1}{3})}{\Gamma(\frac{2}{3})}\right)^3 \pmod{\overline{\mathbb{Q}}^{\times}}$, in conformity with II.7.6.4 (taking into account the fact that $z = 0$ corresponds to a fake elliptic curve with complex multiplication by an order in $M_2(\mathbb{Q}(\sqrt{-3}))$).

6.5.4. For $p \neq 2, 3$, the p -adic radius of convergence of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ and $F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)$ is 1. The unit disk $|z|_p < 1$ is ordinary or supersingular according to whether $p \equiv 1 \pmod{3}$ or not. The ratio $\frac{z^{1/6} F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)}{F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)}$ expresses a local p -adic period mapping.

The p -divisible group of the fake elliptic curve parametrized by z is isogenous to the square of the p -divisible group of an elliptic curve; the period mapping is thus completely analogous to one of those studied in II.6.3.1, 6.3.3 (Dwork, Gross-Hopkins). From the viewpoint of analytic continuation, this is essentially the situation studied in I.3 (for $F(\frac{1}{2}, \frac{1}{2}, 1; z)$).

6.5.5. For $p = 3$ (*resp.* $p = 2$), the situation is radically different; the p -adic radius of convergence of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ and $F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)$ is $3^{-3/2}$ (*resp.* 2^{-6}).

The ratio $\frac{z^{1/6} F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)}{F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)}$ expresses the Drinfeld 3-adic period mapping:

$$\begin{array}{ccc}
\widetilde{\mathcal{X}^+(2)_{\mathbb{C}_3}^{\text{an}}} & \xrightarrow[\sim]{\mathcal{P}} & \mathcal{D} \subset \mathcal{D}^\vee = \mathbb{P}^1 \\
\downarrow & & \downarrow \\
\mathcal{X}^+(2)_{\mathbb{C}_3}^{\text{an}} & & \\
\downarrow & & \\
\mathcal{X}_{\mathbb{C}_3}^{*\text{an}2} & \longrightarrow & \mathcal{D}/\Delta_3(2, 4, 6)
\end{array}$$

Here \mathcal{D} is the image of the Drinfeld space $\Omega_{\mathbb{C}_3}$ by a suitable homography (it would be interesting to determine this homography explicitly). That this homography is actually defined over $\overline{\mathbb{Q}}$ comes from II.7.6.4 (ramified variant), I.4.6.4, 5.3.9 (the point is that the triadic periods, in the sense of I.5.3, of the abelian surface with complex multiplication by $M_2(\mathbb{Q}[\sqrt{-3}])$ parametrized by $z = 0$, belong to $\overline{\mathbb{Q}}$).

Although $F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)/F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ has very small triadic radius of convergence, its pull-back on the Shimura curve $\mathcal{X}^+(2)$ extend to a global multivalued meromorphic function, with poles only above $0, 1, \infty$ (in other words, its pull-back on $\widetilde{\mathcal{X}^+(2)_{\mathbb{C}_3}} \cong \Omega_{\mathbb{C}_3}$ is single-valued).

Note however that the same is not true for $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ and $F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)$ individually, for reasons of exponents at ∞ . Nevertheless, they enjoy a similar property, on replacing $\mathcal{X}^+(2)$ by a suitable ramified finite covering.

Similarly, in the diadic case: the pull-back of $F(\frac{5}{24}, \frac{11}{24}, \frac{7}{6}; z)/F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ on the Shimura curve $\mathcal{X}^+(3)$ extends to a global multivalued meromorphic functions (with poles only above $0, 1, \infty$).

6.6. A triadic analogue of Gauss' formula, and special values of $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ at CM points.

6.6.1. For our hypergeometric function, Gauss' formula reads

$$F\left(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; 1\right) = \frac{\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})}{\Gamma(\frac{19}{24})\Gamma(\frac{13}{24})}.$$

If one considers $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)$ as a multivalued analytic function on $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, the values at 1 generate a $\overline{\mathbb{Q}}$ -subspace of \mathbb{C} of dimension ≤ 2 , namely:

$$\frac{\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})}{\Gamma(\frac{19}{24})\Gamma(\frac{13}{24})}\overline{\mathbb{Q}} + \frac{\Gamma(\frac{5}{6})\Gamma(-\frac{1}{2})}{\Gamma(\frac{1}{24})\Gamma(\frac{7}{24})}\overline{\mathbb{Q}}.$$

Similarly, its values at ∞ generate the $\overline{\mathbb{Q}}$ -subspace

$$\frac{\Gamma(\frac{5}{6})\Gamma(\frac{1}{4})}{\Gamma(\frac{19}{24})\Gamma(\frac{7}{24})}\overline{\mathbb{Q}} + \frac{\Gamma(\frac{5}{6})\Gamma(-\frac{1}{4})}{\Gamma(\frac{1}{24})\Gamma(\frac{13}{24})}\overline{\mathbb{Q}}.$$

6.6.2. Let us consider our hypergeometric function as a triadic function $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; 1)_3$, and more precisely as a multivalued 3-adic analytic function on a suitable finite étale covering of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$.

The values taken by $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)_3$ at 1 belong to $\overline{\mathbb{Q}}$: this follows from 5.6.3, together with the fact that the “periods” (in the sense of I.5.3) attached to the abelian surface with complex multiplication by $M_2(\mathbb{Q}[\sqrt{-24}])$ parametrized by $z = 1$, belong to $\overline{\mathbb{Q}}$, by I.4.6.4, I.5.3.9.

In the same way, one shows that the values taken by $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; z)_3$ at ∞ belong to $(\Gamma_3(\frac{3}{4})/\Gamma_3(\frac{1}{4}))^{-1/2}\overline{\mathbb{Q}} = \Gamma_3(\frac{1}{4})\overline{\mathbb{Q}}$. Conjecture I.5.3.10 predicts that $\Gamma_3(\frac{1}{4})$ is a transcendental number.

6.6.3. In [Elk98, Table 2], one finds a table of rational CM points z (rational points which correspond to abelian surfaces with complex multiplication by an order in $M_2(\mathbb{Q}(\sqrt{-d}))$ for some fundamental discriminant $-d$); with the notation of *loc. cit.*, they are given by $z = B/(B - A) = \pm B/|C|$. Some of them are:

z	$d = -\text{disc}$
$-3^3/2^27^2$	$d = 2^2 \cdot 3 \cdot 7$
$-5^3/3^7$	$d = 2^3 \cdot 5$
$2^{10}/7^4$	$d = 3 \cdot 17$
$-2^{10}/3^7$	$d = 19$.

Notice that for the first value of z , $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; \frac{-3^3}{2^27^2})_3$ converges. Moreover, since $(\frac{2^2 \cdot 7}{3}) = +1$, it follows from I.4.6.4, I.5.3.9, that the triadic “periods” attached to the abelian surface $A_{-3^3/2^27^2}$ with complex multiplication by $M_2(\mathbb{Q}[\sqrt{-d}])$ parametrized by $z = 1$, is algebraic (like the periods of A_0). By 5.6.3, it follows that the triadic number $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; \frac{-3^3}{2^27^2})_3$ is algebraic (can one compute this algebraic number?).

On the other hand since $\mathbb{Q}(\sqrt{-d}) \neq \mathbb{Q}(\sqrt{-3})$ in that case ($d = 2^2 \cdot 3 \cdot 7$), the real number $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; \frac{-3^3}{2^27^2})$ is transcendental; this follows from Wolfart’s theorem [Wol88] (whose proof has been recently completed by work of Cohen-Wüstholz [CW01] and Edixhoven-Yafaev).

It can be proved, a contrario, that if the complex evaluation $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; \xi)$ is algebraic for some algebraic value ξ in the disc of convergence, then any p -adic evaluation $F(\frac{1}{24}, \frac{7}{24}, \frac{5}{6}; \xi)_p$ which makes sense is also algebraic: indeed, by [Wol88], [CW01], such a ξ parametrizes a CM abelian surface by some order in $M_2(\mathbb{Q}[\sqrt{-3}])$, and one concludes again using p -adic Betti lattices as in I.5.3.

This is actually a very special case of a general, but conjectural, “principle of global relations” for special values of solutions of Gauss-Manin connections, *cf.* [And95].

APPENDIX A

Rapid Course in p -adic Analysis.

By F. KATO

1. Introduction.

In this appendix, K always denotes a complete field with respect to a non-archimedean valuation $|\cdot|: K \rightarrow \mathbb{R}_{\geq 0}$. The norm $|\cdot|$ is almost always assumed to be non-trivial, unless otherwise stated.

Let us assume for a while that K is algebraically closed. We view the field K as an affine space, as we do in complex analysis; K is a metrized space with the metric induced from the valuation $|\cdot|$. We can then follow the lines of classical complex analysis and define convergence of power series, Taylor and Laurent expansions, etc...

For instance let us define “holomorphic” functions to be K -valued functions which are locally expressed by convergent power series⁽¹⁾ (cf. [Gou97, Chap. 4]). This approach drives us however to several problems, which come mainly from the fact that K with the metric topology is totally disconnected; for example:

- (i) for an open set $U \subseteq K$ the ring of all such holomorphic functions on U is huge. So is already the subring of all locally constant functions.
- (ii) It can be shown that the sheaf of germs of such functions satisfies the principle of unique continuation; but not in a satisfactory way. In particular, Lemma 1.1.2 of Chapter I holds but is of no use, because there is no non-empty connected open subset.

The trouble becomes more apparent when we think the *local* representability by convergent power series in terms of *coverings*. In the complex analytic situation, when we speak of the holomorphy of a function f defined over a connected open set U , we tacitly take a open covering $\{U_i\}_{i \in I}$ of U consisting of sufficiently small open neighborhoods such that f restricted on each member U_i can be seen as evaluation of a power series convergent in U_i . The point is that, since U is connected, members of $\{U_i\}$ must have overlaps so that the local properties can be transmitted to whole U . But, in the non-archimedean situation, such coverings may be refined in such a way that there are no overlap. This is why the analytic continuation does

⁽¹⁾The reader may wonder whether there is another approach by means of Cauchy-Riemann type differential equations. This is unlikely because the differential calculus does not go well, see [Gou97, Chap. 4].

not work well. So, for the sake of a reasonable analytic theory, we have to “limit” coverings so that we cannot take arbitrarily fine refinements.

A first method is due to Krasner. He introduced the so-called *quasi-connected sets* and, on such subsets, defined analytic functions as uniform convergent limits of rational functions.

A more modern and systematic treatment was introduced by Tate, with the so-called *Rigid Analysis*. Let us briefly view the main idea: we first set

$$K\{t_1, \dots, t_n\} = \left\{ \sum_{\nu_1, \dots, \nu_n \geq 0} a_{\nu_1, \dots, \nu_n} t_1^{\nu_1} \cdots t_n^{\nu_n} \in K[[t_1, \dots, t_n]] \mid \begin{array}{l} a_{\nu_1, \dots, \nu_n} \rightarrow 0 \text{ for} \\ \nu_1 + \cdots + \nu_n \rightarrow \infty \end{array} \right\},$$

which is now called the *Tate algebra* over K . This is the ring of all functions expressed by power series convergent on the closed disk $D(0, 1^+) = \{z \in K^n \mid |z_i| \leq 1\}$. The algebra $K\{t_1, \dots, t_n\}$ is endowed with the sup-norm $\|\cdot\|$, also called the *Gauss norm*. We list up some known properties (cf. [BGR84, 5.2], [FvdP81, II.3]):

- (i) The ring $K\{t_1, \dots, t_n\}$ satisfies the Weierstrass Preparation Theorem.
- (ii) The ring $K\{t_1, \dots, t_n\}$ is Noetherian and factorial.
- (iii) Every ideal of $K\{t_1, \dots, t_n\}$ is closed.
- (iv) For any maximal ideal \mathfrak{m} of $K\{t_1, \dots, t_n\}$, the field $K\{t_1, \dots, t_n\}/\mathfrak{m}$ is a finite extension of K .

If, moreover, K is algebraically closed, the set of all maximal ideals, endowed with the natural topology, coincides with $D(0, 1^+)$. So the situation is analogous to classical algebraic geometry; $D(0, 1^+)$ to the affine n -space, and $K\{t_1, \dots, t_n\}$ to the coordinate ring. In fact, quotients $\mathcal{A} = K\{t_1, \dots, t_n\}/I$ by ideals I (called *affinoid algebras*), and their associated maximal spectra, $\text{Spm}(\mathcal{A})$, called *affinoids*, forms the fundamental patches from which the non-archimedean function theory will be developed. Although the natural metric topology on $\text{Spm}(\mathcal{A})$ is terrible for the same reason as above, the situation becomes much better as far as we deal only with those functions coming from \mathcal{A} . The trick is that, if the algebra \mathcal{A} is integral, then we can pretend that the space $\text{Spm}(\mathcal{A})$ is “connected”; for example, the closed disk $D(0, 1^+)$ is “connected” in this sense. So, basically according to this point of view, we can globalize the situation by gluing affinoid patches to obtain a reasonable theory of analytic functions. The actual recipe to do it is furnished by the notion of Grothendieck topology, which censors the plethora of coverings. We will see this a little more precisely in what follows.

Remark 1.1. Although the rigid analysis seems to provide the reasonable topological and analytical framework, one must not expect it to be as nice as the complex case. Firstly, in classical complex analysis, when we expand a holomorphic function centered at a point inside its region of convergence, the resulting Taylor expansion may have a different region of convergence

than the original one. This is an important method of analytic continuation. But in the non-archimedean case, this never happens (cf. [Gou97, 4.4]).

Secondly, the topological framework given as above is still not powerful enough to handle paths. So it is hopeless to treat, for example, fundamental groups and monodromy in an intuitive way as in the classical complex case. This difficulty is, in fact, remedied by Berkovich's theory which we shall see later on.

2. Rigid analytic spaces.

Let K be as in the previous section; we do not assume in general that K is algebraically closed.

As mentioned above, an *affinoid algebra* A over K is a quotient $K\{t_1, \dots, t_n\}/I$ for some n by an ideal I . Since I is closed, A has a norm $|\cdot|$ induced from the sup-norm $\|\cdot\|$ of $K\{t_1, \dots, t_n\}$ (“*residue norm*”). With the residue norm, A is a Banach K -algebra. The norm $|\cdot|$ itself depends on the presentation $\alpha: K\{t_1, \dots, t_n\}/I \xrightarrow{\sim} A$, while the induced topology does not. So, strictly speaking, we should write it like $|\cdot|_\alpha$. Clearly, A is Noetherian and, for any maximal ideal \mathfrak{m} of A , the residue field A/\mathfrak{m} is a finite extension of K , and hence, the valuation $|\cdot|$ of K naturally extends to that of it, denoted again by $|\cdot|$.

The associated *affinoid* $\mathrm{Spm}(A)$ is the set of all maximal ideals of A . For $x \in \mathrm{Spm}(A)$ and $f \in A$ the value of f at x , denoted by $f(x)$, is the class of f in A/x . The set $\mathrm{Spm}(A)$ has the topology generated by the subsets of form $\{x \in \mathrm{Spm}(A) \mid |f(x)| \leq 1\}$ for $f \in A$. But, as we pointed out above, this topology is not very interesting since it makes $\mathrm{Spm}(A)$ totally disconnected. So we should specify the reasonable family of “admissible” open sets (and coverings), on which the function theory will be built.

Let A be an affinoid algebra over K , and f_i ($i = 0, \dots, n$) a collection of elements in A which have no common zeros on $\mathrm{Spm}(A)$. The subspace

$$R = \{x \in \mathrm{Spm}(A) \mid |f_i(x)| \leq |f_0(x)|, i = 1, \dots, n\}$$

can be identified with the affinoid $\mathrm{Spm}(A_R)$, where

$$A_R = A \widehat{\otimes}_K K\{t_1, \dots, t_n\}/(f_1 - t_1 f_0, \dots, f_n - t_n f_0).$$

A subset of this form is called a *rational subdomain*. The identification comes as follows: We first note that the morphism of affinoids $\mathrm{Spm}(A_R) \rightarrow \mathrm{Spm}(A)$ induced by $A \rightarrow A_R$ maps $\mathrm{Spm}(A_R)$ to R . Then A_R is the unique solution of the following universal property (hence, it can be determined up to canonical isomorphism): for any morphism of affinoids $\phi: \mathrm{Spm}(B) \rightarrow \mathrm{Spm}(A)$ such that $\phi(\mathrm{Spm}(B)) \subset R$, there exists a unique K -homomorphism $A_R \rightarrow B$ such that the resulting diagram

$$\begin{array}{ccc}
 \mathrm{Spm}(\mathbf{B}) & \xrightarrow{\phi} & \mathrm{Spm}(\mathbf{A}) \\
 \vdots & \nearrow & \\
 & \mathrm{Spm}(\mathbf{A}_R) &
 \end{array}$$

is commutative.

Lemma 2.1. (1) If R and S are rational subdomains in $\mathrm{Spm}(\mathbf{A})$, then so is $R \cap S$.

(2) If R_1 is a rational subdomain in $\mathrm{Spm}(\mathbf{A})$, and R_2 is a rational subdomain in R_1 , then R_2 is a rational subdomain in $\mathrm{Spm}(\mathbf{A})$.

PROOF. (1) If $R = \{|f_i(x)| \leq |f_0(x)|, i = 1, \dots, n\}$ and $S = \{|g_j(x)| \leq |g_0(x)|, j = 1, \dots, m\}$, then one sees easily that $R \cap S = \{|f_i(x)g_j(x)| \leq |f_0(x)g_0(x)|, i = 1, \dots, n, j = 1, \dots, m\}$.

(2) Exercise. \square

Corollary 2.2. Any subspace of the form

$$\{|f_i(x)| \leq 1, |g_j(x)| \geq 1, i = 1, \dots, n, j = 1, \dots, m\}$$

is a rational subdomain.

PROOF. It is the intersection of rational subdomains $\{|f_i(x)| \leq 1\}$ for $i = 1, \dots, n$ and $\{1 \leq |g_j(x)|\}$ for $j = 1, \dots, m$. \square

Example 2.3 (Rational subdomains of the unit polydisk). Let us assume that K is algebraically closed. We consider rational subdomains of the unit polydisk $\mathbf{D}(0, 1^+) = \mathrm{Spm}(K\{t_1, \dots, t_n\})$.

(1) *Closed polydisk:* $\mathbf{D}(0, |\pi|^+) = \{x \in K^n \mid |x_i| \leq |\pi_i|, i = 1, \dots, n\}$, for $0 \neq \pi_i \in K, |\pi_i| \leq 1$, is a rational subdomain. The corresponding affinoid algebra is

$$K \left\{ \frac{t_1}{\pi_1}, \dots, \frac{t_n}{\pi_n} \right\}.$$

(2) *Annulus:* $\mathbf{C}(0, |\pi^{(1)}|^+, |\pi^{(2)}|^+) = \{x \in K^n \mid |\pi_i^{(1)}| \leq |x_i| \leq |\pi_i^{(2)}|, i = 1, \dots, n\}$, for $0 \neq \pi_i^{(j)} \in K, |\pi_i^{(j)}| \leq 1$, is a rational subdomain. The corresponding affinoid algebra is

$$K \left\{ \frac{\pi_1^{(1)}}{t_1}, \dots, \frac{\pi_n^{(1)}}{t_n}, \frac{t_1}{\pi_1^{(2)}}, \dots, \frac{t_n}{\pi_n^{(2)}} \right\}.$$

Let us next review the definition of Grothendieck topology (in a narrow sense). Let X be a topological space. A *Grothendieck topology* (G-topology, in short) on X is a pair $(\mathcal{T}, \mathcal{Cov})$ consisting of

- a collection \mathcal{T} of open subsets in X ,
- an assignment $U \mapsto \mathcal{Cov}(U)$ for any $U \in \mathcal{T}$, where $\mathcal{Cov}(U)$ is a collection of coverings by elements in \mathcal{T} ,

such that the following conditions are satisfied:

- (1) $\emptyset \in \mathcal{T}; U, V \in \mathcal{T} \Rightarrow U \cap V \in \mathcal{T}$.
- (2) $U \in \mathcal{T} \Rightarrow \{U\} \in \mathcal{Cov}(U)$.
- (3) $\{U_i\}_{i \in I} \in \mathcal{Cov}(U), V \subseteq U, V \in \mathcal{T} \Rightarrow \{U_i \cap V\}_{i \in I} \in \mathcal{Cov}(V)$.
- (4) $\{U_i\}_{i \in I} \in \mathcal{Cov}(U), \{U_{i,j}\}_{j \in J_i} \in \mathcal{Cov}(U_i) \Rightarrow \{U_{i,j}\}_{i \in I, j \in J} \in \mathcal{Cov}(U)$.

Elements in \mathcal{T} are called *admissible open sets*, and elements in $\mathcal{Cov}(U)$ are called *admissible coverings* of U .

Let $X = \text{Spm}(\mathbb{A})$ be an affinoid over K . We introduce a Grothendieck topology to X by the following recipe:

- An admissible open set is a rational subdomain.
- For a rational subdomain, an admissible covering is a finite covering consisting of rational subdomains.

This is valid, due to Lemma 2.1.

We define the structure presheaf \mathcal{O}_X , with respect to the Grothendieck topology, by assigning for each rational subdomain R the corresponding affinoid algebra A_R . Due to the following weak form of *Tate's acyclicity theorem*, the presheaf \mathcal{O}_X is, in fact, a sheaf (a sheaf of local rings).

Theorem 2.4. Let U_1, \dots, U_m be rational subdomains of $X = \text{Spm}(\mathbb{A})$. Set $U = U_1 \cup \dots \cup U_m$. Then the sequence

$$0 \longrightarrow \mathcal{O}_X(U) \longrightarrow \prod_{i=1}^m \mathcal{O}_X(U_i) \longrightarrow \prod_{i,j=1}^m \mathcal{O}_X(U_i \cap U_j)$$

is exact, where the last arrow is the difference of the two possible restriction morphisms.

For the proof, see, for example, [BGR84, 8.2] or [FvdP81, III.2.2].

Definition 2.5 (Rigid Analytic Space). A *rigid analytic space* is a locally ringed space (X, \mathcal{O}_X) , with a Grothendieck topology in the above sense, locally isomorphic to an affinoid; more precisely, there exists a covering $\{X_i\}_{i \in I}$ (possibly infinite) of X by admissible open sets such that $(X_i, \mathcal{O}_X|_{X_i})$ is isomorphic to a certain affinoid for each i .

Let us see some examples of rigid analytic spaces:

Example 2.6 (Projective Space). The projective space $\mathbb{P}^n(K)$ has the natural rigid analytic structure. Here we limit ourselves to demonstrate it only in the case $n = 1$, and leave the general case to the reader. Let $(X : Y)$ be the homogeneous coordinate in $\mathbb{P}^1(K)$ and set $z = X/Y$. Set

$$\mathbf{U}^+ = \{(X : Y) \mid |X| \leq |Y|\} \text{ and } \mathbf{U}^- = \{(X : Y) \mid |X| \geq |Y|\}.$$

Then \mathbf{U}^+ is isomorphic to the closed disk $\{z \in K \mid |z| \leq 1\}$ (the Southern Hemisphere with the equator), and \mathbf{U}^- to $\{z \in K \cup \{\infty\} \mid |z| \geq 1\}$ (the Northern Hemisphere with the equator). The intersection $\mathbf{U}^+ \cap \mathbf{U}^-$ is thus

isomorphic to the circle $\{z \in K \mid |z| = 1\}$. Hence $\{\mathbf{U}^+, \mathbf{U}^-\}$ gives the admissible covering of $\mathbb{P}^1(K)$ which induces, in the obvious way, the structure of a rigid analytic space.

More generally, an algebraic variety, separated and of finite type over K , carries the canonical structure of a rigid analytic space.

Any open set (not necessarily admissible) U in a rigid analytic space X will be again a rigid analytic space (“*open subanalytic space*”); more precisely, U has the induced analytic structure from that of X . The admissible open sets of U , for example, are those of X contained in U .

3. Relation with Formal Geometry.

In this section we need to consider the valuation ring R of K . Let us denote by \mathfrak{m} the maximal ideal of R , and fix a topological generator $0 \neq \pi \in \mathfrak{m}$. The residue field R/\mathfrak{m} is denoted by k .

Let A be an affinoid algebra over K . The *spectral semi-norm* on A is the function $|\cdot|_{\text{sup}}: A \rightarrow \mathbb{R}_{\geq 0}$ defined, for any $f \in A$, by

$$|f|_{\text{sup}} = \sup_{x \in \text{Spm}(A)} |f(x)|.$$

It is known that for any representation $\alpha: K\{t_1, \dots, t_n\}/I \xrightarrow{\sim} A$, we have $|f|_{\text{sup}} \leq |f|_{\alpha}$ for every $f \in A$ (cf. [BGR84, 6.2.1]).

Theorem 3.1 (Maximal modulus principle). For an affinoid algebra A and an element $f \in A$, there exists a point $x \in \text{Spm}(A)$ such that $|f(x)| = |f|_{\text{sup}}$.

We refer [BGR84, 6.2.1] for the proof.

Given an affinoid algebra A over K , we introduce the following notation:

$$\begin{aligned} A^\circ &= \{f \in A \mid |f|_{\text{sup}} \leq 1\}, \\ A^{\circ\circ} &= \{f \in A \mid |f|_{\text{sup}} < 1\}, \\ \overline{A} &= A^\circ/A^{\circ\circ}; \end{aligned}$$

A° is a R -subalgebra of A , and $A^{\circ\circ}$ is an ideal of it. For instance, $K^\circ = R$, $K^{\circ\circ} = \mathfrak{m}$, and $\overline{K} = k$.

Example 3.2. We can immediately calculate:

$$\begin{aligned} \mathbb{T}_K^{n,\circ} &= R\{t_1, \dots, t_n\} \\ &: = \left\{ \sum_{\nu_1, \dots, \nu_n \geq 0} a_{\nu_1, \dots, \nu_n} t_1^{\nu_1} \cdots t_n^{\nu_n} \mid \begin{array}{l} |a_{\nu_1, \dots, \nu_n}| \rightarrow 0 \text{ for} \\ \nu_1 + \cdots + \nu_n \rightarrow \infty \end{array} \right\} \\ \overline{\mathbb{T}_K^n} &= k[t_1, \dots, t_n]. \end{aligned}$$

Note that $\mathbb{T}_K^{n,\circ}$ is the (π) -adic completion of the polynomial ring $R[t_1, \dots, t_n]$ over R . It should be noticed that $\overline{\mathbb{T}_K^n}$ is a *polynomial ring* over k ; indeed, each element in $R\{t_1, \dots, t_n\}$ lies, by the convergence condition, in $\mathbb{T}_K^{n,\circ}$ modulo finitely many terms.

Proposition 3.3. Let A be an affinoid algebra over K . Then,

- (i) A° is a model of A ; i.e. $A^\circ \otimes_R K \cong A$,

- (ii) A° is (π) -adically completed,
- (iii) A° is topologically of finite type over R ; i.e. $R\{t_1, \dots, t_n\}/\mathfrak{a} \cong A$ for some ideal \mathfrak{a} in $R\{t_1, \dots, t_n\}$,
- (iv) A° is flat over R ,
- (v) \overline{A} is a k -algebra of finite type.

PROOF. (i) and (ii) are easy to see. The flatness of R is now equivalent to the lack of R -torsion. So (iv) can be seen immediately.

Before proving (iii) and (v), we need some terminology: let $\|\cdot\|_\alpha$ be a Banach norm of A ($\alpha: T_K^n \rightarrow A$: surjective). We say an element f of A is *power-bounded* if $\{\|f^n\|_\alpha \mid n \in \mathbb{N}\}$ is bounded. This does not depend on the choice of the presentation α since the equivalence class of the resulting norms do not change. An element f of A is said to be *topologically nilpotent* if $\lim f^n = 0$; here the limit is taken with respect to the Banach norm $\|\cdot\|_\alpha$, and is not dependent of the choice.

Let us prove (iii). By [BGR84, 6.2.3], A° is the set of all power-bounded elements of A , and $A^{\circ\circ}$ is the set of all topologically nilpotent elements. Hence a choice of a presentation $\alpha: T_K^n \rightarrow A$ induces surjections

$$\begin{aligned} T_K^{n,\circ} &\longrightarrow A^\circ \\ T_K^{n,\circ\circ} &\longrightarrow A^{\circ\circ}. \end{aligned}$$

(iii) is due to the surjectivity of the first arrow. By the surjectivity of these arrows and 3.2, we have (v). □

Definition 3.4. An R -algebra A is said to be *admissible* if A is (π) -adically completed, flat over R , and topologically of finite type over R .

Remark 3.5. By [BL93, 1.1 (c)], any admissible R -algebra A is topologically of finite presentation, i.e., there exists a finitely generated ideal \mathfrak{a} in $R\{t_1, \dots, t_n\}$ such that $R\{t_1, \dots, t_n\}/\mathfrak{a} \cong A$.

Definition 3.6 (Formal model and Analytic reduction). Let $X = \text{Spm}A$ be an affinoid over K . Then the formal scheme $\mathcal{X} = \text{Spf} A^\circ$ over R is called the *formal model* of X , and the algebraic scheme $\overline{X} = \text{Spec} \overline{A}$ over k is called the *analytic reduction* of X .

We define the so-called *reduction map*

$$\text{Red}_X: X = \text{Spm}A \longrightarrow \overline{X} = \text{Spec} \overline{A}$$

by $\mathfrak{a} \mapsto \mathfrak{a} \cap A^\circ / \mathfrak{a} \cap A^{\circ\circ}$.

Proposition 3.7. The map Red_X is continuous with respect to the Zariski topology of \overline{X} and the G -topology of X ; it maps X surjectively onto the set of all closed points of \overline{X} .

PROOF. Let us consider the affine open set $U_{\overline{f}} = \text{Spec} \overline{A}_{\overline{f}}$ of $\text{Spec} \overline{A}$ relevant to $\overline{f} \in \overline{A}$. Take $f \in A^\circ$ from the residue class \overline{f} . The function f takes values less than equal to 1 on $\text{Spm}A$. So $U_f = \{x \in \text{Spm}A \mid |f(x)| = 1\}$ is a rational subdomain whose associated affinoid is $\text{Spm}A\{f, f^{-1}\}$. Since the condition

$|f(x)| = 1$ is equivalent to $\bar{f}(\text{Red}_X(x)) \neq 0$, we have $\text{Red}_X^{-1}(U_{\bar{f}}) = U_f$. Then the first assertion follows. The last one is almost trivial. \square

Example 3.8 (Analytic reduction of the unit disk). Let us consider the analytic reduction of the unit disk $D^1 = D(0, 1^+) = \text{Spm}K\{T\}$ (here we assume that K is algebraically closed for simplicity). The analytic reduction is given by the affine line $\mathbb{A}_k^1 = \text{Spec}k[T]$. The reduction map Red_{D^1} maps all “interior” points, i.e. those points lying in $D(0, 1^-)$ to the origin in \mathbb{A}_k^1 , and the other points to the non-zero points. If we identify D^1 and R , \mathbb{A}_k^1 and k , then this is nothing but the mapping obtained by the reduction modulo \mathfrak{m} .

It is very important to see how the formal models behave under localization of affinoids. Let $X = \text{Spm}A$ be an affinoid over K and $f_0, \dots, f_n \in A$ a sequence of elements which does not have common zeros on $\text{Spm}A$. We consider the admissible covering $X = \bigcup_i U_i$ consisting of rational subdomains

$$U_i = \{x \in X \mid |f_j(x)| \leq |f_i(x)| \text{ for } j \neq i\};$$

by multiplying a suitable power of π , we may assume that each f_i belongs to A° . For each i ,

$$U_i = \text{Spm}A \left\{ \frac{f_0}{f_i}, \dots, \frac{f_n}{f_i} \right\},$$

and then the corresponding formal model is given by

$$\mathcal{U}_i = \text{Spf} A^\circ \left\{ \frac{f_0}{f_i}, \dots, \frac{f_n}{f_i} \right\} / ((\pi)\text{-torsions}).$$

These formal schemes glue and the resulting morphism

$$\bigcup_{0 \leq i \leq n} \mathcal{U}_i \longrightarrow \mathcal{X} = \text{Spf} A^\circ$$

is the formal blow-up along the ideal $\mathfrak{a} = (f_0, \dots, f_n) \subset A^\circ$. We note here that the ideal \mathfrak{a} contains a power of π , or what amounts to the same, \mathfrak{a} is an open ideal, for $\mathfrak{a}A = A$.

This simple observation indicates:

Slogan (vague): *Refinements of admissible coverings corresponds to formal blow-up's along open coherent ideals.*

Example 3.9. Let us assume in this example that K is algebraically closed. We consider the admissible covering $D^1 = U_1 \cup U_2$ of the unit disk $D^1 = \text{Spm}K\{T\}$ given by

$$\begin{aligned} U_1 &= \{z \in K \mid |z| \leq |\pi|\} &= \text{Spm}K \left\{ \frac{T}{\pi} \right\}, \\ U_2 &= \{z \in K \mid |\pi| \leq |z| \leq 1\} &= \text{Spm}K \left\{ T, \frac{\pi}{T} \right\}. \end{aligned}$$

The corresponding formal models are

$$\begin{aligned} \mathcal{U}_1 &= \text{Spf} R \left\{ \frac{T}{\pi} \right\} &= \text{Spf} R \{T, U\} / (\pi U - T), \\ \mathcal{U}_2 &= \text{Spf} R \left\{ T, \frac{\pi}{T} \right\} &= \text{Spf} R \{T, V\} / (TV - \pi) \end{aligned}$$

respectively. We glue them as $U = V^{-1}$ and get $\mathcal{U}_1 \cup \mathcal{U}_2 \rightarrow \mathcal{D}^1 = \mathrm{Spf} R\{T\}$, which is the formal blow-up along the ideal (π, T) . Meanwhile, the analytic reduction is obtained by gluing the following two affine sets:

$$\begin{aligned}\overline{U}_1 &= \mathrm{Spec} k[T], \\ \overline{U}_2 &= \mathrm{Spec} k[T, V]/(TV),\end{aligned}$$

this is the normal crossing of the affine line with coordinate T and the projective line with the inhomogeneous coordinate V .

We are going to state the Raynaud's theorem, which provides a close relationship between rigid geometry and formal geometry; we need some terminology:

Definition 3.10 (Admissible formal scheme). An R -formal scheme⁽²⁾ \mathcal{X} is said to be *admissible* if it is Zariski locally isomorphic to admissible $\mathrm{Spf} A$ with A admissible in the sense of Definition 3.4.

Definition 3.11 (Admissible formal blow-up). Let \mathcal{X} be an admissible formal scheme. An *admissible formal blow-up* of \mathcal{X} is the formal blow-up along a coherent open ideal $\mathcal{J} \subset \mathcal{O}_{\mathcal{X}}$, i.e.,

$$\mathcal{X}' = \varinjlim_{\lambda} \mathrm{Proj} \bigoplus_{n=0}^{\infty} \left(\mathcal{J}^n \otimes_{\mathcal{O}_{\mathcal{X}}} \mathcal{O}_{\mathcal{X}}/\pi^\lambda \right) \longrightarrow \mathcal{X}.$$

We are going to construct the functor from the category of admissible formal schemes to the category of rigid analytic spaces; let us start by observing the affine case. From an affine admissible formal scheme $X = \mathrm{Spf} A$, it is easy to get the corresponding rigid analytic space: we just set

$$A_{\mathrm{rig}} := A \otimes_R K$$

which is an affinoid algebra over K . If A is presented as $A = R\{t_1, \dots, t_n\}/\mathfrak{a}$, then $K\{t_1, \dots, t_n\}/\mathfrak{a}K\{t_1, \dots, t_n\}$ gives a presentation of A_{rig} .

To globalize, we need to see the compatibility with localization. Let us consider the completed localization $A\{f^{-1}\}$ of A with respect to $f \in A$. The corresponding affinoid algebra is

$$\begin{aligned}A\{f^{-1}\} \otimes_R K &= A\{T\}/(1 - Tf) \otimes_R K \\ &= A_{\mathrm{rig}}\{T\}/(1 - Tf) = A_{\mathrm{rig}}\{f^{-1}\},\end{aligned}$$

and the latter is the affinoid algebra attached to the rational subdomain

$$\{x \in \mathrm{Spm} A_{\mathrm{rig}} \mid |f| \geq 1\};$$

this means that the completed localization of formal schemes just corresponds to the localization of rigid spaces with respect to the G-topology. So we globalize the recipe as above to obtain the functor

$$\mathrm{Rig}: \mathcal{X} \mapsto \mathcal{X}_{\mathrm{rig}}$$

from admissible formal R -schemes to K -rigid analytic spaces. The rigid analytic space $\mathcal{X}_{\mathrm{rig}}$ is called the *Raynaud generic fiber* of \mathcal{X} .

⁽²⁾We follow [EGA I, §10] for generalities of formal schemes.

Proposition 3.12. The functor Rig maps admissible formal blow-up's of admissible formal schemes to isomorphisms of rigid analytic spaces.

PROOF. This has already been essentially shown just before the Slogan. Indeed, we may limit ourselves to the affine case, for the centers of blow-up's are coherent. We have seen in this case that the Raynaud generic fiber of admissible formal blow-up's is the decomposition into rational subdomains, which does not change the analytic structure. \square

Now we state the theorem of Raynaud:

Theorem 3.13 (Raynaud 1972). The functor gives the equivalence of the following categories,

- (i) the category of quasi-compact admissible formal R -schemes, localized by admissible formal blow-up's,
- (ii) the category of quasi-compact and quasi-separated rigid analytic spaces over K .

The reader finds a nice comprehensive account of the proof of this theorem in [BL93].

4. Topology of Rigid Analytic Space.

In this section, we assume, for simplicity, that K is algebraically closed.

Definition 4.1. (1) A rigid analytic space X is said to be *quasi-compact* if every admissible covering of X has a finite admissible refinement.

- (2) A rigid analytic space X is said to be *connected* if there is no admissible covering $\{U_i\}_{i \in I}$ of X such that

$$\bigcup_{i \in I_1} U_i \cap \bigcup_{i \in I_2} U_i = \emptyset \text{ and } \bigcup_{i \in I_1} U_i \neq \emptyset \neq \bigcup_{i \in I_2} U_i$$

for some non-empty subsets $I_1, I_2 \subseteq I$ with $I = I_1 \sqcup I_2$.

Since admissible coverings of affinoids are fixed as finite, every affinoid is quasi-compact. So, we see that a rigid analytic space is quasi-compact if and only if it is a finite union of affinoids. It is easy to see that a rigid analytic space X is connected if and only if the ring $\Gamma(X, \mathcal{O}_X)$ has no other idempotent than 0 and 1.

Definition 4.2. A morphism $\pi: Y \rightarrow X$ of rigid analytic spaces over K is said to be an *analytic covering* if there exists an admissible covering⁽³⁾ $\{X_i\}_{i \in I}$ of X such that for each $i \in I$ $\pi^{-1}(X_i)$ is isomorphic to the disjoint union of copies of X_i . A connect rigid analytic space X is said to be *simply connected* if there is no other connected analytic covering over X than the trivial one $\text{id}: X \rightarrow X$ (cf. [vdP87]).

⁽³⁾or else a topological covering, referring to the Grothendieck topology; cf. Chapter I, §§1.4

Lemma 4.3. Let X be a connect rigid analytic space X .

(1) The space X is simply connected if and only if every locally constant sheaf of sets, with respect to the Grothendieck topology, is constant.

(2) If there exists an admissible covering $\{X_i\}_{i \in \mathbb{N}}$ of X such that (a) every X_i is simply connected, and (b) $X_i \subseteq X_{i+1}$ for all i , then X is simply connected.

PROOF. Both are straightforward. \square

Example 4.4 (Topology of the closed disk). Let $D = D(0, 1^+)$ be the 1-dimensional closed disk of radius 1. A rational subdomain in D of form

$$\{x \in D \mid |x - a| \leq \rho \text{ and } |x - a_i| \geq \rho_i \text{ for } i = 1, \dots, s\},$$

where $\rho, \rho_i \in |K^*|$ and $a, a_i \in D$, is called a *standard domain*. This is the complement of finite union of open disks. The following properties are easy to verify:

- If S_1 and S_2 are standard domains such that $S_1 \cap S_2 \neq \emptyset$, then so are $S_1 \cup S_2$ and $S_1 \cap S_2$.
- Every finite union of standard domains is uniquely decomposed as a disjoint union of standard domains.

What is more interesting is the following proposition:

Proposition 4.5. Every rational subdomain in D is a finite union of standard domains. (Hence standard domains generate the Grothendieck topology of D .)

SKETCH OF PROOF. (cf. [GvdP80, III.1.18]). Let R be the rational subdomain given by $|f_i(x)| \leq |f_0(x)|$ for $i = 1, \dots, n$. Deforming those functions slightly, if necessary, we may assume f_i and f_j have no common zero for any $i \neq j$, and thus, we may concentrate to the single inequality $|f_1(x)| \leq |f_0(x)|$. By Weierstrass Preparation Theorem we may assume f_0 and f_1 are polynomials having all their roots in D . Then the proposition follows from an easy calculation. \square

Corollary 4.6. The Grothendieck topology of the projective line $\mathbb{P}_K^{1, \text{an}}$ is generated by standard domains.

Proposition 4.7. Standard domains are connected. Moreover, every connected rational subdomain in $\mathbb{P}_K^{1, \text{an}}$ is simply connected.

PROOF. The first assertion is obvious since the corresponding affinoid algebra

$$K\{t, t_0, \dots, t_s\}/((t - a) - t_0\pi, \pi_i - t_i(t - a_i) \text{ for } i = 1, \dots, s),$$

where $a, a_i \in K$ and $\pi, \pi_i \in K$, is integral. (Note that $K\{t, t_0, \dots, t_s\}$ is Noetherian and factorial.) The second one is proved by using 4.3 (1). Let S be a standard domain and \mathcal{F} a constant sheaf of sets on S . We can take an admissible covering $\{S_i\}$ of S which trivializes \mathcal{F} . The point is that any

non-disjoint union of standard domains is standard, and hence, connected. So the restriction maps $\mathcal{F}(S_i) \rightarrow \mathcal{F}(S_i \cap S_j)$ must be bijective as far as $S_i \cap S_j \neq \emptyset$. It follows that $\mathcal{F}(S_i \cup S_j) \rightarrow \mathcal{F}(S_i)$ are bijective, and repeating this argument, we concludes \mathcal{F} is constant. \square

The following corollary may seen at first:

Corollary 4.8. Every connected open subanalytic space⁽⁴⁾ in $\mathbb{P}_K^{1,\text{an}}$ is simply connected.

PROOF. A connected open subanalytic space U has an admissible covering consisting of standard domains. By [GvdP80, III,2.6] and the easy fact that a non-disjoint union of connected sets is again connected, we see that a finite non-disjoint union of standard domain is a connected rational subdomain. Hence the corollary follows from Lemma 4.3 (2) and Proposition 4.7. \square

Finally we quote, without proof, a theorem by van der Put which indicates a close relation between the topology of a rigid analytic space and that of its analytic reduction:

Theorem 4.9. Let X be a quasi-compact rigid analytic space over K which has the irreducible and smooth analytic reduction. Then X is simply connected.

For the proof, see [vdP87].

Remark 4.10. By the corollary, we know, for example, that the space $\mathbb{G}_K^{\text{an}} = K^\times$ is simply connected. But, on the other hand, it is easy to see that the map $K^\times \rightarrow K^\times$ by $x \mapsto x^n$ is an analytic morphism. As far as the characteristic of K does not divide n , we are tempted to think of this morphisms also as an analytic covering. But it is not. Certainly, this morphism induces isomorphisms between stalks of the structure sheaf, but there is no *admissible* covering which trivializes this morphism! This kind of morphisms is said to be *étale*; analytic coverings are étale, but not vice versa. Simply connected analytic spaces may have many étale coverings.

5. Berkovich' approach to non-archimedean analysis

Berkovich' viewpoint provides an innovative approach to topological problems of p -adic geometry and analysis, which is much closer to the familiar intuition derived from the complex context [Ber90]⁽⁵⁾. His analysis, in fact, generalizes, or properly speaking, “completes” the rigid analysis. This situation is, according to what he says in the introduction of his book, somehow analogous to that of \mathbb{R} completing \mathbb{Q} . The standard archimedean metric space \mathbb{Q} is totally disconnected. The rigid analytic viewpoint corresponds, to some extent, to regarding every rational interval $\{r \in \mathbb{Q} \mid a \leq r \leq b\}$ as

⁽⁴⁾We refer [BGR84, 9.3.1] for the definition of open subanalytic spaces.

⁽⁵⁾Already in [vdP82] the *esquisse* of Berkovich' idea can be seen.

“connected”, and considering only those functions which come from analytic functions on the corresponding real interval. Berkovich' non-archimedean analysis can be compared, in this context, to analysis on \mathbb{R} itself. Namely, his presentation of analytic spaces complements those of rigid analysis by putting some “generic points”, and consequently, simplifies their topology.

Let us be a little more precise about his theory. Let A be a Banach ring, i.e., a ring together with a norm $\|\cdot\|$ with respect to which A is complete.

Definition 5.1. (1) A semi-norm $|\cdot|$ on A is said to be *bounded* if there exists $C > 0$ such that $|f| \leq C\|f\|$ for any $f \in A$.

(2) A semi-norm $|\cdot|$ on A is said to be *multiplicative* if $|fg| = |f||g|$ for all $f, g \in A$.

Replacing the maximal spectra $\text{Spm}(A)$ of rigid geometry, we have:

Definition 5.2. The Berkovich' *spectrum* $M(A)$ for a Banach ring A is the set of all bounded multiplicative semi-norms on A , endowed with the weakest topology so that the real valued functions of form $|\cdot| \mapsto |f|$ for $f \in A$ are continuous.

In case A is an affinoid algebra (a *strict* affinoid algebra, in Berkovich' term) over K , the affinoid space $\text{Spm}(A)$ is naturally viewed as a subspace, with the relative topology, of the Berkovich' spectrum $M(A)$. This is done by identifying an element \mathfrak{m} in $\text{Spm}(A)$ with the unique semi-norm $|\cdot|$ on A such that $\mathfrak{m} = \text{Ker } |\cdot|$. The space $\text{Spm}(A)$ is identified, in terms of this correspondence, with the subspace of $M(A)$ consisting of semi-norms $|\cdot|$ such that $\dim_K A/\text{Ker } |\cdot|$ is finite (the “classical points”).

Example 5.3. Let us consider the usual absolute-value norm $|\cdot|_\infty$ on \mathbb{Z} . The spectrum $M(\mathbb{Z})$ consists of the following points:

- (i) $|\cdot|_{\infty, \epsilon} = |\cdot|_\infty^\epsilon$ ($0 < \epsilon \leq 1$).
- (ii) The p -adic norm $|\cdot|_{p, \epsilon}$ with $|p|_{p, \epsilon} = \epsilon$ ($0 < \epsilon \leq 1$).
- (iii) The semi-norm $|\cdot|_p$ induced from the trivial norm on $\mathbb{Z}/p\mathbb{Z}$.
- (iv) The trivial norm $|\cdot|_0$.

As a topological space, $M(\mathbb{Z})$ is a tree with end points $|\cdot|_p$ and $|\cdot|_\infty$. Each of these points is connected by a single edge with $|\cdot|_0$.

On the Berkovich' spectrum, the *affinoid subdomains* (incl. *rational subdomains*) are defined in the similar way to rigid analysis. Thus one can define the structure sheaves on spectra, which one calls affinoid spaces. The local data consisting of Berkovich' affinoids and the structure sheaves can glue to locally ringed spaces; thus we obtain *analytic spaces* in Berkovich' sense.

Example 5.4. (Cf. [Ber90, 1.5].) Let K be a non archimedean complete field. An affine space over K is defined by

$$\mathbb{A}_K^n = \{\text{multiplicative seminorm } |\cdot| \text{ on } K[t_1, \dots, t_n] \mid |\cdot|_{|_K} \text{ is bounded}\}$$

together with the weakest topology so that $|\cdot| \mapsto |f|$ ($f \in K[t_1, \dots, t_n]$) is continuous. This has a structure of analytic space so that the analytic functions over it is characterized by limits of rational functions. When $K = \mathbb{C}$ the space \mathbb{A}_K^n is nothing but the usual affine space in the complex analytic sense. If K is a non-archimedean field, then \mathbb{A}_K^n is a union of balls; i.e., the spectrum of Tate algebra.

Here are some general properties of Berkovich' analytic spaces:

- (i) Every connected K -analytic space is arcwise connected.
- (ii) If $X = M(A)$ with A an affinoid algebra over K , then $\text{Krull-dim} A = \dim |X|$, where $|X|$ is the underlying topological space of X .
- (iii) If a K -analytic space X is smooth, then it is locally contractible.

Relation between Berkovich' analytic spaces and rigid analytic spaces is as follows. If a Berkovich' K -analytic space X is formed only by spectra of (strict) affinoid algebras, we say X is strict. There exists a functor from the category of separated strict K -analytic spaces to that of separated rigid analytic spaces over K by

$$X \mapsto X_0 = \{x \in X \mid [K(x): K] < \infty\},$$

where $K(x)$ is the residue field at x .

Theorem 5.5. This functor is fully-faithful, and preserves fiber products.

APPENDIX B

Overview of theory of p -adic uniformization.

By F. KATO

In this appendix we always denote by K , R , and π a complete discrete valuation field, its valuation ring, and a prime element of R , respectively. We assume that the residue field $k = R/\pi R$ is finite and consists of q elements. The Mumford-Kurihara-Mustafin uniformization is a procedure to construct nice analytic (and in many cases algebraic) varieties by taking discrete Schottky-type quotients of a certain p -adic analogue of symmetric space, so-called *Drinfeld symmetric space* Ω , or its variants.

1. Bruhat-Tits building.

First we survey the construction of a certain simplicial complex, called Bruhat-Tits building (attached to $\mathrm{PGL}(n+1, K)$), which will be closely related with the Drinfeld symmetric space. Let n be a positive integer and V an $n+1$ dimensional vector space over K . A *lattice* in V is a finitely generated R -submodule of V which spans V over K . Every lattice is therefore a free R -module of rank $n+1$. Let $\tilde{\Delta}_0$ be the set of all lattices in V . We say that two lattices M_1 and M_2 are *similar* if there exists $\lambda \in K^\times$ such that $M_1 = \lambda M_2$. The similarity is obviously an equivalence relation. We denote by Δ_0 the set of all similarity classes of lattices in V .

Definition 1.1. The *Bruhat-Tits building* (attached to $\mathrm{PGL}(n+1, K)$) is the finite dimensional simplicial complex Δ with the vertex set Δ_0 defined as follows: A finite subset $\{\Lambda_0, \dots, \Lambda_l\}$ of Δ_0 forms an l -simplex if and only if, after permuting indices if necessary, one can choose $M_i \in \Lambda_i$ for $0 \leq i \leq n+1$ such that

$$M_0 \supsetneq M_1 \supsetneq \cdots \supsetneq M_l \supsetneq \pi M_0.$$

The role of M_0 is by no means important; for example, one can shift the indices like $M_1 \supsetneq \cdots \supsetneq M_l \supsetneq \pi M_0 \supsetneq \pi M_1$.

To understand the structure of Δ , let us fix one vertex $\Lambda = [M]$. Suppose $\{\Lambda_0, \dots, \Lambda_l\}$ is an l -simplex having Λ as a vertex; we may assume $\Lambda = \Lambda_0$, and can take $M_i \in \Lambda_i$ for $1 \leq i \leq l$ such that $M \supsetneq M_1 \supsetneq \cdots \supsetneq M_l \supsetneq \pi M$. Set $\overline{M}_i := M_i/\pi M$. Then we get a flag

$$\overline{M} \supsetneq \overline{M}_1 \supsetneq \cdots \supsetneq \overline{M}_l \supsetneq 0$$


of length $l + 1$ consisting of subspaces of the $n + 1$ -dimensional k -vector space $\overline{M} = M/\pi M$. Moreover, by the elementary algebra one finds that this yields a one-to-one correspondence between l -simplices containing Λ and length $l + 1$ frags in $\overline{M} = M/\pi M$. By this quite easy observation one knows

- (i) Δ is an n -dimensional locally finite simplicial complex,
- (ii) every simplex in Δ is a face of a chamber, a simplex having maximal ($= n$) dimension.

Furthermore, by the construction,

- (iii) the group $\text{Aut}_K(V)/K^\times (\cong \text{PGL}(n + 1, K))$ naturally acts on Δ simplicially.

Example 1.2 ($n = 1$ case). In $n = 1$ case, the vertices adjacent to a fix vertex Λ are in one-to-one correspondence with the set of all k -rational points of the projective line \mathbb{P}_k^1 ; in particular, each vertex has exactly $q + 1$ adjacent vertices. Moreover, it can be seen that the simplicial complex $\Delta = \Delta^1$ is a *tree*. The following picture shows how it looks like in case $q = 2$:



In general, the geometrical realization $|\Delta|$ of Δ is known to be a contractible topological space.

Example 1.3 ($n = 2$ case). Already in case $n = 2$, the structure of $\Delta = \Delta^2$ is fairly complicated. For a fixed vertex Λ , the set of all vertices adjacent of Λ is divided in two; one for those corresponding to $q^2 + q + 1$ k -rational points in \mathbb{P}_k^2 and the other for those corresponding to $q^2 + q + 1$ k -rational lines in \mathbb{P}_k^2 . So it is convenient to consider the surface B which is obtained by blowing-up of \mathbb{P}_k^2 at every k -rational points. Two vertices adjacent to Λ forms with Λ a 2-simplex if and only if the corresponding lines in B intersect.

2. Drinfeld symmetric space.

To each lattice M in V we associate an R -scheme

$$\mathbb{P}(M) := \text{Proj}(\text{Sym}_R M).$$

Obviously the scheme $\mathbb{P}(M)$ depends only on the similarity class; so we should write as $\mathbb{P}(\Lambda)$, where $\Lambda = [M]$. The scheme $\mathbb{P}(M)$ is isomorphic to \mathbb{P}_R^n ; but the isomorphism cannot be taken canonically, while there exists the

canonical isomorphism $\mathbb{V} \cong \mathbb{P}(M) \otimes_R K$ due to the canonical isomorphism $M \otimes_R K \cong V$. So we get a bijection:

$$\Delta_0 \xrightarrow{\sim} \left\{ (\mathbb{P}, \phi) \left| \begin{array}{l} \mathbb{P} \xrightarrow{\sim} \mathbb{P}_R^n, \\ \phi: \mathbb{P} \otimes_R K \xrightarrow{\sim} \mathbb{P}(V) \end{array} \right. \right\} / \sim,$$

where $(\mathbb{P}_1, \phi_1) \sim (\mathbb{P}_2, \phi_2)$ if and only if there exists an R -isomorphism $\Phi: \mathbb{P}_1 \xrightarrow{\sim} \mathbb{P}_2$ which makes the diagram

$$\begin{array}{ccc} \mathbb{P}_1 \otimes_R K & \xrightarrow{\Phi \otimes_R K} & \mathbb{P}_2 \otimes_R K \\ \phi_1 \downarrow & & \downarrow \phi_2 \\ \mathbb{P}(V) & = & \mathbb{P}(V) \end{array}$$

commutative.

Next we consider two vertices Λ_1 and Λ_2 which are adjacent with each other. The canonical isomorphism between the generic fibers induce a natural birational map between $\mathbb{P}(\Lambda_1)$ and $\mathbb{P}(\Lambda_2)$. More explicitly, taking representatives $M_i \in \Lambda_i$ ($i = 1, 2$) so that $M_1 \supsetneq M_2 \supsetneq \pi M_1$, we have the birational map from $\mathbb{P}(\Lambda_1)$ to $\mathbb{P}(\Lambda_2)$ induced by $M_2 \hookrightarrow M_1$. We define the join $\mathbb{P}(\Lambda_1) \vee \mathbb{P}(\Lambda_2)$ to be the closure of the graph in $\mathbb{P}(\Lambda_1) \times_R \mathbb{P}(\Lambda_2)$ of this rational map. It is easily seen that the join $\mathbb{P}(\Lambda_1) \vee \mathbb{P}(\Lambda_2)$ is R -isomorphic to the blow-up of $\mathbb{P}(\Lambda_1)$ at the closed subscheme $\text{Proj}(\text{Sym}_k M_1/M_2)$. In particular, the closed fiber of $\mathbb{P}(\Lambda_1) \vee \mathbb{P}(\Lambda_2)$ consists of two rational varieties intersecting transversally.

Lemma 2.1. The operation \vee is associative and commutative.

The proof is straightforward, and left to the reader. By the lemma, we can extend the construction to any finite subcomplex of Δ . More precisely, for a finite subcomplex S , we take the joins of all vertices in S ; the resulting R -scheme is denoted by $\mathbb{P}(S)$. The scheme $\mathbb{P}(S)$ is a regular projective scheme over R having the following properties:

- (i) The generic fiber is canonically isomorphic to $\mathbb{P}(V)$.
- (ii) The closed fiber is a reduced normal crossing of non-singular rational varieties of which the dual graph coincides with S .

We want to do this for a general convex subcomplex of Δ ; this cannot be done in terms of schemes because we have to consider a limit of blow-up's of \mathbb{P}_R^n centered in the closed fiber. But we can do this in terms of formal schemes, or rigid analytic spaces. Let Δ_* be a convex subcomplex of Δ . For any finite subcomplex S of Δ_* we define $\widehat{\Omega}(S)$ to be the completion of $\mathbb{P}(S)$ along the closed fiber. Consider the maximal Zariski open subset $\widehat{\Omega}(S)'$ of $\widehat{\Omega}(S)$ such that, for any finite subcomplex T of Δ_* containing S , the induced morphism $\rho_S^T: \widehat{\Omega}(T) \rightarrow \widehat{\Omega}(S)$ gives an isomorphism restricted to the pull-back of $\widehat{\Omega}(S)'$.

Definition 2.2. For a convex subcomplex Δ_* of Δ we define

$$\widehat{\Omega}(\Delta_*) := \bigcup_{S: \text{finite}} \widehat{\Omega}(S)'$$

where S runs through all finite subcomplex of Δ_* . The corresponding rigid analytic space (Raynaud generic fiber) is denoted by $\Omega(\Delta_*)$. In case $\Delta_* = \Delta$ we simply write $\widehat{\Omega} = \widehat{\Omega}(\Delta)$ and $\Omega = \Omega(\Delta)$. The rigid analytic space Ω is called the *Drinfeld symmetric space*.

The formal scheme $\widehat{\Omega}(\Delta_*)$ is formally locally of finite type, of which the closed fiber is the normal crossings of non-singular rational varieties having the dual graph isomorphic to Δ_* . By definition, the group $\mathrm{PGL}(n+1, K)$ naturally acts on $\widehat{\Omega}$, and hence Ω .

Example 2.3. Let us consider $\widehat{\Omega}$ in $n = 2$. The closed fiber $\widehat{\Omega}_0$ consists of countably many components which are all isomorphic to the surface B considered in 1.3. In each component every k -rational line is a double curve in $\widehat{\Omega}_0$, and every k -rational point is a triple point.

Structure of Ω . First we note that, as a point set, we have an equality

$$\Omega = \mathbb{P}_K^{n,\mathrm{an}} \setminus \bigcup_{H \in \mathcal{H}} H,$$

where \mathcal{H} is the set of all K -rational hyperplanes.

Here we do not attempt to give a proof of this fact, but rather give a sketchy explanation in order to get a feeling. We limit ourselves to $n = 1$ case just for simplicity. The formal scheme $\widehat{\Omega}$ is, roughly speaking, the *limit* of successive blow-up's

$$\cdots \longrightarrow U_i \xrightarrow{\rho_i} U_{i-1} \longrightarrow \cdots \longrightarrow U_0$$

with $U_0 = \mathbb{P}_R^1$ and ρ_i the blow-up centered at all the k -rational points in the closed fiber of U_{i-1} . Let z be a point in the generic fiber \mathbb{P}_K^1 of each U_i . We consider the closure \bar{z} of z in each U_i . Since U_i is a regular scheme, the section \bar{z} intersects the closed fiber at a smooth point transversally. If z is *not* a K -rational point, then for sufficiently large N , \bar{z} intersects the closed fiber of U_i at a non- k -rational smooth point for $i > N$. But, if z is K -rational, \bar{z} intersects the closed fiber always at a k -rational point which is in the next blow-up center. So if we consider the limit, such formal sections determined by K -rational points does not exist any more, while those by non- K -rational points certainly exist. This explains the above equality in $n = 1$. For higher n we can apply the similar argument.

Next we discuss the analytic structure of Ω ; viz. we show how to introduce an admissible covering to Ω ; since we have constructed the rigid analytic space Ω from the formal model, we must have one such covering such that the associated formal model recovers $\widehat{\Omega}$. This can be actually described in an elegant way in terms of norms: A non-negative real valued function $\alpha: V \rightarrow \mathbb{R}_{\geq 0}$ is said to be a *norm* over K if it satisfies:

- (i) $\alpha(x) > 0$ whenever $x \neq 0$.
- (ii) For $a \in K$, $\alpha(ax) = |a|\alpha(x)$.
- (iii) $\alpha(x + y) \leq \max\{\alpha(x), \alpha(y)\}$.

Two norms α and α' over K is said to be *similar* if $\alpha' = \lambda\alpha$ for some $\lambda \in \mathbb{R}_{>0}$. The set of all similarity classes of norms on V over K forms in an obvious way a topological space. The following fact is well-known ([GI63]):

Proposition 2.4. The topological space of similarity classes of norms on V over K is $\mathrm{PGL}(n + 1, K)$ -equivariantly homeomorphic to the topological realization $|\Delta|$ of the Bruhat-Tits building Δ .

The homeomorphism is given as follows: To any lattice M we need to associate a norm α_M . For a non-zero $x \in V$, the set $\{a \in K \mid ax \in M\}$ is a fractional ideal of R , and hence is of form (π^m) . We then define $\alpha_M(x) = q^m$ for non-zero $x \in V$, and $\alpha_M(0) = 0$. It is easily verified that α_M is a norm. The similarity class of α_M depends only on the similarity class of M . Thus we get a mapping from Δ_0 to the set of similarity classes of norms. Let Λ_0 and Λ_1 be vertices adjacent with each other. We can choose a basis e_0, \dots, e_n of V such that $M_0 = \bigoplus_{i=0}^n Re_i$ and $M_1 = \bigoplus_{i=0}^{n-1} Re_i \oplus R\pi e_n$ give representatives of Λ_0 and Λ_1 , respectively. Then the norms α_{M_0} and α_{M_1} are given by

$$\begin{aligned} \alpha_{M_0}\left(\sum_{i=0}^n a_i e_i\right) &= \max\{|a_0|, \dots, |a_n|\}, \\ \alpha_{M_1}\left(\sum_{i=0}^n a_i e_i\right) &= \max\{|a_0|, \dots, |a_{n-1}|, |\pi|^{-1}|a_n|\}, \end{aligned}$$

respectively. For $0 < t < 1$, the class of the norm

$$\alpha_t\left(\sum_{i=0}^n a_i e_i\right) = \max\{|a_0|, \dots, |a_{n-1}|, |\pi|^{-t}|a_n|\}$$

corresponds to the point $t\Lambda_0 + (1 - t)\Lambda_1$ on the edge connecting Λ_0 and Λ_1 .

Now, to any point $z = (z_0 : \dots : z_n) \in \mathbb{P}(V)$ (subject to some K -basis $\{e_0, \dots, e_n\}$ of V) we associate, up to invertible factor, a non-negative real valued function

$$\alpha_z\left(\sum_{i=0}^n a_i e_i\right) = \left| \sum_{i=0}^n a_i z_i \right|$$

on V . Note that *this determines a class of norm if and only if z is not lying in any K -rational hyperplane*. Thus we get a mapping

$$\rho: \Omega \longrightarrow |\Delta|.$$

The admissible covering in question is actually given by the collection of subsets of Ω consisting of pull-backs of each simplices of $|\Delta|$ by ρ . For example, notation being as above,

$$\rho^{-1}(\Lambda_0) = \{|z_0| = \dots = |z_n| = 1\} \setminus \bigcup_H \{(z_0 : \dots : z_n) \bmod \pi \in H\},$$

where H runs through all hyperplanes in \mathbb{P}_k^n , and

$$\rho^{-1}(t\Lambda_0 + (1 - t)\Lambda_1) = \{|z_0| = \dots = |z_{n-1}| = 1, |z_n| = |\pi|^{-t}\}$$

for $0 < t < 1$.

3. Uniformization.

Recall that, once we fix a K -basis of V , the group $\mathrm{PGL}(n+1, K)$ canonically acts on the Bruhat-Tits building Δ and the Drinfeld symmetric space Ω . We shall consider discrete groups in $\mathrm{PGL}(n+1, K)$ and the quotients of Δ and Ω by them. A subgroup $\Gamma \in \mathrm{PGL}(n+1, K)$ is said to be *hyperbolic* if it acts on Δ discretely and freely.

To a hyperbolic subgroup Γ we associate a convex subcomplex Δ_Γ of Δ by the following recipe: In general, a convex subcomplex generated by vertices of form

$$\left[\sum_{i=0}^n R \cdot \pi^{\sigma_i} Y_i \right], \quad \sigma_i \in \mathbb{Z}, \quad 0 \leq i \leq n,$$

subject to a basis Y_0, \dots, Y_n of V , is said to be an *apartment*. The geometrical realization of each apartment is a triangulation of \mathbb{R}^n of A_n -type, and Δ is the union of all apartments. There exists an obvious bijection between the set of all apartments and the set of all K -split maximal tori in $\mathrm{PGL}(n+1, K)$. For any K -split maximal tori T the intersection $\Gamma \cap T$ is a commutative discrete group of rank at most n . If this rank is n we call the apartment corresponding to T a Γ -*apartment*, and we define Δ_Γ to be the union of all Γ -apartments. Clearly the group Γ acts on Δ_Γ ; if the quotient Δ_Γ/Γ is a finite simplicial complex, we say Γ is *normal hyperbolic*.

Now let us be given a normal hyperbolic subgroup Γ . For each $\gamma \in \Gamma$ and each finite subcomplex S of Δ_Γ , γS is again a finite subcomplex of Δ_Γ , and then, γ induces an R -isomorphism $\mathbb{P}(S) \cong \mathbb{P}(\gamma S)$ which induces on generic fibers ($= \mathbb{P}_K^n$) exactly the linear automorphism γ . Considering all S , gluing, and taking Raynaud generic fiber give a rigid analytic automorphism of $\Omega(\Delta_\Gamma)$ which we denote again by γ . Thus we get an homomorphism

$$\Gamma \longrightarrow \mathrm{Aut}_{K\text{-rig}} \Omega(\Delta_\Gamma).$$

Theorem 3.1 (Mustafin [Mus78], Kurihara [Kur80]). Let Γ be a normal hyperbolic subgroup of $\mathrm{PGL}(n+1, K)$ and Δ_Γ as above. Then there exists, unique up to isomorphisms, a rigid analytic space X_Γ , smooth, proper and of finite type over K , and a topological covering map $\mathfrak{p}: \Omega(\Delta_\Gamma) \rightarrow X_\Gamma$ such that

- (i) $\mathfrak{p} \circ \gamma = \mathfrak{p}$ for any $\gamma \in \Gamma$,
- (ii) for $x, y \in \Omega(\Delta_\Gamma)$, $\mathfrak{p}(x) = \mathfrak{p}(y)$ if and only if $x = \gamma(y)$ for some $\gamma \in \Gamma$.

In the references, this theorem has been stated in terms of formal schemes; viz. they took quotients of $\widehat{\Omega}(\Delta_\Gamma)$ to get a formal scheme \widehat{X}_Γ flat and formally of finite type over R . We have an admissible covering of X_Γ induced from that of $\Omega(\Delta_\Gamma)$; in terms of formal scheme, the closed fiber of \widehat{X}_Γ is a divisor of normal crossings with rational components having the dual group isomorphic to Δ_Γ/Γ .

Algebraicity. In $n = 1$ case the rigid analytic curve X_Γ is algebraizable for any Γ ([**Mu72b**]). In higher dimensions, we know the following sufficient condition: Let Γ be a torsion-free *uniform lattice* in $\mathrm{PGL}(n + 1, K)$, viz. a discrete co-compact subgroup of finite co-volume without elements of finite order. Then, since it is finite co-volume, we have $\Delta_\Gamma = \Delta$. Hence in this case, we are going to take a quotient of the Drinfeld symmetric space Ω .

Theorem 3.2 (loc. cit.). If Γ is a torsion-free uniform lattice, then the quotient rigid analytic variety X_Γ can be algebraizable to a non-singular projective variety having the ample canonical class.

In terms of formal scheme, this means that the relative canonical sheaf of $\widehat{X}_\Gamma/\mathrm{Spf}R$ is ample.

4. Examples.

Tate curve. Let T be the maximal torus consisting of all invertible diagonal matrices. We take n elements $q_i \in K^\times$ ($1 \leq i \leq n$) with $|q_i| < 1$ and set $q_0 = 1$. Define $\Gamma \subset T$ to be the set of all diagonal matrices with (i, i) -entry an integer power of q_i for $0 \leq i \leq n$. Then Γ is a normal hyperbolic subgroup of $\mathrm{PGL}(n + 1, K)$ with Δ_Γ the apartment corresponding to T . If $n = 1$, the curve $\Omega(\Delta_\Gamma)/\Gamma$ is known to be the so-called *Tate curve*; in this case $\Omega(\Delta_\Gamma)$ is simply \mathbb{P}_K^1 minus two points.

Mumford curve. Consider again the $n = 1$ case. The rigid analytic curve X_Γ with Γ generated by at least two elements is called a *Mumford curve*. A Mumford curve is, in fact, algebraized to a non-singular projective curve of genus greater than 1. Moreover, the formal model provided by the admissible covering inherited from $\Omega(\Delta_\Gamma)$ gives a stable k -split multiplicative reduction of that curve. The converse is also known:

Theorem 4.1 (Mumford [**Mu72b**]). Let X be a smooth and proper rigid analytic curve over K , and suppose X admits a formal model with k -split multiplicative reduction. Then X is isomorphic to X_Γ for a unique Γ .

Surface case. In higher dimension it becomes extremely difficult to find normal hyperbolic groups; but there are several known examples of torsion-free uniform lattices. Let Γ is a torsion-free uniform lattice, and N the number of Γ -orbits in the vertex set Δ_0 . Mumford [**Mu79**] calculated the several numerical invariants of the quotient surface X_Γ :

- (i) $\chi(\mathcal{O}_{X_\Gamma}) = N(q - 1)^2(q + 1)/3$.
- (ii) $(c_{1, X_\Gamma})^2 = 3c_{2, X_\Gamma} = 3N(q - 1)^2(q + 1)$.
- (iii) $q(X_\Gamma) = 0$,

where q denotes the irregularity. It is very interesting that the Mumford-Kurihara-Mustafin uniformization in two dimension always produces the surfaces which satisfies the equality in Miyaoka-Yau inequality. Mumford [**Mu79**] constructed one example of Γ in $q = 2$ such that $N = 1$; the resulting surface is one of the so-called “fake projective planes”. Recently,

Cartwright, Mantero, Steger, and Zappa constructed several examples of lattices in $q = 2$ and $q = 3$. Making use of these groups, in [IK98], Ishida and the author discussed other possible fake projective planes, and showed that there are at least three fake projective planes (incl. Mumford's one) which are not isomorphic to among others. We will see in more detail the construction of Mumford's fake projective plane in the next section.

Shimura variety case. Let D be a quaternion algebra over a totally real field F . We assume that D ramifies at every infinite place except one ∞_0 . Let Γ_∞ be a subgroup of D^\times/F^\times defined by some congruence condition; we shall assume that it is "sufficiently small". Then it is classically known that Γ_∞ acts discontinuously on the upper $\frac{1}{2}$ plane \mathbb{H} such that the quotient \mathbb{H}/Γ_∞ is a *projective* curve (Shimura curve of PEL-type). The canonical model Sh is defined over some ray-class field of F .

There must be a finite place ν at which D ramifies. Let p be the residue characteristic of ν . We assume that Γ_∞ is maximal at ν . Then the Shimura curve Sh has a bad reduction at ν , and due to Cherednik [Čer76] and Drinfeld [Dri76], the associated rigid analytic space over F_ν^{ur} has a p -adic uniformization, where F_ν^{ur} is the maximal unramified extension of F_ν . Let \overline{D} be the quaternion algebra which has local invariants obtained precisely by switching those of D at ν and ∞_0 . Then there exists a subgroup Γ_ν in $\overline{D}^\times/F^\times$ defined basically by the same congruence relation as that of Γ_∞ such that $\Omega \otimes F^{\text{ur}}/\Gamma_\nu$ is isomorphic to the associated analytic space of Sh at ν . (This isomorphism actually descends to some finite unramified extension of F_ν , depending on the data of connected components of Sh .)

This story has been generalized to higher dimension by Rapoport-Zink [RZ96] and Boutot-Zink [BZ95]. The next section is devoted to explain one special example of the p -adic uniformization of a Shimura surface.

5. Mumford's fake projective plane.

Our example of p -adically uniformizable Shimura surface is constructed by means of some division algebra D . The following construction is inspired by the Mumford's original paper [Mu79].

Let $\zeta = \zeta_7$ be a primitive 7th-root of unity, and set $L := \mathbb{Q}(\zeta)$; L is a cyclic extension of \mathbb{Q} of degree 6 having an intermediate quadratic extension $K := \mathbb{Q}(\lambda)$ ($\cong \mathbb{Q}(\sqrt{-7})$), where $\lambda = \zeta + \zeta^2 + \zeta^4$. The Galois group of L/K is generated by the Frobenius map $\sigma: \zeta \mapsto \zeta^2$, and that of K/\mathbb{Q} is generated by the complex conjugation $z \mapsto \bar{z}$. Note that the prime 2 decomposes on K , more explicitly, $2 = \lambda\bar{\lambda}$ gives the prime factorization, and the prime 7 ramifies. We fix an infinite place $\varepsilon: K \hookrightarrow \mathbb{C}$ by $\lambda \mapsto \frac{-1+\sqrt{-7}}{2}$.

We set $\mu := \lambda/\bar{\lambda}$, and define the central division algebra D over K of dimension 9 by

$$D := \bigoplus_{i=0}^2 L\Pi^i$$

with $\Pi^3 = \mu$ and $\Pi z = z^\sigma \Pi$ for $z \in L$. The involution $*$ on D is defined by $z^* = \bar{z}$ for $z \in L$ and $\Pi^* = \bar{\mu} \Pi^2$. It is easily verified that $*$ is positive.

We introduce a non-degenerate anti-symmetric \mathbb{Q} -bilinear form $\psi = \psi_b$ on $V = D$ by putting $b = (\lambda - \bar{\lambda}) - \bar{\lambda} \Pi + \bar{\lambda} \Pi^2$. Then define an algebraic group G over \mathbb{Q} as follows: Given a \mathbb{Q} -algebra R , we set

$$G(R) := \{g \in ((D \otimes R)^{\text{opp}})^\times \mid \psi(xg, yg) = c(g)\psi(x, y), c(g) \in R^\times, x, y \in V \otimes R\}.$$

The algebraic group $G_{\mathbb{R}}$ is isomorphic to $\text{GU}(2, 1)$, which acts on the two dimensional complex unit-ball \mathbb{B} . In the following we will construct an open compact subgroup C in $G(\mathbb{A}_f)$. By these data, we can define a Shimura variety

$$\mathcal{S}_C := G(\mathbb{Q}) \backslash \mathbb{B} \times G(\mathbb{A}_f)/C.$$

The canonical model of \mathcal{S}_C is defined over $E := \varepsilon(K)$, which we denote by Sh_C .

The congruence condition. First we define an algebraic group I over \mathbb{Q} by

$$I(R) := \{g \in (\text{M}_d(K) \otimes R)^\times \mid H^{-1} {}^t \bar{g} H \in R^\times\}$$

for any \mathbb{Q} -algebra R , where

$$H = \begin{bmatrix} 3 & \bar{\lambda} & \bar{\lambda} \\ \lambda & 3 & \bar{\lambda} \\ \lambda & \lambda & 3 \end{bmatrix}.$$

Then I is an inner form of G so that $G(\mathbb{A}_f^2) \cong I(\mathbb{A}_f^2)$.

To define the level structure, it suffices to define the prime-2-part C^2 . We define C^2 in the form $C^2 = C^{2,7} C_7$ with $C_7 \in G(\mathbb{Q}_7) \cong I(\mathbb{Q}_7)$ and $C^{2,7}$ being the maximal one:

$$C^{2,7} := \{g \in G(\mathbb{A}_f^{2,7}) \mid \Gamma^{2,7} g \subseteq \Gamma^{2,7}\},$$

with $\Gamma^{2,7} = \Gamma \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}^{2,7}$, where $\widehat{\mathbb{Z}}^{2,7} = \varprojlim \mathbb{Z}/m\mathbb{Z}$ with the projective limit taken over m such that $(2, m) = (7, m) = 1$.

Now the component C_7 is defined as follows: we have an isomorphism

$$G(\mathbb{Q}_7) \xrightarrow{\sim} \{g \in \text{GL}_3(\tilde{\mathbb{Q}}_7) \mid g^* H g = c(g) H, c(g) \in \mathbb{Q}_7^\times\},$$

where $\tilde{\mathbb{Q}}_7$ is the ramified quadratic extension of \mathbb{Q}_7 , and $*$ is the matrix transposition followed by the Galois action. Let \tilde{C}_7 be the maximal open compact subgroup of $G(\mathbb{Q}_7)$ consisting of g which maps $\Gamma \otimes_{\mathbb{Z}} \mathbb{Z}_7$ to itself. We consider modulo $\sqrt{-7}$ reduction of \tilde{C}_7 , which is a subgroup G_0 in $\text{GL}_3(\mathbb{F}_7)$. The matrix $H \bmod \sqrt{-7}$ is of rank 1, and has 2 dimensional null space N_0 . Restricting elements in G_0 to N_0 , we obtain a homomorphism

$$\pi: \tilde{C}_7 \longrightarrow \text{GL}_2(\mathbb{F}_7).$$

The last group has $2^6 3^2 7$ elements. Let S be a 2-Sylow subgroup of $\text{GL}_2(\mathbb{F}_7) \cap \{g \mid \det g = \pm 1\}$. Now we set $C_7 := \pi^{-1}(S)$.

Theorem 5.1. The canonical model Sh_C of the Shimura variety obtained by the data as above has exactly three geometrically connected components defined over $\mathbb{Q}(\zeta_7)$. Moreover, for any connected component S , its base change $S \otimes_{\mathbb{Q}(\zeta_7)} \mathbb{Q}_2(\zeta_7)$ is isomorphic to the Mumford's fake projective plane [Mu79] tensored by $\mathbb{Q}_2(\zeta_7)$.

More precisely, the associated rigid analytic space to $\mathrm{Sh}_C \otimes_E E_\lambda$ is isomorphic to Ω/Γ with

$$\Gamma = I(\mathbb{Q}) \bigcap (J(\mathbb{Q}_2) \times C^2) \subset J(\mathbb{Q}_2) \cong \mathrm{PGL}(3, \mathbb{Q}_2),$$

where $I(\mathbb{Q})$ is considered as a subgroup of $I(\mathbb{A}_f) \cong J(\mathbb{Q}_2) \times G(\mathbb{A}_f^2)$. The proof of the theorem is in [Kat98], where two more fake projective planes are discussed as examples of Shimura varieties.

APPENDIX C

p -adic symmetric domains and Totaro's theorem

By TSUZUKI NOBUO

This appendix is a short exposition of M. Rapoport and T. Zink's construction of p -adic symmetric domains [RZ96] and of B. Totaro's theorem [Tot96]. Let G be a connected reductive algebraic group over \mathbb{Q}_p . The set \mathcal{F} of filtrations on an F -isocrystal with G -structure has a structure of a homogeneous space. Rapoport and Zink introduced a p -adic rigid analytic structure on the set \mathcal{F}^{wa} of weakly admissible points in \mathcal{F} . They conjectured that the point in \mathcal{F}^{wa} is characterized by the semistability in the sense of the geometric invariant theory [MFK94] and Totaro proved this conjecture.

1. Weakly admissible filtered isocrystals.

We recall J.-M. Fontaine's definition of weakly admissible filtered F -isocrystals [Fon79].

1.1. Let p be a prime number, k a perfect field of characteristic p , K_0 an absolutely unramified discrete valuation field of mixed characteristics $(0, p)$ with residue field k , \overline{K}_0 an algebraic closure of K_0 , and σ the Frobenius automorphism on K_0 .

Definition 1.2. (1) An F -isocrystal over k , (we simply say "isocrystal"), is a finite dimensional K_0 -vector space V with a bijective σ -linear endomorphism $\Phi : V \rightarrow V$. We denote the category of isocrystals over k by $\text{Isoc}(K_0)$.

(2) For a totally ramified finite extension K of K_0 in \overline{K}_0 , a filtered isocrystal (V, Φ, F^\bullet) over K is an isocrystal (V, Φ) with a decreasing filtration F^\bullet on the K -vector space $V \otimes_{K_0} K$ such that $F^r = V \otimes_{K_0} K$ for $r \ll 0$ and $F^s = 0$ for $s \gg 0$. We denote the category of filtered isocrystals over K by $MF(K)$.

Fontaine also introduced a filtered isocrystal with nilpotent operator N [Fon94]. In this appendix we restrict our attention to filtered isocrystals with $N = 0$.

The category $MF(K)$ is a \mathbb{Q}_p -linear additive category with \otimes and internal Hom 's, but not abelian. A subobject (V', Φ', F'^\bullet) of a filtered isocrystal

(V, Φ, F^\bullet) is a Φ -stable K_0 -subspace V' such that $\Phi' = \Phi|_{V'}$ and $F'^i = (V' \otimes_{K_0} K) \cap F^i$ for all i .

Definition 1.3. Let K be a totally ramified finite extension of K_0 in $\overline{K_0}$. A filtered isocrystal (V, Φ, F^\bullet) over K is weakly admissible if, for any subobject $(V', \Phi', F'^\bullet) \neq 0$, we have

$$\sum_i i \dim_{F'} \mathrm{gr}_{F'}^i(V' \otimes_{K_0} K) \leq \mathrm{ord}_p(\det(\Phi'))$$

and the equality holds for $(V', \Phi', F'^\bullet) = (V, \Phi, F^\bullet)$. Here ord_p is an additive valuation of K_0 normalized by $\mathrm{ord}_p(p) = 1$.

The category of weakly admissible filtered isocrystals is an abelian category which is closed under duals in the category of filtered isocrystals. Fontaine proved that an admissible filtered isocrystal over K , that means a filtered isocrystal arising from a crystalline representation of the absolute Galois group of K via Fontaine's functor, is weakly admissible and conjectured that a weakly admissible filtered isocrystal is admissible in [Fon79]. The category of admissible filtered isocrystals is a \mathbb{Q}_p -linear abelian category with \otimes and duals. Hence, he also conjectured that the category of weakly admissible filtered isocrystals is closed under \otimes , and this was proved by G. Faltings in [Fal95]. (See also [Tot96].)

In [CF00] P. Colmez and Fontaine proved a weakly admissible filtered isocrystal is admissible.

2. Filtered isocrystals with G -structure.

2.1. Let G be a linear algebraic group over \mathbb{Q}_p and denote by $\mathrm{Rep}_{\mathbb{Q}_p}(G)$ the category of finite dimensional \mathbb{Q}_p -rational representations of G . An exact faithful \otimes -functor $\mathrm{Rep}_{\mathbb{Q}_p}(G) \rightarrow \mathrm{Isoc}(K_0)$ is called an isocrystal with G -structure over K_0 .

Let $b \in G(K_0)$. Then, the functor

$$\mathrm{Rep}_{\mathbb{Q}_p}(G) \rightarrow \mathrm{Isoc}(K_0)$$

associated to b , defined by $V \mapsto (V \otimes K_0, b(\mathrm{id} \otimes \sigma))$, is an isocrystal with G -structure over K_0 . [Kot85] Two elements b and b' in $G(K_0)$ are conjugate if and only if there is an element $g \in G(K_0)$ such that $gb\sigma(g)^{-1} = b'$. In this case, g defines an isomorphism between the isocrystals with G -structure associated to b and b' .

If G is connected and k is algebraically closed, then any isocrystal with G -structure over K_0 is associated to an element $b \in G(K_0)$ as above. [RR96]

2.2. Let $\mathbb{D} = \varprojlim \mathbb{G}_m$ be the pro-algebraic group over \mathbb{Q} whose character group is \mathbb{Q}_p . For an element $b \in G(K_0)$, R.E. Kottwitz defined a morphism

$$\nu : \mathbb{D} \rightarrow G_{K_0}$$

of algebraic groups over K_0 which is characterized by the property that, for any object V in $\mathrm{Rep}_{\mathbb{Q}_p}(G)$, the \mathbb{Q} -grading of $V \otimes K_0$ associated to $-\nu$ is the

slope grading of the isocrystal $(V \otimes K_0, b(\text{id} \otimes \sigma))$. (The sign of our ν is different from the one in [Kot85].) For a suitable positive integer s , $s\nu$ is regarded as a one-parameter subgroup of G over K_0 .

Definition 2.3. A σ -conjugacy class \bar{b} of $G(K_0)$ is decent if there is an element $b \in \bar{b}$ such that

$$(b\sigma)^s = s\nu(p)\sigma^s$$

for some positive integer s .

One knows that, for a decent σ -conjugacy class \bar{b} , b and ν as above are defined over \mathbb{Q}_p^s . If G is connected and k is algebraically closed, then any σ -conjugacy class is decent. [Kot85]

2.4. Let K be a totally ramified finite extension of K_0 in $\overline{K_0}$. For a one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow G$ over K and an element $b \in G(K_0)$, we have an exact \otimes -functor

$$\mathcal{I} : \text{Rep}_{\mathbb{Q}_p}(G) \rightarrow MF(K)$$

which is defined by $V \mapsto (V \otimes K_0, b(\text{id} \otimes \sigma), F_\lambda)$. Here $V_{K,\lambda,j}$ is the subspace of $V \otimes K$ of weight j with respect to λ and

$$F_\lambda^i = \bigoplus_{j \geq i} V_{K,\lambda,j}$$

is the weight filtration associated to λ .

Definition 2.5. A pair (λ, b) as above is weakly admissible if and only if the filtered isocrystal $\mathcal{I}(V)$ over K is so for any object V in $\text{Rep}_{\mathbb{Q}_p}(G)$.

To see the weak admissibility for (λ, b) , it is enough to check the weak admissibility of $\mathcal{I}(V)$ for a faithful representation V of G . Indeed, any representation of G appears as a direct summand of $V^{\otimes m} \otimes (V^\vee)^{\otimes n}$ and $\mathcal{I}(V)^{\otimes m} \otimes (\mathcal{I}(V)^\vee)^{\otimes n}$ is weakly admissible by Faltings (see 1.3). Here V^\vee (resp. $\mathcal{I}(V)^\vee$) is the dual of V (resp. $\mathcal{I}(V)$).

3. Totaro's theorem.

In this section we assume that k is algebraically closed.

3.1. Let G be a reductive algebraic group over \mathbb{Q}_p . We fix a conjugacy class of a one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow G$ over $\overline{K_0}$. Here two one-parameter subgroups λ, λ' are conjugate if and only if $g\lambda g^{-1} = \lambda'$ for some element $g \in G(\overline{K_0})$. Then, there is a finite extension E of \mathbb{Q}_p in $\overline{K_0}$ such that the conjugacy class of λ is defined over E . Let us suppose that λ is defined over E and denote by \check{E} the composite field EK_0 in $\overline{K_0}$

3.2. Two one-parameter subgroups of G over $\overline{K_0}$ are equivalent if and only if they define the same weight filtration for any object in $\text{Rep}_{\mathbb{Q}_p}(G)$. Note that, if two one-parameter subgroups are equivalent, then they belong to the same conjugacy class.

Consider the functor

$$R \mapsto \{\text{the equivalence classes in the conjugacy class of } \lambda \text{ defined over } R\}$$

on the category of E -algebras. If one defines an algebraic subgroup of G over E by

$$P(\lambda)(\overline{K}_0) = \{g \in G(\overline{K}_0) \mid g\lambda g^{-1} \text{ is equivalent to } \lambda\},$$

then $P(\lambda)$ is parabolic and the functor above is represented by the projective variety $G_E/P(\lambda)$. We denote this homogeneous space over E by \mathcal{F}_λ . If V is a faithful representation in $\text{Rep}_{\mathbb{Q}_p}(G)$ and if we denote by $\text{Flag}_\lambda(V)$ the flag variety over \mathbb{Q}_p which represents the functor

$$R \mapsto \left\{ \begin{array}{l} \text{the filtrations } F^\bullet \text{ of } V \otimes R \text{ as } R\text{-modules such that} \\ F^i \text{ is a direct summand and } \text{rank}_R F^i = \dim_{\overline{K}_0} F_\lambda^i(V \otimes \overline{K}_0) \end{array} \right\}$$

on the category of \mathbb{Q}_p -algebras, then there is a natural E -closed immersion

$$\mathcal{F}_\lambda \rightarrow \text{Flag}_\lambda(V) \otimes_{\mathbb{Q}_p} E.$$

3.3. Let $b \in G(K_0)$. For a finite extension K of \check{E} , a point ξ in $\mathcal{F}_\lambda(K)$ is called weakly admissible if and only if the pair (ξ, b) is weakly admissible. This condition is independent of the choice of the representative in the equivalence class ξ . We denote by $\mathcal{F}_{\lambda,b}^{wa}(K)$ the subset of weakly admissible points. Totaro gave a characterization of $\mathcal{F}_{\lambda,b}^{wa}$ in the sense of geometric invariant theory. [Tot96] We explain his theory in the rest of this section.

3.4. For a maximal torus T in $G_{\overline{K}_0}$, let $X^*(T)$ be the free abelian group of characters, $X_*(T)$ the free abelian group of one-parameter subgroups, and $\langle , \rangle : X^*(T) \times X_*(T) \rightarrow \mathbb{Z}$ the perfect pairing with $\chi(\xi(t)) = t^{\langle \chi, \xi \rangle}$. If $N(T)$ is the normalizer of T in G , the Weyl group $W(T) = N(T)/T$ acts $X_*(T)$ via inner automorphisms.

Now we fix an invariant norm $\| \cdot \|$ on G , a non-negative real valued function on the set of one-parameter subgroups of $G_{\overline{K}_0}$, such that

a) $\|g\xi g^{-1}\| = \|\xi\|$ for any $g \in G(\overline{K}_0)$,

b) for any maximal torus T , there is a positive definite rational valued bilinear form (\cdot, \cdot) on $X_*(T) \otimes \mathbb{Q}$ with $(\xi, \xi) = \|\xi\|^2$,

c) $\|\gamma(\xi)\| = \|\xi\|$ for $\gamma \in \text{Gal}(\overline{K}_0/K_0)$, where $\gamma(\xi)(t) = \gamma(\xi(t))$.

The bilinear form on $X_*(T) \otimes \mathbb{Q}$ as above is invariant under the action of the Weyl group by (a). For any maximal torus T , invariant norms are in one-to-one correspondence with $(\text{Gal}(\overline{K}_0/K_0), W(T))$ -invariant positive definite rational valued bilinear forms on $X_*(T) \otimes \mathbb{Q}$ since all maximal tori are conjugate and, if $g\xi g^{-1} \in X_*(T)$ for $\xi \in X_*(T)$ and $g \in G(\overline{K}_0)$, then there is $h \in W(T)$ with $g\xi g^{-1} = h\xi h^{-1}$ by [MFK94]. (See also [Kem78] and [Tot96].) Hence, such an invariant norm exists.

3.5. Now we assume that G is connected. Let $U(\lambda)$ be the unipotent radical of $P(\lambda)$, whose elements act on the graded space $\text{gr } F_\lambda^*$ trivially. Then there is a bijection between the set of maximal tori of G in $P(\lambda)$ and the set of

maximal tori of $P(\lambda)/U(\lambda)$ by the natural projection $T \mapsto \bar{T}$. Hence, the invariant norm on G induces the one on $P(\lambda)/U(\lambda)$. Fix a maximal torus T of G . Since the image of λ is contained in the center of $P(\lambda)/U(\lambda)$, the perfect pairing associated to the invariant norm determines the dual of λ in $X^*(\bar{T})$. This dual can extend to a character $\otimes \mathbb{Q}$ of $P(\lambda)/U(\lambda)$. Now we define a G -line bundle $\otimes \mathbb{Q}, L_\lambda$, on \mathcal{F}_λ by the associated one to the negative of the dual character $\otimes \mathbb{Q}$ of λ . By construction, the line bundle $\otimes \mathbb{Q}, L_\lambda$, depends only on the conjugacy class of λ and is ample.

Let J be a smooth affine group scheme over \mathbb{Q}_p such that

$$J(\mathbb{Q}_p) = \{g \in G(K_0) \mid g(b\sigma) = (b\sigma)g\}$$

(which is introduced in [RZ96]). Since $J_{K_0} \subset G_{K_0}$, the pull back $L_{\lambda\check{E}}$ of L_λ on $\mathcal{F}_{\lambda\check{E}}$ is an ample $J_{\check{E}}$ -line bundle.

By the same construction as above, ν in 2.2 gives a character $\otimes \mathbb{Q}$ of $P(\nu)$. The opposite of this character $\otimes \mathbb{Q}$ determines a $J_{\check{E}}$ -action $\otimes \mathbb{Q}$ on the trivial line bundle on $\mathcal{F}_{\lambda\check{E}}$ since $J_{K_0} \subset P(\nu)$. We denote it by L_ν^0 .

We put a $J_{\check{E}}$ -line bundle $\otimes \mathbb{Q}, L = L_{\lambda\check{E}} \otimes L_\nu^0$, on $\mathcal{F}_{\lambda\check{E}}$. Then it is ample and depends only on b and the conjugacy class of λ . We denote by $\mathcal{F}_{\lambda\check{E}}^{ss}(L)$ the set of semistable points in \mathcal{F}_λ with respect to L in the sense of D. Mumford [MFK94].

Theorem 3.6. [Tot96] *Suppose that G is connected and reductive. For any finite extension K of \check{E} , we have*

$$\mathcal{F}_{\lambda,b}^{wa}(K) = \mathcal{F}_{\lambda\check{E}}^{ss}(L)(K).$$

We shall sketch Totaro's proof. First, let $G = GL(n)$ and let us consider the invariant norm induced by the pairing

$$(\alpha, \beta) = \sum_{i,j} ij \dim_{\bar{K}_0} \text{gr}_{F_\alpha^i}^i \text{gr}_{F_\beta^j}^j (K^n)$$

for one-parameter subgroups $\otimes \mathbb{Q}, \alpha, \beta$ of $GL(n)$ over K . If one puts $\mu_\alpha(V) = (\sum_i i \dim_K \text{gr}_{F_\alpha^i}^i(V \otimes K)) / \dim V$, then one has

$$(\alpha, \beta) = \int (\mu_\alpha(F_\beta^j) - \mu_\alpha(V)) \dim_K F_\beta^j dj + \mu_\alpha(V) \mu_\beta(V) \dim V.$$

So, (ξ, b) is weakly admissible if and only if $(\xi, \alpha) + (\nu, \alpha) \leq 0$ for any one-parameter subgroup α of $GL(n)$ over K_0 with the filtration F_α^\bullet as subsocrystals. In other words, (ξ, b) is weakly admissible if and only if $(\xi, \alpha) + (\nu, \alpha) \leq 0$ for any one-parameter subgroup α of J over \mathbb{Q}_p . Hence, the assertion follows from the calculation of Mumford's numerical invariant below.

Lemma 3.7. *If $\mu(\xi, \alpha, L)$ is Mumford's numerical invariant of $\xi \in \mathcal{F}_\lambda$ for a one-parameter subgroup α of J over \mathbb{Q}_p , then*

$$\mu(\xi, \alpha, L) = -(\xi, \alpha) - (\nu, \alpha).$$

Next, let G be arbitrary, V a faithfully representation of G , and consider the invariant norm on G induced from the above norm by the natural immersion $G \rightarrow GL(V)$. The Mumford's numerical invariant of weakly admissible points is non-negative for any one-parameter subgroup of $J(GL(V))$. Hence it is so for any one-parameter subgroup of J , and the weak admissibility implies the semistability. To see the converse, one needs to show that, if $\xi \in \mathcal{F}_\lambda^{ss}(L)(K)$, $(\xi, \alpha) + (\nu, \alpha) \leq 0$ for any one-parameter subgroup α of $J(GL(V))$ over \mathbb{Q}_p . If α is semistable for the G_K -line bundle L_α on \mathcal{F}_α , the assertion follows from the first part. In the case where α is not semistable, one can use Kempf's filtration [Kem78] and Ramanan and Ramanathan's work [RR84], and obtains the required inequality.

Finally one needs to prove the independence of the choice of the norm. Suppose that the identity is valid for the particular norm. Since G is a quotient of a product of a torus and some simple algebraic groups by a finite central subgroup [BT65], one can reduce the assertion in the case of tori and simple groups. In the case of tori it was proved in [RZ96], and in the case of simple groups it is true since the norm of the simple group comes from the Killing form up to a positive rational multiple. \square

4. p -adic symmetric domains.

Let k be the algebraic closure of the prime field \mathbb{F}_p , \mathbb{C}_p the p -adic completion of a fixed algebraic closure $\overline{\mathbb{Q}_p}$ of \mathbb{Q}_p , and $K_0 = \widehat{\mathbb{Q}_p}^{ur}$ the p -adic completion of the unramified extension in \mathbb{C}_p .

4.1. Let G be a reductive group over \mathbb{Q}_p , $b \in G(K_0)$, and fix a conjugacy class $\{\lambda\}$ of a one-parameter subgroup λ of G over $\overline{\mathbb{Q}_p}$.

Rapoport and Zink gave a rigid analytic structure on $\mathcal{F}_{\lambda,b}^{wa}$ as an admissible open subset in $\mathcal{F}_{\lambda,\check{E}}$ and call it the p -adic symmetric domain associated to the triple $(G, \{\lambda\}, b)$ in [RZ96]. This notion of p -adic symmetric domains is different from that of M. van der Put and H. Voskuil in [vdPV92]. Indeed, for any discrete co-compact subgroup Γ of $G(\check{E})$, the quotient $\mathcal{F}_{\lambda,b}^{wa}/\Gamma$ is not always a proper analytic space over \check{E} .

Theorem 4.2. [RZ96] *The set $\mathcal{F}_{\lambda,b}^{wa}$ of weakly admissible points with respect to b in $\mathcal{F}_\lambda(\mathbb{C}_p)$ is an admissible open subset of $\mathcal{F}_{\lambda,\check{E}}$ as a rigid analytic space.*

Now we sketch the proof of the theorem in [RZ96]. By [Kot85] one may assume that the σ -conjugacy class of b is decent with the decent equation $(b\sigma)^s = s\nu(p)\sigma^s$ as in 2.3. Let V be a faithful representation in $\text{Rep}_{\mathbb{Q}_p}(G)$, $V_s = V \otimes \mathbb{Q}_{p^s}$, and $\Phi_s = b(\text{id} \otimes \sigma)$. Then $(V \otimes K_0, b(\text{id} \otimes \sigma)) = (V_s, \Phi_s) \otimes_{\mathbb{Q}_{p^s}} K_0$. Put $V_s = \bigoplus_\lambda V_{s,j}$ to be the isotypical decomposition for Φ_s . The functor

$$R \mapsto \{V' \subset V_s \otimes_{\mathbb{Q}_{p^s}} R \mid V' \text{ is a direct summand with } V' = \bigoplus_j V' \cap (V_{s,j} \otimes_{\mathbb{Q}_{p^s}} R)\}$$

on the category of \mathbb{Q}_{p^s} -algebras is represented by a disjoint sum T' of closed subschemes of Grassmannians of V_s . T' descends to a \mathbb{Q}_p -variety T and one

has

$$T(\mathbb{Q}_p) = \{\Phi_s\text{-stable subspaces of } V_s\}.$$

Indeed, Φ_s gives a descent datum $\alpha : T' \rightarrow T'^\sigma$, where $T'^\sigma(R)$ is a set of direct summands of $V_{s,\lambda} \otimes_{\mathbb{Q}_{p^s},\sigma} R$ with the isotypical decomposition, and $\alpha^{s-1} \circ \dots \circ \alpha : T' \rightarrow T'$ is the identity by the decent equation.

Consider the closed subscheme over \mathbb{Q}_{p^s}

$$\mathcal{H} \subset (\text{Flag}_\lambda(V) \times T) \otimes_{\mathbb{Q}_p} \mathbb{Q}_{p^s}$$

which consists of pairs (F^\bullet, V') such that

$$\sum_i i \text{rank gr}_{F^\bullet \cap V'}^i(V') > \text{ord}_p(\det(\Phi_s|_{V'})).$$

Then, by the definition of weak admissibility, one has

$$\mathcal{F}_{\lambda,b}^{wa}(\mathbb{C}_p) = \mathcal{F}_\lambda(\mathbb{C}_p) \cap \left(\text{Flag}_\lambda(V)(\mathbb{C}_p) - \bigcup_{t \in T(\mathbb{Q}_p)} \mathcal{H}_t \right),$$

where $\mathcal{F}_\lambda(\mathbb{C}_p)$ is identified with the image of the immersion $\mathcal{F}_\lambda(\mathbb{C}_p) \subset \text{Flag}_\lambda(\mathbb{C}_p)$.

Fix embeddings of $\text{Flag}_\lambda(V)$ and T in projective spaces over \mathbb{Q}_p and a finite set $\{f_j\}$ of bi-homogeneous polynomials of definition of $\text{Flag}_\lambda(V) \times T$ with integral coefficients. For $\epsilon > 0$, consider a tubular neighbourhood

$$\mathcal{H}_t(\epsilon) = \{x \in \text{Flag}_\lambda(V)(\mathbb{C}_p) \mid |f_j(x, t)| < \epsilon \text{ for all } j\}$$

of \mathcal{H}_t . Here we choose unimodular representatives for x and t 's and $|\cdot|$ is an absolute value on \mathbb{C}_p . Then there is a finite set $S \subset T(\mathbb{Q}_p)$ such that $\bigcup_{t \in T(\mathbb{Q}_p)} \mathcal{H}_t(\epsilon) = \bigcup_{t \in S} \mathcal{H}_t(\epsilon)$ for the local compactness. $\mathcal{F}_\epsilon = \mathcal{F}_\lambda(\mathbb{C}_p) \cap (\text{Flag}_\lambda(V)(\mathbb{C}_p) - \bigcup_{t \in T(\mathbb{Q}_p)} \mathcal{H}_t(\epsilon))$ is an admissible open subset of $\mathcal{F}_{\lambda,b\check{E}}^{wa}$, hence

$$\mathcal{F}_1 \subset \mathcal{F}_{\frac{1}{2}} \subset \mathcal{F}_{\frac{1}{3}} \subset \dots$$

is an admissible covering of $\mathcal{F}_{\lambda,b\check{E}}^{wa}$. □

Bibliography

- [AB01] Y. André and F. Baldassarri, *De Rham cohomology of differential modules on algebraic varieties*, Progress in Mathematics, vol. 189, Birkhäuser, 2001.
- [And90] Y. André, *p-adic Betti lattices*, in: *p-adic analysis*, Proc. Trento conf. on p-adic analysis (F. Baldassarri, S. Bosch, and B. Dwork, eds.), Lecture Notes in Math, vol. 1454, Springer, 1990, 23–63.
- [And92] Y. André, *Réalisation de Betti des motifs p-adiques*, preprint IHES, 1992.
- [And95] Y. André, *Théorie des motifs et interprétation géométrique des valeurs p-adiques de G-fonctions*, Number theory (S. David, ed.), Cambridge Univ. Press., 1995, 37–60.
- [And96] Y. André, *G-fonctions et transcendance*, J. Reine Angew. Math. **476** (1996), 95–125.
- [And98] Y. André, *p-adic orbifolds and p-adic triangle groups*, R.I.M.S. Kokyuroku, vol. 1073, 1998, 136–159.
- [And02a] Y. André, *Représentations galoisiennes et opérateurs de Bessel p-adiques*, preprint, to appear in Ann. Inst. Fourier, 2002.
- [And02b] Y. André, *Filtrations de type Hasse-Arf et monodromie p-adique*, Invent. Math. **148** (2002), 285–317.
- [And] Y. André, *A geometric description of $Gal(\overline{\mathbb{Q}}_p/\mathbb{Q}_p)$ and a p-adic avatar of $\hat{G}T$* , preprint, 2002.
- [Bal82] F. Baldassarri, *Differential modules and singular points of p-adic differential equations*, Adv. in Math. **44** (1982), 155–179.
- [BBM82] P. Berthelot, L. Breen, and W. Messing, *Théorie de Dieudonné cristalline II*, Lecture Notes in Math., vol. 930, Springer, 1982.
- [BC91] J.F. Boutot and H. Carayol, *Uniformisation p-adique des courbes de Shimura: les théorèmes de Čerednik et de Drinfeld*, in: Astérisque, vol. 196–197, S.M.F., 1991, 45–158.
- [BD98] M. Bertolini and H. Darmon, *Heegner points, p-adic l-functions, and the Čerednik-Drinfeld uniformization*, Invent. Math. **131** (1998), 453–491.
- [BDR80] S. Bosch, B. Dwork, and P. Robba, *Un théorème de prolongement pour des fonctions analytiques*, Math. Ann. **252** (1980), 165–173.
- [Ber90] V.G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Mathematical Surveys and Monographs, no. 33, A.M.S., 1990.
- [Ber93] V.G. Berkovich, *Étale cohomology for non-Archimedean analytic spaces*, Inst. Hautes Études Sci. Publ. Math. **78** (1993), 5–161.
- [Ber99] V.G. Berkovich, *Smooth p-adic analytic spaces are locally contractible*, Invent. Math. **137** (1999), no. 1, 1–84.
- [Be84] P. Berthelot, *Cohomologie rigide et théorie de Dwork: le cas des sommes exponentielles*, S.M.F. Astérisque **119–120** (1984), 17–49.
- [Be86] P. Berthelot, *Géométrie rigide et cohomologie des variétés algébriques de caractéristique $p > 0$* , Mémoire de la SMF **23** (1986), 7–32.
- [Be96] P. Berthelot, *Cohomologie rigide et cohomologie rigide à supports propres*, preprint IRMAR 96-03 (1996).

- [BGR84] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean Analysis*, Springer-Verlag, 1984, Grundlehren der Math. Wiss.
- [BL93] S. Bosch and W. Lütkebohmert, *Formal and rigid geometry I, Rigid spaces*, Math. Ann. **295** (1993), 291–317.
- [BLR90] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Math., vol. 21, Springer, 1990.
- [BM79] P. Berthelot and W. Messing, *Théorie de Dieudonné cristalline I*, Astérisque **63** (1979), 17–38.
- [BeO83] P. Berthelot and A. Ogus, *F-isocrystals and De Rham cohomology I*, Invent. Math. **72** (1983), 159–199.
- [Bou64] N. Bourbaki, *Algèbre I*, deuxième ed., Hermann, 1964.
- [Bou74] N. Bourbaki, *Topologie générale*, Hermann/CCLS, 1974.
- [Bou83] N. Bourbaki, *Variétés différentielles et analytiques*, fascicule de résultats, CCLS, Paris, 1983.
- [Bre99] C. Breuil, *Representations semi-stables et modules fortement divisibles*, Invent. Math. **136** (1999), 89–122.
- [Bre00] C. Breuil, *Groupes p -divisibles, groupes finis et modules filtrés*, Annals of Math. **152** (2000), 489–549.
- [BT65] A. Borel and J. Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. **27** (1965), 55–151.
- [BZ95] J.F. Boutot and T. Zink, *The p -adic uniformization of Shimura curves*, preprint 95-107, Sonderforschungsbereich 343, Univ. Bielefeld., 1995.
- [Car61] C. Carathéodory, *Funktionentheorie II*, Birkhäuser, 1961.
- [CC89] D. Chudnovsky and G. Chudnovsky, *Computational problems in arithmetic of linear differential equations. some diophantine applications*, Number Theory, Lecture Notes in Math., vol. 1383, Springer, 1989, 12–49.
- [Čer76] I.V. Čerednik, *Uniformization of algebraic curves by discrete subgroup of $\mathrm{PGL}_2(k_w)$ with compact quotients*, Math. USSR-Sb. **29** (1976), 55–78.
- [CF00] P. Colmez and J.-M. Fontaine, *Construction des représentations p -adiques semi-stables*, Invent. Math. **140** (2000), no. 1, 1–43.
- [Cho96] F. Choucroun, *Groupes d'automorphismes et frontières d'arbres: le cas homogène*, Progress in probability, vol. 40, Birkhäuser, 1996, 209–221.
- [C86] G. Christol, *Fonctions et éléments algébriques*, Pacific J. Math. **125** (1986), 1–37.
- [Chl98] Chambert-Loir, *Cohomologie cristalline: un survol*, Exposition. Math. **16** (1998), 333–382.
- [CM97] G. Christol and Z. Mebkhout, *Sur le théorème de l'indice des équations différentielles p -adiques II*, Ann. of Math. **146** (1997), 345–410.
- [CM00] G. Christol and Z. Mebkhout, *Sur le théorème de l'indice des équations différentielles p -adiques III*, Annals of Math. **151** (2000), 385–457.
- [Col90] R. Coleman, *On the Frobenius matrices of Fermat curves*, in: p -adic analysis, Proc. Trento conf. on p -adic analysis (F Baldassarri, S. Bosch, and B. Dwork, eds.), Lecture Notes in Math, vol. 1454, 1990, 173–193.
- [Co91] P. Colmez, *Périodes des variétés abéliennes à multiplication complexe*, Ann. of Math. **138** (1993), 625–683.
- [Co92] P. Colmez, *Périodes p -adiques des variétés abéliennes*, Math. Ann. **292** (1992), 629–644.
- [Co94] P. Colmez, *Les nombres algébriques sont denses dans \mathbf{B}_{dR}^+* , Périodes p -adiques, Astérisque, vol. 223, S.M.F., 1994.
- [Co98] P. Colmez, *Intégration p -adique*, Astérisque, vol. 248, S.M.F., 1998.
- [Cox74] H.M.S. Coxeter, *Regular complex polytopes*, 1991, 2nd ed., Cambridge Univ. Press, 1974.

- [Cox79] H.M.S. Coxeter, *The non-euclidian symmetry of Escher's picture "circle limit III"*, Leonardo (1979), 19–25, Pergamon Press
- [CR94] G. Christol and P. Robba, *Equations différentielles p -adiques; applications aux sommes exponentielles*, Hermann, 1994.
- [Cr85] R. Crew, *F-isocrystals and p -adic representations*, Algebraic Geometry - Bowdoin, Proc. Symp. Pure Math. **17** (1985), 111–138.
- [Cr00] R. Crew, *Canonical Extensions, Irregularities, and the Swan Conductor*, Math. Ann. **316** (2000), 19–37.
- [CW01] P. Cohen and G. Wüstholz, *A panorama in Number Theory*, ch. Applications of the André-Oort conjecture to transcendence, Cambridge Univ. Press, 2001.
- [Del71] P. Deligne, *Travaux de Shimura*, Sémin. Bourbaki 389, Lecture Notes in Math., vol. 244, Springer, 1971.
- [Del74] P. Deligne, *Théorie de Hodge III*, Inst. Hautes Études Sci. Publ. Math. **44** (1974), 5–78.
- [Del89] P. Deligne, *Le groupe fondamental de la droite projective moins trois points*, Galois groups over \mathbb{Q} , Publ. Math. Sci. Res. Inst., vol. 16, Springer Verlag, 1989, Proc. Workshop, Berkeley/CA (USA) 1987, 79–297.
- [dJ95a] A.J. de Jong, *Crystalline Dieudonné module theory via formal and rigid geometry*, Inst. Hautes Études Sci. Publ. Math. **82** (1995), 5–96, erratum ibid. **87** (1998), 175.
- [dJ95b] A.J. de Jong, *Étale fundamental groups of non-archimedean analytic spaces*, Compositio Math. **97** (1995), 89–118.
- [dJvdP96] A.J. de Jong and M. van der Put, *Étale cohomology of rigid analytic spaces*, Doc. Math. **1** (1996), 1–56.
- [DM93] P. Deligne and G. Mostow, *Commensurabilities among lattices in $PU(1, n)$* , Princeton Univ. press, 1993.
- [Dri76] V.G. Drinfeld, *Coverings of p -adic symmetric regions*, Functional. Anal. Appl. **10** (1976), 29–40.
- [dSh87] E. de Shalit, *Iwasawa theory of elliptic curves with complex multiplication*, Acad. Press, 1987.
- [Dw69] B. Dwork, *p -adic cycles*, Inst. Hautes Études Sci. Publ. Math. **37** (1969), 27–115.
- [Dw74] B. Dwork, *Bessel functions as p -adic functions of the argument*, Duke Math. J. **41** (1974), 711–738.
- [Dw82] B. Dwork, *Lectures on p -adic differential equations*, Springer, NY, 1982.
- [Dw83] B. Dwork, *On the Boyarsky principle*, Amer. J. Math. **285** (1983), 115–156.
- [Elk98] N. Elkies, *Shimura curve computations*, Algorithmic number theory (J.P. Buhler, ed.), L. N. Comput. Sci., vol. 1423, Springer, 1998, 3rd international symposium, ANTS-III, Portland, O, USA, June 21–25 Proceedings, 1–47.
- [Esc71] M.C. Escher, *The world of M.C. Escher*, New York H. N. Abrams, 1971.
- [Fal95] G. Faltings, *Mumford-Stabilität in der algebraischen Geometrie*, Proceedings of ICM 1994 (Zürich), Birkhäuser, 1995, 648–655.
- [FM87] J.-M. Fontaine and W. Messing, *p -adic periods and p -adic étale cohomology*, Contemporary Mathematics **67** (1987), 179–207.
- [Fon79] J.-M. Fontaine, *Modules galoisiens, modules filtrés et anneaux de Barsotti-Tate*, Astérisque, vol. 65, Société Mathématique de France, 1979.
- [Fon94] J.-M. Fontaine, *Représentations p -adiques semi-stables*, Périodes p -adiques, Séminaire de Bures, 1988, Astérisque, **223**, S.M.F., (1994), 113–184.
- [For29] L. Ford, *Automorphic functions*, 1951, 2nd ed., Chelsea, New York, 1929.
- [FvdP81] J. Fresnel and M. van der Put, *Géométrie analytique rigide et applications*, Progress in Math, vol. 18, Birkhäuser, Boston, Basel, Stuttgart, 1981.
- [GH94a] B. Gross and M. Hopkins, *Equivariant vector bundles on the Lubin-Tate moduli space*, A.M.S. Contemp. Math. **158** (1994), 23–88.


- [GH94b] B. Gross and M. Hopkins, *The rigid analytic period mapping, Lubin-Tate space, and stable homotopy theory*, Bull. Amer. Math. Soc. (N.S.) **30** (1994), 76–86.
- [GI63] O. Goldmann and N. Iwahori, *The space of p -adic norms*, Acta. Math. **109** (1963), 137–177.
- [Gor01] U. Görtz, *On the flatness of models of certain Shimura varieties of PEL type*, Math. Ann. **321** (2001), 689–727.
- [Gou97] F.G. Gouvêa, *p -adic numbers, An Introduction*, 2nd ed., Universitext, Springer-Verlag, 1997.
- [Gra86] J. Gray, *Linear differential equations and group theory from Riemann to Poincaré*, Birkhäuser, 1986.
- [Gre81] L. Greenberg, *Homomorphisms of triangle groups into $PSL(2, \mathbb{C})$* , Riemann surfaces and related topics, Ann. Math. Studies, vol. 97, 1981, Proc. 1978 Stony Brook Conf., 167–181.
- [Gri71] P. Griffiths, *Periods of integrals on algebraic manifolds III*, Inst. Hautes Études Sci. Publ. Math. **38** (1971), 125–180.
- [Gro70] A. Grothendieck, *Groupes de Barsotti-Tate et cristaux*, Actes, Congrès Intern. Math., no. 1, 1970, 431–436.
- [Gro74] A. Grothendieck, *Groupes de Barsotti-Tate et cristaux de Dieudonné*, Presses de l'Univ. Montréal, 1974.
- [Gun67] R. Gunning, *Special coordinate covering of Riemann surfaces*, Math. Ann. **170** (1967), 67–86.
- [GvdP80] L. Gerritzen and M. van der Put, *Schottky groups and Mumford curves*, Lecture Notes in Math., vol. 817, Springer-Verlag, 1980.
- [H81] E. Hecke, *Vorlesungen über die Theorie der algebraischen Zahlen, (Lectures on the theory of algebraic numbers)*, vol. 77, Springer verlag, 1981.
- [Heh75] D. Hejhal, *On Schottky and Teichmüller spaces*, Adv. in Math. **15** (1975), 133–156.
- [Her80a] F. Herrlich, *Die Ordnung der Automorphismengruppe einer p -adischen Schottkykurve*, Math. Ann. **246** (1980), 125–130.
- [Her80b] F. Herrlich, *Endlich erzeugbare p -adische diskontinuierliche Gruppen*, Arch. Math. **35** (1980), 505–515.
- [Hub96] R. Huber, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, vol. 30, Vieweg-Verlag, Wiesbaden, 1996.
- [Hu87] D. Husemöller, *Elliptic curves*, Graduate Texts in Mathematics, vol. 111, Springer Verlag, 1987.
- [Ic97] T. Ichikawa, *Schottky uniformization theory on Riemann surfaces and Mumford curves of infinite genus*, J. reine angew. Math. **486** (1997), 45–68.
- [IK98] M.-N. Ishida and F. Kato, *The strong rigidity theorem for non-archimedean uniformization*, Tôhoku Math. J. (2) **50** (1998), 537–555.
- [Iv86] B. Iversen, *Cohomology of sheaves*, Universitext, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1986.
- [Ka72] N. Katz, *Algebraic solutions of differential equations. p -curvature and the Hodge filtration*, Invent. Math. **18** (1972), 1–118.
- [Ka73] N. Katz, *Travaux de Dwork, Séminaire Bourbaki*, Lecture Notes in Math **317** (1973), 167–200.
- [Ka81] N. Katz, *Serre-Tate local moduli*, Surface Algébriques, Lecture Notes in Math., vol. 868, Springer, 1981, 138–202.
- [Ka90] N. Katz, *Exponential sums and differential equations*, Annals of Math. Studies, vol. 124, Princeton University press, 1990.
- [Kat98] F. Kato, *On p -adic uniformization of fake projective planes*, preprint: Max-Planck-Institut für Mathematik Bonn (1998).
- [Kat01] F. Kato, *p -adic schwarzian triangle groups of mumford type*, preprint, 2001.


- [Kem78] G. Kempf, *Instability in invariant theory*, Ann. of Math. **108** (1978), 299–316.
- [Kie67a] R. Kiehl, *Der Endlichkeitssatz für eigentliche Abbildungen in der nichtarchimedischen Funktionentheorie*, Invent. Math. **2** (1967), 191–214.
- [Kie67b] R. Kiehl, *Die de Rham Kohomologie algebraischer Mannigfaltigkeiten über einem bewerteten Körper*, Inst. Hautes Études Sci. Publ. Math. **33** (1967), 5–20.
- [Kie67c] R. Kiehl, *Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie*, Invent. Math. **2** (1967), 256–273.
- [Ko79] N. Koblitz, *The hypergeometric function with p -adic parameters*, Proceedings of the Queen's Number Theory Conference (Ontario), 1979, 319–328.
- [Ko80] N. Koblitz, *p -adic analysis: a short course on recent works*, London L.N.M., vol. 42, Cambridge Univ. Press, 1980.
- [Kot85] R.E. Kottwitz, *Isocrystals with additional structure*, Compositio Math. **56** (1985), 201–220.
- [Kot92] R.E. Kottwitz, *Points on some Shimura varieties over finite fields*, J. Amer. Math. Soc. **5** (1992), 373–444.
- [Kra96] D. Kramer, *An example of an arithmetic Fuchsian group*, J. reine angew. Math. **473** (1996), 69–85.
- [Kur79] A. Kurihara, *On some examples of equations defining Shimura curves and the Mumford uniformization*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. (1979), 277–300.
- [Kur80] A. Kurihara, *Construction of p -adic unit balls and the Hirzebruch proportionality*, Amer. J. Math. **24** (1980), 129–174.
- [Laf79] G. Laffaille, *Construction de groupes p -divisibles: le cas de dimension 1*, Astérisque **65** (1979), 103–124.
- [La90] S. Lang, *Cyclotomic Fields I and II*, Graduate Text in Math., Springer-Verlag, 1990.
- [LiO98] K. Li and F. Oort, *Moduli of supersingular abelian varieties*, Lecture Notes in Math., vol. 1680, Springer, 1998.
- [Lüt90] W. Lütkebohmert, *Formal-algebraic and rigid-analytic geometry*, Math. Ann. **286** (1990), 341–371.
- [Lüt93] W. Lütkebohmert, *Riemann's existence problem for a p -adic field*, Invent. Math. **111** (1993), 309–330.
- [Mag74] W. Magnus, *Non-euclidian tessellations and their groups*, vol. 61, Acad. Press, 1974.
- [May91] D. Mayer, *Continued fractions and related transformations*, Ergodic theory, symbolic dynamics, and hyperbolic spaces (Ser. C., ed.), Oxford University Press, 1991, 175–222.
- [Mer54] S.N. Mergelyan, *Uniform approximations to functions of a complex variable*, Trans. Amer. Math. Soc. **101** (1954), 31–122.
- [Mes72] W. Messing, *The crystals attached to Barsotti-Tate groups*, Lecture Notes in Math., vol. 264, Springer, 1972.
- [Mes76] W. Messing, $q_{\text{Serre-Tate}} = q_{\text{Dwork}}$, Notices Amer. Math. Soc., 1976.
- [MFK94] D. Mumford, J. Fogarty, and F. Kirwan, *Geometric Invariant Theory*, third enlarged edition ed., Ergebnisse der Math., vol. 34, Springer-Verlag, 1994.
- [Mic81] J.F. Michon, *Courbes de Shimura hyperelliptiques*, Bull. Soc. Math. France **109** (1981), 217–225.
- [Mic84] J.F. Michon, *Courbes de Shimura*, Séminaire de Théorie des Nombres Paris 1982–1983, Progress in Mathematics, vol. 51, Birkhäuser, 1984.
- [MM74] B. Mazur and W. Messing, *Universal extensions and one-dimensional crystalline cohomology*, Lecture Notes in Math., vol. 370, Springer, 1974.
- [Moc96] S. Mochizuki, *A theory of ordinary p -adic curves*, Publ. R.I.M.S. Kyoto **32** (1996), 957–1151.


- [MoRo69] E. Motzkin and P. Robba, *Prolongement analytique en analyse p -adique*, Séminaire de Théorie des Nombre, Bordeaux, 1968–1969, multigraphed.
- [Mu72a] D. Mumford, *An analytic construction of degenerating abelian varieties over complete rings*, Compositio Math. **24** (1972), 239–272.
- [Mu72b] D. Mumford, *An analytic construction of degenerating curves over complete local rings*, Compositio Math. **24** (1972), 129–174.
- [Mu79] D. Mumford, *An algebraic surface with K ample, $(K^2) = 9$, $p_g = q = 0$* , Amer. J. of Math. **101** (1979), 233–244.
- [Mus78] G.A. Mustafin, *Nonarchimedean uniformization*, Math. USSR-Sb. **34** (1978), 187–214.
- [Oes84] J. Oesterlé, *Nombres de Tamagawa et groupes unipotents en caractéristique*, Invent. Math. **78** (1984), 13–88.
- [Ogg83] A. Ogg, *Real points on Shimura curves*, papers dedicated to Shafarevich, Progress in Math, vol. 35, Birkhäuser, 1983.
- [O90] A. Ogus, *A p -adic analogue of the Chowla-Selberg formula*, in: p -adic analysis, Proc. Trento conf. on p -adic analysis (F Baldassarri, S. Bosch, and B. Dwork, eds.), Lecture Notes in Math, vol. 1454, Springer, 1990, 319–341.
- [O00] A. Ogus, *Elliptic crystals and modular motives*, preprint (2000).
- [P00] G. Pappas, *On the arithmetic moduli schemes of PEL Shimura varieties*, J. of Alg. Geom. **9** (2000), no. 3, 577–605.
- [Ram98] L. Ramero, *On a class of étale analytic sheaves*, J. Algebraic Geom. **7** (1998), no. 3, 405–504.
- [Rap92] A. Rapinchuk, *Congruence subgroup problem for algebraic groups: Old and new*, S.M.F. Astérisque **209** (1992), 73–84.
- [Ray74] M. Raynaud, *Géométrie analytique rigide d’après Tate, Kiehl,.. table ronde d’analyse non-archimédienne*, Bull. Soc. math. France **Mém. 39–40** (1974), 319–327.
- [Ray94] Raynaud, *Revêtements de la droite affine en caractéristique $p > 0$ et conjecture d’Abhyankar*, Invent. Math. **116** (1994), 425–462.
- [Re89] E. Reyssat, *Quelques aspects des surfaces de Riemann*, Progress in Math., vol. 77, Birkhäuser, 1989.
- [Ro75] P. Robba, *Solutions bornées des systèmes différentiels linéaires. Application aux fonctions hypergéométriques*, Groupe d’étude d’analyse ultramétrique, exp. 5, 1975.
- [Ro86] P. Robba, *Une introduction naïve aux cohomologies de Dwork*, Mém. Soc. Math. France (N.S.) **23** (1986), 61–105.
- [RR84] S. Ramanan and A. Ramanathan, *Some remarks on the instability flag*, Tôhoku Math. J. **36** (1984), 269–291.
- [RR96] M. Rapoport and M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, Compositio Math. **103** (1996), 153–181.
- [RZ96] M. Rapoport and Th. Zink, *Period space for p -divisible groups*, Ann. of Math. Study, vol. 141, Princeton University press, 1996.
- [Sai94] K. Saito, *Algebraic representation of the Teichmüller spaces*, The Grothendieck Theory of Dessins d’Enfants (L. Schneps, ed.), vol. 200, London Math. Soc., 1994.
- [Schn37] T. Schneider, *Arithmetische Untersuchungen elliptischer Integrale*, Math. Ann. **113** (1937), 1–13.
- [Scho85] A.J. Scholl, *A trace formula for F -crystals*, Invent. Math. **79** (1985), 31–48.
- [Sch99] H. Schoutens, *Embedded resolution of singularities in rigid analytic geometry*, Ann. Fac. Sci. Toulouse, VI. Ser. **8** (1999), no. 2, 297–330.
- [Ser67] J.-P. Serre, *Complex multiplication*, Algebraic Number Theory (J. Cassels and A. Fröhlich, eds.), Academic Press, 1967.
- [Ser68a] J.-P. Serre, *Abelian l -adic representations and elliptic curves*, Benjamin, 1968.


- [Ser68b] J.-P. Serre, *Corps locaux*, Hermann, 1968.
- [Ser77] J.-P. Serre, *Arbres, amalgames, sl_2* , Astérisque, vol. 46, S.M.F., 1977.
- [Ser94] J.-P. Serre, *Cohomologie galoisienne*, 5ème ed., vol. 5, Springer, 1994.
- [Shi67] G. Shimura, *Construction of class fields and zeta functions of algebraic curves*, Ann. of Math. **85** (1967), 58–159.
- [Shi71] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Iwanami Shoten–Princeton Univ. Press, 1971, Kanô memorial lectures.
- [Si90] C. Simpson, *Transcendental aspects of the Riemann–Hilbert correspondence*, Illinois J. Math. **34** (1990), no. 2.
- [Si94] C. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety I*, Inst. Hautes Études Sci. Publ. Math. **79** (1994), 47–129.
- [SvdP97] M. Singer and M. van der Put, *Galois theory of difference equations*, Lecture Notes in Math., vol. 1666, Springer, 1997.
- [SPM91] J. Stienstra, M. van der Put, and B. van der Marel, *On p -adic monodromy*, Math. Z. **208** (1991), 309–325.
- [Tak77] K. Takeuchi, *Commensurability classes of arithmetic triangle groups*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **24** (1977), 201–212.
- [Ta66] J. Tate, *p -divisible groups*, Proc. of the conference on Local Fields, 1966, 158–183.
- [Ta71] J. Tate, *Rigid analytic spaces*, Invent Math. **12** (1971), 257–289.
- [Ta93] J. Tate, *A review of non-archimedean elliptic functions*, Proc. of the conference on Elliptic Curves and Modular Forms (Hong-Kong), International Press, 1993, 162–184.
- [Ter73] T. Terada, *Problème de Riemann et fonctions automorphes provenant des fonctions hypergéométriques de plusieurs variables*, J. Math. Kyoto Univ. **13** (1973), 557–578.
- [Tot96] B. Totaro, *Tensor products in p -adic Hodge theory*, Duke Math. J **56** (1996), 79–104.
- [Tsu98] N. Tsuzuki, *Finite local monodromy of overconvergent unit-root F -isocrystal on a curve*, Amer. J. of Math. **120** (1998), 1165–1190.
- [Uen87] K. Ueno, *Compact rigid analytic spaces — with special regard to surfaces*, Advanced Studies in Pure Math., vol. 10, 1987, Algebraic Geometry, Sendai, 1985, 765–794.
- [Var98a] Y. Varshavsky, *p -adic uniformization of unitary Shimura varieties*, Inst. Hautes Études Sci. Publ. Math. **57** (1998), 57–119.
- [Var98b] Y. Varshavsky, *p -adic uniformization of unitary Shimura varieties II*, J. Differential Geom. **49** (1998), 75–113.
- [vdM89] B. van der Marel, *Formal groups and differential equations*, Master’s thesis, Groningen, 1989.
- [vdP82] M. van der Put, *Cohomology on affinoid spaces*, Compositio Math. **45** (1982), 165–198.
- [vdP83] M. van der Put, *Étale coverings of a Mumford curve*, Ann. Inst. Fourier **33** (1983), no. 1, 29–52.
- [vdP87] M. van der Put, *A note on p -adic uniformization*, Indag. Math. **49** (1987), 313–318.
- [vdP88] M. van der Put, *Monodromie d’une équation différentielle p -adique*, Groupe d’étude d’analyse ultramétrique, Publ. Université Paris 7, 1987/88.
- [vdP89] M. van der Put, *Les courbes de shimura*, J. Th. Nombres Bordeaux, Ser. II **1** (1989), no. 1, 89–102.
- [vdP92a] M. van der Put, *Discrete groups, Mumford curves and theta functions*, Ann. Fac. Sci. Toulouse, Ser. Math. 1 **VI** (1992), no. 3, 399–438.
- [vdP92b] M. van der Put, *Uniformization of varieties over local fields*, Nieuw Arch. Wiskd. **IV** (1992), no. 1-2, 57–75, Ser. 10.


- [vdPV92] M. van der Put and H. Voskuil, *Symmetric spaces associated to split algebraic groups over a local field*, J. Reine Angew. Math. **433** (1992), 69–100.
- [vdPV00] M. van der Put and H. Voskuil, *Discontinuous subgroups of $PGL_2(K)$* , preprint University Groningen, 2000.
- [Vig80] M.F. Vignéras, *Arithmétique des algèbres de quaternions*, Lecture Notes in Math., vol. 800, Springer, 1980.
- [Wa69] W. Waterhouse, *Abelian varieties over finite fields*, Ann. Sci. École Norm. Sup. (4) **2** (1969), 521–560.
- [Wol88] J. Wolfart, *Wertehypergeometrischer Funktionen*, Invent. Math. **92** (1988), 187–216.
- [Yo87] M. Yoshida, *Fuchsian differential equations*, Aspects of Math., vol. 11, Vieweg, 1987.
- [Yo97] M. Yoshida, *Hypergeometric functions, my love (modular interpretation of configuration spaces)*, Aspects of Math., vol. 32, Vieweg, Publ. Max Planck Inst., 1997.
- [You92] P.T. Young, *Apéry numbers, Jacobi sums, and special values of generalized p -adic hypergeometric functions*, J. Number Theory **41** (1992), 231–255.
- [Yu95] J.K. Yu, *On the moduli of quasi-canonical liftings*, Compositio Math. **96** (1995), no. 3, 293–321.
- [EGA I] A. Grothendieck and J. Dieudonné, *Éléments de Géométrie Algébrique I*, Inst. Hautes Études Sci. Publ. Math., vol. 4, 1960.
- [EGA IV] A. Grothendieck and J. Dieudonné, *Éléments de Géométrie Algébrique IV, Étude locale des schémas et des morphismes de schémas (quatrième partie)*, Inst. Hautes Études Sci. Publ. Math., vol. 32, 1967.
- [SGA 1] A. Grothendieck, *Revêtements étales et groupe fondamental (SGA1)*, Lecture Notes in Math., vol. 224, Springer, 1971.











Grothendieck's
Mysterious Functor

