# Projective normality of special scrolls.

Luis Fuentes García<sup>\*</sup> Manuel Pedreira Pérez

Authors' address: Departamento de Algebra, Universidad de Santiago de Compostela. 15706 Santiago de Compostela. Galicia. Spain. e-mail: pedreira@zmat.usc.es;

luisfg@usc.es

**Abstract:** We study the projective normality of a linearly normal special scroll R of degree d and speciality i over a smooth curve X of genus g. If  $d \ge 4g - 2i + 1$ , we prove that the projective normality of the scroll is equivalent to the projective normality of its directrix curve of minimum degree.

Mathematics Subject Classifications (1991): Primary, 14J26; secondary, 14H25, 14H45. Key Words: Ruled Surfaces, projective normality.

### Introduction.

Let  $R \subset \mathbf{P}^N$  be a scroll, that is, a surface with a one-dimensional family of lines. We say that R is linearly normal when R is not the projection of any other scroll with the same degree and genus. It is known that there are a unique geometrically ruled surface  $\pi : S = \mathbf{P}(\mathcal{E}) \longrightarrow X$  and a unique base point free unisecant linear system |H| on S such that R is the birational image of  $S = \mathbf{P}(\mathcal{E})$  by the regular map defined by the complete linear system |H|. We define the speciality of R as  $i = h^1(\mathcal{O}_S(H))$ . The scroll is special when i > 0.

In this paper we study the projective normality of a special scroll R. We say that R is projectively normal, or equivalently, that the line bundle  $\mathcal{O}_S(H)$  on Sis normally generated, if and only if the natural maps  $Sym_k(H^0(\mathcal{O}_S(H))) \longrightarrow$  $H^0(\mathcal{O}_S(kH))$  are surjective for any  $k \ge 1$ . We say that the cokernel of these maps are the speciality of R respect to hypersurfaces of degree k.

The question of the normal generation of a line bundle  $\mathcal{L}$  on a smooth curve X has been studied with detail. The classical result of Noether says that the canonical bundle is normally generated; the Theorem of Castelnuovo states that any line bundle of degree at least 2g + 1 is normally generated. More recently, H. Lange and G. Martens in [9] and M. Green and R. Lazarsfeld in [7] gave bounds, depending of the Clifford index of X, over the degree of a line bundle to be normally generated.

There are not complete generalizations of these theorems to higer dimension.

<sup>\*</sup>Supported by an F.P.U. fellowship of Spanish Government

If V is a projective variety of dimension n, most of the results study the normal generation of adjoint bundles, that is, line bundles  $\omega_V \otimes \mathcal{L}$  where  $\mathcal{L}$  is an ample line bundle.

M. Andreatta and J. Sommese in [1] and L.Ein and R.Lazarsfled in [3] proved that  $\omega_V \otimes \mathcal{L}^{\otimes n} \otimes \mathcal{F}$  is normally generated if it is very ample,  $\mathcal{L}$  is very ample and  $\mathcal{F}$  is numerically effective. D. Butler in [2] has stabilished that if V is a ruled variety of rank n over a smooth curve X, then  $\omega_V \otimes \mathcal{L}^{\otimes 2n+1}$  is normally generated if  $\mathcal{L}$  is ample.

However, the canonical divisor of a ruled surface S is  $\mathcal{K}_S \sim -2X_0 + (\mathcal{K} + \mathfrak{e})f$ , so in general, it is not easy to write an unisecant linear system like  $\mathcal{K}_S + 2A + B$  with |A| very ample and |B| numerically effective. Furthermore, an ample divisor on S must be at least unisecant so the result of D. Butler applies at least to 3-secant linear systems on S. Then, we will use more particular strategies to the case of ruled surfaces. We try to reduce the study of the normal generation of a line bundle  $\mathcal{O}_S(H)$  to the normal generation of line bundles on the base curve X. We will prove that:

The projective normality of a special scroll of genus g, speciality i and degree  $d \ge 4g - 2i + 1$  is equivalent to the projective normality of its directrix curve of minimum degree.

We refer to [4] for a systematic development of the projective theory of scrolls and ruled surfaces that we will use in this paper and to [5] to study the special scrolls. Anyway, in the first section we recall some basic facts that we will use along the paper.

## 1 Preliminars.

A geometrically ruled surface, or simply a ruled surface, will be a  $\mathbf{P}^1$ -bundle over a smooth curve X of genus g > 0. It will be denoted by  $\pi : S = \mathbf{P}(\mathcal{E}_0) \longrightarrow X$ . We will suppose that  $\mathcal{E}_0$  is a normalized sheaf and  $X_0$  is the section of minimum self-intersection that corresponds to the surjection  $\mathcal{E}_0 \longrightarrow \mathcal{O}_X(\mathfrak{e}) \longrightarrow 0$ ,  $\bigwedge^2 \mathcal{E} \cong \mathcal{O}_X(\mathfrak{e})$  (see [8],V, §2 and [4]).

If  $|H| = |X_0 + \mathbf{b}f|$  is a base-point-free linear system on a ruled surface S, |H| defines a regular map  $\phi_H : S \longrightarrow \mathbf{P}(H^o(\mathcal{O}_S(H)^{\vee}))$ . The image of S is a scroll R. If  $\phi_H$  is a birational map we say that S and H are the ruled surface and the linear system associated to the scroll R. We denote de image of a curve  $C \subset S$  by  $\overline{C} \subset R$ . The curve  $\overline{X_0}$  is the curve of minimum degree of R. It is embedded by the linear system  $|\mathbf{b} + \mathbf{c}|$  on X. We will use the well known fact that, if  $m \ge 0$  and  $i \ge 0$ :

$$h^{i}(\mathcal{O}_{S}(mX_{0}+\mathfrak{b}f)) \leq \sum_{k=0}^{m} h^{i}(\mathcal{O}_{X}(\mathfrak{b}+k\mathfrak{e}))$$

Furthermore, if  $\mathfrak{b}$  is nonspecial

$$h^{i}(\mathcal{O}_{S}(X_{0} + \mathfrak{b}f)) = h^{i}(\mathcal{O}_{X}(\mathfrak{b})) + h^{i}(\mathcal{O}_{X}(\mathfrak{b} + \mathfrak{e}))$$

(see [4]).

**Definition 1.1** Let V be a projective variety. Let  $\mathcal{F}_i$ , i = 1, ..., s, coherent sheaves on V. We call  $s(\mathcal{F}_1, ..., \mathcal{F}_s)$  the cokernel of the map:

$$H^0(\mathcal{F}_1) \otimes \ldots \otimes H^0(\mathcal{F}_s) \longrightarrow H^0(\mathcal{F}_1 \otimes \ldots \otimes \mathcal{F}_s)$$

If  $\mathcal{F}_i$  are invertible sheaves  $\mathcal{O}_V(D_i)$  where  $D_i$  are divisors on V, we will write  $s(D_1, \ldots, D_s)$ .

**Lemma 1.2** If  $s(\mathcal{F}_1, \mathcal{F}_2) = 0$ , then:

$$s(\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3, \dots, \mathcal{F}_s) = s(\mathcal{F}_1 \otimes \mathcal{F}_2, \mathcal{F}_3, \dots, \mathcal{F}_s)$$

**Proof:** It is sufficient note that  $s(\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3, \dots, \mathcal{F}_s)$  is the cokernel of the composition:

$$H^{0}(\mathcal{F}_{1})\otimes\ldots\otimes H^{0}(\mathcal{F}_{s})\to H^{0}(\mathcal{F}_{1}\otimes\mathcal{F}_{2})\otimes H^{0}(\mathcal{F}_{3})\otimes\ldots\otimes H^{0}(\mathcal{F}_{s})\to H^{0}(\mathcal{F}_{1}\otimes\ldots\otimes\mathcal{F}_{s})$$

**Definition 1.3** Let V be a projective variety and let |H| be a complete unisecant base-point-free linear system defining a birational map:

$$\phi_H: V \longrightarrow \overline{V} \subset \mathbf{P}^N$$

We say that  $(V, \mathcal{O}_V(H))$  is projectively normal or  $\mathcal{O}_V(H)$  normally generated or  $\overline{V}$  projectively normal if and only if the natural maps:

$$\operatorname{Sym}_k(H^0(\mathcal{O}_V(H))) \longrightarrow H^0(\mathcal{O}_V(kH))$$

are surjective for all  $k \geq 1$ .

**Remark 1.4** Thus,  $(V, \mathcal{O}_V(H))$  is projectively normal if and only if  $s(H, .\frac{k}{\cdot}, H) = 0$ , for all  $k \geq 2$ . Moreover,  $\dim(s(H, .\frac{k}{\cdot}, H))$  is the speciality of  $\overline{V}$  respect to hypersurfaces of degree k.

**Lemma 1.5 (Green)** Let  $\mathfrak{a},\mathfrak{b}$  be effective divisors on a smooth curve X. Let  $\mathfrak{b}$  be base-point-free. If  $h^1(\mathcal{O}_X(\mathfrak{a}-\mathfrak{b})) \leq h^0(\mathcal{O}_X(\mathfrak{b})) - 2$  then  $s(\mathfrak{a},\mathfrak{b}) = 0$ .

**Proof:** This is a particular case of  $H^{o}$ -Lemma in [6].

### 2 Projective normality of special scrolls.

**Proposition 2.1** Let S be a ruled surface and  $H \sim X_0 + \mathfrak{b}f$  an unisecant linear system, such that  $\mathfrak{b}$  is a nonspecial divisor and  $\mathfrak{b} + \mathfrak{e}$  is effective. Let  $k \geq 2$ . If

 $s(\mathbf{b} + \mathbf{e}, \mathbf{i}, \mathbf{b} + \mathbf{e}, \mathbf{b}, \mathbf{k}, \mathbf{i}, \mathbf{b}) = 0$  for all i, with  $0 \le i \le k - 1$ 

then:

$$s(H, \overset{k}{\ldots}, H) \cong s(\mathfrak{b} + \mathfrak{e}, \overset{k}{\ldots}, \mathfrak{b} + \mathfrak{e})$$

**Proof:** Let i, j be non negative integers, with i + j > 0. We will denote by  $W_{i,j}$  the tensor product:

$$H^0(\mathcal{O}_S(H))\otimes . ! . \otimes H^0(\mathcal{O}_S(H))\otimes H^0(\mathcal{O}_S(H-X_0))\otimes . ! . \otimes H^0(\mathcal{O}_S(H-X_0))$$

With this notation it verifies that:

- 1.  $W_{i,j} \otimes H^0(\mathcal{O}_S(H)) = W_{i+1,j}$ .
- 2.  $W_{i,j} \otimes H^0(\mathcal{O}_S(H X_0)) = W_{i,j+1}$
- 3. We have an exact sequence:

$$W_{i,j} \to H^0(\mathcal{O}_S((i+j)H - jX_0))) \to s(H, .., H, H - X_0, .., H - X_0) \to 0$$

4. The map  $H^0(\mathcal{O}_S(H)) \longrightarrow H^0(\mathcal{O}_{X_0}(H))$  is a surjection, because  $\mathfrak{b}$  is nonspecial. Moreover, we know that  $H^0(\mathcal{O}_S(S-X_0)) \cong H^0(\mathcal{O}_X(\mathfrak{b}))$  and  $H^0(\mathcal{O}_{X_0}(H)) \cong H^0(\mathcal{O}_X(\mathfrak{b} + \mathfrak{e}))$ . From this, we have the following exact sequence:

$$W_{i-1,j} \otimes H^0(\mathcal{O}_{X_0}(H)) \longrightarrow H^0(\mathcal{O}_X((i+j)\mathfrak{b}+i\mathfrak{e})) \longrightarrow$$
$$\longrightarrow s(\mathfrak{b}+\mathfrak{e}, ..., \mathfrak{b}+\mathfrak{e}, \mathfrak{b}, ..., \mathfrak{b}) \to 0$$

We will prove that  $s(H, :i., H, H - X_0, :::, H - X_0) = 0$  if  $0 \le i \le k - 1$ .

We proceed by induction on i:

1. If i = 0, because  $H^0(\mathcal{O}_S(H - X_0)) \cong H^0(\mathcal{O}_X(\mathfrak{b}))$  we immediately obtain that:

$$s(H - X_0, \overset{k}{\ldots}, H - X_0) = s(\mathfrak{b}, \overset{k}{\ldots}, \mathfrak{b})$$

and this is zero by hypotesis.

2. Now, suppose that the assertion is true for i - 1, with  $1 \le i \le k - 1$ . Consider the exact sequence:

$$0 \longrightarrow H^0(\mathcal{O}_S(H - X_0)) \longrightarrow H^0(\mathcal{O}_S(H)) \longrightarrow H^0(\mathcal{O}_{X_0}(H)) \longrightarrow 0$$

Taking the tensor product with  $W_{i-1,k-i}$  we obtain the following commutative diagram:

where

$$coker(\alpha_{1i}) = s(H, \stackrel{i-1}{\dots}, H, H - X_0, \stackrel{k-(i-1)}{\dots}, H - X_0)$$
$$coker(\alpha_{2i}) = s(H, \stackrel{i}{\dots}, H, H - X_0, \stackrel{k-i}{\dots}, H - X_0)$$
$$coker(\alpha_{3i}) = s(\mathfrak{b} + \mathfrak{e}, \stackrel{i}{\dots}, \mathfrak{b} + \mathfrak{e}, \mathfrak{b}, \stackrel{k-i}{\dots}, \mathfrak{b})$$

Moreover,

$$h^{1}(\mathcal{O}_{S}(kH - (k - (i - 1))X_{0})) \leq \sum_{l=0}^{i-1} h^{1}(\mathcal{O}_{X}((k - l)\mathfrak{b} + l(\mathfrak{b} + \mathfrak{e})))$$

Because  $\mathbf{b}$  is nonspecial and  $\mathbf{b} + \mathbf{c}$  is effective, if  $i \leq k$  and  $0 \leq l \leq i-1$  then  $(k-l)\mathbf{b} + l(\mathbf{b} + \mathbf{c})$  is nonspecial. Therefore,  $h^1(\mathcal{O}_S(kH - (k-(i-1))X_0))) = 0$ .

By induction hypothesis,  $s(H, \stackrel{i-1}{\ldots}, H, H - X_0, \stackrel{k-(i-1)}{\ldots}, H - X_0) = 0$  and by the hypothesis of the theorem  $s(\mathbf{b} + \mathbf{e}, \stackrel{i}{\ldots}, \mathbf{b} + \mathbf{e}, \mathbf{b}, \stackrel{k-i}{\ldots}, \mathbf{b}) = 0$  when  $i \leq k-1$ . Thus, since  $\alpha_{1i}$  and  $\alpha_{3i}$  are surjections, we deduce that  $\alpha_{2i}$  is a surjection, and then  $s(H, \stackrel{i}{\ldots}, H, H - X_0, \stackrel{k-i}{\ldots}, H - X_0) = 0$ .

Finally, from the above diagram for i = k, we see that  $\alpha_{1k}$  is a surjection, because  $s(H, \stackrel{k-1}{\ldots}, H, H - X_0) = 0$ . Therefore,  $coker(\alpha_{2k}) \cong coker(\alpha_{3k})$  and,

$$s(H, \stackrel{k}{\ldots}, H) \cong s(\mathfrak{b} + \mathfrak{e}, \stackrel{k}{\ldots}, \mathfrak{b} + \mathfrak{e})$$

**Corollary 2.2** Let S be a geometrically ruled surface. Let  $|H| = |X_0 + \mathfrak{b}f|$  an unisecant complete linear system on S, such that  $\mathfrak{b}$  is a nonspecial divisor and  $\mathfrak{b} + \mathfrak{e}$  is effective. If

 $s(\mathbf{b} + \mathbf{e}, \mathbf{i}, \mathbf{b} + \mathbf{e}, \mathbf{b}, \mathbf{k}, \mathbf{i}, \mathbf{b}) = 0$  for all i, with  $0 \le i \le k$ 

for all  $k \geq 2$  then  $(S, \mathcal{O}_S(H))$  is projectively normal.

**Proposition 2.3** Let  $R \subset \mathbf{P}^N$  be a special linearly normal scroll of genus g, degree d and speciality i. Let S and  $|H| = |X_0 + \mathfrak{b}f|$  be the ruled surface and the linear systema asocciated to R.

If  $\mathfrak{b}$  is nonspecial then the curve of minimum degree of R is special, linearly normal and its speciality coincides with the specialty of R. Moreover, if  $deg(\mathfrak{b}) \geq 2g - 1$  then this is the unique special curve of the scroll.

**Proof:** If  $\mathfrak{b}$  is nonspecial we know that:

 $h^1(\mathcal{O}_S(H)) = h^1(\mathcal{O}_X(\mathfrak{b})) + h^1(\mathcal{O}_X(\mathfrak{b} + \mathfrak{e}))$ 

Thus, the speciality of the curve of minimum degree  $\overline{X_0}$  is  $h^1(\mathcal{O}_X(\mathfrak{b} + \mathfrak{e})) = h^1(\mathcal{O}_S(H)) = i$ . Furthermore,  $\overline{X_0}$  is linearly normal because we have a surjective map:

$$H^0(\mathcal{O}_S(H)) \longrightarrow H^0(\mathcal{O}_{X_0}(H)) \longrightarrow H^1(\mathcal{O}_S(\mathfrak{b})) = 0$$

If  $deg(\mathfrak{b}) \geq 2g - 1$ , the degree of the scroll satisfies  $d \geq 2g - 1 + deg(\overline{X_0})$ . Any other directrix curve of R different from  $\overline{X_0}$  has degree  $d' \geq d - deg(\overline{X_0}) = 2g - 1 > 2g - 2$ , so it is nonspecial.

**Theorem 2.4** Let  $R \subset \mathbf{P}^N$  be a special linearly normal scroll of genus g, degree d and speciality i. Suppose that  $d \geq 4g - 2i + 1$ . Then:

- 1. R has an unique special directrix curve  $\overline{X_0}$ . Moreover,  $\overline{X_0}$  is the curve of minimum degree, it is linearly normal and it has the speciality of R.
- R and X<sub>0</sub> have the same specialty respect to hypersurfaces of degree m. In particular the scroll is projectively normal if and only if the curve of minimum degree if projectively normal.

**Proof:** Let S be the ruled surface and  $|H| = |X_0 + \mathfrak{b}f|$  the linear system corresponding to the scroll R.

1. Since R is special, it has a special directrix curve (see [5]) so the curve  $\overline{X_0}$  of minimum degree of the scroll verifies  $deg(\mathbf{b} + \mathbf{e}) \leq 2g - 2$ .

Suppose that  $d \ge 4g - 2i + 1$ . We know that:

$$d - 2g + 2 + i = h^0(\mathcal{O}_S(H)) \le h^0(\mathcal{O}_X(\mathfrak{b} + \mathfrak{e})) + h^0(\mathcal{O}_X(\mathfrak{b}))$$

If  $deg(\mathfrak{b}) \leq 2g$ , then we can apply Clifford Theorem ([8], page 343) to the divisors  $\mathfrak{b}$  and  $\mathfrak{b} + \mathfrak{e}$ . From the above inequiality we obtain:

$$d - 2g + 2 + i \le \frac{\deg(\mathbf{b} + \mathbf{c})}{2} + 1 + \frac{\deg(\mathbf{b})}{2} + 1 = \frac{d}{2} + 2$$

and then  $d \leq 4g - 2i$ .

Thus, we see that  $deg(\mathfrak{b}) \geq 2g+1$  and now we can apply the Proposition above.

2. We will apply the Proposition 2.1. We have to proof that:

$$s(\mathbf{b} + \mathbf{e}, \mathbf{i}, \mathbf{b} + \mathbf{e}, \mathbf{b}, \mathbf{k}, \mathbf{i}, \mathbf{b}) = 0$$
 with  $0 \le i \le k - 1$ 

for all  $k \geq 2$ .

If  $\mathbf{b} + \mathbf{e} \sim 0$ , then the scroll is a cone and it verifies the condition trivialy. Thus, we will suppose that  $\mathbf{b} + \mathbf{e} \nsim 0$ .

Since  $deg(\mathfrak{b}) \geq 2g+1$ , we can use the Theorem of Castelnuovo to see that  $\mathcal{O}_X(\mathfrak{b})$  is normally generated and then  $s(\mathfrak{b}, .k., \mathfrak{b}) = 0$  for any  $k \geq 2$ . From this, applying Lemma 1.2:

$$s(\mathfrak{b}, ..i., \mathfrak{b}, \mathfrak{b} + \mathfrak{e}, \frac{k-i}{\ldots}, \mathfrak{b} + \mathfrak{e}) = s(i\mathfrak{b}, \mathfrak{b} + \mathfrak{e}, \frac{k-i}{\ldots}, \mathfrak{b} + \mathfrak{e})$$

when  $1 \leq i \leq k$ .

Let us see that  $s(\mathbf{b}, \mathbf{b} + \mathbf{c}) = 0$ . We apply the Lemma 1.5. We need that:

$$h^1(\mathcal{O}_X(\mathfrak{b} - (\mathfrak{b} + \mathfrak{e}))) < h^0(\mathcal{O}_X(\mathfrak{b} + \mathfrak{e})) - 1$$

Note that  $e = -deg(\mathfrak{e}) \ge 0$  because  $deg(\mathfrak{b}) \ge 2g+1$  and  $deg(\mathfrak{b}+\mathfrak{e}) \le 2g-2$ . We distinguish to cases:

- (a) If  $-\mathfrak{e}$  is nonspecial, it is sufficient to check that  $h^0(\mathcal{O}_X(\mathfrak{b}+\mathfrak{e})) > 1$ . But  $|\mathfrak{b}+\mathfrak{e}|$  is base point free and  $\mathfrak{b}+\mathfrak{e} \not\sim 0$ , so  $h^0(\mathcal{O}_X(\mathfrak{b}+\mathfrak{e})) > 1$ .
- (b) If  $-\mathfrak{e}$  is special, we apply Clifford Theorem ([8], page 343):

$$|-\mathfrak{e}| \leq \frac{1}{2}e$$
, or equivalently,  $h^1(\mathcal{O}_X(-\mathfrak{e})) \leq g - \frac{e}{2}$ 

But,

$$h^0(\mathcal{O}_X(\mathfrak{b}+\mathfrak{e})) - 1 = deg(\mathfrak{b}) - e - g + 1 + i - 1 = deg(\mathfrak{b}) - e - g + i$$

By hypotesis  $d \ge 4g - 2i + 1$ , so  $2deg(\mathfrak{b}) - e \ge 4g - 2i + 1$  and  $deg(\mathfrak{b}) \ge \frac{4g - 2i + 1 + e}{2}$ . Then:

$$\begin{split} h^0(\mathcal{O}_X(\mathfrak{b}+\mathfrak{e})) - 1 &\geq \frac{4g - 2i + 1 + e}{2} - e - g + i \geq g - \frac{e}{2} + \frac{1}{2} \geq \\ &\geq h^1(\mathcal{O}_X(-\mathfrak{e})) + \frac{1}{2} > h^1(\mathcal{O}_X(-\mathfrak{e})) \end{split}$$

so we can apply the Lemma of Green 1.5 and we obtain  $s(\mathfrak{b}, \mathfrak{b}+\mathfrak{e}) = 0$ .

Now, let us see that  $s(\lambda \mathbf{b} + \mu \mathbf{c}, \mathbf{b} + \mathbf{c}) = 0$  when  $\lambda \ge 1$  and  $0 \le \mu \le \lambda - 1$ :

- (a) If  $\lambda = 1$ ,  $\mu = 0$  and it is proved.
- (b) Suppose that  $\lambda > 1$ . Note that  $(\lambda 1)\mathbf{b} + (\mu 1)\mathbf{c}$  is a nonspecial divisor, because  $\mathbf{b}$  is nonspecial and  $\mathbf{b} + \mathbf{c}$  is effective. Therefore,  $h^1(\mathcal{O}_X((\lambda 1)\mathbf{b} + (\mu 1)\mathbf{c})) < h^0(\mathcal{O}_X(\mathbf{b} + \mathbf{c})) 1$ , because  $|\mathbf{b} + \mathbf{c}|$  is base point free and different from 0. Applying the Lemma of Green 1.5 we deduce that  $s(\lambda\mathbf{b} + \mu\mathbf{c}, \mathbf{b} + \mathbf{c}) = 0$ .

Finally, applying the Lemma 1.2, we see that:

 $s(i\mathfrak{b},\mathfrak{b}+\mathfrak{e},\overset{k-i}{\ldots},\mathfrak{b}+\mathfrak{e}) = s(i\mathfrak{b}+(k-i-1)\mathfrak{b}+\mathfrak{e},\mathfrak{b}+\mathfrak{e}) = 0$ 

when  $1 \leq i \leq k-1$ ,  $k \geq 2$  and the conclusion follows.

**Remark 2.5** If d < 4g - 2i + 1 the conclusions of the Theorem could fail. Let us see an example:

Suppose that X is an hyperelliptic curve of genus  $g \ge 3$ . Let  $\mathfrak{b}$  a very ample nonspecial divisor of degree 2g and let  $\mathfrak{b} + \mathfrak{e} = \sum_{1}^{n} g_{2}^{1}$  be a special divisor, with  $0 \le n \le g-1$ . Consider the decomposable ruled surface  $\mathbf{P}(\mathcal{O}_{X}(\mathfrak{b}) \oplus \mathcal{O}_{X}(\mathfrak{b} + \mathfrak{e}))$ and the uniscant linear system  $H \sim X_{0} + \mathfrak{b}f$ . We obtain an scroll of degree d = 2g + 2m and speciality i = g - n. In this way d = 4g - 2i. Since the scroll is decomposable, we know that:

 $dim(s(H,H)) = dim(s(\mathfrak{b},\mathfrak{b})) + dim(s(\mathfrak{b}+\mathfrak{e},\mathfrak{b}+\mathfrak{e})) + dim(s(\mathfrak{b}+\mathfrak{e},\mathfrak{b}))$ 

But X is hyperelliptic and  $deg(\mathbf{b}) = 2g$ , so  $(X, \mathcal{O}_X(\mathbf{b})$  is not projectively normal (see [7]) and  $dim(s(\mathbf{b}, \mathbf{b})) > 0$ . Thus, the scroll and the curve of minimum degree have not the same specialty respect to quadrics.

## References

- [1] ANDREATTA, M; SOMMESE, A. On the projective normality of the adjunction bundles. Comment. Math. Helvetici **66** (1991), 162-167.
- BUTLER, D Normal generation of vector bundles over a curve. J.Differential Geometry 39 (1994), 1-34.
- [3] EIN, L; LAZARSFELD, R Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension. Invent. math. 111 (1993), 51-67.
- [4] FUENTES, L.; PEDREIRA, M. The Projective Theory of Ruled Surfaces. Preprint. math.AG/0006204.

- [5] FUENTES, L.; PEDREIRA, M. Canonical geometrically ruled surfaces. Preprint. math.AG/0107114.
- [6] GREEN, M Koszul cohomology and the geometry of projective varieties.
  J. Differential Geometry 19 (1984), 125-171.
- [7] GREEN, M; LAZARSFELD, R. On the projective normality of complete linear series on an algebraic curve. Invent. Math. 83 (1986), pp. 73-90.
- [8] HARTSHORNE, R. Algebraic Geometry. GTM, 52. Springer-Verlag, 1977.
- [9] LANGE, H.; MARTENS, G. Normal generation and presentation of line bundles of low degree. J. Reine Angew Math 356 (1985), pp. 1-18.