

Dimension of the Moduli Space of curves with an involution.

Luis Fuentes García* Manuel Pedreira Pérez

Authors' address: Departamento de Algebra, Universidad de Santiago de Compostela.
15706 Santiago de Compostela. Galicia. Spain. e-mail: pedreira@zmat.usc.es;
luisfg@usc.es

Abstract: Given a smooth curve X of genus g we compute the dimension of the family of curves C which have an involution over X . Moreover we distinguish when the curve C is hyperelliptic.

Mathematics Subject Classifications (1991): Primary, 14H10; secondary, 14H30, 14H37.

Key Words: Curves, involution.

Introduction. Let \mathcal{M}_π be the Moduli of smooth curves of genus π . It is well known that a generic smooth curve $C \in \mathcal{M}_\pi$ with $\pi \geq 3$ does not have nontrivial automorphism. In particular a generic smooth curve does not have involutions.

However, given an smooth curve X of genus g and a divisor $\mathbf{b} \in \text{Div}(X)$ verifying that $2(\mathbf{b} - \mathcal{K})$ is smooth we can construct a double cover $\gamma : C \longrightarrow X$, where C is a smooth curve of genus $\pi \geq 2g - 1$. C is not generic because it has an involution induced by γ . Moreover, we can characterize the divisors \mathbf{b} which provides double covers $\gamma : C \longrightarrow X$ with C hyperelliptic (see [2]). Thus if we denote by \mathcal{C}_π^g the family of curves of genus π with an involution of genus g and by \mathcal{Ch}_π^g the family of hyperelliptic curves of genus π with an involution of genus g , we have that:

$$\mathcal{Ch}_\pi^g \subset \mathcal{C}_\pi^g \subset \mathcal{M}_\pi.$$

In this paper we compute the dimension of \mathcal{C}_π^g and \mathcal{Ch}_π^g . Let $r = 2(\pi - 1) - 4(g - 1)$. The main results are:

1. If $r \geq 0$ then $\dim(\mathcal{C}_\pi^g) = 2\pi - g - 1$. Moreover, the dimension of the family of curves with an involution over a hyperelliptic curve of genus g is $2\pi - 2g + 1$.
2. If $r < 0$ then $\dim(\mathcal{C}_\pi^g) = \emptyset$.

*Supported by an F.P.U. fellowship of Spanish Government

3. If $r = 0, 2, 4$ then $\dim(\mathcal{C}h_\pi^g) = \pi$.
4. If $r < 0$ or $r > 4$ then $\mathcal{C}h_\pi^g = \emptyset$.

Moreover, we prove that $\mathcal{C}h_\pi^g = \mathcal{C}h_\pi^{\pi-g}$. In fact we prove that given an hyperelliptic curve C of genus $\pi \geq 2$, an involution of C of genus g composed with the canonical involution provides an involution of genus $\pi - g$.

1 Preliminaries.

Let C, X be two smooth curves of genus π and g respectively. Let $\gamma : C \longrightarrow X$ be a double cover. We know the following facts (see [2]):

1. $\gamma_*\mathcal{O}_C \cong \mathcal{O}_X \oplus \mathcal{O}_X(\mathcal{K} - \mathfrak{b}) = S_{\mathfrak{b}}$ is a decomposable geometrically ruled surface over the curve X . We call it canonical geometrically ruled surface. \mathfrak{b} is a nonspecial divisor on X verifying $2\mathfrak{b} - 2\mathcal{K} \sim \mathcal{B}$, where \mathcal{B} is the branch divisor. If X_0 is the curve of minimum self-intersection of $S_{\mathfrak{b}}$ and $X_1 \sim X_0 + (\mathfrak{b} - \mathcal{K})f$ then $C \sim 2X_1$. In particular $C \in \langle 2X_0 + \mathcal{B}f, 2X_1 \rangle$.
2. Conversely, let $\mathcal{O}_X \oplus \mathcal{O}_X(\mathcal{K} - \mathfrak{b})$ be a decomposable geometrically ruled surface over the curve X , such that $2\mathfrak{b} - 2\mathcal{K}$ is a smooth divisor. Then the generic curve C in the linear system $|2X_1|$ is smooth and then we have a double cover $C \longrightarrow X$. Moreover, if $C \in \langle 2X_0 + \mathcal{B}f, 2X_1 \rangle$, with $\mathcal{B} \sim 2\mathfrak{b} - 2\mathcal{K}$ then the branch divisor of the cover is \mathcal{B} .

In order to compute the dimension of the curves with an involution over a fixed curve X , we have to study when two curves in the linear system $|2X_1|$ are isomorphic. In this way we have the following proposition:

Proposition 1.1 *Let X be a smooth curve of genus g . Let \mathfrak{b} be a nonspecial divisor of degree $b \geq 2g - 2$ defining a canonical ruled surface $S_{\mathfrak{b}}$. Let $\mathcal{B} \sim 2\mathfrak{b} - 2\mathcal{K}$ be different points. Then, there is a unique curve $C \in |2X_1|$ up to isomorphism with a $2 : 1$ map $\gamma : C \longrightarrow X$ whose ramification points over X are the points of \mathcal{B} .*

Proof: We know that given a curve $X_1 \in |X_0 + (\mathfrak{b} - \mathcal{K})f|$, a generic curve C of the pencil $L = \langle 2X_0 + \mathcal{B}f, 2X_1 \rangle$ is a curve with an involution $\gamma : C \longrightarrow X$ and ramification points at \mathcal{B} (see Lemma 1.9 and Theorem 1.10 in [2]). This curve is invariant by the unique involution of $S_{\mathfrak{b}}$ that fixes the curves X_0 and X_1 . In this way, C meets each generator in two points and these points are related by the involution.

Let C and C' be two curves of the pencil L . Let Pf be a generic generator. We can define an automorphism of $S_{\mathfrak{b}}$ that fixes $X_0 \cap Pf$, $X_1 \cap Pf$ and takes a point of $C \cap Pf$ into a point of $C' \cap Pf$. If we consider the restriction of this automorphism to the pencil L we see that it takes C into C' and the two curves are isomorphic.

Now, let X_1 and X'_1 two irreducible curves of the linear system $|X_0 + (\mathfrak{b} - \mathcal{K})f|$. Since $S_{\mathfrak{b}}$ is a decomposable ruled surface we can define an automorphism of $S_{\mathfrak{b}}$ that takes X_1 into X'_1 . In this way the curves of the pencils $L = \langle 2X_0 + \mathcal{B}f, 2X_1 \rangle$ and $L = \langle 2X_0 + \mathcal{B}f, 2X'_1 \rangle$ are isomorphic and our claim follows. ■

2 Computing the dimensions.

Proposition 2.1 *Let \mathcal{C}_π^X be the family of smooth curves of genus $\pi \geq 1$ with an involution over a smooth curve X of genus $g > 0$. Let $r = 2(\pi - 1) - 4(g - 1)$.*

1. *If $r > 0$ then $\dim(\mathcal{C}_\pi^X) = r - \dim(\text{Aut}(X))$.*
2. *If $r = 0$ then $\dim(\mathcal{C}_\pi^X) = 0$.*
3. *If $r < 0$ then $\mathcal{C}_\pi^X = \emptyset$.*

Proof: Let C a smooth curve with an involution over the curve X of genus g . Let $\gamma : C \rightarrow X$ be the $2 : 1$ map. By Hurwitz Theorem we know that the map X has $r = 2(\pi - 1) - 4(g - 1)$ ramifications. If r is negative $\mathcal{C}_\pi^X = \emptyset$.

1. Suppose that $r > 0$. Consider the following incidence variety:

$$J = \{(C, \mathcal{B}) \in \mathcal{C}_\pi^X \times U^r / C \text{ has an involution branched at } \mathcal{B} \in X\}$$

where $U_r \subset S^r X$ are the open set of r unordered different points. We have two projection maps: $p : J \rightarrow \mathcal{C}_\pi^X$ and $q : J \rightarrow U^r$.

Given a curve $C \in \mathcal{C}_\pi^X$ there is a $2 : 1$ map $C \rightarrow X$ with r ramifications. From this, p is a surjection. Moreover, since $\pi > 1$ the group of automorphisms of C is finite, so there are a finite number of involutions of C over X . In this way we obtain a finite number of possible ramification points of X up to automorphisms of X . But if $g = 1$, $\dim(\text{Aut}(X)) = 1$ and $r \geq 1$ and if $g > 1$, $\dim(\text{Aut}(X)) = 0$. From this there is at most a finite number of automorphism fixing r generic points. Therefore, $\dim(p^{-1}(C)) = \dim(\text{Aut}(X))$.

On the other hand, given a set $\beta \in X$ of r different points we can take a divisor $\mathfrak{b} \in \text{Div}(X)$ such that $2\mathfrak{b} - 2\mathcal{K} \sim \beta$. Let $S_{\mathfrak{b}}$ be the corresponding

canonical ruled surface. By Proposition 1.1, there is a unique curve $C \in |2X_1| \subset S_{\mathbf{b}}$ with an involution over X with branch points over the set β , so q is a surjection. Furthermore, we know that a curve C with an involution over the curve X lays on the linear system $|2X_1|$ of a canonical ruled surface. Since there are a finite number of divisors \mathbf{b} satisfying $2\mathbf{b} - 2\mathcal{K} \sim \mathbf{b}$ (see [2]), we see that $\dim(q^{-1}(C)) = 0$ and $\dim(J) = \dim(U^r) = r$.

Thus, we have:

$$\dim(\mathcal{C}_\pi^X) = \dim(J) - \dim(p^{-1}(C)) = r - \dim(\text{Aut}(X)).$$

2. Suppose that $r = 0$. In this case there are not ramification points. Given a curve $C \in \mathcal{C}_\pi^X$ we know that lays on the canonical system $|2X_1|$ of a canonical ruled surface $S_{\mathbf{b}}$ with $2\mathbf{b} - 2\mathcal{K} \sim 0$. All curves of this system are isomorphic (Proposition 1.1). Moreover, there are a finite number of divisors \mathbf{b} verifying $2\mathbf{b} \sim 2\mathcal{K}$. Thus $\dim(\mathcal{C}_\pi^X) = 0$. ■

Proposition 2.2 *Let \mathcal{C}_π^X be the family of smooth hyperelliptic curves of genus $\pi \geq 1$ with an involution over a smooth curve X of genus g . Let $r = 2(\pi - 1) - 4(g - 1)$. Then:*

1. *If X is neither elliptic nor hyperelliptic then $\mathcal{C}_\pi^X = \emptyset$.*
2. *If X is elliptic or hyperelliptic then:*
 - (a) *If $r > 4$ ($\pi > 2g + 1$) or $r < 0$ ($\pi < 2g - 1$) then $\mathcal{C}_\pi^X = \emptyset$.*
 - (b) *if $r = 4$ ($\pi = 2g + 1$) then $\dim(\mathcal{C}_\pi^X) = 2$*
 - (c) *If $r = 2$ ($\pi = 2g$) then $\dim(\mathcal{C}_\pi^X) = 1$.*
 - (d) *If $r = 0$ ($\pi = 2g - 1$) then $\dim(\mathcal{C}_\pi^X) = 0$.*

Proof: We apply Theorem 3.6 of [2]. We see that $\mathcal{C}_\pi^X = \emptyset$ except when X is elliptic or hyperelliptic and $r = 0, 2, 4$.

1. Suppose that $r = 2$ or $r = 4$. By Theorem 3.6 of [2], the branch divisor β verifies:
 - (a) If X is hyperelliptic and $r = 4$ then $\beta \sim 2g_2^1$.
 - (b) If X is elliptic and $r = 4$ then $\beta \sim a_1 + a_2 + a_3 + a_4$ with $a_1 + a_2 \sim a_3 + a_4$.
 - (c) If X is hyperelliptic and $r = 2$ then $\beta \sim g_2^1$.
 - (d) If X is elliptic and $r = 2$ then $\beta \sim a_1 + a_2$ for any $a_1, a_2 \in X$, $a_1 \neq a_2$.

Thus, if we consider the incidence variety:

$$J_h = \{(C, \mathcal{B}) \in \mathcal{Ch}_\pi^X \times U^r / C \text{ has an involution branched at } \mathcal{B} \in X\}$$

in this case the projection map $q_h : J_h \longrightarrow U^r$ is not a surjection. In fact we have:

- (a) If X is hyperelliptic and $r = 4$ then $\dim(\text{Im}(q_h)) = 2$.
- (b) If X is elliptic and $r = 4$ then $\dim(\text{Im}(q_h)) = 3$.
- (c) If X is hyperelliptic and $r = 2$ then $\dim(\text{Im}(q_h)) = 1$.
- (d) If X is elliptic and $r = 2$ then $\dim(\text{Im}(q_h)) = 2$.

Now, reasoning as in the proposition above we obtain:

$$\dim(\mathcal{Ch}_\pi^X) = \dim(J_h) - \dim(p^{-1}(C)) = \dim(\text{Im}(q_h)) - \dim(\text{Aut}(X)).$$

and the result follows.

- 2. Suppose that $r = 0$. By Theorem 3.6 of [2], $\mathcal{Ch}_\pi^X \neq \emptyset$ when X is elliptic or hyperelliptic and then $\dim(\mathcal{Ch}_\pi^X) = 0$. ■

Proposition 2.3 *Let \mathcal{C}_π^g be the family of smooth curves of genus $\pi > 1$ with an involution over a curve of genus $g \geq 0$. Let $r = 2(\pi - 1) - 4(g - 1)$.*

- 1. *If $r > 0$ ($\pi > 2g - 1$) then $\dim(\mathcal{C}_\pi^g) = r - \dim(\text{Aut}(X)) + \dim(\mathcal{M}_g)$.*
- 2. *If $r = 0$ ($\pi = 2g - 1$) then $\dim(\mathcal{C}_\pi^g) = \dim(\mathcal{M}_g)$.*
- 3. *If $r < 0$ ($\pi < 2g - 1$) then $\dim(\mathcal{C}_\pi^g) = \emptyset$.*

From this, $\dim(\mathcal{C}_\pi^g) = 2\pi - g - 1$. Moreover, the dimension of the family of curves with an involution over a hyperelliptic curve of genus g is $2\pi - 2g + 1$.

Proof: Since a curve of genus $\pi > 1$ has at most a finite number of involutions, such curve only has involutions over a finite number of curves of genus g . Therefore, $\dim(\mathcal{C}_\pi^g) = \dim(\mathcal{C}_\pi^X) + \dim(\mathcal{M}_g)$.

Moreover, we know that:

When $g = 0$ then $\dim(\mathcal{M}_g) = 0$ and $\dim(\text{Aut}(X)) = 3$.

When $g = 1$ then $\dim(\mathcal{M}_g) = 1$ and $\dim(\text{Aut}(X)) = 1$.

When $g > 1$ then $\dim(\mathcal{M}_g) = 3(g - 1)$ and $\dim(\text{Aut}(X)) = 0$.

If $g \geq 2$ the dimension of the family of hyperelliptic curves of genus g is $\dim(\mathcal{C}_g^0) = 2g - 1$.

We have supposed that $\pi > 1$ so $r > 0$ when $g = 1$ or $g = 0$. Now, applying Proposition 2.1 the result follows. ■

Proposition 2.4 *Let Ch_π^g be the family of smooth hyperelliptic curves of genus $\pi > 1$ with an involution over a curve of $g \geq 0$. Let $r = 2(\pi - 1) - 4(g - 1)$.*

1. *If $r = 0, 2, 4$ ($\pi = 2g - 1, 2g, 2g + 1$) then $\dim(Ch_\pi^g) = \pi$.*
2. *If $r < 0$ ($\pi < 2g - 1$) or $r > 4$ ($\pi > 2g + 1$) then $C_\pi^g = \emptyset$.*

Proof: Since a curve of genus $\pi > 1$ has at most a finite number of involutions, such curve only has involutions over a finite number of curves of genus g . Therefore, $\dim(Ch_\pi^g) = \dim(Ch_\pi^X) + \dim(\mathcal{C}_g^0)$. Moreover, we know that $\dim(\mathcal{C}_g^0) = 2g - 1$. Applying the Proposition 2.2 the result follows. ■

Remark 2.5 *Note that the dimension of Ch_π^g does not depend of g . The reason is that an involution of a hyperelliptic curve C of genus g provides an involution of genus $\pi - g$, if we compose it with the canonical involution. We will study this situation in next section.*

3 The hyperelliptic case.

Let C be an hyperelliptic curve of genus $\pi \geq 2$. C has a canonical involution defined by its unique π_2^1 . We will denote it by $\delta : C \rightarrow C$, with $\delta(P) = \pi_2^1 - P$. Note that any automorphism $\nu : C \rightarrow C$ of C verifies that $\nu(\pi_2^1) = \pi_2^1$, so it commutes with δ .

Let $\sigma : C \rightarrow C$ be an involution of genus g and $\gamma : C \rightarrow X$ the corresponding double cover. Let \mathfrak{b} the divisor of X such that $\gamma_*\mathcal{O}_C \sim \mathcal{O}_X \oplus \mathcal{O}_X(\mathcal{K} - \mathfrak{b})$.

Lemma 3.1 *Let P be a point of C . Then $\delta(P) = \sigma(P)$ if and only if $x = \gamma(\mathbf{P})$ is a base point of \mathfrak{b} .*

Proof: Let $x = \gamma(\mathbf{P})$. Then $\delta(P) = \sigma(P)$ if and only if $\gamma^*(x) \sim \pi_2^1$. But

$$\gamma^*(x) \sim \pi_2^1 \iff h^0(\mathcal{O}_C(\mathcal{K}_C - \gamma^*(x))) = h^0(\mathcal{O}_C(\mathcal{K}_C)) - 1$$

and

$$\begin{aligned} h^0(\mathcal{O}_C(\mathcal{K}_C - \gamma^*(x))) &= h^0(\mathcal{O}_{S_{\mathfrak{b}}}(X_0 + (\mathfrak{b} - x)f)) = \\ &= h^0(\mathcal{O}_X(\mathcal{K} - x)) + h^0(\mathcal{O}_X(\mathfrak{b} - x)) \\ h^0(\mathcal{O}_C(\mathcal{K}_C)) &= h^0(\mathcal{O}_{S_{\mathfrak{b}}}(X_0 + \mathfrak{b}f)) = h^0(\mathcal{O}_X(\mathcal{K})) + h^0(\mathcal{O}_X(\mathfrak{b})) \end{aligned}$$

Because \mathcal{K}_X is base-point-free the conclusion follows. ■

Theorem 3.2 *Let C an hyperelliptic curve of genus $\pi \geq 2$. Let $\sigma : C \rightarrow C$ be an involution of genus g . Then $\delta\sigma$ is an involution of C of genus $\pi - g$.*

Proof: Note that $\delta\sigma$ is an involution, because σ commutes with any automorphism of C .

Let us study the ramification points of $\delta\sigma$. We have that $\delta\sigma(P) = P \iff \sigma(P) = \delta(P)$. By Lemma 3.1, this happens when $\gamma(x)$ is a base point of \mathfrak{b} . Thus, the ramification points of $\delta\sigma$ are $\{\gamma^{-1}(x)/x \mid x \text{ is a base point of } \mathfrak{b}\}$. By Theorem 3.6 of [2] we know:

1. If $\pi = 2g + 1$, $\mathfrak{b} = \mathcal{K} + g_2^1$ and \mathfrak{b} is base-point-free.
2. If $\pi = 2g$, $\mathfrak{b} = \mathcal{K} + P$, with $2P \sim g_2^1$ and \mathfrak{b} has one base point.
3. If $\pi = 2g - 1$, $\mathfrak{b} = \sum_1^{g-2} g_2^1 + P + Q$, with $2P \sim 2Q \sim g_2^1$, and \mathfrak{b} has two base points except when X is elliptic. But we have supposed $\pi \geq 2$ so in this case $g > 1$.

From this we see that the number of ramifications of $\delta\sigma$ is 0, 2, 4 when $\pi = 2g + 1, 2g, 2g - 1$ respectively. Applying Hurwitz's formula we obtain that the genus of $\delta\sigma$ is $\pi - g$. ■

Corollary 3.3 *If $\pi \geq 2$ and $g \geq 1$ then $Ch_\pi^g = Ch_\pi^{\pi-g}$.* ■

References

- [1] ARBARELLO, E.; CORNALBA, M.; GRIFFITHS, P. A.; HARRIS, J. *Geometry of Algebraic Curves. Volume I.* Grundlehren der mathematischen Wissenschaften **267**. Springer-Verlag, 1985
- [2] FUENTES, L.; PEDREIRA, M. *Canonical geometrically ruled surfaces.* Preprint. math.AG/0107114.
- [3] HARTSHORNE, R. *Algebraic Geometry.* GTM, 52. Springer-Verlag, 1977.