

ON  $K_1$  OF A SELF-PRODUCT OF A CURVE

KENICHIRO KIMURA

Institute of Mathematics, University of Tsukuba,  
Tsukuba, 305-8571, Japan email: kimurak@math.tsukuba.ac.jp

ABSTRACT. We present elements of  $H^1(C \times C, \mathcal{K}_2)$  for certain specific curves  $C$ . The image of the element under the boundary map arising from the localization sequence of  $K$ -theory is the graph of frobenius endomorphism of the reduction of the curve modulo 3.

**§1 Introduction.** Let  $X$  be a projective smooth variety over  $\mathbb{Q}$ . Let  $S$  be a finite set of primes such that  $X$  has a projective smooth model  $\mathcal{X}$  over  $U := \text{spec} \mathbb{Z}[\frac{1}{S}]$ . Then from the works of Bloch([B1]) and Sherman, we have the following exact localization sequence in algebraic  $K$ -theory, using Gersten's conjecture for  $\mathcal{K}_2$  in a mixed characteristic setting that was proven by Bloch([B2])

$$H^1(\mathcal{X}, \mathcal{K}_2) \rightarrow H^1(X, \mathcal{K}_2) \xrightarrow{\partial} \bigoplus_{p \in U} \text{Pic}(X_p) \rightarrow CH^2(\mathcal{X}) \rightarrow CH^2(X) \rightarrow 0$$

where  $X_p$  is the fiber of  $\mathcal{X}$  at the prime  $p$ .

A conjecture of Beilinson on the special values of  $L$ -function and Tate conjecture tell us that the cokernel of  $\partial$  is torsion. See the remark after Theorem 2.5 in [La] for an account of this. This means there should be enough elements in  $H^1(X, \mathcal{K}_2)$  so that any element of  $\text{Pic}(X_p) \otimes \mathbb{Q}$  is in the image of the map  $\partial \otimes \mathbb{Q}$ . The purpose of this paper is to give a potential method to construct elements in  $H^1(X, \mathcal{K}_2)$  in the case where  $X$  is the product of a curve.

In this case a typical element of  $\text{Pic}(X_p)$  which usually does not lift to the generic fiber is the graph of frobenius endomorphism of the fiber of the curve. For a certain curve  $C$  of special type we construct a divisor  $D \subset C \times C$  such that the reduction  $D_p$  of  $D$  modulo  $p$  contains the graph of frobenius endomorphism  $\Gamma_\varphi$  as an irreducible component (Lemma 2.1).

Next task is to find a nice function  $g$  on  $D$  such that  $g$  vanishes on  $\Gamma_\varphi$  and not on other components of  $D_p$ , and that one can modify  $(g, D)$  to an element

---

<sup>1</sup>The author was partially supported by JSPS Postdoctoral Fellowships for Research Abroad for 2000-2001.

$\Sigma \in H^1(X, \mathcal{K}_2)$  by adding some elements. In this we have succeeded for the curve  $C$  and for the prime 3. This is the main result of this paper (Theorem 2.3).

Although in this paper we are able to construct elements only for curves of very specific type, the idea of construction can be applied to more general curves, and we hope it gives a clue to construct more elements in more general situation.

When  $X$  is the product of a modular curve there is a beautiful construction of elements in  $H^1(X, \mathcal{K}_2)$  by Flach([Fl]) and Mildenhall([Mi]). However their elements are still not enough to show that  $\text{coker}(\partial)$  is torsion for the products of general modular curves.

The author thanks S. Bloch and R. Sreekantan for discussions. This work was carried out while the author stayed at Department of Mathematics, University of Chicago. He is thankful to the people at Chicago for their hospitality.

## §2 Construction of an element in $H^1(X, \mathcal{K}_2)$ .

Let  $C$  be a curve over  $\mathbb{Q}$  defined as the normalization of the projective closure of the affine curve

$$y^m = t^n + 1.$$

Here  $m$  and  $n$  are integers such that  $n \geq 5$ ,  $(n, 6) = 1$ ,  $(m, 3) = 1$  and  $(m, n) = 1$ . We denote its point at infinity by  $\infty$ . For a prime  $p \nmid mn$ , the curve  $C$  naturally extends to a projective smooth model  $\mathcal{C}$  over  $\text{spec } \mathbb{Z}_{(p)}$ . We denote the special fiber of  $\mathcal{C}$  by  $C_p$ . Let

$$\pi : \mathcal{C} \rightarrow \mathbb{P}_{\mathbb{Z}_{(p)}}^1$$

be a map given by  $(y, t) \mapsto (t)$ . Let  $F : \mathbb{P}_{\mathbb{Z}_{(p)}}^1 \rightarrow \mathbb{P}_{\mathbb{Z}_{(p)}}^1$  be a morphism given by  $t \mapsto t^p$ .

Let  $f := F \circ \pi$  and  $\Gamma_f \subset \mathcal{C} \times_{\mathbb{Z}_{(p)}} \mathbb{P}^1$  be its graph. Let  $\mathcal{D} = (id \times \pi)^* \Gamma_f \subset \mathcal{C} \times \mathcal{C}$ . We denote the generic fiber of  $\mathcal{D}$  by  $D$  and the special fiber of  $\mathcal{D}$  by  $D_p$ .

**Lemma 2.1.** *The intersection*

$$\mathcal{D} \cap (\mathcal{C} \times \infty \cup \infty \times \mathcal{C})$$

*is supported on  $\infty \times \infty$ .*

*proof.* If  $x \times \infty \in D_p$  (resp.  $x \times \infty \in D$ ) for a closed point  $x$  of  $C_p$  (resp. of  $C$ ) then  $x \times \pi(\infty) = x \times \infty \in \Gamma_f$  so  $F \circ \pi(x) = \infty$ , hence  $\pi(x) = \infty$ ,  $x = \infty$ .

Similarly, if  $\infty \times x \in D_p$  (resp.  $\in D$ )  $\Rightarrow \infty \times \pi(x) \in \Gamma_f$ ,  $\pi(x) = f(\infty) = \infty$ ,  $x = \infty$ .  $\square$

**Lemma 2.2.** *Let  $\kappa$  be the field generated by a primitive  $m$ -th root of unity  $\xi$  over  $\mathbb{F}_p$ . Then the divisor*

$$D_p \otimes_{\mathbb{F}_p} \kappa = \sum_{j=0}^{m-1} \Gamma_{\xi^j \circ \varphi}.$$

2

Here  $\varphi$  is the Frobenius endomorphism of  $C_p$  and  $\xi$  is the automorphism of  $C$  given by  $y \mapsto \xi y$ .

*proof.* Let  $\mathcal{V} := (C \setminus \{\infty\}) \times_{\mathbb{Z}_{(p)}} (C \setminus \{\infty\})$ . We denote the generic fiber of  $\mathcal{V}$  by  $V$  and its special fiber by  $V_p$ .

By lemma 2.1 it suffices to look at  $D_p \cap V_p$ . On  $V_p$ ,  $D_p$  is defined by the equation  $t_2 - t_1^p = 0$ . Here  $(y_1, t_1) \times (y_2, t_2)$  is a set of coordinates on  $V_p$ . One can see that  $y_2^m - (y_1^p)^m \in (t_2 - t_1^p)$ .  $\square$

*Remark.* This construction of the divisor  $D$  can be applied in general to curves which are cyclic cover of  $\mathbb{P}^1$ .

In the following we specify  $p$  to 3. When  $m = 2$  and  $n = 5$  the curve  $C_3$  over  $\mathbb{F}_3$  is isomorphic to the curve  $t^5 = y(y - 1)$ . In this case the characteristic polynomial of the action of  $\varphi$  on the Tate module of Jacobian of  $C_3$  is  $T^4 + 9$  ([Gr-Ro] Lemma 1.1). Since  $\text{End}(\text{Jac}(C)) = \mathbb{Z}$ ,  $\Gamma_\varphi$  does not lift to the generic fiber.

**Theorem 2.3.** *There is an element  $\Sigma \in H^1(C \times C, \mathcal{K}_2) \otimes \mathbb{Q}$  such that the factor of*

$$\partial(\Sigma) \text{ at } p = \begin{cases} \Gamma_\varphi & p = 3 \\ 0 & p \in \text{spec} \mathbb{Z}[\frac{1}{mn}], p \neq 3. \end{cases}$$

*proof.* Consider the function  $g := y_2 - y_1^3$  on  $D$ . We calculate  $\text{div}(g) \cap V$ .

$$y_2 - y_1^3 = 0 \Rightarrow t_2^n + 1 - (t_1^n + 1)^3 = 0.$$

We write  $T = t_1^n$ . Then it follows  $T^3 + 1 - (T + 1)^3 = 0$ ,  $-3(T^2 + T) = 0$ ,  $T = 0$  or  $-1$ .

So  $t_1 = 0, -\zeta^\alpha$ ,  $\alpha = 0, \dots, n - 1$ .

Here  $\zeta$  is a primitive  $n$ -th root of unity.

Hence  $\text{div}(g)$  is supported on the set  $\{(\xi^j, 0) \times (\xi^{3j}, 0) | j = 0, \dots, m - 1\} \cup \{(0, -\zeta^\alpha) \times (0, -\zeta^{3\alpha}) | \alpha = 0, \dots, n - 1\} \cup \{\infty\}$ .

Here  $\xi$  is a primitive  $m$ -th root of unity.

Let  $\mathcal{S}$  be the set of points on  $C$  given by

$$\mathcal{S} = \{(y, t) \in C \setminus \{\infty\} | y = 0 \text{ or } t = 0\} \cup \{\infty\}.$$

We see that the support of  $\text{div}(g) \subset \mathcal{S} \times \mathcal{S}$ .

For any point  $x \in \mathcal{S}$  the divisor  $(x) - (\infty) \in \text{Pic}^0 C$  is torsion, so we can construct an element  $\Sigma \in H^1(C \times C, \mathcal{K}_2) \otimes \mathbb{Q}$  of the form

$$(g, D) + \sum_{x_i \in \mathcal{S}} (f_i, C \times x_i) + (f'_i, x_i \times C).$$

Since  $g = 0$  at the generic point of  $\Gamma_\varphi$  and  $g \neq 0$  at the generic points of other irreducible components of  $D_3$ , we see that the factor of  $\partial(\Sigma)$  at  $p = 3$  is equal to  $\Gamma_\varphi$ .

**Lemma 2.4.** *For any prime  $l \in \text{spec}\mathbb{Z}[\frac{1}{mn}]$  other than 3, the reduction  $D_l$  of the divisor  $\mathcal{D}$  mod  $l$  is irreducible.*

*proof.* One can show that

$$D_l \cap (\infty \times C_l \cup C_l \times \infty)$$

is supported on  $\infty \times \infty$  in the same way as the proof of Lemma 2.1. So it suffices to show that  $D_l \cap (C_l \setminus \{\infty\}) \times (C_l \setminus \{\infty\})$  is irreducible. On  $(C_l \setminus \{\infty\}) \times (C_l \setminus \{\infty\})$ ,  $D_l$  is the scheme

$$\text{spec } \mathbb{F}_l[y_1, t_1, y_2] / (y_1^m - t_1^n - 1, y_2^m - (y_1^m - 1)^3 - 1).$$

So  $D_l$  is reducible if and only if  $y_2^m - (y_1^m - 1)^3 - 1$  is reducible. There is a point  $s \in C_l(\overline{\mathbb{F}}_l)$  at which the function  $(y_1^m - 1)^3 + 1$  vanishes to order 1. It follows that  $y_2^m - (y_1^m - 1)^3 - 1$  is irreducible from the following well known fact:

*Let  $\mathcal{O}$  be a discrete valuation ring and denote its maximal ideal by  $\mathfrak{m}$ . Let  $f(x) = x^k + a_{k-1}x^{k-1} + \cdots + a_0 \in \mathcal{O}[x]$  be a polynomial. If  $a_j \in \mathfrak{m}$  for  $j = 0, \dots, k-1$  and  $a_0$  generates  $\mathfrak{m}$ , then  $f(x)$  is irreducible.  $\square$*

The above argument and Lemma 2.4 show the existence of  $\Sigma$  as stated in the theorem.  $\square$

## REFERENCES

- [Bl1] Bloch, S., *Lectures on Algebraic Cycles*, Duke University Press, 1980.
- [Bl2] Bloch, S., "A note on Gersten's conjecture in the mixed characteristic case" in *Applications of Algebraic K-theory to Algebraic Geometry and Number Theory, Part I, II*, Contemp. Math. **55** (1986), Amer. Math. Soc. Providence.
- [Fl] Flach, M., *A finiteness theorem for the symmetric square of an elliptic curve*, Invent. Math **109** (1992), 307-327.
- [Gr-Ro] Gross, B., Rohrlich, D., *Some results on the Mordell-Weil group of the Jacobian of the Fermat curve*, Invent. Math **44** (1978), 201-224.
- [La] Langer, A., *Zero cycles on Hilbert-Blumenthal surfaces*, Duke Math. J **103** (2000), 131-163.
- [Mi] Mildenhall, S., *Cycles in a product of elliptic curves, and a group analogous to the class group*, Duke Math. J **67** (1992), 387-406.