Stable systolic inequalities and cohomology products

Victor Bangert

Mathematisches Institut Universität Freiburg, Eckerstr. 1,79104 Freiburg, Germany

AND

Mikhail Katz*

Department of Mathematics and Statistics Bar Ilan University, Ramat Gan, 52900, Israel

Dedicated to the memory of Jürgen Moser

Abstract

Multiplicative relations in the cohomology ring of a manifold impose constraints upon its stable systoles. Given a compact Riemannian manifold (X, g), its real homology $H_*(X, \mathbb{R})$ is naturally endowed with the stable norm. Briefly, if $h \in H_k(X, \mathbb{R})$ then the stable norm of h is the infimum of the Riemannian kvolumes of real cycles representing h. The stable k-systole is the minimum of the stable norm over nonzero elements in the lattice of integral classes in $H_k(X, \mathbb{R})$. Relying on results from the geometry of numbers due to W. Banaszczyk, and extending work by M. Gromov and J. Hebda, we prove metric-independent inequalities for products of stable systoles, where the product can be as long as the real cup length of X.

1 Introduction

Given a compact Riemannian manifold (X, g), its real homology $H_*(X, \mathbb{R})$ is naturally endowed with the stable norm. Briefly, if $h \in H_k(X, \mathbb{R})$ then the stable norm of h is the infimum of the k-volumes, defined in terms of the metric g, of real cycles representing h. The stable k-systole is the minimum of the stable norm

^{*}Supported by The Israel Science Foundation (grant no. 620/00-10.0). Partially supported by the Emmy Noether Research Institute and the Minerva Foundation of Germany.

over nonzero elements in the lattice of integral classes in $H_k(X, \mathbb{R})$, cf. (1.4) below.

Our goal is to explore the hidden power of the seminal calculation contained in lines 7-12, p. 262 of M. Gromov's book [23], cf. (4.4) below. We rely on results of W. Banaszczyk [7, 8] from the geometry of numbers, which provide a bound for certain products of the successive minima of a pair of dual lattices, see (4.1) and (7.1). We thus extend the work of M. Gromov [21] and J. Hebda [24] to prove metric-independent inequalities for products of stable systoles, where the product can be as long as the real cup length of X. In particular, for the pair of stable systoles of dimension and codimension 1, we prove an optimal inequality, where the boundary case of equality is attained, e.g. by flat tori defined as quotients of the so-called dual-critical lattices of A.-M. Bergé and J. Martinet [11].

Geometric Measure Theory provides a framework in which the existence of minimal representatives in homology classes can be proved. If $h \in H_k(X, \mathbb{Z})$ then $\operatorname{vol}_k(h)$ (cf. Definition 1.1 below) is the minimal mass of a closed, integer-multiplicity, rectifiable k-current representing h, cf. [17, 5.1.6]. If $h \in H_k(X, \mathbb{R})$ then the stable norm (or mass norm) of h (cf. Definition 1.2 below) is the minimal mass of a closed (normal) k-current representing h.

In the cases k = 1 and k = n - 1 the transition from the Riemannian length resp. (n-1)-volume functional to the corresponding stable norm is closely related to the process of *homogenization* studied in Analysis. For k = n - 1 this shows up in J. Moser's work [32, Sect. 7], [33], see [35, 36] for more details. It is through these remarks by J. Moser that the first-named author originally came in contact with the subject of this paper.

While the present paper studies systolic ramifications of multiplicative relations in the cohomology ring, the work [27] explores the systolic influence of Massey products.

Let X be a compact manifold of dimension n. We will now provide detailed definitions of the systolic invariants involved. A choice of a Riemannian metric g on X allows us to define the total volume $\operatorname{vol}_n(g)$, as well as the k-volumes of k-dimensional submanifolds of X. More generally, given an integer Lipschitz chain $c = \sum_i n_i \sigma_i$, one defines its volume

$$\operatorname{vol}_k(c) = \sum_i |n_i| \operatorname{vol}_k(\sigma_i).$$

Here the volume $\operatorname{vol}_k(\sigma)$ of a Lipschitz k-simplex $\sigma : \Delta^k \to X$ is the integral over the k-simplex Δ^k of the "volume form" of the pullback $\sigma^*(g)$.

DEFINITION 1.1 The (minimal) volume,

 $\operatorname{vol}_k(h),$

of an integer homology class $h \in H_k(M, \mathbb{Z})$ is the infimum of $\operatorname{vol}_k(c)$ over all integer Lipschitz cycles representing h.

Let k be an integer satisfying $1 \le k \le n$. We define the k-systole $\operatorname{sys}_k(g)$ of (X, g) as the minimum of the volumes of nonzero integer k-homology classes:

$$\operatorname{sys}_k(g) = \min\{\operatorname{vol}_k(h) | h \in H_k(X, \mathbb{Z}) \setminus \{0\}\}.$$

In particular, $\operatorname{sys}_n(g) = \operatorname{vol}_n(g)$ and, formally, $\operatorname{sys}_k(g) = \infty$ if $H_k(X, \mathbb{Z}) = 0$. A good introduction to systoles is M. Berger's survey [13], for further references see [28], sections 2 and 3.

Replacing integer cycles by real cycles we define the stable norm as follows.

DEFINITION 1.2 The stable norm ||h|| of $h \in H_k(X, \mathbb{R})$ is the infimum of the volumes $\operatorname{vol}_k(c) = \sum_i |r_i| \operatorname{vol}_k(\sigma_i)$ over all real Lipschitz cycles $c = \sum_i r_i \sigma_i$ representing h.

Note that $\parallel \parallel$ is indeed a norm, cf. [18] and [23], 4.C.

We denote by $H_k(X, \mathbb{Z})_{\mathbb{R}}$ the image of $H_k(X, \mathbb{Z})$ in $H_k(X, \mathbb{R})$ and by $h_{\mathbb{R}}$ the image of $h \in H_k(X, \mathbb{Z})$ in $H_k(X, \mathbb{R})$. Recall that $H_k(X, \mathbb{Z})_{\mathbb{R}}$ is a lattice in $H_k(X, \mathbb{R})$. Obviously

$$\|h_{\mathbb{R}}\| \le \operatorname{vol}_k(h)$$

for all $h \in H_k(X, \mathbb{Z})$. Moreover, $||h_{\mathbb{R}}|| = \operatorname{vol}_n(h)$ if $h \in H_n(X, \mathbb{Z})$. H. Federer [18] investigated the relations between $||h_{\mathbb{R}}||$ and $\operatorname{vol}_k(h)$ and proved:

(1.2) If
$$h \in H_k(X, \mathbb{Z}), 1 \le k < n$$
, then

$$\|h_{\mathbb{R}}\| = \lim_{i \to \infty} \frac{1}{i} \operatorname{vol}_k(ih).$$

(1.3) If X is orientable and $h \in H_{n-1}(X, \mathbb{Z})$, then

$$\|h_{\mathbb{R}}\| = \operatorname{vol}_{n-1}(h)$$

(see also [38]). We define the stable k-systole $\operatorname{stsys}_k(g)$ of (X, g) by

(1.4)
$$\operatorname{stsys}_k(g) = \min\left\{ \|h\| | h \in H_k(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\} \right\}.$$

If the k'th Betti number $b_k(X) = \dim H_k(X, \mathbb{R})$ of X is positive and $H_k(X, \mathbb{Z})$ is free abelian, then (1.1) implies

$$\operatorname{stsys}_k(g) \leq \operatorname{sys}_k(g).$$

If X is orientable then

$$\operatorname{stsys}_n(g) = \operatorname{sys}_n(g) = \operatorname{vol}_n(g)$$

and (1.3) implies

$$\operatorname{stsys}_{n-1}(g) = \operatorname{sys}_{n-1}(g).$$

2 Statement of main results

Assuming that the fundamental class of a compact, oriented manifold X can be written as a cup product of classes of dimensions k_1, \ldots, k_m , M. Gromov proved an upper bound for the product of the stable systoles $\Pi_j \operatorname{stsys}_{k_j}(g)$ in terms of the volume of X, see [21, 7.4.C] and also [23, 4.38]. We extend this result, provide a simpler proof and analyze the dependence of the constant on the Betti numbers involved.

THEOREM 2.1 Let X be a compact manifold and let $k \ge 1$ be an integer such that $H^k(X, \mathbb{R})$ is not zero and spanned by cup products of classes of dimensions $k_1, \ldots, k_m, \Sigma_j k_j = k$. Then, for every Riemannian metric g on X, we have

$$\prod_{j=1}^{m} \operatorname{stsys}_{k_{j}}(g) \leq C(k) \left(\prod_{j} b_{k_{j}}(X)(1 + \log b_{k_{j}}(X)) \right) \operatorname{stsys}_{k}(g)$$

for a constant C(k) only depending on k.

Using Poincaré duality, J. Hebda [24, Proposition 6] bounds the product of the stable systoles in complementary dimensions by the volume. We generalize his result as follows:

THEOREM 2.2 Let X be a compact manifold and let p, q be integers, $p+q \leq \dim X$. Suppose there exists $h \in H_{p+q}(X,\mathbb{Z})_{\mathbb{R}}$ such that the cap product with h induces an injective map

$$\alpha \in H^p(X,\mathbb{Z})_{\mathbb{R}} \to h \cap \alpha \in H_q(X,\mathbb{Z})_{\mathbb{R}}$$

Then, for every Riemannian metric g on X, we have

 $\operatorname{stsys}_p(g) \operatorname{stsys}_q(g) \le C(p,q)b_p(X) \|h\|$

for a constant C(p,q) only depending on p and q. If our assumption is satisfied for all $h \in H_{p+q}(X,\mathbb{Z})_{\mathbb{R}} \setminus \{0\}$ then

(2.1)
$$\operatorname{stsys}_p(g)\operatorname{stsys}_q(g) \le C(p,q)b_p(X)\operatorname{stsys}_{p+q}(g).$$

Examples of flat tori show that an inequality of type (2.1) has to depend on the Betti number b_p linearly. We do not know if in Theorem 2.1 the dependence on the Betti numbers is optimal. Explicit values for the constants C(p,q) can be computed. Unless $\{p,q\} = \{1, \dim X - 1\}$ the sharp constants C(p,q) are unknown to us. In the case $\{p,q\} = \{1, \dim X - 1\}$ we have the following sharp result (Corollary 2.3 below) that generalizes [24, Theorem A] to the case $b_1(X) > 1$. Given a lattice L in euclidean space, we set

 $\lambda_1(L) = \min\{|v| \mid v \in L \setminus \{0\}\}.$

Let $b \in \mathbb{N}$, and consider the Bergé-Martinet constant γ'_b , see [11],

 $\gamma'_b = \sup\{\lambda_1(L)\lambda_1(L^*) \mid L \text{ a lattice in euclidean space } \mathbb{R}^b\}$

where L^* is the lattice dual to L. Thus, the constant γ'_b is bounded above by the Hermite constant γ_b , cf. e.g. [30], p. 334, and satisfies $\gamma'_1 = 1$ and the inequalities

(2.2)
$$\gamma'_b \le \gamma_b \le \frac{2}{3} b \text{ for all } b \ge 2$$

and

(2.3)
$$\frac{b}{2\pi e}(1+o(1)) \le \gamma'_b \le \frac{b}{\pi e}(1+o(1)) \text{ for } b \to \infty,$$

cf. e.g. [30], pp. 334 and 337.

COROLLARY 2.3 Let X be a compact, orientable manifold, dim X = n, with positive first Betti number $b = b_1(X)$. Then, for every Riemannian metric g on X, we have

$$\operatorname{stsys}_1(g)\operatorname{sys}_{n-1}(g) \le \gamma'_b \operatorname{vol}_n(g).$$

Equality is attained for a flat torus \mathbb{R}^n/L where $L \subseteq \mathbb{R}^n$ is a lattice with $\lambda_1(L)\lambda_1(L^*) = \gamma'_n$.

These inequalities can be seen as analogues of the optimal inequalities of C. Loewner, P. Pu [34], and C. Bavard [10], and as generalizations of the results of R. Accola [1] and C. Blatter [14] (cf. [13], p. 290) in dimension 2. A relative version of Corollary 2.3 is studied in [4].

Note that we have examples X with equality in Corollary 2.3 only if dim $X = n \ge b = \dim H_1(X, \mathbb{R})$. If we fix $n = \dim X$ there might exist a better estimate, maybe even one independent of b. In particular we ask, see also [21], 7.4.C:

QUESTION 2.4 Does there exist a constant C such that

$$\operatorname{stsys}_1(g) \operatorname{sys}_2(g) \le C \operatorname{vol}_3(g)$$

for all 3-dimensional, compact, orientable, Riemannian manifolds (X, g)?

The manifolds (X, g) for which equality holds in Corollary 2.3 will be investigated in [9]. In the case $b_1(X) = 1$, J. Hebda [24, Theorem A] proved that one has equality $\operatorname{stsys}_1(g) \operatorname{sys}_{n-1}(g) = \operatorname{vol}_n(g)$ if and only if X admits a Riemannian submersion $F: X \to S^1$ onto a circle, such that all fibers $F^{-1}(s), s \in S^1$, are minimal hypersurfaces in X.

Consider a *b*-dimensional normed real vector space (V, || ||) and a lattice $L \subseteq V$, in particular $\operatorname{rk}(L) = b$. For $1 \leq i \leq b$ we define the *i*-th successive minimum $\lambda_i(L, || ||)$ as the minimal $\lambda > 0$ for which there exist *i* linearly independent vectors in *L* of norm smaller than or equal to λ .

Now let n = 2p, and consider the L^2 -norm $| |_{L^2}$ in homology $H_p(X, \mathbb{R})$ dual to the one on harmonic forms defined by

$$|f|_{L^2}^2 = \int_X f \wedge *f,$$

7

where * is the Hodge star operator of the metric g. Consider also the conformally invariant norm $\| \|_{L^2}^*$ in $H^p_{dR}(X)$ defined by taking the infimum, over all representatives ω , of the quantity

$$\|\omega\|_{L^2} = \left(\int_X \|\omega_x\|^2 d\operatorname{vol}_n(x)\right)^{\frac{1}{2}},$$

where $\|\omega_x\|$ is the pointwise comass. Let $\|\|_{L^2}$ be the dual norm in homology, cf. (7.4) below. We introduce the conformal invariant

$$\operatorname{conf}_p(g) = \min\{\|h\|_{L^2} \mid h \in H_p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}\}\$$

COROLLARY 2.5 Let $p = \frac{n}{2}$ and $b_p = b_p(X)$. Then

$$\operatorname{conf}_p(g)^2 \le \frac{\lambda_1}{\lambda_{b_p}} \binom{n}{p} b_p$$

where $\lambda_i = \lambda_i (H_p(X, \mathbb{Z})_{\mathbb{R}}, | |_{L^2}).$

Note that [26] studies *lower* bounds for the quantity $\sup_g \operatorname{conf}_2(g)$ for 4-manifolds (X, g), where X is the blowup of the complex projective plane, the bound being *polynomial* of order $n^{\frac{1}{4}}$ in the number n of blown-up points. Whether or not such lower bounds exist for the stable 2-systole is unknown.

QUESTION 2.6 Does there exist a constant C such that

$$\operatorname{stsys}_2(g) \le C \sqrt{\operatorname{vol}_4(g)}$$

for all 4-dimensional, compact, orientable, Riemannian manifolds (X, g)?

Note that this is so in the case of a definite intersection form, cf. [26], (3.5).

1

For the first stable systole M. Gromov proved

(2.4)
$$\operatorname{stsys}_1(g) \le \sqrt{\gamma_n} \operatorname{vol}_n(g)^{\frac{1}{n}}$$

if g is a Riemannian metric on a compact Riemannian manifold X such that $b_1(X) = n = \dim X$, for which there are classes $\alpha_1, \ldots, \alpha_n \in$ $H^1(X, \mathbb{R})$ with $\alpha_1 \cup \ldots \cup \alpha_n \neq 0$. Here γ_n is the Hermite constant, cf. (2.2) above. See [22, 1.A₁], [23], pp. 259-260, [13], p. 283 and additionally [2], Theorem 8.8. Combining the methods used to prove Theorem 2.1 and Theorem 2.2, we obtain a result similar to (2.4) for higher-dimensional stable systoles:

THEOREM 2.7 Let X be a compact, orientable manifold, of dimension n = kp. Suppose there are classes $\beta_1, \ldots, \beta_k \in H^p(X, \mathbb{R})$ such that $\beta_1 \cup \ldots \cup \beta_k \neq 0$. Then, for every Riemannian metric g on X, we have

$$\operatorname{stsys}_p(g) \le C(n) \left(b_p(X)(1 + \log b_p(X)) \right)^{\frac{k-1}{k}} \operatorname{vol}_n(g)^{\frac{1}{k}}$$

for a constant C(n) only depending on n.

It is a natural question whether similar inequalities exist if one replaces the stable systoles by systoles. The most prominent example is Gromov's [22, $3.C_1$] systolic inequality

(2.5)
$$\operatorname{sys}_1(g) \le c_n(\operatorname{vol}_n(g))^{\frac{1}{n}}$$

which holds for all Riemannian metrics on compact *n*-dimensional manifolds X of cuplength n, i.e. if for some field F there are classes $\alpha_1, \ldots, \alpha_n \in H^1(X, F)$ such that $\alpha_1 \cup \ldots \cup \alpha_n \neq 0$. See (7.6) below for a refined inequality of this type for surfaces.

For k-systoles with $k \geq 2$, however, systolic freedom tends to prevail, i.e. there are many examples where inequalities as in Theorem 2.1 and Theorem 2.2 are not true if one replaces stable systoles by systoles. Based on examples by M. Gromov [22, 4.A₅], see also section 6 of M. Berger's survey article [13], this phenomenon has been studied by I. Babenko, A. Suciu and the second author, cf. [5], [6], [23, Appendix D], [3], [28, 29] and [25]. Finally there is the striking example by M. Freedman [19] of a sequence of Riemannian metrics g_j on $S^1 \times S^2$ such that $\lim_j \operatorname{vol}_3(g_j) = 0$ while $\operatorname{length}(\gamma) \operatorname{area}(\Sigma) \geq 1$ (for all j) whenever γ is a noncontractible loop in X and $\Sigma \subset X$ is a closed surface which is not nullhomologous with coefficients in $\mathbb{Z}/2\mathbb{Z}$.

Some topological preliminaries are presented in section 3. The inequality of W. Banaszczyk, a crucial ingredient in our technique, appears in section 4, together with the proof of Theorem 2.1. Theorem 2.2 is proved in section 5, Corollary 2.3 in section 6, Corollary 2.5 in section 7, and Theorem 2.7 in section 8.

3 Some topological preliminaries

In this section we collect some facts from homology theory that will be used in the sequel.

For a compact manifold X we consider the homology modules derived from the singular (co-)chain complexes with Z- or R-coefficients. We assume that the singular simplices are Lipschitz. Since we will often use results that are actually proved for the (continuous) singular or the C^{∞} -singular (co-)homology theories we note that these are naturally isomorphic to the ones based on Lipschitz simplices. The bilinear pairing (Kronecker product) between homology and cohomology will be denoted by [,], cup products by \cup and cap products by \cap . By de Rham's Theorem, see e.g. [37, 5.45], the singular cohomology algebra $H^*(X, \mathbb{R})$ is naturally isomorphic to the de Rham cohomology algebra $H^*_{dR}(X)$. In de Rham theory the cup product is defined by the wedge product of closed forms representing the cohomology classes. The (nondegenerate) bilinear pairing with real homology is given by integrating representing forms over representing Lipschitz cycles, cf. [37, 4.17].

REMARK 3.1 The maps induced from the universal coefficient theorem are compatible with Kronecker pairing, cup and cap products.

More specifically, the map $\iota_* : H_*(X, \mathbb{Z}) \to H_*(X, \mathbb{R}), h \to h_{\mathbb{R}}$ is defined by inclusion on the level of chains. The map $\iota^* : H^*(X, \mathbb{Z}) \to$ $H^*(X, \mathbb{R}) \simeq H^*_{dR}(X), \alpha \to \alpha_{\mathbb{R}}$ corresponds to the natural extension of a cochain $c \in \text{Hom}(C(X, \mathbb{Z}), \mathbb{Z})$ to a cochain in $\text{Hom}(C(X, \mathbb{R}), \mathbb{R})$. According to the universal coefficient theorem in homology, cf. [20, 29.12], the image $H_*(X, \mathbb{Z})_{\mathbb{R}} := \text{Im}(\iota_*)$ is a lattice in $H_*(X, \mathbb{R})$. Let $H^*(X, \mathbb{Z})_{\mathbb{R}} := \text{Im}(\iota^*)$.

LEMMA 3.2 The lattice $H^*(X, \mathbb{Z})_{\mathbb{R}}$ is dual to $H_*(X, \mathbb{Z})_{\mathbb{R}}$ under [,].

PROOF: By naturality mentioned in Remark 3.1 above, we have $[h_{\mathbb{R}}, \alpha_{\mathbb{R}}] = [h, \alpha] \in \mathbb{Z}$ for all $h \in H_*(X, \mathbb{Z}), \alpha \in H^*(X, \mathbb{Z})$. Conversely, suppose $\tilde{\alpha} \in H^*(X, \mathbb{R})$ and $[h_{\mathbb{R}}, \tilde{\alpha}] \in \mathbb{Z}$ for all $h \in H_*(X, \mathbb{Z})$. By the surjectivity result [20, 23.9] for a PID coefficient ring, there exists $\alpha \in H^*(X, \mathbb{Z})$ such that $[h, \alpha] = [h_{\mathbb{R}}, \tilde{\alpha}]$ for all $h \in H_*(X, \mathbb{Z})$. The nondegeneracy of [,] now implies that $\tilde{\alpha} = \alpha_{\mathbb{R}} \in H^*(X, \mathbb{Z})_{\mathbb{R}}$.

Thus the set of de Rham classes $\alpha \in H^*_{dR}(X)$, that are represented by closed forms whose integrals over integer cycles are integers, coincides with $\operatorname{Im}(\iota^*)$. These classes will henceforth be called *integer* classes.

If X is a compact, oriented manifold, dim X = n, then the Poincaré duality isomorphism $H^p(X, \mathbb{R}) \to H_{n-p}(X, \mathbb{R})$ is defined by

$$\operatorname{PD}_{\mathbb{R}} : \alpha \in H^p(X, \mathbb{R}) \to \xi_{\mathbb{R}} \cap \alpha \in H_{n-p}(X, \mathbb{R})$$

where ξ denotes the fundamental class of X. Since the corresponding map PD : $\alpha \in H^p(X, \mathbb{Z}) \to \xi \cap \alpha \in H_{n-p}(X, \mathbb{Z})$ is an isomorphism, cf. [20, 26.6], we see that PD_R induces an isomorphism between $H^p(X, \mathbb{Z})_{\mathbb{R}}$ and $H_{n-p}(X, \mathbb{Z})_{\mathbb{R}}$. In particular, Lemma 3.2 and the preceding argument show that a de Rham class $\alpha \in H^p_{dR}(X)$ is an integer class if $[PD_{\mathbb{R}}(\alpha), \beta] \in \mathbb{Z}$ for all integer classes $\beta \in H^{n-p}_{dR}(X)$. This fact was introduced as a hypothesis in [12], p. 253, and named a "dual lattice condition" in [24], p. 344. The preceding discussion shows that this condition is always satisfied.

4 Banaszczyk's inequality and proof of Theorem 2.1

The following result by W. Banaszczyk [8] is a crucial ingredient in the proof of Theorem 2.1.

Consider a *b*-dimensional normed real vector space (V, || ||) and a lattice $L \subseteq V$, in particular $\operatorname{rk}(L) = b$. For $1 \leq i \leq b$ we let $\lambda_i(L, || ||)$ denote the minimal $\lambda > 0$ for which there exist *i* linearly independent vectors in *L* of norm smaller than or equal to λ . Let L^* denote the lattice dual to *L* in the dual space V^* and let $|| ||^*$ denote the norm dual to || ||. Corollary 2 in [8] implies:

There exists a constant C > 0 such that

(4.1)
$$\lambda_i(L, \| \|) \lambda_{b-i+1}(L^*, \| \|^*) \le C b(1 + \log b)$$

whenever $b \in \mathbb{N}$, $i \in \{1, \ldots, b\}$ and L is a lattice in a b-dimensional normed space $(V, \| \|)$.

In our application of this result V will be $H_p(X, \mathbb{R})$, L will be $H_p(X, \mathbb{Z})_{\mathbb{R}}$ and $\| \|$ will be the stable norm on $H_p(X, \mathbb{R})$. So V^*

11

can be identified with the de Rham cohomology $H^p_{dR}(X)$, and then Lemma 3.2 implies that $L^* = H^p(X, \mathbb{Z})_{\mathbb{R}}$ is the lattice of integer de Rham classes. The norm on $H^p_{dR}(X)$ dual to the stable norm on $H_p(X, \mathbb{R})$ is the comass norm, cf. [18, 4.10] and [23, 4.35]. Here the comass of a *p*-form ω on a compact Riemannian manifold X is

$$\|\omega\|_{\infty} = \max\{\omega_x(e_1,\ldots,e_p) \mid x \in M, e_i \in \mathrm{TM}_x\},\$$

where $|e_i| = 1$ for $1 \le i \le p$, while the comass of $\alpha \in H^p_{dR}(X)$ is

$$\|\alpha\|^* = \inf\{\|\omega\|_{\infty} \mid \omega \text{ a closed } p\text{-form representing } \alpha\}$$

EXAMPLE 4.1 For a flat torus (T^n, g) the calculation of the stable norm and the comass norm reduces to the calculation of the corresponding pointwise quantities, cf. the work [31] by H.B. Lawson, in particular Corollary 4.4. If we normalize g so that $\operatorname{vol}_n(g) = 1$, and represent (T^n, g) as the quotient of euclidean space \mathbb{R}^n by a lattice L, then the standard isomorphisms

$$H_p(T^n, \mathbb{R}) \simeq \Lambda^p \mathbb{R}^n$$
$$H^p(T^n, \mathbb{R}) \simeq \Lambda^p \mathbb{R}^n$$

convert the stable norm on $H_p(T^n, \mathbb{R})$ into the "mass norm" on $\Lambda^p \mathbb{R}^n$ (induced by the euclidean structure on \mathbb{R}^n), and the comass norm on $H^p(T^n, \mathbb{R})$ into the comass norm on $\Lambda^p \mathbb{R}^n$. The cases p = 1and p = n - 1 are particularly simple, since then the mass and comass norms on $\Lambda^1 \mathbb{R}^n \simeq \Lambda^{n-1} \mathbb{R}^n \simeq \mathbb{R}^n$ coincide with the euclidean norm. Moreover the lattices $H_1(T^n, \mathbb{Z})_{\mathbb{R}} \subset H_1(T^n, \mathbb{R}) \simeq \mathbb{R}^n$ and $H_{n-1}(T^n, \mathbb{Z})_{\mathbb{R}} \subset H_{n-1}(T^n, \mathbb{R}) \simeq \mathbb{R}^n$ correspond to L resp. L^* under these isomorphisms. The cases 1 are considerably morecomplicated, see [31].

We are now in a position to prove our first theorem.

Theorem 2.1 Let X be a compact manifold and let $k \ge 1$ be an integer such that $H^k(X, \mathbb{R})$ is not zero and spanned by cup products

of classes of dimensions k_1, \ldots, k_m , $\Sigma_j k_j = k$. Then, for every Riemannian metric g on X, we have

$$\prod_{j=1}^{m} \mathrm{stsys}_{k_{j}}(g) \leq C(k) \left(\prod_{j} b_{k_{j}}(X)(1 + \log b_{k_{j}}(X)) \right) \mathrm{stsys}_{k}(g)$$

for a constant C(k) only depending on k.

PROOF OF THEOREM 2.1: For notational reasons we present the proof for the case that $H^k(X, \mathbb{R})$ is non-zero, and is spanned by cup products of classes of dimensions j and l, with j + l = k. The generalization to more factors is obvious. Since $H^k(X, \mathbb{R}) \neq \{0\}$, there exists $h \in H_k(X, \mathbb{Z})_{\mathbb{R}}$ such that

$$(4.2) ||h|| = \operatorname{stsys}_k(g).$$

We set $\lambda^* = \lambda_{b_j}(H^j(X,\mathbb{Z})_{\mathbb{R}}, \| \|^*)$ and $\mu^* = \lambda_{b_l}(H^l(X,\mathbb{Z})_{\mathbb{R}}, \| \|^*)$. Then we can find integral classes $\alpha_1, \ldots, \alpha_{b_j}$ in $H^j(X,\mathbb{Z})_{\mathbb{R}}$ spanning $H^j(X,\mathbb{R})$ and $\beta_1, \ldots, \beta_{b_l}$ in $H^l(X,\mathbb{Z})_{\mathbb{R}}$ spanning $H^l(X,\mathbb{R})$ such that

(4.3)
$$\|\alpha_s\|^* \leq \lambda^*$$
 for $1 \leq s \leq b_j$, and $\|\beta_t\|^* \leq \mu^*$ for $1 \leq t \leq b_l$.

By assumption $H^k(X, \mathbb{R})$ is spanned by the cup products $\alpha_s \cup \beta_t$, $1 \leq s \leq b_j$ and $1 \leq t \leq b_l$. Hence there exist indices s and t such that

$$[h, \alpha_s \cup \beta_t] = a \neq 0$$

According to Remark 3.1, we have $a \in \mathbb{Z}$. Now we complete the proof by a calculation analogous to Gromov's calculation in [23, Theorem 4.36], p. 262. If c is a real Lipschitz cycle representing h and if ω and π are closed forms representing α_s and β_t , then

(4.4)
$$1 \le |a| = \left| \int_c \omega \wedge \pi \right| \le \frac{k!}{j! \, l!} \operatorname{vol}_k(c) \|\omega\|_{\infty} \|\pi\|_{\infty},$$

cf. [17, 1.8.1] for the factor $\frac{k!}{j!l!} = \binom{k}{j}$. Using this and (4.2) and (4.3) we obtain

$$1 \le |a| \le \frac{k!}{j! \, l!} \operatorname{stsys}_k(g) \lambda^* \mu^*.$$

Since $H^p(X,\mathbb{Z})_{\mathbb{R}}$ is the lattice dual to $H_p(X,\mathbb{Z})_{\mathbb{R}}$ by Lemma 3.2, we can use Banaszczyk's inequality (4.1) to conclude

$$\operatorname{stsys}_{j}(g)\operatorname{stsys}_{l}(g) \leq \frac{k!}{j!\,l!}C^{2}\,b_{j}(1+\log b_{j})b_{l}(1+\log b_{l})\operatorname{stsys}_{k}(g).$$

This is the claim of Theorem 2.1 in the case m = 2. For arbitrary $m \ge 2$ we obtain the constant

$$\frac{k!}{\prod_j k_j!} C^m$$

in this inequality.

5 Inequalities based on cap products and Poincaré duality

In this section we prove Theorem 2.2. The proof of our Theorem 2.1 was based on Banaszczyk's inequality (4.1) applied to the stable norm. For Euclidean norms, Banaszczyk proved a sharper estimate which is linear in the dimension, cf. (7.1) below. Much older, linear estimates for the Hermite constant itself were already mentioned above, cf. (2.2) and (2.3). Replacing the stable norm by the L^2 -norm, J. Hebda [24] was able to apply such estimates in complementary dimensions: if $vol_n(q)$ is normalized to one, then the L^2 -norm on forms is bounded above by (a constant times) the comass norm and consequently the stable norm is bounded above by (a constant times) the dual L^2 -norm. A simple estimate shows that the stable norm of the Poincaré dual of a cohomology class is bounded above by (a constant times) its L^2 -norm. Putting these facts together we see that every inequality of type (4.1) leads to a systolic inequality in complementary dimensions. In Theorem 2.2 we generalize this procedure, by replacing the fundamental cycle used in Poincaré duality by a cycle of arbitrary dimension.

First we discuss a generalization of the L^2 -norm that is appropriate in this context. We stay in the well-known realm of Lipschitz cycles, although at some points the presentation would be more elegant if we used closed currents of finite mass.

V. BANGERT, M. KATZ

If $z = \sum_i r_i \sigma_i$ is a real Lipschitz (p + q)-cycle in the compact Riemannian manifold (X, g), we define a positive semidefinite, symmetric bilinear form $\langle , \rangle_{L^2(z)}$ on the space of *p*-forms as follows. Let $g_i = \sigma_i^* g$ denote the pull-back of *g* by the Lipschitz simplex σ_i . So g_i is a positive semidefinite, symmetric (0,2)-tensor field that is bounded, measurable and defined almost everywhere on the standard (p+q)-simplex Δ^{p+q} . Let \langle , \rangle_{g_i} denote the bilinear form induced by g_i on the bundle of alternating *p*-tensors over Δ^{p+q} . Let $d \operatorname{vol}_{g_i}$ denote the "volume element" induced by g_i on Δ^{p+q} , in particular $\operatorname{vol}_{g_i}(\Delta^{p+q}) = \operatorname{vol}_{p+q}(\sigma_i)$. If ω and $\overline{\omega}$ are *p*-forms on *M* we set

$$\langle \omega, \overline{\omega} \rangle_{L^2(z)} = \sum_i |r_i| \int_{\Delta^{p+q}} \langle \sigma_i^* \omega, \sigma_i^* \overline{\omega} \rangle_{g_i} d\operatorname{vol}_{g_i}$$

Since $\langle (\sigma_i^*\omega)_x, (\sigma_i^*\omega)_x \rangle_{g_i} \leq {p+q \choose p} \|\omega_{\sigma_i(x)}\|^2$ for $x \in \Delta^{p+q}$, cf. [17, 1.8.1], we have

(5.1)
$$\langle \omega, \omega \rangle_{L^2(z)} \le \binom{p+q}{p} \|\omega\|_{\infty}^2 \operatorname{vol}_{p+q}(z)$$

More generally, if π is an additional q-form, we can estimate

(5.2)
$$\left(\int_{z} \omega \wedge \pi\right)^{2} \leq \binom{p+q}{p} \|\omega\|_{\infty}^{2} \langle \pi, \pi \rangle_{L^{2}(z)} \operatorname{vol}_{p+q}(z)$$

Obviously, we can interchange the roles of ω and π in this inequality. If $p + q < n = \dim X$, there will be many *p*-forms ω such that the support of ω is disjoint from all $\sigma_i(\Delta^{p+q})$, and hence $\langle \omega, \omega \rangle_{L^2(z)} = 0$. However, under appropriate homological conditions on *z* the semidefinite form $\langle , \rangle_{L^2(z)}$ induces a scalar product on $H^p(X, \mathbb{R})$.

LEMMA 5.1 Suppose $h \in H_{p+q}(X,\mathbb{R})$ satisfies $h \cap \alpha \neq 0$ for all $\alpha \in H^p(X,\mathbb{R}) \setminus \{0\}$. If z is a Lipschitz cycle representing h, then there exists a scalar product on $H^p(X,\mathbb{R})$, denoted by $\langle , \rangle_{L^2(z)}$ as well, such that for all $\alpha \in H^p(X,\mathbb{R})$

$$\langle \alpha, \alpha \rangle_{L^2(z)} = \inf \left\{ \langle \omega, \omega \rangle_{L^2(z)} \mid \omega \text{ a closed p-form representing } \alpha \right\}.$$

PROOF: The form $\langle , \rangle_{L^2(z)}$ on the space $Z^p(X)$ of closed *p*-forms descends to a scalar product $\langle , \rangle_{L^2(z)}$ on $Z^p(X)/N$, where

$$N = \left\{ \omega \in Z^p(X) \mid \langle \omega, \omega \rangle_{L^2(z)} = 0 \right\}.$$

Now N is contained in the space $B^p(X)$ of exact forms. Indeed, if $\omega \in Z^p(X) \setminus B^p(X)$ represents $\alpha \in H^p(X, \mathbb{R}) \setminus \{0\}$ then $h \cap \alpha \neq 0$ by assumption. Hence there exists $\beta \in H^p(X, \mathbb{R})$ such that $[h \cap \alpha, \beta] \neq 0$. It follows from [20, 24.19] that $[h \cap \alpha, \beta] = [h, \alpha \cup \beta]$. If π is a closed q-form representing β , then by (5.2)

$$0 < [h, \alpha \cup \beta]^2 \le \binom{p+q}{p} \|\pi\|_{\infty}^2 \langle \omega, \omega \rangle_{L^2(z)} \operatorname{vol}_{p+q}(z)$$

and hence $\langle \omega, \omega \rangle_{L^2(z)} > 0$.

Arguing in the completion of $(Z^p(X)/N, \langle , \rangle_{L^2(z)})$ and using the fact that $Z^p(X)/B^p(X) \simeq (Z^p(X)/N)/(B^p(X)/N)$ is finitedimensional, one can easily complete the proof.

We are now in a position to prove our second theorem.

Theorem 2.2 Let X be a compact manifold and let p, q be integers, $p+q \leq \dim X$. Suppose there exists $h \in H_{p+q}(X,\mathbb{Z})_{\mathbb{R}}$ such that the cap product with h induces an injective map

$$\alpha \in H^p(X, \mathbb{Z})_{\mathbb{R}} \to h \cap \alpha \in H_q(X, \mathbb{Z})_{\mathbb{R}}.$$

Then, for every Riemannian metric g on X, we have

$$\operatorname{stsys}_p(g) \operatorname{stsys}_q(g) \le C(p,q)b_p(X) \|h\|$$

for a constant C(p,q) only depending on p and q. If our assumption is satisfied for all $h \in H_{p+q}(X,\mathbb{Z})_{\mathbb{R}} \setminus \{0\}$ then

(2.1)
$$\operatorname{stsys}_p(g)\operatorname{stsys}_q(g) \le C(p,q)b_p(X)\operatorname{stsys}_{p+q}(g).$$

PROOF OF THEOREM 2.2: Let z be a Lipschitz cycle representing h. By the preceding lemma we can consider the scalar product $\langle , \rangle_{L^2(z)}$ on $H^p_{dR}(X)$. From (5.1) we obtain

(5.3)
$$|\alpha|_{L^{2}(z)} \leq {\binom{p+q}{p}}^{\frac{1}{2}} \operatorname{vol}_{p+q}(z)^{\frac{1}{2}} ||\alpha||^{*}$$

for $\alpha \in H^p_{dR}(X)$. Hence the dual euclidean norm on $H_p(X, \mathbb{R})$, also denoted by $||_{L^2(z)}$, satisfies

(5.4)
$$||k|| \le {\binom{p+q}{p}}^{\frac{1}{2}} \operatorname{vol}_{p+q}(z)^{\frac{1}{2}} |k|_{L^{2}(z)}$$

for $k \in H_p(X, \mathbb{R})$. If $\alpha \in H^p_{dR}(X)$ then

16

$$\|h \cap \alpha\| = \sup \left\{ [h, \alpha \cup \beta] \mid \beta \in H^q(X, \mathbb{R}), \|\beta\|^* \le 1 \right\}$$

since the stable norm $\| \|$ and the comass norm $\| \|^*$ are dual to each other and $[h \cap \alpha, \beta] = [h, \alpha \cup \beta]$. Now (5.2) implies

(5.5)
$$||h \cap \alpha|| \le {\binom{p+q}{p}}^{\frac{1}{2}} \operatorname{vol}_{p+q}(z)^{\frac{1}{2}} |\alpha|_{L^2(z)}.$$

Set $b = b_p(X)$. By the definition of the constant γ'_b and Lemma 3.2 there exist $k \in H_p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}$ and $\alpha \in H^p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}$ such that

$$|k|_{L^2(z)} |\alpha|_{L^2(z)} \le \gamma'_b.$$

Using (5.4) and (5.5) we conclude

$$||k|| ||h \cap \alpha|| \le {p+q \choose p} \gamma'_b \operatorname{vol}_{p+q}(z).$$

Since $h \cap \alpha \neq 0$ by assumption and $h \cap \alpha \in H_q(X, \mathbb{Z})_{\mathbb{R}}$ by Remark 3.1 we conclude

(5.6)
$$\operatorname{stsys}_p(g) \operatorname{stsys}_q(g) \le {\binom{p+q}{p}} \gamma_b' \|h\|.$$

Together with (2.2) this proves Theorem 2.2.

REMARK 5.2 In [24], Proposition 6, J. Hebda essentially proved Theorem 2.2 in the case n = p+q. Indeed, his arguments imply that

$$\operatorname{stsys}_p(g) \operatorname{stsys}_{n-p}(g) \le \binom{n}{p} \gamma_b \operatorname{vol}_n(g)$$

17

for every Riemannian metric g on a compact, orientable manifold X with $b = b_p(X) > 0$. Theorem 2.2, however, can be applied in many situations that are not covered by [24, Proposition 6]. In general, the resulting inequality will be sharper than the inequality provided by Theorem 2.1. We give a simple example. We choose integers $p \ge 1, q \ge 1$ and m with m > p + q and apply (5.6) to the manifold $X = T^{p+q} \times S^m$. Recalling that $\gamma'_b \le \frac{2}{3}b$ for $b \ge 2$, cf. (2.2), we obtain

(5.7)
$$\operatorname{stsys}_p(g)\operatorname{stsys}_q(g) \le \frac{2}{3} \binom{p+q}{p}^2 \operatorname{stsys}_{p+q}(g)$$

for every metric g on $X = T^{p+q} \times S^m$. Here, the dependence on $b_p(X) = b_q(X) = {p+q \choose p}$ is quadratic, while the inequality following from the proof of Theorem 2.1 is

$$\operatorname{stsys}_p(g)\operatorname{stsys}_q(g) \le C^2 \binom{p+q}{p} b_p(X)^2 (1+\log b_p(X))^2 \operatorname{stsys}_{p+q}(g)$$

where C is the universal constant in Banaszczyk's inequality (4.1).

6 A sharp inequality in codimension 1

We apply the arguments of section 5 to a fundamental cycle $h = \xi_{\mathbb{R}} \in H_n(X, \mathbb{Z})_{\mathbb{R}}$ of X.

Corollary 2.3 Let X be a compact, orientable manifold, dim X = n, with positive first Betti number $b = b_1(X)$. Then, for every Riemannian metric g on X, we have

 $\operatorname{stsys}_1(g)\operatorname{sys}_{n-1}(g) \le \gamma'_b \operatorname{vol}_n(g).$

Equality is attained for a flat torus \mathbb{R}^n/L where $L \subseteq \mathbb{R}^n$ is a lattice with $\lambda_1(L)\lambda_1(L^*) = \gamma'_n$.

PROOF OF COROLLARY 2.3: Since p = 1, q = n - 1 we can replace the factors $\binom{p+q}{p}^{\frac{1}{2}}$ in (5.4) and (5.5) by one, cf. [17, 1.8.1]. So our final statement is

(6.1)
$$\operatorname{stsys}_1(g)\operatorname{sys}_{n-1}(g) \le \gamma'_b \operatorname{vol}_n(g).$$

V. BANGERT, M. KATZ

There exist lattices L in b-dimensional euclidean space such that

$$\lambda_1(L)\lambda_1(L^*) = \gamma_b',$$

cf. [11]. Consider the corresponding flat torus (T^b, g) . Then Example 4.1 shows that for such tori equality holds in (6.1).

The boundary case of equality in this optimal inequality will be studied in [9].

7 A conformally invariant inequality in the middle dimension

Instead of using the constant γ'_b together with inequality (2.2) in the proof of Theorem 2.2, we can also employ the following estimate by W. Banaszczyk [7], Theorem 2.1, in the case i = b:

If L is a lattice in b-dimensional euclidean space (with the standard innner product norm) and $i \in \{1, \ldots, b\}$, then

(7.1)
$$1 \le \lambda_i(L)\lambda_{b-i+1}(L^*) \le b.$$

Specialized to the case p + q = n, $h = \xi_{\mathbb{R}}$, the proof of Theorem 2.2, combined with (7.1), leads to the estimate

(7.2)
$$\operatorname{stsys}_p(g) \operatorname{stsys}_{n-p}(g) \le \frac{\lambda_1}{\lambda_{b_p}} \binom{n}{p} b_p \operatorname{vol}_n(g)$$

where $\lambda_i = \lambda_i (H_p(X, \mathbb{Z})_{\mathbb{R}}, | |_{L^2})$ and $b_p = b_p(X)$.

In the case of middle dimension $p = \frac{n}{2}$ a similar argument leads to a conformally invariant estimate, cf. [21, 7.4.A]. This case is special in the following sense: the L^2 -norms of p-forms ω

$$|w|_{L^2} = \left(\int_X \langle w, w \rangle d \operatorname{vol}_n\right)^{\frac{1}{2}}$$

and

$$\|\omega\|_{L^2} = \left(\int_X (\|\omega_x\|)^2 d\operatorname{vol}_n(x)\right)^{\frac{1}{2}}$$

are invariant under conformal changes of the metric g. They induce respectively the usual L^2 -norm $||_{L^2}$ on $H^p(X, \mathbb{R})$, for which the

harmonic forms are minimizing representatives of the cohomology classes, and a norm $\| \|_{L^2}^*$ on $H^p(X, \mathbb{R})$ such that

$$\binom{n}{p}^{-\frac{1}{2}} |\alpha|_{L^2} \le ||\alpha||_{L^2}^* \le |\alpha|_{L^2}.$$

We also consider the conformally invariant dual norms on $H_p(X, \mathbb{R})$ that satisfy

(7.3)
$$|h|_{L^2} \le ||h||_{L^2} \le {\binom{n}{p}}^{\frac{1}{2}} |h|_{L^2}.$$

It is not difficult to prove that

(7.4)
$$||h||_{L^2} = \sup\{||h||_{g'} \operatorname{vol}_n(g')^{-\frac{1}{2}} | g' \text{ conformal to } g\},$$

cf. [21, 7.4.A]. Always assuming $p = \frac{n}{2}$, we introduce the conformal invariant

$$\operatorname{conf}_p(g) = \min\{\|h\|_{L^2} \mid h \in H_p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}\}.$$

Note that (7.4) implies that

(7.5)
$$\operatorname{stsys}_p(g')\operatorname{vol}_n(g')^{-\frac{1}{2}} \le \operatorname{conf}_p(g)$$

for every pair g', g of conformal Riemannian metrics. However, it can happen that

$$\sup\{\operatorname{stsys}_p(g')\operatorname{vol}_n(g')^{-\frac{1}{2}} \mid g' \text{ conformal to } g\} < \operatorname{conf}_p(g).$$

This can be seen by comparing Gromov's universal upper bound contained in inequality (+) of [22, 2.C] for the 1-systole to the lower bound for conf₁ of surfaces which follows from the work by P. Buser and P. Sarnak [15]. Indeed, Gromov asserts the existence of a universal constant C such that

(7.6)
$$\operatorname{sys}_1(g) \operatorname{vol}_2(g)^{-\frac{1}{2}} \le C \frac{\log \gamma}{\sqrt{\gamma}}$$

whenever g is a Riemannian metric on a closed, orientable surface of genus $\gamma \geq 2$, cf. (2.5) above. On the other hand [15, 1.13] states that the supremum of $\operatorname{conf}_1(g)^2$ over all Riemannian metrics on a closed, orientable surface of genus γ is bounded below by $c \log(\gamma)$ for some universal constant c > 0. Thus, a priori, an upper bound for $\operatorname{conf}_p(g)$ as given in Corollary 2.5 is stronger than the same upper bound for $\operatorname{stsys}_p(g) \operatorname{vol}_n(g)^{-\frac{1}{2}}$, cf. also [26].

Corollary 2.5 Let $p = \frac{n}{2}$ and $b_p = b_p(X)$. Then

$$\operatorname{conf}_p(g)^2 \le \frac{\lambda_1}{\lambda_{b_p}} \binom{n}{p} b_p$$

where $\lambda_i = \lambda_i (H_p(X, \mathbb{Z})_{\mathbb{R}}, | |_{L^2}).$

20

PROOF OF COROLLARY 2.5: Note that the map

 $\mathrm{PD}_{\mathbb{R}}: (H^p(X,\mathbb{R}), |\mid_{L^2}) \to (H_{n-p}(X,\mathbb{R}), |\mid_{L^2})$

is an isometry. Indeed, if $\alpha \in H^p(X, \mathbb{R})$ and if ω is the harmonic *p*-form representing α , then

$$|\operatorname{PD}_{\mathbb{R}}(\alpha)|_{L^{2}} = \sup\{\int_{M} \omega \wedge \pi \mid \pi \in Z^{n-p}(M), |\pi|_{L^{2}} = 1\} \\ = \sup\{\langle \omega, *\pi \rangle_{L^{2}} \mid \pi \in Z^{n-p}(M), |\pi|_{L^{2}} = 1\} = |\alpha|_{L^{2}}.$$

Setting $\lambda_1^* = \lambda_1(H^p(X, \mathbb{Z})_{\mathbb{R}}, ||_{L^2})$ and $b = b_p(X)$, we conclude $\lambda_1 = \lambda_1^*$ and $\lambda_1^2 \leq \frac{\lambda_1}{\lambda_b} b$ by Banaszczyk's inequality (7.1). Now (7.3) implies our claim.

One can also apply inequality (7.2) in the opposite direction:

COROLLARY 7.1 Let X be a compact, orientable manifold of dimension n = 2p, with $b_p(X) > 0$. For D > 0 let \mathcal{G}_D denote the set of Riemannian metrics g on X such that $\operatorname{vol}_n(g) \leq D$ and $\operatorname{stsys}_p(g) \geq D^{-1}$. Then the set of flat Finsler metrics defined by the stable norms of metrics $g \in \mathcal{G}_D$ on the Jacobian torus $H_p(X, \mathbb{R})/H_p(X, \mathbb{Z})_{\mathbb{R}}$ is contained in a compact part of the set of all flat Finsler metrics on the Jacobian torus.

PROOF: First we will prove the existence of a basis of $H_p(X, \mathbb{Z})_{\mathbb{R}}$ whose elements have stable norm bounded by some function of D,

p and $b_p(X)$. We use Banaszczyk's inequality (4.1) in the case $i = b = b_p(X)$ for the stable norm || || on $L = H_p(X, \mathbb{Z})_{\mathbb{R}}$ and the comass norm $|| ||^*$ on $L^* = H^p(X, \mathbb{Z})_{\mathbb{R}}$ to obtain

(7.7)
$$\lambda_b \lambda_1^* \le Cb(1 + \log b)$$

where $\lambda_i = \lambda_i (H_p(X, \mathbb{Z})_{\mathbb{R}}, || ||)$, and $\lambda_i^* = \lambda_i (H^p(X, \mathbb{Z})_{\mathbb{R}}, || ||^*)$. If $\alpha \in H^p(X, \mathbb{Z})_{\mathbb{R}}$ and $\|\alpha\|^* = \lambda_1^*$ then (5.3) and (5.5) imply

$$\|\operatorname{PD}_{\mathbb{R}}(\alpha)\| \le {\binom{n}{p}}\operatorname{vol}_{n}(g)\|\alpha\|^{*}$$

and hence

$$\operatorname{stsys}_p(g) \le \binom{n}{p} \operatorname{vol}_n(g) \lambda_1^*$$

Since $\operatorname{stsys}_p(g) = \lambda_1$, we can use (7.7) to conclude

(7.8)
$$\operatorname{stsys}_p(g)^2 \le \frac{\lambda_1}{\lambda_b} {n \choose p} C b(1 + \log b) \operatorname{vol}_n(g).$$

Now assume $g \in \mathcal{G}_D$. Then (7.8) implies $\lambda_b \leq {n \choose p} C b(1 + \log b) D^2$. Using [16], p.135, Lemma 8, we obtain a basis v_1, \ldots, v_b of $H_p(X, \mathbb{Z})_{\mathbb{R}}$ such that

(7.9)
$$||v_i|| \le {\binom{n}{p}} C b^2 (1 + \log b) D^2 = E$$

for $1 \leq i \leq b$. Consider the isomorphism $H_p(X, \mathbb{R}) \to \mathbb{R}^b$ mapping (v_1, \ldots, v_b) to the standard basis (e_1, \ldots, e_b) and consider the induced norm on \mathbb{R}^b , denoted by || || as well. Then the unit || ||-ball $B = \{x \in \mathbb{R}^b \mid ||x|| \leq 1\}$ satisfies

(7.10)
$$\left\{ x \in \mathbb{R}^b \mid \sum_{i=1}^b |x_i| \le E^{-1} \right\} \subseteq B$$

On the other hand $\min \{ ||x|| | x \in \mathbb{Z}^b \setminus \{0\} \} = \lambda_1 \ge D^{-1}$ and hence Minkowski's Theorem, cf. [16], p.71, Theorem II, implies

(7.11)
$$\operatorname{vol}_b(B) \le (2D)^b$$

where vol_b denotes the usual Lebesgue measure on \mathbb{R}^b . Now our claim follows from the fact that for fixed numbers D and E > 0 the set of convex bodies in \mathbb{R}^b satisfying (7.10) and (7.11) is compact with respect to the Hausdorff metric.

V. BANGERT, M. KATZ

8 A sublinear estimate for a single systole

If X is a compact, orientable manifold, if dim X = n = kp and if there are classes $\alpha_1, \ldots, \alpha_k \in H^p(X, \mathbb{R})$ such that $\alpha_1 \cup \ldots \cup \alpha_k \neq 0$, then Theorem 2.1 implies

$$\operatorname{stsys}_p(g) \leq \tilde{C}(n) \, b_p(X) (1 + \log b_p(X)) \operatorname{vol}_n(g)^{\frac{1}{k}}$$

for every Riemannian metric g on X. Using Poincaré duality as in Theorem 2.2 we can improve this estimate to sublinear dependence on the Betti number $b = b_p(X)$.

Theorem 2.7 Let X be a compact, orientable manifold, of dimension n = kp. Suppose there are classes $\beta_1, \ldots, \beta_k \in H^p(X, \mathbb{R})$ such that $\beta_1 \cup \ldots \cup \beta_k \neq 0$. Then, for every Riemannian metric g on X, we have

$$\operatorname{stsys}_p(g) \le C(n) \left(b_p(X)(1 + \log b_p(X)) \right)^{\frac{k-1}{k}} \operatorname{vol}_n(g)^{\frac{1}{k}}$$

for a constant C(n) only depending on n.

PROOF OF THEOREM 2.7: We start by choosing linearly independent classes $\alpha_1, \ldots, \alpha_b$ in $H^p(X, \mathbb{Z})_{\mathbb{R}}$ such that $\|\alpha_i\|^* \leq \lambda_b^* = \lambda_b(H^p(X, \mathbb{Z})_{\mathbb{R}}, \|\|^*)$. Our hypothesis implies that there exist indices i_1, \ldots, i_k in $\{1, \ldots, b\}$ such that

$$\alpha_{i_1} \cup \ldots \cup \alpha_{i_k} \neq 0.$$

We may assume that X is connected and oriented. Let $h \in H_p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}$ denote the Poincaré dual of $\alpha_{i_1} \cup \ldots \cup \alpha_{i_{k-1}}$. Let $\omega_1, \ldots, \omega_b$ be p-forms representing $\alpha_1, \ldots, \alpha_b$, and let ω be an arbitrary closed p-form representing some $\alpha \in H^p(X, \mathbb{R})$. Then

$$[h, \alpha] = \left[\xi_{\mathbb{R}}, \alpha_{i_1} \cup \ldots \cup \alpha_{i_{k-1}} \cup \alpha \right] = \int_X \omega_{i_1} \wedge \ldots \wedge \omega_{i_{k-1}} \wedge \omega$$

$$\leq \frac{n!}{(p!)^k} \left(\prod_{j=1}^{k-1} \|\omega_{i_j}\|_{\infty} \right) \|\omega\|_{\infty} \operatorname{vol}_n(X).$$

Since $||h|| = \sup_{\|\alpha\|^* \le 1} [h, \alpha]$, we obtain

$$\|h\| \le \frac{n!}{(p!)^k} \left(\prod_{j=1}^{k-1} \|\alpha_{i_j}\|^* \right) \operatorname{vol}_n(X) \le \frac{n!}{(p!)^k} \left(\lambda_b^*\right)^{k-1} \operatorname{vol}_n(X).$$

Now Banaszczyk's inequality (4.1) implies $\operatorname{stsys}_p(g)\lambda_b^* \leq Cb(1 + \log b)$, and hence

$$\operatorname{stsys}_p(g)^k \le \operatorname{stsys}_p(g)^{k-1} ||h|| \le C^{k-1} \frac{n!}{(p!)^k} (b(1+\log b))^{k-1} \operatorname{vol}_n(X).$$

This proves Theorem 2.7.

Note that for every $r > 1 - \frac{1}{k}$ the factor $(b(1 + \log b))^{\frac{k-1}{k}}$ grows less fast than b^r when $b \to \infty$.

If dim X = n = 3p we can improve Theorem 2.7 as follows:

THEOREM 8.1 Let X be a compact, orientable manifold, of dimension n = 3p. Suppose there exist classes $\beta_1, \beta_2, \beta_3 \in H^p(X, \mathbb{R})$ such that $\beta_1 \cup \beta_2 \cup \beta_3 \neq 0$. Then for every Riemannian metric g on X we have

stsys_p(g)
$$\leq \left(\binom{n}{p} \frac{n!}{(p!)^3}\right)^{\frac{1}{3}} b_p(X)^{\frac{2}{3}} \operatorname{vol}_n(g)^{\frac{1}{3}}.$$

PROOF: Set $\lambda_1 = \lambda_1(H_p(X,\mathbb{Z})_{\mathbb{R}}, ||_{L^2})$. Let $b = b_p(X)$, and set also $\lambda_b^* = \lambda_b(H^p(X,\mathbb{Z})_{\mathbb{R}}, ||_{L^2})$. We choose linearly independent classes $\alpha_1, \ldots, \alpha_b$ in $H^p(X,\mathbb{Z})_{\mathbb{R}}$ such that $|\alpha_i|_{L^2} \leq \lambda_b^*$. As in the proof of Theorem 2.7 we choose indices i_1, i_2, i_3 in $\{1, \ldots, b\}$ such that

$$\alpha_{i_1} \cup \alpha_{i_2} \cup \alpha_{i_3} \neq 0.$$

We may assume that X is connected and oriented. We consider $h = \operatorname{PD}_{\mathbb{R}}(\alpha_{i_1} \cup \alpha_{i_2}) \in H_p(X, \mathbb{Z})_{\mathbb{R}} \setminus \{0\}$, harmonic forms $\omega_{i_1}, \omega_{i_2}$ representing $\alpha_{i_1}, \alpha_{i_2}$ and a closed *p*-form ω representing an arbitrary $\alpha \in H^p(X, \mathbb{R})$. Then we have

$$[h,\alpha] = \int_X \omega_{i_1} \wedge \omega_{i_2} \wedge \omega \le \frac{n!}{(p!)^3} \|\omega\|_{\infty} \int_X |\omega_{i_1}| |\omega_{i_2}| d\operatorname{vol}_n$$

cf. [17], 1.7.5 and 1.8.1 for the constant. Using the definition of comass $||h|| = \sup_{\|\alpha\|^* \le 1} [h, \alpha]$, we obtain

$$||h|| \le \frac{n!}{(p!)^3} |\alpha_{i_1}|_{L^2} |\alpha_{i_2}|_{L^2} \le \frac{n!}{(p!)^3} (\lambda_b^*)^2.$$

V. BANGERT, M. KATZ

We apply (5.4) to the fundamental cycle z of M to obtain

$$\operatorname{stsys}_p(g) \le {\binom{n}{p}}^{\frac{1}{2}} \lambda_1 \operatorname{vol}_n(g)^{\frac{1}{2}}.$$

The preceding inequalities imply

$$\operatorname{stsys}_p(g)^3 \le \operatorname{stsys}_p(g)^2 ||h|| \le \binom{n}{p} \frac{n!}{(p!)^3} (\lambda_1 \lambda_b^*)^2 \operatorname{vol}_n(g).$$

Now we apply Banaszczyk's inequality $\lambda_1 \lambda_b^* \leq b$, cf. (7.1), to complete the proof.

Acknowledgment. The authors have benefited from helpful discussions with M. Kreck and express appreciation to L. Ambrosio, G. Dula, J. Lagarias, C. LeBrun, F. Morgan, S. Weinberger and B. White for insightful comments.

Bibliography

- Accola, R. Differentials and extremal length on Riemann surfaces. Proc. Nat. Acad. Sci. USA 46 (1960), 540–543.
- [2] Babenko, I. Asymptotic invariants of smooth manifolds. Russian Acad. Sci. Izv. Math. 41 (1993), 1–38.
- [3] Babenko, I. Forte souplesse intersystolique de variétés fermées et de polyèdres. Annales de l'Institut Fourier **52** (2002), in press.
- [4] Babenko, I. Loewner's conjecture, Besicovich's example, and relative systolic geometry. [Russian]. Mat. Sbornik 193 (2002), in press.
- [5] Babenko, I.; Katz, M. Systolic freedom of orientable manifolds. Ann. Sci. Ecole Norm. Sup. (Sér. 4) 31 (1998), 787–809.
- [6] Babenko, I.; Katz, M.; Suciu, A. Volumes, middle-dimensional systoles, and Whitehead products. *Math. Res. Lett.* 5 (1998), 461–471.
- [7] Banaszczyk, W. New bounds in some transference theorems in the geometry of numbers. *Math. Ann.* **296** (1993), 625–635.
- [8] Banaszczyk, W. Inequalities for convex bodies and polar reciprocal lattices in ℝⁿ II: Application of K-convexity. Discrete Comput. Geom. 16 (1996), 305–311.

- [9] Bangert, V.; Katz, M. Riemannian manifolds with harmonic 1-forms of constant norm. In preparation.
- [10] Bavard, C. Inégalités isosystoliques conformes pour la bouteille de Klein. Geom. Dedicata 27 (1988), 349–355.
- [11] Bergé, A.-M.; Martinet, J. Sur un problème de dualité lié aux sphères en géométrie des nombres. J. Number Theory 32 (1989), 14–42.
- [12] Berger, M. A l'ombre de Loewner. Ann. Scient. Ec. Norm. Sup. (Sér. 4) 5 (1972), 241–260.
- [13] Berger, M. Systoles et applications selon Gromov. Séminaire N. Bourbaki, exposé 771, Astérisque 216 (1993), 279–310.
- [14] Blatter, C. Uber Extremallängen auf geschlossenen Flächen. Comment. Math. Helv. 35 (1961), 153–168.
- [15] Buser, P.; Sarnak, P. On the period matrix of a Riemann surface of large genus. With an appendix by J. H. Conway and N. J. A. Sloane. *Invent. Math.* **117** (1994), 27–56.
- [16] Cassels, J. W. S. An introduction to the geometry of numbers. Second printing. Grundlehren der mathematischen Wissenschaften, 99. Springer-Verlag, Berlin-Heidelberg-NewYork, 1971.
- [17] Federer, H. Geometric measure theory. Grundlehren der mathematischen Wissenschaften, 153. Springer-Verlag, Berlin, 1969.
- [18] Federer, H. Real flat chains, cochains, and variational problems. Indiana Univ. Math. J. 24 (1974), 351–407.
- [19] Freedman, M. Z₂ Systolic Freedom. Geometry and Topology Monographs, vol. 2, Proceedings of the Kirbyfest (J. Hass, M. Scharlemann eds.), 113– 123, Geometry & Topology, Coventry, 1999.
- [20] Greenberg, M. Lectures on Algebraic Topology. W.A. Benjamin, Inc., New York, Amsterdam, 1967.
- [21] Gromov, M. Filling Riemannian manifolds. J. Differential Geom. 18 (1983), 1–147.
- [22] Gromov, M. Systoles and intersystolic inequalities. Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992), 291–362, Sémin. Congr., vol. 1, Soc. Math. France, Paris, 1996.
- [23] Gromov, M. Metric structures for Riemannian and non-Riemannian spaces. Progress in Mathematics 152, Birkhäuser, Boston, 1999.
- [24] Hebda, J. The collars of a Riemannian manifold and stable isosystolic inequalities. *Pacific J. Math.* **121** (1986), 339-356.
- [25] Katz, M. Local calibration of mass and systolic geometry. Geom. Funct. Anal. (GAFA), in press.

- [26] Katz, M. Four-manifold systoles and surjectivity of period map. Preprint 2002.
- [27] Katz, M.; Kreck, M.; Suciu, A. Massey products and systolic inequalities. In preparation.
- [28] Katz, M.; Suciu, A. Volume of Riemannian manifolds, geometric inequalities, and homotopy theory. *Tel Aviv Topology Conference: Rothenberg Festschrift (M. Farber, W. Lück, and S. Weinberger, eds.)*, 113–136. Contemp. Math., vol. 231, Amer. Math. Soc., Providence, RI, 1999.
- [29] Katz, M.; Suciu, A. Systolic freedom of loop space. Geom. Funct. Anal. (GAFA) 11 (2001), 60–73.
- [30] Lagarias, J.C.; Lenstra, H.W., Jr.; Schnorr, C.P. Bounds for Korkin-Zolotarev reduced bases and successive minima of a lattice and its reciprocal lattice. *Combinatorica* **10** (1990), 343–358.
- [31] Lawson, H.B. The stable homology of a flat torus. Math. Scand. 36 (1975), 49–73.
- [32] Moser, J. Minimal solutions of variational problems on a torus. Ann. Inst. H. Poincaré-Analyse non linéaire 3 (1986), 229–272.
- [33] Moser, J.; Struwe, M. On a Liouville-type theorem for linear and nonlinear elliptic differential equations on a torus. *Bol. Soc. Brasil. Mat.* 23 (1992), 1–20.
- [34] Pu, P.M. Some inequalities in certain nonorientable Riemannian manifolds. Pacific J. Math. 2 (1952), 55–71.
- [35] Senn, W. Strikte Konvexität für Variationsprobleme auf dem ndimensionalen Torus. manuscripta math. 71 (1991), 45–65.
- [36] Senn, W. Uber Mosers regularisiertes Variationsproblem für minimale Blätterungen des n-dimensionalen Torus. J. Appl. Math. Physics (ZAMP) 42 (1991), 527–546.
- [37] Warner, F.W. Foundations of Differentiable Manifolds and Lie Groups. Graduate Texts in Mathematics, 94. Springer-Verlag, New York-Heidelberg-Berlin, 1983.
- [38] White, B. Regularity of area-minimizing hypersurfaces at boundaries with multiplicity. *Seminar on minimal submanifolds (E. Bombieri ed.)*, 293-301. Annals of Mathematics Studies 103. Princeton University Press, Princeton N.J., 1983.

Received Month 200X.