

The Action of Adeles on the Residue Complex

Amnon Yekutieli
 Department of Mathematics
 Ben Gurion University
 Be'er Sheva 84105, ISRAEL
 amyekut@math.bgu.ac.il

Dedicated to Professor Steven Kleiman on the Occasion of his Sixtieth Birthday

Abstract

Let X be a scheme of finite type over a perfect field \mathbb{k} . In this paper we study the relation between two objects associated to X : the *Grothendieck residue complex* \mathcal{K}_X and the *Beilinson adeles complex* $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$. The latter is a differential graded algebra (DGA). Our first main result (Theorem 0.1) is that \mathcal{K}_X is a right differential graded (DG) module over $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$.

We give an application to de Rham theory. Define graded sheaves $\mathcal{F}_X := \mathcal{H}om_{\mathcal{O}_X}(\Omega_{X/\mathbb{k}}, \mathcal{K}_X)$ and $\mathcal{A}_X := \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X) \otimes_{\mathcal{O}_X} \Omega_{X/\mathbb{k}}$. It is known that \mathcal{A}_X is a DGA. Our second main result (Theorem 0.2) is that \mathcal{F}_X is a right DG \mathcal{A}_X -module. When X is smooth then \mathcal{F}_X calculates de Rham homology, \mathcal{A}_X calculates cohomology, and the action induces the cap product. We extend these constructions to singular schemes in characteristic 0 using smooth formal embeddings.

0 Introduction

Let X be a scheme of finite type over a field \mathbb{k} . In this paper we study the relation between two objects associated to X : the *Grothendieck residue complex* \mathcal{K}_X and the *Beilinson adeles complex* $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$.

Grothendieck duality theory, developed around 1960 (cf. [RD]), is a vast generalization of Serre duality. It is a deep and complicated theory, fully expressible only in the language of derived categories. Attempts to simplify it or find some explicit presentation of it attracted a considerable amount

Mathematics Subject Classification 2000. Primary: 14F05; Secondary: 14F40, 13B35, 14B15.

of research (a partial list of references is [AK], [KI], [Li], [HK], [KW], [LS], [Ye1], [Ne], [AJL] and [Co]).

In our situation denote by $\pi : X \rightarrow \text{Spec } \mathbb{k}$ the structural morphism. Then there is a functor $\pi^! : D_c^+(\text{Mod } \mathbb{k}) \rightarrow D_c^+(\text{Mod } \mathcal{O}_X)$ between derived categories, called the twisted inverse image. The object $\pi^! \mathbb{k}$ is a dualizing complex on X . It has a canonical representative, namely its Cousin complex \mathcal{K}_X , which is called the *residue complex* of X . This is a bounded complex of quasi-coherent injective \mathcal{O}_X -modules, and as a sheaf $\mathcal{K}_X = \bigoplus_{x \in X} \mathcal{K}_X(x)$, where $\mathcal{K}_X(x)$ is a constant sheaf with support $\overline{\{x\}}$. \mathcal{K}_X enjoys some remarkable properties, that are deduced from corresponding properties of $\pi^!$.

Almost twenty years later Beilinson introduced his scheme theoretic adèles (see [Be]). This high dimensional generalization of the classical adèles of a curve is actually pretty easy to define (see Section 1). Given any quasi-coherent \mathcal{O}_X -module \mathcal{M} , the complex adèles with values in \mathcal{M} is a bounded complex $\mathbb{A}_{\text{red}}(\mathcal{M})$ of flasque \mathcal{O}_X -modules, and there is a canonical quasi-isomorphism $\mathcal{M} \rightarrow \mathbb{A}_{\text{red}}(\mathcal{M})$. The sheaf $\mathbb{A}_{\text{red}}^q(\mathcal{M})$ is a “restricted product” of local factors, each such local factor corresponding to the geometric data of a chain (x_0, \dots, x_q) of points in X . Taking $\mathcal{M} = \mathcal{O}_X$ we obtain a DGA (differential graded algebra) $\mathbb{A}_{\text{red}}(\mathcal{O}_X)$.

Here is the first main result of this paper.

Theorem 0.1. *Suppose \mathbb{k} is a perfect field and X is a finite type \mathbb{k} -scheme. Then \mathcal{K}_X is a right DG $\mathbb{A}_{\text{red}}(\mathcal{O}_X)$ -module.*

The theorem is restated in more detail in Section 1 (Theorem 1.14) and proved there, using the explicit construction of \mathcal{K}_X described in [Ye3]. This construction is based on the theory of *Beilinson completion algebras* (BCAs) developed in [Ye2]. The action of the adèles is by “taking residues”: multiplication by an adèle supported on a chain (x_0, \dots, x_q) is a map from $\mathcal{K}_X(x_0)$ to $\mathcal{K}_X(x_q)$. In Question 1.20 we speculate on a generalization of Theorem 0.1.

In Section 2 we move on to de Rham theory. Let $(\Omega_{X/\mathbb{k}}, d)$ be the algebraic de Rham complex of X , which is a sheaf of commutative DGAs over \mathbb{k} . The graded sheaf

$$\mathcal{F}_X := \text{Hom}_{\mathcal{O}_X}(\Omega_{X/\mathbb{k}}, \mathcal{K}_X)$$

is a graded $\Omega_{X/\mathbb{k}}$ -module. According to [Ye3] \mathcal{F}_X has a coboundary operator D that’s a differential operator of order ≤ 1 over \mathcal{O}_X , and (\mathcal{F}_X, D) is a DG $\Omega_{X/\mathbb{k}}$ -module.

On the other hand we have the graded algebra

$$\mathcal{A}_X := \bigoplus_{p,q} \mathbb{A}_{\text{red}}^q(\Omega_{X/\mathbb{k}}^p) \cong \mathbb{A}_{\text{red}}(\mathcal{O}_X) \otimes_{\mathcal{O}_X} \Omega_{X/\mathbb{k}}.$$

By [HY1] there is a differential D that makes (\mathcal{A}_X, D) into a DGA, and

moreover $\Omega_{X/\mathbb{k}} \rightarrow \mathcal{A}_X$ is a DGA quasi-isomorphism. Theorem 0.1 implies that \mathcal{F}_X is a graded right \mathcal{A}_X -module. Our second main result is:

Theorem 0.2. *Suppose \mathbb{k} is a perfect field and X is a finite type \mathbb{k} -scheme. Then \mathcal{F}_X is a DG right \mathcal{A}_X -module.*

Theorem 0.2 is used in [HY2] to state the adelic Gauss-Bonnet formula, which is proved there. See also Remark 2.6 below.

Consider a finite type \mathbb{k} -scheme Y and a closed subscheme $X \subset Y$. Define $\mathfrak{X} := Y_{/X}$, the formal completion of Y along X (cf. [EGA] I). Let X_i be the i th infinitesimal neighborhood of $X = X_0$ in Y . Define $\mathcal{A}_{\mathfrak{X}} := \lim_{\leftarrow i} \mathcal{A}_{X_i}$ and $\mathcal{F}_{\mathfrak{X}} := \lim_{i \rightarrow} \mathcal{F}_{X_i}$.

Corollary 0.3. *$\mathcal{F}_{\mathfrak{X}}$ is a right DG $\mathcal{A}_{\mathfrak{X}}$ -module. If $\text{char } \mathbb{k} = 0$ and Y is smooth over \mathbb{k} then de Rham cohomology is $H_{\text{DR}}^p(X) = H^p \Gamma(X, \mathcal{A}_{\mathfrak{X}})$, de Rham homology is $H_p^{\text{DR}}(X) = H^{-p} \Gamma(X, \mathcal{F}_{\mathfrak{X}})$, and the right action of $\mathcal{A}_{\mathfrak{X}}$ on $\mathcal{F}_{\mathfrak{X}}$ induces the cap product.*

The embedding $X \subset \mathfrak{X}$ is an instance of a smooth formal embedding in the sense of [Ye4].

Theorem 0.2 and Corollary 0.3 are proved at the end of Section 2.

Acknowledgments. The paper is dedicated to Steven Kleiman in gratitude for his longtime interest and support of my work on residues. I also wish to thank Reinhold Hübl, Joseph Lipman and Pramathanath Sastry for helpful discussions. Finally I wish to express thanks to the referee for reading the paper carefully and suggesting improvements.

1 The Action

Let us begin with a review of *Beilinson adèles* on a noetherian scheme X . A chain of length q of points in X is a sequence $\xi = (x_0, \dots, x_q)$ of points with $x_{i+1} \in \overline{\{x_i\}}$. Denote by $S(X)_q$ the set of length q chains, so $\{S(X)_q\}_{q \geq 0}$ is a simplicial set. For a subset $T \subset S(X)_q$ and a point $x \in X$ let

$$\hat{x}T := \{(x_1, \dots, x_q) \mid (x, x_1, \dots, x_q) \in T\}.$$

Denote by $\mathfrak{m}_x \subset \mathcal{O}_{X,x}$ the maximal ideal. If \mathcal{M} is a coherent \mathcal{O}_X -module then for any $n \geq 1$ the $\mathcal{O}_{X,x}$ -module $\mathcal{M}_x/\mathfrak{m}_x^n \mathcal{M}_x$ can be thought of as a quasi-coherent sheaf, constant on the closed set $\overline{\{x\}}$. According to [Be] there is a unique collection of functors $\mathbb{A}(T, -) : \text{QCoh } \mathcal{O}_X \rightarrow \text{Ab}$, indexed by subsets $T \subset S(X)_q$, each of which commuting with direct limits, and satisfying

$$\mathbb{A}(T, \mathcal{M}) = \begin{cases} \prod_{(x) \in T} \lim_{\leftarrow n} \mathcal{M}_x/\mathfrak{m}_x^n \mathcal{M}_x & \text{if } q = 0 \\ \prod_{x \in X} \lim_{\leftarrow n} \mathbb{A}(\hat{x}T, \mathcal{M}_x/\mathfrak{m}_x^n \mathcal{M}_x) & \text{if } q > 0 \end{cases} \quad (1.1)$$

for \mathcal{M} coherent. Furthermore each $\mathbb{A}(T, -)$ is exact.

For a single chain ξ one also writes $\mathcal{M}_\xi := \mathbb{A}(\{\xi\}, \mathcal{M})$, and this is the *Beilinson completion* of \mathcal{M} along ξ . Then

$$\mathbb{A}(T, \mathcal{M}) \subset \prod_{\xi \in T} \mathcal{M}_\xi. \quad (1.2)$$

In view of this we shall say that $\mathbb{A}(T, \mathcal{M})$ is the group of adeles combinatorially supported on T and with values in \mathcal{M} .

Observe that for $q = 0$ and \mathcal{M} coherent we have $\mathcal{M}_{(x)} = \widehat{\mathcal{M}}_x$, the \mathfrak{m}_x -adic completion, and (1.2) is an equality.

Define a presheaf $\underline{\mathbb{A}}(T, \mathcal{M})$ by

$$\Gamma(U, \underline{\mathbb{A}}(T, \mathcal{M})) := \mathbb{A}(T \cap S(U)_q, \mathcal{M}) \quad (1.3)$$

for $U \subset X$ open. Then $\underline{\mathbb{A}}(T, \mathcal{M})$ is in fact a flasque sheaf. Also $\underline{\mathbb{A}}(T, \mathcal{O}_X)$ is a flat \mathcal{O}_X -algebra, and $\underline{\mathbb{A}}(T, \mathcal{M}) \cong \underline{\mathbb{A}}(T, \mathcal{O}_X) \otimes_{\mathcal{O}_X} \mathcal{M}$. Given a local section $m \in \underline{\mathbb{A}}(T, \mathcal{M})$ we shall often use the inclusion (1.2) to write $m = (m_\xi)$ where ξ runs over T and $m_\xi \in \mathcal{M}_\xi$.

Let $S(X)_q^{\text{red}}$ be the set of reduced chains (i.e. without repeated points), and define

$$\underline{\mathbb{A}}_{\text{red}}^q(\mathcal{M}) := \underline{\mathbb{A}}(S(X)_q^{\text{red}}, \mathcal{M}).$$

For $0 \leq i \leq q$ the i th face map ∂_i , which omits the point x_i from a chain (x_0, \dots, x_q) , induces a homomorphism

$$\partial^i : \underline{\mathbb{A}}_{\text{red}}^{q-1}(\mathcal{M}) \rightarrow \underline{\mathbb{A}}_{\text{red}}^q(\mathcal{M}).$$

Then $\underline{\mathbb{A}}_{\text{red}}^\bullet(\mathcal{M})$ is a complex with coboundary operator $\partial := \sum (-1)^i \partial^i$, and $\mathcal{M} \rightarrow \underline{\mathbb{A}}_{\text{red}}^\bullet(\mathcal{M})$ is a quasi-isomorphism. Since $\underline{\mathbb{A}}_{\text{red}}^\bullet(\mathcal{M})$ is a complex of flasque sheaves we get

$$H^q \Gamma(X, \underline{\mathbb{A}}_{\text{red}}^\bullet(\mathcal{M})) = H^q(X, \mathcal{M}).$$

The complex $\underline{\mathbb{A}}_{\text{red}}^\bullet(\mathcal{O}_X)$ is a DGA, with the Alexander-Whitney product. For local sections $a \in \underline{\mathbb{A}}_{\text{red}}^q(\mathcal{O}_X)$ and $b \in \underline{\mathbb{A}}_{\text{red}}^{q'}(\mathcal{O}_X)$ the product is

$$a \cdot b := \partial^-(a) \cdot \partial^+(b) \in \underline{\mathbb{A}}_{\text{red}}^{q+q'}(\mathcal{O}_X),$$

where ∂^- and ∂^+ correspond respectively to the initial and final segments of $(0, \dots, q, \dots, q+q')$. This algebra is not (graded) commutative. For proofs and more details see [Hr], [Ye1] Chapter 3 and [HY1] Section 1.

Example 1.4. Suppose X is a nonsingular curve. The relation to the classical ring of adeles $\mathbb{A}(X)$ of Chevalley and Weil associated to X is

$$\mathbb{A}(X) = \Gamma(X, \underline{\mathbb{A}}_{\text{red}}^1(\mathcal{O}_X)).$$

From now we assume X is a finite type scheme over a perfect field \mathbb{k} .

Next let us recall the construction of the residue complex \mathcal{K}_X in [Ye3]. It starts with the theory of Beilinson completion algebras (BCAs) developed in [Ye2]. A BCA A is a semilocal \mathbb{k} -algebra with a topology and with valuations on its residue fields. Each local factor of A is a quotient of $K((s_1, \dots, s_n))[[t_1, \dots, t_m]]$ where K is some finitely generated extension field of \mathbb{k} , and $K((s_1, \dots, s_n)) = K((s_n)) \cdots ((s_1))$ is the field of iterated Laurent series. One considers two kinds of homomorphisms between BCAs: morphisms $f : A \rightarrow B$ and intensifications $u : A \rightarrow \widehat{A}$.

Each BCA A has a dual module $\mathcal{K}(A)$, which is functorial w.r.t. these homomorphisms; namely there are maps $\text{Tr}_f : \mathcal{K}(B) \rightarrow \mathcal{K}(A)$ and $q_u : \mathcal{K}(A) \rightarrow \mathcal{K}(\widehat{A})$. Algebraically $\mathcal{K}(A)$ is an injective hull of A/\mathfrak{t} , where \mathfrak{t} is the Jacobson radical.

Example 1.5. Let $A := \mathbb{k}[[s_1, s_2]]$, $B := \mathbb{k}(s_1, s_2)$ and $\widehat{B} := \mathbb{k}((s_1, s_2))$. The inclusions $f : A \rightarrow \widehat{B}$ and $u : B \rightarrow \widehat{B}$ are respectively a morphism and an intensification. The dual modules are

$$\mathcal{K}(A) = \text{Hom}_{\mathbb{k}}^{\text{cont}}(A, \mathbb{k}),$$

$$\mathcal{K}(B) = \Omega_{B/\mathbb{k}}^2 = B \cdot ds_1 \wedge ds_2$$

and

$$\mathcal{K}(\widehat{B}) = \Omega_{\widehat{B}/\mathbb{k}}^{2, \text{sep}} = \widehat{B} \cdot ds_1 \wedge ds_2.$$

The homomorphism $q_u : \mathcal{K}(B) \rightarrow \mathcal{K}(\widehat{B})$ is the inclusion. For a form $\beta \in \Omega_{\widehat{B}/\mathbb{k}}^{2, \text{sep}}$ the functional $\text{Tr}_f(\beta) \in \mathcal{K}(A)$ is described as follows. Given an element $a \in A$ write

$$a\beta = \sum_{i,j} \lambda_{i,j} s_1^i s_2^j \cdot ds_1 \wedge ds_2 \in \Omega_{\widehat{B}/\mathbb{k}}^{2, \text{sep}}$$

with $\lambda_{i,j} \in \mathbb{k}$. Then

$$\text{Tr}_f(\beta)(a) = \lambda_{-1, -1},$$

namely the residue of $a\beta$.

Suppose $\xi = (x, \dots, y)$ is a saturated chain of points in X (i.e. each point is an immediate specialization of the previous one). Then the Beilinson completion $\mathcal{O}_{X, \xi}$ is a BCA. The natural algebra homomorphisms $\partial^- : \mathcal{O}_{X, (x)} \rightarrow \mathcal{O}_{X, \xi}$ and $\partial^+ : \mathcal{O}_{X, (y)} \rightarrow \mathcal{O}_{X, \xi}$ are an intensification and a morphism, respectively. So there are homomorphisms on dual modules

$$q_{\partial^-} : \mathcal{K}(\mathcal{O}_{X, (x)}) \rightarrow \mathcal{K}(\mathcal{O}_{X, \xi})$$

and

$$\mathrm{Tr}_{\partial^+} : \mathcal{K}(\mathcal{O}_{X,\xi}) \rightarrow \mathcal{K}(\mathcal{O}_{X,(y)}).$$

The composition

$$\mathrm{Tr}_{\partial^+} \circ \mathrm{q}_{\partial^-} : \mathcal{K}(\mathcal{O}_{X,(x)}) \rightarrow \mathcal{K}(\mathcal{O}_{X,(y)})$$

is denoted by δ_ξ . We regard $\mathcal{K}_X(x) := \mathcal{K}(\mathcal{O}_{X,(x)})$ as a quasi-coherent \mathcal{O}_X -module, constant on the closed set $\overline{\{x\}}$. Define

$$\mathcal{K}_X^{-q} := \bigoplus_{\dim \overline{\{x\}}=q} \mathcal{K}_X(x) \quad (1.6)$$

and

$$\delta := (-1)^{q+1} \sum_{(x,y)} \delta_{(x,y)} : \mathcal{K}_X^{-q} \rightarrow \mathcal{K}_X^{-q+1}. \quad (1.7)$$

Then the pair (\mathcal{K}_X, δ) is the residue complex of X . That is to say, in the notation of the introduction, there is a canonical isomorphism $\mathcal{K}_X \cong \pi^! \mathbb{k}$ in the derived category $\mathrm{D}(\mathrm{Mod} \mathcal{O}_X)$ (see [Ye3] Corollary 2.5).

Let x be a point of dimension q in X , and consider a local section $\phi_x \in \mathcal{K}_X(x) \subset \mathcal{K}_X^{-q}$. Let $\xi = (x_0, \dots, x_{q'})$ be any chain of length q' in X , and let $a_\xi \in \mathcal{O}_{X,\xi}$. Define an element $\phi_x \cdot a_\xi \in \mathcal{K}_X^{-q+q'}$ as follows. If $x = x_0$ and ξ is saturated then there are homomorphisms $\mathrm{q}_{\partial^-} : \mathcal{K}_X(x) \rightarrow \mathcal{K}(\mathcal{O}_{X,\xi})$ and $\mathrm{Tr}_{\partial^+} : \mathcal{K}(\mathcal{O}_{X,\xi}) \rightarrow \mathcal{K}_X(x_{q'})$. Since $\mathcal{K}(\mathcal{O}_{X,\xi})$ is an $\mathcal{O}_{X,\xi}$ -module the product $a_\xi \cdot \mathrm{q}_{\partial^-}(\phi_x) \in \mathcal{K}(\mathcal{O}_{X,\xi})$ exists, and we set

$$\phi_x \cdot a_\xi := \mathrm{Tr}_{\partial^+}(a_\xi \cdot \mathrm{q}_{\partial^-}(\phi_x)) \in \mathcal{K}_X(x_{q'}). \quad (1.8)$$

Otherwise we set $\phi_x \cdot a_\xi := 0$.

Remark 1.9. In order to apply the Koszul sign rule we consider the “local” objects $\mathcal{K}_X(x)$ and $\mathcal{O}_{X,\xi}$ as ungraded; whereas the “global” objects \mathcal{K}_X and $\underline{\mathbb{A}}_{\mathrm{red}}(\mathcal{O}_X)$ are considered as graded. Ungraded local elements shall be decorated with suitable subscripts, such as $a_\xi \in \mathcal{O}_{X,\xi}$ or $\phi_x \in \mathcal{K}_X(x)$.

Suppose $\xi = (x_0, \dots, x_p)$ and $\eta = (y_0, \dots, y_q)$ are chains such that $y_0 \in \overline{\{x_p\}}$. Then we denote by $\xi \vee \eta$ the concatenated chain $(x_0, \dots, x_p, y_0, \dots, y_q)$.

Lemma 1.10. *Let $\xi = (x, \dots, y)$ and $\eta = (y, \dots, z)$ be saturated chains in X and let $\phi_x \in \mathcal{K}_X(x)$, $a_\xi \in \mathcal{O}_{X,\xi}$ and $b_\eta \in \mathcal{O}_{X,\eta}$ be some elements. Writing $\xi \vee \partial_0(\eta) := (x, \dots, y, \dots, z)$, one has $a_\xi \cdot b_\eta \in \mathcal{O}_{X,\xi \vee \partial_0(\eta)}$. Then*

$$(\phi_x \cdot a_\xi) \cdot b_\eta = \phi_x \cdot (a_\xi \cdot b_\eta) \in \mathcal{K}_X(z).$$

Proof. Consider the commutative diagram with intensifications u, v, w, u' and morphisms f, f', g, h :

$$\begin{array}{ccccc}
\mathcal{O}_{X,(x)} & \xrightarrow{w} & \mathcal{O}_{X,\xi} & \xleftarrow{f} & \mathcal{O}_{X,(y)} \\
& \searrow v & \downarrow u' & & \downarrow u \\
& & \mathcal{O}_{X,\xi \vee \partial_0(\eta)} & \xleftarrow{f'} & \mathcal{O}_{X,\eta} \\
& & & \swarrow h & \uparrow g \\
& & & & \mathcal{O}_{X,(z)}
\end{array}$$

According to [Ye3] Lemma 1.12 there is an isomorphism

$$\mathcal{O}_{X,\xi \vee \partial_0(\eta)} \cong \mathcal{O}_{X,\xi} \otimes_{\mathcal{O}_{X,(y)}}^{(\wedge)} \mathcal{O}_{X,\eta}$$

(intensification base change), so by [Ye2] Thm. 7.4 (ii) one has

$$\mathrm{Tr}_{f'} \circ \mathrm{q}_{u'} = \mathrm{q}_u \circ \mathrm{Tr}_f.$$

Now the BCA $\mathcal{O}_{X,\xi \vee \partial_0(\eta)}$ is commutative, $\mathrm{Tr}_{f'}$ is $\mathcal{O}_{X,\eta}$ -linear and $\mathrm{q}_{u'}$ is $\mathcal{O}_{X,\xi}$ -linear. Hence

$$\begin{aligned}
\phi_x \cdot (a_\xi \cdot b_\eta) &= \mathrm{Tr}_h((a_\xi \cdot b_\eta) \cdot \mathrm{q}_v(\phi_x)) \\
&= \mathrm{Tr}_g(\mathrm{Tr}_{f'}(f'(b_\eta) \cdot u'(a_\xi) \cdot \mathrm{q}_{u'}(\mathrm{q}_w(\phi_x)))) \\
&= \mathrm{Tr}_g(b_\eta \cdot \mathrm{Tr}_{f'}(\mathrm{q}_{u'}(a_\xi \cdot \mathrm{q}_w(\phi_x)))) \\
&= \mathrm{Tr}_g(b_\eta \cdot \mathrm{q}_u(\mathrm{Tr}_f(a_\xi \cdot \mathrm{q}_w(\phi_x)))) \\
&= (\phi_x \cdot a_\xi) \cdot b_\eta.
\end{aligned}$$

□

Suppose $x \in X$ is a point and $b \in \mathcal{O}_{X,(x)}$ is some element. Using the inclusion $\mathcal{O}_{X,(x)} \subset \underline{\mathbb{A}}_{\mathrm{red}}^0(\mathcal{O}_X)$ we consider b as an adèle.

Lemma 1.11 (Approximation). *Let $U \subset X$ be an affine open set, $x \in U$ a point and $q \geq 1$. Define*

$$T := \{\xi \in S(U)_q^{\mathrm{red}} \mid \xi = (x, \dots)\}.$$

Let $a \in \mathbb{A}(T, \mathcal{O}_X)$ be some adèle.

- (1) *Given an integer $n \geq 1$ there exists adèles $b \in \mathcal{O}_{X,(x)}$ and $c = (c_\eta) \in \mathbb{A}(\partial_0(T), \mathcal{O}_X)$ such that*

$$a - \partial^1(b) \cdot c \in \mathfrak{m}_x^n \cdot \mathbb{A}(T, \mathcal{O}_X).$$

- (2) Given an element $\phi_x \in \mathcal{K}_X(x)$ there is an integer $n \geq 1$ such that $\mathfrak{m}_x^n \cdot \phi_x = 0$. For such n the adeles b and c from part (1) satisfy

$$\phi_x \cdot a_\xi = \phi_x \cdot (b_{(x,y)} \cdot c_\eta) \in \mathcal{K}_X(z)$$

for all $\xi \in T$, where we write $\eta = (y, \dots, z) := \partial_0(\xi)$, and $b_{(x,y)}$ is the (x, y) component of $\partial^1(b)$.

Proof. (1) There is a ring homomorphism $\mathcal{O}_{X,(x)} \rightarrow \mathbb{A}(T, \mathcal{O}_X)$ and via this homomorphism we obtain an ideal $\mathfrak{m}_x^n \cdot \mathbb{A}(T, \mathcal{O}_X)$. Because the functor $\mathbb{A}(T, -)$ is exact we have

$$\frac{\mathbb{A}(T, \mathcal{O}_X)}{\mathfrak{m}_x^n \cdot \mathbb{A}(T, \mathcal{O}_X)} \cong \mathbb{A}(T, \mathcal{O}_{X,x}/\mathfrak{m}_x^n).$$

By the definition of adeles with values in a quasi-coherent sheaf there is an isomorphism

$$\mathbb{A}(T, \mathcal{O}_{X,x}/\mathfrak{m}_x^n) \cong \lim_{\rightarrow} \mathbb{A}(T, \mathcal{L})$$

where the limit runs over the coherent \mathcal{O}_U -submodules $\mathcal{L} \subset \mathcal{O}_{X,x}/\mathfrak{m}_x^n$. We note that $\partial_0(T) = \hat{x}T$, so for \mathcal{L} large enough (i.e. $\mathcal{L}_x = \mathcal{O}_{X,x}/\mathfrak{m}_x^n$)

$$\begin{aligned} \mathbb{A}(T, \mathcal{L}) &= \lim_{\leftarrow i} \mathbb{A}(\partial_0(T), \mathcal{L}_x/\mathfrak{m}_x^i \mathcal{L}_x) \\ &\cong \mathbb{A}(\partial_0(T), \mathcal{O}_{X,x}/\mathfrak{m}_x^n). \end{aligned}$$

The conclusion is that

$$\mathbb{A}(T, \mathcal{O}_{X,x}/\mathfrak{m}_x^n) \cong \mathbb{A}(\partial_0(T), \mathcal{O}_{X,x}/\mathfrak{m}_x^n).$$

Again we go to coherent subsheaves. Write $C := \Gamma(U, \mathcal{O}_X)$ and $\mathcal{O}_Z := \text{Im}(\mathcal{O}_X \rightarrow \mathcal{O}_{X,x}/\mathfrak{m}_x^n)$. Then

$$\begin{aligned} \mathbb{A}(\partial_0(T), \mathcal{O}_{X,x}/\mathfrak{m}_x^n) &= \lim_{\rightarrow} \mathbb{A}(\partial_0(T), \mathcal{L}) \\ &\cong (\mathcal{O}_{X,x}/\mathfrak{m}_x^n) \otimes_C \mathbb{A}(\partial_0(T), \mathcal{O}_Z) \\ &\cong (\mathcal{O}_{X,x}/\mathfrak{m}_x^n) \otimes_C \mathbb{A}(\partial_0(T), \mathcal{O}_X). \end{aligned}$$

So there is a ring surjection

$$\mathcal{O}_{X,x} \otimes_C \mathbb{A}(\partial_0(T), \mathcal{O}_X) \rightarrow \mathbb{A}(T, \mathcal{O}_{X,x}/\mathfrak{m}_x^n). \quad (1.12)$$

Consider the image \bar{a} of the adele a in $\mathbb{A}(T, \mathcal{O}_{X,x}/\mathfrak{m}_x^n)$. Using the surjection (1.12) we can write $\bar{a} = \sum_{i=1}^r b_i \otimes c_i$ with $b_i \in \mathcal{O}_{X,x}$ and $c_i \in \mathbb{A}(\partial_0(T), \mathcal{O}_X)$. By bringing the b_i to a common denominator we can assume $r = 1$.

- (2) Because $\mathcal{K}(\mathcal{O}_{X,(x)})$ is an \mathfrak{m}_x -torsion module the element ϕ_x is annihilated by some power \mathfrak{m}_x^n . Pick adeles b and c as in part (1). Then for any chain $\xi = (x) \vee \eta = (x, y, \dots) \in T$ we obtain

$$a_\xi - b_{(x,y)} \cdot c_\eta \in \mathfrak{m}_x^n \cdot \mathcal{O}_{X,\xi}.$$

Therefore

$$\phi_x \cdot a_\xi = \phi_x \cdot (b_{(x,y)} \cdot c_\eta).$$

□

Lemma 1.13. *Let $U \subset X$ be an open set, $x \in U$ a point, $\phi_x \in \mathcal{K}_X(x)$ an element and $a = (a_\xi) \in \mathbb{A}(S(U)_q^{\text{red}}, \mathcal{O}_X)$ an adele. Then for all but finitely many chains $\xi \in S(U)_q^{\text{red}}$ one has $\phi_x \cdot a_\xi = 0$.*

Proof. The proof is by induction on q . For $q = 0$ there is nothing to prove, so take $q \geq 1$. Because U is covered by finitely many affine open sets, and we are only interested in establishing finiteness, we might as well assume U itself is affine. Moreover by the definition of the product we can neglect those chains in the combinatorial support of a that do not begin with x . Thus we can assume $a \in \mathbb{A}(T, \mathcal{O}_X)$, where T is the set defined in Lemma 1.11. According to this lemma we can find adeles $b \in \mathcal{O}_{X,(x)}$ and $c = (c_\eta) \in \mathbb{A}(\partial_0(T), \mathcal{O}_X)$ such that

$$\phi_x \cdot a_\xi = \phi_x \cdot (b_{(x,y)} \cdot c_\eta) \in \mathcal{K}_X(z)$$

for all $\xi = (x) \vee \eta = (x, y, \dots, z) \in T$.

The only way to get a nonzero product $\phi_x \cdot a_\xi$ is when ξ is a saturated chain. Consider such a chain ξ . According to Lemma 1.10 we have

$$\phi_x \cdot (b_{(x,y)} \cdot c_\eta) = (\phi_x \cdot b_{(x,y)}) \cdot c_\eta.$$

For any point y occurring one has

$$\phi_x \cdot b_{(x,y)} = \delta_{(x,y)}(b\phi_x) \in \mathcal{K}_X(y).$$

It follows that the product $\psi_y := \phi_x \cdot b_{(x,y)}$ vanishes for all but finitely many points y . Fixing y , the induction hypothesis applied to the element $\psi_y \in \mathcal{K}_X(y)$ says that $\psi_y \cdot c_\eta = 0$ for all but finitely many chains $\eta \in S(U)_{q-1}^{\text{red}}$. □

Theorem 1.14. \mathcal{K}_X is a right DG $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$ -module, with product

$$\phi \cdot a = \sum_{x,\xi} \phi_x \cdot a_\xi$$

for local sections $\phi = \sum_x \phi_x \in \mathcal{K}_X$ and $a = (a_\xi) \in \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$.

Proof. According to Lemmas 1.13 and 1.10 this is a well defined associative product. It remains to verify that

$$\delta(\phi \cdot a) = \delta(\phi) \cdot a + (-1)^q \phi \cdot \partial(a) \tag{1.15}$$

for $\phi \in \mathcal{K}_X^{-q}$ and $a \in \underline{\mathbb{A}}_{\text{red}}^{q'}(\mathcal{O}_X)$. We may assume $\phi = \phi_x$ and $a = a_\xi$ for some point x of dimension q and a chain $\xi = (x_0, \dots, x_{q'})$.

Now there are only 3 ways to get any nonzero term in equation (1.15):
(i) ξ is saturated and $x_0 = x$; (ii) ξ is saturated and (x, x_0) is saturated; or
(iii) $x_0 = x$, and for some index $0 \leq i < q'$ and some point $y \in X$, the chain

$$\eta := (x_0, \dots, x_i, y, x_{i+1}, \dots, x_{q'})$$

is saturated.

In case (i), $\delta(\phi) \cdot a = 0$ and

$$\begin{aligned} \phi \cdot \partial(a) &= \phi_x \cdot (-1)^{q'+1} \partial^{q'+1}(a_\xi) \\ &= (-1)^{q'+1} \sum_y \delta_{(x_{q'}, y)}(\phi_x \cdot a_\xi) \\ &= (-1)^q \delta(\phi \cdot a). \end{aligned}$$

In case (ii), $\phi \cdot a = 0$ and

$$\begin{aligned} \phi \cdot \partial(a) &= \phi_x \cdot \partial^0(a)_{(x) \vee \xi} \\ &= \delta_{(x, x_0)}(\phi_x) \cdot a_\xi \\ &= (-1)^{q+1} \delta(\phi) \cdot a. \end{aligned}$$

In this equation $(x) \vee \xi$ is the concatenated chain $(x, x_0, \dots, x_{q'})$.

Finally in case (iii), $\delta(\phi) \cdot a = 0$, $\delta(\phi \cdot a) = 0$, and it remains to show that also $\phi \cdot \partial(a) = 0$. We note that

$$\phi \cdot \partial(a) = \sum_y \phi_x \cdot (-1)^{i+1} \partial^{i+1}(a)_\eta$$

where y runs over the points such that (x_i, y, x_{i+1}) is a saturated chain, and η is as above (and depends on y).

For any index $0 \leq j \leq q'$ let us write $\xi_j := (x_j, \dots, x_{q'})$. We shall use an approximation trick of Lemma 1.11 to define recursively elements $\phi_{x_j} \in \mathcal{K}_X(x_j)$ and $b_{(x_j)} \in \mathcal{O}_{X, (x_j)}$ for $0 \leq j \leq i$, and $a_{\xi_j} \in \mathcal{O}_{X, \xi_j}$ for $0 \leq j \leq i+1$. Pick some affine open set $U \subset X$ containing $x_{q'}$ (U does not play an essential role, since our computation is local anyhow; but it appears in Lemma 1.11). For $j = 0$ we note that $x_0 = x$ and $\xi_0 = \xi$, and the elements ϕ_{x_0} and a_{ξ_0} are already defined. Now suppose $j \leq i$ and ϕ_{x_j} and a_{ξ_j} have been defined. By Lemma 1.11 we can find elements $b_{(x_j)} \in \mathcal{O}_{X, (x_j)}$ and $a_{\xi_{j+1}} \in \mathcal{O}_{X, \xi_{j+1}}$ such that

$$\phi_{x_j} \cdot a_{\xi_j} = \phi_{x_j} \cdot (\partial^1(b_{(x_j)})_{(x_j, x_{j+1})} \cdot a_{\xi_{j+1}}) \in \mathcal{K}(x_{q'}).$$

If $j < i$ we also define

$$\phi_{x_{j+1}} := \phi_{x_j} \cdot \partial^1(b_{(x_j)})_{(x_j, x_{j+1})} \in \mathcal{K}(x_{j+1}).$$

We thus have for any of the points y under consideration:

$$\phi_{x_j} \cdot \partial^{(i+1)-j}(a_{\xi_j})_{\eta_j} = \phi_{x_{j+1}} \cdot \partial^{(i+1)-(j+1)}(a_{\xi_{j+1}})_{\eta_{j+1}},$$

where

$$\eta_j := (x_j, \dots, x_i, y, x_{i+1}, \dots, x_{q'}).$$

Putting it all together we obtain

$$\begin{aligned} \phi_{x_0} \cdot \partial^{i+1}(a_{\xi_0})_{\eta_0} &= \phi_{x_i} \cdot \partial^1(a_{\xi_i})_{\eta_i} \\ &= (\phi_{x_i} \cdot b_{(x_i)}) \cdot (\partial^0 \circ \partial^0)(a_{\xi_{i+1}})_{\eta_i} \\ &= \delta_{(x_i, y, x_{i+1})}(\phi_{x_i} \cdot b_{(x_i)}) \cdot a_{\xi_{i+1}}, \end{aligned}$$

and hence

$$\phi \cdot \partial(a) = (-1)^{i+1} \sum_y \delta_{(x_i, y, x_{i+1})}(\phi_{x_i} \cdot b_{(x_i)}) \cdot a_{\xi_{i+1}}.$$

But according to [Ye3] Lemma 2.15(3), which is a variant of the Parshin-Lomadze Residue Theorem, we have

$$\sum_y \delta_{(x_i, y, x_{i+1})}(\phi_{x_i} \cdot b_{(x_i)}) = 0.$$

□

Example 1.16. The global section $a := \partial^1(1) \in \Gamma(X, \underline{\mathbb{A}}_{\text{red}}^1(\mathcal{O}_X))$ acts on \mathcal{K}_X^q like $(-1)^{q+1}\delta$, namely for any $\phi \in \mathcal{K}_X^q$ we have

$$\phi \cdot a = \phi \cdot \partial^1(1) = (-1)^{q+1}\delta(\phi).$$

Example 1.17. If X is a nonsingular curve then the theorem takes on a very simple form. Here $\mathcal{K}_X^{-1} = \Omega_{K/\mathbb{k}}^1$, where K is the function field of X ; and $\mathcal{K}_X^0 = \text{Coker}(\Omega_{X/\mathbb{k}}^1 \rightarrow \Omega_{K/\mathbb{k}}^1)$. And there is actually an isomorphism of complexes

$$\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X) \cong \mathcal{H}om_{\mathcal{O}_X}(\mathcal{K}_X, \mathcal{K}_X)$$

(cf. Example 1.4).

If X is integral of dimension n , let ω_X be the coherent sheaf $\mathbf{H}^{-n}\mathcal{K}_X$. This is a subsheaf of the constant sheaf $\mathcal{K}_X^{-n} = \Omega_{K/\mathbb{k}}^n$, where K is the function field of X . In [Li] and [KW] ω_X is called the sheaf of regular differential forms (cf. [Ye1] Theorem 4.4.16). Since \mathcal{K}_X is a complex of injectives and $\omega_X \rightarrow \underline{\mathbb{A}}_{\text{red}}(\omega_X)$ is a quasi-isomorphism, there is a map of complexes $\underline{\mathbb{A}}_{\text{red}}(\omega_X) \rightarrow \mathcal{K}_X[-n]$ inducing the identity in \mathbf{H}^0 . Lipman asked for an explicit formula for such a homomorphism of complexes. Producing such a formula was the main result of [HY1]. The following corollary gives essentially the same formula but in terms of DG modules.

Corollary 1.18. *Suppose X is integral of dimension n . Then the map*

$$\underline{\mathbb{A}}_{\text{red}}(\omega_X) \cong \omega_X \otimes_{\mathcal{O}_X} \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X) \rightarrow \mathcal{K}_X[-n],$$

sending $\alpha \otimes a \mapsto \alpha \cdot a$, is a homomorphism of complexes, which induces the identity on $\omega_X = H^0 \underline{\mathbb{A}}_{\text{red}}(\omega_X) = H^0 \mathcal{K}_X[-n]$.

Proof. In $\omega_X \otimes_{\mathcal{O}_X} \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$ one has $\partial(\alpha \otimes a) = \alpha \otimes \partial(a)$. On the other hand in $\mathcal{K}_X[-n]$ the section α has degree 0, and $\delta(\alpha) = 0$, so

$$\delta(\alpha \cdot a) = \delta(\alpha) \cdot a + \alpha \cdot \partial(a) = \alpha \cdot \partial(a).$$

□

Given a morphism $f : X \rightarrow Y$ there is a natural DGA homomorphism $f^* : \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_Y) \rightarrow f_* \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$. There is also a map of graded \mathcal{O}_Y -modules $\text{Tr}_f : f_* \mathcal{K}_X \rightarrow \mathcal{K}_Y$, and this is a map of complexes if f is proper (see [Ye3] Definition 2.11 and Theorem 3.4).

Theorem 1.19. *If $f : X \rightarrow Y$ is proper then $\text{Tr}_f : f_* \mathcal{K}_X \rightarrow \mathcal{K}_Y$ is a homomorphism of DG $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_Y)$ -modules.*

Proof. We have to show that $\text{Tr}_f(\phi \cdot a) = \text{Tr}_f(\phi) \cdot a$ for local sections $\phi \in \mathcal{K}_X$ and $a \in \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_Y)$. For this we may as well assume $\phi = \phi_{x_0} \in \mathcal{K}_X(x_0)$ and $a = a_\eta \in \mathcal{O}_{Y,\eta}$ for a saturated chain $\eta = (y_0, \dots, y_q)$ in Y and a point x_0 which is closed in the fiber $f^{-1}(y_0)$. According to [Ye3] Proposition 2.1 we have an isomorphism of BCAs

$$\prod_{\xi} \mathcal{O}_{X,\xi} \cong \mathcal{O}_{X,(x_0)} \otimes_{\mathcal{O}_{Y,(y_0)}}^{(\wedge)} \mathcal{O}_{Y,\eta}$$

where $\xi = (x_0, \dots, x_q)$ runs over the finitely many chains in X satisfying $f(x_i) = y_i$. Therefore by [Ye2] Theorem 7.4 the left square in the diagram

$$\begin{array}{ccccc} \mathcal{K}(\mathcal{O}_{X,(x_0)}) & \xrightarrow{\mathfrak{q}} & \bigoplus_{\xi} \mathcal{K}(\mathcal{O}_{X,\xi}) & \xrightarrow{\text{Tr}} & \bigoplus_{x_q} \mathcal{K}(\mathcal{O}_{X,(x_q)}) \\ \text{Tr} \downarrow & & \text{Tr} \downarrow & & \text{Tr} \downarrow \\ \mathcal{K}(\mathcal{O}_{Y,(y_0)}) & \xrightarrow{\mathfrak{q}} & \mathcal{K}(\mathcal{O}_{Y,\eta}) & \xrightarrow{\text{Tr}} & \mathcal{K}(\mathcal{O}_{Y,(y_q)}) \end{array}$$

is commutative. The functoriality of Tr with respect to morphisms of BCAs implies the commutativity of the right square. □

Question 1.20. It is plausible to assume that an action as in Theorem 1.14 exists even without the explicit construction of the residue complex. Suppose X is a finite dimensional noetherian scheme endowed with a dimension function $d : X \rightarrow \mathbb{Z}$ (e.g. if X has a dualizing complex). Let \mathcal{L} be a quasi-coherent Cousin complex on X , so that $\mathcal{L}^{-q} \cong \bigoplus_{d(x)=q} H_x^{-q} \mathcal{L}$. Is \mathcal{L} a right DG $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$ -module?

2 de Rham Complexes, Residues and Adeles

Let us continue with the setup of Section 1. Consider the algebraic de Rham complex $(\Omega_{X/\mathbb{k}}, d)$ of X . For any integers p, q define

$$\mathcal{A}_X^{p,q} := \underline{\mathbb{A}}_{\text{red}}^q(\Omega_{X/\mathbb{k}}^p).$$

Since $\mathcal{M} \mapsto \underline{\mathbb{A}}_{\text{red}}^q(\mathcal{M})$ is functorial with respect to differential operators $\mathcal{M} \rightarrow \mathcal{N}$ (cf. [HY1]), the sheaves $\mathcal{A}_X^{p,q}$ make up a double complex, with commuting operators d and ∂ . Let

$$\mathcal{A}_X^n := \bigoplus_{p+q=n} \mathcal{A}_X^{p,q}$$

with coboundary operators

$$\begin{aligned} D' &:= d : \mathcal{A}_X^{p,q} \rightarrow \mathcal{A}_X^{p+1,q}, \\ D'' &:= (-1)^p \partial : \mathcal{A}_X^{p,q} \rightarrow \mathcal{A}_X^{p,q+1} \end{aligned}$$

and $D := D' + D''$. The Alexander-Whitney product $\mathcal{A}_X^{p,q} \otimes \mathcal{A}_X^{p',q'} \rightarrow \mathcal{A}_X^{p+p',q+q'}$ makes (\mathcal{A}_X, D) into a DGA. To be explicit, write $\mathcal{A}_X^{p,q} \cong \Omega_{X/\mathbb{k}}^p \otimes_{\mathcal{O}_X} \underline{\mathbb{A}}_{\text{red}}^q(\mathcal{O}_X)$. Then taking local sections $a \in \underline{\mathbb{A}}_{\text{red}}^q(\mathcal{O}_X)$, $b \in \underline{\mathbb{A}}_{\text{red}}^{q'}(\mathcal{O}_X)$, $\alpha \in \Omega_{X/\mathbb{k}}^p$ and $\beta \in \Omega_{X/\mathbb{k}}^{p'}$, one has

$$(\alpha \otimes a) \cdot (\beta \otimes b) = (-1)^{qp'} \alpha \wedge \beta \otimes \partial^-(a) \cdot \partial^+(b) \in \mathcal{A}_X^{p+p',q+q'}$$

and

$$\begin{aligned} D(\alpha \otimes a) &= D(\alpha) \cdot a + (-1)^p \alpha \cdot D(a) \\ &= d(\alpha) \cdot a + (-1)^p \alpha \cdot d(a) + (-1)^p \alpha \cdot \partial(a). \end{aligned}$$

Each $\mathcal{A}_X^{p,q}$ is a flasque sheaf, and the natural homomorphism of DGAs $\Omega_{X/\mathbb{k}} \rightarrow \mathcal{A}_X$ is a quasi-isomorphism.

In [Ye3] it is proved that any differential operator $D : \mathcal{M} \rightarrow \mathcal{N}$ between \mathcal{O}_X -modules induces a dual operator

$$\text{Dual}(D) : \text{Dual}\mathcal{N} \rightarrow \text{Dual}\mathcal{M},$$

where by definition $\text{Dual}\mathcal{M}$ is the complex $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{M}, \mathcal{K}_X)$. The operator $\text{Dual}(D)$ is defined locally, in terms of Beilinson completion algebras. Let

$$\mathcal{F}_X^{p,q} := \mathcal{H}om_X(\Omega_{X/\mathbb{k}}^{-p}, \mathcal{K}_X^q).$$

We see that \mathcal{F}_X^\bullet is a double complex, with two commuting operators δ and $\text{Dual}(d)$. Set

$$\begin{aligned}\mathcal{F}_X^n &:= \bigoplus_{p+q=n} \mathcal{F}_X^{p,q}, \\ D' &:= (-1)^{p+q+1} \text{Dual}(d) : \mathcal{F}_X^{p,q} \rightarrow \mathcal{F}_X^{p+1,q}, \\ D'' &:= \delta : \mathcal{F}_X^{p,q} \rightarrow \mathcal{F}_X^{p,q+1} \quad \text{and} \\ D &:= D' + D'' : \mathcal{F}_X^n \rightarrow \mathcal{F}_X^{n+1}.\end{aligned}\tag{2.1}$$

Proof of Theorem 0.2. The left action of $\Omega_{X/\mathbb{k}}^\bullet$ on itself makes \mathcal{F}_X^\bullet into a graded right $\Omega_{X/\mathbb{k}}^\bullet$ -module. By Theorem 1.14, \mathcal{F}_X^\bullet is a graded right $\underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$ -module. The formula for the product $\phi \cdot a$ which makes the signs correct is

$$(\phi \cdot a)(\beta) := (-1)^{pq'} \phi(\beta) \cdot a$$

for local sections $\phi \in \mathcal{F}_X^{p,q}$, $a \in \underline{\mathbb{A}}_{\text{red}}^{q'}(\mathcal{O}_X)$ and $\beta \in \Omega_{X/\mathbb{k}}^{-p}$. Since $\mathcal{A}_X \cong \Omega_{X/\mathbb{k}}^\bullet \otimes_{\mathcal{O}_X} \underline{\mathbb{A}}_{\text{red}}(\mathcal{O}_X)$ as graded algebras, we obtain a structure of graded right \mathcal{A}_X -module on \mathcal{F}_X^\bullet .

It remains to check the coboundaries, which we break up into four steps, calculating $D'(\phi \cdot a)$, $D'(\phi \cdot \alpha)$, $D''(\phi \cdot a)$ and $D''(\phi \cdot \alpha)$ separately, with a, ϕ be as above, and $\alpha \in \Omega_{X/\mathbb{k}}^{p'}$.

Since $D''(\alpha) = 0$ we get

$$\begin{aligned}D''(\phi \cdot \alpha)(\beta) &= \delta((\phi \cdot \alpha)(\beta)) = \delta(\phi(\alpha \wedge \beta)) \\ &= D''(\phi)(\alpha \wedge \beta) = (D''(\phi) \cdot \alpha)(\beta) \\ &= (D''(\phi) \cdot \alpha + (-1)^{p+q} \phi \cdot D''(\alpha))(\beta)\end{aligned}$$

for every $\beta \in \Omega_{X/\mathbb{k}}^{-p}$.

Next, by Thm. 1.14

$$\begin{aligned}(D''(\phi \cdot a))(\beta) &= \delta((\phi \cdot a)(\beta)) = \delta((-1)^{pq'} \phi(\beta) \cdot a) \\ &= (-1)^{pq'} D''(\phi(\beta) \cdot a) \\ &= (-1)^{pq'} (D''(\phi(\beta)) \cdot a + (-1)^q \phi(\beta) \cdot D''(a)) \\ &= (D''(\phi) \cdot a + (-1)^{p+q} \phi \cdot D''(a))(\beta).\end{aligned}$$

This takes care of D'' .

As for D' , by [Ye3] Proposition 5.2 we have

$$D'(\phi \cdot \alpha) = D'(\phi) \cdot \alpha + (-1)^{p+q} \phi \cdot D'(\alpha).\tag{2.2}$$

Finally we will prove that

$$D'(\phi \cdot a) = D'(\phi) \cdot a + (-1)^{p+q} \phi \cdot D'(a)$$

by induction on q' . As before we may assume

$$\phi = \phi_x \in \mathcal{H}om_{\mathcal{O}_X}(\Omega_{X/\mathbb{k}}^{-p}, \mathcal{K}_X(x)) \subset \mathcal{F}_X^{p,q}$$

for some point $x = x_0$, and

$$a = a_\xi \in \mathcal{O}_{X,\xi} \subset \underline{\mathbb{A}}_{\text{red}}^{q'}(\mathcal{O}_X)$$

for some saturated chain $\xi = (x_0, \dots, x_{q'})$. Choose an integer n such that $\mathfrak{m}_x^n \cdot \phi_x = 0$. Then we also have $\mathfrak{m}_x^{n+1} \cdot D'(\phi_x) = 0$.

For $q' = 0$ we have $a_{(x)} \in \mathcal{O}_{X,(x)}$. Choose some local section $b \in \mathcal{O}_X$ near x such that $b \equiv a_{(x)} \pmod{\mathfrak{m}_x^{n+1}}$. So by Lemma 1.11 one has $\phi \cdot a = \phi \cdot b$, $D'(\phi) \cdot a = D'(\phi) \cdot b$ and $\phi \cdot D'(a) = \phi \cdot D'(b)$. But by equation (2.2), with $\alpha = b \in \Omega_{X/\mathbb{k}}^0$, we know that

$$D'(\phi \cdot b) = D'(\phi) \cdot b + (-1)^{p+q} \phi \cdot D'(b). \quad (2.3)$$

Now let us handle the case $q' = 1$, i.e. $\xi = (x_0, x_1)$. By Lemma 1.11 there exist adeles $b \in \mathcal{O}_{X,(x_0)}$ and $c \in \mathcal{O}_{X,(x_1)}$ such that

$$a_\xi = b_{(x_0, x_1)} \cdot c_{(x_1)} \in \mathcal{O}_{X,\xi} / \mathfrak{m}_{x_0}^{n+1} \mathcal{O}_{X,\xi},$$

where $b_{(x_0, x_1)}$ is the (x_0, x_1) component of $\partial^1(b)$. Let $d := 1_{(x_0, x_1)} \in \mathcal{O}_{X,(x_0, x_1)} \subset \underline{\mathbb{A}}_{\text{red}}^1(\mathcal{O}_X)$. We then get $\partial^1(b) = b \cdot d$,

$$a - b \cdot d \cdot c \in \mathfrak{m}_{x_0}^{n+1} \mathcal{O}_{X,\xi},$$

$\phi \cdot a = \phi \cdot (b \cdot d \cdot c)$, $D'(\phi) \cdot a = D'(\phi) \cdot (b \cdot d \cdot c)$ and $\phi \cdot D'(a) = \phi \cdot D'(b \cdot d \cdot c)$. By the $q' = 0$ case we know that equation (2.3) holds, and also that

$$D'((\phi \cdot b \cdot d) \cdot c) = D'(\phi \cdot b \cdot d) \cdot c + (-1)^{p+q+1} (\phi \cdot b \cdot d) \cdot D'(c).$$

It remains to verify that

$$D'(\psi \cdot d) = D'(\psi) \cdot d + (-1)^{p+q} \psi \cdot D'(d),$$

where we define $\psi := \phi \cdot b$. But for every $\beta \in \Omega_{X/\mathbb{k}}^{-p}$ we have

$$(\psi \cdot d)(\beta) = (-1)^p \psi(\beta) \cdot d = (-1)^p \delta_{(x_0, x_1)}(\psi(\beta)) = (-1)^{p+q+1} D''(\psi)(\beta).$$

Therefore $\psi \cdot d = (-1)^{p+q+1} D''(\psi)$. Since $D' \circ D'' = -D'' \circ D'$ and $D'(d) = 0$ we arrive at

$$\begin{aligned} D'(\psi \cdot d) &= (-1)^{p+q+1} (D' \circ D'')(\psi) = (-1)^{p+q} (D'' \circ D')(\psi) \\ &= (-1)^{p+q} D''(D'(\psi)) = (-1)^{(p+q)+(p+q+2)} D'(\psi) \cdot d \\ &= D'(\psi) \cdot d + (-1)^{p+q} \psi \cdot D'(d). \end{aligned}$$

To conclude we consider $q' \geq 2$. Since we are working locally, we can assume by the approximation trick (Lemma 1.11) that $a = b \cdot c$ with $b \in \underline{\mathbb{A}}_{\text{red}}^1(\mathcal{O}_X)$ and $c \in \underline{\mathbb{A}}_{\text{red}}^{q'-1}(\mathcal{O}_X)$. The induction hypothesis applies to b and c , and we have

$$\begin{aligned}
D'(\phi \cdot a) &= D'(\phi \cdot b \cdot c) \\
&= D'(\phi \cdot b) \cdot c + (-1)^{p+q+1} \phi \cdot b \cdot D'(c) \\
&= D'(\phi) \cdot b \cdot c + (-1)^{p+q} \phi \cdot D'(b) \cdot c + (-1)^{p+q+1} \phi \cdot b \cdot D'(c) \\
&= D'(\phi) \cdot b \cdot c + (-1)^{p+q} \phi \cdot D'(b \cdot c) \\
&= D'(\phi) \cdot a + (-1)^{p+q} \phi \cdot D'(a).
\end{aligned}$$

□

Proof of Corollary 0.3. By [Ye3] Proposition 5.4 the trace homomorphisms $\text{Tr} : \mathcal{K}_{X_n} \rightarrow \mathcal{K}_{X_{n+1}}$ give rise to DG $\Omega_{X_{n+1}/\mathbb{k}}$ -linear homomorphisms $\text{Tr} : \mathcal{F}_{X_n} \rightarrow \mathcal{F}_{X_{n+1}}$. Therefore $\mathcal{F}_{\mathfrak{X}}$ is a right DG $\mathcal{A}_{\mathfrak{X}}$ -module.

Let $\widehat{\Omega}_{\mathfrak{X}/\mathbb{k}}$ be the formal completion of $\Omega_{Y/\mathbb{k}}$ along X . According to [HY1] the DGA quasi-isomorphism $\widehat{\Omega}_{\mathfrak{X}/\mathbb{k}} \rightarrow \mathcal{A}_{\mathfrak{X}}$ is a flasque resolution, so

$$H^i(X, \widehat{\Omega}_{\mathfrak{X}/\mathbb{k}}) = H^i\Gamma(X, \mathcal{A}_{\mathfrak{X}}).$$

By [Ye3] Proposition 6.8 we get $\mathcal{F}_{\mathfrak{X}} \cong \text{R}\Gamma_X \Omega_{Y/\mathbb{k}}[2m]$, where $m := \dim Y$, and hence

$$H_X^{2m-i}(Y, \Omega_{Y/\mathbb{k}}) = H^{-i}\Gamma(X, \mathcal{F}_{\mathfrak{X}}),$$

compatible with the cap product. But when $\text{char } \mathbb{k} = 0$ one has by definition $H_{\text{DR}}^i(X) := H^i(X, \widehat{\Omega}_{\mathfrak{X}/\mathbb{k}})$ and $H_i^{\text{DR}}(X) := H_X^{2m-i}(Y, \Omega_{Y/\mathbb{k}})$, see [Ha]. □

Remark 2.4. The role of the residue complex as a canonical flasque resolution of $\Omega_{X/\mathbb{k}}$ (for X smooth) appears already in [Ha] and in [EZ]. In order to define the operator dual to d these authors use the fact that $\mathcal{F}_X \cong \Omega_{X/\mathbb{k}} \otimes_{\mathcal{O}_X} \mathcal{K}_X$ is the Cousin resolution of $\Omega_{X/\mathbb{k}}$ in the category of abelian sheaves on X . This should be compared to our approach (in [Ye3] and here) where the dual operator $\text{Dual}(d)$ is defined in terms of BCAs, i.e. by algebraic-analytic methods.

Remark 2.5. Suppose X is smooth irreducible of dimension n . One can regard the sheaf $\mathcal{A}_X^{p,q}$ as an analog of the Dolbeault sheaf of smooth (p, q) forms on a complex manifold (cf. [GH] Ch. 0). The adelic resolution

$$0 \rightarrow \Omega_{X/\mathbb{k}}^p \rightarrow \mathcal{A}_X^{p,0} \xrightarrow{D''} \mathcal{A}_X^{p,1} \rightarrow \cdots \rightarrow \mathcal{A}_X^{p,n} \rightarrow 0$$

corresponds to the $\bar{\partial}$ resolution on the manifold. Any section $\phi \in \Gamma(X, \mathcal{F}_X^{-p, -q})$ determines a functional on $\Gamma(X, \mathcal{A}_X^{p, q})$, namely $\alpha \mapsto \text{Tr}_{X/\mathbb{k}}(\phi \cdot \alpha) \in \mathbb{k}$. Here

$$\text{Tr}_{X/\mathbb{k}} : \Gamma(X, \mathcal{F}_X^{0, 0}) = \Gamma(X, \mathcal{K}_X^0) \rightarrow \mathbb{k}$$

is the “sum of residues” (cf. [Ye3] Def. 1.16). In this way we can think of $\mathcal{F}_X^{-p, -q}$ as an analog of the sheaf of (p, q) currents on a manifold (cf. [GH] Ch. 3).

Remark 2.6. Suppose X is smooth irreducible and $\text{char } \mathbb{k} = 0$. Let $Z \subset X$ be an irreducible closed subset of codimension d , and let

$$0 \rightarrow \mathcal{E}_l \rightarrow \cdots \rightarrow \mathcal{E}_1 \rightarrow \mathcal{E}_0 \rightarrow \mathcal{O}_Z \rightarrow 0$$

be a finite locally free resolution. Using the adelic Chern-Weil theory of [HY2] it is possible to construct a Chern character form $\text{ch}(\mathcal{E}; \nabla) \in \mathcal{A}_X$, depending on adelic connections ∇_i on the \mathcal{E}_i , whose component $\text{ch}(\mathcal{E}; \nabla)_{2d} \in \mathcal{A}_X^{2d}$ satisfies

$$C_X \cdot \text{ch}(\mathcal{E}; \nabla)_{2d} = C_Z \in \mathcal{F}_X.$$

Here $C_X \in \mathcal{F}_X^{-2n}$ and $C_Z \in \mathcal{F}_X^{-2(n-d)}$ are the fundamental classes. This generalizes [HY2] Theorem 6.5.

References

- [AJL] L. Alonso, A. Jeremías and J. Lipman, Duality and flat base change on formal schemes, in “Studies in Duality on Noetherian Formal Schemes and Non-Noetherian Ordinary Schemes,” Contemp. Math. **244**, Amer. Math. Soc., 1999, 3-90.
- [AK] A. Altman and S. Kleiman, “Introduction to Grothendieck Duality,” Lecture Notes in Math. **20**, Springer, 1970.
- [Be] A.A. Beilinson, Residues and adeles, Funkt. Anal. Pril. **14** (1980) no. 1, 44-45; English trans. in Func. Anal. Appl. **14** (1980) no. 1, 34-35.
- [Co] B. Conrad, “Grothendieck Duality and Base Change,” Lecture Notes in Math. **1750**, Springer, 2000.
- [EGA] A. Grothendieck and J. Dieudonné, “Éléments de Géométrie Algébrique”; [EGA] I: Springer-Verlag, Berlin, 1971; [EGA] II-IV: Publ. Math. IHES nos. **8**, **11**, **17**, **20**, **24**, **28**, **32**.
- [EZ] F. El Zein, Complexe Dualisant et Applications à la Classe Fondamentale d’un Cycle, Bull. Soc. Math. France, Mémoire **58**, 1978.

- [Fu] W. Fulton, “Intersection Theory”, Springer-Verlag, Berlin, 1984.
- [GH] P. Griffiths and J. Harris, “Principles of Algebraic Geometry”, Wiley, New York, 1978.
- [Gr] A. Grothendieck, On the de Rham cohomology of algebraic varieties, Publ. Math. IHES **29** (1966), 95-103.
- [Ha] R. Hartshorne, On the de Rham Cohomology of algebraic varieties, Publ. Math. IHES **45** (1975), 5-99.
- [HK] R. Hübl and E. Kunz, Regular differential forms and duality for projective morphisms, J. reine angew. Math. **410** (1990), 84-108.
- [Hr] A. Huber, On the Parshin-Beilinson Adeles for Schemes, Abh. Math. Sem. Univ. Hamburg **61** (1991), 249-273.
- [HY1] R. Hübl and A. Yekutieli, Adeles and differential forms, J. reine angew. Math. **471** (1996), 1-22.
- [HY2] R. Hübl and A. Yekutieli, Adelic Chern forms and applications, Amer. J. Math. **121** (1999), 797-839.
- [Kl] S.L. Kleiman, Relative duality for quasi-coherent sheaves, Compositio Math. **41** (1980), 39-60.
- [KS] M. Kashiwara and P. Schapira, “Sheaves on Manifolds”, Springer-Verlag, Berlin, 1990.
- [KW] E. Kunz and R. Waldi, “Regular Differential Forms,” Contemp. Math. **79** (1988), AMS, Providence.
- [Li] J. Lipman, Dualizing sheaves, differentials and residues on algebraic varieties, Asterisque **117** (1984).
- [LS] J. Lipman and P. Sastry, Regular differentials and equidimensional scheme-maps, J. Alg. Geom. **1** (1992), 101-130.
- [ML] S. Mac Lane, “Homology,” Springer-Verlag, Berlin, 1995 (reprint of 1975 edition).
- [RD] R. Hartshorne, “Residues and Duality”, Lecture Notes in Math. 20, Springer-Verlag, Berlin, 1966.
- [Ne] A. Neeman, The Grothendieck duality theorem via Bousfield’s techniques and Brown representability, Jour. Amer. Math. Soc. **9** (1996), 205-236.
- [SY] P. Sastry and A. Yekutieli, On residue complexes, dualizing sheaves and local cohomology modules, Israel J. Math. **90** (1995), 325-348.

- [Ye1] A. Yekutieli, “An Explicit Construction of the Grothendieck Residue Complex” (with an appendix by P. Sastry), *Astérisque* **208** (1992).
- [Ye2] A. Yekutieli, Traces and differential operators over Beilinson completion algebras, *Compositio Math.* **99** (1995), 59-97.
- [Ye3] A. Yekutieli, Residues and differential operators on schemes, *Duke Math. J.* **95** (1998), 305-341.
- [Ye4] A. Yekutieli, Smooth formal embeddings and the residue complex, *Canadian J. Math.* **50** (1998), 863-896.