

SPACES OF COINVARIANTS AND FUSION PRODUCT I. FROM EQUIVALENCE THEOREM TO KOSTKA POLYNOMIALS

B. FEIGIN, M. JIMBO, R. KEDEM, S. LOKTEV, T. MIWA

ABSTRACT. The fusion rule gives the dimensions of spaces of conformal blocks in the WZW theory. We prove a dimension formula similar to the fusion rule for spaces of coinvariants of affine Lie algebras $\widehat{\mathfrak{g}}$. An equivalence of filtered spaces is established between spaces of coinvariants of two objects: highest weight $\widehat{\mathfrak{g}}$ -modules and tensor products of finite-dimensional evaluation representations of $\mathfrak{g} \otimes \mathbb{C}[t]$.

In the $\widehat{\mathfrak{sl}}_2$ case we prove that their associated graded spaces are isomorphic to the spaces of coinvariants of fusion products, and that their Hilbert polynomials are the level-restricted Kostka polynomials.

1. INTRODUCTION

First recall the notion of conformal blocks in the Wess-Zumino-Witten conformal field theory. Let \mathfrak{g} be a simple Lie algebra over \mathbb{C} , let $\tilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$ be the loop algebra, and $\tilde{\mathfrak{g}}^{\{0\}}$ its completion with respect to power series in t . As a geometric part of the data consider a compact complex curve C with chosen distinct points p_0, \dots, p_N . We take these points to be regular, and choose local coordinates t_0, \dots, t_N . By $\mathfrak{g}^{C \setminus \{p_0, \dots, p_N\}}$ denote the Lie algebra of \mathfrak{g} -valued meromorphic functions on C with possible poles at p_i . Local coordinates provide us with the inclusion

$$(1.1) \quad \mathfrak{g}^{C \setminus \{p_0, \dots, p_N\}} \rightarrow \tilde{\mathfrak{g}}^{\{0\}} \oplus \dots \oplus \tilde{\mathfrak{g}}^{\{0\}} \oplus \mathbb{C}K$$

sending a function into its Laurent expansion at each point. Here $\mathbb{C}K$ is the one-dimensional center. The central extension in the right hand side is given by the sum of canonical cocycles of each summand $\tilde{\mathfrak{g}}^{\{0\}}$. It is trivial on the image of $\mathfrak{g}^{C \setminus \{p_0, \dots, p_N\}}$.

Let L_0, \dots, L_N be irreducible highest weight representations of a fixed level k of $\widehat{\mathfrak{g}}^{\{0\}} = \tilde{\mathfrak{g}}^{\{0\}} \oplus \mathbb{C}K \oplus \mathbb{C}d$. Let v_i be their highest weight vectors.¹

Define the space of conformal blocks by

$$(1.2) \quad \langle L_0, \dots, L_N \rangle_{p_0, \dots, p_N}^C = L_0 \boxtimes \dots \boxtimes L_N / \mathfrak{g}^{C \setminus \{p_0, \dots, p_N\}}(L_0 \boxtimes \dots \boxtimes L_N).$$

Date: November 3, 2018.

¹In the main text, we will consider reducible highest weight representations with cyclic vectors v_i . In this introduction, we restrict to irreducible representations in order to ease the presentation.

In this paper, we call it the space of conformal coinvariants. Here we use the symbol \boxtimes in place of the tensor product symbol \otimes in order to emphasize that we consider the action of the Lie algebra (1.1).

If L_i are integrable, then it was proved by [TUY] that these spaces are finite dimensional, and that their dimensions are given by the fusion rule, in particular, it depends only on representations and the genus of the curve (see Theorem 2.7 for the fusion rule when the curve is \mathbb{CP}^1).

Introduce a filtration on the space of conformal coinvariants in the following way. Let v_1, \dots, v_N be the highest weight vectors of representations L_1, \dots, L_N , respectively. Then we have a map from L_0 to the space of conformal coinvariants given by

$$(1.3) \quad L_0 \simeq L_0 \otimes v_1 \otimes \dots \otimes v_N \rightarrow \langle L_0, \dots, L_N \rangle_{p_0, \dots, p_N}^C.$$

By a standard argument (see Subsection 2.4) one can show that the map (1.3) is surjective. This gives us a realization of the space of conformal coinvariants as a quotient of L_0 . Since the representation L_0 is graded by degree of t_0^{-1} , our space inherits the corresponding increasing filtration.

In this paper, we study this filtration for rational C and irreducible highest weight representations L_0, \dots, L_N of level $k \in \mathbb{Z}_{>0}$.

From this point we restrict to the case of rational C . We take $p_0 = \infty$, and $p_i = z_i \in \mathbb{C}$ ($1 \leq i \leq N$). We use the local coordinates $t_0 = 1/t$ and $t_i = t - z_i$. We take L_0 to be the level k irreducible highest weight module whose degree zero subspace is the irreducible \mathfrak{g} -module π_λ with highest weight λ . We denote this L_0 specifically by $L_\lambda^{(k)}(\infty)$. Similarly, we take highest weights λ_i for L_i ($1 \leq i \leq N$).

The space $L_\lambda^{(k)}(\infty)$ is generated from the highest weight vector v_0 by the action of $\mathfrak{g}[t] = \mathfrak{g} \otimes \mathbb{C}[t]$. Consider the mapping

$$\Delta_{\mathcal{Z}} = (\phi_{z_1}^{-1} \otimes \dots \otimes \phi_{z_N}^{-1}) \circ \Delta^{(N-1)} : U(\mathfrak{g}[t]) \rightarrow U(\mathfrak{g}[t]) \otimes \dots \otimes U(\mathfrak{g}[t]),$$

where ϕ_z is the automorphism of $U(\mathfrak{g}[t])$ changing t to $t - z$. Here \mathcal{Z} denotes the set of points $\{z_1, \dots, z_N\}$.

For $x \in \mathfrak{g}[t]$ and $v^{(i)} \in L_i$, we have

$$(1.4) \quad (xv^{(0)}) \otimes v^{(1)} \otimes \dots \otimes v^{(N)} = v^{(0)} \otimes \Delta_{\mathcal{Z}}(S(x))(v^{(1)} \otimes \dots \otimes v^{(N)})$$

in the quotient space (1.2). Here S is the antipode.

Let $A \subset U(\mathfrak{g}[t]) \otimes \dots \otimes U(\mathfrak{g}[t])$ be the left ideal which annihilates $v_1 \otimes \dots \otimes v_N$. The kernel of the map (1.3) contains $\Delta_{\mathcal{Z}}^{-1}(S(A))L_\lambda^{(k)}(\infty)$. In fact, we will show that they are equal (see Proposition 2.19).

In order to prove this statement, we introduce the notion of fusion right ideal $X_1 \otimes \dots \otimes X_N(\mathcal{Z}) = \Delta_{\mathcal{Z}}^{-1}(B)$ for right ideals $X_1, \dots, X_N \subset U(\mathfrak{g}[t])$ by replacing $S(A)$ with

$$B = \sum_{i=1}^N U(\mathfrak{g}[t])^{\otimes(i-1)} \otimes X_i \otimes U(\mathfrak{g}[t])^{\otimes(N-i)}.$$

It is known that the dimension of the space of conformal blocks is given in terms of the multiplication rule in the Verlinde algebra (see Section 2.3 for the definition of the Verlinde algebra). In Theorem 2.9, we give a similar multiplicative formula for the dimension of the space of coinvariants

$$(1.5) \quad L_\lambda^{(k)}(\infty)/(X_1 \otimes \dots \otimes X_N(\mathbb{Z}))L_\lambda^{(k)}(\infty)$$

under a certain condition on the right ideals X_i . In particular, this gives a proof of the equality of the kernel of (1.3) with the subspace $(X_1 \otimes \dots \otimes X_N(\mathbb{Z}))L_\lambda^{(k)}(\infty)$ by taking $X_i = S(\mathfrak{X}'_{\lambda_i}) + B_1$, where the left ideal $\mathfrak{X}'_{\lambda_i} \subset U(\mathfrak{g})$ is defined by $U(\mathfrak{g})/\mathfrak{X}'_{\lambda_i} \simeq \pi_{\lambda_i}$ and $B_1 = (\mathfrak{g} \otimes t\mathbb{C}[t])U(\mathfrak{g}[t])$. Therefore, the filtration of the conformal coinvariants (1.2) can be studied in the form (1.5) as the coinvariants with respect to the fusion right ideal.

Let us consider another version of this construction. By $\pi(L_i)$ denote the finite-dimensional representation of \mathfrak{g} generated by the highest weight vector v_i . Then one can easily deduce from (1.3) that the following map is also surjective (see Subsection 2.4)

$$(1.6) \quad v_0 \otimes \pi(L_1) \otimes \dots \otimes \pi(L_N) \rightarrow \langle L_0, \dots, L_N \rangle_{p_0, \dots, p_N}^C.$$

Let $I_\lambda^{(k)} \subset \mathfrak{g} \otimes \mathbb{C}[t]$ be the left ideal which annihilates v_0 . Then a similar argument shows that we have a surjective map

$$\pi(L_1) \otimes \dots \otimes \pi(L_N)/S(I_\lambda^{(k)}) \rightarrow \langle L_0, \dots, L_N \rangle_{p_0, \dots, p_N}^C.$$

We will show also that these spaces are in fact isomorphic (see Example 1 in Subsection 3.2).

To describe the filtration on the space of conformal coinvariants in this setting, we use the notion of filtered tensor product introduced in [FL]. It is the tensor product of cyclic \mathfrak{g} -modules V_1, \dots, V_N considered as the evaluation $\mathfrak{g}[t]$ -modules with distinct parameters z_i . Upon choosing cyclic vectors $v_i \in V_i$ we can define a \mathfrak{g} -equivariant filtration. Namely, the vector $v_1 \otimes \dots \otimes v_N$ is cyclic in the tensor product with respect to the action of $\mathfrak{g}[t]$, so the whole representation is a quotient of $U(\mathfrak{g}[t])$. We have a grading of $\mathfrak{g}[t]$ and therefore of $U(\mathfrak{g}[t])$ by the degree in t , so the quotient inherits the corresponding filtration.

In our case we should choose the highest vectors of $\pi(L_i)$ as cyclic to obtain the same filtration as given by (1.5). The associated graded space of the filtered tensor product [FL]

$$V_1 * \dots * V_N(\mathbb{Z}) = \text{gr } \mathcal{F}_{\mathbb{Z}}(V_1, \dots, V_N)$$

is called in [FL] the fusion product.

For $\mathfrak{g} = \mathfrak{sl}_2$ we prove that the Hilbert polynomial of this graded space is independent of the choice of points and coincides with the level-restricted Kostka polynomial (for definition see [SS]). This calculation uses the fusion product. The result of [FF] enables us to give an upper estimate of the Hilbert polynomial for the quotient of the filtered tensor product. The

resulting formula coincides with the known fermionic formula for the level-restricted Kostka polynomial. Since at $q = 1$, the level-restricted Kostka polynomial gives the dimension of the conformal blocks, which is given by the fusion rule, we can conclude that the fermionic formula is exact.

The plan of the paper is as follows. In Section 2, we give the definition of the fusion right ideals in $U(\mathfrak{g}[t])$, and prove the fusion rule for the dimensions of the spaces of coinvariants of integrable highest weight $\widehat{\mathfrak{g}}$ -modules. In Section 3, we establish an equivalence between spaces of coinvariants of irreducible highest weight $\widehat{\mathfrak{g}}$ -modules and those of filtered tensor products of cyclic \mathfrak{g} -modules. We give a brief review on the Kostka polynomials, and propose the definition of fusion Kostka polynomials. In Section 4, we prove that in the case of $\mathfrak{g} = \widehat{\mathfrak{sl}}_2$, the associated graded space of the space of conformal coinvariants is independent of the choice of insertion points, and that its Hilbert polynomial is the level-restricted Kostka polynomial. In Appendix we discuss behavior of fusion ideals and spaces of coinvariants when points collide.

2. FORMULA FOR THE DIMENSION

In this section we introduce the notion of fusion right ideals and consider spaces of coinvariants with respect to them. We give a formula for the dimension of the latter in terms of the Verlinde algebra (see Theorem 2.9).

2.1. Fusion of ideals. Let \mathfrak{g} be a complex Lie algebra. By $\mathfrak{g}[t]$ denote the Lie algebra $\mathfrak{g} \otimes \mathbb{C}[t]$. When there is no fear of confusion, we will identify \mathfrak{g} with $\mathfrak{g} \otimes t^0 \subset \mathfrak{g}[t]$. For $z \in \mathbb{C}$ we denote by ϕ_z the automorphism of $\mathfrak{g}[t]$ sending $x \otimes t^n$ into $x \otimes (t - z)^n$ ($x \in \mathfrak{g}$).

Let $\mathcal{Z} = (z_1, \dots, z_N)$ be a set of complex numbers. Throughout this paper we assume that

$$(2.1) \quad z_i \neq z_j \quad (i \neq j).$$

Consider the map $\mathfrak{g}[t] \rightarrow \bigoplus^N \mathfrak{g}[t]$ which sends $x \in \mathfrak{g}[t]$ to $\phi_{z_1}^{-1}(x) \oplus \dots \oplus \phi_{z_N}^{-1}(x)$. By the universal property it induces the map $U(\mathfrak{g}[t]) \rightarrow U(\mathfrak{g}[t])^{\otimes N}$ given by

$$\Delta_{\mathcal{Z}} = (\phi_{z_1} \otimes \dots \otimes \phi_{z_N})^{-1} \circ \Delta^{(N-1)},$$

where the iterated coproduct is defined by $\Delta^{(n-1)} = (\Delta \otimes \text{id}) \circ \Delta^{(n-2)}$, $\Delta^{(1)} = \Delta$.

Definition 1. Let X_1, \dots, X_N be right ideals in $U(\mathfrak{g}[t])$. Define the fusion right ideal by

$$X_1 \otimes \dots \otimes X_N(\mathcal{Z}) = (\Delta_{\mathcal{Z}})^{-1} \left(\sum_{i=1}^N U(\mathfrak{g}[t])^{\otimes(i-1)} \otimes X_i \otimes U(\mathfrak{g}[t])^{\otimes(N-i)} \right).$$

Throughout this paper, for simplicity of notation, we denote as

$$I^{(N)}(X_1, \dots, X_N) = \sum_{i=1}^N U(\mathfrak{g}[t])^{\otimes(i-1)} \otimes X_i \otimes U(\mathfrak{g}[t])^{\otimes(N-i)}.$$

The following Proposition validates this definition.

Proposition 2.1. *If the right ideals $X_i = \mathfrak{a}_i U(\mathfrak{g}[t])$ are generated by Lie subalgebras $\mathfrak{a}_i \subset U(\mathfrak{g}[t])$, then the fusion right ideal is given by*

$$X_1 \otimes \dots \otimes X_N(\mathcal{Z}) = \left(\bigcap_{i=1}^N \phi_{z_i}(\mathfrak{a}_i) \right) U(\mathfrak{g}[t]).$$

As an example, consider the two-sided ideal of $U(\mathfrak{g}[t])$ consisting of elements which have M -fold zeros at the origin,

$$(2.2) \quad B_M = (\mathfrak{g} \otimes t^M \mathbb{C}[t]) U(\mathfrak{g}[t]).$$

Let

$$(2.3) \quad B_{M,\mathcal{Z}} = (\mathfrak{g} \otimes (t - z_1)^M \dots (t - z_N)^M \mathbb{C}[t]) U(\mathfrak{g}[t])$$

be the two-sided ideal consisting of elements which have M -fold zeros at the points in \mathcal{Z} . Then we have

$$B_M \otimes \dots \otimes B_M(\mathcal{Z}) = B_{M,\mathcal{Z}}.$$

For the proof of Proposition 2.1 we use the following fact, which is a corollary of the Poincaré-Birkhoff-Witt (PBW) theorem.

Lemma 2.2. *Let \mathfrak{a} be a Lie algebra, and let $\mathfrak{b} \subset \mathfrak{a}$ be its Lie subalgebra. Take a basis of \mathfrak{b} , $\{B_i\}_{i \in I}$, and extend it to a basis of \mathfrak{a} by adjoining $\{A_j\}_{j \in J}$. We assume that the index set I and J are ordered. Then the set of elements in $U(\mathfrak{a})$*

$$(2.4) \quad B_{i_1} \dots B_{i_m} A_{j_1} \dots A_{j_n} \quad (i_1 \geq \dots \geq i_m; j_1 \geq \dots \geq j_n)$$

with $m \geq 1, n \geq 0$ forms a basis of $\mathfrak{b}U(\mathfrak{a})$, and the one with $m = 0, n \geq 0$ forms a basis of $U(\mathfrak{a})/\mathfrak{b}U(\mathfrak{a})$.

Lemma 2.3. *Let V_i be a vector space with countable basis, and let W_i be a subspace of V_i , $i = 1, \dots, N$. Then the kernel of the map*

$$V_1 \otimes \dots \otimes V_N \rightarrow (V_1/W_1) \otimes \dots \otimes (V_N/W_N)$$

is equal to

$$\sum_{i=1}^N V_1 \otimes \dots \otimes V_{i-1} \otimes W_i \otimes V_{i+1} \otimes \dots \otimes V_N.$$

Proof. Choosing a filtration

$$V_i = \bigcup_{a=0}^{\infty} V_{i,a}, \quad V_{i,a} \subset V_{i+1,a}$$

such that $\dim V_{i,a}$ is finite, one can reduce the proof to the case where all V_i are finite-dimensional. Then the statement follows by dimension counting. \square

Proposition 2.1 is a consequence of the following fact.

Proposition 2.4. *Let \mathfrak{a} be a Lie algebra with a countable basis, and let $\mathfrak{a}_1, \dots, \mathfrak{a}_N$ be its Lie subalgebras. Then,*

$$\left(\Delta^{(N-1)}\right)^{-1} \left(\sum_{i=1}^N U^{\otimes(i-1)} \otimes \mathfrak{a}_i U \otimes U^{\otimes(N-i)} \right) = (\mathfrak{a}_1 \cap \dots \cap \mathfrak{a}_N)U,$$

where $U = U(\mathfrak{a})$.

Proof. The inclusion \supset is clear. Therefore, by using Lemma 2.3, it is enough to show that the mapping

$$p : U/(\mathfrak{a}_1 \cap \dots \cap \mathfrak{a}_N)U \rightarrow (U/\mathfrak{a}_1 U) \otimes \dots \otimes (U/\mathfrak{a}_N U),$$

which is induced from $\Delta^{(N-1)}$, is injective. We prove this statement by induction on N . The case $N = 1$ is obvious. We prove the case $N = 2$. Take a basis $\{A_\alpha\} \sqcup \{B_\beta\} \sqcup \{C_\gamma\} \sqcup \{D_\delta\}$ of \mathfrak{a} such that $\{A_a\}$ is a basis of $\mathfrak{a}_1 \cap \mathfrak{a}_2$, $\{A_a\} \sqcup \{B_b\}$ is a basis of \mathfrak{a}_1 , and $\{A_a\} \sqcup \{C_c\}$ is a basis of \mathfrak{a}_2 . We fix an ordering of each set $\{A_\alpha\}, \{B_\beta\}, \{C_\gamma\}, \{D_\delta\}$. Write $x \in U$ as a linear combination of the PBW basis using the ordering:

$$(2.5) \quad x = \sum c_{\alpha_1, \dots, \delta_d} A_{\alpha_1} \cdots A_{\alpha_a} B_{\beta_1} \cdots B_{\beta_b} C_{\gamma_1} \cdots C_{\gamma_c} D_{\delta_1} \cdots D_{\delta_d}$$

Here and after, we do not bother to mention a necessary ordering for the indices.

Let

$$\pi : U \otimes U \rightarrow (U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U)$$

be the canonical surjection. We want to show that if $\pi \circ \Delta(x) = 0$ then x belongs to $(\mathfrak{a}_1 \cap \mathfrak{a}_2)U$.

Let $F^l U$ be the filtration of U defined as follows.

$$F^0 U = \mathbb{C} \cdot 1, \quad F^{l+1} U = F^l U + \mathfrak{a} F^l U.$$

One can define a filtration of $U \otimes U$ by setting

$$F^l(U \otimes U) = \sum_{l_1 + l_2 = l} F^{l_1} U \otimes F^{l_2} U,$$

and further induce a filtration of $(U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U)$. By Lemma 2.2, the elements of the form $C_{\gamma_1} \cdots C_{\gamma_c} D_{\delta_{i_1}} \cdots D_{\delta_{i_{d'}}}$ (resp., $B_{\beta_1} \cdots B_{\beta_b} D_{\delta_{j_1}} \cdots D_{\delta_{j_{d''}}}$) form a basis of $U/\mathfrak{a}_1 U$ (resp., $U/\mathfrak{a}_2 U$). We have the basis of $(U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U)$ consisting of the tensor product of these bases. Moreover, those which satisfies $c + b + d' + d'' = l$, form a basis of $(F^l/F^{l-1})((U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U))$. Let us denote them by $X_{\gamma_1, \dots, \gamma_c, \delta_{i_1}, \dots, \delta_{i_{d'}}, \beta_1, \dots, \beta_b, \delta_{j_1}, \dots, \delta_{j_{d''}}}$.

Note that $\pi \circ \Delta(1) \neq 0$. Suppose that $x \in F^l U$ ($l \geq 1$) and $\pi \circ \Delta(x) = 0$. We will show that $x \in (\mathfrak{a}_1 \cap \mathfrak{a}_2)U + F^{l-1} U$. Write x as (2.5). If a monomial in this expression is such that $a \geq 1$ then it belongs to $(\mathfrak{a}_1 \cap \mathfrak{a}_2)U$ and the kernel of $\pi \circ \Delta$. Therefore, one can assume that only the terms with $a = 0$ appear in (2.5).

We want to show that

$$(2.6) \quad c_{\beta_1, \dots, \delta_d} = 0$$

if $b + c + d = l$. Consider the 2^l monomials that are obtained by expanding the product $\Delta(B_{\beta_1}) \cdots \Delta(B_{\beta_b}) \Delta(C_{\gamma_1}) \cdots \Delta(C_{\gamma_c}) \Delta(D_{\delta_1}) \cdots \Delta(D_{\delta_d})$. If a monomial is such that either one of B_{β} is sent to the first component of the tensor product $U \otimes U$, or one of C_{γ} to the second, then the image of that monomial by the surjection π belongs to $F^{l-1}((U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U))$. Therefore, $\pi \circ \Delta(B_{\beta_1} \cdots D_{\delta_d}) \in (F^l/F^{l-1})((U/\mathfrak{a}_1 U) \otimes (U/\mathfrak{a}_2 U))$ is the sum of 2^d elements $X_{\gamma_1, \dots, \gamma_c, \delta_{i_1}, \dots, \delta_{i_{d'}}, \beta_1, \dots, \beta_b, \delta_{j_1}, \dots, \delta_{j_{d''}}}$ where $\{i_1, \dots, i_{d'}\} \sqcup \{j_1, \dots, j_{d''}\}$ is a partition of the index set $\{1, \dots, d\}$. Let us denote this sum by $Y_{\beta_1, \dots, \delta_d}$. The elements $Y_{\beta_1, \dots, \delta_d}$ that are obtained from x are linearly independent. Therefore, we have (2.6).

Finally, suppose that the statement is true for $N - 1$ ($N \geq 3$). It means that the mapping

$$U/(\mathfrak{a}_1 \cap \cdots \cap \mathfrak{a}_{N-1})U \rightarrow (U/\mathfrak{a}_1 U) \otimes \cdots \otimes (U/\mathfrak{a}_{N-1} U)$$

is injective. From this follows that

$$p_1 : (U/(\mathfrak{a}_1 \cap \cdots \cap \mathfrak{a}_{N-1})U) \otimes (U/\mathfrak{a}_N U) \rightarrow (U/\mathfrak{a}_1 U) \otimes \cdots \otimes (U/\mathfrak{a}_N U)$$

is also injective. Consider the mapping

$$p_2 : U/(\mathfrak{a}_1 \cap \cdots \cap \mathfrak{a}_N)U \rightarrow (U/(\mathfrak{a}_1 \cap \cdots \cap \mathfrak{a}_{N-1})U) \otimes (U/\mathfrak{a}_N U)$$

induced from Δ . Since $\Delta^{(N-1)} = (\Delta^{(N-2)} \otimes \text{id}) \circ \Delta$, we have $p = p_1 \circ p_2$. Therefore, it is enough to show that p_2 is injective. This is nothing but the case $N = 2$. \square

Note that $\mathfrak{g}[t]$ and therefore $U(\mathfrak{g}[t])$ are graded by the degree in t .

Definition 2. We call a right ideal $X \subset U(\mathfrak{g}[t])$ a congruence ideal if X is graded and $X \supset B_M$ for a sufficiently large M .

Quite generally, let V be a left module over a Lie algebra \mathfrak{a} . For a left ideal $X \subset U(\mathfrak{a})$, we denote by $V^X = \{v \in V \mid xv = 0 \ \forall x \in X\}$ the space of X -invariants. Likewise, for a right ideal $Y \subset U(\mathfrak{a})$, we denote by $V/Y = V/YV$ the space of Y -coinvariants. We have a canonical isomorphism

$$(2.7) \quad (V/Y)^* \simeq (V^*)^{S(Y)},$$

where $V^* = \text{Hom}_{\mathbb{C}}(V, \mathbb{C})$ and $S : U(\mathfrak{a}) \rightarrow U(\mathfrak{a})$ stands for the antipode (the anti-automorphism such that $S(x) = -x$ for $x \in \mathfrak{a}$). We will consider coinvariant spaces with respect to fusion right ideals.

2.2. Affine Lie algebras. In this subsection we fix our notation concerning affine Lie algebras and integrable modules.

Let \mathfrak{g} be a simple Lie algebra. Fix a triangular decomposition $\mathfrak{g} = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$, where $\mathfrak{h} \subset \mathfrak{g}$ is the Cartan subalgebra and \mathfrak{n}_+ (resp. \mathfrak{n}_-) is the nilpotent subalgebra of all positive (resp. negative) root vectors. The invariant bilinear form (\mid) on \mathfrak{g} is normalized as $(\theta|\theta) = 2$, where θ is the maximal root. Let P be the weight lattice and P_+ be the set of dominant integral weights. For a non-negative integer $k \in \mathbb{Z}_{\geq 0}$, we set

$P_+^{(k)} = \{\lambda \in P_+ \mid \langle \lambda, \theta^\vee \rangle \leq k\}$, where θ^\vee denotes the maximal coroot. For $\lambda \in P_+$, we denote by π_λ the irreducible representation with highest weight λ . We will write $\lambda^* = -w_0(\lambda)$ for the highest weight of the dual representation π_λ^* , w_0 being the longest element of the Weyl group W . If $\lambda \in P_+^{(k)}$, then $\lambda^* \in P_+^{(k)}$.

Let $\widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d$ be the affine Lie algebra of non-twisted type. Here K is a central element, $d = td/dt$,

$$[x \otimes f, y \otimes g] = [x, y] \otimes fg + \Omega_0(x \otimes f, y \otimes g)K$$

$$(x, y \in \mathfrak{g}, f, g \in \mathbb{C}[t, t^{-1}]),$$

and the canonical cocycle is given by

$$\Omega_0(x \otimes f, y \otimes g) = (x|y) \operatorname{Res}_{t=0} df \cdot g.$$

Denote by Λ_i the fundamental weights ($0 \leq i \leq r; r = \dim \mathfrak{h}$) and by δ the null root. Let $\Lambda = \lambda + k\Lambda_0 + a\delta$ ($\lambda \in P$, $a \in \mathbb{C}$) be an affine weight of level k . For an element w of the affine Weyl group W_{aff} , we set

$$(2.8) \quad w \circ (\lambda + k\Lambda_0 + a\delta) = w(\lambda, k) + k\Lambda_0 + (a - d_w(\lambda, k))\delta,$$

$$w(\lambda, k) \in P, d_w(\lambda, k) \in \mathbb{Z},$$

where $w \circ \Lambda = w(\Lambda + \rho) - \rho$ stands for the shifted action and $\rho = \sum_{i=0}^r \Lambda_i$.

We say a $\widehat{\mathfrak{g}}$ -module has level k if K acts as k times the identity on it. For $k \in \mathbb{Z}_{\geq 0}$ and $\lambda \in P_+^{(k)}$, let $L_\lambda^{(k)} = U(\widehat{\mathfrak{g}})v_\lambda^{(k)}$ be the unique level k integrable module such that

$$(2.9) \quad (\mathfrak{g} \otimes t^n)v_\lambda^{(k)} = 0 \quad (n > 0),$$

$$(2.10) \quad \mathfrak{n}_+ v_\lambda^{(k)} = 0,$$

$$(2.11) \quad hv_\lambda^{(k)} = \langle \lambda, h \rangle v_\lambda^{(k)} \quad (h \in \mathfrak{h}), \quad dv_\lambda^{(k)} = 0.$$

We also consider modules ‘placed at infinity’. To be precise, define $\widehat{\mathfrak{g}}(\infty) = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}d$ by the same equations as above, except that we change the cocycle to

$$\Omega_\infty(x \otimes f, y \otimes g) = (x|y) \operatorname{Res}_{t=\infty} df \cdot g$$

$$= -\Omega_0(x \otimes f, y \otimes g).$$

Let $L_\lambda^{(k)}(\infty) = U(\widehat{\mathfrak{g}}(\infty))v_\lambda^{(k)}(\infty)$ be the integrable $\widehat{\mathfrak{g}}(\infty)$ -module characterized by (2.9)–(2.11), with $v_\lambda^{(k)}(\infty)$ in place of $v_\lambda^{(k)}$ and (2.9) being changed to

$$(\mathfrak{g} \otimes t^n)v_\lambda^{(k)}(\infty) = 0 \quad (n < 0).$$

The degree zero part of $L_\lambda^{(k)}$ and $L_\lambda^{(k)}(\infty)$ are both isomorphic to π_λ as \mathfrak{g} -module.

Consider the isomorphism of Lie algebra $\iota : \widehat{\mathfrak{g}} \rightarrow \widehat{\mathfrak{g}}(\infty)$ given by $x \otimes t^n \mapsto x \otimes t^n$, $K \mapsto -K$, $d \mapsto d$. By this isomorphism, $L_\lambda^{(k)}(\infty)$ can be considered

as the level $-k$ lowest weight module with the lowest weight $-\lambda$. We have a non-degenerate pairing

$$(2.12) \quad L_\lambda^{(k)}(\infty) \times L_{\lambda^*}^{(k)} \longrightarrow \mathbb{C}$$

such that $\langle \iota(x)u, v \rangle = -\langle u, xv \rangle$ ($x \in \widehat{\mathfrak{g}}, u \in L_\lambda^{(k)}(\infty), v \in L_{\lambda^*}^{(k)}$), which extends the canonical coupling $\pi_\lambda \times \pi_{\lambda^*} \rightarrow \mathbb{C}$.

Proposition 2.5. *Let $X \subset U(\mathfrak{g}[t])$ be a congruence right ideal. Then $L_\lambda^{(k)}(\infty)/X$ is finite-dimensional. We have an isomorphism*

$$(2.13) \quad (L_\lambda^{(k)}(\infty)/X)^* \simeq (L_{\lambda^*}^{(k)})^{S(X)}.$$

Proof. First we prove the finite dimensionality of $L_\lambda^{(k)}(\infty)/X$. Choose a set of nilpotent generators x_1, \dots, x_m of \mathfrak{g} , and let $\tilde{B}_{M,i}$ be the right ideal generated by $x_j \otimes t^n$ ($1 \leq j \leq m, n \geq M$) and $x_1 \otimes t^{M-1}, \dots, x_i \otimes t^{M-1}$. Note that $\tilde{B}_{M,0} = \tilde{B}_{M+1,m}$. We have $\tilde{B}_{M,i} = \tilde{B}_{M,i-1} + (x_i \otimes t^{M-1})U(\mathfrak{g}[t])$. Since the congruence ideal X contains $\tilde{B}_{M,0}$ for a large M , in order to show the finite-dimensionality of $L_\lambda^{(k)}(\infty)/X$, it suffices to prove it for $L_\lambda^{(k)}(\infty)/\tilde{B}_{M,i}$ for all M, i . Since $L_\lambda^{(k)}(\infty)/\tilde{B}_{1,0} \simeq \pi_\lambda$, the statement is valid for $M = 1, i = 0$. The general case can be shown by induction on M, i , by applying the following lemma to $V = L_\lambda^{(k)}(\infty)/\tilde{B}_{M,i-1}$, $x = x_i \otimes t^{M-1}$.

Eq.(2.12) states that $L_{\lambda^*}^{(k)}$ is the restricted dual of $L_\lambda^{(k)}(\infty)$. Since X is graded and $L_\lambda^{(k)}(\infty)/X$ is finite-dimensional, (2.13) follows from (2.7). \square

Lemma 2.6. *Let V be a vector space and $x \in \text{End}(V)$. Assume that x is locally nilpotent, $\bigcap_{n>0} \text{Im } x^n = \{0\}$, and that $V/\text{Im } x$ is finite dimensional. Then V is also finite dimensional.*

Proof. Choose vectors $v_1, \dots, v_n \in V$ which span $V/\text{Im } x$, and let M be such that $x^M v_i = 0$ for all i . Then the set $\{x^j v_i\}$ with $0 \leq j \leq M-1, 1 \leq i \leq n$ spans V . \square

2.3. The multiplicative formula for dimensions. The aim of this section is to prove the multiplicative formula for the dimension of the space of coinvariants with respect to the fusion right ideals. Before discussing it, we recall the original version of the fusion rule for the space of conformal coinvariants.

From now on, we fix a level $k \in \mathbb{Z}_{\geq 0}$. First recall the definition of the Verlinde algebra. Let $\text{Rep } \mathfrak{g} = \bigoplus_{\lambda \in P_+} \mathbb{Z} \cdot [\lambda]$ be the Grothendieck ring of \mathfrak{g} , where $[\lambda]$ stands for the class of irreducible representation π_λ . Let $\mathcal{R}^{(k)}$ denote the ideal spanned by elements of the form

$$[\lambda] - (-1)^{l(w)}[w(\lambda, k)] \quad (w \in W_{\text{aff}}, \lambda, w(\lambda, k) \in P_+),$$

where $l(w)$ signifies the length of w . The Verlinde algebra is the quotient ring $\mathcal{V}^{(k)} = \text{Rep } \mathfrak{g}/\mathcal{R}^{(k)}$. We use the same letter $[\lambda]$ to represent its class

in $\mathcal{V}^{(k)}$. The set $\{[\lambda]\}_{\lambda \in P_+^{(k)}}$ constitutes a \mathbb{Z} -basis of $\mathcal{V}^{(k)}$. For an element $a \in \mathcal{V}^{(k)}$, we write the coefficient of $[\lambda]$ in this basis as $(a : [\lambda])_k$, i.e.,

$$a = \sum_{\lambda \in P_+^{(k)}} (a : [\lambda])_k [\lambda].$$

We now fix a set of distinct points $\mathcal{Z} = \{z_1, \dots, z_N\}$ in \mathbb{C} .

Theorem 2.7. [TUY] *Let L_i ($i = 0, \dots, N$) be the irreducible highest weight representation of highest weight λ_i . Define the space of conformal coinvariants by (1.2). Then, we have the dimension formula.*

$$(2.14) \quad \dim \langle L_0, \dots, L_N \rangle_{\infty, z_1, \dots, z_N}^{\mathbb{C}P^1} = ([\lambda_1^*] \cdots [\lambda_N^*] : [\lambda_0])_k.$$

Now, consider a set of congruence right ideals X_1, \dots, X_N of $U(\mathfrak{g}[t])$.

Lemma 2.8. *The space of coinvariants*

$$(2.15) \quad L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathcal{Z})$$

is finite dimensional.

Proof. A standard deformation argument (e.g. [FKLMM1], Lemma 20) shows that $\dim L_\lambda^{(k)}(\infty)/B_{M, \mathcal{Z}} \leq \dim L_\lambda^{(k)}(\infty)/B_{NM} < \infty$, where we used Proposition 2.5. Since $X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \supset B_{M, \mathcal{Z}}$ for some M , we obtain the assertion. \square

The main result of this section is the following.

Theorem 2.9. *Notation being as above, we have*

$$\begin{aligned} & \sum_{\lambda \in P_+^{(k)}} \dim(L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathcal{Z})) \cdot [\lambda] \\ &= \prod_{i=1}^N \left(\sum_{\lambda \in P_+^{(k)}} \dim(L_\lambda^{(k)}(\infty)/X_i) \cdot [\lambda] \right). \end{aligned}$$

For the proof, we make use of a construction due to [FFu] which we recall below. See also appendix to [FKLMM1], where a detailed account is given in the case $\mathfrak{g} = \mathfrak{sl}_2$.

Set

$$\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z}) = L_\lambda^{(k)}(\infty)/B_{M, \mathcal{Z}}$$

where $B_{M, \mathcal{Z}}$ is defined in (2.3). Since $B_{M, \mathcal{Z}}$ is a two-sided ideal, $\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z})$ is a $U(\mathfrak{g}[t])$ module. Taking the dual module, we obtain an inductive system $\cdots \subset (\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z}))^* \subset (\mathcal{C}_{\lambda, M+1}^{(k)}(\mathcal{Z}))^* \subset \cdots$. The direct limit

$$\mathcal{L}_\lambda^{(k)}(\mathcal{Z}) = \varinjlim_M \left(\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z})^* \right)$$

admits an action of the algebra

$$U_{\mathcal{Z}}(\mathfrak{g}[t]) = \varprojlim_M U(\mathfrak{g}[t])/B_{M,\mathcal{Z}}.$$

Note that, in the special case where $\mathcal{Z} = \{0\}$,

$$U_{\{0\}}(\mathfrak{g}[t]) = \varprojlim_M U(\mathfrak{g}[t]) / (t^M \mathfrak{g}[t] U(\mathfrak{g}[t]))$$

is the completion by formal power series in t . We denote the completion of $U(\mathfrak{g}[t])^{\otimes N}$ as

$$(2.16) \quad \begin{aligned} & U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes} N} \\ &= \varprojlim_M U(\mathfrak{g}[t])^{\otimes N} / I^{(N)}(B_M, \dots, B_M). \end{aligned}$$

We use the following lemmas.

Lemma 2.10. *Suppose \mathfrak{b} is an ideal of a Lie algebra \mathfrak{a} . Then, we have*

$$U(\mathfrak{a}/\mathfrak{b}) \simeq U(\mathfrak{a})/\mathfrak{b}U(\mathfrak{a}).$$

Lemma 2.11. *Suppose $\mathfrak{a} = \mathfrak{a}_1 \oplus \dots \oplus \mathfrak{a}_N$ is a direct sum decomposition of a Lie algebra. Then, we have an isomorphism of algebras*

$$U(\mathfrak{a}) \simeq U(\mathfrak{a}_1) \otimes \dots \otimes U(\mathfrak{a}_N).$$

Now, we state a basic proposition about localization of the universal enveloping algebra.

Proposition 2.12. (i) *We have a canonical isomorphism*

$$U_{\mathcal{Z}}(\mathfrak{g}[t]) \simeq U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes} N}.$$

(ii) *The canonical inclusion $U = U(\mathfrak{g}[t]) \rightarrow U_{\mathcal{Z}}(\mathfrak{g}[t]) \simeq U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes} N}$ is the composition of $\Delta_{\mathcal{Z}}$ followed by the inclusion $U(\mathfrak{g}[t])^{\otimes N} \rightarrow U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes} N}$.*

(iii) *The action of $U^{\otimes N}$ on $\mathcal{L}_{\lambda}^{(k)}(\mathcal{Z})$ can be extended to an action of $U(\widehat{\mathfrak{g}})^{\otimes N}$.*

Proof. Since the points in \mathcal{Z} are distinct, the restriction of $\Delta_{\mathcal{Z}}$ to $\mathfrak{g}[t]$ induces an isomorphism

$$\iota_{\mathcal{Z}} : \mathfrak{g}[t] / \prod_{i=1}^N (t - z_i)^M \mathfrak{g}[t] \simeq \oplus^N (\mathfrak{g}[t] / t^M \mathfrak{g}[t]).$$

By Lemma 2.11 we have

$$\Delta_{\mathcal{Z}} : U \left(\mathfrak{g}[t] / \prod_{i=1}^N (t - z_i)^M \mathfrak{g}[t] \right) \xrightarrow{\simeq} U(\mathfrak{g}[t] / t^M \mathfrak{g}[t])^{\otimes N}$$

By Lemma 2.10, we have

$$U/B_{M,\mathcal{Z}} \simeq (U/t^M \mathfrak{g}[t]U)^{\otimes N}.$$

For (i) and (ii), it is enough to show that the kernel of the canonical surjection $U^{\otimes N} \rightarrow (U/t^M \mathfrak{g}[t]U)^{\otimes N}$ coincides with $I^{(N)}(B_M, \dots, B_M)$. This follows from Lemma 2.3.

The action of $U(\widehat{\mathfrak{g}})^{\otimes N}$ is defined as follows. Consider an element $x(t) = (x_1(t), \dots, x_N(t))$ of $\oplus^N \widehat{\mathfrak{g}}$. Choose an m such that $t^m x_i(t) \in \mathfrak{g}[t]$ for all i . Then $\prod_{j=1}^N (t - z_j)^m \iota_z^{-1}(x(t)) \in \mathfrak{g}[t]$, so that $\iota_z^{-1}(x(t))$ maps $\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z})$ to $\mathcal{C}_{\lambda, M-m}^{(k)}(\mathcal{Z})$ for $M \geq m$. Therefore $x(t)$ acts on $\mathcal{L}_{\lambda}^{(k)}(\mathcal{Z})$. \square

We will use the following fact.

Theorem 2.13. [FFu] *We have an isomorphism of $U(\widehat{\mathfrak{g}})^{\otimes N}$ -modules.*

$$\mathcal{L}_{\lambda}^{(k)}(\mathcal{Z}) \simeq \bigoplus_{\lambda_1, \dots, \lambda_N \in P_+^{(k)}} ([\lambda_1] \cdots [\lambda_N] : [\lambda])_k \cdot L_{\lambda_1^*}^{(k)} \boxtimes \cdots \boxtimes L_{\lambda_N^*}^{(k)}.$$

For $I \subset U(\mathfrak{g}[t])$, denote by $I^{\mathcal{Z}}$ its closure in $U_{\mathcal{Z}}(\mathfrak{g}[t])$ in the topology of the inverse limit.

Lemma 2.14.

$$\left(L_{\lambda}^{(k)}(\infty) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \right)^* \simeq \left(\mathcal{L}_{\lambda}^{(k)}(\mathcal{Z}) \right)^{S(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))^{\mathcal{Z}}}.$$

Proof. Choose M such that $X_i \supset B_M$ for all i . Then

$$L_{\lambda}^{(k)}(\infty) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \simeq \mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z}) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}).$$

We have

$$\left(\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z}) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \right)^* \simeq \left(\mathcal{C}_{\lambda, M}^{(k)}(\mathcal{Z})^* \right)^{S(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))}.$$

Taking the direct limit in M we obtain the statement of the lemma. \square

Lemma 2.15. *Suppose that $z_1, \dots, z_N \in \mathbb{C}$ are distinct. For any $L > 0$ and $i = 1, \dots, N$, there exists an $f_i(t) \in \mathbb{C}[t]$ satisfying*

$$f_i(t) \equiv \begin{cases} 1 \bmod (t - z_i)^L; \\ 0 \bmod (t - z_j)^L \end{cases} \quad \text{for } j \neq i.$$

Lemma 2.16. *Under the identification of Proposition 2.12 we have*

$$(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))^{\mathcal{Z}} \simeq \sum_{i=1}^N U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes}(i-1)} \widehat{\otimes} X_i^{\{0\}} \widehat{\otimes} U_{\{0\}}(\mathfrak{g}[t])^{\widehat{\otimes}(N-i)}.$$

Here, the right hand side is the completion in (2.16).

Proof. For a sufficiently large L , we have

$$(2.17) \quad I^{(N)}(B_L, \dots, B_L) \subset I^{(N)}(X_1, \dots, X_N).$$

Recall that $X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) = \Delta_{\mathcal{Z}}^{-1}(I^{(N)}(X_1, \dots, X_N))$.

We want to show that for a sufficiently large L and for any element $a \in I^{(N)}(X_1, \dots, X_N)$, there exists an element $x \in X_1 \otimes \dots \otimes X_N(\mathbb{Z})$ satisfying

$$(2.18) \quad \Delta_{\mathbb{Z}}(x) \equiv a \bmod I^{(N)}(B_L, \dots, B_L).$$

From (2.17) it is enough to find $x \in U(\mathfrak{g}[t])$ satisfying (2.18) because it implies $x \in X_1 \otimes \dots \otimes X_N(\mathbb{Z})$. The existence of such an x follows from

$$\Delta_{\mathbb{Z}}(f_i(t)g) \equiv 1 \otimes \dots \otimes g^{i\text{-th}} \otimes \dots \otimes 1 \bmod I^{(N)}(B_L, \dots, B_L),$$

where $g \in \mathfrak{g}[t]$ and $f_i(t)$ is given in Lemma 2.15. \square

Proof of Theorem 2.9. Set $X = X_1 \otimes \dots \otimes X_N(\mathbb{Z})$. Collecting together the statements above, we find

$$\begin{aligned} (L_{\lambda}^{(k)}(\infty)/X)^* &\simeq (\mathcal{L}_{\lambda}^{(k)}(\mathbb{Z}))^{S(X)^{\mathbb{Z}}} && \text{(by Lemma 2.14)} \\ &\simeq \bigoplus_{\lambda_1, \dots, \lambda_N \in P_+^{(k)}} ([\lambda_1] \dots [\lambda_N] : [\lambda])_k \cdot (L_{\lambda_1}^{(k)} \boxtimes \dots \boxtimes L_{\lambda_N}^{(k)})^{S(X)^{\mathbb{Z}}} \\ &&& \text{(by Theorem 2.13)} \\ &\simeq \bigoplus_{\lambda_1, \dots, \lambda_N \in P_+^{(k)}} ([\lambda_1] \dots [\lambda_N] : [\lambda])_k \cdot (L_{\lambda_1}^{(k)})^{S(X_1)} \boxtimes \dots \boxtimes (L_{\lambda_N}^{(k)})^{S(X_N)} \\ &&& \text{(by Lemma 2.16)} \\ &\simeq \bigoplus_{\lambda_1, \dots, \lambda_N \in P_+^{(k)}} ([\lambda_1] \dots [\lambda_N] : [\lambda])_k \\ &\quad \times (L_{\lambda_1}^{(k)}(\infty)/X_1 \boxtimes \dots \boxtimes L_{\lambda_N}^{(k)}(\infty)/X_N)^* \\ &&& \text{(by Proposition 2.5).} \end{aligned}$$

Therefore

$$\dim L_{\lambda}^{(k)}(\infty)/X = \sum_{\lambda_1, \dots, \lambda_N \in P_+^{(k)}} ([\lambda_1] \dots [\lambda_N] : [\lambda])_k \cdot \prod_{i=1}^N \dim L_{\lambda_i}^{(k)}(\infty)/X_i,$$

which proves the theorem. \square

2.4. Space of conformal coinvariants. We give a sketch of proof for the surjectivity statements in Introduction, which motivate the definition of spaces of coinvariants in two different settings. In Section 3 these two spaces will be identified. We also conclude Section 2 by stating the isomorphism between the space of conformal coinvariants and the space of coinvariants by the fusion right ideal.

The proof of the surjectivity of (1.3) and (1.6) is based on the equality (1.4) where we take $x \in \mathfrak{g}^{C \setminus \{p_0, \dots, p_N\}}$. There exists a meromorphic function f on the curve C which has poles only at p_0 and p_i ($i \neq 0$). Moreover, the

singular part of the pole at p_i can be chosen arbitrarily. By using (1.4) for $x = X \otimes f$ ($X \in \mathfrak{g}$), one can prove inductively the surjectivity of the map

$$L_0 \otimes \pi(L_1) \otimes \cdots \otimes \pi(L_N) \rightarrow \langle L_0, \dots, L_N \rangle_{p_0, \dots, p_N}^C.$$

Now, for (1.3), taking f which has a pole only at p_0 and such that $f(p_j) = \delta_{i,j}$, we can reduce the vectors in $\pi(L_i)$ to the highest weight vector v_i . For (1.6), we assume that the curve C is rational. Then, taking f which has a pole only at p_0 (we can choose an arbitrary singular part of the pole), we can reduce the vectors in L_0 to the highest weight vector v_0 .

Note that in the above proof we used only the property of the highest weight vector v_i that it is annihilated by $\mathfrak{g} \otimes t_i \mathbb{C}[t_i]$ and that it is a cyclic vector in L_i . Therefore, we can replace L_i by a direct sum of highest weight representations with a cyclic vector v_i in the degree (with respect to d) zero component. The surjectivity (1.3) is still valid.

For $\lambda \in P_+$, we define the left ideal $\mathfrak{X}_\lambda \subset U(\mathfrak{g})$ by

$$(2.19) \quad \mathfrak{X}_\lambda = U(\mathfrak{g})\mathfrak{n}_+ + \sum_{h \in \mathfrak{h}} U(\mathfrak{g})(h - \langle \lambda, h \rangle).$$

We also define $\mathfrak{X}'_\lambda \subset U(\mathfrak{g})$ by

$$(2.20) \quad \mathfrak{X}'_\lambda = U(\mathfrak{g})\mathfrak{n}_+ + \sum_{h \in \mathfrak{h}} U(\mathfrak{g})(h - \langle \lambda, h \rangle) + \sum_j U(\mathfrak{g})f_j^{\langle \lambda, \alpha_j^\vee \rangle + 1}$$

where α_j and α_j^\vee denote the simple roots and coroots respectively, and the f_j are root vectors corresponding to $-\alpha_j$. Namely \mathfrak{X}'_λ is the annihilating ideal of the highest weight vector $v_\lambda \in \pi_\lambda$.

Lemma 2.17. *We have*

$$\dim \pi_\mu / S(\mathfrak{X}_\lambda) = \dim \pi_\mu / S(\mathfrak{X}'_\lambda) = \delta_{\mu, \lambda^*}.$$

Proof. Let \mathfrak{X} be \mathfrak{X}_λ or \mathfrak{X}'_λ .

Then the right ideal $S(\mathfrak{X})$ contains \mathfrak{n}_+ and $h + \langle \lambda, h \rangle$ ($h \in \mathfrak{h}$). Since $-\lambda = w_0(\lambda^*)$, this implies $\pi_\mu / S(\mathfrak{X}) = 0$ unless $\mu = \lambda^*$.

For $\mu = \lambda^*$, denote by v_λ^* the lowest weight vector of π_{λ^*} such that $\langle v_\lambda^*, v_\lambda \rangle = 1$ with the highest weight vector $v_\lambda \in \pi_\lambda$. We have $\pi_{\lambda^*} / S(\mathfrak{X}) = \mathbb{C}v_\lambda^*$ if one can show that v_λ^* is not contained in the image of $S(\mathfrak{X})$. Suppose that $v_\lambda^* = S(g)w$ for some $g \in \mathfrak{X}$ and $w \in \pi_{\lambda^*}$. Then, we have

$$1 = \langle v_\lambda^*, v_\lambda \rangle = \langle S(g)w, v_\lambda \rangle = \langle w, gv_\lambda \rangle = 0$$

as elements of \mathfrak{X} annihilate the highest weight vector. This is a contradiction. \square

Corollary 2.18. *Set*

$$X_{\lambda_i} = S(\mathfrak{X}_{\lambda_i}) + B_1, \quad X'_{\lambda_i} = S(\mathfrak{X}'_{\lambda_i}) + B_1$$

for $\lambda_i \in P_+^{(k)}$. Then, we have

$$\dim L_\lambda^{(k)}(\infty) / X_{\lambda_1} \otimes \cdots \otimes X_{\lambda_N}(\mathbb{Z}) = ([\lambda_1^*] \cdots [\lambda_N^*] : [\lambda])_k,$$

$$\dim L_\lambda^{(k)}(\infty)/X'_{\lambda_1} \otimes \dots \otimes X'_{\lambda_N}(\mathcal{Z}) = ([\lambda_1^*] \cdots [\lambda_N^*] : [\lambda])_k.$$

Proof. As above, let \mathfrak{X} be \mathfrak{X}_λ or \mathfrak{X}'_λ and let X be X_λ or X'_λ respectively. Then

$$L_\lambda^{(k)}(\infty)/X = L_\lambda^{(k)}(\infty)/(B_1 + S(\mathfrak{X})) \simeq \pi_\lambda/S(\mathfrak{X}).$$

The statement follows from Lemma 2.17 and Theorem 2.9. \square

From Corollary 2.18 we have

Proposition 2.19. *Let z_1, \dots, z_N be distinct points in \mathbb{C} . Set*

$$X_{\lambda_i} = S(\mathfrak{X}_{\lambda_i}) + B_1, \quad X'_{\lambda_i} = S(\mathfrak{X}'_{\lambda_i}) + B_1$$

(i) *We have the canonical isomorphism given by (1.3)*

$$(2.21) \quad L_\lambda^{(k)}(\infty)/X'_{\lambda_1} \otimes \dots \otimes X'_{\lambda_N}(\mathcal{Z}) \simeq \left\langle L_\lambda^{(k)}, L_{\lambda_1}^{(k)}, \dots, L_{\lambda_N}^{(k)} \right\rangle_{\infty, z_1, \dots, z_N}^{\mathbb{C}P^1}$$

(ii) *We also have the canonical isomorphism*

$$(2.22) \quad L_\lambda^{(k)}(\infty)/X_{\lambda_1} \otimes \dots \otimes X_{\lambda_N}(\mathcal{Z}) \simeq \left\langle L_\lambda^{(k)}, L_{\lambda_1}^{(k)}, \dots, L_{\lambda_N}^{(k)} \right\rangle_{\infty, z_1, \dots, z_N}^{\mathbb{C}P^1}$$

Proof. First let us prove (i). The map (1.3) gives a surjection from the left hand side to the right hand side of (2.21). The injectivity follows since their dimensions are equal by Theorem 2.9 and Corollary 2.18.

To prove (ii) note that as $X_\lambda \subset X'_\lambda$ we have the natural surjection

$$(2.23) \quad L_\lambda^{(k)}(\infty)/X_{\lambda_1} \otimes \dots \otimes X_{\lambda_N}(\mathcal{Z}) \rightarrow L_\lambda^{(k)}(\infty)/X'_{\lambda_1} \otimes \dots \otimes X'_{\lambda_N}(\mathcal{Z}).$$

Theorem 2.9 and Corollary 2.18 imply that the dimensions of these spaces are equal, so (2.23) and therefore (2.22) is an isomorphism. \square

3. SPACES OF COINVARIANTS IN TERMS OF FILTERED TENSOR PRODUCT

In this section we establish a connection between spaces of coinvariants (2.15) and the filtered tensor product \mathcal{F}_Z of finite dimensional \mathfrak{g} -modules introduced in [FL]. Motivated by Corollary 2.18, we restrict our considerations to right ideals X_1, \dots, X_N containing B_1 , so that

$$X_i = S(\mathfrak{X}_i) + B_1$$

where \mathfrak{X}_i is a left ideal of $U(\mathfrak{g})$. The main statement is Theorem 3.6.

3.1. Annihilating ideals. For a \mathfrak{g} -module V , we denote by $\text{Ann}V \subset U(\mathfrak{g})$ the kernel of the structure map $U(\mathfrak{g}) \rightarrow \text{End}(V)$. $\text{Ann}V$ is a two-sided ideal and $S(\text{Ann}V) = \text{Ann}V^*$. If V is irreducible, then $U(\mathfrak{g})/\text{Ann}V \simeq \text{End}(V) \simeq V^* \otimes V$. The left \mathfrak{g} -action (resp., the right \mathfrak{g} -action) on $U(\mathfrak{g})/\text{Ann}V$ translates to the left \mathfrak{g} -action on V (resp., the right \mathfrak{g} -action on V^*).

The representation $L_\lambda^{(k)}(\infty)$ is a quotient of $U(\mathfrak{g}[t])$ by the left ideal which describes the conditions on the highest weight vector $v_\lambda^{(k)}$. Hence the space of coinvariants may be viewed as a bi-quotient of the algebra $U(\mathfrak{g}[t])$. More

precisely, let $e_\theta \in \mathfrak{n}_+$ be the root vector for the maximal root θ . Introduce the following left ideals of $U(\mathfrak{g}[t])$, which annihilate $v_\lambda^{(k)}$:

$$(3.1) \quad I_\lambda = U(\mathfrak{g}[t]) \otimes_{U(\mathfrak{g})} \mathfrak{X}_\lambda = U(\mathfrak{g}[t])\mathfrak{n}_+ + \sum_{h \in \mathfrak{h}} U(\mathfrak{g}[t])(h - \langle \lambda, h \rangle),$$

$$(3.2) \quad I_\lambda^{(k)} = I_\lambda + U(\mathfrak{g}[t])(e_\theta \otimes t)^{k - \langle \lambda, \theta^\vee \rangle + 1}.$$

These are not enough for the characterization of $v_\lambda^{(k)}$. However, we have

Lemma 3.1. *We have*

$$U(\mathfrak{g})/(\mathfrak{X}_\lambda + \text{Ann}\pi_\lambda) \simeq \pi_\lambda.$$

In other words, we have the equality of the left ideals (see (2.20))

$$\mathfrak{X}'_\lambda = \mathfrak{X}_\lambda + \text{Ann}\pi_\lambda.$$

Proof. We have

$$U(\mathfrak{g})/\text{Ann}\pi_\lambda \simeq \pi_\lambda^* \otimes \pi_\lambda.$$

Taking the quotient by the left ideal \mathfrak{X}_λ we have

$$(\pi_\lambda^* \otimes \pi_\lambda)/(\pi_\lambda^* \mathfrak{X}_\lambda) \otimes \pi_\lambda \simeq \mathbb{C} v_\lambda^* \otimes \pi_\lambda \simeq \pi_\lambda.$$

□

We have the following characterization of $L_\lambda^{(k)}(\infty)$ as a quotient of $U(\mathfrak{g}[t])$.

Proposition 3.2. *We have*

$$L_\lambda^{(k)}(\infty) \simeq U(\mathfrak{g}[t])/(I_\lambda^{(k)} + U(\mathfrak{g}[t])\text{Ann}\pi_\lambda).$$

Proof. Using Lemma 3.1, we obtain

$$\begin{aligned} L_\lambda^{(k)}(\infty) &\simeq (U(\mathfrak{g}[t]) \otimes_{U(\mathfrak{g})} \pi_\lambda) / U(\mathfrak{g}[t])(e_\theta \otimes t)^{k - \langle \lambda, \theta^\vee \rangle + 1} \\ &\simeq U(\mathfrak{g}[t]) \otimes_{U(\mathfrak{g})} (U(\mathfrak{g})/(\mathfrak{X}_\lambda + \text{Ann}\pi_\lambda)) / U(\mathfrak{g}[t])(e_\theta \otimes t)^{k - \langle \lambda, \theta^\vee \rangle + 1} \\ &\simeq U(\mathfrak{g}[t]) / (I_\lambda^{(k)} + U(\mathfrak{g}[t])\text{Ann}\pi_\lambda). \end{aligned}$$

□

Set

$$\mathfrak{J}^{(k)} = \text{Ann} \bigoplus_{\lambda \in P_+^{(k)}} \pi_\lambda.$$

Clearly $\mathfrak{J}^{(k)}$ is a two-sided ideal and $S(\mathfrak{J}^{(k)}) = \mathfrak{J}^{(k)}$. We have an isomorphism of $U(\mathfrak{g})$ -bimodules

$$(3.3) \quad U(\mathfrak{g})/\mathfrak{J}^{(k)} \simeq \bigoplus_{\lambda \in P_+^{(k)}} \pi_\lambda^* \otimes \pi_\lambda.$$

For a left ideal $\mathfrak{X} \subset U(\mathfrak{g})$, introduce the left \mathfrak{g} -module

$$(3.4) \quad \pi^{(k)}(\mathfrak{X}) = \bigoplus_{\lambda \in P_+^{(k)}} \left(\pi_\lambda^* \right)^* \otimes \pi_\lambda,$$

and the left ideal

$$\tilde{\mathfrak{X}} = \mathfrak{X} + \mathfrak{J}^{(k)} \subset U(\mathfrak{g}).$$

Proposition 3.3. *We have*

$$(3.5) \quad \pi^{(k)}(\mathfrak{X}) \simeq U(\mathfrak{g})/\tilde{\mathfrak{X}}.$$

In particular, $\pi^{(k)}(\mathfrak{X})$ is a cyclic \mathfrak{g} -module, the cyclic vector being the sum of canonical vectors of $(\pi_\lambda^\mathfrak{X})^ \otimes \pi_\lambda \simeq \text{Hom}_\mathbb{C}(\pi_\lambda^\mathfrak{X}, \pi_\lambda)$ corresponding to the inclusion $\pi_\lambda^\mathfrak{X} \subset \pi_\lambda$.*

Proof. The isomorphism (3.5) is clear from (3.3):

$$U(\mathfrak{g})/\tilde{\mathfrak{X}} \simeq (U(\mathfrak{g})/\mathfrak{J}^{(k)})/\mathfrak{X} \simeq \bigoplus_{\lambda \in P_+^{(k)}} (\pi_\lambda^*/\pi_\lambda^*\mathfrak{X}) \otimes \pi_\lambda \simeq \pi^{(k)}(\mathfrak{X}).$$

Since $1 \in U(\mathfrak{g})$ is mapped in (3.3) to the canonical vector in the right hand side, the statement about the cyclic vector follows. \square

The statement of Lemma 3.1 can be improved as follows.

Corollary 3.4. *For $\lambda \in P_+^{(k)}$ we have*

$$U(\mathfrak{g})/(\mathfrak{X}_\lambda + \mathfrak{J}^{(k)}) \simeq \pi^{(k)}(\mathfrak{X}_\lambda) \simeq \pi_\lambda.$$

In other words, we have the equality of the left ideals

$$(3.6) \quad \mathfrak{X}'_\lambda = \mathfrak{X}_\lambda + \text{Ann}\pi_\lambda = \mathfrak{X}_\lambda + \mathfrak{J}^{(k)} = \widetilde{\mathfrak{X}_\lambda}$$

3.2. Isomorphism with filtered tensor product.

Proposition 3.5. *Let $\tilde{\mathfrak{X}}_i \subset U(\mathfrak{g})$ be left ideals such that $U(\mathfrak{g})/\tilde{\mathfrak{X}}_i$ is finite dimensional, $i = 1, \dots, N$. Define right ideals of $U(\mathfrak{g}[t])$ by $\tilde{X}_i = S(\tilde{\mathfrak{X}}_i) + B_1$. Then we have a canonical isomorphism*

$$(3.7) \quad U(\mathfrak{g}[t])/\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathbb{Z}) \simeq \mathcal{F}_\mathbb{Z}(U(\mathfrak{g})/\tilde{\mathfrak{X}}_1, \dots, U(\mathfrak{g})/\tilde{\mathfrak{X}}_N).$$

In this isomorphism the right action of an element $x \in U(\mathfrak{g}[t])$ in the left hand side is translated to the left action of $S(x)$ in the right hand side.

Proof. Set $U = U(\mathfrak{g}[t])$, and let $p_i : U \rightarrow U/\phi_{z_i}(\tilde{X}_i)$ be the projection. Consider the composition of maps

$$\begin{aligned} U &\xrightarrow{\Delta^{(N-1)}} U \otimes \cdots \otimes U \\ &\xrightarrow{p_1 \otimes \cdots \otimes p_N} U/\phi_{z_1}(\tilde{X}_1) \otimes \cdots \otimes U/\phi_{z_N}(\tilde{X}_N) \\ &\xrightarrow{S \otimes \cdots \otimes S} U/\phi_{z_1}(\tilde{\mathfrak{X}}_1 + B_1) \otimes \cdots \otimes U/\phi_{z_N}(\tilde{\mathfrak{X}}_N + B_1). \end{aligned}$$

Since $U/\phi_{z_i}(\tilde{\mathfrak{X}}_i + B_1)$ is isomorphic to the evaluation module $U(\mathfrak{g})/\tilde{\mathfrak{X}}_i$ at $t = z_i$, the last member is nothing but

$$\mathcal{F}_\mathbb{Z}(U(\mathfrak{g})/\tilde{\mathfrak{X}}_1, \dots, U(\mathfrak{g})/\tilde{\mathfrak{X}}_N).$$

The map $U \rightarrow \mathcal{F}_\mathbb{Z}(U(\mathfrak{g})/\tilde{\mathfrak{X}}_1, \dots, U(\mathfrak{g})/\tilde{\mathfrak{X}}_N)$ is surjective since the filtered tensor product is cyclic.

Proposition is proved if we show that the kernel of $(p_1 \otimes \cdots \otimes p_N) \circ \Delta^{(N-1)}$ coincides with $\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathbb{Z})$. This is shown similarly as in the proof of Proposition 2.12 by using Lemma 2.3. \square

We are now in a position to state the main result of this section.

Theorem 3.6. *Let $\mathfrak{X}_i \subset U(\mathfrak{g})$ ($i = 1, \dots, N$) be a set of left ideals. Define right ideals of $U(\mathfrak{g}[t])$ by $X_i = S(\mathfrak{X}_i) + B_1$. Then we have a canonical isomorphism of filtered vector space*

$$(3.8) \quad L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathbb{Z}) \simeq \mathcal{F}_Z\left(\pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N)\right) / S\left(I_\lambda^{(k)}\right)$$

where $\pi^{(k)}(\mathfrak{X})$ is defined in (3.4) and the right ideal $S(I_\lambda^{(k)})$ is given by (3.2):

$$S(I_\lambda^{(k)}) = \mathfrak{n}_+ U(\mathfrak{g}[t]) + \sum_{h \in \mathfrak{h}} (h + \langle \lambda, h \rangle) U(\mathfrak{g}[t]) + (e_\theta \otimes t)^{k - \langle \lambda, \theta^\vee \rangle + 1} U(\mathfrak{g}[t]).$$

Proof. Set

$$\tilde{\mathfrak{X}}_i = \mathfrak{X}_i + \mathfrak{I}^{(k)}.$$

We have

$$\tilde{X}_i \stackrel{\text{def}}{=} S(\tilde{\mathfrak{X}}_i) + B_1 = X_i + \mathfrak{I}^{(k)} U(\mathfrak{g}[t]).$$

Since $\mathfrak{I}^{(k)}$ annihilates the \mathfrak{g} -module $L_\lambda^{(k)}(\infty)/B_1$, we have $L_\lambda^{(k)}(\infty)/X_i \simeq L_\lambda^{(k)}(\infty)/\tilde{X}_i$. Comparing dimensions using Theorem 2.9, we find that the canonical projection

$$(3.9) \quad L_\lambda^{(k)}(\infty)/X_1 \otimes \cdots \otimes X_N(\mathbb{Z}) \longrightarrow L_\lambda^{(k)}(\infty)/\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathbb{Z})$$

is an isomorphism.

We have

$$\begin{aligned} U(\mathfrak{g}[t])/\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathbb{Z}) &\simeq \mathcal{F}_Z(U(\mathfrak{g})/\tilde{\mathfrak{X}}_1, \dots, U(\mathfrak{g})/\tilde{\mathfrak{X}}_N) \\ &\quad (\text{by Proposition 3.5}) \\ &\simeq \mathcal{F}_Z(\pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N)). \\ &\quad (\text{by Proposition 3.3}) \end{aligned}$$

Therefore, from Proposition 3.2 follows that

$$(3.10) \quad \begin{aligned} &L_\lambda^{(k)}(\infty)/\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathbb{Z}) \\ &\simeq \mathcal{F}_Z(\pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N)) / \left(S(I_\lambda^{(k)}) + \text{Ann} \pi_{\lambda^*} U(\mathfrak{g}[t]) \right). \end{aligned}$$

The filtered tensor product $V = \mathcal{F}_Z(\pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N))$ is a finite dimensional \mathfrak{g} -module. Hence it has the form $V = \oplus_\mu V_\mu$ where V_μ is a direct sum of copies of π_μ . Noting that $S(I_\lambda) = \mathfrak{n}_+ U(\mathfrak{g}[t]) + \sum_{h \in \mathfrak{h}} (h + \langle \lambda, h \rangle) U(\mathfrak{g}[t])$, we have $V/S(I_\lambda) \simeq V_{\lambda^*}/S(I_\lambda)$, and $\text{Ann}(\pi_{\lambda^*})$ acts as 0 on it. Therefore

$$(3.11) \quad V / \left(S(I_\lambda^{(k)}) + \text{Ann}(\pi_{\lambda^*}) U(\mathfrak{g}[t]) \right) \simeq V/S(I_\lambda^{(k)}).$$

Combining (3.9), (3.10) and (3.11), we obtain the assertion. \square

For filtered tensor product of irreducible modules, we obtain the following rule for the dimension.

Corollary 3.7. *Suppose that $\lambda_1, \dots, \lambda_N \in P_+^{(k)}$. Then, we have*

$$(3.12) \quad \dim \mathcal{F}_{\mathcal{Z}}(\pi_{\lambda_1}, \dots, \pi_{\lambda_N}) / S(I_{\lambda}^{(k)}) = ([\lambda_1^*] \cdots [\lambda_N^*] : [\lambda])_k.$$

Proof. Set $\mathfrak{X}_i = \mathfrak{X}_{\lambda_i}$. From Corollary 3.4 we have $\pi^{(k)}(\mathfrak{X}_i) \simeq \pi_{\lambda_i}$. Therefore, we have

$$\mathcal{F}_{\mathcal{Z}}(\pi_{\lambda_1}, \dots, \pi_{\lambda_N}) / S(I_{\lambda}^{(k)}) \simeq \mathcal{F}_{\mathcal{Z}}(\pi^{(k)}(\mathfrak{X}_1), \dots, \pi^{(k)}(\mathfrak{X}_N)) / S(I_{\lambda}^{(k)})$$

Using Theorem 3.6, and applying Theorem 2.9 and Corollary 2.18, we have the assertion. \square

Example 1. For $\lambda_i \in P_+^{(k)}$, set $\mathfrak{X}_i = \mathfrak{X}_{\lambda_i}$ or $\mathfrak{X}_i = \mathfrak{X}'_{\lambda_i}$ (see (2.19), (2.20)). Then, from Corollary 3.4, we have

$$L_{\lambda}^{(k)}(\infty) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \simeq \mathcal{F}_{\mathcal{Z}}(\pi_{\lambda_1}, \dots, \pi_{\lambda_N}) / S(I_{\lambda}^{(k)}).$$

By Proposition 2.19, we have an isomorphism to the space of conformal coinvariants on \mathbb{CP}^1 :

$$L_{\lambda}^{(k)}(\infty) / X_1 \otimes \cdots \otimes X_N(\mathcal{Z}) \simeq L_{\lambda} \boxtimes L_{\lambda_1} \boxtimes \cdots \boxtimes L_{\lambda_N} / \mathfrak{g}^{\mathbb{CP}^1 \setminus \{\infty, z_1, \dots, z_N\}}.$$

Example 2. Let $\mathfrak{X}_i = 0$ for all i . Then $\pi^{(k)}(0) = \sum_{\lambda \in P_+^{(k)}} \pi_{\lambda}^* \otimes \pi_{\lambda}$, and

$$L_{\lambda}^{(k)}(\infty) / B_{1, \mathcal{Z}} \simeq \mathcal{F}_{\mathcal{Z}}(\pi^{(k)}(0), \dots, \pi^{(k)}(0)) / S(I_{\lambda}^{(k)}).$$

The left hand side has dimension $\left(\left(\sum_{\mu \in P_+^{(k)}} \dim \pi_{\mu} \cdot [\mu] \right)^N : [\lambda] \right)_k$.

Example 3. Let $\mathfrak{X}_i = \mathfrak{X} = U(\mathfrak{g})\mathbf{n}_+$ for all i . We choose the sum of highest weight vectors of π_{λ} as cyclic vector of the module $\pi^{(k)}(\mathfrak{X}) \simeq \bigoplus_{\lambda \in P_+^{(k)}} \pi_{\lambda}$. Set $X = \mathbf{n}_+ U(\mathfrak{g}) + B_1 = (\mathbf{n}_+ \otimes \mathbb{C}[t] \oplus \mathfrak{h} \otimes t\mathbb{C}[t] \oplus \mathbf{n}_- \otimes t\mathbb{C}[t])U(\mathfrak{g}[t])$. From Proposition 2.1, the fusion right ideal is generated by the Lie subalgebra

$$\mathfrak{a}(\mathcal{Z}) = \mathbf{n}_+ \otimes \mathbb{C}[t] \oplus \mathfrak{h} \otimes \prod_{i=1}^N (t - z_i)\mathbb{C}[t] \oplus \mathbf{n}_- \otimes \prod_{i=1}^N (t - z_i)\mathbb{C}[t].$$

Then, we have

$$L_{\lambda}^{(k)}(\infty) / \mathfrak{a}(\mathcal{Z}) \simeq \mathcal{F}_{\mathcal{Z}}(\pi^{(k)}(\mathfrak{X}), \dots, \pi^{(k)}(\mathfrak{X})) / S(I_{\lambda}^{(k)}).$$

The left hand side has dimension $\left(\left(\sum_{\mu \in P_+^{(k)}} [\mu] \right)^N : [\lambda] \right)_k$.

Example 4. Let $\mathfrak{g} = \mathfrak{sl}_2$ with standard generators e, f, h . Let π_l ($0 \leq l \leq k$) denote the $(l+1)$ -dimensional irreducible representation, $[l]$ the corresponding element of $\mathcal{V}^{(k)}$, and l the highest weight. For $0 \leq m \leq k$, set

$$\begin{aligned} \mathfrak{V}_m &= U(\mathfrak{g})e^{m+1} + U(\mathfrak{g})f, \\ \overline{\mathfrak{V}}_m &= U(\mathfrak{g})f^{m+1} + U(\mathfrak{g})e, \end{aligned}$$

and $Y_m = S(\mathfrak{Y}_m) + B_1$, $\overline{Y}_m = S(\overline{\mathfrak{Y}}_m) + B_1$. Then

$$\pi^{(k)}(\mathfrak{Y}_m) = \oplus_{l=0}^k \left(\pi_l^{\mathfrak{Y}_m} \right)^* \otimes \pi_l \simeq \oplus_{l=0}^m \pi_l,$$

with the cyclic vector being the sum of lowest weight vectors. Likewise $\pi^{(k)}(\overline{\mathfrak{Y}}_m) \simeq \oplus_{l=0}^m \pi_l$ has the sum of highest weight vectors as cyclic vector. Let Y be the fusion right ideal of

$$\overbrace{Y_1, \dots, Y_1}^{m_1}, \dots, \overbrace{Y_k, \dots, Y_k}^{m_k}, \overbrace{\overline{Y}_1, \dots, \overline{Y}_1}^{n_1}, \dots, \overbrace{\overline{Y}_k, \dots, \overline{Y}_k}^{n_k}.$$

Then the space of coinvariants $L_l^{(k)}(\infty)/Y$ has dimension

$$\left(([0] + [1])^{m_1+n_1} \dots ([0] + \dots + [k])^{m_k+n_k} : [l] \right)_k.$$

For $m = k$, one can replace \mathfrak{Y}_k by $U(\mathfrak{g})f$ without changing the space of coinvariants (3.8) since $\pi^{(k)}(\mathfrak{Y}_k) = \pi^{(k)}(U(\mathfrak{g})f)$. Similarly, one can replace $\overline{\mathfrak{Y}}_k$ by $U(\mathfrak{g})e$. Therefore, instead of Y_k and \overline{Y}_k , one can use

$$\begin{aligned} Z &= (e \otimes t\mathbb{C}[t] \oplus h \otimes t\mathbb{C}[t] \oplus f \otimes \mathbb{C}[t])U(\mathfrak{g}), \\ \overline{Z} &= (e \otimes \mathbb{C}[t] \oplus h \otimes t\mathbb{C}[t] \oplus f \otimes t\mathbb{C}[t])U(\mathfrak{g}), \end{aligned}$$

respectively.

Consider the special case where $m_i = n_i = 0$ for $1 \leq i \leq k-1$ and $m_k = M$, $n_k = N$. In this case, the fusion right ideal

$$\overbrace{Z \otimes \dots \otimes Z}^M \otimes \overbrace{\overline{Z} \otimes \dots \otimes \overline{Z}}^N (z_1, \dots, z_M, z'_1, \dots, z'_N)$$

is generated by the Lie subalgebra

$$\begin{aligned} &\mathfrak{a}^{(M,N)}(\mathcal{Z}, \mathcal{Z}') \\ &= e \otimes \prod_{i=1}^M (t - z_i) \mathbb{C}[t] \oplus h \otimes \prod_{i=1}^M (t - z_i) \prod_{i=1}^N (t - z'_i) \mathbb{C}[t] \oplus f \otimes \prod_{i=1}^N (t - z'_i) \mathbb{C}[t]. \end{aligned}$$

The space of coinvariants with respect to $\mathfrak{a}^{(M,N)}(\mathcal{Z}, \mathcal{Z}')$, in particular in the degeneration limit $z_1 = \dots = z_M = z'_1 = \dots = z'_N = 0$, is the subject of the works [FKLMM2, FKLMM3].

3.3. Fusion Kostka polynomials. In this subsection, we first recall the definition of q -supernomials, Kostka polynomials and their generalizations from the literature, and then propose their counterparts in the theory of fusion product.

For two partitions λ, μ such that $|\lambda| = |\mu|$, the Kostka-Foulkes polynomials $K_{\lambda, \mu}(q)$ are the entries of the transition matrix from Schur functions $s_\lambda(x)$ to Hall-Littlewood functions $P_\mu(x; q)$ (see [Mac] p. 239):

$$(3.13) \quad s_\lambda(x) = \sum_{\mu} K_{\lambda, \mu}(q) P_\mu(x, q).$$

If $|\lambda| \neq |\mu|$, $K_{\lambda, \mu}(q) = 0$.

From the theory of Schur functions, it is known that the Kostka number $K_{\lambda,\mu}(1)$ is the cardinality of the set of semi-standard tableaux $\text{Tab}(\lambda, \mu)$ of shape λ and weight μ . Therefore it was conjectured by [Fo] that there should exist a charge statistic $c : \text{Tab}(\lambda, \mu) \rightarrow \mathbb{Z}_{\geq 0}$ on semi-standard tableaux such that

$$K_{\lambda,\mu}(q) = \sum_{t \in \text{Tab}(\lambda, \mu)} q^{c(t)}.$$

This charge statistic was indeed found in [LS].

In the representation theory of \mathfrak{gl}_n , the Kostka number appears in two ways. In this connection, we have the restriction on the length of λ : $\ell(\lambda) \leq n$. First, for μ satisfying $\ell(\mu) \leq n$, it is equal to the multiplicity of the weight μ in the \mathfrak{gl}_n representation V_λ with highest weight λ .

Secondly, for general μ such that $\ell(\mu) = N$, it is equal to the multiplicity of the \mathfrak{gl}_n -representation π_λ in the tensor product of the N symmetric tensor representations $V_{(\mu_i)}$ of \mathfrak{gl}_n :

$$(3.14) \quad \pi_{(\mu_1)} \otimes \cdots \otimes \pi_{(\mu_N)} \simeq \bigoplus_{\lambda: \ell(\lambda) \leq n} K_{\lambda,\mu}(1) \pi_\lambda.$$

From the latter viewpoint, the Kostka polynomial can be regarded as the q -multiplicity of the representation π_λ in the tensor product of symmetric tensor representations. Namely, it is the Hilbert polynomial in the variable q of the graded space

$$\text{Hom}_{\mathfrak{gl}_n}(\pi_\lambda, \pi_{(\mu_1)} \otimes \cdots \otimes \pi_{(\mu_N)}),$$

where the grading is given by the charge statistic.

This interpretation first appeared in [KR], where the completeness problem for Bethe ansatz solutions of the XXX model was considered. This is also the place where fermionic formulas for the Kostka polynomials first appeared. In that approach, the fermionic formulas were proven by constructing a (charge preserving) bijection between rigged configurations appearing in the completeness problem for the Bethe equations, and Young tableaux. The term “fermionic formulas” were invented in the works of Stony Brook group [KKMM],[KM], in which it was found that the Bethe ansatz solution gives, in the conformal limit, the characters of conformal field theory in fermionic form, where the q -grading is the linearized quasi-particle energy function. This gives a physical interpretation to the q -grading used in [KR].

Kostka polynomials were generalized [KS, SW1, SW2] to the case where the representations in the tensor product correspond to highest weight representations for \mathfrak{sl}_n with highest weight $m\lambda_i$ (rectangular highest weight) where λ_i ($1 \leq i \leq n-1$) are the fundamental weights.

There are two variants of Kostka polynomials: the q -supernomials and the level-restricted Kostka polynomials.

The q -supernomials are related to the Kostka polynomials by

$$S_{\lambda,\mu}(q) = \sum_{\nu} K_{\nu,\lambda}(1) K_{\nu,\mu}(q).$$

Namely, it is the q -multiplicity of the weight λ in the tensor product (3.14). Fermionic formulas for the q -supernomial $S_{\lambda,\mu}(q)$ is known [HKKOTY], and recursion equations for the generalized version in the sense of rectangular diagram are studied [SW2].

The level restriction has an origin in the corner transfer matrix method of Baxter. It was systematically studied as one-dimensional configuration sums in connection with the characters of integrable highest weight representations of affine Lie algebras. In this context, the level parameter appears as the level of representation. Kashiwara's theory of crystal base gave a solid basis to this approach. Quite generally, the crystal base of level k integrable highest weight representations of affine Lie algebras are given in terms of paths, i.e., elements in semi-infinite product $B^{\otimes \infty/2}$ of a finite dimensional affine crystal B . The q -statistics for paths are related to the eigenvalues of corner transfer matrix in the zero temperature limit.

The connection between Kostka polynomials and crystal base was discovered and generalized in [NY, SW1, SW2, HKKOTY]. In these works were considered finite and inhomogeneous paths, i.e., elements in a finite tensor product $B_1 \otimes \cdots \otimes B_N$ of affine crystals B_i . There are actions of Kashiwara operators \tilde{e}_i, \tilde{f}_i ($i = 0, 1, \dots, r$) on affine crystals corresponding to the Chevalley generators e_i, f_i of the affine Lie algebras. In this language, the generalized Kostka polynomial is the q -statistical sum over paths restricted by the highest weight conditions.

$$\tilde{e}_i(b_1 \otimes \cdots \otimes b_N) = 0 \quad (i = 1, \dots, r).$$

The q -supernomial is nothing but the unrestricted sum, and the level k restricted Kostka polynomial is the sum with the integrability conditions

$$\tilde{f}_0^{k - \langle \lambda, \theta^\vee \rangle + 1}(b_1 \otimes \cdots \otimes b_N) = 0.$$

For the generalized Kostka polynomials, fermionic formulas are known in the case of rectangular weights [KSS]. To our knowledge, no formulas are available at present for an arbitrary sequence of highest weights. Fermionic formulas for the level restricted (generalized) Kostka polynomials were obtained in [SS] (see also [HKKOTY]).

Now let us formulate the fusion product analogs of the Kostka and related polynomials. In what follows we consider only left ideals $\mathfrak{X} \subset U(\mathfrak{g})$ which are also P -graded,

$$\mathfrak{X} = \bigoplus_{\nu \in P} \mathfrak{X}_\nu,$$

where $\mathfrak{X}_\nu = \{x \in \mathfrak{X} \mid [h, x] = \langle \nu, h \rangle x \ \forall h \in \mathfrak{h}\}$. The space of coinvariants are then P -graded, as well as filtered by the degree of t . In the isomorphism (3.7) of Proposition 3.5, the left action by $h \in \mathfrak{h}$ on the left hand side is translated to the right action by $S(h) = -h$ on the filtered tensor product (the right hand side). The latter commutes with the action of \mathfrak{g} , and is not to be confused with the left action of $h \in \mathfrak{g}$. To be concrete, let us take a basis $\{u_{\lambda,i}\}$ of $\pi_\lambda^{\mathfrak{X}}$ and the dual basis $\{u_\lambda^i\}$ of $(\pi_\lambda^{\mathfrak{X}})^*$. The cyclic vector

of $\pi^{(k)}(\mathfrak{X})$ is $u^{(k)}(\mathfrak{X}) = \sum_{\lambda \in P_+^{(k)}} \sum_i u_\lambda^i \otimes u_{\lambda,i}$, and the right action of $S(h)$ ($h \in \mathfrak{h}$) on a vector $xu^{(k)}(\mathfrak{X})$ ($x \in \mathfrak{g}$) is given by

$$(3.15) \quad xu^{(k)}(\mathfrak{X})S(h) = - \sum_{\lambda \in P_+^{(k)}} \sum_i (u_\lambda^i h) \otimes (xu_{\lambda,i}).$$

The P -grading on the space of coinvariants (3.8) of Theorem 3.6 is induced from the above P -grading.

In general, let $W = \oplus_{\nu \in P} W_\nu = \varinjlim F^j$ be a P -graded vector space equipped with a filtration $0 = F^{-1} \subset F^0 \subset F^1 \subset \dots$ by P -graded subspaces F^j . We define the character of the associated graded space by

$$\text{ch gr } W = \sum_{j \in \mathbb{Z}_{\geq 0}, \nu \in P} \dim(F^j / F^{j-1})_\nu q^j e^\nu,$$

where e^ν stands for the formal exponential symbol. When the P -grading is trivial, we use also the notation ch_q .

The foregoing discussions lead us to consider three kinds of polynomials in q , $\mathcal{S}_\lambda(V_1, \dots, V_N)$, $\mathcal{K}_\lambda(V_1, \dots, V_N)$ and $\mathcal{K}_\lambda^{(k)}(V_1, \dots, V_N)$.

Fusion q -supernomial coefficient

$$(3.16) \quad \text{ch gr } \mathcal{F}_Z(V_1, \dots, V_N) = \sum_{\lambda \in P} e^\lambda \mathcal{S}_\lambda(V_1, \dots, V_N).$$

Fusion Kostka polynomial²

$$(3.17) \quad \mathcal{K}_\lambda(V_1, \dots, V_N) = \text{ch gr} \left(\mathcal{F}_Z(V_1, \dots, V_N) / S(I_\lambda) \right),$$

Fusion restricted Kostka polynomial

$$(3.18) \quad \mathcal{K}_\lambda^{(k)}(V_1, \dots, V_N) = \text{ch gr} \left(\mathcal{F}_Z(V_1, \dots, V_N) / S(I_\lambda^{(k)}) \right).$$

The first two are simply related by an alternating sum over the Weyl group as

$$\mathcal{K}_\lambda(V_1, \dots, V_N) = \sum_{w \in W} (-1)^{l(w)} \mathcal{S}_{w \circ \lambda}(V_1, \dots, V_N).$$

It is also clear that $\mathcal{K}_\lambda(V_1, \dots, V_N)$ is obtained from $\mathcal{K}_\lambda^{(k)}(V_1, \dots, V_N)$ in the limit $k \rightarrow \infty$.

We expect that, in the case when V_1, \dots, V_N are tensor powers of the fundamental representations discussed in [HKKOTY],[SW2], the ‘fusion-product’ version defined above reduces to the corresponding object known in the literature. We expect also the following relation:

²The fusion Kostka polynomial defined in [BKMW] is different from ours.

Conjecture.

$$(3.19) \quad \mathcal{K}_\lambda^{(k)}(V_1, \dots, V_N) = \sum_{\substack{w \in W_{\text{aff}} \\ w \circ \lambda \in P_+}} (-1)^{l(w)} q^{d_w(\lambda, k)} \mathcal{K}_{w \circ \lambda}(V_1, \dots, V_N).$$

The quantity $d_w(\lambda, k)$ is defined in (2.8). For the cases corresponding to Examples 2,3, (3.19) was put forward in [FL] as the ‘main conjecture’.

4. THE CASE $\mathfrak{g} = \mathfrak{sl}_2$

In this section we restrict ourselves to the case $\mathfrak{g} = \mathfrak{sl}_2$ as mentioned in (A.6). In this special case, we shall identify the fusion-product version of Kostka/level-restricted Kostka polynomials with the ordinary ones and prove the identity (3.19). As before, we write π_l for the $(l+1)$ -dimensional irreducible representation and $[l]$ for the corresponding element of $\mathcal{V}^{(k)}$. We also write the left ideal (3.2) as $I_l^{(k)}$, and $S(I_l^{(k)})$ as $\langle h[0] + l, e[0], e[1]^{k-l+1} \rangle$, where $x[i] = x \otimes t^i$.

4.1. level-restricted Kostka polynomials. First we recall a few basic facts about Kostka and level-restricted Kostka polynomials for \mathfrak{sl}_2 , following the exposition of [SS]. We fix the notation as follows. Let $\mathbf{m} = (m_1, \dots, m_k)$ be a k -tuple of non-negative integers, and let $0 \leq l \leq k$. Here and after, we use boldface letters to represent a vector. The transposition symbol is omitted. Set

$$\begin{aligned} K_{l, \mathbf{m}}(q) &= q^{\|\mathbf{m}\|} K_{\lambda R(\mathbf{m})}(q^{-1}), \\ K_{l, \mathbf{m}}^{(k)}(q) &= q^{\|\mathbf{m}\|} K_{\lambda R(\mathbf{m})}^k(q^{-1}). \end{aligned}$$

Here the right hand sides stand for the Kostka and level-restricted Kostka polynomials in the notation of [SS],

$$\begin{aligned} \lambda &= \left(\frac{|\mathbf{m}| + l}{2}, \frac{|\mathbf{m}| - l}{2} \right), \\ R(\mathbf{m}) &= (k^{m_k}, \dots, 2^{m_2}, 1^{m_1}), \end{aligned}$$

and $|\mathbf{m}| = \sum_{i=1}^k i m_i$, $2\|\mathbf{m}\| = \sum_{1 \leq i, j \leq k} \min(i, j) m_i m_j - |\mathbf{m}|$. Set further

$$(4.1) \quad A_{ab} = \min(a, b), \quad v_a = \max(a - k + l, 0).$$

We use the q -binomial symbol

$$\begin{aligned} \begin{bmatrix} \mathbf{m} \\ \mathbf{n} \end{bmatrix} &= \prod_{i=1}^k \begin{bmatrix} m_i \\ n_i \end{bmatrix}, \\ \begin{bmatrix} m \\ n \end{bmatrix} &= \begin{cases} \frac{[m]!}{[n]![m-n]!} & (0 \leq n \leq m), \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

where $[n]! = \prod_{j=1}^n ((1 - q^j)/(1 - q))$. We shall use the following facts.

Fermionic formula:

$$(4.2) \quad K_{l,\mathbf{m}}^{(k)}(q) = \sum_{\substack{\mathbf{s} \in \mathbb{Z}_{\geq 0}^k \\ 2|\mathbf{s}| = |\mathbf{m}| - l}} q^{\mathbf{s}A\mathbf{s} + \mathbf{v}\mathbf{s}} \begin{bmatrix} A(\mathbf{m} - 2\mathbf{s}) - \mathbf{v} + \mathbf{s} \\ \mathbf{s} \end{bmatrix}.$$

Alternating sum formula:

$$(4.3) \quad \begin{aligned} K_{l,\mathbf{m}}^{(k)}(q) &= \sum_{i \geq 0} q^{(k+2)i^2 + (l+1)i} K_{2(k+2)i+l,\mathbf{m}}(q) \\ &\quad - \sum_{i > 0} q^{(k+2)i^2 - (l+1)i} K_{2(k+2)i-l-2,\mathbf{m}}(q). \end{aligned}$$

Verlinde number: In the Verlinde algebra $\mathcal{V}^{(k)}$, we have

$$(4.4) \quad [1]^{m_1} \cdots [k]^{m_k} = \sum_{l=0}^k K_{l,\mathbf{m}}^{(k)}(1)[l].$$

Eq. (4.2) and (4.3) are taken from Theorem 6.2 and eq. (6.8) in [SS], respectively (after simplifications due to $n = 2$). In [HKKOTY], $K_{l,\mathbf{m}}^{(k)}(q)$ is defined as a generating sum over level k restricted paths in affine crystals. With this definition, (4.4) follows from the fact that the structure constants of the Verlinde algebra are the same as the number of successive triples in a path satisfying the level-restriction conditions.

4.2. The goal. The fusion product is defined to be the associated graded vector space of the filtered tensor product

$$V_1 * \cdots * V_N(\mathcal{Z}) = \text{gr } \mathcal{F}_{\mathcal{Z}}(V_1, \dots, V_N).$$

It gives a meaning to the ‘limit’ of the filtered tensor product where all points z_i tend to 0, as explained in Appendix. We consider the filtered tensor product and fusion product of cyclic modules $V_i = \pi_{l_i}$, choosing the highest weight vector as cyclic vector. Let $\mathbf{m} = (m_1, \dots, m_k)$ and set

$$(4.5) \quad V_{\mathbf{m}} = \overbrace{\pi_k * \cdots * \pi_k}^{m_k} * \cdots * \overbrace{\pi_1 * \cdots * \pi_1}^{m_1}(\mathcal{Z}).$$

It is known (for $\mathfrak{g} = \mathfrak{sl}_2$) that (4.5) is independent of \mathcal{Z} [FL, FF]. Our goal is to show the following.

Theorem 4.1. (i) *Notations being as above, we have*

$$\text{ch}_q V_{\mathbf{m}} / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle = K_{l,\mathbf{m}}^{(k)}(q).$$

(ii) *The identity (3.19) is true in this case.*

In the next subsection, we prove the inequality

$$(4.6) \quad \text{ch}_q V_{\mathbf{m}} / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle \leq K_{l,\mathbf{m}}^{(k)}(q).$$

Namely, each coefficient of powers in q in the right hand side is greater than or equal to that in the left hand side.

Assuming (4.6), let us prove Theorem 4.1.

Proof of Theorem 4.1. From Corollary 3.7 and (A.3), we have for any $V_i = \pi_{l_i}$ that

$$\begin{aligned} ([l_1] \cdots [l_N] : [l])_k &= \dim \mathcal{F}_{\mathcal{Z}}(V_1, \dots, V_N) / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle \\ &\leq \dim V_1 * \cdots * V_N / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle. \end{aligned}$$

Combining this with (4.4), we obtain

$$K_{l, \mathbf{m}}^{(k)}(1) \leq \dim V_{\mathbf{m}} / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle.$$

Therefore, it follows from (4.6) that equality takes place in all intermediate steps. In particular, assertion (i) is implied by the equality in (4.6). Assertion (ii) is then an immediate consequence of the known alternating sum formula (4.3) for level-restricted Kostka polynomials. \square

4.3. Gordon filtration. It remains to prove the inequality (4.6). For this purpose we invoke the ‘functional’ description of the dual space [FS].

It was shown in [FF] that, as graded vector space, the dual of $V_{\mathbf{m}}$ is isomorphic to the space of all symmetric polynomials $f(x_1, \dots, x_s)$ ($s \geq 0$) satisfying the condition

$$(4.7) \quad \deg_x f(\overbrace{x, \dots, x}^a, x_{a+1}, \dots, x_s) \leq M_a - a$$

for any $a = 1 \cdots s$,

where

$$M_a = \sum_{i=1}^k \min(a, i) m_i.$$

Therefore the dual to the quotient space of (4.5) by $\langle h[0] + l, e[0], e[1]^{k-l+1} \rangle$ is realized as follows.

Lemma 4.2. *As a graded vector space, $(V_{\mathbf{m}} / \langle h[0] + l, e[0], e[1]^{k-l+1} \rangle)^*$ is isomorphic to the space \mathcal{F} of all symmetric polynomials satisfying (4.7) as well as the conditions*

- (i) *The number of variables is $s = \frac{1}{2}(|\mathbf{m}| - l)$,*
- (ii) *$f(0, x_2, \dots, x_s) = 0$,*
- (iii) *If $s \geq k - l + 1$, then setting $f(x_1, \dots, x_s) = (\prod_{i=1}^s x_i) g(x_1, \dots, x_s)$ we have*

$$(4.8) \quad g(\overbrace{0, \dots, 0}^{k-l+1}, x_{k-l+2}, \dots, x_s) = 0.$$

Lemma 4.3. *Assume that $s \geq k + 1$. For any $f \in \mathcal{F}$ we have*

$$f(\overbrace{x, \dots, x}^a, x_{a+1}, \dots, x_s) = 0 \quad (\text{if } a \geq k + 1).$$

Proof. Let $g(x_1, \dots, x_s)$ be as in Lemma 4.2 (iii). From (4.8), we find

$$(4.9) \quad \left(\frac{\partial}{\partial x}\right)^j g(\overbrace{x, \dots, x}^a, x_{a+1}, \dots, x_s) \Big|_{x=0} = 0$$

$$(0 \leq j \leq l, a \geq k+1).$$

Namely we have

$$(4.10) \quad g(\overbrace{x, \dots, x}^a, x_{a+1}, \dots, x_s) = O(x^{l+1}).$$

We prove the assertion by descending induction on a . First let $a = s$. Since $s \geq k+1$, we have $M_a = |\mathbf{m}|$. Then (4.7) and Lemma 4.2 (i) imply $\deg_x g(x, \dots, x) \leq |\mathbf{m}| - 2s = l$. Hence we have $g(x, \dots, x) = 0$ by (4.10). Suppose the assertion is true for $a+1$ ($a \geq k+1$). Then $g(x_1, \dots, x_s)$ can be written as

$$g(\overbrace{x, \dots, x}^a, x_{a+1}, \dots, x_s) = \prod_{i=a+1}^s (x - x_i)^2 h(x; x_{a+1}, \dots, x_s).$$

From (4.7) we find $\deg_x h \leq l$. Therefore (4.10) implies $h = 0$, and the proof is over. \square

In order to estimate the character of \mathcal{F} , we apply the standard technique of Gordon filtration [FS]. Let λ be a partition of width $\lambda_1 \leq k$, and write $\lambda = (k^{s_k} \dots 2^{s_2} 1^{s_1})$. For a symmetric polynomial f we define

$$\begin{aligned} \varphi_\lambda(f)(x_1^{(1)}, \dots, x_{s_1}^{(1)}; \dots; x_1^{(k)}, \dots, x_{s_k}^{(k)}) \\ = f(x_1^{(1)}, \dots, x_{s_1}^{(1)}, \dots, \overbrace{x_1^{(k)}, \dots, x_1^{(k)}}^k, \dots, \overbrace{x_{s_k}^{(k)}, \dots, x_{s_k}^{(k)}}^k), \end{aligned}$$

where each variable $x_i^{(a)}$ appears a times in the right hand side. Introduce the lexicographic ordering $\mu > \lambda$ by $\mu_1 = \lambda_1, \dots, \mu_{i-1} = \lambda_{i-1}$ and $\mu_i > \lambda_i$ for some i , and set

$$\mathcal{F}_\lambda = \bigcap_{\mu > \lambda} \text{Ker } \varphi_\mu.$$

Notice that $\mathcal{F}_{(k)} = \mathcal{F}$ by Lemma 4.3. The family of subspaces $\{\mathcal{F}_\lambda\}_\lambda$ defines a filtration of \mathcal{F} , whose associated graded space has the λ -th component isomorphic to $\varphi_\lambda(\mathcal{F}_\lambda)$. An element of the latter has the form

$$\begin{aligned} & \left(\prod_{\substack{1 \leq i \leq s_a \\ 1 \leq a \leq k}} x_i^{(a)} \right)^{a+(a-k+l)_+} \prod_{\substack{1 \leq i < j \leq s_a \\ 1 \leq a \leq k}} (x_i^{(a)} - x_j^{(a)})^{2a} \\ & \times \prod_{\substack{1 \leq i \leq s_a, 1 \leq j \leq s_b \\ 1 \leq a < b \leq k}} (x_i^{(a)} - x_j^{(b)})^{2 \min(a,b)} \cdot h, \end{aligned}$$

with a partially symmetric polynomial

$$h \in \mathbb{C}[x_1^{(1)}, \dots, x_{s_1}^{(1)}, \dots, x_1^{(k)}, \dots, x_{s_k}^{(k)}]^{\mathfrak{S}_{s_1} \times \dots \times \mathfrak{S}_{s_k}},$$

\mathfrak{S}_s denoting the symmetric group on s letters. We have the relations for the degree

$$\begin{aligned} \deg_{x_i^{(a)}} h &\leq (A(\mathbf{m} - 2\mathbf{s}) - \mathbf{v})_a, \\ \text{tot. deg } f &= \text{tot. deg } h + \mathbf{s}A\mathbf{s} + \mathbf{v}\mathbf{s}, \end{aligned}$$

where we used the vector notation (4.1), and tot. deg signifies the total degree. We thus obtain for the character $\text{ch}_q \text{gr } \mathcal{F}$ an upper bound

$$\sum_{\substack{\mathbf{s} \in \mathbb{Z}_{\geq 0}^k \\ 2|\mathbf{s}| = |\mathbf{m}| - l}} q^{\mathbf{s}A\mathbf{s} + \mathbf{v}\mathbf{s}} \begin{bmatrix} A(\mathbf{m} - 2\mathbf{s}) - \mathbf{v} + \mathbf{s} \\ \mathbf{s} \end{bmatrix},$$

which coincides with the fermionic form of the level-restricted Kostka polynomial (4.2). This completes the proof of (4.6).

APPENDIX A. DEGENERATION OF COINVARIANTS AND FUSION PRODUCT

Here we give a motivation for the notion of fusion product introduced in [FL] and used in Section 4.

In this paper we have considered fusion of congruence right ideals of $U(\mathfrak{g}[t])$ attached to distinct points. They are parameterized by the configuration space of complex points. In general, it is not clear what is the analog of fusion when points collide, but we can extend the definition to the boundary of the configuration space $\mathbb{C}^N \setminus \bigcup \Delta_{[a,b]}$ (see [FM]) in the following way.

For $a \in \mathbb{C} \setminus \{0\}$ we define an automorphism $l_a : \mathfrak{g}[t] \rightarrow \mathfrak{g}[t]$ multiplying the variable t by a^{-1} . Clearly we have

$$l_a(X_1 \otimes \dots \otimes X_N(\mathcal{Z})) = X_1 \otimes \dots \otimes X_N(a\mathcal{Z}).$$

These spaces form a family over $\mathbb{C} \setminus \{0\}$, and we want to extend it to the origin.

Definition 3. For a subspace W' of a graded vector space $W = \bigoplus_{j \geq 0} W_j$, we denote by W'^{top} the graded subspace of W spanned by homogeneous elements $w \in W_r$, such that for some $w_j \in W_j$ ($0 \leq j \leq r$) we have $w = w_r$ and $w_r + w_{r-1} + \dots + w_0 \in W'$.

Note that $\mathfrak{g}[t]$ and therefore $U(\mathfrak{g}[t])$ has a natural grading by degree of t . Hence, for any subspace $X \subset U(\mathfrak{g}[t])$ we can consider the subspace $X^{\text{top}} \subset U(\mathfrak{g}[t])$. In particular, the space $(X_1 \otimes \dots \otimes X_N(\mathcal{Z}))^{\text{top}}$ can be thought of as the limit of $X_1 \otimes \dots \otimes X_N(a\mathcal{Z})$ when the parameter a tends to 0.

More generally, combining the operation of \otimes and taking $(\dots)^{\text{top}}$ in various ways, one can extend the definition of fusion to the compactification

$\mathbb{C}[N]$ of the configuration space in the sense of [FM]. For example, corresponding to Figure 4 in [FM], we have the following fusion of right ideals:

$$\begin{aligned} Y_1 &= (X_1 \otimes X_3 \otimes X_8(z_1, z_3, z_8))^{\text{top}}, \\ Y_2 &= (X_4 \otimes X_6(z_4, z_6))^{\text{top}}, \\ Y_3 &= (X_2 \otimes X_5(z_2, z_5))^{\text{top}}, \\ Y_4 &= (Y_1 \otimes Y_2(z_{138}, z_{46}))^{\text{top}}, \\ Y_5 &= (Y_4 \otimes Y_3 \otimes X_7(z_{13468}, z_{25}, z_7))^{\text{top}}. \end{aligned}$$

Question 1. When is the ideal $(X_1 \otimes \dots \otimes X_N(\mathcal{Z}))^{\text{top}}$ independent of \mathcal{Z} ? More generally, when do the extended fusion right ideals depend only on the image of the point in $\mathbb{C}[N]$ under the natural map $\mathbb{C}[N] \rightarrow \mathbb{C}^N$?

In the following situation the top part of the right ideal is given rather explicitly.

Proposition A.1. *Let $\mathfrak{b} \subset \mathfrak{g}[t]$ be a Lie subalgebra. Then, we have*

$$(A.1) \quad \mathfrak{b}^{\text{top}} U(\mathfrak{g}[t]) = (\mathfrak{b} U(\mathfrak{g}[t]))^{\text{top}}.$$

Proof. Let $\mathbb{C}[t]_{\leq d}$ be the space of polynomials of degree less than or equal to d . Set $\mathfrak{b}_{\leq d} = \mathfrak{b} \cap \mathbb{C}[t]_{\leq d}$. One can construct a basis of \mathfrak{b} , $\{B_i\}$, in such a way that for each d the subset $\{B_i\} \cap \mathfrak{b}_{\leq d}$ constitutes a basis of $\mathfrak{b}_{\leq d}$. Then, $\{B_i^{\text{top}}\}$ is a basis of $\mathfrak{b}^{\text{top}}$. One can construct a homogeneous basis of $\mathfrak{g}[t]$ consisting of $\{B_i^{\text{top}}\}$ and new members $\{A_i\}$. Since \mathfrak{b} is a Lie subalgebra, by Lemma 2.2 $B_{i_1} \dots B_{i_m} A_{j_1} \dots A_{j_n}$ where $m \geq 1, i_1 \leq \dots \leq i_m$ and $n \geq 0, j_1 \leq \dots \leq j_n$ constitute a basis of the right ideal $\mathfrak{b} U(\mathfrak{g}[t])$. Note that

$$(B_{i_1} \dots B_{i_m} A_{j_1} \dots A_{j_n})^{\text{top}} = B_{i_1}^{\text{top}} \dots B_{i_m}^{\text{top}} A_{j_1} \dots A_{j_n}$$

constitute a basis of $(\mathfrak{b} U(\mathfrak{g}[t]))^{\text{top}}$. Since $\mathfrak{b}^{\text{top}}$ is also a Lie subalgebra, these elements constitute a basis of $\mathfrak{b}^{\text{top}} U(\mathfrak{g}[t])$. Therefore, we have the equality of the right ideals (A.1). \square

In particular, we have a positive answer to Question 1 for $X_i = B_{n_i}$, since

$$(B_{n_1} \otimes \dots \otimes B_{n_N})^{\text{top}} = B_{n_1 + \dots + n_N}.$$

Now let us proceed to coinvariants. By the definition, there is a canonical isomorphism of graded vector space

$$(A.2) \quad \text{gr}(W/W') \simeq W/W'^{\text{top}}.$$

On the other hand, if X is a right ideal of a graded algebra U acting on W , the dimension of the space of coinvariants may jump in general.

$$(A.3) \quad \dim W/XW \leq \dim W/X^{\text{top}}W.$$

Proposition A.2. *There is a surjective map*

$$(A.4) \quad L_{\lambda}^{(k)}(\infty) / (X_1 \otimes \dots \otimes X_N(\mathcal{Z}))^{\text{top}} \rightarrow \text{gr } L_{\lambda}^{(k)}(\infty) / (X_1 \otimes \dots \otimes X_N(\mathcal{Z}))$$

Question 2. When is the map (A.4) an isomorphism? More generally, when is the dimension of coinvariants independent of the point in $\mathbb{C}[N]$?

The answer to this question for $X_i = B_{n_i}$ is positive for $\mathfrak{g} = \mathfrak{sl}_2$ (see [FKLMM1]). On the other hand, if $\mathfrak{g} = E_8$, it is negative even for $X_i = B_1$ with $k = 1$. This is because for level 1 the only integrable representation is the vacuum representation, i.e., the highest weight is 0. Therefore, the right hand side is always one-dimensional, while the left hand side becomes infinite in the limit $N \rightarrow \infty$.

Similar considerations for the filtered tensor product of modules can be found in [FL]. Recall that the associated graded vector space of the filtered tensor product

$$(A.5) \quad V_1 * \cdots * V_N(\mathcal{Z}) = \text{gr } \mathcal{F}_{\mathcal{Z}}(V_1, \dots, V_N)$$

is called the fusion product. It gives a meaning to the similar limit of the filtered tensor product where all points z_i tend to 0. In [FL] it was conjectured that, for finite-dimensional cyclic modules over a simple Lie algebra \mathfrak{g} , the fusion product $V_1 * \cdots * V_N(\mathcal{Z})$ is independent of the choice of \mathcal{Z} . This conjecture was proved in [FL] in the simplest case where

$$(A.6) \quad \mathfrak{g} = \mathfrak{sl}_2, \quad V_i : \text{irreducible module} \\ \text{with highest weight vector as cyclic vector,}$$

and in [FF] by a different method. Indeed, the fusion product can also be extended to $\mathbb{C}[N]$ and it can be proved by the method of [FF] that in this case it depends only on the image in \mathbb{C}^N .

An advantage of this definition is that we can split the map (A.4) into two.

Theorem A.3. (i) *The map (A.4) is a composition of the following surjective maps*

$$(A.7) \quad L_{\lambda}^{(k)}(\infty)/(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))^{\text{top}} \rightarrow \pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)/S(I_{\lambda}^{(k)})$$

$$(A.8) \quad \pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)/S(I_{\lambda}^{(k)}) \rightarrow \text{gr } L_{\lambda}^{(k)}(\infty)/(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))$$

(ii) *Assume that $\mathfrak{X}_i \supset \mathfrak{J}^{(k)}$. Then the map (A.7) is an isomorphism.*

Proof. To prove (i) we just follow the proof of Theorem 3.6.

Let $\tilde{\mathfrak{X}}_i = \mathfrak{X}_i + \mathfrak{J}^{(k)}$. As usual, set $X_i = B_1 + S(\mathfrak{X}_i)$, $\tilde{X}_i = B_1 + S(\tilde{\mathfrak{X}}_i)$. Then the natural projection

$$L_{\lambda}^{(k)}(\infty)/(X_1 \otimes \cdots \otimes X_N(\mathcal{Z})) \rightarrow L_{\lambda}^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))$$

is an isomorphism.

So we can split the map (A.4) into composition of the projection

$$L_{\lambda}^{(k)}(\infty)/(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))^{\text{top}} \rightarrow L_{\lambda}^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}}$$

and the map (A.4) for \tilde{X}_i :

$$L_\lambda^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}} \rightarrow \text{gr } L_\lambda^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z})).$$

To prove (i) it remains to identify $L_\lambda^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}}$ with the space $\pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)/S(I_\lambda^{(k)})$.

We have

$$\begin{aligned} & L_\lambda^{(k)}(\infty)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}} \\ \simeq & U(\mathfrak{g}[t])/(I_\lambda^{(k)} + U(\mathfrak{g}[t])\text{Ann}\pi_\lambda)/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}} \\ & \quad (\text{by Proposition 3.2}) \\ \simeq & U(\mathfrak{g}[t])/(\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z}))^{\text{top}}/(I_\lambda^{(k)} + U(\mathfrak{g}[t])\text{Ann}\pi_\lambda) \\ & \quad (\text{as a bi-quotient by the left and right ideals}) \\ \simeq & \text{gr}\left(U(\mathfrak{g}[t])/\tilde{X}_1 \otimes \cdots \otimes \tilde{X}_N(\mathcal{Z})\right)/(I_\lambda^{(k)} + U(\mathfrak{g}[t])\text{Ann}\pi_\lambda) \\ & \quad (\text{by (A.2) and using that } I_\lambda^{(k)} \text{ is homogeneous}) \\ \simeq & \pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)/(S(I_\lambda^{(k)}) + \text{Ann}\pi_\lambda * U(\mathfrak{g}[t])) \\ & \quad (\text{by Proposition 3.5 and by (A.5)}) \\ \simeq & \pi^{(k)}(\mathfrak{X}_1) * \cdots * \pi^{(k)}(\mathfrak{X}_N)/S(I_\lambda^{(k)}). \\ & \quad (\text{by repeating the last part of the proof of Theorem 3.6}) \end{aligned}$$

To prove (ii) just note that $\mathfrak{X}_i \supset \mathfrak{I}^{(k)}$ implies $X_i = \tilde{X}_i$, so the first map is an isomorphism. \square

Let us illustrate how it works for the case of conformal blocks. For simplicity suppose $\mathfrak{g} = \mathfrak{sl}_2$.

For this purpose we need ideals $\mathfrak{X} \subset U(\mathfrak{g})$ such that $\pi^{(k)}(\mathfrak{X}) = \pi_l$. Actually, we can choose them in different ways. Following (2.19) and (2.20) set

$$\mathfrak{X}_l = U(\mathfrak{sl}_2)e + U(\mathfrak{sl}_2)(h - l),$$

$$\mathfrak{X}'_l = U(\mathfrak{sl}_2)e + U(\mathfrak{sl}_2)(h - l) + U(\mathfrak{sl}_2)f^{l+1}.$$

Then $\mathfrak{X}'_l \supset \mathfrak{I}^{(k)}$ because \mathfrak{X}'_l is the annihilating ideal of the highest weight vector in π_l where $0 \leq l \leq k$. Therefore, by Theorem A.3 the map (A.7) is an isomorphism. In Section 4 we prove that the map (A.8) is an isomorphism, so the composition map (A.4) is an isomorphism, i.e., the answer to the Question 2 is positive in this case.

But this is not so for \mathfrak{X}_l . Namely, let us show that the map (A.7) can not always be an isomorphism.

Let $X_i = S(\mathfrak{X}_i) + B_1$. In a way similar to the proof of Proposition A.1, one can show that $X_1 \otimes \cdots \otimes X_N(\mathcal{Z})$ is generated by $e \otimes t^j$ and $h \otimes t^j - \sum_{i=1}^N l_i z_i^j$ ($j \geq 0$). Hence $(X_1 \otimes \cdots \otimes X_N(\mathcal{Z}))^{\text{top}}$ is generated by $e \otimes t^j$ ($j \geq 0$), $h \otimes t^j$ ($j > 0$) and $h - \sum_{i=1}^N l_i$. This means that the left hand side of (A.7)

depends only on $\sum_{i=1}^N l_i$, but it is easy to see (e.g. by taking $k \rightarrow \infty$) that the dimension of the right hand side depends on each l_i .

Acknowledgments. This work is partially supported by the Grant-in-Aid for Scientific Research (A1) no.13304010 and (B2) no.14340040, Japan Society for the Promotion of Science. BF is partially supported by grants RFBR 02-01-01015 and INTAS-00-00055. The work of SL is partially supported by the grant RFBR-01-01-00546. The last stage of this work was carried out while the authors were visiting Mathematical Sciences Research Institute, Berkeley, March 2002.

REFERENCES

- [BKMW] L. Begin, A.N. Kirillov, P. Mathieu and M.A. Walton, Berenstein-Zelevinski triangles, elementary coupling and fusion rules, *Lett. Math. Phys.*, **28** (1993), 257-268.
- [FF] B. L. Feigin and E. Feigin, q -characters of the tensor products in \mathfrak{sl}_2 -case, math.QA/0201111 (2002).
- [FFu] B. L. Feigin and D. B. Fuchs, Cohomology of some nilpotent subalgebras of the Virasoro and Kac-Moody algebras, *Geometry and Physics*, Essays in honor of I.M.Gelfand on the occasion of his 75th birthday, eds. S.Gindikin and I.M.Singer, 209–235, North Holland, (1991)
- [FKLMM1] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, Combinatorics of $\widehat{\mathfrak{sl}}_2$ spaces of coinvariants, *Transformation Groups* **6** (2001) No.1 25–52.
- [FKLMM2] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, Combinatorics of the $\widehat{\mathfrak{sl}}_2$ spaces of Coinvariants: Loop Heisenberg modules and recursion, math.QA/0009198
- [FKLMM3] B. Feigin, R. Kedem, S. Loktev, T. Miwa and E. Mukhin, Combinatorics of the $\widehat{\mathfrak{sl}}_2$ spaces of coinvariants: Dual functional realization and recursion, math.QA/0012190
- [FL] B. L. Feigin and S. Loktev, On generalized Kostka polynomials and quantum Verlinde rule, math.QA/9812093, *Amer. Math. Sci. Transl.* **194** (1999) 61–79.
- [FM] W. Fulton and P. MacPherson. A compactification of configuration spaces, *Ann. Math.* **139** (1994) 183–225.
- [Fo] H. O. Foulkes, A survey of some combinatorial aspects of symmetric functions, in *Permutations*, Gauthier-Villars (1974), Paris.
- [FS] B. L. Feigin and A. V. Stoyanovsky, Functional models for representations of current algebras and semi-infinite Schubert cells, *Funct. Anal. and Its Appl.* **28** (1993) 55–72.
- [HKKOTY] G. Hatayama, A. N. Kirillov, A. Kuniba, M. Okado, T. Takagi and Y. Yamada, Character formulae of $\widehat{\mathfrak{sl}}_n$ -modules and inhomogeneous paths, math.QA/9802085, *Nucl. Phys. B* **536** (1999) 575–616.
- [KKMM] R. Kedem, T. R. Klassen, B. M. McCoy and E. Melzer, Fermionic sum representations for conformal field theory characters, *Phys. Lett. B* **307** (1993) 68–76.
- [KM] R. Kedem and B. M. McCoy, Construction of modular branching functions from Bethe’s equations in the 3-state potts chain, *J. Stat. Phys.* **71** (1993) 875–901.
- [KR] A. N. Kirillov and N. Yu. Reshetikhin, The Bethe Ansatz and the combinatorics of Young tableaux, *J. Soviet Math.* **41** (1988) 925-955.

- [KS] A. N. Kirillov and M. Shimozono, A generalization of the Kostka-Foulkes polynomials, *math.QA/9804039*, *J. Algebraic Combin.* **15** (2002) 27–69.
- [KSS] A. N. Kirillov, M. Shimozono and A. Schilling, A bijection between Littlewood-Richardson tableaux and rigged configurations. (Preprint *math.CO/9901037*).
- [LS] A. Lascoux and M.P. Schützenberger, Sur une conjecture de H. O. Foulkes, *C. R. Acad. Sci. Paris Ser. A-B* **286** (1978) A323–A324.
- [Mac] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford University Press, New York, 1995.
- [NY] A. Nakayashiki and Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, *Selecta Math.* **3** (1997) 547–599.
- [SS] A. Schilling and M. Shimozono, Fermionic formulas for level-restricted generalized Kostka polynomials and coset branching functions, *math.QA/0001114*, *Commun. Math. Phys.* **220** (2001) 105–164.
- [SW1] A. Schilling and O. Warnaar, Supernomial coefficients, polynomial identities and q -series, *Ramanujan J.* **2** (1998) 459–494.
- [SW2] A. Schilling and S. O. Warnaar, Inhomogeneous lattice paths, generalized Kostka polynomials and A_{n-1} supernomials, *math.QA/9802111*, *Commun. Math. Phys.* **202** (1999) 359–401.
- [TUY] A. Tsuchiya, K. Ueno and Y. Yamada, Conformal field theory on the universal family of stable curves with gauge symmetries, *Adv. Stud. Pure Math.* **19** (1989) 459–566.

BF: LANDAU INSTITUTE FOR THEORETICAL PHYSICS, CHERNOGOLOVKA, 142432, RUSSIA

E-mail address: feigin@feigin.mccme.ru

MJ: GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, TOKYO 153-8914, JAPAN

E-mail address: jimbomic@ms.u-tokyo.ac.jp

RK: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 W. GREEN ST. URBANA, IL 61801, USA

E-mail address: rinat@math.uiuc.edu

SL: INSTITUTE FOR THEORETICAL AND EXPERIMENTAL PHYSICS, B. CHEREMUSHKINSKAYA, 25, MOSCOW 117259, RUSSIA

INDEPENDENT UNIVERSITY OF MOSCOW, B. VLASIEVSKY PER, 11, MOSCOW 121002, RUSSIA

E-mail address: loktev@mccme.ru

TM: DIVISION OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606-8502 JAPAN

E-mail address: tetsuji@kusm.kyoto-u.ac.jp