MAXIMAL COMPLEXIFICATIONS OF CERTAIN RIEMANNIAN HOMOGENEOUS MANIFOLDS

S. HALVERSCHEID AND A. IANNUZZI*

ABSTRACT. Let M = G/K be a Riemannian homogeneous manifold with $dim_{\mathbb{C}}G^{\mathbb{C}} = dim_{\mathbb{R}}G$, where $G^{\mathbb{C}}$ denotes the universal complexification of G. Under certain extensibility assumptions on the geodesic flow of M, we give a characterization of the maximal domain of definition in TM for the adapted complex structure and show that it is unique. For instance, this can be done for generalized Heisenberg groups and naturally reductive Riemannian homogeneous spaces. As an application it is shown that the case of generalized Heisenberg groups yields examples of maximal domains of definitions for the adapted complex structure which are neither holomorphically separable, nor holomorphically convex.

1. Introduction

It is well known that complexifications of a real-analytic manifold M exist and are equivalent near M, but differ usually very much in nature. If a complete real-analytic metric on M is given, one can construct canonical complexifications of M compatible with the metric by defining an adapted complex structure on a domain Ω of the tangent bundle TM (see [GS] and [LS]). This can be characterized by the condition that the "complexification" $(x+iy)\mapsto y\gamma'(x)\in\Omega$ of any geodesic $x\mapsto\gamma(x)$ of M be a complex submanifold near the zero section. By the results of Guillemin-Stenzel and Lempert-Szőke cited above, the adapted complex structure exists and is unique on a sufficiently small neighborhood of M. Here M is identified with its zero section in TM.

In particular it is natural to ask for maximal domains around M on which it exists which, by functoriality of the definition, may be regarded as invariants of the metric, i.e., isometric manifolds have biholomorphic maximal domains. For instance examples are known for symmetric spaces of non-compact type ([BHH]), compact normal Riemannian Homogeneous spaces ([Sz2]), compact symmetric spaces ([Sz1]) and spaces obtained by Kählerian reduction of these ([A]). Note that in the mentioned cases maximal domains turn out to be Stein.

The aim of this work is to characterize maximal domains for the adapted complex structure for a class of Riemannian homogeneous spaces with "big" isometry

²⁰⁰⁰ Mathematics Subject Classification. Primary 32C09, 53C30, 32M05; Secondary 32Q28. Key words: adapted complex structure, Riemannian homogeneous spaces, Stein manifold.

^{*}Partially supported by University of Bologna, funds for selected research topics.

group. Let M = G/K, with G a Lie group of isometries and K compact, and assume that $dim_{\mathbb{C}}G^{\mathbb{C}} = dim_{\mathbb{R}}G$, where $G^{\mathbb{C}}$ is the universal complexification of G. Then $K^{\mathbb{C}}$ acts on $G^{\mathbb{C}}$, the left action on M induces a natural G-action on TM and under certain extensibility assumptions on the geodesic flow of M one obtains a real-analyti and G-equivariant map $P:TM \to G^{\mathbb{C}}/K^{\mathbb{C}}$ such that (see Theorem 3.2 for the precise statement)

The connected component of the non-singular locus of DP containing M is the unique maximal domain on which the adapted complex structure exists.

This applies to the case of naturally reductive Riemannian homogeneous spaces (corollary 3.3) and of generalized Heisenberg groups (see Sect. 4 and 5).

As an application it is shown that for all generalized Heisenberg groups such maximal domain is neither holomorphically separable, nor holomorphically convex (Proposition 5.1). We are not aware of previous non-Stein examples. In the case of the 3-dimensional Heisenberg group we determine its envelope of holomorphy as well as a certain maximal Stein subdomain (Proposition 4.3).

2. Preliminaries

Here we introduce notations and briefly recall basic results we will need in the present paper. Let M be a complete real-analytic Riemannian manifold which will be often identified with the zero section in its tangent bundle TM. Following [LS] (see also [GS] for an equivalent characterization) we say that a real-analytic complex structure defined on a domain Ω of TM is adapted if all complex leaves of the Riemannian foliation are submanifolds with their natural complex structure, i. e., for any geodesic $\gamma: \mathbb{R} \to \mathbb{C}$ the induced map $f: \mathbb{C} \to TM$ defined by $(x+iy) \mapsto y \gamma'(x)$ is holomorphic on $f^{-1}(\Omega)$ with respect to the adapted complex structure. Here $y \gamma'(t) \in T_{\gamma(t)}M$ is the scalar multiplication in the vector space $T_{\gamma(t)}M$.

The adapted complex structure exists and is unique on a sufficiently small neighborhood of M and if Ω is a domain around M in TM on which it is defined, we refer to it as an adapted complexification. Sometimes these are called Grauert tubes. For later use we need the following

Lemma 2.1. Let $F:TM \to \mathbb{C}$ be a real-analytic map which is holomorphic on any complex leaf of TM in a neighbourhood of M. Then F is holomorphic on every adapted complexification.

Proof. Following the proof of [Sz1, Prop. 3.2 p. 416] one checks that the restriction of F to M extends to a holomorphic map \hat{F} in a neighborhood

U of $M \subset TM$ where the adapted complex structure J_0 exists and, in order to have connected leaves, U may be assumed to be starshaped. By assumption we can also assume that for any geodesic $\gamma: \mathbb{R} \to M$ the map $x+iy \mapsto F(y\cdot\gamma'(x))$ is holomorphic for all x+iy such that $y\gamma'(x)\in U$. Now $F=\hat{F}$ on $\gamma(\mathbb{R})\subset M\subset TM$, therefore $F=\hat{F}$ on every complex leaf, i.e., on U. In particular $DF\circ J_0=iDF$ on U and since all maps are real-analytic the statement follows from the identity principle.

A real Lie group G acts on a complex manifold X, i.e., X is a G-manifold, if there exists a real-analytic surjective map $G \times X \to X$ given by $(g,x) \mapsto g \cdot x$ such that for fixed $g \in G$ the map $x \mapsto g \cdot x$ is holomorphic and $(gh) \cdot x = g \cdot (h \cdot x)$ for all $h, g \in G$ and $x \in X$. Furthermore if $\dim_{\mathbb{R}} G = \dim_{\mathbb{C}} G^{\mathbb{C}}$, where $\iota : G \to G^{\mathbb{C}}$ is the universal complexification of G (see e.g. [Ho]), then $Lie(G^{\mathbb{C}}) = \mathfrak{g}^{\mathbb{C}}$ and one obtains an induced local holomorphic $G^{\mathbb{C}}$ -action by integrating the holomorphic vector fields given by the G-action. Here \mathfrak{g} denotes the Lie algebra of G.

Let M = G/K be a Riemannian homogeneous manifold with G a connected Lie group of isometries and K compact, and consider the induced G-action on TM defined by $g \cdot w := g_* w$ for all $g \in G$ and $w \in TM$. Then if Ω is a G-invariant adapted complexification, as an easy consequence of the definitions g_* is a biholomorphic extension of the isometry g, i.e., $G \subset Aut(\Omega)$.

If one assumes that $\dim_{\mathbb{R}} G = \dim_{\mathbb{C}} G^{\mathbb{C}}$, then the natural map $\iota: G \to G^{\mathbb{C}}$ is an immersion and from the universality property of the universal complexification $K^{\mathbb{C}}$ of K it follows that the restriction $\iota|_K$ of ι to K extends to an immersion $\iota^{\mathbb{C}}: K^{\mathbb{C}} \to G^{\mathbb{C}}$. Moreover the subgroup $\iota^{\mathbb{C}}(K^{\mathbb{C}})$ acts by right multiplication on $G^{\mathbb{C}}$ and one has a commutative diagram

$$G \xrightarrow{\iota} G^{\mathbb{C}}$$

$$\downarrow \qquad \qquad \downarrow$$

$$G/K \rightarrow G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}}).$$

Also notice the G-action on $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ defined by $g \cdot h \iota^{\mathbb{C}}(K^{\mathbb{C}}) := \iota(g) h \iota^{\mathbb{C}}(K^{\mathbb{C}})$ for all $g \in G$ and $h \in G^{\mathbb{C}}$.

Lemma 2.2. Let G be a connected Lie group, K a compact subgroup and assume that $\dim_{\mathbb{C}} G^{\mathbb{C}} = \dim_{\mathbb{R}} G$. Then $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ is a complex G-manifold and $\dim_{\mathbb{C}} G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}}) = \dim_{\mathbb{R}} G/K$.

Proof. One needs to show that $\iota^{\mathbb{C}}(K^{\mathbb{C}})$ is closed in $G^{\mathbb{C}}$. Note that $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ is the orbit space with respect to the $K^{\mathbb{C}}$ -action on $G^{\mathbb{C}}$ defined by $k \cdot h := h \iota(k^{-1})$ for all $k \in K^{\mathbb{C}}$ and $h \in G^{\mathbb{C}}$. Since $G^{\mathbb{C}}$ is Stein ([He]) and $K^{\mathbb{C}}$ is reductive it

follows that every fiber of the categorical quotient $G^{\mathbb{C}} \to G^{\mathbb{C}}/\!/K^{\mathbb{C}}$ is equivariantly biholomorphic to an affine algebraic variety on which $K^{\mathbb{C}}$ acts algebraically ([Sn]). In particular there exists at least one closed $K^{\mathbb{C}}$ -orbit and consequently $\iota^{\mathbb{C}}(K^{\mathbb{C}})$ is closed in $G^{\mathbb{C}}$. Thus $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ is a complex G-manifold and by construction its complex dimension is $\dim_{\mathbb{R}} G/K$.

3. A CHARACTERIZATION OF MAXIMAL ADAPTED COMPLEXIFICATIONS

If M=G/K is a symmetric space of the non-compact type, then $G^{\mathbb{C}}/K^{\mathbb{C}}$ is a natural candidate for a complexification of M and there exists a G-equivariant map $P:TM\to G^{\mathbb{C}}/K^{\mathbb{C}}$ embedding holomorphically a maximal adapted complexification of M (see [BHH], [Ha], [AG]). As a matter of fact one may show that DP is singular on the border $\partial\Omega$ of Ω .

Here we consider a Riemannian homogeneous manifold M = G/K endowed with the additional data of a certain real-analytic G-equivariant map P from TM to a suitable complex G-manifold, characterizing a maximal adapted complexification Ω_M as the connected component of $\{DP \text{ not singular }\}$ containing M. Unicity of Ω_M follows.

The existence of such a data is proved when $\dim_{\mathbb{C}} G^{\mathbb{C}} = \dim_{\mathbb{R}} G$ and the geodesic flow "extends" holomorphically on $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ (cf. Lemma 2.2). As a consequence the characterization applies to the case of naturally reductive Riemannian homogeneous spaces and of generalized Heisenberg groups.

Proposition 3.1. Let M = G/K be an n-dimensional Riemannian homogeneous space and X a G-complex manifold of complex dimension n such that the induced local $G^{\mathbb{C}}$ -action is locally transitive. Assume there exists a real-analytic map $P: TM \to X$ which is

- i) G-equivariant and
- ii) holomorphic on every complex leaf of TM.

Then the connected component Ω_M of $\{p \in TM : DP_p \text{ not singular }\}$ containing M is the unique maximal adapted complexification and $P|_{\Omega_M}$ is locally biholomorphic.

Proof. First we show that Ω_M is well defined, i.e., DP has maximal rank along M. Since from Lemma 2.1 it follows that P is holomorphic on M with respect to the adapted complex structure, this is a consequence of the following

Claim: Assume that P is holomorphic in $p \in TM$. Then DP_p has maximal rank.

Proof of the claim: Since $G^{\mathbb{C}}$ acts locally transitively on X, there exist elements ξ_1, \dots, ξ_n of \mathfrak{g} such that the induced vector fields $\xi_{X,1}, \dots, \xi_{X,n}$ on X span a totally real and maximal dimensional subspace $V_{P(p)}$ of $T_{P(p)}X$, where

$$\xi_{X,j}(x) := \frac{d}{dt}\Big|_{0} \exp_{G^{\mathbb{C}}}(t\xi_{j}) \cdot x$$

for $j=1,\cdots,n$ and all $x\in X$. By equivariance it follows that $DP_p(V_p)=V_{P(p)}$, where V_p is the subspace of T_pTM spanned by $\xi_{TM,1},\cdots,\xi_{TM,n}$, here $\xi_{TM,j}(q):=\frac{d}{dt}\bigg|_0\exp_G(t\xi)\cdot q$ for all $q\in TM$. In particular $\dim_{\mathbb{R}}V_p=n$ and since P is holomorphic in p, V_p is totally real and DP_p has maximal rank, proving the claim.

Now we see that the pulled-back complex structure J_o on Ω_M of the complex structure J on X is the adapted complex structure. For this consider a complex leaf $f: \mathbb{C} \to TM$ defined by $f(x+iy) := y \cdot \gamma'(x)$, where γ is a geodesic of M, and note that by ii

$$DP \circ Df(i\eta) = DP \circ J_o \circ Df(\eta)$$

for all η tangent in $f^{-1}(\Omega_M)$. Since DP has maximal rank on Ω_M , then

$$Df(i\eta) = J_o \circ Df(\eta)$$

showing that J_o is the adapted complex structure. In particular $P|_{\Omega_M}$ is locally biholomorphic.

In order to prove maximality, assume that J_o extends analytically in a neighborhood of a certain $p \in \partial \Omega_M \subset TM$. By construction $DP \circ J_o = J \circ DP$ on Ω_M and since all maps are real-analytic P is holomorphic in p. Then the above claim shows that DP_p has maximal rank, contradicting the definition of Ω_M .

Finally we want to show that any adapted complexification Ω is contained in Ω_M . If this is not the case, there exists a point p in $\Omega \cap \partial \Omega_M$ and from Lemma 2.1 it follows that $P|_{\Omega}$ is holomorphic. In particular P is holomorphic in p and one obtains a contradiction arguing as above. Thus Ω_M is unique and this concludes the statement.

Now we determine a class of Riemannian homogeneous spaces to which Proposition 3.1 may be applied in order to determine the maximal adapted complexification.

Theorem 3.2. Let M = G/K be a Riemannian homogeneous space with $\dim_{\mathbb{R}} G = \dim_{\mathbb{C}} G^{\mathbb{C}}$ and assume there exists a map $\varphi : \mathbb{C} \times T_K M \mapsto \mathfrak{g}^{\mathbb{C}}$ real-analytic and holomophic on the first component such that $\varphi(\mathbb{R} \times T_K M) \subset \mathfrak{g}$ and $t \mapsto \exp_G \circ \varphi(t, v) K$ is the unique geodesic tangent to v at 0 for all $v \in T_K M$. Then the map

$$P:TM\mapsto G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$$

defined by

$$P(g_*(v)) := \iota(g) \, \exp_{G^{\mathbb{C}}}(\varphi(i, v)) \, \iota^{\mathbb{C}}(K^{\mathbb{C}})$$

for all $g \in G$ and $v \in T_KM$ is as in Proposition 3.1. In particular the connected component Ω_M of $\{p \in TM : DP_p \text{ not singular }\}$ containing M is the maximal adapted complexification and $P|_{\Omega_M}$ is locally biholomorphic.

Proof. In order to prove that P is well defined we need to show that if $w = k_*v$ for some $k \in K$ and $v \in T_KM$ then $P(w) = P(k_*v)$, i. e.,

(1)
$$\exp_{G^{\mathbb{C}}}(\varphi(i,w)) \iota^{\mathbb{C}}(K^{\mathbb{C}}) = \iota(k) \exp_{G^{\mathbb{C}}}(\varphi(i,v)) \iota^{\mathbb{C}}(K^{\mathbb{C}}).$$

For this note that $t \mapsto k \exp_G \circ \varphi(t, v) K$ is the unique geodesic tangent to w at 0 in K, thus

$$\exp_G \circ \varphi(t, w) K = k \exp_G \circ \varphi(t, v) K$$
.

Then the commutativity of the diagram

$$\mathfrak{g} \longrightarrow \mathfrak{g}^{\mathbb{C}}$$
 $\downarrow \exp_G \qquad \downarrow \exp_{G^{\mathbb{C}}}$
 $G \stackrel{\iota}{\longrightarrow} G^{\mathbb{C}}$

implies that

$$\exp_{G^{\mathbb{C}}} \circ \varphi(t, w) \iota^{\mathbb{C}}(K^{\mathbb{C}}) = \iota(k) (\exp_{G^{\mathbb{C}}} \circ \varphi(t, v)) \iota^{\mathbb{C}}(K^{\mathbb{C}})$$

for all $t \in \mathbb{R}$ and equation (1) is a consequence of the identity principle for holomorphic maps.

Now define $\Phi(z,v) := \exp_{G^{\mathbb{C}}} \circ \varphi(z,v)$ for all $z \in \mathbb{C}$ and $v \in T_K M$ and, in order to simplify notations, assume that the canonical immersion $\iota: G \to G^{\mathbb{C}}$ is injective so that once we identify G with $\iota(G)$ the curve $\gamma(t) := \Phi(t,v)K$ is the unique geodesic tangent to v at 0. In what follows it is easy to check that all arguments apply to the case where ι is a non-injective immersion.

Fix $x \in \mathbb{R}$, let $g := \Phi(x, v)$ and note that

$$y \mapsto g \Phi(y, g_*^{-1} \gamma'(x)) K$$

is the unique geodesic tangent to $\gamma'(x)$ at 0. Therefore one has

$$\Phi(x,v)\Phi(y,g_*^{-1}\gamma'(x))K = \Phi(x+y,v)K$$

for all $y \in \mathbb{R}$ and by the identity principle it follows that

(2)
$$\Phi(x,v)\,\Phi(z,g_*^{-1}\gamma'(x))\,\iota^{\mathbb{C}}(K^{\mathbb{C}}) = \Phi(x+z,v)\,\iota^{\mathbb{C}}(K^{\mathbb{C}})$$

for all $v \in T_K M$, $x \in \mathbb{R}$ and $z \in \mathbb{C}$. For $h \in G$ and $v \in T_K M$ consider the unique geodesic $\tilde{\gamma}(t) := h \cdot \gamma(t)$ tangent to $h_*(v)$ at 0. One has

$$P(y\tilde{\gamma}'(x)) = P(h_*y\gamma'(x)) = h P(g_*g_*^{-1}y\gamma'(x)) = h g \Phi(i, y g_*^{-1}\gamma'(x)) \iota^{\mathbb{C}}(K^{\mathbb{C}}) = h \Phi(x, v) \Phi(iy, g_*^{-1}\gamma'(x)) \iota^{\mathbb{C}}(K^{\mathbb{C}}) = h \Phi(x + iy, v) \iota^{\mathbb{C}}(K^{\mathbb{C}}),$$

where we used (2) and the fact that $\Phi(z,yv) \iota^{\mathbb{C}}(K^{\mathbb{C}}) = \Phi(zy,v) \iota^{\mathbb{C}}(K^{\mathbb{C}})$ for all $z \in \mathbb{C}$, since this holds for all $z \in \mathbb{R}$. As a consequence the map $(x+iy) \mapsto P(y\tilde{\gamma}'(x))$ is holomorphic for all geodesics $\tilde{\gamma}$ of M, i.e., P is holomorphic on every complex leaf of TM.

Finally the map P is G-equivariant by construction and the G-action on $G^{\mathbb{C}}/\iota^{\mathbb{C}}(K^{\mathbb{C}})$ induces a holomorphic $G^{\mathbb{C}}$ -action which may be obtained through left multiplication on $G^{\mathbb{C}}$. Thus it is obviously transitive and this yields the statement.

Now let M be a naturally reductive Riemannian homogeneous space and M = G/K be a natural realization of M, i.e., there exists a reductive decomposition $\mathfrak{g} = Lie(K) \oplus \mathfrak{m}$ of the Lie algebra of G such that every geodesic in M is the orbit of a one parameter subgroup of G generated by an element of \mathfrak{m} (see e.g. [BTV]). Consider the natural projection $\Pi: G \to M$ and note that $D\Pi_e(\mathfrak{m}) = T_K M$, where e is the neutral element of G. Denote by $L: T_K M \to \mathfrak{m}$ the inverse of the restriction of $D\Pi_e$ to \mathfrak{m} . Since L is linear it extends \mathbb{C} -linearly from $(T_K M)^{\mathbb{C}}$ to $\mathfrak{m}^{\mathbb{C}}$ and the map $\varphi: \mathbb{C} \times T_K M \to \mathfrak{g}^{\mathbb{C}}$ defined by $\varphi(z,v):=zL(v)$ is as in the above Theorem. Therefore one has

Corollary 3.3. Let M = G/K be a natural realization of a naturally reductive Riemannian homogeneous space and assume that $\dim_{\mathbb{R}} G = \dim_{\mathbb{C}} G^{\mathbb{C}}$. Then the map

$$P:TM\to G^{\mathbb{C}}/\iota(K^{\mathbb{C}})$$

defined by

$$P(q_*(v)) := \iota(q) \exp_{C^{\mathbb{C}}}(iL(v)) \iota(K^{\mathbb{C}})$$

for all $g \in G$ and $v \in T_K M$ meets the conditions of Proposition 3.1. In particular the connected component Ω_M of $\{p \in TM : DP_p \text{ not singular }\}$ containing M is the maximal adapted complexification and $P|_{\Omega_M}$ is locally biholomorphic.

4. The 3-dimensional Heisenberg group

Here we apply results of the previous section in order to give a concrete description of the unique maximal adapted complexification for the 3-dimensional Heisenberg group. It turns out that such domain is neither holomorphically separable, nor holomorphically convex. We also determine its envelope of holomorphy and a particular maximal Stein subdomain. We remark that in all previous examples we are aware of, maximal adapted complexifications are Stein.

Consider the 3-dimensional Heisenberg group defined as a subgroup of $GL_3(\mathbb{R})$ by

$$H:=\left\{\left(\begin{array}{ccc} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array}\right) \ : \ \alpha,\beta,\gamma\in\mathbb{R}\right\},$$

fix the inner product of the tangent space T_eH in the neutral element e for which the canonical basis determined by the global natural chart (α, β, γ) is orthonormal and let (a, b, c) be coordinates of T_eH with respect to this basis. Endow H with the induced H-invariant metric

$$(d\alpha)^2 + (d\beta)^2 + (d\gamma - \alpha \cdot d\beta)^2,$$

let $\mathfrak{h} = Lie(H)$ and define $\varphi : \mathbb{R} \times T_eH \to \mathfrak{h}$ by

$$\varphi(t,(a,b,c)) \,:=\, \left(\, a\frac{\sin(tc)}{c} \,-\, b\frac{1-\cos(tc)}{c} \,\,,\,\, b\frac{\sin(tc)}{c} \,+\, a\frac{1-\cos(tc)}{c} \,\,,\,\, \left(\, t+\frac{a^2+b^2}{2c^2}\left(t-\frac{\sin(tc)}{c}\right)\right)c\,\right),$$

where the coordinates of \mathfrak{h} are induced by those of T_eH via the natural identification $\mathfrak{h} \cong T_eH$. Note that all singularities are removable and consequently φ is real-analytic. Following [BTV, Th. p. 31] one checks that $t \mapsto \exp_H \circ \varphi(t, (a, b, c))$ is the unique geodesic tangent to (a, b, c) at 0. Furthermore by expanding the power series it is easy to verify that $\varphi(\cdot, (a, b, c))$ extends holomorphically on \mathbb{C} to $(T_eH)^{\mathbb{C}}$ and by considering the polar decomposition $H \times \mathfrak{h} \to H^{\mathbb{C}}$ of $H^{\mathbb{C}}$ given by $(g, \xi) \mapsto g \exp_{H^{\mathbb{C}}}(i\xi)$, one obtains real-analytic functions $(a, b, c) \mapsto h_{(a,b,c)} \in H$ and $(a, b, c) \to \xi_{(a,b,c)} \in \mathfrak{h}$ such that

$$\exp_{H^{\mathbb{C}}} \circ \varphi(i,(a,b,c)) \ = \ h_{(a,b,c)} \, \exp_{H^{\mathbb{C}}} (i\xi_{(a,b,c)}).$$

Define $P: TH \to H^{\mathbb{C}} \cong H \times \mathfrak{h}$ by

$$g_*(a,b,c) \mapsto g \exp_{H^{\mathbb{C}}} \circ \varphi(i,(a,b,c)) \cong (gh_{(a,b,c)}, \xi_{(a,b,c)}).$$

Then Theorem 3.2 implies that the connected component Ω_H containing H of $\{DP \text{ not singular }\}$ is the maximal adapted complexification. Note that since P is H-equivariant, Ω_H is H-invariant. Moreover T_eH is a global slice for the H-action on TH, i.e., the map $H \times T_eH \to TH$ given by $(g, (a, b, c)) \to g_*(a, b, c)$ is

a H-equivariant real-analytic diffeomorphism, thus Ω_H is completely determined by its slice $\Omega_H \cap T_eH$.

Furthermore H acts freely on the first component of $H \times \mathfrak{h}$, then H-equivariance of P implies that $DP_{g_*(a,b,c)}$ has maximal rank if and only if $D\tilde{P}_{(a,b,c)}$ has maximal rank, where $\tilde{P}:=p_2\circ P|_{T_eH}:T_eH\to\mathfrak{h}$ is given by

$$\tilde{P}(a,b,c) \,=\, \xi_{(a,b,c)} \,=\, \left(\, a \frac{\sinh(c)}{c} \,\,,\,\, b \frac{\sinh(c)}{c} \,\,,\,\, \left(1 + \frac{a^2 + b^2}{2c^3} \left(c - \sinh(c) \, \cosh(c)\right)\right)c\,\right).$$

Here $p_2: H \times \mathfrak{h} \to \mathfrak{h}$ is the canonical projection. It follows that $\Omega_H = H \cdot O_0$, where O_0 is the connected component of $\{\det(D\tilde{P}) \neq 0\}$ containing 0 in T_eH . Now a straightforward computation shows that

$$\det(D\tilde{P}_{(a,b,c)}) = \frac{\sinh(c)}{c} \left(\frac{\sinh(c)}{c} + (a^2 + b^2) \left(\frac{\sinh(c) - c \cosh(c)}{c^3} \right) \right),$$

therefore

$$O_0 = \left\{ (a, b, c) \in T_e H : a^2 + b^2 < \frac{c^2 \sinh(c)}{c \cosh(c) - \sinh(c)} \right\}.$$

We want to discuss injectivity of $P|_{\Omega_H}:\Omega_H\to H^{\mathbb{C}}\cong H\times\mathfrak{h}$ and again this is equivalent to injectivity of $\tilde{P}|_{O_0}$.

Note that \tilde{P} is equivariant with respect to rotations around the c-axis as well as to the reflection σ with respect to the plane $\{c=0\}$. In particular for any

$$(a,b,c) \in \left\{ 1 + \frac{a^2 + b^2}{2c^3} \left(c - \sinh(c) \cosh(c) \right) = 0 \right\}$$

one has $\tilde{P}(a,b,c) = \tilde{P}(a,b,-c) = \left(a \frac{\sinh(c)}{c}, b \frac{\sinh(c)}{c}, 0\right)$. Therefore we are induced to investigate the domain

$$O_1 := \left\{ (a, b, c) \in T_e H : a^2 + b^2 < \frac{2c^3}{\sinh(c)\cosh(c) - c} \right\}.$$

Lemma 4.1. The domain O_1 is the maximal σ -invariant subdomain of O_0 containing 0 on which \tilde{P} is injective. In particular $\tilde{P}|_{O_0}$ is not injective.

Proof. Let f_i be the real function defining

$$O_i = \{ (a, b, c) \in T_e H : a^2 + b^2 < f_i(c) \}$$

for $j=0,1\,.$ First we want to show that $\,O_1\,$ is a subdomain of $\,O_0\,,$ i.e,

$$f_1(c) \le f_0(c)$$

for all $c \in \mathbb{R}$, which is equivalent to

$$2\cosh(c) \le \frac{\sinh(c)}{c} + \frac{\sinh^2(c)}{c^2}\cosh(c).$$

Expanding in power series one obtains

$$2 + c^2 + \frac{1}{12}c^4 + \dots \le \left(1 + \frac{1}{6}c^2 + \frac{1}{120}c^4 + \dots\right) + \left(1 + \frac{5}{6}c^2 + \left(\frac{5}{6} + \frac{1}{24} + \dots\right)c^4 + \dots\right).$$

All coefficients are non-negative and one easily checks that for $k \geq 2$ the coefficient of c^{2k} in the last series on the right side is strictly greater than that in the series on the left, hence $O_1 \subset O_0$. Moreover $\partial O_1 \cap \partial O_0 = \{(a, b, 0) \in T_eH : a^2 + b^2 = 3\}$, thus O_1 is a proper subdomain of O_0 .

Furthermore by the previous remarks any σ -invariant domain containing 0 on which \tilde{P} is injective is necessarily contained in O_1 .

Assume that there exist (a',b',c'), $(a'',b'',c'') \in O_1$ such that $\tilde{P}(a',b',c') = \tilde{P}(a'',b'',c'') =: (A,B,C)$. If C=0 then c'=c''=0 and consequently a'=a''=A and b=b''=B. If $C\neq 0$ by eventually acting with σ and a rotation around the c-axis we may assume that $a,A\geq 0$, b=B=0 and c>0. Now one has

$$a'\frac{\sinh(c')}{c'} = a''\frac{\sinh(c'')}{c''} = A$$

therefore (a',0,c') and (a'',0,c'') lie on the same level curve $\rho_A: \mathbb{R} \to T_e H$ given by

$$\rho_A(t) := \left(A \frac{t}{\sinh(t)}, 0, t \right).$$

One has the following

Claim: Let $A \geq 0$ and $t_0 \in \mathbb{R}^{\geq 0}$ such that $\rho_A(t_0) \in \overline{O}_0$. Then $\rho_A(t) \in O_0$ for all $t > t_0$.

Proof of the claim: One needs to show that $A^2 \frac{t^2}{\sinh^2(t)} < f_0(t)$ for all $t > t_0$, that is

(3)
$$A^2 < \frac{\sinh^2(t)}{t^2} f_0(t).$$

By expanding in power series as above one has the estimate

$$2t \cosh^2(t) - 3\cosh(t)\sinh(t) + t > 0,$$

for all t > 0, which by a straightforward computation implies that the derivative of the function at the right hand side of (3) is positive for all t > 0, proving the claim.

Now let $t_0 := \min(c', c'')$ and note that since $O_1 \subset O_0$, then as a consequence of the above claim there exists $\epsilon > 0$ such that $\rho_A(t) \in O_0$ for $t > t_0 - \epsilon$. In

particular (a', 0, c') and (a'', 0, c'') lie in the same connected real one dimensional submanifold $N := \rho_A(t_0 - \epsilon, \infty)$ of O_0 and $\tilde{P}|_N : N \to \{(A, 0, \cdot) \in T_e H\} \cong \mathbb{R}$ is locally diffeomorphic. Then a classical argument implies that $\tilde{P}|_N$ is injective, thus (a', 0, c') = (a'', 0, c'') as wished.

We also want to determine the image of $P|_{\Omega_H}$ in $H^{\mathbb{C}}$. Note that $P(\Omega_H)$ is H-invariant and the polar decomposition implies that $\exp_{H^{\mathbb{C}}}(i\mathfrak{g})$ is a global slice for the H-action on $H^{\mathbb{C}}$. Then this can be achieved by describing $\exp_{H^{\mathbb{C}}}(i\tilde{P}(O_0)) = P(\Omega_H) \cap \exp_{H^{\mathbb{C}}}(i\mathfrak{g})$.

Lemma 4.2.
$$\tilde{P}(O_0) = \mathfrak{h} \setminus \{ (A, B, C) \in \mathfrak{h} : A^2 + B^2 = 3, C = 0 \}.$$

Proof. Let $(a,0,c) \in \{a^2 = f_1(c)\} \subset \partial O_1$ with c > 0. From the proof of Lemma 4.1 it follows that $(a,0,c) \in O_0$. Since $\tilde{P}(a,0,c) = (a\frac{\sinh(c)}{c},0,0)$ and

$$\sqrt{f_1(0)} = \sqrt{3}$$
 and $\lim_{c \to \infty} \sqrt{f_1(c)} \frac{\sinh(c)}{c} = \infty$

it follows that $(A,0,0) \in \tilde{P}(O_0)$ for all $A > \sqrt{3}$.

For $A > \sqrt{3}$ let $(a,0,c) \in O_0$ such that $\tilde{P}(a,0,c) = (A,0,0)$. By the claim in Lemma 4.1 one has $\rho_A(t) \in O_0$ for all $t \geq c$. Moreover one sees that

(4)
$$\tilde{P}(\rho_A(t)) = (A, 0, C_A(t))$$
 with $\lim_{t \to \infty} C_A(t) = \infty$.

Then by σ -invariance of O_0 and σ -equivariance of \tilde{P} it follows that $(A,0,C) \in \tilde{P}(O_0)$ for all $C \in \mathbb{R}$, and $A > \sqrt{3}$.

Now note that $\tilde{P}(a,0,0)=(a,0,0)$ and $f_0(0)=3$, thus $(A,0,0)\in \tilde{P}(O_0)$ for all $A<\sqrt{3}$ and arguing as above it follows that $(A,0,C)\in \tilde{P}(O_0)$ for all $C\in\mathbb{R}$ and $A<\sqrt{3}$.


Finally $\rho_{\sqrt{3}}(0) \in \partial O_0$, thus $\rho_{\sqrt{3}}(t) \in O_0$ for all t>0. It follows that $t\mapsto \tilde{P}(\rho_{\sqrt{3}}(t))$ is injective for t>0 and since

$$\lim_{t\to 0^+} C_{\sqrt{3}}(t) = 0 \quad \text{ and } \quad \lim_{t\to \infty} C_{\sqrt{3}}(t) = \infty$$

then $(\sqrt{3}, 0, C) \in \tilde{P}(O_0)$ if and only if $C \neq 0$.

The statement follows from the invariance of O_0 and the equivariance of \tilde{P} with respect to the group of rotations around the c-axis.

In the picture below one sees the border of O_0 and O_1 determined by f_0 and f_1 respectively as well as the level curve ρ_2 in the upper half-plane $\{b=0, a \geq 0\}$ of T_eH . Since O_0 and O_1 are invariant with respect to rotations around



the c-axis this completely determine their shape and, by H-equivariance, that of Ω_H .

Proposition 4.3. The maximal domain Ω_H is neither holomorphically separable, nor holomorphically convex, its envelope of holomorphy is biholomorphic to \mathbb{C}^3 . The maximal σ -invariant Stein subdomain of Ω_H is biholomorphic to $\{(z_1, z_2, z_3) \in \mathbb{C}^3 : (\operatorname{Im} z_1)^2 + (\operatorname{Im} z_2)^2 < 3\}.$

Proof. From [CIT, Prop. p. 543] it follows that any holomorphically separable Riemann H-domain over $H^{\mathbb{C}}$ is univalent. Moreover Lemma 4.1 implies that $P|_{\Omega_H}:\Omega_H\to H^{\mathbb{C}}$ is not injective, therefore Ω_H is not holomorphically separable

By a result of Loeb ([L, Th. p. 186]), a Stein H-invariant domain U of $H^{\mathbb{C}}$ is "geodesically" convex, i.e., it is convex with respect to all curves of the form $t\mapsto g\exp_{H^{\mathbb{C}}}(it\xi)$, with $g\in U$ and $\xi\in\mathfrak{h}$. Since $H^{\mathbb{C}}$ admits polar decomposition and U is H-invariant it is enough to consider curves of the form $\exp_{H^{\mathbb{C}}}(i\eta)\exp_{H^{\mathbb{C}}}(it\xi)$, with $\exp_{H^{\mathbb{C}}}(i\eta)\in U$ and $\xi\in\mathfrak{h}$. Furthermore for a two steps nilpotent Lie group one has

$$\exp_{H^{\mathbb{C}}}(i\eta) \exp_{H^{\mathbb{C}}}(it\xi) = \exp_{H^{\mathbb{C}}}(i\eta + it\xi - \frac{t}{2}[\eta, \xi]) =$$
$$\exp_{H^{\mathbb{C}}}(-\frac{t}{2}[\eta, \xi]) \exp_{H^{\mathbb{C}}}(i(\eta + t\xi))$$

and using H-invariance one more time we conclude that if $U = H \cdot \exp_{H^{\mathbb{C}}}(iD)$, with D a domain in \mathfrak{h} , is Stein then D is convex in the usual affine sense.

Since $P(\Omega_H) = H \cdot exp_{H^{\mathbb{C}}}(i\tilde{P}(O_0))$ and as a consequence of Lemma 4.2 the domain $\tilde{P}(O_0)$ is not convex, then $P(\Omega_H)$ is not Stein. Now $H^{\mathbb{C}}$ is Stein

and $P|_{\Omega_H}$ is locally biholomorphic, therefore by [R] there exists a commutative diagram

$$\Omega_H \stackrel{j}{\longrightarrow} \hat{\Omega}_H
P \searrow \downarrow \hat{P}
H^{\mathbb{C}}$$

where $\hat{\Omega}_H$ is the envelope of holomorphy of Ω_H . Moreover H acts on $\hat{\Omega}_H$ and all maps are H-equivariant. Furthermore \hat{P} is injective by [CIT, Th. p. 543] and if Ω_H is holomorphically convex then j is surjective and consequently $\hat{\Omega}_H$ is biholomorphic to $P(\Omega_H)$, giving a contradiction. Hence Ω_H is not holomorphically convex.

Notice that $\hat{\Omega} \cong \hat{P}(\hat{\Omega})$ contains $P(\Omega_H) = H \cdot \exp_{H^{\mathbb{C}}}(i\tilde{P}(O_0))$ and the convex envelope of $\tilde{P}(O_0)$ is \mathfrak{h} , thus by the above arguments the envelope of holomorphy $\hat{\Omega}$ is biholomorphic to $H^{\mathbb{C}} \cong \mathbb{C}^3$.

Finally the maximal convex σ -invariant subdomain of $\tilde{P}(O_0)$ is $\{(A, B, C) \in \mathfrak{h} : A^2 + B^2 < 3\} = \tilde{P}(O_2)$, where $O_2 := \{(a, b, c) \in \mathfrak{h} : a^2 + b^2 < 3\frac{c^2}{\sinh^2(c)}\}$. One checks that $O_2 \subset O_1$, thus $\tilde{P}|_{O_2}$ is injective and $H \cdot O_2$ is biholomorphic to $H \cdot \exp_{H^{\mathbb{C}}}(i\tilde{P}(O_2))$. Moreover one has

$$\begin{split} \exp_{H^{\mathbb{C}}}(\,(A',B',C') + i(A,B,C)\,) \, = \\ \exp_{H}(A',B',C') \, \exp_{H^{\mathbb{C}}}(\,i(A,B,C - \frac{1}{2}(A'B - AB')\,)\,). \end{split}$$

It follows that

$$\exp_{H^{\mathbb{C}}}^{-1}(H \cdot \exp_{H^{\mathbb{C}}}(i\tilde{P}(O_2))) = \{ (\operatorname{Im} z_1)^2 + (\operatorname{Im} z_2)^2 < 3 \},\$$

where $(z_1, z_2, z_3) = (A' + iA, B' + iB, C' + iC)$ are natural complex coordinates of $h^{\mathbb{C}} \cong \mathbb{C}^3$ and this yields the statement.

Remark. Since \tilde{P} is injective on O_1 , then the H-invariant domain defined by O_1 is holomorphically separable. As a matter of fact one may show that $\tilde{P}(O_1) = \mathfrak{h} \setminus \{ (A, B, C) \in \mathfrak{h} : A^2 + B^2 \geq 3, C = 0 \}$ and analogous arguments as above show that such H-invariant domain is not holomorphically convex.

5. Generalized Heisenberg groups

Here we apply results of the previous section to generalized Heisenberg groups exhibiting additional examples of non-Stein maximal domains of existence for the adapted complex structure. We refer to [BTV] for the basic properties of generalized Heisenberg groups.

Let G be a generalized Heisenberg group with Lie algebra \mathfrak{g} , consider its abelian subalgebra $\mathfrak{z} := [\mathfrak{g}, \mathfrak{g}]$ and the subspace \mathfrak{v} orthogonal to \mathfrak{z} with respect to the G-invariant metric (\cdot, \cdot) of G. Then for all $V + Y \in \mathfrak{v} \oplus \mathfrak{z} = \mathfrak{g} \cong T_e G$ the unique geodesic tangent to V + Y at 0 can be explicitly given by $t \mapsto \exp_G \circ \varphi_G(t, V + Y)$ for a certain real-analytic map $\varphi_G : \mathbb{R} \times T_e G \to \mathfrak{g}$ (see [BTV, Th. p. 31]).

A straightforward computation shows that φ_G extends holomorphically on $\mathbb{C} \times T_e G$ to $\mathfrak{g}^{\mathbb{C}}$ and analogous arguments as in the previous section imply that $\Omega_G := G \cdot O_G \subset TG$ is the maximal the adapted complexification, where O_G is the connected component of

$$\{V+Y\in T_eG: \det(D\tilde{P}_G)_{V+Y}\neq 0\}$$

containing 0 and $\tilde{P}_G: T_eG \to \mathfrak{g}$ is given by

$$V + Y \mapsto l(|Y|)V + (1 + |V|^2 m(|Y|))Y$$
.

Here $|\cdot|$ denotes the norm induced by (\cdot,\cdot) and the real-analytic functions $l,m:\mathbb{R}\to\mathbb{R}$ are defined by

$$l(t) := \frac{\sinh(t)}{t}, \qquad m(t) := \frac{t - \sinh(t) \cosh(t)}{2t^3}.$$

Now for $V + Y \in \mathfrak{v} \oplus \mathfrak{z} = T_e G$ with $Y \neq 0$ and $U + X \in T_{V+Y} T_e G \cong T_e G$ one has

$$(D\tilde{P}_G)_{V+Y}(U+X) = \frac{\partial}{\partial t} \tilde{P}_G((V+tU)+(Y+tX))|_{t=0} = l'(|Y|)\frac{(Y,X)}{|Y|}V + l(|Y|)U$$
$$+ \left(2(V,U)m(|Y|) + |V|^2m'(|Y|)\frac{(Y,X)}{|Y|}\right)Y + (1+|V|^2m(|Y|))X.$$

Note that the equation is written according to the splitting $\mathfrak{v} \oplus \mathfrak{z}$ and since l(|Y|) never vanishes, the \mathfrak{v} -part vanishes if and only if

$$U = -\frac{(Y,X) l'(|Y|)}{|Y| l(|Y|)} V.$$

It follows that the central 3-part also vanishes if and only if

$$(1+|V|^2m(|Y|))X = |V|^2 \frac{(X,Y)}{|Y|} \left(2\frac{l'(|Y|)}{l(|Y|)}m(|Y|) - m'(|Y|)\right)Y.$$

In particular Y and X have to be proportional. Since both sides are homogeneous of degree 1 in X, then $(D\varphi)|_{(V+Y)}$ is singular if and only if

(5)
$$1 + |V|^2 m(|Y|) = |V|^2 |Y| \left(2 \frac{l'(|Y|)}{l(|Y|)} m(|Y|) - m'(|Y|) \right)$$

An analogous computation shows that $(D\tilde{P}_G)_{V+Y}$ has maximal rank if Y=0, thus equation (5) describes the singular locus of $D\tilde{P}_G$. It is remarkable that this identity is independent of the fine structure of the generalized Heisenberg group, e. g. of its dimension or the dimension of its centre. In particular if H is the 3-dimensional Heisenberg group considered in the previous section, equation (5) determines the border of $O_H = \Omega_H \cap T_e H$. Using this fact we are now going to show that for a generalized Heisenberg group G there exist many copies of Ω_H embedded as closed submanifolds in Ω_G .

Let G be a generalized Heisenberg group and choose non zero elements $\bar{V}_1 \in \mathfrak{v}$ and $\bar{Y} \in \mathfrak{z}$. Then there exists an element $\bar{V}_2 \in \mathfrak{v}$ such that the closed subgroup $\exp_G(\operatorname{span}\{\bar{V}_1,\bar{V}_2,\bar{Y}\})$ is a totally geodesically embedded 3-dimensional Heisenberg group (see [BTV, p. 30]). Denote by $I: H \to G$ such an embedding and note that since $\exp_{G^{\mathbb{C}}}: \mathfrak{g}^{\mathbb{C}} \to G^{\mathbb{C}}$ is a biholomorphism, then I extends to a holomorphic embedding $I^{\mathbb{C}}: H^{\mathbb{C}} \to G^{\mathbb{C}}$ of the universal complexification of H into the universal complexification of G such that the diagram

$$\mathfrak{h}^{\mathbb{C}} \xrightarrow{DI^{\mathbb{C}}} \mathfrak{g}^{\mathbb{C}}$$

$$\downarrow \exp_{H^{\mathbb{C}}} \qquad \downarrow \exp_{G^{\mathbb{C}}}$$

$$H^{\mathbb{C}} \xrightarrow{I^{\mathbb{C}}} G^{\mathbb{C}}$$

commutes. Now $I: H \to G$ is totally geodesic, thus $t \mapsto I \circ \exp_H \circ \varphi_H(t, v)$ is the unique geodesic of G tangent to DI(v) at 0 for all $v \in \mathfrak{h}$. Then

$$I \circ exp_H \circ \varphi_H(t, v) = exp_G \circ \varphi_G(t, DI(v))$$

and by the identity principle

(6)
$$I^{\mathbb{C}} \circ \exp_{H^{\mathbb{C}}} \circ \varphi_H(z, v) = \exp_{G^{\mathbb{C}}} \circ \varphi_G(z, DI(v))$$

for all $z \in \mathbb{C}$, since this holds for all $z \in \mathbb{R}$. Commutativity of the diagram

(7)
$$\begin{array}{ccc}
TH & \stackrel{DI}{\rightarrow} & TG \\
\downarrow P_H & & \downarrow P_G \\
H^{\mathbb{C}} & \stackrel{I^{\mathbb{C}}}{\rightarrow} & G^{\mathbb{C}}
\end{array}$$

follows. For this note that being I a group homomorphism then $DI: TH \to TG$ is H-equivariant, i.e., $DI(g_*w) = I(g)_*DI(w)$ for all $g \in H$ and $w \in TH$. In particular

$$P_G \circ DI(g_*v) \, = \, P_G(\, I(g)_*DI(v) \,) = I(g) \, \exp_{G^{\mathbb{C}}} \circ \varphi_G(i,DI(v))$$

for all $g \in H$ and $v \in T_eH$. On the other hand using equation (6) one obtains

$$I^{\mathbb{C}} \circ P_{H}(g_{*}v) = I^{\mathbb{C}}(g \exp_{H^{\mathbb{C}}} \circ \varphi_{H}(i,v)) = I^{\mathbb{C}}(g) I^{\mathbb{C}}(\exp_{H^{\mathbb{C}}} \circ \varphi_{H}(i,v)) =$$
$$I(g) \exp_{G^{\mathbb{C}}} \circ \varphi_{G}(i,DI(v)),$$

showing that the above diagram is commutative.

From the equivariance of DI it follows that $DI(\Omega_H) = DI(H \cdot O_H) = I(H) \cdot DI(O_H)$ and since DI is isometric and the border of O_H is defined by equation (5) which also describes the singular locus of \tilde{P}_G one has

$$DI(O_H) \subset O_G \cap \operatorname{span}\{\bar{V}_1, \bar{V}_2, \bar{Y}\}, \qquad DI(\partial O_H) \subset \partial O_G.$$

Thus $DI(\Omega_H) \cong I(H) \times DI(O_H)$ is closed in $\Omega_G \cong G \times O_G$.

Furthermore DI is injective and P_H , P_G are locally biholomorphic where the adapted complex structure is defined (cf. Theorem 3.2), thus diagram (7) shows that $DI(\Omega_H) \cong \Omega_H$ is a closed complex submanifold of Ω_G . Finally by Proposition (4.3) the domain Ω_H is neither holomorphically separable, nor holomorphically convex, thus one has

Proposition 5.1. Let G be a generalized Heisenberg group. Then the maximal adapted complexification Ω_G is neither holomorphically separable, nor holomorphically convex.

REFERENCES

- [A] AGUILAR, R. Symplectic reduction and the homogeneous complex Monge-Ampère equation Ann. Global Anal. Geom. 19 (2001), no. 4, 327–353
- [AG] AKHIEZER, D.-N.; GINDIKIN, S.-G. On Stein extension of real symmetric spaces Math. Ann. 286 (1990), 1–12
- [B] Burns, D. On the uniqueness and characterization of Grauert tubes Complex analysis and geometry (Trento, 1993), 119–133, Lecture Notes in Pure and Applied Math. 173
- [BHH] Burns, D.; Halverscheid, S.; Hind, R. The Geometry of Grauert Tubes and Complexification of Symmetric Spaces Preprint, 2001, math.CV/0109186
- [BTV] BERNDT, J.; TRICERRI, F.; VANHECKE, L. Generalized Heisenberg Groups and Damek-Ricci Harmonic Spaces LNM 1598, Springer-Verlag 1995
- [CIT] CASADIO-TARABUSI, E.; IANNUZZI, A.; TRAPANI, S. Globalizations, fiber bundles and envelopes of holomorphy Math. Z. 233 (2000), 535–551
- [GS] Guillemin, V.; Stenzel, M. Grauert tubes and the homogeneous Monge-Ampère equation J. Differential Geom. 34 (1991), no. 2, 561–570 (first part); J. Differential Geom. 35 (1992), no. 3, 627–641 (second part)
- [He] Heinzner, P. Equivariant holomorphic extensions of real analytic manifolds Bull. Soc. Math. France 121 (1993), 445–463
- [Ha] HALVERSCHEID, S. On Maximal Domains of Definition of Adapted Complex Structures for Symmetric Spaces of Non-compact Type Thesis, Ruhr-Universität Bochum, 2001
- [Ho] Hochschild, G. The structure of Lie groups Holden-Day, San Francisco, 1965
- [LS] LEMPERT, L.; SZŐKE, R. Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundles of Riemannian manifolds Math. Ann. 290 (1991), 689–712
- [L] LOEB, J. J. Pseudo-convexité des ouverts invariants et convexité géodésique de certains espaces symétriques Séminaire d'Analyse (P. Lelong, P. Dolbeault, H. Skoda), Lecture Notes, Springer-Verlag (1983-84)
- [R] Rossi, H. On envelopes of holomorphy Communications on pure and applied mathematics 16 (1963), 9–17
- [PW] Patrizio, G.; Wong, P.-M. Stein manifolds with compact symmetric center Math. Ann. 289 (1991), no. 4, 355–382
- [Sn] Snow, D-M. Reductive Group Actions on Stein Spaces Math. Ann. 259 (1982), 79–97
- [Sz1] Szőke, R. Complex structures on tangent bundles of Riemannian manifolds Math. Ann. 291 (1991), 409–428
- [Sz2] Szőke, R. Adapted complex structures and Riemannian homogeneous spaces Complex analysis and applications (Warsaw, 1997) Ann. Polon. Math. 70 (1998), 215–220.

Stefan Halverscheid

Math. Fakultät, Ruhr-Universität Bochum

Universitätsstr. 150, D-44780 Bochum, Germany

E- mail: sth@cplx.ruhr-uni-bochum.de

Andrea Iannuzzi

Dip. di Matematica, Università di Bologna

P. zza di Porta S. Donato 5, I-40126 Bologna, Italy

E-mail: iannuzzi@dm.unibo.it