

# Bases of the contact-order filtration of derivations of Coxeter arrangements

HIROAKI TERAO \*

*Tokyo Metropolitan University, Mathematics Department  
Minami-Ohsawa, Hachioji, Tokyo 192-0397, Japan*

## Abstract

In [6], we constructed a basis for the contact-order filtration of the module of derivations on the orbit space of a finite real reflection group acting on an  $\ell$ -dimensional Euclidean space. Recently M. Yoshinaga constructed another basis for the contact-order filtration in [8]. In this note we give an explicit formula relating Yoshinaga's basis to the basis given in [6]. The two bases turn out to be equal (up to a constant matrix). *Mathematics Subject Classification (2000): 32S22*

## 1 The setup and the main result

Let  $V$  be an  $\ell$ -dimensional Euclidean vector space with inner product  $I$ . Then its dual space  $V^*$  is equipped with the inner product  $I^*$  which is induced by  $I$ . Let  $S$  be the symmetric algebra of  $V^*$  over  $\mathbb{R}$ . Identify  $S$  with the algebra of polynomial functions on  $V$ . Let  $\text{Der}_S$  be the  $S$ -module of  $\mathbb{R}$ -linear derivations of  $S$ . When  $X_1, \dots, X_\ell$  denote a basis for  $V^*$ , the partial derivations  $\partial_i := \partial/\partial X_i$  with respect to  $X_i$  ( $1 \leq i \leq \ell$ ) naturally form a basis for  $\text{Der}_S$  over  $S$ . Let  $K$  be the field of quotients of  $S$  and  $\text{Der}_K$  be the  $K$ -vector space of  $\mathbb{R}$ -linear derivations of  $K$ . Then the partial derivations  $\partial_i$  ( $1 \leq i \leq \ell$ ) naturally form a basis for  $\text{Der}_K$  over  $K$ .

Let  $W$  be a finite irreducible orthogonal reflection group (a **Coxeter group**) acting on  $V$ . The Coxeter group  $W$  naturally acts on  $V^*$ ,  $S$  and  $\text{Der}_S$ . The  $W$ -invariant subring of  $S$  is denoted by  $R$ . Then it is classically known [1, V.5.3, Theorem 3] that there exist algebraically independent homogeneous polynomials  $P_1, \dots, P_\ell \in R$  with  $\deg P_1 \leq \dots \leq \deg P_\ell$ , which are called **basic invariants**,

---

\*partially supported by the Grant-in-aid for scientific research (No. 14340018 and 13874005), the Ministry of Education, Sports, Science and Technology, Japan

such that  $R = \mathbb{R}[P_1, \dots, P_\ell]$ . The **primitive derivation**  $D \in \text{Der}_R$  is characterized by

$$DP_i = \begin{cases} 1 & \text{for } i = \ell, \\ 0 & \text{otherwise.} \end{cases}$$

Write  $\mathbf{P} = (P_1, \dots, P_\ell)$ . Let  $J(\mathbf{P})$  denote the Jacobian matrix  $[\partial P_j / \partial X_i]_{ij}$ . Define  $\Delta := \det J(\mathbf{P})$ . Then  $\Delta$  is an anti-invariant [1, V.5.4] and  $\Delta^2 \in R$ . Let

$$I : \text{Der}_R \times \text{Der}_R \rightarrow \frac{1}{\Delta^2} R$$

be the symmetric  $R$ -bilinear form induced by  $I$ . Let

$$\begin{aligned} \nabla : \text{Der}_R \times \text{Der}_R &\longrightarrow \frac{1}{\Delta^2} \text{Der}_R \\ (X, Y) &\longmapsto \nabla_X Y \end{aligned}$$

be the **Levi-Civita connection** with respect to  $I$ . Since  $\frac{\partial}{\partial P_j} \in \frac{1}{\Delta} \text{Der}_S$  by Cramer's rule, one can embed  $\text{Der}_R$  into  $\text{Der}_K$ . Extend the Levi-Civita connection naturally to

$$\nabla : \text{Der}_K \times \text{Der}_K \longrightarrow \text{Der}_K.$$

Note  $\nabla_\xi(\eta) = \sum_i \xi(\eta(X_i)) \partial_i$  because each  $I(\partial_i, \partial_j)$  is constant.

In [6], we introduced derivations  $\xi_1^{(m)}, \dots, \xi_\ell^{(m)} \in \text{Der}_S$  for  $m \geq 0$  by

$$(\xi_1^{(m)}, \dots, \xi_\ell^{(m)}) := \begin{cases} (\frac{\partial}{\partial P_1}, \dots, \frac{\partial}{\partial P_\ell}) J(\mathbf{P})^T A J(D^k[\mathbf{X}])^{-1} & \text{if } m = 2k, \\ (\frac{\partial}{\partial P_1}, \dots, \frac{\partial}{\partial P_\ell}) J(\mathbf{P})^T A J(D^k[\mathbf{X}])^{-1} J(\mathbf{P}) & \text{if } m = 2k + 1. \end{cases}$$

Here  $A := [I^*(X_i, X_j)]_{ij}$  and  $J(D^k[\mathbf{X}]) := [\partial(D^k(X_j) / \partial X_i)]_{ij}$  ( $D^k := D \circ \dots \circ D$  ( $k$  times)). Let  $\mathcal{A}$  be the Coxeter arrangement determined by the Coxeter group  $W$ :  $\mathcal{A}$  is the set of reflecting hyperplanes. Choose for each hyperplane  $H \in \mathcal{A}$  a linear form  $\alpha_H \in V^*$  such that  $H = \ker(\alpha_H)$ . The derivations  $\xi_1^{(m)}, \dots, \xi_\ell^{(m)} \in \text{Der}_S$  were constructed in [6] so that they may form a basis for the  $S$ -module [10]

$$D^{(m)}(\mathcal{A}) := \{\theta \in \text{Der}_S \mid \theta(\alpha_H) \in S\alpha_H^m \text{ for any } H \in \mathcal{A}\}$$

for each nonnegative integer  $m$ . The filtration

$$\text{Der}_S = D^{(0)}(\mathcal{A}) \supset D^{(1)}(\mathcal{A}) \supset D^{(2)}(\mathcal{A}) \supset \dots$$

of  $\text{Der}_S$  is called the **contact-order filtration**. Define

$$\mathcal{H}^{(k)} = D^{(2k-1)}(\mathcal{A}) \cap \text{Der}_R$$

for each  $k \geq 1$ . Let  $\mathcal{H}^{(0)} = \text{Der}_R$ . Then we have a filtration

$$\text{Der}_R = \mathcal{H}^{(0)} \supset \mathcal{H}^{(1)} \supset \mathcal{H}^{(2)} \supset \dots$$

of  $\text{Der}_R$ . This filtration is known [7] to be equal to the **Hodge filtration** introduced by K. Saito [4]. The derivations  $\xi_1^{(2k-1)}, \dots, \xi_\ell^{(2k-1)} \in \text{Der}_R$  form an  $R$ -basis for  $\mathcal{H}^{(k)}$  for  $k \geq 1$ . Therefore the contact-order filtration, restricted to  $\text{Der}_R$ , gives the Hodge filtration. Define

$$T := \{f \in R \mid Df = 0\} = \mathbb{R}[P_1, \dots, P_{\ell-1}].$$

Then the covariant derivative  $\nabla_D : \text{Der}_K \rightarrow \text{Der}_K$  is  $T$ -linear. The Hodge filtration was originally defined so that the  $T$ -linear map

$$\nabla_D^k := \nabla_D \circ \dots \circ \nabla_D \text{ (} k \text{ times)} : \mathcal{H}^{(k)} \rightarrow \mathcal{H}^{(0)} = \text{Der}_R$$

is bijective [4]. Thus we may define  $\nabla_D^{-k} \xi \in \mathcal{H}^{(k)}$  for any  $\xi \in \text{Der}_R$  and  $k \geq 0$ . In [8] M. Yoshinaga proved the following

**Theorem 1.1.** ([8, Theorem 6]) *Let  $k \geq 0$  and*

$$E := \sum_i X_i \partial_i = \sum_i (\deg P_i)^{-1} P_i (\partial / \partial P_i)$$

*be the Euler derivation. Suppose that  $\xi_1, \dots, \xi_\ell$  are a basis for  $\mathcal{D}^{(1)}(\mathcal{A})$ . Then*

(1) *the derivations  $\nabla_{\xi_1} \nabla_D^{-k} E, \dots, \nabla_{\xi_\ell} \nabla_D^{-k} E$  form a basis for  $\mathcal{D}^{(2k-1)}(\mathcal{A})$  over  $S$ , and*

(2) *the derivations  $\nabla_{\partial_1} \nabla_D^{-k} E, \dots, \nabla_{\partial_\ell} \nabla_D^{-k} E$  form a basis for  $\mathcal{D}^{(2k)}(\mathcal{A})$  over  $S$ .*

It thus seems natural to ask how the basis above constructed by Yoshinaga is related to the basis  $\xi_1^{(m)}, \dots, \xi_\ell^{(m)}$  given in [6]. The following theorem answers this question:

**Theorem 1.2.** *Let  $k \geq 0$ . Then*

$$(\xi_1^{(2k+1)}, \dots, \xi_\ell^{(2k+1)}) = (-1)^k (\nabla_{\xi_1^{(1)}} \nabla_D^{-k} E, \dots, \nabla_{\xi_\ell^{(1)}} \nabla_D^{-k} E)$$

and

$$(\xi_1^{(2k)}, \dots, \xi_\ell^{(2k)}) = (-1)^k (\nabla_{\partial_1} \nabla_D^{-k} E, \dots, \nabla_{\partial_\ell} \nabla_D^{-k} E)A.$$

So the two bases turn out to be equal up to a constant matrix.

The significance of Theorem 1.2 is as follows: Recently M. Yoshinaga [9] affirmatively settled the Edelman-Reiner conjecture which asserts that the cones of the extended Shi/Catalan arrangements are free. In other words, there exist basic derivations for each extended Shi/Catalan arrangement. However, a formula for basic derivations is still unknown. Since the derivations  $\xi_i^{(m)}$  ( $i = 1, \dots, \ell$ ) are the principal (=highest degree) parts of basic derivations, Theorem 1.2 can be interpreted as a differential-geometric formula for the ‘‘principal parts.’’ So it may suggest the existence of a differential-geometric formula for the *whole* basic derivations, including the ‘‘non-principal part.’’ One may also regard Theorem 1.2 as a very explicit algebraic description of the derivations in the right hand side. They are, as mentioned above, when  $m$  is odd, bases for the Hodge filtration which is the key to define the flat structure on the orbit space  $V/W$  in [5]. The flat structure is called the *Frobenius manifold structure* from the view point of topological field theory [2].

## 2 Proof

We will prove Theorem 1.2 in this section. First we show the following

**Lemma 2.1.** *For  $k \geq 1$  and  $\xi \in \text{Der}_K$ , we have*

$$\nabla_D^k \circ \nabla_\xi - \nabla_\xi \circ \nabla_D^k = k \nabla_D^{k-1} \circ \nabla_{[D, \xi]}.$$

*Proof.* We use an induction on  $k$ . When  $k = 1$ , the lemma asserts

$$\nabla_D \circ \nabla_\xi - \nabla_\xi \circ \nabla_D = \nabla_{[D, \xi]},$$

which is the integrable property of the Levi-Civita connection  $\nabla$ . Let  $k > 1$ . We have

$$\begin{aligned} \nabla_D^k \circ \nabla_\xi &= \nabla_D^{k-1} \circ (\nabla_\xi \circ \nabla_D + \nabla_{[D, \xi]}) \\ &= (\nabla_D^{k-1} \circ \nabla_\xi) \circ \nabla_D + \nabla_D^{k-1} \circ \nabla_{[D, \xi]} \\ &= (\nabla_\xi \circ \nabla_D^{k-1} + (k-1)\nabla_D^{k-2} \circ \nabla_{[D, \xi]}) \circ \nabla_D + \nabla_D^{k-1} \circ \nabla_{[D, \xi]} \\ &= \nabla_\xi \circ \nabla_D^k + (k-1)\nabla_D^{k-2} \circ \nabla_{[D, \xi]} \circ \nabla_D + \nabla_D^{k-1} \circ \nabla_{[D, \xi]} \end{aligned}$$

by using the induction assumption. Since  $[D, [D, \xi]] = 0$ , we obtain

$$\nabla_D \circ \nabla_{[D, \xi]} = \nabla_{[D, \xi]} \circ \nabla_D.$$

This implies

$$\nabla_D^k \circ \nabla_\xi = \nabla_\xi \circ \nabla_D^k + k \nabla_D^{k-1} \circ \nabla_{[D, \xi]}.$$

□

Recall the  $W$ -invariant inner product  $I^* : V^* \times V^* \rightarrow \mathbb{R}$ . Let  $\Omega_R^1$  denote the  $R$ -module of Kähler differentials. Let

$$I^* : \Omega_R^1 \times \Omega_R^1 \rightarrow R$$

be the symmetric  $R$ -bilinear form induced by  $I^*$ . Let

$$G := [I^*(dP_i, dP_j)]_{ij} = J(\mathbf{P})^T A J(\mathbf{P}).$$

Define

$$B^{(k)} := -J(\mathbf{P})^T A J(D^k[\mathbf{X}]) J(D^{k-1}[\mathbf{X}])^{-1} J(\mathbf{P})$$

for  $k \geq 1$  as in [6]. Then we have

- Lemma 2.2.** (1) *Every entry of  $B^{(k)}$  lies in  $T$ ;  $D[B^{(k)}] = 0$ ,*  
(2)  *$\det B^{(k)} \in \mathbb{R}^*$ ,*  
(3)  *$D[G] = B^{(1)} + (B^{(1)})^T$ ,*  
(4)  *$B^{(k+1)} = B^{(1)} + kD[G]$ .*

*Proof.* By [6, Lemmas 3.3 - 3.6, Remark 3.7].

□

The following proposition is one of the main results in [7]:

**Proposition 2.3.** For  $k \geq 1$ ,

$$(\nabla_D^k \xi_1^{(2k-1)}, \dots, \nabla_D^k \xi_\ell^{(2k-1)}) = (-1)^{k-1} \left( \frac{\partial}{\partial P_1}, \dots, \frac{\partial}{\partial P_\ell} \right) B^{(k)}.$$

In the rest of this note, let  $(\xi_1, \dots, \xi_\ell)$  denote  $(\xi_1^{(1)}, \dots, \xi_\ell^{(1)})$  for simplicity. Note

$$(\xi_1, \dots, \xi_\ell) = \left( \frac{\partial}{\partial P_1}, \dots, \frac{\partial}{\partial P_\ell} \right) G.$$

**Lemma 2.4.**

$$([D, \xi_1], \dots, [D, \xi_\ell]) = \left( \frac{\partial}{\partial P_1}, \dots, \frac{\partial}{\partial P_\ell} \right) D[G] = (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell) (B^{(1)})^{-1} D[G].$$

*Proof.* The first equality easily follows from

$$[D, \xi_j](P_i) = D \circ \xi_j(P_i) - \xi_j \circ D(P_i) = D \circ \xi_j(P_i) = D[I^*(dP_i, dP_j)].$$

For the second equality, apply Proposition 2.3 when  $k = 1$ .  $\square$

**Lemma 2.5.** For any  $\eta \in \text{Der}_K$ ,

$$(\nabla_{\xi_1} \eta, \dots, \nabla_{\xi_\ell} \eta) = (\nabla_{\partial_1} \eta, \dots, \nabla_{\partial_\ell} \eta) AJ(\mathbf{P}).$$

*Proof.* Since both sides are additive with respect to  $\eta$ , we may assume  $\eta = f\partial_j$  for some  $j$  and  $f \in K$ . Then we have

$$\begin{aligned} (\nabla_{\xi_1} \eta, \dots, \nabla_{\xi_\ell} \eta) &= (\nabla_{\xi_1}(f\partial_j), \dots, \nabla_{\xi_\ell}(f\partial_j)) \\ &= (\xi_1(f)\partial_j, \dots, \xi_\ell(f)\partial_j) \\ &= ((\partial_1 f)\partial_j, \dots, (\partial_\ell f)\partial_j) AJ(\mathbf{P}) \\ &= (\nabla_{\partial_1}(f\partial_j), \dots, \nabla_{\partial_\ell}(f\partial_j)) AJ(\mathbf{P}) \\ &= (\nabla_{\partial_1} \eta, \dots, \nabla_{\partial_\ell} \eta) AJ(\mathbf{P}). \end{aligned}$$

$\square$

**Proof of Theorem 1.2.** When  $k = 0$ , the formulas are obviously true. Let  $k \geq 1$ . Let  $\xi \in \{\xi_1, \dots, \xi_\ell\}$ . Apply the identity in Lemma 2.1 to  $\nabla_D^{-k} E$  to get

$$(\nabla_D^k \circ \nabla_\xi)(\nabla_D^{-k} E) - \nabla_\xi E = k(\nabla_D^{k-1} \circ \nabla_{[D, \xi]})(\nabla_D^{-k} E) = k \nabla_D^{-1}(\nabla_{[D, \xi]} E).$$

Since  $\nabla_\eta E = \eta$  for any  $\eta \in \text{Der}_K$ , one obtains

$$\nabla_D^k(\nabla_\xi \nabla_D^{-k} E) - \xi = k \nabla_D^{-1}[D, \xi]$$

and thus

$$\nabla_D^{k+1}(\nabla_\xi \nabla_D^{-k} E) = \nabla_D \xi + k [D, \xi]$$

On the other hand, Lemma 2.4 asserts

$$([D, \xi_1], \dots, [D, \xi_\ell]) = (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell)(B^{(1)})^{-1} D[G].$$

So we have

$$\begin{aligned} & (\nabla_D^{k+1}(\nabla_{\xi_1} \nabla_D^{-k} E), \dots, \nabla_D^{k+1}(\nabla_{\xi_\ell} \nabla_D^{-k} E)) \\ &= (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell) + k (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell)(B^{(1)})^{-1} D[G] \\ &= (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell)(B^{(1)})^{-1} (B^{(1)} + kD[G]) \\ &= (\nabla_D \xi_1, \dots, \nabla_D \xi_\ell)(B^{(1)})^{-1} B^{(k+1)} \\ &= (\partial/\partial P_1, \dots, \partial/\partial P_\ell) B^{(k+1)} \\ &= (-1)^k (\nabla_D^{k+1} \xi_1^{(2k+1)}, \dots, \nabla_D^{k+1} \xi_\ell^{(2k+1)}) \end{aligned}$$

by Lemma 2.2 and Proposition 2.3. This proves the first formula.

For the second formula, compute

$$(\nabla_{\xi_1} \nabla_D^{-k} E, \dots, \nabla_{\xi_\ell} \nabla_D^{-k} E) = (\nabla_{\partial_1} \nabla_D^{-k} E, \dots, \nabla_{\partial_\ell} \nabla_D^{-k} E) A J(\mathbf{P})$$

by Lemma 2.5. Apply the first formula and we get

$$\begin{aligned} (\xi_1^{(2k)}, \dots, \xi_\ell^{(2k)}) J(\mathbf{P}) &= (\xi_1^{(2k+1)}, \dots, \xi_\ell^{(2k+1)}) \\ &= (-1)^k (\nabla_{\xi_1} \nabla_D^{-k} E, \dots, \nabla_{\xi_\ell} \nabla_D^{-k} E) \\ &= (-1)^k (\nabla_{\partial_1} \nabla_D^{-k} E, \dots, \nabla_{\partial_\ell} \nabla_D^{-k} E) A J(\mathbf{P}). \end{aligned}$$

This completes the proof of Theorem 1.2. □

## References

- [1] Bourbaki, N.: *Groupes et Algèbres de Lie*. Chapitres 4,5 et 6, Hermann, Paris 1968
- [2] Dubrovin, B.: Geometry of 2D topological field theories. In: *“Integrable systems and quantum groups”* (ed. Francaviglia, M., Greco, S.), Lectures at C.I.M.E., 1993, LNM **1620**, Springer, Berlin-Heidelberg-New York, 1996, pp. 120–348
- [3] Orlik, P., Terao, H.: *Arrangements of Hyperplanes*. Grundlehren der Math. Wiss. **300**, Springer Verlag, 1992
- [4] Saito, K.: On a linear structure of a quotient variety by a finite reflexion group. RIMS Kyoto preprint **288**, 1979 = Publ. Res. Inst. Math. Sci. **29** (1993) 535–579
- [5] Saito, K.: Finite reflection groups and related geometry (A motivation to the period mapping for primitive forms). preprint, 2000

- [6] Terao, H.: Multiderivations of Coxeter arrangements. *Inventiones math.*, **148** (2002) 659–674
- [7] Terao, H.: The Hodge filtration and the contact-order filtration of derivations of Coxeter arrangements. preprint 2002 (math.CO/0205058)
- [8] Yoshinaga, M.: The primitive derivation and freeness of multi-Coxeter arrangements. *Proc. Japan Acad. Ser. A Math. Sci.* **78** (2002), no. 7, 116–119
- [9] Yoshinaga, M.: Characterization of a free arrangement and conjecture of Edelman and Reiner. to appear in *Inventiones math.*
- [10] Ziegler, G. M.: Multiarrangements of hyperplanes and their freeness. In: *Singularities*. Contemporary Math. **90**, Amer. Math. Soc., 1989, pp. 345–359