Twisted K-Theory and Loop G roups

Daniel S.Freed

D epartm ent of M athem atics U niversity of Texas at A ustin

June 23, 2002

A bstract. Twisted K -theory has received much attention recently in both mathematics and physics. W e describe som emodels of twisted K -theory, both topological and geometric. Then we state a theorem which relates representations of loop groups to twisted equivariant K -theory. This is joint work with M ichael Hopkins and Constantin Teleman.

The loop group of a compact Lie group G is the space of sm ooth m aps S¹! G with multiplication de ned pointwise. Loop groups have been around in topology for quite some time [Bo], and in the 1980s were extensively studied from the point of view of representation theory [Ka], [PS]. In part this was driven by the relationship to conform alleld theory. The interesting representations of loop groups are projective, and with xed projective cocycle there is a nite number of irreducible representations up to isom orphism. Considerations from conform alleld theory [V] led to a ring structure on the abelian group R (G) they generate, at least for transgressed twistings. This is the Verlinde ring. For G simply connected R (G) is a quotient of the representation ring of G, but that is not true in general. At about this time W itten [W] introduced a three-dimensional topological quantum eld theory in which the Verlinde ring plays an important role. Eventually it was understood that the fundamental object in that theory is a 'm odular tensor category" whose G rothendieck group is the Verlinde ring. Typically it is a category of representations of a loop group or quantum group.

For the special case of a nite group G the topological eld theory is specied by a certain cocycle on G and the category can be calculated explicitly [F1]. We identied it as a category of representations of a Hopf algebra constructed from G, thus directly linking the Chem-Sim ons lagrangian and quantum groups. Only recently did we realize that this category has a description in terms of twisted equivariant K -theory, and it was natural to guess that the Verlinde ring for arbitrary G has a similar description. Ongoing joint work with M ichael Hopkins and Constantin

The author is supported by NSF grant DMS-0072675.

Telem an has con med this description. We can speculate further and hope that twisted K – theory provides a construction of the modular tensor category, and perhaps even more of the three-dimensional topological eld theory. In another direction the use of K - theory may shed light on Verlinde's formula for certain R iem ann-R och numbers. In any case our result to swell with other uses of K - theory in representation theory [CG], for example in the geometric Langlands program.

The physics motivation for the main theorem is discussed in [F2], [F3]. Here, in x3, we explain the statement of our result in mathematical terms; the proof will appear elsewhere. A s background we describe some concrete topological models of twisted K -theory in x1, and give a twisted version of the Chern-W eil construction in x2.

As mentioned above the work I am discussing is being carried out with M ichael Hopkins and C onstantin Teleman. I thank them for a most enjoyable collaboration.

x1 Twistings of K-theory

Let X be a reasonable compact space, say a nite CW complex. Then isomorphism classes of complex vector bundles over X form a sem igroup whose group completion is the K -theory group K⁰ (X). This basic idea was introduced by G rothendieck in the context of algebraic geom etry [BS], and was subsequently transported to topology by Atiyah and Hirzebruch [AH]. Vector bundles are local they can be cut and glued and in the topological realm this leads to a cohom ology theory. In particular, there are groups Kⁿ (X) de ned for n 2 Z. Historically, K -theory was the rst example of a generalized cohomology theory, and it retains the features of ordinary cohomology with one notable exception: the cohomology of a point is nontrivial in all even degrees, as determ ined by Bott periodicity. There are many nice spaces B which represent com plex K -theory in the sense that K⁰(X) is the set of hom otopy classes of maps from X to B. A particularly nice choice [A1, Appendix], [J] is the space B = Fred (H) of Fredholm operators on a separable complex Hilbert space H. Thus a map T:X ! Fred (H) determ ines a K-theory class on X. It is convenient to generalize and allow the Hilbert space H to vary as follows. A Fredholm complex¹ is a graded Hilbert space bundle $E = E^0 E^1$! X together with a berwise Fredholm map E^0 ! E^1 , and it also represents an element of K 0 (X). Another innovation was the introduction by A tiyah and Segal [S2] of the equivariant K -theory groups K $_{\rm G}$ (X) for a compact space X which carries the action of a compact Lie group G. The basic objects are G-equivariant vector bundles E! X, and K_{G}^{0} (X) is the group completion of the set of equivalence classes. For example, if X is a point then the equivariant K -theory is the representation ring K $_{\rm G}\,$ of the compact Lie group G; in general, K_G (X) is a K_G -m odule.

As a rst example of twisted K-theory we consider twisted versions of K_G. A twisting is a

 $^{^{1}}$ O ne [S1] can allow more general topological vector spaces and complexes which are nonzero in degrees other than 0 and 1.

central extension

where T C is the circle group of unit norm complex numbers. Then the twisted representation $\ring" K_G$ is the group completion of equivalence classes of complex representations of G on which the central T acts by scalar multiplication. Twisted K -theory is not a ring, but rather K_G is a K_G - m odule. A map of twistings, i.e., an isom orphism of central extensions, determ ines an isom orphism of twisted K -groups. So twisted K -theory is determ ined up to noncanonical isom orphism by the equivalence class of the twisting. For example, $K_{SO(3)} = Z$ [s] is the polynom ial ring on a single generator, the 3-dimensional de ning representation. Up to equivalence there is a single nontrivial central extension

= f1 ! T ! U(2) ! SO(3) ! 1g

which is induced by the inclusion Z = 2Z ,! T from the extension

1 ! Z=2Z ! SU(2) ! SO(3) ! 1:

V intual representations of U (2) on which the central element acts naturally correspond 1:1 with virtual representations of SU (2) on which the central element acts as 1. Now $K_{SU(2)} = Z[t]$, where t is the dening 2-dimensional representation, and we identify $K_{SO(3)} = K_{SU(2)}$ as the subgroup of odd polynom ials in t; the $K_{SO(3)} = Z[s]$ -m odule structure is structure is structure is structure is a free m odule of rank one.

M one generally, on a G-space X a \cocycle" for the equivariant cohom ology group $H_G^3(X;Z)$ de ness a twisted K -theory group $K_G(X)$ which is a module over $K_G^0(X)$. A better point of view is that is a cocycle, or geometric representative, of a class in $H_G^1(X;A_G)$, where A_G is a group of autom orphism s of equivariant K -theory. W e will not try to make \autom orphism of equivariant K -theory. W e will not try to make \autom orphism of equivariant K -theory " precise here, but content ourselves with a concrete model, rst in the nonequivariant case. Take Fred (H) to be the classifying space of K -theory. Then the group A = PGL(H) acts as autom orphism s by conjugation. Since A = GL(H) is contractible [K], the quotient A = A = C is hom otopy equivalent to CP^1 , and so $H^1(X;A) = H^3(X;Z)$. A twisting can be taken to be a principal A-bundle :P ! X. The action of A on Fred (H) de ness an associated bundle Fred (H)_P ! X, and the twisted K -group K (X) is the group of hom otopy classes of sections of this bundle [A 2]. A section is an A-equivariant m ap T :P ! Fred (H), and as before it is convenient to let the H ilbert space vary. Thus de ne a P -twisted Fredholm com plex to be an A-equivariant graded H ilbert space bundle E ! P, where the center C A acts by scalar multiplication, together with a berw ise A-equivariant Fredholm operator $E^0 \stackrel{\text{F}}{!} E^1$. Then (E ;T) represents an element of K (X).

It is perhaps unsettling that the model is in nite dimensional, but that is unavoidable unless the class of the twisting in H 3 (X ;Z) is torsion.

There are many other models of twistings and twisted K -theory. For example, gerbes are geometric representatives of elements in degree three integral cohom obgy; see [H], [B], [M] for example. In a Cech description we have a covering $X = {S \atop i} U_i$ of X by open sets and a complex line bundle L_{ij} ! $U_i \setminus U_j$ on double intersections. There is further cocycle data on triple intersections. In a model of K -theory on which line bundles act as automorphisms this can be used to de netwisted K -theory; see [BCM M S] for example. In fact, the \group" Z=2Z CP¹ of graded lines act as autom orphisms of K -theory, so there is a larger group of (equivalence classes of) twistings

(1)
$$H^{1}(X;Z=2Z CP^{1}) = H^{1}(X;Z=2Z) H^{3}(X;Z):$$

W e rem ark that there is a natural group structure on (1), but it is not the product. In topology these twisted versions of K -theory, at least for torsion twistings, were introduced by D onovan and K aroubi [D K], who also considered real versions. There is another view point and generalization of K -theory using C -algebras, and in that context twisted K -theory was discussed by R osenberg [R] for both torsion and nontorsion twistings. Twisted versions of K -theory have appeared recently in various parts of geom etry and index theory, for example in [LU], [A R], [To], [N T], [M M S].

A generalization of the previous m odel is useful. Here a twisting is a quartet = (G; ;G;P):

G topological group :G!Z=2Z hom om orphism term ed the grading G!G central extension by T P!X principal G-bundle

We require the existence of a hom on orphism $G \,!\, GL(H)$ which is the identity on the central T. Let $G_0 \,!\, G_0$ be the restriction of the central extension over the subgroup $G_0 = {}^1$ (0). Then the equivalence class of f is the obstruction in (1) to restricting/lifting P to a principal G_0 bundle. Our previous construction is the special case G = P G L(H) and trivial. An element of K (X) is represented by a G-equivariant Z=2Z-graded Fredholm com plex over P, where the action of G is compatible with the grading . If a compact Lie group G acts on X then it is easy to extend this to a model of equivariant twistings and equivariant K -theory.

As an illustration of a nontrivial H¹-twisting, consider X = pt and = 0 (2); ;0 (2) T;0 (2) with nontrivial grading. The representation ring of 0 (2) may be written

$$K_{O(2)} = Z[;]$$
 ((1); ² 1);

where is the one-dimensional sign representation and the standard two dimensional representation. Then the twisted K-group K₀₍₂₎ is a module over K₀₍₂₎ with a single generator t and the

relation t = t. There is a nontrivial odd twisted K -group $K_{0}^{+1}_{(2)}$ which is also a module with a single generator u; the relations are u = u and u = 0.

M any topological properties of K -theory, including exact and spectral sequences, have straightforward analogs in the twisted case. This is easiest to see from the hom otopy-theoretic view of cohom ology theories, and so applies to twisted cohom ology theories in general. C om putations are usually based on these sequences. A more specialized result is the completion theorem in equivariant K -theory [AS]; its generalization to the twisted case has some new features [FHT]. The Thom isom orphism theorem ts naturally into the twisted theory [DK]. Let V ! X be a real vector bundle of nite rank, which for convenience we suppose endowed with a metric. There is an associated twisting

$$(V) = O(n); ;Pinc(n);O(V);$$

where O(V)! X is the orthonormal frame bundle of V and : O(n)! Z=2Z is the nontrivial grading. The isomorphism class of (V) is the pair of Stiefel-W hitney classes $w_1(V)$; $W_3(V)$. Denote

(3)
$$K^{V}(X) = K^{\operatorname{rank}(V) + (V)}(X)$$

as the degree-shifted twisted K -theory. Then the Thom isomorphism theorem asserts that the naturalm ap

(4)
$$K^{q+V}(X) ! K^{q+2 \operatorname{rank}(V)}(V)$$

is an isomorphism. We remark that if X is a smooth compact manifold, then Poincare duality identies K^{TX} (X) with the K-homology group K₀(X). Also, we can de ne K^V (X) for virtual bundles V.

The Chem character maps twisted K -theory to a twisted version of real cohomology, as we explain next in the context of Chem-W eiltheory.

x2 A differential-geometric model

For simplicity we work in the nonequivariant context. Our thoughts here were stimulated by reading BCMMS].

Let a = pgl(H) denote the Lie algebra of $A^{\sim} = P G L (H)$ and a = gl(H) the Lie algebra of A = G L (H). They t into the exact sequence of Lie algebras

A linear map L:a! a is a splitting if the composition with a! a is the identity. A splitting L determ ines a closed right-invariant 2-form on A which represents the generator of H² (A;Z); its value on right-invariant vector elds; is

Let : P ! X be a principal A-bundle. W e now add two pieces of geom etric data:

(5) 2 ¹ (P;a) connection on P ! X L:P ! Hom (a;a) A -invariant m ap into splittings

Both are sections of a ne space bundles over X , so can be constructed using partitions of unity. A susual, de ne the curvature

$$F = d + \frac{1}{2} [^{ }]:$$

It is an a-valued 2-form on P . Introduce the scalar 2-form

=
$$dL() + \frac{1}{2}[L()^{L}()] L(F_{A}):$$

Then one can check that is transgressive. In other words, d = for a closed scalar 3-form 2³ (X). The de Rham cohom ology class of in H³ (X;R) represents the image in real cohom ology of the isomorphism class of the twisting P! X.

Now let $E^0 \stackrel{!}{!} E^1$ be a twisted Fredholm complex over P. Thus E^i ! P are A'-equivariant Hilbert space bundles, with C A' acting by scalar multiplication, and T is an A'-equivariant Fredholm map. The A' action determines an A'-invariant partial covariant derivative on E ! P along the bers of : P ! X. A gain we introduce di erential-geometric data:

r A - invariant extension of the partial covariant derivative

Such an extension is a section of an a ne space bundle over X, so can be constructed via a partition of unity. Introduce a form al parameter u of degree 2 and its inverse u¹ of degree 2, and so the graded ring of R [[u;u¹]]-valued di erential form s. (O ne can identify u as a generator of the K-theory of a point.) Following Q uillen [Q] de ne the A-invariant superconnection

$$D = \begin{array}{c} r^{0} & uT \\ T & r^{1} \end{array}$$

on E ! X. Its usual curvature $D^2 2 P$; End E R $[[u; u^1]]^2$ is an A-invariant form of total degree 2. However, it does not descend to the base X. Instead, one can check that the twisted curvature

$$F(E;T;r) = D^2$$
 id

is A-invariant and basic, so descends to an element of X; EndE R $[[u; u^1]]^2$. (Note that EndE ! P descends to a graded vector bundle on X, since the center of A acts trivially.) It does not, how ever, satisfy a B ianchi identity, since is not closed.

The Chern character form

(6)
$$ch(E;T;r) := Trexp(u F)$$

is an element in X; R [[u; u¹]]⁰ of total degree 0. Here we assume favorable circum stances in which the graded trace Tr is nite. For example, if the twisting class is torsion then we can take E ! P nite dimensional. Or, if the superconnection comes from a family of elliptic operators as in [Bi] then the graded trace exists. The Chern character form (6) is not closed in the usual sense, but rather

$$(d + u^{1}) ch (E; T; r) = 0:$$

The dimensional d+ u¹ on X; R [[u; u¹]] computes a twisted version of real cohomology which is the codom ain of the Chern character.

This construction works with little change for the more general twistings (2).

The di erential geom etric model we have outlined not only gives geom etric representatives of twisted topological K -theory classes, but also geom etric representatives of twisted di erential K - theory classes. Sim ilarly, the geom etric twistings (5) give geom etric models for di erential cohomology classes. See [HS] for foundations of (untwisted) di erential cohomology theories in general, and [L] for the basics di erential K -theory.

x3 Loop groups

Let G be a compact Lie group. The bop group LG of G is the space of smooth maps $S^1 \\! G$. There is a twisted description we use instead. Namely, let R $! S^1$ be a principal G-bundle and LG_R the group of gauge transform ations, i.e., the space of smooth sections of the bundle of groups $G_R \\! S^1$ associated to R. A trivialization of R gives an isomorphism $LG_R = LG$. Note that R is necessarily trivializable if G is connected, and in general the topological class of R is labeled by a conjugacy class in $_0G$. Let $G(R) \\= G_R$ is described in $\mathbb{P}S$, Ka, and some speci c further developments appear in $\mathbb{F}F$, W e]. One salient feature of loop groups is the existence of nontrivial central extensions

$$1 ! T ! LG_R ! LG_R ! 1:$$

As in (2) we may also consider gradings $: LG_R ! Z=2Z . We call the pair = (LG_R;) a graded central extension and denote it simply as LG_R. It has an invariant in$

(7)
$$H_{G}^{1} G(\mathbb{R}); Z = 2Z \qquad H_{G}^{3} G(\mathbb{R}); Z$$

as follows. Fix a basepoint s 2 S and consider the product

$$P = A_R \qquad R_s$$

of the space of connections A_R on R and the ber R_s at the basepoint. Then LG_R acts freely on P with quotient the holonom y m ap

Furtherm ore, the G action on G (R) by conjugation lifts to the G action on the R_s factor of P, where it commutes with the LG_R action. Thus we have an equivariant twisting

$$\sim = (LG_{R}; ; LG_{R}; P)$$

whose isom orphism class, called the level, lies in (7). (The basepoint is not necessary. It is more natural to replace (8) by A_R R and LG_R by the group of bundle autom orphism s which cover rotations in the base S^1 .)

As a warm -up to representations of loop groups, recall the Borel-W eil construction of representations of a compact connected Lie group G. Let T G be a maximal torus, and F = G=T the agmanifold. Then F admits G-invariant complex structures. Fix one. A character of T determ ines a holom orphic line bundle L ! F, and the standard construction takes the induced virtual representation of G to be $_{\rm q}(1)^{\rm q} {\rm H}^{\rm q}$ (F;L), which is the G-equivariant index of the @ operator. We can use the D irac operator instead of the @ operator. This has the advantage that no complex structure need be chosen, though it can be described in holom orphic terms as the -shift L ! L $K_{\rm F}^{1=2}$, where $K_{\rm F}^{1=2}$! F is a square root of the canonical bundle. M ore generally, if G is the centralizer of any subtorus of T, and F⁰ = G=Z the corresponding generalized ag manifold, then there is a D irac induction map

(9)
$$K_{Z} = K_{G} (F^{0}) ! K_{G} :$$

A representation of Z de nes a G -equivariant vector bundle on F 0 , and (9) is the equivariant index of the D irac operator with coe cients in this bundle.

A similar construction works for loop groups. Fix a conjugacy class C G(R). The group LG_R G acts transitively on hol¹ (C). Let Z_C denote the stabilizer at some point; it embeds isom orphically into either factor of LG_R G. Introduce the agmanifold $F_C = hol¹$ (C)=G; then the loop group LG_R acts transitively F_C with stabilizer the image of Z_C in LG_R . On the other hand, the image of Z_C in G is the centralizer of an element in C. The geometry of F_C is similar to that of nite dimensional agmanifolds [F4]. An important special case is C = flg and R ! S¹ trivial. Then the agmanifold is the \loop grassmannian" F = LG = G.

There is a D irac induction for loop groups which generalizes (9). For this we introduce spinors on LG_R and a canonical graded central extension. Consider the principal SO (g R)-bundle associated to R ! S^1 via the twisted adjoint hom on orphism

It is trivializable, and any trivialization induces an isomorphism LG_R ! LSO (N) for N = dim (G) + 1. There is a canonical graded central extension of LSO (N) from the spin representation, and it pulls back to the desired canonical graded central extension $LG_R^{R^{(G)}}$! LG_R . The spin representation itself de nes (projective) spinors on LG_R . For a conjugacy class C G (C) the embedding $i_C : Z_C$, LG_R induces a graded central extension $Z_C^{i_C R^{(G)}}$ which may be described instead by the real adjoint representation z_C , viewed as a Z_C -equivariant vector bundle over a point (see (3)). In this form it carries a degree shift, and so it is natural to also include a degree shift in $_R$ (G). We do not specify it precisely, but remark that its parity agrees with that of dim Z_C . If G is simple, connected, and simply connected and R ! S^1 is trivial, then $_R$ (G) has degree shift dim G and the H³ component of the level in (7) is the dual Coxeter number of G times a generator; the H¹ component vanishes.

For a xed graded central extension LG_R of LG_R there is a nite set of isom orphism classes of irreducible positive energy representations of LG_R on which the center acts by scalar multiplication. Let R (G) denote the abelian group generated by these equivalence classes. It is natural to extend this to a Z-graded group with m od 2 periodicity and possibly nontrivial groups in odd degrees. Now we describe D irac induction for loop groups. As in the rst m ap of (9) a representation of Z_C de nes an LG_R -equivariant vector bundle over the ag manifold F_C . However, we are interested in LG_R -equivariant vector bundles, so need to start with a representation of the central extension of Z_C de ned by i_C . Finally, spinors on the ag manifold F_C may be constructed from spinors on LG_R by \subtracting" spinors on the adjoint representation of Z_C , and this imposes an additional twisting. A ltogether, then, D irac induction is a map

(10)
$$K_{Z_{c}}^{i_{c} z_{c}} ! R_{Q}^{R} G(R) :$$

The adjoint shift $_{R}$ (G) on the right hand side means that we obtain representations of the ber product of LG_{R} ! LG_{R} with the inverse of LG_{R}^{R} (G) ! LG_{R} , including a degree shift. For connected, simply connected G it su ces to consider C = flg, since then (10) is surjective, but this is not true in general.

The inclusion $i_c : C , C (R)$ induces a pushforward in twisted K -theory (cf. (4)):

(11)
$$K_{Z_{C}}^{i_{C}} = K_{G}^{i_{C} \to TC TG j_{C}}$$
 (C) ! K_{G}^{c} G (R) :

The maps (10) and (11) give, for each conjugacy class C, a correspondence between certain representations of the loop group and a twisted K -theory group.Our main result is

Theorem 1. These correspondences induce an isom orphism of abelian groups

(12)
$$R = {}^{R} (G) G (R) ! K_{G} G (R) :$$

There is a transgression from H 4 (BG;Z) to levels (7) with trivial rst component. M ore generally, there is an extension

$$0 ! H^{4}(BG;Z) ! E^{4}(BG) ! H^{2}(BG;Z=2Z) ! 0;$$

and elements of E⁴ (BG) transgress to general levels. For transgressed levels there is a ring structure on both sides of (12). The multiplication on representations is the fusion product [V], [F], [I]; on twisted K -theory it is the pushforward by multiplication G G! G or equivalently the Pontrjagin product in K -hom ology.

Theorem 2. If the level of the twisting $\sim +$ (TG) is transgressed, then (12) is an isomorphism of rings.

Our proof reduces both sides of (12) to the statem ent for tori, where there is a direct argum ent.

References

- [AR] A.Adem, Y.Ruan, Twisted orbifold K-theory, math.AT/0107168.
- [A1] M.F.Atiyah, K-Theory, Benjamin, New York, 1967.
- [A2] M.F.Atiyah, K -theory past and present, math.KT/0012213.
- [AH] M.F.Atiyah, F.Hirzebruch, Vector bundles and hom ogeneous spaces, Proc. Symp. Pure M ath. 3 (1961), 7{38.

- [AS] M.F.Atiyah, G.B.Segal, Equivariant K -theory and completion, J.Di.Geom.3 (1969), 1{18.
- [Bi] J.M.Bismut, The Atiyah-Singer Index Theorem for families of Dirac operators: two heat equation proofs, Invent. math. 83 (1986), 91{151.
- [BS] A. Borel, J.-P. Serre, Le theorem e de Riem ann-Roch, Bull. Soc. M ath. France 86 (1958), 97{136.
- [Bo] R.Bott, The space of bops on a Lie group, Michigan Math. J.5 (1958), 35{61.
- BCMMS] P.Bouwknegt, A.L.Carey, V.M athai, M.K.Murray, D.Stevenson, Twisted K-theory and K-theory of bundle gerbes, hep-th/0106194.
- [B] J.-L.Brylinski, Loop spaces, characteristic classes and geom etric quantization, Birkhauser Boston Inc., Boston, MA, 1993.
- [CG] N.Chriss, V.G inzburg, Representation Theory and Complex Geometry, Birkhauser, Boston, 1997.
- [DK] P.Donovan, M.Karoubi, G raded B rauer groups and K -theory with local coe cients, Inst. Hautes Etudes Sci. Publ. M ath. 38 (1970), 5{25.
- [F] G.Faltings, A proof for the Verlinde form ula, J.Algebraic Geom. 3 (1994), 347{374.
- [FF] B.L.Fe gin and E.V.Frenkel, A ne Kac-M cody algebras and sem i-in nite agm anifolds, Comm.M ath.Phys.128 (1990), 161{189.
- [F1] D.S.Freed, Higher algebraic structures and quantization, Commun.Math.Phys.159 (1994), 343{398, hep-th/9212115.
- [F2] D.S.Freed, The Verlinde algebra is twisted equivariant K -theory, Turkish J.M ath. 25 (2001), 159{167, math.RT/0101038.
- [F3] D.S.Freed, K -theory in quantum eld theory, Current D evelopm ents in M athem atics 2001, math-ph/0206031.
- [F4] D.S.Freed, The geometry of bop groups, J.Di.Geo.28 (1988), 223{276.
- [H] N.J.Hitchin, Lectures on special Lagrangian submanifolds, W inter School on M irror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), Amer.Math.Soc., Providence, RI, 2001, pp. 151{182, math.DG/9907034.
- [HS] M.J.Hopkins, I.M. Singer, Quadratic functions in geometry, topology, and M-theory (in preparation).
- [J] K.Janich, Vektorraum bundel und der Raum der Fredholm Operatoren, Math. Annalen
 161 (1965), 129{42.
- [Ka] V.G.Kac, In nite dim ensional Lie algebras, Cambridge University Press, Cambridge, 1990.
- [K] N.H.Kuiper, The hom otopy type of the unitary group of Hilbert space, Topology 3 (1965), 19{30.

- [LU] E.Lupercio, B.Uribe, Gerbes over orbifolds and twisted K-theory, math.AT/0105039.
- [MMS] V.Mathai, R.B.Melrose, IM. Singer, The index of projective families of elliptic operators, math.DG/0206002.
- [M] M.K.Murray, Bundle gerbes, J.London Math. Soc. 54 (1996), 403{416.
- [NT] V.N istor, E.Troitsky, An index for gauge-invariant operators and the D ixm ier-D ouady invariant, math.KT/0201207.
- [PS] A. Pressley, G. Segal, Loop Groups, Oxford University Press, Oxford, 1986.
- [Q] D.Quillen, Superconnections and the Chem character, Topology 24 (1985), 89{95.
- [R] J.Rosenberg, Continuous trace algebras from the bundle theoretic point of view, Jour.
 Austr.Math.Soc.47 (1989), 368{381.
- [S1] G.B. Segal, Fredholm com plexes, Quart. J. M ath. Oxford 21 (1970), 385{402.
- [S2] G.B. Segal, Equivariant K -theory, Publ. M ath.Inst. H autes. Etudes Sci. 34 (1968), 129(151.
- [T] C. Telem an, Lie algebra cohom ology and the fusion rules, Comm. M ath. Phys. 173 (1995), 265{311.
- [[o] B.Toen, Notes on G-theory of Deligne-Mum ford stacks, math.AG/9912172.
- [V] E.Verlinde, Fusion rules and modular transformations in 2D conformal eld theory, Nuclear Phys. B 300 (1988), 360{376.
- [We] R.Wendt, Weyl's character formula for non-connected Lie groups and orbital theory for twisted a ne Lie algebras, J.Funct. Anal. 180 (2001), 31{65, math.RT/9909059.
- [W] E.W itten, Quantum eld theory and the Jones polynomial, Commun.Math.Phys. 121 (1989), 351{399.

D epartment of M athematics, University of Texas, Austin, TX 78712 E-m ailaddress: dafr@math.utexas.edu

[[]L] J.Lott, R=Z index theory, Comm.Anal.Geom.2 (1994), 279{311.